

$\mathcal{N} = 2$ resonant superalgebra for supergravity

Remigiusz Durka ^{*1} and Krzysztof M. Graczyk ^{†1}

¹*University of Wrocław, Institute for Theoretical Physics, pl. M. Borna 9, 50-204 Wrocław, Poland*

May 13, 2022

Abstract

We present new superalgebra for $\mathcal{N} = 2$ $D = 3, 4$ supergravity theory endowed with the $U(1)$ generator. The superalgebra is rooted in the so-called Soroka-Soroka algebra and spanned by the Lorentz J_{ab} and Lorentz-like Z_{ab} , translation P_a and T generators, as well as two supercharges Q_α^I . We construct a corresponding 3D Chern-Simons supergravity realization of the superalgebra and discuss its relevance.

1 Introduction

Supergravity incorporates the interchangeability of the fermionic and bosonic fields under the so-called supersymmetry. Consequently, the graviton (spin-2) comes with its superpartner gravitino (spin-3/2). A natural generalization is obtained in \mathcal{N} -extended supergravity model containing \mathcal{N} gravitinos. Obviously $\mathcal{N} = 0$ gives general relativity while $\mathcal{N} = 1$ corresponds to standard supergravity [1, 2, 3].

An effort has been made to construct the unified theory of gravity with other interactions, e.g., such as electromagnetism. In the latter, the unification is realized within an algebraic framework of supergravity. An example of such theory is $\mathcal{N} = 2$ supergravity [4]. In this model, the graviton (spin-2) and gravitino (spin-3/2) gauge action couples with another gravitino (spin-3/2) and spin-1 vector field. Note that the action is obtained from the supersymmetry group, which includes the central charges [5] and rotation of two gravitinos [6].

We distinguish two basic initial algebraic structures for supergravity construction, namely, Poincaré and AdS (Anti de-Sitter) algebras. In this paper we do not consider the Poincaré scenario [4, 7] but we focus on the AdS $\mathcal{N} = 2$ case [8, 9, 10]. The corresponding superalgebra in $D = 4$ with the Lorentz generator J , translation P , $U(1)$ generator T , and two supercharge Q^I generators is expressed as:

$$\begin{aligned}
 [J_{ab}, J_{cd}] &= \eta_{bc}J_{ad} - \eta_{ac}J_{bd} + \eta_{ad}J_{bc} - \eta_{bd}J_{ac}, \\
 [J_{ab}, P_b] &= \eta_{bc}P_a - \eta_{ac}P_b, \quad [P_a, P_b] = J_{ab}, \\
 [J_{ab}, Q_\alpha^I] &= \frac{1}{2}(\Gamma_{ab})_\alpha^\beta Q_\beta^I, \quad [P_a, Q_\alpha^I] = \frac{1}{2}(\Gamma_a)_\alpha^\beta Q_\beta^I, \quad [T, Q_\alpha^I] = \epsilon^{IJ}Q_\alpha^J, \\
 \{Q_\alpha^I, Q_\beta^J\} &= -\delta^{IJ} \left((\Gamma^a C)_{\alpha\beta} P_a - \frac{1}{2}(\Gamma^{ab} C)_{\alpha\beta} J_{ab} \right) + \epsilon^{IJ} C_{\alpha\beta} T.
 \end{aligned} \tag{1}$$

^{*}remigiusz.durka@uwr.edu.pl

[†]krzysztof.graczyk@uwr.edu.pl

By α, β we denote spinorial indices, whereas $a, b = 0, 1, 2, 3$ are the Lorentz indices, and $I = 1, 2$. By $\eta_{ab} = \text{diag}(-, +, +, +)$ we denote flat Minkowski metric, and by $C_{\alpha\beta}$ the anti-symmetric charge conjugation matrix.

By gauging the superalgebra, one can define the connection, curvature, corresponding gauge transformations, and Killing forms. With these definitions, we can proceed with constructing the action being invariant under Lorentz and supersymmetry transformations.

The Poincaré $\mathcal{N} = 2$ supergravity contains $SO(2)$ automorphisms group generated by T . Generator T rotates supercharges. Simultaneously there is another $U(1)$ generator playing the role of the central charge in the anticommutator of $\{Q_\alpha^I, Q_\beta^J\}$. The Killing form between them is non-zero [7], leading to a mixed $U(1)$ term in the gauge action. While in the AdS case, we have a single generator T in both mentioned the (anti)-commutation rules. Hence we can not call it a central charge. However, this way, we achieve the non-zero Killing form $\langle TT \rangle$ and a standard $U(1)$ term in the gauge action, which is not possible in the Poincaré theory.

The \mathcal{N} -extended theories are obtained by enlarging the number of fermionic generators. A similar mechanism has been used for the bosonic counterpart in recent years. It is achieved by including additional Lorentz-like and translational-like generators to the base of Poincaré/AdS algebras. This research direction takes its origin from the so-called Maxwell algebra [11, 12], and further development obtained by the so-called Semigroup expansion (*S-expansion*) procedure [13, 14, 15, 16, 17]. It provides a consistent way of deriving new (super)algebras. The essence of the semigroup expansion method comes down to maintaining the same pattern of structure constants in the (anti)-commutation rules for the newly added generators as for primary (supersymmetric) AdS algebra. The algebras which obey this property we shall call “resonant”.

Some of the resonant/semigroup algebras were analyzed in the context of various applications [18, 19, 20, 21, 22, 23, 24, 25, 26]. However, still, the interpretation of the new fields remains unclear.

The discussion of all algebraic possibilities for a given set of generators in the bosonic sector was completed in papers [27, 28]. After a series of particular examples [29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39] of supersymmetric versions, the resonant $\mathcal{N} = 1$ cases were concluded in Ref. [40]. In our recent paper [41], after an extensive search, we provided the complete list of resonant superalgebras up to two fermionic generators.

In the present paper, similarly to AdS scenario, we investigate the $\mathcal{N} = 2$ framework with T generator while still considering the resonant enlargement on the bosonic counterpart. Using the method described in Ref. [41], we consider the algebra structure, $JPZ + QY + T$, that contains the Lorentz J_{ab} , translation P_a and the additional Lorentz-like generator Z_{ab} . Note that this superalgebra contains two fermionic Majorana supercharges Q_α^I , ($Q_\alpha^1 \equiv Q_\alpha$ and $Q_\alpha^2 \equiv Y_\alpha$), and the $U(1)$ generator, denoted as T . Interestingly, we have obtained only one possible realization of the superalgebra in such a case.

In further sections, we will construct the 3D Chern-Simons (CS) Lagrangian [42, 43, 44] based on the 3D version of the derived superalgebra. We discuss the differences between the newly obtained supergravity model and the previous AdS result. Similar studies can be found in literature, [30, 40]. However, we provide a much simpler algebraic structure, which originates from the so-called Soroka-Soroka enlargement [14, 29, 45] instead of the one based on Maxwell algebra [11, 12]. In our approach, we avoided considering additional bosonic $U(1)$ generators.

The paper is organized as follows: in Sec. 2 we give new $\mathcal{N} = 2$ superalgebra; in Sec. 3 we construct the respective action for the 3D Chern-Simons model. We complete the paper with the summary and outlook in Sec. 4. The paper also contains three appendixes.

2 $\mathcal{N} = 2$ Soroka-Soroka superalgebra

Following the “resonant“ method introduced in Ref. [41], we consider, besides Lorentz and translations generators, also bosonic Lorentz-like Z_{ab} . We include the T generator, incorporating the $U(1)$ field (associated with T) into the supergravity framework. The obtained results are valid for 3D and 4D; for more explanation see [41].

The structure of superalgebras with T generator is more complicated than the ones without it. Similarly, as in Ref. [41], to find the superalgebras, we consider computer-generated thousands of algebra candidates (14580). The successful superalgebra must satisfy the Jacobi identities, therefore the Jacobi identities have to be computed and verified for each candidate.

Ultimately we arrive at one distinct realization of $JPZ + QY + T$ superalgebra, which we shall discuss below. It is based on the Soroka-Soroka \mathfrak{C}_4 algebra [29, 45] and has the following form:

$$\begin{aligned}
[J_{ab}, J_{cd}] &= \eta_{bc}J_{ad} - \eta_{ac}J_{bd} + \eta_{ad}J_{bc} - \eta_{bd}J_{ac}, \\
[J_{ab}, Z_{cd}] &= \eta_{bc}Z_{ad} - \eta_{ac}Z_{bd} + \eta_{ad}Z_{bc} - \eta_{bd}Z_{ac}, \\
[Z_{ab}, Z_{cd}] &= \eta_{bc}Z_{ad} - \eta_{ac}Z_{bd} + \eta_{ad}Z_{bc} - \eta_{bd}Z_{ac}, \\
[J_{ab}, P_b] &= \eta_{bc}P_a - \eta_{ac}P_b, \\
[Z_{ab}, P_b] &= \eta_{bc}P_a - \eta_{ac}P_b, \\
[P_a, P_b] &= Z_{ab}, \\
[J_{ab}, Q_\alpha^I] &= \frac{1}{2} (\Gamma_{ab})_\alpha^\beta Q_\beta^I, & [P_a, Q_\alpha^I] &= \frac{1}{2} (\Gamma_a)_\alpha^\beta Q_\beta^I, \\
[Z_{ab}, Q_\alpha^I] &= \frac{1}{2} (\Gamma_{ab})_\alpha^\beta Q_\beta^I, & [T, Q_\alpha^I] &= \epsilon^{IJ} Q_\alpha^J \\
\{Q_\alpha^I, Q_\beta^J\} &= -\delta^{IJ} \left((\Gamma^a C)_{\alpha\beta} P_a - \frac{1}{2} (\Gamma^{ab} C)_{\alpha\beta} Z_{ab} \right) + \epsilon^{IJ} C_{\alpha\beta} T. \tag{2}
\end{aligned}$$

Derivation was based on the consistency with commutation relations $[T, Q^I] = \epsilon^{IJ} Q^J$ (rotating $[T, Q] \sim Y$ and $[T, Y] \sim Q$), commutator $[J, X] \sim X$ (with X being any generator, in particular, $[J, Q^I] \sim Q^I$), and the necessity of non-vanishing anticommutator $\{Q, Q\} \neq 0$, as well as T appearing in the anticommutator $\{Q, Y\} \sim T$. Note that Soroka-Soroka \mathfrak{C}_4 algebra can be expressed as the direct sum of the AdS and Lorentz algebras [27, 29], which is achieved by an appropriate change of the basis. However, the resulting generators lose their physical role; see comment in Appendix C.

We remind that there are six available bosonic JPZ algebras altogether [41], where besides the Soroka-Soroka algebra [29, 45] (also called *AdS-Maxwell* or *AdS-Lorentz*), we have the Maxwell algebra \mathfrak{B}_4 [11, 12] and four Poincaré-like algebras [27]. The bosonic subalgebra for newly obtained example turned out to be solely given by the Soroka-Soroka algebra.

Some effort has been made to achieve algebra based on Maxwell \mathfrak{B}_4 [11, 12]. However, they have ended with a much more complicated setup with additional $U(1)$ generators, or a modification, which includes the new Lorentz-like generator Z_{ab} in $\{Q_\alpha, Y_\beta\}$ [40]. Note that $\gamma_5 C_{\alpha\beta}$ and central charge T_C could be included in $\{Q_\alpha^I, Q_\beta^J\}$ [30], but we do not consider this possibility for direct clarity. In the future, we will discuss a much broader scope and incorporate the resonant framework within the Poincaré $\mathcal{N} = 2$ superalgebra.

3 Supergravity action in $D = 3$

As the construction of the 3D Chern-Simons model is straightforward and assures the full symmetry invariance, we give the 3D CS model based on the new superalgebra. In this case, we transit to dual definitions of generators $J_a = \frac{1}{2}\epsilon_a{}^{bc} J_{bc}$, spin connection $\omega_a = \frac{1}{2}\epsilon_a{}^{bc} \omega_{bc}$, as well as definitions like Lorentz curvature $\mathcal{R}^a(\omega) = d\omega^a + \frac{1}{2}\epsilon^{abc}\omega_b\omega_c$. We define also covariant derivative, which acting on dreibein gives $D_\omega e^a = de^a + \epsilon^{abc}\omega_b e_c$, and acting on the Majorana spinor gives $D_\omega\psi = d\psi + \frac{1}{2}\omega^a\Gamma_a\psi$.

The necessary elements to construct Lagrangian are the gauge connection one-form \mathbb{A} ; the super-curvature two-form \mathbb{F} ; the gauge parameter Θ , along with the gauge transformations; and the Killing form $\langle \dots \rangle$ given in the form of the invariant tensors.

The connection contains additional Lorentz-like generator $Z_a = \frac{1}{2}\epsilon_a{}^{bc} Z_{bc}$ with a corresponding field $h_a = \frac{1}{2}\epsilon_a{}^{bc} h_{bc}$, and two gravitinos, ψ associated with Q and χ associated with Y ,

$$\mathbb{A} = \omega^a J_a + \frac{1}{\ell} e^a P_a + h^a Z_a + \frac{1}{\sqrt{\ell}} \psi^\alpha Q_\alpha + \frac{1}{\sqrt{\ell}} \chi^\alpha Y_\alpha + \frac{1}{\ell} a T. \quad (3)$$

We also introduce the ℓ parameter to make dimensions right. The super-curvature two-form \mathbb{F} is built straightforwardly from the \mathbb{A} connection

$$\mathbb{F} = F^a J_a + \frac{1}{\ell} T^a P_a + H^a Z_a + \frac{1}{\sqrt{\ell}} \mathcal{F}^\alpha Q_\alpha + \frac{1}{\sqrt{\ell}} \mathcal{G}^\alpha Y_\alpha + \frac{1}{\ell} f(a) T. \quad (4)$$

Indeed, for the superalgebra (2) we have:

$$\begin{aligned} F^a &= \mathcal{R}^a(\omega), \\ H^a &= D_\omega h^a + \frac{1}{2}\epsilon^{abc} h_b h_c + \frac{1}{2\ell^2} \epsilon^{abc} e_b e_c + \frac{1}{2\ell} (\bar{\psi}\Gamma^a\psi + \bar{\chi}\Gamma^a\chi), \\ T^a &= D_\omega e^a + \frac{1}{2} (\bar{\psi}\Gamma^a\psi + \bar{\chi}\Gamma^a\chi), \\ \mathcal{F} &= \mathcal{D}_\omega\psi + \frac{1}{2\ell} e^a \Gamma_a \psi + \frac{1}{2} h^a \Gamma_a \psi - \frac{1}{\ell} a \chi, \\ \mathcal{G} &= \mathcal{D}_\omega\chi + \frac{1}{2\ell} e^a \Gamma_a \chi + \frac{1}{2} h^a \Gamma_a \chi + \frac{1}{\ell} a \psi, \\ f &= da - \frac{1}{2} (\psi^\alpha C_{\alpha\beta} \chi^\beta - \chi^\alpha C_{\alpha\beta} \psi^\beta) = da - \frac{1}{2} (\bar{\psi}\chi - \bar{\chi}\psi). \end{aligned} \quad (5)$$

The general the three-dimensional Chern-Simons action reads

$$I_{CS} = \frac{k}{4\pi} \int_{\mathcal{M}} \left\langle \mathbb{A} \wedge d\mathbb{A} + \frac{1}{3} \mathbb{A} \wedge [[\mathbb{A}, \mathbb{A}]] \right\rangle, \quad (6)$$

where $[[\dots]]$ is the generalized commutator/anticommutator, and $\langle \dots \rangle$ is Killing form being used to contract all the group indices. Depending on the particular algebra [27, 41] we calculate Killing forms as the direct consequence of the evaluation of identity with (anti)-commutation relations:

$$\langle [[X_i, X_j]] X_k \rangle = \langle X_i [[X_j, X_k]] \rangle, \quad (7)$$

where X_i is any generator. As a result the Killing metric takes the form of the invariant tensor given for any combination of two generators. As for the bosonic part:

$$\langle J_a J_b \rangle = \alpha_0 \eta_{ab}, \quad (8)$$

$$\langle J_a P_b \rangle = \langle Z_a P_b \rangle = \alpha_1 \eta_{ab}, \quad (9)$$

$$\langle J_a Z_b \rangle = \langle Z_a Z_b \rangle = \langle P_a P_b \rangle = \alpha_2 \eta_{ab}. \quad (10)$$

For the fermionic part we have:

$$\langle Q_\alpha Q_\beta \rangle = \langle Y_\alpha Y_\beta \rangle = 2(\alpha_1 + \alpha_2) C_{\alpha\beta}. \quad (11)$$

The most important here, although $[T, T] = 0$, is the non-degenerate

$$\langle T T \rangle = 2(\alpha_1 + \alpha_2), \quad (12)$$

allowing for the ada term in the action.

Note that the resonant/semigroup framework introduces the sub-invariant sectors by the arbitrary valued α 's constants in front of invariant tensors.

The $JPZ + QY + T$ superalgebra 2 has the corresponding 3D Chern-Simons action:

$$\begin{aligned} I = \frac{k}{4\pi} \int \left[\alpha_0 \left(\omega^a d\omega_a + \frac{1}{3} \epsilon^{abc} \omega_a \omega_b \omega_c \right) \right. \\ \left. + \alpha_1 \left(\frac{2}{\ell} \mathcal{R}^a e_a + \frac{1}{3\ell^3} \epsilon^{abc} e_a e_b e_c + \frac{2}{\ell} e_a D_\omega h^a + \frac{1}{\ell} \epsilon^{abc} e_a h_b h_c + \frac{2}{\ell} \bar{\psi} \mathcal{F} + \frac{2}{\ell} \bar{\chi} \mathcal{G} + \frac{2}{\ell^2} a f(a) \right) \right. \\ \left. + \alpha_2 \left(\frac{1}{\ell^2} e_a D_\omega e^a + 2h_a \mathcal{R}^a + \frac{1}{\ell^2} \epsilon^{abc} e_a e_b h_c + h_a D_\omega h^a + \frac{1}{3} \epsilon^{abc} h_a h_b h_c + \frac{2}{\ell} \bar{\psi} \mathcal{F} + \frac{2}{\ell} \bar{\chi} \mathcal{G} + \frac{2}{\ell^2} a f(a) \right) \right]. \end{aligned} \quad (13)$$

Similarly, as in previous works [26, 40, 41, 46] the action can be decomposed into several sub-invariant sectors defined by dimensionless α_i 's constants. The separation of terms into the sectors directly depends on the superalgebra.

Let us underline that our model's supergravity action (α_1 sector) is very similar to the AdS one. Indeed, the fermionic contributions are the same. A difference is induced by the presence of the field h^a corresponding to the Z generator.

In comparison to the $\mathcal{N} = 2$ AdS case (see Appendix (B)), due to the generator Z , we have a new sector governed by the α_2 constant. Moreover, the torsion contributes to α_2 sector in the case of new algebra, whereas for the AdS, it appears in the "exotic" α_0 sector. Hence, the full new supergravity model leads to modification of the fermionic sector.

Besides the dynamical α_1 sector, our new action contains additional fermionic terms. The fermionic terms contribute to α_2 sector, whereas α_0 sector contains purely $CS(\omega)$.

We note that besides the term $a(\bar{\psi}\chi - \bar{\chi}\psi)$, resulting directly from a sum of $a f(a)$ and $\bar{\psi}\mathcal{F} + \bar{\chi}\mathcal{G}$ part, we got cancellations in the actions of other mixed fermionic terms due to $\langle Q_\alpha Y_\beta \rangle = 0$. As a result, in the final action we have pure $U(1)$ term $\frac{2}{\ell^2} a da$ plus the mentioned $\frac{1}{\ell^2} a(\bar{\psi}\chi - \bar{\chi}\psi)$ contribution.

4 Outlook

We have introduced a new $\mathcal{N} = 2$ superalgebra rooted in the S -expansion procedure. This superalgebra, originating in the Soroka-Soroka algebra (with generator Z_{ab} besides J_{ab} and P_a), represents a much simpler configuration than found in Refs. [30, 40]). Similar to our previous paper [41], this results from extensive computer-assisted validation of the super Jacobi identities among thousands of possible algebra candidates.

We provide Chern-Simons 3D action corresponding to new superalgebra 2. Compared to the known $\mathcal{N} = 2$ AdS example, the new CS model provides modified term distribution into sub-invariant sectors.

The action construction could be carried out in 4D, but we leave it for future work. The discussion of a broader perspective on the closing of resonant algebras, including the Poincaré-like scenarios, will be a subject of separate studies.

Acknowledgments

We thank prof. Jerzy Lukierski, for reading the manuscript and for many valuable insights and comments. We also thank prof. Dragoljub Gočanin for pointing out helpful literature concerning $\mathcal{N} = 2$ Poincaré/AdS superalgebras.

The research project for both authors was partly supported by the program "Excellence initiative - research university" for the years 2020-2026 for the University of Wrocław.

A 3D AdS $\mathcal{N} = 2$ superalgebra

For completeness, we provide $JPZ + QY + T$ configuration for the AdS $D = 3$ $\mathcal{N} = 2$:

$$\begin{aligned} [J_a, J_b] &= \epsilon_{abc} J^c, & [J_a, P_b] &= \epsilon_{abc} P^c, & [P_a, P_b] &= \epsilon_{abc} J^c, \\ [J_a, Q_\alpha^I] &= \frac{1}{2} (\Gamma_a)^\beta_\alpha Q_\beta^I, & [P_a, Q_\alpha^I] &= \frac{1}{2} (\Gamma_a)^\beta_\alpha Q_\beta^I, & [T, Q_\alpha^I] &= \epsilon^{IJ} Q_\alpha^J, \\ \{Q_\alpha^I, Q_\beta^J\} &= -\delta^{IJ} \left((\Gamma^a C)_{\alpha\beta} P_a + (\Gamma^a C)_{\alpha\beta} J_a \right) + \epsilon^{IJ} C_{\alpha\beta} T. \end{aligned} \quad (14)$$

The algebraic structure can be highlighted by the convenient table description shown in [41, 46], which allows us to focus on the type of the generator outcomes, leaving aside the particular structure constants. For the Majorana supercharges $Q^1 \equiv Q$ and $Q^2 \equiv Y$ written explicitly, the superalgebra (14) in the table form is given by

$$\begin{array}{c|ccc} [,] & J & P & T \\ \hline J & J & P & 0 \\ P & P & J & 0 \\ \hline T & 0 & 0 & 0 \end{array} \quad \begin{array}{c|cc} [,] & Q & Y \\ \hline J & Q & Y \\ P & Q & Y \\ \hline T & Y & Q \end{array} \quad \begin{array}{c|cc} \{, \} & Q & Y \\ \hline Q & P+J & T \\ Y & T & P+J \end{array} \quad (15)$$

We highlight the difference between the AdS (1) and our new $\mathcal{N} = 2$ superalgebra, by presenting (2) in the table-form:

$$\begin{array}{c|ccc} [,] & J & P & Z \\ \hline J & J & P & Z \\ P & P & Z & P \\ Z & Z & P & Z \\ \hline T & 0 & 0 & 0 \end{array} \quad \begin{array}{c|cc} [,] & Q & Y \\ \hline J & Q & Y \\ P & Q & Y \\ Z & Q & Y \\ \hline T & Y & Q \end{array} \quad \begin{array}{c|cc} \{, \} & Q & Y \\ \hline Q & P+Z & T \\ Y & T & P+Z \end{array} \quad (16)$$

B 3D AdS $\mathcal{N} = 2$ supergravity

The connection corresponding to the supersymmetric AdS is given by

$$\mathbb{A} = \omega^a J_a + \frac{1}{\ell} e^a P_a + \frac{1}{\sqrt{\ell}} \psi^\alpha Q_\alpha + \frac{1}{\sqrt{\ell}} \chi^\alpha Y_\alpha + \frac{1}{\ell} a T, \quad (17)$$

where we assure proper dimensionality by the introducing length parameter ℓ related to the cosmological constant $\Lambda = -1/\ell^2$.

The super-curvature two-form $\mathbb{F} = d\mathbb{A} + \frac{1}{2} [[\mathbb{A}, \mathbb{A}]]$ is built straightforwardly from \mathbb{A} connection,

$$\mathbb{F} = F^a J_a + \frac{1}{\ell} T^a P_a + \frac{1}{\sqrt{\ell}} \mathcal{F}^\alpha Q_\alpha + \frac{1}{\sqrt{\ell}} \mathcal{G}^\alpha Y_\alpha + \frac{1}{\ell} f(a) T, \quad (18)$$

where

$$\begin{aligned}
F^a &= \mathcal{R}^a(\omega) + \frac{1}{2\ell^2}\epsilon^{abc}e_b e_c + \frac{1}{2\ell}\bar{\psi}\Gamma^a\psi + \frac{1}{2\ell}\bar{\chi}\Gamma^a\chi, \\
T^a &= D_\omega e^a + \frac{1}{2}\bar{\psi}\Gamma^a\psi + \frac{1}{2}\bar{\chi}\Gamma^a\chi, \\
f &= da - \frac{1}{2}(\bar{\psi}\chi - \bar{\chi}\psi), \\
\mathcal{F} &= \mathcal{D}_\omega\psi + \frac{1}{2\ell}e^a\Gamma_a\psi - \frac{1}{\ell}a\chi, \\
\mathcal{G} &= \mathcal{D}_\omega\chi + \frac{1}{2\ell}e^a\Gamma_a\chi + \frac{1}{\ell}a\psi.
\end{aligned} \tag{19}$$

Evaluation of the invariant tensors eventually leads to the corresponding action:

$$\begin{aligned}
I_{CS}^{AdS} &= \frac{k}{4\pi} \int \left[\alpha_0 \left(\omega^a d\omega_a + \frac{1}{3}\epsilon^{abc}\omega_a\omega_b\omega_c + \frac{1}{\ell^2}e_a D_\omega e^a + \frac{2}{\ell}\bar{\psi}\mathcal{F} + \frac{2}{\ell}\bar{\chi}\mathcal{G} + \frac{2}{\ell^2}a f(a) \right) \right. \\
&\quad \left. + \alpha_1 \left(\frac{2}{\ell}\mathcal{R}^a e_a + \frac{1}{3\ell^3}\epsilon^{abc}e_a e_b e_c + \frac{2}{\ell}\bar{\psi}\mathcal{F} + \frac{2}{\ell}\bar{\chi}\mathcal{G} + \frac{2}{\ell^2}a f(a) \right) \right].
\end{aligned} \tag{20}$$

with emergence of the sub-invariant sectors given by the arbitrary constants: α_0 and α_1 . In the first line we recognize “exotic” Lagrangian, whereas other is a standard ”gravitational” one.

C $\mathcal{N} = 2$ super $AdS \oplus Lorentz$

The Soroka-Soroka \mathfrak{C}_4 algebra can be expressed as the direct sum of the AdS and Lorentz algebras [27, 29]. It is done by the change of basis where the generators are rearranged as follows $L_{AB} = (L_{ab} = Z_{ab}, L_{a4} = P_a)$ and $N_{ab} = J_{ab} - Z_{ab}$. Then the superalgebra (2) reads

$$\begin{array}{c|ccc|c}
[,] & L_{..} & L & N_{..} & T \\
\hline
L_{..} & L_{..} & L & 0 & 0 \\
L & L & L_{..} & 0 & 0 \\
N_{..} & 0 & 0 & N_{..} & 0 \\
\hline
T & 0 & 0 & 0 & 0
\end{array}
\quad
\begin{array}{c|cc}
[,] & Q & Y \\
\hline
L_{..} & Q & Y \\
L & Q & Y \\
N_{..} & 0 & 0 \\
\hline
T & Y & Q
\end{array}
\quad
\begin{array}{c|cc}
\{ , \} & Q & Y \\
\hline
Q & L_{..} + L_{..} & T \\
Y & T & L_{..} + L_{..}
\end{array} \tag{21}$$

Evidently the above tables show that (2) represents the direct sum of the super AdS $\mathcal{N} = 2$ (spanned by $L_{ab}, P_a, Q_\alpha^I, T$) and the Lorentz algebra given by N_{ab} . Note, however, that neither the L_{ab} nor N_{ab} are the physical Lorentz generator as they break general covariance. Therefore, such change of basis is convenient only for computational reasons. Eventually, let us underline the special role of the Lorentz J generator that introduces the spin connection ω and preserves all of other generators in the (anti)-commutation relations.

References

- [1] Daniel Z. Freedman, Peter van Nieuwenhuizen, and Sergio Ferrara. Progress Toward a Theory of Supergravity. *Phys. Rev. D*, 13:3214–3218, 1976.
- [2] Stanley Deser and Bruno Zumino. Consistent Supergravity. *Phys. Lett. B*, 62:335, 1976.
- [3] Paul K. Townsend. Cosmological Constant in Supergravity. *Phys. Rev. D*, 15:2802–2804, 1977.

- [4] Sergio Ferrara and Peter van Nieuwenhuizen. Consistent Supergravity with Complex Spin 3/2 Gauge Fields. *Phys. Rev. Lett.*, 37:1669, 1976.
- [5] Rudolf Haag, Jan T. Lopuszanski, and Martin Sohnius. All Possible Generators of Supersymmetries of the s Matrix. *Nucl. Phys. B*, 88:257, 1975.
- [6] Peter Van Nieuwenhuizen. Supergravity. *Phys. Rept.*, 68:189–398, 1981.
- [7] Paul S. Howe, Jose. M. Izquierdo, Georgios Papadopoulos, and Paul K. Townsend. New supergravities with central charges and Killing spinors in (2+1)-dimensions. *Nucl. Phys. B*, 467:183–214, 1996.
- [8] Paul K. Townsend and Peter van Nieuwenhuizen. Geometrical Interpretation of Extended Supergravity. *Phys. Lett. B*, 67:439–442, 1977.
- [9] Tomas Ortin. *Gravity and Strings*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, 2nd ed. edition, 7 2015.
- [10] Dragoljub Gočanin and Voja Radovanović. Canonical deformation of $N = 2$ AdS_4 supergravity. *Phys. Rev. D*, 100(9):095019, 2019.
- [11] Robert Schrader. The maxwell group and the quantum theory of particles in classical homogeneous electromagnetic fields. *Fortsch. Phys.*, 20:701–734, 1972.
- [12] Henri Bacry, Philippe Combe, and Jean-Louis Richard. Group-theoretical analysis of elementary particles in an external electromagnetic field. 1. the relativistic particle in a constant and uniform field. *Nuovo Cim. A*, 67:267–299, 1970.
- [13] Fernando Izaurieta, Eduardo Rodriguez, and Patricio Salgado. Expanding Lie (super)algebras through Abelian semigroups. *J. Math. Phys.*, 47:123512, 2006.
- [14] Patricio Salgado and Sebastian Salgado. $\mathfrak{so}(D - 1, 1) \otimes \mathfrak{so}(D - 1, 2)$ algebras and gravity. *Phys. Lett. B*, 728:5–10, 2014.
- [15] Laura Andrianopoli, Nelson Merino, Felip Nadal, and Mario Trigiante. General properties of the expansion methods of Lie algebras. *J. Phys. A*, 46:365204, 2013.
- [16] Patrick K. Concha, Remigiusz Durka, Nelson Merino, and Evelyn. K. Rodríguez. New family of Maxwell like algebras. *Phys. Lett. B*, 759:507–512, 2016.
- [17] Patrick K. Concha, Remigiusz Durka, Carlos. Inostroza, Nelson Merino, and Evelyn K. Rodríguez. Pure Lovelock gravity and Chern-Simons theory. *Phys. Rev. D*, 94(2):024055, 2016.
- [18] Jose A. de Azcarraga, Kiyoshi Kamimura, and Jerzy Lukierski. Generalized cosmological term from Maxwell symmetries. *Phys. Rev. D*, 83:124036, 2011.
- [19] Laura Andrianopoli and Lucrezia Ravera. On the Geometric Approach to the Boundary Problem in Supergravity. *Universe*, 7(12):463, 2021.
- [20] Remigiusz Durka and Jerzy Kowalski-Glikman. Local Maxwell symmetry and gravity. 10 2011.

- [21] Patricio Salgado, Richard J. Szabo, and Omar Valdivia. Topological gravity and transgression holography. *Phys. Rev. D*, 89(8):084077, 2014.
- [22] Patrick Concha, Nelson Merino, Olivera Miskovic, Evelyn Rodríguez, Patricio Salgado-Rebolledo, and Omar Valdivia. Asymptotic symmetries of three-dimensional Chern-Simons gravity for the Maxwell algebra. *JHEP*, 10:079, 2018.
- [23] Patrick Concha, Nelson Merino, Evelyn Rodríguez, Patricio Salgado-Rebolledo, and Omar Valdivia. Semi-simple enlargement of the \mathfrak{bms}_3 algebra from a $\mathfrak{so}(2, 2) \oplus \mathfrak{so}(2, 1)$ Chern-Simons theory. *JHEP*, 02:002, 2019.
- [24] Patricio Salgado-Rebolledo. The Maxwell group in 2+1 dimensions and its infinite-dimensional enhancements. *JHEP*, 10:039, 2019.
- [25] Giandomenico Palumbo. Geometric Model of Topological Insulators from the Maxwell Algebra. *Annals Phys.*, 386:15–24, 2017.
- [26] Remigiusz Durka and Jerzy Kowalski-Glikman. Resonant algebras in Chern-Simons model of topological insulators. *Phys. Lett. B*, 795:516–520, 2019.
- [27] Remigiusz Durka. Resonant algebras and gravity. *J. Phys. A*, 50(14):145202, 2017.
- [28] Remigiusz Durka and Kamil Grela. On the number of possible resonant algebras. *J. Phys. A*, 53(35):355202, 2020.
- [29] Dmitriy V. Soroka and Vyacheslav A. Soroka. Semi-simple extension of the (super)Poincare algebra. *Adv. High Energy Phys.*, 2009:234147, 2009.
- [30] Sotirios Bonanos, Joaquim Gomis, Kiyoshi Kamimura, and Jerzy Lukierski. Maxwell Superalgebra and Superparticle in Constant Gauge Backgrounds. *Phys. Rev. Lett.*, 104:090401, 2010.
- [31] Sotirios Bonanos, Joaquim Gomis, Kiyoshi Kamimura, and Jerzy Lukierski. Deformations of Maxwell Superalgebras and Their Applications. *J. Math. Phys.*, 51:102301, 2010.
- [32] Kiyoshi Kamimura and Jerzy Lukierski. Supersymmetrization Schemes of D=4 Maxwell Algebra. *Phys. Lett. B*, 707:292–297, 2012.
- [33] Jose. A. de Azcarraga and Jose M. Izquierdo. Minimal D = 4 supergravity from the superMaxwell algebra. *Nucl. Phys. B*, 885:34–45, 2014.
- [34] Remigiusz Durka, Jerzy Kowalski-Glikman, and Michal Szczachor. AdS-Maxwell superalgebra and supergravity. *Mod. Phys. Lett. A*, 27:1250023, 2012.
- [35] Remigiusz Durka. *Deformed BF theory as a theory of gravity and supergravity*. PhD thesis, Wrocław U., 2012.
- [36] Jose A. de Azcarraga, Jose M. Izquierdo, Jerzy Lukierski, and Mariusz Woronowicz. Generalizations of Maxwell (super)algebras by the expansion method. *Nucl. Phys. B*, 869:303–314, 2013.
- [37] Octavio Fierro, Fernando. Izaurieta, Patricio Salgado, and Omar Valdivia. Minimal AdS-Lorentz supergravity in three-dimensions. *Phys. Lett. B*, 788:198–205, 2019.

- [38] Diego M. Peñafiel and Lucrezia Ravera. Generalized cosmological term in $D = 4$ supergravity from a new AdS–Lorentz superalgebra. *Eur. Phys. J. C*, 78(11):945, 2018.
- [39] Patrick K. Concha and Evelyn K. Rodríguez. $N = 1$ Supergravity and Maxwell superalgebras. *JHEP*, 09:090, 2014.
- [40] Patrick Concha. N -extended Maxwell supergravities as Chern-Simons theories in three spacetime dimensions. *Phys. Lett. B*, 792:290–297, 2019.
- [41] Remigiusz Durka and Krzysztof M. Graczyk. Resonant superalgebras for supergravity. *Eur. Phys. J. C*, 82(3):254, 2022.
- [42] Ana Achúcarro and Paul K. Townsend. A Chern-Simons Action for Three-Dimensional anti-De Sitter Supergravity Theories. *Phys. Lett. B*, 180:89, 1986.
- [43] Mokhtar Hassaine and Jorge Zanelli. *Chern-Simons (super)gravity*, volume 2 of *100 years of general relativity*. World Scientific, Hackensack, 2016 edition, 2016.
- [44] Patrick K. Concha, Octavio Fierro, Evelyn K. Rodríguez, and Patricio Salgado. Chern–Simons supergravity in $D=3$ and Maxwell superalgebra. *Phys. Lett. B*, 750:117–121, 2015.
- [45] Dmitriy V. Soroka and Vyacheslav A. Soroka. Gauge semi-simple extension of the Poincaré group. *Phys. Lett. B*, 707:160–162, 2012.
- [46] Patrick Concha, Remigiusz Durka, and Evelyn Rodríguez. Resonant superalgebras and $\mathcal{N} = 1$ supergravity theories in three spacetime dimensions. *Phys. Lett. B*, 808:135659, 2020.