

Engineering steady-state correlations in uncorrelated thermal states using stochastic driving

Armin Rahmani^{1,2}

¹*Department of Physics and Astronomy and Advanced Materials Science and Engineering Center,
Western Washington University, Bellingham, Washington 98225, USA*

²*Kavli Institute for Theoretical Physics, University of California, Santa Barbara, California 93106, USA*

(Dated: May 13, 2022)

Nonequilibrium quantum dynamics can give rise to the emergence of novel steady states. We propose a scheme for driving an initially uncorrelated thermal state to generate customized correlation functions by determining and reverse-engineering the steady-state two-point functions for a class of Markov processes. We also extend the formalism to the calculation of four-point functions. We then apply our method to generating power-law correlated fermionic Green's functions. Furthermore, we find that the power-law patterns emerge at much shorter times than the convergence to the steady-state, at which point the disorder in the two-point correlations disappears. On the other hand, the density-density correlations exhibit steady-state disorder while following a power-law trendline distinct from Luttinger liquids.

I. INTRODUCTION

There has been significant progress in driving nonequilibrium quantum matter to prepare desired quantum states, particularly in the context of quantum simulations [1, 2]. While the objective is often creating states with equilibrium counterparts, such as the ground state of interacting Hamiltonians [3, 4], it has been realized that quantum driving may give rise to the emergence of states without equilibrium counterparts. Floquet engineering, i.e., controlling the evolution by periodic driving, has been a fertile ground for generating such states [5–7]. Furthermore, dissipation engineering can be similarly effective in quantum control for open systems [8–10].

In this work, we focus on a class of stochastic drives [11–15], which can be viewed as time-averaged evolution of a closed quadratic system [16], with formal similarities to open systems [17–20], and propose a scheme for creating almost arbitrary two-point correlation functions starting from uncorrelated thermal states. Our method utilizes a temporally uncorrelated noisy Hamiltonian drive with long-range hopping and correlated noise. The Lindblad master equation governing the dynamics does not have a Heisenberg term and only contains one double commutator, analogous to a fully dissipative open system. We show that it is possible to engineer a broad class of two-point correlation functions with some freedom over the drive structure. Long-range couplings [21–23] are necessary for creating long-range power-law correlations in the shortest possible time.

There is much less information in two-point correlations than in a quantum system's full many-body wave function. Therefore the nature of the steady states created by our engineered stochastic drive designed only to make a specific two-point function is unknown. To investigate whether our steady states fall into a known universality class with similar two-point functions, we calculated the long-time limit of the four-point density-density correlation function. We found a novel behavior, where the two-point function decays as a clean power law with the density correlator remaining highly disordered but following a power-law trendline with a larger exponent than x^{-2} , suggesting a distinct universality class.

The outline of this paper is as follows. In Sec. II, we ex-

tend the results of Ref. [24] on the steady-state two-point functions to steady-state four-point functions under fully stochastic dissipative driving. In Sec. III, we present our method for reverse-engineering the drive to provide the desired matrix of fermionic Green's functions. In Sec. IV, we apply our method to the case of power-law correlated Green's functions. In Sec. V, we examine the steady-state density-density correlation functions for the drive used in engineering the Green's function. Finally, we present our conclusions in Sec. VI.

II. STEADY-STATE TWO- AND FOUR-POINT FUNCTIONS FOR FULLY STOCHASTIC QUADRATIC DRIVING

Here we present the theoretical framework for stochastic dynamics capable of generating nontrivial noise-averaged correlations in fermionic systems, initially in an uncorrelated thermal state. For a system of N spinless fermions, where the creation/annihilation operators satisfy the anticommutation relation $\{c_i^\dagger, c_j\} = \delta_{ij}$, consider a quadratic Hamiltonian that stochastically fluctuates around zero. We write $H(t)$ as

$$H(t) = \xi(t)V, \quad V = \sum_{ij} J_{ij}c_i^\dagger c_j, \quad t > 0. \quad (1)$$

Here, J is an $N \times N$ Hermitian matrix, and $\xi(t)$ is Gaussian white noise with $\overline{\xi(t)} = 0$ and $\overline{\xi(t)\xi(t')} = \delta(t-t')$. The overline symbol represents averaging over the realizations of noise. The noise strength only sets the equilibration timescale and does not affect the nature of the steady-state. As we are interested in the long-time limit, we have set the noise strength to unity.

The evolution by $H(t)$ describes an isolated system with a vanishing average Hamiltonian driven by temporally uncorrelated and spatially correlated noise, with J encoding the spatial correlations. The system will reach a steady-state density matrix in the long-time limit $t \rightarrow \infty$, given by

$$\rho(t \rightarrow \infty) = \lim_{t \rightarrow \infty} \text{tr} \left[e^{-i \int_0^t H(r) dr} \rho(0) e^{i \int_0^t H(r) dr} \right], \quad (2)$$

where $\rho(0)$ is the initial density matrix. Notice that the Hamiltonians at different times commute and time-ordering is not

necessary. We can view the dynamics as a Markov process. The long-time limit of the dynamics can be alternatively considered as a time average over random times after a quench to Hamiltonian V , as the white noise is simply creating a random walk for the total time.

The equation of motion for the density matrix is then a Lindblad master equation $\partial_t \rho(t) = -\frac{1}{2}[[\rho, V], V]$. Similar to Ref. [24], we write a Heisenberg-picture analog of the master equation for a noise-averaged Heisenberg operator $O(t)$ as

$$\partial_t O(t) = -\frac{1}{2}[[O(t), V], V]. \quad (3)$$

An important observation is that under evolution with Eq. (3), generic quadratic (quartic) operators remain quadratic (quartic). In contrast, the evolution can spread operators into nontrivial linear combinations of other quadratic (quartic) operators. The above observation follows from the fact that in the commutator of two-quadratic forms, the quartic terms cancel out and we are left with another quadratic form $[\sum_{\alpha\beta} A_{\alpha\beta} c_\alpha^\dagger c_\beta, \sum_{ij} B_{ij} c_i^\dagger c_j] = \sum_{\alpha\beta} [A, B]_{\alpha\beta} c_\alpha^\dagger c_\beta$ [25].

We now consider a general quadratic operator

$$O(t) = \sum_{\alpha\beta} \mathcal{O}_{\alpha\beta}(t) c_\alpha^\dagger c_\beta, \quad (4)$$

where \mathcal{O} is an $N \times N$ matrix. In terms of the $N \times N$ matrix J [see Eq. (1)], we can then write the solution of the master equation (3) as [24]

$$\mathcal{O}_{\alpha\beta}(t) = \sum_{\sigma\lambda\eta\gamma} \mathcal{U}_{\alpha\sigma} \mathcal{U}_{\lambda\beta}^\dagger e^{-\frac{1}{2}(\mathcal{D}_\sigma - \mathcal{D}_\lambda)^2 t} \mathcal{U}_{\sigma\eta}^\dagger \mathcal{U}_{\eta\gamma} \mathcal{O}_{\eta\gamma}(0), \quad (5)$$

where column i of the unitary matrix \mathcal{U} is the eigenvector of J with eigenvalue \mathcal{D}_i . In other words,

$$J = \mathcal{U} \text{diag}(\mathcal{D}_1, \mathcal{D}_2, \dots, \mathcal{D}_N) \mathcal{U}^\dagger. \quad (6)$$

Assuming J has a nondegenerate spectrum, we can then write the long-time limit of the noise averaged Heisenberg operator for $t \rightarrow \infty$ as

$$\mathcal{O}_{\alpha\beta}(\infty) = \sum_{\eta\gamma} \left(\sum_{\lambda} \mathcal{U}_{\alpha\lambda} \mathcal{U}_{\lambda\beta}^\dagger \mathcal{U}_{\lambda\eta}^\dagger \mathcal{U}_{\eta\gamma} \right) \mathcal{O}_{\eta\gamma}(0), \quad (7)$$

We now extend this formalism to quartic operators. Of particular interest are density-density correlation functions. To find the steady-state correlation functions, again we use the Heisenberg-picture approach. Due to the cancellation of the sextic terms in the commutator of a quartic form and a quadratic from $V = \sum_{ij} J_{ij} c_i^\dagger c_j$, we find

$$\left[\sum_{\alpha\beta\gamma\delta} \mathcal{R}_{\alpha\beta\gamma\delta}(t) c_\alpha^\dagger c_\beta c_\gamma^\dagger c_\delta, V \right] = \sum_{\alpha\beta\gamma\delta} \mathcal{S}_{\alpha\beta\gamma\delta}(t) c_\alpha^\dagger c_\beta c_\gamma^\dagger c_\delta, \quad (8)$$

where

$$\mathcal{S}_{\alpha\beta\gamma\delta} = \sum_j \left(\mathcal{R}_{\alpha\beta\gamma j} J_{j\delta} - \mathcal{R}_{\alpha\beta j\delta} J_{\gamma j} + \mathcal{R}_{\alpha j\gamma\delta} J_{j\beta} - \mathcal{R}_{j\beta\gamma\delta} J_{\alpha j} \right). \quad (9)$$

Therefore all quartic operators of the form above retain this structure when evolving with the master equation (3). By using Eq. (9) twice, we can write the equation of motion for the tensor \mathcal{R} as

$$\partial_t \mathcal{R}_{\alpha\beta\gamma\delta} = -\frac{1}{2} \sum_{xyzw} \left(Q_{\alpha\beta\gamma\delta}^{xyzw} + Q_{\beta\gamma\delta\alpha}^{yzwx} + Q_{\gamma\delta\alpha\beta}^{zwx y} + Q_{\delta\alpha\beta\gamma}^{wxyz} \right) \mathcal{R}_{xyzw} \quad (10)$$

where

$$Q_{\alpha\beta\gamma\delta}^{xyzw} = -J_{\alpha x} J_{y\beta} \delta_{\gamma z} \delta_{w\delta} + J_{\alpha x} \delta_{y\beta} J_{\gamma z} \delta_{w\delta} - J_{\alpha x} \delta_{y\beta} \delta_{\gamma z} J_{w\delta} + J_{\alpha x}^2 \delta_{y\beta} \delta_{\gamma z} \delta_{w\delta}. \quad (11)$$

We now use the diagonalization of J [Eq. (6)] to find an exact solution to the equation above. We write the Kronecker deltas as elements of $\mathcal{U}^\dagger \mathcal{U}$ to obtain

$$\mathcal{R}_{\alpha\beta\gamma\delta}(t) = \sum_{xyzw,abcd} \mathcal{U}_{\alpha a} \mathcal{U}_{y b} \mathcal{U}_{\gamma c} \mathcal{U}_{w d} e^{-\frac{1}{2}(\mathcal{D}_a + \mathcal{D}_c - \mathcal{D}_b - \mathcal{D}_d)^2 t} \mathcal{U}_{d\delta}^\dagger \mathcal{U}_{c\gamma}^\dagger \mathcal{U}_{b\beta}^\dagger \mathcal{U}_{a\alpha}^\dagger \mathcal{R}_{xyzw}(0). \quad (12)$$

In the absence of any accidental vanishing of the exponent, we can calculate the long-time limit of the \mathcal{R} tensor by noting that we must have either $b = a$ and $d = c$ or $b = c$ and $d = a$ (which double counts the case with $a = b = c = d$ [26, 27]). Thus we find

$$\mathcal{R}_{\alpha\beta\gamma\delta}(\infty) = \sum_{xyzw} \left[\sum_{ac} \mathcal{U}_{\alpha a} \mathcal{U}_{\gamma c} \mathcal{U}_{c\gamma}^\dagger \mathcal{U}_{a\alpha}^\dagger \left(\mathcal{U}_{y a} \mathcal{U}_{w c} \mathcal{U}_{c\delta}^\dagger \mathcal{U}_{\alpha\beta}^\dagger + \mathcal{U}_{y c} \mathcal{U}_{w a} \mathcal{U}_{a\delta}^\dagger \mathcal{U}_{c\beta}^\dagger \right) - \sum_a \mathcal{U}_{\alpha a} \mathcal{U}_{\gamma a} \mathcal{U}_{a\gamma}^\dagger \mathcal{U}_{a\alpha}^\dagger \mathcal{U}_{y a} \mathcal{U}_{w a} \mathcal{U}_{a\delta}^\dagger \mathcal{U}_{\alpha\beta}^\dagger \right] \mathcal{R}_{xyzw}(0). \quad (13)$$

For the density-density correlator $n_1 n_j = c_i^\dagger c_i c_j^\dagger c_j$, we have

$$\mathcal{R}_{xyzw}(0) = \delta_{xi} \delta_{yi} \delta_{zj} \delta_{wj}.$$

However, we need to work with the more general tensors above as they appear in the Heisenberg-picture operator.

III. ENGINEERING STEADY-STATE GREEN'S FUNCTIONS IN UNCORRELATED THERMAL STATES

An important class of correlation functions, i.e., the equal-time single-particle Green's functions (also known as the

single-particle density matrix [28]) are given by the expectation values

$$G_{ij}(t) = \langle c_i^\dagger c_j \rangle = \text{tr}[\rho(t)c_i^\dagger c_j], \quad (14)$$

where $\rho(t)$ is the density matrix of the system at time t . We can collect these correlators into an $N \times N$ Hermitian matrix G , with elements $G_{ij} = G_{ji}^*$.

Let us consider a system initially prepared in a trivial thermal state at inverse temperature β with the density matrix

$$\rho(0) = \exp(-\beta H_0) / Z, \quad H_0 = \sum_i \mu_i \hat{n}_i, \quad (15)$$

with the partition function $Z = \text{tr}[\exp(-\beta H_0)]$. As H_0 is diagonal in the occupation-number basis, the different sites are uncorrelated in the thermal state above and we can write the initial Green's function as

$$G_{ij}(0) = f(\mu_j) \delta_{ij} = \frac{1}{1 + e^{\beta \mu_i}} \delta_{ij}, \quad (16)$$

where f represents the Fermi-Dirac function.

Suppose we drive this system by a stochastic quadratic Hamiltonian as in Eq. (1). Can we engineer custom-ordered Green's functions in the steady-state of the system that emerges in the long time limit? More precisely, suppose our goal is choosing μ_i and J_{ij} such that

$$\lim_{t \rightarrow \infty} \overline{\text{tr}[U(t)\rho_0 U^\dagger(t)c_i^\dagger c_j]} = G_{ij}, \quad (17)$$

for given (perhaps even arbitrary) Hermitian correlation matrix G and initial inverse temperature β .

For the $c_i^\dagger c_j$ operator, we have $\mathcal{O}_{\eta\gamma}(0) = \delta_{\eta i} \delta_{\gamma j}$, which leads to the following matrix elements when inserted in Eq. (5)

$$\mathcal{O}_{\alpha\beta}(t) = \sum_{\sigma\lambda} \mathcal{U}_{\alpha\sigma} \mathcal{U}_{\lambda\beta}^\dagger e^{-\frac{1}{2}(\mathcal{D}_\sigma - \mathcal{D}_\lambda)^2 t} \mathcal{U}_{\sigma i}^\dagger \mathcal{U}_{j\lambda} \quad (18)$$

Since we are engineering the matrix J , we can require it to have a nondegenerate spectrum. Thus, analogously to Eq. (7), the Heisenberg-picture matrix corresponding to the long-time limit of the $c_i^\dagger c_j$ operator is given by

$$\mathcal{O}_{\alpha\beta}(\infty) = \sum_\lambda \mathcal{U}_{\alpha\lambda} \mathcal{U}_{\lambda\beta}^\dagger \mathcal{U}_{\lambda i}^\dagger \mathcal{U}_{j\lambda} \quad (19)$$

The above expressions lead to the relationships below between the time-dependent and steady-state correlation functions and the initial correlation functions through the eigenvectors of J :

$$G_{ij}(t) = \sum_{\alpha\beta\lambda\sigma} \mathcal{U}_{j\lambda} \mathcal{U}_{\lambda\beta}^\dagger G_{\alpha\beta}(0) e^{-\frac{1}{2}(\mathcal{D}_\sigma - \mathcal{D}_\lambda)^2 t} \mathcal{U}_{\alpha\sigma} \mathcal{U}_{\sigma i}^\dagger \quad (20)$$

$$G_{ij} = G_{ij}(\infty) = \sum_\lambda \mathcal{U}_{j\lambda} \left(\sum_{\alpha\beta} \mathcal{U}_{\lambda\beta}^\dagger G_{\alpha\beta}(0) \mathcal{U}_{\alpha\lambda} \right) \mathcal{U}_{\lambda i}^\dagger \quad (21)$$

The above equation implies that $G^T = \mathcal{U} \mathcal{A} \mathcal{U}^\dagger$, where \mathcal{A} is a diagonal matrix with

$$\mathcal{A}_{\lambda\sigma} = \delta_{\lambda\sigma} \sum_{\alpha\beta} \mathcal{U}_{\lambda\beta}^\dagger G_{\alpha\beta}(0) \mathcal{U}_{\alpha\lambda}. \quad (22)$$

Two conditions follow immediately: (i) G^T and J must have the same eigenvectors since the same unitary operator \mathcal{U} diagonalizes both of them, and (ii) The eigenvalues g_σ of G^T are the diagonal elements of \mathcal{A} and related to the initial correlation functions as follows

$$g_\sigma = \sum_\alpha |\mathcal{U}_{\alpha\sigma}|^2 f(\mu_\alpha) \equiv \sum_\alpha W_{\sigma\alpha} f(\mu_\alpha), \quad (23)$$

where the matrix W of transition probabilities is defined through

$$W_{\sigma\alpha} = |\mathcal{U}_{\alpha\sigma}|^2. \quad (24)$$

We have inserted Eq. (16) into Eq. (22) to obtain the condition (23). The matrix W is a doubly stochastic (it has nonnegative elements and the sum of every row and every column is equal to unity), which implies the sum rule $\sum_\alpha g_\alpha = \sum_\alpha f(\mu_\alpha)$. Given a matrix G of desired steady-state correlators, we can find g_σ and \mathcal{U} by diagonalizing G . The matrix W can then be explicitly constructed. For a given initial inverse temperature β , we can then write the chemical potentials in the initial Hamiltonian as

$$\mu_j = \frac{1}{\beta} \ln \left(\frac{1}{\sum_\sigma W_{j\sigma}^{-1} g_\sigma} - 1 \right). \quad (25)$$

The columns of the matrix $\mathcal{U} = (|g_1\rangle, |g_2\rangle, \dots)$ are eigenvectors $|g_\alpha\rangle$ of $G = \sum_\alpha g_\alpha |g_\alpha\rangle \langle g_\alpha|$. The matrix J can then be any linear combination

$$J = \sum_\alpha d_\alpha |g_\alpha\rangle \langle g_\alpha| \quad (26)$$

as long as the eigenvalues $\{d_\alpha\}$ are nondegenerate.

It may appear that the Hermitian matrix G can be completely arbitrary, and we can create any correlation functions we choose. Hermiticity is not the only requirement of a physical matrix of Green's functions, however. For example, all diagonal elements are the occupation number of a site and must be between zero and one, so creating states with nonphysical occupation numbers should be impossible. Furthermore, the method fails for many physically allowed Green's functions, as discussed below.

First, the matrix W must be invertible, which is not guaranteed. Indeed we have found that W tends to be singular for uniform systems, and some disorder in the occupation numbers is necessary for its invertibility. Second, the Fermi-Dirac functions are bounded as $0 \leq f(\mu_\alpha) \leq 1$, which is equivalent to the condition that the chemical potentials written in Eq. (25) are real numbers. Therefore, the eigenvalues and eigenvectors of G must be consistent with the following constraint

$$0 \leq \sum_\sigma W_{j\sigma}^{-1} g_\sigma \leq 1 \quad (27)$$

for all j .

The constraint of Eq. (27) is not too restrictive if we are only interested in a pattern of correlations between different sites (off-diagonal elements of G) without simultaneously




FIG. 1. (a) An example of the target Green's function G with power-law off-diagonal elements for a system of $N = 50$ sites, with $c = 0.02$, and $\nu = 1.5$ [see Eq. (30)]. Only the off-diagonal elements are illustrated in panel (a). (b) The diagonal disordered elements of G , i.e., the target steady-state occupation numbers, and the corresponding initial occupation numbers $f(\mu_i)$ form Eq. (25).

specifying onsite occupation numbers (diagonal elements of G). This observation is because we can change the eigenvalues of G by adding a multiple of the identity matrix to G , which only changes the occupation numbers, or multiplying G by a constant, which only rescales the Green's functions, without changing their overall pattern. More specifically, suppose for some matrix G , $a = \min(\sum_{\sigma} W_{j\sigma}^{-1} g_{\sigma}) < 0$ and $b = \max(\sum_{\sigma} W_{j\sigma}^{-1} g_{\sigma}) > 1$. Transforming $G \rightarrow \frac{1}{b-a}(G - aI)$ by a shift to the diagonal elements and a rescaling guarantees the constraint (27) is satisfied. Here, I is the identity matrix.

While the initial chemical potentials μ_j are uniquely determined by G and β as shown in Eq. (25), we have many choices for J_{ij} . Essentially, the only constraint on J is through its eigenvectors, and we can choose the eigenvalues at will as long as they are nondegenerate. The absence of degeneracy is essential for our method. For example., in the extreme case where all eigenvalues d_{α} are degenerate, the matrix J becomes proportional to the identity matrix, which will not change the initial Green's functions. The hopping matrix J is also entirely local in the limit of all degenerate eigenvalues. There is indeed competition between locality and equilibration time. To speed up the emergence of the steady-state, we spread out these eigenvalues and make the gaps $d_{\alpha} - d_{\sigma}$ as large as possible.

The timescale for the equilibration goes as

$$t_{\text{eq}} = \frac{1}{[\min(d_{\alpha} - d_{\sigma})]^2}, \quad (28)$$

according to Eq. (5). Of course, we can arbitrarily shorten the timescale if we allow the energy scale of the Hamiltonian to




FIG. 2. (a) The elements of the drive Hamiltonian J . The largest elements are local in the vicinity of the diagonal, but J exhibits non-local power-law decay in off-diagonal elements. (b) The elements of J decay with the same ν exponent (in this example $\nu = 1.5$ as the two-point function).

run to infinity. To find an optimal J with fixed energy scale, we set the spectral norm of J , i.e., the square root of the largest eigenvalue of the matrix $J^{\dagger}J$ to unity, $\|J\| = 1$, which implies $-1 \leq d_{\alpha} \leq 1$. To minimize t_{eq} , we can make all the gaps identical by choosing

$$d_{\alpha} = -1 + 2 \frac{\alpha - 1}{N - 1}, \quad \alpha = 1, \dots, N. \quad (29)$$

While using all permutations of the above values of d_{α} give the same equilibration time, we have found that the smoothest site-dependence in matrix J corresponds to the case where eigenvectors of G are ordered according to their eigenvalues, i.e., $g_{\alpha} \leq g_{\alpha+1}$.

IV. ENGINEERING POWER-LAW TWO-POINT FUNCTIONS

We now consider an example of creating off-diagonal Green's functions

$$G_{ij} = \frac{c}{|i - j|^{\nu}}, \quad i \neq j, \quad (30)$$

for an arbitrary exponent ν . An example of a target G_{ij} for $i \neq j$ is shown in Fig. 1(a) for $\nu = 1.5$. We found that requiring a uniform average density gives a singular W matrix. Thus, to make the matrix W invertible, we choose random values for the diagonal elements G_{ii} (note that G_{ii} is the expectation value of the occupation number of fermionic mode and must be between 0 and 1 for any physical system).




FIG. 3. The time-dependent approach of the two-point function to the engineered steady-state values.

To apply our method, we find the eigenvalues and eigenvector of G , which in turn yield the initial chemical potentials according to Eq. (25). As long as the initial Fermi-Dirac functions $f(\mu_i)$, which only depend on the eigenvalues and eigenvectors of G , are between 0 and 1, we can find real chemical potential for any β . As an example, we show in Fig. 1(b) the $f(\mu_i)$ that yield the Green's functions shown in Fig. 1, together with the corresponding G_{ii} . The initial occupation numbers $f(\mu_i)$ are close to the target occupation numbers G_{ii} but not identical. The smallest difference for $i = 47$ is 0.13%, while the largest difference for $i = 28$ is 53%. We note that while the initial occupation numbers are redistributed in the final steady-state, the total charge is conserved, and we have $\sum_i G_{ii} = \sum_i f(\mu_i)$.

In Fig. 2, we show the elements of the J matrix of Eq. (26) obtained by eigenvalues (29). We find that the elements of J exhibit similar critical correlations as the correlations we seek to create. We examined many different permutations of the eigenvalues. While the J can become highly disordered, it appears that the correlation structure does not change from critical as long as we keep the eigenvalues separated for optimal

The long-time limit for $t \rightarrow \infty$ follows from Eq. (13) as shown below

$$\langle \hat{n}_i \hat{n}_j \rangle_{(\infty)} = \sum_{\alpha\beta\gamma\delta} \left[\sum_{ac} \mathcal{U}_{\alpha a} \mathcal{U}_{\gamma c} \mathcal{U}_c^\dagger \mathcal{U}_{ai}^\dagger (\mathcal{U}_{ia} \mathcal{U}_{jc} \mathcal{U}_{cd}^\dagger \mathcal{U}_{\alpha\beta}^\dagger + \mathcal{U}_{ic} \mathcal{U}_{ja} \mathcal{U}_{\alpha\delta}^\dagger \mathcal{U}_{c\beta}^\dagger) - \sum_a \mathcal{U}_{\alpha a} \mathcal{U}_{\gamma a} \mathcal{U}_{aj}^\dagger \mathcal{U}_{ai}^\dagger \mathcal{U}_{ia} \mathcal{U}_{ja} \mathcal{U}_{\alpha\delta}^\dagger \mathcal{U}_{\alpha\beta}^\dagger \right] \langle c_\alpha^\dagger c_\beta c_\gamma^\dagger c_\delta \rangle_{(0)}. \quad (33)$$

It is convenient to define column vectors $F = (f(\mu_1), f(\mu_2), \dots, f(\mu_N))$, $K = (1 - f(\mu_1), 1 - f(\mu_2), \dots, 1 - f(\mu_N))$, and matrices $A^{ij} = \mathcal{U} \text{diag}(\mathcal{U}_{i,1} \mathcal{U}_{1,j}^\dagger, \mathcal{U}_{i,2} \mathcal{U}_{2,j}^\dagger, \dots, \mathcal{U}_{i,N} \mathcal{U}_{N,j}^\dagger) \mathcal{W}^\dagger$, where $f(\mu_\alpha)$ is defined in Eq. (16). Inserting Eq. (32) into the above expression, upon some matrix algebra, leads to

$$\langle \hat{n}_i \hat{n}_j \rangle_{(\infty)} = \sum_{\alpha\beta} \left\{ [A_{\alpha\beta}^{ii} A_{\beta\alpha}^{jj} + A_{\alpha\alpha}^{jj} A_{\beta\beta}^{ii}] F_\alpha K_\beta + [A_{\alpha\alpha}^{ii} A_{\beta\beta}^{jj} + A_{\alpha\beta}^{jj} A_{\beta\alpha}^{ii}] F_\alpha F_\beta - W_{\beta,i} W_{\beta\alpha} F_\alpha W_{\beta,j} \right\} \quad (34)$$

where the W matrices are defined in Eq. (24). The subtraction of the background $\langle \hat{n}_i \rangle \langle \hat{n}_j \rangle$ is subtle. It might appear that each $\langle \hat{n}_i \rangle$ in the long-time limit is simply a diagonal element of G . However, that is only correct if we first take the noise average of each expectation value of density and multiply the noise-averaged results. It is more natural to take the quantum averages of density for various realizations of noise first, multiply them, and then take the noise average, keeping the

equilibration.

It is illuminating to examine the time dependence of the two-point function as it approaches the engineered critical correlations. They start from zero and quickly acquire a noisy pattern aligned with the overall power-law dependence. The evolution continuously shifts the correlators up in a log-log plot while reducing the noise until it converges to the target disorder-free power law. The exponent of the critical correlation emerges much faster than the quantitative emergence of those correlation functions.

V. DENSITY-DENSITY CORRELATIONS IN THE STEADY STATE

To understand the nature of the steady-state we have generated, we compute the density-density correlation functions, as the two-point function does not fully characterize the steady-state. A widely studied class of correlation functions is the connected density-density correlator

$$R_{ij}(t) = \langle \hat{n}_i \hat{n}_j \rangle - \langle \hat{n}_i \rangle \langle \hat{n}_j \rangle, \quad (31)$$

We treat the two terms in the above expression separately. To find $\langle \hat{n}_i \hat{n}_j \rangle$, we use the formalism of Sec. II to obtain the long-time limit of the rank-4 tensor \mathcal{R} for the density-density operator. We can then obtain the time dependence as well as the long-time limit of the $\langle \hat{n}_i \hat{n}_j \rangle$ correlator in terms of the initial 4-point-function expectation values $\langle c_\alpha^\dagger c_\beta c_\gamma^\dagger c_\delta \rangle_{(0)}$ at $t = 0$, which using the Wick's theorem is given by

$$\langle c_\alpha^\dagger c_\beta c_\gamma^\dagger c_\delta \rangle_{(0)} = \delta_{\beta\gamma} \delta_{\alpha\delta} [1 - f(\mu_\beta)] f(\mu_\alpha) + \delta_{\alpha\beta} \delta_{\gamma\delta} f(\mu_\alpha) f(\mu_\gamma). \quad (32)$$

noise averaging consistently in the last step. The long-time limit can be obtained by time-averaging the product of expectation values upon a quantum quench to Hamiltonian V . In this approach, the background term yields two terms that are identical to terms in $\langle \hat{n}_i \hat{n}_j \rangle_{(\infty)}$. We find

$$\langle \hat{n}_i \rangle \langle \hat{n}_j \rangle = \sum_{\alpha\beta} [A_{\alpha\alpha}^{ii} A_{\beta\beta}^{jj} + A_{\alpha\beta}^{jj} A_{\beta\alpha}^{ii}] F_\alpha F_\beta - \sum_\alpha W_{i,\alpha} W_{j,\alpha} (\bar{F}_\alpha)^2,$$




FIG. 4. The density-density correlation functions in the steady-state with a $j^{-1.5}$ Green's function. Despite strong disorder, the correlation decays with a trendline of j^{-3} .

where the matrix $\tilde{F} = \mathcal{U}^\dagger \text{diag}(F_1, \dots, F_N) \mathcal{U}$. Using these expressions we calculated the long-time limit of R_{ij} for the same drive that produces power-law two-point functions of Eq. (30). While the long-time limit of G has no disorder in the off-diagonal terms, the density-density correlation functions are highly disordered. A universality class for one-dimensional quantum systems with power-law correlators is the Luttinger liquid, in which $R_{i,j}$ decays as $|j - i|^{-2}$. Despite the strong disorder, we see that our $R_{i,j}$ does not trend as $|j - i|^{-2}$. The minima in $|R_{i,j}|$ decay as $|j - i|^{-3}$ for $\nu = 1.5$. It seems that the steady states emerging upon engineering clean $x^{-\nu}$ two-point functions have highly disordered density-density four-

point functions with a trendline scaling as $x^{-2\nu}$.

VI. CONCLUSIONS

In this paper, we extended the Heisenberg-picture formalism of stochastic driving to the case of fermionic four-point functions. We then proposed a scheme of reverse-engineering the drive to generate the desired matrix of fermionic two-point functions from a completely uncorrelated initial thermal state. Finally, we examined the case of generating critical correlations and found that the steady-state exhibits clean engineered two-point functions and highly disordered density-density correlation functions. Such steady states with custom-ordered correlations provide novel examples of drive engineering where the system evades the generic infinite-temperature fixed point [29] in the long-time limit.

VII. ACKNOWLEDGMENTS

I thank Zohar Nussinov for helpful discussions. I acknowledge support from NSF Award DMR-1945395 and thank the Kavli Institute for Theoretical Physics for hospitality, acknowledging support by the NSF under Grant PHY1748958.

-
- [1] I. M. Georgescu, S. Ashhab, and F. Nori, *Rev. Mod. Phys.* **86**, 153 (2014).
 - [2] J. D. Biamonte, V. Bergholm, J. D. Whitfield, J. Fitzsimons, and A. Aspuru-Guzik, *AIP Advances* **1**, 022126 (2011).
 - [3] E. Torrontegui, S. Ibáñez, S. Martínez-Garaot, M. Modugno, A. del Campo, D. Guéry-Odelin, A. Ruschhaupt, X. Chen, and J. G. Muga, in *Advances in Atomic, Molecular, and Optical Physics*, Advances In Atomic, Molecular, and Optical Physics, Vol. 62, edited by E. Arimondo, P. R. Berman, and C. C. Lin (Academic Press, 2013) pp. 117 – 169.
 - [4] J. Werschnik and E. K. U. Gross, *Journal of Physics B: Atomic, Molecular and Optical Physics* **40**, R175 (2007).
 - [5] N. Goldman and J. Dalibard, *Phys. Rev. X* **4**, 031027 (2014).
 - [6] T. Oka and S. Kitamura, *Annual Review of Condensed Matter Physics* **10**, 387 (2019).
 - [7] T. Qin and W. Hofstetter, *Phys. Rev. B* **97**, 125115 (2018).
 - [8] B. Kraus, H. P. Büchler, S. Diehl, A. Kantian, A. Micheli, and P. Zoller, *Phys. Rev. A* **78**, 042307 (2008).
 - [9] S. Diehl, A. Micheli, A. Kantian, B. Kraus, H. P. Büchler, and P. Zoller, *Nat. Phys.* **4**, 878 (2008).
 - [10] D. M. Busiello, C. Jarzynski, and O. Raz, *New Journal of Physics* **20**, 093015 (2018).
 - [11] T. Grotz, L. Heaney, and W. T. Strunz, *Phys. Rev. A* **74**, 022102 (2006).
 - [12] J. Marino and A. Silva, *Phys. Rev. B* **86**, 060408 (2012).
 - [13] H. Pichler, J. Schachenmayer, A. J. Daley, and P. Zoller, *Phys. Rev. A* **87**, 033606 (2013).
 - [14] L. D'Alessio and A. Rahmani, *Phys. Rev. B* **87**, 174301 (2013).
 - [15] A. Chenu, M. Beau, J. Cao, and A. del Campo, *Phys. Rev. Lett.* **118**, 140403 (2017).
 - [16] A. Barchielli and M. Gregoratti, *Quantum Trajectories and Measurements in Continuous Time* (Springer Berlin, Heidelberg, 2009).
 - [17] T. Prosen, *New Journal of Physics* **10**, 043026 (2008).
 - [18] M. Žnidarič, *Journal of Statistical Mechanics: Theory and Experiment* **2010**, L05002 (2010).
 - [19] V. Eisler, *Journal of Statistical Mechanics: Theory and Experiment* **2011**, P06007 (2011).
 - [20] M. Žnidarič, *Phys. Rev. E* **83**, 011108 (2011).
 - [21] J. W. Britton, B. C. Sawyer, A. C. Keith, C.-C. J. Wang, J. K. Freericks, H. Uys, M. J. Biercuk, and J. J. Bollinger, *Nature* **484**, 489 (2012).
 - [22] P. Richerme, Z.-X. Gong, A. Lee, C. Senko, J. Smith, M. Foss-Feig, S. Michalakis, A. V. Gorshkov, and C. Monroe, *Nature* **511**, 198 (2014).
 - [23] P. Jurcevic, B. P. Lanyon, P. Hauke, C. Hempel, P. Zoller, R. Blatt, and C. F. Roos, *Nature* **511**, 202 (2014).
 - [24] A. Rahmani, *Phys. Rev. A* **92**, 042110 (2015).
 - [25] I. Klich, in *Quantum Noise in Mesoscopic Systems*, edited by Y. V. Nazarov (Kluwer, 2003).
 - [26] S. Ziraldo, A. Silva, and G. E. Santoro, *Phys. Rev. Lett.* **109**, 247205 (2012).
 - [27] A. Rahmani and S. Vishveshwara, *Phys. Rev. B* **97**, 245116 (2018).
 - [28] L. A. Peña Ardila, M. Heyl, and A. Eckardt, *Phys. Rev. Lett.* **121**, 260401 (2018).
 - [29] G. Bunin, L. D'Alessio, Y. Kafri, and A. Polkovnikov, *Nature Physics* **7**, 913 (2011).