NONLINEAR BSDES WITH TWO OPTIONAL DOOB'S CLASS BARRIERS SATISFYING WEAK MOKOBODZKI'S CONDITION AND EXTENDED DYNKIN GAMES

TOMASZ KLIMSIAK^{1,2} MAURYCY RZYMOWSKI²

¹ Institute of Mathematics, Polish Academy of Sciences, Śniadeckich 8, 00-656 Warsaw, Poland

² Faculty of Mathematics and Computer Science, Nicolaus Copernicus University, Chopina 12/18, 87-100 Toruń, Poland

ABSTRACT. We study reflected backward stochastic differential equations (RBSDEs) on the probability space equipped with a Brownian motion. The main novelty of the paper lies in the fact that we consider the following weak assumptions on the data: barriers are optional of class (D) satisfying weak Mokobodzki's condition, generator is continuous and non-increasing with respect to the value-variable (no restrictions on the growth) and Lipschitz continuous with respect to the control-variable, and the terminal condition and the generator at zero are supposed to be merely integrable. We prove that under these conditions on the data there exists a solution to corresponding RBSDE. In the second part of the paper, we apply the theory of RBSDEs to solve basic problems in Dynkin games driven by nonlinear expectation based on the generator mentioned above. We prove that the main component of a solution to RBSDE represents the value process in corresponding extended nonlinear Dynkin game. Moreover, we provide sufficient conditions on the barriers guaranteeing the existence of the value for nonlinear Dynkin games and the existence of a saddle point.

1. INTRODUCTION

Let *B* be a standard *d*-dimensional Brownian motion on a given probability space (Ω, \mathcal{F}, P) , *T* be a strictly positive real number (horizon time) and let $\mathbb{F} \coloneqq (\mathcal{F}_t)_{0 \le t \le T}$ be the standard augmentation of the filtration generated by *B*. In the present paper, we study Reflected Backward Stochastic Differential Equations (RBSDEs for short) of the following form

$$\begin{cases} Y_t = \xi + \int_t^T f(r, Y_r, Z_r) \, dr + R_T - R_t - \int_t^T Z_r \, dB_r, & t \in [0, T], \\ L_t \le Y_t \le U_t, & t \in [0, T], \end{cases}$$
(1.1)

where ξ (terminal value) is an \mathcal{F}_T -measurable random variable, the mapping $f: \Omega \times [0,T] \times \mathbb{R} \times \mathbb{R}^d \to \mathbb{R}$ (generator) is an \mathbb{F} -progressively measurable process with respect to the first two variables and L, U (barries) are \mathbb{F} -optional processes of class (D). We look for a triple (Y, Z, R) of \mathbb{F} -progressively measurable processes, with R of finite variation and $R_0 = 0$, that satisfies (1.1). Given a solution (Y, Z, R) to (1.1), we call the process Y the main part of the solution. The role of R is to keep Y between barriers L, U, and the role of Z is to keep

e-mail: tomas@mat.umk.pl

e-mail: maurycyrzymowski@mat.umk.pl

Y adapted to F. In order to get the uniqueness for problem (1.1) one requires from R to satisfy the so called *minimality condition* which stands that

$$\int_{0}^{T} (Y_{r-} - \limsup_{s\uparrow r} L_s) dR_r^{*,+} + \int_{0}^{T} (\liminf_{s\uparrow r} U_s - Y_{r-}) dR_r^{*,-} = 0$$

$$\sum_{0 \le r < T} (Y_r - L_r) \max\{R_{r+} - R_r, 0\} + \sum_{0 \le r < T} (U_r - Y_r) \max\{R_r - R_{r+}, 0\} = 0,$$
(1.2)

where R^* is the càdlàg part of R and $R^{*,+}$, $R^{*,-}$ its Jordan decomposition.

Formulation of the problems. In the paper, we merely assume that

- (A1) $\mathbb{E}|\xi| + \mathbb{E}\int_0^T |f(r,0,0)| dr < \infty$,
- (A2) there is $\lambda \ge 0$ such that $|f(t, y, z) f(t, y, z')| \le \lambda |z z'|$ for $t \in [0, T], y \in \mathbb{R}, z, z' \in \mathbb{R}^d$, (A3) $y \mapsto f(t, y, z)$ in non-increasing and continuous for fixed $t \in [0, T], z \in \mathbb{R}^d$,

- (A4) $\int_0^T |f(r, y, 0)| dr < \infty$ for every $y \in \mathbb{R}$, (Z) there exist $\gamma \ge 0, \ \kappa \in [0, 1)$ and a non-negative \mathbb{F} -progressively measurable process g, satisfying $\mathbb{E} \int_0^T g_r dr < \infty$, such that

$$|f(t, y, z) - f(t, y, 0)| \le \gamma (g_t + |y| + |z|)^{\kappa}, \quad t \in [0, T], \ y \in \mathbb{R}, \ z \in \mathbb{R}^d.$$

By [4], under (A1)-(A4),(Z), there exists a solution (Y,Z) to (1.1) without barriers (BSDE), i.e.

$$Y_t = \xi + \int_t^T f(r, Y_r, Z_r) \, dr - \int_t^T Z_r \, dB_r, \quad t \in [0, T].$$
(1.3)

Moreover, it is unique provided Y is of class (D) and $Z \in \mathcal{H}^s_{\mathbb{F}}(0,T)$ - the class of \mathbb{F} progressively measurable processes satisfying $\mathbb{E}[\int_0^T |Z_r|^2 dr]^{s/2} < \infty$ - for some $s > \kappa$. In the present paper, we focus on the existence and uniqueness problem for (1.1)–(1.2) under conditions (A1)-(A4),(Z). We shall also study the representation of the process Y as the value process in nonlinear Dynkin games.

The existence problem. First, observe that by the very definition of a solution to (1.1)its main part is a semimartingale. Consequently, we deduce at once, that the existence of a semimartingale between the barriers L, U is a necessary condition for the existence of a solution to (1.1) (intrinsic condition). The said condition is known in the literature as weak Mokobodzki's condition (see [19]):

(WM) there exists a semimartingale X such that $L_t \leq X_t \leq U_t, t \in [0, T]$.

It is a natural question whether under (A1)-(A4), (Z) the above condition is also sufficient for the existence of a solution to (1.1)-(1.2). We give a positive answer to this question, and prove even more, that condition (Z) can be dropped.

Theorem 1. Assume that (A1)–(A4) hold and weak Mokobodzki's condition (WM) is in force. Then there exist a solution (Y, Z, R) to (1.1)–(1.2).

It appears, and it may seem surprising at first, that the above result does not hold for BSDEs (1.3) (see Remark 3.5). The explanation of this phenomenon is that in the case of reflected BSDEs barriers keep the *main part* of a solution in the class (D). At this point it

is worth mentioning that the following condition (complete separation)

$$L, U$$
 are càdlàg, $L_t < U_t, t \in [0, T], L_{t-} < U_{t-}, t \in (0, T]$ (1.4)

implies (WM) (see [36, Lemma 3.1]). We provide a generalization of this condition, by dropping càdlàg regularity assumption on L, U, and we prove that the following condition implies (WM):

$$L, U$$
 are left-limited, $L_{t-} < U_{t-}$, $\limsup_{s \downarrow t} L_s < \liminf_{s \downarrow t} U_s$, $t \in [0, T]$. (1.5)

The uniqueness problem. An interesting issue is also the problem of the uniqueness for solutions to (1.1)-(1.2). In the proof of the uniqueness for BSDEs (1.3) (see [4] and Theorem 3.6) the crucial roles were played by condition (Z) and the fact that for any solution (Y,Z) to (1.3) we have, under conditions (A1)–(A4),(Z), that $Z \in \mathcal{H}^s_{\mathbb{R}}(0,T)$ for some $s > \kappa$ provided Y is of class (D). For reflected BSDEs this property does not hold even if $f \equiv 0$ (see [24, Example 5.6]). Nevertheless, we are able to prove the following result.

Theorem 2. Assume that (A1)-(A4),(Z) are in force. Then there exists at most one solution to RBSDE (1.1)-(1.2).

Solutions to RBSDEs as value processes in Dynkin games. The above theorem is a consequence of a much deeper result, which is our third main result of the paper. In order to formulate it, we use the notion of the nonlinear expectation introduced by Peng in [35]. For given stopping times $\alpha \leq \beta \leq T$ consider mapping

$$\mathbb{E}^{f}_{\alpha,\beta}: L^{1}(\mathcal{F}_{\beta}) \to L^{1}(\mathcal{F}_{\alpha}),$$

by letting $\mathbb{E}^{f}_{\alpha,\beta}\xi \coloneqq Y^{\beta}_{\alpha}$, where (Y^{β}, Z^{β}) is a (unique) solution to (1.3), with T replaced by β , such that Y^{β} is of class (D). For given stopping times $\tau, \sigma \leq T$ and sets $H \in \mathcal{F}_{\tau}, G \in \mathcal{F}_{\sigma}$, we let

$$J(\tau, H; \sigma, G) \coloneqq (L_{\tau} \mathbf{1}_{H} + \limsup_{h \ge 0} L_{\tau+h} \mathbf{1}_{H^{c}}) \mathbf{1}_{\{\tau \le \sigma < T\}} + (U_{\sigma} \mathbf{1}_{G} + \liminf_{h \ge 0} U_{\sigma+h} \mathbf{1}_{G^{c}}) \mathbf{1}_{\{\sigma < \tau\}} + \xi \mathbf{1}_{\{\tau = \sigma = T\}},$$

with the convention that $L_t = L_{t \wedge T}, U_t := U_{t \wedge T}, t \ge 0$. We prove the following representation theorem.

Theorem 3. Assume that (A1)–(A4), (Z) are in force. If (Y, Z, R) is a solution to RBSDE (1.1)–(1.2), then

$$Y_{\theta} = \underset{\sigma \geq \theta, G \in \mathcal{F}_{\sigma}}{\operatorname{ess \, sup}} \underset{\tau \geq \theta, H \in \mathcal{F}_{\tau}}{\operatorname{ess \, sup}} \mathbb{E}_{\theta, \tau \wedge \sigma}^{J} J(\tau, H; \sigma, G)$$
$$= \underset{\tau \geq \theta, H \in \mathcal{F}_{\tau}}{\operatorname{ess \, sup}} \underset{\sigma \geq \theta, G \in \mathcal{F}_{\sigma}}{\operatorname{ess \, sup}} \mathbb{E}_{\theta, \tau \wedge \sigma}^{f} J(\tau, H; \sigma, G)$$

for any stopping time $\theta \leq T$.

In other words, we show that Y is a value process in an extended nonlinear Dynkin game: "nonlinear" since we consider the nonlinear expectation, and "extended" since players may change payoffs L, U on sets H^c, G^c , respectively, which extends the set of their strategies (in the classical Dynkin games the players are not allowed to choose sets G, H). Observe that

3

the above extended nonlinear Dynkin game reduces to the nonlinear Dynkin game provided L and U are right-continuous. We prove however a stronger result.

Theorem 4. Assume that (A1)–(A4), (Z) are in force. Moreover, suppose that L is right upper semicontinuous and U is right lower semicontinuous. If (Y, Z, R) is a solution to RBSDE (1.1)–(1.2), then

$$Y_{\theta} = \operatorname{ess\,inf}_{\sigma \geq \theta} \operatorname{ess\,sup}_{\tau \geq \theta} \mathbb{E}^{J}_{\theta, \tau \wedge \sigma} J(\tau, \Omega; \sigma, \Omega)$$
$$= \operatorname{ess\,sup\,ess\,inf}_{\tau \geq \theta} \mathbb{E}^{f}_{\theta, \tau \wedge \sigma} J(\tau, \Omega; \sigma, \Omega).$$

for any stopping time $\theta \leq T$.

Thus, Y represents the value process in a nonlinear Dynkin game provided L, U are sufficiently regular as mentioned above. The above result was achieved by Bayraktar and Yao in [3] for continuous barriers L, U satisfying (1.4) and under the following additional conditions: $\mathbb{E} \sup_{t \leq T} |L_t| + \mathbb{E} \sup_{t \leq T} |U_t| < \infty$, generator f admits the linear growth with respect to Y-variable, i.e. $|f(t, y, 0)| \leq g_t + \psi |y|$ for some $\psi \geq 0$. Note that in the present paper growth of f with respect to Y-variable is subject to no restriction.

Finally, we show that further regularity assumptions on barriers L, U allow one to indicate saddle points for nonlinear Dynkin games. For any stopping time $\theta \leq T$ set:

$$\tau_{\theta}^* \coloneqq \inf\{t \ge \theta, Y_t = L_t\} \land T; \quad \sigma_{\theta}^* \coloneqq \inf\{t \ge \theta, Y_t = U_t\} \land T \tag{1.6}$$

and

$$\bar{\tau}_{\theta} \coloneqq \inf\{t \ge \theta, R_t^+ > R_{\theta}^+\} \wedge T; \quad \bar{\sigma}_{\theta} \coloneqq \inf\{t \ge \theta, R_t^- > R_{\theta}^-\} \wedge T.$$

$$(1.7)$$

Theorem 5. Assume that (A1)-(A4), (Z) are in force. Moreover, suppose that L is upper semicontinuous and U is lower semicontinuous. Then

$$\mathbb{E}^{J}_{\theta,\tau_{\theta}^{*}\wedge\sigma_{\theta}^{*}}J(\tau_{\theta}^{*},\Omega;\sigma_{\theta}^{*},\Omega) = \mathbb{E}^{J}_{\theta,\bar{\tau}_{\theta}\wedge\bar{\sigma}_{\theta}}J(\bar{\tau}_{\theta},\Omega;\bar{\sigma}_{\theta},\Omega)$$
$$= \operatorname{ess\,inf}_{\sigma\geq\theta}\operatorname{ess\,sup}_{\tau\geq\theta}\mathbb{E}^{J}_{\theta,\tau\wedge\sigma}J(\tau,\Omega;\sigma,\Omega).$$

for any stopping time $\theta \leq T$.

Proof techniques and relations of main results to the existing literature. First, note that Mokobodzki's condition, additionally to (WM), requires from the semimartingale X, lying between the barriers, some integrability of its finite variation and martingale part (depending on the authors you may find different integrability conditions for the process X, nonetheless it is always assumed to be at least the difference of positive supermartingales). This additional requirement is the reason why the complete separation condition (1.4) does not imply Mokobodzki's condition. The fact that (1.4) implies weak Mokobodzki's condition is an easy calculation and may be found e.g. in [36] and [19] (in the case of continuous barriers). Reflected BSDEs with condition (1.4) imposed on the barriers have been considered in many papers (see [3, 5, 12, 18, 20, 21, 22, 36]). In the papers [5], [18] and [20], L^2 -data and sublinear growth of the generator with respect to Y-variable are required. In [21] authors considered bounded data and continuous generator with quadratic-growth with respect to Z-variable. Some results for RBSDEs with generators subject to sublinear growth with

respect to Y-variable are described also in [3, 22] (L^1 -data) and [12] (L^p -data, $p \in (1, 2)$). L^1 -data and generator being merely monotone and continuous with respect to Y-variable were considered in [36].

In all the mentioned papers (besides [21]), the method of local solutions and pasting local solutions, introduced by Hamadène and Hassani in [18], has been applied to achieve the existence for underlying RBSDEs. This method is rather complicated, and this is perhaps the reason why the development of theory of RBSDEs with barriers satisfying complete separation condition is far from being satisfactory. The second drawback of the method is that it is based on the penalization scheme which is not available for RBSDEs with optional barriers. In [23], the author proposed a different method which applies to RBSDEs with barriers satisfying even more general than (1.4) weak Mokobodzki's condition (WM). We call this method localization procedure. The advantage of the method is its simplicity and wide applicability. The method is based on the following simple observation: for any chain (τ_k) , i.e. non-decreasing sequence of stopping times satisfying

$$P(\tau_k < T, k \ge 1) = 0,$$

we have

$$(Y, Z, R)$$
 solves $\operatorname{RBSDE}^T(\xi, f, L, U)$
iff
 (Y, Z, R) solves $\operatorname{RBSDE}^{\tau_k}(Y_{\tau_k}, f, L, U), \ k \ge 1.$

The method consists of finding a proper regular approximation (Y^n) , on the whole interval [0,T], of a potential solution Y of a given problem (by "proper" we mean an approximation which does not blow up when passing to the limit). The terms of approximating sequence may solve BSDEs or RBSDEs of the generic form

$$Y_t^n = \xi_n + \int_t^T f_n(r, Y_r^n, Z_r^n) \, dr + R_T^n - R_t^n - \int_t^T Z_r^n \, dB_r \quad t \in [0, T],$$

with suitable chosen ξ_n, f_n, R^n . In the first step one shows that (Y^n) converges to a process Y. After that, we show that Y is the main part of a solution to $\text{RBSDE}^{\tau_k}(Y_{\tau_k}, f, L, U)$ for each $k \ge 1$. Since (τ_k) is a *chain*, we conclude that Y is the main part of a solution to $\text{RBSDE}^T(\xi, f, L, U)$.

By using *localization procedure* in [23], the first author of the present paper was able to provide an existence result for RBSDEs with merely càdlàg barriers of class (D) satisfying (WM), L^1 -data, and generator being continuous and non-increasing with respect to Y-variable (with no restrictions on the growth of the generator with respect to Y-variable).

As far as we know the only papers in the literature concerned with RBSDEs of the form (1.1) with non-càdlàg barriers satisfying (WM) are [26, 32]. In [26] RBSDEs on a general filtered space are studied under (WM) but with f independent of Z-variable. In The generalization of (1.4) to the case of làdlàg barriers was presented in [32] the authors considered làdlàg barriers and stochastic Lipschitz generator f (on the Brownian-Poisson filtration).

As to the nonlinear Dynkin games, to the best of our knowledge, there are only few papers in the literature: [9, 10, 11, 15, 17] - all with L^2 -data and Lipschitz generator - and [25, 26] - with L^1 -data and continuous and monotone generator with respect to Y-variable and independent of Z-variable.

Comments on the related literature. Reflected backward stochastic differential equations with two barriers have been introduced by Cvitanić and Karatzas in [7] as a generalization of backward stochastic differential equations introduced by Pardoux and Peng in [33] (analogous results for one reflecting barrier, i.e. in case $U \equiv \infty$, have been presented for the first time by El Karoui et al. in [13]). In [7] the authors considered (1.1) with barriers being continuous processes satisfying *Mokobodzki's condition* i.e. there exists $L_t \leq X_t \leq U_t$, $t \in [0,T]$, such that $X = X^1 - X^2$ and X^i is a positive supermartingale satisfying $\mathbb{E}\sup_{t\leq T} |X^i|_t^2 < \infty$, i = 1, 2. Moreover, they assumed that data are L^2 -integrable (i.e. $\sup_{t\leq T} |L_t|, \sup_{t\leq T} |U_t|, |\xi|, \int_0^T |f(r,0,0)| dr$ have second moments) and f is Lipschitz continuous with respect to (Y, Z)-variable (uniformly in (ω, t)). Under these assumptions a solution to (1.1) has been defined in [7] as a triple (Y, Z, R) of \mathbb{F} -progressively measurable processes such that Y is continuous, and R is a continuous finite variation process, with $R_0 = 0$, satisfying the minimality condition of the form

$$\int_0^T (Y_r - L_r) \, dR_r^+ = \int_0^T (U_r - Y_r) \, dR_r^- = 0,$$

where $R = R^+ - R^-$ is the Jordan decomposition of R. Observe that with continuous Y, L, U, R, condition (1.2) reduces to the above condition.

BSDEs and Reflected BSDEs are of great interest to scientists because of their numerous applications in various fields of mathematics and problems (e.g. partial differential equations, integro-differential equations, variational inequalities, optimization theory, control theory, mathematical finance etc., see [6, 34, 37] and the references therein). Over the past two decades, many interesting results have been obtained regarding RSBDEs. In particular, numerous existence results for RBSDEs, which strengthen the result of [7] by weakening assumptions on generator f, filtration \mathbb{F} , barriers L, U and horizon time T, have been provided.

Despite of intensive research, until 2016, only RBSDEs with càdlàg barriers were considered in the literature. With the work by Grigorova et al. in [14] there was a change in this regard and papers on less regular barriers began to appear. Equations of that type with L^2 -data and Lipschitz generator were studied in [31] (Brownian filtration), in [14, 15] (Brownian-Poisson filtration) and [1, 2, 16] (general filtration). RBSDEs with optional barriers and L^1 -data were considered only in [27, 28], in the case of Brownian filtration. Results on optional barriers, L^1 -data and possibly infinite horizon time were presented in [26] but with f independent of Z-variable. The case of L^2 -data and f being stochastic Lipschitz driver was presented in [30] (Brownian-Poisson filtration) and in [29, 32] (general filtration).

2. Basic notation

We say that a function $y: [0,T] \to \mathbb{R}^d$ is regulated on [0,T] if for any $t \in [0,T)$, there exists the limit $y_{t+} \coloneqq \lim_{u \downarrow t} y_u$ and for any $s \in (0,T]$ there exists the limit $y_{s-} \coloneqq \lim_{u \uparrow s} y_u$. For any regulated function y on [0,T] we define $\Delta^+ y_t \coloneqq y_{t+} - y_t$, $t \in [0,T)$ and $\Delta^- y_s \coloneqq y_s - y_{s-}$, $s \in (0,T]$.

For $x \in \mathbb{R}^d$ by |x| we denote the euclidean norm. As mentioned in Section 1, \mathcal{T} stands for the set of all stopping times taking values in [0, T]. What is more, for $\alpha, \beta \in \mathcal{T}, \mathcal{T}_{\alpha,\beta} \coloneqq \{\tau \in \mathcal{T}, \alpha \leq \tau \leq \beta\}, \mathcal{T}_{\alpha} \coloneqq \mathcal{T}_{\alpha,T}, \mathcal{T}^{\beta} \coloneqq \mathcal{T}_{0,\beta}$.

7

Let $\alpha, \beta \in \mathcal{T}, \alpha \leq \beta$, and $p \geq 1$. By $\mathcal{S}^p_{\mathbb{F}}(\alpha, \beta)$ we denote all \mathbb{F} -progessively measurable, \mathbb{R} -valued processes $Y = (Y_t)_{t \in [0,T]}$ such that

$$||Y||_{\mathcal{S}^p_{\mathbb{F}}(\alpha,\beta)} \coloneqq \left(\mathbb{E}\sup_{\alpha \le t \le \beta} |Y_t|^p\right)^{\frac{1}{p}} < \infty.$$

 $\mathcal{M}_{loc}(\alpha,\beta)$ is the space of all \mathbb{F} -local martingales on $[[\alpha,\beta]]$. Let $q \ge 1$. By $L^{p,q}_{\mathbb{F}}(\alpha,\beta)$ we denote the set of all \mathbb{F} -progressively measurable, \mathbb{R} -valued processes $X = (X_t)_{t \in [0,T]}$ such that

$$||X||_{L^{p,q}_{\mathbb{F}}(\alpha,\beta)} \coloneqq \left(\mathbb{E} \left(\int_{\alpha}^{\beta} |X_r|^p \, dr \right)^{\frac{q}{p}} \right)^{\frac{q}{p}} < \infty.$$

 $L^p_{\mathbb{F}}(\alpha,\beta)$ is a shorthand for $L^{p,p}_{\mathbb{F}}(\alpha,\beta)$.

Let $\mathcal{G} \subset \mathcal{F}$. $L^p(\mathcal{G})$ is the set of all \mathcal{G} -measurable random variables X such that

$$||X||_{L^p} \coloneqq \left(\mathbb{E}|X|^p\right)^{\frac{1}{p}} < \infty.$$

By $\mathcal{H}_{\mathbb{F}}(\alpha,\beta)$, we denote the space of all \mathbb{F} -progensively measurable, \mathbb{R}^d -valued processes $Z = (Z_t)_{t \in [0,T]}$ such that

$$\int_{\alpha}^{\beta} |Z_r|^2 \, dr < \infty \quad P\text{-a.s.}$$

 $\mathcal{H}^s_{\mathbb{F}}(\alpha,\beta), s > 0$, is a subspace of $\mathcal{H}_{\mathbb{F}}(\alpha,\beta)$ consisting of Z satisfying

$$\mathbb{E}\Big(\int_{\alpha}^{\beta} |Z_r|^2 \, dr\Big)^{\frac{s}{2}} < \infty.$$

We say that \mathbb{F} -progessively measurable process $X = (X_t)_{t \in [0,T]}$ is of class (D) on $[[\alpha, \beta]]$ if the family $\{X_{\tau}, \tau \in \mathcal{T}_{\alpha,\beta}\}$ is uniformly integrable. By $\mathcal{D}_{\mathbb{F}}^2(\alpha,\beta)$ we denote the set of all \mathbb{F} -progressively measurable, \mathbb{R} -valued processes $Y = (Y_t)_{t \in [0,T]}$ such that $|Y|^2$ is of class (D) on $[[\alpha, \beta]]$. We equip $\mathcal{D}_{\mathbb{F}}^2(\alpha, \beta)$ with the norm

$$||Y||_{\mathcal{D}^2(\alpha,\beta)} \coloneqq \Big(\sup_{\sigma \in \mathcal{T}_{\alpha,\beta}} \mathbb{E}|Y_{\sigma}|^2\Big)^{\frac{1}{2}}.$$

A sequence $(\tau_k)_{k\geq 1} \subset \mathcal{T}_{\alpha,\beta}$ is called a chain on $[[\alpha,\beta]]$ if

$$\forall_{\omega \in \Omega} \exists_{n \in \mathbb{N}} \forall_{k \ge n} \tau_k(\omega) = \beta(\omega).$$

By $\mathcal{V}_{\mathbb{F}}(\alpha,\beta)$ (resp. $\mathcal{V}_{\mathbb{F}}^{+}(\alpha,\beta)$) we denote a space of \mathbb{F} -progensively measurable, \mathbb{R} -valued processes $V = (V_t)_{t \in [0,T]}$ with finite variation (resp. nondecreasing) on $[[\alpha,\beta]]$ and $\mathcal{V}_{0,\mathbb{F}}(\alpha,\beta)$ (resp. $\mathcal{V}_{0,\mathbb{F}}^{+}(\alpha,\beta)$) is a subspace of $\mathcal{V}_{\mathbb{F}}(\alpha,\beta)$ (resp. $\mathcal{V}_{\mathbb{F}}^{+}(\alpha,\beta)$) consisting of processes V such that $V_{\alpha} = 0$. $\mathcal{V}_{\mathbb{F}}^{p}(\alpha,\beta)$ (resp. $\mathcal{V}_{\mathbb{F}}^{+,p}(\alpha,\beta)$) is the set of all $V \in \mathcal{V}_{\mathbb{F}}(\alpha,\beta)$ (resp. $V \in \mathcal{V}_{\mathbb{F}}^{+}(\alpha,\beta)$) such that $E|V|_{\alpha,\beta}^{p} < \infty$, where $|V|_{\alpha,\beta}$ denotes the total variation of V on $[[\alpha,\beta]]$.

Let $V \in \mathcal{V}_{\mathbb{F}}(0,T)$. By V^* we denote the càdlàg part of the process V, i.e.

$$V_t^* = V_t - \sum_{0 \le r < t} \Delta^+ V_r.$$

Throughout the paper all relations between random variables are supposed to hold *P*-a.s. For processes $X^1 = (X_t^1)_{t \in [0,T]}$ and $X^2 = (X_t^2)_{t \in [0,T]}$ we write $X^1 \leq X^2$ if $X_t^1 \leq X_t^2$, $t \in [0,T]$, *P*-a.s.

Let $V^1, V^2 \in \mathcal{V}_{0,\mathbb{F}}(0,\tau)$. We write $dV^1 \leq dV^2$, if $dV^{1,*} \leq dV^{2,*}$ and $\Delta^+ V^1 \leq \Delta^+ V^2$ on $[0, \tau].$

For an \mathbb{F} -optional process $X = (X_t)_{t \in [0,T]}$ we set $\overrightarrow{X}_s = \limsup_{r \uparrow s} X_r$, $\underline{X}_s = \liminf_{r \uparrow s} X_r$, $s \in (0,T]$ and $\overleftarrow{X}_s = \limsup_{r \downarrow s} X_r, \ \underline{X}_s = \liminf_{r \downarrow s} X_r, \ s \in [0,T], \ s \in [0,T].$

3. BACKWARD SDES

Let $p \ge 1$. We shall need the following hypotheses:

- (H1) there is $\lambda \ge 0$ such that $|f(t, y, z) f(t, y, z')| \le \lambda |z z'|$ for $t \in [0, T]$, $y \in \mathbb{R}$, $z, z' \in \mathbb{R}^d$, (H2) there is $\mu \in \mathbb{R}$ such that $(y y')(f(t, y, z) f(t, y', z)) \le \mu (y y')^2$ for $t \in [0, T]$,
- $y, y' \in \mathbb{R}, z \in \mathbb{R}^d$
- (H3) for every $(t, z) \in [0, T] \times \mathbb{R}^d$ the mapping $\mathbb{R} \ni y \to f(t, y, z)$ is continuous,
- (H4) $\int_0^T |f(r, y, 0)| dr < \infty \text{ for every } y \in \mathbb{R},$ (H5) $\xi \in L^p(\mathcal{F}_T), f(\cdot, 0, 0) \in L^{1,p}_{\mathbb{F}}(0, T),$

Let $\alpha, \beta \in \mathcal{T}, \alpha \leq \beta$, and $\hat{\xi} \in \mathcal{F}_{\beta}$.

Definition 3.1. We say that a pair (Y, Z) of \mathbb{F} -adapted processes is a solution to backward stochastic differential equation on the interval $[[\alpha, \beta]]$ with right-hand side f and terminal value $\hat{\xi}$ (BSDE^{α,β}($\hat{\xi}, f$) for short) if

- (a) Y is a continuous process and $Z \in \mathcal{H}_{\mathbb{F}}(\alpha, \beta)$,
- (b) $\int_{\alpha}^{\beta} |f(r, Y_r, Z_r)| dr < \infty$,
- (c) $Y_t = \hat{\xi} + \int_t^\beta f(r, Y_r, Z_r) dr \int_t^\beta Z_r dB_r, t \in [\alpha, \beta].$

Let $V \in \mathcal{V}_{0,\mathbb{F}}(\alpha,\beta)$.

Definition 3.2. We say that a pair (Y, Z) of \mathbb{F} -adapted processes is a solution to backward stochastic differential equation on the interval $[[\alpha, \beta]]$ with right-hand side f + dV and terminal value $\hat{\xi}$ (BSDE^{α,β}($\hat{\xi}, f + dV$) for short) if (Y - V, Z) is a solution to BSDE^{α,β}($\hat{\xi}, f_V$), where $f_V(t, y, z) = f(t, y + V_t, z)$.

Let us adopt the shorthand $BSDE^{\beta} := BSDE^{0,\beta}$.

The following results follow from [4, Proposition 3.2, Theorem 4.2].

Theorem 3.3. Assume that (H1)–(H4) are in force. Suppose that (H5) holds with p > 1. Then the following assertions hold.

- (i) There exists a solution $(Y, Z) \in \mathcal{S}^p_{\mathbb{R}}(0, T) \times \mathcal{H}^p_{\mathbb{R}}(0, T)$ to $BSDE^T(\xi, f)$.
- (ii) There exists at most one solution (Y,Z) to $BSDE^T(\xi, f)$ such that $Y \in \mathcal{S}^p_{\mathbb{R}}(0,T)$.

Proposition 3.4. Assume that (H5), with p > 1, and (H1),(H2) are satisfied. Let (Y, Z) be a solution to $BSDE^{T}(\xi, f)$ such that $Y \in \mathcal{S}_{\mathbb{F}}^{p}(0,T)$. Then there exists c > 0, depending only on μ, λ, T, p , such that

$$\mathbb{E}\Big[\sup_{0\leq t\leq T}|Y_t|^p + \Big(\int_0^T |Z_r|^2 \, dr\Big)^{\frac{p}{2}} + \Big(\int_0^T |f(r, Y_r, Z_r)| \, dr\Big)^p\Big] \\ \leq c \mathbb{E}\Big[|\xi|^p + \Big(\int_0^T |f(r, 0, 0)| \, dr\Big)^p\Big].$$
(3.1)

In case (H5) is satisfied with p = 1, we shall need for the existence and uniqueness of solutions to BSDEs additional hypothesis.

(Z) There exists an \mathbb{F} -progressively measurable process $g \in L^1_{\mathbb{F}}(0,T)$ and $\gamma \ge 0$, $\kappa \in [0,1)$ such that

$$|f(t, y, z) - f(t, y, 0)| \le \gamma (g_t + |y| + |z|)^{\kappa}, \quad t \in [0, T], \ y \in \mathbb{R}, \ z \in \mathbb{R}^d.$$

Remark 3.5. Condition (Z) says that driver f is allowed to have at most sublinear growth with respect to z-variable. A typical example of a driver satisfying (H1)–(H5), (Z) is of the following form

$$f(t, y, z) \coloneqq f_0(t, y) + b(y)(1 + |z|)^{\kappa},$$

where f_0 satisfies (H2)–(H5), b is continuous, non-increasing and bounded, and $\kappa \in (0, 1)$.

Observe that, in general, under merely (H1)–(H4) and (H5) with p = 1, we cannot expect the existence of a solution (Y, Z) to $BSDE^{T}(\xi, f)$ with positive ξ such that Y is positive and of class (D). Indeed, assume that (Y, Z) is a solution to the following BSDE

$$Y_{t} = \xi + \int_{t}^{T} Z_{r} dr - \int_{t}^{T} Z_{r} dB_{r}, \quad t \in [0, T],$$

with positive $\xi \in L^1(\mathcal{F}_T)$, and Y is positive of class (D). Let (τ_k) be chain such that $Y \in S^2(0, \tau_k), Z \in \mathcal{H}^2_{\mathbb{F}}(0, \tau_k), k \geq 1$. Then, by Itô's formula

$$Y_0 = \mathbb{E}\left[Y_{\tau_k} \exp\left(-\frac{\tau_k}{2} + B_{\tau_k}\right)\right]$$

Therefore, by applying Fatou's lemma, we find that

$$e^T Y_0 \ge \mathbb{E}[\xi \exp(B_T)].$$

If the above inequality was true for any positive $\xi \in L^1(\mathcal{F}_T)$, then $\exp(B_T)$ would be bounded, a contradiction.

Theorem 3.6. Assume that (H1)-(H4), (Z) are in force. Moreover, assume that (H5) is satisfied with p = 1. Then the following assertions hold.

- (i) There exists a solution (Y,Z) of $BSDE^T(\xi, f)$ such that Y is of class (D) and $Z \in \mathcal{H}^s_{\mathbb{F}}(0,T), s \in (0,1).$
- (ii) There exists at most one solution (Y,Z) to $BSDE^{T}(\xi, f)$ such that Y is of class (D).

Proof. The assertion (i) follows from [4, Theorem 6.3]. As to (ii), by [4, Theorem 6.2], there exists at most one solution (Y, Z) to $BSDE^T(\xi, f)$ such that Y is of class (D) and $Z \in \mathcal{H}^s_{\mathbb{F}}(0,T), s \in (0,1)$. So, it is enough to show that if (Y,Z) is a solution to $BSDE^T(\xi, f)$ such that Y is of class (D), then $Z \in \mathcal{H}^s_{\mathbb{F}}(0,T), s \in (0,1)$. This follows at once from [28, Remark 2.1] and [4, Lemma 3.1].

4. Reflected BSDEs with two optional barriers under Mokobodzki's condition

In this section we assume that processes L and U are merely \mathbb{F} -optional. Let $\alpha, \beta \in \mathcal{T}$, $\alpha \leq \beta$, and $\hat{\xi} \in \mathcal{F}_{\beta}$ such that $L_{\beta} \leq \hat{\xi} \leq U_{\beta}$.

9

Definition 4.1. We say that a triple (Y, Z, R) of \mathbb{F} -adapted processes is a solution to reflected backward stochastic differential equation on the interval $[[\alpha, \beta]]$ with right-hand side f, terminal value $\hat{\xi}$, lower barrier L and upper barrier U (RBSDE^{α,β}($\hat{\xi}, f, L, U$) for short) if

- (a) Y is a regulated process and $Z \in \mathcal{H}_{\mathbb{F}}(\alpha, \beta)$,
- (b) $R \in \mathcal{V}_{0,\mathbb{F}}(\alpha,\beta), L_t \leq Y_t \leq U_t, t \in [\alpha,\beta], \text{ and}$

$$\int_{\alpha}^{\beta} (Y_{r-} - \vec{L}_r) dR_r^{*,+} + \sum_{\alpha \le r < \beta} (Y_r - L_r) (\Delta^+ R_r)^+$$

=
$$\int_{\alpha}^{\beta} (\underline{U}_r - Y_{r-}) dR_r^{*,-} + \sum_{\alpha \le r < \beta} (U_r - Y_r) (\Delta^+ R_r)^- = 0,$$

- where $R^* = R^{*,+} R^{*,-}$ is the Jordan decomposition of R^* , (c) $\int_{\alpha}^{\beta} |f(r, Y_r, Z_r)| dr < \infty$, (d) $Y_t = \hat{\xi} + \int_t^{\beta} f(r, Y_r, Z_r) dr + R_{\beta} R_t \int_t^{\beta} Z_r dB_r$, $t \in [\alpha, \beta]$.

In what follows we refer to condition (b) as the *minimality condition*. Let us adopt the shorthand $RBSDE^{\beta} := RBSDE^{0,\beta}$.

We consider the following condition, which we call strong Mokobodzki's condition:

(H6) there exists a process $X \in \mathcal{M}_{loc}(0,T) + \mathcal{V}^p_{\mathbb{F}}(0,T)$ such that $L \leq X \leq U, X \in \mathcal{S}^p(0,T)$ and $f(\cdot, X, 0) \in L^{1,p}_{\mathbb{R}}(0, T)$.

Assume that $L_T \leq \xi \leq U_T$. The following result has been proven in [28, Proposition 3.2, Theorem 3.9].

Theorem 4.2. Let p > 1. Assume (H1)-(H5).

- (i) There exists at most one solution (Y, Z, R) to $\text{RBSDE}^T(\xi, f, L, U)$ such that $Y \in$ $\mathcal{S}^p_{\mathbb{F}}(0,T).$
- (ii) There exists a solution (Y, Z, R) to $\operatorname{RBSDE}^T(\xi, f, L, U)$ such that $Y \in \mathcal{S}^p_{\mathbb{F}}(0, T)$, $Z \in \mathcal{H}^p_{\mathbb{F}}(0,T)$ and $R \in \mathcal{V}^p_{0,\mathbb{F}}(0,T)$ if and only if (H6) holds.

In the case of p = 1, we consider the following version of strong Mokobodzki's condition:

(H6*) there exists a process $X \in \mathcal{M}_{loc}(0,T) + \mathcal{V}^1_{\mathbb{F}}(0,T)$ such that X is of class (D), $L \leq X \leq U$ and $f(\cdot, X, 0) \in L^{1}_{\mathbb{F}}(0, T)$.

The following result has been proven in [28, Theorem 3.8].

Theorem 4.3. Let p = 1. Assume (H1)-(H5), (H6^{*}), (Z). Then there exists a unique solution (Y, Z, R) of $\operatorname{RBSDE}^T(\xi, f, L, U)$ such that Y is of class (D), $Z \in \mathcal{H}^q_{\mathbb{F}}(0, T), q \in (0, 1)$ and $R \in \mathcal{V}^1_{0,\mathbb{F}}(0,T)$.

5. Nonlinear expectation

Let $p \ge 1$. Throughout this section, we assume that either p = 1 and (H1)-(H5), (Z) are in force or p > 1 and (H1)-(H5) are in force. Let $\alpha, \beta \in \mathcal{T}, \alpha \leq \beta$. We define the operator

$$\mathbb{E}_{\alpha,\beta}^{(1),f}:L^1(\mathcal{F}_\beta)\to L^1(\mathcal{F}_\alpha),$$

by letting $\mathbb{E}_{\alpha,\beta}^{(1),f}(\xi) \coloneqq Y_{\alpha}$, where (Y,Z) is a solution to $BSDE^{\beta}(\xi, f)$ such that Y is of class (D). By Theorem 3.6, the operator $\mathbb{E}_{\alpha,\beta}^{(1),f}$ is well defined under conditions (H1)-(H5), (Z). By Theorem 3.3, under (H1)-(H5) (with p > 1), we may define the operator

$$\mathbb{E}_{\alpha,\beta}^{(p),f}: L^p(\mathcal{F}_\beta) \to L^p(\mathcal{F}_\alpha),$$

with $\mathbb{E}_{\alpha,\beta}^{(p),f}(\xi) \coloneqq Y_{\alpha}$, where (Y,Z) is a solution to $BSDE^{\beta}(\xi,f)$ such that $Y \in \mathcal{S}_{\mathbb{F}}^{p}(0,\beta)$. Finally, we define operator

$$\mathbb{E}^{f}_{\alpha,\beta}: L^{1}(\mathcal{F}_{\beta}) \to L^{1}(\mathcal{F}_{\alpha}),$$

by letting

$$\mathbb{E}^{f}_{\alpha,\beta}(\xi) \coloneqq \begin{cases} \mathbb{E}^{(1),f}_{\alpha,\beta}(\xi) & \text{for } \xi \in L^{1}(\mathcal{F}_{\beta}) \smallsetminus \bigcup_{p>1} L^{p}(\mathcal{F}_{\beta}) \\ \mathbb{E}^{(p),f}_{\alpha,\beta}(\xi) & \text{for } \xi \in \bigcup_{p>1} L^{p}(\mathcal{F}_{\beta}). \end{cases}$$
(5.1)

We say that a process X of class (D) is an \mathbb{E}^{f} -supermartingale (resp. \mathbb{E}^{f} -submartingale) on $[[\alpha,\beta]]$, if $\mathbb{E}^{f}_{\sigma,\tau}(X_{\tau}) \leq X_{\sigma}$ (resp. $\mathbb{E}^{f}_{\sigma,\tau}(X_{\tau}) \geq X_{\sigma}$) for every $\sigma, \tau \in \mathcal{T}_{\alpha,\beta}, \sigma \leq \tau$. X is an \mathbb{E}^{f} -martingale on $[[\alpha,\beta]]$, if X is at the same time an \mathbb{E}^{f} -supermartingale and an \mathbb{E}^{f} -submartingale on $[[\alpha,\beta]]$.

Remark 5.1. Note that the process Y of class (D) is an \mathbb{E}^{f} -martingale on $[[\alpha, \beta]]$ if and only if Y is indistinguishable from the first component of the solution to BSDE^{α,β}(Y^{β}, f) on $[[\alpha, \beta]]$. Thus, in order to prove that Y is an \mathbb{E}^{f} -martingale on $[[\alpha, \beta]]$, it suffices to show that $Y_{\sigma} = \mathbb{E}^{f}_{\sigma,\beta}(Y_{\beta})$, for any $\sigma \in \mathcal{T}_{\alpha,\beta}$.

Proposition 5.2. Let $\alpha, \beta \in \mathcal{T}, \alpha \leq \beta$.

- (i) Let $\xi \in L^p(\mathcal{F}_{\beta})$ and let V be an \mathbb{F} -adapted, finite variation process such that $V_{\alpha} = 0$. Let (X, H) be a solution to $BSDE^{\alpha,\beta}(\xi, f + dV)$ such that X is of class (D), in case p = 1, and $X \in \mathcal{S}_{\mathbb{F}}^p(\alpha, \beta)$, in case p > 1. If V (resp. -V) is increasing, then X is \mathbb{E}^f -supermartingale (resp. \mathbb{E}^f -submartingale) on $[[\alpha, \beta]]$.
- (*ii*) If $\xi_1, \xi_2 \in L^p(\mathcal{F}_\beta)$ and $\xi_1 \leq \xi_2$, then $\mathbb{E}^f_{\alpha,\beta}(\xi_1) \leq \mathbb{E}^f_{\alpha,\beta}(\xi_2)$.
- (iii) Let $\xi \in L^p(\mathcal{F}_\beta)$. For every $A \in \mathcal{F}_\alpha$,

$$\mathbf{1}_A \mathbb{E}^f_{\alpha,\beta}(\xi) = \mathbb{E}^{f_A}_{\alpha,\beta}(\mathbf{1}_A \xi)$$

where $f_A(t, y, z) = f(t, y, z) \mathbf{1}_A \mathbf{1}_{\{t \ge \alpha\}}$. (iv) Let $\xi \in L^p(\mathcal{F}_\beta)$. For every $\gamma \in \mathcal{T}$ such that $\gamma \ge \beta$,

$$\mathbb{E}^{f}_{\alpha,\beta}(\xi) = \mathbb{E}^{f^{\beta}}_{\alpha,\gamma}(\xi)$$

where $f^{\beta}(t, y, z) = f(t, y, z) \mathbf{1}_{\{t \le \beta\}}$.

(v) Let p = 1. Assume that f_1, f_2 satisfies (H1)-(H5),(Z) and let $\alpha, \beta_1, \beta_2 \in \mathcal{T}, \alpha \leq \beta_1 \leq \beta_2$. Assume that $\xi_1 \in L^1(\mathcal{F}_{\beta_1})$ and $\xi_2 \in L^1(\mathcal{F}_{\beta_2})$. Moreover, let (Y^1, Z^1) be a solution to $BSDE^{\alpha,\beta_2}(\xi_2, f_1^{\beta_1})$, where $f_1^{\beta_1}(t, y, z) = f_1(t, y, z)\mathbf{1}_{\{t\leq\beta\}}$, and (Y^2, Z^2) be a solution to $BSDE^{\alpha,\beta_2}(\xi_2, f_2)$, such that Y^1, Y^2 are of class (D). If $(Y^1 - Y^2) \in \mathcal{S}_{\mathbb{F}}^2(\alpha, \beta_2)$,

then

$$\begin{split} \|\mathbb{E}_{\alpha,\beta_{1}}^{f_{1}}(\xi_{1}) - \mathbb{E}_{\alpha,\beta_{2}}^{f_{2}}(\xi_{2})\|^{2} &\leq C\mathbb{E}\Big(\int_{\alpha}^{\beta_{1}}|Y_{r}^{1} - Y_{r}^{2}||f_{1} - f_{2}|(r,Y_{r}^{2},Z_{r}^{2})\,dr \\ &+ |\xi_{1} - \xi_{2}|^{2} + \int_{\beta_{1}}^{\beta_{2}}|Y_{r}^{1} - Y_{r}^{2}||f_{2}(r,Y_{r}^{2},Z_{r}^{2})|\,dr|\mathcal{F}_{\alpha}\Big), \end{split}$$

for some C depending only on λ, μ, T .

(vi) Let p > 1. Assume that f_1, f_2 satisfies (H1)-(H5) and let $\alpha, \beta_1, \beta_2 \in \mathcal{T}, \alpha \leq \beta_1 \leq \beta_2$. Assume that $\xi_1 \in L^p(\mathcal{F}_{\beta_1})$ and $\xi_2 \in L^p(\mathcal{F}_{\beta_2})$. Moreover, let (Y^1, Z^1) be a solution to $BSDE^{\beta_2}(\xi_1, f_1^{\beta_1})$, with $Y^1 \in \mathcal{S}_{\mathbb{F}}^p(0, \beta_2)$, where $f_1^{\beta_1}(t, y, z) = f_1(t, y, z)\mathbf{1}_{\{t \leq \beta_1\}}$, and (Y^2, Z^2) be a solution to $BSDE^{\beta_2}(\xi_2, f_2)$, with $Y^2 \in \mathcal{S}_{\mathbb{F}}^p(0, \beta_2)$. Then there exists c > 0, depending only on T, μ, λ, p , such that such that

$$\begin{aligned} \|\mathbb{E}_{\cdot,\beta_{1}}^{f_{1}}(\xi_{1}) - \mathbb{E}_{\cdot,\beta_{2}}^{f_{2}}(\xi_{2})\|_{\mathcal{S}^{p}(0,\beta_{1})} &\leq c \Big[\mathbb{E}\Big(\int_{0}^{\beta_{1}} |f_{1} - f_{2}|(r,Y_{r}^{2},Z_{r}^{2}) dr\Big)^{p} \\ &+ \mathbb{E}|\xi_{1} - \xi_{2}|^{p} + \mathbb{E}\Big(\int_{\beta_{1}}^{\beta_{2}} |f_{2}(r,Y_{r}^{2},Z_{r}^{2})| dr\Big)^{p}\Big]^{1/p}. \end{aligned}$$

Proof. (i) Assume that V is increasing. Let $\sigma, \tau \in \mathcal{T}$ be such that $\alpha \leq \sigma \leq \tau \leq \beta$. Obviously, (X, H) is a solution to $\text{BSDE}^{\alpha,\tau}(X_{\tau}, f + dV)$. Let (\tilde{X}, \tilde{H}) be a solution to $\text{BSDE}^{\alpha,\tau}(X_{\tau}, f)$ such that \tilde{X} is of class (D), in case p = 1, and $\tilde{X} \in \mathcal{S}_{\mathbb{F}}^{p}(\alpha, \tau)$, in case p > 1. By [28, Proposition 3.2, Lemma 3.3] and Theorem 3.6, $X \geq \tilde{X}$ on $[[\alpha, \tau]]$, in particular, $X_{\sigma} \geq \tilde{X}_{\sigma}$. Therefore, we have $\mathbb{E}_{\sigma,\tau}^{f}(X_{\tau}) = \tilde{X}_{\sigma} \leq X_{\sigma}$, hence X is \mathbb{E}^{f} -supermartingale. Analogously, we show that if -V is increasing, then X is \mathbb{E}^{f} -submartingale. This completes the proof of (i). The assertion (ii) follows directly from [28, Proposition 3.2, Lemma 3.3] and Theorem 3.6. As to (iii), let $A \in \mathcal{F}_{\alpha}$. Let (Y, Z), (\bar{Y}, \bar{Z}) be solutions to $\text{BSDE}^{\beta}(\xi, f)$ and $\text{BSDE}^{\beta}(\mathbf{1}_{A}\xi, f_{A})$, respectively, such that $Y_{\tau} = \mathbb{E}_{\tau,\beta}^{f}(\xi)$, $\tau \in \mathcal{T}^{\beta}$ and $\bar{Y}_{\tau} = \mathbb{E}_{\tau,\beta}^{f_{A}}(\mathbf{1}_{A}\xi)$, $\tau \in \mathcal{T}^{\beta}$. It is easy to see that $(\mathbf{1}_{A}Y, \mathbf{1}_{A}Z)$ is a solution to $\text{BSDE}^{\alpha,\beta}(\mathbf{1}_{A}\xi, f_{A})$. Indeed, for $\sigma \in \mathcal{T}_{\alpha,\beta}$,

$$\mathbf{1}_{A}Y_{t} = \mathbf{1}_{A}\xi + \int_{\sigma}^{\beta} \mathbf{1}_{A}f(r, Y_{r}, Z_{r}) dr - \int_{\sigma}^{\beta} \mathbf{1}_{A}Z_{r} dB_{r}$$
$$= \mathbf{1}_{A}\xi + \int_{\sigma}^{\beta} \mathbf{1}_{A}f(r, \mathbf{1}_{A}Y_{r}, \mathbf{1}_{A}Z_{r}) dr - \int_{\sigma}^{\beta} \mathbf{1}_{A}Z_{r} dB_{r}$$

By the uniqueness for BSDEs (see Theorems 3.3,3.6) $\mathbf{1}_A Y = \overline{Y}$ on $[[\alpha, \beta]]$, which implies (iii). For (iv), let (Y, Z) be a solution to $BSDE^{\beta}(\xi, f)$ and let $(\overline{Y}, \overline{Z})$ be a solution to $BSDE^{\gamma}(\xi, f^{\beta})$ such that $Y_{\tau} = \mathbb{E}_{\tau,\beta}^{f}(\xi), \tau \in \mathcal{T}^{\beta}$ and $\overline{Y}_{\tau} = \mathbb{E}_{\tau,\gamma}^{f^{\beta}}(\xi), \tau \in \mathcal{T}^{\gamma}$. Note that, $(\overline{Y}, \overline{Z})$ is a solution to $BSDE^{\alpha,\beta}(\overline{Y}_{\beta}, f)$. What is more, for $\sigma \in \mathcal{T}_{\alpha,\beta}$,

$$\bar{Y}_{\sigma} = \mathbb{E}\Big(\xi + \int_{\sigma}^{\gamma} f^{\beta}(r, \bar{Y}_{r}, \bar{Z}_{r}) dr | \mathcal{F}_{\sigma}\Big) = \mathbb{E}\Big(\xi + \int_{\sigma}^{\beta} f(r, \bar{Y}_{r}, \bar{Z}_{r}) dr | \mathcal{F}_{\sigma}\Big),$$

therefore $\bar{Y}_{\beta} = \xi$. Hence, (\bar{Y}, \bar{Z}) is a solution to $BSDE^{\alpha,\beta}(\xi, f)$, which results, by the uniqueness argument, that $\bar{Y} = Y$ on $[[\alpha, \beta]]$ and $\mathbb{E}^{f}_{\alpha,\beta}(\xi) = \bar{Y}_{\alpha} = \mathbb{E}^{f^{\beta}}_{\alpha,\gamma}, \alpha \leq \beta$. This concludes the proof of (iv). Now, we shall proceed to the proof of (v). Let (Y^{1}, Z^{1}) , (Y^{2}, Z^{2}) be defined as in the assertion (v). By (iv), we know that $\mathbb{E}^{f_{1}}_{\alpha,\beta_{1}}(\xi_{1}) = Y^{1}_{\alpha}$, for $\alpha \in \mathcal{T}^{\beta_{1}}$. Let us define

$$\tau_k = \inf\left\{t \ge 0: \int_0^t |Z_r^1 - Z_r^2|^2 \, dr \ge k\right\} \land \beta_2, \quad k \in \mathbb{N}.$$

From the definition of a solution to BSDE, we have that $\{\tau_k\}_{k\geq 1}$ is a chain. By Ito's formula, for $\alpha \in \mathcal{T}^{\beta_1}$,

$$e^{a\alpha}|Y_{\alpha}^{1} - Y_{\alpha}^{2}|^{2} + \int_{\alpha}^{\tau_{k}} e^{ar}|Z_{r}^{1} - Z_{r}^{2}|^{2} dr + a \int_{\alpha}^{\tau_{k}} e^{ar}|Y_{r}^{1} - Y_{r}^{2}|^{2} dr$$

$$\leq e^{a\tau_{k}}|Y_{\tau_{k}}^{1} - Y_{\tau_{k}}^{2}|^{2} + 2 \int_{\alpha}^{\tau_{k}} e^{ar}(Y_{r}^{1} - Y_{r}^{2})(f_{1}^{\beta_{1}}(r, Y_{r}^{1}, Z_{r}^{1}) - f_{2}(r, Y_{r}^{2}, Z_{r}^{2})) dr \qquad (5.2)$$

$$-2 \int_{\alpha}^{\tau_{k}} e^{ar}(Y_{r}^{1} - Y_{r}^{2})(Z_{r}^{1} - Z_{r}^{2}) dB_{r}, \quad a \geq 0.$$

By (H1), (H2) (without loss of generality we may assume that $\mu = 0$), we have

$$\begin{split} &(Y_r^1 - Y_r^2) (f(r, Y_r^1, Z_r^1) - f(r, Y_r^2, Z_r^2)) \leq \lambda |Y_r^1 - Y_r^2| |Z_r^1 - Z_r^2| \\ &+ |Y_r^1 - Y_r^2| |f_1^{\beta_1} - f_2| (r, Y_r^2, Z_r^2) \leq 4\lambda^2 |Y_r^1 - Y_r^2|^2 + \frac{1}{4} |Z_r^1 - Z_r^2|^2 \\ &+ |Y_r^1 - Y_r^2| |f_1^{\beta_1} - f_2| (r, Y_r^2, Z_r^2). \end{split}$$

Therefore, by (5.2), we have

$$\begin{split} e^{a\alpha}|Y_{\alpha}^{1}-Y_{\alpha}^{2}|^{2} &+ \int_{\alpha}^{\tau_{k}} e^{ar}|Z_{r}^{1}-Z_{r}^{2}|^{2} dr + a \int_{\alpha}^{\tau_{k}} e^{ar}|Y_{r}^{1}-Y_{r}^{2}|^{2} dr \\ &\leq e^{a\tau_{k}}|Y_{\tau_{k}}^{1}-Y_{\tau_{k}}^{2}|^{2} + 8\lambda^{2} \int_{\alpha}^{\beta_{2}} e^{ar}|Y_{r}^{1}-Y_{r}^{2}|^{2} dr + \frac{1}{2} \int_{\alpha}^{\beta_{2}} e^{ar}|Z_{r}^{1}-Z_{r}^{2}|^{2} dr \\ &+ 2 \int_{\alpha}^{\beta_{2}} e^{ar}|Y_{r}^{1}-Y_{r}^{2}||f_{1}^{\beta_{1}}-f_{2}|(r,Y_{r}^{2},Z_{r}^{2}) dr \\ &- 2 \int_{\alpha}^{\tau_{k}} e^{ar}(Y_{r}^{1}-Y_{r}^{2})(Z_{r}^{1}-Z_{r}^{2}) dB_{r}. \end{split}$$

Consequently, by the fact that $\int_0^{\cdot} e^{\alpha r} (Y_r^1 - Y_r^2) (Z_r^1 - Z_r^2) dB_r$ is a martingale on $[[\alpha, \tau_k]]$, we find that for $a \ge 8\lambda^2$,

$$e^{a\alpha}|Y_{\alpha}^{1} - Y_{\alpha}^{2}|^{2} \leq \mathbb{E}\Big(e^{a\tau_{k}}|Y_{\tau_{k}}^{1} - Y_{\tau_{k}}^{2}|^{2} + 2\int_{\alpha}^{\beta_{2}}e^{ar}|Y_{r}^{1} - Y_{r}^{2}||f_{1}^{\beta_{1}} - f_{2}|(r, Y_{r}^{2}, Z_{r}^{2})\,dr|\mathcal{F}_{\alpha}\Big).$$
(5.3)

Since $Y^1 - Y^2 \in \mathcal{S}^2_{\mathbb{F}}(\alpha, \beta_2)$, we may conclude, by letting $k \to \infty$ in the right-hand side of (5.3) and applying the Lebesgue dominated convergence theorem, that

$$e^{a\alpha}|Y_{\alpha}^{1} - Y_{\alpha}^{2}|^{2} \leq \mathbb{E}\left(e^{a\beta_{2}}|\xi_{1} - \xi_{2}|^{2} + 2\int_{\alpha}^{\beta_{2}}e^{ar}|Y_{r}^{1} - Y_{r}^{2}||f_{1}^{\beta_{1}} - f_{2}|(r, Y_{r}^{2}, Z_{r}^{2}) dr|\mathcal{F}_{\alpha}\right),$$

which implies that

$$|Y_{\alpha}^{1} - Y_{\alpha}^{2}|^{2} \leq C\mathbb{E}\Big(|\xi_{1} - \xi_{2}|^{2} + \int_{\alpha}^{\beta_{2}} |Y_{r}^{1} - Y_{r}^{2}||f_{1}^{\beta_{1}} - f_{2}|(r, Y_{r}^{2}, Z_{r}^{2}) dr|\mathcal{F}_{\alpha}\Big)$$
(5.4)

for some C > 0 depending only on λ and β_2 . Finally, note that

$$\begin{split} \int_{\alpha}^{\beta_2} |Y_r^1 - Y_r^2| |f_1^{\beta_1} - f_2|(r, Y_r^2, Z_r^2) \, dr &= \int_{\alpha}^{\beta_1} |Y_r^1 - Y_r^2| |f_1^{\beta_1} - f_2|(r, Y_r^2, Z_r^2) \, dr \\ &+ \int_{\beta_1}^{\beta_2} |Y_r^1 - Y_r^2| |f_2(r, Y_r^2, Z_r^2)| \, dr, \end{split}$$

which combined with (5.4) completes the proof of (v). The inequality asserted in (vi) follows directly from Proposition 3.4.

6. EXTENDED NONLINEAR DYNKIN GAMES

In the whole section, we assume that (H1)-(H5), (Z) are in force and that L and U are \mathbb{F} -optional processes of class (D).

Definition 6.1. Let $\tau \in \mathcal{T}$ and $H \in \mathcal{F}_{\tau}$. A pair $\rho = (\tau, H)$ is called a *stopping system* if $\{\tau = T\} \subset H$. For brevity, we write $\tau \mid H$.

By \mathcal{U} we denote the set of all stoping systems. We then have $\mathcal{T} \subset \mathcal{U}$, by using embedding $\mathcal{T} \ni \tau \mapsto \tau \lfloor \Omega \in \mathcal{U}$. For given $\theta \in \mathcal{T}$, we denote by \mathcal{U}_{θ} the set of all stopping systems $\tau \lfloor H$ such that $\tau \geq \theta$. For an optional, right-limited process ϕ and $\tau \lfloor H \in \mathcal{U}$ we put

$$\phi_{\tau|H} \coloneqq \phi_{\tau} \mathbf{1}_H + \phi_{\tau+} \mathbf{1}_{H^c}$$

In particular, we have $\phi_{\tau|\Omega} = \phi_{\tau}$. For an optional process ϕ we let

$$\phi^{u}_{\tau \lfloor H} \coloneqq \phi_{\tau} \mathbf{1}_{H} + \overleftarrow{\phi}_{\tau} \mathbf{1}_{H^{c}} \quad \text{and} \quad \phi^{l}_{\tau \lfloor H} \coloneqq \phi_{\tau} \mathbf{1}_{H} + \underbrace{\phi}_{\tau} \mathbf{1}_{H^{c}}.$$

Note that, when ϕ is right-limited, then $\phi_{\tau|H}^u = \phi_{\tau|H}^l = \phi_{\tau|H}$.

For two stopping systems $\tau | H, \sigma | G \in \mathcal{U}$ we define the pay-off

$$J(\tau \lfloor H, \sigma \lfloor G) \coloneqq L^u_{\tau \lfloor H} \mathbf{1}_{\{\tau \le \sigma < T\}} + U^l_{\sigma \lfloor G} \mathbf{1}_{\{\sigma < \tau\}} + \xi \mathbf{1}_{\{\tau = \sigma = T\}}.$$
(6.1)

Note that $J(\tau[H, \sigma[G])$ is $\mathcal{F}_{\tau \wedge \sigma}$ -measurable random variable. Now, we shall proceed to the so called *extended Dynkin games*.

Definition 6.2. Let $\theta \in \mathcal{T}$.

(i) Upper and lower value of the game are defined respectively as

$$\overline{V}(\theta) \coloneqq \operatorname{ess\,inf\,ess\,sup}_{\sigma \lfloor G \in \mathcal{U}_{\theta}} \mathbb{E}^{f}_{\theta, \tau \wedge \sigma} J(\tau \lfloor H, \sigma \lfloor G);$$

$$\underline{V}(\theta) \coloneqq \operatorname{ess\,sup\,ess\,inf}_{\tau \mid H \in \mathcal{U}_{\theta}} \mathbb{E}^{f}_{\theta, \tau \wedge \sigma} J(\tau \lfloor H, \sigma \lfloor G).$$
(6.2)

(ii) We say that an extended \mathbb{E}^f -Dynkin game with pay-off function J has a value if $\overline{V}(\theta) = \underline{V}(\theta)$ for any $\theta \in \mathcal{T}$.

Let (Y, Z, R) be a solution to $\text{RBSDE}^T(\xi, f, L, U)$. For every $\theta \in \mathcal{T}$ and $\varepsilon > 0$ we define the following sets

$$A^{\varepsilon} := \{(\omega, t) \in \Omega \times [0, T] : Y_t(\omega) \le L_t^{\xi}(\omega) + \varepsilon\}; B^{\varepsilon} := \{(\omega, t) \in \Omega \times [0, T] : Y_t(\omega) \ge U_t^{\xi}(\omega) - \varepsilon\},\$$

where $L_{\theta}^{\xi} \coloneqq L_{\theta} \mathbf{1}_{\{\theta < T\}} + \xi \mathbf{1}_{\{\theta = T\}}, U_{\theta}^{\xi} \coloneqq U_{\theta} \mathbf{1}_{\{\theta < T\}} + \xi \mathbf{1}_{\{\theta = T\}}$ for $\theta \in \mathcal{T}$. Let us also define the following stopping times

$$\tau^{\varepsilon}_{\theta} \coloneqq \inf\{t \geq \theta, \, Y_t \leq L^{\xi}_t + \varepsilon\} \wedge T, \quad \sigma^{\varepsilon}_{\theta} \coloneqq \inf\{t \geq \theta, \, Y_t \geq U^{\xi}_t - \varepsilon\} \wedge T.$$

We let

$$H^{\varepsilon} \coloneqq \{\omega \in \Omega : (\omega, \tau^{\varepsilon}_{\theta}(\omega)) \in A^{\varepsilon}\}; \quad G^{\varepsilon} \coloneqq \{\omega \in \Omega : (\omega, \sigma^{\varepsilon}_{\theta}(\omega)) \in B^{\varepsilon}\}$$

Consider the following stopping systems

$$\tau_{\theta}^{\varepsilon} [H^{\varepsilon} \text{ and } \sigma_{\theta}^{\varepsilon}] G^{\varepsilon}.$$
 (6.3)

Lemma 6.3. Let (Y, Z, R) be a solution to $\operatorname{RBSDE}^T(\xi, f, L, U)$. Then Y is an \mathbb{E}^f -submartingale on $[[\theta, \tau_{\theta}^{\varepsilon}]]$ and an \mathbb{E}^f -supermartingale on $[[\theta, \sigma_{\theta}^{\varepsilon}]]$.

Proof. We show that Y is \mathbb{E}^f -submartingale on $[[\theta, \tau_{\theta}^{\varepsilon}]]$. The proof of the second assertion runs analogously. By the definition of $\tau_{\theta}^{\varepsilon}$ we have $Y_t > L_t + \varepsilon$ on $[[\theta, \tau_{\theta}^{\varepsilon}][$. This implies, by the minimality condition, that R^+ is constant on $[[\theta, \tau_{\theta}^{\varepsilon}][$. By the last inequality, we also have $Y_{\tau_{\theta}^{\varepsilon}-} \geq \vec{L}_{\tau_{\theta}^{\varepsilon}} + \varepsilon$, so as a result, by the minimality condition again, we get $\Delta^- R_{\tau_{\theta}^{\varepsilon}}^+ = 0$. Therefore, R^+ is constant on $[[\theta, \tau_{\theta}^{\varepsilon}]]$. This implies that

$$Y_t = Y_{\tau_{\theta}^{\varepsilon}} + \int_t^{\tau_{\theta}^{\varepsilon}} f(r, Y_r, Z_r) \, dr - R_{\tau_{\theta}^{\varepsilon}}^- + R_t^- - \int_t^{\tau_{\theta}^{\varepsilon}} Z_r \, dB_r, \quad t \in [\theta, \tau_{\theta}^{\varepsilon}].$$

Thus, by Proposition 5.2 (i), Y is an \mathbb{E}^{f} -submartingale on $[[\theta, \tau_{\theta}^{\varepsilon}]]$.

Lemma 6.4. Let (Y, Z, R) be a solution to $\text{RBSDE}^T(\xi, f, L, U)$.

(i) For any $\theta \in \mathcal{T}$,

$$Y_{\tau_{\theta}^{\varepsilon} \mid H^{\varepsilon}} \leq L_{\tau_{\theta}^{\varepsilon} \mid H^{\varepsilon}}^{\xi, u} + \varepsilon \quad \text{and} \quad Y_{\sigma_{\theta}^{\varepsilon} \mid G^{\varepsilon}} \geq U_{\sigma_{\theta}^{\varepsilon} \mid G^{\varepsilon}}^{\xi, l} - \varepsilon.$$

$$(6.4)$$

(*ii*) For each $\theta \in \mathcal{T}$ and any $\tau \lfloor A, \sigma \lfloor B \in \mathcal{U}_{\theta}$,

$$\mathbb{E}^{f}_{\theta,\tau_{\theta}^{\varepsilon}\wedge\sigma}(Y_{\tau_{\theta}^{\varepsilon}\mid H^{\varepsilon}}\mathbf{1}_{\{\tau_{\theta}^{\varepsilon}\leq\sigma\}}+Y_{\sigma\mid B}\mathbf{1}_{\{\sigma<\tau_{\theta}^{\varepsilon}\}})\geq Y_{\theta} \quad \text{and} \\
\mathbb{E}^{f}_{\theta,\tau\wedge\sigma_{\theta}^{\varepsilon}}(Y_{\tau\mid A}\mathbf{1}_{\{\tau\leq\sigma_{\theta}^{\varepsilon}\}}+Y_{\sigma_{\theta}^{\varepsilon}\mid G^{\varepsilon}}\mathbf{1}_{\{\sigma_{\theta}^{\varepsilon}<\tau\}})\leq Y_{\theta}.$$
(6.5)

Proof. (i) We shall prove the first inequality in (6.4), the proof of the second one runs analogously. Due to the definitions of $\tau_{\theta}^{\varepsilon} [H^{\varepsilon}, Y_{\tau_{\theta}^{\varepsilon}}]_{H^{\varepsilon}}, L_{\tau_{\theta}^{\varepsilon}}^{\xi,u}]_{H^{\varepsilon}}$ and H^{ε} , we have, on the set H^{ε} , that $Y_{\tau_{\theta}^{\varepsilon}}]_{H^{\varepsilon}} = Y_{\tau_{\theta}^{\varepsilon}} \leq L_{\tau_{\theta}^{\varepsilon}}^{\xi} + \varepsilon = L_{\tau_{\theta}^{\varepsilon}}^{\xi,u}]_{H^{\varepsilon}} + \varepsilon$, while on the set $H^{\varepsilon,c}$, we have

$$Y_{\tau_{\theta}^{\varepsilon} \mid H^{\varepsilon}} = Y_{\tau_{\theta}^{\varepsilon}} + \quad \text{and} \quad L_{\tau_{\theta}^{\varepsilon} \mid H^{\varepsilon}}^{\xi, u} = \overleftarrow{L}_{\tau_{\theta}^{\varepsilon}}^{\xi}.$$
(6.6)

On the other hand, by the definition of $\tau_{\theta}^{\varepsilon}$, for P-a.e. $\omega \in \Omega$ there exists nonincreasing sequence (t_n) (depending on $\omega \in \Omega$) such that $t_n \searrow \tau_{\theta}^{\varepsilon}$ and $Y_{t_n} \leq L_{t_n}^{\xi} + \varepsilon$ for all $n \in \mathbb{N}$. Therefore

$$\limsup_{n \to \infty} Y_{t_n} \le \limsup_{n \to \infty} L_{t_n}^{\xi} + \varepsilon.$$

Due to the definiton of \widetilde{L}^{ξ} , we have $\limsup_{n \to \infty} L_{t_n}^{\xi} \leq \widetilde{L}_{\tau_{\theta}^{\varepsilon}}^{\xi}$. Since Y is regulated, $\limsup_{n \to \infty} Y_{t_n} = Y_{\tau_{\theta}^{\varepsilon}+}$. Thus, $Y_{\tau_{\theta}^{\varepsilon}+} \leq \widetilde{L}_{\tau_{\theta}^{\varepsilon}}^{\xi} + \varepsilon$. This inequality combined with (6.6) implies that $Y_{\tau_{\theta}^{\varepsilon}|H^{\varepsilon}} \leq L_{\tau_{\theta}^{\varepsilon}|H^{\varepsilon}}^{\xi,u} + \varepsilon$ on $H^{\varepsilon,c}$.

(ii) First, we shall prove the first inequality in (6.5). We have

$$Y_{\tau_{\theta}^{\varepsilon} \lfloor H^{\varepsilon}} \mathbf{1}_{\{\tau_{\theta}^{\varepsilon} \leq \sigma\}} + Y_{\sigma \lfloor B} \mathbf{1}_{\{\sigma < \tau_{\theta}^{\varepsilon}\}} = Y_{\tau_{\theta}^{\varepsilon}} \mathbf{1}_{H^{\varepsilon}} \mathbf{1}_{\{\tau_{\theta}^{\varepsilon} \leq \sigma\}} + Y_{\tau_{\theta}^{\varepsilon}} \mathbf{1}_{H^{\varepsilon,c}} \mathbf{1}_{\{\tau_{\theta}^{\varepsilon} \leq \sigma\}} + Y_{\sigma} \mathbf{1}_{B} \mathbf{1}_{\{\sigma < \tau_{\theta}^{\varepsilon}\}} + Y_{\sigma+} \mathbf{1}_{B^{c}} \mathbf{1}_{\{\sigma < \tau_{\theta}^{\varepsilon}\}}.$$

$$(6.7)$$

By Lemma 6.3 Y is an \mathbb{E}^f -submartingale on $[[\theta, \tau_{\theta}^{\varepsilon}]]$, which implies that

$$\mathbf{1}_{\{\sigma < \tau_{\theta}^{\varepsilon}\}} Y_{\sigma+} \ge \mathbf{1}_{\{\sigma < \tau_{\theta}^{\varepsilon}\}} Y_{\sigma \land \tau_{\theta}^{\varepsilon}}.$$
(6.8)

By the form of H^{ε} and the minimality condition, $\mathbf{1}_{H^{\varepsilon,c}}\Delta^+ R^+_{\tau^{\varepsilon}_{a}} = 0$. Thus,

$$\mathbf{1}_{H^{\varepsilon,c}} Y_{\tau_{\theta}^{\varepsilon}+} \ge \mathbf{1}_{H^{\varepsilon,c}} Y_{\tau_{\theta}^{\varepsilon}}.$$
(6.9)

By virtue of (6.7)–(6.9), we conclude that $Y_{\tau_{\theta}^{\varepsilon} \mid H^{\varepsilon}} \mathbf{1}_{\{\tau_{\theta}^{\varepsilon} \leq \sigma\}} + Y_{\sigma \mid B} \mathbf{1}_{\{\sigma < \tau_{\theta}^{\varepsilon}\}} \geq Y_{\tau_{\theta}^{\varepsilon} \wedge \sigma}$. Since the operator \mathbb{E}^{f} is nondecreasing (see Proposition 5.2 (ii))

$$\mathbb{E}^{f}_{\theta,\tau^{\varepsilon}_{\theta}\wedge\sigma}(Y_{\tau^{\varepsilon}_{\theta}\mid H^{\varepsilon}}\mathbf{1}_{\{\tau^{\varepsilon}_{\theta}\leq\sigma\}}+Y_{\sigma\mid B}\mathbf{1}_{\{\sigma<\tau^{\varepsilon}_{\theta}\}})\geq\mathbb{E}^{f}_{\theta,\tau^{\varepsilon}_{\theta}\wedge\sigma}(Y_{\tau^{\varepsilon}_{\theta}\wedge\sigma}).$$

Using again the fact that Y is \mathbb{E}^{f} -submartingale on $[[\theta, \tau_{\theta}^{\varepsilon}]]$, we conclude that $\mathbb{E}^{f}_{\theta, \tau_{\theta}^{\varepsilon} \wedge \sigma}(Y_{\tau_{\theta}^{\varepsilon} \wedge \sigma}) \geq Y_{\theta}$, and as a result $\mathbb{E}^{f}_{\theta, \tau_{\theta}^{\varepsilon} \wedge \sigma}(Y_{\tau_{\theta}^{\varepsilon} \mid H^{\varepsilon}} \mathbf{1}_{\{\tau_{\theta}^{\varepsilon} \leq \sigma\}} + Y_{\sigma \lfloor B} \mathbf{1}_{\{\sigma < \tau_{\theta}^{\varepsilon}\}}) \geq Y_{\theta}$.

The proof of the second inequality in (6.5) requires slightly different arguments. We have

$$Y_{\tau \lfloor A} \mathbf{1}_{\{\tau \leq \sigma_{\theta}^{\varepsilon}\}} + Y_{\sigma_{\theta}^{\varepsilon} \lfloor G^{\varepsilon}} \mathbf{1}_{\{\sigma_{\theta}^{\varepsilon} < \tau\}} = Y_{\tau} \mathbf{1}_{A} \mathbf{1}_{\{\tau \leq \sigma_{\theta}^{\varepsilon}\}} + Y_{\tau+} \mathbf{1}_{A^{c}} \mathbf{1}_{\{\tau \leq \sigma_{\theta}^{\varepsilon}\}} + Y_{\sigma_{\theta}^{\varepsilon}} \mathbf{1}_{G^{\varepsilon}} \mathbf{1}_{\{\sigma_{\theta}^{\varepsilon} < \tau\}} + Y_{\sigma_{\theta}^{\varepsilon+}} \mathbf{1}_{G^{\varepsilon,c}} \mathbf{1}_{\{\sigma_{\theta}^{\varepsilon} < \tau\}}.$$
(6.10)

By the analogous argument as in the proof of (6.7) - the form of G^{ε} combined with the definition of a solution to RBSDE give $\mathbf{1}_{G^{\varepsilon,c}}\Delta^+ R^-_{\sigma^{\varepsilon}_a} = 0$ - we find that

$$\mathbf{1}_{G^{\varepsilon,c}} Y_{\sigma_{\rho}^{\varepsilon}+} \leq \mathbf{1}_{G^{\varepsilon,c}} Y_{\sigma_{\rho}^{\varepsilon}}.$$
(6.11)

Using the fact that Y is an \mathbb{E}^{f} -supermartingale on $[[\theta, \sigma_{\theta}^{\varepsilon}]]$ (see Lemma 6.3) we obtain that

$$\mathbf{1}_{\{\tau < \sigma_{\theta}^{\varepsilon}\}} Y_{\tau +} \le \mathbf{1}_{\{\tau < \sigma_{\theta}^{\varepsilon}\}} Y_{\tau}.$$
(6.12)

But we also need

$$\mathbf{1}_{\{\tau=\sigma_{\theta}^{\varepsilon}\}}Y_{\tau+} \leq \mathbf{1}_{\{\tau=\sigma_{\theta}^{\varepsilon}\}}Y_{\tau}.$$
(6.13)

The above inequality may not hold only in case $\Delta^+ R_{\sigma_{\theta}^{\varepsilon}}^- > 0$. The last relation implies, by using the *minimality condition*, that $Y_{\sigma_{\theta}^{\varepsilon}} = U_{\sigma_{\theta}^{\varepsilon}}$. On the other hand, by the definition of $\sigma_{\theta}^{\varepsilon}$, $U_t - Y_t \ge \varepsilon$, $t \in [\theta, \sigma_{\theta}^{\varepsilon})$. This and the previous equation force left positive jump, so $\Delta R_{\sigma_{\theta}^{\varepsilon}}^- > 0$. Consequently, by the *minimality condition*, $Y_{\sigma_{\theta}^{\varepsilon}-} = \bigcup_{\sigma_{\theta}^{\varepsilon}} S$. This contradicts the relation $U_t - Y_t \ge \varepsilon$, $t \in [\theta, \sigma_{\theta}^{\varepsilon})$. Therefore, (6.13) must hold. Combining (6.10)–(6.13) gives

$$Y_{\tau \lfloor A} \mathbf{1}_{\{\tau \leq \sigma_{\theta}^{\varepsilon}\}} + Y_{\sigma_{\theta}^{\varepsilon} \lfloor G^{\varepsilon}} \mathbf{1}_{\{\sigma_{\theta}^{\varepsilon} < \tau\}} \leq Y_{\tau \wedge \sigma_{\theta}^{\varepsilon}}.$$

With the aid of monotonicity of the operator $\mathbb{E}^{f}_{\theta, \tau \wedge \sigma^{\varepsilon}_{\theta}}$ and the fact that Y is \mathbb{E}^{f} -supermartingale on $[[\theta, \sigma^{\varepsilon}_{\theta}]]$, we easily deduce from the above inequality the result. \Box

Lemma 6.5. Let (Y, Z, R) be a solution to $\text{RBSDE}^T(\xi, f, L, U)$. We have the following inequalities:

$$\mathbb{E}^{f}_{\theta,\tau\wedge\sigma^{\varepsilon}_{\theta}}J(\tau\lfloor A,\sigma^{\varepsilon}_{\theta}\lfloor G^{\varepsilon}) - C\varepsilon \leq Y_{\theta} \leq \mathbb{E}^{f}_{\theta,\tau^{\varepsilon}_{\theta}\wedge\sigma}J(\tau^{\varepsilon}_{\theta}\lfloor H^{\varepsilon},\sigma\lfloor B) + C\varepsilon,$$
(6.14)

where C is a constant depending only on $\lambda, \mu, \kappa, T, \|g\|_{L^1}, \gamma$.

Proof. Let $\theta \in \mathcal{T}$ and $\varepsilon > 0$. We shall show the first inequality in (6.14) (the proof of the other one is analogous). By Lemma 6.4

$$Y_{\theta} \leq \mathbb{E}^{f}_{\theta, \tau_{\theta}^{\varepsilon} \wedge \sigma} (Y_{\tau_{\theta}^{\varepsilon} \mid H^{\varepsilon}} \mathbf{1}_{\{\tau_{\theta}^{\varepsilon} \leq \sigma\}} + Y_{\sigma \mid B} \mathbf{1}_{\{\sigma < \tau_{\theta}^{\varepsilon}\}}).$$
(6.15)

We have

$$Y_{\tau_{\theta}^{\varepsilon} \mid H^{\varepsilon}} \mathbf{1}_{\{\tau_{\theta}^{\varepsilon} \le \sigma\}} + Y_{\sigma \mid B} \mathbf{1}_{\{\sigma < \tau_{\theta}^{\varepsilon}\}} = Y_{\tau_{\theta}^{\varepsilon} \mid H^{\varepsilon}} \mathbf{1}_{\{\tau_{\theta}^{\varepsilon} \le \sigma < T\}} + Y_{\sigma \mid B} \mathbf{1}_{\{\sigma < \tau_{\theta}^{\varepsilon}\}} + \xi \mathbf{1}_{\{\tau_{\theta}^{\varepsilon} = \delta = T\}}.$$

By Lemma 6.4, $Y_{\tau_{\theta}^{\varepsilon} \mid H^{\varepsilon}} \leq L_{\tau_{\theta}^{\varepsilon} \mid H^{\varepsilon}}^{u} + \varepsilon$. Moreover, since $Y \leq U$ and Y is right-limited, we have $Y_{\sigma \mid B} = Y_{\sigma \mid B}^{l} \leq U_{\sigma \mid B}^{l}$. Consequently,

$$\begin{split} Y_{\tau_{\theta}^{\varepsilon} \lfloor H^{\varepsilon}} \mathbf{1}_{\{\tau_{\theta}^{\varepsilon} \leq \sigma\}} + Y_{\sigma \lfloor B} \mathbf{1}_{\{\sigma < \tau_{\theta}^{\varepsilon}\}} &\leq \left(L_{\tau_{\theta}^{\varepsilon} \lfloor H^{\varepsilon}}^{u} + \varepsilon \right) \mathbf{1}_{\{\tau_{\theta}^{\varepsilon} \leq \sigma < T\}} + U_{\sigma \lfloor B}^{l} \mathbf{1}_{\{\sigma < \tau_{\theta}^{\varepsilon}\}} \\ &+ \xi \mathbf{1}_{\{\tau_{\theta}^{\varepsilon} = \sigma = T\}} \leq J(\tau_{\theta}^{\varepsilon} \lfloor H^{\varepsilon}, \sigma \lfloor B) + \varepsilon. \end{split}$$

By (6.15) and by properties of the operator \mathbb{E}^{f} (see Proposition 5.2 (ii) and (v)), we get

$$Y_{\theta} \leq \mathbb{E}^{f}_{\theta, \tau^{\varepsilon}_{\theta} \wedge \sigma} (J(\tau^{\varepsilon}_{\theta} \lfloor H^{\varepsilon}, \sigma \lfloor B) + \varepsilon) \leq \mathbb{E}^{f}_{\theta, \tau^{\varepsilon}_{\theta} \wedge \sigma} (J(\tau^{\varepsilon}_{\theta} \lfloor H^{\varepsilon}, \sigma \lfloor B)) + C\varepsilon.$$
(6.16)

Theorem 6.6. Let (Y, Z, R) be a solution to $\text{RBSDE}^T(\xi, f, L, U)$. The extended \mathbb{E}^f -Dynkin game has a value. What is more, for any stopping time $\theta \in \mathcal{T}$,

$$\underline{V}(\theta) = Y_{\theta} = \overline{V}(\theta).$$

Moreover, for every $\theta \in \mathcal{T}$ and $\varepsilon > 0$ the pair of stopping systems $(\tau_{\theta}^{\varepsilon} | H^{\varepsilon}, \delta_{\theta}^{\varepsilon} | G^{\varepsilon})$ defined in (6.3) is ε -saddle point in time θ for extended \mathbb{E}^{f} -Dynkin game, i.e. satisfies inequalities (6.4).

Proof. Since right-hand side inequality in (6.15) is satisfied for all $\sigma | B \in \mathcal{U}_{\theta}$ we have that

$$Y_{\theta} \leq \underset{\sigma \mid B \in \mathcal{U}_{\theta}}{\operatorname{ess inf}} \mathbb{E}^{f}_{\theta, \tau_{\theta}^{\varepsilon} \wedge \sigma} (J(\tau_{\theta}^{\varepsilon} \mid H^{\varepsilon}, \sigma \mid B)) + C\varepsilon$$
$$\leq \underset{\tau \mid A \in \mathcal{U}_{\theta}}{\operatorname{ess sup ess inf}} \mathbb{E}^{f}_{\theta, \tau \wedge \sigma} (J(\tau \mid A, \sigma \mid B)) + C\varepsilon$$

Thus, by the definition of \underline{V}_{θ} (see (6.2)) we have that $Y_{\theta} \leq \underline{V}(\theta) + C\varepsilon$. Similarly, we show that $\overline{V}(\theta) - C\varepsilon \leq Y_{\theta}$ for all $\varepsilon > 0$. In consequence, $\overline{V}(\theta) \leq Y_{\theta} \leq \underline{V}(\theta)$, which combined with the obvious inequality $\underline{V}(\theta) \leq \overline{V}(\theta)$ gives us $\underline{V}(\theta) = Y_{\theta} = \overline{V}(\theta)$. \Box

7. Nonlinear Dynkin games

Throughout the section, we assume that (H1)-(H5), (Z) are in force and that L and U are \mathbb{F} -optional processes of class (D).

For $\tau, \sigma \in \mathcal{T}$ we define the pay-off

$$J_0(\tau,\sigma) \coloneqq L_\tau \mathbf{1}_{\{\tau \le \sigma < T\}} + U_\sigma \mathbf{1}_{\{\sigma < \tau\}} + \xi \mathbf{1}_{\{\tau = \sigma = T\}}.$$
(7.1)

Definition 7.1. Let $\theta \in \mathcal{T}$. Upper and lower value of the game are defined respectively as

$$\overline{V}_{0}(\theta) \coloneqq \operatorname{ess\,sup}_{\sigma \geq \theta} \operatorname{E}_{\sigma \geq \theta}^{f} \mathbb{E}_{\theta, \tau \wedge \sigma}^{J} J_{0}(\tau, \sigma);$$

$$\underline{V}_{0}(\theta) \coloneqq \operatorname{ess\,sup}_{\tau \geq \theta} \operatorname{ess\,sup}_{\sigma \geq \theta} \mathbb{E}_{\theta, \tau \wedge \sigma}^{f} J_{0}(\tau, \sigma).$$
(7.2)

Lemma 7.2. Let (Y, Z, R) be a solution to $\text{RBSDE}^T(\xi, f, L, U)$. Assume that L is right upper semicontinuous and U is right lower semicontinuous. Then

$$Y_{\tau_{\theta}^{\varepsilon}} \le L_{\tau_{\theta}^{\varepsilon}}^{\xi} + \varepsilon, \qquad Y_{\sigma_{\theta}^{\varepsilon}} \ge U_{\sigma_{\theta}^{\varepsilon}}^{\xi} - \varepsilon.$$

$$(7.3)$$

Proof. In case $\tau_{\theta}^{\varepsilon} = T$, (7.3) is obvious. Suppose that $\tau_{\theta}^{\varepsilon} < T$. Suppose by contradiction that $P(Y_{\tau_{\theta}^{\varepsilon}} > L_{\tau_{\theta}^{\varepsilon}}^{\xi} + \varepsilon) > 0$. Without loss of generality we may assume that properties attributed to Y, M, R, L, U and holding *P*-a.s. hold for any $\omega \in \Omega$. Fix $\omega \in \{Y_{\tau_{\theta}^{\varepsilon}} > L_{\tau_{\theta}^{\varepsilon}}^{\xi} + \varepsilon\}$. By the minimality condition $\Delta^{+}R_{\tau_{\theta}^{\varepsilon}}^{+}(\omega) = 0$, and so $\Delta^{+}Y_{\tau_{\theta}^{\varepsilon}}(\omega) = -(\Delta^{+}R_{\tau_{\theta}^{\varepsilon}}^{+} - \Delta^{+}R_{\tau_{\theta}^{\varepsilon}}^{-})(\omega) = \Delta^{+}R_{\tau_{\theta}^{\varepsilon}}^{-}(\omega) > 0$. Therefore

$$Y_{(\tau_{\theta}^{\varepsilon})^{+}}(\omega) > L_{\tau_{\theta}^{\varepsilon}}^{\xi}(\omega) + \varepsilon.$$
(7.4)

Take $\omega \in \Omega$. By the definition of $\tau_{\theta}^{\varepsilon}$ there exists a non-increasing sequence $(t_n(\omega)) \searrow \tau_{\theta}^{\varepsilon}(\omega)$ such that $Y_{t_n}(\omega) \leq L_{t_n}^{\xi}(\omega) + \varepsilon$ for any $n \in \mathbb{N}$. Hence $\limsup_{n \to \infty} Y_{t_n}(\omega) \leq \limsup_{n \to \infty} L_{t_n}^{\xi}(\omega) + \varepsilon$. By the assumptions made L is right upper semicontinuous, thus $\limsup_{n \to \infty} L_{t_n}^{\xi}(\omega) \leq L_{\tau_{\theta}^{\varepsilon}}^{\xi}(\omega)$. On the other hand $t_n(\omega) \searrow \tau_{\theta}^{\varepsilon}(\omega)$ implies $\limsup_{n \to \infty} Y_{t_n}(\omega) = Y_{(\tau_{\theta}^{\varepsilon})+}(\omega)$. Consequently, $Y_{(\tau_{\theta}^{\varepsilon})+}(\omega) \leq L_{\tau_{\theta}^{\varepsilon}}^{\xi}(\omega) + \varepsilon$, which contradicts (7.4). From this we deduce that $Y_{\tau_{\theta}^{\varepsilon}} \leq L_{\tau_{\alpha}^{\varepsilon}}^{\xi} + \varepsilon$ for P-a.e. $\omega \in \Omega$.

Theorem 7.3. Assume that L is right upper semicontinuous and U is right lower semicontinuous. Let (Y, Z, R) be a solution to $\text{RBSDE}^T(\xi, f, L, U)$. Then for any $\theta \in \mathcal{T}$

$$Y_{\theta} = \overline{V}_0(\theta) = \underline{V}_0(\theta). \tag{7.5}$$

Moreover, for any $(\tau, \sigma) \in \mathcal{T}_{\theta} \times \mathcal{T}_{\theta}$

$$\mathbb{E}^{f}_{\theta,\tau\wedge\sigma^{\varepsilon}_{\theta}}J_{0}(\tau,\sigma^{\varepsilon}_{\theta}) - C\varepsilon \leq Y_{\theta} \leq \mathbb{E}^{f}_{\theta,\tau^{\varepsilon}_{\theta}\wedge\sigma}J_{0}(\tau^{\varepsilon}_{\theta},\sigma) + C\varepsilon,$$
(7.6)

where C is a constant depending only on $\lambda, \mu, \alpha, T, \|g\|_{L^1}, \gamma$.

Proof. Let $\theta \in \mathcal{T}$ and $\varepsilon > 0$. We shall prove that $(\tau_{\theta}^{\varepsilon}, \sigma_{\theta}^{\varepsilon})$ satisfies (7.6). By Lemma 6.3 Y is an \mathbb{E}^{f} -submartingale on $[[\theta, \tau_{\theta}^{\varepsilon}]]$. We thus have

$$Y_{\theta} \leq \mathbb{E}^{f}_{\theta, \tau^{\varepsilon}_{\theta} \wedge \sigma} [Y_{\tau^{\varepsilon}_{\theta} \wedge \sigma}].$$
(7.7)

By the assumptions made on L and Lemma 7.2, $Y_{\tau_{\theta}^{\varepsilon}} \leq L_{\tau_{\theta}^{\varepsilon}} + \varepsilon$. From this and the fact that $Y \leq U$ we have

$$Y_{\tau_{\theta}^{\varepsilon} \wedge \sigma} \leq (L_{\tau_{\theta}^{\varepsilon}} + \varepsilon) \mathbf{1}_{\{\tau_{\theta}^{\varepsilon} \leq \sigma < T\}} + U_{\sigma} \mathbf{1}_{\{\sigma < \tau_{\theta}^{\varepsilon}\}} + \xi \mathbf{1}_{\{\tau_{\theta}^{\varepsilon} = \sigma = T\}} \leq J_0(\tau_{\theta}^{\varepsilon}, \sigma) + \varepsilon.$$

Applying (7.7) and properties of the operator \mathbb{E}^{f} (see Proposition 5.2 (ii) and (v)) yields

$$Y_{\theta} \leq \mathbb{E}^{f}_{\theta, \tau^{\varepsilon}_{\theta} \wedge \sigma} (J_{0}(\tau^{\varepsilon}_{\theta}, \sigma) + \varepsilon) \leq \mathbb{E}^{f}_{\theta, \tau^{\varepsilon}_{\theta} \wedge \sigma} J_{0}(\tau^{\varepsilon}_{\theta}, \sigma) + C\varepsilon.$$
(7.8)

By Lemma 6.3 Y is an \mathcal{E}^{f} -supermartingale on $[[\theta, \sigma_{\theta}^{\varepsilon}]]$. As a result

$$Y_{\theta} \ge \mathbb{E}^{f}_{\theta, \tau \land \sigma_{\theta}^{\varepsilon}}(Y_{\tau \land \sigma_{\theta}^{\varepsilon}}).$$

$$(7.9)$$

By the assumptions made on U and Lemma 7.2 we have $Y_{\sigma_{\theta}^{\varepsilon}} \ge U_{\sigma_{\theta}^{\varepsilon}} - \varepsilon$. Applying analogous arguments as in case of L yields $Y_{\theta} \ge \mathbb{E}^{f}_{\theta, \tau \land \sigma_{\theta}^{\varepsilon}} J_{0}(\tau, \sigma_{\theta}^{\varepsilon}) - C\varepsilon$, which combined with (7.8) gives (7.6). Consequently,

$$Y_{\theta} \leq \operatorname{ess\,inf}_{\sigma \geq \theta} \mathbb{E}^{f}_{\theta, \tau^{\varepsilon}_{\theta} \wedge \sigma} J_{0}(\tau^{\varepsilon}_{\theta}, \sigma) + \varepsilon \leq \operatorname{ess\,sup}_{\tau \geq \theta} \operatorname{ess\,inf}_{\sigma \geq \theta} \mathbb{E}^{f}_{\theta, \tau \wedge \sigma} J_{0}(\tau, \sigma) + \varepsilon,$$

which combined with the definition of $\underline{V}_0(\theta)$ yields $Y_{\theta} \leq \underline{V}(\theta) + \varepsilon$. Analogous reasoning gives $\overline{V}(\theta) - \varepsilon \leq Y_{\theta}$. Letting $\varepsilon \to 0$ we find that $\overline{V}(\theta) \leq Y_{\theta} \leq \underline{V}(\theta)$, which combined with the obvious inequality $\underline{V}(\theta) \leq \overline{V}(\theta)$ gives $\underline{V}(\theta) = Y_{\theta} = \overline{V}_{\theta}$.

8. EXISTENCE OF SADDLE POINTS.

In the whole section, we assume that (H1)-(H5), (Z) are in force and that L and U are \mathbb{F} -optional processes of class (D).

Let (Y, Z, R) be a solution to $\operatorname{RBSDE}^T(\xi, f, L, U)$. We shall prove that there exists a saddle point for a nonlinear Dynkin game with sufficiently regular payoffs. For $\theta \in \mathcal{T}$ we define:

$$\tau_{\theta}^* \coloneqq \inf\{t \ge \theta, Y_t = L_t^{\xi}\} \land T; \quad \sigma_{\theta}^* \coloneqq \inf\{t \ge \theta, Y_t = U_t^{\xi}\} \land T$$

$$(8.1)$$

and

$$\bar{\tau}_{\theta} \coloneqq \inf\{t \ge \theta, R_t^+ > R_{\theta}^+\} \wedge T; \quad \bar{\sigma}_{\theta} \coloneqq \inf\{t \ge \theta, R_t^- > R_{\theta}^-\} \wedge T.$$

$$(8.2)$$

Proposition 8.1. Assume that L is right upper semicontinuous and U is right lower semicontinuous. Let (Y, Z, R) be a solution to $\text{RBSDE}^T(\xi, f, L, U)$ and let $\theta \in \mathcal{T}$.

1) If $R^{-,*}$ is continuous, then Y is an \mathbb{E}^{f} -supermartingale on $[[\theta, \overline{\sigma}_{\theta}]]$. Moreover,

$$Y_{\sigma_{\theta}^{*}} = U_{\sigma_{\theta}^{*}}^{\xi} \quad \text{and} \quad Y_{\bar{\sigma}_{\theta}} = U_{\bar{\sigma}_{\theta}}^{\xi}.$$

$$(8.3)$$

2) If $R^{+,*}$ is continuous, then Y is an \mathbb{E}^{f} -submartingale on $[[\theta, \overline{\tau}_{\theta}]]$. Moreover,

$$Y_{\tau_{\theta}^{*}} = L_{\tau_{\theta}^{*}}^{\xi} \quad \text{and} \quad Y_{\bar{\tau}_{\theta}} = L_{\bar{\tau}_{\theta}}^{\xi}.$$

$$(8.4)$$

Proof. Ad 1). Assume that $R^{-,*}$ is continuous. By the definition of $\bar{\sigma}_{\theta}$ we have that $R^{-}_{\bar{\sigma}_{\theta}} = R^{-}_{\theta}$. Thus, for any $a \ge 0$,

$$Y_t = Y_{\bar{\sigma}_{\theta}} + \int_t^{\bar{\sigma}_{\theta}} f(r, Y_r, Z_r) dr + R^+_{\bar{\sigma}_{\theta}} - R^+_t - \int_t^{\bar{\sigma}_{\theta}} Z_r dB_r, \quad t \in [\theta, \bar{\sigma}_{\theta}].$$

By Proposition 5.2, Y is an \mathbb{E}^{f} -supermartingale on $[[\theta, \bar{\sigma}_{\theta}]]$. We shall prove that $Y_{\bar{\sigma}_{\theta}} = U_{\bar{\sigma}_{\theta}}^{\xi}$. Assume that $\bar{\sigma}_{\theta} < T$ (in case $\bar{\sigma}_{\theta} = T$ the desired equality is obvious). Suppose, by contradiction, that $P(Y_{\bar{\sigma}_{\theta}} < U_{\bar{\sigma}_{\theta}}^{\xi}) > 0$. By the minimality condition, $\Delta^{+}R_{\bar{\sigma}_{\theta}}^{-} = 0$ on $\{Y_{\bar{\sigma}_{\theta}} < U_{\bar{\sigma}_{\theta}}^{\xi}\}$. Observe that $\Delta^{+}Y_{\bar{\sigma}_{\theta}} = -(\Delta^{+}R_{\bar{\sigma}_{\theta}}^{-} - \Delta^{+}R_{\bar{\sigma}_{\theta}}^{-}) = -\Delta^{+}R_{\bar{\sigma}_{\theta}}^{+} \leq 0$, which implies that Y is right upper semicontinuous on $\{Y_{\bar{\sigma}_{\theta}} < U_{\bar{\sigma}_{\theta}}^{\xi}\}$. Let $a \in \mathbb{R}$ and $\varepsilon > 0$ (depending on $\omega \in \Omega$) be such that $U_{\bar{\sigma}_{\theta}}^{\xi} > a + \varepsilon$ and $Y_{\bar{\sigma}_{\theta}} < a - \varepsilon$. Since Y is right upper semicontinuous on $\{Y_{\bar{\sigma}_{\theta}} < U_{\bar{\sigma}_{\theta}}^{\xi}\}$, and U is right lower semicontinuous, there exists $\delta > 0$ (depending on $\omega \in \Omega$) such that $U_{\bar{\sigma}_{\theta}+s}^{\xi} > a + \varepsilon$ and $Y_{\bar{\sigma}_{\theta}+s} < a - \varepsilon$, $s \in [0, \delta]$. Furthermore, from the definition of $\bar{\sigma}_{\theta}$ we have $R_{\bar{\sigma}_{\theta}+s}^{-,*} > R_{\bar{\sigma}_{\theta}}^{-,*}$. Consequently, on the set $\{Y_{\bar{\sigma}_{\theta}} < U_{\bar{\sigma}_{\theta}}^{\xi}\}$ the following holds

$$\int_{\bar{\sigma}_{\theta}}^{\bar{\sigma}_{\theta}+\delta} (\underline{U}_{r}^{\xi}-Y_{r-}) dR_{r}^{-,*} > 2\varepsilon (R_{\bar{\sigma}_{\theta}+\delta}^{-,*}-R_{\bar{\sigma}_{\theta}}^{-,*}) > 0$$

This contradicts the minimality condition.

What is left is to show that $Y_{\sigma_{\theta}^{*}} = U_{\sigma_{\theta}^{*}}^{\xi}$. In case $\sigma_{\theta}^{*} = T$ the equation follows at once. Suppose that $\sigma_{\theta}^{*} < T$. If $\Delta^{+}R_{\sigma_{\theta}^{*}}^{-}(\omega) > 0$, then by the very definition of a solution to RBSDE, we have $Y_{\sigma_{\theta}^{*}}(\omega) = U_{\sigma_{\theta}^{*}}^{\xi}(\omega)$. Suppose that $\Delta^{+}R_{\sigma_{\theta}^{*}}^{-}(\omega) = 0$. Observe that

$$\Delta^+ Y_{\sigma_\theta^*}(\omega) = -(\Delta_{\sigma_\theta^*}^+ R_{\sigma_\theta^*}^+(\omega) - \Delta^+ R_{\sigma_\theta^*}^-(\omega)) = -\Delta_{\sigma_\theta^*}^+ R_{\sigma_\theta^*}^+(\omega) \le 0.$$

Thus, $Y_{\sigma_{\theta}^{*}+}(\omega) \leq Y_{\sigma_{\theta}^{*}}(\omega)$. By the definition of σ_{θ}^{*} there exists a non-increasing sequence $(t_{n}(\omega))_{n\geq 1}$ such that $t_{n}(\omega) \searrow \sigma_{\theta}^{*}(\omega)$ and $Y_{t_{n}}(\omega) = U_{t_{n}}^{\xi}(\omega)$. Letting $n \to \infty$ and using right lower semicontinuity of U we find that $Y_{\sigma_{\theta}^{*}+}(\omega) \geq U_{\sigma_{\theta}^{*}}^{\xi}(\omega)$, which combined with $Y_{\sigma_{\theta}^{*}+}(\omega) \leq Y_{\sigma_{\theta}^{*}}(\omega)$ gives the result.

Ad A2). The case when $R^{+,*}$ is supposed to be continuous runs analogously.

Corollary 8.2. Under assumptions of Proposition 8.1 we have that continuity of $R^{-,*}$ (resp. $R^{+,*}$) implies $\sigma_{\theta}^* \leq \bar{\sigma}_{\theta}$ (resp. $\tau_{\theta}^* \leq \bar{\tau}_{\theta}$).

Proposition 8.3. Let (Y, Z, R) be a solution to $\text{RBSDE}^T(\xi, f, L, U)$. If L (resp. U) is left upper semicontinuous (resp. left lower semicontinuous), then $R^{+,*}$ (resp. $R^{-,*}$) is continuous.

Proof. Let $\tau \in \mathcal{T}$ be predictable. We shall prove that $\Delta^{-}R_{\tau}^{+,*} = 0$. We have

$$\Delta Y_{\tau} = -\Delta R_{\tau}^{+,*} + \Delta R_{\tau}^{-,*} = -\Delta R_{\tau}^{+,*} \mathbf{1}_{\{Y_{\tau-} = \overrightarrow{L}_{\tau}\} \cap D} + \Delta R_{\tau}^{-,*} \mathbf{1}_{\{Y_{\tau-} = \underbrace{U}_{\tau}\} \cap D'}, \tag{8.5}$$

where $D := \{\Delta R_{\tau}^{+,*} > 0\}$ and $D' := \{\Delta R_{\tau}^{-,*} > 0\}$. Since $dR^+ \perp dR^-$, $D \cap D' = \emptyset$. Thus, on the set $D, \Delta Y_{\tau} \leq 0$. From this and the regularity assumption on $L, \vec{L}_{\tau} \leq L_{\tau} \leq Y_{\tau} \leq Y_{\tau-}$ on D. Consequently, $\Delta Y_{\tau} = 0$ on $\{Y_{\tau-} = \vec{L}_{\tau}\} \cap D$. This combined with (8.5) implies $\Delta^- R_{\tau}^{+,*} = 0$. Since the last inequality holds for any predictable $\tau \in \mathcal{T}$, we deduce that $R^{+,*}$ is continuous. The similar reasoning may be applied to U.

Theorem 8.4. Suppose that L is upper semicontinuous and U is lower semicontinuous. Let (Y, Z, R) be a solution to $\text{RBSDE}^T(\xi, f, L, U)$. Then for any $\theta \in \mathcal{T}$ couples (8.1) and (8.2) are saddle points at θ for the nonlinear Dynkin game with the payoff function (7.1).

Proof. Let $\theta \in \mathcal{T}$. By Theorem 7.3 $Y_{\theta} = \overline{V}_0(\theta) = \underline{V}_0(\theta)$. By Proposition 8.3 $R^{+,*}$, $R^{-,*}$ are continuous. Let $\tau \in \mathcal{T}_{\theta}$. Since $\sigma_{\theta}^* \leq \overline{\sigma}_{\theta}$ (see Corollary 8.2), by Proposition 8.1 process Y is an \mathbb{E}^f -supermartingale on $[[\theta, \tau \wedge \sigma_{\theta}^*]]$. Therefore,

$$Y_{\theta} \ge \mathbb{E}^{f}_{\theta, \tau \wedge \sigma^{*}_{\theta}} [Y_{\tau \wedge \sigma^{*}_{\theta}}].$$

$$(8.6)$$

Since $Y \ge L$ and $Y_{\sigma_{\alpha}^*} = U_{\sigma_{\alpha}^*}$ (see Proposition 8.1), we also have

$$Y_{\tau \wedge \sigma_{\theta}^*} = Y_{\tau} \mathbf{1}_{\{\tau \leq \sigma_{\theta}^*\}} + Y_{\sigma_{\theta}^*} \mathbf{1}_{\{\sigma_{\theta}^* < \tau\}} \ge L_{\tau} \mathbf{1}_{\{\tau \leq \sigma_{\theta}^*\}} + U_{\sigma_{\theta}^*} \mathbf{1}_{\{\sigma_{\theta}^* < \tau\}} = J_0(\tau, \sigma_{\theta}^*).$$

Using (8.6) and the fact that \mathbb{E}^{f} is a non-decreasing operator, we deduce that $Y_{\theta} \geq \mathbb{E}^{f}_{\theta, \tau \wedge \sigma_{\theta}^{*}} J_{0}(\tau, \sigma_{\theta}^{*})$ for any $\tau \in \mathcal{T}_{\theta}$, in particular $\mathbb{E}^{f}_{\theta, \tau_{\theta}^{*} \wedge \sigma_{\theta}^{*}} J_{0}(\tau_{\theta}^{*}, \sigma_{\theta}^{*}) \leq Y_{\theta}$. In the similar way we arrive at $Y_{\theta} \leq \mathbb{E}^{f}_{\theta, \tau_{\theta}^{*} \wedge \sigma} J_{0}(\tau_{\theta}^{*}, \sigma)$ for any $\sigma \in \mathcal{T}_{\theta}$, and so $Y_{\theta} \leq \mathbb{E}^{f}_{\theta, \tau_{\theta}^{*} \wedge \sigma_{\theta}^{*}} J_{0}(\tau_{\theta}^{*}, \sigma_{\theta}^{*})$. Consequently, $Y_{\theta} = \mathbb{E}^{f}_{\theta, \tau_{\theta}^{*} \wedge \sigma_{\theta}^{*}} J(\tau_{\theta}^{*}, \sigma_{\theta}^{*})$ and $(\tau_{\theta}^{*}, \sigma_{\theta}^{*})$ is a saddle point at θ . Analogously, one shows, by using Proposition 8.1, that $(\bar{\tau}_{\theta}, \bar{\sigma}_{\theta})$ is a saddle point at θ .

9. Existence result

In the whole section, we assume that L, U are \mathbb{F} -optional processes of class (D).

Let us consider the following assumption, which is called in the literature *weak Moko-bodzki's condition*.

(WM) There exists a semimartingale X such that $L \leq X \leq U$.

Proposition 9.1. Assume that L, U are left-limited, and

$$L_t < U_t, \quad L_{t-} < U_{t-}, \quad t \in [0, T].$$
 (9.1)

Then weak Mokobodzki's condition (WM) holds for L, U.

Proof. We let $\tau_0 \coloneqq 0$, and for $n \ge 1$,

$$\tau_n \coloneqq \inf\{t > \tau_{n-1} : (\overleftarrow{L}_{\tau_{n-1}} + \underbrace{U}_{\tau_{n-1}}) < 2L_t \text{ or } (\overleftarrow{L}_{\tau_{n-1}} + \underbrace{U}_{\tau_{n-1}}) > 2U_t\} \land T.$$

Obviously, (τ_n) is nondecreasing. Observe that by the definition of τ_n for each $\omega \in \Omega$ there exists a sequence $\{t_m^n\}$ such that $t_m^n \searrow \tau_n(\omega)$ and for all $m \in \mathbb{N}$

$$\overleftarrow{L}_{\tau_{n-1}(\omega)}(\omega) + \underbrace{U}_{\tau_{n-1}(\omega)}(\omega) < 2L_{t_m^n}(\omega) \quad \text{or} \\
\overleftarrow{L}_{\tau_{n-1}(\omega)}(\omega) + \underbrace{U}_{\tau_{n-1}(\omega)}(\omega) > 2U_{t_m^n}(\omega).$$
(9.2)

Letting $m \to \infty$ yields

$$\begin{aligned}
\overleftarrow{L}_{\tau_{n-1}(\omega)}(\omega) + \underbrace{U}_{\tau_{n-1}(\omega)}(\omega) &\leq 2\overleftarrow{L}_{\tau_n(\omega)}(\omega) \quad \text{or} \\
\overleftarrow{L}_{\tau_{n-1}(\omega)}(\omega) + \underbrace{U}_{\tau_{n-1}(\omega)}(\omega) &\geq 2\underbrace{U}_{\tau_n(\omega)}(\omega), \quad n \geq 1.
\end{aligned}$$
(9.3)

Step 1. We shall prove that (τ_n) is a chain. First, note that

$$P(\tau_{n-1} = \tau_n < T) = 0, \quad n \ge 1.$$
(9.4)

Indeed, suppose that for some $n \ge 1$, $P(\tau_{n-1} = \tau_n < T) > 0$. Let $\omega \in \{\tau_{n-1} = \tau_n < T\}$. Then, by (9.3)

$$\underbrace{U}_{\tau_n(\omega)}(\omega) \leq \overline{L}_{\tau_n(\omega)}(\omega).$$

Therefore, $P(\underbrace{U}_{\tau_n} \leq \overleftarrow{L}_{\tau_n}) > 0$, which contradicts (9.1). Suppose that (τ_n) is not a chain. Then, according to (9.4), there must exist $\tau \in \mathcal{T}$ such that $\tau_n \nearrow \tau$ and $P(\bigcap_{n=1}^{\infty} \{\tau_n < \tau\}) > 0$. Let $\omega \in \bigcap_{n=1}^{\infty} \{\tau_n < \tau\}$. By the second inequality in (9.1) for any $\delta > 0$ there exists $n_{\delta} \geq 1$ such that

$$\overleftarrow{L}_{\tau_{n-1}(\omega)}(\omega) \leq L_{\tau(\omega)-}(\omega) + \delta, \quad U_{\tau(\omega)-}(\omega) - \delta \leq \underbrace{U}_{\tau_{n-1}(\omega)}(\omega), \quad n \geq n_{\delta}.$$

Suppose that the first inequality in (9.3) holds for infinitely many $n \ge 1$ (the proof in the second case is analogous). Then, by the above inequalities, we conclude from (9.3) that

$$L_{\tau(\omega)-}(\omega) + U_{\tau(\omega)-}(\omega) - 2\delta \le L_{\tau(\omega)-}(\omega) + \delta.$$

Letting $\delta \searrow 0$, we obtain that $U_{\tau(\omega)-}(\omega) \le L_{\tau(\omega)-}(\omega)$. Therefore, $P(U_{\tau-} \le L_{\tau-}) > 0$, which contradicts (9.1). Thus, (τ_n) is a chain.

Step 2. We shall construct a semimartingale lying between barriers L, U. Define

$$X_{t} \coloneqq \frac{1}{2} \sum_{n=1}^{\infty} \Big((\overleftarrow{L}_{\tau_{n-1}} + \underbrace{U}_{\tau_{n-1}}) \mathbf{1}_{(\tau_{n-1},\tau_{n})}(t) + (L_{\tau_{n-1}} + U_{\tau_{n-1}}) \mathbf{1}_{\{\tau_{n-1}\}}(t) \Big),$$

for $t \in [0,T]$. Clearly, $L_t \leq X_t \leq U_t$, $t \in [0,T]$ and X is F-adapted. Since (τ_n) is a chain, we get that X is of finite variation, thus a semimartingale. This completes the proof.

Proposition 9.2. Assume that f_1, f_2 satisfy (H1)–(H4), ξ_1, ξ_2, f_1, f_2 satisfy (H5) with p = 2, and $|f^1 - f^2|(\cdot, Y^2, Z^2) \in L^{1,2}_{\mathbb{F}}(0,T)$. Let (Y^i, Z^i, R^i) be a solution to $\text{RBSDE}^T(\xi_i, f^i, L, U)$ such that $Y^i \in S^2_{\mathbb{F}}(0,T)$, i = 1, 2. Then there exists c > 0, depending only on T, μ, λ , such that

$$\|Y^{1} - Y^{2}\|_{\mathcal{D}^{2}(0,T)} \le c \Big(\|\xi_{1} - \xi_{2}\|_{L^{2}} + \||f_{1} - f_{2}|(\cdot, Y^{2}, Z^{2})\|_{L^{1,2}_{\mathbb{F}}(0,T)}\Big).$$

$$(9.5)$$

Proof. We let J^i denote the right-hand side of (6.1) but with ξ replaced by ξ_i , i = 1, 2. Let

$$\tilde{f}(t,y,z) \coloneqq f_1(t,Y_t^1,Z_t^1) - f_2(t,Y_t^1,Z_t^1) + f_2(t,y,z), \quad t \in [0,T], \ y \in \mathbb{R}, \ z \in \mathbb{R}^d.$$

Observe that (Y^1, Z^1, R^1) is a solution to $\text{RBSDE}^T(\xi_1, \tilde{f}, L, U)$. By Theorem 6.6

$$Y_{\theta}^{1} = \operatorname{ess\,sup\,ess\,inf}_{\tau \mid A \in \mathcal{U}_{\theta}} \mathbb{E}_{\theta, \tau \wedge \sigma}^{f} J^{1}(\tau \mid A, \sigma \mid B)$$

and

$$Y_{\theta}^{2} = \operatorname{ess\,sup\,ess\,inf}_{\tau \lfloor A \in \mathcal{U}_{\theta}} \mathbb{E}_{\theta, \tau \wedge \sigma}^{f_{2}} J^{2}(\tau \lfloor A, \sigma \lfloor B).$$

Hence

$$Y_{\theta}^{1} - Y_{\theta}^{2} \leq \underset{\tau \mid A \in \mathcal{U}_{\theta}}{\operatorname{ess \, sup \, ess \, sup \, ess \, sup \, }} \left| \mathbb{E}_{\theta, \tau \wedge \sigma}^{\tilde{f}} J^{1}(\tau \mid A, \sigma \mid B) - \mathbb{E}_{\theta, \tau \wedge \sigma}^{f_{2}} J^{2}(\tau \mid A, \sigma \mid B) \right|.$$
(9.6)

Applying Proposition 5.2(vi) yields

$$\mathbb{E} \Big| \mathbb{E}_{\theta,\tau\wedge\sigma}^{\tilde{f}} J^{1}(\tau \lfloor A, \sigma \lfloor B) - \mathbb{E}_{\theta,\tau\wedge\sigma}^{f_{2}} J^{2}(\tau \lfloor A, \sigma \lfloor B) \Big|^{2} \\ \leq c \mathbb{E} \Big[|\xi_{1} - \xi_{2}|^{2} + \Big(\int_{0}^{T} |f_{1} - f_{2}|(r, Y_{r}^{2}, Z_{r}^{2}) dr \Big)^{2} \Big].$$

Combining the last two inequalities gives at once the result.

Theorem 9.3. Assume that (H1)-(H4),(H7) are in force. Suppose that (H5) is satisfied with p = 1. Then there exists a solution (Y, Z, R) to $\text{RBSDE}^T(\xi, f, L, U)$.

Proof. Let X be the process appearing in (H7). Since X is a special semimartingale, there exists a chain $(\hat{\gamma}_k)$ and processes $H \in \mathcal{H}_{\mathbb{F}}(0,T)$ and $C \in \mathcal{V}_{\mathbb{F}}(0,T)$ such that $H \in \mathcal{H}^2_{\mathbb{F}}(0,\hat{\gamma}_k)$, $C \in \mathcal{V}^2_{\mathbb{F}}(0,\hat{\gamma}_k)$, $k \ge 1$, and

$$X_t = X_0 + C_t + \int_0^t H_r \, dB_r, \quad t \in [0, T].$$

Let

$$f_{n,m}(t,y,z) = \{f(t,y,z) \land n\} \lor (-m).$$

Note that $f_{n,m}$ is nondecreasing with respect to n and nonincreasing with respect to m. Moreover, $f_{n,m}(t,y,z) \nearrow f_m(t,y,z) \coloneqq f(t,y,z) \lor (-m)$, when $n \to \infty$, and $f_m(t,y,z) \searrow f(t,y,z)$, when $m \to \infty$. By [8, page 417] there exist regulated processes \hat{L} , \hat{U} satisfying

$$\hat{L}_{\alpha} = \operatorname{ess\,inf}_{\tau \in \mathcal{T}_{\alpha}} \mathbb{E}(L_{\tau} | \mathcal{F}_{\alpha}), \quad \hat{U}_{\alpha} = \operatorname{ess\,sup}_{\tau \in \mathcal{T}_{\alpha}} \mathbb{E}(U_{\tau} | \mathcal{F}_{\alpha}), \, \alpha \in \mathcal{T}$$

Furthermore, by [27, Proposition 3.8], $-\hat{L}$, \hat{U} are supermartingales of class (D) on [0,T]. As a result, there exist processes $F, G \in \mathcal{H}_{\mathbb{F}}(0,T)$ and $A, D \in \mathcal{V}_{\mathbb{F}}^{+,1}(0,T)$ such that

$$\hat{L}_t = \hat{L}_T - \int_t^T dA_r - \int_t^T F_r \, dB_r, \quad t \in [0, T].$$

and

$$\hat{U}_t = \hat{U}_T - \int_t^T dD_r - \int_t^T G_r \, dB_r, \quad t \in [0, T].$$

Obviously, $\hat{L} \leq L \leq U \leq \hat{U}$. Since \hat{L} and \hat{U} are of class (D), by (H4) there exists a chain (τ_k^1) on [0,T] such that

$$\mathbb{E}\Big(\int_{0}^{\tau_{k}} |f(r,\hat{L}_{r},0)| \, dr\Big)^{2} + \mathbb{E}\Big(\int_{0}^{\tau_{k}} |f(r,\hat{U}_{r},0)| \, dr\Big)^{2} \le k.$$
(9.7)

Moreover, let us consider chain (τ_k^2) on [0,T] such that $\hat{L}, \hat{U} \in \mathcal{S}^2_{\mathbb{F}}(0,\tau_k^2), f(\cdot,0,0) \in L^{1,2}_{\mathbb{F}}(0,\tau_k^2)$, and $A, D \in \mathcal{V}^2_{\mathbb{F}}(0,\tau_k^2), k \ge 1$. We let $\gamma_k \coloneqq \hat{\gamma}_k \wedge \tau_k^1 \wedge \tau_k^2$ Define

$$L_t^n = L_t \mathbf{1}_{\{t \le \gamma_n\}} + \hat{L} \mathbf{1}_{\{t > \gamma_n\}}, \quad U_t^n = U_t \mathbf{1}_{\{t \le \gamma_n\}} + \hat{U}_t \mathbf{1}_{\{t > \gamma_n\}}$$

Note that

$$\hat{L} \le L^n \le L^{n+1} \le L \le U \le U^{n+1} \le U^n \le \hat{U}, \quad n \ge 1.$$

$$(9.8)$$

Moreover, $L^n \nearrow L$ and $U^n \searrow U$. Finally, we define

$$X_t^{n,m} = X_t \mathbf{1}_{\{t \le \gamma_n \land \gamma_m\}} + \hat{L}_t \mathbf{1}_{\{t > \gamma_n \ge \gamma_m\}} + \hat{U}_t \mathbf{1}_{\{t > \gamma_m > \gamma_n\}}$$

Note that $L^n \leq X^{n,m} \leq U^m$ and $X^{n,m}$ is a difference of two supermartingales of class (D). Therefore, by the definition of $f_{n,m}$, strong Mokobodzki's condition (H6^{*}) is satisfied with $L^n, U^m, X^{n,m}$ and $f_{n,m}$. By Theorem 4.3 there exists a unique solution $(Y^{n,m}, Z^{n,m}, R^{n,m})$ to $\text{RBSDE}^T(\xi, f_{n,m}, L^n, U^m)$ such that $Y^{n,m}$ is of class (D), $Z^{n,m} \in \mathcal{H}^q_{\mathbb{F}}(0,T)$, $q \in (0,1)$ and $R^{n,m} \in \mathcal{V}^1_{0,\mathbb{F}}(0,T)$. By [28, Proposition 3.2, Lemma 3.3], $Y^{n,m}$ is nondecreasing with respect to n and nonincreasing with respect to m. Let us define

$$Y^m = \sup_{n \ge 1} Y^{n,m}, \quad Y = \inf_{m \ge 1} Y^m$$

Obviously, Y^m and Y are of class (D). The remainder of the proof, we divide into two steps.

Step 1. We shall prove that or any $k \leq m$, process Y^m is the first component of a solution to $\operatorname{RBSDE}^{\gamma_k}(Y^m_{\gamma_k}, f_m, L, U^m)$. Let $k \leq m \leq n$. Since $\hat{L} \leq Y^{n,m} \leq \hat{U}$, we have $Y^{n,m}, Y^m \in \mathcal{S}^2_{\mathbb{F}}(0,\gamma_k)$. According to Theorem 4.2 - observe that (H5), (H6) are satisfied with L, U^m, f_m, Y^m, X and p = 2 on $[[0, \gamma_k]]$ - there exists a solution $(\tilde{Y}^{k,m}, \tilde{Z}^{k,m}, \tilde{R}^{k,m})$ to $\operatorname{RBSDE}^{\gamma_k}(Y^m_{\gamma_k}, f_m, L, U^m)$ such that $\tilde{Y}^{k,m} \in \mathcal{S}^2_{\mathbb{F}}(0, \gamma_k)$, $\tilde{Z}^{k,m} \in \mathcal{H}^2_{\mathbb{F}}(0, \gamma_k)$ and $\tilde{R}^{k,m} \in \mathcal{V}^2_{0,\mathbb{F}}(0, \gamma_k)$. We shall show that $Y^m = \tilde{Y}^{k,m}$ on $[[0, \gamma_k]]$. By Proposition 9.2

$$\begin{split} \|\tilde{Y}^{k,m} - Y^{n,m}\|_{\mathcal{D}^{2}(0,\gamma_{k})}^{2} &\leq c \mathbb{E}\Big[\Big(\int_{0}^{\gamma_{k}} |f_{m} - f_{n,m}|(r,\tilde{Y}_{r}^{k,m},\tilde{Z}_{r}^{k,m}) dr\Big)^{2}\Big] \\ &+ |\tilde{Y}_{\gamma_{k}}^{k,m} - Y_{\gamma_{k}}^{n,m}|^{2} = c \mathbb{E}\Big[|Y_{\gamma_{k}}^{m} - Y_{\gamma_{k}}^{n,m}|^{2} \\ &+ \Big(\int_{0}^{\gamma_{k}} |f(r,\tilde{Y}_{r}^{k,m},\tilde{Z}_{r}^{k,m})| \mathbf{1}_{\{f(r,\tilde{Y}_{r}^{k,m},\tilde{Z}_{r}^{k,m})>n\}} dr\Big)^{2}\Big]. \end{split}$$
(9.9)

Observe that $0 \leq Y_{\gamma_k}^m - Y_{\gamma_k}^{n,m} \leq Y_{\gamma_k}^m \in L^2(\Omega, \mathcal{F}_{\gamma_k})$ (the last assertion is a consequence of the fact that $Y^m \in \mathcal{S}^2_{\mathbb{F}}(0, \gamma_k)$). Therefore, by the Lebesgue dominated convergence theorem,

the first term on the right-hand side of (9.9) tends to zero as $n \to \infty$. Note that, by the definition of γ_k , (H1), (9.7) and Jensen's inequality

$$\begin{split} \mathbb{E}\Big(\int_{0}^{\gamma_{k}}|f(r,\tilde{Y}_{r}^{k,m},\tilde{Z}_{r}^{k,m})|\,dr\Big)^{2} &\leq \mathbb{E}\Big(\lambda\int_{0}^{\gamma_{k}}|\tilde{Z}_{r}^{k,m}|\,dr\Big)^{2} \\ &+ \mathbb{E}\Big(\int_{0}^{\gamma_{k}}|f(r,\tilde{Y}_{r}^{k,m},0)|\,dr\Big)^{2} \leq T\lambda^{2}\mathbb{E}\int_{0}^{\gamma_{k}}|\tilde{Z}_{r}^{k,m}|^{2}\,dr \\ &+ \mathbb{E}\Big(\int_{0}^{\gamma_{k}}|f(r,\hat{L}_{r},0)|+|f(r,\hat{U}_{r},0)|\,dr\Big)^{2} \\ &\leq T\lambda^{2}\mathbb{E}\int_{0}^{\gamma_{k}}|\tilde{Z}_{r}^{k,m}|^{2}\,dr+T^{2}\cdot k^{2}<\infty. \end{split}$$

Consequently, by the Lebesgue dominated convergence theorem, the most right term in (9.9) tends to zero as $n \to \infty$. As a result, letting $n \to \infty$ in (9.9), we obtain that $Y^{n,m} \to \tilde{Y}^{k,m}$ in $\mathcal{D}^2_{\mathbb{R}}(0,\gamma_k)$. This completes the proof of step 1.

Step 2. We shall prove that Y is the first component of a solution to RBSDE(ξ, f, L, U). Let $k \leq m$. Since $\hat{L} \leq Y^m \leq \hat{U}$, we have that $Y^m, Y \in S^2_{\mathbb{F}}(0, \gamma_k)$. Observe that conditions (H5) and (H6) are met by L, U, f, X on $[[0, \gamma_k]]$ with p = 2. Therefore, by Theorem 4.2, there exists a solution $(\tilde{Y}^k, \tilde{Z}^k, \tilde{R}^k)$ to RBSDE^{γ_k} (Y_{γ_k}, f, L, U) such that $\tilde{Y}^k \in S^2_{\mathbb{F}}(0, \gamma_k)$, $\tilde{Z}^k \in \mathcal{H}^2_{\mathbb{F}}(0, \gamma_k)$, and $\tilde{R}^k \in \mathcal{V}^2_{0,\mathbb{F}}(0, \gamma_k)$. We shall show that $Y = \tilde{Y}^k$ on $[[0, \gamma_k]]$. By Proposition 9.2

$$\begin{split} \|\tilde{Y}^{k} - Y^{m}\|_{\mathcal{D}^{2}(0,\gamma_{k})} &\leq C\mathbb{E}\Big[\Big(\int_{0}^{\gamma_{k}} |f(r,\tilde{Y}^{k}_{r},\tilde{Z}^{k}_{r}) - f_{m}(r,\tilde{Y}^{k}_{r},\tilde{Z}^{k}_{r})|\,dr\Big)^{2} \\ &+ |Y_{\gamma_{k}} - Y^{m}_{\gamma_{k}}|^{2}\Big] = C\mathbb{E}\Big[\Big(\int_{0}^{\gamma_{k}} |f(r,\tilde{Y}^{k}_{r},\tilde{Z}^{k}_{r})|\mathbf{1}_{\{f(r,\tilde{Y}^{k}_{r},\tilde{Z}^{k}_{r})<-m\}}\,dr\Big)^{2} \\ &+ |Y_{\gamma_{k}} - Y^{m}_{\gamma_{k}}|^{2}\Big]. \end{split}$$
(9.10)

By the Lebesgue dominated convergence theorem the first term on the right-hand side of (9.10) tends to zero when $m \to \infty$. By combining (9.7), the definition of γ_k , condition (H1), and Jensen's inequality, we conclude that

$$\mathbb{E}\Big(\int_{0}^{\gamma_{k}} |f(r, \tilde{Y}_{r}^{k}, \tilde{Z}_{r}^{k})| dr\Big)^{2} \leq \mathbb{E}\Big(\lambda \int_{0}^{\gamma_{k}} |\tilde{Z}_{r}^{k}| dr\Big)^{2} + \mathbb{E}\Big(\int_{0}^{\gamma_{k}} |f(r, \tilde{Y}_{r}^{k}, 0)| dr\Big)^{2} \\
\leq T\lambda^{2} \mathbb{E} \int_{0}^{\gamma_{k}} |\tilde{Z}_{r}^{k}|^{2} dr + \mathbb{E}\Big(\int_{0}^{\gamma_{k}} |f(r, \hat{L}_{r}, 0)| + |f(r, \hat{U}_{r}, 0)| dr\Big)^{2} \\
\leq T\lambda^{2} \mathbb{E} \int_{0}^{\gamma_{k}} |\tilde{Z}_{r}^{k}|^{2} dr + T^{2} \cdot k^{2} < \infty.$$

Therefore, by the Lebesgue dominated convergence theorem, the most right term in (9.10) tends to zero as $m \to \infty$. Consequently, letting $m \to \infty$ in (9.10), we deduce that $Y^m \to \tilde{Y}^k$ in $\mathcal{D}^2_{\mathbb{F}}(0,\gamma_k)$. Hence, $\tilde{Y}^k = Y$ on $[0,\gamma_k], k \ge 1$. In other words, for any $k \ge 1$, process Y is the first component of a solution to $\text{RBSDE}^{\gamma_k}(Y_{\gamma_k}, f, L, U)$. This in turn implies, by using the uniqueness argument (see Theorem 4.2), that $\tilde{Z}^k = \tilde{Z}^{k+1}$, and $\tilde{R}^k = \tilde{R}^{k+1}$ on $[0,\gamma_k], k \ge 1$. With the aid of these properties, one easily checks that the triple (Y, M, R) is a solution to $\text{RBSDE}^{\gamma_k}(Y_{\gamma_k}, f, L, U)$ for each $k \ge 1$, where

$$Z_t \coloneqq \sum_{k=0}^{\infty} \tilde{Z}_t^k \mathbf{1}_{(\gamma_k, \gamma_{k+1}]}(t), \quad R_t \coloneqq \sum_{k=0}^{\infty} \tilde{R}_t^k \mathbf{1}_{(\gamma_k, \gamma_{k+1}]}(t).$$

This combined with the fact that (γ_k) is a chain implies that (Y, Z, R) is a solution to $\text{RBSDE}^T(\xi, f, L, U)$.

Acknowledgements. T. Klimsiak is supported by Polish National Science Centre: Grant No. 2017/25/B/ST1/00878. M. Rzymowski acknowledges the support of the Polish National Science Centre: Grant No. 2018/31/N/ST1/00417.

References

- B. Baadi and Y. Ouknine. Reflected BSDEs when the obstacle is not right-continuous in a general filtration. ALEA Lat. Am. J. Probab. Math. Stat., 14(1):201–218, 2017.
- [2] B. Baadi and Y. Ouknine. Reflected BSDEs with optional barrier in a general filtration. Afr. Mat., 29(7-8):1049–1064, 2018.
- [3] E. Bayraktar and S. Yao. Doubly reflected BSDEs with integrable parameters and related Dynkin games. Stochastic Process. Appl., 125(12):4489–4542, 2015.
- [4] P. Briand, B. Delyon, Y. Hu, E. Pardoux, and L. Stoica. L^p solutions of backward stochastic differential equations. Stochastic Process. Appl., 108(1):109–129, 2003.
- [5] R. Buckdahn and J. Li. Probabilistic interpretation for systems of Isaacs equations with two reflecting barriers. NoDEA Nonlinear Differential Equations Appl., 16(3):381–420, 2009.
- [6] S. Crépey. *Financial modeling*. Springer Finance. Springer, Heidelberg, 2013. A backward stochastic differential equations perspective, Springer Finance Textbooks.
- [7] J. Cvitanić and I. Karatzas. Backward stochastic differential equations with reflection and Dynkin games. Ann. Probab., 24(4):2024–2056, 1996.
- [8] C. Dellacherie and P.-A. Meyer. Probabilities and potential. B, volume 72 of North-Holland Mathematics Studies. North-Holland Publishing Co., Amsterdam, 1982. Theory of martingales, Translated from the French by J. P. Wilson.
- [9] R. Dumitrescu, M.-C. Quenez, and A. Sulem. Generalized Dynkin games and doubly reflected BSDEs with jumps. *Electron. J. Probab.*, 21:Paper No. 64, 32, 2016.
- [10] R. Dumitrescu, M.-C. Quenez, and A. Sulem. Game options in an imperfect market with default. SIAM J. Financial Math., 8(1):532–559, 2017.
- [11] R. Dumitrescu, M.-C. Quenez, and A. Sulem. Mixed generalized Dynkin game and stochastic control in a Markovian framework. *Stochastics*, 89(1):400–429, 2017.
- [12] B. El Asri, S. Hamadène, and H. Wang. L^p-solutions for doubly reflected backward stochastic differential equations. Stoch. Anal. Appl., 29(6):907–932, 2011.
- [13] N. El Karoui, C. Kapoudjian, E. Pardoux, S. Peng, and M. C. Quenez. Reflected solutions of backward SDE's, and related obstacle problems for PDE's. Ann. Probab., 25(2):702–737, 1997.
- [14] M. Grigorova, P. Imkeller, E. Offen, Y. Ouknine, and M.-C. Quenez. Reflected BSDEs when the obstacle is not right-continuous and optimal stopping. Ann. Appl. Probab., 27(5):3153–3188, 2017.
- [15] M. Grigorova, P. Imkeller, Y. Ouknine, and M.-C. Quenez. Doubly reflected BSDEs and \mathcal{E}^{f} -Dynkin games: beyond the right-continuous case. *Electron. J. Probab.*, 23:Paper No. 122, 38, 2018.
- [16] M. Grigorova, P. Imkeller, Y. Ouknine, and M.-C. Quenez. Optimal stopping with *f*-expectations: the irregular case. *Stochastic Process. Appl.*, 130(3):1258–1288, 2020.
- [17] M. Grigorova and M.-C. Quenez. Optimal stopping and a non-zero-sum Dynkin game in discrete time with risk measures induced by BSDEs. *Stochastics*, 89(1):259–279, 2017.
- [18] S. Hamadène and M. Hassani. BSDEs with two reflecting barriers: the general result. Probab. Theory Related Fields, 132(2):237–264, 2005.
- [19] S. Hamadène and M. Hassani. BSDEs with two reflecting barriers driven by a Brownian and a Poisson noise and related Dynkin game. *Electron. J. Probab.*, 11:no. 5, 121–145, 2006.
- [20] S. Hamadène, M. Hassani, and Y. Ouknine. Backward SDEs with two *rcll* reflecting barriers without Mokobodski's hypothesis. *Bull. Sci. Math.*, 134(8):874–899, 2010.
- [21] S. Hamadène and I. Hdhiri. Backward stochastic differential equations with two distinct reflecting barriers and quadratic growth generator. J. Appl. Math. Stoch. Anal., pages Art. ID 95818, 28, 2006.

- [22] I. Hassairi. Existence and uniqueness for D-solutions of reflected BSDEs with two barriers without Mokobodzki's condition. Commun. Pure Appl. Anal., 15(4):1139–1156, 2016.
- [23] T. Klimsiak. BSDEs with monotone generator and two irregular reflecting barriers. Bull. Sci. Math., 137(3):268–321, 2013.
- [24] T. Klimsiak. Cauchy problem for semilinear parabolic equation with time-dependent obstacles: a BSDEs approach. *Potential Anal.*, 39(2):99–140, 2013.
- [25] T. Klimsiak. Non-semimartingale solutions of reflected BSDEs and applications to Dynkin games. Stochastic Process. Appl., 134:208–239, 2021.
- [26] T. Klimsiak and M. Rzymowski. Reflected BSDEs with two optional barriers and monotone coefficient on general filtered space. *Electron. J. Probab.*, 26:Paper No. 91, 24, 2021.
- [27] T. Klimsiak, M. Rzymowski, and L. Słomiński. Reflected BSDEs with regulated trajectories. Stochastic Process. Appl., 129(4):1153–1184, 2019.
- [28] T. Klimsiak, M. Rzymowski, and L. Słomiński. Reflected backward stochastic differential equations with two optional barriers. Bull. Sci. Math., 158:102820, 49, 2020.
- [29] M. Marzougue. A note on optional Snell envelopes and reflected backward SDEs. Statist. Probab. Lett., 165:108833, 7, 2020.
- [30] M. Marzougue and M. El Otmani. Non-continuous double barrier reflected BSDEs with jumps under a stochastic Lipschitz coefficient. Commun. Stoch. Anal., 12(4):Art. 1, 359–381, 2018.
- [31] M. Marzougue and M. El Otmani. BSDEs with right upper-semicontinuous reflecting obstacle and stochastic Lipschitz coefficient. Random Oper. Stoch. Equ., 27(1):27–41, 2019.
- [32] M. Marzougue and M. El Otmani. BSDEs with jumps and two completely separated irregular barriers in a general filtration. ALEA Lat. Am. J. Probab. Math. Stat., 18(1):761–792, 2021.
- [33] E. Pardoux and S. G. Peng. Adapted solution of a backward stochastic differential equation. Systems Control Lett., 14(1):55–61, 1990.
- [34] E. Pardoux and A. Răşcanu. Stochastic differential equations, backward SDEs, partial differential equations, volume 69 of Stochastic Modelling and Applied Probability. Springer, Cham, 2014.
- [35] S. Peng. Backward SDE and related g-expectation. In Backward stochastic differential equations (Paris, 1995–1996), volume 364 of Pitman Res. Notes Math. Ser., pages 141–159. Longman, Harlow, 1997.
- [36] M. Topolewski. Reflected BSDEs with general filtration and two completely separated barriers. Probab. Math. Statist., 39(1):199–218, 2019.
- [37] J. Zhang. Backward stochastic differential equations, volume 86 of Probability Theory and Stochastic Modelling. Springer, New York, 2017. From linear to fully nonlinear theory.