

SOLVABILITY OF SOME STEFAN TYPE PROBLEMS

YOUSSEF AKDIM¹, MOHAMMED EL ANSARI² AND SOUMIA LALAOUI RHALI³

ABSTRACT. In this paper, we interest on some class of Stefan type problems. We prove the existence and uniqueness of renormalized solution in anisotropic Sobolev spaces with data belongs to L^1 – data, based on the properties of the renormalized truncations and the generalized monotonicity method in the functional spaces.

1. INTRODUCTION

Anisotropic elliptic equations have received much attention in recent years (see for example, [31], [8], [13], [17], [22] and their references). Time dependent versions of these equations have been used as mathematical models to describe the spread of an epidemic disease, see [9]. such evolution models also arise in fluid dynamics when the media has different conductivities in the different directions (see [6], [6]), and electrorheological fluids (see [31] for more details) as an important class of non-Newtonian fluids.

We are interested in the study of the behavior of solutions for a class of Stefan-type problems of form:

$$(E, f) \begin{cases} \beta(u) - \operatorname{div}(a(x, Du) + F(u)) \ni f & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

with Ω is a bounded domain in $\mathbb{R}^N (N \geq 1)$ and $\partial\Omega$ Lipschitz boundary if $N \geq 2$, a right-hand side f which is assumed to belong to $L^\infty(\Omega)$ or $L^1(\Omega)$ for (E, f) . Furthermore, $F : \mathbb{R} \rightarrow \mathbb{R}^N$ is locally lipschitz continuous and $\beta : \mathbb{R} \rightarrow 2^{\mathbb{R}}$ is a set valued, maximal monotone mapping such that $0 \in \beta(0)$ and $a : \Omega \times \mathbb{R}^N \rightarrow \mathbb{R}^N$ is a Carathéodory function satisfying the following assumptions:

(**H**₁)– Coerciveness: there exists a positive constant λ such that

$$\sum_{i=1}^N a_i(x, \xi) \cdot \xi_i \geq \lambda \sum_{i=1}^N |\xi_i|^{p_i}$$

holds for all $\xi \in \mathbb{R}^N$ and almost every $x \in \Omega$.

(**H**₂)– Growth restriction:

Date: Received: xxxxxx; Revised: yyyyyy; Accepted: zzzzzz

* Corresponding author.

2010 *Mathematics Subject Classification.* 35J60; 35A01.

Key words and phrases. Anisotropic Sobolev spaces, Truncations, maximal monotone graph, Stefan type problems.

$$|a_i(x, \xi)| \leq \gamma(d_i(x) + |\xi_i^{p_i-1}|)$$

for almost every $x \in \Omega$, γ is a positive constant for $i = 1, \dots, N$, d_i is a positive function in $L^{p'_i}(\Omega)$ and every $\xi \in \mathbb{R}^N$.

(\mathbf{H}_3)– Monotonicity in $\xi \in \mathbb{R}^N$:

$$(a(x, \xi) - a(x, \eta)) \cdot (\xi - \eta) \geq 0,$$

for almost every $x \in \Omega$ and for $\xi, \eta \in \mathbb{R}^N$.

Due to the possible jumps of β , problem (E, f) enters to class of stefan problem for wich there exists a large number of references, among them [19], [24].

Here we use the notion of renormalised solution developed by by DiPerna and Lions [21], for first order equations for L^1 -data in [30], and for Radon mesure data in [20]. It was then extended to the study of various problems of partial differential equations of parabolic, elliptic-parabolic and hyperbolic type, we refer to [14], [15].

Our problem has been studied in variable exponents spaces and Orlicz by Wittbold et al. [33], [25] and in the weighted Sobolev spaces by Akdim and Allalou [3]. Other works in this direction can be found in [4],[11],[2]. Our objective in this work is to prove an existence result of (E, f) in anisotropic Sobolev spaces, this notion were introduced by Nikolskii [29], and Troisi [32]. The main tools in our proofs are Poincaré inequality and the embedding for anisotropic Sobolev spaces. It would be interesting to refer to some Embedding theorems for anisotropic Sobolev-Orlicz spaces [27] and a fully anisotropic Sobolev Inequality established by Cianchi in [18].

The paper isorganized as follows: In Section 2, we recall the standard framework of anisotropic Sobolev spaces and some notations which will be used frequently. In Section 3, we introduce the notion of weak and also renormalized solution for the problem (E, f) for any L^1 -data. In Section 4, we give our main results on the existence and uniqueness of renormalized solutions and we discuss the existence of weak solutions. Section 5 is devoted to the case where $f \in L^\infty(\Omega)$, we prove the existence of a renormalized solution. Based on this result, the existence and uniqueness of a renormalized solution in the case where $f \in L^1(\Omega)$ is shown in Section 6. In Section 7, we will prove the existence of a weak solution. Finally, we give an example for illustrating our abstract result.

2. FUNCTION SPACES

2.1. Anisotropic Sobolev spaces. Let Ω be a bounded open subset of \mathbb{R}^N , ($N \geq 2$) and let $1 \leq p_1, \dots, p_N < \infty$ be N real numbers, $p^+ = \max(p_1, \dots, p_N)$, $p^- = \min(p_1, \dots, p_N)$ and $\vec{p} = (p_1, \dots, p_N)$. The anisotropic spaces (see [32])

$$W^{1, \vec{p}}(\Omega) = \{u \in W^{1,1}(\Omega) : \partial_{x_i} u \in L^{p_i}(\Omega), i = 1, \dots, N\}.$$

is a Banach space with respect to norm

$$\|u\|_{W^{1,\vec{p}}(\Omega)} = \|u\|_{L^1(\Omega)} + \sum_{i=1}^N \|\partial_{x_i} u\|_{L^{p_i}(\Omega)}$$

The space $W_0^{1,\vec{p}}(\Omega)$ is the closure of $C_0^\infty(\Omega)$ with respect to this norm.

The dual space of anisotropic Sobolev space $W_0^{1,\vec{p}}(\Omega)$ is equivalent to $W^{-1,\vec{p}'}(\Omega)$, where $\vec{p}' = (p'_1, \dots, p'_N)$ and $p'_i = \frac{p_i}{p_i - 1}$ for all $i = 1, \dots, N$.

We recall now a Poincaré-type inequality:

Let $u \in W_0^{1,\vec{p}}(\Omega)$, then for every $q \geq 1$ there exists a constant C_p (depending on q and i (see [23]), such that

$$\|u\|_{L^q(\Omega)} \leq C_p \|\partial_{x_i} u\|_{L^{p_i}(\Omega)} \quad \text{for } i = 1, \dots, N. \quad (2.1)$$

Moreover a Sobolev-type inequality holds. Let us denote by \bar{p} the harmonic mean of these numbers, i.e. $\frac{1}{\bar{p}} = \frac{1}{N} \sum_{i=1}^N \frac{1}{p_i}$. Let $u \in W_0^{1,\vec{p}}(\Omega)$. It follows from [32] that there exists a constant C_s such that

$$\|u\|_{L^q(\Omega)} \leq C_s \prod_{i=1}^N \|\partial_{x_i} u\|_{L^{p_i}(\Omega)}^{\frac{1}{N}}, \quad (2.2)$$

where $q = \bar{p}^* = \frac{N\bar{p}}{N-\bar{p}}$ if $\bar{p} < N$ or $q \in [1, +\infty[$ if $\bar{p} \geq N$. On the right-hand side of (2.2) it is possible to replace the geometric mean by the arithmetic mean: let a_1, \dots, a_N be positive numbers, it holds

$$\prod_{i=1}^N a_i^{\frac{1}{N}} \leq \frac{1}{N} \sum_{i=1}^N a_i,$$

which implies by 2.2 that

$$\|u\|_{L^q(\Omega)} \leq \frac{C_s}{N} \sum_{i=1}^N \|\partial_{x_i} u\|_{L^{p_i}(\Omega)}. \quad (2.3)$$

Note that when the following inequality holds

$$\bar{p} < N, \quad (2.4)$$

inequality (2.3) implies the continuous embedding of the space $W_0^{1,\vec{p}}(\Omega)$ into $L^q(\Omega)$ for every $q \in [1, \bar{p}^*]$. On the other hand, the continuity of the embedding $W_0^{1,\vec{p}}(\Omega) \hookrightarrow L^{p^+}(\Omega)$ with $p^+ := \max\{p_1, \dots, p_N\}$ relies on inequality 2.1.

It may happen that $\bar{p}^* < p^+$ if the exponents p_i are closed enough, then $p_\infty := \max\{\bar{p}^*, p^+\}$ turns out to be the critical exponent in the anisotropic Sobolev embedding (see [32]).

Proposition 2.1. *If the condition 2.4 holds, then for $q \in [1, p_\infty]$ there is a continuous embedding $W_0^{1,\vec{p}}(\Omega) \hookrightarrow L^q(\Omega)$. For $q < p_\infty$ the embedding is compact.*

$$W_0^{1,\vec{p}}(\Omega) \hookrightarrow\hookrightarrow L^q(\Omega). \quad (2.5)$$

2.2. Notations and functions. Before we discuss the concept of solution we introduce some notations and functions that will be frequently used.

We begin by introducing the truncature operator. For given constant $k > 0$ we define the cut function $T_k : \mathbb{R} \rightarrow \mathbb{R}$ as

$$T_k(r) = \begin{cases} -k, & \text{if } r \leq -k, \\ r, & \text{if } |r| < k, \\ k, & \text{if } r \geq k, \end{cases}$$




FIGURE 1. Truncation function

and for $r \in \mathbb{R}$, let us define the functions $r \rightarrow r^+ := \max(r, 0)$ and $r \rightarrow \text{sign}_0(r)$ the usual sign function which is defined by

$$r \rightarrow \text{sign}_0(r) := \begin{cases} -1, & \text{on }]-\infty, 0[, \\ 1, & \text{on }]0, \infty[, \\ 0, & \text{if } r = 0. \end{cases}$$

and

$$r \rightarrow \text{sign}_0^+(r) := \begin{cases} 1, & \text{if } r > 0, \\ 0, & \text{if } r \leq 0. \end{cases}$$

Let $h_l : \mathbb{R} \rightarrow \mathbb{R}$ be defined by $h_l(r) := \min((l+1-|r|)^+, 1)$ for each $r \in \mathbb{R}$.




FIGURE 2. Function $h_l(r)$

For $\sigma > 0$, we define $H_\sigma^+ : \mathbb{R} \rightarrow \mathbb{R}$ by

$$H_\sigma^+(r) := \begin{cases} 0, & \text{if } r < 0, \\ \frac{1}{\sigma}r, & \text{if } 0 \leq r \leq \sigma, \\ 1, & \text{if } r > \sigma. \end{cases}$$

and $H_\sigma : \mathbb{R} \rightarrow \mathbb{R}$ by

$$H_\sigma(r) := \begin{cases} -1, & \text{if } r < -\sigma, \\ \frac{1}{\sigma}r, & \text{if } -\sigma \leq r \leq \sigma, \\ 1, & \text{if } r > \sigma. \end{cases}$$

3. NOTION OF SOLUTIONS

3.1. Weak solutions.

Definition 3.1. A weak solution of (E, f) is a pair of functions $(u, b) \in W_0^{1, \vec{p}}(\Omega) \times L^1(\Omega)$ satisfying $F(u) \in (L_{loc}^1(\Omega))^N$, $b \in \beta(u)$ almost everywhere in Ω and

$$b - \operatorname{div}(a(x, Du) + F(u)) = f \text{ in } D'(\Omega). \quad (3.1)$$

3.2. Renormalized solutions.

Definition 3.2. A renormalized solution of (E, f) is a pair of functions (u, b) satisfying the following conditions:

(**R**₁) $u : \Omega \rightarrow \mathbb{R}$ is measurable, $b \in L^1(\Omega)$, $u(x) \in D(\beta(x))$ and $b(x) \in \beta(u(x))$ for a.e. $x \in \Omega$.

(**R**₂) For each $k > 0$, $T_k(u) \in W_0^{1, \vec{p}}(\Omega)$ and

$$\int_{\Omega} b \cdot h(u) \phi + \int_{\Omega} (a(x, D(u)) + F(u)) \cdot D(h(u) \phi) = \int_{\Omega} f h(u) \phi, \quad (3.2)$$

holds for all $h \in C_c^1(\mathbb{R})$ and all $\phi \in W_0^{1, \vec{p}}(\Omega) \cap L^\infty(\Omega)$.

(**R**₃) $\int_{\{k < |u| < k+1\}} a(x, Du) \cdot Du \rightarrow 0$ as $k \rightarrow \infty$.

4. MAIN RESULTS

In this section, we will first state the existence and uniqueness of renormalized solutions for (E, f) . Then, we will prove that the renormalized solution of (E, f) is a weak solution.

Theorem 4.1. *For $f \in L^1(\Omega)$, there exists at least one renormalized solution (u, b) of (E, f) .*

Theorem 4.2. *Let $\beta : \Omega \times \mathbb{R} \rightarrow 2^{\mathbb{R}}$ be such that $\beta(x, \cdot)$ is strictly monotone for almost every $x \in \Omega$. For $f \in L^1(\Omega)$, let $(u, b), (\tilde{u}, \tilde{b})$ be renormalized solutions of (E, f) . Then $u = \tilde{u}$ and $b = \tilde{b}$.*

Proposition 4.3. *Let (u, b) be a renormalized solution of (E, f) for $f \in L^\infty(\Omega)$. Then $u \in W^{1, \vec{p}}(\Omega) \cup L^\infty(\Omega)$ and thus, in particular, u is a weak solution of (E, f) .*

To prove Theorem 4.1, we will introduce and solve approximation problems. To this end, for $f \in L^1(\Omega)$ and $m, n \in \mathbb{N}$ we define $f_{m,n} : \Omega \rightarrow \mathbb{R}$ by

$$f_{m,n}(x) = \max(\min(f(x), m), -n)$$

for almost every $x \in \Omega$. Clearly, $f_{m,n} \in L^\infty(\Omega)$ for each $m, n \in \mathbb{N}$, $|f_{m,n}(x)| \leq |f(x)|$ a.e. in Ω , hence $\lim_{n \rightarrow \infty} \lim_{m \rightarrow \infty} f_{m,n} = f$ in $L^1(\Omega)$ and almost everywhere in Ω .

The next theorem will give us existence of renormalized solutions $(u_{m,n}, b_{m,n})$ of $(E, f_{m,n})$ for each $m, n \in \mathbb{N}$.

5. CASE WHERE $f \in L^\infty(\Omega)$

Theorem 5.1. *For $f \in L^\infty(\Omega)$, there exists at least one renormalized solution (u, b) of (E, f) .*

The following section will be devoted to prove of Theorem 5.1, and we will divide it into several steps.

5.1. Approximate solution for L^∞ - data. First we will introduce the approximate problem to (E, f) for $f \in L^\infty(\Omega)$ and for which the existence can be proved by standard variational arguments. For $0 < \varepsilon \leq 1$, let $\beta_\varepsilon : \mathbb{R} \rightarrow \mathbb{R}$ be the Yosida approximation of β (see [16]). We introduce the operators

$$\begin{aligned} A_{1,\varepsilon} &: W_0^{1, \vec{p}}(\Omega) \rightarrow W^{-1, \vec{p}'}(\Omega), \\ u &\rightarrow \beta_\varepsilon(T_{1/\varepsilon}(u)) + \varepsilon \arctan(u) - \operatorname{div}(a(x), Du) \end{aligned}$$

and

$$\begin{aligned} A_{2,\varepsilon} &: W_0^{1, \vec{p}}(\Omega) \rightarrow W^{-1, \vec{p}'}(\Omega), \\ u &\rightarrow -\operatorname{div}F(T_{1/\varepsilon}(u)). \end{aligned}$$

Because of $(\mathbf{H}_2) - (\mathbf{H}_3)$, $A_{1/\varepsilon}$ is well-defined and monotone (see [28] for instance). Since $\beta_\varepsilon \circ T_{1/\varepsilon}$ is bounded and continuous and thanks to the growth condition (H_2) on a , it follows that $A_{1,\varepsilon}$ is hemicontinuous (see [28]). From the continuity and boundedness of $F \circ T_{1/\varepsilon}$, it follows that $A_{2,\varepsilon}$ is strongly continuous. Therefore the operator $A_\varepsilon := A_{1,\varepsilon} + A_{2,\varepsilon}$ is pseudomonotone. Using the monotonicity of β_ε , the Gauss-Green Theorem for Sobolev functions and the boundary condition on the convection term $\int_\Omega F(T_{1/\varepsilon}(u)) \cdot Du$, we show by using similar arguments as in [12] that A_ε is coercive and bounded. Then it follows from [28] Theorem 2.7 that A_ε is surjective, i.e., for each $0 < \varepsilon \leq 1$ and $f \in W^{-1, \vec{p}'}(\Omega)$ there exists a solution $u_\varepsilon \in W_0^{1, \vec{p}}(\Omega)$ of the problem

$$(E_\varepsilon, f) \begin{cases} \beta_\varepsilon(T_{1/\varepsilon}(u_\varepsilon)) + \varepsilon \arctan(u_\varepsilon) - \operatorname{div}(a(x), Du_\varepsilon) + F((T_{1/\varepsilon}(u_\varepsilon))) = f & \text{in } \Omega, \\ u_\varepsilon = 0 & \text{on } \partial\Omega, \end{cases}$$

such that the following inequality holds for all $\phi \in W_0^{1, \vec{p}}(\Omega)$

$$\int_{\Omega} (\beta_{\varepsilon}(T_{1/\varepsilon}(u_{\varepsilon})) + \varepsilon \arctan(u))\phi + \int_{\Omega} (a(x, D(u_{\varepsilon})) + F(T_{1/\varepsilon}(u_{\varepsilon}))).D\phi = \langle f, \phi \rangle \quad (5.1)$$

where $\langle \cdot, \cdot \rangle$ denotes the duality pairing between $W_0^{1, \vec{p}}(\Omega)$ and $W^{-1, \vec{p}'}(\Omega)$.

Proposition 5.2. *For $0 < \varepsilon \leq 1$ fixed and $f, \tilde{f} \in L^{\infty}(\Omega)$, let $u_{\varepsilon}, \tilde{u}_{\varepsilon} \in W_0^{1, \vec{p}}(\Omega)$ be solutions of (E_{ε}, f) and $(E_{\varepsilon}, \tilde{f})$, respectively, then, the following comparison principle holds:*

$$\varepsilon \int_{\Omega} (\arctan(u_{\varepsilon}) - \arctan(\tilde{u}_{\varepsilon}))^{+} \leq \int_{\Omega} (f - \tilde{f}) \text{sign}_0^{+}(u_{\varepsilon} - \tilde{u}_{\varepsilon}). \quad (5.2)$$

Proof. We use the test function $\varphi = H_{\sigma}^{+}(u_{\varepsilon} - \tilde{u}_{\varepsilon})$ in the weak formulation (5.1) for u_{ε} and \tilde{u}_{ε} . Subtracting the resulting inequalities, we obtain

$$I_{l, \delta}^1 + I_{l, \delta}^2 + I_{l, \delta}^3 + I_{l, \delta}^4 = I_{l, \delta}^5,$$

where

$$\begin{aligned} I_{l, \delta}^1 &= \int_{\Omega} (\beta_{\varepsilon} T_{\frac{1}{\varepsilon}}(u_{\varepsilon}) - \beta_{\varepsilon} T_{\frac{1}{\varepsilon}}(\tilde{u}_{\varepsilon})) H_{\delta}^{+}(u_{\varepsilon} - \tilde{u}_{\varepsilon}) \\ I_{l, \delta}^2 &= \int_{\Omega} (\varepsilon \arctan(u_{\varepsilon}) - \varepsilon \arctan(\tilde{u}_{\varepsilon})) H_{\delta}^{+}(u_{\varepsilon} - \tilde{u}_{\varepsilon}), \\ I_{l, \delta}^3 &= \int_{\Omega} a(x, Du_{\varepsilon}) - a(x, D\tilde{u}_{\varepsilon}).DH_{\delta}^{+}(u_{\varepsilon} - \tilde{u}_{\varepsilon}), \\ I_{l, \delta}^4 &= \int_{\Omega} (F(T_{\frac{1}{\varepsilon}}(u_{\varepsilon})) - F(T_{\frac{1}{\varepsilon}}(\tilde{u}_{\varepsilon}))).DH_{\delta}^{+}(u_{\varepsilon} - \tilde{u}_{\varepsilon}) \\ I_{l, \delta}^5 &= \int_{\Omega} (f - \tilde{f}) H_{\delta}^{+}(u_{\varepsilon} - \tilde{u}_{\varepsilon}). \end{aligned}$$

Passing to the limit with $\sigma \rightarrow 0$, (5.2) follows. \square

Remark 5.3. Let $f, \tilde{f} \in L^{\infty}(\Omega)$ be such that $f \leq \tilde{f}$ almost everywhere in Ω , $\varepsilon > 0$ and $u_{\varepsilon}, \tilde{u}_{\varepsilon} \in W_0^{1, \vec{p}}(\Omega)$ be solutions of (E_{ε}, f) and $(E_{\varepsilon}, \tilde{f})$, respectively, then it is an immediate consequence of Proposition 5.2 is that $u_{\varepsilon} \leq \tilde{u}_{\varepsilon}$ almost everywhere in Ω . Furthermore, from the monotonicity of $\beta_{\varepsilon} \circ T_{1/\varepsilon}$ it follows that also

$$\beta_{\varepsilon}(T_{1/\varepsilon}(u_{\varepsilon})) \leq \beta_{\varepsilon}(T_{1/\varepsilon}(\tilde{u}_{\varepsilon}))$$

a.e. in Ω .

5.2. A priori estimates.

Lemma 5.4. *For $0 < \varepsilon \leq 1$ and $f \in L^\infty(\Omega)$ let $u_\varepsilon \in W_0^{1, \vec{p}}(\Omega)$ be a solution of (E_ε, f) . Then*

i) There exists a constant $C_1 = C_1(\|f\|_\infty, \lambda, p_i, N) > 0$, not depending on ε , such that

$$\|u_\varepsilon\| \leq C_1. \quad (5.3)$$

ii) for all $0 < \varepsilon \leq 1$, we have

$$\|\beta_\varepsilon(T_{1/\varepsilon}(u_\varepsilon))\|_\infty \leq \|f\|_\infty \quad (5.4)$$

iii) for all $0 < \varepsilon \leq 1$ and all $l, k > 0$, we have

$$\int_{\{l \leq |u| \leq k+l\}} a(x, Du_\varepsilon) \cdot Du_\varepsilon \leq k \int_{\{|u_\varepsilon| > l\}} |f|. \quad (5.5)$$

Proof. *i)* Taking u_ε as a test function in (5.1) we obtain

$$\int_\Omega (\beta_\varepsilon(T_{1/\varepsilon}(u_\varepsilon)) + \varepsilon \arctan(u_\varepsilon)) u_\varepsilon dx + \int_\Omega a(x, Du_\varepsilon) \cdot Du_\varepsilon dx + \int_\Omega F(T_{1/\varepsilon}(u_\varepsilon)) \cdot Du_\varepsilon dx = \int_\Omega f u_\varepsilon dx$$

As the first term on the left-hand side is nonnegative and the integral over the convection term vanishes by (H_1) , we have

$$\lambda \sum_{i=1}^N \int_\Omega \left| \frac{\partial u_\varepsilon}{\partial x_i} \right|^{p_i} dx \leq \sum_{i=1}^N \int_\Omega a_i(x, Du_\varepsilon) \cdot \frac{\partial u_\varepsilon}{\partial x_i} dx \leq \int_\Omega f u_\varepsilon dx \leq C \|f\|_\infty \left(\sum_{i=1}^N \int_\Omega \left| \frac{\partial u_\varepsilon}{\partial x_i} \right|^{p_i} dx \right)^{1/p_i}.$$

due to the Hölder inequality. Thus $\|u_\varepsilon\|^{p_i} \leq C_2 \|u_\varepsilon\|$, where C_2 is a positive constant. Then we can deduce that u_ε remains bounded in $W_0^{1, \vec{p}}(\Omega)$ i.e.,

$$\|u_\varepsilon\| \leq C_1.$$

ii) Taking $\frac{1}{\sigma} [T_{k+\sigma}(\beta_\varepsilon(T_{1/\varepsilon}(u_\varepsilon))) - T_k(\beta_\varepsilon(T_{1/\varepsilon}(u_\varepsilon)))]$ as a test function in (5.1), passing to the limit with $\sigma \rightarrow 0$ and choosing $k > \|f\|_\infty$, we obtain *ii)*.

iii) For $k, l > 0$ fixed we take $T_k(u_\varepsilon - T_l(u_\varepsilon))$ as a test function in (5.1).

Using $\int_\Omega a(x, Du_\varepsilon) \cdot DT_k(u_\varepsilon - T_l(u_\varepsilon)) dx = \int_{\{l < |u_\varepsilon| < l+k\}} a(x, Du_\varepsilon) \cdot Du_\varepsilon dx$, and as the first term on the left-hand side is nonnegative and the convection term vanishes, we get

$$\int_{\{l < |u_\varepsilon| < k+l\}} a(x, Du_\varepsilon) \cdot Du_\varepsilon \leq \int_\Omega f T_k(u_\varepsilon - T_l(u_\varepsilon)) dx \leq \int_{\{|u_\varepsilon| > l\}} |f| dx. \quad (5.6)$$

□

Remark 5.5. For $k > 0$, from *iii)* in Lemma 5.4, we deduce that

$$|\{|u_\varepsilon| \geq l\}| \leq \frac{C_2}{l^{1-\frac{1}{\vec{p}}}} \quad (5.7)$$

$$\int_{\{l \leq |u_\varepsilon| \leq k+l\}} a(x, Du_\varepsilon) \cdot Du_\varepsilon \leq k \|f\|_\infty |\{|u_\varepsilon| > l\}| \leq \frac{C_2(k)}{l^{\frac{1}{\vec{p}}-1}} \quad (5.8)$$

for any $0 < \varepsilon \leq 1$ and a constant $C_2(k) > 0$ not depending on ε .

Indeed, let $l > 0$ large enough we have:

$$l|\{|u_\varepsilon| \geq l\}| = \int_{\{|u_\varepsilon| \geq l\}} |T_l(u_\varepsilon)| dx \leq C \left(\sum_{i=1}^N \int_{\Omega} \left| \frac{\partial T_l(u_\varepsilon)}{\partial x_i} \right|^{p_i} \right)^{1/p_i} \leq C_2 l^{1/\bar{p}}$$

with implies $|\{|u_\varepsilon| \geq l\}| \leq C_2 l^{1/\bar{p}-1}$. Then

$$\lim_{l \rightarrow +\infty} |\{|u_\varepsilon| \geq l\}| = 0.$$

Therefore, (5.7) follows from (5.8).

5.3. Basic convergence results.

Lemma 5.6. *For $0 < \varepsilon \leq 1$ and $f \in L^\infty(\Omega)$, let $u_\varepsilon \in W_0^{1, \vec{p}}(\Omega)$ be a solution of (E_ε, f) . There exist $u \in W_0^{1, \vec{p}}(\Omega)$, $b \in L^\infty(\Omega)$ such that for a not relabeled subsequence of $(u_\varepsilon)_{0 < \varepsilon \leq 1}$ as $\varepsilon \mapsto 0$:*

$$u_\varepsilon \rightharpoonup u \quad \text{in } W_0^{1, \vec{p}}(\Omega) \text{ and a.e. in } \Omega, \quad (5.9)$$

$$T_k(u_\varepsilon) \rightharpoonup T_k(u) \quad \text{in } W_0^{1, \vec{p}}(\Omega) \text{ and strongly in } L^q(\Omega), \quad (5.10)$$

$$\beta_\varepsilon(T_{1/\varepsilon}(u_\varepsilon)) \rightharpoonup b \quad \text{in } L^\infty(\Omega). \quad (5.11)$$

Moreover, for any $k > 0$,

$$DT_k(u_\varepsilon) \rightharpoonup DT_k(u) \quad \text{in } \prod_{i=1}^N L^{p_i}(\Omega), \quad (5.12)$$

$$a(x, DT_k(u_\varepsilon)) \rightharpoonup a(x, DT_k(u)) \quad \text{in } \prod_{i=1}^N L^{p'_i}(\Omega). \quad (5.13)$$

Proof. By combining Lemma 5.4 and Remark 5.5, we obtain (5.11). From (5.7), (5.3) and (2.5), we deduce with a classical argument (see [1]) that for a subsequence still indexed by ε , (5.9) – (5.10) and (5.12) hold as ε tends to 0, where u is a measurable function defined on Ω .

It is left to prove (5.13). For this, by (H_2) and (5.3) it follows that given any subsequence of $(a(x, DT_k(u_\varepsilon))_\varepsilon)$, there exists a subsequence, still denoted by

$(a(x, DT_k(u_\varepsilon))_\varepsilon)$, such that $a(x, DT_k(u_\varepsilon)) \rightharpoonup \Phi_k$ in $\prod_{i=1}^N L^{p'_i}(\Omega)$. We will prove that

$\Phi_k = a(x, DT_k(u))$ a.e. on Ω . The proof consists of three steps.

Step 1: For every $h \in W^{1, \infty}(\mathbb{R})$, $h \leq 0$ and $\text{supp}(h)$ compact, we will prove that

$$\limsup_{\varepsilon \rightarrow 0} \int_{\Omega} a(x, DT_k(u_\varepsilon)) \cdot D[h(u_\varepsilon)(T_k(u_\varepsilon) - T_k(u))] dx \leq 0. \quad (5.14)$$

Taking $h(u_\varepsilon)(T_k(u_\varepsilon) - T_k(u))$ as a test function in (5.1), we have

$$\begin{aligned} \int_{\Omega} (\beta_\varepsilon(T_{1/\varepsilon}(u_\varepsilon)) + \varepsilon \arctan(u_\varepsilon)) h(u_\varepsilon) (T_k(u_\varepsilon) - T_k(u)) + \int_{\Omega} a(x, D(u_\varepsilon)) \cdot D[h_\varepsilon(T_k(u_\varepsilon) - T_k(u))] \\ + \int_{\Omega} F(T_{1/\varepsilon}(u_\varepsilon)) \cdot D[h_\varepsilon(T_k(u_\varepsilon) - T_k(u))] = \int_{\Omega} f h(u_\varepsilon) (T_k(u_\varepsilon) - T_k(u)). \end{aligned} \quad (5.15)$$

Using $|h_\varepsilon(T_k(u_\varepsilon) - T_k(u))| \leq 2k\|h\|_\infty$, by Lebesgue's dominated convergence theorem we find that $\lim_{\varepsilon \rightarrow 0} \int_\Omega fh(u_\varepsilon)(T_k(u_\varepsilon) - T_k(u)) = 0$ and then $\lim_{\varepsilon \rightarrow 0} \int_\Omega F(T_{1/\varepsilon}(u_\varepsilon)) \cdot D[h_\varepsilon(u_\varepsilon)(T_k(u_\varepsilon) - T_k(u))] = 0$. By using the same arguments as in [4], we can prove that

$$\limsup_{\varepsilon \rightarrow 0} \int_\Omega \beta_\varepsilon(T_{1/\varepsilon}(u_\varepsilon)) \cdot [h(u_\varepsilon)(T_k(u_\varepsilon) - T_k(u))] dx \geq 0.$$

Passing to the limit in (5.15) and using the above results, we obtain (5.14).

Step 2: We now prove that for every $k > 0$,

$$\limsup_{\varepsilon \rightarrow 0} \int_\Omega a(x, DT_k(u_\varepsilon)) \cdot [D(T_k(u_\varepsilon) - DT_k(u))] dx \leq 0. \quad (5.16)$$

Indeed, for $k > l$, take $h_l(u_\varepsilon)(T_k(u_\varepsilon) - T_k(u))$ as a test function in (5.1). Letting $\varepsilon \rightarrow 0$ and then $l \rightarrow \infty$, we obtain

$$\limsup_{\varepsilon \rightarrow 0} \int_\Omega a(x, DT_k(u_\varepsilon)) \cdot D[h_l(u_\varepsilon)(T_k(u_\varepsilon) - T_k(u))] dx = E_1 + E_2 + E_3.$$

where

$$E_1 = \int_{\{|u_\varepsilon| \leq k\}} h_l(u_\varepsilon) a(x, DT_k(u_\varepsilon)) \cdot [DT_k(u_\varepsilon) - DT_k(u)] dx,$$

$$E_2 = \int_{\{|u_\varepsilon| > k\}} h_l(u_\varepsilon) a(x, DT_k(u_\varepsilon)) \cdot (-DT_k(u)) dx,$$

$$E_3 = \int_\Omega h'_l(u_\varepsilon)(T_k(u_\varepsilon) - T_k(u)) a(x, DT_k(u_\varepsilon)) \cdot Du_\varepsilon dx.$$

Since $l > k$, on the set $\{|u_\varepsilon| \leq k\}$ we have $h_l(u_\varepsilon) = 1$ so that we can write

$$\limsup_{\varepsilon \rightarrow 0} E_1 = \limsup_{\varepsilon \rightarrow 0} \int_\Omega a(x, DT_k(u_\varepsilon)) \cdot (DT_k(u_\varepsilon) - DT_k(u)) dx.$$

For E_2 , using Lebesgue's dominated convergence theorem, we get

$$\lim_{\varepsilon \rightarrow 0} E_2 = \int_{\{|u_\varepsilon| > k\}} h_l(u) \Phi_{l+1} \cdot DT_k(u) dx = 0.$$

For E_3 , we have

$$\begin{aligned} & - \int_\Omega h'_l(u_\varepsilon)(T_k(u_\varepsilon) - T_k(u)) a(x, DT_k(u_\varepsilon)) Du_\varepsilon dx \\ & \leq 2k \int_{\{l < |u_\varepsilon| \leq l+1\}} a(x, Du_\varepsilon) Du_\varepsilon dx. \end{aligned}$$

Using (5.8), we deduce that

$$\limsup_{l \rightarrow \infty} \limsup_{\varepsilon \rightarrow 0} \left(- \int_\Omega h'_l(u_\varepsilon)(T_k(u_\varepsilon) - T_k(u)) a(x, DT_k(u_\varepsilon)) \cdot Du_\varepsilon dx \right) \leq 0.$$

Applying (5.14) with h replaced by h_l , $l > k$, we get

$$\begin{aligned} & \limsup_{\varepsilon \rightarrow 0} \int_\Omega a(x, DT_k(u_\varepsilon)) \cdot [DT_k(u_\varepsilon) - DT_k(u)] dx \\ & \leq \limsup_{\varepsilon \rightarrow 0} \left(- \int_\Omega h'_l(u_\varepsilon)(T_k(u_\varepsilon) - T_k(u)) a(x, DT_k(u_\varepsilon)) \cdot Du_\varepsilon dx \right). \end{aligned}$$

Now letting $l \rightarrow \infty$, (5.16) yields.

Step 3: In this step, we prove by monotonicity arguments that for $k > 0$, $\overline{\Phi_k} = a(x, DT_k(u))$ for almost every $x \in \Omega$. Let $\phi \in D(\Omega)$ and $\alpha \in \mathbb{R}$. Using (5.16), we have

$$\alpha \lim_{\varepsilon \rightarrow 0} \int_{\Omega} a(x, DT_k(u_\varepsilon)) \cdot D\phi dx \geq \alpha \int_{\Omega} a(x, D(T_k(u) - \alpha\phi)) \cdot D\phi dx.$$

Dividing by $\alpha > 0$ and $\alpha < 0$ and letting $\alpha \rightarrow 0$, we obtain

$$\lim_{\varepsilon \rightarrow 0} \int_{\Omega} a(x, DT_k(u_\varepsilon)) \cdot D\phi dx = \int_{\Omega} a(x, DT_k(u)) \cdot D\phi dx. \quad (5.17)$$

This means that for all $k > 0$, $\int_{\Omega} \Phi_k \cdot D\phi dx = \int_{\Omega} a(x, DT_k(u)) \cdot D\phi dx$, and then $\Phi_k = a(x, DT_k(u))$ in $D'(\Omega)$ for all $k > 0$. Hence $\Phi_k = a(x, DT_k(u))$ a.e. in Ω and then $a(x, DT_k(u_\varepsilon)) \rightharpoonup a(x, DT_k(u))$ weakly in $\prod_{i=1}^N L^{p_i}(\Omega)$. \square

Remark 5.7. As an immediate consequence of (5.16) and (H_3) we obtain

$$\lim_{\varepsilon \rightarrow 0} \int_{\Omega} a(x, DT_k(u_\varepsilon)) - a(x, DT_k(u)) \cdot (DT_k(u_\varepsilon) - T_k(u)) = 0. \quad (5.18)$$

Let us see finally that

$$\lim_{l \rightarrow \infty} \int_{l < |u| < l+1} a(x, DT(u)) dx = 0. \quad (5.19)$$

Indeed, for any $l \geq 0$ fixed we have

$$\begin{aligned} \int_{l < |u| < l+1} a(x, D(u_\varepsilon)) \cdot D(u_\varepsilon) dx &= \int_{\Omega} a(x, DT_{l+1}(u_\varepsilon)) \cdot (DT_{l+1}(u_\varepsilon) - DT_l(u_\varepsilon)) dx \\ &= \int_{\Omega} a(x, DT_{l+1}(u_\varepsilon)) \cdot DT_{l+1}(u_\varepsilon) dx - \int_{\Omega} a(x, DT_l(u_\varepsilon)) \cdot DT_l(u_\varepsilon) dx. \end{aligned}$$

By (5.18) and passing to the limit as $\varepsilon \rightarrow 0$ for fixed $l \geq 0$ we obtain

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \int_{\{l < |u_\varepsilon| < l+1\}} a(x, D(u_\varepsilon)) \cdot D(u_\varepsilon) dx &= \int_{\Omega} a(x, DT_{l+1}(u)) \cdot DT_{l+1}(u) dx \\ &\quad - \int_{\Omega} a(x, DT_l(u)) \cdot DT_l(u) dx = \int_{\{l < |u| < l+1\}} a(x, Du) \cdot D(u) dx. \end{aligned} \quad (5.20)$$

Therefore, taking $l \rightarrow +\infty$ in (5.20) and using the estimate (5.8) show that satisfies (R_3) .

5.4. Proof of the existence result. We are now in position to conclude the proof of our main result presented in Theorem 5.1:

Proof. Let $h \in C_c^1(\mathbb{R})$ and $\varphi \in W_0^{1, \vec{p}}(\Omega) \cap L^\infty(\Omega)$. Taking $h_l(u_\varepsilon)h(u)\varphi$ as a test function in (5.1), we obtain

$$I_{\varepsilon, l}^1 + I_{\varepsilon, l}^2 + I_{\varepsilon, l}^3 + I_{\varepsilon, l}^4 = I_{\varepsilon, l}^5 \quad (5.21)$$

where

$$I_{\varepsilon, l}^1 = \int_{\Omega} \beta_\varepsilon(T_{1/\varepsilon}(u_\varepsilon)) h_l(u_\varepsilon) h(u) \varphi,$$

$$\begin{aligned}
I_{\varepsilon,l}^2 &= \varepsilon \int_{\Omega} \arctan(u_{\varepsilon}) h_l(u_{\varepsilon}) h(u) \varphi, \\
I_{\varepsilon,l}^3 &= \int_{\Omega} a(x, Du_{\varepsilon}) \cdot D(h_l(u_{\varepsilon}) h(u) \varphi), \\
I_{\varepsilon,l}^4 &= \int_{\Omega} F(T_{1/\varepsilon}(u_{\varepsilon})) \cdot D(h_l(u_{\varepsilon}) h(u) \varphi), \\
I_{\varepsilon,l}^5 &= \int_{\Omega} f h_l(u_{\varepsilon}) h(u) \varphi.
\end{aligned}$$

Step 1: Letting $\varepsilon \rightarrow 0$ obviously, we have

$$\lim_{\varepsilon \rightarrow 0} I_{\varepsilon,l}^2 = 0. \quad (5.22)$$

Using the convergence results (5.9), ((5.11) from Lemma 5.6 we can immediately calculate the following limits:

$$\lim_{\varepsilon \rightarrow 0} I_{\varepsilon,l}^1 = \int_{\Omega} b h_l(u) h(u) \varphi, \quad (5.23)$$

$$\lim_{\varepsilon \rightarrow 0} I_{\varepsilon,l}^5 = \int_{\Omega} f h_l(u) h(u) \varphi. \quad (5.24)$$

We write $I_{\varepsilon,l}^3 = I_{\varepsilon,l}^{3,1} + I_{\varepsilon,l}^{3,2}$ where

$$I_{\varepsilon,l}^{3,1} = \int_{\Omega} h'_l(u_{\varepsilon}) a(x, Du_{\varepsilon}) \cdot Du_{\varepsilon} h(u) \varphi, \quad I_{\varepsilon,l}^{3,2} = \int_{\Omega} h_l(u_{\varepsilon}) a(x, Du_{\varepsilon}) D(h(u) \varphi)..$$

Using (5.8), we get the estimate

$$|\lim_{\varepsilon \rightarrow 0} I_{\varepsilon,l}^{3,1}| \leq \|h\|_{\infty} \|\varphi\|_{\infty} \cdot C_2 l^{-(1-1/\bar{p})}. \quad (5.25)$$

By Lebesgue's dominated convergence theorem it follows that for any $i \in \{1, \dots, N\}$, we have

$$h_l(u_{\varepsilon}) \frac{\partial}{\partial x_i} (h(u) \varphi) \rightarrow h_l(u) \frac{\partial}{\partial x_i} (h(u) \varphi) \quad \text{in } L^{p_i} \text{ as } \varepsilon \rightarrow 0.$$

Keeping in mind that $I_{\varepsilon,l}^{3,2} = \int_{\Omega} h_l(u_{\varepsilon}) a(x, DT_{l+1}(u_{\varepsilon})) \cdot D(h(u) \varphi)$ and by using (5.13), we get

$$\lim_{\varepsilon \rightarrow 0} I_{\varepsilon,l}^{3,2} = \int_{\Omega} h_l(u) a(x, DT_{l+1}(u)) \cdot D(h(u) \varphi). \quad (5.26)$$

Let us write $I_{\varepsilon,l}^4 = I_{\varepsilon,l}^{4,1} + I_{\varepsilon,l}^{4,2}$, where

$$I_{\varepsilon,l}^{4,1} = \int_{\Omega} h'_l(u_{\varepsilon}) F(T_{1/\varepsilon}(u_{\varepsilon})) \cdot Du_{\varepsilon} h(u) \varphi,$$

$$I_{\varepsilon,l}^{4,2} = \int_{\Omega} h_l(u_{\varepsilon}) F(T_{1/\varepsilon}(u_{\varepsilon})) \cdot D(h(u) \varphi).$$

For any $l \in \mathbb{N}$, there exists $\varepsilon_0(l)$ such that for all $\varepsilon < \varepsilon_0(l)$,

$$I_{\varepsilon,l}^{4,1} = \int_{\Omega} h'_l(T_{l+1}(u_\varepsilon))F(T_{l+1}(u_\varepsilon)).h(u)\varphi. \quad (5.27)$$

Using the Gauss-Green Theorem for Sobolev functions in (5.27), we get for all $\varepsilon < \varepsilon_0(l)$,

$$I_{\varepsilon,l}^{4,1} = - \int_{\Omega} \int_0^{T_{l+1}(u_\varepsilon)} h'_l(r)F(r)dr.D(h(u)\varphi). \quad (5.28)$$

Now, using (5.9) and the Gauss-Green Theorem, after letting $\varepsilon \rightarrow 0$, we get

$$\lim_{\varepsilon \rightarrow 0} I_{\varepsilon,l}^{4,1} = \int_{\Omega} h'_l(u)F(u).Duh(u)\varphi. \quad (5.29)$$

Choosing ε small enough, we can write

$$I_{\varepsilon,l}^{4,2} = \int_{\Omega} h_l(u_\varepsilon)F(T_{l+1}(u_\varepsilon)).D(h(u)\varphi), \quad (5.30)$$

and conclude that

$$\lim_{\varepsilon \rightarrow 0} I_{\varepsilon,l}^{4,2} = \int_{\Omega} h_l(u)F(u).D(h(u)\varphi). \quad (5.31)$$

Step 2: Passage to the limit with $l \rightarrow \infty$.

Combining (5.21) and (5.22) – (5.31) we deduce that

$$I_l^1 + I_l^2 + I_l^3 + I_l^4 + I_l^5 = I_l^6 \quad (5.32)$$

where

$$I_l^1 = \int_{\Omega} bh_l(u)h(u)\varphi, \quad I_l^2 = \int_{\Omega} h_l(u)a(x, DT_{l+1}(u)).D(h(u)\varphi),$$

$$|I_l^3| \leq C_2|l^{-(1-1/\bar{p})}||h||_{\infty}||\varphi||_{\infty}, \quad I_l^4 = \int_{\Omega} h_l(u)F(u).D(h(u)\varphi),$$

$$I_l^5 = \int_{\Omega} h'_l(u)F(u).Duh(u)\varphi, \quad I_l^6 = \int_{\Omega} fh_l(u)h(u)\varphi.$$

Obviously, we have

$$\lim_{\varepsilon \rightarrow \infty} I_l^3 = 0. \quad (5.33)$$

Choosing $m > 0$ such that $\text{supp } h \subset [-m, m]$, we can replace u by $T_m(u)$ in $I_l^1, I_l^2, \dots, I_l^6$, and

$$h'_l(u) = h'_l(T_m(u)) = 0 \text{ if } l+1 > m, \quad h_l(u) = h_l(T_m(u)) = 0 \text{ if } l > m.$$

Therefore, letting $l \rightarrow \infty$ and combining (5.32) with (5.33) we obtain

$$\int_{\Omega} bh(u)\varphi + \int_{\Omega} (a(x, Du) + F(u)).D(h(u)\varphi) = \int_{\Omega} fh(u)\varphi \quad (5.34)$$

for all $h \in C_c^1(\mathbb{R})$ and all $\varphi \in W_0^{1, \vec{p}}(\Omega) \cap L^\infty(\Omega)$.

Step 3: Subdifferential argument

It is left to prove that $u(x) \in D(\beta(x))$ and $b(x) \in \beta(u(x))$ for almost all $x \in \Omega$. Since β is a maximal monotone graph, there exist a convex, l.s.c and proper function $j : \mathbb{R} \rightarrow [0, \infty]$, such that

$$\beta(r) = \partial j(r) \quad \text{for all } r \in \mathbb{R}.$$

According to [16], for $0 < \varepsilon \leq 1$, $j_\varepsilon : \mathbb{R} \rightarrow \mathbb{R}$ defined by $j_\varepsilon(r) = \int_0^r \beta_\varepsilon(s) ds$ has the following properties as in [33]

i) For any $0 < \varepsilon \leq 1$, j_ε is convex and differentiable for all $r \in \mathbb{R}$, such that

$$j'_\varepsilon(r) = \beta_\varepsilon(r) \quad \text{for all } r \in \mathbb{R} \text{ and any } 0 < \varepsilon \leq 1.$$

ii) $j_\varepsilon(r) \rightarrow j(r)$ for all $r \in \mathbb{R}$ as $\varepsilon \rightarrow 0$.

From *i)*, it follows that for any $0 < \varepsilon \leq 1$

$$j_\varepsilon(r) \geq j_\varepsilon(T_{1/\varepsilon}(u_\varepsilon)) + (r - T_{1/\varepsilon}(u_\varepsilon))\beta_\varepsilon(T_{1/\varepsilon}(u_\varepsilon)) \quad (5.35)$$

holds for all $r \in \mathbb{R}$ and almost everywhere in Ω .

Let $E \cup \Omega$ be an arbitrary measurable set and χ_E its characteristic function. We fix $\varepsilon_0 > 0$. Multiplying (5.35) by $h_l(u_\varepsilon)\chi_E$, integrating over Ω and using *ii)*, we obtain

$$j(r) \int_E h_l(u_\varepsilon) \geq \int_E j_{\varepsilon_0}(T_{l+1}(u_\varepsilon))h_l(u_\varepsilon) + (r - T_{l+1}(u_\varepsilon))\beta_{\varepsilon_0}(T_{l+1}(u_\varepsilon)) \quad (5.36)$$

for all $r \in \mathbb{R}$ and all $0 < \varepsilon < \min(\varepsilon_0, \frac{1}{l})$.

As $\varepsilon \rightarrow 0$, taking into account that E arbitrary we obtain from (5.36)

$$j(r)h_l(u) \geq j_{\varepsilon_0}(T_{l+1}(u))h_l(u) + bh_l(u)(r - T_{l+1}(u)) \quad (5.37)$$

for all $r \in \mathbb{R}$ and almost everywhere in Ω .

Passing to the limit with $l \rightarrow \infty$ and then with $\varepsilon_0 \rightarrow 0$ in (5.37) finally yields

$$j(r) \geq j(u(x)) + b(x)(r - u(x)) \quad (5.38)$$

for all $r \in \mathbb{R}$ and almost everywhere in Ω , hence $u \in D(\beta)$ and $b \in \beta(u)$ for almost everywhere in Ω . With this last step the proof of Theorem 5.1 is concluded. \square

6. CASE WHERE $f \in L^1(\Omega)$

6.1. Approximate solution for L^1 - data. The comparison principle from proposition will be the tool in second approximation procedure. For $f \in L^1(\Omega)$ and $m, n \in \mathbb{N}$ let $f_{m,n} \in L^\infty(\Omega)$ be defined as in Section 3. Using Proposition 4.3, we deduce that for any $m, n \in \mathbb{N}$, there exists $u_{m,n} \in W_0^{1, \vec{p}}(\Omega)$, $b_{m,n} \in L^\infty(\Omega)$, such that $(u_{m,n}, b_{m,n})$ is a renormalized solution of $(E, f_{m,n})$. Therefore

$$\int_\Omega b_{m,n} h(u_{m,n}) \phi + \int_\Omega (a(x, Du_{m,n}) + F(u_{m,n})) \cdot D(h(u_{m,n}) \phi) = f_{m,n} h(u_{m,n}) \phi \quad (6.1)$$

holds for all $m, n \in \mathbb{N}$, $h \in C_c^1(\mathbb{R})$, $\phi \in W_0^{1, \vec{p}}(\Omega) \cap L^\infty(\Omega)$. In the next lemma, we give a priori estimates that will be important in the the following:

Lemma 6.1. For $m, n \in \mathbb{N}$, let $(u_{m,n}, b_{m,n})$ be a renormalized solution of $(E, f_{m,n})$. Then,

i) For any $k > 0$ we have,

$$\sum_{i=1}^N \int_{\Omega} |DT_k(u_{m,n})|^{p_i} \leq \frac{k}{\gamma} \|f\|_1 \quad (6.2)$$

ii) for any $k > 0$, there exists a constant $C_3(k) > 0$, not depending on $m, n \in \mathbb{N}$, such that

$$\sum_{i=1}^N \int_{\Omega} |DT_k(u_{m,n})|^{p_i} \leq C_3(k). \quad (6.3)$$

iii) For $m, n \in \mathbb{N}$, we have:

$$\|b_{m,n}\|_1 \leq \|f\|_1. \quad (6.4)$$

Proof. For $l, k > 0$, we plug $h_l(u_{m,n})T_k(u_{m,n})$ as a test function in (6.1). Then i) and ii) follows with similar arguments as used in the proof of Lemma 5.4. To prove iii), we neglect the positive term

$$\int_{\Omega} a(x, DT_k(u_{m,n})) DT_k(u_{m,n})$$

and keep

$$\int_{\Omega} b_{m,n} T_k(u_{m,n}) \leq \int_{\Omega} f_{m,n} T(u_{m,n}). \quad (6.5)$$

Since $b_{m,n} \in \beta(u_{m,n})$ a.e. in Ω , it follows from (6.5) that

$$\int_{|u_{m,n}| > k} |b_{m,n}| \leq \int_{\Omega} |f|. \quad (6.6)$$

and we find iii) by passing to the limit with $k \rightarrow 0$. \square

By definition we have

$$f_{m,n} \leq f_{m+1,n} \quad \text{and} \quad f_{m,n+1} \leq f_{m,n} \quad (6.7)$$

From Propostion 5.2 it follows that

$$u_{m,n}^{\varepsilon} \leq u_{m+1,n}^{\varepsilon} \quad \text{and} \quad u_{m,n+1}^{\varepsilon} \leq u_{m,n}^{\varepsilon}, \quad (6.8)$$

almost everywhere in Ω for any $m, n \in \mathbb{N}$ and all $\varepsilon > 0$.

Hence passing to the limit with $\varepsilon \rightarrow 0$ in (6.8) yields

$$u_{m,n} \leq u_{m+1,n} \quad \text{and} \quad u_{m,n+1} \leq u_{m,n}, \quad (6.9)$$

almost everywhere in Ω for any $m, n \in \mathbb{N}$.

Setting $b_{\varepsilon} := \beta_{\varepsilon}(T_{1/\varepsilon}(u_{\varepsilon}))$, using (6.8), Remark 5.3 and the fact that $b_{m,n}^{\varepsilon} \rightarrow b_{m,n}$ in $L^{\infty}(\Omega)$ and since this convergence preserves order we get

$$b_{m,n} \leq b_{m+1,n} \quad \text{and} \quad b_{m,n+1} \leq b_{m,n} \quad (6.10)$$

almost everywhere in Ω for any $m, n \in \mathbb{N}$. By (6.10) and (6.4), for any $n \in \mathbb{N}$ there exist $b^n \in L^1(\Omega)$ such that $b_{m,n} \rightarrow b^n$ and $m \rightarrow \infty$ in $L^1(\Omega)$ and almost everywhere and $b \in L^1(\Omega)$, such that $b^n \rightarrow b$ as $n \rightarrow \infty$ in $L^1(\Omega)$ and almost everywhere in Ω . By (6.9), the sequence $(u_{m,n})_m$ is monotone increasing, hence, for

any $n \in \mathbb{N}$, $u_{m,n} \rightarrow u^n$ almost everywhere in Ω , where $u^n : \Omega \rightarrow \overline{\mathbb{R}}$ is a measurable function. In order to show that u is finite almost everywhere we will give an estimate on the level sets of $u_{m,n}$ in the next lemma:

Lemma 6.2. *For $m, n \in \mathbb{N}$ let $(u_{m,n}, b_{m,n})$ be a renormalized solution of $(E, f_{m,n})$. Then, there exists a constant $C_4 > 0$, not depending on $m, n \in \mathbb{N}$, such that*

$$|\{|u_{m,n}| \geq l\}| \leq C_4 l^{\frac{1}{\bar{p}}-1} \quad (6.11)$$

for all $l \geq 0$.

Proof. With the same arguments as in remark 5.5 we obtain

$$|\{|u_{m,n}| \geq l\}| \leq C(\bar{p}, N) l^{\bar{p}-} \left(\sum_{i=1}^N \int_{\Omega} |DT_k(u_{m,n})|^{p_i} + |\Omega| \right) \quad (6.12)$$

for all $m, n \in \mathbb{N}$ where $C(\bar{p}, N)$ is the constant from Sobolev embedding in (2.5). Now we plug (6.2) into (6.12) to obtain (6.11). Note that, as $(u_{m,n})_m$ is pointwise increasing with respect to m ,

$$\lim_{m \rightarrow \infty} |\{u_{m,n} \geq l\}| = |\{u^n \geq l\}| \quad (6.13)$$

and

$$\lim_{m \rightarrow \infty} |\{u_{m,n} \leq -l\}| = |\{u^n \leq -l\}|. \quad (6.14)$$

Combining (6.11) with (6.13) and (6.14) we get

$$|\{u^n \leq -l\}| + |\{u^n > l\}| \leq C_4 l^{\frac{1}{\bar{p}}-1} \quad (6.15)$$

for any $l \geq 1$, hence u^n is finite almost everywhere for $n \in \mathbb{N}$. By the same arguments we get

$$|\{u < -l\}| + |\{u > l\}| \leq C_4 l^{\frac{1}{\bar{p}}-1} \quad (6.16)$$

from (6.15), hence u is finite almost everywhere. Now, since $b_{m,n} \in \beta(u_{m,n})$ almost everywhere in Ω it follows by a subdifferential argument that $b^n \in \beta(u^n)$ and $b \in \beta(u)$ a.e. in Ω . \square

Remark 6.3. If $(u_{m,n}, b_{m,n})$ is renormalized solution of $(E, f_{m,n})$, using $h_\nu(u_{m,n})T_k(u_{m,n} - T_l(u_{m,n}))$ as a test function in (6.1), neglecting positive terms and passing to the limit with $\nu \rightarrow \infty$ we obtain

$$\int_{\{|l < |u_{m,n}| < l+k\}} a(x, Du_{m,n}) \cdot Du_{m,n} \leq k \left(\int_{\{|u_{m,n}| > l\} \cap \{|f| < \sigma\}} |f| + \int_{\{|f| > \sigma\}} |f| \right) \quad (6.17)$$

for any $k, \sigma > 0, l$. Now applying (6.11) to (6.17), we find that

$$\int_{\{|l < |u_{m,n}| < l+k\}} a(x, Du_{m,n}) \cdot Du_{m,n} \leq \sigma k C_4 l^{\frac{1}{\bar{p}}-1} + k \int_{\{|f| > \sigma\}} |f| \quad (6.18)$$

holds for any $k, \sigma > 0, l \geq 0$ uniformly in $m, n \in \mathbb{N}$.

6.2. Basic convergence results.

Lemma 6.4. *For $m, n \in \mathbb{N}$ let $(u_{m,n}, b_{m,n})$ be a renormalized solution of $(E, F_{m,n})$. There exists a subsequence $(m(n))_n$ such that setting $f_n := f_{m(n),n}$, $b_n := b_{m(n),n}$, $u_n := u_{m(n),n}$ we have*

$$u_n \rightarrow u \quad \text{almost everywhere in } \Omega. \quad (6.19)$$

Moreover, for any $k > 0$,

$$T_k(u_n) \rightarrow T_k(u) \quad \text{in } W_0^{1, \vec{p}}(\Omega), \quad (6.20)$$

$$DT_k(u_n) \rightharpoonup DT_k(u) \quad \text{in } \prod_{i=1}^N L^{p_i}(\Omega), \quad (6.21)$$

$$a(x, DT_k(u_n)) \rightharpoonup a(x, DT_k(u)) \quad \text{in } \prod_{i=1}^N L^{p_i}(\Omega), \quad (6.22)$$

as $n \rightarrow \infty$.

Proof. We construct a subsequence $(m(n))_n$, such that

$$\arctan(u_{m(n),n}) \rightarrow \arctan(u),$$

$$b_n := b_{m(n),n} \rightarrow b,$$

$$f_n := f_{m(n),n} \rightarrow f$$

as $n \rightarrow \infty$ in $L^1(\Omega)$ and almost everywhere in Ω . It follows that (6.19) and (6.20) hold. Combining (6.20) with (6.3) we get $T_k(u) \in W_0^{1, \vec{p}}(\Omega)$, $T_k(u_n) \rightarrow T_k(u) \in W_0^{1, \vec{p}}(\Omega)$ and (6.21) holds for any $k > 0$. From (6.2) and (H_2) , it follows that for fixed $k > 0$, given any subsequence of $(a(x, DT_k(u_n)))_n$ there exists a subsequence, still denoted by such that $a(x, DT_k(u_n))_n$, such that

$$a(x, DT_k(u_n))_n \rightharpoonup \Phi_k \quad \text{in } \prod_{i=1}^N L^{p_i}(\Omega)$$

as $n \rightarrow \infty$. Since $h_l(u_n)(T_k(u_n) - T_k(u))$ is an admissible test function in (6.1),

$$\limsup_{n \rightarrow \infty} \int_{\Omega} a(x, DT_k(u_n)) D(T_k(u_n) - T_k(u)) \leq 0. \quad (6.23)$$

Then, (6.22) follows with the same arguments as in the proof of Lemma 5.6. \square

Remark 6.5. With the same arguments as in Remark 5.7, we have

$$\lim_{n \rightarrow \infty} \int_{\Omega} a(x, DT_k(u_n) - a(x, DT_k(u))) \cdot D(T_k(u_n) - T_k(u)) = 0, \quad (6.24)$$

$$\lim_{l \rightarrow \infty} \int_{\{|l < |u| < l+1\}} a(x, Du) \cdot Du = 0. \quad (6.25)$$

6.3. Conclusion of the proof of Theorem 4.1. It is left to prove that (u, b) satisfies

$$\int_{\Omega} bh(u)\phi + \int_{\Omega} (a(x, Du) + F(u)).D(h(u)\phi) = \int_{\Omega} fh(u)\phi. \quad (6.26)$$

for all $h \in C_c^1(\mathbb{R})$ and $\phi \in W_0^{1, \vec{p}}(\Omega) \cap L^\infty(\Omega)$. To this end, we take $h \in C_c^1(\mathbb{R})$ and $\phi \in W_0^{1, \frac{\vec{p}}{2}}(\Omega) \cap L^\infty(\Omega)$ arbitrary and plug $h_l(u_n)h(u)\phi$ into (6.1) to obtain

$$I_{n,l}^1 + I_{n,l}^2 + I_{n,l}^3 = I_{n,l}^4, \quad (6.27)$$

where

$$\begin{aligned} I_{n,l}^1 &= \int_{\Omega} b_n h_l(u_n) h(u) \phi, \\ I_{n,l}^2 &= \int_{\Omega} a(x, Du_n).D(h_l(u_n)h(u)\phi), \\ I_{n,l}^3 &= \int_{\Omega} F(u_n).D(h_l(u_n)h(u)\phi), \\ I_{n,l}^4 &= \int_{\Omega} f_n h_l(u_n) h(u) \phi. \end{aligned}$$

Step 1. Passing to the limit as $n \rightarrow \infty$, applying the convergence results from Lemma 6.4 we get

$$\lim_{n \rightarrow \infty} I_{n,l}^1 = \int_{\Omega} b h_l(u) h(u) \phi, \quad \lim_{n \rightarrow \infty} I_{n,l}^4 = \int_{\Omega} f h_l(u) h(u) \phi. \quad (6.28)$$

Let us write

$$I_{n,l}^2 = I_{n,l}^{2,1} + I_{n,l}^{2,2}, \quad (6.29)$$

where

$$I_{n,l}^{2,1} = \int_{\Omega} h_l(u_n) a(x, Du_n).D(h(u)\phi), \quad I_{n,l}^{2,2} = \int_{\Omega} h_l'(u_n) a(x, Du_n).Du_n h(u)\phi. \quad (6.30)$$

With similar arguments as in the proof of (5.26) it follows that

$$\lim_{n \rightarrow \infty} I_{n,l}^{2,1} = \int_{\Omega} h_l(u) a(x, Du).D(h(u)\phi). \quad (6.31)$$

By (6.18), we get the estimate

$$\left| \lim_{n \rightarrow \infty} I_{n,l}^{2,2} \right| \leq \|h\|_{\infty} \|\phi\|_{\infty} (\delta C_4 l^{\frac{1}{\vec{p}}-1} + \int_{\{|f|>\delta\}} |f|), \quad (6.32)$$

for all $n \in \mathbb{N}$ and all $l \geq 1$, $\delta > 0$. Next, we write

$$I_{n,l}^3 = I_{n,l}^{3,1} + I_{n,l}^{3,2},$$

where

$$\lim_{n \rightarrow \infty} I_{n,l}^{3,1} = \int_{\Omega} h_l(u) F(u).D(h(u)\phi), \quad \lim_{n \rightarrow \infty} I_{n,l}^{3,2} = \int_{\Omega} h_l'(u) F(u).Du h(u)\phi, \quad (6.33)$$

follows with the same arguments as in (5.27) – (5.31).

Step 2. Passing to the limit as $l \rightarrow \infty$. Combining (6.27) with (6.28)-(6.33) we get for all $\delta > 0$ and all $l \geq 1$

$$I_l^1 + I_l^2 + I_l^3 + I_l^4 + I_l^5 = I_l^6, \quad (6.34)$$

where

$$I_l^1 = \int_{\Omega} bh_l(u)h(u)\phi, \quad I_l^2 = \int_{\Omega} h_l(u)a(x, DT_{l+1}(u)).D(h(u)\phi)$$

$$|I_l^3| \leq \|h\|_{\infty}\|\phi\|_{\infty}(\delta C_4 l^{\frac{1}{p}-1} + \int_{\{|f|>\delta\}} |f|),$$

for any $\delta > 0$ and

$$I_l^4 = \int_{\Omega} h_l'(u)F(u)h(u)\phi Du, \quad I_l^5 = \int_{\Omega} h_l(u)F(u).D(h(u)\phi), \quad |I_l^6| = \int_{\Omega} fh_l(u)h(u)\phi.$$

Choosing $m > 0$ such that $\text{supph} \subset [-m, m]$, we can replace u by $T_m(u)$ in $I_l^1, I_l^2, \dots, I_l^6$ hence

$$\lim_{l \rightarrow \infty} I_l^1 = \int_{\Omega} bh(u)\phi, \quad \lim_{l \rightarrow \infty} I_l^2 = \int_{\Omega} a(x, Du).D(h(u)\phi), \quad (6.35)$$

$$\lim_{l \rightarrow \infty} |I_l^3| \leq \|h\|_{\infty}\|\phi\|_{\infty} \int_{\{|f|>\sigma\}} |f|, \quad \lim_{l \rightarrow \infty} I_l^4 = 0, \quad (6.36)$$

$$\lim_{l \rightarrow \infty} |I_l^5| = \int_{\Omega} F(u).D(h(u)\phi), \quad \lim_{l \rightarrow \infty} |I_l^6| = \int_{\Omega} fh(u)\phi, \quad (6.37)$$

for all $\delta > 0$. Combining (6.34) with (6.35)-(6.37) we finally deduce that (6.1) holds for all $h \in C_C^1(\mathbb{R})$ and all $\phi \in W_0^{1, \vec{p}}(\Omega) \cap L^{\infty}(\Omega)$.

Hence (u, b) satisfies (R1), (R2) and (R3) and the proof of the theorem is completed.

6.4. Proof of Theorem 4.2 (Uniqueness).

Lemma 6.6. For $f, \tilde{f} \in L^1(\Omega)$ let $(u, b), (\tilde{u}, \tilde{b})$ be the renormalized solutions to (E, f) and (E, \tilde{f}) respectively, then

$$\int_{\Omega} (b - \tilde{b}) \text{Sign}_0^+(u - \tilde{u}) dx \leq \int_{\Omega} (f - \tilde{f}) \text{Sign}_0^+(u - \tilde{u}) dx, \quad (6.38)$$

Proof. For $\delta > 0$ let H_{δ}^+ be a Lipschitz approximation of the sign_0^+ function. Since $(u, b), (\tilde{u}, \tilde{b})$ are renormalized solutions, it follows that

$$T_{l+1}(u), T_{l+1}(\tilde{u}) \in W_0^{1, \vec{p}}(\Omega) \cap L^{\infty}(\Omega) \text{ for all } l > 0.$$

Hence $H_{\delta}^+(T_{l+1}(u) - T_{l+1}(\tilde{u})) \in W_0^{1, \vec{p}}(\Omega) \cap L^{\infty}(\Omega)$ for $l, \delta > 0$. Now, we choose $H_{\delta}^+(T_{l+1}(u) - T_{l+1}(\tilde{u}))$ as a test function in the renormalized formulation with $h = h_l$ for (u, b) and for (\tilde{u}, \tilde{b}) respectively. Subtracting the resulting equalities, we obtain

$$I_{l,\delta}^1 + I_{l,\delta}^2 + I_{l,\delta}^3 + I_{l,\delta}^4 + I_{l,\delta}^5 = I_{l,\delta}^6, \quad (6.39)$$

where $K = \{0 < T_{l+1}(u) - T_{l+1}(\tilde{u}) < \delta\}$ and

$$I_{l,\delta}^1 = \int_{\Omega} (bh_l(u) - \tilde{b}h_l(\tilde{u}))H_{\delta}^+(T_{l+1}(u) - T_{l+1}(\tilde{u}))dx,$$

$$I_{l,\delta}^2 = \int_{\Omega} (h'_l(u)a(x, Du).Du - h'_l(\tilde{u})a(x, D\tilde{u}).D\tilde{u}).H_{\delta}^+(T_{l+1}(u) - T_{l+1}(\tilde{u}))dx,$$

$$I_{l,\delta}^3 = \frac{1}{\delta} \int_K (h_l(u)a(x, Du) - h_l(\tilde{u})a(x, D\tilde{u})).D(T_{l+1}(u) - T_{l+1}(\tilde{u}))dx,$$

$$I_{l,\delta}^4 = \int_{\Omega} (h'_l(u)F(u).Du - h'_l(\tilde{u})F(\tilde{u}).D\tilde{u})H_{\delta}^+(T_{l+1}(u) - T_{l+1}(\tilde{u}))dx,$$

$$I_{l,\delta}^5 = \frac{1}{\delta} \int_K (h_l(u)F(u) - h_l(\tilde{u})F(\tilde{u})).D(T_{l+1}(u) - T_{l+1}(\tilde{u}))dx,$$

$$I_{l,\delta}^6 = \int_{\Omega} (fh_l(u) - \tilde{f}h_l(\tilde{u}))H_{\delta}^+(T_{l+1}(u) - T_{l+1}(\tilde{u}))dx.$$

Using the same arguments as in [33] i.e., neglecting the nonnegative part of $I_{l,\delta}^3$ and using that F is locally Lipschitz continuous, we can pass to the limit as $\delta \rightarrow 0$.

Using the energy dissipation condition (R_3) we can pass the limit as $l \rightarrow \infty$ and obtain (6.38).

Now we are in position to give the proof of Theorem 4.2:

Assuming $f = \tilde{f}$, from lemma 6.6 we get

$$\int_{\Omega} (b - \tilde{b})\text{sign}_0^+(u - \tilde{u})dx \leq 0, \quad (6.40)$$

hence $(b - \tilde{b})\text{sign}_0^+(u - \tilde{u}) = 0$ almost everywhere in Ω . Now, let us write $\Omega = \Omega_1 \cup \Omega_2$, where $\Omega_1 := \{x \in \Omega : \text{sign}_0^+(u(x) - \tilde{u}(x)) = 0\}$, $\Omega_2 := \{x \in \Omega : (b(x) - \tilde{b}(x)) = 0\}$. Since $r \mapsto \beta(x, r)$ is strictly increasing for a.e. $x \in \Omega$, we can define the function $\beta^{-1} : \mathbb{R} \rightarrow \mathbb{R}$ such that $\beta^{-1} = s$ for all $(r, s) \in \mathbb{R}^2$ such that $r \in \beta(x, r)$ for a.e. $x \in \Omega$. For a.e. $x \in \Omega_2$ we have $b(x) = \tilde{b}(x)$, hence $u(x) = \beta^{-1}(b(x)) = \beta^{-1}(\tilde{b}(x)) = \tilde{u}(x)$. Therefore, $u(x) = \tilde{u}(x)$ a.e. in Ω_2 and $\text{sign}_0^+(u - \tilde{u}) = 0$. Interchanging the roles of u and \tilde{u} and repeating the arguments, we get $\text{sign}_0^+(\tilde{u} - u) = 0$ a.e. in Ω and we finally arrive at $u = \tilde{u}$ a.e. in Ω . Now, we write the renormalized formulation for (u, b) and (\tilde{u}, \tilde{b}) respectively. Subtracting the resulting equalities, we obtain

$$\int_{\Omega} (b - \tilde{b})h(u)\varphi dx = 0$$

for all $h \in C_c^1(\mathbb{R})$ and all $\varphi \in C_c^\infty(\Omega)$. Choosing $h(u) = h_l(u)$ and passing to the limit with $l \rightarrow \infty$ we find $b = \tilde{b}$ a.e. in Ω .

7. PROOF OF PROPOSITION 4.3

Note that for $\varepsilon, k > 0$, $h_l(u) \frac{1}{\varepsilon} T_\varepsilon(u - T_k(u))$ as a test function in (3.2). Neglecting positive terms and passing to the limit with $l \rightarrow \infty$, we obtain

$$\frac{1}{\varepsilon} \sum_{i=1}^N \int_{k < |u| < k + \varepsilon} |Du|^{p_i} \leq \|f\|_N (\phi(k))^{(N-1)/N}, \quad (7.1)$$

where $\phi(k) := |\{|u| > k\}|$ for $k > 0$. Now we use similar arguments as in [33]. We apply the continuous embedding of $W_0^{1,1}(\Omega)$ into $L^{N/N-1}(\Omega)$ and the Hölder inequality to get

$$\frac{1}{\varepsilon C_N} \|T_\varepsilon(u - T_k(u))\|_{\frac{N}{N-1}} \leq \left(\frac{\phi(k) - \phi(k + \varepsilon)}{\varepsilon} \right)^{1/(p^-)'} \left(\frac{1}{\varepsilon} \int_{k < |u| < k + \varepsilon} |Du|^{p^-} \right)^{1/p^-}, \quad (7.2)$$

where $C_N > 0$ is the constant coming from the Sobolev embedding. Notice that

$$\frac{1}{\varepsilon} \sum_{i=1}^N \int_{k < |u| < k + \varepsilon} |Du|^{p^-} \leq \frac{\phi(k) - \phi(k + \varepsilon)}{\varepsilon} + \frac{1}{\varepsilon} \sum_{i=1}^N \int_{k < |u| < k + \varepsilon} |Du|^{p_i}, \quad (7.3)$$

hence, from (7.1), (7.2) and (7.3) we deduce that

$$\frac{1}{\varepsilon C_N} \|T_\varepsilon(u - T_k(u))\|_{\frac{N}{N-1}} \leq \left(\frac{\phi(k) - \phi(k + \varepsilon)}{\varepsilon} \right)^{1/(p^-)'} \left(\frac{\phi(k) - \phi(k + \varepsilon)}{\varepsilon} + \|f\|_N (\phi(k))^{(N-1)/N} \right)^{1/(p^-)}. \quad (7.4)$$

From (7.4) and Young's inequality with $\alpha > 0$ it follows that

$$\frac{1}{C_N C} (\phi(k + \varepsilon))^{(N-1)/N} - \frac{\alpha^{p^-}}{p^- C} \|f\|_N (\phi(k))^{(N-1)/N} - \frac{\phi(k) - \phi(k + \varepsilon)}{\varepsilon} \leq 0, \quad (7.5)$$

where

$$C := \left(\frac{1}{\alpha^{(p^-)'} (p^-)'} + \frac{\alpha^{p^-}}{p^-} \right) > 0.$$

The mapping $(0, \infty) \ni k \rightarrow \phi(k)$ is non-increasing and therefore of bounded variation, hence it is differentiable almost everywhere on $(0, \infty)$ with $\phi' \in L_{loc}^1(0, \infty)$. Since it is also continuous from the right, we can pass to the limit with $\varepsilon \downarrow 0$ in (7.5) to find

$$C'' (\phi(k))^{(N-1)/N} + \phi'(k) \leq 0 \quad (7.6)$$

for almost every $k > 0$ and $\alpha > 0$ chosen small enough such that

$$C'' := \left(\frac{C_N}{C} - \frac{\alpha^{p^-}}{p^- C} \|f\|_N \right) > 0.$$

Now, the conclusion of the proof follows by contradiction. We assume that $\phi(k) > 0$ for each $k > 0$. For $k > 0$ fixed, we choose $k_0 < k$. From (7.6) it follows that

$$\frac{1}{N}C'' + \frac{d}{ds}((\phi(s))^{(1/N)}) \leq 0 \quad (7.7)$$

for almost all $s \in (k_0, k)$. The left hand side of (7.7) is in $L^1(k_0, k)$, hence we integrate (7.7) over $[k_0, k]$. Moreover, since ϕ is non-increasing, integrating (7.7) over (k_0, k) we get

$$(\phi(k))^{1/N} \leq \phi(k_0)^{1/N} + \frac{1}{N}C''(k_0 - k) \quad (7.8)$$

and from (7.8) the contradiction follows.

8. EXAMPLE

This section is devoted to an example for illustrating our abstract result. Let us consider the special case:

$$\beta(r) = (r - 1)^+ - (r - 1)^-, \quad F : \mathbb{R} \rightarrow (F_i)_{i=1, \dots, N} \in \mathbb{R}^N,$$

where F is locally lipshitz continuous function, and

$$a_i(x, \xi) = \sum_{i=1}^N |\xi_i|^{p_i-1} \text{sgn}(\xi_i), \quad i = 1, \dots, N,$$

the $a_i(x, \xi)$ are Carathéodory function satisfying the growth condition (\mathbf{H}_2) , and the coercivity (\mathbf{H}_1) . On the other the monotonicity condition is verified. In fact

$$\sum_{i=1}^N \left(a_i(x, \xi) - a_i(x, \tilde{\xi}) \right) (\xi_i - \tilde{\xi}_i) = \sum_{i=1}^N \left(|\xi_i|^{p_i-1} \text{sgn}(\xi_i) - |\tilde{\xi}_i|^{p_i-1} \text{sgn}(\tilde{\xi}_i) \right) (\xi_i - \tilde{\xi}_i) \geq 0,$$

for almost all $x \in \Omega$ and for all $\xi, \tilde{\xi} \in \mathbb{R}^N$. This last inequality can not be strict, since for $\xi \neq \tilde{\xi}$ with $\xi_N \neq \tilde{\xi}_N$ and $\xi = \tilde{\xi}, i = 1, \dots, N - 1$. The corresponding expression is zero.

Therefore, for all $f \in L(\Omega)$, the following problem:

$$\left\{ \begin{array}{l} T_k(u) \in W_0^{1, \vec{p}}(\Omega) \text{ for } (k > 0); b \in L^1(\Omega) \text{ and } b(x) \in \beta(u(x)), \\ \lim_{l \rightarrow \infty} \int_{\{|u| < l+1\}} a(x, Du) \cdot D u dx = 0, \\ \int_{\Omega} b h(u) \varphi dx + \int_{\Omega} h(u) \sum_{i=1}^N \left| \frac{\partial u}{\partial x_i} \right|^{p_i-1} \text{sgn} \left(\frac{\partial u}{\partial x_i} \right) \cdot \frac{\partial \varphi}{\partial x_i} dx \\ + \int_{\Omega} h'(u) \sum_{i=1}^N \left| \frac{\partial u}{\partial x_i} \right|^{p_i-1} \text{sgn} \left(\frac{\partial u}{\partial x_i} \right) \cdot \frac{\partial \varphi}{\partial x_i} dx + \int_{\Omega} F(u) \cdot D(h(u) \varphi) dx \\ = \int_{\Omega} f \cdot D(h(u) \varphi), \quad \forall \varphi \in W_0^{1, \vec{p}}(\Omega) \cap L^\infty(\Omega) \text{ and } h \in C_c^1(\mathbb{R}), \end{array} \right.$$

at least one renormalized solution.

REFERENCES

- [1] L. Aharouch, E. Azroul and A. Benkirane, Quasilinear degenerated equations with L^1 datum and without coercivity in perturbation terms, *Electron. J. Qualit. Theory Differential Equations* (2006), 19.18.
- [2] L. Aharouch, A. Benkirane, J. Bennouna and A. Touzani, Existence and uniqueness of solutions of somme nonlinear equations in Orlicz spaces and weithed Sobolev spaces, in: *Recent Developement in Nonlinear Analysis*, World Sci (2010), 170-180.
- [3] Y. Akdim, C. Allalou: Existence of renormalized solutions of nonlinear elliptic problems in weighted variable-exponent space. *J. Math. Study. Vol(48)4* (2015), 375-397.
- [4] F. Andreu, N. Igbida, J. M. Mazón and J. Toledo, L^1 existence and uniqueness results for quasi-linear elliptic equations with nonlinear boundary conditions. *Ann. Inst. H. Poincaré. Anal. Non Linéaire vol(24)(2007)*, 61-89.
- [5] S. Antontsev, S. Shmarev, Elliptic equations and systems with nonstandard growth conditions: existence, uniqueness and localization properties of solutions. *Anal.Nonlinear vol (65)* (2006), 728-761.
- [6] S. N. Antontsev, J. I. Díaz, S. Shmarev, *Energy Methods for Free Boundary Problems. Applications to Nonlinear PDEs and Fluid Mechanics. In: Progress in Nonlinear Differential Equations and Their Applications. vol. 48,* (2002).
- [7] S. Antontsev, S. Shmarev, Elliptic equations and systems with nonstandard growth conditions: existence, uniqueness and localization properties of solutions. *Nonlinear Anal. vol(65)* (2006), 728-761.
- [8] M. Bendahmane, K. H. Karlsen, Anisotropic nonlinear elliptic systems with measure data and anisotropic harmonic maps into spheres. *Electron. J. Differential Equations*, no. vol (46) (2006).
- [9] M. Bendahmane, K. H. Karlsen, Renormalized solutions of an anisotropic reaction-diffusion-advection system with L^1 data. *Commun. Pure Appl. Anal. vol(5)* (2006), 733-762.
- [10] M. Bendahmane and P. Wittbold. Renormalized solutions for nonlinear elliptic equations with variable exponents and L^1 -data. *Nonlinear Anal. TMA. vol(70)2* (2009), 567-583.
- [11] A. Benkirane and J. Bennouna, Existence of solutions for nonlinear elliptic degenerate equations, *Nonlinear Anal. vol(54)(2003)*, 9-37.
- [12] P. Bénilan, L. Boccardo, T. Gallouët, R. Gariepy, M. Pierre, and J.L.Vázquez: An L^1 -theory of existence and uniqueness of solutions of nonlinear equations, *Ann. Scuola Norm. Sup. Pisa, Cl. Sci. vol(22)2* (1995), 241-273.
- [13] L. Boccardo, T. Gallouet, P. Marcellini, Anisotropic equations in " L^1 . *Diff. Int. Eqs. vol(9)* (1996), 209-212.
- [14] L. Boccardo, J.I. Diaz, D. Giachetti, and F. Murat, Existence of a solution for a weaker form of a nonlinear elliptic equation, *Recent advances in nonlinear elliptic and parabolic problems (Nancy 1988)*, Pitman Res. Notes Mat. Ser., vol(208), Longman Sci. Tech., Harlow, (1989), 229-246.
- [15] D. Blanchard, F. Murat, and H. Redwane, Existence and Uniqueness of a Renormalized Solution for a Fairly General Class of Nonlinear Parabolic Problems, *J. Diff. Equ. vol(177)* (2001), 331-347.
- [16] H. Brézis. *Opérateurs Maximaux Monotones*. North-Holland, Amsterdam, (1973).
- [17] A. Cianchi, Symmetrization in anisotropic elliptic problems. *Comm. Part. Diff. Eqs. vol(32)* (2007), 693-717.
- [18] Andrea Cianchi, A Fully anisotropic Sobolev inequality, *Pacific Journal of Mathematic*, No.2, Vol(196) (2000), 283-294.
- [19] A. Damlamian, Some result on the multiphase Stefan problem, *Comm. Partial Diff. Equations vol(2)* (1977), 1017-1044.

- [20] G. Dal Maso, F. Murat, L. Orsina, and A. Prignet, Renormalized solutions of elliptic equations with general measure data, *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, vol(28) (1999), 741-808.
- [21] R.J. DiPerna and P-L. Lions, On the Cauchy problem for Boltzmann equations: global existence and weak stability, *Ann. of Math.* vol(130)2 (1989), 321-366.
- [22] A. El Hamidi, J. M. Rakotoson, Extremal functions for the anisotropic Sobolev inequalities. *Ann. Inst. H. Poincaré Anal. Non Linéaire* vol(24) (2007), 741–756.
- [23] I. Fragalà, F. Gazzola, B. Kawohl, Existence and nonexistence results for anisotropic quasi-linear elliptic equations, *Ann. Inst. H. Poincaré Anal. Non Linéaire* vol(21)5 (2004), 715-734.
- [24] A. Friedman, *Variational Principles and Free-boundary Problems*, Wiley, New York, 1982.
- [25] P. Gwiazda, P. Wittbold, A. Wróblewska and A. Zimmermann, Renormalized solutions of nonlinear elliptic problems in generalized Orlicz spaces, *J. Differential Equations* vol(253)(2012), 635-666.
- [26] H. Hjjaj, Stanislas Ouaro, M. Badr Benboubker: Entropy solutions to nonlinear anisotropic problem with variable exponent. *Journal of Applied Analysis and Computation* Vol(4)3 (2014).
- [27] A.G. Korolev, Embedding theorems for anisotropic Sobolev-Orlicz spaces, *Vestnik Moskovskogo Universiteta, Matematika*, vol(38) (1983), 32-37 .
- [28] J. L. Lions, *Quelques méthodes de résolution des problèmes aux limites non linéaires*, Dunod, Paris, (1969).
- [29] S. M. Nikolskii, An imbedding theorem for functions with partial derivatives considered in different metrics, *Izv. Akad. Nauk SSSR Ser. Mat.*, 22 (1958), 321-336.
- [30] F. Murat, Soluciones renormalizadas de EDP elipticas non lineales, Tech. Report R93023, Laboratoire d'Analyse Numérique, Paris VI, 1993, Cours à l'Université de Séville.
- [31] M. Růžička, *Electrorheological Fluids: Modelling and Mathematical Theory*, Springer, Berlin, (2000).
- [32] M. Troisi, Teoremi di inclusione per spazi di Sobolev non isotropi, *Ricerche Mat.* vol(18)(1969), 3-24.
- [33] P. Wittbold and Zimmermann, Existence and uniqueness of renormalized solutions to nonlinear elliptic equations with variable exponents and L^1 – data, *Nonlinear Anal.* vol(72)(2010), 2990-3008.

¹ SIDI MOHAMED BEN ABDELLAH UNIVERSITY, LAMA LABORATORY, FACULTY OF SCIENCES D'HAR EL MAHRAZ, P.O. BOX 1796 ATLAS FEZ, MOROCCO.

Email address: ¹youssef.akdim@usmba.ac.ma

^{2,3} SIDI MOHAMED BEN ABDELLAH UNIVERSITY, LABORATORY OF ENGINEERING SCIENCES, MULTIDISCIPLINARY FACULTY OF TAZA, P.O. BOX 1223 TAZA, MOROCCO.

Email address: ²mohammed.elansari1@usmba.ac.ma, ³soumia.lalaoui@usmba.ac.ma