

HEAT KERNEL ESTIMATE IN A CONICAL SINGULAR SPACE

XIAOQI HUANG AND JUNYONG ZHANG

ABSTRACT. Let (X, g) be a product cone with the metric $g = dr^2 + r^2h$, where $X = C(Y) = (0, \infty)_r \times Y$ and the cross section Y is a $(n-1)$ -dimensional closed Riemannian manifold (Y, h) . We study the upper boundedness of heat kernel associated with the operator $\mathcal{L}_V = -\Delta_g + V_0 r^{-2}$, where $-\Delta_g$ is the positive Friedrichs extension Laplacian on X and $V = V_0(y)r^{-2}$ and $V_0 \in C^\infty(Y)$ is a real function such that the operator $-\Delta_h + V_0 + (n-2)^2/4$ is a strictly positive operator on $L^2(Y)$. The new ingredient of the proof is the Hadamard parametrix and finite propagation speed of wave operator on Y .

1. INTRODUCTION

Let (Y, h) be a $(n-1)$ -dimensional closed Riemannian manifold, we consider the product cone $X = C(Y) = (0, \infty)_r \times Y$ and the metric $g = dr^2 + r^2h$. The product cone is an incomplete manifold, however one can complete it to $C^*(Y) = C(Y) \cup P$ where P is its cone tip, see Cheeger [5, 6]. Let Δ_g denote the Friedrichs' self-adjoint extension of Laplace-Beltrami operator from the domain $C_c^\infty(X)$ that consist of the compactly supported smooth functions on the interior of the metric cone. One can write

$$(1.1) \quad -\Delta_g = -\partial_r^2 - \frac{n-1}{r}\partial_r + \frac{-\Delta_h}{r^2},$$

where $-\Delta_h$ is the positive Laplacian on the closed Riemannian manifold Y , see [7, p. 302] and [20, Theorem 2.1]. The heat kernel associated with the operator Δ_g has been investigated, we refer to Moer [20] and Nagase [21] for asymptotic expansion, to Li [17] for upper boundedness and to Coulhon-Li [3] for lower boundedness.

In this paper, we consider the heat kernel associated with the Schrödinger operator

$$(1.2) \quad \mathcal{L}_V = -\Delta_g + V, \quad V = V_0(y)r^{-2}$$

where $V_0(y)$ is a smooth function on the section Y such that the operator $-\Delta_h + V_0 + (n-2)^2/4$ is a strictly positive operator on $L^2(Y)$ space. The decay of the perturbation potential considered is scaling critical and is closely related to the angular momentum as $r \rightarrow \infty$, hence the Schrödinger operator \mathcal{L}_V has attracted interest from other topics. For examples, we refer to [2, 25] for the asymptotical behavior of the Schrödinger propagator, to [13, 16] for the Riesz transform, to [10, 28–30] for the Strichartz estimates and the restriction estimates. In the present paper, we focus on the upper boundedness of heat kernel. More precisely, we prove

Theorem 1.1. *Let \mathcal{L}_V be the operator on metric cone of dimension $n \geq 2$ given in (1.2) and suppose $\{\lambda_k, \varphi_k\}_{k=0}^\infty$ to be the eigenvalues and eigenfunctions of the operator*

$-\Delta_h + V_0(y) + (n-2)^2/4$, which satisfies

$$(1.3) \quad \begin{cases} (-\Delta_h + V_0(y) + (n-2)^2/4)\varphi_k(y) = \lambda_k\varphi_k(y), \\ \int_Y |\varphi_k(y)|^2 dy = 1. \end{cases}$$

and the eigenvalues $\{\lambda_k\}_{k=0}^\infty$ enumerated such that

$$(1.4) \quad 0 < \lambda_0 \leq \lambda_1 \leq \dots$$

repeating each eigenvalue as many times as its multiplicity. Then, for $t > 0$ and $(r, y), (s, y') \in X$, the heat kernel can be written as

$$(1.5) \quad e^{-t\mathcal{L}_V}(r, y; s, y') = \frac{e^{-\frac{r^2+s^2}{4t}}}{2t} (rs)^{-\frac{n-2}{2}} \sum_{k \in \mathbb{N}} \varphi_k(y) \overline{\varphi_k(y')} I_{\mu_k} \left(\frac{rs}{2t} \right),$$

where $\mu_k = \sqrt{\lambda_k}$ and I_μ is the modified Bessel function of the first kind of order μ . Furthermore, there exist positive constants c and C such that

$$(1.6) \quad |e^{-t\mathcal{L}_V}(r, y; s, y')| \leq C \left[\min \left\{ 1, \left(\frac{rs}{2t} \right) \right\} \right]^{-\sigma} t^{-\frac{n}{2}} e^{-\frac{d^2((r,y),(s,y'))}{ct}},$$

where $\sigma = \frac{n-2}{2} - \mu_0$. Here $d((r, y), (s, y'))$ is the distance between two points $(r, y), (s, y') \in X$

$$(1.7) \quad d((r, y), (s, y')) = \begin{cases} \sqrt{r^2 + s^2 - 2rs \cos(d_h(y, y'))}, & d_h(y, y') \leq \pi; \\ r + s, & d_h(y, y') \geq \pi, \end{cases}$$

and d_h is the distance on the section Y .

Remark 1.1. From (1.6), the square root of the smallest eigenvalue λ_0 plays a non-trivial role. In particular, when $Y = \mathbb{S}^{n-1}$ and $V_0(y) = a$ with $a \geq -(n-2)^2/4$, then

$$\sigma = \frac{n-2}{2} - \sqrt{\frac{(n-2)^2}{4} + a},$$

which is positive when $-(n-2)^2/4 \leq a < 0$ while is nonpositive when $a \geq 0$. Hence if $-(n-2)^2/4 \leq a < 0$, then

$$\left[\min \left\{ 1, \left(\frac{rs}{2t} \right) \right\} \right]^{-\sigma} \leq C \left[\min \left\{ \frac{r}{\sqrt{t}}, 1 \right\} \right]^{-\sigma} \left[\min \left\{ \frac{s}{\sqrt{t}}, 1 \right\} \right]^{-\sigma}$$

together with (1.6) shows

$$|e^{-t(-\Delta+a|z|^{-2})}(z, z')| \leq C \left[\min \left\{ \frac{|z|}{\sqrt{t}}, 1 \right\} \right]^{-\sigma} \left[\min \left\{ \frac{|z'|}{\sqrt{t}}, 1 \right\} \right]^{-\sigma} t^{-\frac{n}{2}} e^{-\frac{|z-z'|}{ct}},$$

which consists with the results of Liskevich-Sobol [18] and Milman-Semenov [19].

Remark 1.2. The proof is based on the Hadamard parametrix and finite propagation speed of wave operator on Y , which is a bit different from [17] due to the perturbation of the potential V . In [17], Li used a theorem of Grigor'yan [11, Theorem 1.1] which claims that if the heat kernel $H(t, r, y; s, y')$ on Riemannian manifolds satisfies on-diagonal bounds

$$H(t, r, y; r, y) \lesssim t^{-\frac{n}{2}}, \quad H(t, s, y'; s, y') \lesssim t^{-\frac{n}{2}},$$

then we have

$$H(t, r, y; s, y') \lesssim t^{-\frac{n}{2}} \exp\left(-\frac{d(r, y; s, y')^2}{ct}\right).$$

However, we do not know whether the analogs hold or not for the Laplacian with the perturbation of Hardy type potential. Therefore, we have to use a different argument instead.

Remark 1.3. As an application of the estimate (1.6) of the heat kernel, one can use the argument of [14] and the Hardy inequality in [2] to establish the Littlewood-Paley theory, Sobolev embedding associated with the operator \mathcal{L}_V . For example, we can prove the Mihlin Multipliers estimates: Suppose $m : [0, \infty) \rightarrow \mathbb{C}$ satisfies

$$(1.8) \quad |\partial^j m(\lambda)| \lesssim \lambda^{-j} \quad \text{for all } j \geq 0$$

Then $m(\sqrt{\mathcal{L}_V})$ is a bounded operator on $L^p(\mathbb{R}^n)$ provided that either

- $\mu_0 > (n-2)/2$ and $1 < p < \infty$, or
- $0 < \mu_0 \leq \frac{n-2}{2}$ and $p_0 < p < p'_0 := \frac{n}{\sigma}$.

Acknowledgement. The authors would like to thank Hongquan Li for his discussions and comments that improved the exposition. JZ acknowledges support from National Natural Science Foundation of China (12171031, 11831004).

2. CONSTRUCTION OF THE HEAT KERNEL

In this section, in spirit of functional calculus in Cheeger-Taylor [7, 8], we construct the heat kernel associated with the operator \mathcal{L}_V . Since the metric $g = dr^2 + r^2 h(y, dy)$, we write the operator \mathcal{L}_V in the coordinate (r, y) ,

$$(2.1) \quad \mathcal{L}_V = -\partial_r^2 - \frac{n-1}{r} \partial_r + \frac{1}{r^2} (-\Delta_h + V_0(y)).$$

where Δ_h is the Laplace-Beltrami operator on (Y, h) .

From classical spectral theory, the spectrum of $-\Delta_h + V_0(y) + (n-2)^2/4$ is formed by a countable family of real eigenvalues $\{\lambda_k\}_{k=0}^\infty$ enumerated such that

$$(2.2) \quad 0 < \lambda_0 \leq \lambda_1 \leq \dots$$

where we repeat each eigenvalue as many times as its multiplicity, and $\lim_{k \rightarrow \infty} \lambda_k = +\infty$. Let $\{\varphi_k(y)\}$ be the eigenfunctions of $-\tilde{\Delta}_h = -\Delta_h + V_0(y) + (n-2)^2/4$, that is

$$(2.3) \quad \begin{cases} (-\Delta_h + V_0(y) + (n-2)^2/4)\varphi_k(y) = \lambda_k \varphi_k(y), \\ \int_Y |\varphi_k(y)|^2 dy = 1. \end{cases}$$

Therefore, we obtain an orthogonal decomposition of the $L^2(Y)$ in a sense that

$$L^2(Y) = \bigoplus_{k \in \mathbb{N}} \mathcal{H}^k, \quad \mathbb{N} = \{0, 1, 2, \dots\}.$$

where

$$\mathcal{H}^k = \text{span}\{\varphi_k\}.$$

Define the orthogonal projection π_k of f onto \mathcal{H}^k by

$$(2.4) \quad \pi_k f = \int_Y f(r, y') H_k(y, y') dh, \quad f \in L^2(X),$$

where dh is the measure on Y under the metric h and the kernel

$$(2.5) \quad H_k(y, y') = \varphi_k(y) \overline{\varphi_k(y')},$$

then

$$(2.6) \quad f(r, y) = \sum_{k \in \mathbb{N}} \pi_k f = \sum_{k \in \mathbb{N}} a_k(r) \varphi_k(y), \quad a_k(r) = \int_Y f(r, y') \overline{\varphi_k(y')} dh.$$

Let $\mu = \mu_k = \sqrt{\lambda_k}$, for $f \in L^2(X)$, as [1, Page 523], we define the Hankel transform of order μ

$$(2.7) \quad (\mathcal{H}_\mu f)(\rho, y) = \int_0^\infty (r\rho)^{-\frac{n-2}{2}} J_\mu(r\rho) f(r, y) r^{n-1} dr,$$

where the Bessel function of order μ is given by

$$(2.8) \quad J_\mu(r) = \frac{(r/2)^\mu}{\Gamma(\mu + \frac{1}{2}) \Gamma(1/2)} \int_{-1}^1 e^{isr} (1 - s^2)^{(2\mu-1)/2} ds, \quad \mu > -1/2, r > 0,$$

which satisfies the following equation

$$r^2 \frac{d^2}{dr^2} (J_\mu(r)) + r \frac{d}{dr} (J_\mu(r)) + (r^2 - \mu^2) J_\mu(r) = 0.$$

Following the [24, (8.45)], for well-behaved functions F , we have by the functional calculus

$$(2.9) \quad F(\mathcal{L}_V) f(r, y) = \int_0^\infty \int_0^{2\pi} K(r, y, s, y') f(s, y') s^{n-1} ds dh(y')$$

the kernel

$$K(r, y, s, y') = (rs)^{-\frac{n-2}{2}} \sum_{k \in \mathbb{N}} \varphi_k(y) \overline{\varphi_k(y')} K_{\mu_k}(r, s) = (rs)^{-\frac{n-2}{2}} \sum_{k \in \mathbb{N}} H_k(y, y') K_{\mu_k}(r, s),$$

and

$$(2.10) \quad K_{\mu_k}(r, s) = \int_0^\infty F(\rho^2) J_{\mu_k}(r\rho) J_{\mu_k}(s\rho) \rho d\rho.$$

In particular, $F(\rho^2) = e^{-t\rho^2}$, by using the Weber's second exponential integral [26, Section 13.31 (1)], we obtain

$$(2.11) \quad K_{\mu_k}(r, s) = \int_0^\infty e^{-t\rho^2} J_{\mu_k}(r\rho) J_{\mu_k}(s\rho) \rho d\rho = (2t)^{-1} e^{-\frac{r^2+s^2}{4t}} I_\mu\left(\frac{rs}{2t}\right), \quad t > 0.$$

where $I_\mu(x)$ is the modified Bessel function of the first kind in series version

$$I_\mu(x) = \sum_{j=0}^{\infty} \frac{1}{j! \Gamma(\mu + j + 1)} (x/2)^{\mu+2j},$$

or in the integral representation

$$I_\mu(x) = \frac{1}{\sqrt{\pi} \Gamma(\mu + \frac{1}{2})} \left(\frac{x}{2}\right)^\mu \int_{-1}^1 e^{-x\tau} (1 - \tau^2)^{\mu-\frac{1}{2}} d\tau.$$

Therefore we obtain (1.5)

$$e^{-t\mathcal{L}_V}(r, y; s, y') = (2t)^{-1} e^{-\frac{r^2+s^2}{4t}} (rs)^{-\frac{n-2}{2}} \sum_{k \in \mathbb{N}} \varphi_k(y) \overline{\varphi_k(y')} I_{\mu_k} \left(\frac{rs}{2t} \right).$$

3. THE PROOF OF THE UPPER BOUNDEDNESS

In this section, we prove the upper boundedness (1.6). To this end, by observing (1.7) and scaling (r, s) , it suffices to show

$$(3.1) \quad \left| e^{-\frac{r^2+s^2}{4}} (rs)^{-\frac{n-2}{2}} \sum_{k \in \mathbb{N}} \varphi_k(y) \overline{\varphi_k(y')} I_{\mu_k} \left(\frac{rs}{2} \right) \right| \leq C \left[\min \left\{ 1, \left(\frac{rs}{2} \right) \right\} \right]^{-\sigma} e^{-\frac{d^2((r,y),(s,y'))}{c}},$$

which is the consequence of the following lemma

Lemma 3.1. *Let $z = \frac{1}{2}rs$, $\sigma = \frac{n-2}{2} - \mu_0$ and $\delta = d_h(y, y')$, then there exist constants C and N only depending on n such that*

- either for $0 < z \leq 1$,

$$(3.2) \quad \left| z^{-\frac{n-2}{2}} \sum_{k \in \mathbb{N}} \varphi_k(y) \overline{\varphi_k(y')} I_{\mu_k}(z) \right| \leq C z^{-\sigma} \times \begin{cases} e^{z \cos \delta}, & 0 \leq \delta \leq \pi, \\ 1 & \pi \leq \delta, \end{cases}$$

- or for $z \gtrsim 1$

$$(3.3) \quad \left| z^{-\frac{n-2}{2}} \sum_{k \in \mathbb{N}} \varphi_k(y) \overline{\varphi_k(y')} I_{\mu_k}(z) \right| \leq C \times \begin{cases} e^{z \cos \delta} + z^N e^{z \cos(\frac{1}{2}\epsilon_0)}, & 0 \leq \delta \leq \frac{1}{2}\epsilon_0, \\ z^N e^{z \cos \delta}, & \frac{1}{2}\epsilon_0 \leq \delta \leq \pi, \\ e^{\frac{z}{2}} & \pi \leq \delta, \end{cases}$$

where ϵ_0 is the injective radius of the manifold Y with $0 < \epsilon_0 \leq \pi$.

Remark 3.1. *If the injective radius ϵ_0 of the manifold Y is larger than π , we do not need to introduce the cutoff function χ in the **Step 1** below, and we can follow the same argument to obtain, for $z \gtrsim 1$*

$$(3.4) \quad \left| z^{-\frac{n-2}{2}} \sum_{k \in \mathbb{N}} \varphi_k(y) \overline{\varphi_k(y')} I_{\mu_k}(z) \right| \leq C \times \begin{cases} e^{z \cos \delta}, & 0 \leq \delta \leq \pi, \\ e^{\frac{z}{2}} & \pi \leq \delta, \end{cases}$$

which is better than (3.3). Hence we omit this easier case in the following argument.

Now we assume Lemma 3.1 to prove (3.1). We divide into two cases $z \leq 1$ and $z \geq 1$ where $z = \frac{rs}{2}$. When $z \leq 1$, we recall the distance (1.7) and put (3.2) into the left hand side of (3.1) to obtain

$$(3.5) \quad \left| e^{-\frac{r^2+s^2}{4}} (rs)^{-\frac{n-2}{2}} \sum_{k \in \mathbb{N}} \varphi_k(y) \overline{\varphi_k(y')} I_{\mu_k} \left(\frac{rs}{2} \right) \right| \leq C \left(\frac{rs}{2} \right)^{-\sigma} e^{-\frac{d^2((r,y),(s,y'))}{c}}.$$

Next we consider the case that $z \geq 1$. In the subcase that $0 \leq \delta \leq \frac{1}{2}\epsilon_0$ of (3.3), we have

$$\begin{aligned} \text{LHS of (3.1)} &\lesssim e^{-\frac{r^2-2rs\cos\delta+s^2}{4}} + z^N e^{-\frac{r^2-2rs\cos(\frac{1}{2}\epsilon_0)+s^2}{4}} \\ &\lesssim e^{-\frac{r^2-2rs\cos\delta+s^2}{8}} \left(1 + z^N e^{-\frac{z}{4}(1-\cos(\frac{1}{2}\epsilon_0))}\right) \lesssim e^{-\frac{d^2((r,y),(s,y'))}{c}}. \end{aligned}$$

In the subcase that $\frac{1}{2}\epsilon_0 \leq \delta \leq \pi$ of (3.3), we have

$$\text{LHS of (3.1)} \lesssim z^N e^{-\frac{r^2-2rs\cos\delta+s^2}{4}} \lesssim z^N e^{-\frac{z}{4}(1-\cos\delta)} e^{-\frac{r^2-2rs\cos\delta+s^2}{8}}.$$

Since $\frac{1}{2}\epsilon_0 \leq \delta \leq \pi$, one has $1 - \cos\delta \geq \epsilon^2/4 > 0$. Hence, for $z \geq 1$, no matter how large N is, there exists a constant C independent of z such that

$$\text{LHS of (3.1)} \leq C e^{-\frac{r^2-2rs\cos\delta+s^2}{8}} \lesssim e^{-\frac{d^2((r,y),(s,y'))}{c}}.$$

In the last subcase $\delta \geq \pi$, we have

$$\text{LHS of (3.1)} \leq C e^{-\frac{r^2+s^2}{8}} e^{-\frac{r^2-2rs+s^2}{8}} \leq C e^{-\frac{(r+s)^2}{16}}.$$

Therefore we prove (3.1) once we could prove (3.2) and (3.3), which are our main tasks from now on. To this end, we first claim that

$$(3.6) \quad |H_k(y, y')| \leq \|\varphi_k(y)\|_{L^\infty(Y)}^2 \leq C(1 + \lambda_k)^{\frac{n-2}{2}} \leq C(1 + k)^{\frac{n-2}{n-1}},$$

where we used the eigenfunction estimate (see [23, (3.2.5)-(3.2.6)]) and the Weyl's asymptotic formula (e.g. see [27])

$$(3.7) \quad \lambda_k \sim (1 + k)^{\frac{2}{n-1}}, \quad k \geq 1, \implies \mu_k \sim (1 + k)^{\frac{1}{n-1}}.$$

3.1. The proof of (3.2). For the case $0 \leq \delta \leq \pi$, we first notice that $e^z \lesssim e^{z \cos \delta}$ if $0 \leq z \leq 1$, then the modified Bessel function satisfies

$$(3.8) \quad |I_\mu(z)| \leq \sqrt{\pi} e^z \frac{(\frac{z}{2})^\mu}{\Gamma(\mu + \frac{1}{2})} \lesssim e^{z \cos \delta} \frac{z^{\mu_0}}{2^\mu \Gamma(\mu + \frac{1}{2})}, \quad \mu \geq \mu_0.$$

Therefore, from (3.6), we have

$$\text{LHS of (3.2)} \lesssim z^{-\frac{n-2}{2}} e^{z \cos \delta} \sum_{k \in \mathbb{N}} (1 + k)^{\frac{n-2}{n-1}} \frac{z^{\mu_0}}{2^{\mu_k} \Gamma(\mu_k + \frac{1}{2})}.$$

Note that $\mu_k \geq \mu_0$ and $\mu_k = \sqrt{\lambda_k} \sim (1 + k)^{\frac{1}{n-1}}$, then the summation converges. Hence we show

$$(3.9) \quad \text{LHS of (3.2)} \lesssim e^{z \cos \delta} z^{\mu_0 - \frac{n-2}{2}}, \quad 0 \leq \delta \leq \pi,$$

as desired. For the case that $\delta \geq \pi$, we replace (3.8) by

$$|I_\mu(z)| \leq \sqrt{\pi} e^z \frac{(\frac{z}{2})^{\mu_0}}{\Gamma(\mu + \frac{1}{2})} \lesssim \frac{z^{\mu_0}}{2^\mu \Gamma(\mu + \frac{1}{2})}, \quad \mu \geq \mu_0.$$

Therefore, as before the summation converges, we obtain

$$\text{LHS of (3.2)} \lesssim z^{\mu_0 - \frac{n-2}{2}}.$$

which complete the proof of (3.2).

3.2. The proof of (3.3). In this case, we need the integral representation (see [26], [21, p. 419] or [9, (10.32.4)]) of the modified Bessel function

$$(3.10) \quad I_\mu(z) = \frac{1}{\pi} \int_0^\pi e^{z \cos(\tau)} \cos(\mu\tau) d\tau - \frac{\sin(\mu\pi)}{\pi} \int_0^\infty e^{-z \cosh \tau} e^{-\tau\mu} d\tau.$$

We divide into three steps to prove (3.3).

Step 1: we consider the case $0 \leq \delta \leq \frac{1}{2}\epsilon_0$ which is the most difficult case. We first introduce a function $\chi \in C_c^\infty([0, \pi])$ such that

$$(3.11) \quad \chi(\tau) = \begin{cases} 1, & \tau \in [0, \frac{1}{2}\epsilon_0], \\ 0, & \tau \in [\epsilon_0, \pi]. \end{cases}$$

Lemma 3.2. For fixed $0 \leq \delta \leq \pi$ and $m \geq 0$, let χ be in (3.11), we have

$$(3.12) \quad \begin{aligned} & \frac{1}{\pi} \int_0^\pi e^{z(\cos \tau - \cos \delta)} \cos(\mu\tau) d\tau - \frac{\sin(\mu\pi)}{\pi} \int_0^\infty e^{-z(\cosh \tau + \cos \delta)} e^{-\tau\mu} d\tau \\ &= \frac{1}{\pi} \int_0^\pi e^{z(\cos \tau - \cos \delta)} \chi(\tau) \cos(\mu\tau) d\tau \\ &+ \frac{(-1)^m}{\pi} \int_0^\pi \left(\frac{\partial}{\partial \tau}\right)^{2m} (e^{z(\cos \tau - \cos \delta)} (1 - \chi(\tau))) \frac{\cos(\mu\tau)}{\mu^{2m}} d\tau \\ &- \frac{\sin(\mu\pi)}{\pi} \int_0^\infty \left(\frac{\partial}{\partial \tau}\right)^{2m} (e^{-z(\cosh \tau + \cos \delta)}) \frac{e^{-\tau\mu}}{\mu^{2m}} d\tau. \end{aligned}$$

Proof. This is a variant of [21, (5.30)]. We prove this lemma by using induction argument and the argument of [21]. We first verify $m = 1$. By integration by parts, we have

$$\begin{aligned} & \frac{1}{\pi} \int_0^\pi e^{z(\cos \tau - \cos \delta)} \cos(\mu\tau) d\tau - \frac{\sin(\mu\pi)}{\pi} \int_0^\infty e^{-z(\cosh \tau + \cos \delta)} e^{-\tau\mu} d\tau \\ &= \frac{1}{\pi} \int_0^\pi e^{z(\cos \tau - \cos \delta)} \chi(\tau) \cos(\mu\tau) d\tau + \frac{1}{\pi} (e^{z(\cos \tau - \cos \delta)} (1 - \chi(\tau))) \frac{\sin(\mu\tau)}{\mu} \Big|_{\tau=0}^{\tau=\pi} \\ &+ \frac{(-1)}{\pi} \int_0^\pi \left(\frac{\partial}{\partial \tau}\right) (e^{z(\cos \tau - \cos \delta)} (1 - \chi(\tau))) \frac{\sin(\mu\tau)}{\mu} d\tau \\ &+ \frac{\sin(\mu\pi)}{\pi} (e^{-z(\cosh \tau + \cos \delta)}) \frac{e^{-\tau\mu}}{\mu} \Big|_{\tau=0}^\infty - \frac{\sin(\mu\pi)}{\pi} \int_0^\infty \left(\frac{\partial}{\partial \tau}\right) (e^{-z(\cosh \tau + \cos \delta)}) \frac{e^{-\tau\mu}}{\mu} d\tau. \end{aligned}$$

Note the boundary term

$$\begin{aligned} & \frac{1}{\pi} (e^{z(\cos \tau - \cos \delta)} (1 - \chi(\tau))) \frac{\sin(\mu\tau)}{\mu} \Big|_{\tau=0}^{\tau=\pi} + \frac{\sin(\mu\pi)}{\pi} (e^{-z(\cosh \tau + \cos \delta)}) \frac{e^{-\tau\mu}}{\mu} \Big|_{\tau=0}^\infty \\ &= \frac{1}{\pi} e^{z(\cos \tau - \cos \delta)} \frac{\sin(\mu\tau)}{\mu} \Big|_{\tau=\pi} - \frac{\sin(\mu\pi)}{\pi} (e^{-z(\cosh \tau + \cos \delta)}) \frac{e^{-\tau\mu}}{\mu} \Big|_{\tau=0} = 0. \end{aligned}$$

By integration by parts again, we have

$$\begin{aligned}
& \frac{1}{\pi} \int_0^\pi e^{z(\cos \tau - \cos \delta)} \cos(\mu \tau) d\tau - \frac{\sin(\mu \pi)}{\pi} \int_0^\infty e^{-z(\cosh \tau + \cos \delta)} e^{-\tau \mu} d\tau \\
&= \frac{1}{\pi} \int_0^\pi e^{z(\cos \tau - \cos \delta)} \chi(\tau) \cos(\mu \tau) d\tau + \frac{1}{\pi} \left(\frac{\partial}{\partial \tau} \right) \left(e^{z(\cos \tau - \cos \delta)} (1 - \chi(\tau)) \right) \frac{\cos(\mu \tau)}{\mu^2} \Big|_{\tau=0}^{\tau=\pi} \\
&\quad + \frac{(-1)}{\pi} \int_0^\pi \left(\frac{\partial}{\partial \tau} \right)^2 \left(e^{z(\cos \tau - \cos \delta)} (1 - \chi(\tau)) \right) \frac{\cos(\mu \tau)}{\mu^2} d\tau \\
&\quad + \frac{\sin(\mu \pi)}{\pi} \left(\frac{\partial}{\partial \tau} \right) \left(e^{-z(\cosh \tau + \cos \delta)} \right) \frac{e^{-\tau \mu}}{\mu^2} \Big|_{\tau=0}^\infty - \frac{\sin(\mu \pi)}{\pi} \int_0^\infty \left(\frac{\partial}{\partial \tau} \right) \left(e^{-z(\cosh \tau + \cos \delta)} \right) \frac{e^{-\tau \mu}}{\mu^2} d\tau.
\end{aligned}$$

Again we observe that the boundary term

$$\begin{aligned}
& \frac{1}{\pi} \left(\frac{\partial}{\partial \tau} \right) \left(e^{z(\cos \tau - \cos \delta)} (1 - \chi(\tau)) \right) \frac{\cos(\mu \tau)}{\mu^2} \Big|_{\tau=0}^{\tau=\pi} + \frac{\sin(\mu \pi)}{\pi} \left(\frac{\partial}{\partial \tau} \right) \left(e^{-z(\cosh \tau + \cos \delta)} \right) \frac{e^{-\tau \mu}}{\mu^2} \Big|_{\tau=0}^\infty \\
&= \frac{1}{\pi} \left(\frac{\partial}{\partial \tau} \right) \left(e^{z(\cos \tau - \cos \delta)} (1 - \chi(\tau)) \right) \frac{\cos(\mu \tau)}{\mu^2} \Big|_{\tau=\pi} - \frac{\sin(\mu \pi)}{\pi} \left(\frac{\partial}{\partial \tau} \right) \left(e^{-z(\cosh \tau + \cos \delta)} \right) \frac{e^{-\tau \mu}}{\mu^2} \Big|_{\tau=0}^\infty
\end{aligned}$$

vanishes due to the fact $\sin \pi = \sinh 0 = 0$. Therefore, we have proved (3.12) with $m = 1$. Now we assume (3.12) holds for $m = k$, that is,

$$\begin{aligned}
& \frac{1}{\pi} \int_0^\pi e^{z(\cos \tau - \cos \delta)} \cos(\mu \tau) d\tau - \frac{\sin(\mu \pi)}{\pi} \int_0^\infty e^{-z(\cosh \tau + \cos \delta)} e^{-\tau \mu} d\tau \\
&= \frac{1}{\pi} \int_0^\pi e^{z(\cos \tau - \cos \delta)} \chi(\tau) \cos(\mu \tau) d\tau \\
&\quad + \frac{(-1)^k}{\pi} \int_0^\pi \left(\frac{\partial}{\partial \tau} \right)^{2k} \left(e^{z(\cos \tau - \cos \delta)} (1 - \chi(\tau)) \right) \frac{\cos(\mu \tau)}{\mu^{2k}} d\tau \\
&\quad - \frac{\sin(\mu \pi)}{\pi} \int_0^\infty \left(\frac{\partial}{\partial \tau} \right)^{2k} \left(e^{-z(\cosh \tau + \cos \delta)} \right) \frac{e^{-\tau \mu}}{\mu^{2k}} d\tau,
\end{aligned}$$

we aim to prove (3.12) when $m = k + 1$. To this end, it suffices to check the boundary terms vanish. Indeed,

$$\begin{aligned}
& \frac{(-1)^k}{\pi} \left(\frac{\partial}{\partial \tau} \right)^{2k} \left(e^{z(\cos \tau - \cos \delta)} (1 - \chi(\tau)) \right) \frac{\sin(\mu \tau)}{\mu^{2k+1}} \Big|_{\tau=0}^{\tau=\pi} \\
&\quad + \frac{\sin(\mu \pi)}{\pi} \left(\frac{\partial}{\partial \tau} \right)^{2k} \left(e^{-z(\cosh \tau + \cos \delta)} \right) \frac{e^{-\tau \mu}}{\mu^{2k+1}} \Big|_{\tau=0}^\infty \\
&= \frac{(-1)^k}{\pi} \left(\frac{\partial}{\partial \tau} \right)^{2k} \left(e^{z(\cos \tau - \cos \delta)} (1 - \chi(\tau)) \right) \frac{\sin(\mu \tau)}{\mu^{2k+1}} \Big|_{\tau=\pi} \\
&\quad - \frac{\sin(\mu \pi)}{\pi} \left(\frac{\partial}{\partial \tau} \right)^{2k} \left(e^{-z(\cosh \tau + \cos \delta)} \right) \frac{e^{-\tau \mu}}{\mu^{2k+1}} \Big|_{\tau=0}^\infty = 0,
\end{aligned}$$

and

$$\begin{aligned}
& \frac{(-1)^{k+1}}{\pi} \left(\frac{\partial}{\partial \tau} \right)^{2k+1} \left(e^{z(\cos \tau - \cos \delta)} (1 - \chi(\tau)) \right) \frac{\cos(\mu \tau)}{\mu^{2k+2}} \Big|_{\tau=0}^{\tau=\pi} \\
& + \frac{\sin(\mu \pi)}{\pi} \left(\frac{\partial}{\partial \tau} \right)^{2k+1} \left(e^{-z(\cosh \tau + \cos \delta)} \right) \frac{e^{-\tau \mu}}{\mu^{2k+2}} \Big|_{\tau=0}^{\infty} \\
& = \frac{(-1)^{k+1}}{\pi} \left(\frac{\partial}{\partial \tau} \right)^{2k+1} \left(e^{z(\cos \tau - \cos \delta)} (1 - \chi(\tau)) \right) \frac{\cos(\mu \tau)}{\mu^{2k+2}} \Big|_{\tau=\pi} \\
& \quad - \frac{\sin(\mu \pi)}{\pi} \left(\frac{\partial}{\partial \tau} \right)^{2k+1} \left(e^{-z(\cosh \tau + \cos \delta)} \right) \frac{e^{-\tau \mu}}{\mu^{2k+2}} \Big|_{\tau=0} = 0,
\end{aligned}$$

where we use the facts derived from [21, Pag. 420]

$$(-1)^{k+1} \left(\frac{\partial}{\partial \tau} \right)^{2k} \left(e^{-z(\cos \delta - \cos \tau)} (1 - \chi(\tau)) \right) \Big|_{\tau=\pi} = \left(\frac{\partial}{\partial \tau} \right)^{2k} \left(e^{-z(\cos \delta + \cosh \tau)} \right) \Big|_{\tau=0},$$

and

$$(-1)^{k+1} \left(\frac{\partial}{\partial \tau} \right)^{2k+1} \left(e^{-z(\cos \delta - \cos \tau)} (1 - \chi(\tau)) \right) \Big|_{\tau=\pi} = \left(\frac{\partial}{\partial \tau} \right)^{2k+1} \left(e^{-z(\cos \delta + \cosh \tau)} \right) \Big|_{\tau=0} = 0.$$

□

Let $P = \sqrt{-\Delta_h + V_0(y) + \frac{(n-2)^2}{4}}$, then the left hand side of (3.3) can be regarded as the operator

$$\begin{aligned}
(3.13) \quad & e^{z \cos \delta} z^{-\frac{n-2}{2}} \left(\frac{1}{\pi} \int_0^\pi e^{z(\cos \tau - \cos \delta)} \chi(\tau) \cos(\tau P) d\tau \right. \\
& + \frac{(-1)^m}{\pi} \int_0^\pi \left(\frac{\partial}{\partial \tau} \right)^{2m} \left(e^{z(\cos \tau - \cos \delta)} (1 - \chi(\tau)) \right) \frac{\cos(\tau P)}{P^{2m}} d\tau \\
& \left. - \frac{\sin(\pi P)}{\pi} \int_0^\infty \left(\frac{\partial}{\partial \tau} \right)^{2m} \left(e^{-z(\cosh \tau + \cos \delta)} \right) \frac{e^{-\tau P}}{P^{2m}} d\tau \right).
\end{aligned}$$

The $\cos(tP)f$ in the first term is the unique solutions of wave equation

$$(3.14) \quad \begin{cases} (\partial_t^2 + (-\Delta_h + V_0(y) + \frac{(n-2)^2}{4}))u = 0, \\ u|_{t=0} = f, \quad \partial_t u|_{t=0} = 0. \end{cases}$$

By the finite speed of propagation [4, Theorem 3.3], $\cos(\tau P)(y, y')$ vanishes if $\tau < d_h(y, y')$ where d_h denotes the distance in Y , then (3.13) equals

$$\begin{aligned}
(3.15) \quad & e^{z \cos \delta} z^{-\frac{n-2}{2}} \left(\frac{1}{\pi} \int_\delta^\pi e^{z(\cos \tau - \cos \delta)} \chi(\tau) \cos(\tau P) d\tau \right. \\
& + \frac{(-1)^m}{\pi} \int_\delta^\pi \left(\frac{\partial}{\partial \tau} \right)^{2m} \left(e^{z(\cos \tau - \cos \delta)} (1 - \chi(\tau)) \right) \frac{\cos(\tau P)}{P^{2m}} d\tau \\
& \left. - \frac{\sin(\pi P)}{\pi} \int_0^\infty \left(\frac{\partial}{\partial \tau} \right)^{2m} \left(e^{-z(\cosh \tau + \cos \delta)} \right) \frac{e^{-\tau P}}{P^{2m}} d\tau \right).
\end{aligned}$$

Then the first case of (3.3) is a consequence of the following lemma.

Lemma 3.3. For $0 \leq \delta \leq \frac{1}{2}\epsilon_0$ and $z \geq 1$, there exists a constant N such that

$$(3.16) \quad \left| z^{-\frac{n-2}{2}} \int_{\delta}^{\pi} e^{z(\cos \tau - \cos \delta)} \chi(\tau) \cos(\tau P)(y, y') d\tau \right| \lesssim 1$$

and

$$(3.17) \quad \left| z^{-\frac{n-2}{2}} \sum_{k \in \mathbb{N}} \varphi_k(y) \overline{\varphi_k(y')} \mu_k^{-2m} \left(\int_{\delta}^{\pi} \left(\frac{\partial}{\partial \tau} \right)^{2m} (e^{z(\cos \tau - \cos \delta)} (1 - \chi(\tau))) \cos(\tau \mu_k) d\tau \right. \right. \\ \left. \left. - \sin(\pi \mu_k) \int_0^{\infty} \left(\frac{\partial}{\partial \tau} \right)^{2m} (e^{-z(\cosh \tau + \cos \delta)}) e^{-\tau \mu_k} d\tau \right) \right| \lesssim z^N e^{z(\cos(\frac{1}{2}\epsilon_0) - \cos \delta)}.$$

Proof. We first prove (3.17). Notice that $z \geq 1$, by the chain rule, we obtain

$$(3.18) \quad \left| \left(\frac{\partial}{\partial \tau} \right)^{2m} (e^{-z(\cos \delta - \cos \tau)} (1 - \chi(\tau))) \right| \leq C e^{-z(\cos \delta - \cos \tau)} z^{2m},$$

and

$$(3.19) \quad \left| \left(\frac{\partial}{\partial \tau} \right)^{2m} (e^{-z(\cos \delta + \cosh \tau)}) \right| \leq C e^{-z(\cos \delta + \frac{1}{2} \cosh \tau)} z^{2m}.$$

Therefore, by (3.6) and the support of $1 - \chi$, we show that the LHS of (3.17) is bounded by

$$z^{2m - \frac{n-2}{2}} \sum_{k \in \mathbb{N}} (1+k)^{\frac{n-2}{n-1}} \mu_k^{-2m} \left(\int_{\frac{\epsilon_0}{2}}^{\pi} e^{-z(\cos \delta - \cos \tau)} d\tau + e^{-z \cos \delta} \int_0^{\infty} e^{-\frac{z}{2} \cosh \tau} d\tau \right) \\ \lesssim z^{2m - \frac{n-2}{2}} e^{z(\cos(\frac{1}{2}\epsilon_0) - \cos \delta)} \sum_{k \in \mathbb{N}} (1+k)^{\frac{n-2}{n-1}} \mu_k^{-2m}.$$

Note $\mu_k = \sqrt{\lambda_k} \sim (1+k)^{\frac{1}{n-1}}$ again, we choose m large enough to ensure that

$$\sum_{k \in \mathbb{N}} (1+k)^{\frac{n-2}{n-1}} \mu_k^{-2m} \lesssim 1.$$

Therefore, we obtain (3.17) by choosing $N = 2n - 3$.

We next prove (3.16). Due to the compact support of χ , for $|\tau| \leq \epsilon_0$, it suffices to prove

$$(3.20) \quad \left| z^{-\frac{n-2}{2}} \int_{\delta}^{\epsilon_0} e^{z(\cos \tau - \cos \delta)} \chi(\tau) \cos(\tau P)(y, y') d\tau \right| \lesssim 1.$$

By using the Hadamard parametrix (e.g. [23, Theorem 3.1.5]), for $\tau < \epsilon_0$, we have

$$(3.21) \quad \cos(\tau P)(y, y') = K_N(\tau, y, y') + R_N(\tau; y, y'), \quad \forall N > n + 3$$

where $R_N(\tau, y, y') \in \mathcal{C}^{N-n-3}([0, \epsilon_0] \times Y \times Y)$ and

$$(3.22) \quad K_N(\tau, y, y') = (2\pi)^{n-1} \int_{\mathbb{R}^{n-1}} e^{id_h(y, y') \cdot \mathbf{1} \cdot \xi} a(\tau, y, y'; |\xi|) \cos(\tau |\xi|) d\xi$$

where $\mathbf{1} = (1, 0, \dots, 0)$ and $a \in S^0$ zero order symbol satisfies

$$(3.23) \quad |\partial_{\tau, y, y'}^{\alpha} \partial_{\rho}^k a(\tau, y, y'; \rho)| \leq C_{\alpha, k, V_0} (1 + \rho)^{-k}.$$

We remark here that even though $P = \sqrt{-\Delta_h + V_0(y) + \frac{(n-2)^2}{4}}$ is disturbed by a smoothing potential which is not exact same as the Laplacian on a compact manifold in [23, Theorem 3.1.5], the perturbation is harmless for the parametrix (3.21).

Actually, in the construction of Hadamard parametrix, the original transport equation for non-perturbed operator is [23, (2.4.15)-(2.4.16)], which are

$$\rho\alpha_0 = 2\langle x, \nabla_x \alpha_0 \rangle, \alpha_0(0) = 1$$

as well as $\alpha_\nu(x)$, $\nu = 1, 2, 3..$ so that

$$2\nu\alpha_\nu - \rho\alpha_\nu + 2\langle x, \nabla_x \alpha_\nu \rangle - 2\Delta_g \alpha_{\nu-1} = 0$$

In our case, the new transport equation for the perturbed operator would be

$$\rho\alpha_0 = 2\langle x, \nabla_x \alpha_0 \rangle, \alpha_0(0) = 1$$

as well as $\alpha_\nu(x)$, $\nu = 1, 2, 3..$ so that

$$2\nu\alpha_\nu - \rho\alpha_\nu + 2\langle x, \nabla_x \alpha_\nu \rangle - 2\Delta_g \alpha_{\nu-1} - 2V_0 \cdot \alpha_{\nu-1} = 0$$

Thus two solutions to the transport equations only differ in $\alpha_\nu(x)$, $\nu = 1, 2, 3..$, which is the reason why (3.23) is true. We also refer the reader to Hörmander [12, §17.4] for the parametrix of a general second order differential operator with low order perturbations.

We choose N large enough such that

$$|R_N(\tau, y, y)| \leq \tau^{2N+2-n} \leq 1, \quad \tau \leq \epsilon_0,$$

then it is easy to see

$$(3.24) \quad \left| z^{-\frac{n-2}{2}} \int_\delta^\pi e^{z(\cos \tau - \cos \delta)} \chi(\tau) R_N(\tau; y, y') d\tau \right| \leq C z^{-\frac{n-2}{2}} \lesssim 1.$$

Therefore it suffices to prove

$$(3.25) \quad \left| z^{-\frac{n-2}{2}} \int_\delta^\pi e^{z(\cos \tau - \cos \delta)} \chi(\tau) \int_{\mathbb{R}^{n-1}} e^{id_h(y, y') \mathbf{1} \cdot \xi} a(\tau, y, y'; |\xi|) \cos(\tau|\xi|) d\xi d\tau \right| \leq C.$$

From [22, Theorem 1.2.1], we also note that

$$(3.26) \quad \int_{\mathbb{S}^{n-2}} e^{id_h(y, y') \rho \mathbf{1} \cdot \omega} d\omega d\tau = \sum_{\pm} a_{\pm}(\rho d_h(y, y')) e^{\pm i \rho d_h(y, y')}$$

where

$$(3.27) \quad |\partial_r^k a_{\pm}(r)| \leq C_k (1+r)^{-\frac{n-2}{2}-k}, \quad k \geq 0.$$

Then we are reduce to estimate the integral

$$\begin{aligned} & z^{-\frac{n-2}{2}} \int_\delta^\pi e^{z(\cos \tau - \cos \delta)} \chi(\tau) \int_{\mathbb{R}^{n-1}} e^{id_h(y, y') \mathbf{1} \cdot \xi} a(\tau, y, y'; |\xi|) \cos(\tau|\xi|) d\xi d\tau \\ &= z^{-\frac{n-2}{2}} \int_\delta^\pi e^{z(\cos \tau - \cos \delta)} \chi(\tau) \int_0^\infty a_{\pm}(\rho d_h(y, y')) e^{\pm i \rho d_h(y, y')} a(\tau, y, y'; \rho) \cos(\tau\rho) \rho^{n-2} d\rho d\tau. \end{aligned}$$

To prove (3.20), we shall divide the discussion into two cases.

Case 1. $z \leq 2\delta^{-2} = 2d_h^{-2}(y, y')$. In this case, we do not need to make use of the finite propagation speed property, by the above argument using Hadamard parametrix, our goal is to show

$$z^{-\frac{n-2}{2}} \int_0^\pi e^{z(\cos \tau - \cos \delta)} \chi(\tau) \int_0^\infty a_\pm(\rho d_h(y, y')) e^{\pm i \rho d_h(y, y')} a(\tau, y, y'; \rho) \cos(\tau \rho) \rho^{n-2} d\rho d\tau \lesssim 1$$

Since the above integral is even in τ , we can further reduced to showing that

$$(3.28) \quad z^{-\frac{n-2}{2}} \int_{\mathbb{R}} e^{z(\cos \tau - \cos \delta)} \chi(\tau) \int_0^\infty b(\rho d_h(y, y')) a(\tau, y, y'; \rho) \cos(\tau \rho) \rho^{n-2} d\rho d\tau \lesssim 1,$$

where we set $b(\rho d_h(y, y')) = a_\pm(\rho d_h(y, y')) e^{\pm i \rho d_h(y, y')}$.

Let us fix a Littlewood-Paley bump function $\beta \in C_0^\infty((1/2, 2))$ satisfying

$$\sum_{\ell=-\infty}^{\infty} \beta(2^{-\ell} s) = 1, \quad s > 0,$$

and we set

$$\beta_0(s) = \sum_{\ell \leq 0} \beta(2^{-\ell} |s|) \in C_0^\infty((-2, 2)).$$

Subcase 1.1 the sub-case that $|\tau| \leq 4z^{-1/2}$. In this case, we want to show that

$$(3.29) \quad z^{-\frac{n-2}{2}} \int_{\mathbb{R}} e^{z(\cos \tau - \cos \delta)} \chi(\tau) (\beta_0(z^{1/2} \tau) + \beta(2^{-1} z^{1/2} \tau)) \\ \times \int_0^\infty b(\rho d_h(y, y')) a(\tau, y, y'; \rho) \cos(\tau \rho) \rho^{n-2} d\rho d\tau \lesssim 1.$$

If we also have $\rho \leq 4z^{1/2}$, then we do not do any integration by parts, note that by using the condition $z \leq 2\delta^{-2}$, the power on the exponential $z(\cos \tau - \cos \delta) \lesssim 1$ as long as $|\tau| \leq 4z^{-1/2}$. Thus the integral in (3.29) is always bounded by

$$z^{-\frac{n-2}{2}} z^{-1/2} z^{\frac{n-1}{2}} \lesssim 1$$

If on the other hand, we have $\rho \geq 4z^{1/2}$, we do integration by parts in $d\tau$, then each time we gain a factor of ρ^{-1} from the function $\cos(\tau \rho)$, and we at most loss a factor of $z \sin \tau$ or $z^{1/2}$, which is always less than $z^{1/2}$ up to a constant, so after integration by parts N times for $N \geq n$, the integral in (3.29) is bounded by

$$z^{-\frac{n-2}{2}} z^{-1/2} z^{N/2} \int_{z^{1/2}}^\infty \rho^{n-2-N} d\rho \lesssim 1$$

Subcase 1.2 $\tau \approx 2^j z^{-1/2}$, $j \geq 2$ and $2^j \lesssim \epsilon_0 z^{1/2}$. In this case, we want to show that

$$(3.30) \quad z^{-\frac{n-2}{2}} \int_{-\infty}^{+\infty} e^{z(\cos \tau - \cos \delta)} \chi(\tau) \beta(z^{1/2} 2^{-j} |\tau|) \\ \times \int_0^\infty b(\rho d_h(y, y')) a(\tau, y, y'; \rho) \cos(\tau \rho) \rho^{n-2} d\rho d\tau \lesssim e^{-j}.$$

which would give us desired bounds after summing over j . Since $\tau \approx 2^j z^{-1/2}$ and $z \leq 2d_h^{-2}(y, y')$ imply that $|\tau| \geq 2d_h(y, y')$, it is straightforward to check that $\cos \tau - \cos \delta \approx -\tau^2$ in this case, which implies

$$e^{z(\cos \tau - \cos \delta)} \lesssim e^{-2^j}.$$

Now we can repeat the previous argument, if in this case we have $\rho \leq 2^{-j} z^{1/2}$, then we do not do any integration by parts, the integral in (3.30) is always bounded by

$$z^{-\frac{n-2}{2}} z^{-1/2} 2^j 2^{-(n-1)j} z^{\frac{n-1}{2}} e^{-2^j} \lesssim e^{-j}$$

If on the other hand, we have $\rho \geq 2^{-j} z^{1/2}$, we do integration by parts in $d\tau$, then each time we gain a factor of ρ^{-1} from the function $\cos(\tau\rho)$, and we at most loss a factor of $z \sin \tau \lesssim 2^j z^{1/2}$, so after integration by parts N times for $N \geq n$, the integral in (3.30) is bounded by

$$e^{-2^j} z^{-\frac{n-2}{2}} z^{-1/2} 2^j z^{Nz/2} 2^{Nj} \int_{2^{-j} z^{1/2}}^{\infty} \rho^{n-2-N} d\rho \lesssim e^{-j}$$

Case 2. $z \geq 2\delta^{-2} = 2d_h^{-2}(y, y')$. In this case, we need to make use of the finite propagation speed property, in order to avoid the blow up of the function $e^{z(\cos \tau - \cos \delta)}$ when τ is close to 0. We choose a smooth cut off function $\eta(\tau) = \beta_0(z\delta|\tau - \delta|) + \sum_{\ell \geq 1} \beta(z\delta 2^{-\ell}(\tau - \delta))$. It is easy to see that $\text{supp } \eta \subset (\delta - 2(z\delta)^{-1}, +\infty)$ which is a subset of $(0, \infty)$ since $(z\delta)^{-1} < \delta/2$ in our case. Also note that $\eta(\tau) \equiv 1, \forall \tau \geq \delta$, by using the finite propagation speed property of $\cos(\tau P)$ and Hadamard parametrix as above, we are reduced to show that

$$z^{-\frac{n-2}{2}} \int_0^\pi e^{z(\cos \tau - \cos \delta)} \chi(\tau) \eta(\tau) \int_0^\infty b(\rho d_h(y, y')) a(\tau, y, y'; \rho) \cos(\tau\rho) \rho^{n-2} d\rho d\tau \lesssim 1$$

Subcase 2.1 $|\tau - \delta| \leq (z\delta)^{-1} < \frac{\delta}{2}$. In this case, we want to show that

$$(3.31) \quad z^{-\frac{n-2}{2}} \int_0^\pi e^{z(\cos \tau - \cos \delta)} \chi(\tau) \beta_0(z\delta|\tau - \delta|) \\ \times \int_0^\infty b(\rho d_h(y, y')) a(\tau, y, y'; \rho) \cos(\tau\rho) \rho^{n-2} d\rho d\tau \lesssim 1.$$

If we also have $\rho \leq z\delta$, then we do not do any integration by parts, note that since in this case $\tau \approx \delta$, we have $\cos \tau - \cos \delta = 2 \sin(\frac{\delta+\tau}{2}) \sin(\frac{\delta-\tau}{2}) \approx \delta(\delta - \tau)$, the power on the exponential $z(\cos \tau - \cos \delta) \lesssim 1$ as long as $|\tau - \delta| \leq (z\delta)^{-1}$. Thus the integral in (3.31) is always bounded by

$$z^{-\frac{n-2}{2}} (z\delta)^{-1} (z\delta)^{\frac{n-2}{2}+1} \delta^{-\frac{n-2}{2}} \lesssim 1,$$

where we used the fact that $b(\rho d_h(y, y')) \leq C(\rho\delta)^{-\frac{n-2}{2}}$.

If on the other hand, we have $\rho \geq z\delta$, we do integration by parts in $d\tau$, then each time we gain a factor of ρ^{-1} from the function $\cos(\tau\rho)$, and we at most loss a factor of $z \sin \tau \lesssim z\delta$, so after integration by parts N times for $N \geq n$, the integral in (3.31) is bounded by

$$z^{-\frac{n-2}{2}} (z\delta)^{-1} (z\delta)^{N/2} \int_{z\delta}^\infty \rho^{\frac{n-2}{2}-N} \delta^{-\frac{n-2}{2}} d\rho \lesssim 1$$

Subcase 2.2 $\tau - \delta \approx 2^j(z\delta)^{-1}$, $j \geq 1$ and $2^j \lesssim \epsilon_0 z \delta$. In this case, we want to show that

$$(3.32) \quad z^{-\frac{n-2}{2}} \int_{-\infty}^{+\infty} e^{z(\cos \tau - \cos \delta)} \chi(\tau) \beta(z\delta 2^{-j}(\tau - \delta)) \\ \times \int_0^\infty b(\rho d_h(y, y')) a(\tau, y, y'; \rho) \cos(\tau \rho) \rho^{n-2} d\rho d\tau \lesssim e^{-j}.$$

which would give us desired bounds after summing over j . In this case, it is straightforward to check that $z(\cos \tau - \cos \delta) = 2z \sin(\frac{\delta+\tau}{2}) \sin(\frac{\delta-\tau}{2}) \lesssim -2^j$ in this case, which implies

$$e^{z(\cos \tau - \cos \delta)} \lesssim e^{-2^j}.$$

Now we can repeat the previous argument. To begin with, we shall further assume that

(i) $\tau \leq 2\delta$, in other words $2^j(z\delta)^{-1} \lesssim \delta$.

In this case, if we have $\rho \leq 2^{-j}z\delta$, then we do not do any integration by parts, the integral in (3.32) is always bounded by

$$z^{-\frac{n-2}{2}} 2^j (z\delta)^{-1} (2^{-j}z\delta)^{\frac{n-2}{2}+1} \delta^{-\frac{n-2}{2}} e^{-2^j} \lesssim e^{-j}$$

On the other hand, if we have $\rho \geq 2^{-j}z\delta$, we do integration by parts in $d\tau$, then each time we gain a factor of ρ^{-1} from the function $\cos(\tau\rho)$, and we at most loss a factor of $z \sin \tau \lesssim z\delta$, so after integration by parts N times for $N \geq n$, the integral in (3.32) is bounded by

$$e^{-2^j} z^{-\frac{n-2}{2}} 2^j (z\delta)^{-1} (z\delta)^{N/2} \int_{2^{-j}z\delta}^\infty \rho^{\frac{n-2}{2}-N} \delta^{-\frac{n-2}{2}} d\rho \lesssim e^{-j}$$

(ii) $\tau \geq 2\delta$, in other words $2^j(z\delta)^{-1} \gtrsim \delta$.

In this case, if we have $\rho \leq \delta^{-1}2^j \approx z\tau$, then we do not do any integration by parts, the integral in (3.32) is always bounded by

$$z^{-\frac{n-2}{2}} 2^j (z\delta)^{-1} (\delta^{-1}2^j)^{\frac{n-2}{2}+1} \delta^{-\frac{n-2}{2}} e^{-2^j} \lesssim e^{-j},$$

where we used the fact that $z^{-n/2} \delta^{-n} \lesssim 1$.

On the other hand, if we have $\rho \geq \delta^{-1}2^j$, we do integration by parts in $d\tau$, then each time we gain a factor of ρ^{-1} from the function $\cos(\tau\rho)$, and we at most loss a factor of $z \sin \tau \lesssim \delta^{-1}2^j$ or $z\delta 2^{-j} \lesssim \delta^{-1}$, so after integration by parts N times for $N \geq n$, the integral in (3.32) is bounded by

$$e^{-2^j} z^{-\frac{n-2}{2}} 2^j (z\delta)^{-1} (\delta^{-1}2^j)^{N/2} \int_{2^j\delta^{-1}}^\infty \rho^{\frac{n-2}{2}-N} \delta^{-\frac{n-2}{2}} d\rho \lesssim e^{-j}.$$

□

Step 2: we consider the case $\frac{1}{2}\epsilon_0 \leq \delta \leq \pi$. In this case, it is more easier than the above step. Indeed, we do not need the cut function χ and we modify the above

argument. By integration by parts and arguing as Lemma 3.2, for fixed $0 \leq \delta \leq \pi$ and $m \geq 0$, we compute that

$$\begin{aligned}
(3.33) \quad e^{-z \cos \delta} I_\mu(z) &= \frac{1}{\pi} \int_0^\pi e^{z(\cos \tau - \cos \delta)} \cos(\mu \tau) d\tau - \frac{\sin(\mu \pi)}{\pi} \int_0^\infty e^{-z(\cosh \tau + \cos \delta)} e^{-\tau \mu} d\tau \\
&= \frac{(-1)^m}{\pi} \int_0^\pi \left(\frac{\partial}{\partial \tau} \right)^{2m-1} (e^{z(\cos \tau - \cos \delta)}) \frac{\sin(\mu \tau)}{\mu^{2m-1}} d\tau \\
&\quad - \frac{\sin(\mu \pi)}{\pi} \int_0^\infty \left(\frac{\partial}{\partial \tau} \right)^{2m-1} (e^{-z(\cosh \tau + \cos \delta)}) \frac{e^{-\tau \mu}}{\mu^{2m-1}} d\tau \\
&= \frac{(-1)^m}{\pi} \int_0^\pi \left(\frac{\partial}{\partial \tau} \right)^{2m} (e^{z(\cos \tau - \cos \delta)}) \frac{\cos(\mu \tau)}{\mu^{2m}} d\tau \\
&\quad - \frac{\sin(\mu \pi)}{\pi} \int_0^\infty \left(\frac{\partial}{\partial \tau} \right)^{2m} (e^{-z(\cosh \tau + \cos \delta)}) \frac{e^{-\tau \mu}}{\mu^{2m}} d\tau.
\end{aligned}$$

By the finite speed of propagation and similar argument as above, it suffices to prove

Lemma 3.4. *For $\frac{1}{2}\epsilon_0 \leq \delta \leq \pi$ and $z \geq 1$, there exists a constant N such that*

$$\begin{aligned}
(3.34) \quad & \left| z^{-\frac{n-2}{2}} \sum_{k \in \mathbb{N}} \varphi_k(y) \overline{\varphi_k(y')} \mu_k^{-2m} \left(\int_\delta^\pi \left(\frac{\partial}{\partial \tau} \right)^{2m} (e^{z(\cos \tau - \cos \delta)}) \cos(\tau \mu_k) d\tau \right. \right. \\
& \left. \left. - \sin(\pi \mu_k) \int_0^\infty \left(\frac{\partial}{\partial \tau} \right)^{2m} (e^{-z(\cosh \tau + \cos \delta)}) e^{-\tau \mu_k} d\tau \right) \right| \lesssim z^N.
\end{aligned}$$

Proof. Notice that $z \geq 1$, we further obtain

$$(3.35) \quad \left| \left(\frac{\partial}{\partial \tau} \right)^{2m} (e^{-z(\cos \delta - \cos \tau)}) \right| \leq C e^{-z(\cos \delta - \cos \tau)} z^{2m},$$

and

$$(3.36) \quad \left| \left(\frac{\partial}{\partial \tau} \right)^{2m} (e^{-z(\cos \delta + \cosh \tau)}) \right| \leq C e^{-z(\cos \delta + \frac{1}{2} \cosh \tau)} z^{2m}.$$

Therefore, by (3.6), we show that the LHS of (3.34) is bounded by

$$(3.37) \quad z^{2m - \frac{n-2}{2}} \sum_{k \in \mathbb{N}} (1+k)^{\frac{n-2}{n-1}} \mu_k^{-2m} \left(\int_\delta^\pi e^{-z(\cos \delta - \cos \tau)} d\tau + e^{-z \cos \delta} \int_0^\infty e^{-\frac{z}{2} \cosh \tau} d\tau \right).$$

Note $\mu_k = \sqrt{\lambda_k} \sim (1+k)^{\frac{1}{n-1}}$ again, we choose m large enough to ensure that

$$\sum_{k \in \mathbb{N}} (1+k)^{\frac{n-2}{n-1}} \mu_k^{-2m} \lesssim 1.$$

Therefore, we obtain (3.34) when $\frac{1}{2}\epsilon_0 \leq \delta \leq \pi$ by choosing $N = 2n - 3$. \square

Step 3: We consider the last case that $\delta > \pi$. Arguing as above, to prove (3.1), it suffices to prove

Lemma 3.5. *For $\pi \leq \delta$ and $z \geq 1$, we have the estimate*

$$(3.38) \quad \left| z^{-\frac{n-2}{2}} \sum_{k \in \mathbb{N}} \varphi_k(y) \overline{\varphi_k(y')} \mu_k^{-2m} \left(\int_0^\pi \left(\frac{\partial}{\partial \tau} \right)^{2m} (e^{z(\cos \tau - 1)}) \cos(\tau \mu_k) d\tau \right. \right. \\ \left. \left. - \sin(\pi \mu_k) \int_0^\infty \left(\frac{\partial}{\partial \tau} \right)^{2m} (e^{-z(\cosh \tau + 1)}) e^{-\tau \mu_k} d\tau \right) \right| \lesssim e^{-z}.$$

Proof. By the finite speed of propagation, the first term of (3.38) vanishes due to $\delta > \pi$. Arguing as before, it is easy to see

$$\text{LHS of (3.38)} \lesssim z^{-\frac{n-2}{2}} z^{2m} e^{-\frac{5}{4}z} \sum_{k \in \mathbb{N}} (1+k)^{\frac{n-2}{n-1}} \mu_k^{-2m} \int_0^\infty e^{-\frac{z}{4} \cosh \tau} d\tau \leq C_2 e^{-z}.$$

We choose m large enough to ensure the summation converges, and the factor z^{2m} can be absorbed by $e^{-\frac{z}{4}}$. \square

REFERENCES

- [1] N. BURQ, F. PLANCHON, J. STALKER AND A. S. TAHVILDAR-ZADEH, Strichartz estimates for the wave and Schrödinger equations with the inverse-square potential, *J. Funct. Anal.*, 203 (2003), 519-549. 4
- [2] G. CARRON, Le saut en zéro de la fonction de décalage spectral, *J. Funct. Anal.*, 212 (2004), 222-260. 1, 3
- [3] T. COUHON AND H. LI, Estimations inférieures du noyau de la chaleur sur les variétés coniques et transformée de Riesz, *Arch. Math.* 83(2004) 229-242. 1
- [4] T. COULHON AND A. SIKORA, Gaussian heat kernel upper bounds via the Phragmén-Lindelöf theorem. *Proc. Lond. Math. Soc.* (3), 96(2):507-544, 2008. 9
- [5] J. CHEEGER, On the spectral geometry of spaces with cone-like singularities, *Proceedings of the National Academy of Sciences of the United States of America*, 76(1979), 2103-2106. 1
- [6] J. CHEEGER, Spectral geometry of singular Riemannian spaces, *J. Diff. Geo.* 18(1983), 575-657. 1
- [7] J. CHEEGER AND M. TAYLOR, On the diffraction of waves by conical singularities, I, *Comm. Pure Appl. Math.*, 35(1982), 275-331. 1, 3
- [8] J. CHEEGER AND M. TAYLOR, On the diffraction of waves by conical singularities, II, *Comm. Pure Appl. Math.*, 35(1982), 487-529. 3
- [9] <https://dlmf.nist.gov/10.32> 7
- [10] X. GAO, J. ZHANG AND J. ZHENG, Restriction estimates in a conical singular space: wave equation, *J. Fourier Anal and Appl.* DOI : 10.1007/s00041-022-09941-7. 1
- [11] A. GRIGOR'YAN, Gaussian upper bounds for the heat kernel on arbitrary manifolds, *J. Differ. Geom.* 45(1997), 33-52. 2
- [12] L. HÖRMANDER, *The analysis of linear partial differential operators III: pseudo-differential operators.* Berlin-Heidelberg 1985. 11
- [13] A. HASSELL AND P. LIN, The Riesz transform for homogeneous Schrödinger operators on metric cones, *Rev. Mat. Iberoamericana*, 30(2014), 477-522. 1
- [14] R. KILLIP, C. MIAO, M. VISAN, J. ZHANG AND J. ZHENG, Sobolev spaces adapted to the Schrödinger operator with inverse-square potential, *Math. Z.* 288(2018), 1273-1298. 3
- [15] H. LI, L^p -estimates for the wave equation on manifolds with conical singularities, *Math. Z.* 272 (2012), 551-575.
- [16] H. LI, La transformation de Riesz sur les variétés coniques, *J. Funct. Anal.* 168 (1999), 145-238. 1
- [17] H. LI, Estimations du noyau de la chaleur sur les variétés coniques et ses applications, *Bull. Sci. math.* 124(2000) 365-384. 1, 2
- [18] V. LISKEVICH AND Z. SOBOL, Estimates of integral kernels for semigroups associated with second order elliptic operators with singular coefficients. *Potential Anal.* 18 (2003), 359-390. 2
- [19] P. D. MILMAN AND YU. A. SEMENOV, Global heat kernel bounds via desingularizing weights. *J. Funct. Anal.* 212 (2004), 373-398. 2

- [20] E. MOOER, Heat kernel asymptotics on manifolds with conic singularities, *J. Anal. Math.*, 78(1999), 1-36. 1
- [21] M. NAGASE, The fundamental solutions of the heat equations on Riemannian spaces with cone-like singular points, *Kodai Math. J.*, 7(1984), 382-455. 1, 7, 9
- [22] C. D. SOGGE, *Fourier Integrals in Classical Analysis*, *Cambridge Tracts in Mathematics*, vol. 105, Cambridge University Press, Cambridge, 1993. 11
- [23] C. D. SOGGE, *Hangzhou Lectures on Eigenfunctions of the Laplacian*, Princeton University press, Princeton and Oxford, 2014. 6, 10, 11
- [24] M. TAYLOR, *Partial Differential Equations, Vol II*, Springer, 1996. 4
- [25] X. WANG, Asymptotic expansion in time of the Schrödinger group on conical manifolds, *Ann. Inst. Fourier*, 56(2006), 1903-1945. 1
- [26] G. N. WATSON, *A Treatise on the Theory of Bessel Functions. Second Edition*, Cambridge University Press, 1944. 4, 7
- [27] S. T. YAU, Nonlinear Analysis in Geometry, Monographie No. 33 de l'Enseignement Mathématique, Série des Conférences de l'Union Mathématique Internationale, No. 8. 6
- [28] J. ZHANG, Linear restriction estimates for Schrödinger equation on metric cones, *Comm. in PDE*, 40(2015), 995-1028. 1
- [29] J. ZHANG AND J. ZHENG, Global-in-time Strichartz estimates and cubic Schrödinger equation in a conical singular space, arXiv:1702.05813 1
- [30] J. ZHANG AND J. ZHENG, Strichartz estimates and wave equation in a conic singular space, *Math. Ann.*, 376(2020), 525-581. 1

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MARYLAND, COLLEGE PARK, MD, 20742
Email address: xhuang49@umd.edu

DEPARTMENT OF MATHEMATICS, BEIJING INSTITUTE OF TECHNOLOGY, BEIJING 100081
Email address: zhang-junyong@bit.edu.cn