

A DYNAMIC APPROACH TO HETEROGENEOUS ELASTIC WIRES

ANNA DALL'ACQUA, LEONIE LANGER, AND FABIAN RUPP

ABSTRACT. We consider closed planar curves with fixed length whose elastic energy depends on an additional density variable and a spontaneous curvature. Working with the inclination angle, the associated L^2 -gradient flow is a nonlocal quasilinear coupled parabolic system of second order. We show local well-posedness and global existence of solutions for initial data in a weak regularity class and with arbitrary winding number.

1. INTRODUCTION AND MAIN RESULTS

In this work we consider a sufficiently smooth closed planar curve γ with density ρ describing a heterogeneous elastic wire. The density ρ can be used to model a distribution of mass or temperature, for instance. We assume that the bending stiffness of the wire depends on the density and consider the energy

$$(1.1) \quad \mathcal{E}_\mu(\gamma, \rho) = \frac{1}{2} \int_\gamma \left(\beta(\rho)(\kappa - c_0)^2 + \mu (\partial_s \rho)^2 \right) ds,$$

with κ the curvature of γ , the arc-length element $ds = |\partial_x \gamma(x)| dx$ and $\partial_s = \frac{1}{|\partial_x \gamma|} \partial_x$. Here $\mu > 0$ models the diffusivity of the density, the smooth positive function $\beta: \mathbb{R} \rightarrow \mathbb{R}$ is the density-modulated stiffness and $c_0 \in \mathbb{R}$ gives the spontaneous curvature. This energy is invariant under orientation-preserving reparametrizations. The first part of the energy can be seen as a Helfrich-type energy for curves with modulated stiffness. The second part, the Dirichlet energy of ρ , is needed to get control of the density.

Modulo isometries of \mathbb{R}^2 , the planar curve γ together with its orientation is uniquely determined by its inclination angle. If the curve has length $L > 0$ and is parametrized by arc-length, the energy can be expressed in terms of an inclination angle $\theta: [0, L] \rightarrow \mathbb{R}$ and the density $\rho: [0, L] \rightarrow \mathbb{R}$ by

$$(1.2) \quad E_\mu(\theta, \rho) = \frac{1}{2} \int_0^L \left(\beta(\rho)(\partial_s \theta - c_0)^2 + \mu (\partial_s \rho)^2 \right) ds,$$

using that $\kappa = \partial_s \theta$. More precisely, we have

$$\mathcal{E}_\mu(\gamma, \rho) = E_\mu(\theta, \rho),$$

where θ is an inclination angle for an orientation-preserving arc-length parametrization of γ . Note that in (1.2), the differential operator ∂_s is just the ordinary derivative with respect to s , which is applied to the functions $\theta = \theta(s)$ and $\rho = \rho(s)$. Similarly, ds just denotes integration with respect to the 1-dimensional Lebesgue measure in the variable $s \in [0, L]$. A given function $\theta: [0, L] \rightarrow \mathbb{R}$ represents the

2020 *Mathematics Subject Classification.* 35K45 (primary), 53E10 (secondary).

Key words and phrases. Euler–Bernoulli bending energy, Canham–Helfrich functional, heterogeneous material, elastic flow, nonlocal parabolic system, time-weighted Hölder spaces.

inclination angle of a C^1 -closed curve if and only if

$$(1.3) \quad \int_0^L \cos \theta \, ds = \int_0^L \sin \theta \, ds = 0,$$

$$(1.4) \quad \theta(L) - \theta(0) = 2\pi\omega,$$

for some $\omega \in \mathbb{Z}$, the rotation index of the curve. Notice that allowing ω to be negative we keep track of the orientation.

In the case of zero spontaneous curvature, this energy has previously been considered in [3], where the minimization problem under the constraints of fixed length, fixed total mass (of the density) and fixed rotation index equal to one is studied. Note that especially in the case of $c_0 \neq 0$ it makes sense to fix the length $L > 0$ of the curve, as otherwise a circle with suitable radius and a constant density is the trivial minimizer.

The goal of this article is to introduce a dynamic approach to minimize the elastic energy E_μ by evolving (θ, ρ) by the associated L^2 -gradient flow. In order to guarantee that θ describes a closed curve and to preserve the total mass of ρ , the evolution equations include nonlocal Lagrange multipliers. This results in the initial boundary value problem

$$(1.5) \quad \begin{cases} \partial_t \theta = \partial_s [\beta(\rho)(\partial_s \theta - c_0)] + \lambda_{\theta 1} \sin \theta - \lambda_{\theta 2} \cos \theta & \text{in } (0, T) \times [0, L], \\ \partial_t \rho = \mu \partial_s^2 \rho - \frac{1}{2} \beta'(\rho)(\partial_s \theta - c_0)^2 - \lambda_\rho & \text{in } (0, T) \times [0, L], \\ \theta(\cdot, L) - \theta(\cdot, 0) = 2\pi\omega, \quad \rho(\cdot, L) = \rho(\cdot, 0) & \text{on } [0, T], \\ \partial_s \theta(\cdot, L) = \partial_s \theta(\cdot, 0), \quad \partial_s \rho(\cdot, L) = \partial_s \rho(\cdot, 0) & \text{on } [0, T], \\ \theta(0, \cdot) = \theta_0, \quad \rho(0, \cdot) = \rho_0 & \text{on } [0, L], \end{cases}$$

where θ_0 satisfies (1.3). The Lagrange multipliers $\lambda_{\theta 1}$, $\lambda_{\theta 2}$ and λ_ρ are chosen such that

$$(1.6) \quad \int_0^L \cos \theta(t, s) \, ds = 0, \quad \int_0^L \sin \theta(t, s) \, ds = 0, \quad \int_0^L \rho(t, s) \, ds = \int_0^L \rho_0(s) \, ds$$

for all $t \in (0, T)$ (see Section 2 for the explicit formulas), so that the evolution indeed corresponds to an evolution of closed curves with fixed total mass. The first boundary condition in (1.5) ensures that the rotation index is constant along the flow. The other boundary conditions are imposed in order to achieve that the corresponding curve is C^2 -closed and that the density is C^1 -periodic. Observe that the evolution equations are coupled so that we cannot treat them separately.

An essential feature of working with the angle function θ instead of the curve γ is that the corresponding L^2 -gradient flow of the energy (1.2) is a parabolic system of second order in (θ, ρ) . In contrast, the Euler–Lagrange equation of (1.1) as a functional of γ and ρ is of fourth order in γ , like the classical elastica equation. Such an approach using the angle function has been used for instance in [20] to prove well-posedness and global existence for a flow towards elastica. Local and global existence of solutions for the classical elastic flow, given by a parabolic fourth order equation in \mathbb{R}^n , has been shown in several works, for instance [10, 16, 8, 7, 15, 21, 6, 19, 17, 18]. For a more detailed overview, see the recent survey [14]. A parabolic system related to (1.5) is studied in [1] and numerically elaborated in [9].

Before summarizing our results, we state that during this article

$$L > 0, \quad \mu > 0, \quad \omega \in \mathbb{Z} \text{ and } c_0 \in \mathbb{R}$$

are fixed. Moreover, $\beta: \mathbb{R} \rightarrow \mathbb{R}$ is a C^∞ -function satisfying $\beta(x) > 0$ for all $x \in \mathbb{R}$.

In many results, we will refer to the following Hypothesis 1.1, which summarizes the compatibility conditions we impose on our initial datum.

Hypothesis 1.1. Let $\theta_0, \rho_0: [0, L] \rightarrow \mathbb{R}$ be C^1 -functions such that

$$(1.7) \quad \begin{aligned} \theta_0(L) - \theta_0(0) &= 2\pi\omega, & \partial_s \theta_0(L) &= \partial_s \theta_0(0), \\ \rho_0(L) &= \rho_0(0), & \partial_s \rho_0(L) &= \partial_s \rho_0(0). \end{aligned}$$

Moreover, let θ_0 satisfy (1.3).

Our first main result is the following local well-posedness theorem. Here $h^{1+\alpha}([0, L])$ denotes the class of little Hölder functions, see Appendix A.

Theorem 1.2 (Local well-posedness). *Assume Hypothesis 1.1 and suppose that $(\theta_0, \rho_0) \in h^{1+\alpha}([0, L])$ for some $\alpha \in (0, 1)$. Then there exists $T_0 > 0$ and a unique solution $(\theta, \rho) \in C^\infty((0, T_0) \times [0, L])$ of (1.5) on $(0, T_0) \times [0, L]$ which satisfies the initial condition in the sense that*

$$\lim_{t \rightarrow 0} (\theta(t), \rho(t)) = (\theta_0, \rho_0) \quad \text{in } C^{1+\alpha}([0, L]).$$

Moreover, the solution depends continuously on the initial datum.

The idea of the proof is to get rid of the boundary conditions by transforming the problem to an equivalent one in a periodic setting. The key ingredient is then to apply the Inverse Function Theorem between appropriate time-dependent little Hölder spaces, yielding continuous dependence on the initial datum as a consequence. We deduce the local invertibility of the Fréchet derivative from the maximal regularity theory developed in [12] and a perturbation argument. The precise formulation of the continuous dependence is given in Proposition 3.3 and Remark 3.4 below. In order to show smoothness for positive times, we rely on the theory of parabolic systems by Ladyženskaja–Solonnikov–Ural’ceva [11] and a bootstrap argument. Finally, exploiting the dissipative structure of the system, uniqueness follows from an energy argument.

Next, we examine global existence. For the classical elastic flow, one expects long-time existence of the solution. We prove such a result in this more general model.

Theorem 1.3 (Global existence). *Assume Hypothesis 1.1 and suppose that $(\theta_0, \rho_0) \in h^{1+\alpha}([0, L])$ for some $\alpha \in (0, 1)$. Then the unique solution (θ, ρ) of (1.5) which satisfies $\lim_{t \rightarrow 0} (\theta(t), \rho(t)) = (\theta_0, \rho_0)$ in $C^{1+\alpha}([0, L])$ exists up to $T = \infty$. Moreover, as $t_i \rightarrow \infty$ the solution $(\theta(t_i), \rho(t_i))$ subconverges in $C^{2+\tilde{\alpha}}([0, L])$ for any $\tilde{\alpha} \in (0, \frac{1}{2})$ to a stationary solution of (1.2) as $i \rightarrow \infty$, i.e. to a solution of*

$$(1.8) \quad \begin{cases} 0 = \partial_s [\beta(\rho)(\partial_s \theta - c_0)] + \lambda_{\theta 1} \sin \theta - \lambda_{\theta 2} \cos \theta & \text{in } [0, L], \\ 0 = \mu \partial_s^2 \rho - \frac{1}{2} \beta'(\rho) (\partial_s \theta - c_0)^2 - \lambda_\rho & \text{in } [0, L], \\ \theta(L) - \theta(0) = 2\pi\omega, & \rho(L) = \rho(0), \\ \partial_s \theta(L) = \partial_s \theta(0), & \partial_s \rho(L) = \partial_s \rho(0). \end{cases}$$

Since we are facing a parabolic system, we cannot use maximum principles to show global existence. Instead, we rely on energy estimates to prove that suitable Sobolev norms along the solution cannot blow up in finite time. To that end, we first show boundedness of the velocity $\partial_t(\theta, \rho)$ in order to deal with the nonlinear coupling which limits the direct application of interpolation inequalities.

Geometrically, the angle function θ in Theorem 1.3 describes a closed curve for each fixed time t , and hence there exists a smooth family of closed planar curves with density ρ that evolve in time decreasing the energy in (1.1). We plan to address further geometric and analytic properties of the flow in a future work.

This article is structured as follows. In Chapter 2, we briefly review some basic geometric concepts concerning curves and give the formulas for the Lagrange

multipliers. The goal of Section 3 is to prove Theorem 1.2. First, we prove a local well-posedness result in the periodic setting and then show that this periodic solution gives a solution to (1.5). Section 4 is devoted to prove Theorem 1.3.

2. PRELIMINARIES

We consider an arc-length parametrization of a closed regular curve $\gamma: [0, L] \rightarrow \mathbb{R}^2$ in the plane, which we assume to be sufficiently smooth. Here $L > 0$ is the length of the curve. By a standard result in elementary differential geometry, there exists a function $\theta: [0, L] \rightarrow \mathbb{R}$ such that

$$\partial_s \gamma = \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix}.$$

Such an *inclination angle* θ is not unique, in fact it is only unique up to addition of an integer multiple of 2π . Nevertheless, many geometric quantities of the curve can be expressed in terms of an inclination angle; the (signed) *curvature* κ of the curve γ is given by $\kappa = \partial_s \theta$ and its *rotation index* $\omega \in \mathbb{Z}$ satisfies $2\pi\omega = \theta(L) - \theta(0)$, i.e. it measures the number of complete turns of the tangent vector $\partial_s \gamma$.

In (1.5) three Lagrange multipliers appear. In the first equation, λ_{θ_1} and λ_{θ_2} ensure that γ remains closed along the flow if the initial curve is closed. Indeed, defining

$$(2.1) \quad \begin{pmatrix} \lambda_{\theta_1}(t) \\ \lambda_{\theta_2}(t) \end{pmatrix} := \Pi^{-1}(\theta)(t) \int_0^L \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix} \partial_s (\beta(\rho)(\partial_s \theta - c_0)) \, ds,$$

where $\Pi^{-1}(\theta)(t)$ denotes the inverse of the matrix

$$(2.2) \quad \Pi(\theta)(t) := \begin{pmatrix} \int_0^L \sin^2 \theta \, ds & -\int_0^L \cos \theta \sin \theta \, ds \\ -\int_0^L \cos \theta \sin \theta \, ds & \int_0^L \cos^2 \theta \, ds \end{pmatrix},$$

for a sufficiently smooth solution (θ, ρ) of (1.5) we have

$$(2.3) \quad \partial_t \int_0^L \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} \, ds = 0.$$

Remark 2.1. As already observed in [20, Lemma 2.1] we have

$$\det \Pi(\theta)(t) = \frac{1}{2} \int_0^L \int_0^L \sin^2 \psi(t, s, \tilde{s}) \, ds \, d\tilde{s},$$

where $\psi(t, s, \tilde{s}) := \theta(t, s) - \theta(t, \tilde{s})$. Hence, as long as $\theta(t, \cdot)$ is not constant, the matrix $\Pi(\theta)(t)$ is invertible. In particular, the Lagrange multipliers in (2.1) are well-defined, as long as the angle function describes a closed curve.

Remark 2.2. The boundary conditions in (1.5) make it possible to use integration by parts to write the Lagrange multipliers λ_{θ_1} and λ_{θ_2} in the form

$$\begin{pmatrix} \lambda_{\theta_1}(t) \\ \lambda_{\theta_2}(t) \end{pmatrix} = \Pi^{-1}(\theta)(t) \int_0^L \begin{pmatrix} -\cos \theta \\ -\sin \theta \end{pmatrix} \partial_s \theta \beta(\rho)(\partial_s \theta - c_0) \, ds.$$

The third Lagrange multiplier λ_ρ is chosen so that the total mass remains fixed during the flow. Defining

$$(2.4) \quad \lambda_\rho(t) := -\frac{1}{2L} \int_0^L \beta'(\rho)(\partial_s \theta - c_0)^2 \, ds,$$

a sufficiently smooth solution (θ, ρ) of (1.5) satisfies

$$(2.5) \quad \partial_t \int_0^L \rho \, ds = 0.$$

From the fact that λ_{θ_1} and λ_{θ_2} ensure that the curve remains closed during the flow, we can already deduce a first property of a solution — namely the preservation of the integral over the inclination angle.

Lemma 2.3. *Let $T > 0$ and (θ, ρ) be a sufficiently smooth solution of (1.5) on $(0, T) \times [0, L]$. Then for all $t \in (0, T)$ we have*

$$\partial_t \int_0^L \theta \, ds = 0.$$

Proof. Integration of the evolution equation for θ in (1.5) yields

$$\partial_t \int_0^L \theta \, ds = \beta(\rho) (\partial_s \theta - c_0) \Big|_0^L + \lambda_{\theta_1} \int_0^L \sin \theta \, ds - \lambda_{\theta_2} \int_0^L \cos \theta \, ds.$$

By (1.6), the integrals vanish. The boundary conditions in (1.5) yield the claim. \square

As a last preliminary consideration, we note that the energy (1.2) decreases along the flow. In fact, we have a L^2 -gradient flow structure

$$(2.6) \quad \frac{d}{dt} E_\mu(\theta, \rho) = - \int_0^L (\partial_t \theta)^2 \, ds - \int_0^L (\partial_t \rho)^2 \, ds$$

which is easily verified by a direct computation for a sufficiently smooth solution (θ, ρ) of (1.5).

3. LOCAL WELL-POSEDNESS AND REGULARITY

To prove Theorem 1.2, we will consider a periodic setting. This is quite natural due to the geometric interpretation of the problem. Doing so, we get rid of the boundary conditions which enables us to apply an existence result in [12] for a suitable linearization of our problem. We first prove short-time existence and continuous dependence on the data as well as uniqueness and smoothness of the solution to the periodic problem. Then, by a suitable transformation, we can transfer the results to problem (1.5).

3.1. Transformation to a periodic setting. For the moment, assume that (θ, ρ) is a sufficiently smooth function in (t, s) . We observe that the boundary conditions in (1.5) imply that $\partial_s \theta$ is a C^0 -periodic function, i.e. it can be extended to a continuous L -periodic function on \mathbb{R} . Similarly, ρ is a C^1 -periodic function. The angle function θ is itself not periodic if the rotation number is different from zero. We hence first perform a suitable transformation. Define the function

$$\phi: [0, L] \rightarrow \mathbb{R}, \quad \phi(s) := \frac{2\pi\omega s}{L}$$

and consider the function $u(t, s) := \theta(t, s) - \phi(s)$. The boundary conditions in (1.5) imply that u and ρ are C^1 -periodic functions on $[0, L]$. We thus consider the L -periodic extensions of $u_0 := \theta_0 - \phi$ and ρ_0 to all of \mathbb{R} and look for L -periodic solutions (u, ρ) defined on $(0, T) \times \mathbb{R}$ for some $T > 0$. Now, instead of Hypothesis 1.1, it is sufficient to assume that

$$(3.1) \quad \int_0^L \cos(u_0 + \phi) \, ds = \int_0^L \sin(u_0 + \phi) \, ds = 0.$$

In terms of u , (1.5) can be equivalently written as

$$(3.2) \quad \begin{cases} \partial_t u = \partial_s [\beta(\rho) (\partial_s u + \partial_s \phi - c_0)] + \Lambda_u(u, \rho) & \text{in } (0, T) \times \mathbb{R}, \\ \partial_t \rho = \mu \partial_s^2 \rho - \frac{1}{2} \beta'(\rho) (\partial_s u + \partial_s \phi - c_0)^2 + \Lambda_\rho(u, \rho) & \text{in } (0, T) \times \mathbb{R}, \\ u(0, \cdot) = u_0, \quad \rho(0, \cdot) = \rho_0 & \text{on } \mathbb{R}. \end{cases}$$

Here, to simplify the notation, instead of working with $(\lambda_{\theta_1}, \lambda_{\theta_2})$ (see Remark 2.2) we consider Λ_u given by

$$\begin{aligned} \Lambda_u(u, \rho)(t, s) &:= \begin{pmatrix} \lambda_{\theta_1}(u + \phi, \rho)(t) \\ \lambda_{\theta_2}(u + \phi, \rho)(t) \end{pmatrix} \cdot \begin{pmatrix} \sin(u(t, s) + \phi(s)) \\ -\cos(u(t, s) + \phi(s)) \end{pmatrix} \\ &= \Pi^{-1}(u + \phi) \int_0^L \begin{pmatrix} \cos(u + \phi) \\ \sin(u + \phi) \end{pmatrix} (\partial_s u + \partial_s \phi) \beta(\rho) (\partial_s u + \partial_s \phi - c_0) ds \\ (3.3) \quad &\cdot \begin{pmatrix} \sin(u + \phi) \\ -\cos(u + \phi) \end{pmatrix} =: \Pi^{-1}(u + \phi) \mathcal{J}(u, \rho) \cdot \begin{pmatrix} \sin(u + \phi) \\ -\cos(u + \phi) \end{pmatrix} \end{aligned}$$

for $(t, s) \in (0, T) \times \mathbb{R}$ and, instead of λ_ρ , Λ_ρ given by

$$\Lambda_\rho(u, \rho)(t) := \lambda_\rho(u + \phi, \rho)(t) = \frac{1}{2L} \int_0^L \beta'(\rho) (\partial_s u + \partial_s \phi - c_0)^2 ds,$$

for $t \in (0, T)$, see (2.4).

We now define our concept of a solution. To that end, we work with time-weighted little Hölder spaces, which turns out to be an appropriate framework for the local well-posedness of the problem. A precise definition of the function spaces used in the following can be found in Appendix A.

Definition 3.1. Let $T > 0$, $\alpha \in (0, 1)$ and $\eta \in (\frac{1}{2}, 1)$ be such that $2\eta + \alpha \notin \mathbb{Z}$. Moreover, let $(u_0, \rho_0) \in h_L^{2\eta+\alpha}(\mathbb{R}; \mathbb{R}^2)$ be such that u_0 satisfies (3.1). We call (u, ρ) a *solution of (3.2) on $(0, T) \times \mathbb{R}$ with initial datum (u_0, ρ_0)* if

$$(u, \rho) \in \text{BUC}_{1-\eta}^1([0, T']; h_L^\alpha(\mathbb{R}; \mathbb{R}^2)) \cap \text{BUC}_{1-\eta}([0, T']; h_L^{2+\alpha}(\mathbb{R}; \mathbb{R}^2))$$

for all $T' < T$, if (u, ρ) solves the flow equations in (3.2) on $(0, T) \times [0, L]$ and if

$$\lim_{t \rightarrow 0} (u, \rho)(t, \cdot) = (u_0, \rho_0) \quad \text{in } C^{2\eta+\alpha}(\mathbb{R}; \mathbb{R}^2).$$

The time weight (described by the parameter η) allows us to consider less regular initial data. For readability, for $T > 0$, $\alpha \in (0, 1)$ and $\eta \in (\frac{1}{2}, 1)$ such that $2\eta + \alpha \notin \mathbb{Z}$, we set

$$(3.4) \quad \mathbb{E}_0([0, T]) := \text{BUC}_{1-\eta}([0, T]; h_L^\alpha(\mathbb{R}; \mathbb{R}^k))$$

and

$$(3.5) \quad \mathbb{E}_1([0, T]) := \text{BUC}_{1-\eta}^1([0, T]; h_L^\alpha(\mathbb{R}; \mathbb{R}^k)) \cap \text{BUC}_{1-\eta}([0, T]; h_L^{2+\alpha}(\mathbb{R}; \mathbb{R}^k))$$

for $k = 1, 2$. For $k = 2$, \mathbb{E}_1 is the space of solutions by Definition 3.1. For $(u, \rho) \in \mathbb{E}_1$ the right hand side of the evolution equations in (3.2) belongs to \mathbb{E}_0 , cf. Remark A.5. We define $\gamma\mathbb{E}_1 := h_L^{2\eta+\alpha}(\mathbb{R}; \mathbb{R}^k)$, the space for initial data (see (A.3)). In the following, we will omit the index k since it will be clear whether $k = 1$ or $k = 2$ is meant.

We introduce the open set

$$\mathbb{U}([0, T]) := \left\{ (u, \rho) \in \mathbb{E}_1([0, T]) : \inf_{[0, T]} \det \Pi(u + \phi) > 0 \right\}$$

and define the nonlinear differential operator

$$\mathbf{F}: \mathbb{U}([0, T]) \rightarrow \mathbb{E}_0([0, T]), \quad \mathbf{F}(u, \rho) = (\mathbf{F}_1(u, \rho), \mathbf{F}_2(u, \rho)),$$

where

$$(3.6) \quad \begin{aligned} \mathbf{F}_1(u, \rho) &:= \partial_s [\beta(\rho) (\partial_s u + \partial_s \phi - c_0)] + \Lambda_u(u, \rho), \\ \mathbf{F}_2(u, \rho) &:= \mu \partial_s^2 \rho - \frac{1}{2} \beta'(\rho) (\partial_s u + \partial_s \phi - c_0)^2 + \Lambda_\rho(u, \rho). \end{aligned}$$

The evolution equations in (3.2) can thus be written as

$$\partial_t (u, \rho) = \mathbf{F}(u, \rho) \quad \text{on } (0, T) \times \mathbb{R}.$$

Remark 3.2. The choice $\eta > \frac{1}{2}$ ensures that the initial datum has finite energy, but it also turns out to be essential to control the nonlinearities in the equation (cf. Lemma B.3 and Proposition B.6) and to show uniqueness (cf. Lemmas 3.6 and A.6).

3.2. Short-time existence and continuous dependence. To prove short-time existence and continuous dependence on the data, we essentially follow the ideas of [2, Theorem 2]. However, since we work with periodic little Hölder spaces, we have to justify some steps differently.

Proposition 3.3 (Existence and continuous dependence). *Let $\alpha \in (0, 1)$ and $\eta \in (\frac{1}{2}, 1)$. Let $(u_0, \rho_0) \in \gamma\mathbb{E}_1$ and let u_0 satisfy (3.1). Then there exist $T_0 \in (0, \infty)$ and $r > 0$ such that for every $(\tilde{u}_0, \tilde{\rho}_0) \in \gamma\mathbb{E}_1$ with $\|(u_0, \rho_0) - (\tilde{u}_0, \tilde{\rho}_0)\|_{C^{2\eta+\alpha}([0, L])} < r$ the problem*

$$(3.7) \quad \begin{cases} \partial_t \tilde{u} = \partial_s [\beta(\tilde{\rho}) (\partial_s \tilde{u} + \partial_s \phi - c_0)] + \Lambda_u(\tilde{u}, \tilde{\rho}) & \text{in } (0, T_0) \times \mathbb{R}, \\ \partial_t \tilde{\rho} = \mu \partial_s^2 \tilde{\rho} - \frac{1}{2} \beta'(\tilde{\rho}) (\partial_s \tilde{u} + \partial_s \phi - c_0)^2 + \Lambda_\rho(\tilde{u}, \tilde{\rho}) & \text{in } (0, T_0) \times \mathbb{R}, \\ \tilde{u}(0, \cdot) = \tilde{u}_0, \quad \tilde{\rho}(0, \cdot) = \tilde{\rho}_0 & \text{on } \mathbb{R}, \end{cases}$$

admits a solution $(\tilde{u}, \tilde{\rho}) \in \mathbb{E}_1([0, T_0])$. Moreover, there is a C^1 -map which associates to the initial datum a solution of the problem. In particular, there exists a solution $(u, \rho) \in \mathbb{E}_1([0, T_0])$ of (3.2) on $(0, T_0) \times \mathbb{R}$ with initial datum (u_0, ρ_0) .

Proof. We divide the proof into several steps. For the sake of readability, the details are partially moved to Appendix B. We will refer to this at the respective points.

Step 1: Construction of a “good” function. We show that there exist $\bar{T} > 0$ and $(\bar{u}, \bar{\rho}) \in \mathbb{E}_1([0, \bar{T}])$ with $(\bar{u}, \bar{\rho})(0, \cdot) = (u_0, \rho_0)$, such that $(\bar{u}, \bar{\rho}) \in \mathcal{U}([0, \bar{T}])$, i.e.

$$(3.8) \quad \inf_{[0, \bar{T}]} \det(\Pi(\bar{u} + \phi)) > 0,$$

and having some extra regularity. More precisely, $\partial_s(\bar{u}, \bar{\rho}) \in \tilde{\mathbb{E}}_1([0, \bar{T}])$ where

$$\tilde{\mathbb{E}}_1([0, \bar{T}]) := \text{BUC}_{1-\tilde{\eta}}^1([0, \bar{T}]; h_L^\alpha(\mathbb{R})) \cap \text{BUC}_{1-\tilde{\eta}}([0, \bar{T}]; h_L^{2+\alpha}(\mathbb{R}))$$

for $\tilde{\eta} := \eta - \frac{1}{2} > 0$.

To begin with we take the solution $(\bar{u}, \bar{\rho}) \in \mathbb{E}_1([0, \hat{T}])$, for $\hat{T} > 0$ arbitrary, of the evolution problem

$$\begin{cases} \partial_t(v, \sigma) - \partial_s^2(v, \sigma) = (0, 0) & \text{in } (0, \hat{T}) \times \mathbb{R}, \\ (v, \sigma)(0, \cdot) = (u_0, \rho_0) & \text{on } \mathbb{R}. \end{cases}$$

Such a solution exists by [12, Theorem 6.4]. As solution of the heat equation, we know that $(\bar{u}, \bar{\rho}) \in C^\infty((0, \hat{T}) \times \mathbb{R})$. Due to the fact that $\eta > \frac{1}{2}$, $\partial_s(u_0, \rho_0) \in h_L^{2\tilde{\eta}+\alpha}(\mathbb{R}) = \gamma\tilde{\mathbb{E}}_1$ and hence it is an admissible initial datum. We look at

$$(3.9) \quad \begin{cases} \partial_t(v, \sigma) - \partial_s^2(v, \sigma) = (0, 0) & \text{in } (0, \hat{T}) \times \mathbb{R}, \\ (v, \sigma)(0, \cdot) = \partial_s(u_0, \rho_0) & \text{on } \mathbb{R}. \end{cases}$$

Again, we use [12, Theorem 6.4] and obtain a unique solution in $\tilde{\mathbb{E}}_1([0, \hat{T}])$ that is smooth for positive times. Since this solution coincides with $\partial_s(\bar{u}, \bar{\rho})$ for $t = 0$ and $\partial_s(\bar{u}, \bar{\rho})$ solves the evolution equation in (3.9), we conclude by uniqueness that $\partial_s(\bar{u}, \bar{\rho})$ is the solution of (3.9), so that $\partial_s(\bar{u}, \bar{\rho}) \in \tilde{\mathbb{E}}_1([0, \hat{T}])$.

It remains to check (3.8). Since $u_0 + \phi$ satisfies Hypothesis 1.1, using Remark 2.1 and the embedding in Lemma A.3, by continuity there exists $\bar{T} \leq \hat{T}$ such that (3.8) is satisfied.

Step 2: We claim that there exists $T' \in (0, \bar{T}]$ such that the operator

$$\begin{aligned} \Phi: \mathbb{U}([0, T']) &\rightarrow \mathbb{E}_0([0, T']) \times \gamma\mathbb{E}_1, \\ \Phi(\tilde{u}, \tilde{\rho}) &:= \left(\partial_t(\tilde{u}, \tilde{\rho}) - \mathbf{F}(\tilde{u}, \tilde{\rho}), (\tilde{u}, \tilde{\rho})(0, \cdot) \right) \end{aligned}$$

is well-defined and a local diffeomorphism at $(\bar{u}, \bar{\rho})$ (restricted to $[0, T']$). Here \mathbf{F} is the map defined in (3.6).

For the claim it is sufficient to show the existence of $T' \in (0, \bar{T}]$ such that the Fréchet derivative

$$(3.10) \quad \begin{aligned} D\Phi(\bar{u}, \bar{\rho}): \mathbb{E}_1([0, T']) &\rightarrow \mathbb{E}_0([0, T']) \times \gamma\mathbb{E}_1, \\ D\Phi(\bar{u}, \bar{\rho})(v, \sigma) &:= \left(\partial_t(v, \sigma) - D\mathbf{F}(\bar{u}, \bar{\rho})(v, \sigma), (v, \sigma)(0, \cdot) \right), \end{aligned}$$

is a linear isomorphism. It then follows, since $\Phi \in C^1$, by the Local Inverse Function Theorem that Φ is a local diffeomorphism at $(\bar{u}, \bar{\rho})$. The proof that the map in (3.10) is a linear isomorphism is given in Proposition B.6 (with $\tilde{\mathbf{J}} = D\Phi(\bar{u}, \bar{\rho})$). The idea is to use a maximal regularity result in time-weighted little Hölder spaces [12], where the time weight allows for rough initial data. Since the results in [12] are applicable to linear systems with time-independent coefficients, we first consider an auxiliary linear problem, cf. Proposition B.1. A perturbation argument then gives that the map in (3.10) is an isomorphism for small times.

Step 3: By Step 2, there is a neighborhood $V \subset \mathbb{U}([0, T'])$ of $(\bar{u}, \bar{\rho})$ and a neighborhood $W \subset \mathbb{E}_0([0, T']) \times \gamma\mathbb{E}_1$ of $\Phi(\bar{u}, \bar{\rho}) =: ((\bar{f}_1, \bar{f}_2), (u_0, \rho_0))$ such that $\Phi: V \rightarrow W$ is a diffeomorphism. Choose $r > 0$ small enough such that $B_r(\bar{f}_1, \bar{f}_2) \times B_r(u_0, \rho_0) \subset W$. Moreover, choose $T'' \in (0, T')$ small enough so that

$$(3.11) \quad \|(\tilde{f}_1, \tilde{f}_2)\|_{\mathbb{E}_0([0, T''])} < r.$$

Here it is crucial that the \mathbb{E}_0 -norm becomes arbitrary small for small times (see the definition of $\text{BUC}_{1-\eta}$ in Section A.2).

Step 4: Define $(\tilde{f}_1, \tilde{f}_2) \in \mathbb{E}_0([0, T'])$ by

$$(\tilde{f}_1, \tilde{f}_2) := \begin{cases} (0, 0) & \text{on } [0, \frac{1}{2}T''], \\ \frac{2}{T''}(t - \frac{1}{2}T'')(f_1, f_2) & \text{on } (\frac{1}{2}T'', T''), \\ (\bar{f}_1, \bar{f}_2) & \text{on } [T'', T']. \end{cases}$$

Then by a short computation and the choice of T'' in (3.11) we have

$$\|(\tilde{f}_1, \tilde{f}_2) - (\bar{f}_1, \bar{f}_2)\|_{\mathbb{E}_0([0, T'])} < r.$$

This implies that, with $(\tilde{u}_0, \tilde{\rho}_0) \in \gamma\mathbb{E}_1$ as in the statement,

$$((\tilde{f}_1, \tilde{f}_2), (\tilde{u}_0, \tilde{\rho}_0)) \in B_r(\bar{f}_1, \bar{f}_2) \times B_r(u_0, \rho_0) \subset W.$$

Since $\Phi: V \rightarrow W$ is a diffeomorphism, there exists $(\tilde{u}, \tilde{\rho}) \in V \subset \mathbb{U}([0, T'])$ such that $\Phi(\tilde{u}, \tilde{\rho}) = ((\tilde{f}_1, \tilde{f}_2), (\tilde{u}_0, \tilde{\rho}_0))$. We restrict $(\tilde{u}, \tilde{\rho})$ to $[0, \frac{1}{2}T''] \times \mathbb{R}$, set $T_0 := \frac{1}{2}T''$ and obtain a solution of (3.7) on $(0, T_0) \times \mathbb{R}$.

Step 5: We constructed a mapping which maps every $(\tilde{u}_0, \tilde{\rho}_0) \in B_r(u_0, \rho_0)$ to a solution $(\tilde{u}, \tilde{\rho}) \in \mathbb{E}_1([0, T_0])$ of the problem. It consists essentially of the composition of the C^1 -mapping Φ^{-1} and a restriction operator and is thus continuously differentiable. \square

Remark 3.4. In the following section we prove uniqueness of solutions. Together with Proposition 3.3 this yields the continuity — or more precisely even the continuous differentiability — of the mapping which maps any initial datum $(u_0, \rho_0) \in \gamma\mathbb{E}_1$ to the unique solution $(u, \rho) \in \mathbb{E}_1([0, T_0])$ of (3.2).

3.3. Smoothness and uniqueness of solutions. So far, we have proven existence of a solution as well as continuous dependence on the data. We now show that a solution is smooth for positive times.

Lemma 3.5 (Smoothing). *Let $0 < T \leq \infty$ and (u, ρ) be a solution, in the sense of Definition 3.1, of (3.2) on $(0, T) \times \mathbb{R}$. Then $(u, \rho) \in C^\infty((0, T) \times \mathbb{R})$.*

The idea is to define a smooth cut-off function, which vanishes at $t = 0$ and outside of a spatial interval of length bigger than L . The product of u with this cut-off function is the solution of a new initial value problem — this time with boundary values and zero initial data. This enables us to use [11, Theorem IV.5.2], which gives us higher regularity of u in the scale of parabolic Hölder spaces, cf. Appendix A.3. We proceed similarly for ρ and iterate this procedure to conclude smoothness.

Proof of Lemma 3.5. By Definition 3.1 the solution (u, ρ) satisfies $(u, \rho) \in \mathbb{E}_1([0, T'])$ for all $T' < T$. Let $0 < \delta < T' < T$ and let $\psi_1 : [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$ be a smooth cut-off function satisfying

$$\begin{aligned} \psi_1(t, s) &\equiv 1 \text{ for } (t, s) \in [\delta/2, T'] \times [-L/4, 5L/4], \\ \psi_1(t, s) &\equiv 0 \text{ for } (t, s) \in ([0, \delta/4] \times \mathbb{R}) \cup ([0, T'] \times (\mathbb{R} \setminus [-L/2, 3L/2])). \end{aligned}$$

We define $u_1 := \psi_1 u \in \mathbb{E}_1([0, T'])$. By (3.2) we have

$$\partial_t u_1 - a_{11} \partial_s^2 u_1 + a_1 \partial_s u_1 = f_1,$$

where

$$\begin{aligned} a_{11}(t, s) &:= \beta(\rho), \quad a_1(t, s) := -\beta'(\rho) \partial_s \rho, \\ f_1(t, s) &:= -2\beta(\rho) \partial_s u \partial_s \psi_1 - \beta(\rho) u \partial_s^2 \psi_1 - \beta'(\rho) u \partial_s \rho \partial_s \psi_1 + \psi_1 \beta'(\rho) \partial_s \rho (\partial_s \phi - c_0) \\ &\quad + \psi_1 \Lambda_u(u, \rho) + u \partial_t \psi_1. \end{aligned}$$

Defining $\zeta := \eta + \frac{\alpha}{2} - \frac{1}{2} \in (0, 1)$, Lemma A.7 in the appendix yields

$$\partial_s u, \partial_s \rho \in H^{\frac{\zeta}{2}, \zeta}([0, T'] \times [-L, 2L]).$$

Since $\beta \in C^\infty(\mathbb{R})$, we conclude $a_{11}, a_1, f_1 \in H^{\frac{\zeta}{2}, \zeta}([0, T'] \times [-L, 2L])$. By [11, Theorem IV.5.2] there exists a unique solution $w \in H^{\frac{\zeta+2}{2}, \zeta+2}([0, T'] \times [-L, 2L])$ of

$$(3.12) \quad \begin{cases} \partial_t w - a_{11} \partial_s^2 w + a_1 \partial_s w = f_1 & \text{in } (0, T') \times (-L, 2L), \\ w = 0 & \text{on } (0, T') \times \{-L, 2L\}, \\ w(0, \cdot) = 0 & \text{on } (-L, 2L). \end{cases}$$

By uniqueness of weak solutions, $u_1 = w$ and $u_1 \in H^{\frac{\zeta+2}{2}, \zeta+2}([0, T'] \times [-L, 2L])$. Proceeding analogously for ρ , we obtain

$$u, \rho \in H^{\frac{\zeta+2}{2}, \zeta+2}([\delta/2, T'] \times [-L/4, 5L/4]).$$

Since u and ρ are periodic with period length L we even have

$$(3.13) \quad u, \rho \in H^{\frac{\zeta+2}{2}, \zeta+2}([\delta/2, T'] \times [-L, 2L]).$$

Let $\psi_2 : [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$ be another smooth cut-off function satisfying

$$\begin{aligned} \psi_2(t, s) &\equiv 1 \text{ for } (t, s) \in [3\delta/4, T'] \times [-L/4, 5L/4], \\ \psi_2(t, s) &\equiv 0 \text{ for } (t, s) \in ([0, 2\delta/3] \times \mathbb{R}) \cup ([0, T'] \times \mathbb{R} \setminus [-L/2, 3L/2]). \end{aligned}$$

We set $u_2 := \psi_2 u \in H^{\frac{\zeta+2}{2}, \zeta+2}([\delta/2, T'] \times [-L, 2L])$. Now, u_2 solves an initial boundary value problem similar to (3.12) where f_2 is given by f_1 with ψ_1 replaced by ψ_2 and where the initial condition is prescribed at $t = \delta/2$. By (3.13), we find that $a_{11}, a_1, f_2 \in H^{\frac{\zeta+1}{2}, \zeta+1}([\delta/2, T'] \times [-L, 2L])$, so again applying [11, Theorem

IV.5.2], we conclude $u_2 \in H^{\frac{\zeta+3}{2}, \zeta+3}([\delta/2, T'] \times [-L, 2L])$. Again, proceeding similarly for ρ , we find

$$u, \rho \in H^{\frac{\zeta+3}{2}, \zeta+3}([3\delta/4, T'] \times [-L, 2L]).$$

Iterating the above procedure, by periodicity and since $0 < \delta < T' < T$ were arbitrary, it follows $(u, \rho) \in C^\infty((0, T) \times \mathbb{R})$. \square

We are now able to prove that solutions are unique.

Lemma 3.6 (Uniqueness). *Let $(u_0, \rho_0) \in \gamma\mathbb{E}_1$. Let $0 < T_1, T_2 \leq \infty$ and let (u_1, ρ_1) and (u_2, ρ_2) be solutions of (3.2) on $(0, T_1) \times \mathbb{R}$ and $(0, T_2) \times \mathbb{R}$ with initial datum (u_0, ρ_0) . Then $(u_1, \rho_1) \equiv (u_2, \rho_2)$ on $[0, \min\{T_1, T_2\})$.*

Proof. Without loss of generality we assume $0 < T_1 \leq T_2$. Let $0 < T < T_1$. We show that the difference $(u_d, \rho_d) := (u_1, \rho_1) - (u_2, \rho_2)|_{[0, T]}$ of the two solutions is identically zero with an argument based on Gronwall's lemma. Note that by Lemma 3.5 $(u_d, \rho_d) \in C^\infty((0, T] \times \mathbb{R})$. In particular, the function

$$\varphi(t) := \frac{1}{2} \int_0^L \left((\partial_s u_d)^2 + (\partial_s \rho_d)^2 + u_d^2 + \rho_d^2 \right) ds$$

is differentiable on $(0, T]$. In the following, C denotes a constant, only depending on u_1, u_2, ρ_1, ρ_2 and the model parameters $\beta, c_0, L, \mu, \omega$, which may change from line to line. Other dependencies will be given explicitly. By Lemma A.3 and Remark A.4, we have

$$(3.14) \quad \sup_{t \in [0, T]} \|(u_i, \rho_i)\|_{C^1} \leq C, \quad \sup_{[0, T] \times \mathbb{R}} |\beta^{(k)}(\rho_i)| \leq C, \quad \sup_{[0, T]} \|\Pi^{-1}(u_i + \phi)\| \leq C$$

for $i = 1, 2, k = 0, 1, 2$. In particular, there exists a compact interval $J \subset \mathbb{R}$ such that $\rho_i(t, s) \in J$ for all $(t, s) \in [0, T] \times \mathbb{R}$ and $i = 1, 2$. Consequently, we have

$$(3.15) \quad \inf_{[0, T] \times \mathbb{R}} \beta(\rho_i) \geq c > 0 \quad \text{for } i = 1, 2.$$

After integration by parts and using $\int_0^L \rho_d ds = 0$ by (2.5), we get

$$\begin{aligned} \frac{d}{dt} \varphi(t) &= - \int_0^L \beta(\rho_1) (\partial_s^2 u_d)^2 ds - \mu \int_0^L (\partial_s^2 \rho_d)^2 ds - \mu \int_0^L (\partial_s \rho_d)^2 ds \\ &\quad - \int_0^L \partial_s^2 u_d (\beta(\rho_1) - \beta(\rho_2)) \partial_s^2 u_2 ds \\ &\quad - \int_0^L \partial_s^2 u_d \left(\beta'(\rho_1) \partial_s \rho_1 (\partial_s u_1 + \partial_s \phi - c_0) + \Lambda_u(u_1, \rho_1) \right. \\ &\quad \quad \left. - \beta'(\rho_2) \partial_s \rho_2 (\partial_s u_2 + \partial_s \phi - c_0) - \Lambda_u(u_2, \rho_2) \right) ds \\ &\quad + \int_0^L \partial_s^2 \rho_d \left(\frac{1}{2} \beta'(\rho_1) (\partial_s u_1 + \partial_s \phi - c_0)^2 - \frac{1}{2} \beta'(\rho_2) (\partial_s u_2 + \partial_s \phi - c_0)^2 \right) ds \\ &\quad + \int_0^L u_d \left(\beta(\rho_1) \partial_s^2 u_1 + \beta'(\rho_1) \partial_s \rho_1 (\partial_s u_1 + \partial_s \phi - c_0) + \Lambda_u(u_1, \rho_1) \right. \\ &\quad \quad \left. - \beta(\rho_2) \partial_s^2 u_2 - \beta'(\rho_2) \partial_s \rho_2 (\partial_s u_2 + \partial_s \phi - c_0) - \Lambda_u(u_2, \rho_2) \right) ds \\ (3.16) \quad &+ \int_0^L \rho_d \left(-\frac{1}{2} \beta'(\rho_1) (\partial_s u_1 + \partial_s \phi - c_0)^2 + \frac{1}{2} \beta'(\rho_2) (\partial_s u_2 + \partial_s \phi - c_0)^2 \right) ds. \end{aligned}$$

The first three terms have a negative sign. The remaining terms can be estimated using Young's inequality to absorb higher order derivatives and bound the remaining ones by $\varphi(t)$ using (3.14). Indeed, since $\int_0^L \rho_d ds = 0$, for each t there is $s_t \in \mathbb{R}$

with $\rho_d(t, s_t) = 0$ by the Mean Value Theorem for Integrals. The Fundamental Theorem of Calculus thus yields

$$(3.17) \quad |\rho_d(t, s)| \leq \int_0^L |\partial_s \rho_d| ds.$$

Hence, the term on the second line in (3.16) can be estimated by

$$\sup_{\mathbb{R}} |\rho_d| \int_0^L |\partial_s^2 u_d| |\partial_s^2 u_2| ds \leq \delta \int_0^L (\partial_s^2 u_d)^2 ds + C(\delta) \int_0^L (\partial_s^2 u_2)^2 ds \int_0^L (\partial_s \rho_d)^2 ds.$$

We now examine the third line in (3.16). After adding and subtracting to create differences, using Young's inequality we find on the one hand

$$(3.18) \quad \begin{aligned} & - \int_0^L \partial_s^2 u_d (\beta'(\rho_1) \partial_s \rho_1 (\partial_s u_1 + \partial_s \phi - c_0) - \beta'(\rho_2) \partial_s \rho_2 (\partial_s u_2 + \partial_s \phi - c_0)) ds \\ &= - \int_0^L \partial_s^2 u_d \beta'(\rho_1) \partial_s \rho_d (\partial_s u_1 + \partial_s \phi - c_0) ds - \int_0^L \partial_s^2 u_d \beta'(\rho_1) \partial_s \rho_2 \partial_s u_d ds \\ & \quad - \int_0^L \partial_s^2 u_d (\beta'(\rho_1) - \beta'(\rho_2)) \partial_s \rho_2 (\partial_s u_2 + \partial_s \phi - c_0) ds \\ & \leq \delta \int_0^L (\partial_s^2 u_d)^2 ds + C(\delta) \left(\int_0^L (\partial_s \rho_d)^2 + (\partial_s u_d)^2 + \rho_d^2 ds \right). \end{aligned}$$

On the other hand, for the term in the third line of (3.16) involving the Lagrange multiplier we have

$$(3.19) \quad \begin{aligned} & - \int_0^L \partial_s^2 u_d (\Lambda_u(u_1, \rho_1) - \Lambda_u(u_2, \rho_2)) ds \\ &= - \int_0^L \partial_s^2 u_d \left[\Pi^{-1}(u_1 + \phi) \mathcal{J}(u_1, \rho_1) \cdot \begin{pmatrix} \sin(u_1 + \phi) \\ -\cos(u_1 + \phi) \end{pmatrix} \right. \\ & \quad \left. - \Pi^{-1}(u_2 + \phi) \mathcal{J}(u_2, \rho_2) \cdot \begin{pmatrix} \sin(u_2 + \phi) \\ -\cos(u_2 + \phi) \end{pmatrix} \right] ds. \end{aligned}$$

We write the term in square brackets as

$$\begin{aligned} & [\Pi^{-1}(u_1 + \phi) - \Pi^{-1}(u_2 + \phi)] \mathcal{J}(u_1, \rho_1) \cdot \begin{pmatrix} \sin(u_1 + \phi) \\ -\cos(u_1 + \phi) \end{pmatrix} \\ & + \Pi^{-1}(u_2 + \phi) (\mathcal{J}(u_1, \rho_1) - \mathcal{J}(u_2, \rho_2)) \cdot \begin{pmatrix} \sin(u_1 + \phi) \\ -\cos(u_1 + \phi) \end{pmatrix} \\ & + \Pi^{-1}(u_2 + \phi) \mathcal{J}(u_2, \rho_2) \cdot \begin{pmatrix} \sin(u_1 + \phi) - \sin(u_2 + \phi) \\ -\cos(u_1 + \phi) + \cos(u_2 + \phi) \end{pmatrix}. \end{aligned}$$

Using the explicit formula for the inverse matrix, it can be shown that

$$\|\Pi^{-1}(u_1 + \phi) - \Pi^{-1}(u_2 + \phi)\| \leq C \int_0^L |u_d| ds.$$

Proceeding as in (3.18), after adding and subtracting, we have

$$|\mathcal{J}(u_1, \rho_1) - \mathcal{J}(u_2, \rho_2)| \leq C \int_0^L (|u_d| + |\partial_s u_d| + |\rho_d|) ds.$$

Consequently, (3.19) may be estimated by

$$\delta \int_0^L (\partial_s^2 u_d)^2 ds + C(\delta) \int_0^L u_d^2 ds + C(\delta) \int_0^L (\partial_s u_d)^2 + C(\delta) \int_0^L \rho_d^2 ds.$$

For the remaining terms, we proceed similarly. Using (3.15) and choosing $\delta > 0$ sufficiently small, the first two terms in (3.16) can be used to absorb all terms involving a factor of δ . Altogether, we obtain the differential inequality

$$\frac{d}{dt}\varphi(t) \leq C \left(1 + \int_0^L (\partial_s^2 u_2)^2 ds \right) \varphi(t) \quad \text{for } t \in (0, T).$$

Using Gronwall's lemma, we conclude that for all $t' \in [0, T]$ we have

$$\varphi(t') \leq C\varphi(0) \exp \left(\int_0^{t'} \left(1 + \int_0^L (\partial_s^2 u_2)^2 ds \right) dt \right).$$

By Lemma A.6, the integral is finite and with $\varphi(0) = 0$, it follows that $(u_1, \rho_1) \equiv (u_2, \rho_2)$ on $[0, T]$. \square

3.4. Proof of Theorem 1.2. It remains to transfer the results of Section 3.2 and Section 3.3 to the initial nonperiodic setting and to prove Theorem 1.2.

Proof of Theorem 1.2. We define (u_0, ρ_0) as the L -periodic extension of $(\theta_0 - \phi, \rho_0)$ to all of \mathbb{R} . Then $(u_0, \rho_0) \in h_L^{1+\alpha}(\mathbb{R})$. Now we choose $\bar{\alpha} \in (0, 1)$ and $\bar{\eta} \in (\frac{1}{2}, 1)$ such that $2\bar{\eta} + \bar{\alpha} = 1 + \alpha$. Since $(u_0, \rho_0) \in h_L^{2\bar{\eta} + \bar{\alpha}}(\mathbb{R})$, Proposition 3.3 yields $\bar{T} > 0$ and a solution

$$(3.20) \quad (u, \rho) \in \text{BUC}_{1-\bar{\eta}}^1([0, \bar{T}]; h_L^{\bar{\alpha}}(\mathbb{R})) \cap \text{BUC}_{1-\bar{\eta}}([0, \bar{T}]; h_L^{2+\bar{\alpha}}(\mathbb{R}))$$

of (3.2) on $(0, \bar{T}) \times \mathbb{R}$ that satisfies $\lim_{t \rightarrow 0} (u, \rho)(t) = (u_0, \rho_0)$ in $C^{2\bar{\eta} + \bar{\alpha}}(\mathbb{R})$ by Lemma A.3. According to Proposition 3.6, the solution is unique and due to Lemma 3.5 it holds that $(u, \rho) \in C^\infty((0, \bar{T}) \times \mathbb{R})$. Now we define (θ, ρ) as the restriction of $(u + \phi, \rho)$ on $[0, \bar{T}] \times [0, L]$. By construction (θ, ρ) solves (1.5) on $(0, \bar{T}) \times [0, L]$. Moreover, the C^1 -periodicity of (u, ρ) ensures, that (θ, ρ) satisfies the boundary conditions in (1.5). Furthermore, the choice of $\bar{\alpha}$ and $\bar{\eta}$ as $2\bar{\eta} + \bar{\alpha} = 1 + \alpha$ gives us $\lim_{t \rightarrow 0} (\theta(t), \rho(t)) = (\theta_0, \rho_0)$ in $C^{1+\alpha}([0, L]; \mathbb{R}^2)$. The uniqueness of the solution, its smoothness and the continuous dependence on the initial datum transfer from (u, ρ) to (θ, ρ) . \square

Remark 3.7. In the above proof of Theorem 1.2 we chose $\bar{\alpha} \in (0, 1)$ and $\bar{\eta} \in (\frac{1}{2}, 1)$ such that $2\bar{\eta} + \bar{\alpha} = 1 + \alpha \in (1, 2)$. This yields the convergence of the solution (θ, ρ) to the initial datum (θ_0, ρ_0) in $C^{1+\alpha}([0, L])$. But in fact, in Proposition 3.3 we can allow for arbitrary $\bar{\alpha} \in (0, 1)$ and $\bar{\eta} \in (\frac{1}{2}, 1)$ with just the condition that $\bar{\alpha} \in (0, 1)$ and $\bar{\eta} \in (\frac{1}{2}, 1)$ and $2\bar{\eta} + \bar{\alpha} \notin \mathbb{Z}$. By exploiting this freedom we observe the following. If we even have $(\theta_0, \rho_0) \in h^{2+\alpha}([0, L])$ for $\alpha \in (0, 1)$, we get convergence of the solution (θ, ρ) to the initial datum (θ_0, ρ_0) in $C^{2+\alpha}([0, L])$. This is an important ingredient in proving global existence, cf. Section 4.3 below.

4. GLOBAL EXISTENCE

Let $(\theta_0, \rho_0) \in h^{1+\alpha}([0, L])$, $\alpha \in (0, 1)$ be such that Hypothesis 1.1 is satisfied. As a consequence of Theorem 1.2, a standard argument for initial value problems yields the existence of a *maximal existence time* $T_{\max} \in (0, \infty]$ such that the unique solution (θ, ρ) to (1.5) with this initial datum can be smoothly extended to $(0, T_{\max})$. The goal of this section is to prove Theorem 1.3, i.e. that $T_{\max} = \infty$.

4.1. Controlling the matrix Π^{-1} . As a first step, we show that the inverse of the matrix in (2.2), which defines the Lagrange-multipliers in (2.1), does not degenerate if the L^2 -norm of the curvature $\kappa = \partial_s \theta$ is bounded. In contrast to [20, Lemma 2.1], our proof relies on variational methods and does not depend on the winding number.

Lemma 4.1. *Let $K > 0$. Then there exists $M = M(K, L) > 0$ such that for all functions $\theta \in W^{1,2}([0, L])$ satisfying (1.3) and*

$$(4.1) \quad \int_0^L (\partial_s \theta)^2 ds \leq K$$

we have

$$\|\Pi^{-1}(\theta)\| := \sup_{\substack{X \in \mathbb{R}^2, \\ |X|=1}} |\Pi^{-1}(\theta)(X)| \leq M.$$

Proof. By (2.2), all entries of the matrix $\Pi(\theta)$ are bounded by L , thus it suffices to bound $\det \Pi(\theta)$ from below for all $\theta \in W^{1,2}([0, L])$ satisfying (1.3) and (4.1) by some constant $m = m(K, L) > 0$. Let $(\theta_k)_{k \in \mathbb{N}} \subset W^{1,2}([0, L])$ be a minimizing sequence for

$$m := \inf \{ \det \Pi(\theta) \mid \theta \in W^{1,2}([0, L]) \text{ satisfying (1.3) and (4.1)} \}.$$

Note that $m \geq 0$ by Remark 2.1. Without loss of generality, we may assume $\theta_k(0) = 0$, after replacing θ_k by $\theta_k - \theta_k(0)$. Indeed, by direct computation (4.1) and also (1.3) are invariant under addition of a constant. Moreover, Remark 2.1 directly implies that $\det \Pi(\theta_k - \theta_k(0)) = \det \Pi(\theta_k)$.

Consequently, $(\theta_k)_{k \in \mathbb{N}} \subset W^{1,2}([0, L])$ is bounded, so that we may assume $\theta_k \rightharpoonup \theta$ in $W^{1,2}([0, L])$ and $\theta_k \rightarrow \theta$ in $C([0, L])$ by Sobolev embedding. The limit θ clearly satisfies (1.3) and, by weak lower semicontinuity of the L^2 -norm, also (4.1). From (2.2), we find $\det \Pi(\theta_k) \rightarrow \det \Pi(\theta)$ and thus conclude that $m = \det \Pi(\theta)$ is a minimum. Since θ is continuous, by Remark 2.1 we find that $m = 0$ if and only if θ is constant. However, the latter is impossible due to (1.3), so $m > 0$ and the claim follows. \square

4.2. Energy estimates. We now prove norm estimates for our solution, which by Lemma 3.5 instantaneously becomes smooth. Since we are only interested in long-time existence here, for the rest of this subsection we assume that we have a fixed *smooth* solution $(\theta, \rho) \in C^\infty([0, T] \times [0, L])$ of (1.5). This means that the initial datum (θ_0, ρ_0) is smooth and satisfies Hypothesis 1.1.

Furthermore, throughout this subsection $C \in (0, \infty)$ denotes a generic constant that may vary from line to line. The constant is only allowed to depend on the fixed model parameters as well as the initial datum (θ_0, ρ_0) . In particular, C does not depend on the (not necessarily maximal) existence time T .

This is essential for proving the subconvergence part of Theorem 1.3. First, we gather some direct consequences of the constrained gradient flow structure.

Proposition 4.2. *We have*

- (i) $\|\theta\|_{L^\infty([0, T] \times [0, L])} \leq C, \quad \|\rho\|_{L^\infty([0, T] \times [0, L])} \leq C,$
- (ii) $\sup_{t \in [0, T]} \|\partial_s \theta\|_{L^2(0, L)} \leq C, \quad \sup_{t \in [0, T]} \|\partial_s \rho\|_{L^2(0, L)} \leq C,$
- (iii) $\|\partial_t \theta\|_{L^2([0, T] \times [0, L])} \leq C, \quad \|\partial_t \rho\|_{L^2([0, T] \times [0, L])} \leq C.$

Proof. Statement (iii) follows from integrating (2.6) in time and using $E_\mu(\theta, \rho) \geq 0$. Moreover, for any $t \in [0, T]$ we obtain

$$E_\mu(\theta_0, \rho_0) \geq \frac{\mu}{2} \int_0^L (\partial_s \rho)^2 ds$$

and the second estimate in (ii) is proven.

We will now use this to get an L^∞ -bound on ρ . Since the total mass is conserved along the evolution by (1.6), the Mean Value Theorem for integrals implies the

existence of $s_t \in [0, L]$ with $\rho(t, s_t) = \frac{1}{L} \int_0^L \rho_0 ds$ for any $t \in [0, T]$. Similar to (3.17), the Fundamental Theorem of Calculus yields

$$(4.2) \quad |\rho(t, s) - \rho(t, s_t)| \leq \int_0^L |\partial_s \rho| ds \leq L^{\frac{1}{2}} \left(\int_0^L |\partial_s \rho|^2 ds \right)^{\frac{1}{2}} \leq C$$

and the second estimate of (i) is proven.

Consequently, there exists a compact interval $J \subset \mathbb{R}$ only depending on θ_0, ρ_0 and the model parameters such that

$$(4.3) \quad \rho(t, s) \in J \quad \text{for all } t \in [0, T], s \in [0, L].$$

By continuity, we find that

$$(4.4) \quad \inf_J \beta \geq C > 0.$$

Therefore, using (2.6) again, we obtain

$$E_\mu(\theta_0, \rho_0) \geq \inf_J \beta \int_0^L (\partial_s \theta - c_0)^2 ds \geq C \left(\int_0^L (\partial_s \theta)^2 ds - 2c_0 \int_0^L \partial_s \theta ds + c_0^2 L \right).$$

By (1.4), we have $\int_0^L \partial_s \theta ds = 2\pi\omega$, and hence the first part of (ii) follows.

This together with the fact that $\int_0^L \theta ds = \int_0^L \theta_0 ds$ by Lemma 2.3 yields the first estimate in (i) by proceeding in the same fashion as in (4.2). \square

Proposition 4.3. *For the Lagrange multipliers we have*

$$\sup_{t \in [0, T]} |\lambda_{\theta_1}| \leq C, \quad \sup_{t \in [0, T]} |\lambda_{\theta_2}| \leq C, \quad \sup_{t \in [0, T]} |\lambda_\rho| \leq C.$$

Proof. Because of Proposition 4.2, Lemma 4.1 can be applied with $K = C$, only depending on the model parameters and the initial datum. Hence, for the Lagrange multipliers λ_{θ_1} and λ_{θ_2} by Remark 2.2 we have

$$\left| \begin{pmatrix} \lambda_{\theta_1} \\ \lambda_{\theta_2} \end{pmatrix} \right| \leq C \sup_J \beta \left(\int_0^L (\partial_s \theta)^2 ds + \int_0^L |c_0 \partial_s \theta| ds \right),$$

where J is as in (4.3). The right hand side is bounded for all $t \in [0, T]$ by Proposition 4.2 and the Cauchy-Schwarz inequality. This way, we obtain the desired bounds for λ_{θ_1} and λ_{θ_2} . For λ_ρ we use (2.4) and proceed similarly, using that $\sup_J |\beta'| \leq C$ by continuity. \square

As a next step, we would like to bound the L^2 -norm of $\partial_s^2(\theta, \rho)$ uniformly in time. Directly pursuing this idea, we encounter difficulties in controlling the nonlinear coupling of the system (1.5), which seems to be incompatible with a direct application of interpolation inequalities. Instead, we control the L^2 -norm of the velocity $\partial_t(\theta, \rho)$ first.

Proposition 4.4. *We have*

$$\sup_{t \in [0, T]} \|\partial_t \theta\|_{L^2(0, L)} \leq C, \quad \sup_{t \in [0, T]} \|\partial_t \rho\|_{L^2(0, L)} \leq C.$$

Proof. By continuity, Proposition 4.2 and with J as in (4.3) we have that

$$(4.5) \quad \sup_J |\beta'| \leq C, \quad \sup_J |\beta''| \leq C.$$

We consider the smooth function

$$\varphi(t) := \frac{1}{2} \int_0^L ((\partial_t \theta)^2 + (\partial_t \rho)^2) ds$$

for $t \in [0, T]$. We now use (1.5) and integration by parts to differentiate φ . The boundary terms disappear as a consequence of the boundary conditions in (1.5). We have

$$\begin{aligned} \frac{d}{dt}\varphi(t) &= \int_0^L \partial_t \theta \partial_t \left[\partial_s (\beta(\rho)(\partial_s \theta - c_0)) + \lambda_{\theta 1} \sin \theta - \lambda_{\theta 2} \cos \theta \right] ds \\ &\quad + \int_0^L \partial_t \rho \partial_t \left(\mu \partial_s^2 \rho - \frac{1}{2} \beta'(\rho)(\partial_s \theta - c_0)^2 - \lambda_\rho \right) ds \\ &= - \int_0^L \beta(\rho)(\partial_t \partial_s \theta)^2 ds - \int_0^L \mu (\partial_t \partial_s \rho)^2 ds \\ &\quad - 2 \int_0^L \partial_t \partial_s \theta \beta'(\rho) \partial_t \rho (\partial_s \theta - c_0) ds - \frac{1}{2} \int_0^L (\partial_t \rho)^2 \beta''(\rho) (\partial_s \theta - c_0)^2 ds \\ &\quad + \int_0^L (\partial_t \theta)^2 \lambda_{\theta 1} \cos \theta ds + \int_0^L (\partial_t \theta)^2 \lambda_{\theta 2} \sin \theta ds. \end{aligned}$$

Here no time derivatives of the Lagrange multipliers appear by (2.3) and (2.5). Using (4.5), Young's inequality, Proposition 4.2 and Proposition 4.3, we can estimate this by

$$\begin{aligned} \frac{d}{dt}\varphi(t) &\leq - \inf_J \beta \int_0^L (\partial_t \partial_s \theta)^2 ds - \mu \int_0^L (\partial_t \partial_s \rho)^2 ds + \delta \int_0^L (\partial_t \partial_s \theta)^2 ds \\ (4.6) \quad &\quad + C(\delta) \sup_{s \in [0, L]} (\partial_t \rho)^2 + C \int_0^L (\partial_t \theta)^2 ds \end{aligned}$$

for $\delta > 0$ to be chosen and some correspondingly adjusted $C(\delta) > 0$. Using (2.5) and arguing as in (4.2), we find there exists $s_t \in [0, L]$ with $\partial_t \rho(t, s_t) = 0$ and thus

$$\begin{aligned} \sup_{s \in [0, L]} |\partial_t \rho|^2 &= \sup_{s \in [0, L]} |(\partial_t \rho)^2 - (\partial_t \rho(t, s_t))^2| \leq \int_0^L \left| \partial_s (\partial_t \rho)^2 \right| ds \\ (4.7) \quad &\leq 2 \|\partial_t \rho\|_{L^2(0, L)} \|\partial_t \partial_s \rho\|_{L^2(0, L)} \leq \delta \|\partial_t \partial_s \rho\|_{L^2(0, L)}^2 + C(\delta) \|\partial_t \rho\|_{L^2(0, L)}^2, \end{aligned}$$

using Young's inequality in the last step. Using (4.4) and choosing $\delta > 0$ small enough in (4.6) and (4.7), we obtain

$$(4.8) \quad \frac{d}{dt}\varphi(t) \leq C\varphi(t) \quad \text{for all } t \in [0, T].$$

By Proposition 4.2, we find that $\int_0^T \varphi(t) dt$ is bounded independently of T and thus the claim follows from integrating (4.8). \square

Proposition 4.5. *We have*

$$\sup_{t \in [0, T]} \|\partial_s^2 \theta\|_{L^2(0, L)} \leq C, \quad \sup_{t \in [0, T]} \|\partial_s^2 \rho\|_{L^2(0, L)} \leq C.$$

Proof. To prove the required estimates, we need to consider simultaneously the evolution equation of θ and the evolution equation of ρ . Recall from (1.5) that we have

$$\begin{aligned} \beta(\rho) \partial_s^2 \theta &= \partial_t \theta - \beta'(\rho) \partial_s \rho (\partial_s \theta - c_0) - \lambda_{\theta 1} \sin \theta + \lambda_{\theta 2} \cos \theta, \\ \mu \partial_s^2 \rho &= \partial_t \rho + \frac{1}{2} \beta'(\rho) (\partial_s \theta - c_0)^2 + \lambda_\rho \end{aligned}$$

on $[0, T)$. Using (4.4) and arguing as in (4.5), by Young's inequality we find

$$\begin{aligned} & \|\partial_s^2 \theta\|_{L^2(0,L)}^2 + \|\partial_s^2 \rho\|_{L^2(0,L)}^2 \\ & \leq C \left(\int_0^L (\partial_t \theta)^2 ds + \int_0^L (\partial_s \rho)^2 (\partial_s \theta - c_0)^2 ds + \lambda_{\theta 1}^2 \int_0^L \sin^2 \theta ds + \lambda_{\theta 2}^2 \int_0^L \cos^2 \theta ds \right. \\ & \quad \left. + \int_0^L (\partial_t \rho)^2 ds + \int_0^L (\partial_s \theta - c_0)^4 ds + \lambda_\rho^2 L \right) \end{aligned}$$

for all $t \in [0, T)$. By Propositions 4.3 and 4.4 it is sufficient to consider the second term and the sixth term on the right side of the inequality. Using Hölder's inequality, interpolation [4, p. 233] and $\int_0^L \partial_s \rho ds = 0$, we find

$$\begin{aligned} & \|(\partial_s \rho)^2 (\partial_s \theta)^2\|_{L^1(0,L)} \leq \|\partial_s \rho\|_{L^4(0,L)}^2 \|\partial_s \theta\|_{L^4(0,L)}^2 \\ & \leq C \|\partial_s \rho\|_{L^2(0,L)}^{\frac{1}{2}} \|\partial_s \rho\|_{L^2(0,L)}^{\frac{3}{2}} \left(\|\partial_s^2 \theta\|_{L^2(0,L)}^{\frac{1}{4}} \|\partial_s \theta\|_{L^2(0,L)}^{\frac{3}{4}} + \|\partial_s \theta\|_{L^2(0,L)} \right)^2. \end{aligned}$$

Due to Proposition 4.2 the norm $\|\partial_s \theta\|_{L^2(0,L)}$ is bounded for all $t \in [0, T)$ and similarly for ρ . Therefore, for $\delta > 0$ to be chosen, using twice Young's inequality we estimate

$$\|(\partial_s \rho)^2 (\partial_s \theta)^2\|_{L^1(0,L)} \leq \delta \|\partial_s^2 \rho\|_{L^2(0,L)}^2 + \delta \|\partial_s^2 \theta\|_{L^2(0,L)}^2 + C(\delta).$$

In an analogous way we also get

$$\|(\partial_s \theta)^4\|_{L^1(0,L)} \leq \delta \|\partial_s^2 \theta\|_{L^2(0,L)}^2 + C(\delta).$$

Finally, taking $\delta > 0$ sufficiently small and absorbing, the claim follows. \square

Proposition 4.6. *We have*

$$\sup_{t \in [0, T)} \|\partial_s^3 \theta\|_{L^2(0,L)} \leq C, \quad \sup_{t \in [0, T)} \|\partial_s^3 \rho\|_{L^2(0,L)} \leq C.$$

Proof. Again, we use a Gronwall argument. To that end, we define

$$\varphi(t) := \frac{1}{2} \int_0^L ((\partial_s^3 \theta)^2 + (\partial_s^3 \rho)^2) ds$$

for $t \in [0, T)$ and obtain after integration by parts

$$(4.9) \quad \frac{d}{dt} \varphi(t) + \varphi(t) = - \int_0^L \partial_s^4 \theta \partial_t \partial_s^2 \theta + \partial_s^4 \rho \partial_t \partial_s^2 \rho ds - \frac{1}{2} \int_0^L \partial_s^4 \theta \partial_s^2 \theta + \partial_s^4 \rho \partial_s^2 \rho ds.$$

Differentiating twice the evolution equations (1.5) with respect to s yields

$$\begin{aligned} \partial_s^2 \partial_t \theta &= 3\beta'(\rho) \partial_s \rho \partial_s^3 \theta + \beta(\rho) \partial_s^4 \theta + 3\beta''(\rho) (\partial_s \rho)^2 \partial_s^2 \theta + 3\beta'(\rho) \partial_s^2 \rho \partial_s^2 \theta \\ & \quad + \beta'''(\rho) (\partial_s \rho)^3 (\partial_s \theta - c_0) + 3\beta''(\rho) \partial_s^2 \rho \partial_s \rho (\partial_s \theta - c_0) + \beta'(\rho) \partial_s^3 \rho (\partial_s \theta - c_0) \\ & \quad - \lambda_{\theta 1} \sin \theta (\partial_s \theta)^2 + \lambda_{\theta 1} \cos \theta \partial_s^2 \theta + \lambda_{\theta 2} \cos \theta (\partial_s \theta)^2 + \lambda_{\theta 2} \sin \theta \partial_s^2 \theta \end{aligned}$$

and

$$\begin{aligned} \partial_s^2 \partial_t \rho &= \mu \partial_s^4 \rho - \frac{1}{2} \beta'''(\rho) (\partial_s \rho)^2 (\partial_s \theta - c_0)^2 - \frac{1}{2} \beta''(\rho) \partial_s^2 \rho (\partial_s \theta - c_0)^2 \\ & \quad - 2\beta''(\rho) \partial_s \rho (\partial_s \theta - c_0) \partial_s^2 \theta - \beta'(\rho) (\partial_s^2 \theta)^2 - \beta'(\rho) (\partial_s \theta - c_0) \partial_s^3 \theta. \end{aligned}$$

By continuity, Proposition 4.2 and (4.3), we find that $|\beta^{(k)}(\rho)|$ is bounded by C for $k = 0, \dots, 3$. Moreover, due to the embedding $W^{2,2}(0, L) \hookrightarrow C^1([0, L])$, Proposition 4.2 and Proposition 4.5 we know that $\sup_{s \in [0, L]} |\partial_s \theta|$ is bounded by C for all $t \in$

$[0, T)$. The same applies to $\sup_{s \in [0, L]} |\partial_s \rho|$. This implies that after plugging the last two equations into (4.9) we get

$$(4.10) \quad \frac{d}{dt} \varphi(t) + \varphi(t) \leq - \inf_J \beta(\rho) \int_0^L (\partial_s^4 \theta)^2 ds - \mu \int_0^L (\partial_s^4 \rho)^2 ds + CR(t),$$

where $R(t)$ is a sum of terms of the form

$$(4.11) \quad \int_0^L |\partial_s^4 f| |\partial_s^3 g| ds, \int_0^L |\partial_s^4 f| |\partial_s^2 g| ds, \int_0^L |\partial_s^4 f| ds, \int_0^L |\partial_s^4 f| |\partial_s^2 f| |\partial_s^2 g| ds$$

for $f, g \in \{\theta, \rho\}$. For the first term, we use Young's inequality, integration by parts and Proposition 4.5 and estimate

$$\begin{aligned} \int_0^L |\partial_s^4 f| |\partial_s^3 g| ds &\leq \delta \int_0^L (\partial_s^4 f)^2 ds + C(\delta) \int_0^L (\partial_s^3 g)^2 ds \\ &= \delta \int_0^L (\partial_s^4 f)^2 ds - C(\delta) \int_0^L \partial_s^4 g \partial_s^2 g ds \\ &\leq \delta \int_0^L (\partial_s^4 f)^2 ds + \delta \int_0^L (\partial_s^4 g)^2 ds + C(\delta). \end{aligned}$$

Similarly, for terms of the second and third type we have

$$\int_0^L (|\partial_s^4 f| |\partial_s^2 g| + |\partial_s^4 f|) ds \leq \delta \int_0^L (\partial_s^4 f)^2 ds + C(\delta).$$

The terms of the fourth type in (4.11) can be controlled by interpolation, cf. [4, p. 233] (using $\int_0^L \partial_s^k f ds = 0$ for $k \geq 2$, $f \in \{\theta, \rho\}$) to estimate

$$\begin{aligned} \int_0^L |\partial_s^4 f| |\partial_s^2 f| |\partial_s^2 g| ds &\leq \|\partial_s^4 f\|_{L^2(0, L)} \|\partial_s^2 f\|_{L^4(0, L)} \|\partial_s^2 g\|_{L^4(0, L)} \\ &\leq C \|\partial_s^4 f\|_{L^2(0, L)} \|\partial_s^4 f\|_{L^2(0, L)}^{\frac{1}{8}} \|\partial_s^2 f\|_{L^2(0, L)}^{\frac{7}{8}} \|\partial_s^4 g\|_{L^2(0, L)}^{\frac{1}{8}} \|\partial_s^2 g\|_{L^2(0, L)}^{\frac{7}{8}} \\ &\leq \delta \|\partial_s^4 f\|_{L^2(0, L)}^2 + \delta \|\partial_s^4 g\|_{L^2(0, L)}^2 + C(\delta) \end{aligned}$$

as above by Proposition 4.5 and Young's inequality for $\delta > 0$ to be chosen. This way, we have estimated the remainder term $R(t)$ in (4.10). Consequently, if we now use $\inf_J \beta \geq C$ by (4.4) and choose $\delta > 0$ sufficiently small, after absorbing we get

$$\frac{d}{dt} \varphi(t) + \varphi(t) \leq C.$$

With Gronwall's lemma, we conclude that $\varphi(t) \leq C$ for all $t \in [0, T)$. \square

4.3. Proof of Theorem 1.3. Using the previous propositions, we are now able to prove long-time existence.

Proof of Theorem 1.3. Let $0 < T_{\max} \leq \infty$ be the maximal existence time of the smooth solution of (1.5) with initial datum $(\theta, \rho)(0, \cdot) = (\theta_0, \rho_0)$ in $C^{1+\alpha}([0, L])$. We first want to show that the assumption $T_{\max} < \infty$ yields a contradiction as we can extend the solution past T_{\max} .

Assume $T_{\max} < \infty$ and let $\delta \in (0, T_{\max})$. We have $(\theta, \rho) \in C^\infty([\delta, T_{\max}) \times [0, L])$. By Propositions 4.2 and 4.6, there is a constant $C > 0$ depending only on $(\theta, \rho)(\delta, \cdot)$ and the model parameters such that

$$(4.12) \quad \sup_{t \in [\delta, T_{\max})} \|\theta\|_{W^{3,2}(0, L)} \leq C \quad \text{and} \quad \sup_{t \in [\delta, T_{\max})} \|\rho\|_{W^{3,2}(0, L)} \leq C.$$

We consider an arbitrary sequence $(t_n)_{n \in \mathbb{N}}$ with $t_n \nearrow T_{\max}$. Let $\tilde{\alpha} \in (0, \frac{1}{2})$. The bounds in (4.12) together with the compact embedding $W^{3,2}(0, L) \hookrightarrow C^{2+\tilde{\alpha}}([0, L])$

and the closedness of $h^{2+\bar{\alpha}}([0, L])$ yield the existence of $\tilde{\theta}_0 \in h^{2+\bar{\alpha}}([0, L])$ and $\tilde{\rho}_0 \in h^{2+\bar{\alpha}}([0, L])$ such that after passing to a subsequence

$$(4.13) \quad \theta(t_n, \cdot) \rightarrow \tilde{\theta}_0 \quad \text{and} \quad \rho(t_n, \cdot) \rightarrow \tilde{\rho}_0 \quad \text{in } C^{2+\bar{\alpha}}([0, L]) \quad \text{as } n \rightarrow \infty.$$

Suppose $t'_n \nearrow T_{\max}$ denotes another sequence such that (4.13) holds with t'_n instead of t_n and $(\tilde{\theta}'_0, \tilde{\rho}'_0)$ instead of $(\tilde{\theta}_0, \tilde{\rho}_0)$. Using (1.5), Propositions 4.2, 4.3 and (4.12), we find $\|\partial_t \theta\|_{W^{1,2}(0,L)} \leq C$ for $t \in [\delta, T_{\max})$, where $C > 0$ again only depends on the model parameters and $(\theta, \rho)(\delta, \cdot)$. Hence by the Sobolev embedding $W^{1,2}(0, L) \hookrightarrow C^{\bar{\alpha}}([0, L])$, we find

$$(4.14) \quad \|\theta(t_n, \cdot) - \theta(t'_n, \cdot)\|_{C^{\bar{\alpha}}([0,L])} \leq \left\| \int_{t_n}^{t'_n} \partial_t \theta(\tau, \cdot) \, d\tau \right\|_{C^{\bar{\alpha}}([0,L])} \leq C |t_n - t'_n|,$$

after suitably modifying $C > 0$. Taking $n \rightarrow \infty$ yields $\tilde{\theta}'_0 = \tilde{\theta}_0$. With exactly the same argument we find $\tilde{\rho}'_0 = \tilde{\rho}_0$. Consequently

$$(4.15) \quad \theta(t, \cdot) \rightarrow \tilde{\theta}_0 \quad \text{and} \quad \rho(t, \cdot) \rightarrow \tilde{\rho}_0 \quad \text{in } C^{2+\bar{\alpha}}([0, L]) \quad \text{as } t \nearrow T_{\max}.$$

It is clear that $(\tilde{\theta}_0, \tilde{\rho}_0)$ satisfies (1.7). Thus, Theorem 1.2 and Remark 3.7 yield the existence of $\tilde{T} > 0$ and a unique solution $(\tilde{\theta}, \tilde{\rho}) \in C^\infty((0, \tilde{T}) \times [0, L])$ of (1.5) which satisfies

$$(4.16) \quad \tilde{\theta}(t, \cdot) \rightarrow \tilde{\theta}_0 \quad \text{and} \quad \tilde{\rho}(t, \cdot) \rightarrow \tilde{\rho}_0 \quad \text{in } C^{2+\bar{\alpha}}([0, L]) \quad \text{as } t \searrow 0.$$

We take this solution to extend the solution (θ, ρ) past T_{\max} by defining

$$(\bar{\theta}, \bar{\rho})(t, s) := \begin{cases} (\theta, \rho)(t, s) & \text{for } t \in (0, T_{\max}), \\ (\tilde{\theta}_0, \tilde{\rho}_0)(s) & \text{for } t = T_{\max}, \\ (\tilde{\theta}, \tilde{\rho})(t - T_{\max}, s) & \text{for } t \in (T_{\max}, T_{\max} + \tilde{T}). \end{cases}$$

We now claim that

$$(4.17) \quad (\bar{\theta}, \bar{\rho}) \in C^\infty((0, T_{\max} + \tilde{T}) \times [0, L]).$$

Fix $T \in (T_{\max}, T_{\max} + \tilde{T})$. Examining $t \nearrow T_{\max}$ and using (4.15), respectively $t \searrow T_{\max}$ and using (4.16), we conclude

$$(\bar{\theta}, \bar{\rho}) \in \text{BUC}([\delta, T]; C^{2+\bar{\alpha}}([0, L])).$$

By (1.5), we may write $\partial_t(\theta, \rho)$ in terms of $\partial_s^2(\theta, \rho)$, $\partial_s(\theta, \rho)$ and (θ, ρ) on $(0, T_{\max})$ and similarly for $\partial_t(\tilde{\theta}, \tilde{\rho})$ on $(0, \tilde{T})$. Hence $\partial_t(\bar{\theta}, \bar{\rho})$ exists on $[\delta, T_{\max}) \cup (T_{\max}, T]$ and possesses a continuous extension at $t = T_{\max}$, so that we find

$$\partial_t(\bar{\theta}, \bar{\rho}) \in \text{BUC}([\delta, T]; C^{\bar{\alpha}}([0, L])).$$

Consequently, we obtain

$$(4.18) \quad (\bar{\theta}, \bar{\rho}) \in \text{BUC}^1([\delta, T]; C^{\bar{\alpha}}([0, L])) \cap \text{BUC}([\delta, T]; C^{2+\bar{\alpha}}([0, L])).$$

By (3.20), for every $T' < T_{\max}$ the solution (θ, ρ) satisfies

$$(4.19) \quad (\theta, \rho) \in \text{BUC}_{1-\bar{\eta}}^1([0, T']; h^{\bar{\alpha}}([0, L])) \cap \text{BUC}_{1-\bar{\eta}}([0, T']; h^{2+\bar{\alpha}}([0, L])),$$

where $\bar{\alpha} \in (0, 1)$ and $\bar{\eta} \in (\frac{1}{2}, 1)$ are chosen such that $2\bar{\eta} + \bar{\alpha} = 1 + \alpha$. Since the time weight $t^{1-\bar{\eta}}$ only plays a role near $t = 0$, from (4.18) and (4.19) we conclude that

$$(\bar{\theta}, \bar{\rho}) \in \text{BUC}_{1-\bar{\eta}}^1([0, T]; h^{\bar{\alpha}}([0, L])) \cap \text{BUC}_{1-\bar{\eta}}([0, T]; h^{2+\bar{\alpha}}([0, L])),$$

for $0 < \hat{\alpha} < \min\{\bar{\alpha}, \bar{\alpha}\}$, using Lemma A.1. If we transfer this back to the periodic setting as in Section 3.1, we may apply Lemma 3.5 to deduce that $(\bar{\theta}, \bar{\rho}) \in C^\infty((0, T) \times [0, L])$. Since $T \in (T_{\max}, T_{\max} + \tilde{T})$ was arbitrary, we conclude that (4.17) is satisfied.

Consequently, we have extended the solution (θ, ρ) smoothly past T_{\max} , a contradiction, so $T_{\max} = \infty$ has to hold.

For the subconvergence part, let $(t_n)_{n \in \mathbb{N}}$ be any sequence with $t_n \rightarrow \infty$. Note that the constant C in (4.12) does not depend on the existence time, so (4.12) remains valid for $T_{\max} = \infty$. The compact embedding $W^{3,2}(0, L) \hookrightarrow C^{2+\tilde{\alpha}}([0, L])$, where $\tilde{\alpha} \in (0, \frac{1}{2})$, and a subsequence argument yields that there exists $(t'_n)_{n \in \mathbb{N}}$, a subsequence of $(t_n)_{n \in \mathbb{N}}$, and $(\theta_\infty, \rho_\infty)$ such that

$$(4.20) \quad \lim_{n \rightarrow \infty} (\theta(t'_n), \rho(t'_n)) = (\theta_\infty, \rho_\infty) \quad \text{in } C^{2+\tilde{\alpha}}([0, L])$$

for all $\tilde{\alpha} \in (0, \frac{1}{2})$.

It remains to prove that $(\theta_\infty, \rho_\infty)$ is stationary. We consider the function

$$\varphi(t) = \frac{1}{2} \int_0^L ((\partial_t \theta)^2 + (\partial_t \rho)^2) \, ds, \quad t \in [\delta, \infty),$$

which is bounded by Proposition 4.4, and thus by (4.8) its derivative $\frac{d}{dt} \varphi$ is bounded from above. Since φ is integrable on $[\delta, \infty)$ by Proposition 4.2, this is sufficient to guarantee that $\lim_{t \rightarrow \infty} \varphi(t) = 0$. In particular by (4.20), we have $\partial_t \theta(t'_n, \cdot) \rightarrow 0$, $\partial_t \rho(t'_n, \cdot) \rightarrow 0$ in $C^{\tilde{\alpha}}([0, L])$ for all $\tilde{\alpha} \in (0, \frac{1}{2})$, so $(\theta_\infty, \rho_\infty)$ is stationary, thus a solution to (1.8). \square

Remark 4.7. Instead of only bounding the $W^{3,2}$ -norm of a solution globally in time (cf. Proposition 4.6), it is possible to bound $\sup_{t \in [0, T]} \|(\theta, \rho)\|_{W^{m,2}(0, L)} \leq C(m)$, $m \in \mathbb{N}$, for a smooth solution by essentially the same arguments as in Proposition 4.6. This way, one can deduce that the subconvergence in (4.20) is even smooth.

In general, the limit $(\theta_\infty, \rho_\infty)$ in Theorem 1.3 might depend on the choice of the (sub)sequence and the argument in (4.14) does not transfer to the case $T_{\max} = \infty$.

APPENDIX A. FUNCTION SPACES

Already in the introduction in Theorem 1.2 little Hölder functions appear. Later in Chapter 3 we work with time-weighted little Hölder spaces and in the proof of Lemma 3.5 additionally the parabolic Hölder spaces are used. Here we collect the definitions of these spaces and list some properties.

A.1. Little Hölder functions. For $\alpha \in \mathbb{R}_+ \setminus \mathbb{Z}$, we denote by $[\alpha]$ the largest integer less than α and $\{\alpha\} := \alpha - [\alpha] \in (0, 1)$. Let $k \in \mathbb{N}$. The space of L -periodic \mathbb{R}^k -valued functions over \mathbb{R} which are continuous and $[\alpha]$ -times continuously differentiable is denoted by $C_L^{[\alpha]}(\mathbb{R}; \mathbb{R}^k)$. The *periodic Hölder space with exponent α* is defined as

$$C_L^\alpha(\mathbb{R}; \mathbb{R}^k) := \left\{ f \in C_L^{[\alpha]}(\mathbb{R}; \mathbb{R}^k) : [f^{([\alpha])}]_{\alpha, \mathbb{R}} := \sup_{\substack{x, y \in \mathbb{R}, \\ x \neq y}} \frac{|f^{([\alpha])}(x) - f^{([\alpha])}(y)|}{|x - y|^{\{\alpha\}}} < \infty \right\}$$

and equipped with the norm $\|f\|_{C^\alpha(\mathbb{R})} := \|f\|_{C^{[\alpha]}(\mathbb{R})} + [f^{([\alpha])}]_{\alpha, \mathbb{R}}$. Therewith, we define the *periodic little-Hölder space with exponent α* over \mathbb{R} by

$$(A.1) \quad h_L^\alpha(\mathbb{R}; \mathbb{R}^k) := \left\{ f \in C_L^\alpha(\mathbb{R}; \mathbb{R}^k) : \lim_{\delta \rightarrow 0} \sup_{\substack{x, y \in \mathbb{R}, \\ 0 < |x - y| < \delta}} \frac{|f^{([\alpha])}(x) - f^{([\alpha])}(y)|}{|x - y|^{\{\alpha\}}} = 0 \right\}.$$

This is a closed subspace of $C_L^\alpha(\mathbb{R}; \mathbb{R}^k)$ and therefore a Banach space when equipped with the same norm.

In the following we collect some properties of periodic little Hölder functions. This type of functions plays a fundamental role in the classical maximal regularity theory, see for example [13].

Lemma A.1. [12, Prop. 1.2] *Let $\alpha_1, \alpha_2 \in \mathbb{R}_+ \setminus \mathbb{Z}$ such that $\alpha_1 < \alpha_2$. Then $h_L^{\alpha_1}(\mathbb{R})$ is the closure of $C_L^{\alpha_2}(\mathbb{R})$ in $(C_L^{\alpha_1}(\mathbb{R}), \|\cdot\|_{C^{\alpha_1}})$ and $h_L^{\alpha_2}(\mathbb{R}) \hookrightarrow h_L^{\alpha_1}(\mathbb{R})$ densely.*

Lemma A.2 (Product of periodic little Hölder functions). *Let $\alpha_1, \alpha_2 \in \mathbb{R}_+ \setminus \mathbb{Z}$. Let $g_1 \in h_L^{\alpha_1}(\mathbb{R})$ and let $g_2 \in h_L^{\alpha_2}(\mathbb{R})$. Then*

$$g_1 g_2 \in h_{per, L}^{\min\{\alpha_1, \alpha_2\}}(\mathbb{R}) \quad \text{and} \quad \|g_1 g_2\|_{C^{\min\{\alpha_1, \alpha_2\}}} \leq C(\alpha_1, \alpha_2, L) \|g_1\|_{C^{\alpha_1}} \|g_2\|_{C^{\alpha_2}}.$$

Proof. This follows from (A.1) and a tedious computation. \square

A.2. Time-weighted BUC-spaces. Let E be an arbitrary Banach space. Let $\eta \in (0, 1)$ and let $T > 0$. We define

$$\text{BUC}_{1-\eta}([0, T]; E) := \left\{ \xi \in C([0, T]; E) : [t \mapsto t^{1-\eta} \xi(t)] \text{ bounded and uniformly continuous on } (0, T] \text{ with } \lim_{t \rightarrow 0^+} t^{1-\eta} \|\xi(t)\|_E = 0 \right\}.$$

Moreover, we define the subspace

$$\text{BUC}_{1-\eta}^1([0, T]; E) := \left\{ \xi \in C^1([0, T]; E) : \xi, \partial_t \xi \in \text{BUC}_{1-\eta}([0, T]; E) \right\}.$$

In Section 3, we use this space with E a periodic little Hölder space. In particular, the spaces \mathbb{E}_0 and \mathbb{E}_1 , defined in (3.4) and (3.5) are Banach spaces with the norms

$$\begin{aligned} \|\xi\|_{\mathbb{E}_0} &= \|\xi\|_{\mathbb{E}_0([0, T])} := \sup_{t \in (0, T]} t^{1-\eta} \|\xi(t)\|_{C^\alpha(\mathbb{R})} \quad \text{and} \\ \|\xi\|_{\mathbb{E}_1} &= \|\xi\|_{\mathbb{E}_1([0, T])} := \sup_{t \in (0, T]} t^{1-\eta} \left(\|\partial_t \xi(t)\|_{C^\alpha(\mathbb{R})} + \|\xi(t)\|_{C^{2+\alpha}(\mathbb{R})} \right), \end{aligned}$$

respectively. We use that the mapping

$$\gamma: \mathbb{E}_1([0, T]) \rightarrow h_L^\alpha(\mathbb{R}; \mathbb{R}^k), \quad \xi(t, \cdot) \mapsto \xi(0, \cdot)$$

is well defined, linear and continuous. This follows from the observation that for $\xi \in \mathbb{E}_1([0, T])$ and $0 < s < t$

$$(A.2) \quad \xi(t) - \xi(s) = \int_s^t \partial_t \xi(\tau) \, d\tau,$$

and hence there exists $\lim_{t \rightarrow 0} \xi(t)$ in $h_L^\alpha(\mathbb{R})$, see [5, Remark 2.1]. We introduce the trace space $\gamma \mathbb{E}_1 := \text{Im}(\gamma)$. In particular, $\gamma \mathbb{E}_1$ does not depend on $T > 0$. In [12, Section 5.1] it is shown that up to equivalent norms

$$(A.3) \quad \gamma \mathbb{E}_1 = h_L^{2\eta+\alpha}(\mathbb{R}; \mathbb{R}^k)$$

provided $2\eta + \alpha \notin \mathbb{Z}$. For simplicity, we will mostly omit the index k .

We state here an important embedding result, yielding that if $\eta > \frac{1}{2}$ we have continuity in space and time of the first order s -derivative. This is essential in Section 3.

Lemma A.3. [5, Lemma 2.2] *Let $T > 0$. We have*

$$\mathbb{E}_1([0, T]) \hookrightarrow \text{BUC}([0, T]; h_L^{2\eta+\alpha}(\mathbb{R})).$$

In particular, this implies that for $\xi \in \mathbb{E}_1([0, T])$ there exists $\lim_{t \rightarrow 0} \xi(t)$ in $C_L^{2\eta+\alpha}(\mathbb{R})$ and $\sup_{t \in [0, T]} \|\xi\|_{C^{2\eta+\alpha}} \leq C(T) \|\xi\|_{\mathbb{E}_1} < \infty$.

Remark A.4. Lemma A.3 implies that if $\eta \geq \frac{1}{2}$, for $\xi \in \mathbb{E}_1([0, T])$, there exists $\lim_{t \rightarrow 0} \partial_s \xi(t)$ in $C_L^\alpha(\mathbb{R})$ and $\partial_s \xi \in \text{BUC}([0, T]; h_L^\alpha(\mathbb{R}))$.

Remark A.5. We observe that from the definition of the spaces $\mathbb{E}_0([0, T])$ and $\mathbb{E}_1([0, T])$ we can derive the following properties.

- (1) If $f \in \mathbb{E}_1([0, T])$, then $\partial_s f, \partial_s^2 f \in \mathbb{E}_0([0, T])$.
- (2) If $f_1 \in \mathbb{E}_1([0, T])$ and $f_2 \in \mathbb{E}_0([0, T])$, then $f_1 f_2 \in \mathbb{E}_0([0, T])$.

(3) If $f \in \mathbb{E}_1([0, T])$ and $g \in C^\infty(\mathbb{R})$, then $g \circ f \in \mathbb{E}_1([0, T])$.

Lemma A.6. *Let $T > 0$, $\eta \in (\frac{1}{2}, 1)$, $\xi \in \mathbb{E}_1([0, T])$. Then $\partial_s^2 \xi \in L^2([0, T] \times [0, L])$.*

Proof. Since the \mathbb{E}_1 -norm is finite, there exists $K > 0$ such that

$$\sup_{s \in \mathbb{R}} |\partial_s^2 \xi(t, s)| \leq K t^{\eta-1} \quad \text{for all } t \in [0, T].$$

It thus follows

$$\int_0^T \int_0^L (\partial_s^2 \xi)^2 ds dt \leq K^2 L \int_0^T t^{2\eta-2} dt = \frac{K^2 L}{2\eta-1} T^{2\eta-1} < \infty. \quad \square$$

A.3. Parabolic Hölder spaces. At the end of Section 3 we show that the solution of (3.2) from Proposition 3.3 is smooth. For this purpose we consider two additional function spaces. First, we set for $\alpha \in \mathbb{R}_+ \setminus \mathbb{Z}$, $\zeta \in [0, 1)$ and $T > 0$

$$\text{BUC}^\zeta([0, T]; h_L^\alpha(\mathbb{R})) := \left\{ \xi : [0, T] \rightarrow h_L^\alpha(\mathbb{R}) \text{ bounded and uniformly continuous} \right. \\ \left. \text{on } [0, T], \sup_{\substack{t_1, t_2 \in [0, T] \\ t_1 \neq t_2}} \frac{\|\xi(t_1, s) - \xi(t_2, s)\|_{C^\alpha}}{|t_1 - t_2|^\zeta} < \infty \right\}.$$

Moreover, we work with parabolic Hölder spaces, cf. [11]. Let $l \in \mathbb{R}_+ \setminus \mathbb{Z}$, $a, b \in \mathbb{R}$ with $a < b$, $T > 0$ and let $Q_T := (0, T) \times (a, b)$. The parabolic Hölder space $H^{\frac{l}{2}, l}(\overline{Q_T})$ is the space of all functions $f : \overline{Q_T} \rightarrow \mathbb{R}$ with continuous derivatives $\partial_t^m \partial_s^n f$ for $2m + n < l$ and finite norm

$$\|f\|_{H^{\frac{l}{2}, l}(\overline{Q_T})} := \sum_{j=0}^{\lfloor l \rfloor} \sum_{2m+n=j} \sup_{(t,s) \in Q_T} |\partial_t^m \partial_s^n f| \\ + \sum_{2m+n=\lfloor l \rfloor} \sup_{\substack{(t,s_1), (t,s_2) \in Q_T \\ s_1 \neq s_2}} \frac{|\partial_t^m \partial_s^n f(t, s_1) - \partial_t^m \partial_s^n f(t, s_2)|}{|s_1 - s_2|^{l-\lfloor l \rfloor}} \\ + \sum_{0 < l-2m-n < 2} \sup_{\substack{(t_1,s), (t_2,s) \in Q_T \\ t_1 \neq t_2}} \frac{|\partial_t^m \partial_s^n f(t_1, s) - \partial_t^m \partial_s^n f(t_2, s)|}{|t_1 - t_2|^{\frac{l-2m-n}{2}}}.$$

The following lemma shows how the space \mathbb{E}_1 relates to these parabolic Hölder spaces.

Lemma A.7. *Let $\alpha \in (0, 1)$ and $\eta \in [\frac{1}{2}, 1)$ such that $2\eta + \alpha \notin \mathbb{Z}$. Let $a, b \in \mathbb{R}$ with $a < b$ and let $T > 0$. Let $f \in \mathbb{E}_1([0, T])$. Then*

$$\partial_s f \in H^{\frac{\zeta}{2}, \zeta}([0, T] \times [a, b])$$

for $\zeta := \eta + \frac{\alpha}{2} - \frac{1}{2} \in (0, 1)$.

Proof. Since $0 < \frac{\eta}{2} - \frac{\alpha}{4} + \frac{1}{4} < \eta$, [5, Lemma 2.2 d)] together with [12, Prop. 1.2 b)] give

$$\mathbb{E}_1([0, T]) \hookrightarrow \text{BUC}^{\eta - (\frac{\eta}{2} - \frac{\alpha}{4} + \frac{1}{4})} \left([0, T]; h_L^{2(\frac{\eta}{2} - \frac{\alpha}{4} + \frac{1}{4}) + \alpha}(\mathbb{R}) \right).$$

With $\zeta = \eta + \frac{\alpha}{2} - \frac{1}{2} \in (0, 1)$, we thus have (in particular) $\partial_s f \in \text{BUC}^{\frac{\zeta}{2}}([0, T]; h_L^\zeta(\mathbb{R}))$. The continuity of $\partial_s f$ in $[0, T] \times [a, b]$ (remember $\eta \geq \frac{1}{2}$) and the finiteness of the

norm

$$\begin{aligned} \|\partial_s f\|_{H^{\frac{\zeta}{2}, \zeta}(\overline{Q_T})} &= \sup_{(t,s) \in Q_T} |\partial_s f| + \sup_{\substack{(t,s_1), (t,s_2) \in Q_T \\ s_1 \neq s_2}} \frac{|\partial_s f(t, s_1) - \partial_s f(t, s_2)|}{|s_1 - s_2|^\zeta} \\ &\quad + \sup_{\substack{(t_1,s), (t_2,s) \in Q_T \\ t_1 \neq t_2}} \frac{|\partial_s f(t_1, s) - \partial_s f(t_2, s)|}{|t_1 - t_2|^{\frac{\zeta}{2}}}. \end{aligned}$$

imply that $\partial_s f \in H^{\frac{\zeta}{2}, \zeta}([0, T] \times [a, b])$. \square

APPENDIX B. AUXILIARY RESULTS FOR SECTION 3.2

In the following, we show Step 2 of the proof of Proposition 3.3, i.e. that $D\Phi(\bar{u}, \bar{\rho})$ (see (3.10)) is a linear isomorphism. This is equivalent to show existence and uniqueness of a solution for a certain inhomogeneous linear initial value problem. In this section, $\alpha \in (0, 1)$ and $\eta \in (\frac{1}{2}, 1)$ are fixed.

B.1. The linearization. We compute here $D\mathbf{F}[\bar{u}, \bar{\rho}]$ needed in (3.10), where \mathbf{F} is given as in (3.6). By a direct calculation one finds

$$\begin{aligned} D\mathbf{F}_1(\bar{u}, \bar{\rho})(v, \sigma) &= \beta(\bar{\rho}) \partial_s^2 v + \beta'(\bar{\rho}) \partial_s^2 \bar{u} \sigma + \beta'(\bar{\rho}) \partial_s \bar{\rho} \partial_s v \\ &\quad + \beta'(\bar{\rho}) (\partial_s \bar{u} + \partial_s \phi - c_0) \partial_s \sigma + \beta''(\bar{\rho}) \partial_s \bar{\rho} (\partial_s \bar{u} + \partial_s \phi - c_0) \sigma \\ &\quad + D\Lambda_u(\bar{u}, \bar{\rho})(v, \sigma), \end{aligned} \tag{B.1}$$

$$\begin{aligned} D\mathbf{F}_2(\bar{u}, \bar{\rho})(v, \sigma) &= \mu \partial_s^2 \sigma - \frac{1}{2} \beta''(\bar{\rho}) (\partial_s \bar{u} + \partial_s \phi - c_0)^2 \sigma - \beta'(\bar{\rho}) (\partial_s \bar{u} + \partial_s \phi - c_0) \partial_s v \\ &\quad + D\Lambda_\rho(\bar{u}, \bar{\rho})(v, \sigma). \end{aligned} \tag{B.2}$$

Let $T > 0$. We first observe that by (3.3), in the Lagrange multipliers only spatial derivatives of first order appear. More specifically, Λ_u is well defined with $\Lambda_u(u, \rho) \in \mathbb{E}_0([0, T])$ for $(u, \rho) \in \mathbb{V}([0, T])$ where

$$\begin{aligned} \mathbb{V}([0, T]) &:= \left\{ (u, \rho) \in \mathbb{F}([0, T]) := \text{BUC}_{1-\eta}^1([0, T]; h_L^\alpha(\mathbb{R})) \cap \text{BUC}_{1-\eta}([0, T]; h_L^{1+\alpha}(\mathbb{R})) \right. \\ &\quad \left. \text{such that } \inf_{[0, T]} \det \Pi(u + \phi) > 0 \right\}. \end{aligned} \tag{B.3}$$

This is an open subset of $\mathbb{F}([0, T])$ by continuity. Moreover, using $\eta > \frac{1}{2}$ and (A.3), any initial datum $(v_0, \sigma_0) \in \gamma \mathbb{E}_1$ is in $\mathbb{F}([0, T])$ if we identify it with the function $(t, s) \mapsto (v_0(s), \sigma_0(s))$. It is not difficult to see that

$$\mathbb{V}([0, T]) \ni (u, \rho) \mapsto \Lambda_u(u, \rho) \in \mathbb{E}_0([0, T]) \tag{B.4}$$

is of class C^1 . Moreover, it follows from the product rule that

$$\begin{aligned} D\Lambda_u(\bar{u}, \bar{\rho})(v, \sigma) &= D\Pi^{-1}(\bar{u} + \phi)(v) \mathcal{J}(\bar{u}, \bar{\rho}) \cdot \begin{pmatrix} \sin(\bar{u} + \phi) \\ -\cos(\bar{u} + \phi) \end{pmatrix} \\ &\quad + \Pi^{-1}(\bar{u} + \phi) D\mathcal{J}(\bar{u}, \bar{\rho})(v, \sigma) \cdot \begin{pmatrix} \sin(\bar{u} + \phi) \\ -\cos(\bar{u} + \phi) \end{pmatrix} \\ &\quad + \Pi^{-1}(\bar{u} + \phi) \mathcal{J}(\bar{u}, \bar{\rho}) \cdot \begin{pmatrix} \cos(\bar{u} + \phi) \\ \sin(\bar{u} + \phi) \end{pmatrix} \end{aligned} \tag{B.5}$$

for all $(v, \sigma) \in \mathbb{F}([0, T])$. A similar argument yields that $\Lambda_\rho: \mathbb{V}([0, T]) \rightarrow \mathbb{E}_0([0, T])$ is well-defined and C^1 with derivative

$$D\Lambda_\rho(\bar{u}, \bar{\rho})(v, \sigma) = \frac{1}{2L} \int_0^L \beta''(\bar{\rho}) (\partial_s \bar{u} + \partial_s \phi - c_0)^2 \sigma + 2\beta'(\bar{\rho}) (\partial_s \bar{u} + \partial_s \phi - c_0) \partial_s v \, ds.$$

B.2. Linear evolution problem with time-independent coefficients. We consider an auxiliary linear evolution problem. Since we rely on [12], where time-independent coefficients are used, we would like to freeze the coefficients of the terms in (B.1) and (B.2) in $t = 0$. For the second term in (B.1), the regularity of the initial datum does not allow this. Therefore, we treat this term differently from the others and move it together with $\text{D}\Lambda_u$ and $\text{D}\Lambda_\rho$, which are nonlocal, to the right hand side of the inhomogeneous initial value problem we want to solve. This explains the structure of the differential operator chosen below.

Proposition B.1. *Let $(u_0, \rho_0) \in \gamma\mathbb{E}_1$ such that u_0 satisfies (3.1). Let $(\bar{u}, \bar{\rho}) \in \mathbb{E}_1([0, \bar{T}])$ be the function constructed in the proof of Proposition 3.3. Then for all $T \in (0, \bar{T}]$ the mapping*

$$\mathbf{J}: \mathbb{E}_1([0, T]) \rightarrow \mathbb{E}_0([0, T]) \times \gamma\mathbb{E}_1, \quad (v, \sigma) \mapsto (\partial_t(v, \sigma) - \mathcal{A}(v, \sigma) - \Psi, (v, \sigma)(0, \cdot))$$

where

$$\begin{aligned} \mathcal{A}(v, \sigma) &:= \left(\beta(\rho_0) \partial_s^2 v + \beta'(\rho_0) \partial_s \rho_0 \partial_s v + \beta'(\rho_0) (\partial_s u_0 + \partial_s \phi - c_0) \partial_s \sigma \right. \\ &\quad \left. + \beta''(\rho_0) \partial_s \rho_0 (\partial_s u_0 + \partial_s \phi - c_0) \sigma, \right. \\ &\quad \left. \mu \partial_s^2 \sigma - \frac{1}{2} \beta''(\rho_0) (\partial_s u_0 + \partial_s \phi - c_0)^2 \sigma - \beta'(\rho_0) (\partial_s u_0 + \partial_s \phi - c_0) \partial_s v \right), \\ \Psi &= \Psi(v_0, \sigma_0) := \left(\beta'(\bar{\rho}) \partial_s^2 \bar{u} \sigma_0 + \text{D}\Lambda_u(\bar{u}, \bar{\rho})(v_0, \sigma_0), \text{D}\Lambda_\rho(\bar{u}, \bar{\rho})(v_0, \sigma_0) \right) \end{aligned}$$

is well-defined and an isomorphism. Here $(v_0, \sigma_0) = (v, \sigma)(0, \cdot) \in \mathbb{F}([0, T])$, cf. (B.3). Equivalently¹, for any pair $(\varphi_1, \varphi_2) \in \mathbb{E}_0([0, T])$ and $(v_0, \sigma_0) \in \gamma\mathbb{E}_1$ there exists a unique solution $(u, \rho) \in \mathbb{E}_1([0, T])$ of the linear problem

$$(B.6) \quad \begin{cases} \partial_t(v, \sigma) - \mathcal{A}(v, \sigma) = (\varphi_1, \varphi_2) + \Psi & \text{in } (0, T) \times \mathbb{R}, \\ (v, \sigma)(0, \cdot) = (v_0, \sigma_0) & \text{on } \mathbb{R}. \end{cases}$$

Proof. We want to use [12], which works with functions on the one-dimensional torus, or equivalently for periodic functions, cf. [12, Proposition 1.1]. With (A.3) and Lemma A.2, the time-independent coefficients of the operator \mathcal{A} are in $h_L^\alpha(\mathbb{R})$. Because $\beta(\rho_0)$ is bounded away from zero and $\mu > 0$, \mathcal{A} is a uniformly elliptic operator. Apart from this we have to check that $\Psi \in \mathbb{E}_0([0, T])$. Using Lemma A.2 and Remark A.5, we find $\beta'(\bar{\rho}) \in \mathbb{E}_1([0, T])$ and thus $\beta'(\bar{\rho}) \partial_s^2 \bar{u} \sigma_0 \in \mathbb{E}_0([0, T])$. From (B.4), we conclude that $\text{D}\Lambda_u(\bar{u}, \bar{\rho})(v_0, \sigma_0) \in \mathbb{E}_0([0, T])$ and similarly for Λ_ρ . Applying [12, Theorem 6.4] yields a unique solution $(v, \sigma) \in \mathbb{E}_1([0, T])$ of (B.6). \square

From the fact that the mapping \mathbf{J} from Proposition B.1 is an isomorphism, an estimate of the norm of the solution follows implicitly. We now show that the constant in this estimate can be chosen independent of T .

Lemma B.2. *Let $(u_0, \rho_0) \in \gamma\mathbb{E}_1$ such that u_0 satisfies (3.1). Let $(\bar{u}, \bar{\rho}) \in \mathbb{E}_1([0, \bar{T}])$ be the map constructed in the proof of Proposition 3.3. Let $T \in (0, \bar{T}]$. Then the solution operator \mathbf{J}^{-1} satisfies the estimate*

$$\|\mathbf{J}^{-1}((\varphi_1, \varphi_2), (v_0, \sigma_0))\|_{\mathbb{E}_1([0, T])} \leq C \left(\|(\varphi_1, \varphi_2)\|_{\mathbb{E}_0([0, T])} + \|(v_0, \sigma_0)\|_{\gamma\mathbb{E}_1} \right)$$

for all $(\varphi_1, \varphi_2) \in \mathbb{E}_0([0, T])$ and $(v_0, \sigma_0) \in \gamma\mathbb{E}_1$ with $C = C(u_0, \rho_0, \bar{T})$.

Notice that here we do not write explicitly the dependence on T of the solution operator. This abuse of notation is justified in the proof below.

¹Here we slightly abuse notation, denoting by (v_0, σ_0) both the initial datum of the problem (B.6) and the evaluation at $t = 0$ of $(v, \sigma) \in \mathbb{E}_1([0, T])$. However, for solving (B.6), this does not make a difference.

Proof of Lemma B.2. Proposition B.1 already yields the claimed inequality with $C = C(u_0, \rho_0, T)$. We show that the constant C can be chosen only depending on $\bar{T} > T$ by an extension argument. Let $(v, \sigma) = \mathbf{J}^{-1}((\varphi_1, \varphi_2), (v_0, \sigma_0)) \in \mathbb{E}_1([0, T])$. Consider

$$(\varphi_1^*, \varphi_2^*)(t, s) = \begin{cases} (\varphi_1, \varphi_2)(t, s) & \text{for } (t, s) \in [0, T] \times \mathbb{R}, \\ (\varphi_1, \varphi_2)(T, s) & \text{for } (t, s) \in (T, \bar{T}] \times \mathbb{R}. \end{cases}$$

Then $(\varphi_1^*, \varphi_2^*) \in \mathbb{E}_0([0, \bar{T}])$ and we can apply Proposition B.1 and obtain $(v^*, \sigma^*) = \mathbf{J}^{-1}((\varphi_1^*, \varphi_2^*), (v_0, \sigma_0)) \in \mathbb{E}_1([0, \bar{T}])$, the solution of (B.6) on $(0, \bar{T}]$ with $(\varphi_1^*, \varphi_2^*)$ instead of (φ_1, φ_2) . The same proposition gives that \mathbf{J}^{-1} is an isomorphism from $\mathbb{E}_1([0, \bar{T}])$ to $\mathbb{E}_0([0, \bar{T}])$. Hence, there exists a constant $C = C(u_0, \rho_0, \bar{T})$ such that

$$\|(v^*, \sigma^*)\|_{\mathbb{E}_1([0, \bar{T}])} \leq C \left(\|(\varphi_1^*, \varphi_2^*)\|_{\mathbb{E}_0([0, \bar{T}])} + \|(v_0, \sigma_0)\|_{\gamma\mathbb{E}_1} \right).$$

Because of the uniqueness of the solution and using that (B.6) is local in time, we find $(v^*, \sigma^*)|_{[0, T]} = (v, \sigma)$. The claim follows using that $\|(v, \sigma)\|_{\mathbb{E}_1([0, T])} \leq \|(v^*, \sigma^*)\|_{\mathbb{E}_1([0, \bar{T}])}$, together with $\|(\varphi_1^*, \varphi_2^*)\|_{\mathbb{E}_0([0, \bar{T}])} = \|(\varphi_1, \varphi_2)\|_{\mathbb{E}_0([0, T])}$. \square

B.3. Some auxiliary estimates. We collect here some technical results, which we use to examine the differences appearing when comparing the linear operators in (3.10) and \mathbf{J} from Proposition B.1.

Throughout this subsection, we fix $(u_0, \rho_0) \in \gamma\mathbb{E}_1$ such that u_0 satisfies (3.1) and denote by $(\bar{u}, \bar{\rho}) \in \mathbb{E}_1([0, \bar{T}])$ the function constructed in the proof of Proposition 3.3. In particular, $\eta \in (\frac{1}{2}, 1)$, $\tilde{\eta} = \eta - \frac{1}{2}$ and $\partial_s(\bar{u}, \bar{\rho}) \in \tilde{\mathbb{E}}_1([0, \bar{T}])$.

Lemma B.3. *Let $t \in (0, \bar{T}]$. Then we have*

$$\|\bar{u}(t) - u_0\|_{C^\alpha} \leq \|\bar{u}\|_{\mathbb{E}_1} \frac{t^\eta}{\eta}, \quad \|\partial_s \bar{u}(t) - \partial_s u_0\|_{C^\alpha} \leq \|\partial_s \bar{u}\|_{\tilde{\mathbb{E}}_1} \frac{t^{\tilde{\eta}}}{\tilde{\eta}},$$

and similarly for $\bar{\rho}$.

Proof. Inspired by (A.2), we compute

$$\begin{aligned} \|\bar{u}(t) - u_0\|_{C^\alpha} &= \left\| \lim_{t' \rightarrow 0^+} \bar{u}(t) - \bar{u}(t') \right\|_{C^\alpha} = \left\| \lim_{t' \rightarrow 0^+} \int_{t'}^t \partial_\tau \bar{u}(\tau) \, d\tau \right\|_{C^\alpha} \\ &\leq \lim_{t' \rightarrow 0^+} \int_{t'}^t \|\tau^{\eta-1} \tau^{1-\eta} \partial_\tau \bar{u}(\tau)\|_{C^\alpha} \, d\tau \leq \|\bar{u}\|_{\mathbb{E}_1} \lim_{t' \rightarrow 0^+} \int_{t'}^t \tau^{\eta-1} \, d\tau = \|\bar{u}\|_{\mathbb{E}_1} \frac{t^\eta}{\eta}. \end{aligned}$$

For $\partial_s \bar{u}$, we proceed completely analogously. \square

In the sequel, we need to estimate terms of the form $\beta^{(i)}(\bar{\rho})$. Since we have $\bar{\rho} \in \text{BUC}([0, \bar{T}]; h_L^{2\eta+\alpha}(\mathbb{R}))$, see Lemma A.3, we find that there exists $J \subset \mathbb{R}$, a compact interval, such that $\bar{\rho}(t, s) \in J$ for all $(t, s) \in [0, \bar{T}] \times \mathbb{R}$. Since $\beta \in C^\infty(\mathbb{R})$, we find that there exist a constant $M \in (0, \infty)$ such that

$$(B.7) \quad \begin{aligned} \sup_{t \in [0, \bar{T}]} \|\Pi^{-1}(\bar{u} + \phi)\| &\leq M, \quad \sup_J |\beta^{(i)}| \leq M \quad \text{for } i = 0, \dots, 4, \\ \sup_{t \in [0, \bar{T}]} \|\bar{u}\|_{C^1}, \|\bar{u}\|_{\mathbb{E}_1([0, \bar{T}])}, \sup_{t \in [0, \bar{T}]} \|\bar{\rho}\|_{C^1} &\leq M. \end{aligned}$$

Lemma B.4. *Let $M > 0$ be as in (B.7). There is $C = C(M)$ such that for all $i \in \{0, 1, 2\}$ we have*

$$\|\beta^{(i)}(\rho_0) - \beta^{(i)}(\bar{\rho}(t))\|_{C^\alpha} \leq C(M) \|\bar{\rho}\|_{\mathbb{E}_1} \frac{t^\eta}{\eta} \quad \text{for all } t \in [0, \bar{T}].$$

Proof. For the supremum norm we find

$$(B.8) \quad \|\beta(\rho_0) - \beta(\bar{\rho})\|_{C^0} = \sup_{s \in \mathbb{R}} |\beta(\rho_0) - \beta(\bar{\rho})| \leq \sup_J |\beta'| \sup_{s \in \mathbb{R}} |\rho_0 - \bar{\rho}|.$$

To estimate the Hölder semi-norm, we note that for $s_1, s_2 \in \mathbb{R}$

$$\begin{aligned} & |\beta(\rho_0(s_1)) - \beta(\bar{\rho}(s_1)) - \beta(\rho_0(s_2)) + \beta(\bar{\rho}(s_2))| \\ &= \left| \int_0^1 \partial_\lambda [\beta(\lambda \rho_0(s_1) + (1-\lambda)\bar{\rho}(s_1)) - \beta(\lambda \rho_0(s_2) + (1-\lambda)\bar{\rho}(s_2))] \, d\lambda \right| \\ &\leq \int_0^1 |\beta'(\lambda \rho_0(s_1) + (1-\lambda)\bar{\rho}(s_1))| |\rho_0(s_1) - \bar{\rho}(s_1) - \rho_0(s_2) + \bar{\rho}(s_2)| \, d\lambda \\ &+ \int_0^1 |\beta'(\lambda \rho_0(s_1) + (1-\lambda)\bar{\rho}(s_1)) - \beta'(\lambda \rho_0(s_2) + (1-\lambda)\bar{\rho}(s_2))| |\rho_0(s_2) - \bar{\rho}(s_2)| \, d\lambda \\ &\leq \sup_J |\beta'| [\rho_0 - \bar{\rho}]_\alpha |s_1 - s_2|^\alpha \\ &\quad + \sup_J |\beta''| (|\rho_0(s_1) - \rho_0(s_2)| + |\bar{\rho}(s_1) - \bar{\rho}(s_2)|) |\rho_0(s_2) - \bar{\rho}(s_2)| \\ &\leq \left[\sup_J |\beta'| [\rho_0 - \bar{\rho}]_\alpha + \sup_J |\beta''| ([\rho_0]_\alpha + [\bar{\rho}]_\alpha) \sup_{s \in \mathbb{R}} |\rho_0 - \bar{\rho}| \right] |s_1 - s_2|^\alpha. \end{aligned}$$

This estimate together with (B.8) and Lemma B.3 yields the claim for $i = 0$. Since β is smooth, we can proceed in the same way for $i \geq 1$. \square

Lemma B.5. *Let $M > 0$ be as in (B.7). Then there exists a constant C , only depending on M , η , α and the model parameters, such that for all $T \in (0, \bar{T}]$, $(v, \sigma) \in \mathbb{E}_1([0, T])$ and $(v_0, \sigma_0) := (v, \sigma)(0, \cdot) \in \gamma\mathbb{E}_1$ we have*

$$(B.9) \quad \begin{aligned} \|\mathrm{DA}_u(\bar{u}, \bar{\rho})(v, \sigma) - \mathrm{DA}_u(\bar{u}, \bar{\rho})(v_0, \sigma_0)\|_{\mathbb{E}_0} &\leq C (\|v\|_{\mathbb{E}_1} + \|\sigma\|_{\mathbb{E}_1}) \max\{T, T^\eta\}, \\ \|\mathrm{DA}_\rho(\bar{u}, \bar{\rho})(v, \sigma) - \mathrm{DA}_\rho(\bar{u}, \bar{\rho})(v_0, \sigma_0)\|_{\mathbb{E}_0} &\leq C (\|v\|_{\mathbb{E}_1} + \|\sigma\|_{\mathbb{E}_1}) \max\{T, T^\eta\}. \end{aligned}$$

Here DA_u is the Fréchet derivative of the map defined in (B.4) and may thus be applied to (v_0, σ_0) , identified with an element of $\mathbb{F}([0, T])$, cf. (B.3). Similarly for DA_ρ .

Proof of Lemma B.5. In the following, C denotes a constant which varies from line to line but only depends on M and the model parameters. We start by proving (B.9) considering each term in (B.5) separately. Firstly, by (2.2) with $\theta = \bar{u} + \phi$ we find

$$(B.10) \quad \mathrm{DII}(\bar{u} + \phi)(v) = \begin{pmatrix} \int_0^L \sin(2(\bar{u} + \phi)) v \, ds & - \int_0^L \cos(2(\bar{u} + \phi)) v \, ds \\ - \int_0^L \cos(2(\bar{u} + \phi)) v \, ds & - \int_0^L \sin(2(\bar{u} + \phi)) v \, ds \end{pmatrix}$$

and obtain using (B.7) and (B.10) (considering the operator norm of the matrix)

$$\begin{aligned} & \|\mathrm{DII}^{-1}(\bar{u} + \phi)(v) - \mathrm{DII}^{-1}(\bar{u} + \phi)(v_0)\| \\ &= \|-\Pi^{-1}(\bar{u} + \phi)(\mathrm{DII}(\bar{u} + \phi)(v) - \mathrm{DII}(\bar{u} + \phi)(v_0))\Pi^{-1}(\bar{u} + \phi)\| \\ &\leq M^2 \sqrt{2}L \sup_{s \in [0, L]} |v - v_0| \leq C \|v - v_0\|_{C^\alpha}. \end{aligned}$$

Hence, with Lemma B.3 the difference coming from the first term on the right hand side of (B.5) can be estimated by

$$\begin{aligned} & \left\| (\mathrm{DII}^{-1}(\bar{u} + \phi)(v) - \mathrm{DII}^{-1}(\bar{u} + \phi)(v_0)) \mathcal{J}(\bar{u}, \bar{\rho}) \cdot \begin{pmatrix} \sin(\bar{u} + \phi) \\ -\cos(\bar{u} + \phi) \end{pmatrix} \right\|_{\mathbb{E}_0} \\ &\leq C \sup_{t \in (0, T]} t^{1-\eta} \|v - v_0\|_{C^\alpha} \leq C \|v\|_{\mathbb{E}_1} T. \end{aligned}$$

For the second term of (B.5) we compute

$$\begin{aligned}
\mathrm{D}\mathcal{J}(\bar{u}, \bar{\rho})(v, \sigma) &= \int_0^L \begin{pmatrix} -\sin(\bar{u} + \phi) \\ \cos(\bar{u} + \phi) \end{pmatrix} v (\partial_s \bar{u} + \partial_s \phi) \beta(\bar{\rho}) (\partial_s \bar{u} + \partial_s \phi - c_0) \, ds \\
&\quad + \int_0^L \begin{pmatrix} \cos(\bar{u} + \phi) \\ \sin(\bar{u} + \phi) \end{pmatrix} (\partial_s \bar{u} + \partial_s \phi) \beta'(\bar{\rho}) \sigma (\partial_s \bar{u} + \partial_s \phi - c_0) \, ds \\
&\quad - \int_0^L \begin{pmatrix} -\sin(\bar{u} + \phi) \\ \cos(\bar{u} + \phi) \end{pmatrix} (\partial_s \bar{u} + \partial_s \phi) (2\partial_s \bar{u} + 2\partial_s \phi - c_0) \beta(\bar{\rho}) v \, ds \\
&\quad - \int_0^L \begin{pmatrix} \cos(\bar{u} + \phi) \\ \sin(\bar{u} + \phi) \end{pmatrix} 2\partial_s^2 \bar{u} \beta(\bar{\rho}) v \, ds \\
&\quad - \int_0^L \begin{pmatrix} \cos(\bar{u} + \phi) \\ \sin(\bar{u} + \phi) \end{pmatrix} (2\partial_s \bar{u} + 2\partial_s \phi - c_0) \beta'(\bar{\rho}) \partial_s \bar{\rho} v \, ds.
\end{aligned}$$

where we have integrated once by parts. So the difference we need to consider can be estimated by

$$\begin{aligned}
&\| \mathrm{D}\mathcal{J}(\bar{u}, \bar{\rho})(v_0, \sigma_0) - \mathrm{D}\mathcal{J}(\bar{u}, \bar{\rho})(v, \sigma) \|_{C^0} \\
&\leq C (1 + \|\partial_s^2 \bar{u}\|_{C^\alpha}) \|v - v_0\|_{C^\alpha} + C \|\sigma - \sigma_0\|_{C^\alpha} \\
&\leq C t^\eta (1 + \|\partial_s^2 \bar{u}\|_{C^\alpha}) (\|v\|_{\mathbb{E}_1} + \|\sigma\|_{\mathbb{E}_1}).
\end{aligned}$$

In the last step we used Lemma B.3. Using (B.7), we obtain

$$\begin{aligned}
&\left\| \Pi^{-1}(\bar{u} + \phi) \left(\mathrm{D}\mathcal{J}(\bar{u}, \bar{\rho})(v, \sigma) - \mathrm{D}\mathcal{J}(\bar{u}, \bar{\rho})(v_0, \sigma_0) \right) \cdot \begin{pmatrix} \sin(\bar{u} + \phi) \\ -\cos(\bar{u} + \phi) \end{pmatrix} \right\|_{\mathbb{E}_0} \\
&\leq C \sup_{t \in (0, T]} t^{1-\eta} t^\eta (1 + \|\partial_s^2 \bar{u}\|_{C^\alpha}) (\|v\|_{\mathbb{E}_1} + \|\sigma\|_{\mathbb{E}_1}) \leq C (\|v\|_{\mathbb{E}_1} + \|\sigma\|_{\mathbb{E}_1}) (T + T^\eta).
\end{aligned}$$

Finally, we look at the difference coming from the third term of (B.5). We have

$$\left\| \begin{pmatrix} \cos(\bar{u} + \phi) \\ \sin(\bar{u} + \phi) \end{pmatrix} (v - v_0) \right\|_{C^\alpha} \leq C \|v - v_0\|_{C^\alpha}.$$

Thereby we obtain using (B.7) and Lemma B.3

$$\begin{aligned}
&\left\| \Pi^{-1}(\bar{u} + \phi) \mathcal{J}(\bar{u}, \bar{\rho}) \cdot \begin{pmatrix} \cos(\bar{u} + \phi) \\ \sin(\bar{u} + \phi) \end{pmatrix} (v - v_0) \right\|_{\mathbb{E}_0} \\
&\leq C \sup_{t \in (0, T]} t^{1-\eta} \|v - v_0\|_{C^\alpha} \leq C \|v\|_{\mathbb{E}_1} T.
\end{aligned}$$

This proves the first part of the claim. The second part of the claim follows with the same strategy. \square

B.4. Solving the linearized problem. We show that the map in (3.10) is an isomorphism, now considering a linear initial value problem with time-dependent coefficients.

Proposition B.6. *Let $(u_0, \rho_0) \in \gamma\mathbb{E}_1$ such that u_0 satisfies (3.1) and let $(\bar{u}, \bar{\rho}) \in \mathbb{E}_1([0, \bar{T}])$ be the function constructed in the proof of Proposition 3.3. Then there exists $T' \in (0, \bar{T}]$ such that the mapping*

$$\tilde{\mathbf{J}}: \mathbb{E}_1([0, T']) \rightarrow \mathbb{E}_0([0, T']) \times \gamma\mathbb{E}_1, (v, \sigma) \mapsto (\partial_t(v, \sigma) - \mathrm{DF}(\bar{u}, \bar{\rho})(v, \sigma), (v, \sigma)(0, \cdot))$$

is an isomorphism. Equivalently, for any right-hand side $(\varphi_1, \varphi_2) \in \mathbb{E}_0([0, T'])$ and any initial datum $(v_0, \sigma_0) \in \gamma\mathbb{E}_1$ there is a unique solution $(v, \sigma) \in \mathbb{E}_1([0, T'])$ of

$$\begin{cases} \partial_t(v, \sigma) - \mathrm{DF}(\bar{u}, \bar{\rho})(v, \sigma) = (\varphi_1, \varphi_2) & \text{in } (0, T') \times \mathbb{R}, \\ (v, \sigma)(0, \cdot) = (v_0, \sigma_0) & \text{on } \mathbb{R}. \end{cases}$$

Proof. Let $T \in (0, \bar{T}]$ to be chosen. We compare $\tilde{\mathbf{J}}$ with \mathbf{J} from Proposition B.1, which we already know is an isomorphism. We have

$$\begin{aligned}\tilde{\mathbf{J}}(v, \sigma) &= \mathbf{J}(v, \sigma) + (\mathcal{A}(v, \sigma) + \Psi(v_0, \sigma_0) - \mathbf{DF}(\bar{u}, \bar{\rho})(v, \sigma), (0, 0)) \\ &= \mathbf{J}(\mathbf{I}(v, \sigma) + \mathbf{J}^{-1}(v, \sigma) (\mathbf{S}(v, \sigma), (0, 0))),\end{aligned}$$

where \mathcal{A}, Ψ are given as in Proposition B.1, \mathbf{I} is the operator identity and

$$\begin{aligned}\mathbf{S} &:= (\mathbf{S}_1, \mathbf{S}_2): \mathbb{E}_1([0, T]) \rightarrow \mathbb{E}_0([0, T]), \\ \mathbf{S}(v, \sigma) &= \mathcal{A}(v, \sigma) + \Psi(v_0, \sigma_0) - \mathbf{DF}(\bar{u}, \bar{\rho})(v, \sigma).\end{aligned}$$

Thus, by a Neumann series argument, it is sufficient to show that the operator norm of \mathbf{S} becomes arbitrarily small for T getting small and at the same time the operator norm $\|\mathbf{J}^{-1}\|$ remains bounded, independent of $T \leq \bar{T}$. This last part has already been proven in Lemma B.2. Hence, the statement follows if we show

$$(B.11) \quad \|\mathbf{S}\| \leq C \max \{T, T^\eta, T^{\bar{\eta}}\}.$$

By definition of \mathcal{A}, Ψ and using (B.1), (B.2) we find

$$\begin{aligned}(B.12) \quad \mathbf{S}_1(v, \sigma) &= (\beta(\rho_0) - \beta(\bar{\rho})) \partial_s^2 v + \beta'(\bar{\rho}) \partial_s^2 \bar{u} (\sigma_0 - \sigma) + (\beta'(\rho_0) \partial_s \rho_0 - \beta'(\bar{\rho}) \partial_s \bar{\rho}) \partial_s v \\ &\quad + (\beta'(\rho_0) (\partial_s u_0 + \partial_s \phi - c_0) - \beta'(\bar{\rho}) (\partial_s \bar{u} + \partial_s \phi - c_0)) \partial_s \sigma \\ &\quad + (\beta''(\rho_0) \partial_s \rho_0 (\partial_s u_0 + \partial_s \phi - c_0) - \beta''(\bar{\rho}) \partial_s \bar{\rho} (\partial_s \bar{u} + \partial_s \phi - c_0)) \sigma \\ &\quad + \mathbf{D}\Lambda_u(\bar{u}, \bar{\rho})(v_0, \sigma_0) - \mathbf{D}\Lambda_u(\bar{u}, \bar{\rho})(v, \sigma)\end{aligned}$$

and

$$\begin{aligned}(B.13) \quad \mathbf{S}_2(v, \sigma) &= \left(-\frac{1}{2} \beta''(\rho_0) (\partial_s u_0 + \partial_s \phi - c_0)^2 + \frac{1}{2} \beta''(\bar{\rho}) (\partial_s \bar{u} + \partial_s \phi - c_0)^2 \right) \sigma \\ &\quad + (-\beta'(\rho_0) (\partial_s u_0 + \partial_s \phi - c_0) + \beta'(\bar{\rho}) (\partial_s \bar{u} + \partial_s \phi - c_0)) \partial_s v \\ &\quad + \mathbf{D}\Lambda_\rho(\bar{u}, \bar{\rho})(v_0, \sigma_0) - \mathbf{D}\Lambda_\rho(\bar{u}, \bar{\rho})(v, \sigma).\end{aligned}$$

We now take a look at the terms of (B.12) and (B.13). In the following, C denotes a constant, only depending on the model parameters, α, η and M as in (B.7), which is allowed to change from line to line. The \mathbb{E}_0 -norm of the first term of (B.12) can be estimated using Lemma A.2 and Lemma B.4 by

$$\|(\beta(\rho_0) - \beta(\bar{\rho})) \partial_s^2 v\|_{\mathbb{E}_0} \leq C \sup_{t \in (0, T]} t^{1-\eta} \|\beta(\rho_0) - \beta(\bar{\rho})\|_{C^\alpha} \|v\|_{C^{2+\alpha}} \leq C \|v\|_{\mathbb{E}_1} T^\eta.$$

Here we used the time-weight of the \mathbb{E}_0 -norm to bound the second derivative of v . For the second term of (B.12) we have to use the time-weight to bound the second derivative of \bar{u} . An additional factor depending on T and going to zero for T going to zero is given by Lemma B.3. This motivates the choice of the first term in Ψ in Proposition B.1. We obtain

$$\begin{aligned}\|\beta'(\bar{\rho}) \partial_s^2 \bar{u} (\sigma_0 - \sigma)\|_{\mathbb{E}_0} &\leq C \sup_{t \in (0, T]} t^{1-\eta} \|\beta'(\bar{\rho})\|_{C^\alpha} \|\bar{u}\|_{C^{2+\alpha}} \|\sigma_0 - \sigma\|_{C^\alpha} \\ &\leq C \|\sigma\|_{\mathbb{E}_1} T^\eta.\end{aligned}$$

The third term of (B.12) can be estimated using Lemma B.4 and Lemma B.3 by

$$\begin{aligned}&\|(\beta'(\rho_0) \partial_s \rho_0 - \beta'(\bar{\rho}) \partial_s \bar{\rho}) \partial_s v\|_{\mathbb{E}_0} \\ &\leq C \sup_{t \in (0, T]} t^{1-\eta} \|\beta'(\rho_0) - \beta'(\bar{\rho})\|_{C^\alpha} \|\rho_0\|_{C^{1+\alpha}} \|v\|_{C^{1+\alpha}} \\ &\quad + C \sup_{t \in (0, T]} t^{1-\eta} \|\beta'(\bar{\rho})\|_{C^\alpha} \|\partial_s \rho_0 - \partial_s \bar{\rho}\|_{C^\alpha} \|v\|_{C^{1+\alpha}} \leq C \|v\|_{\mathbb{E}_1} T^\eta + C \|v\|_{\mathbb{E}_1} T^{\bar{\eta}}.\end{aligned}$$

The same also applies for the fourth term of (B.12) and the second term of (B.13). For the fifth term of (B.12) we proceed similarly and estimate

$$\begin{aligned} & \|(\beta''(\rho_0)\partial_s\rho_0(\partial_s u_0 + \partial_s\phi - c_0) - \beta''(\bar{\rho})\partial_s\bar{\rho}(\partial_s\bar{u} + \partial_s\phi - c_0))\sigma\|_{\mathbb{E}_0} \\ & \leq C \sup_{t \in (0, T]} t^{1-\eta} \|\beta''(\rho_0) - \beta''(\bar{\rho})\|_{C^\alpha} \|\rho_0\|_{C^{1+\alpha}} \|\partial_s u_0 + \partial_s\phi - c_0\|_{C^\alpha} \|\sigma\|_{C^\alpha} \\ & \quad + C \sup_{t \in (0, T]} t^{1-\eta} \|\beta''(\bar{\rho})\|_{C^\alpha} \|\partial_s\rho_0 - \partial_s\bar{\rho}\|_{C^\alpha} \|\partial_s u_0 + \partial_s\phi - c_0\|_{C^\alpha} \|\sigma\|_{C^\alpha} \\ & \quad + C \sup_{t \in (0, T]} t^{1-\eta} \|\beta''(\bar{\rho})\|_{C^\alpha} \|\bar{\rho}\|_{C^{1+\alpha}} \|\partial_s u_0 - \partial_s\bar{u}\|_{C^\alpha} \|\sigma\|_{C^\alpha} \leq C \|\sigma\|_{\mathbb{E}_1} (T^\eta + T^{\bar{\eta}}). \end{aligned}$$

Analogously, we can treat the first term of (B.13). Together with Lemma B.5, (B.11) and thus the claim follows for $T' = T < \bar{T}$ sufficiently small. \square

ACKNOWLEDGEMENTS

The authors acknowledge support by the DFG (German Research Foundation), project no. 404870139. The third author is additionally supported by the Austrian Science Fund (FWF) project/grant P 32788-N.

REFERENCES

- [1] H. Abels, F. Bürger, and H. Garcke. Short time existence for coupling of scaled mean curvature flow and diffusion. *arXiv*, 2204.07626, 2022.
- [2] S. Boussandel, R. Chill, and E. Fasangova. Maximal regularity, the local inverse function theorem, and local well-posedness for the curve shortening flow. *Czechoslovak Mathematical Journal*, 62(2):335–346, 2012.
- [3] K. Brazda, G. Jankowiak, C. Schmeiser, and U. Stefanelli. Bifurcation of elastic curves with modulated stiffness. *European Journal of Applied Mathematics*, page 1–27, 2022.
- [4] H. Brezis. *Functional analysis, Sobolev spaces and partial differential equations*. Universitext. Springer, New York, 2011.
- [5] P. Clément and G. Simonett. Maximal regularity in continuous interpolation spaces and quasilinear parabolic equations. *J. Evol. Equ.*, 1(1):39–67, 2001.
- [6] A. Dall'Acqua, C.-C. Lin, and P. Pozzi. A gradient flow for open elastic curves with fixed length and clamped ends. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 17(3):1031–1066, 2017.
- [7] A. Dall'Acqua and P. Pozzi. A Willmore–Helfrich L^2 -flow of curves with natural boundary conditions. *Communications in Analysis and Geometry*, 22:617–669, 11 2012.
- [8] G. Dziuk, E. Kuwert, and R. Schätzle. Evolution of elastic curves in \mathbb{R}^n : existence and computation. *SIAM J. Math. Anal.*, 33(5):1228–1245, 2002.
- [9] C. M. Elliott, H. Garcke, and B. Kovács. Numerical analysis for the interaction of mean curvature flow and diffusion on closed surfaces. *arXiv*, 2202.03302, 2022.
- [10] N. Koiso. On the motion of a curve towards elastica. In *Actes de la Table Ronde de Géométrie Différentielle (Luminy, 1992)*, volume 1 of *Sémin. Congr.*, pages 403–436. Soc. Math. France, Paris, 1996.
- [11] O. A. Ladyženskaja, V. A. Solonnikov, and N. N. Ural'ceva. *Linear and quasilinear equations of parabolic type*. Translations of Mathematical Monographs, Vol. 23. American Mathematical Society, Providence, R.I., 1968. Translated from the Russian by S. Smith.
- [12] J. LeCrone. Elliptic operators and maximal regularity on periodic little-Hölder spaces. *J. Evol. Equ.*, 12(2):295–325, 2012.
- [13] A. Lunardi. *Analytic semigroups and optimal regularity in parabolic problems*. Modern Birkhäuser Classics. Birkhäuser/Springer Basel AG, Basel, 1995.
- [14] C. Mantegazza, A. Pluda, and M. Pozzetta. A survey of the elastic flow of curves and networks. *Milan J. Math.*, 89(1):59–121, 2021.
- [15] M. Novaga and S. Okabe. Curve shortening-straightening flow for non-closed planar curves with infinite length. *J. Differential Equations*, 256(3):1093–1132, 2014.
- [16] A. Polden. *Curves and Surfaces of Least Total Curvature and Fourth Order Flows*. PhD thesis, Universität Tübingen, 1996.
- [17] P. Pozzi and B. Stinner. Elastic flow interacting with a lateral diffusion process: the one-dimensional graph case. *IMA J. Numer. Anal.*, 39(1):201–234, 2019.
- [18] F. Rupp and A. Spener. Existence and convergence of the length-preserving elastic flow of clamped curves. *arXiv*, 2009.06991, 2020.

- [19] A. Spener. Short time existence for the elastic flow of clamped curves. *Math. Nachr.*, 290(13):2052–2077, 2017.
- [20] Y. Wen. L^2 flow of curve straightening in the plane. *Duke Math. J.*, 70(3):683–698, 1993.
- [21] G. Wheeler. Global analysis of the generalised Helfrich flow of closed curves immersed in \mathbb{R}^n . *Trans. Amer. Math. Soc.*, 367(4):2263–2300, 2015.

(A. Dall’Acqua) INSTITUTE OF APPLIED ANALYSIS, HELMHOLTZSTRASSE 18, 89081 ULM, GERMANY.
Email address: `anna.dallacqua@uni-ulm.de`

(L. Langer) INSTITUTE OF APPLIED ANALYSIS, HELMHOLTZSTRASSE 18, 89081 ULM, GERMANY.
Email address: `leonie.langer@uni-ulm.de`

(F. Rupp) FACULTY OF MATHEMATICS, UNIVERSITY OF VIENNA, OSKAR-MORGENSTERN-PLATZ 1,
1090 VIENNA, AUSTRIA.
Email address: `fabian.rupp@univie.ac.at`