Improved Upper Bound on Independent Domination Number for Hypercubes

Debabani Chowdhury & Debesh K. Das

Department of Computer Sc. & Engg. Jadavpur University, Kolkata - 700 032, India debabani.chowdhury@gmail.com debeshkdas@gmail.com

Bhargab B. Bhattacharya

Department of Computer Sc. & Engg. Indian Institute of Technology Kharagpur WB - 721 302, India bbbiitkgp@gmail.com

May 16, 2022

Abstract

We revisit the problem of determining the independent domination number in hypercubes for which the known upper bound is still not tight for general dimensions. We present here a constructive method to build an independent dominating set S_n for the n-dimensional hypercube Q_n , where n=2p+1, p being a positive integer ≥ 1 , provided an independent dominating set S_p for the p-dimensional hypercube Q_p , is known. The procedure also computes the minimum independent dominating set for all $n=2^k-1$, k>1. Finally, we establish that the independent domination number $\alpha_n \leq 3 \times 2^{n-k-2}$ for $7 \times 2^{k-2} - 1 \leq n < 2^{k+1} - 1$, k>1. This is an improved upper bound for this range as compared to earlier work.

Keywords: Hypercubes, independent dominating set, minimum independent dominating set, independent domination number, Boolean functions, Hamming distance

1 Introduction and related work

Hypercubes are classical structures that are well-studied in many important areas such as set theory, graph theory, combinatorics and Boolean algebra [15]. Given an n-dimensional hypercube graph $Q_n = (V_n, E_n)$, where V_n be the set of vertices (nodes) and E_n be the set of edges in Q_n , and a subset $S_n \subset V_n$, we say that S_n dominates its neighbourhood $N(S_n)$, i.e., the set of all nodes that

are adjacent to at least one node in S_n [1,15]. If S_n dominates $V_n \setminus S_n$, then S_n is called a dominating set of Q_n . S_n is said to be independent if no two nodes in S_n are adjacent. An independent dominating set with minimum cardinality is called the minimum independent dominating set. The independent domination number α_n is the cardinality of the minimum independent dominating set of Q_n .

Many results are known on independent domination for simple regular and other graphs [3]- [8]. However, there are many unsettled questions regarding minimum dominating sets in hypercubes as stated by Harary et al. [1]. They derived the minimum independent dominating sets of Q_n for dimension n=1,2,3,4,5,6: $\alpha_1=1,\alpha_2=2,\alpha_3=2,\alpha_4=4,\alpha_5=8,\alpha_6=12$. It was also reported that $\alpha_n=2^{n-k}$ for $n=(2^k-1)$ and $n=2^k, k>0$. Yet, the upper bound on α_n is still not tight for many other values of n. Mane et al. [2] showed that $\alpha_n \leq 2^{n-k}$ for $2^k < n < 2^{k+1} - 1$, for any n. In this work, we improve this bound for certain values of n. We split the range $2^k-1\leq n<2^{k+1}-1$ into two non-overlapping cases:

Case 1:
$$2^k - 1 \le n < 7 \times 2^{k-2} - 1$$
, $k > 1$;
Case 2: $7 \times 2^{k-2} - 1 \le n < 2^{k+1} - 1$, $k > 1$.

In this paper, we show that $\alpha_n \leq 3 \times 2^{n-k-2}$ for $7 \times 2^{k-2} - 1 \leq n < 2^{k+1} - 1, k > 1$. This tightens the upper bound for Case 2 in comparison to the result $\alpha_n \leq 2^{n-k}$ proved earlier for both cases by Mane et al. [2]. Given an independent dominating set S_p , our procedure constructs an independent dominating set S_n , where $|S_n| = 2^p \times |S_p|$, n = 2p+1, p being a positive integer ≥ 1 . Our result follows from a procedure that iteratively constructs a minimal independent dominating set in a hypercube from that of a smaller-dimension hypercube.

2 Concepts and techniques

The *n*-dimensional hypercube $Q_n = (V_n, E_n)$ is an undirected regular graph whose vertex set where V_n and E_n denote the set of nodes and set of edges in Q_n , respectively. The set V_n can be envisaged as the set of all binary *n*-tuples of zeros and ones, i.e. $\{0,1\}^n$. Thus, $|V_n| = 2^n$. Two nodes of Q_n are adjacent if their binary *n*-tuples differ in exactly one place, i.e., if they are unit Hamming distance apart. Let the Hamming distance between two binary *n*-tuples corresponding to vertices $v_1, v_2 \in V_n$ be represented as $d_H(v_1, v_2)$.

Lemma 1
$$\alpha_n \ge |(2^n/n+1)|$$

Proof: For any $Q_n = (V_n, E_n)$, $|V_n| = 2^n$ and every vertex in V_n has n adjacent vertices. That is, a vertex of V_n can dominate at most n vertices uniquely. Thus, the minimum number of vertices required to dominate all 2^n vertices of Q_n will be $2^n/(n+1)$. As the number of vertices should be an integer, $\alpha_n > \lfloor (2^n/n+1) \rfloor$.

3 Iterative construction of an independent dominating set in a hypercube

The following procedure constructs an independent dominating set S_{n+1} of Q_{n+1} , given an independent dominating set S_n of Q_n , where $|S_{n+1}| = 2 \times |S_n|$. We use the following notation in Procedure 1.

Let S'_n be an independent dominating set of Q_n such that $|S'_n| = |S_n|$. Let a vertex $v_{k,n} \in S_n$, $v'_{k,n} \in S'_n$ be represented by n-bit binary vectors $b^k_{n-1}b^k_{n-2}\cdots b^k_1b^k_0$, $b^k_{n-1}b^k_{n-2}\cdots b^k_1b^k_0$, respectively, where $b^k_l \in \{0,1\}$, $b^k_l \in \{0,1\}$, $(1 \le k \le |S_n|)$, $(0 \le l < n)$.

Procedure 1:

Input: S_n Output: S_{n+1}

Step 1. Assign $S'_n \leftarrow \phi, S_{n+1} \leftarrow \phi$.

Step 2. For every vertex $v_{k,n} \in S_n$ get a vertex $v'_{k,n}$, such that $v_{k,n}, v'_{k,n}$ differ in LSB, i.e., $v_{k,n} = b^k_{n-1}b^k_{n-2}\cdots b^k_1b^k_0$, and $v'_{k,n} = b'^k_{n-1}b'^k_{n-2}\cdots b'^k_1b'^k_0 = b^k_{n-1}b^k_{n-2}\cdots b^k_1\overline{b^k_0}$.

Step 3. $\forall v'_{k,n}, S'_n = S'_n \cup \{v'_{k,n}\}.$

Step 4. For every $v_{k,n} \in S_n$, and for every $v'_{k,n} \in S'_n$ get $v_{k,n+1}, v'_{k,n+1}$ respectively, such that $v_{k,n+1} = b^k_n.v_{k,n} = b^k b^k_{n-1} b^k_{n-2} \cdot \cdot \cdot b^k_1 b^k_0, \ b^k_n \in \{0,1\},$ and $v'_{k,n+1} = b'^k_n.v'_{k,n} = b'^k b'^k_{n-1} b'^k_{n-2} \cdot \cdot \cdot b'^k_1 b^k_0 = \overline{b^k_n} b^k_{n-1} b^k_{n-2} \cdot \cdot \cdot b^k_1 \overline{b^k_0},$ $b'^k_n \in \{0,1\}.$

Step 5. $S_{n+1} = S_{n+1} \cup \{v_{k,n+1}, v'_{k,n+1}\}, \forall v_{k,n} \in S_n, \forall v'_{k,n} \in S'_n$.

Step 6. Output S_n .

Step 7. End.

Lemma 2 $|S_{n+1}| = 2 \times |S_n|$

Proof: It follows from Procedure 1.

Lemma 3 S_{n+1} as constructed by Procedure 1 is an independent dominating set of Q_n .

Proof: Follows from [9].

4 Constructing an independent dominating set for Q_n , where $n = (2p + 1), p \ge 1$

Let S_p, S_n be independent dominating sets for p-, and n-dimensional hypercubes $Q_p = (V_p, E_p)$ and $Q_n = (V_n, E_n)$, respectively, where n = 2p + 1. Clearly, $S_p \subset V_p$, $S_n \subset V_n$, $|V_p| = 2^p$, $|V_n| = 2^n$. Let s_k^p , s_l^n be vertices of Q_p and Q_n , respectively, such that $s_k^p \in S_p$, $(1 \le k \le |S_p|)$, $s_l^n \in S_n$, $(1 \le l \le |S_n|)$. The

following procedure constructs S_n from S_p .

Procedure 2:

Input: S_p where every vector $s_k^p \in S_p$ is represented as p-bit binary $s_k^p = b_{(p-1)}^k b_{(p-2)}^k \cdots b_i^k b_0^k$, where $b_i^k \in \{0,1\}, (0 \le i \le p-1)$. Output: S_n where every vector $s_l^n \in S_n$ is represented as n-bit binary $s_l^n = c_{(n-1)}^l c_{(n-2)}^l \cdots c_i^l \cdots c_1^l c_0^l$, where $c_i^l \in \{0,1\}, (0 \le i \le n-1)$.

Step 1. Assign $N \leftarrow 0, S_n \leftarrow \phi$

Step 2. Get p-bit binary tuple of N as $A^N = a_{p-1}^N a_{p-2}^N \cdots a_i^N \cdots a_0^N$, where $a_i^N \in \{0,1\}, (0 \le i \le p-1)$.

Step 3. Set k = 1.

Step 4. Set $l = N \times |S_p| + k$.

Step 5. $j \leftarrow 0$.

Step 6. $c_{j+p}^l = a_j^N$.

Step 7. If $a_j^N = 1$, then $c_j^l = \overline{b_j^k}$ else $c_j^l = b_j^k$.

Step 8. j = j + 1; If j < p then goto Step 6.

Step 9. If A^N is of even parity then $c_{n-1}^l = 0$ else $c_{n-1}^l = 1$.

Step 10. $S_n = S_n \cup \{s_l^n\}.$

Step 11. k = k + 1; If $k \le |S_p|$, then goto Step 4.

Step 12. N = N + 1; If $N < 2^p$, then goto Step 2.

Step 13. Output S_n .

Step 14. End.

Example 1 Table 1 shows how Procedure 2 is executed, given a minimum independent dominating set $S_3 = \{000, 111\}$ for the 3-dimensional hypercube Q_3 to obtain an independent dominating set S_7 for the 7-dimensional hypercube Q_7 . Also, set S_7 produced by Procedure 2 is a minimum independent dominating set in Q_7 . Figure 1 shows how a vector of S_3 is modified to produce a vector of S_7 . It also includes, in general, the concatenation process that leads to an n=(2p+1)-dimensional vector of the dominating set S_n starting from a p-dimensional vector of the dominating set S_p . Given, p=3, n=7, $1 \le k \le |S_p|$, i.e., $1 \le k \le 2$, $k=\{k_1,k_2\}=\{1,2\}$ and $S_p=\{s_{k_1}^p,s_{k_2}^p\}=\{s_1^3,s_2^3\}=\{010,101\}$.

Lemma 4 S_n as produced by Procedure 2 has cardinality $|S_n| = 2^p \times |S_p|$.

Proof: Immediate from Procedure 2.

Theorem 1 S_n as obtained by Procedure 2, is an independent dominating set of Q_n .

Proof: First, we prove that S_n is an independent set of Q_n . Next, we will show that S_n is also a dominating set of Q_n .

Let $s_{l_1}^n, s_{l_2}^n \in S_n$ be two vertices of Q_n obtained by Procedure 2. Clearly,

$$\begin{split} s_{l_1}^n &= c_{n-1}^{l_1} c_{n-2}^{l_1} c_{n-3}^{l_1} \cdots c_p^{l_1} c_{p-1}^{l_1} c_{p-2}^{l_2} \cdots c_0^{l_1} \\ &= c_{n-1}^{l_1} a_{p-1}^{N_1} a_{p-2}^{N_1} \cdots a_0^{N_1} b_{p-1}^{k_1^*} b_{p-2}^{k_2^*} \cdots b_0^{k_1^*}, \text{ where either } b_{p-1}^{k_1^*} = b_{p-1}^{k_1} \text{ or } \overline{b_{p-1}^{k_1}} \\ \text{and} \\ s_{l_2}^n &= c_{n-1}^{l_2} c_{n-2}^{l_2} c_{n-3}^{l_2} \cdots c_p^{l_2} c_{p-1}^{l_2} c_{p-2}^{l_2} \cdots c_0^{l_2} \\ &= c_{n-1}^{l_2} a_{p-1}^{N_2} a_{p-2}^{N_2} \cdots a_0^{N_2} b_{p-1}^{k_2^*} b_{p-2}^{k_2^*} \cdots b_0^{k_2^*}, \text{ where either } b_{p-1}^{k_2^*} = b_{p-1}^{k_2} \text{ or } \overline{b_{p-1}^{k_2}}, \\ (1 \leq l_1, l_2 \leq |S_n|), |S_n| &= 2^p \times |S_n|, l_1 \neq l_2, 0 \leq N_1, N_2 \leq (2^p - 1), 1 \leq k_1, k_2 \leq |S_p| \end{split}$$

For any two vetices $s_{l_1}^n, s_{l_2}^n \in S_n$, there are three possible cases;

Case 2: $N_1 \neq N_2$, $c_{n-1}^{l_1} = c_{n-1}^{l_2}$, i.e., both A^{N_1}, A^{N_2} have the same parity, either both even or both odd;

Case 3: $c_{n-1}^{l_1} \neq c_{n-1}^{l_2}$, i.e., if A^{N_1} has even (odd) parity, then A^{N_2} has odd (even)

For Case 1, the Hamming distance between n-bit binary vectors of $s_{l_1}^n, s_{l_2}^n$ is

$$\begin{split} &d_H(s_{l_1}^n,s_{l_2}^n)\\ &=d_H(c_{n-1}^{l_1},c_{n-1}^{l_2})+d_H(A^{N_1},A^{N_2})+d_H(b_{p-1}^{k_1^*}b_{p-2}^{k_1^*}\cdots b_0^{k_1^*},b_{p-1}^{k_2^*}b_{p-2}^{k_2^*}\cdots b_0^{k_2^*})\\ &=(d_H(c_{n-1}^{l_1},c_{n-1}^{l_2})+d_H(A^{N_1},A^{N_2})+d_H(s_{k_1}^p,s_{k_2}^p))\geq 2\\ &\text{Note that } d_H(c_{n-1}^{l_1},c_{n-1}^{l_2})=d_H(A^{N_1},A^{N_2})=0, \text{ and } d_H(s_{k_1}^p,s_{k_2}^p)\geq 2 \text{ as } s_{k_1}^p,s_{k_2}^p\in S_p \end{split}$$

For Case 2,

$$\begin{split} &d_{H}(s_{l_{1}}^{n}, s_{l_{2}}^{n}) \\ &= (d_{H}(c_{n-1}^{l_{1}}, c_{n-1}^{l_{2}}) + d_{H}(A^{N_{1}}, A^{N_{2}}) + d_{H}(b_{p-1}^{k_{1}^{*}} b_{p-2}^{k_{1}^{*}} \cdots b_{0}^{k_{1}^{*}}, b_{p-1}^{k_{2}^{*}} b_{p-2}^{k_{2}^{*}} \cdots b_{0}^{k_{2}^{*}})) \geq 2 \\ &\text{when } d_{H}(c_{n-1}^{l_{1}}, c_{n-1}^{l_{2}}) = 0, \ d_{H}(A^{N_{1}}, A^{N_{2}}) \geq 2 \end{split}$$

For Case 3,

$$\begin{split} &d_{H}(s_{l_{1}}^{n},s_{l_{2}}^{n})\\ &=(d_{H}(c_{n-1}^{l_{1}},c_{n-1}^{l_{2}})+d_{H}(A^{N_{1}},A^{N_{2}})+d_{H}(b_{p-1}^{k_{1}^{*}}b_{p-2}^{k_{1}^{*}}\cdots b_{0}^{k_{1}^{*}},b_{p-1}^{k_{2}^{*}}b_{p-2}^{k_{2}^{*}}\cdots b_{0}^{k_{2}^{*}}))\geq 2\\ &\text{when }d_{H}(c_{n-1}^{l_{1}},c_{n-1}^{l_{2}})=1,\ d_{H}(A^{N_{1}},A^{N_{2}})\geq 1 \end{split}$$

Thus, for any two vertices $s_{l_1}^n, s_{l_2}^n \in S_n$, we have $d_H(s_{l_1}^n, s_{l_2}^n) \geq 2$. That is, no two vertices in S_n are adjacent. Hence, S_n is an independent set of Q_n .

To prove that S_n is a dominating set, we argue as follows. We have proved that S_n is an independent set, i.e., its no two vertices are adjacent, i.e., $\forall v_1, v_2 \in$ $S_n, d_H(v_1, v_2) \geq 2$. Thus, for every vertex $v_n \in S_n$, its all n adjacent vertices are in S'_n . Thus, we have the following four cases: Case 1: S'_n contains only $N(S_n)$;

Case 2: $S_n \cap S_n' = \phi$; Case 3: $\forall v_n' \in S_n', \exists \text{ a } v_n \in S_n$, such that v_n' is adjacent to v_n .

Let $N(v_1), N(v_2)$ be the neighbourhood of $v_1, v_2 \in S_n$, respectively. Now, $\forall v_1, v_2 \in S_n$, if every $N(v_1) \cap N(v_2) = \phi$, then

$$|S_n| + |S'_n|$$

$$= (2^p \times |S_p|) + (n \times 2^p \times |S_p|)$$

$$= 2^p \times |S_p| \times (n+1)$$

$$= |S_p| \times 2^p \times (2p+2)$$

$$= 2^{p+1} \times (p+1) \times |S_p|$$

$$= 2^{p+1} \times (|S_p| + p \times |S_p|)$$

$$= 2^{p+1} \times (|S_p| + |S'_p|)$$

$$= 2^{p+1} \times 2^p = 2^{2p+1} = 2^n$$
So,
$$|S_n| + |S'_n| = |V_n|$$

Cace 4: $S_n \cup S'_n = V_n$.

By combining Cases (1), (2) and (4), we have $S'_n = V_n \setminus S_n$, i.e., S'_n contains all those vertices of Q_n which are not in S_n . Again, from Case 3, $\forall v'_n \in S'_n$, \exists a $v_n \in S_n$, such that v'_n is adjacent to v_n . Thus, S_n is a dominating set of Q_n .

Procedure 2 leads to the following known result [1].

Lemma 5 $\alpha_n = 2^{n-k}$ for $n = 2^k - 1, k > 0$.

Proof: From Lemma 1, $\alpha_n \ge \lfloor (2^n/n+1) \rfloor$. From Lemma 1, it follows that $\alpha_n > \lfloor (2^n/n+1) \rfloor$. When $n = 2^k - 1$, $2^n/(n+1) = 2^{n-k}$ is an integer, and hence, $\alpha_n = 2^{n-k}$.

5 Upper bound on the independent domination number for hypercubes

Theorem 2 For $n = 7 \times 2^{k-2} - 1$, where k > 1, $\alpha_n < 3 \times 2^{n-k-2}$

Proof: From [1], we know that $\alpha_6 = 12$. In this case, k = 2. Following Lemma 4 and Theorem 1, for n = 13, we obtain $|S_{13}| = 12 \times 2^6$, where k = 3. By iteration, we can then derive results for $n = 27, 55, 111, 223, \cdots$, corresponding to $k = 4, 5, 6, 7, \cdots$, respectively. Following Procedure 2, we obtain $|S_n| = 12 \times 2^6, 12 \times 2^{19}, 12 \times 2^{46}, \cdots$ for $n = 13, 27, 55, \cdots$, respectively. The k^{th} term in the series can be evaluated as $7 \times 2^{k-2} - 1$. Hence, the proof.

Theorem 3 For $7 \times 2^{k-2} - 1 \le n < 2^{k+1} - 1$, where k > 1, $\alpha_n \le 3 \times 2^{n-k-2}$

Proof: If S_n is given, we can construct S_{n+1} following Procedure 1. Thus, when $|S_n| = 3 \times 2^{n-k-2}$, $|S_{n+1}| = 2 \times |S_n| = 3 \times 2^{(n+1)-k-2}$. Hence, the proof.

Thus it improves the previous upper bound $\alpha_n \leq 2^{n-k}$ [2] for the range of $7 \times 2^{n-k} \leq n < 2^{k+1} - 1$.

Theorem 4 For $2^k - 1 \le n < 7 \times 2^{k-2} - 1$, k > 0, $\alpha_n \le 2^{n-k}$.

Proof: If S_n is given, we can construct S_{n+1} following Procedure 1. Thus, when $|S_n| = 2^{n-k}$, $|S_{n+1}| = 2 \times |S_n| = 2^{(n+1)-k}$. Hence, the proof.




Figure 1: Construction of an n=(2p+1)-dimensional vector of the independent dominating set S_n from a p-dimensional vector of the dominating set S_p , where n=(2p+1). It illustatrates the construction of a vector for p=3, $S_3=\{010,101\}$, i.e., the construction of a vector of S_7 from a vector of S_3 .

N	A^N	k	l	c_{n-1}^l	$s_k^p = s_k^3 = b_{p-1}^k b_{p-2}^k \cdots b_0^k = b_2^k b_1^k b_0^k$	$c_{p-1}^l c_{p-2}^l \cdots c_0^l$	$s_{ S_n \times N+k}^n$
	$\begin{vmatrix} = a_{p-1}^N a_{p-1}^N \cdots a_0^N \\ = a_2^N a_1^N a_0^N \end{vmatrix}$			$= c_6^l$	$=b_{p-1}^k b_{p-2}^k \cdots b_0^k$	$=c_2^l c_1^l c_0^l$	$=s_{2\times N+k}^{7}$
	$=a_2^N a_1^N a_0^N$			(parity	$=b_2^k b_1^k b_0^k$		$=c_{n-1}^{l}A^{N}c_{p-1}^{l}$
				of A^N)			$\cdots c_1^l c_0^l$
0	000	1	1	0	000	000	0000000
0	000	2	2	0	111	111	0 000111
1	001	1	3	1	000	001	1001001
1	001	2	4	1	111	110	1001110
2	010	1	5	1	000	010	1010010
2	010	2	6	1	111	101	1010101
3	011	1	7	0	000	011	0 011011
3	011	2	8	0	111	100	0 011100
4	100	1	9	1	000	100	1100100
4	100	2	10	1	111	011	1100011
5	101	1	11	0	000	101	0 101101
5	101	2	12	0	111	010	0 101010
6	110	1	13	0	000	110	0110110
6	110	2	14	0	111	001	0110001
7	111	1	15	1	000	111	1111111
7	111	2	16	1	111	000	1111000

Table 1: Construction of S_7 from S_3

6 Independent domination number for hypercubes

Here, we summarize our results and compare them with previously known bounds.

6.1 Previous results

For
$$n = 2^k - 1$$
, $k \ge 1$, $\alpha_n = 2^{n-k}$ [1,10–12]
For $n = 2^k$, $k \ge 1$, $\alpha_n = 2^{n-k}$ [1,2,13]
For $2^k < n < 2^{k+1} - 1$, $k \ge 1$, $\alpha_n \le 2^{n-k}$ [2]

6.2 Proposed work

For
$$n=2^k-1, \ k>0, \ \alpha_n=2^{n-k}$$

For $7\times 2^{k-2}-1\le n<2^{k+1}-1, \ k>1, \ \alpha_n\le 3\times 2^{n-k-2}$
For $2^k-1\le n<7\times 2^{k-2}-1, \ k>0, \ \alpha_n\le 2^{n-k}$

6.3 Numerical comparison for some values of n

We report the values of α_n for different values of n in Table 2. Results in Table 3 are obtained from Table 2, which show improved upper bound for some values on $n \leq 62$.

	k	> 0		k > 1
n	$2^{k} - 1$	2^k	$2^k < n < 7 \times 2^{k-2} - 1$	$7 \times 2^{k-2} - 1 \le n < 2^{k+1} - 1$
$\alpha_n [1, 2, 10-13]$	2^{n-k}	2^{n-k}		
α_n [2]		2^{n-k}	$\leq 2^{n-k}$	$\leq 2^{n-k}$
α_n (this work)	2^{n-k}	$\leq 2^{n-k}$	$\leq 2^{n-k}$	$\leq 3 \times 2^{n-k-2}$

Table 2: α_n of Q_n for $2^k-1\leq n\leq 2^{k+1}-1, k\geq 1$

k	n	α_n	α_n	α_n
		[1, 2, 10]	[2]	(this work)
		[11, 12]		
		[13, 14]		
1	2	2	2	≤ 2
	3	2		2
2	4	4	2^{2}	$\begin{array}{l} \leq 2^2 \\ \leq 2^3 \end{array}$
	5	8	$\leq 2^3$	$\leq 2^3$
	6	12	$2^{2} \le 2^{3} \le 2^{4}$	$\leq 3 \times 2^2$
	7	$\frac{2^4}{2^5}$		2^{4}
	7 8 9	2^{5}	2^{5}	$\leq 2^5$
	9		$\leq 2^6$	$\leq 2^6$
3	10		$\leq 2^7$	$\leq 2^7$
0	11		$\leq 2^8$	$\leq 2^8$
	12		$\le 2^{9}$	$ \leq 2^{5} \leq 2^{6} \leq 2^{7} \leq 2^{8} \leq 2^{9} $
	13			$\begin{array}{l} \leq 3 \times 2^8 \\ \leq 3 \times 2^9 \end{array}$
	14		$\leq 2^{11}$	$\leq 3 \times 2^9$

k	n	$ \begin{array}{c} \alpha_n \\ [1,2,10] \\ [11,12] \\ [13,14] \end{array} $	α_n [2]	$\begin{array}{c} \alpha_n \\ \text{(this work)} \end{array}$
4	27 28 29 30			
5	55 56 57 58 59 60 61 62			

Table 3: α_n for $n \leq 62$

7 Conclusion

In this work, we have derived an improved upper bound on independent domination number in an n-dimensional hypercube for certain range of values of n.

References

- [1] F. Harary and M. Livingston, "Independent domination in hypercubes", Appl. Math. Lett., vol. 6, no. 3, pp. 27 28, 1993, https://doi.org/10.1016/0893-9659(93)90027-K.
- [2] S. A. Mane В. N. Waphare, "On independent and and hypercubes", (d, n)-domination numbers of AKCE Int. Vol. 9, 2012, J. Graphs Comb., No. 2, 161-168, pp. https://www.tandfonline.com/doi/abs/10.1080/09728600.2012.12088957
- [3] J. Haviland, "On minimum maximal independent sets of a graph", Discrete Mathematics, Volume 94, Issue 2, pp. 95-101, 28 November 1991, https://doi.org/10.1016/0012-365X(91)90318-V

- [4] Jerzy Topp, Lutz Volkmann, "On graphs with equal domination and independent domination numbers, Discrete Mathematics, Vol. 96, pp. 75-80, 1991, https://doi.org/10.1016/0012-365X(91)90473-F
- [5] J. Haviland, "Independent domination in regular graphs", Discrete Mathematics, Vol. 143, Issues 1-3, pp. 275-280, 16 August 1995, https://doi.org/10.1016/0012-365X(94)00022-B
- [6] J. Haviland, "Independent dominating sets in regular graphs", Journal of Combinatorial Optimization, Vol. 26, pp. 120-126, 2013, https://doi.org/10.1007/s10878-011-9439-6
- [7] Peter Che Bor Lam, Wai Chee Shiu, Liang Sun. "On independent domination number of regular graphs, Discrere Mathematics, Vol. 202,Issues 1-3,pp. 135-144, 6th May 1999,https://doi.org/10.1016/S0012-365X(98)00350-1
- [8] W. Goddard, M.A. Henning, "Independent domination in graphs: A survey and recent results", Discrete Mathematics, Vol. 313, Issue 7, pp. 839-854, 6 April 2013, https://doi.org/10.1016/j.disc.2012.11.031
- [9] E. Pasalic, A. Chattapadhyay, D. Chowdhury, "An analysis of root-functions A subclass of the Impossible Class of Faulty Functions (ICFF)", Discrete Applied Mathematics, Vol. 222, pp. 1-13, 2017, http://dx.doi.org//10.1016/J.dam.2016.12.001
- [10] T.M. Thompson, From error-correcting codes through sphere packings to simple groups, Math. Assoc. Amer., Washington, Vol. 21, pp. 228, 1983, https://doi.org/10.5948/UP09781614440215
- [11] E.R. Berlekamp, Algebraic Coding Theory, (revised), Aegean Park Press, Laguna Hills, pp. 500, 1984, https://doi.org/10.1142/9407
- [12] J. Lee, "Independent perfect dominating sets in Cayley Graphs", J. Graph Theory, Vol. 37, pp. 213-219, 2001, https://doi.org/10.1002/jgt.1016.
- [13] G.J.M. van Wee, "Improved sphere bounds on the covering radius of codes", IEEE Trans. Inform. Theory, Vol. 34, pp. 237-245, 1988, https://doi.org/10.1109/18.2632.
- [14] R. G. Stanton and J. G. Kalbfleisch, "Intersection inequalities for the covering problem", SIAM J. Appl. Math., Vol. 11, pp. 1311-1316, 1969, https://doi.org/10.1137/0117123
- [15] F. Harary, J.P. Hays, H.-J. Wu, "A survey of the theory of hypercube graphs, Comput. Math. Applic., Vol. 15, No. 4, pp. 277-289, 1988, https://doi.org/10.1016/0898-1221(88)90213-1