

# ON WEIGHTED GREEDY-TYPE BASES

HÙNG VIỆT CHU

**ABSTRACT.** In this paper, we study weights for the Thresholding Greedy Algorithm (TGA). While previous work focused on sequential weights  $\varsigma = (s_n)_{n \in \mathbb{N}}$  on each positive integer, we study a more general weight  $\omega = (w_A)_{A \subset \mathbb{N}}$  on each set  $A \subset \mathbb{N}$ . We define and characterize  $\omega$ -(almost) greedy bases. Furthermore, we leverage existing results to show that there exists an  $\omega$ -greedy unconditional basis that is not  $\varsigma$ -almost greedy for any weight sequence  $\varsigma$ . Last but not least, we show the equivalence between  $\omega$ -semi-greedy bases and  $\omega$ -almost greedy bases when  $\omega$  is a so-called structured weight, thus considerably extending the equivalence previously known to hold for sequential weights.

## 1. INTRODUCTION

**1.1. Background.** Let  $(\mathbb{X}, \|\cdot\|)$  be a Banach space over the field  $\mathbb{K} = \mathbb{R}$  or  $\mathbb{C}$  with a semi-normalized Schauder basis  $\mathcal{B} = (e_n)_{n=1}^\infty$  satisfying

$$0 < c_1 := \inf_n \|e_n\| \leq \sup_n \|e_n\| =: c_2 < \infty. \quad (1.1)$$

Let  $(e_n^*)_{n=1}^\infty \subset \mathbb{X}^*$  be the biorthogonal functionals such that  $e_n^*(e_m) = \delta_{n,m}$ . Every  $x \in \mathbb{X}$  can be uniquely written as the series  $\sum_{n=1}^\infty e_n^*(x)e_n$ . Recall that the partial sum operators, defined as  $S_m(x) = \sum_{n=1}^m e_n^*(x)e_n$ , are uniformly bounded. We let  $K_b := \sup_m \|S_m\|$ . It is easy to verify that for a semi-normalized basis, the corresponding biorthogonal functionals are also semi-normalized, i.e.,

$$0 < c_1^* := \inf_n \|e_n^*\| \leq \sup_n \|e_n^*\| =: c_2^* < \infty. \quad (1.2)$$

In 1999, Konyagin and Temlyakov [13] introduced the *Thresholding Greedy Algorithm* (TGA) to approximate each vector  $x$  using finite linear combinations of basis vectors. In particular, for each  $x \in \mathbb{X}$  and  $m \in \mathbb{N}$ , a set  $\Lambda_m(x)$  is called a *greedy set* of order  $m$  of  $x$  if  $|\Lambda_m(x)| = m$  and

$$\min_{n \in \Lambda_m(x)} |e_n^*(x)| \geq \max_{n \notin \Lambda_m(x)} |e_n^*(x)|.$$

For  $x \in \mathbb{X}$ , the TGA produces a sequence of approximating greedy sums  $(G_m(x))_{m=1}^\infty$ , where  $G_m(x) := \sum_{n \in \Lambda_m(x)} e_n^*(x)e_n$ . Here  $G_m(x)$  depends on  $\Lambda_m(x)$ . A basis is *quasi-greedy* if there exists  $C \geq 1$  such that

$$\|x - G_m(x)\| \leq C\|x\|, \forall x \in \mathbb{X}, \forall m \in \mathbb{N}, \forall \Lambda_m(x). \quad (1.3)$$

---

*Date:* February 10, 2023.

*2020 Mathematics Subject Classification.* 41A65; 46B15.

*Key words and phrases.* Thresholding greedy algorithm, greedy, almost greedy, semi-greedy, partially greedy, weight.

The author is thankful to Timur Oikhberg for helpful feedback on earlier drafts of this paper.

The least constant  $C$  satisfying (1.3) is denoted by  $C_\ell$ , and we say  $\mathcal{B}$  is  $C_\ell$ -suppression quasi-greedy. A basis is *greedy* if the TGA gives essentially the best approximation, i.e., there exists a constant  $C \geq 1$  such that

$$\|x - G_m(x)\| \leq C\sigma_m(x), \forall x \in \mathbb{X}, \forall m \in \mathbb{N}, \forall \Lambda_m(x),$$

where

$$\sigma_m(x) := \inf \left\{ \left\| x - \sum_{n \in A} a_n e_n \right\| : A \subset \mathbb{N}, |A| = m, a_n \in \mathbb{K} \right\}.$$

A basis is *almost greedy* if the TGA gives essentially the best projection approximation, i.e., there exists a constant  $C \geq 1$  such that

$$\|x - G_m(x)\| \leq C\tilde{\sigma}_m(x), \forall x \in \mathbb{X}, \forall m \in \mathbb{N}, \forall \Lambda_m(x),$$

where

$$\tilde{\sigma}_m(x) := \inf \left\{ \left\| x - \sum_{n \in A} e_n^*(x) e_n \right\| : A \subset \mathbb{N}, |A| = m \right\}.$$

A beautiful theorem of Konyagin and Temlyakov [13] characterizes greedy bases as being unconditional and democratic (defined later.) In the same spirit, Dilworth et al. [9] characterized almost greedy bases as being quasi-greedy and democratic.

As a variant of (almost) greedy bases, one can introduce the weighted version, where a weight sequence  $\varsigma = (s_n)_{n=1}^\infty \in (0, \infty)^\mathbb{N}$  is involved. Given a set  $A \subset \mathbb{N}$ , the weight of  $A$  is  $s(A) := \sum_{n \in A} s_n$ . For  $\alpha \geq 0$ , we define the error  $\sigma_\alpha^\varsigma(x)$  and  $\tilde{\sigma}_\alpha^\varsigma(x)$

$$\begin{aligned} \sigma_\alpha^\varsigma(x) &:= \inf \left\{ \left\| x - \sum_{n \in A} a_n e_n \right\| : |A| < \infty, s(A) \leq \alpha, a_n \in \mathbb{K} \right\}, \text{ and} \\ \tilde{\sigma}_\alpha^\varsigma(x) &:= \inf \left\{ \left\| x - \sum_{n \in A} e_n^*(x) e_n \right\| : |A| < \infty, s(A) \leq \alpha \right\}. \end{aligned}$$

**Definition 1.1.** A basis  $\mathcal{B}$  is

- (1)  $\varsigma$ -greedy if there exists a constant  $C \geq 1$  such that

$$\|x - G_m(x)\| \leq C\sigma_{s(\Lambda_m(x))}^\varsigma(x), \forall x \in \mathbb{X}, \forall m \in \mathbb{N}, \forall \Lambda_m(x). \quad (1.4)$$

- (2)  $\varsigma$ -almost greedy if there exists a constant  $C \geq 1$  such that

$$\|x - G_m(x)\| \leq C\tilde{\sigma}_{s(\Lambda_m(x))}^\varsigma(x), \forall x \in \mathbb{X}, \forall m \in \mathbb{N}, \forall \Lambda_m(x). \quad (1.5)$$

Weighted greedy-type bases have received much attention and witnessed progresses in various directions: see [2, 4, 5, 10, 12]. Specifically, [4, 10] characterized weighted greedy and weighted almost greedy bases; [2, 5, 10] studied weighted weak semi-greedy bases and weighted semi-greedy bases; [12] investigated weighted partially greedy and weighted reverse partially greedy bases. It follows trivially from [12, Remark 3.3 and Theorem 3.11] that there exists a basis that is weighted partially greedy but is not partially greedy. Furthermore, [10, Remark 4.10] gives an example of a basis that is weighted greedy but is not almost greedy.

**1.2. Weights on sets and main goals.** This paper generalizes sequential weights  $\varsigma$  to weights on sets  $\omega$ . First, we characterize weighted (almost) greedy bases (Section 2). We show that weights on sets are very general in the sense that a basis is unconditional if and only if it is  $\omega$ -greedy for some  $\omega$ . Similarly, a basis is quasi-greedy if and only if it is  $\omega$ -almost greedy for some  $\omega$  (see Corollary 2.11.)

For our next result, it is worth noting that S. J. Dilworth et al. [10] briefly discussed a generalization of sequential weights  $\varsigma$ , denoted by  $\nu$ . Here  $\nu$  satisfies

$$\nu(\emptyset) = 0 \text{ and } \nu(A) \leq \nu(B) \implies \nu(A \setminus B) \leq \nu(B \setminus A).$$

In [10, Remark 2.7], the authors provided an example of a weight  $\nu$  on subsets of  $\mathbb{N}$  that cannot be obtained by any sequential weight  $\varsigma$ . However, it was not known if a  $\nu$ -weighted basis must be  $\varsigma$ -weighted for some sequential weight  $\varsigma$ . Motivated by this, we leverage recent results to obtain an example of an  $\omega$ -greedy basis that is not  $\varsigma$ -greedy for any  $\varsigma$  (Theorem 3.4.)

Last but not least, we study  $\omega$ -semi-greedy and  $\omega$ -partially greedy bases. One notable result is the equivalence between  $\omega$ -semi-greedy and  $\omega$ -almost greedy bases (Theorem 4.2), which considerably extends the same equivalence known to hold for sequential weights  $\varsigma$ .

To prepare for the next section, we give a formal definition of a general weight  $\omega$  on sets and  $\omega$ -(almost) greedy bases.

**Definition 1.2.** Let  $\mathcal{P}(\mathbb{N})$  be the power set of  $\mathbb{N}$ . A weight on sets is a nonnegative function  $\omega : \mathcal{P}(\mathbb{N}) \rightarrow [0, \infty]$  such that

- $w(\emptyset) = 0$ ,
- $w(A) \in (0, \infty]$  for each nonempty  $A \subset \mathbb{N}$ .

For  $B \subset \mathbb{N}$ , define

$$\sigma_B^\omega(x) := \inf \left\{ \left\| x - \sum_{n \in A} a_n e_n \right\| : A \in \mathbb{N}^{<\infty}, w(A \setminus B) \leq w(B \setminus A), a_n \in \mathbb{K} \right\},$$

where  $\mathbb{N}^{<\infty}$  is the set of all finite subsets of  $\mathbb{N}$ .

The following definition of  $\omega$ -greedy bases generalizes the classical weighted bases (see [4, Definition 1.1].)

**Definition 1.3.** A basis  $\mathcal{B}$  is  $\omega$ -greedy if there exists a constant  $C \geq 1$  such that for all  $x \in \mathbb{X}$ ,  $m \in \mathbb{N}$ , and  $\Lambda_m(x)$ ,

$$\|x - G_m(x)\| \leq C \sigma_{\Lambda_m(x)}^\omega(x).$$

The least constant  $C$  is denoted by  $C_g^\omega$ .

In a similar manner, we define  $\omega$ -almost greedy. For  $B \subset \mathbb{N}$ , let

$$\tilde{\sigma}_B^\omega(x) := \inf \{ \|x - P_A(x)\| : A \in \mathbb{N}^{<\infty}, w(A \setminus B) \leq w(B \setminus A) \},$$

where  $P_A(x) = \sum_{n \in A} e_n^*(x) e_n$ .

**Definition 1.4.** A basis  $\mathcal{B}$  is  $\omega$ -almost greedy if there exists a constant  $C \geq 1$  such that for all  $x \in \mathbb{X}$ ,  $m \in \mathbb{N}$ , and  $\Lambda_m(x)$ ,

$$\|x - G_m(x)\| \leq C \tilde{\sigma}_{\Lambda_m(x)}^\omega(x).$$

The least constant  $C$  is denoted by  $C_{al}^\omega$ .

**Remark 1.5.** It is easy to check that (1.4) and (1.5) are special cases of Definitions 1.3 and 1.4, respectively.

## 2. CHARACTERIZATIONS OF $\omega$ -(ALMOST) GREEDY BASES

In order to characterize  $\omega$ -(almost) greedy bases, we need the notion of unconditionality and  $\omega$ -Property (A).

**Definition 2.1.** A basis  $\mathcal{B}$  is unconditional if there exists  $C \geq 1$  such that

$$\|P_A(x)\| \leq C\|x\|, \forall x \in \mathbb{X}, \forall A \subset \mathbb{N}.$$

In this case, we say that  $\mathcal{B}$  is  $C$ -suppression unconditional. The least such  $C$  is denoted by  $\mathbf{K}_s$ . For an unconditional basis, there also exists a constant  $\mathbf{K}_u$  such that

$$\left\| \sum_{n=1}^N a_n e_n \right\| \leq \mathbf{K}_u \left\| \sum_{n=1}^N b_n e_n \right\|,$$

for all  $N \geq 1$  and for all scalars  $a_n, b_n$  with  $|a_n| \leq |b_n|$ .

Let

$$1_A = \sum_{n \in A} e_n \text{ and } 1_{\varepsilon A} = \sum_{n \in A} \varepsilon_n e_n,$$

where  $\varepsilon = (\varepsilon_n)_{n=1}^\infty \subset \mathbb{K}$  and  $|\varepsilon_n| = 1$ . For  $x \in \mathbb{X}$ ,  $\text{supp}(x) := \{n : e_n^*(x) \neq 0\}$ ,  $\|x\|_\infty := \sup_n |e_n^*(x)|$ , and we write  $A \sqcup B \sqcup x$  to indicate that  $A, B$ , and  $\text{supp}(x)$  are pairwise disjoint. Finally,  $\text{sgn}(e_n^*(x)) = \begin{cases} 1 & \text{if } e_n^*(x) = 0, \\ e_n^*(x)/|e_n^*(x)| & \text{if } e_n^*(x) \neq 0. \end{cases}$

**Definition 2.2.** A basis  $\mathcal{B}$  has  $\omega$ -Property (A) if there exists  $C \geq 1$  such that

$$\|x + 1_{\varepsilon A}\| \leq C\|x + 1_{\delta B}\|,$$

for all  $A, B \in \mathbb{N}^{<\infty}$  with  $w(A) \leq w(B)$ , signs  $(\varepsilon), (\delta)$ , and  $x \in \mathbb{X}$  with  $\|x\|_\infty \leq 1$  and  $A \sqcup B \sqcup x$ . The least constant  $C$  is denoted by  $\mathbf{C}_b^\omega$ .

**Theorem 2.3.** Let  $\mathcal{B}$  be a basis and  $\omega$  be a weight on subsets of  $\mathbb{N}$ .

- (1) If  $\mathcal{B}$  is  $\mathbf{C}_g^\omega$ - $\omega$ -greedy, then  $\mathcal{B}$  is  $\mathbf{C}_g^\omega$ -suppression unconditional and satisfies  $\mathbf{C}_g^\omega$ - $\omega$ -Property (A).
- (2) If  $\mathcal{B}$  is  $\mathbf{K}_s$ -suppression unconditional and satisfies  $\mathbf{C}_b^\omega$ - $\omega$ -Property (A), then  $\mathcal{B}$  is  $\mathbf{K}_s \mathbf{C}_b^\omega$ - $\omega$ -greedy.

First, we need an useful reformulation of  $\omega$ -Property (A).

**Lemma 2.4.** A basis  $\mathcal{B}$  has  $\mathbf{C}_b^\omega$ - $\omega$ -Property (A) if and only if

$$\|x\| \leq \mathbf{C}_b^\omega \|x - P_A(x) + 1_{\varepsilon B}\|, \tag{2.1}$$

for all  $x \in \mathbb{X}$  with  $\|x\|_\infty \leq 1$ ,  $A, B \in \mathbb{N}^{<\infty}$  with  $w(A) \leq w(B)$  and  $B \cap (A \cup \text{supp}(x)) = \emptyset$ , and  $\text{sign}(\varepsilon)$ .

*Proof.* Assume that  $\mathcal{B}$  has  $\mathbf{C}_b^\omega$ - $\omega$ -Property (A). Let  $x, A, B, (\varepsilon)$  be chosen as in (2.1). We have

$$\begin{aligned} \|x\| &= \left\| x - P_A(x) + \sum_{n \in A} e_n^*(x) e_n \right\| \leq \sup_{(\delta)} \|x - P_A(x) + 1_{\delta A}\| \\ &\leq \mathbf{C}_b^\omega \|x - P_A(x) + 1_{\varepsilon B}\|, \end{aligned}$$

as desired.

Next, assume that  $\mathcal{B}$  satisfies (2.1). Let  $x, A, B, (\varepsilon), (\delta)$  be chosen as in Definition 2.2. Let  $y = x + 1_{\varepsilon A}$ . By (2.1),

$$\|x + 1_{\varepsilon A}\| = \|y\| \leq \mathbf{C}_b^\omega \|y - P_A(y) + 1_{\delta B}\| = \mathbf{C}_b^\omega \|x + 1_{\delta B}\|.$$

This completes our proof.  $\square$

**Proposition 2.5.** *Let  $\mathcal{B} = (e_n)_{n=1}^\infty$  be a  $\mathbf{K}_s$ -suppression unconditional basis. Fix  $N \in \mathbb{N}$ . For any scalars  $a_1, \dots, a_N, b_1, \dots, b_N$  so that either  $a_0 = 0$  or  $\text{sgn}(a_n) = \text{sgn}(b_n)$  and  $|a_n| \leq |b_n|$  for all  $1 \leq n \leq N$ , we have*

$$\left\| \sum_{n=1}^N a_n e_n \right\| \leq \mathbf{K}_s \left\| \sum_{n=1}^N b_n e_n \right\|.$$

*Proof.* See [1, Proposition 2.1].  $\square$

*Proof of Theorem 2.3.* Assume that  $\mathcal{B}$  is  $\mathbf{C}_g^\omega$ - $\omega$ -greedy. Let  $x \in \mathbb{X}$  and  $B \in \mathbb{N}^{<\infty}$ . Write

$$y = \sum_{n \in B} (\alpha + e_n^*(x)) e_n + P_{B^c}(x),$$

where  $\alpha$  is chosen sufficiently large such that  $B$  is a greedy set of  $y$ . Then

$$\|P_{B^c}(x)\| = \|y - G_{|B|}(y)\| \leq \mathbf{C}_g^\omega \sigma_B^\omega(y) \leq \mathbf{C}_g^\omega \|y - \alpha 1_B\| = \mathbf{C}_g^\omega \|x\|.$$

Hence,  $\mathcal{B}$  is  $\mathbf{C}_g^\omega$ -suppression unconditional. Next, we prove  $\omega$ -Property (A). We choose  $x, A, B, (\varepsilon), (\delta)$  as in Definition 2.2. Set  $y = x + 1_{\varepsilon A} + 1_{\delta B}$ . Then  $B$  is a greedy set of  $y$ . We have

$$\|x + 1_{\varepsilon A}\| = \|y - G_{|B|}(y)\| \leq \mathbf{C}_g^\omega \sigma_B^\omega(y) \leq \mathbf{C}_g^\omega \|y - P_A(y)\| = \mathbf{C}_g^\omega \|x + 1_{\delta B}\|.$$

This completes the proof.

Now we assume that  $\mathcal{B}$  is  $\mathbf{K}_s$ -suppression unconditional and satisfies  $\mathbf{C}_b^\omega$ - $\omega$ -Property (A). Let  $x \in \mathbb{X}$  have a greedy set  $A \subset \mathbb{N}$ . Let  $B \in \mathbb{N}^{<\infty}$  with  $w(B \setminus A) \leq w(A \setminus B)$ . Also, choose arbitrary  $(b_n)_{n \in B} \subset \mathbb{K}$ . Let  $\alpha := \min_{n \in A} |e_n^*(x)|$ . By Lemma 2.4 and

Proposition 2.5, we have

$$\begin{aligned}
\|x - P_A(x)\| &\leq \mathbf{C}_b^\omega \left\| x - P_A(x) - P_{B \setminus A}(x) + \alpha \sum_{n \in A \setminus B} \text{sgn}(e_n^*(x)) e_n \right\| \\
&\leq \mathbf{C}_b^\omega \left\| P_{(A \cup B)^c}(x) + \alpha \sum_{n \in A \setminus B} \text{sgn}(e_n^*(x)) e_n \right\| \\
&\leq \mathbf{K}_s \mathbf{C}_b^\omega \left\| P_{(A \cup B)^c}(x) + \sum_{n \in B} (e_n^*(x) - b_n) e_n + P_{A \setminus B}(x) \right\| \\
&= \mathbf{K}_s \mathbf{C}_b^\omega \left\| x - \sum_{n \in B} b_n e_n \right\|.
\end{aligned}$$

This completes our proof that  $\mathcal{B}$  is  $\mathbf{K}_s \mathbf{C}_b^\omega$ - $\omega$ -greedy.  $\square$

When we do not need tight estimates, the notion of  $\omega$ -disjoint (super)democracy can play the role of  $\omega$ -Property (A), providing other characterizations of  $\omega$ -greedy bases.

**Definition 2.6.** A basis  $\mathcal{B}$  is  $\omega$ -disjoint democratic ( $\omega$ -disjoint superdemocratic, respectively) if there exists  $C \geq 1$  such that

$$\|1_A\| \leq C \|1_B\|, (\|1_{\varepsilon A}\| \leq C \|1_{\delta B}\|, \text{ respectively}),$$

for all  $A, B \in \mathbb{N}^{<\infty}$  with  $w(A) \leq w(B)$ ,  $A \cap B = \emptyset$  and signs  $(\varepsilon), (\delta)$ . The least constant  $C$  is denoted by  $\mathbf{C}_{d, \sqcup}^\omega$  (and  $\mathbf{C}_{sd, \sqcup}^\omega$ , respectively.)

**Remark 2.7.** A basis  $\mathcal{B}$  is said to be  $\omega$ -(super)democratic if in Definition 2.6, we drop the requirement  $A \cap B = \emptyset$ ;  $\mathcal{B}$  is said to be (super)democratic if it is  $\omega$ -(super)democratic for  $\omega$  being the cardinality weight, i.e.,  $w(A) = |A|, \forall A \subset \mathbb{N}$ .

**Theorem 2.8.** Let  $\mathcal{B}$  be a basis and  $\omega$  be a weight on subsets of  $\mathbb{N}$ . The following are equivalent:

- (1)  $\mathcal{B}$  is  $\omega$ -greedy,
- (2)  $\mathcal{B}$  is unconditional and satisfies  $\omega$ -Property (A),
- (3)  $\mathcal{B}$  is unconditional and  $\omega$ -disjoint superdemocratic,
- (4)  $\mathcal{B}$  is unconditional and  $\omega$ -disjoint democratic.

*Proof.* By Theorem 2.3, we know that (1)  $\iff$  (2). It follows immediately from definitions that  $\omega$ -Property (A)  $\implies \omega$ -disjoint superdemocratic  $\implies \omega$ -disjoint democratic. Hence, (2)  $\implies$  (3)  $\implies$  (4). It remains to show that (4)  $\implies$  (2). Let  $x, A, B, (\varepsilon), (\delta)$  be chosen as in Definition 2.2. We have

$$\begin{aligned}
\|x + 1_{\varepsilon A}\| &\leq \|x\| + \|1_{\varepsilon A}\| \leq \|x\| + \mathbf{K}_u \|1_A\| \\
&\leq \|x\| + \mathbf{K}_u \mathbf{C}_{d, \sqcup}^\omega \|1_B\| \\
&\leq \mathbf{K}_s \|x + 1_{\delta B}\| + \mathbf{K}_u^2 \mathbf{C}_{d, \sqcup}^\omega \|x + 1_{\delta B}\| \\
&= (\mathbf{K}_s + \mathbf{K}_u^2 \mathbf{C}_{d, \sqcup}^\omega) \|x + 1_{\delta B}\|.
\end{aligned}$$

This completes our proof.  $\square$

For  $\omega$ -almost greedy bases, corresponding results hold. We include the proof of the next theorem in the Appendix.

**Theorem 2.9.** *Let  $\mathcal{B}$  be a basis and  $\omega$  be a weight on subsets of  $\mathbb{N}$ .*

- (1) *If  $\mathcal{B}$  is  $\mathbf{C}_{al}^\omega$ - $\omega$ -almost greedy, then  $\mathcal{B}$  is  $\mathbf{C}_{al}^\omega$ -suppression quasi-greedy and satisfies  $\mathbf{C}_{al}^\omega$ - $\omega$ -Property (A).*
- (2) *If  $\mathcal{B}$  is  $\mathbf{C}_\ell$ -suppression quasi-greedy and satisfies  $\mathbf{C}_b^\omega$ - $\omega$ -Property (A), then  $\mathcal{B}$  is  $\mathbf{C}_\ell \mathbf{C}_b^\omega$ - $\omega$ -almost greedy.*

**Theorem 2.10.** *Let  $\mathcal{B}$  be a basis and  $\omega$  be a weight on subsets of  $\mathbb{N}$ . The following are equivalent:*

- (1)  *$\mathcal{B}$  is  $\omega$ -almost greedy,*
- (2)  *$\mathcal{B}$  is quasi-greedy and satisfies  $\omega$ -Property (A),*
- (3)  *$\mathcal{B}$  is quasi-greedy and  $\omega$ -disjoint superdemocratic,*
- (4)  *$\mathcal{B}$  is quasi-greedy and  $\omega$ -disjoint democratic.*

**Corollary 2.11.** (1) *A basis  $\mathcal{B}$  is unconditional if and only if  $\mathcal{B}$  is  $\omega$ -greedy for some weight  $\omega$ .*

(2) *A basis  $\mathcal{B}$  is quasi-greedy if and only if  $\mathcal{B}$  is  $\omega$ -almost greedy for some weight  $\omega$ .*

*Proof.* (1) If  $\mathcal{B}$  is  $\omega$ -greedy for some weight  $\omega$ , then  $\mathcal{B}$  is unconditional by Theorem 2.3. Conversely, suppose that  $\mathcal{B}$  is unconditional. Define the weight  $\omega$  on a set  $A$

$$\omega(A) = \begin{cases} \|1_A\| & \text{if } A \text{ is finite,} \\ \infty & \text{if } A \text{ is infinite.} \end{cases}$$

By Theorem 2.8, it suffices to show that  $\mathcal{B}$  is  $\omega$ -disjoint democratic. This is clearly true since for two finite sets  $A, B$  with  $\omega(A) \leq \omega(B)$ , we get  $\|1_A\| \leq \|1_B\|$  by the definition of  $\omega$ .

The proof of (2) is similar to that of (1). □

### 3. A SET-WEIGHTED-GREEDY BASIS THAT IS NOT SEQUENCE-WEIGHTED-GREEDY

The following theorem provides a necessary condition for a basis to be  $\varsigma$ -greedy for some weight sequence  $\varsigma$ .

**Theorem 3.1.** *If a basis  $\mathcal{B} = (e_n)_{n=1}^\infty$  is  $\varsigma$ -(almost) greedy for some weight sequence  $\varsigma$ , then either  $\mathcal{B}$  is (almost) greedy or there exists a subsequence  $(e_{n_k})_{k=1}^\infty$  equivalent to the canonical basis of  $c_0$ .*

Observe that  $\varsigma$ -Property (A) is  $\omega$ -Property (A) when the weight  $\omega$  on sets is determined by a weight sequence  $\varsigma$ . Particularly, Property (A) (first introduced in [1] and later generalized in [8]) is  $\varsigma$ -Property (A) when  $\varsigma = (1, 1, \dots)$ .

**Definition 3.2.** A basis  $\mathcal{B}$  has  $\varsigma$ -Property (A) if there exists  $C \geq 1$  such that

$$\|x + 1_{\varepsilon A}\| \leq C\|x + 1_{\delta B}\|,$$

for all  $A, B \in \mathbb{N}^{<\infty}$  with  $s(A) \leq s(B)$ , signs  $(\varepsilon), (\delta)$ , and  $x \in \mathbb{X}$  with  $\|x\|_\infty \leq 1$  and  $A \sqcup B \sqcup x$ . The least constant  $C$  is denoted by  $\mathbf{C}_b^\varsigma$ . As a special case, a basis  $\mathcal{B}$  is said to have Property (A) if it has  $\varsigma$ -Property (A) for  $\varsigma = (1, 1, \dots)$ .



*Proof of Theorem 3.1.* We assume that  $\mathcal{B}$  is  $\varsigma$ -greedy for some weight sequence  $\varsigma = (s(n))_{n=1}^\infty$ . By [4, Theorem 4.1],  $\mathcal{B}$  is unconditional and has  $\varsigma$ -Property (A).

If  $0 < \inf s(n) \leq \sup s(n) < \infty$ , then [4, Proposition 3.5] implies that  $\mathcal{B}$  has Property (A). According to [8, Theorem 2], we know that  $\mathcal{B}$  is greedy.

If  $\sup s(n) = \infty$ , then [4, Proposition 3.10] states that  $\mathcal{B}$  is equivalent to the canonical basis of  $c_0$  and thus, is greedy.

If  $\inf s(n) = 0$ , then by [4, Proposition 3.10],  $(e_n)_{n=1}^\infty$  has a subsequence  $(e_{n_k})_{k=1}^\infty$  that is equivalent to the canonical basis of  $c_0$ .

The proof of the almost greedy case is similar.  $\square$

We now state the existence of an  $\omega$ -greedy basis that is not  $\varsigma$ -almost greedy for any weight sequence  $\varsigma$ . We can, in particular, require the weight  $\omega$  to have a more rigid structure than in Definition 1.2. For conciseness, we let  $w_n := w(\{n\})$ .

**Definition 3.3.** A structured weight is a nonnegative function  $\omega : \mathcal{P}(\mathbb{N}) \rightarrow [0, \infty]$  such that

- (a)  $w(\emptyset) = 0$ ,
- (b)  $w(A) < \infty$  if  $|A| < \infty$ ,
- (c)  $w(A) \in (0, \infty]$  for each nonempty  $A \subset \mathbb{N}$ ,
- (d)  $w(A) \rightarrow 0$  as  $\sum_{n \in A} w_n \rightarrow 0$ ,
- (e)  $w(A) \rightarrow \infty$  as  $\sum_{n \in A} w_n \rightarrow \infty$ ,
- (f) There exists an arbitrarily large number  $N \in \mathbb{N}$  such that there exists an  $\varepsilon > 0$  satisfying  $w(\{N, n\}) - w_n > \varepsilon$  for all  $n \in \mathbb{N}, n \neq N$ .

Conditions (a), (b), and (c) are almost the same as what we have in Definition 1.2, except that we now require the weight on a finite set to be finite. Conditions (d) and (e) are reasonable. Condition (d) states that the weight on a set approaches 0 when the sum of weights of its singletons approaches 0, while (e) states the same condition with 0 replaced by  $\infty$ . Throughout this paper, we will specify whether we need structured weights in our results. If we state a result without mentioning structured weights, then the result holds for weights in Definition 1.2.

**Theorem 3.4.** *There exists a basis that is  $\omega$ -greedy for some structured weight  $\omega$  on sets but is not  $\varsigma$ -almost greedy for any weight sequence  $\varsigma$  on positive integers.*

*Proof.* By Theorem 3.1, an unconditional basis that is neither democratic nor has a subsequence equivalent to the canonical basis of  $c_0$  is not  $\varsigma$ -almost greedy on any  $\varsigma$ .

Set  $P := \{2^k : k \geq 1\}$ ,  $a_n = 1/n^{1/2}$  and  $b_n = 1/n$  for  $n \geq 1$ . Let  $\mathbb{X}$  be the completion of  $c_{00}$  under the following norm: for  $x = (x_1, x_2, \dots) \in c_{00}$ , define

$$\|x\| := \left( \sup_{\sigma} \sum_{i \in P} a_{\sigma(i)} |x_i| \right) + \left( \sup_{\pi} \sum_{i \notin P} b_{\pi(i)} |x_i| \right),$$

where  $\sigma : P \rightarrow \mathbb{N}$  and  $\pi : \mathbb{N} \setminus P \rightarrow \mathbb{N}$  are bijections. Let  $\mathcal{B} = (e_n)_{n=1}^\infty$  be the canonical basis. Clearly,  $\mathcal{B}$  is unconditional and normalized. However,  $\mathcal{B}$  is not democratic. Indeed, fix  $N \in \mathbb{N}$  and set  $A = \{3^1, 3^2, \dots, 3^N\}$ ,  $B = \{2^1, 2^2, \dots, 2^N\}$ . We have

$$\|1_A\| = \sum_{n=1}^N \frac{1}{n} \sim \ln(N) \text{ and } \|1_B\| = \sum_{n=1}^N \frac{1}{\sqrt{n}} \sim \sqrt{N}.$$



Since  $\|1_B\|/\|1_A\| \sim \sqrt{N}/\ln(N) \rightarrow \infty$  as  $N \rightarrow \infty$ , we know that  $\mathcal{B}$  is not democratic and thus, not almost greedy.

By the proof of Corollary 2.11,  $\mathcal{B}$  is  $\omega$ -greedy for the following weight  $\omega$

$$\omega(A) = \begin{cases} \|1_A\| & \text{if } A \text{ is finite,} \\ \infty & \text{if } A \text{ is infinite.} \end{cases}$$

It is easy to check that  $\omega$  is a structured weight.

We claim that there is no subsequence of  $\mathcal{B} = (e_n)_{n=1}^\infty$  that is equivalent to the canonical basis of  $c_0$ . Indeed, pick any subsequence  $(e_{n_k})_{k=1}^\infty$  of  $\mathcal{B}$ . For  $N \in \mathbb{N}$ , we have

$$\left\| \sum_{k=1}^N e_{n_k} \right\| \geq \sum_{n=1}^N \frac{1}{n} \sim \ln(N).$$

Hence,  $(e_{n_k})_{k=1}^\infty$  is not equivalent to the canonical basis of  $c_0$ . Therefore, Theorem 3.1 and the fact that  $\mathcal{B}$  is not almost greedy tell us that  $\mathcal{B}$  is not  $\varsigma$ -almost greedy for any weight sequence  $\varsigma$ .  $\square$

#### 4. $\omega$ -SEMI-GREEDY BASES

First, we define the  $\omega$ -version of the classical semi-greedy bases (first introduced in [11]). Corresponding to each greedy set  $\Lambda_m(x)$ , there is a so-called *Chebyshev greedy sum* of order  $m$ , denoted by  $CG_m(x)$ , such that

- (1)  $\text{supp}(CG_m(x)) \subset \Lambda_m(x)$  and
- (2) we have

$$\|x - CG_m(x)\| = \min \left\{ \left\| x - \sum_{n \in \Lambda_m(x)} a_n e_n \right\| : (a_n) \subset \mathbb{K} \right\}.$$

**Definition 4.1.** A basis  $\mathcal{B}$  is  $\omega$ -semi-greedy if there exists a constant  $C \geq 1$  such that for all  $x \in \mathbb{X}$ ,  $m \in \mathbb{N}$ , and  $\Lambda_m(x)$ ,

$$\|x - CG_m(x)\| \leq C \sigma_{\Lambda_m(x)}^\omega(x).$$

The least constant  $C$  is denoted by  $C_s^\omega$ .

The main goal of this section is to establish the following theorem, which, by Theorem 3.4, is a nontrivial extension of [5, Theorem 1.10].

**Theorem 4.2.** *Let  $\omega$  be a structured weight. Then  $\mathcal{B}$  is  $\omega$ -semi-greedy if and only if it is  $\omega$ -almost greedy.*

**Proposition 4.3.** *Let  $\mathcal{B}$  be a  $C_s^\omega$ - $\omega$ -semi-greedy basis, where  $\omega$  is structured.*

- (1) *Let  $B \in \mathbb{N}^{<\infty}$  and  $w(B) \leq \limsup_{n \rightarrow \infty} w_n$ . Then we have*

$$\sup_{(\varepsilon)} \|1_{\varepsilon B}\| \leq 2K_b C_s^\omega c_2,$$

where  $c_2$  is in (1.1).

- (2) *If  $\sup_n w_n = \infty$  or  $\sum_n w_n < \infty$ , then  $\mathcal{B}$  is equivalent to the canonical basis of  $c_0$ .*

(3) If  $\inf_n w_n = 0$ , then  $\mathcal{B}$  contains a subsequence equivalent to the canonical basis of  $c_0$ .

**Remark 4.4.** The conclusions in Proposition 4.3 still hold if our basis  $\mathcal{B}$  is  $\mathbf{C}_{sd, \sqcup}^\omega$ -disjoint superdemocratic. The proof is left for interested readers.

*Proof.* (1) Pick  $B \in \mathbb{N}^{<\infty}$  and  $(\varepsilon)$ . Choose  $N_1 > \max B$  be the number in condition (f) of a structured weight such that there exists  $\varepsilon > 0$  satisfying  $w(\{N_1, n\}) > w_n + \varepsilon$  for all  $n \neq N_1$ . It follows that

$$\limsup_{n \rightarrow \infty} w(\{N_1, n\}) \geq \limsup_{n \rightarrow \infty} w_n + \varepsilon \geq w(B) + \varepsilon.$$

Pick  $N_2 > N_1$  such that  $w(\{N_1, N_2\}) > w(B)$ . (This is possible due to condition (b).) Set  $x := 1_{\varepsilon B} + e_{N_1} + e_{N_2}$ . Then  $\{N_1, N_2\}$  is a greedy set of  $x$ . Let  $\|x - CG_2(x)\| = \|1_{\varepsilon B} + \alpha_1 e_{N_1} + \alpha_2 e_{N_2}\|$  for some  $\alpha_1, \alpha_2 \in \mathbb{K}$ . We have

$$\begin{aligned} \|1_{\varepsilon B}\| &\leq \mathbf{K}_b \|1_{\varepsilon B} + \alpha_1 e_{N_1} + \alpha_2 e_{N_2}\| \leq \mathbf{K}_b \mathbf{C}_s^\omega \sigma_{\{N_1, N_2\}}^\omega(x) \\ &\leq \mathbf{K}_b \mathbf{C}_s^\omega \|e_{N_1} + e_{N_2}\| \leq 2\mathbf{K}_b \mathbf{C}_s^\omega c_2. \end{aligned}$$

(2) If  $\sup_n w_n = \infty$ , then by (1),  $\sup_{(\varepsilon)} \|1_{\varepsilon B}\| \leq 2\mathbf{K}_b \mathbf{C}_s^\omega c_2, \forall B \in \mathbb{N}^{<\infty}$ . Hence, the basis is equivalent to the canonical basis of  $c_0$ . If  $\sum_n w_n < \infty$ , then choose  $N \in \mathbb{N}$  such that  $\sum_{n=N+1}^\infty w_n$  is so small that  $w(E) < w_1$  for all  $E \subset \mathbb{N}_{\geq N+1}$ . This can be done due to condition (d) of  $\omega$ . We claim that for any  $B \in \mathbb{N}^{<\infty}$  and any sign  $(\varepsilon)$ , we have  $\|1_{\varepsilon B}\| = O(1)$ . Let  $B_1 = B \cap [1, N]$  and  $B_2 = B \cap [N+1, \infty)$ . Observe that

$$\|1_{\varepsilon B}\| \leq \|1_{\varepsilon B_1}\| + \|1_{\varepsilon B_2}\| \leq Nc_2 + \|1_{\varepsilon B_2}\|.$$

Set  $x := e_1 + 1_{\varepsilon B_2}$ . Then  $\{1\}$  is a greedy set of  $x$ . Let  $\|x - CG_1(x)\| = \|\alpha e_1 + 1_{\varepsilon B_2}\|$  for some  $\alpha \in \mathbb{K}$ . Since  $w(B_2) < w_1$ , we have

$$\begin{aligned} \|1_{\varepsilon B_2}\| &\leq (\mathbf{K}_b + 1) \|\alpha e_1 + 1_{\varepsilon B_2}\| \leq (\mathbf{K}_b + 1) \mathbf{C}_s^\omega \sigma_{\{1\}}^\omega(x) \\ &\leq (\mathbf{K}_b + 1) \mathbf{C}_s^\omega \|e_1\| \leq (\mathbf{K}_b + 1) \mathbf{C}_s^\omega c_2. \end{aligned}$$

This completes our proof that  $\|1_{\varepsilon B}\| = O(1)$  and so,  $\mathcal{B}$  is equivalent to the canonical basis of  $c_0$ .

(3) Choose a subsequence  $(n_k)_{k=1}^\infty$  such that  $\sum_{k=1}^\infty w_{n_k} < \infty$  and apply (2).  $\square$

**Theorem 4.5.** Let  $\omega$  be a structured weight. If a basis  $\mathcal{B}$  is  $\omega$ -semi-greedy, then it is quasi-greedy and  $\omega$ -superdemocratic.

*Proof.* Suppose that  $\sum_{n=1}^\infty w_n < \infty$  or  $\sup_n w_n = \infty$ . By Proposition 4.3, we know that  $\mathcal{B}$  is equivalent to the canonical basis of  $c_0$ , and the desired conclusion follows trivially. For the rest of the proof, let us assume that  $\sum_{n=1}^\infty w_n = \infty$  and  $\sup_n w_n < \infty$ .

Quasi-greedy: Let  $x \in \mathbb{X}$  with  $\|x\|_\infty \leq 1$ ,  $|\text{supp}(x)| < \infty$ , and a greedy set  $\Lambda_m(x)$ .

Case 1:  $w(\Lambda_m(x)) \leq \limsup_{n \rightarrow \infty} w_n$ . By Proposition 4.3, we have

$$\sup_{(\varepsilon)} \|1_{\varepsilon \Lambda_m(x)}\| \leq 2\mathbf{K}_b \mathbf{C}_s^\omega c_2.$$

By norm convexity,

$$\begin{aligned} \|P_{\Lambda_m(x)}(x)\| &\leq \max_n |e_n^*(x)| \sup_{(\varepsilon)} \|1_{\varepsilon\Lambda_m(x)}\| \\ &\leq \sup_n \|e_n^*\| \|x\| \cdot 2\mathbf{K}_b \mathbf{C}_s^\omega c_2 \leq 2\mathbf{K}_b \mathbf{C}_s^\omega c_2 c_2^* \|x\|, \end{aligned}$$

where  $c_2$  and  $c_2^*$  are in (1.1) and (1.2), respectively.

Case 2:  $w(\Lambda_m(x)) > \limsup_{n \rightarrow \infty} w_n$ . We build a finite set  $E$  as follows: choose  $N > \max \text{supp}(x)$  such that  $w_N \leq w(\Lambda_m(x))$ . Let  $k$  be the smallest positive integer verifying

$$w(\{N, N+1, \dots, N+k\}) \leq w(\Lambda_m(x)) < w(\{N, N+1, \dots, N+k, N+k+1\}).$$

We know such  $k$  exists due to  $\sum_n w_n = \infty$  and condition (e) of a structured weight. Let  $A = \{N, N+1, \dots, N+k\}$  and  $B = A \cup \{N+k+1\}$ . Define

$$y := x - P_{\Lambda_m(x)}(x) + \alpha 1_B,$$

where  $\alpha := \min_{n \in \Lambda_m(x)} |e_n^*(x)|$ . Since  $B$  is a greedy set of  $y$ , by  $\mathbf{C}_s^\omega$ - $\omega$ -semi-greediness, there exist  $(b_n)_{n \in B} \subset \mathbb{K}$  such that

$$\begin{aligned} \|x - P_{\Lambda_m(x)}(x)\| &\leq \mathbf{K}_b \left\| x - P_{\Lambda_m(x)}(x) + \sum_{n \in B} b_n e_n \right\| \leq \mathbf{K}_b \mathbf{C}_s^\omega \sigma_B^\omega(y) \\ &\leq \mathbf{K}_b \mathbf{C}_s^\omega \|x + \alpha 1_B\| \leq \mathbf{K}_b \mathbf{C}_s^\omega (\|x\| + \alpha \|1_A\| + \alpha \|e_{N+k+1}\|). \end{aligned} \quad (4.1)$$

Pick  $j \in \Lambda_m(x)$ . We have

$$\alpha \|e_{N+k+1}\| \leq \alpha c_2 \leq c_2 |e_j^*(x)| \leq c_2 \|e_j^*\| \|x\| \leq c_2 c_2^* \|x\|. \quad (4.2)$$

It remains to bound  $\alpha \|1_A\|$ . Let  $z := x + \alpha 1_A$ . Since  $\Lambda_m(x)$  is a greedy set of  $z$ ,  $\mathbf{C}_s^\omega$ - $\omega$ -semi-greediness gives  $(t_n)_{n \in \Lambda_m(x)} \subset \mathbb{K}$  such that

$$\begin{aligned} \|\alpha 1_A\| &\leq (\mathbf{K}_b + 1) \left\| \sum_{n \in \Lambda_m(x)} t_n e_n + P_{\Lambda_m(x)^c}(x) + \alpha 1_A \right\| \\ &\leq (\mathbf{K}_b + 1) \mathbf{C}_s^\omega \sigma_{\Lambda_m(x)}^\omega(z) \leq (\mathbf{K}_b + 1) \mathbf{C}_s^\omega \|x\|. \end{aligned} \quad (4.3)$$

From (4.1), (4.2), and (4.3), we have shown that

$$\|x - P_{\Lambda_m(x)}(x)\| = O(\|x\|).$$

This completes our proof that  $\mathcal{B}$  is quasi-greedy.

$\omega$ -superdemocratic: Let  $A, B \in \mathbb{N}^{<\infty}$  with  $w(A) \leq w(B)$ . Pick signs  $(\varepsilon), (\delta)$ .

Case 1:  $w(A) \leq w(B) \leq \limsup_{n \rightarrow \infty} w_n$ . By Proposition 4.3, we know that

$$\|1_{\varepsilon A}\| \leq 2\mathbf{K}_b \mathbf{C}_s^\omega c_2.$$

On the other hand, if  $j = \min B$ , then

$$\|1_{\delta B}\| \geq \|e_j\| / \mathbf{K}_b \geq c_1 / \mathbf{K}_b,$$

where  $c_1$  is in (1.1). Therefore,

$$\|1_{\varepsilon A}\| \leq 2\mathbf{K}_b^2 \mathbf{C}_s^\omega \frac{c_2}{c_1} \|1_{\delta B}\|.$$

Case 2:  $w(B) > \limsup_{n \rightarrow \infty} w_n$ . As when we prove quasi-greediness, choose  $E$  and  $F = E \cup \{N\}$  such that  $A \cup B < E < \{N\}$  and  $w(E) \leq w(B) < w(F)$ . Set  $x := 1_{\varepsilon A} + 1_F$ . Then  $F$  is a greedy set of  $x$ . By  $\mathbf{C}_s^\omega$ - $\omega$ -semi-greediness, there exist  $(a_n)_{n \in F} \subset \mathbb{K}$  such that

$$\|1_{\varepsilon A}\| \leq \mathbf{K}_b \left\| 1_{\varepsilon A} + \sum_{n \in F} a_n e_n \right\| \leq \mathbf{K}_b \mathbf{C}_s^\omega \sigma_F^\omega(x) \leq \mathbf{K}_b \mathbf{C}_s^\omega \|1_F\|. \quad (4.4)$$

Now, let  $y = 1_{\delta B} + 1_E$ . Since  $B$  is a greedy set of  $y$ , by  $\mathbf{C}_s^\omega$ - $\omega$ -semi-greediness, we obtain

$$\|1_E\| \leq (\mathbf{K}_b + 1) \left\| \sum_{n \in B} b_n e_n + 1_E \right\| \leq \mathbf{C}_s^\omega (\mathbf{K}_b + 1) \sigma_B^\omega(y) \leq \mathbf{C}_s^\omega (\mathbf{K}_b + 1) \|1_{\delta B}\|, \quad (4.5)$$

for some  $(b_n)_{n \in B} \subset \mathbb{K}$ . Furthermore, if  $u = \min E$ ,

$$\|1_F\| \leq \|1_E\| + \|e_N\| \leq \|1_E\| + c_2 \leq \|1_E\| + \frac{c_2}{c_1} \|e_u\| \leq \left( \frac{c_2}{c_1} \mathbf{K}_b + 1 \right) \|1_E\|. \quad (4.6)$$

From (4.4), (4.5), and (4.6), we obtain

$$\|1_{\varepsilon A}\| \leq (\mathbf{C}_s^\omega)^2 \mathbf{K}_b (\mathbf{K}_b + 1) \left( \frac{c_2}{c_1} \mathbf{K}_b + 1 \right) \|1_{\delta B}\|.$$

Hence,  $\mathcal{B}$  is  $\omega$ -superdemocratic.  $\square$

The proof of the next theorem is similar to that of [11, Theorem 3.2] with obvious modifications, so we move the proof to the Appendix.

**Theorem 4.6.** *If a basis  $\mathcal{B}$  is quasi-greedy and  $\omega$ -disjoint superdemocratic, then it is  $\omega$ -semi-greedy.*

*Proof of Theorem 4.2.* The theorem follows from Theorems 2.8, 4.5, and 4.6.  $\square$

## 5. $\omega$ -PARTIALLY GREEDY BASES

Partially greedy bases were first introduced and characterized in [9] to compare the performance of the TGA to that of the partial sum operators  $(S_m)_{m=1}^\infty$ . In this section, we characterize  $\omega$ -partially greedy bases and prove the existence of  $\omega$ -partially greedy bases that are not  $\varsigma$ -partially greedy for any sequence weight  $\varepsilon$ . For each  $m \geq 0$ , let  $L_m := \{1, 2, \dots, m\}$ . The following is a generalization of [4, Definition 6.1] and [6, Definition 3.4], which defines  $(\varsigma)$ -partial greediness.

**Definition 5.1.** A basis is said to be  $\omega$ -partially greedy if there exists  $C \geq 1$  such that for all  $x \in \mathbb{X}$ ,  $m \in \mathbb{N}$ , and  $\Lambda_m(x)$ , we have

$$\|x - G_m(x)\| \leq C \bar{\sigma}_{\Lambda_m(x)}^\omega(x),$$

where

$$\bar{\sigma}_A^\omega(x) := \inf \{ \|x - S_k(x)\| : w(L_k \setminus A) \leq w(A \setminus L_k) \}.$$

The least such  $C$  is denoted by  $\mathbf{C}_p^\omega$ .

We shall characterize  $\omega$ -partial greediness, generalizing existing characterizations of  $\varsigma$ -partially greedy bases. In [3], the authors introduce partial symmetry for largest coefficients (PSLC).

**Definition 5.2.** A basis is  $C$ - $\omega$ -PSLC if

$$\|x + 1_{\varepsilon A}\| \leq C\|x + 1_{\delta B}\|,$$

for all  $x \in \mathbb{X}$  with  $\|x\|_\infty \leq 1$ , for all finite sets  $A, B \subset \mathbb{N}$  with  $w(A) \leq w(B)$  and  $A < \text{supp}(x) \sqcup B$ , and for all signs  $(\varepsilon), (\delta)$ . The least constant  $C$  is denoted by  $C_{pl}^\omega$ .

**Theorem 5.3.** Let  $\mathcal{B}$  be a basis and  $\omega$  be a weight on subsets of  $\mathbb{N}$ .

- (1) If  $\mathcal{B}$  is  $C_p^\omega$ - $\omega$ -partially greedy, then  $\mathcal{B}$  is  $C_p^\omega$ -suppression quasi-greedy and is  $C_p^\omega$ - $\omega$ -PSLC.
- (2) If  $\mathcal{B}$  is  $C_\ell$ -suppression quasi-greedy and is  $C_{pl}^\omega$ - $\omega$ -PSLC, then  $\mathcal{B}$  is  $C_\ell C_{pl}^\omega$ - $\omega$ -partially greedy.

*Proof.* Similar to the proof of Theorem 2.9.  $\square$

**Definition 5.4.** A basis  $\mathcal{B}$  is  $\omega$ -conservative ( $\omega$ -superconservative, respectively) if there exists  $C \geq 1$  such that

$$\|1_A\| \leq C\|1_B\|, (\|1_{\varepsilon A}\| \leq C\|1_{\delta B}\|, \text{ respectively}),$$

for all  $A, B \in \mathbb{N}^{<\infty}$  with  $A < B$ ,  $w(A) \leq w(B)$ , and signs  $(\varepsilon), (\delta)$ .

We have the following equivalences, whose proof is similar to that of Theorem 2.8 and thus, is left for interested readers.

**Theorem 5.5.** Let  $\mathcal{B}$  be a basis and  $\omega$  be a weight on subsets of  $\mathbb{N}$ . The following are equivalent:

- (1)  $\mathcal{B}$  is  $\omega$ -partially greedy,
- (2)  $\mathcal{B}$  is quasi-greedy and is  $\omega$ -PSLC,
- (3)  $\mathcal{B}$  is quasi-greedy and  $\omega$ -superconservative.
- (4)  $\mathcal{B}$  is quasi-greedy and  $\omega$ -conservative.

The following is an analog of Theorem 3.1.

**Theorem 5.6.** If a basis  $\mathcal{B} = (e_n)_{n=1}^\infty$  is  $\varsigma$ -partially greedy for some weight sequence  $\varsigma = (s(n))_{n=1}^\infty$  with  $\inf s(n) > 0$ , then  $\mathcal{B}$  is partially greedy.

*Proof.* We assume that  $\mathcal{B}$  is  $\varsigma$ -partially greedy for some weight sequence  $\varsigma = (s(n))_{n=1}^\infty$ . By [4, Theorem 6.4],  $\mathcal{B}$  is quasi-greedy and is  $\varsigma$ -conservative.

If  $0 < \inf s(n) \leq \sup s(n) < \infty$ , then [12, Proposition 4.5] implies that  $\mathcal{B}$  is conservative. According to [9, Theorem 3.4],  $\mathcal{B}$  is partially greedy.

If  $\sup s(n) = \infty$ , then [12, Proposition 4.1] states that  $\mathcal{B}$  is equivalent to the canonical basis of  $c_0$  and thus, is greedy.  $\square$

**Theorem 5.7.** There exists a Schauder basis that is  $\omega$ -partially greedy for some structured weight  $\omega$  on sets but is not  $\varsigma$ -partially greedy for any weight sequence  $\varsigma = (s(n))_{n=1}^\infty$  with  $\inf s(n) > 0$ .

*Proof.* The basis  $\mathcal{B}$  in Section 3 is not conservative. To see this, simply pick  $A = \{2, 2^2, \dots, 2^N\}$  and  $B = \{3^{N+1}, \dots, 3^{2N}\}$ . We have  $\|1_A\|/\|1_B\| \sim \sqrt{N}/\ln(N) \rightarrow \infty$  as  $N \rightarrow \infty$ . Hence,  $\mathcal{B}$  is not partially greedy due to [9, Theorem 3.4]. Applying Theorem 5.6, we obtain the desired conclusion.  $\square$

**Remark 5.8.** Theorem 5.7 is sharp in the sense that we cannot drop the requirement  $\inf s(n) > 0$ . Indeed, Khurana [12] characterized  $\varsigma$ -partially greedy bases by quasi-greediness and the so-called  $\varsigma$ -left-Property (A). By [12, Remark 3.3], any basis trivially satisfies  $\varsigma$ -left-Property (A) with  $\varsigma = (s(n))_{n=1}^\infty = (2^{-n})_{n=1}^\infty$ . Hence, if we have an  $\omega$ -partially greedy, it is quasi-greedy by Theorem 5.3 and has  $\varsigma$ -left-Property (A) for  $s(n) = 2^{-n}$ . Therefore, the basis is automatically  $\varsigma$ -partially greedy.

## 6. QUESTIONS AND DISCUSSION

We list several open questions for future research.

- Q1 We show that for a structured weight  $\omega$ , a basis is  $\omega$ -almost greedy if and only if it is  $\omega$ -semi-greedy. Does the result hold for a larger class of weights?
- Q2 For weights in Definition 1.2, is an  $\omega$ -disjoint superdemocratic basis also  $\omega$ -superdemocratic? If not, what minimal condition(s) to put on  $\omega$  so that the two properties are equivalent.

For the second question, we know that for a structured weight, an  $\omega$ -disjoint superdemocratic is  $\omega$ -superdemocratic.

**Proposition 6.1.** *For a structured weight  $\omega$ , a basis  $\mathcal{B}$  is  $\omega$ -superdemocratic if and only if  $\mathcal{B}$  is  $\omega$ -disjoint superdemocratic.*

*Proof.* Assume that  $\mathcal{B}$  is  $\omega$ -disjoint superdemocratic. Let  $A, B \in \mathbb{N}^{<\infty}$  with  $w(A) \leq w(B)$ . Pick signs  $(\varepsilon), (\delta)$ . If  $\sum_{n=1}^\infty w_n < \infty$  or  $\sup_n w_n = \infty$ , then by Proposition 4.3 and Remark 4.4,  $\mathcal{B}$  is equivalent to the canonical basis of  $c_0$ , and the desired conclusion follows trivially. For the rest of the proof, we assume that  $\sum_{n=1}^\infty w_n = \infty$  and  $\sup_n w_n < \infty$ .

Case 1:  $w(A) \leq \limsup_{n \rightarrow \infty} w_n$ . We proceed as in the proof of Theorem 4.5 to show

$$\|1_{\varepsilon A}\| \lesssim \|1_{\delta B}\|.$$

Case 2:  $w(A) > \limsup_{n \rightarrow \infty} w_n$ . Choose  $E$  and  $F = E \cup \{N\}$  such that  $A \cup B \subset E \subset \{N\}$  and  $w(E) \leq w(A) < w(F)$ . By  $\mathbf{C}_{sd, \sqcup}^\omega$ - $\omega$ -disjoint superdemocracy and (4.6), we have

$$\|1_{\varepsilon A}\| \leq \mathbf{C}_{sd, \sqcup}^\omega \|1_F\| \leq \mathbf{C}_{sd, \sqcup}^\omega \left( \frac{c_2}{c_1} \mathbf{K}_b + 1 \right) \|1_E\| \leq (\mathbf{C}_{sd, \sqcup}^\omega)^2 \left( \frac{c_2}{c_1} \mathbf{K}_b + 1 \right) \|1_{\delta B}\|.$$

Therefore,  $\mathcal{B}$  is  $\omega$ -superdemocratic.  $\square$

## 7. APPENDIX

**7.1. Proof of Theorem 2.9.** The key input is the uniform boundedness of the truncation function. For each  $\alpha > 0$ , we define the truncation function  $T_\alpha$  as follows: for  $b \in \mathbb{K}$ ,

$$T_\alpha(b) = \begin{cases} \operatorname{sgn}(b)\alpha, & \text{if } |b| > \alpha, \\ b, & \text{if } |b| \leq \alpha. \end{cases}$$

We define the truncation operator  $T_\alpha : \mathbb{X} \rightarrow \mathbb{X}$  as

$$T_\alpha(x) = \sum_{n=1}^{\infty} T_\alpha(e_n^*(x))e_n = \alpha 1_{\Gamma_\alpha(x)} + P_{\Gamma_\alpha^c(x)}(x),$$

where  $\Gamma_\alpha(x) = \{n : |e_n^*(x)| > \alpha\}$  and  $\varepsilon_n = \text{sgn}(e_n^*(x))$  for all  $n \in \Gamma_\alpha(x)$ . The operator  $T_\alpha$  is well-defined as  $|\Gamma_\alpha(x)| < \infty$  for all  $\alpha > 0$  and  $x \in \mathbb{X}$ .

**Theorem 7.1.** [7, Lemma 2.5] *Let  $\mathcal{B}$  be  $\mathbf{C}_\ell$ -suppression quasi-greedy. Then for any  $\alpha > 0$ ,  $\|T_\alpha\| \leq \mathbf{C}_\ell$ .*

*Proof of Theorem 2.9.* Assume that  $\mathcal{B}$  is  $\mathbf{C}_{al}^\omega$ - $\omega$ -almost greedy. Let  $x \in \mathbb{X}$  and  $A$  be a greedy set of  $x$ . We have

$$\|x - P_A(x)\| \leq \mathbf{C}_{al}^\omega \tilde{\sigma}_A^\omega(x) \leq \mathbf{C}_{al}^\omega \|x - P_\emptyset(x)\| = \mathbf{C}_{al}^\omega \|x\|.$$

The proof of  $\mathbf{C}_{al}^\omega$ - $\omega$ -Property (A) uses the exact argument as in the proof of Theorem 2.3, so we skip it.

Now assume that  $\mathcal{B}$  is  $\mathbf{C}_\ell$ -suppression quasi-greedy and satisfies  $\mathbf{C}_b^\omega$ - $\omega$ -Property (A). Let  $x \in \mathbb{X}$  and  $A$  be a greedy set of  $x$ . Let  $B \in \mathbb{N}^{<\infty}$  such that  $w(B \setminus A) \leq w(A \setminus B)$ . By Lemma 2.4 and Theorem 7.1, we have

$$\begin{aligned} \|x - P_A(x)\| &\leq \mathbf{C}_b^\omega \left\| x - P_A(x) - P_{B \setminus A}(x) + \alpha \sum_{n \in A \setminus B} \text{sgn}(e_n^*(x))e_n \right\| \\ &\leq \mathbf{C}_b^\omega \left\| P_{(A \cup B)^c}(x) + \alpha \sum_{n \in A \setminus B} \text{sgn}(e_n^*(x))e_n \right\| \\ &= \mathbf{C}_b^\omega \|T_\alpha(P_{(A \cup B)^c}(x) + P_{A \setminus B}(x))\| \\ &\leq \mathbf{C}_\ell \mathbf{C}_b^\omega \|x - P_B(x)\|. \end{aligned}$$

This completes our proof that  $\mathcal{B}$  is  $\mathbf{C}_\ell \mathbf{C}_b^\omega$ - $\omega$ -almost greedy.

## 7.2. Proof of Theorem 4.6.

**Lemma 7.2.** [7, Lemma 2.3]. *Let  $\mathcal{B}$  be a  $\mathbf{C}_\ell$ -suppression quasi-greedy basis and  $x \in \mathbb{X}$ . If  $A$  is a greedy set of  $x$ , then*

$$\min_{n \in A} |e_n^*(x)| \left\| \sum_{n \in A} \varepsilon_n e_n \right\| \leq 2\mathbf{C}_\ell \|x\|, \quad (7.1)$$

where  $\varepsilon_n = \text{sgn}(e_n^*(x))$ .

*Proof of Theorem 4.6.* Let us assume that  $\mathcal{B}$  is  $\mathbf{C}_\ell$ -suppression quasi-greedy and  $\mathbf{C}_{sd, \sqcup}^\omega$ - $\omega$ -disjoint superdemocratic. Let  $x \in \mathbb{X}$  with  $|\text{supp}(x)| < \infty$  and  $\Lambda_m(x)$  be a greedy set. Fix  $\varepsilon > 0$ . Let  $y = \sum_{n \in A} a_n e_n$ , where  $A \in \mathbb{N}^{<\infty}$ ,  $w(A \setminus \Lambda_m(x)) \leq w(\Lambda_m(x) \setminus A)$  and  $\|x - y\| < \sigma_{\Lambda_m(x)}^\omega(x) + \varepsilon$ . Write  $x - y = \sum_{n=1}^{\infty} b_n e_n$ , where  $b_n = e_n^*(x) - a_n$  if  $n \in A$  and  $b_n = e_n^*(x)$  if  $n \notin A$ . We shall find a vector  $w$  with  $\text{supp}(w) \subset \Lambda_m(x)$  such that

$$\|x - w\| \leq \mathbf{C}_\ell (1 + 4\mathbf{C}_{sd, \sqcup}^\omega \mathbf{C}_\ell) (\sigma_{\Lambda_m(x)}^\omega(x) + \varepsilon). \quad (7.2)$$



Set  $\alpha := \max_{n \notin \Lambda_m(x)} |e_n^*(x)|$ . If  $\alpha = 0$ , then choose  $w = x$  and we are done. Assume that  $\alpha > 0$ . Consider the following vector:

$$z := \sum_{n \in \Lambda_m(x)} T_\alpha(b_n)e_n + P_{\Lambda_m(x)^c}(x) \quad (7.3)$$

$$\begin{aligned} &= \sum_{n \in \Lambda_m(x)} T_\alpha(b_n)e_n + \sum_{n \notin A \cup \Lambda_m(x)} T_\alpha(b_n)e_n + \sum_{n \in A \setminus \Lambda_m(x)} e_n^*(x)e_n \\ &= \sum_{n \notin A \setminus \Lambda_m(x)} T_\alpha(b_n)e_n + \sum_{n \in A \setminus \Lambda_m(x)} e_n^*(x)e_n \\ &= \sum_{n=1}^{\infty} T_\alpha(b_n)e_n + \sum_{n \in A \setminus \Lambda_m(x)} (e_n^*(x) - T_\alpha(b_n))e_n. \end{aligned} \quad (7.4)$$

We claim that  $x - z$  is a choice for  $w$ . Indeed, using (7.3), we know that  $\text{supp}(w) = \text{supp}(x - z) \subset \Lambda_m(x)$ . By Theorem 7.1, we have

$$\left\| \sum_{n=1}^{\infty} T_\alpha(b_n)e_n \right\| \leq \mathbf{C}_\ell \|x - y\|. \quad (7.5)$$

Note that  $|e_n^*(x) - T_\alpha(b_n)| \leq 2\alpha$  for all  $n \in A \setminus \Lambda_m(x)$ . Let  $\eta = (\text{sgn}(e_n^*(x - y)))_{n=1}^\infty$ . We have

$$\begin{aligned} \left\| \sum_{n \in A \setminus \Lambda_m(x)} (e_n^*(x) - T_\alpha(b_n))e_n \right\| &\leq 2\alpha \sup_{(\delta)} \|1_{\delta A \setminus \Lambda_m(x)}\| \\ &\leq 2\mathbf{C}_{sd, \sqcup}^\omega \min_{n \in \Lambda_m(x) \setminus A} |e_n^*(x - y)| \|1_{\eta \Lambda_m(x) \setminus A}\|. \end{aligned}$$

Let  $B := \{n : |e_n^*(x - y)| \geq \min_{n \in \Lambda_m(x) \setminus A} |e_n^*(x - y)|\}$ . Then  $B$  is a greedy set of  $x - y$  and  $\Lambda_m(x) \setminus A \subset B$ . Therefore, we obtain

$$\|1_{\eta \Lambda_m(x) \setminus A}\| \leq \mathbf{C}_\ell \|1_{\eta B}\|$$

and so, by (7.1),

$$\begin{aligned} \left\| \sum_{n \in A \setminus \Lambda_m(x)} (e_n^*(x) - T_\alpha(b_n))e_n \right\| &\leq 2\mathbf{C}_{sd, \sqcup}^\omega \mathbf{C}_\ell \min_{n \in \Lambda_m(x) \setminus A} |e_n^*(x - y)| \|1_{\eta B}\| \\ &\leq 4\mathbf{C}_{sd, \sqcup}^\omega \mathbf{C}_\ell^2 \|x - y\|. \end{aligned} \quad (7.6)$$

Using (7.4), (7.5), and (7.6), we obtain (7.2). Therefore,

$$\|x - CG_m(x)\| \leq \mathbf{C}_\ell (1 + 4\mathbf{C}_{sd, \sqcup}^\omega \mathbf{C}_\ell) (\sigma_{\Lambda_m(x)}^\omega(x) + \varepsilon).$$

Letting  $\varepsilon \rightarrow 0$  completes the proof.  $\square$

$\square$

## REFERENCES

- [1] F. Albiac and P. Wojtaszczyk, Characterization of 1-greedy bases, *J. Approx. Theory* **138** (2006), 65–86.
- [2] M. Berasategui and S. Lassalle, Weak weight-semi-greedy Markushevich bases, preprint. Available at <https://arxiv.org/pdf/2111.00693>.
- [3] M. Berasategui, P. M. Berná, and S. Lassalle, Strong partially greedy bases and Lebesgue-type inequalities, *Constr. Approx.* **54** (2021), 507–528.
- [4] P. M. Berná, S. J. Dilworth, D. Kutzarova, T. Oikhberg, and B. Wallis, The weighted property (A) and the greedy algorithm, *J. Approx. Theory* **248** (2019), 1–18.
- [5] P. M. Berná, Characterization of weight-semi-greedy bases, *J. Fourier Anal. Appl.* **26** (2020), 1–21.
- [6] P. M. Berná, A note on partially-greedy bases in quasi-Banach spaces, *Studia Math.* **259** (2021), 225–239.
- [7] P. M. Berná, O. Blasco, and G. Garrigós, Lebesgue inequalities for greedy algorithm in general bases, *Rev. Mat. Complut.* **30** (2017), 369–392.
- [8] S. J. Dilworth, D. Kutzarova, E. Odell, Th. Schlumprecht, and A. Zsák, Renorming spaces with greedy bases, *J. Approx. Theory* **188** (2014), 39–56.
- [9] S. J. Dilworth, N. J. Kalton, D. Kutzarova, and V. N. Temlyakov, The thresholding greedy algorithm, greedy bases, and duality, *Constr. Approx.* **19** (2003), 575–597.
- [10] S. J. Dilworth, D. Kutzarova, V. N. Temlyakov, and B. Wallis, Weight-almost greedy bases, *Proc. Steklov. Inst. Math.* **303** (2018), 109–128.
- [11] S. J. Dilworth, N. J. Kalton, and D. Kutzarova, On the existence of almost greedy bases in Banach spaces, *Studia Math.* **159** (2003), 67–101.
- [12] D. Khurana, Weight-partially greedy bases and weight-property (A), *Ann. Funct. Anal.* **11** (2020), 101–117.
- [13] S. V. Konyagin and V. N. Temlyakov, A remark on greedy approximation in Banach spaces, *East J. Approx.* **5** (1999), 365–379.

Email address: [hungchu2@illinois.edu](mailto:hungchu2@illinois.edu)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN, URBANA, IL 61820, USA