

PATTERN RECONSTRUCTION WITH RESTRICTED BOLTZMANN MACHINES

GIUSEPPE GENOVESE

ABSTRACT. We show that the ability of a restricted Boltzmann machine to reconstruct a random pattern depends on the tail of the hidden prior distribution: hidden priors with strictly sub-Gaussian tails give only a logarithmic loss in pattern retrieval, while an efficient retrieval is much harder with hidden units with strictly super-Gaussian tails; reconstruction with sub-Gaussian hidden prior is regulated by the number of hidden units (as in the Hopfield model). This is proved by localisation estimates for the local minima of the energy function.

1. INTRODUCTION

This paper investigates restricted Boltzmann machines (RBMs) from the point of view of Hebbian learning. RBMs are widely studied generative models of machine learning, introduced long ago in [18] and their relation with models of associative memory, such as the ones proposed by Little [14] or Hopfield [12], was noted at the very early stage of the theory, see [11]. Simple in formulation yet very rich in structure, RBMs represent a nice prototype of energy model, on which theoretical progresses could be made and new ideas could be tested.

Here we address the following natural question from the viewpoint of associative networks: is it possible to reconstruct a pattern just looking at the typical configurations of the visible layer of a RBM? Since the hidden prior determines an energy function on the visible layer, see [22], [4], [8], we can analyse the corresponding associative memory, for which the natural term of comparison is the Hopfield model, obtained by a Bernoulli-Gaussian RBM via marginalisation over the hidden layer. The question is relevant for the understanding of the RBM probability distribution and in particular of how the hidden layer affects the configurations of the visible units. Nowadays the main focus on RBMs is on learning the probability distribution underlying a given dataset. In a sense, this amounts to fitting the true law of the data by tuning the weights in the density of the model, in such a way that the datapoints fall close to deep local minima of the energy function. Therefore the analysis of learning in RBMs consists of two tasks: understanding the complex landscape of the energy in the vicinity of the datapoints at given weights and devising good optimisation algorithms to fit the data. This steps, at least from a mathematical angle, are very challenging and the problem is widely open. In these respects pattern retrieval represents a simplified setting to study at first instance, as the roles of the datapoints and of the weights is undertaken by the same objects, the patterns. So there is no optimisation, but one only has to look at the energy landscape in the vicinity of the patterns.

In fact we study the retrieval of i.i.d. binary patterns with RBMs with binary visible units, as the distribution of the hidden layer varies. We do it looking at the local minima of the energy function, in what is called in statistical physics a zero temperature limit, as an analogous rigorous analysis of the RBM density (see (1.2) below) is still prohibitive. Anyway for what concerns pattern reconstruction the zero-temperature regime is particularly relevant, since it is exactly where retrieval is maximised (see [2]). We prove that the tail of the hidden prior distribution determines the retrieval capability of RBMs. More precisely, for priors with tails ranging from exponential to Gaussian we prove that deep local minima are well localised about the patterns.

Date: May 17, 2022.

On the other hand, if the tails of the hidden prior decay faster than Gaussian, we show that the patterns cannot be retrieved well in any case. RBMs whose hidden priors have Gaussian tails (a class including the Hopfield model) represent special threshold cases which we treat separately in either the positive (Theorem 1.1) and the negative (Theorem 1.2) result below.

1.1. Setting. We consider RBMs with i.i.d. Bernoulli ± 1 visible units $\sigma_1, \dots, \sigma_{N_1}$ and symmetric i.i.d. hidden units z_1, \dots, z_{N_2} and we allow a certain freedom in the choice of the hidden prior π , for which we only require that

$$\pi(|z| \geq t) \simeq e^{-|t|^q} \quad \text{for some } q > 1. \quad (1.1)$$

Let $\xi^{(1)}, \dots, \xi^{(N_2)}$ denote independent patterns with N_1 i.i.d. ± 1 components. The (unnormalised) density of the RBM probability distribution reads

$$p(\sigma, z; \xi) := \exp \left(\frac{\beta}{N_1^{q_-}} \sum_{\mu \in [N_2]} (\sigma, \xi^{(\mu)}) z_\mu \right), \quad (1.2)$$

where $\beta > 0$ is a parameter usually called *inverse temperature* and $q_- := \min(q, 2)$. We shall consider the ratio between the number of visible and hidden units as follows

$$\alpha := \frac{N_2}{N_1^{\frac{p_+}{2}}}, \quad (1.3)$$

where $\frac{1}{p_+} + \frac{1}{q_-} = 1$. The normalisation factors in (1.2) and (1.3) ensure that either the energy of the single pattern and the global maximum of (1.2) as $\beta \rightarrow \infty$ scale linearly with α (with constants depending on the prior distribution); more details on that in Section 2.

1.2. Main results. The main object of interest in this paper will be the following two sets. These definitions are slightly formal and a more precise characterisation of it will be given in the next section.

$$\begin{aligned} \text{LM}_{N_1} &:= \{ \text{local maxima of (1.2) as } \beta \rightarrow \infty \}, \\ \text{dLM}_{N_1}^{(\mu)} &:= \{ \text{local maxima of (1.2) as } \beta \rightarrow \infty \text{ higher than the value at } \xi^{(\mu)} \}. \end{aligned} \quad (1.4)$$

Below $\text{Hamming}(a, b)$ denotes the Hamming distance between a, b (*i.e.* the number of different entries) and $\widehat{B}_{\mu, R}^{N_1}$ is the ball in this metric centred at the μ -th pattern with radius R . Throughout the paper we will repeatedly use that two patterns are typically separated by $N_1/2$ flips, so that $\widehat{B}_{\mu, \lfloor N_1/2 \rfloor}^{N_1}$ and $\widehat{B}_{\mu', \lfloor N_1/2 \rfloor}^{N_1}$ typically do not overlap.

Our first result states that the error of reconstructing a given pattern is very small in terms of the size of the network if the decay of the hidden prior (1.1) is slower than Gaussian, while a finite fraction of bits cannot be retrieved for $q = 2$.

Theorem 1.1. *Let $q \in (1, 2)$. There exists $r_0 \in (0, \frac{1}{2}]$ such that with high probability*

$$\max_{\mu \in [N_2]} \max_{\sigma \in \text{dLM}_{N_1}^{(\mu)} \cap \widehat{B}_{\mu, \lfloor r_0 N_1 \rfloor}^{N_1}} \text{Hamming}(\sigma, \xi^{(\mu)}) \lesssim (\log N_1)^{\frac{q-1}{2-q}} \quad (1.6)$$

Let $q = 2$. There exists $r_0 \in (0, \frac{1}{2}]$ such that for all $r \in (0, r_0)$ if $\alpha \lesssim \sqrt{r}$ then with high probability

$$\max_{\mu \in [N_2]} \max_{\sigma \in \text{dLM}_{N_1}^{(\mu)} \cap \widehat{B}_{\mu, \lfloor r_0 N_1 \rfloor}^{N_1}} \text{Hamming}(\sigma, \xi^{(\mu)}) \lesssim \lfloor r N_1 \rfloor. \quad (1.7)$$

Theorem 1.1 characterises the models with $q < 2$ as excellent in pattern reconstruction: there are high maxima located few flips away from the patterns (in fact polylog flips, see (1.6)) and no higher maxima appear in an extended region. Specifying the radius r_0 and the basin of attraction of the patterns would require a careful optimisation of the constants in the proof that we do not attempt.

Local maxima are not directly related to the typical configurations of (1.2). However it is well known that any algorithmic search of typical configurations will finish to find a hopefully representative local maximum. This can be done by the usual FLIP algorithm, that is greedy flipping of one units at time decreasing the energy until no more decreasing is possible. Therefore $\text{dLM}_{N_1}^{(\mu)}$ has a direct interpretation in terms of retrieval. Take for instance for $q < 2$. By Theorem 1.1 any FLIP search initialised for instance at $\xi^{(\mu)}$ will end up in a point of $\text{dLM}_{N_1}^{(\mu)}$ falling only $(\log N_1)^{\frac{q-1}{2-q}}$ flips away from the pattern, which means that only few bits are mis-retrieved.

Somewhat in the opposite direction, the next result shows that for $q > 2$ in (1.1), the local maxima of the density as $\beta \rightarrow \infty$ are quite far from the patterns.

Theorem 1.2. *Let $q \geq 2$, $r \in [0, \frac{1}{2}]$ and let $\alpha_q(r) = \sqrt{S(r)}$ for $q \neq 2$ and $\alpha_2(r) := S(r)/(1-2r)^2$. For every $r \in (0, \frac{1}{2})$ and for all $\alpha \gtrsim \alpha_q(r)$ we have with high probability that*

$$\text{Hamming}(\text{LM}_{N_1}, \xi^{(\mu)}) \gtrsim \lfloor rN_1 \rfloor. \quad (1.8)$$

According to Theorem 1.2, if $q > 2$ one could still hope for retrieval with a very small amount of hidden variables, *i.e.* for α small enough (indeed for $\alpha = 0$ reconstruction is possible, see [4]), but for α larger than a given constant no recovery is allowed. For $q = 2$ the situation improves a bit in the sense that pattern reconstruction becomes less and less efficient as α grows.

The paper [4] showed that RBMs with hidden prior interpolating between a Gaussian and a bimodal symmetric distribution exhibit retrieval at finite $\alpha > 0$, which disappears in the degenerate case when the Gaussian part is switched off. It is also argued that such a lack of retrieval should persist at least for any compactly supported hidden prior. This is demonstrated using non-rigorous replica computations and numerics. We give here the first mathematical confirm of these findings, as Theorem 1.1 implies pattern retrieval if in the above interpolating prior the Gaussian part is present, whatever small, and Theorem 1.2 is a strong indication for lack of retrieval for hidden prior with a Bernoulli ± 1 distribution (for which we should read $q = \infty$).

1.3. Related literature. The results here presented mark a neat difference in the retrieval capabilities of RBMs with hidden priors (1.1) with $q < 2$ (very good capabilities) and $q > 2$ (not so good) with a transition at the sub-Gaussian case $q = 2$. As already remarked, a notable instance of the case $q = 2$ is the Hopfield model, for which a similar analysis at zero temperature was done in [16] (analog of Theorem 1.1), [15] (analog of Theorem 1.2) and [20] in the attempt of proving the picture of [2]. When comparing these papers to ours, we underline that due to the generality of the models we treat, we do not seek to characterise our estimate with the best possible constants (we leave all the numerical constants in fact undetermined), which was instead a very relevant component of all these previous papers.

We exploit and make mathematically precise the heuristics of [4]. Namely we use that the tail of the hidden prior determines the behaviour for large argument of the energy function of the associative network (around zero it is always quadratic). It is exactly this asymptotic that governs retrieval: the more convex the better. More details on that in Section 2. Mathematically speaking the introduction of the hidden layer is a way to linearise the energy function (over the visible units) and different prior distributions for the hidden layer correspond to different associative networks. Similar ideas have been used by [22], [4], [21], [8], [17] to study the performance of the RBMs with varying hidden unit statistics.

The FLIP algorithm is a very natural choice to explore the energy landscape of RBMs and indeed it is essentially the original network dynamics proposed by [12]. This gives a nice connection with the local max-cut problem as analysed for instance by [10] and [3], even though here we exploit crucially the presence of the patterns, which constitute a special class of local minima. This is even more clear by comparing with the analysis for the Sherrington-Kirkpatrick model of [1].

Many other dynamics have been proposed alternative to the FLIP algorithm mainly for the Hopfield model and we will not give here an account on that (see the recent work [6] and the references therein). We just mention that the dynamics analysed by [5], which is a zero-temperature version of the alternate Gibbs sampling typically used to train RBMs, is in spirit very close to our zero-temperature reduction.

1.4. Notations. Throughout the paper $p, q \geq 1$ will always be Hölder conjugate, that is $\frac{1}{p} + \frac{1}{q} = 1$ and similarly for q_-, p_+ , with $q_- := \min(2, q)$, $p_+ := \max(2, p)$. C, c everywhere denote positive absolute constants which may change from formula to formula. We write $X \lesssim Y$ if $X \leq CY$ and $X \simeq Y$ if $Y \lesssim X \lesssim Y$. Sometimes we write \simeq_a or \lesssim_a to stress the dependence of the constants C above on a parameter a . We indicate by (\cdot, \cdot) the inner product in \mathbb{R}^{N_1} or \mathbb{R}^{N_2} and the meaning will be always clear from the context and by $\|\cdot\|_p$ the ℓ_p -norms. $\mathbf{1}$ may represent the vector in \mathbb{R}^{N_1} or in \mathbb{R}^{N_2} with all entries equal to 1. $B_N^{(q)}$ is the ℓ_q centred ball of radius one in \mathbb{R}^N . $\widehat{S}_{\mu, R}^{N_1-1}$, $\widehat{B}_{\mu, R}^{N_1}$ denote respectively the N_1 -dimensional Hamming ball and sphere centred at $\xi^{(\mu)}$ of radius R . If $v \in \mathbb{R}^N$ and $J \subset [N]$ we denote by v_J a vector in $\mathbb{R}^{|J|}$ such that $(v_J)_i = v_{j_i}$ if $J = \{j_1, \dots, j_{|J|}\}$. For any $J \subset [N]$ we also associate a FLIP operator F_J defined by $(F_J v)_i = -v_i$ if $i \in J$ and $(F_J v)_i = v_i$ if $i \notin J$. We will use the following ψ_r -norms:

$$\|Z\|_{\psi_r} := \inf \left\{ c > 0 : E \left[\psi_r \left(\frac{|Z|}{c} \right) \right] < 2 \right\}, \quad r > 0, \quad (1.9)$$

where $\psi_r(x) = e^{x^r}$ for any $x > 0$ for $r \geq 1$, while for $r \in (0, 1)$ there are $c(r), x(r)$ such that for $x \in (0, x(r))$ it is $\psi_r(x) = c(r)x$. We underline that, setting $q_s := \sup\{q' > 1 : \|Z\|_{\psi_{q'}} < \infty\}$ we have $P(|Z| \geq t) \simeq e^{-|t|^{q_s}}$ (we convey that bounded random variables have finite ψ_∞ -norm). We will often deal with the following function:

$$S(r) := -r \log r - (1-r) \log(1-r)$$

and use the standard bound

$$\text{Card } \widehat{S}_{\mu, \lfloor r N_1 \rfloor}^{N_1-1} = \binom{\lfloor r N_1 \rfloor}{N_1} \leq e^{N_1 S(r)}. \quad (1.10)$$

We denote the transpose patterns $\tilde{\xi}^{(i)}$ by $\tilde{\xi}_\mu^{(i)} := \xi_i^{(\mu)}$, $i \in [N_1]$, $\mu \in [N_2]$. Sometimes we write $\widehat{\xi} := \xi/\sqrt{N_1}$. A^c is the complement of the set A . We say that the event A occurs with high probability if $P(A) \geq 1 - N^{-x}$ for all $x > 0$.

Acknowledgements. The author thanks David Belius for many valuable suggestions regarding the presentation of the results.

2. ZERO TEMPERATURE REDUCTION

Here we show how to single out an effective energy function which captures the RBM landscape at zero temperature. We exploit a reduction argument introduced in [4].

Integrating out the hidden layer in (1.2) we get a probability distribution over the visible units. This writes as

$$\int p(\sigma, z; \xi) \pi(dz_1) \dots \pi(dz_{N_2}) = \exp \left(\sum_{\mu \in [N_2]} u \left(\frac{\beta}{N_1^{\frac{1}{q}}} (\xi^\mu, \sigma) \right) \right), \quad (2.1)$$

where

$$u(x) := \log E[e^{xz_1}].$$

We are interested in studying the local minima of this distribution as β is very large. The following simple observation starts our considerations: for any z_1 such that (1.1) holds for some $q > 1$, it is

$$\lim_{x \rightarrow \infty} \frac{u(x)}{|x|^p} \simeq_p \|z_1\|_{\psi_q}^p. \quad (2.2)$$

The proof is immediate: we write

$$\begin{aligned} E[e^{xz_1}] &= \int_0^\infty d\lambda P(z_1 \geq x^{-1} \log \lambda) \simeq \int_0^\infty d\lambda e^{-\frac{1 \log \lambda|^q}{\|z_1\|_{\psi_q}^q |x|^q}} \\ &= \|z_1\|_{\psi_q} |x| \int_0^\infty d\lambda e^{\lambda \|z_1\|_{\psi_q} |x| - |\lambda|^q} \\ &= \|z_1\|_{\psi_q} |x| e^{\frac{\|z_1\|_{\psi_q}^p |x|^p}{p}} \int_0^\infty d\lambda e^{\lambda \|z_1\|_{\psi_q} x - |\lambda|^q - \frac{\|z_1\|_{\psi_q}^p |x|^p}{p}}, \end{aligned}$$

and the last integral is finite by the Young inequality. Taking the log on both sides and passing to the limit we get (2.2).

Therefore by (2.2), as $\beta \rightarrow \infty$ we are led to consider the following energy function (reminiscent of the p -spin Hopfield model of [7], but with a different normalisation)

$$H^{(p)}(\sigma; \xi) := -\frac{1}{N_1^{\kappa(p)}} \sum_{\mu \in [N_2]} |(\xi^{(\mu)}, \sigma)|^p, \quad (2.3)$$

with $\kappa(p) := 1 + p - \frac{p}{p_+}$ (here we include the usual normalisation factor $1/N_1$ of the internal energy directly in the definition of $H^{(p)}$). To fix the ideas, $\kappa(p) = p$ for $p \geq 2$ and $\kappa(p) = 1 + \frac{p}{2}$ for $p \leq 2$.

The focus of this paper will be to study the location of the minima of (2.3) on $\{-1, 1\}^{N_1}$ close to the pattern configurations, in the limit $N_1, N_2 \rightarrow \infty$ while α_p remains constant. Hence the following are more precise definitions of the sets (1.5) and (1.4) previously introduced:

$$\text{LM}_{N_1} := \{\text{local minima of (2.3)}\}, \quad (2.4)$$

$$\text{dLM}_{N_1}^{(\mu)} := \{\text{local minima } \bar{\sigma} \text{ of (2.3), s.t. } H^{(p)}(\xi^{(\mu)}; \xi) - H^{(p)}(\bar{\sigma}; \xi) > 0\}. \quad (2.5)$$

Next we show that the single pattern energy is close to the ground state, so providing a motivation for the normalisation factors in (1.2) and (1.3). First of all we give a lower bound for the ground state energy linear in α . To do so we do not actually need binary patterns.

Proposition 2.1. *Let $\xi^{(1)}, \dots, \xi^{(N_2)}$ be independent vectors in \mathbb{R}^{N_1} with i.i.d. centred sub-Gaussian entries. It holds for $p \geq 1$*

$$\inf_{\sigma \in \{-1, 1\}^{N_1}} H^{(p)}(\sigma; \xi) \gtrsim_p -(1 + \alpha) \quad (2.6)$$

with probability larger than $1 - e^{-c\alpha^{\frac{2}{p}} N_1}$.

To prove Proposition 2.1 we need the following auxiliary lemma.

Lemma 2.1. *Let $\xi^{(1)}, \dots, \xi^{(N_2)}$ be independent vectors in \mathbb{R}^{N_1} with i.i.d. centred sub-Gaussian entries. Let $p \geq 1$. For all $t \gtrsim_p (1 + \alpha)^{\frac{1}{p}}$*

$$P\left(\frac{1}{N_1^{\max(0, \frac{q-2}{2q})}} \sup_{\sigma \in \frac{1}{\sqrt{N_1}} \{-1, 1\}^{N_1}} \sup_{\tau \in B_{N_2}^{(q)}} \frac{(\xi^{(\mu)}, \sigma) \tau_\mu}{\sqrt{N_1}} \geq t\right) \leq 2e^{-ct^2 N_1} \quad (2.7)$$

where $c > 0$ depends only on the distribution of $\xi_1^{(1)}$.

Proof. We introduce the transpose patterns $\tilde{\xi}^{(i)}$ by $\tilde{\xi}_\mu^{(i)} := \xi_i^{(\mu)}$, $i \in [N_1]$, $\mu \in [N_2]$. First of all we note that

$$\frac{1}{N_1^{\max(0, \frac{q-2}{2q})}} \sup_{\sigma \in \frac{1}{\sqrt{N}} \{-1, 1\}^{N_1}} \sum_{i \in [N_1]} \sum_{\mu \in [N_2]} \frac{\xi_i^{(\mu)} \sigma_i \tau_\mu}{\sqrt{N_1}} = \frac{1}{N_1} \sum_{i \in [N_1]} \frac{|\langle \tilde{\xi}^{(i)}, \tau \rangle|}{N_1^{\max(0, \frac{q-2}{2q})}}. \quad (2.8)$$

Moreover, since for all $\tau \in B_{N_2}^{(q)}$

$$\|\tau\|_2 \leq N_2^{\max(0, \frac{q-2}{2q})},$$

we have

$$P \left(\frac{1}{N_1^{\max(0, \frac{q-2}{2q})}} \left| \langle \tilde{\xi}^{(i)}, \tau \rangle \right| \geq t \right) \leq 2e^{-\frac{t^2 N_1^{\max(0, \frac{q-2}{2q})}}{2N_2^{\max(0, \frac{q-2}{2q})} \|\xi_1^{(1)}\|_{\psi_2}^2}}. \quad (2.9)$$

The r.h.s. of (2.8) is the sum of independent sub-Gaussian random variables, thus for all $\tau \in B_{N_2}^{(q)}$

$$P \left(\frac{1}{N_1} \sum_{i \in [N_1]} \frac{|\langle \tilde{\xi}^{(i)}, \tau \rangle|}{N_1^{\max(0, \frac{q-2}{2q})}} \geq t \right) \leq 2e^{-\frac{t^2 N_1^{1+\max(0, \frac{q-2}{2q})}}{2N_2^{\max(0, \frac{q-2}{2q})} \|\xi_1^{(1)}\|_{\psi_2}^2}} \quad (2.10)$$

for some $c > 0$ depending only on the distribution of $\xi_1^{(1)}$.

Next we cover $B_{N_2}^{(q)}$ with a number of ℓ_r -balls in \mathbb{R}^{N_2} with some small radius $\varepsilon > 0$. For $p \geq 2$ we can use Euclidean balls and the Sudakov inequality gives a bound on the minimal number $N(B_q^{N_2}, \varepsilon B_2^{N_2})$ of such balls

$$N(B_q^{N_2}, \varepsilon B_2^{N_2}) \leq e^{cN_2^{\frac{2}{p}}}$$

(here we used that for a gaussian vector g $E[\max_{\tau \in B_q^{N_2}} \langle \tau, g \rangle] = E[\|g\|_p] \simeq N_2^{\frac{1}{p}}$).

For $p \in [1, 2)$ we use ℓ_q -balls and we have

$$N(B_q^{N_2}, \varepsilon B_q^{N_2}) \leq e^{cN_2}$$

(in the two estimates above the constants c depends on ε in a way we do not keep track of).

Assume now $(1 + \alpha)^{\frac{1}{p}} \lesssim_p t$ (this is to take into account also the behaviour for small α). By the union bound for $p \geq 2$ we get

$$P \left(\frac{1}{N_1^{\max(0, \frac{q-2}{2q})}} \sup_{\sigma \in \frac{1}{\sqrt{N}} \{-1, 1\}^{N_1}} \sup_{\tau \in B_{N_2}^{(q)}} (\widehat{\xi} \sigma, \tau) \geq t \right) \leq 2e^{cN_2^{\frac{2}{p}} - cN_1 t^2} \leq 2e^{-ct^2 N_1}. \quad (2.11)$$

Similarly for $p \in [1, 2)$

$$P \left(\frac{1}{N_1^{\frac{q-2}{2q}}} \sup_{\sigma \in \frac{1}{\sqrt{N}} \{-1, 1\}^{N_1}} \sup_{\tau \in B_{N_2}^{(q)}} (\widehat{\xi} \sigma, \tau) \geq t \right) \leq 2e^{cN_2 - c \left(\frac{N_1}{N_2}\right)^{\frac{q-2}{q}} N_1 t^2} \leq 2e^{-ct^2 N_1}. \quad (2.12)$$

□

Proof of Proposition 2.1. The role of hidden variables at zero temperature is played by duality:

$$\sum_{\mu \in [N_2]} |(\xi^{(\mu)}, \sigma)|^p = \left| \sup_{\tau \in B_{N_2}^{(q)}} \sum_{\mu \in [N_2]} (\xi^{(\mu)}, \sigma) \tau_\mu \right|^p, \quad p \geq 1. \quad (2.13)$$

Therefore (here we shorten $\hat{\xi} := \xi/\sqrt{N_1}$)

$$\begin{aligned} \inf_{\sigma \in \{-1,1\}^{N_1}} H^{(p)}(\sigma; \xi) &= - \sup_{\sigma \in \frac{1}{\sqrt{N_1}}\{-1,1\}^{N_1}} \left| \frac{1}{N_1^{\max(0, \frac{q-2}{2q})}} \sup_{\tau \in B_{N_2}^{(q)}} \sum_{\mu \in [N_2]} (\hat{\xi}^{(\mu)}, \sigma) \tau_\mu \right|^p \\ &= - \left| \frac{1}{N_1^{\max(0, \frac{q-2}{2q})}} \sup_{\sigma \in \frac{1}{\sqrt{N_1}}\{-1,1\}^{N_1}} \sup_{\tau \in B_{N_2}^{(q)}} \sum_{\mu \in [N_2]} (\hat{\xi}^{(\mu)}, \sigma) \tau_\mu \right|^p \end{aligned} \quad (2.14)$$

by symmetry. It suffices to focus on the quantity inside the modulus above, which is dealt in Lemma 2.1 of Appendix ?? below. We have for all $t \geq p(1+\alpha)^{\frac{1}{p}}$

$$P \left(\frac{1}{N_1^{\max(0, \frac{q-2}{2q})}} \sup_{\sigma \in \frac{1}{\sqrt{N_1}}\{-1,1\}^{N_1}} \sup_{\tau \in B_{N_2}^{(q)}} (\hat{\xi}\sigma, \tau) \geq t \right) \leq 2e^{-ct^2 N_1}. \quad (2.15)$$

Combining (2.14) with the bound above we obtain the statement. \square

Now we shortly show that the patterns have energy of the same order in α of the global minimum, even though we can already observe a difference between the models with $p \geq 2$ and $p < 2$. We deal with ± 1 binary patterns for simplicity, but a similar argument can be easily repeated for symmetric patterns with minor modifications. We have

$$H^{(p)}(\xi^{(1)}; \xi) = - \frac{\|\xi^{(1)}\|_2^p}{N_1^{1+p-p/p_+}} - \sum_{j=2}^{N_2} \frac{|(\xi^{(1)}, \xi^{(j)})|^p}{N_1^{1+p-p/p_+}}.$$

This quantity concentrates around its average as N_1 grows:

Lemma 2.2. *Take $t > 0$ uniformly in N_1 , small enough. It holds*

$$P \left(\left| H^{(p)}(\xi^{(1)}; \xi) - E[H^{(p)}(\xi^{(1)}; \xi)] \right| \geq t \right) \leq \begin{cases} \exp(-cN_1 t^2) & p \in [1, 2] \\ \exp(-cN_1 t^{\frac{2}{p}}) & p > 2 \end{cases} \quad (2.16)$$

Proof. We write

$$H^{(p)}(\xi^{(1)}; \xi) = - \frac{\|\xi^{(1)}\|_2^p}{N_1^{1+p-p/p_+}} - \sum_{\mu=2}^{N_2} \frac{|(\xi^{(1)}, \xi^{(\mu)})|^p}{N_1^{1+p-p/p_+}}.$$

It suffices to focus on the second summand on the r.h.s. above. We have by independence

$$\begin{aligned} &P \left(\left| \sum_{\mu=2}^{N_2} \frac{|(\xi^{(1)}, \xi^{(\mu)})|^p}{N_1^{\frac{p}{2}}} - E \left[\frac{|(\xi^{(1)}, \xi^{(\mu)})|^p}{N_1^{\frac{p}{2}}} \right] \right| \geq t N_1^{1+p/2-p/p_+} \right) \\ &= P \left(\left| \sum_{\mu=2}^{N_2} \frac{|(1, \xi^{(\mu)})|^p}{N_1^{\frac{p}{2}}} - E \left[\frac{|(1, \xi^{(\mu)})|^p}{N_1^{\frac{p}{2}}} \right] \right| \geq t N_1^{1+p/2-p/p_+} \right), \end{aligned} \quad (2.17)$$

where 1 is the constant vector with all entries equal to 1. The random variables

$$T_p^{(\mu)} := \frac{|(1, \xi^{(\mu)})|^p}{N_1^{\frac{p}{2}}} - E \left[\frac{|(1, \xi^{(\mu)})|^p}{N_1^{\frac{p}{2}}} \right], \quad \mu = 2, \dots, N_2$$

are i.i.d. with $\|T_p^{(\mu)}\|_{\psi_{\frac{2}{p}}} \simeq 1$. Thus by Proposition A.1 with $\ell = 2/p$ we have

$$(2.17) \leq \begin{cases} \exp\left(-cN_1 \min(t^2, t^{\frac{2}{p}})\right) & p \in [1, 2] \\ \exp\left(-cN_1 \min(t^2 N_1^{p-2}, t^{\frac{2}{p}})\right) & p > 2 \end{cases} \quad (2.18)$$

and the proof is complete. \square

Moreover we have

$$|E[H^{(p)}(\xi^{(1)}; \xi)]| \leq \frac{N_1^p + N_1^{\frac{p}{2}} N_2}{N_1^{1+p-p/p_+}} \simeq -\alpha \frac{p}{2} \Gamma\left(\frac{p}{2}\right) - \frac{1}{N_1^{1-p/p_+}}. \quad (2.19)$$

In fact

$$\begin{aligned} E[|(\xi^{(1)}, \xi^{(\mu)})|^p] &= E\left[\int d\lambda P(\xi^{(\mu)} : |(\xi^{(1)}, \xi^{(\mu)})| \geq \lambda^{1/p})\right] \\ &\leq E[\|\xi^{(1)}\|_1^p] \int_0^\infty d\lambda e^{-\lambda^{\frac{2}{p}}} = \frac{p}{2} \Gamma\left(\frac{p}{2}\right) N_1^{\frac{p}{2}}. \end{aligned} \quad (2.20)$$

Therefore

$$H^{(p)}(\xi^{(1)}; \xi) \simeq -\frac{p}{2} \Gamma\left(\frac{p}{2}\right) \alpha - \frac{1}{N_1^{1-p/p_+}}$$

with very high probability. We see that if $p \geq 2$ this value is really of the same order of the ground state, while if $p \in (1, 2)$ for α small and N_1 large the patterns have higher energy.

3. RETRIEVAL FOR $p \geq 2$

In this section we prove Theorem 1.1. We look at all configurations reachable from $\xi^{(\mu)}$ by $\lfloor rN_1 \rfloor$ flips and compare the energy of the pattern with the minimal energy of such configurations. By symmetry of the patters we can reduce to look at $\mu = 1$ and we may and will assume that $\mathbf{1}$, *i.e.* the vector with entries all equal to one, lies in $\widehat{S}_{1, \lfloor rN_1 \rfloor}^{N_1-1}$.

Without further explanation, we introduce some more notations. Given $v \in \widehat{S}_{1, \lfloor rN_1 \rfloor}^{N_1-1}$ we set

$$J_\mu(v) = J := \{i \in [N_1] : v_i \neq \xi^{(\mu)}\}.$$

and

$$X^{(\mu)}(v) := \frac{1}{\sqrt{N_1}}(\xi_J^{(\mu)}, v_J), \quad Y^{(\mu)}(v) := \frac{1}{\sqrt{N_1}}(\xi_{J^c}^{(\mu)}, v_{J^c}). \quad (3.1)$$

and $X^{(\mu)}(\mathbf{1}) =: X^{(\mu)}$, $Y^{(\mu)}(\mathbf{1}) =: Y^{(\mu)}$. We conveniently let

$$F_p(x, y) := |x + y|^p - |x - y|^p, \quad \bar{F}_p(r) := F_p(r, 1 - r) = 1 - (1 - 2r)^p > 0. \quad (3.2)$$

For all $p \geq 1$ it is

$$|F_p(x, y)| \leq 2^p |x|^{\frac{p}{2}} |y|^{\frac{p}{2}}. \quad (3.3)$$

The proof of (3.3) is easy. By convexity

$$||x - y|^p - |x + y|^p| \leq 2^p |x|^p, \quad ||x + y|^p - |x - y|^p| \leq 2^p |y|^p,$$

therefore squaring $|F_p(x, y)|$ (3.3) follows (in fact we have $F_2(x, y) = 4xy$).

We have the following useful representation.

Lemma 3.1. *Let $r \in (0, \frac{1}{2})$, $J \subset [N_1]$ with $|J| = \lfloor rN_1 \rfloor$. It is*

$$H^{(p)}(\xi^{(1)}) - H^{(p)}(F_J \xi^{(1)}) = -\frac{1}{N_1^{\frac{p_+-p}{2}}} \bar{F}_p(r) - \frac{1}{N_1^{\frac{p_+}{2}}} \sum_{\mu=2}^{N_2} F_p(X_J^{(\mu)}(\xi^{(1)}), Y_J^{(\mu)}(\xi^{(1)})). \quad (3.4)$$

Proof. Compute

$$\begin{aligned} H^{(p)}(v) - H^{(p)}(F_J v) &= -\frac{1}{N_1^{\kappa(p)}} \sum_{\mu \in [N_2]} |(\xi_J^{(\mu)}, v_J) + (\xi_{J^c}^{(\mu)}, v_{J^c})|^p - |-(\xi_J^{(\mu)}, v_J) + (\xi_{J^c}^{(\mu)}, v_{J^c})|^p \\ &= -\frac{1}{N_1^{\kappa(p) - \frac{p}{2}}} \sum_{\mu \in [N_2]} F_p(X^{(\mu)}(v), Y^{(\mu)}(v)), \end{aligned}$$

by the definitions (3.1), (3.2). We have $\kappa(p) - \frac{p}{2} = 1 + \frac{p}{2} - \frac{p}{p_+} = \frac{p_+}{2}$. Take now $v = \xi^{(1)}$. An easy computation gives

$$X_J^{(1)}(\xi^{(1)}) = \frac{\|\xi_J^{(1)}\|_2^2}{\sqrt{N_1}} = \frac{|J|}{\sqrt{N_1}} = r\sqrt{N_1}, \quad Y_{J^c}^{(1)}(\xi^{(1)}) = \frac{\|\xi_{J^c}^{(1)}\|_2^2}{\sqrt{N_1}} = \frac{|J^c|}{\sqrt{N_1}} = (1-r)\sqrt{N_1}. \quad (3.5)$$

Thus

$$\frac{1}{N_1^{\frac{p_+}{2}}} F_p(X^{(1)}(\xi^{(1)}), Y^{(1)}(\xi^{(1)})) = \frac{1}{N_1^{\frac{p_+ - p}{2}}} F_p(r, 1-r)$$

and (3.4) follows. \square

The necessary tail estimates in order to prove Theorem 1.1 are given in the next lemma. We shorten $P_\xi(\cdot) = P(\cdot | \xi)$.

Lemma 3.2. *Let $r \in (0, \frac{1}{2})$, $p \geq 2$, $0 < y \leq 1 - (1 - 2r)^p$, $t := 1 - (1 - 2r)^p - y$. Take any $\sigma \in \widehat{S}_{1, [rN_1]}^{N_1 - 1}$. If $\alpha \lesssim N^{\frac{p-2}{2}} r^{\frac{p-1}{2}}$ then*

$$P_{\xi^{(1)}} \left(H^{(p)}(\xi^{(1)}) - H^{(p)}(\sigma) \geq -t \right) \leq \exp \left(-\frac{cN_1 |y|^{\frac{2}{p}}}{\sqrt{r(1-r)}} \right), \quad (3.6)$$

and otherwise

$$P_{\xi^{(1)}} \left(H^{(p)}(\xi^{(1)}) - H^{(p)}(\sigma) \geq -t \right) \leq \exp \left(-\frac{cN_1^{\frac{p}{2}} y^2}{\alpha (r(1-r))^{\frac{p}{2}}} \right). \quad (3.7)$$

Remark 3.1. If $r > 0$ uniformly in N_1 we flip a number of bits proportional to N_1 and the tail exponent is sub-linear for all α if $p > 2$ and for $\alpha \lesssim \sqrt{r}$ if $p = 2$. A sub-linear number of flips corresponds to take $r \simeq N^{-x}$ for some $x > 0$ (modulo log-corrections, see below). In this case we see that if $x \in (\frac{p-2}{p-1}, 1)$ the sub-exponential estimate still prevails, while otherwise we have a Gaussian tail.

The following result is used for proving Lemma 3.2. We omit the proof, which is standard.

Lemma 3.3. *$\{X^{(\mu)}\}_{\mu \in [N_2] \setminus \{1\}}$ and $\{Y^{(\mu)}\}_{\mu \in [N_2] \setminus \{1\}}$ are independent sub-Gaussian random variables, independent one from each other, with*

$$\|X^{(\mu)}\|_{\psi_2} \leq \sqrt{\frac{r}{\log 2}} \quad \|Y^{(\mu)}\|_{\psi_2} \leq \sqrt{\frac{1-r}{\log 2}}. \quad (3.8)$$

Moreover $\{F_p(X^{(\mu)}, Y^{(\mu)})\}_{\mu \in [N_2]}$ are i.i.d. $\psi_{2/p}$ r.vs with

$$\|F_p(X^{(\mu)}, Y^{(\mu)})\|_{\psi_{2/p}}^{\frac{2}{p}} \leq \frac{\sqrt{r(1-r)}}{\log 2}. \quad (3.9)$$

Proof of Lemma 3.2. By (3.4) we have

$$\begin{aligned}
& P_{\xi^{(1)}} \left(H^{(p)}(\xi^{(1)}) - H^{(p)}(\sigma) \geq -t \right) \\
&= P_{\xi^{(1)}} \left(- \sum_{\mu=2}^{N_2} F_p(X_J^{(\mu)}(\xi^{(1)}), Y_J^{(\mu)}(\xi^{(1)})) \geq N_1^{\frac{p}{2}} (\bar{F}_p(r) - t) \right) \\
&= P_{\xi^{(1)}} \left(- \sum_{\mu=2}^{N_2} F_p(X_J^{(\mu)}, Y_J^{(\mu)}) \geq N_1^{\frac{p+}{2}} y \right) \\
&= P \left(- \sum_{\mu=2}^{N_2} F_p(X_J^{(\mu)}, Y_J^{(\mu)}) \geq N_1^{\frac{p+}{2}} y \right), \tag{3.10}
\end{aligned}$$

because of independence of the patterns and $F_p(r, 1-r) \geq 0$.

Note that $y > 0$ is equivalent to $0 \leq t < \bar{F}_p(r)$. By Lemma 3.3 $\{F_p(X_J^{(\mu)}, Y_J^{(\mu)})\}_{\mu \in [N_2]}$ are centred i.i.d. r.v.s which fit the assumptions of Proposition A.1 below (with $\ell = 2/p \in (0, 1]$). Therefore

$$(3.10) \leq \exp \left(-c \min \left(\frac{N_1^p y^2}{N_2(r(1-r))^{\frac{p}{2}}}, \frac{N_1 |y|^{\frac{2}{p}}}{\sqrt{r(1-r)}} \right) \right). \tag{3.11}$$

The value of this minimum depends on α . Bearing in mind that $y < \bar{F}_p(r) \lesssim r^{\frac{p}{2}}$ we have that the sub-linear term prevails if

$$\alpha \lesssim N_1^{\frac{p-2}{2}} r^{\frac{p-1}{2}},$$

otherwise we have a Gaussian tail. \square

Now we are ready for the main proof.

Proof of Theorem 1.1. We shorten

$$\mathsf{D}_{\mu, N_1}(r_0) := \{ \sigma \in \widehat{B}_{\mu, \lfloor r_0 N_1 \rfloor}^{N_1} : H^{(p)}(\xi^{(\mu)}; \xi) \geq H^{(p)}(\sigma; \xi) \} \tag{3.12}$$

and note that since for any $\mu \in [N_2]$

$$\mathsf{dLM}_{N_1}^{(\mu)} \cap \widehat{B}_{\mu, \lfloor r_0 N_1 \rfloor}^{N_1} \subseteq \mathsf{D}_{\mu, N_1}(r_0),$$

it is

$$P \left(\forall \mu \in [N_2] \quad \mathsf{dLM}_{N_1}^{(\mu)} \cap \widehat{B}_{\mu, \lfloor r_0 N_1 \rfloor}^{N_1} \subseteq \widehat{B}_{\mu, R}^{N_1} \right) \geq P \left(\forall \mu \in [N_2] \quad \mathsf{D}_{\mu, N_1}(r_0) \subseteq \widehat{B}_{\mu, R}^{N_1} \right). \tag{3.13}$$

Let us introduce the sets

$$\text{Bar}_{N_1, N_2, p}(n) := \left\{ \min_{\mu \in [N_2]} \min_{\sigma \in \widehat{S}_{\mu, n}^{N_1-1}} H^{(p)}(\sigma) - H^{(p)}(\xi^{(\mu)}) \geq t(n) \right\} \tag{3.14}$$

on which the minimal energy gap of n flips from the patterns is a given $t(n)$. Write now for $r \in (0, \frac{1}{2})$ $n = \lfloor r N_1 \rfloor$ and $t(n) = t(r) = t$ and choose, according to Lemma 3.2, $t \simeq r^{\frac{p}{2}}$. Also we take a $r_0 \in (0, \frac{1}{2}]$ such that $S(r) \leq c\sqrt{r}$, for a constant $c > 0$ to be specified below, for all $r \in [0, r_0]$. Then, bearing in mind (3.13), the crux is

$$\begin{aligned}
& P \left(\forall \mu \in [N_2] \quad \mathsf{D}_{\mu, N_1}(r_0) \subseteq \widehat{B}_{\mu, R}^{N_1} \right) \geq P \left(\bigcap_{n=\lfloor R \rfloor}^{\lfloor r_0 N_1 \rfloor} \text{Bar}_{N_1, N_2, p}(n) \right) \\
& \geq 1 - \sum_{n=\lfloor R \rfloor}^{\lfloor r_0 N_1 \rfloor} P(\text{Bar}_{N_1, N_2, p}^c(n)) \gtrsim 1 - N_1 \min_{\lfloor R \rfloor \leq n \leq \lfloor r_0 N_1 \rfloor} P(\text{Bar}_{N_1, N_2, p}^c(n)). \tag{3.15}
\end{aligned}$$

By the standard estimate (1.10) and the union bound we have

$$\begin{aligned} P(\text{Bar}_{N_1, N_2, p}^c(n)) &= P\left(\min_{\mu \in [N_2]} \min_{\sigma \in \widehat{S}_{\mu, n}^{N_1-1}} H^{(p)}(\sigma) - H^{(p)}(\xi_\mu) \leq t\right) \\ &\leq \exp(\log N_2 + N_1 S(r)) E \left[\sup_{\sigma \in \widehat{S}_{1, [rN_1]}^{N_1-1}} P_{\xi^{(1)}} \left(H^{(p)}(\xi^{(1)}) - H^{(p)}(\sigma) \geq -t \right) \right]. \end{aligned}$$

The probabilities appearing in the last line are evaluated using Lemma 3.2 with $y \simeq t$.

Let us first deal with $p > 2$. We consider different regimes. If $r \gtrsim N^{-x}$ (i.e. $n \gtrsim N_1^{1-x}$) with $x > x_p := \frac{p-2}{p-1}$ then Lemma 3.2 yields for all $\alpha > 0$

$$P(\text{Bar}_{N_1, N_2, p}^c(n)) \leq N_2 \exp(N_1(S(r) - c\sqrt{r})) \lesssim e^{-c\sqrt{r}N_1} \simeq e^{-c\sqrt{nN_1}}. \quad (3.16)$$

The above estimate must hold for all $r \in [0, r_0]$, which determines r_0 . Thus

$$\min_{\lfloor N_1^{1-x_p} \rfloor < n \leq \lfloor r_0 N_1 \rfloor} P(\text{Bar}_{N_1, N_2, p}^c(n)) \leq e^{-cN_1^{1-\frac{x_p}{2}}}. \quad (3.17)$$

For $r < N^{-x}$ with $x \in (x_p, 1)$ Lemma 3.2 gives for all $\alpha > 0$

$$P(\text{Bar}_{N_1, N_2, p}^c(n)) \leq N_2 e^{N_1 S(r) - \frac{N_1^{\frac{p}{2}} r^{\frac{p}{2}}}{\alpha}} \leq N_2 e^{n|\log N_1| - \frac{n^{\frac{p}{2}}}{\alpha}}. \quad (3.18)$$

Thus for all $\varepsilon > 0$ sufficiently small

$$\min_{\lfloor N_1^{2\varepsilon} \rfloor < n < \lfloor N_1^{1-x_p} \rfloor} P(\text{Bar}_{N_1, N_2, p}^c(n)) \leq e^{-cN_1^{\varepsilon p}}. \quad (3.19)$$

Moreover by (3.18) we see that also a poly-log number of flips is allowed:

$$\min_{\lfloor (\log N_1)^{\frac{2}{p-2}} \rfloor < n \leq \lfloor N_1^{2\varepsilon} \rfloor} P(\text{Bar}_{N_1, N_2, p}^c(n)) \leq e^{-c(\log N_1)^{1+\frac{2}{2-p}}}. \quad (3.20)$$

Finally we look at $n \simeq N_1^{1-x_p}$. In this case we have to fix some $\alpha_0 > 0$ and we use for $\alpha \leq \alpha_0$ the bound (3.16) and for $\alpha > \alpha_0$ the bound (3.18). We have

$$\min_{n \simeq \lfloor N_1^{1-x_p} \rfloor} P(\text{Bar}_{N_1, N_2, p}^c(n)) \lesssim e^{-cN_1^{1-\frac{x_p}{2}}}. \quad (3.21)$$

Combining (3.15) with $R = (\log N_1)^{\frac{2}{p-2}}$ and (3.17), (3.19), (3.20), (3.21) we get

$$P\left(\forall \mu \in [N_2] \quad \text{dLM}_{N_1}^{(\mu)} \cap \widehat{B}_{\mu, [r_0 N_1]}^{N_1} \subseteq \widehat{B}_{\mu, [(\log N_1)^{\frac{2}{p-2}}]}^{N_1}\right) \gtrsim 1 - N_1 e^{-c(\log N_1)^{1+\frac{2}{2-p}}} \quad (3.22)$$

whence (1.6) follows.

Now we look at $p = 2$. If $\alpha \lesssim \sqrt{r}$ then Lemma 3.2 gives

$$P(B_{N_1, N_2, p}^c(n)) \leq N_2 \exp(N_1(S(r) - c\sqrt{r})) \lesssim e^{-c\sqrt{nN_1}} \quad r \in [0, r_0], \quad (3.23)$$

where again the above estimate determines r_0 . Thus by (3.15) with $R = \lfloor rN_1 \rfloor$

$$P\left(\forall \mu \in [N_2] \quad \text{dLM}_{N_1}^{(\mu)} \cap \widehat{B}_{\mu, [r_0 N_1]}^{N_1} \subseteq \widehat{B}_{\mu, [rN_1]}^{N_1}\right) \gtrsim 1 - N_1 e^{-c\sqrt{r}N_1}, \quad (3.24)$$

whence the $p = 2$ part of Theorem 1.1 follows. \square

4. ABSENCE OF RETRIEVAL FOR $p \in (1, 2]$

In this section we present the proof of Theorem 1.2.

We set for brevity for $\mu \in [N_2] \setminus \{1\}$, $p \in [1, 2]$, $k \in [N_1]$, $J \subseteq [N_1]$, $v \in \{-1, 1\}^{N_1}$

$$W_{p,k,J}^{(\mu)}(v) := \frac{2p}{\sqrt{N_1}} \xi_k^{(\mu)} v_k \text{sign}(Z_J^{(\mu)}(v)) |Z_J^{(\mu)}(v)|^{p-1} \quad (4.1)$$

where

$$Z_J^{(\mu)}(v) := \frac{1}{\sqrt{N_1}} (\xi^{(\mu)}, F_J v). \quad (4.2)$$

Next we give the central technical lemma employed in the proof of Theorem 1.2. In the sequel we shorten $J+k := J \cup \{k\}$ if $k \notin J$ and $J-k := J \setminus \{k\}$ if $k \in J$.

Lemma 4.1. *Let $r \in (0, \frac{1}{2}]$, $J \subseteq [N_1]$ with $|J| = \lfloor rN_1 \rfloor$. For any $p \in [1, 2]$ we have*

$$\begin{aligned} N_1(H^{(p)}(F_{J \pm k} \xi^{(1)}) - H^{(p)}(F_J \xi^{(1)})) &= \mp \sum_{\mu=2}^{N_2} W_{p,k,J}^{(\mu)}(\xi^{(1)}) - \alpha \varsigma N_1^{1-\frac{p}{2}} \\ &\pm \frac{2p(1-2r)^{p-1}}{N_1^{1-\frac{p}{2}}} + O\left(\frac{1}{N_1^{2-\frac{p}{2}}}\right) \end{aligned} \quad (4.3)$$

where ς is a positive and uniformly bounded random variable depending on $\{\xi_k^{(1)} \xi_k^{(\mu)}, Z_J^{(\mu)}(\xi^{(1)})\}_{\mu=2 \dots N_2}$. In particular

$$\frac{N_1}{4} (H^{(2)}(F_{J \pm k} \xi^{(1)}) - H^{(2)}(F_J \xi^{(1)})) = \mp \sum_{\mu \geq 2} \frac{\xi_k^{(1)} \xi_k^{(\mu)}}{\sqrt{N_1}} Z_J^{(\mu)}(\xi^{(1)}) - \alpha \pm (1-2r) \mp \frac{1}{N_1}. \quad (4.4)$$

Proof. By Lemma 3.1 we have for any $k \notin J$

$$\begin{aligned} N_1(H^{(p)}(F_{J+k} \xi^{(1)}) - H^{(p)}(F_J \xi^{(1)})) &= N_1(H(F_{J+k} \xi^{(1)}) - H(\xi^{(1)}) - (H(F_J \xi^{(1)}) - H(\xi^{(1)}))) \\ &= N_1^{\frac{p}{2}} \left(\bar{F}_p \left(r + \frac{1}{N_1} \right) - \bar{F}_p(r) \right) \\ &+ \sum_{\mu=2}^{N_2} F_p(X_{J+k}^{(\mu)}(\xi^{(1)}), Y_{J+k}^{(\mu)}(\xi^{(1)})) \\ &- \sum_{\mu=2}^{N_2} F_p(X_J^{(\mu)}(\xi^{(1)}), Y_J^{(\mu)}(\xi^{(1)})) \\ &= N_1^{\frac{p}{2}} \left(|1-2r|^p - \left| 1-2r - \frac{2}{N_1} \right|^p \right) \\ &+ \sum_{\mu=2}^{N_2} \left(|Z_J^{(\mu)}(\xi^{(1)})|^p - \left| Z_J^{(\mu)}(\xi^{(1)}) + 2 \frac{\xi_k^{(\mu)} \xi_k^{(1)}}{\sqrt{N_1}} \right|^p \right). \end{aligned} \quad (4.5)$$

Similarly for all $k \in J$

$$\begin{aligned} N_1(H^{(p)}(F_{J-k} \xi^{(1)}) - H^{(p)}(F_J \xi^{(1)})) &= N_1^{\frac{p}{2}} \left(|1-2r|^p - \left| 1-2r + \frac{2}{N_1} \right|^p \right) \\ &+ \sum_{\mu=2}^{N_2} \left(|Z_J^{(\mu)}(\xi^{(1)})|^p - \left| Z_J^{(\mu)}(\xi^{(1)}) - 2 \frac{\xi_k^{(\mu)} \xi_k^{(1)}}{\sqrt{N_1}} \right|^p \right) \end{aligned} \quad (4.6)$$

For $p = 2$ a straightforward computation gives (4.4) from (4.5) and (4.6).

In general for $p \in [1, 2)$ we have to use Taylor expansion. Let $(p)_0 := 1$ and $(p)_k := \prod_{j=0}^{k-1} (p-j)$ for $k \geq 1$. Assuming $r \in (0, \frac{1}{2})$, N_1 large enough (*i.e.* $N_1(1-2r) > 2$) we have

$$\left| 1 - 2r \pm \frac{2}{N_1} \right|^p = |1 - 2r|^p \pm \frac{2p}{N_1} |1 - 2r|^{p-1} + \frac{4}{N_1^2} \sum_{k \geq 2} \frac{(p)_k}{k!} \frac{2^{k-2} (1-2r)^{p-k}}{N_1^{k-2}} \quad (4.7)$$

and using $|(p)_k| \leq k!$ we get

$$\left| \frac{4}{N_1^2} \sum_{k \geq 2} \frac{(p)_k}{k!} \frac{2^{k-2} (1-2r)^{p-k}}{N_1^{k-2}} \right| \leq \frac{4}{N_1^2 (1-2r)^{2-p}} \sum_{k \geq 0} \frac{2^k}{((1-2r)N_1)^k} \lesssim \frac{1}{N_1^2 (1-2r)^{2-p}}. \quad (4.8)$$

Therefore

$$N_1^{\frac{p}{2}} \left(|1 - 2r|^p - \left| 1 - 2r \pm \frac{2}{N_1} \right|^p \right) = \mp \frac{2p}{N_1^{1-\frac{p}{2}}} |1 - 2r|^{p-1} + O\left(\frac{1}{N_1^{2-\frac{p}{2}}}\right). \quad (4.9)$$

On the other hand, for $r = \frac{1}{2}$ this correction term is trivially of order $N^{-\frac{p}{2}}$.

Recall now (4.1) and compute

$$\begin{aligned} & \left| Z_J^{(\mu)}(\xi^{(1)}) \pm 2 \frac{\xi_k^{(\mu)} \xi_k^{(1)}}{\sqrt{N_1}} \right|^p \\ &= \mathbf{1}_{\{|Z_J^{(\mu)}(\xi^{(1)})| \geq 2N_1^{-\frac{1}{2}}\}} \left(|Z_J^{(\mu)}(\xi^{(1)})|^p \pm W_{p,k,J}^{(\mu)}(\xi^{(1)}) \right) \\ &+ \sum_{\ell \geq 2} \frac{(p)_\ell}{\ell!} \frac{|Z_J^{(\mu)}(\xi^{(1)})|^p}{(Z_J^{(\mu)}(\xi^{(1)}))^\ell} \frac{(\pm 2 \xi_k^{(\mu)} \xi_k^{(1)})^\ell}{N_1^{\frac{\ell}{2}}} \\ &+ \mathbf{1}_{\{|Z_J^{(\mu)}(\xi^{(1)})| < 2N_1^{-\frac{1}{2}}\}} \frac{2^p}{N_1^{\frac{p}{2}}} \left(1 \pm \frac{p}{2} \xi_k^{(\mu)} \xi_k^{(1)} Z_J^{(\mu)}(\xi^{(1)}) \sqrt{N_1} \right) \\ &+ \sum_{\ell \geq 2} \frac{(p)_\ell}{\ell!} \left(\pm \frac{\xi_k^{(\mu)} \xi_k^{(1)}}{2} Z_J^{(\mu)}(\xi^{(1)}) \sqrt{N_1} \right)^\ell. \end{aligned}$$

Thus

$$\begin{aligned} & \left(|Z_J^{(\mu)}(\xi^{(1)})|^p - \left| Z_J^{(\mu)}(\xi^{(1)}) \pm 2 \frac{\xi_k^{(\mu)} \xi_k^{(1)}}{\sqrt{N_1}} \right|^p \right) = \mp W_{p,k,J}^{(\mu)}(\xi^{(1)}) \\ &- \mathbf{1}_{\{|Z_J^{(\mu)}(\xi^{(1)})| \geq 2N_1^{-\frac{1}{2}}\}} \sum_{\ell \geq 2} \frac{(p)_\ell}{\ell!} \frac{|Z_J^{(\mu)}(\xi^{(1)})|^p}{(Z_J^{(\mu)}(\xi^{(1)}))^\ell} \frac{(\pm 2 \xi_k^{(\mu)} \xi_k^{(1)})^\ell}{N_1^{\frac{\ell}{2}}} \quad (4.10) \end{aligned}$$

$$\begin{aligned} &- \mathbf{1}_{\{|Z_J^{(\mu)}(\xi^{(1)})| < 2N_1^{-\frac{1}{2}}\}} \left(\frac{2^p}{N_1^p} \pm \frac{2^p}{N_1^p} \frac{\xi_k^{(\mu)} \xi_k^{(1)}}{2} Z_J^{(\mu)}(\xi^{(1)}) \sqrt{N_1} - |Z_J^{(\mu)}(\xi^{(1)})|^p \mp W_{p,k,J}^{(\mu)}(\xi^{(1)}) \right) \quad (4.11) \end{aligned}$$

$$\begin{aligned} &- \mathbf{1}_{\{|Z_J^{(\mu)}(\xi^{(1)})| < 2N_1^{-\frac{1}{2}}\}} \frac{2^p}{N_1^{\frac{p}{2}}} \sum_{\ell \geq 2} \frac{(p)_\ell}{\ell!} 2^{-\ell} (\pm Z_J^{(\mu)}(\xi^{(1)}) \xi_k^{(\mu)} \xi_k^{(1)} \sqrt{N_1})^\ell. \quad (4.12) \end{aligned}$$

The contributions (4.10) and (4.12) are very similar and will be dealt together. Using that for $\ell \geq 2$ it is $\ell - p > 0$ and $\ell^2(p)_\ell \leq 2p^2 \ell!$ we have

$$|(4.10)| \leq \frac{2p^2}{N_1^{\frac{p}{2}}} \sum_{\ell \geq 2} \ell^{-2}. \quad (4.13)$$

Similarly

$$|(4.12)| \leq \frac{2^{p+1}p^2}{N_1^{\frac{p}{2}}} \sum_{\ell \geq 2} \ell^{-2}. \quad (4.14)$$

Furthermore, depending on the value of $\text{sign}(Z_J^{(\mu)}(\xi^{(1)}))\xi_k^{(1)}\xi_k^{(\mu)}$ we can have either

$$\begin{aligned} (4.10) &= 1_{\{|Z_J^{(\mu)}(\xi^{(1)})| < 2N_1^{-\frac{1}{2}}\}} \frac{1}{N_1^{\frac{p}{2}}} \sum_{\ell \geq 2} \frac{(p)_\ell}{\ell!} \frac{|Z_J^{(\mu)}(\xi^{(1)})|^p}{|Z_J^{(\mu)}(\xi^{(1)})|^\ell} \frac{1}{N_1^{\frac{\ell}{2}}} \\ &= 1_{\{|Z_J^{(\mu)}(\xi^{(1)})| < 2N_1^{-\frac{1}{2}}\}} \frac{|Z_J^{(\mu)}(\xi^{(1)})|^p}{N_1^{\frac{p}{2}}} \left[(1 + |Z_J^{(\mu)}(\xi^{(1)})\sqrt{N_1}|^{-1})^p - (1 + |Z_J^{(\mu)}(\xi^{(1)})\sqrt{N_1}|^{-1}) \right] \geq 0 \end{aligned} \quad (4.15)$$

and

$$\begin{aligned} (4.12) &= 1_{\{|Z_J^{(\mu)}(\xi^{(1)})| \geq 2N_1^{-\frac{1}{2}}\}} \frac{2^p}{N_1^{\frac{p}{2}}} \sum_{\ell \geq 2} \sum_{\ell \geq 2} \frac{(p)_\ell}{\ell!} 2^{-\ell} |Z_J^{(\mu)}(\xi^{(1)})\sqrt{N_1}|^\ell \\ &= 1_{\{|Z_J^{(\mu)}(\xi^{(1)})| \geq 2N_1^{-\frac{1}{2}}\}} \frac{|Z_J^{(\mu)}(\xi^{(1)})|^p}{N_1^{\frac{p}{2}}} \left[(1 + |2^{-1}Z_J^{(\mu)}(\xi^{(1)})\sqrt{N_1}|)^p - (1 + |2^{-1}Z_J^{(\mu)}(\xi^{(1)})\sqrt{N_1}|) \right] \geq 0 \end{aligned} \quad (4.16)$$

(equality is achieved only if $p = 1$) or

$$\begin{aligned} (4.10) &= 1_{\{|Z_J^{(\mu)}(\xi^{(1)})| < 2N_1^{-\frac{1}{2}}\}} \frac{1}{N_1^{\frac{p}{2}}} \sum_{\ell \geq 2} (-1)^\ell \frac{(p)_\ell}{\ell!} \frac{|Z_J^{(\mu)}(\xi^{(1)})|^p}{|Z_J^{(\mu)}(\xi^{(1)})|^\ell} \frac{2^\ell}{N_1^{\frac{\ell}{2}}} \\ &= 1_{\{|Z_J^{(\mu)}(\xi^{(1)})| < 2N_1^{-\frac{1}{2}}\}} \frac{|Z_J^{(\mu)}(\xi^{(1)})|^p}{N_1^{\frac{p}{2}}} \sum_{\ell \geq 1} \left(\frac{\sqrt{N_1}Z_J^{(\mu)}(\xi^{(1)})}{2} \right)^{-2\ell} \left(\frac{(p)_{2\ell}}{2\ell!} - \frac{(p)_{2\ell+1}}{2\ell+1!} \left| \frac{\sqrt{N_1}Z_J^{(\mu)}(\xi^{(1)})}{2} \right|^{-1} \right) \end{aligned} \quad (4.17)$$

(split the sum over even and odd $\ell \geq 2$ and rename the indices) and

$$\begin{aligned} (4.12) &= 1_{\{|Z_J^{(\mu)}(\xi^{(1)})| \geq 2N_1^{-\frac{1}{2}}\}} \frac{2^p}{N_1^{\frac{p}{2}}} \sum_{\ell \geq 2} (-1)^\ell \frac{(p)_\ell}{\ell!} 2^{-\ell} |Z_J^{(\mu)}(\xi^{(1)})\sqrt{N_1}|^\ell \\ &= 1_{\{|Z_J^{(\mu)}(\xi^{(1)})| \geq 2N_1^{-\frac{1}{2}}\}} \frac{2^p}{N_1^{\frac{p}{2}}} \sum_{\ell \geq 1} \left(\frac{\sqrt{N_1}Z_J^{(\mu)}(\xi^{(1)})}{2} \right)^{2\ell} \left(\frac{(p)_{2\ell}}{2\ell!} - \frac{(p)_{2\ell+1}}{2\ell+1!} \frac{|\sqrt{N_1}Z_J^{(\mu)}(\xi^{(1)})|}{2} \right). \end{aligned} \quad (4.18)$$

The quantities in the two displays above are positive, since for $\ell \geq 1$ $(p)_{2\ell} \geq 0$ and $(p)_{2\ell+1} \leq 0$, which can be shown by observing that $(p)_\ell \geq 3 = (-1)^\ell p(p-1) \prod_{j=2}^{\ell-1} (j-p)$ (again the equality is achieved only for $p = 1$). We conclude

$$(4.10) < 0, \quad (4.12) < 0 \quad \text{for } p \in (1, 2) \quad \text{and} \quad (4.10) = (4.12) = 0 \quad \text{for } p = 1.$$

With a bit of algebra we rewrite the term in (4.11) as

$$\begin{aligned} &1_{\{|Z_J^{(\mu)}(\xi^{(1)})| < 2N_1^{-\frac{1}{2}}\}} \frac{2^p}{N_1^{\frac{p}{2}}} \left(\left(1 - \frac{|2Z_J^{(\mu)}(\xi^{(1)})|^p}{N_1^{\frac{p}{2}}} \right) \right. \\ &\quad \left. \mp p\xi_k^{(\mu)}\xi_k^{(1)} \text{sign}(Z_J^{(\mu)}(\xi^{(1)})) \left(\frac{|2Z_J^{(\mu)}(\xi^{(1)})|^{p-1}}{N_1^{\frac{p-1}{2}}} - \frac{|2Z_J^{(\mu)}(\xi^{(1)})|}{N_1^{\frac{1}{2}}} \right) \right) \end{aligned} \quad (4.19)$$

According to the value of $\xi_k^{(\mu)} \xi_k^{(1)} \text{sign}(Z_J^{(\mu)}(\xi^{(1)}))$ the term inside the parenthesis can be either

$$1 - |x|^p + p(|x|^{p-1} - |x|) \quad \text{or} \quad 1 - |x|^p - p(|x|^{p-1} - |x|),$$

where we shortened $|x| := \frac{|2Z_J^{(\mu)}(\xi^{(1)})|}{N_1^{\frac{1}{2}}} < 1$. The first expression above is clearly positive, while the second one is positive thanks to Lemma B.1. From the representation (4.19) we also get the bound

$$(4.11) \leq \frac{C}{N_1^{\frac{p}{2}}}. \quad (4.20)$$

Therefore there is a positive and uniformly bounded random variable

$$\varsigma := \varsigma(\{\xi_k^{(1)} \xi_k^{(\mu)}, Z_J^{(\mu)}(\xi^{(1)})\}_{\mu=2 \dots N_2})$$

such that

$$\sum_{\mu=2}^{N_2} \left(|Z_J^{(\mu)}(\xi^{(1)})|^p - \left| Z_J^{(\mu)}(\xi^{(1)}) \pm 2 \frac{\xi_k^{(\mu)} \xi_k^{(1)}}{\sqrt{N_1}} \right|^p \right) = \mp \sum_{\mu=2}^{N_2} W_{p,k,J}^{(\mu)}(\xi^{(1)}) + \alpha \varsigma N_1^{1-\frac{p}{2}}. \quad (4.21)$$

This and (4.9) give (4.3). \square

Now we turn to the proof of Theorem 1.2, which we conveniently split in several steps.

Step 1: reduction. Due to the exchangeability of the patterns and their entries we have

$$\begin{aligned} & P\left(\mathbf{LM}_{N_1}^{(\mu)} \cap \widehat{B}_{\mu, \lfloor rN_1 \rfloor}^{N_1} \neq \emptyset\right) \\ & \leq N_2 \sum_{\ell=1}^{\lfloor rN_1 \rfloor} \sum_{\substack{J \subset [N_1] \\ |J|=\ell}} P\left(\bigcap_{k \notin J} \{H(F_{J+k}\xi^{(1)}) - H(F_J\xi^{(1)}) > 0\}, \bigcap_{k \in J} \{H(F_{J-k}\xi^{(1)}) - H(F_J\xi^{(1)}) > 0\}\right) \\ & = N_2 \sum_{\ell=1}^{\lfloor rN_1 \rfloor} \binom{N_1}{\ell} P\left(\bigcap_{k>\ell} \{H(F_{[\ell]+k}\xi^{(1)}) - H(F_{[\ell]}\xi^{(1)}) > 0\}, \bigcap_{k \leq \ell} \{H(F_{[\ell]-k}\xi^{(1)}) - H(F_{[\ell]}\xi^{(1)}) > 0\}\right). \end{aligned} \quad (4.22)$$

We set for brevity

$$Z_\ell^{(\mu)}(\xi^{(1)}) := \frac{1}{\sqrt{N_1}} \left(\sum_{k=1}^{\ell} \xi_k^{(1)} \xi_k^{(\mu)} - \sum_{k \geq \ell+1} \xi_k^{(1)} \xi_k^{(\mu)} \right), \quad (4.23)$$

$$W_{p,k,\ell}^{(\mu)}(\xi^{(1)}) := \frac{2p}{\sqrt{N_1}} \xi_k^{(\mu)} \xi_k^{(1)} \text{sign}(Z_{[\ell]}^{(\mu)}(\xi^{(1)})) |Z_{[\ell]}^{(\mu)}(\xi^{(1)})|^{p-1}, \quad W_{p,k,\ell}^{(\mu)} := W_{p,k,\ell}^{(\mu)}(1), \quad (4.24)$$

$$M^{(\mu)} := \frac{1}{N_1} \sum_{i \in [N_1]} \xi_i^{(\mu)}, \quad Q^{(\mu)} := \text{sign}(M^{(\mu)}) |M^{(\mu)}|^{p-1}. \quad (4.25)$$

By Lemma B.1 in Appendix B we have for N_1 large enough ($r' := \ell/N_1$)

$$\begin{aligned}
(4.22) &= N_2 \sum_{\ell=1}^{\lfloor rN_1 \rfloor} \binom{N_1}{\ell} P \left(\forall k > \ell - \sum_{\mu=2}^{N_2} W_{p,k,\ell}^{(\mu)}(\xi^{(1)}) \gtrsim \alpha N_1^{1-\frac{p}{2}} - \frac{2p(1-2r')^{p-1}}{N_1^{1-\frac{p}{2}}}, \right. \\
&\quad \left. \forall k \in [\ell] \sum_{\mu=2}^{N_2} W_{p,k,\ell}^{(\mu)}(\xi^{(1)}) \gtrsim \alpha N_1^{1-\frac{p}{2}} + \frac{2p(1-2r')^{p-1}}{N_1^{1-\frac{p}{2}}} \right) \\
&= N_2 \sum_{\ell=1}^{\lfloor rN_1 \rfloor} \binom{N_1}{\ell} P \left(\forall k > \ell - \sum_{\mu=2}^{N_2} W_{p,k,\ell}^{(\mu)} \gtrsim \alpha N_1^{1-\frac{p}{2}} - \frac{2p(1-2r')^{p-1}}{N_1^{1-\frac{p}{2}}}, \right. \\
&\quad \left. \forall k \in [\ell] \sum_{\mu=2}^{N_2} W_{p,k,\ell}^{(\mu)} \gtrsim \alpha N_1^{1-\frac{p}{2}} + \frac{2p(1-2r')^{p-1}}{N_1^{1-\frac{p}{2}}} \right) \\
&= N_2 \sum_{\ell=1}^{\lfloor rN_1 \rfloor} \binom{N_1}{\ell} P \left(\forall k > \ell (Q, \tilde{\xi}^{(k)}) \gtrsim \alpha N_1^{1-\frac{p}{2}} - \frac{2p(1-2r')^{p-1}}{N_1^{1-\frac{p}{2}}}, \right. \\
&\quad \left. \forall k \in [\ell] (Q, \tilde{\xi}^{(k)}) \gtrsim \alpha N_1^{1-\frac{p}{2}} + \frac{2p(1-2r')^{p-1}}{N_1^{1-\frac{p}{2}}} \right) \tag{4.26}
\end{aligned}$$

(recall that $\tilde{\xi}^{(k)}$ denotes the k -th transposed pattern). In the second identity above we have exploited independence of the pattern $\xi^{(1)}$ to replace $W_{p,k,\ell}^{(\mu)}(\xi^{(1)})$ by $W_{p,k,\ell}^{(\mu)}$ and in the third one the independence of the first ℓ entries from all the others and the flip-symmetry to replace $Z_\ell^{(\mu)}$ by $M^{(\mu)}$. We also used that the independent random variables $Q^{(2)} \dots Q^{(N_2)}$ are symmetric.

Step 2: disentangling by the FKG inequality. Let us now call \mathcal{M} the σ -field generated by $M^{(2)}, \dots, M^{(N_2)}$ and notice that $Q^{(2)} \dots Q^{(N_2)}$ are \mathcal{M} -measurable. We shorten $P_{\mathcal{M}}(\cdot) := P(\cdot | \mathcal{M})$. The crucial observation here (first done for the Hopfield model in [15]) is that for each $\mu \in [N_2]$ the law of $\xi^{(\mu)}$ conditionally on \mathcal{M} is a permutation distribution (as the increments of a simple random walk given the position). Therefore the FKG inequality applies (see for instance [9]) and we have

$$\begin{aligned}
&P_{\mathcal{M}} \left(\forall k > \ell (Q, \tilde{\xi}^{(k)}) \gtrsim \alpha N_1^{1-\frac{p}{2}} - \frac{2p(1-2r')^{p-1}}{N_1^{1-\frac{p}{2}}}, \right. \\
&\quad \left. \forall k \in [\ell] (Q, \tilde{\xi}^{(k)}) \gtrsim \alpha N_1^{1-\frac{p}{2}} + \frac{2p(1-2r')^{p-1}}{N_1^{1-\frac{p}{2}}} \right) \\
&\leq \prod_{k>\ell} P_{\mathcal{M}} \left((Q, \tilde{\xi}^{(k)}) \gtrsim \alpha N_1^{1-\frac{p}{2}} - \frac{2p(1-2r')^{p-1}}{N_1^{1-\frac{p}{2}}} \right) \\
&\quad \prod_{k \in [\ell]} P_{\mathcal{M}} \left((Q, \tilde{\xi}^{(k)}) \gtrsim \alpha N_1^{1-\frac{p}{2}} + \frac{2p(1-2r')^{p-1}}{N_1^{1-\frac{p}{2}}} \right) \tag{4.27}
\end{aligned}$$

Note that $E[\xi_k^{(\mu)} | \mathcal{M}] = M^{(\mu)}$ and $\text{Var}[\xi_k^{(\mu)} | \mathcal{M}] = 1 - (M^{(\mu)})^2$. Therefore introducing $\overline{Q} \in \mathbb{R}^{N_2}$ with components $\overline{Q}^{(\mu)} := Q^{(\mu)} \sqrt{1 - (M^{(\mu)})^2}$, we get

$$P_{\mathcal{M}} \left((Q, \tilde{\xi}^{(k)}) \geq t \right) \leq \exp \left(-c \frac{(t - \|M\|_p^p)^2}{\|\overline{Q}\|_2^2} \right), \tag{4.28}$$

by the Hoeffding inequality. Note that this quantity is independent on k and also we have the simple bound $\|\widehat{Q}\|_2^2 \leq \|Q\|_2^2 = \|M\|_{2p-2}^{2p-2}$. Then

$$(4.27)_{p \in [1,2]} \leq \exp \left(-cN_1 \frac{(\alpha N_1^{1-\frac{2}{p}} - \|M\|_p^p)^2}{\|M\|_{2p-2}^{2p-2}} \right), \quad (4.29)$$

$$(4.27)_{p=2} \leq \exp \left(-\frac{cN_1}{\|M\|_2^2} \left(r'(\alpha - (1-2r') - \|M\|_2^2)^2 + (1-r')(\alpha + (1-2r') - \|M\|_2^2)^2 \right) \right). \quad (4.30)$$

Step 3: concentration. Now we have to take the global expectations of the r.h.s. above. A similar computation as in (2.20) gives $E[\|M\|_p^p] \simeq \alpha N_1^{1-\frac{2}{p}}$ (this is an identity for $p = 2$). Moreover $\|(M^{(\mu)})^p - E[(M^{(\mu)})^p]\|_{\psi_{\frac{2}{p}}} \simeq N^{-\frac{2}{p}}$. Hence by Proposition A.1 in Appendix A (with $\ell = 2/p \geq 1$)

$$P \left(\left| \sum_{\mu \geq 2} (M^{(\mu)})^p - E[(M^{(\mu)})^p] \right| \geq t \right) \leq \exp \left(-c \min \left(\frac{t^2 N_1^{p-1}}{\alpha}, \frac{t^{\frac{2}{p}} N^{2-\frac{2}{p}}}{\alpha^{\frac{2}{p}-1}} \right) \right). \quad (4.31)$$

Step 4: finalising the argument for $p \in (1, 2)$. Using (4.31) with $t = \frac{1}{2}(\alpha N_1^{1-\frac{2}{p}} - E[\|M\|_p^p])$ in the first line below and likewise but for $2p-2$ in the second line we have

$$\begin{aligned} E[\text{r.h.s. of (4.29)}] &\leq E \left[\exp \left(-c \frac{\alpha^2 N_1^{3-p}}{\|M\|_{2p-2}^{2p-2}} \right) \right] + e^{-c \min(\alpha, \alpha^2) N_1} \\ &\leq e^{-c\alpha N_1} + 2e^{-c \min(\alpha, \alpha^2) N_1} \end{aligned} \quad (4.32)$$

Combining (4.22), (4.27), (4.29), (4.32) and using the standard bound (1.10) we have for $p \in [1, 2)$

$$\begin{aligned} P \left(\text{LM}_{N_1}^{(\mu)} \cap \widehat{B}_{\mu, \lfloor rN_1 \rfloor}^{N_1} \neq \emptyset \right) &\leq N_2 \sum_{\ell=1}^{\lfloor rN_1 \rfloor} \binom{N_1}{\ell} \left(e^{-c\alpha N_1} + 2e^{-c \min(\alpha, \alpha^2) N_1} \right) \\ &\lesssim rN_2 N_1 e^{N_1(S(r) - c \min(\alpha, \alpha^2))}, \end{aligned} \quad (4.33)$$

which yields the assertion for $p \in [1, 2)$.

Step 5: finalising the argument for $p = 2$. Using (4.31) _{$p=2$} with $t = \tau := (1+\alpha)(1-2r')$ in the first line in the following display and $t = C\alpha$ in the second line, we have

$$\begin{aligned} E[\text{r.h.s. of (4.30)}] &\leq E \left[\exp \left(-cN_1 \frac{((1-2r') - \tau)^2}{\|M\|_2^2} \right) \right] + e^{-cN_1(1+\alpha)(1-2r') \min(1, \frac{(1+\alpha)(1-2r')}{\alpha})} \\ &\leq e^{-c\alpha N_1(1-2r')^2} + e^{-C\alpha N_1} + e^{-cN_1(1+\alpha)(1-2r') \min(1, \frac{(1+\alpha)(1-2r')}{\alpha})} \\ &\leq e^{-c\alpha N_1(1-2r')^2} + e^{-C\alpha N_1} + e^{-cN_1(1+\alpha)(1-2r') \min(1, \frac{(1+\alpha)(1-2r')}{\alpha})}, \end{aligned} \quad (4.34)$$

as r' ranges from $1/N_1$ to r . So combining (4.22), (4.27), (4.29), (4.34) and using again (1.10) we obtain

$$\begin{aligned} P \left(\text{LM}_{N_1}^{(\mu)} \cap \widehat{B}_{\mu, \lfloor rN_1 \rfloor}^{N_1} \neq \emptyset \right) &\leq N_2 \sum_{\ell=1}^{\lfloor rN_1 \rfloor} \binom{N_1}{\ell} \left(e^{-c\alpha N_1(1-2r)^2} + e^{-c\alpha N_1} + e^{-cN_1(1+\alpha)(1-2r) \min(1, \frac{(1+\alpha)(1-2r)}{\alpha})} \right) \\ &\lesssim rN_2 N_1 \left(e^{N_1(S(r) - c\alpha(1-2r)^2)} + e^{N_1(S(r) - c\alpha)} \right. \\ &\quad \left. + e^{N_1(S(r) - c(1+\alpha)(1-2r) \min(1, \frac{(1+\alpha)(1-2r)}{\alpha}))} \right). \end{aligned}$$

The first two summands are negative if $\alpha > S(r)/(1-2r)^2$. The fact that also the third one is so is verified in Lemma B.2, Appendix B. The proof is complete.

APPENDIX A. A TAIL ESTIMATE

Here we present a tail estimate for sums of i.i.d. r.v.s which is used in the main text. The following statement is not new and we give the proof here mainly for the reader's convenience. In fact the proof of the subsequent formula (A.1) for $\ell \in [1, 2]$ is classical and can be found for instance in [19, Corollaries 2.9, 2.10]. So we focus on the case $\ell \in (0, 1)$. For similar statements, see [16, Proposition 3.2] and [13, Theorem 6.21].

Proposition A.1. *Let $\ell \in (0, 2]$, X_1, \dots, X_N i.i.d. r.v.s with $\|X_1\|_{\psi_\ell} < \infty$. There is $c = c(\ell) > 0$ such that*

$$P\left(\left|\sum_{i \in [N]} X_i\right| \geq t\right) \leq \exp\left(-c \min\left(\frac{t^2}{\|X_1\|_{\psi_\ell}^2 N}, \frac{t^\ell}{\|X_1\|_{\psi_\ell}^\ell N^{\max(\ell-1, 0)}}\right)\right). \quad (\text{A.1})$$

only for $\ell \in (0, 1)$. We have by assumption

$$P(|X_1| \geq t) \leq e^{-\frac{t^\ell}{2\|X_1\|_{\psi_\ell}^\ell}}. \quad (\text{A.2})$$

Let now $s := N^{\frac{1}{2-\ell}}$ and set

$$X_i^s := X_i 1_{\{|X_i| < s\}}. \quad (\text{A.3})$$

Then we have

$$\begin{aligned} P\left(\left|\sum_{i \in [N]} X_i\right| \geq t\right) &\leq P\left(\sum_{i \in [N]} X_i \geq t, \sup_{i \in [N]} |X_i| < s\right) + P\left(\sup_{i \in [N]} |X_i| \geq s\right) \\ &\leq P\left(\left|\sum_{i \in [N]} X_i^s\right| \geq t\right) + e^{-\frac{s^\ell}{4\|X_1\|_{\psi_\ell}^\ell}}. \end{aligned} \quad (\text{A.4})$$

Set now

$$\bar{\mu} := \frac{1}{10s^{1-\ell}\|X\|_{\psi_\ell}^\ell},$$

and for $0 \leq \mu \leq \bar{\mu}$ we note that

$$(\mu X_i^s)^n \leq \left(\frac{(X_i^s)^\ell}{10\|X\|_{\psi_\ell}^\ell}\right)^n.$$

Using the bound $x^2 \leq e^{\frac{x}{10}}$ we compute

$$\begin{aligned} E[e^{\mu X_i^s}] &= 1 + \mu^2 \|X_i^s\|_{\psi_\ell}^2 \sum_{n \geq 0} \mu^n \frac{E[(X_i^s)^{n+2}]}{\|X_i^s\|_{\psi_\ell}^2 (n+2)!} \\ &\leq 1 + \mu^2 \|X_i^s\|_{\psi_\ell}^2 E\left[e^{\frac{(X_i^s)^\ell}{10\|X\|_{\psi_\ell}^\ell}} \sum_{n \geq 0} \frac{1}{n!} \left(\frac{(X_i^s)^\ell}{10\|X\|_{\psi_\ell}^\ell}\right)^n\right] \\ &\leq 1 + \mu^2 \|X_i^s\|_{\psi_\ell}^2 E\left[e^{\frac{(X_i^s)^\ell}{5\|X\|_{\psi_\ell}^\ell}}\right] \\ &\leq \exp\left(\mu^2 \|X_i^s\|_{\psi_\ell}^2 E\left[e^{\frac{(X_i^s)^\ell}{5\|X\|_{\psi_\ell}^\ell}}\right]\right) \leq 2\mu^2 \|X_i^s\|_{\psi_\ell}^2. \end{aligned} \quad (\text{A.5})$$

It follows that

$$P \left(\left| \sum_{i \in [N]} X_i^s \right| \geq t \right) \leq 2e^{-\mu t + 2N\mu^2 \|X_i^s\|_{\psi_\ell}^2} \leq \begin{cases} 2 \exp \left(-\frac{t^2}{8 \|X_i^s\|_{\psi_\ell}^2 N} \right) & 0 < t < 4N\bar{\mu}; \\ 2 \exp \left(-\bar{\mu}t + 2N\bar{\mu}^2 \|X_i^s\|_{\psi_\ell}^2 \right) & t \geq 4N\bar{\mu}. \end{cases} \quad (\text{A.6})$$

With our choice of parameters the above formula rewrites as

$$P \left(\left| \sum_{i \in [N]} X_i^s \right| \geq t \right) \leq \begin{cases} 2 \exp \left(-\frac{t^2}{8 \|X_i^s\|_{\psi_\ell}^2 N} \right) & 0 < t < 4s; \\ 2 \exp \left(-\frac{t^\ell}{10(4 \|X_1\|_{\psi_\ell}^\ell)^\ell} + \frac{t^\ell}{50(4 \|X_1\|_{\psi_\ell}^\ell)^{\ell-1}} \right) & t \geq 4s. \end{cases} \quad (\text{A.7})$$

Combining (A.4) and (A.7) gives the assertion. \square

APPENDIX B. TWO TECHNICAL LEMMAS

The following two results are basically calculus.

Lemma B.1. *Let $f(x, p) := 1 - x^p - px^{p-1} + px$. It is $f \geq 0$ for all $p \in [1, 2]$ and $x \in [0, 1]$.*

Proof. First we note that

$$1 + (x + p) \log x \leq x, \quad \forall x \in [0, 1]. \quad (\text{B.1})$$

The proof is simple: we compare the function $\log x$ with $\frac{x-1}{x+p}$ for $x \in [0, 1]$ and since

$$\frac{1}{x} = \frac{d}{dx} \log x \geq \frac{d}{dx} \frac{x-1}{x+p} = \frac{p+1}{(x+p)^2} \quad \forall x \in [0, 1]$$

and in $x = 1$ the two functions intersect, (B.1) follows.

Next we note that $f(x, 1) = 0$ and $f(x, 2) \geq 0$ for all $x \in [0, 1]$. Then it suffices to show that f is not decreasing in p uniformly in $x \in [0, 1]$. We compute

$$\frac{\partial}{\partial x} f(x, p) = x \left(1 - x^{p-1} \left(\frac{1 + (x+p) \log x}{x} \right) \right) \geq x (1 - x^{p-1}) \geq 0$$

thanks to (B.1). \square

Lemma B.2. *Let $r \in [0, \frac{1}{2}]$, $c_1 > 0$. There is $c_2 > 0$ such that for all $\alpha \geq c_2 S(r)/(1-2r)^2$ it holds*

$$S(r) \leq c_1(1+\alpha)(1-2r) \min \left(1, \frac{(1+\alpha)(1-2r)}{\alpha} \right). \quad (\text{B.2})$$

Proof. (B.2) selects two conditions, namely either

$$\alpha \leq \frac{1-2r}{2r}, \quad c_1 \alpha \geq \frac{S(r)}{1-2r} - c_1 \quad \text{or} \quad \alpha > \frac{1-2r}{2r}, \quad c_1 \frac{(1+\alpha)^2}{\alpha} \geq \frac{S(r)}{(1-2r)^2}. \quad (\text{B.3})$$

For $r \in [0, 1/2]$ the function $S(r)$ increases and $c_1(1-2r)$ decreases. Let us denote \bar{r} their unique intersection point in $[0, 1/2]$. Clearly \bar{r} depends on c_1 and $\bar{r} \rightarrow 0$ as $c_1 \rightarrow 0$. If $r \in [0, \bar{r}]$ then for every $\alpha < (1-2r)/2r$ it holds

$$c_1 \alpha \geq \frac{S(r)}{1-2r} - c_1.$$

Moreover there is $C > 0$ such that

$$\frac{1-2r}{2r} \geq C \frac{S(r)}{(1-2r)^2} \quad \forall r \in [0, \bar{r}].$$

Indeed by definition of \bar{r} the above condition is implied by

$$\frac{1-2r}{2r} \geq \frac{c_1 C}{1-2r} \quad \forall r \in [0, \bar{r}],$$

therefore it suffices to take $c_2 := \frac{(1-2\bar{r})^2}{2c_1\bar{r}}$ and we have the statement for $r \in [0, \bar{r}]$.

For $r \in [\bar{r}, 1/2]$ we use the second condition in (B.3). First we observe that, since $(1+\alpha)^2/\alpha > \alpha$, the condition $c_1 \frac{(1+\alpha)^2}{\alpha} \geq \frac{S(r)}{(1-2r)^2}$ is implied by $\alpha \geq CS(r)/(1-2r)^2$ for all $C > c_1^{-1}$. It remains to show that there is $C > c_1^{-1}$ such that

$$\frac{1-2r}{2r} \leq C \frac{S(r)}{(1-2r)^2} \quad \text{or equivalently} \quad \frac{(1-2r)^2}{2r} \leq C \frac{S(r)}{(1-2r)}. \quad (\text{B.4})$$

By definition of \bar{r}

$$\frac{S(r)}{(1-2r)} \geq c_1 \quad \forall r \in [\bar{r}, 1/2].$$

The l.h.s. of the second inequality in (B.4) is decreasing and its r.h.s. is increasing, whence it suffices to require

$$\frac{(1-2\bar{r})^2}{2\bar{r}} \leq Cc_1.$$

Thus taking $c_2 := \max(\frac{1}{c_1}, \frac{(1-2\bar{r})^2}{2c_1\bar{r}})$ we have proved the statement also for $r \in [\bar{r}, 1/2]$. \square

REFERENCES

1. Louigi Addario-Berry, Luc Devroye, Gábor Lugosi, and Roberto I. Oliveira, *Local optima of the Sherrington-Kirkpatrick Hamiltonian*, Journal of Mathematical Physics **60** (2019), no. 4, 043301.
2. Daniel J. Amit, Hanoch Gutfreund, and H. Sompolinsky, *Storing infinite numbers of patterns in a spin-glass model of neural networks*, Phys. Rev. Lett. **55** (1985), 1530–1533.
3. Omer Angel, Sebastien Bubeck, Yuval Peres, and Fan Wei, *Local max-cut in smoothed polynomial time*, Proceedings of the 49th Annual ACM SIGACT Symposium on Theory of Computing, ACM, jun 2017.
4. Adriano Barra, Giuseppe Genovese, Peter Sollich, and Daniele Tantari, *Phase diagram of restricted Boltzmann machines and generalized Hopfield networks with arbitrary priors*, Physical Review E **97** (2018), no. 2, 022310—.
5. Marco Benedetti, Victor Dotsenko, Giulia Fischetti, Enzo Marinari, and Gleb Oshanin, *Recognition capabilities of a hopfield model with auxiliary hidden neurons*, Phys. Rev. E **103** (2021), L060401.
6. Marco Benedetti, Enrico Ventura, Enzo Marinari, Giancarlo Ruocco, and Francesco Zamponi, *Supervised versus unsupervised learning: approaching optimal memory retrieval in hopfield networks*, (2022).
7. Anton Bovier and Beat Niederhauser, *The spin-glass phase-transition in the Hopfield model with p-spin interactions*, Advances in Theoretical and Mathematical Physics **5** (2001), no. 6, 1001–1046.
8. Nicola Bulso and Yasser Roudi, *Restricted Boltzmann Machines as Models of Interacting Variables*, Neural Computation **33** (2021), no. 10, 2646–2681.
9. Desh Ranjan Devdatt Dubhashi, Volker Priebe, *Negative dependence through the FKG inequality*, Tech. Report RS - 96-27, University of Aarhus, July 1996.
10. Michael Etscheid and Heiko Röglin, *Smoothed analysis of local search for the maximum-cut problem*, ACM Transactions on Algorithms **13** (2017), no. 2, 1–12.
11. Geoffrey E Hinton and Terrence J Sejnowski, *Optimal perceptual inference*, Proceedings of the IEEE conference on Computer Vision and Pattern Recognition, vol. 448, Citeseer, 1983.
12. J. J. Hopfield, *Neural networks and physical systems with emergent collective computational abilities*, Proc. Nat. Acad. Sci. **79** (1982), 2554–2558.
13. Michel Ledoux and Michel Talagrand, *Probability in banach spaces*, Springer Berlin Heidelberg, 1991.
14. W.A. Little, *The existence of persistent states in the brain*, Mathematical Biosciences **19** (1974), no. 1, 101–120.
15. Daria Loukianova, *Lower bounds on the restitution error in the Hopfield model*, Probability Theory and Related Fields **107** (1997), no. 2, 161–176.
16. Charles M Newman, *Memory capacity in neural network models: Rigorous lower bounds*, Neural Networks **1** (1988), no. 3, 223–238.
17. Francesca Schönsberg, Yasser Roudi, and Alessandro Treves, *Efficiency of local learning rules in threshold-linear associative networks*, Phys. Rev. Lett. **126** (2021), 018301.
18. P. Smolensky, *Information processing in dynamical systems: Foundations of harmony theory*, p. 194?281, MIT Press, Cambridge, MA, USA, 1986.

19. Michel Talagrand, *The supremum of some canonical processes*, American Journal of Mathematics **116** (1994), no. 2, 283.
20. ———, *Rigorous results for the Hopfield model with many patterns*, Probability Theory and Related Fields **110** (1998), no. 2, 177—275.
21. J. Tubiana, S. Cocco, and R. Monasson, *Learning Compositional Representations of Interacting Systems with Restricted Boltzmann Machines: Comparative Study of Lattice Proteins*, Neural Computation **31** (2019), no. 8, 1671–1717.
22. J. Tubiana and R. Monasson, *Emergence of compositional representations in restricted boltzmann machines*, Phys. Rev. Lett. **118** (2017), 138301.

INSTITUTE OF MATHEMATICS, UNIVERSITY OF ZURICH, WINTERTHURERSTRASSE 190, 8057 ZURICH, SWITZERLAND.

Email address: `giuseppe.genovese@math.uzh.ch`