

ANALYTIC SMOOTHING EFFECT OF LINEAR LANDAU EQUATION WITH SOFT POTENTIAL

HAO-GUANG LI AND CHAO-JIANG XU

ABSTRACT. In this work, we study the linear Landau equation with soft potential and show that the solution to the Cauchy problem with initial datum in $L^2(\mathbb{R}^3)$ enjoys an analytic regularizing effect, and the evolution of analytic radius is same as heat equations.

1. INTRODUCTION

In this work, we study the Cauchy problem of spatially homogeneous Landau equation :

$$\begin{cases} \partial_t F = Q(F, F), \\ F|_{t=0} = F_0, \end{cases} \quad (1.1)$$

where $F = F(t, v) \geq 0$ is the density distribution function depending on the velocity variables $v \in \mathbb{R}^3$ and the time $t \geq 0$. The Landau bilinear collision operator is given by

$$Q(G, F)(v) = \sum_{j,k=1}^3 \partial_j \left(\int_{\mathbb{R}^3} a_{jk}(v - v_*) [G(v_*)(\partial_k F)(v) - (\partial_k G)(v_*)F(v)] dv_* \right),$$

where

$$a_{jk}(v) = (\delta_{jk}|v|^2 - v_j v_k) |v|^\gamma, \quad \gamma \geq -3. \quad (1.2)$$

One calls hard potentials if $\gamma > 0$, Maxwellian molecules if $\gamma = 0$, soft potentials if $\gamma \in]-3, 0[$ and Coulombian potential if $\gamma = -3$. We shall study the linearization of the Landau equation (1.1) near the absolute Maxwellian distribution

$$\mu(v) = (2\pi)^{-\frac{3}{2}} e^{-\frac{|v|^2}{2}}.$$

Considering the fluctuation of density distribution function

$$F(t, v) = \mu(v) + \sqrt{\mu}(v)f(t, v),$$

since $Q(\mu, \mu) = 0$, the Cauchy problem (1.1) is reduced to the Cauchy problem

$$\begin{cases} \partial_t f + \mathcal{L}(f) + \mathcal{K}(f) = \mathbf{\Gamma}(f, f), \quad t > 0, v \in \mathbb{R}^3, \\ f|_{t=0} = f_0, \end{cases} \quad (1.3)$$

with $F_0(v) = \mu + \sqrt{\mu}f_0(v)$, where

$$\mathbf{\Gamma}(f, f) = \mu^{-\frac{1}{2}} Q(\sqrt{\mu}f, \sqrt{\mu}f),$$

Date: May 17, 2022.

2010 Mathematics Subject Classification. 35B65, 76P05, 82C40.

Key words and phrases. linear Landau equation; analytic smoothing effect; soft potential.

and

$$\mathcal{L}f = -\Gamma(\sqrt{\mu}, f), \quad \mathcal{K}f = -\Gamma(f, \sqrt{\mu}). \quad (1.4)$$

In this work, we study Cauchy problem of the linear Landau equation, such as

$$\begin{cases} \partial_t f = \mathcal{L}f, \\ f|_{t=0} = f_0 \in L^2(\mathbb{R}^3). \end{cases} \quad (1.5)$$

Using the references [9] and [19], we show that the diffusion part \mathcal{L} is written as follows

$$\mathcal{L}f = -\nabla_v \cdot [A(v)\nabla_v f] + (A(v)\frac{v}{2} \cdot \frac{v}{2})f - \nabla_v \cdot [A(v)\frac{v}{2}]f, \quad (1.6)$$

with $A(v) = (\bar{a}_{ij})_{1 \leq i, j \leq 3}$ is a symmetric matrix where

$$\bar{a}_{ij} = a_{ij} * \mu = \int_{\mathbb{R}^3} (\delta_{jk}|v - v'|^2 - (v_j - v'_j)(v_k - v'_k))|v - v'|^\gamma \mu(v') dv'. \quad (1.7)$$

For the hard potential case, the existence, uniqueness of the solution to Cauchy problem for the spatially homogeneous Landau equation has already been treated in [6],[18] under rather weak assumption on the initial datum. Moreover, they prove the smoothness of the solution in $C^\infty([0, +\infty[; \mathcal{S}(\mathbb{R}^3))$. In [4], Chen-Li-Xu improve this smoothing property and prove that the solution is in fact analytic for any $t > 0$ (See [2, 3] for the Gevrey regularity).

In the Maxwellian molecules case, in [10], Lerner, Morimoto, Pravda-Starov and Xu study the spatially homogeneous non-cutoff Boltzmann equation and Landau equation in a close-to-equilibrium framework and show that the solution enjoys the Gelfand-Shilov smoothing effect (see also [15], [12] and [11]). This implies that the nonlinear spatial homogeneous Landau equation has the same smoothing effect properties as the classic heat equation or harmonic oscillators heat equation. In addition, starting from a L^2 initial datum at $t = 0$, the solution of Cauchy problem is spatial analytic for any $t > 0$ and the analytic radius is $c_0 t^{\frac{1}{2}}$. In the non-Maxwellian case, we can't use the Fourier transformation and spectral decomposition as in [10, 11, 12, 15]. Recently, Li and Xu in [13] proved the analytic smoothing effect of the solution to the nonlinear Landau equation with hard potentials and Maxwellian molecules case, that is $\gamma \geq 0$.

In this work, we study the Cauchy problem of the linear Landau equation with soft potential $-3 < \gamma < 0$, we show the solution enjoys an analytic smoothing effect with the analytic radius $c_0 t^{\frac{1}{2}}$. The main theorem is in the following.

Theorem 1.1. *For soft potential $-3 < \gamma < 0$ and for any $T > 0$, the initial datum $f_0 \in L^2(\mathbb{R}^3)$, the Cauchy problem (1.5) admits a unique weak solution*

$$f \in L^\infty([0, T]; L^2(\mathbb{R}^3)).$$

Moreover, for any $\alpha \in \mathbb{N}^3$, $\tilde{t} = \min\{t, 1\}$, we have,

$$\| \tilde{t}^{\frac{|\alpha|}{2}} \langle v \rangle^{\frac{\gamma|\alpha|}{2}} \partial^\alpha f(t) \|_{L^2(\mathbb{R}^3)} \leq C^{|\alpha|+1} \alpha!, \quad t \in [0, T].$$

Remark 1.2. *Equivalently, for any $m \in \mathbb{N}$, we have, for soft potential $-3 < \gamma < 0$,*

$$\begin{aligned} & \| \tilde{t}^{\frac{m}{2}} \langle v \rangle^{\frac{\gamma m}{2}} \nabla^m f(t) \|_{L^2(\mathbb{R}^3)}^2 \\ &= \sum_{|\alpha|=m} \frac{(m!)^2}{(\alpha!)^2} \| \tilde{t}^{\frac{|\alpha|}{2}} \langle v \rangle^{\frac{\gamma|\alpha|}{2}} \partial^\alpha f(t) \|_{L^2(\mathbb{R}^3)}^2 \leq (3C)^{2m+2} (m!)^2, \quad t \in [0, T]. \end{aligned}$$

This paper is arranged as follows : We prove the ultra-analytic for the coefficient of the Landau operator in Section 2. In Section 3, we deal with estimation of the commutators and prove the coercivity property of the linear Landau operator. In Section 4, we study the Cauchy problem for linear Landau equation, and show that the existence and uniqueness properties of the weak solution. The analytic smoothing effect of the weak solution for the linear Landau equation with soft potential will be proved in Section 5. In the Appendix 6, we introduce the Hermite operator and related results.

2. ULTRA-ANALYTIC FOR THE COEFFICIENT OF LANDAU OPERATOR

For $\gamma \in \mathbb{R}$, denote

$$\|\langle v \rangle^\gamma f\|_{L^2(\mathbb{R}^3)} = \|f\|_{2,\gamma}, \quad \|\langle v \rangle^\gamma f\|_{L^\infty(\mathbb{R}^3)} = \|f\|_{\infty,\gamma}$$

where we use the notations $\langle v \rangle = (1 + |v|^2)^{1/2}$.

In addition, for the matrix A defined in (1.7), we denote

$$\|u\|_A^2 = \sum_{j,k=1}^3 \int_{\mathbb{R}^3} (\bar{a}_{jk} \partial_j u \partial_k u + \frac{1}{4} \bar{a}_{jk} v_j v_k u^2) dv, \quad (2.1)$$

and weighted norm, for $\theta \in \mathbb{R}$,

$$\|u\|_{A,\theta}^2 = \sum_{j,k=1}^3 \int_{\mathbb{R}^3} \langle v \rangle^{2\theta} (\bar{a}_{jk} \partial_j u \partial_k u + \frac{1}{4} \bar{a}_{jk} v_j v_k u^2) dv. \quad (2.2)$$

From formula (21) of Corollary 1 in [9], for any $\theta \in \mathbb{R}$, there exist $C_1 > 0$, such that

$$\|f\|_{A,\theta}^2 \geq C_1 (\|P_v \nabla f\|_{2,\frac{\gamma}{2}+\theta}^2 + \|(I - P_v) \nabla f\|_{2,1+\frac{\gamma}{2}+\theta}^2 + \|f\|_{2,1+\frac{\gamma}{2}+\theta}^2). \quad (2.3)$$

where for any vector-valued function $G(v) = (G_1, G_2, G_3)$, we define the projection to the vector $v = (v_1, v_2, v_3)$ as

$$P_v G_i = \sum_{j=1}^3 G_j v_j \frac{v_i}{|v|^2}, \quad 1 \leq i \leq 3.$$

Notice that $\nabla f = P_v \nabla f + (I - P_v) \nabla f$, we have

$$\|f\|_{A,\theta} \geq C_1 (\|\nabla f\|_{2,\frac{\gamma}{2}+\theta} + \|f\|_{2,1+\frac{\gamma}{2}+\theta}). \quad (2.4)$$

We can also refer to [5] and their references. Remark that the weighted of f and ∇f are different in the norm L_A^2 ,

Firstly, for any $\gamma > -3$ and $\delta > 0$, we have

$$\int_{\mathbb{R}^3} |v - w|^\gamma e^{-\delta|w|^2} dw \lesssim \langle v \rangle^\gamma. \quad (2.5)$$

Which imply

$$|\bar{a}_{ij}(v)| \lesssim \langle v \rangle^{\gamma+2}. \quad (2.6)$$

In the following, we prove that the coefficients of linear Landau operator are ultra-analytic.

Lemma 2.1. For any $\beta \in \mathbb{N}^3$ with $|\beta| \geq 1$ and \bar{a}_{ij} was defined in (1.7) with $-3 < \gamma < 0$, then we have,

$$|\partial^\beta \bar{a}_{ij}(v)| \lesssim \langle v \rangle^{\gamma+1} \sqrt{\beta!}. \quad (2.7)$$

Moreover, for any $\beta \in \mathbb{N}^3$,

$$\begin{aligned} |\partial^\beta (\sum_{i,j=1}^3 \partial_i a_{ij} * (v_j \mu))| &\lesssim \langle v \rangle^{\gamma+1} (|\beta| + 1) \sqrt{\beta!}, \\ |\partial^\beta (\sum_{i,j=1}^3 \bar{a}_{ij} v_i v_j)| &\lesssim \langle v \rangle^{\gamma+1} (|\beta| + 1) \sqrt{\beta!}. \end{aligned} \quad (2.8)$$

Proof. For $\beta \in \mathbb{N}^3$ with $|\beta| \geq 1$, without loss of generality, we set

$$\beta_1 = \max(\beta_1, \beta_2, \beta_3).$$

Notice that $\bar{a}_{ij} = a_{ij} * \mu$, then

$$\partial^\beta \bar{a}_{ij} = \partial^\beta [a_{ij} * \mu] = \partial_1 a_{ij} * \partial^{\beta - e_1} \mu$$

Direct calculation shows that, for any $1 \leq i \leq 3$,

$$\partial_i (\sqrt{\mu} f) = \sqrt{\mu} (\partial_i - \frac{v_i}{2}) f = -\sqrt{\mu} A_{+,i} f. \quad (2.9)$$

For more details of the operators $A_{\pm,i}$, we can refer to the Appendix 6. By using the fact

$$|\partial_1 a_{ij}(v)| \lesssim |v|^{\gamma+1},$$

it follows from Cauchy-Schwartz's inequality and (2.5) that

$$\begin{aligned} |\partial^\beta \bar{a}_{ij}(v)| &\leq \int_{\mathbb{R}^3} |\partial_1 a_{ij}(v-w) (\sqrt{\mu} A_{+,1}^{\beta_1-1} A_{+,2}^{\beta_2} A_{+,3}^{\beta_3} \sqrt{\mu})(w)| dw \\ &\lesssim \left(\int_{\mathbb{R}^3} |v-w|^{\frac{3}{2}(\gamma+1)} |\sqrt{\mu}(w)| dw \right)^{\frac{2}{3}} \|A_{+,1}^{\beta_1-1} A_{+,2}^{\beta_2} A_{+,3}^{\beta_3} \sqrt{\mu}\|_{L^3(\mathbb{R}^3)} \\ &\lesssim (1+|v|)^{\gamma+1} \|A_{+,1}^{\beta_1-1} A_{+,2}^{\beta_2} A_{+,3}^{\beta_3} \sqrt{\mu}\|_{L^3(\mathbb{R}^3)} \end{aligned}$$

where we use the fact

$$\frac{3(\gamma+1)}{2} > -3, \quad \forall -3 < \gamma < 0.$$

Due to $\sqrt{\mu}(v) = \Psi_0(v)$, we can deduce from (6.3) that

$$A_{+,1}^{\beta_1-1} A_{+,2}^{\beta_2} A_{+,3}^{\beta_3} \sqrt{\mu} = \sqrt{(\beta_1-1)! \beta_2! \beta_3!} \Psi_{\beta-e_1}, \quad (2.10)$$

where $\{\Psi_\alpha\}_{\alpha \in \mathbb{N}^3}$ is the orthonormal basis in $L^2(\mathbb{R}^3)$. The Hölder's inequality and Poincaré's inequality implies

$$\|f\|_{L^3(\mathbb{R}^3)}^2 \leq \|f\|_{L^2(\mathbb{R}^3)} \|f\|_{L^6(\mathbb{R}^3)} \lesssim \|f\|_{L^2(\mathbb{R}^3)} \|\nabla f\|_{L^2(\mathbb{R}^3)},$$

along with the equalities (6.2) and (6.3) shows that,

$$\begin{aligned} |\partial^\beta \bar{a}_{ij}(v)| &\lesssim \langle v \rangle^{\gamma+1} \|\nabla \Psi_{\beta-e_1}\|_{L^2(\mathbb{R}^3)}^{\frac{1}{2}} \\ &\lesssim \langle v \rangle^{\gamma+1} \sum_{k=1}^3 \sqrt{(\beta - e_1 + e_k)!} \end{aligned}$$

Consider that $\beta_1 = \max(\beta_1, \beta_2, \beta_3)$, we get that

$$|\partial^\beta \bar{a}_{ij}(v)| \lesssim \langle v \rangle^{\gamma+1} \sqrt{\beta!}.$$

For the estimate of the remaining inequalities (2.8), an integration by parts inside the convolution show that,

$$\begin{aligned} \sum_{i,j=1}^3 \partial_i a_{ij} * (v_j \mu) &= - \sum_{i,j=1}^3 \partial_i a_{ij} * (\sqrt{\mu} A_{+,j} \sqrt{\mu}), \\ \sum_{i,j=1}^3 a_{ij} * (v_i v_j \mu) &= \sum_{i,j=1}^3 a_{ij} * (\sqrt{\mu} (A_{+,i} A_{+,j} \sqrt{\mu} - \delta_{i,j} A_{+,j} \sqrt{\mu})) \\ &= - \sum_{i,j=1}^3 \partial_j a_{ij} * (\sqrt{\mu} (A_{+,i} \sqrt{\mu} - \delta_{i,j} \sqrt{\mu})). \end{aligned}$$

By using the fact

$$|\partial_i a_{ij}(v)| \lesssim |v|^{\gamma+1}, \quad |\partial_j a_{ij}(v)| \lesssim |v|^{\gamma+1}.$$

A calculation similar to that as above, it follows that

$$\begin{aligned} &|\partial^\beta (\sum_{i,j=1}^3 \partial_i a_{ij} * (v_j \mu))| \\ &\lesssim \sum_{j=1}^3 \left(\int_{\mathbb{R}^3} |v-w|^{\frac{3}{2}(\gamma+1)} e^{-\frac{|w|^2}{2}} dw \right)^{\frac{1}{2}} \sqrt{(\beta + e_j)!} \|\Psi_{\beta+e_j}\|_{L^3(\mathbb{R}^3)} \\ &\lesssim \langle v \rangle^{\gamma+1} \sum_{j=1}^3 \sqrt{(\beta + e_j)!} \|\Psi_{\beta+e_j}\|_{L^3(\mathbb{R}^3)} \\ &\lesssim \langle v \rangle^{\gamma+1} \sum_{j=1}^3 \sqrt{(\beta + e_j)!} \|\nabla \Psi_{\beta+e_j}\|_{L^2(\mathbb{R}^3)}^{\frac{1}{2}} \\ &\lesssim \langle v \rangle^{\gamma+1} \sum_{k,j=1}^3 \sqrt{(\beta + e_j + e_k)!} \\ &\leq \langle v \rangle^{\gamma+1} (|\beta| + 1) \sqrt{\beta!}. \end{aligned}$$

The same estimate holds true for the last term, such that

$$|\partial^\beta (\sum_{i,j=1}^3 \bar{a}_{ij} v_i v_j)| \lesssim \langle v \rangle^{\gamma+1} (|\beta| + 1) \sqrt{\beta!}.$$

We end the proof of Lemma 2.1. \square

In order to prove the coercivity of the linear Landau operator, we need one more estimate to control the weighted \bar{a}_{ij} .

Lemma 2.2. *For $f, g \in \mathcal{S}(\mathbb{R}^3)$, for any $\beta \in \mathbb{N}^3$ and $\theta \in \mathbb{R}$, we have*

$$\left| \sum_{i,j=1}^3 \langle v \rangle^{2\theta} \partial^\beta \bar{a}_{ij} \partial_j f, \partial_i g \right| \lesssim \sqrt{\beta!} \|f\|_{A,\theta} \|g\|_{A,\theta}. \quad (2.11)$$

where \bar{a}_{ij} was defined in (1.7) with $-3 < \gamma < 0$.

Proof. In fact, the inner product

$$\langle \langle v \rangle^{2\theta} \partial^\beta \bar{a}_{ij} \partial_j f, \partial_i g \rangle = \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \langle v \rangle^{2\theta} a_{ij}(v-w) \partial^\beta \mu(w) \partial_j f(v) \partial_i g(v) dw dv.$$

We decompose the integration region $[v, w] \in \mathbb{R}^3 \times \mathbb{R}^3$ into three parts:

$$\{|v| \leq 1\}, \quad \{2|w| \geq |v|, |v| \geq 1\}, \quad \text{and} \quad \{2|w| \leq |v|, |v| \geq 1\}.$$

For the first part $\{|v| \leq 1\}$, by Lemma 2.1, we have

$$\begin{aligned} & \left| \sum_{i,j=1}^3 \iint_{\{|v| \leq 1\} \times \mathbb{R}^3} \langle v \rangle^{2\theta} a_{ij}(v-w) \partial^\beta \mu(w) \partial_j f(v) \partial_i g(v) dw dv \right| \\ & \lesssim \sum_{i,j=1}^3 \int_{|v| \leq 1} \langle v \rangle^{\gamma+2} \sqrt{\beta!} \langle v \rangle^{2\theta} |\partial_j f \partial_i g| dv \\ & \leq \sqrt{\beta!} \|\langle v \rangle^{\frac{\gamma}{2}} \nabla f\|_{2,\theta} \|\langle v \rangle^{\frac{\gamma}{2}} \nabla g\|_{2,\theta} \lesssim \sqrt{\beta!} \|f\|_{A,\theta} \|g\|_{A,\theta}. \end{aligned}$$

For the second part $\{2|w| \geq |v|, |v| \geq 1\}$, we have

$$e^{-\frac{|w|^2}{4}} \leq e^{-\frac{|w|^2}{8}} e^{-\frac{|v|^2}{32}}.$$

Similar to the proof as Lemma 2.1, one can verify that

$$\begin{aligned} & \left| \sum_{i,j=1}^3 \iint_{\{2|w| \geq |v|, |v| \geq 1\}} \langle v \rangle^{2\theta} a_{ij}(v-w) \sqrt{\mu(w)} \Psi_\beta(w) \partial_j f(v) \partial_i g(v) dw dv \right| \\ & \lesssim \sum_{i,j=1}^3 \int_{\mathbb{R}^3} \langle v \rangle^{\gamma+2} e^{-\frac{|v|^2}{32}} \sqrt{\beta!} \langle v \rangle^{2\theta} |\partial_j f \partial_i g| dv \\ & \lesssim \sqrt{\beta!} \|f\|_{A,\theta} \|g\|_{A,\theta}. \end{aligned}$$

We finally consider the third part $\{2|w| \leq |v|, |v| \geq 1\}$. Expanding $a_{ij}(v-w)$ to get

$$a_{ij}(v-w) = a_{ij}(v) + \sum_{k=1}^3 \partial_k a_{ij}(v) w_k + \frac{1}{2} \sum_{k,l=1}^3 \left(\int_0^1 \partial_{kl} a_{ij}(v-sw) ds \right) w_k w_l.$$

Along with the fact that

$$\sum_{i=1}^3 a_{ij} v_i = 0, \quad \sum_{i,j=1}^3 \partial_k a_{ij}(v) v_i v_j = -2 \sum_{j=1}^3 a_{kj}(v) v_j = 0,$$

show immediately

$$\begin{aligned} & \sum_{i,j=1}^3 \iint_{\{2|w| \leq |v|, |v| \geq 1\}} \langle v \rangle^{2\theta} a_{ij}(v-w) \partial^\beta \mu(w) \partial_j f(v) \partial_i g(v) dw dv \\ & = \sum_{i,j=1}^3 \iint_{2|w| \leq |v|, |v| \geq 1} \langle v \rangle^{2\theta} \partial^\beta \mu(w) a_{ij}(v) [(I - P_v) \partial_j f] [(I - P_v) \partial_i g] dw dv \\ & \quad + \sum_{i,j=1}^3 \sum_{k=1}^3 \iint_{2|w| \leq |v|, |v| \geq 1} \langle v \rangle^{2\theta} w_k \partial^\beta \mu(w) \partial_k a_{ij}(v) [P_v \partial_j f] [(I - P_v) \partial_i g] dw dv \end{aligned}$$

$$\begin{aligned}
& + \sum_{i,j=1}^3 \sum_{k=1}^3 \iint_{2|w| \leq |v|, |v| \geq 1} \langle v \rangle^{2\theta} w_k \partial^\beta \mu(w) \partial_k a_{ij}(v) [(I - P_v) \partial_j f] [\partial_i g] dw dv \\
& + \frac{1}{2} \sum_{k,l=1}^3 \sum_{i,j=1}^3 \int_0^1 \iint_{2|w| \leq |v|, |v| \geq 1} \langle v \rangle^{2\theta} w_k w_l \partial^\beta \mu(w) \partial_{kl} a_{ij}(v - sw) \partial_j f \partial_i g dw dv ds.
\end{aligned}$$

Since $2|w| \leq |v|, |v| \geq 1, 0 < s < 1$, for $-3 < \gamma < 0$, we have

$$\begin{aligned}
|a_{ij}(v)| & \lesssim |v|^{\gamma+2} \lesssim \langle v \rangle^{\gamma+2}; \quad |\partial_k a_{ij}(v)| \lesssim \langle v \rangle^{\gamma+1}; \\
|\partial_{kl} a_{ij}(v - sw)| & \leq C|v - sw|^\gamma \leq C4^{-\gamma}(1 + |v|)^\gamma.
\end{aligned}$$

It follows from the inequality (2.10) and the norm equality (2.3) that

$$\begin{aligned}
& \left| \sum_{i,j=1}^3 \iint_{\{2|w| \leq |v|, |v| \geq 1\}} \langle v \rangle^{2\theta} a_{ij}(v - w) \partial^\beta \mu(w) \partial_j f(v) \partial_i g(v) dw dv \right| \\
& \lesssim \sqrt{\beta!} \|(I - P_v) \nabla f\|_{2, \frac{\gamma+2}{2} + \theta} \|(I - P_v) \nabla g\|_{2, \frac{\gamma+2}{2} + \theta} \\
& \quad + \sqrt{\beta!} \|(I - P_v) \nabla f\|_{2, \frac{\gamma+2}{2} + \theta} \|P_v \nabla g\|_{2, \frac{\gamma}{2} + \theta} \\
& \quad + \sqrt{\beta!} \|(I - P_v) \nabla f\|_{2, \frac{\gamma+2}{2} + \theta} \|\nabla g\|_{2, \frac{\gamma}{2} + \theta} + \sqrt{\beta!} \|\nabla f\|_{2, \frac{\gamma}{2} + \theta} \|\nabla g\|_{2, \frac{\gamma}{2} + \theta} \\
& \lesssim \sqrt{\beta!} \|f\|_{A, \theta} \|g\|_{A, \theta}.
\end{aligned}$$

This is the inequality (2.11). We end the proof of Lemma 2.2. \square

3. ESTIMATIONS OF COMMUTATORS

Proposition 3.1. *Let $f \in \mathcal{S}(\mathbb{R}^3)$, \mathcal{L} was defined in (1.5), for any $\alpha \in \mathbb{N}^3$ and $\theta \in \mathbb{R}$, there exist a positive constant $C_0 > 0$ which is independent on α and θ , such that,*

$$\begin{aligned}
& \left(\langle v \rangle^{2\theta} \partial^\alpha \mathcal{L} f, \partial^\alpha f \right)_{L^2(\mathbb{R}^3)} \leq -\|\partial^\alpha f\|_{A, \theta}^2 + C_0 \langle \theta \rangle \|\partial^\alpha f\|_{A, \theta} \|\partial^\alpha f\|_{2, \frac{\gamma}{2} + \theta} \\
& \quad + C_0 \sum_{\substack{|\beta| \geq 1 \\ \beta \leq \alpha}} C_\alpha^\beta \sqrt{\beta!} \|\partial^{\alpha-\beta} f\|_{A, \theta} \left(\|\partial^\alpha f\|_{A, \theta} + |\theta| \|\partial^\alpha f\|_{2, \theta + \frac{\gamma}{2}} \right) \\
& \quad + C_0 \sum_{\substack{|\beta| \geq 1 \\ \beta \leq \alpha}} C_\alpha^\beta |\beta| \sqrt{\beta!} \|\partial^{\alpha-\beta} f\|_{A, \theta} \|\partial^\alpha f\|_{2, \theta + \frac{\gamma}{2}}.
\end{aligned}$$

Remark 3.2. *Remark that, we have*

(1) *For $\alpha = 0, \theta = 0$, we have*

$$\left(\mathcal{L} f, f \right)_{L^2(\mathbb{R}^3)} \leq -\frac{1}{2} \|f\|_A^2 + C_0 \|f\|_{\frac{\gamma}{2}}^2. \quad (3.1)$$

(2) *For $|\alpha| \geq 1$,*

$$\|\partial^\alpha f\|_{2, \frac{\gamma}{2} + \theta} \lesssim \|\nabla \partial^{\alpha - e_{j_0}} f\|_{2, \frac{\gamma}{2} + \theta} \lesssim \|\partial^{\alpha - e_{j_0}} f\|_{A, \theta}, \quad (3.2)$$

where $\alpha_{j_0} = \max\{\alpha_1, \alpha_2, \alpha_3\}$.

Proof. Recalled the formula $\mathcal{L}f$ in (1.5), for the smooth function f , integrated by parts, we have

$$\begin{aligned}
- (\langle v \rangle^{2\theta} \partial^\alpha \mathcal{L}f, \partial^\alpha f)_{L^2(\mathbb{R}^3)} &= \sum_{i,j=1}^3 \int_{\mathbb{R}^3} \langle v \rangle^{2\theta} \{ \partial^\alpha (\bar{a}_{ij} \partial_j f) \} \partial^\alpha \partial_i f dv \\
&+ \sum_{i,j=1}^3 \int_{\mathbb{R}^3} \{ \partial_i \langle v \rangle^{2\theta} \} \{ \partial^\alpha (\bar{a}_{ij} \partial_j f) \} \partial^\alpha f dv \\
&+ \frac{1}{4} \sum_{i,j=1}^3 \int_{\mathbb{R}^3} \langle v \rangle^{2\theta} \{ \partial^\alpha (\bar{a}_{ij} v_i v_j f) \} \partial^\alpha f dv \\
&- \frac{1}{2} \sum_{i,j=1}^3 \int_{\mathbb{R}^3} \langle v \rangle^{2\theta} \{ \partial^\alpha \partial_i ((\bar{a}_{ij} v_j) f) \} \partial^\alpha f dv.
\end{aligned}$$

Then, using (2.2), we have,

$$- (\langle v \rangle^{2\theta} \partial^\alpha \mathcal{L}f, \partial^\alpha f)_{L^2(\mathbb{R}^3)} = \|\partial^\alpha f\|_{A,\theta}^2 + \mathbf{R}(f), \quad (3.3)$$

with

$$\begin{aligned}
\mathbf{R}(f) &= \sum_{i,j=1}^3 \int_{\mathbb{R}^3} \langle v \rangle^{2\theta} \{ [\partial^\alpha, \bar{a}_{ij}] \partial_j f \} \partial^\alpha \partial_i f dv \\
&+ \sum_{i,j=1}^3 \int_{\mathbb{R}^3} \{ \partial_i \langle v \rangle^{2\theta} \} \{ \partial^\alpha (\bar{a}_{ij} \partial_j f) \} \partial^\alpha f dv \\
&+ \frac{1}{4} \sum_{i,j=1}^3 \int_{\mathbb{R}^3} \langle v \rangle^{2\theta} \{ [\partial^\alpha, \bar{a}_{ij} v_i v_j] f \} \partial^\alpha f dv \\
&- \frac{1}{2} \sum_{i,j=1}^3 \int_{\mathbb{R}^3} \langle v \rangle^{2\theta} \{ \partial^\alpha \partial_i ((\bar{a}_{ij} v_j) f) \} \partial^\alpha f dv.
\end{aligned}$$

By using Leibniz formula,

$$\partial^\alpha (gh) = g \partial^\alpha h + \sum_{\substack{|\beta|=1 \\ \beta \leq \alpha}} C_\alpha^\beta \partial^\beta g \partial^{\alpha-\beta} h + \sum_{\substack{|\beta| \geq 2 \\ \beta \leq \alpha}} C_\alpha^\beta \partial^\beta g \partial^{\alpha-\beta} h,$$

and

$$\bar{a}_{ij} = a_{ij} * \mu, \quad \sum_j \bar{a}_{ij} v_j = \sum_j a_{ij} * (v_j \mu), \quad \sum_{i,j} \bar{a}_{ij} v_i v_j = \sum_j a_{ij} * (v_i v_j \mu),$$

it follows that

$$\mathbf{R}(f) = \mathbf{R}_0(f) + \mathbf{R}_1(f)$$

where

$$\begin{aligned}
\mathbf{R}_0(f) &= \sum_{i,j=1}^3 \int_{\mathbb{R}^3} (\partial_i \langle v \rangle^{2\theta}) (\bar{a}_{ij} \partial_j \partial^\alpha f) \partial^\alpha f dv \\
&- \frac{1}{2} \sum_{i,j=1}^3 \int_{\mathbb{R}^3} \langle v \rangle^{2\theta} (\partial_i a_{ij}) * (v_j \mu) |\partial^\alpha f|^2 dv = \mathbf{R}_{01}(f) + \mathbf{R}_{02}(f);
\end{aligned}$$

$$\begin{aligned}
\mathbf{R}_1(f) &= \sum_{\substack{|\beta| \geq 1 \\ \beta \leq \alpha}} C_\alpha^\beta \sum_{i,j=1}^3 \int_{\mathbb{R}^3} \langle v \rangle^{2\theta} (\partial^\beta \bar{a}_{ij}) (\partial^{\alpha-\beta} \partial_j f) \partial^\alpha \partial_i f dv \\
&+ \sum_{\substack{|\beta| \geq 1 \\ \beta \leq \alpha}} C_\alpha^\beta \sum_{i,j=1}^3 \int_{\mathbb{R}^3} \{ \partial_i \langle v \rangle^{2\theta} \} (\partial^\beta \bar{a}_{ij}) \{ \partial_j \partial^{\alpha-\beta} f \} \partial^\alpha f dv \\
&+ \frac{1}{4} \sum_{\substack{|\beta| \geq 1 \\ \beta \leq \alpha}} C_\alpha^\beta \sum_{i,j=1}^3 \int_{\mathbb{R}^3} \langle v \rangle^{2\theta} \{ \partial^\beta (a_{ij} * (v_i v_j \mu)) \} \partial^{\alpha-\beta} f \partial^\alpha f dv \\
&- \frac{1}{2} \sum_{\substack{|\beta| \geq 1 \\ \beta \leq \alpha}} C_\alpha^\beta \sum_{i,j=1}^3 \int_{\mathbb{R}^3} \langle v \rangle^{2\theta} (\partial_i \partial^\beta a_{ij}) * (v_j \mu) \partial^{\alpha-\beta} f \partial^\alpha f dv \\
&= \mathbf{R}_{11} + \mathbf{R}_{12} + \mathbf{R}_{13} + \mathbf{R}_{14}.
\end{aligned}$$

So that the proof of the Proposition 3.1 is reduced to the estimations of terms $\mathbf{R}_0(f)$ and $\mathbf{R}_1(f)$, which will give by the following two lemmas. \square

Lemma 3.3. *We have, for $\alpha \in \mathbb{N}^3$,*

$$|\mathbf{R}_0(f)| \lesssim \langle \theta \rangle \|\partial^\alpha f\|_{A,\theta} \|\partial^\alpha f\|_{2, \frac{\gamma}{2} + \theta}. \quad (3.4)$$

Proof. By using (2.5), we get

$$|(\partial_i a_{ij}) * (v_j \mu)| \lesssim \langle v \rangle^{\gamma+1}, \quad |(a_{ij}) * (v_j \mu)| \lesssim \langle v \rangle^{\gamma+2}$$

then (2.4) imply

$$\begin{aligned}
|\mathbf{R}_{02}(f)| &= \left| \sum_{i,j=1}^3 \int_{\mathbb{R}^3} \langle v \rangle^{2\theta} (\partial_i a_{ij}) * (v_j \mu) |\partial^\alpha f|^2 dv \right| \\
&\lesssim \int_{\mathbb{R}^3} \langle v \rangle^{\gamma+1+2\theta} |\partial^\alpha f|^2 dv \\
&\lesssim \|\partial^\alpha f\|_{2, \frac{\gamma}{2} + \theta} \|\partial^\alpha f\|_{2, \frac{\gamma}{2} + 1 + \theta} \lesssim \|\partial^\alpha f\|_{2, \frac{\gamma}{2} + \theta} \|\partial^\alpha f\|_{A,\theta}.
\end{aligned}$$

For the term \mathbf{R}_{01} , we use

$$\partial_i \langle v \rangle^{2\theta} = 2\theta \langle v \rangle^{2\theta-2} v_i,$$

then,

$$\begin{aligned}
\mathbf{R}_{01}(f) &= \sum_{i,j=1}^3 \int_{\mathbb{R}^3} (\partial_i \langle v \rangle^{2\theta}) (\bar{a}_{ij} \partial_j \partial^\alpha f) \partial^\alpha f dv \\
&= 2\theta \sum_{i,j=1}^3 \int_{\mathbb{R}^3} \langle v \rangle^{2\theta-2} (\bar{a}_{ij} v_i \partial_j \partial^\alpha f) \partial^\alpha f dv \\
&= 2\theta \sum_{i,j=1}^3 \int_{\mathbb{R}^3} \langle v \rangle^{2\theta-2} (a_{ij} * (v_i \mu) \partial_j \partial^\alpha f) \partial^\alpha f dv.
\end{aligned}$$

It follows that

$$|\mathbf{R}_{01}(f)| \leq 2|\theta| \sum_{j=1}^3 \int_{\mathbb{R}^3} \langle v \rangle^{2\theta+\gamma} |\partial_j \partial^\alpha f| |\partial^\alpha f| dv$$

$$\begin{aligned}
&\lesssim |\theta| \sum_{j=1}^3 \|\partial_j \partial^\alpha f\|_{2, \frac{\gamma}{2} + \theta} \|\partial^\alpha f\|_{2, \frac{\gamma}{2} + \theta} \\
&\lesssim |\theta| \|\partial^\alpha f\|_{A, \theta} \|\partial^\alpha f\|_{2, \frac{\gamma}{2} + \theta}.
\end{aligned}$$

Which give the estimate of $\mathbf{R}_0(f)$. \square

Lemma 3.4. *We have, for $\alpha \in \mathbb{N}^3, |\alpha| \geq 1$,*

$$|\mathbf{R}_1(f)| \lesssim \sum_{\substack{|\beta|=1 \\ \beta \leq \alpha}} C_\alpha^\beta \|\partial^{\alpha-\beta} f\|_{A, \theta} \left(\|\partial^\alpha f\|_{A, \theta} + |\theta| \|\partial^\alpha f\|_{2, \theta + \frac{\gamma}{2}} \right). \quad (3.5)$$

Proof. Now we estimate \mathbf{R}_1 . For the term $\mathbf{R}_{11}(f)$, by using the inequality (2.11) in Lemma 2.2 directly, we obtain that

$$\begin{aligned}
|\mathbf{R}_{11}(f)| &= \left| \sum_{\substack{|\beta| \geq 1 \\ \beta \leq \alpha}} C_\alpha^\beta \sum_{i,j=1}^3 \int_{\mathbb{R}^3} \langle v \rangle^{2\theta} (\partial^\beta \bar{a}_{ij}) (\partial^{\alpha-\beta} \partial_j f) \partial^\alpha \partial_i f dv \right| \\
&\lesssim \sum_{\substack{|\beta| \geq 1 \\ \beta \leq \alpha}} C_\alpha^\beta \sqrt{|\beta|} \|\partial^{\alpha-\beta} f\|_{A, \theta} \|\partial^\alpha f\|_{A, \theta}.
\end{aligned}$$

For the two terms $\mathbf{R}_{13}(f), \mathbf{R}_{14}(f)$, we can deduce from the inequality (2.8) in Lemma 2.1 and (2.4) that

$$\begin{aligned}
|\mathbf{R}_{13}(f)| + |\mathbf{R}_{14}(f)| &= \left| \frac{1}{4} \sum_{\substack{|\beta| \geq 1 \\ \beta \leq \alpha}} C_\alpha^\beta \sum_{i,j=1}^3 \int_{\mathbb{R}^3} \langle v \rangle^{2\theta} \{ \partial^\beta (a_{ij} * (v_i v_j \mu)) \} \partial^{\alpha-\beta} f \partial^\alpha f dv \right| \\
&\quad + \left| \frac{1}{2} \sum_{\substack{|\beta| \geq 1 \\ \beta \leq \alpha}} C_\alpha^\beta \sum_{i,j=1}^3 \int_{\mathbb{R}^3} \langle v \rangle^{2\theta} (\partial_i \partial^\beta a_{ij}) * (v_j \mu) \partial^{\alpha-\beta} f \partial^\alpha f dv \right| \\
&\lesssim \sum_{\substack{|\beta| \geq 1 \\ \beta \leq \alpha}} C_\alpha^\beta \int_{\mathbb{R}^3} \langle v \rangle^{2\theta + \gamma + 1} |\beta| \sqrt{|\beta|} \|\partial^{\alpha-\beta} f\| \|\partial^\alpha f\| dv \\
&\lesssim \sum_{\substack{|\beta| \geq 1 \\ \beta \leq \alpha}} C_\alpha^\beta |\beta| \sqrt{|\beta|} \|\partial^{\alpha-\beta} f\|_{2, \theta + \frac{\gamma+2}{2}} \|\partial^\alpha f\|_{2, \theta + \frac{\gamma}{2}} \\
&\lesssim \sum_{\substack{|\beta| \geq 1 \\ \beta \leq \alpha}} C_\alpha^\beta |\beta| \sqrt{|\beta|} \|\partial^{\alpha-\beta} f\|_{A, \theta} \|\partial^\alpha f\|_{2, \theta + \frac{\gamma}{2}}.
\end{aligned}$$

For the term $\mathbf{R}_{12}(f)$, for $|\beta| \geq 1$, it follows from (2.7) in Lemma 2.1 that

$$|\partial_i \langle v \rangle^{2\theta} \partial^\beta \bar{a}_{ij}| \lesssim |\theta| \langle v \rangle^{2\theta-1} \langle v \rangle^{\gamma+1} \sqrt{|\beta|} \leq |\theta| \langle v \rangle^{2\theta+\gamma} \sqrt{|\beta|}.$$

Then we have

$$|\mathbf{R}_{12}(f)| = \left| \sum_{\substack{|\beta| \geq 1 \\ \beta \leq \alpha}} C_\alpha^\beta \sum_{i,j=1}^3 \int_{\mathbb{R}^3} \{ \partial_i \langle v \rangle^{2\theta} \} (\partial^\beta \bar{a}_{ij}) \{ \partial_j \partial^{\alpha-\beta} f \} \partial^\alpha f dv \right|$$

$$\begin{aligned}
&\lesssim |\theta| \sum_{\substack{|\beta| \geq 1 \\ \beta \leq \alpha}} C_\alpha^\beta \sqrt{\beta!} \sum_{j=1}^3 \|\partial^{\alpha-\beta} \partial_j f\|_{2, \frac{\gamma}{2} + \theta} \|\partial^\alpha f\|_{2, \frac{\gamma}{2} + \theta} \\
&\lesssim |\theta| \sum_{\substack{|\beta| \geq 1 \\ \beta \leq \alpha}} C_\alpha^\beta \sqrt{\beta!} \|\partial^{\alpha-\beta} f\|_{A, \theta} \|\partial^\alpha f\|_{2, \frac{\gamma}{2} + \theta}.
\end{aligned}$$

We conclude that

$$\begin{aligned}
|\mathbf{R}_1(f)| &\lesssim \sum_{\substack{|\beta| \geq 1 \\ \beta \leq \alpha}} C_\alpha^\beta \sqrt{\beta!} \|\partial^{\alpha-\beta} f\|_{A, \theta} \|\partial^\alpha f\|_{A, \theta} \\
&\quad + |\theta| \sum_{\substack{|\beta| \geq 1 \\ \beta \leq \alpha}} C_\alpha^\beta \sqrt{\beta!} \|\partial^{\alpha-\beta} f\|_{A, \theta} \|\partial^\alpha f\|_{2, \theta + \frac{\gamma}{2}} \\
&\quad + \sum_{\substack{|\beta| \geq 1 \\ \beta \leq \alpha}} C_\alpha^\beta |\beta| \sqrt{\beta!} \|\partial^{\alpha-\beta} f\|_{A, \theta} \|\partial^\alpha f\|_{2, \theta + \frac{\gamma}{2}}.
\end{aligned}$$

□

Substituting the estimates of $\mathbf{R}_0(f)$, $\mathbf{R}_1(f)$ into the decomposition (3.3), we can end the proof of Proposition 3.1.

4. EXISTENCE AND UNIQUENESS OF LINEAR LANDAU EQUATION.

Proposition 4.1. *For $-3 < \gamma < 0$, $T > 0$, $f_0 \in L^2(\mathbb{R}^3)$, the Cauchy problem (1.5) admits a unique weak solution*

$$f \in L^\infty([0, T]; L^2(\mathbb{R}^3))$$

satisfying

$$\|f(t)\|_{L^2(\mathbb{R}^3)} \leq 2e^{2C_0 T} \|f_0\|_{L^2(\mathbb{R}^3)}$$

where C_0 was defined in Proposition 3.1.

Proof. The existence of the weak solution is similar to that in [1, 14, 9]. We consider the operator

$$\mathcal{P} = -\partial_t + \mathcal{L}.$$

For any $\varphi \in C^\infty([0, T]; \mathcal{S}(\mathbb{R}^3))$ with $\varphi(T) = 0$, it follows from (3.1) that

$$\begin{aligned}
(\varphi(t), \mathcal{P}^* \varphi(t))_{L^2(\mathbb{R}^3)} &= (\partial_t \varphi, \varphi)_{L^2(\mathbb{R}^3)} + (\mathcal{L} \varphi, \varphi)_{L^2(\mathbb{R}^3)} \\
&\leq \frac{1}{2} \frac{d}{dt} \|\varphi\|_{L^2(\mathbb{R}^3)}^2 - \|\varphi\|_A^2 + C_0 \|\varphi\|_{2, \frac{\gamma}{2}}^2.
\end{aligned}$$

Since $\gamma < 0$, we have

$$-\frac{1}{2} \frac{d}{dt} \|\varphi\|_{L^2(\mathbb{R}^3)}^2 + \|\varphi\|_A^2 \leq (\varphi(t), \mathcal{P}^* \varphi(t))_{L^2(\mathbb{R}^3)} + C_0 \|\varphi\|_{L^2(\mathbb{R}^3)}^2.$$

Which implies that

$$-\frac{d}{dt} \left(e^{2C_0 t} \|\varphi\|_{L^2(\mathbb{R}^3)}^2 \right) + e^{2C_0 t} \|\varphi\|_A^2 \leq 2e^{2C_0 t} \|\varphi\|_{L^2(\mathbb{R}^3)} \|\mathcal{P}^* \varphi(t)\|_{L^2(\mathbb{R}^3)},$$

Then one can verify that

$$\sup_{0 \leq t \leq T} \|\varphi(t)\|_{L^2(\mathbb{R}^3)}^2 \leq 2e^{2C_0 T} \int_0^T \|\mathcal{P}^* \varphi(s)\|_{L^2(\mathbb{R}^3)} ds. \quad (4.1)$$

In the following, we set the vector subspace,

$$\mathbb{U} = \{u = \mathcal{P}\varphi : \varphi \in C^\infty([0, T], \mathcal{S}(\mathbb{R}_{x,v}^6)), \varphi(T) = 0\} \subset L^1([0, T], L^2(\mathbb{R}^3)).$$

Since $f_0 \in L^2(\mathbb{R}^3)$, we define the linear functional

$$\begin{aligned} \mathcal{G} : \mathbb{U} &\rightarrow \mathbb{R} \\ h &= \mathcal{P}^*\varphi \mapsto (f_0, \varphi(0))_{L^2(\mathbb{R}^3)}, \end{aligned}$$

where $\varphi \in C^\infty([0, T], \mathcal{S}(\mathbb{R}^3))$ with $\varphi(T) = 0$. From (4.1), the operator \mathcal{P}^* is injective. Therefore the linear functional \mathcal{G} is well-defined and moreover, we obtain

$$\begin{aligned} |\mathcal{G}(h)| &\leq \|f_0\|_{L^2} \|\varphi(0)\|_{L^2(\mathbb{R}^3)} \leq \|f_0\|_{L^2} \|\varphi\|_{L^\infty([0, T]; L^2(\mathbb{R}^3))} \\ &\leq 2e^{2C_0T} \|f_0\|_{L^2} \|\mathcal{P}^*\varphi\|_{L^1([0, T], L^2(\mathbb{R}^3))} = 2e^{2C_0T} \|f_0\|_{L^2} \|h\|_{L^1([0, T]; L^2(\mathbb{R}^3))}. \end{aligned}$$

This shows that \mathcal{G} is a continuous linear form on $(\mathbb{U}, \|\cdot\|_{L^1([0, T], L^2(\mathbb{R}^3))})$. By using the Hahn-Banach theorem, \mathcal{G} can be extended as a continuous linear form on $L^1([0, T]; L^2(\mathbb{R}^3))$ with a norm smaller than $2e^{2C_0T} \|f_0\|_{L^2(\mathbb{R}^3)}$. It follows the Riesz representation theorem that there exists a unique $f \in L^\infty([0, T]; L^2(\mathbb{R}^3))$ satisfying

$$\|f\|_{L^\infty([0, T]; L^2(\mathbb{R}^3))} \leq 2e^{2C_0T} \|f_0\|_{L^2(\mathbb{R}^3)},$$

such that

$$\forall h \in L^1([0, T]; L^2(\mathbb{R}^3)), \quad \mathcal{G}(h) = \int_0^T (f(t), h(t))_{L^2(\mathbb{R}^3)} dt.$$

Which implies that for all $\varphi \in C_0^\infty((0, T), \mathcal{S}(\mathbb{R}^3))$,

$$\mathcal{G}(\mathcal{P}^*\varphi) = \int_0^T (f(t), \mathcal{P}^*\varphi(t))_{L^2(\mathbb{R}^3)} dt = (f_0, \varphi(0))_{L^2(\mathbb{R}^3)}.$$

Therefore, $f \in L^\infty([0, T]; L^2(\mathbb{R}^3))$ is a weak solution of the Cauchy problem (1.5). Let $\tilde{f} \in L^\infty([0, T]; L^2(\mathbb{R}^3))$ be another weak solution of the Cauchy problem (1.5) satisfying

$$\|\tilde{f}\|_{L^\infty([0, T]; L^2(\mathbb{R}^3))} \leq 2e^{2C_0T} \|f_0\|_{L^2(\mathbb{R}^3)},$$

Set $w(t) = f(t) - \tilde{f}(t)$, we have that for all $\varphi \in C_0^\infty((0, T), \mathcal{S}(\mathbb{R}^3))$,

$$\int_0^T (w(t), \mathcal{P}^*\varphi(t))_{L^2(\mathbb{R}^3)} dt = 0.$$

Which shows that $w(t) = 0$ in $L^\infty([0, T]; L^2(\mathbb{R}^3))$. The proof of Proposition 4.1 is completed. \square

5. ANALYTIC SMOOTHING EFFECT FOR LINEAR LANDAU EQUATION

Let f be the solution of Cauchy problem (1.5), that is,

$$\partial_t f = \mathcal{L}f, \quad f|_{t=0} = f_0,$$

then similar to the estimate of \mathbf{R}_0 in Proposition 3.1 with $\theta = 0$, we have

$$\begin{aligned} \frac{d}{dt} \|f\|_{L^2(\mathbb{R}^3)}^2 &= 2(\partial_t f, f)_{L^2(\mathbb{R}^3)} = 2(\mathcal{L}f, f)_{L^2(\mathbb{R}^3)} \\ &\leq -2\|f\|_A^2 + 2C_0\|f\|_{2, \frac{3}{2}}^2. \end{aligned}$$

This implies that (since $\gamma < 0$),

$$\frac{d}{dt} \|f\|_{L^2(\mathbb{R}^3)}^2 + 2\|f\|_A^2 \leq 2C_0 \|f\|_{L^2(\mathbb{R}^3)}^2.$$

By using the Gronwall inequality, for any $T > 0$ and $0 < t < T$, we have

$$\|f(t)\|_{L^2(\mathbb{R}^3)}^2 + \int_0^t \|f(s)\|_A^2 ds \leq e^{2C_0 T} \|f_0\|_{L^2(\mathbb{R}^3)}^2. \quad (5.1)$$

Let $|\alpha| = 1$, it follows from Proposition 3.1 with $\theta = \gamma$ that

$$\begin{aligned} \frac{d}{dt} \|t^{\frac{1}{2}} \langle v \rangle^{\frac{\gamma}{2}} \partial^\alpha f\|_{L^2(\mathbb{R}^3)}^2 &= 2\langle \langle v \rangle^\gamma \partial^\alpha \partial_t f, t \partial^\alpha f \rangle + \|\langle v \rangle^{\frac{\gamma}{2}} \partial^\alpha f\|_{L^2(\mathbb{R}^3)}^2 \\ &= 2\langle \langle v \rangle^\gamma \partial^\alpha \mathcal{L}f, t \partial^\alpha f \rangle_{L^2(\mathbb{R}^3)} + \|\langle v \rangle^{\frac{\gamma}{2}} \partial^\alpha f\|_{L^2(\mathbb{R}^3)}^2 \\ &\leq -2\|t^{\frac{1}{2}} \partial^\alpha f\|_{A, \frac{\gamma}{2}}^2 + 2C_0 t \|f\|_{A, \frac{\gamma}{2}} \|\partial^\alpha f\|_{A, \frac{\gamma}{2}} + \|\langle v \rangle^{\frac{\gamma}{2}} \partial^\alpha f\|_{L^2(\mathbb{R}^3)}^2. \end{aligned} \quad (5.2)$$

By using the inequality (2.4), one can verify that, for any $0 < t < T$ and $\gamma < 0$,

$$\|f\|_{A, \frac{\gamma}{2}} \leq \|f\|_A, \quad \|\langle v \rangle^{\frac{\gamma}{2}} \partial^\alpha f\|_{L^2(\mathbb{R}^3)}^2 \leq \frac{1}{C_1} \|f\|_A^2.$$

Substituting back to the estimate (5.2), we have

$$\frac{d}{dt} \|t^{\frac{1}{2}} \partial^\alpha f\|_{2, \frac{\gamma}{2}}^2 + \|t^{\frac{1}{2}} \partial^\alpha f\|_{A, \frac{\gamma}{2}}^2 \leq \left(C_0^2 t + \frac{1}{C_1} \right) \|f\|_A^2.$$

Integral on $[0, t]$, and using (5.1), one can verify that, for $|\alpha| = 1$,

$$\|t^{\frac{1}{2}} \partial^\alpha f\|_{2, \frac{\gamma}{2}}^2 + \int_0^t \|\tau^{\frac{1}{2}} \partial^\alpha u\|_A^2 d\tau \leq \left(C_0^2 T + \frac{1}{C_1} \right) e^{2C_0 T} \|f_0\|_{L^2(\mathbb{R}^3)}^2 \leq C^4. \quad (5.3)$$

Proposition 5.1. *For any $m \in \mathbb{N}$ and $\alpha \in \mathbb{N}^3$, $|\alpha| = m$, we have, for $0 < t \leq 1$,*

$$\|t^{\frac{|\alpha|}{2}} \partial^\alpha f\|_{2, \frac{\gamma|\alpha|}{2}}^2 + \int_0^t \|\tau^{\frac{|\alpha|}{2}} \partial^\alpha f\|_{A, \frac{\gamma|\alpha|}{2}}^2 d\tau \leq C^{2|\alpha|+2} (\alpha!)^2. \quad (5.4)$$

where C is independent on α .

This Proposition imply Theorem 1.1.

Proof. In fact, we have proved that the assumption (5.4) holds true for $m = 0, 1$ by (5.1) and (5.3).

Now assume that the assumption (5.4) holds true for $|\alpha| \leq m - 1$, that means, for any $|\alpha| \leq m - 1$, for $0 < t \leq 1$,

$$\|t^{\frac{|\alpha|}{2}} \partial^\alpha f\|_{2, \frac{\gamma|\alpha|}{2}}^2 + \int_0^t \|\tau^{\frac{|\alpha|}{2}} \partial^\alpha f\|_{A, \frac{\gamma|\alpha|}{2}}^2 d\tau \leq C^{2|\alpha|+2} (\alpha!)^2. \quad (5.5)$$

We intend to prove the validity of (5.4) for m . Firstly

$$\begin{aligned} \frac{d}{dt} \|t^{\frac{|\alpha|}{2}} \langle v \rangle^{\frac{\gamma|\alpha|}{2}} \partial^\alpha f\|_{L^2(\mathbb{R}^3)}^2 &= 2t^{|\alpha|} \langle \langle v \rangle^{\gamma|\alpha|} \partial^\alpha \partial_t f, \partial^\alpha f \rangle_{L^2(\mathbb{R}^3)} + |\alpha| t^{|\alpha|-1} \|\langle v \rangle^{\frac{\gamma|\alpha|}{2}} \partial^\alpha f\|_{L^2(\mathbb{R}^3)}^2 \\ &= 2t^{|\alpha|} \langle \langle v \rangle^{\gamma|\alpha|} \partial^\alpha \mathcal{L}f, \partial^\alpha f \rangle_{L^2(\mathbb{R}^3)} + |\alpha| t^{|\alpha|-1} \|\langle v \rangle^{\frac{\gamma|\alpha|}{2}} \partial^\alpha f\|_{L^2(\mathbb{R}^3)}^2 \end{aligned}$$

Let $\theta = \frac{\gamma|\alpha|}{2}$ in Proposition 3.1, it follows that,

$$\frac{d}{dt} \|t^{\frac{|\alpha|}{2}} \langle v \rangle^{\frac{\gamma|\alpha|}{2}} \partial^\alpha f\|_{L^2(\mathbb{R}^3)}^2 \leq -2\|t^{\frac{|\alpha|}{2}} \partial^\alpha f\|_{A, \frac{\gamma|\alpha|}{2}}^2 + \mathbf{B}(f),$$

and

$$\sum_{|\alpha|=m} \|t^{\frac{|\alpha|}{2}} \langle v \rangle^{\frac{\gamma|\alpha|}{2}} \partial^\alpha f\|_{L^2(\mathbb{R}^3)}^2 + 2 \sum_{|\alpha|=m} \int_0^t \|\tau^{\frac{|\alpha|}{2}} \partial^\alpha f\|_{A, \frac{\gamma|\alpha|}{2}}^2 d\tau \leq \int_0^t \mathbf{B}(f) d\tau, \quad (5.6)$$

with

$$\begin{aligned} \mathbf{B}(f) &= |\alpha| t^{|\alpha|-1} \|\langle v \rangle^{\frac{\gamma|\alpha|}{2}} \partial^\alpha f\|_{L^2(\mathbb{R}^3)}^2 \\ &\quad + 2C_0 \sum_{\substack{|\beta| \geq 1 \\ \beta \leq \alpha}} C_\alpha^\beta \sqrt{\beta!} t^{|\alpha|} \|\partial^{\alpha-\beta} f\|_{A, \frac{\gamma|\alpha|}{2}} \|\partial^\alpha f\|_{A, \frac{\gamma|\alpha|}{2}} \\ &\quad + 2C_0 \sum_{\substack{|\beta| \geq 2 \\ \beta \leq \alpha}} C_\alpha^\beta |\beta| \sqrt{\beta!} t^{|\alpha|} \|\partial^{\alpha-\beta} f\|_{A, \frac{\gamma|\alpha|}{2}} \|\partial^\alpha f\|_{2, \frac{\gamma|\alpha|}{2} + \frac{\gamma}{2}} \\ &\quad + C_0 |\gamma| |\alpha| \sum_{\beta \leq \alpha} C_\alpha^\beta \sqrt{\beta!} t^{|\alpha|} \|\partial^{\alpha-\beta} f\|_{A, \frac{\gamma|\alpha|}{2}} \|\partial^\alpha f\|_{\frac{\gamma|\alpha|}{2} + \frac{\gamma}{2}} \\ &= \mathbf{B}_1(f) + \mathbf{B}_2(f) + \mathbf{B}_3(f) + \mathbf{B}_4(f). \end{aligned}$$

It follows from the inequality (2.4) again,

$$\begin{aligned} \mathbf{B}_1(f) &= |\alpha| t^{|\alpha|-1} \|\langle v \rangle^{\frac{\gamma|\alpha|}{2}} \partial^\alpha f\|_{L^2(\mathbb{R}^3)}^2 \\ &\leq |\alpha| t^{|\alpha|-e_{k_0}} \|\langle v \rangle^{\frac{\gamma|\alpha|-e_{k_0}|}{2}} \partial^{\alpha-e_{k_0}} \nabla f\|_{L^2(\mathbb{R}^3)}^2 \\ &\leq \frac{1}{C_1} |\alpha| \|t^{\frac{|\alpha|-e_{k_0}|}{2}} \langle v \rangle^{\frac{\gamma(|\alpha|-e_{k_0}|)}{2}} \partial^{\alpha-e_{k_0}} f\|_A^2, \end{aligned}$$

where k_0 is choose with $\alpha_{k_0} = \max\{\alpha_1, \alpha_2, \alpha_3\}$. So that, by induction assumption (5.5) for $|\alpha - e_{k_0}| = m - 1$

$$\begin{aligned} \int_0^t \mathbf{B}_1(f) d\tau &\leq \frac{1}{C_1} |\alpha| \int_0^t \|\tau^{\frac{|\alpha|-e_{k_0}|}{2}} \langle v \rangle^{\frac{\gamma(|\alpha|-e_{k_0}|)}{2}} \partial^{\alpha-e_{k_0}} f\|_A^2 d\tau \\ &\leq \frac{|\alpha|}{C_1} C^{2|\alpha|-e_{k_0}|+2} (\alpha - e_k!)^2 \leq \frac{|\alpha|}{C_1 (\alpha_k)^2} C^{2|\alpha|} (\alpha!)^2. \end{aligned}$$

We get then

$$\int_0^t \mathbf{B}_1(f) d\tau \leq \frac{3}{C_1} C^{2|\alpha|} (\alpha!)^2 \quad (5.7)$$

For the term $\mathbf{B}_2(f)$, using the fact, for any $\beta \leq \alpha$, $\gamma < 0$,

$$\|\partial^{\alpha-\beta} f\|_{A, \frac{\gamma|\alpha|}{2}} \leq \|\partial^{\alpha-\beta} f\|_{A, \frac{\gamma|\alpha-\beta|}{2}},$$

by using induction assumption (5.5) for $|\alpha - \beta| \leq m - 1$, for $0 < t \leq 1$,

$$\begin{aligned} \int_0^t \mathbf{B}_2(f) d\tau &\leq 2C_0 T^m \sum_{\substack{|\beta| \geq 1 \\ \beta \leq \alpha}} C_\alpha^\beta \sqrt{\beta!} \int_0^t \|\tau^{\frac{|\alpha-\beta|}{2}} \partial^{\alpha-\beta} f\|_{A, \frac{\gamma|\alpha-\beta|}{2}} \|\tau^{\frac{|\alpha|}{2}} \partial^\alpha f\|_{A, \frac{\gamma|\alpha|}{2}} d\tau \\ &\leq 2C_0 \sum_{\substack{|\beta| \geq 1 \\ \beta \leq \alpha}} C_\alpha^\beta \sqrt{\beta!} \left(\int_0^t \|\tau^{\frac{|\alpha-\beta|}{2}} \partial^{\alpha-\beta} f\|_{A, \frac{\gamma|\alpha-\beta|}{2}}^2 d\tau \right)^{1/2} \left(\int_0^t \|\tau^{\frac{|\alpha|}{2}} \partial^\alpha f\|_{A, \frac{\gamma|\alpha|}{2}}^2 d\tau \right)^{1/2} \\ &\leq 4 \left(2C_0 \sum_{|\beta| \geq 1, \beta \leq \alpha} C_\alpha^\beta \sqrt{\beta!} C^{|\alpha-\beta|+1} (\alpha - \beta)! \right)^2 + \frac{1}{2} \int_0^t \|\tau^{\frac{|\alpha|}{2}} \partial^\alpha f\|_{A, \frac{\gamma|\alpha|}{2}}^2 d\tau. \end{aligned}$$

We get then, for $0 < t \leq 1$,

$$\int_0^t \mathbf{B}_2(f) d\tau \leq \tilde{C}_1^2 C^{2|\alpha|} (\alpha!)^2 + \frac{1}{2} \int_0^t \|\tau^{\frac{|\alpha|}{2}} \partial^\alpha f\|_{A, \frac{\gamma|\alpha|}{2}}^2 d\tau \quad (5.8)$$

with

$$\tilde{C}_1 \geq 4C_0 \sum_{|\beta| \geq 1} \frac{1}{\sqrt{\beta!}} C^{1-|\beta|}. \quad (5.9)$$

For the term $\mathbf{B}_3(f)$, by using induction assumption (5.5) for $|\alpha - \beta| \leq m - 2$, $|\alpha - e_{k_0}| = m - 1$,

$$\begin{aligned} \int_0^t \mathbf{B}_3(f) d\tau &\leq 2C_0 \sum_{\substack{|\beta| \geq 2 \\ \beta \leq \alpha}} C_\alpha^\beta |\beta| \sqrt{\beta!} \int_0^t \tau^{|\alpha|} \|\partial^{\alpha-\beta} f\|_{A, \frac{\gamma|\alpha|}{2}} \|\partial^\alpha f\|_{2, \frac{\gamma|\alpha|}{2} + \frac{\gamma}{2}} d\tau \\ &\leq 2C_0 \sum_{\substack{|\beta| \geq 2 \\ \beta \leq \alpha}} C_\alpha^\beta |\beta| \sqrt{\beta!} \left(\int_0^t \|\tau^{\frac{|\alpha-\beta|}{2}} \partial^{\alpha-\beta} f\|_{A, \frac{\gamma|\alpha-\beta|}{2}}^2 d\tau \right)^{1/2} \\ &\quad \times \left(\int_0^t \|\tau^{\frac{|\alpha|-1}{2}} \partial^{\alpha-e_{k_0}} \nabla f\|_{2, \frac{\gamma|\alpha|-1}{2} + \frac{\gamma}{2}}^2 d\tau \right)^{1/2} \\ &\leq 2C_0 \sum_{\substack{|\beta| \geq 2 \\ \beta \leq \alpha}} C_\alpha^\beta |\beta| \sqrt{\beta!} \left(\int_0^t \|\tau^{\frac{|\alpha-\beta|}{2}} \partial^{\alpha-\beta} f\|_{A, \frac{\gamma|\alpha-\beta|}{2}}^2 d\tau \right)^{1/2} \\ &\quad \times \left(\int_0^t \|\tau^{\frac{|\alpha|-1}{2}} \partial^{\alpha-e_{k_0}} f\|_{A, \frac{\gamma|\alpha|-1}{2}}^2 d\tau \right)^{1/2} \\ &\leq 2C_0 \sum_{\substack{|\beta| \geq 2 \\ \beta \leq \alpha}} C_\alpha^\beta |\beta| \sqrt{\beta!} C^{|\alpha-\beta|+1} (\alpha - \beta)! C^{|\alpha|} (\alpha - e_{k_0})!, \end{aligned}$$

we get then

$$\int_0^t \mathbf{B}_3(f) d\tau \leq \tilde{C}_2^2 C^{2|\alpha|} (\alpha!)^2 \quad (5.10)$$

with

$$\tilde{C}_2^2 \geq 6C_0 \sum_{|\beta| \geq 2, \beta \leq \alpha} \frac{1}{\sqrt{\beta!}} C^{1-|\beta|} \geq 2C_0 \sum_{|\beta| \geq 2, \beta \leq \alpha} \frac{|\beta|}{\sqrt{\beta!} \alpha_{k_0}} C^{1-|\beta|}. \quad (5.11)$$

Finally for the term $\mathbf{B}_4(f)$, by using induction assumption (5.5) for $|\alpha - e_{k_0}| = m - 1$,

$$\begin{aligned} \int_0^t \mathbf{B}_4(f) d\tau &\leq C_0 |\gamma| |\alpha| \sum_{0 \leq \beta \leq \alpha} C_\alpha^\beta \sqrt{\beta!} \int_0^t \left(\|\tau^{\frac{|\alpha-\beta|}{2}} \|\partial^{\alpha-\beta} f\|_{A, \frac{\gamma|\alpha|}{2}} \right. \\ &\quad \left. \times \|\tau^{\frac{|\alpha-e_{k_0}|}{2}} \partial^{\alpha-e_{k_0}} \nabla f\|_{\frac{\gamma|\alpha-e_{k_0}|}{2} + \frac{\gamma}{2}} \right) d\tau \\ &\leq C_0 |\gamma| |\alpha| \sum_{0 \leq \beta \leq \alpha} C_\alpha^\beta \sqrt{\beta!} \left(\int_0^t \|\tau^{\frac{|\alpha-\beta|}{2}} \|\partial^{\alpha-\beta} f\|_{A, \frac{\gamma|\alpha-\beta|}{2}}^2 d\tau \right)^{1/2} \\ &\quad \times \left(\int_0^t \|\tau^{\frac{|\alpha-e_{k_0}|}{2}} \partial^{\alpha-e_{k_0}} f\|_{A, \frac{\gamma|\alpha-e_{k_0}|}{2}}^2 d\tau \right)^{1/2} \end{aligned}$$

$$\begin{aligned} &\leq C_0|\gamma||\alpha| \left(\int_0^t \|\tau^{\frac{|\alpha|}{2}} \|\partial^\alpha f\|_{A, \frac{\gamma|\alpha|}{2}}^2 d\tau \right)^{1/2} C^{|\alpha-e_{k_0}|+1}(\alpha-e_{k_0})! \\ &\quad + C_0|\gamma||\alpha| \sum_{0<\beta\leq\alpha} C_\alpha^\beta \sqrt{\beta!} C^{|\alpha-\beta|+1}(\alpha-\beta)! C^{|\alpha-e_{k_0}|+1}(\alpha-e_{k_0})!, \end{aligned}$$

so that

$$\begin{aligned} \int_0^t \mathbf{B}_4(f) d\tau &\leq \frac{1}{2} \int_0^t \|\tau^{\frac{|\alpha|}{2}} \|\partial^\alpha f\|_{A, \frac{\gamma|\alpha|}{2}}^2 d\tau + 4C_0^2|\gamma|^2|\alpha|^2 C^{2|\alpha|}((\alpha-e_{k_0})!)^2 \\ &\quad + C_0|\gamma||\alpha| \sum_{0<\beta\leq\alpha} C_\alpha^\beta \sqrt{\beta!} C^{|\alpha-\beta|+1}(\alpha-\beta)! C^{|\alpha-e_{k_0}|+1}(\alpha-e_{k_0})!, \end{aligned}$$

We get then, for $0 < t \leq 1$,

$$\int_0^t \mathbf{B}_4(f) d\tau \leq \tilde{C}_3^2 C^{2|\alpha|}(\alpha!)^2 + \frac{1}{2} \int_0^t \|\tau^{\frac{|\alpha|}{2}} \partial^\alpha f\|_{A, \frac{\gamma|\alpha|}{2}}^2 d\tau \quad (5.12)$$

with

$$\tilde{C}_3^2 \geq 4C_0^2|\gamma|^2 \frac{|\alpha|^2}{\alpha_{k_0}^2} + \sum_{|\beta|\geq 1} \frac{C_0|\gamma||\alpha|}{\sqrt{\beta!}\alpha_{k_0}} C^{1-|\beta|}. \quad (5.13)$$

Taking the constant C satisfying (5.3), and

$$C^2 \geq \frac{3}{C_1} + \tilde{C}_1^2 + \tilde{C}_2^2 + \tilde{C}_3^2,$$

where the constants are defined by (5.9), (5.11) and (5.13). Combine (5.6), (5.7), (5.8), (5.10) and (5.12), we ends the proof of Proposition 5.1. \square

6. APPENDIX

The standard Hermite functions $(\varphi_n)_{n \in \mathbb{N}}$ are defined for $v \in \mathbb{R}$,

$$\varphi_n(v) = \frac{(-1)^n}{\sqrt{2^n n!} \sqrt{\pi}} e^{\frac{v^2}{2}} \frac{d^n}{dv^n} (e^{-\frac{v^2}{2}}) = -\frac{1}{\sqrt{2^n n!} \sqrt{\pi}} (v - \frac{d}{dv})^n (e^{-\frac{v^2}{2}}) = \frac{a_+^n \varphi_0}{\sqrt{n!}},$$

where a_+ is the creation operator

$$a_+ = \frac{1}{\sqrt{2}} \left(v - \frac{d}{dv} \right).$$

The family $(\varphi_n)_{n \in \mathbb{N}}$ is an orthonormal basis of $L^2(\mathbb{R})$. we set for $n \geq 0$, $\alpha = (\alpha_1, \alpha_2, \alpha_3) \in \mathbb{N}^3$, $x \in \mathbb{R}, v \in \mathbb{R}^3$,

$$\psi_n(x) = 2^{-1/4} \varphi_n(2^{-1/2}x), \quad \psi_n = \frac{1}{\sqrt{n!}} \left(\frac{x}{2} - \frac{d}{dx} \right)^n \psi_0,$$

$$\Psi_\alpha(v) = \prod_{j=1}^3 \psi_{\alpha_j}(v_j), \quad \mathcal{E}_k = \text{Span}(\Psi_\alpha)_{\alpha \in \mathbb{N}^3, |\alpha|=k},$$

with $|\alpha| = \alpha_1 + \alpha_2 + \alpha_3$. The family $(\Psi_\alpha)_{\alpha \in \mathbb{N}^3}$ is an orthonormal basis of $L^2(\mathbb{R}^3)$ composed by the eigenfunctions of the 3-dimensional harmonic oscillator

$$\mathcal{H} = -\Delta_v + \frac{|v|^2}{4} = \sum_{k \geq 0} (k + \frac{3}{2}) \mathbb{P}_k, \quad Id = \sum_{k \geq 0} \mathbb{P}_k, \quad (6.1)$$

where \mathbb{P}_k stands for the orthogonal projection

$$\mathbb{P}_k f = \sum_{|\alpha|=k} (f, \Psi_\alpha)_{L^2(\mathbb{R}^v)} \Psi_\alpha.$$

In particular,

$$\Psi_0(v) = \frac{1}{(2\pi)^{\frac{3}{4}}} e^{-\frac{|v|^2}{4}} = \mu^{1/2}(v),$$

where $\mu(v)$ is the Maxwellian distribution. Setting

$$A_{\pm,j} = \frac{v_j}{2} \mp \partial_j, \quad 1 \leq j \leq 3, \quad (6.2)$$

we have

$$\Psi_\alpha = \frac{1}{\sqrt{\alpha_1! \alpha_2! \alpha_3!}} A_{+,1}^{\alpha_1} A_{+,2}^{\alpha_2} A_{+,3}^{\alpha_3} \Psi_0, \quad \alpha = (\alpha_1, \alpha_2, \alpha_3) \in \mathbb{N}^3,$$

and

$$A_{+,j} \Psi_\alpha = \sqrt{\alpha_j + 1} \Psi_{\alpha + e_j}, \quad A_{-,j} \Psi_\alpha = \sqrt{\alpha_j} \Psi_{\alpha - e_j} (= 0 \text{ if } \alpha_j = 0), \quad (6.3)$$

where (e_1, e_2, e_3) stands for the canonical basis of \mathbb{R}^3 . For more details of the Hermite functions, we can refer to [15] and the reference theorem.

ACKNOWLEDGMENTS

The first author is supported by the Fundamental Research Funds for the Central Universities of China, South-Central University for Nationalities (No. CZT20007). The second author is supported by the NSFC (No.12031006) and the Fundamental Research Funds for the Central Universities of China.

REFERENCES

- [1] R. Alexandre, Y. Morimoto, S. Ukai, C.-J. Xu and T. Yang, Uncertainty principle and kinetic equations, *J. Funct. Anal.*, **255** (2008), 2013–2066.
- [2] Hua Chen, Wei-Xi Li and Chao-Jiang Xu, Analytic smoothness effect of solutions for spatially homogeneous Landau equation, *Journal of Differential Equations* 248 (2010), 77-94.
- [3] Hua Chen, Wei-Xi Li and Chao-Jiang Xu, Propagation of Gevrey regularity for solutions of Landau equations, *Kinetic and Related Models* (2008), no. 3, 355- 368.
- [4] Hua Chen, Wei-Xi Li and Chao-Jiang Xu, Gevrey regularity for solution of the spatially homogeneous Landau equation. *Acta Mathematica Scientia* 29(2009), 673-686.
- [5] P. Degond, M. Lemou, Dispersion Relations for the Linearized Fokker-Planck Equation, *Arch. Rat. Mech. Anal.*, 138(1997), 137-167.
- [6] L. Desvillettes, C. Villani, On the spatially homogeneous Landau equation for hard potentials. I. Existence, uniqueness and smoothness. *Comm. Partial Differential Equations* 25, (2000), 8 179-259.
- [7] L. Glangetas, H.-G. Li, C.-J. Xu, Sharp regularity properties for the non-cutoff spatially homogeneous Boltzmann equation, *Kinet. Relat. Models* 9 (2016), 299-371.
- [8] T. Gramchev, S. Pilipović, L. Rodino, Classes of degenerate elliptic operators in Gelfand-Shilov spaces. *New Developments in Pseudo-Differential Operators*. Birkhäuser Basel, 2009: 15-31.
- [9] Y. Guo, The Landau Equation in a Periodic Box, *Comm. Math. Phys.* 231(2002), 391-434.
- [10] N. Lerner, Y. Morimoto, K. Pravda-Starov, C.-J. Xu, Phase space analysis and functional calculus for the linearized Landau and Boltzmann operators, *Kinet. Relat. Models* 6 (2013), no. 3, 625-648.
- [11] H.-G. Li, The Gelfand-Shilov smoothing effect for the radially symmetric homogeneous Landau equation with Shubin initial datum, *Comptes rendus Mathématique*, 356(2018), 613-625.

- [12] H.-G. Li, C.-J. Xu, Cauchy problem for the spatially homogeneous Landau equation with Shubin class initial datum and Gelfand-Shilov smoothing effect, *Siam J. Math. Anal.*, 51 (2019), 532-564.
- [13] H.-G. Li, C.-J. Xu, Analytic smoothing effect of the non-linear spatially homogeneous Landau equation with hard potentials, *Science China Mathematics* (2022), inpress.
- [14] Y. Morimoto, C.-J. Xu, Ultra-analytic effect of Cauchy problem for a class of kinetic equations, *Journal of Differential Equations*, 247 (2009), no.2, 596-617.
- [15] Y. Morimoto, K. Pravda-Starov, C.-J. Xu, A remark on the ultra-analytic smoothing properties of the spatially homogeneous Landau equation. *Kinetic and Related Models* 6 (2013), 715-727.
- [16] K. A. Ross, *Elementary analysis: The theory of calculus*, Undergraduate texts in Mathematics, 2013, 230-240.
- [17] C. Villani, On a new class of weak solutions to the spatially homogeneous Boltzmann and Landau equations, *Arch. Rational Mech. Anal.*, 143 (1998), 273-307.
- [18] C. Villani, On the spatially homogeneous Landau equation for Maxwellian molecules, *Mathematical Models and Methods in Applied Sciences*, 8(1998), no. 6, 957-983.
- [19] H. Wang, K.-C. Wu, Solving linearized Landau equation pointwisely. arXiv preprint arXiv:1709.00839 (2017).

HAO-GUANG LI,
 SCHOOL OF MATHEMATICS AND STATISTICS, SOUTH-CENTRAL UNIVERSITY FOR NATIONALITIES
 430074, WUHAN, CHINA
Email address: lihaoguang@scuec.edu.cn

CHAO-JIANG XU,
 DEPARTMENT OF MATHEMATICS, NANJING UNIVERSITY OF AERONAUTICS AND ASTRONAUTICS,
 NANJING 211106, CHINA
 KEY LABORATORY OF MATHEMATICAL MODELLING AND HIGH PERFORMANCE COMPUTING OF
 AIR VEHICLES (NUAA), MIIT, NANJING 210016, CHINA
 UNIVERSITÉ DE ROUEN, CNRS UMR 6085, LABORATOIRE DE MATHÉMATIQUES
 76801 SAINT-ETIENNE DU ROUVRAY, FRANCE
Email address: xuchaojiang@nuaa.edu.cn