

PREPARED FOR SUBMISSION TO JHEP

# Lense–Thirring effects from on-shell amplitudes

---

Wei-Ming Chen<sup>1</sup> Ming-Zhi Chung<sup>2</sup> Yu-tin Huang<sup>2,3</sup> Jung-Wook Kim<sup>4</sup>

<sup>1</sup>*Department of Physics, Kobe University, Kobe 657-8501, Japan*

<sup>2</sup>*Department of Physics and Astronomy, National Taiwan University, Taipei 10617, Taiwan*

<sup>3</sup>*Physics Division, National Center for Theoretical Sciences, National Tsing-Hua University, No.101, Section 2, Kuang-Fu Road, Hsinchu, Taiwan*

<sup>4</sup>*Centre for Theoretical Physics, Department of Physics and Astronomy, Queen Mary University of London, Mile End Road, London E1 4NS, United Kingdom*

*E-mail:* [tainist@gmail.com](mailto:tainist@gmail.com), [dchung0741@gmail.com](mailto:dchung0741@gmail.com),  
[yutin@phys.ntu.edu.tw](mailto:yutin@phys.ntu.edu.tw), [jung-wook.kim@qmul.ac.uk](mailto:jung-wook.kim@qmul.ac.uk)

ABSTRACT: Effects of massive object’s spin on massive-massless  $2 \rightarrow 2$  classical scattering is studied. Focus is set on the less-considered dimensionless expansion parameter  $\lambda/b$ , where  $\lambda$  is the massless particle’s wavelength and  $b$  is the impact parameter. Corrections in  $\lambda/b$  start to appear from  $\mathcal{O}(G^2)$ , with leading correction terms tied to the gravitational Faraday effect, which is a special case of the Lense–Thirring effect. We compute the eikonal phase up to  $\mathcal{O}(G^2)$  and extract spin effect on the scattering angle and time delay up to 14th order in spin. The gravitational Faraday effect at linear order in spin [1] is reproduced by  $\lambda/b$  correction terms, which we compute to higher orders in spin. We find that the equivalence principle, or universality, holds up to NLO for general spinning bodies, i.e. away from geometric optics limit. Furthermore, in the black hole limit, we confirm the absence of particular spin structure observed and conjectured to hold to arbitrary spin order [2–8].

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>General discussion of <math>\lambda/b</math> corrections</b>	<b>6</b>
2.1	Dimensional analysis	7
2.2	Subleading $\lambda/b$ corrections and rotations of polarization planes	9
<b>3</b>	<b>The <math>\mathcal{O}(G^1)</math> Compton eikonal phase for spinning objects</b>	<b>10</b>
3.1	The tree amplitude for eikonal phase	11
3.2	The $\mathcal{O}(G^1)$ eikonal phase and observables	13
<b>4</b>	<b>The <math>\mathcal{O}(G^2)</math> eikonal phase and observables</b>	<b>15</b>
4.1	Extraction of scalar integral coefficients	16
4.1.1	Extraction of box and crossed box coefficients	18
4.1.2	Extraction of triangle coefficients	18
4.2	eikonal phase and its exponentiation	19
4.3	results	22
4.3.1	leading order	22
4.3.2	subleading order	25
4.4	Discussions: equivalence principle, spin structures and massless limits	26
<b>5</b>	<b>Conclusion</b>	<b>28</b>
<b>A</b>	<b>Eikonal phase normalisation</b>	<b>29</b>
<b>B</b>	<b>Linking eikonal phase to geometric optics</b>	<b>30</b>
<b>C</b>	<b>Simplifying the spin factor for long-distance physics</b>	<b>32</b>
<b>D</b>	<b>Eikonal phase at NLO to <math>\mathcal{O}(S^4)</math></b>	<b>33</b>
D.1	Cubic order in spin	34
D.2	Quartic order in spin	34

---

# 1 Introduction

The fact that the classical limit of scattering amplitudes can correctly capture black hole dynamics, is on itself a fascinating subject irrespective of its remarkable success in applications to gravitational wave physics, which has helped brought into the spotlight since the land mark detection by the LIGO/Virgo collaboration [9, 10]. That point-like sources can faithfully reproduce effects of gravitational field in the weak field limit was appreciated long ago in the seminal work of ref.[11], where it was shown that the Schwarzschild metric can be perturbatively reconstructed order by order in  $G$  through Feynman diagrams. This was later bolstered by the computations of perturbative stress-energy form factors [12, 13]. Later development focused on the Post Minkowskian (PM) expansion [14–23] for conservative two-body potential of binary black holes, which can be extracted from the gravitational scattering amplitude of massive scalars [24–27]. By matching the field theory amplitude to effective field theory (EFT) description [28], the on-shell simplification led to rapid progress, culminating in the state of the art 4 PM conservative potential [29–31]. As the PM expansion is exact in velocity ( $v^2$ ), the results can be directly compared and matched to the post-Newtonian (PN) expansion [32–39].

The extension to spinning black holes brings in a new angle to the correspondence. It was already shown in the early work of refs.[40, 41] that the spinning Hamiltonian can be captured by the scattering of elementary spinning particle. However, the absence of elementary particle beyond spin-2 casts an intriguing conundrum for the program. This was partially overcome by the kinematic definition of minimal coupling for arbitrary spin particle [42] whose classical spin-limit ( $\hbar \rightarrow 0$ ,  $s \rightarrow \infty$  and  $s\hbar$  held fixed) successfully reproduced the 1 PM dynamics of Kerr black hole [43–50]. For non black hole spinning bodies, in the PN EFT approach one introduces spins as extra degrees of freedom [51] on the worldline for the point particle effective action, accompanied by an infinite number of spin-induced multipole operators whose strengths are parametrised by the corresponding Wilson coefficients [52, 53]; see refs.[54–57] for recent works, ref.[58] in the context of PM EFT, and refs.[59–61] for worldline QFT. On the amplitude side, this can be incorporated by matching the Wilson coefficients to those of non-minimal couplings [46, 62], or to introduce a field theory effective lagrangian [2]. The latter approach has been implemented at 2 PM to obtain the spin-dependent part of the conservative hamiltonian up to fifth order in spin [4, 7].

In going beyond the quartic order in spin at 2 PM, from the on-shell approach one needs the gravitational Compton amplitude for the interaction of higher spin states which are not unique [42, 45]. A proposal utilizing appropriately conserved off-shell current was given for spin-5/2 in ref.[63] (see ref.[64] for a compact expression of the ambiguity). When translated to the effective field theory Lagrangian approach in ref.[2], the issue becomes the determination of black hole Wilson coefficients at 2 PM. In a nutshell, the question is *can one identify the underlying principles that determine the Kerr black hole limit ?*

A natural starting point is to examine the 1 PM dynamics and to take note of any special aspects emerging in the Kerr limit. It was observed in ref.[65] that the change in

spin-entanglement entropy is minimized for minimal couplings, which was later identified as the suppression of spin-flipping sector in the eikonal phase [66]. The absence of spin-flipping sectors for the 1 PM eikonal amplitude is not surprising as minimal coupling for massive spinning higher spins are defined such that it has optimal power counting in the UV, which corresponds to helicity preserving interactions. Besides optimal UV behaviour, an interesting aspect of the 1 PM potential is the presence of shift symmetry. For general spinning bodies, the 1 PM potential depends on the spin vector through the combination  $\epsilon(q, p_2, p_1, S)$  [62], which exhibits the independence on the longitudinal part of the spin vector along the impact parameter plane, i.e. the potential is invariant under the shift symmetry proposed in ref.[7],

$$S_i^\mu \rightarrow S_i^\mu + \xi_i \frac{q^\mu}{q^2}. \quad (1.1)$$

On the other hand, preservation of this symmetry at 2 PM becomes a nontrivial constraint. Earlier, it was noted that in the spin bilinear sector the spin-spin interaction  $(S_1 \cdot S_2)$  only occurs through a special combination [2, 40, 67, 68]

$$(q \cdot S_1)(q \cdot S_2) - q^2(S_1 \cdot S_2),$$

which is secretly related to shift symmetry, as will be explained. This combination can be generalised to the case where spin vectors are taken from the same body, i.e.

$$(q \cdot S_i)(q \cdot S_j) - q^2(S_i \cdot S_j), \quad (1.2)$$

where  $i = j$  is also allowed. It was noted in refs.[3, 4] that for 2 PM classical amplitude, from which the conservative Hamiltonian is extracted, the black hole limit (defined by absence of tidal terms or by correct spin-quadrupole moments) is special in that the structure eq.(1.2) appears, at least to *quadratic or bilinear* order in spin. A related observation is that when the spin operators are expanded on the basis of Lorentz invariants


$$\{(q \cdot S_i), (p_j \cdot S_i), \epsilon(q, p_1, p_2, S_i)\}, \quad (1.3)$$

the black hole limit is special in that  $(q \cdot S_i)$  terms all vanish, at least to *quartic* order in spin [5](appendix B).<sup>1</sup> The two observations are related since the structure eq.(1.2) appears when the product  $\epsilon(q, p_1, p_2, S_i)\epsilon(q, p_1, p_2, S_j)$  is reorganised using Levi-Civita identities. Also, the latter observation is equivalent to shift symmetry eq.(1.1);  $\epsilon(q, p_2, p_1, S)$  is manifestly invariant, while  $(p_j \cdot S_i)$  is invariant classically. Recently, these equivalent conditions on the spin structures of the Hamiltonian were used with an additional assumption of favorable high energy behaviour to fix Wilson coefficients that determine spinning-spinless sectors at 2 PM [6–8].<sup>2</sup>

---

<sup>1</sup>The validity of the exponential form of the Kerr Compton amplitude used in refs.[5, 44, 47, 49, 69] to  $\mathcal{O}(S^4)$  has been confirmed by comparing to black hole perturbation theory. The authors would like to thank Justin Vines for sharing conclusions of their upcoming work [70].

<sup>2</sup>The authors would like to thank Kays Haddad for clarification on the relation of the spin structure eq.(1.2) to the Kerr limit.



**Figure 1:** A cartoon of gravitational Faraday effect. The polarisation direction (black double arrow) of the electromagnetic wave propagating along the direction of the spinning axis gets rotated (red double arrow) by the frame-dragging effect of the spinning body.

In this paper, we revisit the classical limit of the gravitational Compton amplitude with spin effects included. The goal is twofold: on the one hand a rotating object will induce Lense–Thirring (or gravitomagnetic frame-dragging) effects, which can rotate the polarization plane of propagating electromagnetic or gravitational waves as in fig.1. This effect should be visible through the classical Compton amplitude describing the scattering of photons or gravitons off a spinning object. At the same time, the role of shift symmetry or the distinctive spin-spin interaction in determining the black hole limit can be tested in this context, as the amplitude will be parameterised by the worldline Wilson coefficients for linear coupling to gravitons.

Earlier studies of gravitational Compton amplitude focused on the photon/graviton and massive scalar system, where one is interested in the corrections to the scattering angle (light bending) of null geodesics [71–74] and Shapiro time delay [75, 76]. An important tool is the eikonal approximation, corresponding to large center of mass energy and small scattering angle ( $s \gg t$ ), where the amplitude exponentiates [77–84]. Importantly, since the amplitude depends on the external helicity states, instead of the eikonal phase we have the eikonal phase matrix [75, 76]

$$\begin{pmatrix} \delta_{+,+} & \delta_{-,+} \\ \delta_{+,-} & \delta_{-,-} \end{pmatrix} = \bar{\delta} \mathbb{I} + \alpha_i \sigma_i, \quad (1.4)$$

where we use  $2 \rightarrow 2$  convention for the helicity states, and the matrix is written in a manifest Hermitian form. The phases  $\bar{\delta}, \alpha_i$ , which are functions of impact parameter  $b$  and frequency  $\omega$  of the massless mode, can be perturbatively computed in powers of  $G$ . Previous computations focused on  $\bar{\delta} = (\delta_{+,+} + \delta_{-,-})/2$ , from which one extracts the scattering angle and time delay, via

$$\theta = \frac{1}{\omega} \frac{\partial}{\partial b} \bar{\delta}(b, \omega), \quad \delta t = \frac{\partial}{\partial \omega} \bar{\delta}(b, \omega). \quad (1.5)$$

For scalar Compton  $\alpha_3 = 0$  due to the lack of parity violating interactions, while  $\alpha_{1,2}$  was studied for the effects of higher derivative corrections to Einstein-Hilbert term, which generate non-trivial same helicity sector at tree and one-loop [75, 76]. These helicity non-preserving components modify circular polarisation states to elliptical polarised states.

We will compute the Compton eikonal matrix for general spinning bodies up to  $\mathcal{O}(G^2)$ . The resulting scattering angle and time delay will be a function of Wilson coefficients. In contrast with the eikonal phase for massive scattering amplitudes, there is a new dimensionless parameter in the classical regime:  $1/\omega b = \lambda/b$  where  $\lambda$  is the wavelength of the gravitational/EM wave. Thus for spinning bodies we have a double expansion in  $\lambda/b$  and  $a/b$ , where  $a = S/m$  is the spin length. From dimensional analysis, we can see that  $\lambda/b$  corrections in the classical limit only occurs at  $\mathcal{O}(G^2)$ :

$$\begin{aligned} \mathcal{O}(G^2) : \quad \bar{\delta} &= \frac{15\pi G^2 m^2 \omega}{4b} \left[ 1 - \frac{(\hat{k}_2 \times \vec{a}) \cdot \vec{b}}{3b^2} + \mathcal{O}\left(\frac{a^2}{b^2}\right) \right], \\ \alpha := -\frac{\alpha_3}{|\hbar|} &= \frac{5\pi G^2 m^2 (\hat{k}_2 \cdot \vec{a})}{4b^3} \left[ 1 - (1+C_{S^2}) \frac{9(\hat{k}_2 \times \vec{a}) \cdot \vec{b}}{8b^2} + \mathcal{O}\left(\frac{a^2}{b^2}\right) \right], \end{aligned} \quad (1.6)$$

where  $\hat{k}_2$  is the spatial direction of the incoming massless plane wave. Here we are giving the result in terms of an  $a/b$  expansion. Note that LO for  $\alpha_3$  is subleading for the individual eikonal phase  $\delta_{+,+}$  and  $\delta_{-,-}$ . We identify  $\alpha$  as the rotation angle of the gravitational Faraday effect [1, 85–92] whose name is an analogy to the magneto-optical effect bearing the same name; for linearly polarised EM waves, the polarisation direction rotates due to the “magnetic” background. Indeed we see that  $\alpha$  is only present when there is a non-zero spin and the leading linear in spin term matches with that computed using parallel transport along null geodesics [1]. While the leading term for  $\alpha$  is universal, the subleading terms depend on the Wilson coefficients. We provide the result up to 14th order in spin, for both photons and gravitons.

The spin dependence of the classical Compton amplitude can also be cast into a Lorentz invariance basis  $(q \cdot S)$ ,  $(k \cdot S)$  and  $\epsilon(q, k, p, S)$  where  $k$  is the momentum of the incoming massless state. We find that at leading order in  $\lambda/b$  expansion, the  $\mathcal{O}(G^2)$   $\bar{\delta}$  and  $\alpha_3$  also exhibits vanishing  $(q \cdot S)$  term in the black hole limit. Note that the naive extension of the shift symmetry in eq.(1.1) no longer applies here due to different  $\hbar$  counting  $(k \cdot S) = \mathcal{O}(\hbar^0)$ : the shift operation acting on  $(k \cdot S) \rightarrow (k \cdot S) + (k \cdot q)/q^2 = (k \cdot S) - 1/2$  generates a classical term. Similar violation was in fact already seen in the helicity non-conserving Compton amplitude in ref.[6]. Here we observe it at  $\mathcal{O}(G^2)$  in the helicity conserving sector. However, since the invariant result is the absence of  $(q \cdot S)$  terms, which in the 2→2 massive scattering is equivalent to the shift symmetry, our observation should be interpreted as raising the interesting prospect of generalizing the shift for massless-massive scattering. Similarly by considering the normalized quantity  $\alpha_3/|\hbar|$  along with  $\bar{\delta}$ , one can show that equivalence principle holds at leading order in  $\lambda/b$  expansion. At subleading order in  $\lambda/b$  both of these properties begin to deviate. The deviations can be understood as the eikonal limit being the geometric optics limit, where one ignores the wave nature of electric-magnetic/gravitational waves and treat it as null rays. This is valid when the wave length is much smaller than

the impact parameter  $\frac{\lambda}{b} \ll 1$ . Thus expansion in  $\frac{\lambda}{b}$  would correspond to corrections to the approximation.

This paper is organized as follows: we begin with a brief discussion of  $\lambda/b$  corrections to the eikonal phase and their physical interpretation in sec.2. Then in sec.3 we compute the  $\mathcal{O}(G^1)$  eikonal phase for general spinning body, and derive physical observables. In sec.4, we present the computation and result for  $\mathcal{O}(G^2)$  eikonal phase.

## 2 General discussion of $\lambda/b$ corrections

The kinematics of interest has four length scales; the Compton wavelength  $\lambda_C \sim m^{-1}$  controlling the “quantumness” of the massive body, the wavelength  $\lambda = \omega^{-1}$  of the massless particle,<sup>3</sup> the spin length  $a \sim |\vec{S}|m^{-1}$  controlling the “size” of the massive body and the impact parameter  $b$  controlling the separation. In the (classical) geometric optics regime the scale hierarchy is<sup>4</sup>

$$\lambda_C \ll \lambda \ll a \ll b. \quad (2.1)$$

Treating the internal nature of the massive body as quantum, the separation between “quantum” and “classical” effects amounts to utilizing the scale hierarchy between  $\lambda_C$  and all other scales:

$$\frac{\lambda_C}{b} \sim \frac{\hbar}{mb}, \quad \frac{\lambda_C}{a} \sim \frac{\hbar}{|\vec{S}|}, \quad \frac{\lambda_C}{\lambda} \sim \frac{\hbar\omega}{m}, \quad (2.2)$$

where  $\hbar\omega$  is the energy of the massless particle of momentum. Considering that the transfer momentum  $q \sim b^{-1}$  is the Fourier dual of the impact parameter  $b$ , the classical limit corresponds to the following  $\hbar$  scaling for the variables:

$$q^\mu \rightarrow \hbar q^\mu, \quad k^\mu \rightarrow \hbar k^\mu, \quad S^\mu \rightarrow \frac{S^\mu}{\hbar}, \quad m \rightarrow m, \quad \omega \rightarrow \hbar\omega, \quad G \rightarrow \frac{G}{\hbar}. \quad (2.3)$$

Due to the separation of scale in eq.(2.1), we have two classical expansion parameters:


$$\frac{a}{b} \sim \frac{|\vec{S}|}{mb}, \quad \frac{\lambda}{b} = \frac{1}{b\omega}. \quad (2.4)$$

Thus we will only compute the leading term in  $\hbar \rightarrow 0$  limit, with the spin and energy expansion  $a/|b|$ ,  $(\omega b)^{-1}$  being an additional expansion *in the classical regime*. From now on we will suppress  $\hbar$  dependence for brevity by setting  $\hbar = 1$ .

Let’s consider the Compton amplitude where the incoming massless state (with  $k_2$ ) scatters off a massive object (with momenta  $p_1$ ) as in fig.2. Due to vanishing/suppressed helicity-flip amplitudes in general relativity, the outgoing massless state (with  $k_3$ ) will

<sup>3</sup>For massless particles we consider the wavenumber four-vector  $\bar{k}^\mu := k^\mu/\hbar$  as the classical quantity.

<sup>4</sup>The hierarchy  $\lambda \ll a$  is only needed for the conceptual foundation of geometric optics and is irrelevant for any of the analyses in this paper. In terms of wavepackets, there is an additional scale hierarchy  $\lambda \ll \xi \ll b$  where  $\xi$  is the wavepacket size [93]. JWK would like to thank Donal O’Connell for discussions on this point.



**Figure 2:** The diagram shows the configuration of the Compton amplitudes in our consideration. The wavy line represents the massless particle and solid line is the massive spinning object. The arrows show the directions of momentum flow.

be of the same helicity as the incoming state. The Compton amplitude of interest is  $M(\mathbf{1}2^{\pm h}\mathbf{3}^{\pm h}\mathbf{4})$ , where  $h = 1, 2$  denotes photon and graviton respectively. The eikonal approximation is an all-order resummation of  $M_4$  where the amplitude is reorganised as a phase factor  $iM_4 = e^{i\delta} - 1$ , which has been extensively studied in the literature [77–84]. The eikonal phase  $\delta(b)$  is obtained by Fourier transforming the  $2 \rightarrow 2$  amplitude to impact parameter space<sup>5</sup>

$$\begin{aligned} e^{i\delta_{\pm,\pm}} - 1 &:= \int \frac{d^D q}{(2\pi)^D} \hat{\delta}(2p_1 \cdot q) \hat{\delta}(2k_2 \cdot q) \left[ e^{-ib \cdot q} iM_4(\mathbf{1}2^{\pm h}\mathbf{3}^{\pm h}\mathbf{4}) \right] \\ &= \frac{i}{4m\omega} \int \frac{d^{D-2} q}{(2\pi)^{D-2}} \left[ e^{i\vec{b} \cdot \vec{q}} M_4(\mathbf{1}2^{\pm h}\mathbf{3}^{\pm h}\mathbf{4}) \right], \end{aligned} \quad (2.5)$$

where  $\hat{\delta}(x) = 2\pi\delta(x)$  and Mandelstam invariants are parametrised as:

$$s = (p_1 + k_2)^2 = 2m\omega + m^2, \quad t = (k_3 - k_2)^2 = -|\vec{q}|^2, \quad u = (p_1 - k_3)^2 = m^2 - 2m\omega + |\vec{q}|^2. \quad (2.6)$$

This parametrises  $\omega = (p_1 \cdot k_2)/m$  as the frequency of the incoming massless particle in the rest frame of  $p_1$ . The transfer momentum from the massive to massless state is  $q^\mu = k_3^\mu - k_2^\mu$  and its spatial part in centre of momentum (COM) frame is the three-vector  $\vec{q}$ .

$$\begin{aligned} p_1^\mu &= (\sqrt{m^2 + k^2}, -\vec{k}), \quad p_4^\mu = (\sqrt{m^2 + k^2}, -\vec{k} - \vec{q}), \\ k_2^\mu &= (k, \vec{k}), \quad k_3^\mu = (k, \vec{k} + \vec{q}), \end{aligned} \quad (2.7)$$

where the COM frame variable  $k$  is given by  $\omega$  through the relation

$$(k^2/m^2) = \frac{\sqrt{1 + 4(\omega^2/m^2)} - 1}{2} = (\omega^2/m^2) \times [1 + \mathcal{O}(\hbar^2)], \quad (2.8)$$

and can be considered as  $\omega$  in the classical limit.

## 2.1 Dimensional analysis

Before we begin, let's first perform a dimension analysis to determine the allowed operators that can accompany various subleading (in  $\lambda/b$ ) corrections. Each element of the eikonal

<sup>5</sup>Different definitions for the impact parameter space are also used in the literature, which is reviewed in appendix A.



phase matrix, which we collectively denote by  $\delta$ , can be expanded as a perturbative series in the coupling  $G$ , the spin  $S$ , and the dimensionless ratio  $\lambda/b$ .

$$\delta = \sum_{L=0}^{\infty} G^{L+1} \delta^{(L)} = \sum_{L=0}^{\infty} G^{L+1} \left( \sum_{j=0}^{\infty} \delta_{S^j}^{(L)} \frac{a^j}{b^j} \right) = \sum_{L=0}^{\infty} G^{L+1} \left( \sum_{j=0}^{\infty} \frac{a^j}{b^j} \left( \sum_{i=0}^{\infty} \delta_{S^j}^{(L,i)} \frac{\lambda^i}{b^i} \right) \right), \quad (2.9)$$

where the superscript  $(L)$  denotes the  $L$ -loop ( $\mathcal{O}(G^{L+1})$ ) contribution and subscript  $S^n$  denotes the  $\mathcal{O}(S^n)$  contribution of the given  $L$ -loop contribution. The second subscript  $i$  of the paired subscript  $(L, i)$  denotes subleading  $(\lambda/b)^i$  order contribution to the given  $L$ -loop contribution.

Let's define  $M_4^{(L,0,0)}$  as the terms of the Compton amplitude  $M_4$  that determine the leading spin-less sector  $\delta_{S^0}^{(L,0)}$  of the  $L$ -loop eikonal phase  $\delta^{(L)}$ , and  $M_4^{(L,i,j)}$  as that which determines  $(\lambda/b)^i (a/b)^j$  corrections ( $\delta_{S^j}^{(L,i)}$ ). The classical limit of background-test mass  $2 \rightarrow 2$  scattering has only one master scalar integral (the  $L$ -loop fan integral) at each loop order, while other remaining integrals contribute to the iteration terms required from the exponentiation in the eikonal approximation [94]. Thus as long as the eikonal approximation is valid and iteration holds, only one integral is relevant at each loop. Defining the ratio:

$$F_{(L,i,j)} := \frac{M_4^{(L,i,j)}}{M_4^{(L,0,0)}}, \quad (2.10)$$

becomes simply the ratios of the integral coefficient of the master scalar integral, and thus functions of Lorentz invariants (the helicity weight is cancelled in the ratio). To analyse  $F_{(L,i,j)}$  we first write all possible Lorentz invariants that can be constructed from kinematics. There are Mandelstam invariants which we can combine into a dimensionless expansion parameter  $\lambda/b$ .<sup>6</sup>

$$(s - m^2) \sim m\omega \sim \frac{m}{\lambda}, \quad t = -|\vec{q}|^2 \sim \frac{1}{b^2} \quad \Rightarrow \quad \left[ \frac{m\sqrt{-t}}{s - m^2} \right] \sim \frac{\lambda}{b} \quad (2.11)$$

The introduction of the spin vector brings three new invariants due to the transverse condition  $p \cdot S = 0$ . The three Lorentz invariants can be combined into the independent dimensionless expansion parameter  $a/b$  as:

$$\begin{aligned} (q \cdot S) \sim \frac{ma}{b} &\Rightarrow \left[ \frac{q \cdot S}{m} \right] \sim \frac{a}{b} \\ (n \cdot S) \sim \frac{m^2 a}{\lambda b} &\Rightarrow \left[ \frac{n \cdot S}{m(s - m^2)} \right] \sim \frac{a}{b} \\ (k_2 \cdot S) \sim \frac{ma}{\lambda} &\Rightarrow \left[ \frac{\sqrt{-t}(k_2 \cdot S)}{s - m^2} \right] \sim \frac{a}{b}. \end{aligned} \quad (2.12)$$

---

<sup>6</sup>We ignore  $\mathcal{O}(1)$  numerical factors in power counting analysis.

The orthogonal vector  $n^\mu$  is defined as  $n^\mu = \epsilon^{\mu\alpha\beta\gamma} p_{1\alpha} k_{2\beta} q_\gamma$ . These invariants form a basis for  $F_{(L,i,j)}$

$$F_{(L,i,j)} = \sum_{\substack{\alpha+\beta+\gamma=j \\ \alpha,\beta,\gamma \geq 0}} c_{i,\alpha,\beta,\gamma} \left[ \frac{m\sqrt{-t}}{s-m^2} \right]^i \left[ \frac{q \cdot S}{m} \right]^\alpha \left[ \frac{n \cdot S}{m(s-m^2)} \right]^\beta \left[ \frac{\sqrt{-t}(k_2 \cdot S)}{s-m^2} \right]^\gamma, \quad (2.13)$$

where  $c_{i,\alpha,\beta,\gamma}$  are numerical coefficients. Since  $F_{(L,i,j)}$  is a rational function, the powers of  $\sqrt{-t}$  must add up to an even number and we have the constraint  $i + \gamma \in 2\mathbb{Z}$ . We list the consequences of this constraint:

1. There are no  $\lambda/b$  corrections at  $\mathcal{O}(G^1)$ . The scalar term  $M_4^{(0,0,0)}$  contains a  $t^{-1}$  pole prefactor. As any expansion in  $\lambda/b$  must be combined to form integer powers of  $-t = |\vec{q}|^2$ , the corrections will cancel the  $t^{-1}$  pole and result in derivatives of delta functions in impact parameter space, which do not contribute to long range effects.
2. At  $\mathcal{O}(G^2)$ , the scalar term  $M_4^{(1,0,0)}$  contains a  $1/|\vec{q}|$  prefactor from the scalar triangle integral. The non-analytic structure of the  $1/|\vec{q}|$  prefactor yields long range contributions which cannot be cancelled by integer powers of  $-t$ , allowing arbitrary order  $\lambda/b$  corrections. The condition  $i + \gamma \in 2\mathbb{Z}$  forces odd powers of  $\lambda/b$  corrections to accompany odd powers of  $(k_2 \cdot S)$ , and the first  $\lambda/b$  correction to be at  $(\lambda/b)^2$  order for spinless case ( $j = 0$ ). The corrections at this order are dependent on conventions for the eikonal phase as explained in appendix A.

As we will argue in the following, the polarisation plane rotation is an  $\mathcal{O}(\lambda/b)$  correction from the leading eikonal approximation, thus it is first observed at one-loop order. This is consistent with the conclusion in ref.[95] that gravitational Faraday effect cannot be observed at  $\mathcal{O}(G^1)$  order.

## 2.2 Subleading $\lambda/b$ corrections and rotations of polarization planes

Previous studies on eikonal phases of gravitational Compton scattering mostly ignored subleading effects in the  $\lambda/b$  expansion. These subleading corrections are related to the wave-like nature of the massless particle since they depend on the wavelength  $\lambda$  of the particle, but which physical properties are these terms reflecting?

To answer this question, we consider geometric optics. The propagation of light (or gravitational wave) is treated as a classical curve where the tangent vector at each point of the curve is normal to the wavefront. For an introduction, see e.g. §53 of ref.[96]. We consider the propagation of waves as a classical trajectory in the limit  $\lambda \rightarrow 0$  or  $\lambda/b \ll 1$ . Selecting a component of the wave (e.g. any component of  $\vec{E}$  or  $\vec{B}$  for an electromagnetic wave) and denoting it as  $f$ , we can write an ansatz<sup>7</sup>

$$f = ae^{i\psi}, \quad (2.14)$$

---

<sup>7</sup>The terminology *WKB approximation* is also used in the literature for this ansatz, e.g. ref.[97].

where  $a = a(x, t)$  is the amplitude of the particular component and  $\psi = \psi(x, t)$  is the phase of the wave.  $\psi$  is called the *eikonal* and its derivative  $\bar{k}_\mu = -\partial_\mu \psi$  is regarded as the local wavenumber four-vector of the ray, which is normal to the wavefront and tangent to the propagation direction [96]. Since  $\psi$  changes by  $2\pi$  as one transverses one  $\lambda$ , the limit  $\lambda \rightarrow 0$  implies large gradient of  $\psi$ . Now recall that the wave equation is given by:

$$\frac{\partial^2 f}{\partial x_\mu \partial x^\mu} = \left[ \frac{\partial^2 a}{\partial x_\mu \partial x^\mu} e^{i\psi} + 2i \frac{\partial a}{\partial x_\mu} \frac{\partial \psi}{\partial x^\mu} e^{i\psi} + \frac{\partial^2 \psi}{\partial x_\mu \partial x^\mu} f \right] - \frac{\partial \psi}{\partial x_\mu} \frac{\partial \psi}{\partial x^\mu} f = 0. \quad (2.15)$$

Since  $\partial \psi \sim \frac{1}{\lambda}$  is large, we expect the first three terms in square brackets to be subleading compared to the last by a factor of  $\lambda/b$ . Amongst the three terms in the square brackets, since  $\frac{\partial a}{\partial x_\mu}$  is accompanied by a factor of  $\frac{\partial \psi}{\partial x^\mu}$ , the subleading corrections at this order will be dominated by the change in the wave amplitude denoted as  $a$ , which also encodes polarisation data. Thus if we can identify  $\psi$  with  $\delta(b)$ , the  $\lambda/b$  corrections reflect polarisation dependence. We provide an argument in appendix B on why we can identify the eikonal  $\psi$  in eq.(2.14) as the eikonal phase  $\delta(b)$  in eq.(2.5). Indeed in section 4, our explicit results show that

$$\delta_{+,+} = \delta_{-,-} \times [1 + \mathcal{O}(\lambda/b)]. \quad (2.16)$$

that is, the polarization dependence occurs at subleading order in  $\lambda/b$ .

The difference  $2\alpha_3 \equiv (\delta_{+,+} - \delta_{-,-})$ , which characterises the first subleading order in  $\lambda/b$ , has an interpretation as *the rotation angle of the polarisation plane*. To see this, consider positive and negative helicity one-particle states propagating along the  $z$  direction. The two states will acquire opposite phases under a rotation of angle  $\alpha$  around the  $z$ -axes, which is simply a little group rotation.

$$|+\rangle' \rightarrow e^{-i\alpha J^{12}} |+\rangle = e^{-i\hbar\alpha} |+\rangle, \quad |-\rangle' \rightarrow e^{-i\alpha J^{12}} |-\rangle = e^{+i\hbar\alpha} |-\rangle. \quad (2.17)$$

The difference of the phase gained by positive and negative helicity photons ( $|h| = 1$ ) is tied to the rotation angle of the polarisation plane  $\alpha$  by

$$\alpha = -\frac{\delta_{\gamma,+,+} - \delta_{\gamma,-,-}}{2}. \quad (2.18)$$

There is an additional factor of 2 for gravitons ( $|h| = 2$ ),

$$\alpha = -\frac{\delta_{g,+,+} - \delta_{g,-,-}}{4}, \quad (2.19)$$

which compensates for the spin of the graviton. We show that indeed the difference in eikonal phase reproduces the result of ref.[1], where the polarisation plan rotation angle was computed by considering a parallel transport along the light ray's null geodesic. Finally, we see that the angle can be identified with the  $\sigma_3$  component of the eikonal phase matrix:  $\alpha = -\alpha_3/|h|$ .

### 3 The $\mathcal{O}(G^1)$ Compton eikonal phase for spinning objects

We begin with the  $\mathcal{O}(G^1)$  Compton eikonal phase for general spinning body. This will allow us to introduce the setup which will carry over to the  $\mathcal{O}(G^2)$  computation in the next

section. Furthermore, the  $\mathcal{O}(G^1)$  result will be important in checking whether the box and the cross box contributions at  $\mathcal{O}(G^2)$  matches with the square of the former, as required from exponentiation pattern of the eikonal amplitude.

### 3.1 The tree amplitude for eikonal phase

Here we will only consider the helicity conserving configuration  $\widetilde{M}_4(\mathbf{1}2^{+h}\mathbf{3}^{+h}\mathbf{4})$  of the classical amplitude. Here we use  $\widetilde{\phantom{x}}$  to make a distinction between the classical amplitude from “semi”-classical amplitude. From the previous discussion we’ve seen that helicity dependence of the eikonal phase can only be subleading in  $\lambda/b$ , while dimensional analysis tells us that such corrections start at  $\mathcal{O}(G^2)$ . Thus this tell us that at  $\mathcal{O}(G^1)$ , the amplitudes  $\widetilde{M}_4(\mathbf{1}2^{\pm h}\mathbf{3}^{\pm h}\mathbf{4})$  must be identical. Indeed this is verified through our explicit computation. Note that there are also helicity non-preserving amplitudes for the Einstein-Hilbert action [76], which is suppressed by  $\mathcal{O}((\lambda/b)^4)$  compared to its helicity preserving counterpart. Such a high suppression in  $\lambda/b$  also occurs at one-loop, and as we are not interested in  $\lambda/b$  expansion to such high orders, we will solely focus on helicity preserving processes. The gravitational coupling constant is normalised as  $\kappa = \sqrt{32\pi G}$ , where  $G$  is the Newton’s constant.

The eikonal phase is given by the fourier transform of the Compton amplitude in the  $q^2 \rightarrow 0$  limit. At  $\mathcal{O}(G^1)$  this limit is given by the product of two three-point amplitudes, one corresponding to graviton coupling to massive spinning and the other massless helicity states. This can be extracted from the following integral formula, motivated by the exponentiated representation for amplitudes of Kerr black holes [5, 44, 47, 49, 69]<sup>8</sup>

$$M_4^s = M_4^{s=0} \oint \frac{dz}{2\pi iz} \left( \sum_j C_{S^j} z^j \right) \exp(-iK^\mu L^\nu J_{\mu\nu}), \quad (3.1)$$

$$K^\mu = -\frac{q^\mu}{z}, \quad L^\mu = \frac{\langle 3|\sigma^\mu|2\rangle}{\langle 3|p_1|2\rangle},$$

where the *scalar factor*  $M_4^{s=0}$  is the spin-independent part, i.e. it represents the Compton amplitude for two massive scalar and two identical helicity state generated via a graviton exchange. The above can be viewed as a semi-classical amplitude where the spin is taken to the classical limit, i.e.  $s \rightarrow \infty$  and  $\hbar \rightarrow 0$  with  $\hbar s$  held fixed. Indeed taking  $s$  finite, and  $C_{S^j} = 1$ , this reduces to the Compton amplitude for minimally coupled spinning particle [5]. From  $M_4^s$  we can derive the fully classical amplitude  $\widetilde{M}$  by further expanding the kinematic variables with appropriate  $\hbar$  counting.

The *spin factor* refers to the remaining auxiliary residue integral terms in eq.(3.1), where the contour is taken to be encircling the origin counter-clockwise. The spin factor is motivated to capture the full  $t$ -channel (graviton pole) factorization limit, which is modified from the spinless case by the Wilson coefficients  $C_{S^n}$  parametrising the spin-multipole moments [45, 46, 53, 62]. In particular, the  $t = 0$  limit can be attained by either setting

<sup>8</sup>The expression is  $2 \rightarrow 2$  continued from the  $4 \rightarrow 0$  expression. Spinor notations follow that of ref.[45].

$|2\rangle \propto |3\rangle$ , or  $|2\rangle \propto |3\rangle$ . For the former, the scalar factor  $M_4^{s=0}$  factorizes to (in all incoming convention)

$$-tM_4^{s=0} \rightarrow M_3^{+h,-h,+2} \times M_{3-}^{s=0}, \quad (3.2)$$

where  $M_3^{+h,-h,+2}$  is the amplitude of a helicity  $h$  particle coupling to a positive helicity graviton and  $M_{3-}^{s=0} = -\kappa(p_1 \cdot \varepsilon_{3-})^2$  is the amplitude for a massive scalar coupling to a negative helicity graviton. The spin factor in eq.(3.1) becomes

$$K^\mu = -\frac{q^\mu}{z}, \quad L^\mu \rightarrow \frac{\varepsilon_{q-}^\mu}{\varepsilon_{q-} \cdot p_1} \quad \Rightarrow \quad \exp(-iK^\mu L^\nu J_{\mu\nu}) \rightarrow \exp\left(i\frac{q^\mu \varepsilon_{q-}^\nu J_{\mu\nu}}{(p_1 \cdot \varepsilon_{q-})z}\right), \quad (3.3)$$

which reproduces the spin factor of eq.(B.1) in ref.[5].<sup>9</sup> The factorisation behaviour for the case  $|2\rangle \propto |3\rangle$  can be analysed in a similar manner. Although eq.(3.1) has spurious poles due to the factor  $\langle 3|p_1|2\rangle$  in the denominator of the exponent, the spurious poles can be resolved by terms that do not affect the  $t$ -channel pole which determines the  $\mathcal{O}(G^1)$  eikonal phase, therefore only considering the terms in eq.(3.1) is enough to compute the  $\mathcal{O}(G^1)$  eikonal phase.

The scalar amplitudes for massive (uncharged) scalar interacting with massless spin- $|h|$  states through graviton exchange are given as

$$\frac{M_4^{s=0}}{(\kappa/2)^2} = \begin{cases} -\frac{(s-u)^2}{4t} & h=0 \\ -\frac{[2|p_1|3](s-u)}{2t} & h=1/2 \\ -\frac{[2|p_1|3]^2}{t} & h=1 \\ \frac{[2|p_1|3]^3(s-u)}{2(s-m^2)(u-m^2)t} & h=3/2 \\ \frac{[2|p_1|3]^4}{(s-m^2)(u-m^2)t} & h=2 \end{cases}. \quad (3.4)$$

Explicit evaluation of relevant spinor contractions yield


$$\begin{aligned} [2|p_1|3] &= [3|p_1|2] = 2m\omega \sqrt{1 - \frac{|\vec{q}|^2(1 + \frac{2\omega}{m})}{4\omega^2}} = 2m\omega \times [1 + \mathcal{O}(q^2/\omega^2, \hbar)], \\ [23] &= -\langle 23 \rangle = |\vec{q}|, \end{aligned} \quad (3.5)$$

up to phase factors for little group scaling, thus all scalar amplitudes in eq.(3.4) can be treated as

$$M_4^{s=0} \simeq \frac{32\pi Gm^2\omega^2}{\vec{q}^2}, \quad (3.6)$$

---

<sup>9</sup>The  $|2\rangle$  spinors of  $L^\mu$  are interpreted as the auxiliary spinor of  $\varepsilon_{q-}^\mu$ . The relative sign difference is due to outgoing convention for  $q$ .



**Figure 3:** Kinematic set-up of the vectors describing spin orientation.  $\vec{a}$  is the spin-length vector and  $\vec{k}_2$  is the momentum of the incoming massless particle.  $\vec{a}_\perp$  is the projected component of  $\vec{a}$  orthogonal to  $\vec{k}_2$ . The vectors  $\vec{d}$ ,  $\vec{a}_\perp$  and  $\vec{k}_2$  are respectively orthogonal with  $|\vec{d}| = |\vec{a}_\perp|$ .

in the eikonal phase computation, regardless of the helicity. Therefore, the classical part of the eikonal phase at tree order and the observables derived from it are independent of the massless particle's helicity, which can be understood as a manifestation of the equivalence principle.

Before closing we remark that the  $M_4^s$  in eq.(3.1) for  $h = 2$  differs from the classical gravitational Compton amplitude in eq.(B.12) of ref.[5], as the latter was further constrained to have the correct massive factorization channel residues. However for our purpose, the  $\mathcal{O}(G^1)$  eikonal phase, both yield the same result as we demonstrate in appendix C.

### 3.2 The $\mathcal{O}(G^1)$ eikonal phase and observables

Following conventions of refs.[2, 4, 5], we identify the Lorentz generator  $J_{\mu\nu}$  as the spin tensor  $S_{\mu\nu} = -\frac{1}{m}\epsilon_{\mu\nu\alpha\beta}p_1^\alpha S^\beta$ . The exponent in the spin factor eq.(3.1) becomes

$$i\frac{q^\mu\langle 3|\sigma^\nu|2\rangle S_{\mu\nu}}{z\langle 3|p_1|2\rangle} = -\frac{i(n\cdot S)}{m^2\omega z} + \mathcal{O}(|\vec{q}|^2, \hbar^1), \quad (3.7)$$

where  $n^\mu = \epsilon^{\mu\alpha\beta\gamma}p_{1\alpha}k_{2\beta}q_\gamma$  and only the relevant term for  $\mathcal{O}(G^1)$  eikonal phase has been kept. Note that the apparent spurious  $\langle 3|p_1|2\rangle$  pole simply becomes  $\omega$  in the  $q^2 \rightarrow 0$  limit. Furthermore, the  $\omega \rightarrow 0$  limit is non-singular since  $n^\mu$  vanishes. Similar comment applies to our  $\mathcal{O}(G^2)$  analysis. The classical amplitude is then given by:

$$\widetilde{M}_4|_{\mathcal{O}(G^1)} = -\frac{32\pi Gm^2\omega^2}{q^2} \oint \frac{dz}{2\pi iz} \left( \sum_j C_{S^j} z^j \right) \exp\left(-\frac{i(n\cdot S)}{m^2\omega z}\right). \quad (3.8)$$

To obtain the eikonal phase, we Fourier transform to impact parameter space,

$$\delta^{(0)}(b) = \frac{1}{4m\omega} \int \frac{d^{D-2}q}{(2\pi)^{D-2}} \left[ e^{i\vec{b}\cdot\vec{q}} \widetilde{M}_4(q)|_{\mathcal{O}(G^1)} \right], \quad (3.9)$$

which identifies  $\nabla_b \Leftrightarrow i\vec{q}$ . We define the following impact parameter space variables.

$$\vec{a} = \frac{\vec{S}}{m}, \quad \hat{k}_2 = \frac{\vec{k}_2}{\omega}, \quad \vec{d} = \hat{k}_2 \times \vec{a}, \quad \vec{a}_\perp = \vec{a} - (\vec{a} \cdot \hat{k}_2)\hat{k}_2. \quad (3.10)$$

The unit vector  $\hat{k}_2$  defines the direction orthogonal to the impact parameter space ( $\hat{k}_2 \cdot \vec{b} = 0$ ), and  $\vec{a}_\perp$  is the projection of the spin-length vector  $\vec{a}$  onto the impact parameter plane so that  $\vec{a} \cdot \vec{b} = \vec{a}_\perp \cdot \vec{b}$ . Since  $\vec{d} \cdot \vec{a}_\perp = 0$ , the two vectors  $\vec{a}_\perp$  and  $\vec{d}$  span the two-dimensional impact parameter space. These variables are related to the spin Lorentz invariants by

$$\begin{aligned} -\frac{i(n \cdot S)}{m^2\omega} &= (\hat{k}_2 \times \vec{a}) \cdot (i\vec{q}) \Leftrightarrow (\hat{k}_2 \times \vec{a}) \cdot \nabla_b = \vec{d} \cdot \nabla_b \\ -\frac{i(q \cdot S)}{m} &= \vec{a} \cdot (i\vec{q}) \Leftrightarrow \vec{a} \cdot \nabla_b = \vec{a}_\perp \cdot \nabla_b \\ \frac{k_2 \cdot S}{m\omega} &\Leftrightarrow -\hat{k}_2 \cdot \vec{a} \end{aligned} \quad (3.11)$$

The tree order eikonal phase  $\delta^{(0)}$  can be obtained from the tree amplitude eq.(3.1) by expanding eq.(2.5) to leading coupling order,

$$\delta^{(0)}(b) = \frac{1}{4m\omega} \oint \frac{dz}{2\pi iz} \left( \sum_j C_{S^j} z^j \right) \int \frac{d^{D-2}q}{(2\pi)^{D-2}} \left[ e^{i[\vec{b} + (\vec{d}/z)] \cdot \vec{q}} \widetilde{M}_4^{s=0}(q) \right], \quad (3.12)$$

where the order of integration has been changed, and the exponent has been simplified using eq.(3.7). Inserting eq.(3.6) and performing the Fourier integral in  $D = 4$  leads to the eikonal phase

$$\begin{aligned} \delta^{(0)}(b) - \delta_{\text{IR}}^{(0)} &= -\left(\frac{\kappa}{2}\right)^2 \frac{m\omega}{2\pi} \oint \frac{dz}{2\pi iz} \left( \sum_j C_{S^j} z^j \right) \log \left| \vec{b} + \frac{\vec{d}}{z} \right| \\ &= -2Gm\omega \left[ 2 \log b + \frac{2\vec{d} \cdot \vec{b}}{b^2} + C_{S^2} \left( -\frac{2(\vec{d} \cdot \vec{b})^2}{b^4} + \frac{d^2}{b^2} \right) \right. \\ &\quad + C_{S^3} \left( \frac{8(\vec{d} \cdot \vec{b})^3}{3b^6} - \frac{2(\vec{d} \cdot \vec{b})d^2}{b^4} \right) \\ &\quad \left. + C_{S^4} \left( -\frac{4(\vec{d} \cdot \vec{b})^4}{b^8} + \frac{4(\vec{d} \cdot \vec{b})^2 d^2}{b^6} - \frac{d^4}{2b^4} \right) + \dots \right], \quad (3.13) \end{aligned}$$

$$\delta_{\text{IR}}^{(0)} = -\frac{4Gm\omega}{4-D} \quad (3.14)$$

where we have separated the IR divergence  $\delta_{\text{IR}}^{(0)}$ . Setting all Wilson coefficients to Kerr value  $C_{S^n} = 1$  the residue integral localises to  $z = 1$  and we obtain the eikonal phase eq.(5.32) of ref.[98], reproduced below.

$$\delta_{\text{Kerr}}^{(0)}(b) - \delta_{\text{IR}}^{(0)} = -4Gm\omega \log |\vec{b} + \vec{d}|.$$

The time delay can be computed from eq.(3.13) using eq.(1.5), where the IR scale  $b_0 \gg b$  was introduced as the reference so that the time delay becomes a finite quantity [75, 76]

$$\begin{aligned}
\delta t^{(0)} &= -\left(\frac{\kappa}{2}\right)^2 \frac{m}{2\pi} \oint \frac{dz}{2\pi iz} \left( \sum_j C_{S^j} z^j \right) \log \left| \vec{b} + \frac{\vec{d}}{z} \right| \\
&= -2Gm \left[ 2 \log \frac{b}{b_0} + \frac{2\vec{d} \cdot \vec{b}}{b^2} + C_{S^2} \left( -\frac{2(\vec{d} \cdot \vec{b})^2}{b^4} + \frac{d^2}{b^2} \right) \right. \\
&\quad + C_{S^3} \left( \frac{8(\vec{d} \cdot \vec{b})^3}{3b^6} - \frac{2(\vec{d} \cdot \vec{b})d^2}{b^4} \right) \\
&\quad \left. + C_{S^4} \left( -\frac{4(\vec{d} \cdot \vec{b})^4}{b^8} + \frac{4(\vec{d} \cdot \vec{b})^2 d^2}{b^6} - \frac{d^4}{2b^4} \right) + \dots \right]. \quad (3.15)
\end{aligned}$$

The conventional scattering angle is not well-defined in presence of spin, as the motion is no longer restricted to lie on the orbital plane. One way to proceed is to define the scattering angle  $\theta$  as a vectorial quantity  $\vec{\theta}$  that is directed along the direction of the impulse [99]. The impulse for the massless probe  $\vec{q}$  can be computed from the eikonal via

$$\vec{q} = \nabla_b \delta(b), \quad (3.16)$$

therefore the vectorial scattering angle  $\vec{\theta}$  is given as

$$\vec{\theta} := \frac{\nabla_b \bar{\delta}(b, \omega)}{|\vec{k}_2|} = \frac{1}{\omega} \nabla_b \bar{\delta}(b, \omega), \quad (3.17)$$

since the size of the incoming momentum  $\vec{k}_2$  is  $\omega$ . The vectorial scattering angle is


$$\begin{aligned}
\vec{\theta}^{(0)} &= -\left(\frac{\kappa}{2}\right)^2 \frac{m}{2\pi} \oint \frac{dz}{2\pi iz} \left( \sum_j C_{S^j} z^j \right) \frac{\vec{b} + \vec{d}/z}{|\vec{b} + \vec{d}/z|^2} \\
&= -4Gm \left[ \frac{\vec{b}}{b^2} + \left( \frac{\vec{d}}{b^2} - \frac{2(\vec{d} \cdot \vec{b})\vec{b}}{b^4} \right) - C_{S^2} \left( \frac{2(\vec{d} \cdot \vec{b})^2}{b^4} \vec{d} + \frac{b^2 d^2 - 4(\vec{d} \cdot \vec{b})^2}{b^6} \vec{b} \right) \right. \\
&\quad + C_{S^3} \left( \frac{4(\vec{d} \cdot \vec{b})^2 - b^2 d^2}{b^6} \vec{d} + \frac{4(\vec{d} \cdot \vec{b})(b^2 d^2 - 2(\vec{d} \cdot \vec{b})^2)}{b^8} \vec{b} \right) \\
&\quad + C_{S^4} \left( \frac{4(\vec{d} \cdot \vec{b})(b^2 d^2 - 2(\vec{d} \cdot \vec{b})^2)}{b^8} \vec{d} \right. \\
&\quad \left. \left. + \frac{b^4 d^4 - 12b^2 d^2 (\vec{d} \cdot \vec{b})^2 + 16(\vec{d} \cdot \vec{b})^4}{b^{10}} \vec{b} \right) + \dots \right]. \quad (3.18)
\end{aligned}$$

#### 4 The $\mathcal{O}(G^2)$ eikonal phase and observables

In this section, we compute the  $\mathcal{O}(G^2)$  eikonal phase for massless helicity states scattering of a general spinning body. Expanding LHS of eq.(2.5) to  $\mathcal{O}(G^2)$  yields two terms.

$$e^{i\delta} - 1 \Big|_{\mathcal{O}(G^2)} = i\delta^{(1)} - \frac{[\delta^{(0)}]^2}{2}. \quad (4.1)$$





**Figure 4:** The cut diagrams relevant to classical physics. The thick solid lines, wavy lines and double wavy lines represent massive spinning bodies, massless particles and gravitons, respectively. The cut propagators are labeled by the red dotted lines. 1 and 2 are ingoing, 3 and 4 are outgoing, and  $\ell$  labels the loop momentum.

The matching terms on the RHS of eq.(2.5) is the one-loop ( $\mathcal{O}(G^2)$ ) amplitude, which can be expanded on a basis of master scalar integrals, whose coefficients can be computed via generalized unitarity methods [100–104]. Only the scalar triangle integral  $\mathcal{I}_\Delta$  and the box integrals  $\mathcal{I}_\square$  and  $\mathcal{I}_\times$  are relevant for the classical limit [12]

$$\widetilde{M}_4 \Big|_{\mathcal{O}(G^2)} = [c_\Delta \mathcal{I}_\Delta + c_\square \mathcal{I}_\square + c_\times \mathcal{I}_\times] \Big|_{\hbar \rightarrow 0}. \quad (4.2)$$

The contributions from the box integrals,  $c_\square \mathcal{I}_\square + c_\times \mathcal{I}_\times$ , are known to reproduce the iteration terms of the eikonal phase,  $-\delta^{(0)2}/2$ . Therefore for computing the one-loop eikonal phase  $\delta^{(1)}$  it suffices to compute the triangle coefficient  $c_\Delta$  [83, 84]. However, to check consistency, we will also compute the box and cross box coefficients to demonstrate exponentiation.

We will give the leading ( $\lambda/b$ ) corrections from the triangle coefficient as an expansion in the spin  $a/b$ . This will capture the classical gravitational Faraday effect, and reproduce the linear in spin term computed via null geodesic parallel transport computation in ref.[1]. We will give the subleading in  $a/b$  corrections up to  $a^{14}$ . The  $\mathcal{O}(G^2)$  eikonal phase also provides a consistency test to the proposed universal spin-tensor structure conjectured in ref.[6] (and equivalently shift symmetry in ref.[7]) in the black hole limit for the scattering phase.

#### 4.1 Extraction of scalar integral coefficients

It will be more convenient to use the triple cut of the loop integrals as the seed to compute the triangle, box and crossed box coefficients. The triple cut can be used to extract the triangle coefficient and, by imposing one more cut condition, can lead to quadruple-cuts to generate the box and crossed box coefficients. In our case, the triple-cut is formed by gluing a massless-massless-graviton-graviton amplitude to a product of two three-point graviton-massive-massive amplitudes together.

Here we list the relevant tree amplitudes with the ingoing momenta of 1 and 2 and outgoing momenta 3 and 4. The explicit form of the massless-massless-graviton-graviton amplitude is

$$M_4^{EH}(1^{+h}, 2^{+2}, 3^{+2}, 4^{+h}) = \left( -\frac{[24]}{[12]} \right)^{4-2h} \frac{[12]^4 \langle 34 \rangle^4}{stu}, \quad (4.3)$$

where  $\pm h$  are the helicities of the massless particles. The product of two three-point graviton-massive-massive amplitudes can be found in ref.[5],

$$C_4(1^s, 2^+, 3^+, 4^s) \equiv M_3(1^s, 2^+, P)M_3(-P, 3^+, 4^s) = -\frac{[2|p_1|3]^4}{t^2}\mathcal{S}_{\{+++ \}}(1, 2, 3, 4), \quad (4.4)$$

$$C_4(1^s, 2^-, 3^+, 4^s) \equiv M_3(1^s, 2^-, P)M_3(-P, 3^+, 4^s) = -\frac{\langle 23 \rangle^4}{t^2}\mathcal{S}_{\{-+ \}}(1, 2, 3, 4), \quad (4.5)$$

where  $P = p_1 + k_2$  and

$$\mathcal{S}_{\{+++ \}}(1, 2, 3, 4) = \oint \prod_{k=1,2} \frac{dz_k}{2\pi i z_k} \left( \sum_m C_{S^m} z_k^m \right) e^{-i\left(\frac{v_1}{z_1} + \frac{v_2}{z_2}\right)}, \quad (4.6)$$

$$\mathcal{S}_{\{-+ \}}(1, 2, 3, 4) = \oint \prod_{k=1,2} \frac{dz_k}{2\pi i z_k} \left( \sum_m C_{S^m} z_k^m \right) e^{-i\left(\frac{v_1}{z_1} - \frac{v_2^*}{z_2} + \frac{v_3}{z_1 z_2}\right)}, \quad (4.7)$$

$$v_1 = -i(k_2 \cdot S), \quad v_2 = -\frac{i[(2m^2 - t)(k_2 \cdot S) + 2m^2(q \cdot S) + 2i(n \cdot S)]}{2m^2},$$

$$v_3 = \frac{-it(k_2 \cdot S) + 2(n \cdot S)}{2m^2},$$

$v_2^*$  is the complex conjugate of  $v_2$  and the irrelevant quantum contributions from  $f(x)$  factor in ref.[5] were omitted. The  $\mathcal{O}(S^n)$  contribution is obtained by expanding the exponential to  $\mathcal{O}(S^n)$ , and then performing the auxiliary  $z_k$  integrals.

According to the triple-cut diagram shown in fig.4, the cut conditions are

$$\ell^2 = \ell \cdot p_1 = 0, \quad 2\ell \cdot q = -q^2. \quad (4.8)$$

Also, the triple-cut involves two types of cuts, depending on whether the two cut gravitons have equal helicity. For external helicity  $h < 2$  only the opposite helicity cut is non-vanishing, while for gravitons, the equal helicity is non-vanishing for the helicity non-preserving case:

$$C_{\{+++ \}}^{\{h\}} = \sum_{h'=\pm 2} C_4(1^{s_a}, \ell^{+h'}, -\ell_1^{+h'}, 4^{s_a})M_4^{EH}(2^{+h}, \ell_1^{-h'}, -\ell^{-h'}, 3^{+h}) \Big|_{\text{eq.(4.8)}}$$

$$= \frac{[-\text{tr}_-(k_3 \ell p_1 \ell_1)]^{4-2h}}{t^3(2\ell_1 \cdot k_3)(2\ell \cdot k_3)} \left[ \frac{\text{tr}_-(k_2 \ell p_1 \ell_1 k_3 p_1)}{[3|p_1|2]} \right]^{2h} \mathcal{S}_{\{+++ \}}(1, \ell, -\ell_1, 4)$$

$$+ (\ell \longleftrightarrow \ell_1) \Big|_{\text{eq.(4.8)}}, \quad (4.9)$$

$$C_{\{-+ \}}^{\{h=2\}} = \sum_{h'=\pm 2} C_4(1^{s_a}, \ell^{-h'}, -\ell_1^{+h'}, 4^{s_a})M_4^{EH}(2^{-2}, \ell_1^{-h'}, -\ell^{+h'}, 3^{+2}) \Big|_{\text{eq.(4.8)}}$$

$$= \frac{m^4 \langle 23 \rangle^4 t}{(2\ell \cdot k_3)(2\ell_1 \cdot k_3)} \mathcal{S}_{\{-+ \}}(1, \ell, -\ell_1, 4) + (\ell \longleftrightarrow \ell_1) \Big|_{\text{eq.(4.8)}}, \quad (4.10)$$

$$C_{\{-+ \}}^{\{h \neq 2\}} = 0, \quad (4.11)$$

where  $\ell_1 = -\ell - q$ ,  $\text{tr}_-(\dots) = \frac{1}{2}\text{tr}[(1 - \gamma_5)\dots]$  and the subscripts denote the sign of  $\{h_2, h_3\}$ . As mentioned before, we will use these triple-cuts to compute all the relevant integral coefficients in the following subsections.

### 4.1.1 Extraction of box and crossed box coefficients

The box coefficient can be computed from the quadruple cut using unitarity methods [100–104]. We can impose one more cut condition (see fig.4)  $\ell \cdot k_2 = 0$  to get the box cut or  $\ell \cdot k_3 = 0$  to get the crossed box cut. For each quadruple cut, there are two solutions to loop momentum which solve the cut conditions. We use the loop momentum parametrisation of ref.[2] to compute the quadruple cut, with small modifications to account for masslessness of  $k_2^\mu$ .

$$\begin{aligned}\ell^\mu &= \alpha p_1^\mu + \beta k_2^\mu + \gamma q^\mu + \delta \eta^\mu, \\ \eta^\mu &= \langle p_1^\flat | \sigma^\mu | k_2 \rangle, p_1^\flat = p_1 - \frac{m}{2\omega} k_2.\end{aligned}\tag{4.12}$$

$p_1^\flat$  is a null vector computed using Mandelstam parametrisations eq.(2.6). The solutions to the cut conditions,

$$\ell^2 = p_1 \cdot \ell = k_2 \cdot \ell = 0, q \cdot \ell = -\frac{q^2}{2},\tag{4.13}$$

are given as

$$\ell_-^\mu = -\frac{q^2}{2q \cdot \eta} \eta^\mu, \ell_+^\mu = \frac{-(2q^2\omega)p_1^\mu + 2q^2(m + \omega)k_2^\mu - 4m\omega^2 q^\mu}{mq^2 + 2q^2\omega + 4m\omega^2} + \frac{q^2}{2q \cdot \eta} \eta^\mu.\tag{4.14}$$

The coefficient of the box integral can be computed as an average over the two loop momentum solutions  $\ell_\pm$ ,

$$c_{\square, \pm\pm}^{\{h\}} = \frac{1}{2} \sum_{\ell=\ell_\pm} \left[ C_{\square}^{\pm\pm}(\ell, -\ell_1) \mathcal{S}_{\{\pm\pm\}}(1, \ell, -\ell_1, 4) + C_{\square}^{\pm\pm}(\ell_1, -\ell) \mathcal{S}_{\{\pm\pm\}}(1, \ell_1, -\ell, 4) \right]_{\ell \cdot k_2=0},\tag{4.15}$$

where  $C_{\square}^{\pm\pm}$  is the part of  $[C_{\{\pm\pm\}}(-2\ell \cdot k_2)]_{\ell \cdot k_2=0}$  without the spin factor  $\mathcal{S}_{\{\pm\pm\}}$ . The (helicity factor stripped) crossed box coefficient can be obtained from that of the box by the substitution  $k_2 \rightarrow -k_3$ ,  $k_3 \rightarrow -k_2$ ,  $h_2 \rightarrow -h_3$ , and  $h_3 \rightarrow -h_2$ . For example,

$$\left. \frac{c_{\square, ++}^{\{h\}}}{((3|p_1|2])^h} \right|_{\pm k_2 \leftrightarrow \mp k_3} = \frac{c_{\square, --}^{\{h\}}}{((2|p_1|3])^h}.\tag{4.16}$$

Similar relations hold for other helicity sectors such as  $c_{\square, +-}^{\{h\}}$  and  $c_{\square, ++}^{\{h\}}$ .

### 4.1.2 Extraction of triangle coefficients

For the triple-cut as shown in fig.4, the cut-satisfied loop momentum can be parametrized as

$$\ell_\pm^\mu = -\left(\frac{t}{t-4m^2}\right) p_1^\mu + \left(\frac{2m^2}{t-4m^2}\right) q^\mu + z P_\pm^\mu + \left[\frac{m^2 t}{4\gamma_\pm(t-4m^2)}\right] \frac{Q_\pm^\mu}{z},\tag{4.17}$$

where the  $\pm$  labels two independent solutions for the cut conditions and

$$\gamma_{\pm} = \frac{1}{2} \left( -t \pm \sqrt{t(t - 4m^2)} \right). \quad (4.18)$$

The explicit forms of  $P$  and  $Q$  are not important but they satisfy the following relations

$$P_{\pm}^2 = Q_{\pm}^2 = P_{\pm} \cdot p_1 = P_{\pm} \cdot q = Q_{\pm} \cdot p_1 = Q_{\pm} \cdot q = 0, \quad P_{\pm} \cdot Q_{\pm} = 2\gamma_{\pm}, \quad (4.19)$$

which imply a useful formula for the loop computation,

$$P_{\pm}^{\mu} Q_{\pm}^{\nu} = \gamma_{\pm} \eta^{\mu\nu} + \frac{4\gamma_{\pm}}{t - 4m^2} \left[ p_1^{\mu} p_1^{\nu} + \frac{m^2 q^{\mu} q^{\nu}}{t} + \frac{p_1^{\mu} q^{\nu} + q^{\mu} p_1^{\nu}}{2} - \frac{i(2m^2 + \gamma_{\pm})}{2\gamma_{\pm}} \epsilon^{\mu\nu\rho\sigma} (p_1)_{\rho} q_{\sigma} \right]. \quad (4.20)$$

By averaging the residues of the triple-cut with loop momentum solutions eq.(4.17) at  $z = \infty$ , we can obtain the triangle coefficients  $c_{\{+, \pm\}}^{\{h\}}$ , formally as,

$$c_{\{+, \pm\}}^{\{h\}} = \frac{1}{2} \text{Res}_{z=\infty} \left[ \sum_{\ell=\ell_{\pm}} C_{\{+\pm\}}^{\{h\}}(\ell) \right]. \quad (4.21)$$

Note that for the helicity preserving case, we can rewrite eq.(4.9) as

$$C_{\{+, +\}}^{\{h\}} = - \left( \frac{[2\ell]}{[3\ell]} \right)^{2h} \frac{\text{tr} [(1 - \gamma_5) k_3 \ell p_1 \ell_1]^4 \mathcal{S}_{\{++\}}(1, \ell, -\ell_1, 4)}{t^3 (2\ell \cdot k_3) (2\ell_1 \cdot k_3)} \Big|_{\text{eq.(4.8)}} + (\ell \longleftrightarrow \ell_1). \quad (4.22)$$

In the leading order in  $(q/\omega)$ -expansion,  $[2\ell] \simeq [3\ell]$ , so  $C_{\{+, +\}}^{\{h\}}$  is independent of  $|h|$  and this can be understood as a manifestation of the equivalence principle. However, in the subleading order  $C_{\{+, +\}}^{\{h\}}$  is no longer  $|h|$  independent and the equivalence principle ceases to apply.

## 4.2 eikonal phase and its exponentiation

The main contribution to tree order eikonal phase matrix comes from helicity preserving processes, therefore studying  $c_{\square}^{++}$  is enough for checking exponentiation. The box coefficient for the helicity non-preserving case  $c_{\square}^{+-}$  either vanishes ( $|h| < 2$ ) or is suppressed by  $\mathcal{O}((q/\omega)^4)$  compared to  $c_{\square}^{++}$  ( $|h| = 2$ ) which is beyond the regime where exponentiation holds.

An interesting behaviour of the quadruple cut eq.(4.15) is that only one helicity configuration contributes to each loop momentum solution  $\ell_+$  and  $\ell_-$ , thus the sum over all cut solutions and intermediate massless states becomes a sum over two terms

$$c_{\square, ++}^{\{h\}} = \frac{1}{2} \left[ C_{\square}^{++}(\ell_+, -\ell_{1+}) \mathcal{S}_{\{++\}}(1, \ell_+, -\ell_{1+}, 4) + C_{\square}^{++}(\ell_{1-}, -\ell_-) \mathcal{S}_{\{++\}}(1, \ell_{1-}, -\ell_-, 4) \right]_{\ell \cdot k_2=0}, \quad (4.23)$$

where the spin factors are

$$\mathcal{S}_{\{++\}}(\ell_{\pm}) = \oint \prod_{k=1,2} \left[ \frac{dz_k}{2\pi i z_k} \left( \sum_j C_{S^j} z_k^j \right) \right] e^{ib_{\pm}^{\mu\nu} S_{\mu\nu}}, \quad (4.24)$$

where  $\mathcal{S}_{\{++\}}(\ell_+)$  and  $\mathcal{S}_{\{++\}}(\ell_-)$  refer to the spin factors appearing in the first and second line of (4.23) respectively. Note that the *spin factors*  $\mathcal{S}_{\{++\}}(\ell_{\pm})$  are *distinct*. By using the loop momentum solutions (4.14) and the explicit form of spin factor (4.6) subjected to the cut conditions the exponent of the spin factor can be reorganised as

$$ib_{\pm}^{\mu\nu} S_{\mu\nu} = - \left( \frac{1}{z_1} + \frac{1}{z_2} \right) \frac{i(n \cdot S)}{2m^2 \omega} \mp \left( \frac{1}{z_1} - \frac{1}{z_2} \right) \frac{(q \cdot S)}{2m} + \mathcal{O}(|\vec{q}|^2, \hbar^1), \quad (4.25)$$

where the subleading classical terms of order  $\mathcal{O}(|\vec{q}|^2 \hbar^0)$  have been omitted, since they do not contribute to exponentiation. However, the distinct spin factors can be reorganised into a common factor that can be pulled out from the sum eq.(4.23), due to the symmetry of the auxiliary integrals in  $z_k$ ;  $b_{+}^{\mu\nu}$  becomes  $b_{-}^{\mu\nu}$  to the order considered in eq.(4.25), when exchanging the auxiliary variables  $z_1 \leftrightarrow z_2$  for the cut solution  $\ell_+$ . The full box coefficient is now a product of the common spin factor and the scalar factor  $c_{\square, (s=0)}^{++}$ , which is the box integral coefficient for the massive scalar-massless scattering.

$$c_{\square, ++}^{\{h\}} = \oint \prod_{k=1,2} \left[ \frac{dz_k}{2\pi i z_k} \left( \sum_j C_{S^j} z_k^j \right) \right] e^{ib^{\mu\nu} S_{\mu\nu}} \times c_{\square, (s=0)}^{++} [1 + \mathcal{O}(|\vec{q}|^2, \hbar^1)], \quad (4.26)$$

$$c_{\square, (s=0)}^{++} = \frac{C_{\square}^{++}(\ell_+, -\ell_{1+}) + C_{\square}^{++}(\ell_{1-}, -\ell_-)}{2},$$

where we have omitted the subscript in  $b_{-}^{\mu\nu} = b^{\mu\nu}$  for brevity. Up to the considered order, the spin factor  $e^{ib^{\mu\nu} S_{\mu\nu}}$  is the same spin factor eq.(C.2) of the gravitational Compton amplitude eq.(C.1).

Applying the rules (4.16) to eq.(4.26) yields the factorised form of the crossed box coefficient

$$c_{\boxtimes}^{++} = \oint \prod_{k=1,2} \left[ \frac{dz_k}{2\pi i z_k} \left( \sum_j C_{S^j} z_k^j \right) \right] e^{ib^{\mu\nu} S_{\mu\nu}} \times c_{\boxtimes, (s=0)}^{++} [1 + \mathcal{O}(|\vec{q}|^2, \hbar^1)]. \quad (4.27)$$

The box coefficient and the crossed box coefficient for the massive scalar are equal in the classical limit, i.e.

$$c_{\boxtimes, (s=0)}^{++} = c_{\square, (s=0)}^{++} [1 + \mathcal{O}(\hbar^1)], \quad c_{\square, (s=0)}^{++} = 16(\kappa/2)^4 m^4 \omega^4 [1 + \mathcal{O}(|\vec{q}|^2, \hbar^1)], \quad (4.28)$$

which means the sum of eq.(4.26) and eq.(4.27), the full box topology contribution, also factorises into the spin factor and the scalar factor,

$$[c_{\square}^{++} \mathcal{I}_{\square} + c_{\boxtimes}^{++} \mathcal{I}_{\boxtimes}] = \oint \prod_{k=1,2} \left[ \frac{dz_k}{2\pi i z_k} \left( \sum_j C_{S^j} z_k^j \right) \right] e^{ib^{\mu\nu} S_{\mu\nu}} \times c_{\square, (s=0)}^{++} [\mathcal{I}_{\square} + \mathcal{I}_{\boxtimes}], \quad (4.29)$$

up to subleading  $\mathcal{O}(|\vec{q}|^2, \hbar^1)$  terms, which are irrelevant for exponentiation. The validity of approximating box integral contributions by eq.(4.29) has been explicitly checked up to  $\mathcal{O}(S^4)$  order; the difference between the classical box and the classical crossed box coefficient for a specified spin sector appears at quantum subleading order

$$c_{\mathbb{X}}^{++} = c_{\square}^{++} \times [1 + \mathcal{O}(\hbar^1)] . \quad (4.30)$$

The difference between  $c_{\square}^{++}$  and  $c_{\square}^{--}$ , which signifies breaking of exponentiation as  $\alpha_3 = 0$  at tree order, appears at  $\mathcal{O}((q/\omega)^3)$  subleading order.

$$c_{\square}^{--} = c_{\square}^{++} \times [1 + \mathcal{O}((q/\omega)^3, \hbar^1)] . \quad (4.31)$$

The difference term should be added as corrections to one-loop eikonal phase, corresponding to NLO for  $\alpha$  (contributions to  $\alpha^{(1,1)}$ ). Moreover, the  $\mathcal{O}(|\vec{q}|^2 \hbar^0)$  subleading classical terms of the spin factor eq.(4.25) also results in terms not captured by exponentiation. These terms yield NLO corrections to  $\bar{\delta}$  (contributions to  $\bar{\delta}^{(1,1)}$ ).

In impact parameter space, eq.(4.29) scales as  $\propto \omega^2$  and provides the iteration term of tree eikonal phase [76].

$$c_{\square, (s=0)}^{++} [\mathcal{I}_{\square} + \mathcal{I}_{\mathbb{X}}] \Leftrightarrow -\frac{[\delta_{S^0}^{(0)}(b)]^2}{2} . \quad (4.32)$$

The full spinning contribution in impact parameter space can be written as

$$[c_{\square}^{++} \mathcal{I}_{\square} + c_{\mathbb{X}}^{++} \mathcal{I}_{\mathbb{X}}] \Leftrightarrow \oint \prod_{k=1,2} \left[ \frac{dz_k}{2\pi i z_k} \left( \sum_j C_{S^j} z_k^j \right) \right] e^{ib^{\mu\nu} S_{\mu\nu}} \times \left[ -\frac{[\delta_{S^0}^{(0)}(b)]^2}{2} \right] , \quad (4.33)$$

where the spin factor exponent is a derivative operator

$$ib^{\mu\nu} S_{\mu\nu} = \left( \frac{1}{z_1} + \frac{1}{z_2} \right) \frac{\vec{d} \cdot \nabla_b}{2} + \left( \frac{1}{z_1} - \frac{1}{z_2} \right) \frac{i(\vec{a}_{\perp} \cdot \nabla_b)}{2} , \quad (4.34)$$

acting on the “square” of tree eikonal phase. Computation of  $\mathcal{O}(S^n)$  terms follow the same procedure of computing the box coefficients; the exponential is expanded to  $\mathcal{O}(S^n)$ , the  $z_k$  auxiliary integrals are performed, and then the derivative operators  $\nabla_b$  are evaluated.

In the Kerr limit  $C_{S^j} = 1$  the spin factor simplifies to

$$\oint \prod_{k=1,2} \left[ \frac{dz_k}{2\pi i z_k} \left( \sum_j C_{S^j} z_k^j \right) \right] e^{ib^{\mu\nu} S_{\mu\nu}} \xrightarrow{C_{S^j} \rightarrow 1} \exp\left(-\frac{i(n \cdot S)}{m^2 \omega}\right) \Leftrightarrow \exp(\vec{d} \cdot \nabla_b) , \quad (4.35)$$

which is a constant shift by  $\vec{d} = \hat{k}_2 \times \vec{a}$  in impact parameter space. This is consistent with the fact that the tree eikonal phase for Kerr is obtained from the spinless case by a constant shift in impact parameter space [98].

$$\delta_{\text{Kerr}}^{(0)}(\vec{b}) = \delta_{\text{Scalar}}^{(0)}(\vec{b} + \vec{d}) . \quad (4.36)$$

### 4.3 results

We are interested in the helicity-preserving parts of the eikonal phase in the classical limit, which we decompose as

$$\bar{\delta} := \frac{\delta_{+,+} + \delta_{-,-}}{2}, \quad \alpha := \frac{\delta_{-,-} - \delta_{+,+}}{2|h|}. \quad (4.37)$$

The superscripts and subscripts for  $\delta(b)$  defined in eq.(2.9) will be used to denote the same perturbative expansions of  $\bar{\delta}$  and  $\alpha$ . As explained in section 2.2,  $\alpha$  is identified as the polarisation plane rotation angle. For all spin sectors we find

$$\frac{\alpha}{\bar{\delta}} \sim \mathcal{O}\left(\frac{1}{\omega b}\right), \quad (4.38)$$

confirming the expectation eq.(2.16) based on geometric optics analysis.

The leading order terms of  $\bar{\delta}$  and  $\alpha$  have the dependence

$$\text{LO:} \quad \bar{\delta}_{\text{LO}} \sim \frac{G^2 m^2 \omega}{b}, \quad \alpha_{\text{LO}} \sim \frac{G^2 m^2}{b^2} \quad (4.39)$$

where each is multiplied by dimensionless combinations of spin vectors, impact parameter and momenta. The results are independent of helicity, which can be understood as the equivalence principle in the geometric optics limit. In the Kerr limit  $C_{S^n} = 1$ , we find that only particular spin structures appear, which can be connected to the shift symmetry eq.(1.1) considered by Bern et.al. [7]. Furthermore, the leading terms  $\bar{\delta}_{\text{LO}}$  were checked to be consistent with eq.(5.38) of ref.[98]. We report triangle coefficients  $c_{\text{LO}}$  together with  $\bar{\delta}_{\text{LO}}$  and  $\alpha_{\text{LO}}$  to  $\mathcal{O}(S^6)$  in the main text, and report higher spin order terms in the ancillary file.

The NLO terms have the dependence

$$\text{NLO:} \quad \bar{\delta}_{\text{NLO}} \sim \frac{G^2 m^2}{b^2}, \quad \alpha_{\text{NLO}} \sim \frac{G^2 m^2}{b^3 \omega}, \quad (4.40)$$

which do not depend on  $|h|$ ; the equivalence principle still holds. However, the NLO terms are purely imaginary in impact parameter space, and new spin structures that are inconsistent with the shift symmetry eq.(1.1) appear in the Kerr limit. Together with the triangle coefficients  $c_{\bar{\delta}/\alpha}$ , we also report residual contributions from box topology  $c_{\square}^{\bar{\delta}/\alpha}$  that contribute to  $\bar{\delta}_{\text{NLO}}$  and  $\alpha_{\text{NLO}}$ , up to  $\mathcal{O}(S^2)$  in the main text and to  $\mathcal{O}(S^4)$  in appendix D.

#### 4.3.1 leading order

The triangle coefficients are normalised by  $iM_4 = (\kappa/2)^4 c_{\Delta} \mathcal{I}_{\Delta} + \dots$ . Our results can be summarized into the defined coefficient average  $c_{\bar{\delta}} = \frac{c_{++} + c_{--}}{2}$  and coefficient difference  $c_{\alpha} = \frac{c_{--} - c_{++}}{2|h|}$ . Both  $c_{\bar{\delta}}$  and  $c_{\alpha}$  are independent of  $|h|$  at this order.

To linear order in spin,

$$\begin{aligned} c_{\bar{\delta},\text{LO}}^{\text{S}^1} &= 20im^2\omega(n \cdot S), & c_{\alpha,\text{LO}}^{\text{S}^1} &= 5m^3q^2(k_2 \cdot S), \\ \bar{\delta}_{\text{LO}}^{\text{S}^1} &= -\frac{5\pi G^2 m^2 \omega (\vec{d} \cdot \vec{b})}{b^3}, & \alpha_{\text{LO}}^{\text{S}^2} &= \frac{5\pi G^2 m^2 (\hat{k}_2 \cdot \vec{a})}{4b^3}, \end{aligned} \quad (4.41)$$

non-vanishing coefficients in BH limit	vanishing coefficients in BH limit
$Z_{2,1} = 1 + C_{S^2}$	$Z_{2,2} = 1 - C_{S^2}$
$Z_{3,1} = 3C_{S^2} + C_{S^3}$	$Z_{3,2} = C_{S^2} - C_{S^3}$
$Z_{4,1} = 3C_{S^2}^2 + 4C_{S^3} + C_{S^4}$	$Z_{4,3} = C_{S^2}^2 - C_{S^4}$ $Z_{4,4} = 3C_{S^2}^2 - 4C_{S^3} + C_{S^4}$
$Z_{5,1} = 10C_{S^2}C_{S^3} + 5C_{S^4} + C_{S^5}$	$Z_{5,2} = 2C_{S^2}C_{S^3} - C_{S^4} - C_{S^5}$ $Z_{5,3} = 2C_{S^2}C_{S^3} - 3C_{S^4} + C_{S^5}$
$Z_{6,1} = 10C_{S^3}^2 + 15C_{S^2}C_{S^4} + 6C_{S^5} + C_{S^6}$	$Z_{6,2} = 2C_{S^3}^2 + C_{S^2}C_{S^4} - 2C_{S^5} - C_{S^6}$ $Z_{6,3} = 10C_{S^3}^2 - 15C_{S^2}C_{S^4} + 6C_{S^5} - C_{S^6}$ $Z_{6,4} = 2C_{S^3}^2 - C_{S^2}C_{S^4} - 2C_{S^5} + C_{S^6}$

**Table 1:** The table provides the specific combination of Wilson coefficients used in the results of  $\bar{\delta}$  and  $\alpha$ . Note that  $Z_{4,2}$  appears in [7] which does not appear in our results.

where  $\alpha$  matches with the null geodesic parallel transport computation of ref.[1].

At quadratic order in spin

$$\begin{aligned}
c_{\bar{\delta},\text{LO}}^{S^2} &= \frac{5}{16} \left\{ Z_{2,1} [19(n \cdot S)^2 + 7m^2 q^2 (k_2 \cdot S)^2] + 12Z_{2,2} m^2 \omega^2 (q \cdot S)^2 \right\}, \\
\bar{\delta}_{\text{LO}}^{S^2} &= \frac{5\pi G^2 m^2 \omega}{64} \left\{ Z_{2,1} \left[ 19 \left( \frac{3(\vec{d} \cdot \vec{b})^2 - d^2 b^2}{b^5} \right) - 7 \frac{(\hat{k}_2 \cdot \vec{a})^2}{b^3} \right] \right. \\
&\quad \left. + 12 Z_{2,2} \left[ \frac{3(\vec{a} \cdot \vec{b})^2 - a_{\perp}^2 b^2}{b^5} \right] \right\}, \\
c_{\alpha,\text{LO}}^{S^2} &= -\frac{15i Z_{2,1} m q^2 (n \cdot S)(k_2 \cdot S)}{8\omega}, \quad \alpha_{\text{LO}}^{S^2} = -\frac{45 Z_{2,1} \pi G^2 m^2 (\hat{k}_2 \cdot \vec{a})(\vec{d} \cdot \vec{b})}{32b^5}. \tag{4.42}
\end{aligned}$$

At cubic order in spin

$$\begin{aligned}
c_{\bar{\delta},\text{LO}}^{S^3} &= -\frac{i(n \cdot S)}{8m^2 \omega} [Z_{3,1} \{9(n \cdot S)^2 + 7m^2 q^2 (k_2 \cdot S)^2\} + 20Z_{3,2} m^2 \omega^2 (q \cdot S)^2], \\
\bar{\delta}_{\text{LO}}^{S^3} &= -\frac{3\pi G^2 m^2 \omega (\vec{d} \cdot \vec{b})}{32} \left\{ Z_{3,1} \left[ 9 \left( \frac{5(\vec{d} \cdot \vec{b})^2 - 3d^2 b^2}{b^7} \right) - 7 \left( \frac{(\hat{k}_2 \cdot \vec{a})^2}{b^5} \right) \right] \right. \\
&\quad \left. + 20Z_{3,2} \left[ \frac{5(\vec{a} \cdot \vec{b})^2 - a_{\perp}^2 b^2}{b^7} \right] \right\}, \\
c_{\alpha,\text{LO}}^{S^3} &= -\frac{q^2 (k_2 \cdot S)}{96m\omega^2} [Z_{3,1} \{39(n \cdot S)^2 - 7m^2 q^2 (k_2 \cdot S)^2\} + 60Z_{3,2} m^2 \omega^2 (q \cdot S)^2], \\
\bar{\alpha}_{\text{LO}}^{S^3} &= \frac{3\pi G^2 m^2 (\hat{k}_2 \cdot \vec{a})}{128} \left\{ Z_{3,1} \left[ 13 \left( \frac{5(\vec{d} \cdot \vec{b})^2 - d^2 b^2}{b^7} \right) + 7 \frac{(\hat{k}_2 \cdot \vec{a})^2}{b^5} \right] \right. \\
&\quad \left. + 20Z_{3,2} \left[ \frac{5(\vec{a} \cdot \vec{b})^2 - a_{\perp}^2 b^2}{b^7} \right] \right\}. \tag{4.43}
\end{aligned}$$



At quartic order in spin

$$\begin{aligned}
c_{\delta, \text{LO}}^{\text{S}^4} &= \frac{-1}{1536m^4\omega^2} \left\{ Z_{4,1} [294m^2q^2(n \cdot S)^2(k_2 \cdot S)^2 - 21m^4q^4(k_2 \cdot S)^4 + 239(n \cdot S)^4] \right. \\
&\quad \left. + 60Z_{4,3}m^2\omega^2(q \cdot S)^2 [19(n \cdot S)^2 + 7m^2q^2(k_2 \cdot S)^2] + 120Z_{4,4}m^4\omega^4(q \cdot S)^4 \right\}, \\
\bar{\delta}_{\text{LO}}^{\text{S}^4} &= \frac{3\pi G^2 m^2 \omega}{256} \left\{ \frac{7Z_{4,1}}{4} \left[ -7(\hat{k}_2 \cdot \vec{a})^2 \left( \frac{5(\vec{d} \cdot \vec{b})^2 - d^2 b^2}{b^7} \right) \right. \right. \\
&\quad \left. \left. - \frac{3(\hat{k}_2 \cdot \vec{a})^4}{2b^5} + \frac{239}{42} \left( \frac{35(\vec{d} \cdot \vec{b})^4 - 30(\vec{d} \cdot \vec{b})^2 d^2 b^2 + 3d^4 b^4}{b^9} \right) \right] \right. \\
&\quad \left. + \frac{5Z_{4,3}}{2} \left[ -7(\hat{k}_2 \cdot \vec{a})^2 \left( \frac{5(\vec{a} \cdot \vec{b})^2 - a_{\perp}^2 b^2}{b^7} \right) \right. \right. \\
&\quad \left. \left. + 19 \left( \frac{35(\vec{a} \cdot \vec{b})^2 (\vec{d} \cdot \vec{b})^2 - 5(\vec{a} \cdot \vec{b})^2 d^2 b^2 - 5(\vec{d} \cdot \vec{b})^2 a_{\perp}^2 b^2 + d^2 a_{\perp}^2 b^4}{b^9} \right) \right] \right. \\
&\quad \left. + 5Z_{4,4} \left[ \frac{35(\vec{a} \cdot \vec{b})^4 - 30(\vec{a} \cdot \vec{b})^2 a_{\perp}^2 b^2 + 3a_{\perp}^4 b^4}{b^9} \right] \right\}, \tag{4.44}
\end{aligned}$$

$$\begin{aligned}
c_{\alpha, \text{LO}}^{\text{S}^4} &= \frac{iq^2(n \cdot S)(k_2 \cdot S)}{384m^3\omega^3} \left\{ Z_{4,1} [23(n \cdot S)^2 - 7m^2q^2(k_2 \cdot S)^2] + 90Z_{4,3}m^2\omega^2(q \cdot S)^2 \right\}, \\
\alpha_{\text{LO}}^{\text{S}^4} &= -\frac{5\pi G^2 m^2 (\hat{k}_2 \cdot \vec{a})(\vec{d} \cdot \vec{b})}{512} \left\{ Z_{4,1} \left[ 23 \left( \frac{7(\vec{d} \cdot \vec{b})^2 - 3d^2 b^2}{b^9} \right) + 21 \frac{(\hat{k}_2 \cdot \vec{a})^2}{b^7} \right] \right. \\
&\quad \left. + 90Z_{4,3} \left[ \frac{7(\vec{a} \cdot \vec{b})^2 - a_{\perp}^2 b^2}{b^9} \right] \right\}. \tag{4.45}
\end{aligned}$$

At quintic order in spin

$$\begin{aligned}
c_{\delta, \text{LO}}^{\text{S}^5} &= \frac{i(n \cdot S)}{768m^6\omega^3} \left\{ Z_{5,1} [-3m^4q^4(k_2 \cdot S)^4 + 22m^2q^2(k_2 \cdot S)^2(n \cdot S)^2 + 13(n \cdot S)^4] \right. \\
&\quad \left. + 12Z_{5,2}m^2\omega^2(q \cdot S)^2 [7m^2q^2(k_2 \cdot S)^2 + 9(n \cdot S)^2] + 40Z_{5,3}m^4\omega^4(q \cdot S)^4 \right\}, \\
\bar{\delta}_{\text{LO}}^{\text{S}^5} &= -\frac{5\pi G^2 m^2 \omega (\vec{d} \cdot \vec{b})}{1024b^{11}} \left\{ Z_{5,1} \left[ -9b^4(\vec{k}_2 \cdot \vec{a})^4 + 22(\vec{k}_2 \cdot \vec{a})^2 (3b^4 d^2 - 7b^2(\vec{d} \cdot \vec{b})^2) \right. \right. \\
&\quad \left. \left. + 13(-70b^2 d^2(\vec{d} \cdot \vec{b})^2 + 63(\vec{d} \cdot \vec{b})^4 + 15b^4 d^4) \right] \right. \\
&\quad \left. + 12Z_{5,2} \left[ 7(\vec{k}_2 \cdot \vec{a})^2 (a_{\perp}^2 b^4 - 7b^2(\vec{a} \cdot \vec{b})^2) - 189(\vec{a} \cdot \vec{b})^2 (b^2 d^2 - 3(\vec{d} \cdot \vec{b})^2) \right. \right. \\
&\quad \left. \left. + 9a_{\perp}^2 b^2 (3b^2 d^2 - 7(\vec{d} \cdot \vec{b})^2) \right] - 120Z_{5,3} (14a_{\perp}^2 b^2(\vec{a} \cdot \vec{b})^2 - 21(\vec{a} \cdot \vec{b})^4 - a_{\perp}^4 b^4) \right\}, \\
c_{\alpha, \text{LO}}^{\text{S}^5} &= \frac{q^2(k_2 \cdot S)}{15360m^5\omega^4} \left\{ Z_{5,1} [9m^4q^4(k_2 \cdot S)^4 - 50m^2q^2(k_2 \cdot S)^2(n \cdot S)^2 + 105(n \cdot S)^4] \right. \\
&\quad \left. + 20Z_{5,2}m^2\omega^2(q \cdot S)^2 [39(n \cdot S)^2 - 7m^2q^2(k_2 \cdot S)^2] + 200m^4\omega^4 Z_{5,3}(q \cdot S)^4 \right\}, \\
\alpha_{\text{LO}}^{\text{S}^5} &= \frac{15\pi G^2 m^2 (\vec{k}_2 \cdot \vec{a})}{4096b^{11}} \left\{ Z_{5,1} \left[ 9b^4(\vec{k}_2 \cdot \vec{a})^4 - 10(\vec{k}_2 \cdot \vec{a})^2 (b^4 d^2 - 7b^2(\vec{d} \cdot \vec{b})^2) \right. \right.
\end{aligned}$$

$$\begin{aligned}
& +21 \left( -14b^2 d^2 (\vec{d} \cdot \vec{b})^2 + 21(\vec{d} \cdot \vec{b})^4 + b^4 d^4 \right) \Big] \\
& - 4Z_{5,2} \left[ 7(\vec{k}_2 \cdot \vec{a})^2 \left( a_\perp^2 b^4 - 7b^2 (\vec{a} \cdot \vec{b})^2 \right) + 91(\vec{a} \cdot \vec{b})^2 \left( b^2 d^2 - 9(\vec{d} \cdot \vec{b})^2 \right) \right. \\
& \left. - 13a_\perp^2 b^2 \left( b^2 d^2 - 7(\vec{d} \cdot \vec{b})^2 \right) \right] + 40Z_{5,3} \left( a_\perp^4 b^4 - 14a_\perp^2 b^2 (\vec{a} \cdot \vec{b})^2 + 21(\vec{a} \cdot \vec{b})^4 \right) \Big\} . \quad (4.46)
\end{aligned}$$

At sextic order in spin

$$\begin{aligned}
c_{\delta, \text{LO}}^{\text{S}^6} &= \frac{1}{491520m^8 \omega^4} \left\{ Z_{6,1} \left[ 33m^6 q^6 (k_2 \cdot S)^6 - 345m^4 q^4 (k_2 \cdot S)^4 (n \cdot S)^2 \right. \right. \\
& \quad \left. \left. + 1615m^2 q^2 (k_2 \cdot S)^2 (n \cdot S)^4 + 745(n \cdot S)^6 \right] \right. \\
& - 40Z_{6,2} m^2 \omega^2 (q \cdot S)^2 \left[ 21m^4 q^4 (k_2 \cdot S)^4 - 294m^2 q^2 (k_2 \cdot S)^2 (n \cdot S)^2 - 239(n \cdot S)^4 \right] \\
& \left. + 320Z_{6,3} m^6 \omega^6 (q \cdot S)^6 + 400Z_{6,4} m^4 \omega^4 (q \cdot S)^4 \left[ 7m^2 q^2 (k_2 \cdot S)^2 + 19(n \cdot S)^2 \right] \right\} , \\
\bar{\delta}_{\text{LO}}^{\text{S}^6} &= -\frac{15\pi G^2 m^2 \omega}{131072b^{13}} \left\{ Z_{6,1} \left[ 33b^6 (\vec{k}_2 \cdot \vec{a})^6 - 69(\vec{k}_2 \cdot \vec{a})^4 \left( b^6 d^2 - 7b^4 (\vec{d} \cdot \vec{b})^2 \right) \right. \right. \\
& \quad \left. \left. + 323(\vec{k}_2 \cdot \vec{a})^2 \left( -14b^4 d^2 (\vec{d} \cdot \vec{b})^2 + 21b^2 (\vec{d} \cdot \vec{b})^4 + b^6 d^4 \right) \right. \right. \\
& \quad \left. \left. + 149 \left( -105b^4 d^4 (\vec{d} \cdot \vec{b})^2 + 315b^2 d^2 (\vec{d} \cdot \vec{b})^4 - 231(\vec{d} \cdot \vec{b})^6 + 5b^6 d^6 \right) \right] \right. \\
& + 8Z_{6,2} \left[ 98(\vec{k}_2 \cdot \vec{a})^2 \left( a_\perp^2 b^4 \left( b^2 d^2 - 7(\vec{d} \cdot \vec{b})^2 \right) - 7(\vec{a} \cdot \vec{b})^2 \left( b^4 d^2 - 9b^2 (\vec{d} \cdot \vec{b})^2 \right) \right) \right. \\
& - 21(\vec{k}_2 \cdot \vec{a})^4 \left( a_\perp^2 b^6 - 7b^4 (\vec{a} \cdot \vec{b})^2 \right) + 239 \left( a_\perp^2 b^2 \left( -14b^2 d^2 (\vec{d} \cdot \vec{b})^2 + 21(\vec{d} \cdot \vec{b})^4 + b^4 d^4 \right) \right. \\
& \left. \left. - 7(\vec{a} \cdot \vec{b})^2 \left( -18b^2 d^2 (\vec{d} \cdot \vec{b})^2 + 33(\vec{d} \cdot \vec{b})^4 + b^4 d^4 \right) \right) \right] \\
& + 64Z_{6,3} \left[ -105a_\perp^4 b^4 (\vec{a} \cdot \vec{b})^2 + 315a_\perp^2 b^2 (\vec{a} \cdot \vec{b})^4 - 231(\vec{a} \cdot \vec{b})^6 + 5a_\perp^6 b^6 \right] \\
& + 80Z_{6,4} \left[ 19 \left( -14a_\perp^2 b^2 (\vec{a} \cdot \vec{b})^2 \left( b^2 d^2 - 9(\vec{d} \cdot \vec{b})^2 \right) + 21(\vec{a} \cdot \vec{b})^4 \left( b^2 d^2 - 11(\vec{d} \cdot \vec{b})^2 \right) \right. \right. \\
& \left. \left. + a_\perp^4 b^4 \left( b^2 d^2 - 7(\vec{d} \cdot \vec{b})^2 \right) \right) + 7(\vec{k}_2 \cdot \vec{a})^2 \left( -14a_\perp^2 b^4 (\vec{a} \cdot \vec{b})^2 + 21b^2 (\vec{a} \cdot \vec{b})^4 + a_\perp^4 b^6 \right) \right] \Big\} , \\
c_{\alpha, \text{LO}}^{\text{S}^6} &= -\frac{iq^2 (k_2 \cdot S)(n \cdot S)}{737280m^7 \omega^5} \left\{ 3600Z_{6,4} m^4 \omega^4 (q \cdot S)^4 \right. \\
& \quad \left. + Z_{6,1} \left[ 81m^4 q^4 (k_2 \cdot S)^4 - 290m^2 q^2 (k_2 \cdot S)^2 (n \cdot S)^2 + 465(n \cdot S)^4 \right] \right. \\
& \quad \left. + 240Z_{6,2} m^2 \omega^2 (q \cdot S)^2 \left[ 23(n \cdot S)^2 - 7m^2 q^2 (k_2 \cdot S)^2 \right] \right\} , \\
\alpha_{\text{LO}}^{\text{S}^6} &= -\frac{105\pi G^2 m^2 (\vec{k}_2 \cdot \vec{a})(\vec{d} \cdot \vec{b})}{65536b^{13}} \left\{ Z_{6,1} \left[ 27b^4 (\vec{k}_2 \cdot \vec{a})^4 - 58(\vec{k}_2 \cdot \vec{a})^2 \left( b^4 d^2 - 3b^2 (\vec{d} \cdot \vec{b})^2 \right) \right. \right. \\
& \quad \left. \left. + 31 \left( -30b^2 d^2 (\vec{d} \cdot \vec{b})^2 + 33(\vec{d} \cdot \vec{b})^4 + 5b^4 d^4 \right) \right] \right. \\
& + 16Z_{6,2} \left[ 23 \left( (\vec{a} \cdot \vec{b})^2 \left( 33(\vec{d} \cdot \vec{b})^2 - 9b^2 d^2 \right) + a_\perp^2 b^2 \left( b^2 d^2 - 3(\vec{d} \cdot \vec{b})^2 \right) \right) \right. \\
& \left. \left. - 7(\vec{k}_2 \cdot \vec{a})^2 \left( a_\perp^2 b^4 - 9b^2 (\vec{a} \cdot \vec{b})^2 \right) \right] + 240Z_{6,4} \left[ a_\perp^4 b^4 - 18a_\perp^2 b^2 (\vec{a} \cdot \vec{b})^2 + 33(\vec{a} \cdot \vec{b})^4 \right] \right\} . \quad (4.47)
\end{aligned}$$

### 4.3.2 subleading order

The residuals from the box contribution not cancelled by iteration of tree contribute to subleading  $\bar{\delta}$  and  $\alpha$ . The coefficients are normalised by  $iM_4 = (\kappa/2)^4 c_{\square} [\mathcal{I}_{\square} + \mathcal{I}_{\boxtimes}] + \dots$

with coefficient average  $c_{\square}^{\bar{\delta}} = \frac{c_{\square}^{++} + c_{\square}^{--}}{2}$  and coefficient difference  $c_{\square}^{\bar{\alpha}} = \frac{c_{\square}^{--} - c_{\square}^{++}}{2|h|}$ . All integral coefficients ( $c_{\bar{\delta}}$ ,  $c_{\alpha}$ ,  $c_{\square}^{\bar{\delta}}$ , and  $c_{\square}^{\bar{\alpha}}$ ) are independent of  $|h|$  at this order.

Note that  $\bar{\delta}$  and  $\alpha$  are purely imaginary at this order, which are written as a sum of two terms. The first part containing  $\pi$  is from the triangle coefficients and the second part without  $\pi$  is from the box coefficients.

At linear order in spin

$$\begin{aligned} c_{\bar{\delta},\text{NLO}}^{\text{S}^1} &= 0, \quad \bar{c}_{\square,\text{NLO}}^{\text{S}^1,\bar{\delta}} = 4im^2q^2\omega(n \cdot S), \quad \bar{\delta}_{\text{NLO}}^{\text{S}^1} = 0 + i \left[ \frac{8G^2m^2(\vec{d} \cdot \vec{b})}{b^4} \right], \\ c_{\alpha,\text{NLO}}^{\text{S}^1} &= \frac{5m^3q^2(q \cdot S)}{2}, \quad c_{\square,\text{NLO}}^{\text{S}^1,\alpha} = 2m^3q^4(k_2 \cdot S), \\ \alpha_{\text{NLO}}^{\text{S}^1} &= i \left[ \frac{15\pi G^2m^2(\vec{a} \cdot \vec{b})}{8b^5\omega} \right] - i \left[ \frac{8G^2m^2(\hat{k}_2 \cdot \vec{a})}{b^4\omega} \right]. \end{aligned} \quad (4.48)$$

At quadratic order in spin

$$\begin{aligned} c_{\bar{\delta},\text{NLO}}^{\text{S}^2} &= \frac{35(1 + C_{\text{S}^2})m^2q^2(q \cdot S)(k_2 \cdot S)}{16}, \\ \bar{c}_{\square,\text{NLO}}^{\text{S}^2,\bar{\delta}} &= q^2(C_{\text{S}^2} + 1)(m^2q^2(k_2 \cdot S)^2 + 2(n \cdot S)^2), \\ \frac{\bar{\delta}_{\text{NLO}}^{\text{S}^2}}{iG^2m^2} &= -\frac{105(1 + C_{\text{S}^2})\pi(\hat{k}_2 \cdot \vec{a})(\vec{a} \cdot \vec{b})}{64b^5} + \frac{4(C_{\text{S}^2} + 1)[b^2a^2 - 4(\vec{d} \cdot \vec{b})^2]}{b^6}, \\ c_{\alpha,\text{NLO}}^{\text{S}^2} &= -\frac{15i(1 + C_{\text{S}^2})mq^2(q \cdot S)(n \cdot S)}{16\omega}, \\ c_{\square,\text{NLO}}^{\text{S}^1,\alpha} &= -\frac{imq^4(C_{\text{S}^2} + 1)(k_2 \cdot S)(n \cdot S)}{\omega}, \\ \frac{\omega \alpha_{\text{NLO}}^{\text{S}^2}}{iG^2m^2} &= -\frac{225(1 + C_{\text{S}^2})\pi(\vec{a} \cdot \vec{b})(\vec{d} \cdot \vec{b})}{64b^7} + \frac{16(C_{\text{S}^2} + 1)(\vec{d} \cdot \vec{b})(\hat{k}_2 \cdot \vec{a})}{b^6}. \end{aligned} \quad (4.49)$$

$a^2 = d^2 + (\hat{k}_2 \cdot \vec{a})^2 = a_{\perp}^2 + (\hat{k}_2 \cdot \vec{a})^2$  has been used to simplify  $\bar{\delta}_{\text{NLO}}^{\text{S}^2}$ .

#### 4.4 Discussions: equivalence principle, spin structures and massless limits

In the geometric optics limit, the observables considered in this paper—the observables eq.(1.5) derived from  $\bar{\delta}$  and the rotation angle  $\alpha$ —has geometric interpretations in classical general relativity which do not depend on worldline degrees of freedom. This means the observables do not depend on the details of the particle, which can be understood as a manifestation of the equivalence principle. For the computations considered in this paper, the equivalence principle is realised as the independence of  $\bar{\delta}$  and  $\alpha$  from helicity  $|h|$  of the massless particle. The term equivalence principle will be used to denote this helicity independence when referring to eikonal phase variables.

Since  $\lambda/b$  is the classical expansion parameter that parametrises deviations from the geometric optics limit, we expect the equivalence principle to be obeyed by  $\bar{\delta}$  and  $\alpha$  at leading order in  $\lambda/b$ . Indeed this is the case. The LO  $\bar{\delta}$  and  $\alpha$  obeys the equivalence

principle, at least up to one-loop order. The general expectation is that the equivalence principle for  $\bar{\delta}$  and  $\alpha$  is broken at sufficiently high orders in  $\lambda/b$ , since the corrections in  $\lambda/b$  can be interpreted as wave property corrections to the point particle approximation; the equivalence principle only applies to the strict point particle limit where tidal effects can be neglected, while a characteristic property of waves is that they cannot be localised to a point. Interestingly, the equivalence principle is still obeyed at NLO for both  $\bar{\delta}$  and  $\alpha$ .

### spin structures

All LO triangle coefficients  $c_{\bar{\delta},\text{LO}}$  and  $c_{\alpha,\text{LO}}$  have the special property that their dependence on  $(q \cdot S)$  vanish in the Kerr limit  $C_{S^j} \rightarrow 1$ . As discussed in the introduction, this echos the observed spin operator dependence for the 2 PM conservative potential in the black hole limit [5], where in the basis of

$$\{(q \cdot S_i), (p_j \cdot S_i), (n \cdot S_i)\} \quad (4.50)$$

only the latter two appears. However, naive application of the shift proposed in ref.[7]

$$S_i^\mu \rightarrow S_i^\mu + \xi_i \frac{q^\mu}{q^2}, \quad (4.51)$$

no longer leads to a symmetry in our massless-massive scattering. This is because while the change in  $(p_2 \cdot S_1)$  under the shift is a *quantum* effect, it is classical for  $(k_2 \cdot S_1)$ . To see this note that  $(p_2 \cdot S_1) \sim \mathcal{O}(\hbar^{-1})$  is shifted by  $\frac{p_2 \cdot q}{q^2} \sim \mathcal{O}(\hbar^0)$  where we've used momentum conservation  $(p_j \cdot q) = \pm q^2/2$ ;  $\frac{(p_j \cdot q)/q^2}{(p_j \cdot S_i)} = \mathcal{O}(\hbar^1)$ , thus the shift is a quantum effect and the classical result is invariant. On the other hand,  $(k_2 \cdot S) \sim \mathcal{O}(\hbar^0)$  while it is shifted by  $\frac{k_2 \cdot q}{q^2} \sim \mathcal{O}(\hbar^0)$ , i.e. a classical effect. Since the original definition was for the massive-massive scattering, the observation here does not contradict the conjecture in ref.[7].

Note that vanishing of  $(q \cdot S)$  dependence for the triangle coefficients is a LO property; all NLO triangle coefficients  $c_{\bar{\delta},\text{NLO}}$  and  $c_{\alpha,\text{NLO}}$  has *linear dependence* on  $(q \cdot S)$  in the Kerr limit. However, the Kerr limit is still special in that all higher order dependences on  $(q \cdot S)$  vanish.

### leading order terms as the massless limit

It is known that the leading order  $\bar{\delta}_{\text{LO}}$  can be obtained as the massless limit of the massive case.<sup>10</sup> Denoting the massive eikonal phase as  $\chi$ , the explicit map is

$$\bar{\delta}_{\text{LO}} = \lim_{m_2 \rightarrow 0} \chi_{S_1^n S_2^0} \Big|_{m_2 \sigma = \omega}. \quad (4.52)$$

The subscript denotes that we substitute  $\sigma = \omega/m_2$  and then take the  $m_2 \rightarrow 0$  limit. We also find that  $\alpha_{\text{LO}}$  can be obtained as the massless limit of the massive case. We first align the spin of the massive probe particle  $m_2$  so that the probe particle has a well-defined helicity. In the rest frame of  $m_1$ , we can write the relevant four-vectors as

$$p_1^\mu = (m_1, \vec{0}), p_2^\mu = (\omega, k\hat{z}), S_2^\mu = (hk/\omega, h\hat{z}), \quad (4.53)$$

---

<sup>10</sup>JWK would like to thank Francesco Comberiati and Leonarde de la Cruz for pointing this out.

where  $\omega^2 = (k^2 + m_2^2)$  and  $h$  is the helicity defined by  $h = \frac{\vec{p}_2 \cdot \vec{S}_2}{|\vec{p}_2|}$ . The aligned configuration sets  $n \cdot S_2 = 0$ , and “factoring” the helicity  $h$  by setting  $p_1 \cdot S_2 = -m_1 \sigma$  we find that  $\alpha$  corresponds to the massless limit of  $\chi_{S_2^1}$ .<sup>11</sup>

$$\alpha_{\text{LO}} = \lim_{m_2 \rightarrow 0} \chi_{S_1^n S_2^1} \Big|_{m_2 \sigma = \omega, n \cdot S_2 = 0, p_1 \cdot S_2 = -\frac{m_1 \omega}{m_2}}. \quad (4.54)$$

The limit has been found by comparing to the results given in appendix B of ref.[5]. It would be interesting to link the NLO imaginary terms of the eikonal phase variables to massive eikonal phase.

### purely imaginary NLO terms

Both  $\bar{\delta}_{\text{NLO}}$  and  $\alpha_{\text{NLO}}$  are purely imaginary in impact parameter space. Imaginary terms of the eikonal phase are usually interpreted as signs of non-conservative effects such as dissipation or radiation. However, the purely imaginary NLO terms  $\bar{\delta}_{\text{NLO}}$  and  $\alpha_{\text{NLO}}$  cannot be interpreted this way, as  $2 \rightarrow 2$  scattering at one-loop order is purely elastic. On the other hand, the imaginary term is NLO in  $\lambda/b$ , which is the order the wave amplitude  $a$  correction enters the geometric optics eikonal  $\psi$  as can be seen in eq.(2.15). Therefore a possible interpretation is that the imaginary terms encode (de)amplification of the waves due to spin of the massive body, which may have connections with polarising effects discussed in section 3.3 of ref.[98].

## 5 Conclusion

In this paper, we derive the leading order gravitational Faraday effect,  $\alpha := -\alpha_3/|h|$  for polarized electromagnetic and gravitational waves using on-shell scattering amplitude. The result is given as an expansion in the ratio of spin-length and impact parameter,  $a/b$ . While the linear term for Kerr black hole matches with result in [1], we provide the result for general spinning body up to 14th order in spin. We also give the  $\mathcal{O}(G^2)$  eikonal phase  $\bar{\delta}$  for Compton scattering, from which one can derive the scattering angle and Shapiro time delay. We note that in both cases, the leading term satisfies the equivalence principle, as well as the vanishing  $(q \cdot S)$  terms in the Lorentz invariant expansion basis

$$\{(q \cdot S), (k \cdot S), \epsilon(q, k, p, S)\} \quad (5.1)$$

for the black hole limit, observed in the massive scattering amplitude up to quartic order in spin [2, 4, 5], and conjectured to hold to all orders [6–8]. Naive extension of the shift symmetry eq.(1.1) proposed in ref.[2] no longer holds here. Since the original discussion was strictly for massive-massive scattering, this leaves open the tantalizing possibility of modifying the shift to account for massless kinematics.

Although the equivalence principle still applies, both of the spin structure properties no longer hold for NLO corrections in  $\lambda/b$  which are purely imaginary in position space. Given that the expansion in  $\lambda/b$  is an expansion around the geometric optics approximation, the

<sup>11</sup>This scaling keeps Pauli-Lubanski vector  $W_2^\mu = -\frac{1}{2} \epsilon^{\mu\alpha\beta\gamma} k_{2\alpha} J_{\beta\gamma} = m_2 S_2^\mu$  finite in the massless limit.

violation can be interpreted as the breakdown of eikonal approximation. Indeed we see that at this order, the overall box topology contribution no longer matches the square of the tree eikonal phase as required from exponentiation. It would be interesting nonetheless to explore whether sensible observables can be extracted from these terms.

## Acknowledgments

We are grateful for Zvi Bern, Andreas Brandhuber, Kays Haddad, Michèle Levi, Andrés Luna, Donal O’Connell, Gabriele Travaglini, Radu Roiban and Justin Vines for discussions and valuable feedback on the first draft of the manuscript. JWK would like to thank Manuel Accattulli Huber and Stefano De Angelis for sharing computation details of ref.[76] and Andrés Luna for sharing unlisted box coefficients of ref.[4] for comparison. WMC is supported by in part by JSPS KAKENHI Grant Number 21F21317. MZC and YTH is supported by MoST Grant No. 109-2112-M-002 -020 -MY3. JWK was supported by the Science and Technology Facilities Council (STFC) Consolidated Grant ST/T000686/1 “*Amplitudes, Strings and Duality*”.

## A Eikonal phase normalisation

We examine the different conventions and resulting differences in the definition of the eikonal phase. In the  $2 \rightarrow 2$  convention for the  $T$ -matrix element  $M_4$ , where  $S = \mathbb{1} + iT$ , the definitions for the eikonal phase used in the literature generically takes the form

$$\begin{aligned} e^{i\delta(b)} - 1 &:= \int \frac{d^D q}{(2\pi)^D} (2\pi)\delta[V \cdot q] (2\pi)\delta[W \cdot q] e^{-ib \cdot q} iM_4 \\ &= \int \frac{d^{D-2} q}{(2\pi)^{D-2}} \frac{i}{\sqrt{|V^2 W^2 - (V \cdot W)^2|}} \left[ e^{i\vec{b} \cdot \vec{q}} M_4(q) \right], \end{aligned} \quad (\text{A.1})$$

where  $q^\mu = p_3^\mu - p_2^\mu$  and different definitions differ by the vectors  $V^\mu$  and  $W^\mu$  used to define the impact parameter space. For strict exponentiation of the classical terms at the diagram level the vectors are defined as [94]

$$V = p_4 + p_1, \quad W = p_3 + p_2, \quad (\text{A.2})$$

and the independent variable is average three-momentum  $\vec{p}$ . The geometric interpretation of eikonal variables in Fig.1 of ref.[105] becomes natural for this choice of  $V^\mu$  and  $W^\mu$ . In centre of momentum (COM) frame the momenta are given as

$$\begin{aligned} p_1^\mu &= (\sqrt{m_1^2 + \vec{p}^2 + \vec{q}^2/4}, -\vec{p} + \vec{q}/2), \quad p_4^\mu = (\sqrt{m_1^2 + \vec{p}^2 + \vec{q}^2/4}, -\vec{p} - \vec{q}/2), \\ p_2^\mu &= (\sqrt{m_2^2 + \vec{p}^2 + \vec{q}^2/4}, \vec{p} - \vec{q}/2), \quad p_3^\mu = (\sqrt{m_2^2 + \vec{p}^2 + \vec{q}^2/4}, \vec{p} + \vec{q}/2), \end{aligned} \quad (\text{A.3})$$

with  $\vec{p} \cdot \vec{q} = 0$ . The square of the Jacobian factor is

$$(V \cdot W)^2 - V^2 W^2 = 16|\vec{p}|^2 \left[ \sqrt{m_1^2 + \vec{p}^2 + \vec{q}^2/4} + \sqrt{m_2^2 + \vec{p}^2 + \vec{q}^2/4} \right]^2. \quad (\text{A.4})$$

For both massive case ( $m_1 \neq 0$  and  $m_2 \neq 0$ ) we rescale  $\vec{q} \rightarrow \hbar\vec{q}$ . Only the  $q$ -dependent terms carry extra powers of  $\hbar^2$  and the terms can be dropped in the classical limit.

$$(V \cdot W)^2 - V^2W^2 = 16|\vec{p}|^2 \left[ \sqrt{m_1^2 + \vec{p}^2} + \sqrt{m_2^2 + \vec{p}^2} \right]^2 + \mathcal{O}(\hbar^2). \quad (\text{A.5})$$

For  $m_2 = 0$  we rescale  $\vec{p} \rightarrow \hbar\vec{p}$  and  $\vec{q} \rightarrow \hbar\vec{q}$ . The classical limit is given as

$$(V \cdot W)^2 - V^2W^2 = 16\hbar^2 m_1^2 |\vec{p}|^2 [1 + \mathcal{O}(\hbar^2)]. \quad (\text{A.6})$$

When comparing with the Mandelstam parametrisation using  $\omega$ , we have the relation

$$\hbar m_1 \omega = p_1 \cdot p_2 \quad \Rightarrow \quad \omega = |\vec{p}| \sqrt{1 + \frac{\vec{q}^2}{4\vec{p}^2}} [1 + \mathcal{O}(\hbar^1)]. \quad (\text{A.7})$$

Therefore to the first subleading order in  $q/\omega$  or  $q/\bar{p}$ , the two ratios can be considered equal  $q/\omega = q/\bar{p} \times [1 + \mathcal{O}((q/\bar{p})^2)]$ .

We use a simpler prescription used in the literature [76, 84]

$$V = 2p_1, W = 2p_2, (V \cdot W)^2 - V^2W^2 = 16m_1^2 m_2^2 (\sigma^2 - 1), \quad (\text{A.8})$$

where  $\sigma := \frac{p_1 \cdot p_2}{m_1 m_2}$ . A slightly different parametrisation for the COM frame momenta is more useful for this prescription:

$$\begin{aligned} p_1^\mu &= (\sqrt{m_1^2 + \vec{p}^2}, -\vec{p}), p_4^\mu = (\sqrt{m_1^2 + \vec{p}^2}, -\vec{p} - \vec{q}), \\ p_2^\mu &= (\sqrt{m_2^2 + \vec{p}^2}, \vec{p}), p_3^\mu = (\sqrt{m_2^2 + \vec{p}^2}, \vec{p} + \vec{q}), \end{aligned} \quad (\text{A.9})$$

which yields the squared Jacobian factor

$$(V \cdot W)^2 - V^2W^2 = 16m_1^2 m_2^2 (\sigma^2 - 1) = 16|\vec{p}|^2 \left[ \sqrt{m_1^2 + \vec{p}^2} + \sqrt{m_2^2 + \vec{p}^2} \right]^2, \quad (\text{A.10})$$

which is exact. In the classical limit, this factor is equivalent to the factor eq.(A.5) up to the loop order considered in this paper if  $\bar{p}$  is identified with  $p$ . For  $m_2 = 0$  the factor simplifies to

$$(V \cdot W)^2 - V^2W^2 = 16\hbar^2 m_1^2 \omega^2, \quad (\text{A.11})$$

which gives the normalisation factor used in eq.(2.5). The difference of normalisation factors eq.(A.6) and eq.(A.11) starts at  $\mathcal{O}((q/\omega)^2)$  order, therefore the two prescriptions are indistinguishable up to the first subleading order in  $q/\omega$ .

## B Linking eikonal phase to geometric optics

We present an argument why the eikonal phase  $\delta(b)$  can be identified as the geometric optics eikonal  $\psi$ , which can be viewed as a simplified version of the arguments presented

in ref.[106]. A related approach is to identify the eikonal phase as the Hamilton’s principal function for the massive case [31, 94, 107, 108]. We also comment that Hamiltonian mechanics was developed from the correspondence between the eikonal and Hamilton’s principal function.

To build the connection, we identify the field component  $f$  in eq.(2.14) as the following matrix element in a specified gauge, e.g. radiation gauge

$$ae^{i\psi} = f = \langle \vec{p}' | A^\mu(x) | \vec{p}; \vec{k}, \varepsilon \rangle, \quad (\text{B.1})$$

where  $\vec{p}$  is the momentum of the incoming massive particle,  $\vec{p}'$  is the momentum of the outgoing massive particle, and  $\vec{k}$  is the momentum of the incoming photon with polarisation  $\varepsilon$ . This matrix element can be converted to  $2 \rightarrow 2$  scattering amplitude of a photon scattering off a massive particle by the LSZ reduction formula

$$M_4(p; k, \varepsilon; k', \varepsilon'; p') = \int d^d x e^{ik'_\mu x^\mu} [\varepsilon'_\mu]^* \partial^\nu \partial_\nu \langle \vec{p}' | A^\mu(x) | \vec{p}; \vec{k}, \varepsilon \rangle, \quad (\text{B.2})$$

which can be “inverted” to obtain an asymptotic form of the matrix element in eq.(B.1)

$$\langle \vec{p}' | A_\mu(x) | \vec{p}; \vec{k}, \varepsilon \rangle \simeq \int \frac{d^d k'}{(2\pi)^d} \varepsilon'_\mu M_4(p; k, \varepsilon; k', \varepsilon'; p') e^{-ik'_\mu x^\mu}, \quad (\text{B.3})$$

where  $A_4$  is the scattering amplitude. The “sum” over  $k'$  can be decomposed as a sum over transferred momentum  $q = k' - k$  and the “forward momentum”  $\bar{k} = (k + k')/2$ , which we set as  $\bar{k}^\mu = (\omega, \bar{k}_z \hat{z})$ . Combining with the identification eq.(B.1), eq.(B.3) is then recast as

$$ae^{i\psi} \simeq \int \frac{d\omega d\bar{k}_z}{(2\pi)^2} \int \frac{d^{d-2}q}{(2\pi)^{d-2}} J e^{-i(\omega t - \bar{k}_z z)} \left[ e^{i\vec{b} \cdot \vec{q}} M_4(q) \right]. \quad (\text{B.4})$$

where  $\vec{b}$  is the impact parameter space coordinate<sup>12</sup> and  $J$  denotes the unimportant terms; the on-shell condition deltas, the Jacobian, and the polarisation vector. Note that the sign of the Fourier factor  $e^{i\vec{b} \cdot \vec{q}}$  originates from the positive frequency plane wave factor  $e^{-ik'_\mu x^\mu}$  with outgoing momentum  $k' = k + q$ . Using eq.(2.5), eq.(B.4) can be recast as

$$e^{i\psi + [\log a]} \simeq \int \frac{d\omega d\bar{k}_z}{(2\pi)^2} [\delta_{\text{o-s}}] e^{-i(\omega t - \bar{k}_z z) + i\delta(\vec{b}) + [\log N']}, \quad (\text{B.5})$$

where  $N' = J/N$  is a slowly varying factor and  $\delta_{\text{o-s}}$  gives the on-shell conditions. In other words, we can approximate the eikonal  $\psi$  of geometric optics as

$$\psi(t, z, \vec{b}) - i[\log a] \simeq -\omega t + \bar{k}_z z + \delta(\vec{b}) - i[\log N']. \quad (\text{B.6})$$

where terms in the square brackets are “small” compared to other terms. The polarisation dependence is given by the ratio  $\partial[\log a]/\partial\psi$ , and estimating the amplitude variation length scale as  $\partial[\log a] \sim b^{-1}$  we have the ratio  $\partial[\log a]/\partial\psi \sim \lambda/b$ .

<sup>12</sup>In this case, the coordinate values of  $x^\mu$  in eq.(B.3) except the time  $t$  and  $z$  component.



## C Simplifying the spin factor for long-distance physics

The ansatz for the classical amplitude, eq.(3.1), is tailored to correctly capture the  $t$ -channel residue. While the  $t$ -channel is the only physical channel for  $h \leq 1$ , this is no longer true for  $h > 1$  and  $s$ - and  $u$ -channels will also become physical. For example, the gravitational Compton amplitude eq.(B.12) of ref.[5], reproduced below with  $2 \rightarrow 2$  continuation from  $4 \rightarrow 0$ , has a different spin factor from eq.(3.1).

$$M_4^s = M_4^{s=0} \oint \prod_{k=1,2} \left[ \frac{dz_k}{2\pi i z_k} \left( \sum_j C_{S^j} z_k^j \right) \right] \exp(-i\bar{K}^\mu L^\nu J_{\mu\nu}), \quad (\text{C.1})$$

$$\bar{K}^\mu = \frac{k_2^\mu}{z_1} - \frac{k_3^\mu}{z_2}, \quad L^\mu = \frac{\langle 3|\sigma^\mu|2\rangle}{\langle 3|p_1|2\rangle}.$$

The scalar factor  $A_4^{s=0}$  is the  $h = 2$  case of eq.(3.4), and the integration contours of the spin factor encircle the origins  $z_{1,2} = 0$  counter-clockwise. While the spin factor is motivated to capture the  $s$ - and  $u$ -channel poles, it also correctly captures the  $t$ -channel pole. This is because in the  $|2\rangle \otimes |3\rangle$  limit ( $|2\rangle \otimes |3\rangle$  limit) the vector  $L^\mu$  becomes proportional to  $k_2^\mu$  ( $k_3^\mu$ ), so the  $z_1$  ( $z_2$ ) dependence of  $\bar{K}^\mu$  drops out and the spin factor of eq.(C.1) effectively reduces to that of eq.(3.1).

The gravitational Compton amplitude eq.(C.1) was motivated by the fact that all physical channel poles are correctly captured, therefore it will only differ from the “real” Compton amplitude by unknown polynomial terms, at least up to  $\mathcal{O}(J^4)$  since there are no spurious poles. Since polynomial terms become (derivatives of) Dirac delta interactions in impact parameter space and thus cannot affect long-distance physics, it is concluded that the amplitude eq.(C.1) correctly captures the long-distance physics of graviton scattering from a spinning compact object. The conclusion poses a mystery; while eq.(3.1) is *linear* in Wilson coefficients, eq.(C.1) is *quadratic*. How can their long-distance behaviours be equivalent?

In fact, there is a nontrivial cancellation for eq.(C.1) and the long-distance behaviour reduces to that of eq.(3.1). To simplify the analysis, we recast the spin factor exponent of eq.(C.1) in a form similar to eq.(3.7). Substituting the Lorentz generator  $J^{\mu\nu}$  by the spin tensor  $S^{\mu\nu}$  and using the definition  $q^\mu = k_3^\mu - k_2^\mu$ ,

$$-i\bar{K}^\mu L^\nu S_{\mu\nu} = -\left(\frac{1}{z_1} + \frac{1}{z_2}\right) \frac{i(n \cdot S)}{2m^2\omega} + \left(\frac{1}{z_1} - \frac{1}{z_2}\right) \frac{(q \cdot S)}{2m} + \mathcal{O}(|\vec{q}|^2, \hbar^1). \quad (\text{C.2})$$

Due to the antisymmetry of  $z_1$  and  $z_2$ , any terms with odd powers of  $(q \cdot S)$  become irrelevant and only even powers of  $(q \cdot S)$  need to be considered. The even powers of  $(q \cdot S)$  can be traded for even powers of  $(n \cdot S)$

$$\left[\frac{(q \cdot S)}{m}\right]^2 = \left[-\frac{i(n \cdot S)}{m^2\omega}\right]^2 + \mathcal{O}(q^2), \quad (\text{C.3})$$

which is equivalent to the relation  $(\vec{d} \cdot \nabla_b)^2 = a_\perp^2 \nabla_b^2 - (\vec{a}_\perp \cdot \nabla_b)^2$  in impact parameter space. The Laplacian cannot contribute to the long-distance behaviour at tree level, therefore it can be ignored in the analysis.

We now focus on the (effective) coefficient of  $(n \cdot S)^N$  term when the exponentials in eq.(3.1) and eq.(C.1) are expanded to  $\mathcal{O}(S^N)$  order. We will ignore the irrelevant  $(N!)^{-1}$  factor from the exponential function in both cases. For eq.(3.1) we get

$$\oint \frac{dz}{2\pi iz} \left( \sum_j C_{S^j} z^j \right) \left( -\frac{i(n \cdot S)}{zm^2\omega} \right)^N = C_{S^N} \left( -\frac{i(n \cdot S)}{m^2\omega} \right)^N. \quad (\text{C.4})$$

For eq.(C.1) the expansion is rather involved, as all even power contributions from  $(q \cdot S)$  must be included. Using the binomial theorem, collecting even powers of  $(q \cdot S)$ , and using eq.(C.3) to reorganise the expression as  $(n \cdot S)^N$ , the relevant  $\mathcal{O}(S^N)$  term from eq.(C.1) can be written as

$$\oint \prod_{k=1,2} \left[ \frac{dz_k}{2\pi iz_k} \left( \sum_j C_{S^j} z_k^j \right) \right] g_N(z_1, z_2) \left( -\frac{i(n \cdot S)}{m^2\omega} \right)^N, \quad (\text{C.5})$$

$$g_N(z_1, z_2) := \frac{1}{2^N} \sum_{M=0}^{\lfloor N/2 \rfloor} \binom{N}{2M} \left( \frac{1}{z_1} + \frac{1}{z_2} \right)^{N-2M} \left( \frac{1}{z_1} - \frac{1}{z_2} \right)^{2M},$$

where  $\lfloor x \rfloor$  is the floor function. The sum in  $g_N(z_1, z_2)$  can be explicitly evaluated.

$$\begin{aligned} g_N(z_1, z_2) &= \frac{1}{2^N} \sum_{J=0}^N \frac{1 + (-1)^J}{2} \binom{N}{J} \left( \frac{1}{z_1} + \frac{1}{z_2} \right)^{N-J} \left( \frac{1}{z_1} - \frac{1}{z_2} \right)^J \\ &= \frac{1}{2^{N+1}} \left[ \sum_{J=0}^N \binom{N}{J} \left( \frac{1}{z_1} + \frac{1}{z_2} \right)^{N-J} \left( \frac{1}{z_1} - \frac{1}{z_2} \right)^J \right. \\ &\quad \left. + \sum_{J=0}^N \binom{N}{J} \left( \frac{1}{z_1} + \frac{1}{z_2} \right)^{N-J} \left( \frac{1}{z_2} - \frac{1}{z_1} \right)^J \right] \\ &= \frac{1}{2} \left( \frac{1}{z_1^N} + \frac{1}{z_2^N} \right). \end{aligned} \quad (\text{C.6})$$

Therefore eq.(C.5) becomes

$$\oint \prod_{k=1,2} \left[ \frac{dz_k}{2\pi iz_k} \left( \sum_j C_{S^j} z_k^j \right) \right] \left( \frac{1}{2z_1^N} + \frac{1}{2z_2^N} \right) \left( -\frac{i(n \cdot S)}{m^2\omega} \right)^N = C_{S^N} \left( -\frac{i(n \cdot S)}{m^2\omega} \right)^N, \quad (\text{C.7})$$

which is the same as eq.(C.4). In conclusion, the ansatz eq.(3.1) may be used in place of the ‘‘correct’’ gravitational Compton amplitude eq.(C.1) for evaluating  $\mathcal{O}(G^1)$  eikonal phase.

## D Eikonal phase at NLO to $\mathcal{O}(S^4)$

The relation  $a^2 = d^2 + (\hat{k}_2 \cdot \vec{a})^2 = a_\perp^2 + (\hat{k}_2 \cdot \vec{a})^2$  was used to simplify some expressions.

## D.1 Cubic order in spin

$$c_{\delta, \text{NLO}}^{\text{S}^3} = -\frac{7iq^2(3C_{\text{S}^2} + C_{\text{S}^3})(q \cdot S)(k_2 \cdot S)(n \cdot S)}{8\omega}, \quad (\text{D.1})$$

$$\bar{c}_{\square, \text{NLO}}^{\text{S}^3, \delta} = -\frac{iq^2(n \cdot S)}{2m^2\omega} [(C_{\text{S}^3} + 3C_{\text{S}^2})\{m^2q^2(k_2 \cdot S)^2 + (n \cdot S)^2\} \\ + (C_{\text{S}^2} - C_{\text{S}^3})m^2\omega^2(q \cdot S)^2], \quad (\text{D.2})$$

$$\frac{\bar{\delta}_{\text{NLO}}^{\text{S}^3}}{iG^2m^2} = \frac{105\pi(3C_{\text{S}^2} + C_{\text{S}^3})(\hat{k}_2 \cdot \vec{a})(\vec{d} \cdot \vec{b})(\vec{a} \cdot \vec{b})}{32b^7} \\ + \frac{4(\vec{d} \cdot \vec{b})}{b^8} [(C_{\text{S}^3} + 3C_{\text{S}^2})\{6(\vec{d} \cdot \vec{b})^2 - b^2(d^2 + 2a^2)\} \\ - (C_{\text{S}^2} - C_{\text{S}^3})\{b^2a_{\perp}^2 - 6(\vec{a} \cdot \vec{b})^2\}], \quad (\text{D.3})$$

$$c_{\alpha, \text{NLO}}^{\text{S}^3} = -\frac{q^2(q \cdot S)}{64m\omega^2} [(3C_{\text{S}^2} + C_{\text{S}^3})\{13(n \cdot S)^2 - 7m^2q^2(k_2 \cdot S)^2\} \\ - 20(C_{\text{S}^3} - C_{\text{S}^2})m^2\omega^2(q \cdot S)^2], \quad (\text{D.4})$$

$$\bar{c}_{\square, \text{NLO}}^{\text{S}^3, \alpha} = -\frac{q^4(k_2 \cdot S)((C_{\text{S}^2} - C_{\text{S}^3})m^2\omega^2(q \cdot S)^2 + (C_{\text{S}^3} + 3C_{\text{S}^2})(n \cdot S)^2)}{4m\omega^2}, \quad (\text{D.5})$$

$$\frac{\omega \alpha_{\text{NLO}}^{\text{S}^3}}{iG^2m^2} = -\frac{75\pi(\vec{a} \cdot \vec{b})}{64b^9} [(C_{\text{S}^3} - C_{\text{S}^2})\{7(\vec{a} \cdot \vec{b})^2 - 3a_{\perp}^2b^2\} \\ - \frac{(3C_{\text{S}^2} + C_{\text{S}^3})}{20}\{21(\hat{k}_2 \cdot \vec{a})^2b^2 + 13(7(\vec{d} \cdot \vec{b})^2 - d^2b^2)\}] \\ + \frac{4(\hat{k}_2 \cdot \vec{a})}{b^8} [(C_{\text{S}^3} + 3C_{\text{S}^2})\{b^2d^2 - 6(\vec{d} \cdot \vec{b})^2\} \\ + (C_{\text{S}^2} - C_{\text{S}^3})\{b^2a_{\perp}^2 - 6(\vec{a} \cdot \vec{b})^2\}]. \quad (\text{D.6})$$

## D.2 Quartic order in spin

$$c_{\delta, \text{NLO}}^{\text{S}^4} = -\frac{7q^2(q \cdot S)(k_2 \cdot S)}{256m^2\omega^2} [10(C_{\text{S}^2}^2 - C_{\text{S}^4})m^2\omega^2(q \cdot S)^2 \\ + (3C_{\text{S}^2}^2 + 4C_{\text{S}^3} + C_{\text{S}^4})\{7(n \cdot S)^2 - m^2q^2(k_2 \cdot S)^2\}], \quad (\text{D.7})$$

$$\bar{c}_{\square, \text{NLO}}^{\text{S}^4, \delta} = -\frac{q^2}{24m^4\omega^2} [(C_{\text{S}^4} + 4C_{\text{S}^3} + 3C_{\text{S}^2}^2)(n \cdot S)^2\{2(n \cdot S)^2 + 3m^2q^2(k_2 \cdot S)^2\} \\ + 3(C_{\text{S}^2}^2 - C_{\text{S}^4})m^2\omega^2(q \cdot S)^2\{2(n \cdot S)^2 + m^2q^2(k_2 \cdot S)^2\}], \quad (\text{D.8})$$

$$\frac{\bar{\delta}_{\text{NLO}}^{\text{S}^4}}{iG^2m^2} = -\frac{105\pi(\hat{k}_2 \cdot \vec{a})(\vec{a} \cdot \vec{b})}{512b^9} [5(C_{\text{S}^2}^2 - C_{\text{S}^4})\{7(\vec{a} \cdot \vec{b})^2 - 3a_{\perp}^2b^2\} \\ + \frac{(3C_{\text{S}^2}^2 + 4C_{\text{S}^3} + C_{\text{S}^4})}{2}\{3(\hat{k}_2 \cdot \vec{a})^2b^2 + 7(7(\vec{d} \cdot \vec{b})^2 - d^2b^2)\}] \\ - \frac{2}{b^{10}} [(C_{\text{S}^2}^2 - C_{\text{S}^4})\{6(\vec{a} \cdot \vec{b})^2(8(\vec{d} \cdot \vec{b})^2 - b^2a^2) + b^2a_{\perp}^2(b^2a^2 - 6(\vec{d} \cdot \vec{b})^2)\} \\ + (C_{\text{S}^4} + 4C_{\text{S}^3} + 3C_{\text{S}^2}^2)\{b^4d^2a^2 - 6b^2(\vec{d} \cdot \vec{b})^2(d^2 + a^2) + 16(\vec{d} \cdot \vec{b})^4\}], \quad (\text{D.9})$$

$$c_{\alpha, \text{NLO}}^{\text{S}^4} = \frac{iq^2(q \cdot S)(n \cdot S)}{768m^3\omega^3} [(3C_{\text{S}^2}^2 + 4C_{\text{S}^3} + C_{\text{S}^4})\{23(n \cdot S)^2 - 21m^2q^2(k_2 \cdot S)^2\}$$

$$+ 90(C_{S^2}^2 - C_{S^4})m^2\omega^2(q \cdot S)^2] , \quad (\text{D.10})$$

$$\bar{c}_{\square, \text{NLO}}^{S^4, \alpha} = \frac{iq^4(k_2 \cdot S)(n \cdot S)}{24m^3\omega^3} \left[ (C_{S^4} + 4C_{S^3} + 3C_{S^2}^2)(n \cdot S)^2 + 3m^2\omega^2(C_{S^2}^2 - C_{S^4})(q \cdot S)^2 \right] , \quad (\text{D.11})$$

$$\begin{aligned} \frac{\omega \alpha_{\text{NLO}}^{S^4}}{iG^2m^2} = & -\frac{105\pi(\vec{d} \cdot \vec{b})(\vec{a} \cdot \vec{b})}{512b^{11}} \left[ 45(C_{S^2}^2 - C_{S^4}) \{3(\vec{a} \cdot \vec{b})^2 - a_{\perp}^2 b^2\} \right. \\ & \left. + \frac{(3C_{S^2}^2 + 4C_{S^3} + C_{S^4})}{2} \left\{ 21(\hat{k}_2 \cdot \vec{a})^2 b^2 + 23 \left( 3(\vec{d} \cdot \vec{b})^2 - d^2 b^2 \right) \right\} \right] \\ & + \frac{4(\hat{k}_2 \cdot \vec{a})(\vec{d} \cdot \vec{b})}{b^{10}} \left[ (C_{S^4} + 4C_{S^3} + 3C_{S^2}^2) \left( 8(\vec{d} \cdot \vec{b})^2 - 3b^2 d^2 \right) \right. \\ & \left. - 3(C_{S^2}^2 - C_{S^4}) \left( b^2 a_{\perp}^2 - 8(\vec{a} \cdot \vec{b})^2 \right) \right] . \quad (\text{D.12}) \end{aligned}$$

## References

- [1] H. Ishihara, M. Takahashi and A. Tomimatsu, *GRAVITATIONAL FARADAY ROTATION INDUCED BY KERR BLACK HOLE*, *Phys. Rev. D* **38** (1988) 472.
- [2] Z. Bern, A. Luna, R. Roiban, C.-H. Shen and M. Zeng, *Spinning black hole binary dynamics, scattering amplitudes, and effective field theory*, *Phys. Rev. D* **104** (2021) 065014 [[2005.03071](#)].
- [3] R. Aoude, K. Haddad and A. Helset, *Tidal effects for spinning particles*, *JHEP* **03** (2021) 097 [[2012.05256](#)].
- [4] D. Kosmopoulos and A. Luna, *Quadratic-in-spin Hamiltonian at  $\mathcal{O}(G^2)$  from scattering amplitudes*, *JHEP* **07** (2021) 037 [[2102.10137](#)].
- [5] W.-M. Chen, M.-Z. Chung, Y.-t. Huang and J.-W. Kim, *The 2PM Hamiltonian for binary Kerr to quartic in spin*, [2111.13639](#).
- [6] R. Aoude, K. Haddad and A. Helset, *Searching for Kerr in the 2PM amplitude*, [2203.06197](#).
- [7] Z. Bern, D. Kosmopoulos, A. Luna, R. Roiban and F. Teng, *Binary Dynamics Through the Fifth Power of Spin at  $\mathcal{O}(G^2)$* , [2203.06202](#).
- [8] R. Aoude, K. Haddad and A. Helset, *Classical gravitational spinning-spinless scattering at  $\mathcal{O}(G^2S^\infty)$* , [2205.02809](#).
- [9] LIGO SCIENTIFIC, VIRGO collaboration, *Observation of Gravitational Waves from a Binary Black Hole Merger*, *Phys. Rev. Lett.* **116** (2016) 061102 [[1602.03837](#)].
- [10] LIGO SCIENTIFIC, VIRGO collaboration, *GW170817: Observation of Gravitational Waves from a Binary Neutron Star Inspiral*, *Phys. Rev. Lett.* **119** (2017) 161101 [[1710.05832](#)].
- [11] M.J. Duff, *Quantum Tree Graphs and the Schwarzschild Solution*, *Phys. Rev. D* **7** (1973) 2317.
- [12] B.R. Holstein and J.F. Donoghue, *Classical physics and quantum loops*, *Phys. Rev. Lett.* **93** (2004) 201602 [[hep-th/0405239](#)].
- [13] J.F. Donoghue, B.R. Holstein, B. Garbrecht and T. Konstandin, *Quantum corrections to the Reissner-Nordström and Kerr-Newman metrics*, *Phys. Lett. B* **529** (2002) 132 [[hep-th/0112237](#)].
- [14] B. Bertotti, *On gravitational motion*, *Nuovo Cim.* **4** (1956) 898.
- [15] R.P. Kerr, *The Lorentz-covariant approximation method in general relativity I*, *Nuovo Cim.* **13** (1959) 469.
- [16] B. Bertotti and J. Plebanski, *Theory of gravitational perturbations in the fast motion approximation*, *Annals Phys.* **11** (1960) 169.
- [17] M. Portilla, *MOMENTUM AND ANGULAR MOMENTUM OF TWO GRAVITATING PARTICLES*, *J. Phys. A* **12** (1979) 1075.
- [18] K. Westpfahl and M. Goller, *GRAVITATIONAL SCATTERING OF TWO RELATIVISTIC PARTICLES IN POSTLINEAR APPROXIMATION*, *Lett. Nuovo Cim.* **26** (1979) 573.
- [19] M. Portilla, *SCATTERING OF TWO GRAVITATING PARTICLES: CLASSICAL APPROACH*, *J. Phys. A* **13** (1980) 3677.

- [20] L. Bel, T. Damour, N. Deruelle, J. Ibanez and J. Martin, *Poincaré-invariant gravitational field and equations of motion of two pointlike objects: The postlinear approximation of general relativity*, *Gen. Rel. Grav.* **13** (1981) 963.
- [21] K. Westpfahl, *High-Speed Scattering of Charged and Uncharged Particles in General Relativity*, *Fortsch. Phys.* **33** (1985) 417.
- [22] T. Ledvinka, G. Schaefer and J. Bicak, *Relativistic Closed-Form Hamiltonian for Many-Body Gravitating Systems in the Post-Minkowskian Approximation*, *Phys. Rev. Lett.* **100** (2008) 251101 [[0807.0214](#)].
- [23] T. Damour, *High-energy gravitational scattering and the general relativistic two-body problem*, *Phys. Rev. D* **97** (2018) 044038 [[1710.10599](#)].
- [24] J.F. Donoghue, *General relativity as an effective field theory: The leading quantum corrections*, *Phys. Rev. D* **50** (1994) 3874 [[gr-qc/9405057](#)].
- [25] N.E.J. Bjerrum-Bohr, J.F. Donoghue and B.R. Holstein, *Quantum gravitational corrections to the nonrelativistic scattering potential of two masses*, *Phys. Rev. D* **67** (2003) 084033 [[hep-th/0211072](#)].
- [26] N.E.J. Bjerrum-Bohr, J.F. Donoghue and P. Vanhove, *On-shell Techniques and Universal Results in Quantum Gravity*, *JHEP* **02** (2014) 111 [[1309.0804](#)].
- [27] D. Neill and I.Z. Rothstein, *Classical Space-Times from the S Matrix*, *Nucl. Phys. B* **877** (2013) 177 [[1304.7263](#)].
- [28] C. Cheung, I.Z. Rothstein and M.P. Solon, *From Scattering Amplitudes to Classical Potentials in the Post-Minkowskian Expansion*, *Phys. Rev. Lett.* **121** (2018) 251101 [[1808.02489](#)].
- [29] Z. Bern, C. Cheung, R. Roiban, C.-H. Shen, M.P. Solon and M. Zeng, *Scattering Amplitudes and the Conservative Hamiltonian for Binary Systems at Third Post-Minkowskian Order*, *Phys. Rev. Lett.* **122** (2019) 201603 [[1901.04424](#)].
- [30] Z. Bern, C. Cheung, R. Roiban, C.-H. Shen, M.P. Solon and M. Zeng, *Black Hole Binary Dynamics from the Double Copy and Effective Theory*, *JHEP* **10** (2019) 206 [[1908.01493](#)].
- [31] Z. Bern, J. Parra-Martinez, R. Roiban, M.S. Ruf, C.-H. Shen, M.P. Solon et al., *Scattering Amplitudes and Conservative Binary Dynamics at  $\mathcal{O}(G^4)$* , *Phys. Rev. Lett.* **126** (2021) 171601 [[2101.07254](#)].
- [32] A. Einstein, L. Infeld and B. Hoffmann, *The Gravitational equations and the problem of motion*, *Annals Math.* **39** (1938) 65.
- [33] T. Ohta, H. Okamura, T. Kimura and K. Hiida, *Physically acceptable solution of einstein's equation for many-body system*, *Prog. Theor. Phys.* **50** (1973) 492.
- [34] P. Jaranowski and G. Schaefer, *Third postNewtonian higher order ADM Hamilton dynamics for two-body point mass systems*, *Phys. Rev. D* **57** (1998) 7274 [[gr-qc/9712075](#)].
- [35] T. Damour, P. Jaranowski and G. Schaefer, *Dynamical invariants for general relativistic two-body systems at the third postNewtonian approximation*, *Phys. Rev. D* **62** (2000) 044024 [[gr-qc/9912092](#)].
- [36] L. Blanchet and G. Faye, *Equations of motion of point particle binaries at the third postNewtonian order*, *Phys. Lett. A* **271** (2000) 58 [[gr-qc/0004009](#)].

- [37] T. Damour, P. Jaranowski and G. Schafer, *Dimensional regularization of the gravitational interaction of point masses*, *Phys. Lett. B* **513** (2001) 147 [[gr-qc/0105038](#)].
- [38] T. Damour, P. Jaranowski and G. Schäfer, *Nonlocal-in-time action for the fourth post-Newtonian conservative dynamics of two-body systems*, *Phys. Rev. D* **89** (2014) 064058 [[1401.4548](#)].
- [39] P. Jaranowski and G. Schäfer, *Derivation of local-in-time fourth post-Newtonian ADM Hamiltonian for spinless compact binaries*, *Phys. Rev. D* **92** (2015) 124043 [[1508.01016](#)].
- [40] B.R. Holstein and A. Ross, *Spin Effects in Long Range Gravitational Scattering*, [0802.0716](#).
- [41] V. Vaidya, *Gravitational spin Hamiltonians from the S matrix*, *Phys. Rev. D* **91** (2015) 024017 [[1410.5348](#)].
- [42] N. Arkani-Hamed, T.-C. Huang and Y.-t. Huang, *Scattering amplitudes for all masses and spins*, *JHEP* **11** (2021) 070 [[1709.04891](#)].
- [43] B. Maybee, D. O’Connell and J. Vines, *Observables and amplitudes for spinning particles and black holes*, *JHEP* **12** (2019) 156 [[1906.09260](#)].
- [44] A. Guevara, A. Ochirov and J. Vines, *Scattering of Spinning Black Holes from Exponentiated Soft Factors*, *JHEP* **09** (2019) 056 [[1812.06895](#)].
- [45] M.-Z. Chung, Y.-T. Huang, J.-W. Kim and S. Lee, *The simplest massive S-matrix: from minimal coupling to Black Holes*, *JHEP* **04** (2019) 156 [[1812.08752](#)].
- [46] M.-Z. Chung, Y.-T. Huang and J.-W. Kim, *Classical potential for general spinning bodies*, *JHEP* **09** (2020) 074 [[1908.08463](#)].
- [47] A. Guevara, A. Ochirov and J. Vines, *Black-hole scattering with general spin directions from minimal-coupling amplitudes*, *Phys. Rev. D* **100** (2019) 104024 [[1906.10071](#)].
- [48] N. Arkani-Hamed, Y.-t. Huang and D. O’Connell, *Kerr black holes as elementary particles*, *JHEP* **01** (2020) 046 [[1906.10100](#)].
- [49] R. Aoude, K. Haddad and A. Helset, *On-shell heavy particle effective theories*, *JHEP* **05** (2020) 051 [[2001.09164](#)].
- [50] R. Aoude and A. Ochirov, *Classical observables from coherent-spin amplitudes*, *JHEP* **10** (2021) 008 [[2108.01649](#)].
- [51] R.A. Porto, *Post-Newtonian corrections to the motion of spinning bodies in NRGR*, *Phys. Rev. D* **73** (2006) 104031 [[gr-qc/0511061](#)].
- [52] R.A. Porto and I.Z. Rothstein, *Next to Leading Order Spin(1)Spin(1) Effects in the Motion of Inspiralling Compact Binaries*, *Phys. Rev. D* **78** (2008) 044013 [[0804.0260](#)].
- [53] M. Levi and J. Steinhoff, *Spinning gravitating objects in the effective field theory in the post-Newtonian scheme*, *JHEP* **09** (2015) 219 [[1501.04956](#)].
- [54] M. Levi, S. Mougiakakos and M. Vieira, *Gravitational cubic-in-spin interaction at the next-to-leading post-Newtonian order*, *JHEP* **01** (2021) 036 [[1912.06276](#)].
- [55] M. Levi, A.J. McLeod and M. Von Hippel,  *$N^3LO$  gravitational quadratic-in-spin interactions at  $G^4$* , *JHEP* **07** (2021) 116 [[2003.07890](#)].
- [56] M. Levi and F. Teng, *NLO gravitational quartic-in-spin interaction*, *JHEP* **01** (2021) 066 [[2008.12280](#)].

- [57] J.-W. Kim, M. Levi and Z. Yin, *Quadratic-in-spin interactions at fifth post-Newtonian order probe new physics*, [2112.01509](#).
- [58] Z. Liu, R.A. Porto and Z. Yang, *Spin Effects in the Effective Field Theory Approach to Post-Minkowskian Conservative Dynamics*, *JHEP* **06** (2021) 012 [[2102.10059](#)].
- [59] G.U. Jakobsen, G. Mogull, J. Plefka and J. Steinhoff, *Gravitational Bremsstrahlung and Hidden Supersymmetry of Spinning Bodies*, *Phys. Rev. Lett.* **128** (2022) 011101 [[2106.10256](#)].
- [60] G.U. Jakobsen, G. Mogull, J. Plefka and J. Steinhoff, *SUSY in the sky with gravitons*, *JHEP* **01** (2022) 027 [[2109.04465](#)].
- [61] G.U. Jakobsen and G. Mogull, *Conservative and Radiative Dynamics of Spinning Bodies at Third Post-Minkowskian Order Using Worldline Quantum Field Theory*, *Phys. Rev. Lett.* **128** (2022) 141102 [[2201.07778](#)].
- [62] M.-Z. Chung, Y.-t. Huang, J.-W. Kim and S. Lee, *Complete Hamiltonian for spinning binary systems at first post-Minkowskian order*, *JHEP* **05** (2020) 105 [[2003.06600](#)].
- [63] M. Chiodaroli, H. Johansson and P. Pichini, *Compton Black-Hole Scattering for  $s \leq 5/2$* , [2107.14779](#).
- [64] A. Falkowski and C.S. Machado, *Soft Matters, or the Recursions with Massive Spinors*, *JHEP* **05** (2021) 238 [[2005.08981](#)].
- [65] R. Aoude, M.-Z. Chung, Y.-t. Huang, C.S. Machado and M.-K. Tam, *Silence of Binary Kerr Black Holes*, *Phys. Rev. Lett.* **125** (2020) 181602 [[2007.09486](#)].
- [66] B.-T. Chen, M.-Z. Chung, Y.-t. Huang and M.K. Tam, *Minimal spin deflection of Kerr-Newman and supersymmetric black hole*, *JHEP* **10** (2021) 011 [[2106.12518](#)].
- [67] A. Guevara, *Holomorphic Classical Limit for Spin Effects in Gravitational and Electromagnetic Scattering*, *JHEP* **04** (2019) 033 [[1706.02314](#)].
- [68] P.H. Damgaard, K. Haddad and A. Helset, *Heavy Black Hole Effective Theory*, *JHEP* **11** (2019) 070 [[1908.10308](#)].
- [69] Y.F. Bautista and A. Guevara, *From Scattering Amplitudes to Classical Physics: Universality, Double Copy and Soft Theorems*, [1903.12419](#).
- [70] Y.F. Bautista, A. Guevara, C. Kavanagh and J. Vines, *to be published*, .
- [71] N.E.J. Bjerrum-Bohr, J.F. Donoghue, B.R. Holstein, L. Planté and P. Vanhove, *Bending of Light in Quantum Gravity*, *Phys. Rev. Lett.* **114** (2015) 061301 [[1410.7590](#)].
- [72] N.E.J. Bjerrum-Bohr, J.F. Donoghue, B.R. Holstein, L. Plante and P. Vanhove, *Light-like Scattering in Quantum Gravity*, *JHEP* **11** (2016) 117 [[1609.07477](#)].
- [73] D. Bai and Y. Huang, *More on the Bending of Light in Quantum Gravity*, *Phys. Rev. D* **95** (2017) 064045 [[1612.07629](#)].
- [74] H.-H. Chi, *Graviton Bending in Quantum Gravity from One-Loop Amplitudes*, *Phys. Rev. D* **99** (2019) 126008 [[1903.07944](#)].
- [75] X.O. Camanho, J.D. Edelstein, J. Maldacena and A. Zhiboedov, *Causality Constraints on Corrections to the Graviton Three-Point Coupling*, *JHEP* **02** (2016) 020 [[1407.5597](#)].
- [76] M. Accattulli Huber, A. Brandhuber, S. De Angelis and G. Travaglini, *Eikonal phase*



- matrix, deflection angle and time delay in effective field theories of gravity*, *Phys. Rev. D* **102** (2020) 046014 [[2006.02375](#)].
- [77] H. Cheng and T.T. Wu, *High-energy elastic scattering in quantum electrodynamics*, *Phys. Rev. Lett.* **22** (1969) 666.
- [78] H.D.I. Abarbanel and C. Itzykson, *Relativistic eikonal expansion*, *Phys. Rev. Lett.* **23** (1969) 53.
- [79] M. Levy and J. Sucher, *Eikonal approximation in quantum field theory*, *Phys. Rev.* **186** (1969) 1656.
- [80] D. Amati, M. Ciafaloni and G. Veneziano, *Superstring Collisions at Planckian Energies*, *Phys. Lett. B* **197** (1987) 81.
- [81] D. Amati, M. Ciafaloni and G. Veneziano, *Classical and Quantum Gravity Effects from Planckian Energy Superstring Collisions*, *Int. J. Mod. Phys. A* **3** (1988) 1615.
- [82] D.N. Kabat and M. Ortiz, *Eikonal quantum gravity and Planckian scattering*, *Nucl. Phys. B* **388** (1992) 570 [[hep-th/9203082](#)].
- [83] R. Akhoury, R. Saotome and G. Sterman, *High Energy Scattering in Perturbative Quantum Gravity at Next to Leading Power*, *Phys. Rev. D* **103** (2021) 064036 [[1308.5204](#)].
- [84] A. Koemans Collado, P. Di Vecchia and R. Russo, *Revisiting the second post-Minkowskian eikonal and the dynamics of binary black holes*, *Phys. Rev. D* **100** (2019) 066028 [[1904.02667](#)].
- [85] M. Nouri-Zonoz, *Gravoelectromagnetic approach to the gravitational Faraday rotation in stationary space-times*, *Phys. Rev. D* **60** (1999) 024013 [[gr-qc/9901011](#)].
- [86] M. Sereno, *Gravitational Faraday rotation in a weak gravitational field*, *Phys. Rev. D* **69** (2004) 087501 [[astro-ph/0401295](#)].
- [87] A. Brodutch and D.R. Terno, *Polarization rotation, reference frames and Mach's principle*, *Phys. Rev. D* **84** (2011) 121501 [[1107.1274](#)].
- [88] A. Farooqui, N. Kamran and P. Panangaden, *An Exact Expression for Photon Polarization in Kerr Geometry*, *Adv. Theor. Math. Phys.* **18** (2014) 659 [[1306.6292](#)].
- [89] A.A. Shoom, *Gravitational Faraday and spin-Hall effects of light*, *Phys. Rev. D* **104** (2021) 084007 [[2006.10077](#)].
- [90] A.A. Deriglazov, *Massless polarized particle and Faraday rotation of light in the Schwarzschild spacetime*, *Phys. Rev. D* **104** (2021) 025006 [[2103.07794](#)].
- [91] C. Chakraborty, *Gravitational analogue of Faraday rotation in the magnetized Kerr and Reissner-Nordström spacetimes*, [2106.03520](#).
- [92] Z. Li, W. Zhao and X. Er, *Gravitational Faraday rotation of gravitational waves by a Kerr black hole*, [2204.10512](#).
- [93] A. Cristofoli, R. Gonzo, D.A. Kosower and D. O'Connell, *Waveforms from Amplitudes*, [2107.10193](#).
- [94] A. Brandhuber, G. Chen, G. Travaglini and C. Wen, *Classical gravitational scattering from a gauge-invariant double copy*, *JHEP* **10** (2021) 118 [[2108.04216](#)].
- [95] H. Asada and M. Kasai, *Can we see a rotating gravitational lens?*, *Prog. Theor. Phys.* **104** (2000) 95 [[astro-ph/0006157](#)].

- [96] L. Landau, E. Lifshitz and M. Hamermesh, *The Classical Theory of Fields: Volume 2*, Course of theoretical physics, Elsevier Science (1975).
- [97] R.A. Isaacson, *Gravitational Radiation in the Limit of High Frequency. I. The Linear Approximation and Geometrical Optics*, *Phys. Rev.* **166** (1968) 1263.
- [98] Y.F. Bautista, A. Guevara, C. Kavanagh and J. Vines, *From Scattering in Black Hole Backgrounds to Higher-Spin Amplitudes: Part I*, [2107.10179](#).
- [99] J.-W. Kim and M. Shim, *Gravitational Dyonic Amplitude at One-Loop and its Inconsistency with the Classical Impulse*, *JHEP* **02** (2021) 217 [[2010.14347](#)].
- [100] Z. Bern, L.J. Dixon, D.C. Dunbar and D.A. Kosower, *One loop  $n$  point gauge theory amplitudes, unitarity and collinear limits*, *Nucl. Phys. B* **425** (1994) 217 [[hep-ph/9403226](#)].
- [101] Z. Bern, L.J. Dixon, D.C. Dunbar and D.A. Kosower, *Fusing gauge theory tree amplitudes into loop amplitudes*, *Nucl. Phys. B* **435** (1995) 59 [[hep-ph/9409265](#)].
- [102] Z. Bern, L.J. Dixon and D.A. Kosower, *One loop amplitudes for  $e^+ e^-$  to four partons*, *Nucl. Phys. B* **513** (1998) 3 [[hep-ph/9708239](#)].
- [103] D. Forde, *Direct extraction of one-loop integral coefficients*, *Phys. Rev. D* **75** (2007) 125019 [[0704.1835](#)].
- [104] W.B. Kilgore, *One-loop Integral Coefficients from Generalized Unitarity*, [0711.5015](#).
- [105] Z. Bern, H. Ita, J. Parra-Martinez and M.S. Ruf, *Universality in the classical limit of massless gravitational scattering*, *Phys. Rev. Lett.* **125** (2020) 031601 [[2002.02459](#)].
- [106] A. Cristofoli, R. Gonzo, N. Moynihan, D. O’Connell, A. Ross, M. Sergola et al., *The Uncertainty Principle and Classical Amplitudes*, [2112.07556](#).
- [107] P.H. Damgaard, L. Plante and P. Vanhove, *On an Exponential Representation of the Gravitational  $S$ -Matrix*, [2107.12891](#).
- [108] U. Kol, D. O’connell and O. Telem, *The Radial Action from Probe Amplitudes to All Orders*, [2109.12092](#).