

# Scaling limit for a second-order particle system with local annihilation <sup>\*†</sup>

Ruojun Huang

## Abstract

For a second-order particle system in  $\mathbb{R}^d$  subject to locally-in-space pairwise annihilation, we prove a scaling limit for its empirical measure on position and velocity towards a degenerate elliptic partial differential equation. Crucial ingredients are Green's function estimates for the associated hypoelliptic operator and an Itô-Tanaka trick.

## 1 Introduction

Second-order particle systems comprising of dynamics of positions and velocities, in the spirit of Newtonian mechanics, give a more complete description of interacting particles. When the behavior of a first-order system of positions only is understood, or is insufficient for one's purpose, it is natural to look at a second-order system. The technical difficulties associated with second-order systems are in general greater, some coming from the degeneracy of the associated operators. Here, we explore a simplest model of interacting particles *with local interactions* in  $\mathbb{R}^d$  written in second-order, where the interaction occurs not in the equation of the motion but in pairwise annihilation. Besides well related to works on annihilating (or proliferating, coagulating, etc) particle systems (e.g. [18, 27, 21, 22, 19, 13, 12, 10]) and generally relevant for reaction-diffusion phenomena, one motivation is also from the technical side, namely, we aim to derive a scaling limit when the interaction range between particles is truly local (see Condition 2).

Given a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ , for any positive integer  $N \in \mathbb{N}$  and  $d \geq 1$ , consider a system of particles, whose cardinality at time  $t = 0$  is  $N(0) = N$ , and with positions and velocities at any time  $t \geq 0$  denoted by  $x_i^N(t) \in \mathbb{R}^d$ ,  $v_i^N(t) \in \mathbb{R}^d$ ,  $i = 1, \dots, N(t)$ , and the interaction comes from the possibility of pairwise annihilation when spatial positions are close. More specifically, the position and velocity of each particle, while active, obeys the second-order dynamics

$$\begin{cases} x_i^N(t) = x_i(0) + \int_0^t v_i^N(s) ds \\ v_i^N(t) = v_i(0) + B_i(t) \end{cases}, \quad t \geq 0, i \in \mathcal{N}(t), \quad (1)$$

where  $\mathcal{N}(t)$  denotes the set of indices of active particles at time  $t$  whose cardinality  $|\mathcal{N}(t)| = N(t) \leq N$ ,  $\{B_i(t)\}_{i=1}^\infty$  is a collection of independent standard Brownian motions in  $\mathbb{R}^d$ , and the initial conditions  $(x_i(0), v_i(0))_{i=1}^\infty$  are random, chosen i.i.d. with some given probability density  $f_0(x, v) : \mathbb{R}^{2d} \rightarrow \mathbb{R}_+$  satisfying the condition below, independent of the Brownian motions.

**Condition 1** For every  $d \geq 1$ , we assume that there exists some finite constants  $\Gamma$  and  $R$  such that  $\|f_0(x, v)\|_\infty \leq \Gamma$  and  $\text{supp}(f_0) \subset \mathbb{B}_E((0, 0), R)$ , the closed Euclidean ball of radius  $R$  centered at the origin in  $\mathbb{R}^{2d}$ .

The probability space is endowed with the canonical filtration

$$\mathcal{F}_t := \sigma\{\{B_i(s) : 0 \leq s \leq t\}_{i=1}^\infty, (x_i(0), v_i(0))_{i=1}^\infty\}, \quad t \geq 0.$$

Interaction between different particles comes by means of pairwise annihilation, informally described as follows. Let  $\theta(x) : \mathbb{R}^d \rightarrow \mathbb{R}_+$  be given, nonnegative, of class  $C^\infty$ , with  $\int \theta dx = 1$ ,  $\theta(0) = 0$ ,  $\theta(x) = \theta(-x)$  and compact support in  $\mathbb{B}_E(0, 1)$ . Then, for any  $\epsilon \in (0, 1]$ , denote by

$$\theta^\epsilon(x) := \epsilon^{-d} \theta(\epsilon^{-1}x).$$

<sup>\*</sup>Keywords and phrases: second-order system; scaling limit; annihilating particles; local interaction; hypoellipticity.

<sup>†</sup>AMS subject classification (2020): 60K35, 82C21, 82C40, 35J70.

A configuration of the particle system is given by

$$\eta = (x_1, v_1, x_2, v_2, \dots, x_N, v_N) \in (\mathbb{R}^d \cup \emptyset)^{2N}$$

where we adopt the convention that if a particle  $i_0$  has been annihilated, we set  $x_{i_0} = v_{i_0} = \emptyset$ . It will no more participate in the dynamics of the system. Now suppose the current configuration is  $\eta$ , then for any pair of particles  $i, j$  with  $i \neq j$  (each ranging over the index set  $\mathcal{N}(\eta)$  of active particles), with an infinitesimal rate

$$\frac{1}{N} \theta^\epsilon(x_i - x_j)$$

we remove both particles  $i, j$  from the system (i.e. they annihilate with each other) and obtain a new configuration, denoted  $\eta^{-ij}$  which is equal to  $\eta$  in all other slots except for  $x_i, v_i, x_j, v_j$  which have been set to  $\emptyset$ . Since the support of the function  $\theta^\epsilon$  has radius  $\epsilon$ , interaction happens for the pair  $i, j$  only when their spatial distance is within the critical distance  $\epsilon$ . In fact, our interest is in the case where  $\epsilon = \epsilon(N)$  depends on  $N$  and converges to zero as  $N \rightarrow \infty$ . An important case to cover is the regime of local interaction, corresponding to  $\epsilon = N^{-1/d}$  in which case an individual particle typically interacts with a bounded number of others at any given time. Contrary to lattice models [4, 16] where nearest-neighbor or bounded-range interactions are standard, in our continuum case such local interaction has to be enforced by a potential (i.e.  $\theta^\epsilon$ ), which is a source of technical difficulty. One of the key techniques available to date to handle such local interaction, or sometimes even more stringently the mean-free path interaction, is due to Hammond and Rezakhanlou [13], see also [18, 27, 29, 28] for other techniques. On the other hand, mean-field type models correspond to choosing  $\epsilon$  independent of  $N$ , and in that setting techniques for proving scaling limits of (both first and second-order) particle systems have become very advanced, cf. the survey [15], even when interactions are very singular and influence directly the stochastic differential equations. Clearly, there is a big technical gap between mean-field and more local cases, which serves as another motivation for this research.

**Condition 2** For any  $d \geq 1$ , we assume that  $\epsilon = \epsilon(N) \rightarrow 0$  as  $N \rightarrow \infty$  and

$$\limsup_{N \rightarrow \infty} \frac{\epsilon^{-d}}{N} \leq 1.$$

When the equality is achieved, we call the interaction local.

We now give formally the infinitesimal generator of the jointly Markov process  $(x_i^N(t), v_i^N(t))_{i=1}^{\mathcal{N}(t)}$ :

$$\mathcal{L}_N = \mathcal{L}_N^D + \mathcal{L}_N^J$$

where the diffusion and respectively, jump (i.e. annihilation) part of the generator are (acting on test functions  $F$ )

$$\begin{aligned} \mathcal{L}_N^D F(\eta) &:= \sum_{i \in \mathcal{N}(\eta)} [v_i \cdot \nabla_{x_i} F(\eta) + \frac{1}{2} \Delta_{v_i} F(\eta)] \\ \mathcal{L}_N^J F(\eta) &:= \sum_{i \in \mathcal{N}(\eta)} \sum_{j \in \mathcal{N}(\eta); j \neq i} \frac{1}{N} \theta^\epsilon(x_i - x_j) [F(\eta^{-ij}) - F(\eta)]. \end{aligned}$$

For every  $N$  and  $T$ , denote the process of joint empirical measure of positions and velocities of active particles

$$\mu_t^N(dx, dv) := \frac{1}{N} \sum_{i \in \mathcal{N}(t)} \delta_{x_i^N(t)}(dx) \delta_{v_i^N(t)}(dv), \quad t \in [0, T]. \quad (2)$$

For each  $t$ ,  $\mu_t^N(dx, dv)$  is a subprobability measure on  $\mathbb{R}^{2d}$ . Denote by  $\mathcal{M}_{+,1}(\mathbb{R}^{2d})$  the space of subprobability measures on  $\mathbb{R}^{2d}$  endowed with the weak topology, and by

$$\mathcal{D}_T(\mathcal{M}_{+,1}) := \mathcal{D}([0, T]; \mathcal{M}_{+,1}(\mathbb{R}^{2d})) \quad (3)$$

the space of càdlàg functions on  $[0, T]$  taking values in  $\mathcal{M}_{+,1}(\mathbb{R}^{2d})$  endowed with the Skorohod topology [6, Chapter 3]. Our goal is to prove that the  $N \rightarrow \infty$  limit of (2) is given by the unique weak solution (see Definition 20) of the following degenerate elliptic partial differential equation (PDE)

$$\begin{cases} \partial_t f(t, x, v) &= -v \cdot \nabla_x f + \frac{1}{2} \Delta_v f - 2 \int_{\mathbb{R}^d} f(t, x, v) f(t, x, v') dv' \\ f|_{t=0} &= f_0(x, v) \end{cases}, \quad (t, x, v) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d. \quad (4)$$

**Theorem 3** Under Conditions 1 and 2, for any finite  $T$  and  $d \geq 1$ , the sequence of empirical measures  $\{\mu_t^N : t \in [0, T]\}_{N \in \mathbb{N}}$  converges in probability in the space  $\mathcal{D}_T(\mathcal{M}_{+,1})$  towards a limit  $\{\bar{\mu}_t : t \in [0, T]\}$ . For every  $t \in [0, T]$ , the measure  $\bar{\mu}_t(dx, dv)$  has a density  $f(t, x, v)$  with respect to Lebesgue measure and  $\{f(t, x, v) : t \in [0, T], x, v \in \mathbb{R}^d\}$  is the unique weak solution to the PDE (4) in the sense of Definition 20.

**Remark 4** Our setting (1) of a second-order system is almost the simplest possible. The motivation and challenge is to handle the local interaction. One may notice that, if substituting the velocity into the first equation (1), then we get a closed expression for the dynamics of the position

$$x_i^N(t) = x_i(0) + v_i(0)t + \int_0^t B_i(s)ds. \quad (5)$$

One may ask why not study directly a first-order system of  $N$  locally interacting particles with dynamics given by (5), especially if one is mainly interested in the limiting density of positions alone

$$\rho(t, x) := \int_{\mathbb{R}^d} f(t, x, v)dv.$$

However, it is not hard to check that  $t \mapsto \int_0^t B(s)ds$  is not a Markov process, hence working directly with the first-order dynamics is not necessarily easier. Consistent with that, if we integrate the PDE (4) formally in variable  $v$ , we do not obtain a closed equation for  $\rho(t, x)$ . It remains unclear what (if any) equation  $\rho$  should satisfy.

The rest of the article is organized as follows. In Section 2, we derive an identity satisfied by the empirical measure, and identify the nonlinear term whose convergence is the main issue. For this purpose, in Section 3 we set up the Itô-Tanaka trick which involves the solution of an elliptic PDE with a degenerate Kolmogorov (or Fokker-Planck) operator. It turns out that key to our proof are pointwise estimates of the Green's function associated with this operator, which are recalled and developed in Section 4. In Section 5, we apply these estimates to show that various terms coming from the Itô-Tanaka trick are negligible. In Section 6, we check the tightness of the laws of our particle system, which together with previous sections completes the proof of subsequential convergence. In the last section, we prove the uniqueness of the limit PDE. The techniques used here, which are very influenced by Hammond-Rezakhanlou [13], share elements with the author's previous works [10, 11], hence certain arguments are made more succinct.

## 2 Identity from the empirical measure

Fixing any  $\phi(x, v) \in C_c^\infty(\mathbb{R}^{2d})$ , we consider the process

$$\langle \phi, \mu_t^N \rangle = \frac{1}{N} \sum_{i \in \mathcal{N}(t)} \phi(x_i^N(t), v_i^N(t)), \quad t \geq 0.$$

Applying Itô's formula we obtain

$$\begin{aligned} \langle \phi, \mu_t^N \rangle &= \langle \phi, \mu_0^N \rangle + \int_0^t [\langle v \cdot \nabla_x \phi, \mu_t^N \rangle + \frac{1}{2} \langle \Delta_v \phi, \mu_t^N \rangle] dt \\ &\quad - \int_0^t \frac{1}{N^2} \sum_{i \in \mathcal{N}(t)} \sum_{j: j \neq i} \theta^\epsilon(x_i^N(t) - x_j^N(t)) [\phi(x_i^N(t), v_i^N(t)) + \phi(x_j^N(t), v_j^N(t))] dt \\ &\quad + M_T^{\phi, D} + M_T^{\phi, J}, \end{aligned} \quad (6)$$

where  $M_T^{\phi, D}, M_T^{\phi, J}$  are two martingale due to diffusion and jump (i.e. annihilation) respectively. We note that  $v \cdot \nabla_x \phi$  is bounded continuous since  $v$  is in the compact support of  $\phi$ . We can bound the martingales via quadratic variation

$$\begin{aligned} \mathbb{E}[|M_T^{\phi, D}|^2] &= \mathbb{E}\left[\left|\frac{1}{N} \int_0^T \sum_{i \in \mathcal{N}(t)} \nabla_{v_i} \phi(x_i^N(t), v_i^N(t)) \cdot dB_i(t)\right|^2\right] \\ &= \frac{1}{N^2} \mathbb{E} \int_0^T \sum_{i \in \mathcal{N}(t)} \left| \nabla_{v_i} \phi(x_i^N(t), v_i^N(t)) \right|^2 dt \\ &\leq \frac{\|\phi\|_{C^1}^2}{N^2} \mathbb{E} \int_0^T \mathcal{N}(t) dt \leq \frac{\|\phi\|_{C^1}^2 T}{N}, \end{aligned} \quad (7)$$

and cf. [3, Proposition 8.7]

$$\begin{aligned} c_1 \mathbb{E}[|M_T^{\phi, \psi}|^2] &\leq \mathbb{E} \int_0^T \frac{1}{N} \sum_{i \in \mathcal{N}(t)} \sum_{j: j \neq i} \theta^\epsilon(x_i^N(t) - x_j^N(t)) \left[ \frac{1}{N} \phi(x_i^N(t), v_i^N(t)) + \frac{1}{N} \phi(x_j^N(t), v_j^N(t)) \right]^2 dt \\ &= \frac{4\|\phi\|_\infty^2}{N^3} \mathbb{E} \int_0^T \sum_{i \in \mathcal{N}(t)} \sum_{j: j \neq i} \theta^\epsilon(x_i^N(t) - x_j^N(t)) dt, \end{aligned}$$

for some numerical constant  $c_1$ . To bound the remaining quantity, applying Itô's formula to  $\mathcal{N}(t)$  and then taking expectation, we obtain

$$\mathbb{E} \mathcal{N}(T) = \mathbb{E} \mathcal{N}(0) - \mathbb{E} \int_0^T \sum_{i \in \mathcal{N}(t)} \sum_{j: j \neq i} \frac{2}{N} \theta^\epsilon(x_i^N(t) - x_j^N(t)) dt$$

hence we see that

$$\mathbb{E} \int_0^T \sum_{i \in \mathcal{N}(t)} \sum_{j: j \neq i} \theta^\epsilon(x_i^N(t) - x_j^N(t)) dt \leq \frac{N^2}{2},$$

whereby

$$c_1 \mathbb{E}[|M_T^{\phi, \psi}|^2] \leq \frac{\|\phi\|_\infty^2}{N}. \quad (8)$$

To handle the nonlinearity, we first fix another test function  $\psi(x, v) \in C_c^\infty(\mathbb{R}^{2d})$  and consider

$$\int_0^T \frac{1}{N^2} \sum_{i \in \mathcal{N}(t)} \sum_{j: j \neq i} \theta^\epsilon(x_i^N(t) - x_j^N(t)) \phi(x_i^N(t), v_i^N(t)) \psi(x_j^N(t), v_j^N(t)) dt. \quad (9)$$

Let us denote by  $K = K^{\phi, \psi}$  the maximal radius of the compact supports of  $\phi$  and  $\psi$ , and by

$$\chi^K(\xi) : \mathbb{R}^d \rightarrow [0, 1] \quad (10)$$

a compactly-supported function of class  $C^\infty$  that is identically 1 for  $|\xi| \leq 2K$  and 0 for  $|\xi| \geq 4K$ . Then, (9) is equal to

$$\int_0^T \frac{1}{N^2} \sum_{i \in \mathcal{N}(t)} \sum_{j: j \neq i} \theta^\epsilon(x_i^N(t) - x_j^N(t)) \chi^K(v_i^N(t) - v_j^N(t)) \phi(x_i^N(t), v_i^N(t)) \psi(x_j^N(t), v_j^N(t)) dt \quad (11)$$

Indeed, since  $v_i^N(t), v_j^N(t)$  are in the arguments of  $\phi, \psi$ , their moduli are less than  $K$ , whereby  $|v_i^N(t) - v_j^N(t)| \leq 2K$  otherwise one of  $\phi, \psi$  is zero. Hence, adding the function  $\chi^K$  to (9) does not change its value.

In the next two sections we concentrate on proving

**Proposition 5** *For any finite  $T$  and  $d \geq 1$ ,*

$$\begin{aligned} \lim_{|z| \rightarrow 0} \limsup_{N \rightarrow \infty} \mathbb{E} \left| \int_0^T \frac{1}{N^2} \sum_{i \in \mathcal{N}(t)} \sum_{j: j \neq i} [\theta^\epsilon(x_i^N(t) - x_j^N(t) + z) - \theta^\epsilon(x_i^N(t) - x_j^N(t))] \right. \\ \left. \cdot \chi^K(v_i^N(t) - v_j^N(t)) \phi(x_i^N(t), v_i^N(t)) \psi(x_j^N(t), v_j^N(t)) dt \right| = 0. \end{aligned}$$

It should be seen as a form of local equilibrium. Since the function  $\theta^\epsilon$  is very close to singular (for  $\epsilon$  small), we need to apply the Itô-Tanaka trick to (11), where the additional localization by  $\psi$  would be useful.

Given Proposition 5, one can argue as in [13, pp. 42-43] (or [10, pp. 631-633]) to derive the following proposition. Let  $\eta(x) : \mathbb{R}^d \rightarrow \mathbb{R}_+$  be given, nonnegative, of class  $C^\infty$  with  $\int \eta dx = 1$ ,  $\eta(x) = \eta(-x)$  and compactly supported in  $\mathbb{B}_E(0, 1)$ . Then, for any  $\delta \in (0, 1]$ , denote by

$$\eta^\delta(x) := \delta^{-d} \eta(\delta^{-1}x).$$

**Proposition 6** For any finite  $T$  and  $d \geq 1$ ,

$$\begin{aligned} \lim_{\delta \rightarrow 0} \limsup_{N \rightarrow \infty} \mathbb{E} \left| \int_0^T \frac{1}{N^2} \sum_{i \in \mathcal{N}(t)} \sum_{j: j \neq i} \theta^\varepsilon(x_i^N(t) - x_j^N(t)) \phi(x_i^N(t), v_i^N(t)) \psi(x_j^N(t), v_j^N(t)) dt \right. \\ \left. - \int_0^T \int_{\mathbb{R}^d} dw \langle \eta^\delta(w - x_1) \eta^\delta(w - x_2) \phi(w, v_1) \psi(w, v_2), \mu_t^N(dx_1, dv_1) \mu_t^N(dx_2, dv_2) \rangle dt \right| = 0. \end{aligned}$$

**Proof.** We only sketch the proof as it is similar to arguments detailed in [13, 10, 11]. In the sequel, we use  $\text{Err}(N, \delta)$  to denote a stochastic error that can change from line to line and which vanishes in the following limit

$$\lim_{\delta \rightarrow 0} \limsup_{N \rightarrow \infty} \mathbb{E} |\text{Err}(N, \delta)| = 0.$$

By Proposition 5 (we can remove the function  $\chi^K$  without changing the value, as explained near (11)) and  $\theta(0) = 0$ , we have that

$$\begin{aligned} & \int_0^T \frac{1}{N^2} \sum_{i, j \in \mathcal{N}(t)} \theta^\varepsilon(x_i^N(t) - x_j^N(t)) \phi(x_i^N(t), v_i^N(t)) \psi(x_j^N(t), v_j^N(t)) dt \\ &= \int_0^T \iint_{\mathbb{R}^{2d}} \eta^\delta(z_1) \eta^\delta(z_2) dz_1 dz_2 \frac{1}{N^2} \sum_{i, j \in \mathcal{N}(t)} \theta^\varepsilon(x_i^N(t) - x_j^N(t) - z_1 + z_2) \\ & \quad \cdot \phi(x_i^N(t), v_i^N(t)) \psi(x_j^N(t), v_j^N(t)) dt + \text{Err}(N, \delta) \\ &= \int_0^T \iint_{\mathbb{R}^{2d}} \eta^\delta(z_1) \eta^\delta(z_2) dz_1 dz_2 \frac{1}{N^2} \sum_{i, j \in \mathcal{N}(t)} \theta^\varepsilon(x_i^N(t) - x_j^N(t) - z_1 + z_2) \\ & \quad \cdot \phi(x_i^N(t) - z_1, v_i^N(t)) \psi(x_j^N(t) - z_2, v_j^N(t)) dt + \text{Err}(N, \delta) \\ &= \int_0^T \iint_{\mathbb{R}^{2d}} \eta^\delta(x_i^N(t) - w_1) \eta^\delta(x_j^N(t) - w_2) dw_1 dw_2 \frac{1}{N^2} \sum_{i, j \in \mathcal{N}(t)} \theta^\varepsilon(w_1 - w_2) \\ & \quad \cdot \phi(w_1, v_i^N(t)) \psi(w_2, v_j^N(t)) dt + \text{Err}(N, \delta) \\ &= \int_0^T \iint_{\mathbb{R}^{2d}} \eta^\delta(x_i^N(t) - w_1) \eta^\delta(x_j^N(t) - w_1) dw_1 dw_2 \frac{1}{N^2} \sum_{i, j \in \mathcal{N}(t)} \theta^\varepsilon(w_1 - w_2) \\ & \quad \cdot \phi(w_1, v_i^N(t)) \psi(w_1, v_j^N(t)) dt + \text{Err}(N, \delta) \\ &= \int_0^T \int_{\mathbb{R}^d} \eta^\delta(x_i^N(t) - w_1) \eta^\delta(x_j^N(t) - w_2) dw_1 \frac{1}{N^2} \sum_{i, j \in \mathcal{N}(t)} \phi(w_1, v_i^N(t)) \psi(w_1, v_j^N(t)) dt + \text{Err}(N, \delta) \\ &= \int_0^T \int_{\mathbb{R}^d} dw_1 \langle \eta^\delta(x_1 - w_1) \eta^\delta(x_2 - w_1) \phi(w_1, v_1) \psi(w_1, v_2), \mu_t^N(dx_1, dv_1) \mu_t^N(dx_2, dv_2) \rangle dt + \text{Err}(N, \delta), \end{aligned}$$

where the second equality is due to a shift of the arguments of  $\phi, \psi$  by  $z_1, z_2$  causing an additional error on the order of  $\delta$ , the third equality is a change of variables, the fourth equality is a shift in the arguments of  $\eta^\delta$  and  $\psi$  from  $w_2$  to  $w_1$  causing an additional error on the order of  $\varepsilon \delta^{-2d-1}$ , the fifth equality is due to  $\int \theta^\varepsilon(w_1 - w_2) dw_2 = 1$  for any  $w_1$ , and the last equality is a rewriting in terms of the empirical measure  $\mu_t^N$ . ■

### 3 The Itô-Tanaka trick

We apply the Itô-Tanaka trick to (11). The trick itself is a classical one in stochastic analysis, see e.g. [9], but its use in a particle system context we learned from Hammond and Rezakhanlou [13]. For the above-fixed test functions  $\phi, \psi$ , and for any fixed  $z \in \mathbb{R}^d$ , consider the process

$$\mathcal{Y}_t := \int_0^t \frac{1}{N^2} \sum_{i \in \mathcal{N}(s)} \sum_{j: j \neq i} H^{\varepsilon, z}(x_i^N(s) - x_j^N(s), v_i^N(s) - v_j^N(s)) \phi(x_i^N(s), v_i^N(s)) \psi(x_j^N(s), v_j^N(s)) ds, \quad t \geq 0$$

where

$$H^{\varepsilon,z}(x, v) := U^\varepsilon(x + z, v) - U^\varepsilon(x, v)$$

and  $U^\varepsilon(x, v) : \mathbb{R}^{2d} \rightarrow \mathbb{R}$  is a function of class  $C^\infty$  to be defined in (30). Applying Itô's formula to  $\mathcal{Y}_t$ , we obtain that

$$\mathcal{Y}_T - \mathcal{Y}_0 \tag{12}$$

$$= \int_0^T \frac{1}{N^2} \sum_{i \in \mathcal{N}(t)} \sum_{j: j \neq i} (v_i^N(t) - v_j^N(t)) \cdot \nabla_x H^{\varepsilon,z}(x_i^N(t) - x_j^N(t), v_i^N(t) - v_j^N(t)) \phi(x_i^N(t), v_i^N(t)) \psi(x_j^N(t), v_j^N(t)) dt \tag{13}$$

$$+ \int_0^T \frac{1}{N^2} \sum_{i \in \mathcal{N}(t)} \sum_{j: j \neq i} \Delta_v H^{\varepsilon,z}(x_i^N(t) - x_j^N(t), v_i^N(t) - v_j^N(t)) \phi(x_i^N(t), v_i^N(t)) \psi(x_j^N(t), v_j^N(t)) dt \tag{14}$$

$$+ \int_0^T \frac{1}{N^2} \sum_{i \in \mathcal{N}(t)} \sum_{j: j \neq i} \left[ v_i^N(t) \cdot \nabla_x \phi(x_i^N(t), v_i^N(t)) \psi + v_j^N(t) \cdot \nabla_x \psi(x_j^N(t), v_j^N(t)) \phi \right] H^{\varepsilon,z} dt \tag{15}$$

$$+ \int_0^T \frac{1}{2N^2} \sum_{i \in \mathcal{N}(t)} \sum_{j: j \neq i} \left[ \Delta_v \phi(x_i^N(t), v_i^N(t)) \psi(x_j^N(t), v_j^N(t)) + \Delta_v \psi(x_j^N(t), v_j^N(t)) \phi(x_i^N(t), v_i^N(t)) \right] H^{\varepsilon,z} dt \tag{16}$$

$$- \int_0^T \frac{2}{N^3} \sum_{i \in \mathcal{N}(t)} \sum_{k: k \neq i} \theta^\varepsilon(x_i^N(t) - x_k^N(t)) \sum_{j: j \neq i, k} H^{\varepsilon,z}(x_i^N(t) - x_j^N(t), v_i^N(t) - v_j^N(t)) \phi(x_i^N(t), v_i^N(t)) \psi(x_j^N(t), v_j^N(t)) dt \tag{17}$$

$$- \int_0^T \frac{2}{N^3} \sum_{i \in \mathcal{N}(t)} \sum_{k: k \neq i} \theta^\varepsilon(x_i^N(t) - x_k^N(t)) \sum_{j: j \neq i, k} H^{\varepsilon,z}(x_j^N(t) - x_i^N(t), v_j^N(t) - v_i^N(t)) \phi(x_j^N(t), v_j^N(t)) \psi(x_i^N(t), v_i^N(t)) dt \tag{18}$$

$$- \int_0^T \frac{2}{N^3} \sum_{i \in \mathcal{N}(t)} \sum_{k: k \neq i} \theta^\varepsilon(x_i^N(t) - x_k^N(t)) H^{\varepsilon,z}(x_i^N(t) - x_k^N(t), v_i^N(t) - v_k^N(t)) \phi(x_i^N(t), v_i^N(t)) \psi(x_k^N(t), v_k^N(t)) dt \tag{19}$$

$$+ M_T^D + M_T^J,$$

where we omitted some arguments for brevity as they are self-evident. The martingale associated with diffusion is

$$M_T^D := \frac{1}{N^2} \int_0^T \sum_{i \in \mathcal{N}(t)} \sum_{j: j \neq i} (\nabla_v H^{\varepsilon,z} \phi \psi + H^{\varepsilon,z} \psi \nabla_v \phi) dB_j(t) + (-\nabla_v H^{\varepsilon,z} \phi \psi + H^{\varepsilon,z} \phi \nabla_v \psi) dB_j(t),$$

hence

$$\frac{1}{2} \mathbb{E}[|M_T^D|^2] \leq \frac{1}{N^4} \int_0^T \sum_{i \in \mathcal{N}(t)} \left| \sum_{j: j \neq i} (\nabla_v H^{\varepsilon,z} \phi \psi + H^{\varepsilon,z} \psi \nabla_v \phi) \right|^2 dt \tag{20}$$

$$+ \frac{1}{N^4} \int_0^T \sum_{j \in \mathcal{N}(t)} \left| \sum_{i: i \neq j} (-\nabla_v H^{\varepsilon,z} \phi \psi + H^{\varepsilon,z} \phi \nabla_v \psi) \right|^2 dt. \tag{21}$$

The martingale associated with jumps can be bounded via

$$c_1 \mathbb{E}[|M_T^J|^2] \leq \int_0^T \sum_{i \in \mathcal{N}(t)} \sum_{k: k \neq i} \frac{1}{N} \theta^\varepsilon(x_i^N(t) - x_k^N(t)) \left[ \frac{1}{N} \sum_{j: j \neq i, k} H^{\varepsilon,z}(x_i^N(t) - x_j^N(t), v_i^N(t) - v_j^N(t)) \phi \psi \right]^2 dt \tag{22}$$

$$+ \int_0^T \sum_{i \in \mathcal{N}(t)} \sum_{k: k \neq i} \frac{1}{N} \theta^\varepsilon(x_i^N(t) - x_k^N(t)) \left[ \frac{1}{N^2} \sum_{j: j \neq i, k} H^{\varepsilon,z}(x_j^N(t) - x_i^N(t), v_j^N(t) - v_i^N(t)) \phi \psi \right]^2 dt \tag{23}$$

$$+ \int_0^T \sum_{i \in \mathcal{N}(t)} \sum_{k: k \neq i} \frac{1}{N} \theta^\varepsilon(x_i^N(t) - x_k^N(t)) \left[ \frac{1}{N^2} H^{\varepsilon,z}(x_i^N(t) - x_k^N(t), v_i^N(t) - v_k^N(t)) \phi \psi \right]^2 dt, \tag{24}$$

for some numerical constant  $c_1$ , cf. [3, Proposition 8.7].

The only term with second derivatives of  $H^{\varepsilon,z}$  from the above development is (14). In order to substitute the highest derivative (also together with (13)), we set up the following elliptic equation for  $U^\varepsilon : \mathbb{R}^{2d} \rightarrow \mathbb{R}$ :

$$v \cdot \nabla_x U^\varepsilon(x, v) + \Delta_v U^\varepsilon(x, v) = -\theta^\varepsilon(x) \chi^K(v). \quad (25)$$

We will choose in (30) a particular solution of (25). It yields, by linearity, the equation for  $H^{\varepsilon,z} : \mathbb{R}^{2d} \rightarrow \mathbb{R}$  and any fixed  $z$ :

$$v \cdot \nabla_x H^{\varepsilon,z}(x, v) + \Delta_v H^{\varepsilon,z}(x, v) = [\theta^\varepsilon(x) - \theta^\varepsilon(x+z)] \chi^K(v). \quad (26)$$

Consequently, we can substitute (13)-(14) as follows

$$(13) + (14) = \int_0^T \frac{1}{N^2} \sum_{i \in \mathcal{N}(t)} \sum_{j: j \neq i} \\ \cdot [\theta^\varepsilon(x_i^N(t) - x_j^N(t)) - \theta^\varepsilon(x_i^N(t) - x_j^N(t) + z)] \chi^K(v_i^N(t) - v_j^N(t)) \phi(x_i^N(t), v_i^N(t)) \psi(x_j^N(t), v_j^N(t)).$$

The RHS is precisely the expression appearing in Proposition 5.

We remark at this stage that  $\nabla_x H^{\varepsilon,z}$  does not appear in the rest of terms coming from the Itô formula applied to  $\mathcal{Y}_t$ . Its estimate is expected to be bad due to the degeneracy of the operator in (26) in the direction of  $x$ . What do appear in the rest of the terms are  $\nabla_v H^{\varepsilon,z}$  and  $H^{\varepsilon,z}$  itself. We will show that all of them (i.e. apart from (13)-(14)) are negligible in the order of limits as in Proposition 5.

## 4 The hypoelliptic Green's function

Denote the operator

$$L := v \cdot \nabla_x + \Delta_v \quad (27)$$

acting on appropriate functions on  $\mathbb{R}^{2d}$ , and we rewrite the PDE (25) more compactly as

$$LU^\varepsilon = -\theta^\varepsilon(x) \chi^K(v). \quad (28)$$

It is well-known that  $L$  (27) is an hypoelliptic operator in the sense of Hörmander [14], and its parabolic version (i.e.  $\partial_t - L$ ) was earlier studied by Kolmogorov [17] who wrote down its kernel explicitly. cf. [1] for a recent survey. Crucial for our purpose are pointwise estimates for the Green's function of  $-L$ .

To briefly recall that  $L$  is hypoelliptic, let us write

$$L = A_0 + \sum_{i=1}^d A_i^2 \quad (29)$$

where

$$A_i = \partial_{v_i}, \quad i = 1, \dots, d; \quad A_0 = \sum_{j=1}^d v_j \partial_{x_j}.$$

Since the commutators

$$[A_0, A_i] = \sum_{j=1}^d v_j \partial_{x_j} (\partial_{v_i}) - \partial_{v_i} (\sum_{j=1}^d v_j \partial_{x_j}) = -\partial_{x_i}, \quad i = 1, \dots, d$$

we see that at every point in  $\mathbb{R}^{2d}$ ,

$$\text{span}\{A_i, [A_0, A_i], i = 1, \dots, d\} = \mathbb{R}^{2d}.$$

Thus  $L$  is hypoelliptic, and since the RHS of (28) is of class  $C^\infty$  with compact support, we can find a solution of the form

$$U^\varepsilon(x, v) = \int_{\mathbb{R}^{2d}} G((x, v), (y, w)) \theta^\varepsilon(y) \chi^K(w) dy dw, \quad (30)$$

where  $G = G_L$  is a distribution of class  $C^\infty$  away from the diagonal, called the Green's function of  $-L$ , see e.g. [25, pp. 143], [8, pp. 247]. Hypoellipticity further implies that  $U^\epsilon$  is of class  $C^\infty$ . In particular, this is the function that is used in Section 3 to define  $\mathcal{Y}_i$ .

To state the classical results of [20, 25, 8] on the estimates for the Green's function of hypoelliptic operators, we recall that they are expressed in terms of a control metric (or Carnot-Carathéodory metric)  $\rho_L(X, Y)$  canonically attached to  $L$  (to avoid confusion, we use  $X, Y, \dots$  to denote variables in  $\mathbb{R}^{2d}$ ). We also set

$$D := 2d.$$

**Definition 7 ([20, Definition 1.1])** *Let  $C(\delta)$  denote the class of absolutely continuous mappings  $\phi : [0, 1] \rightarrow \mathbb{R}^D$  which almost everywhere satisfy the differential equation*

$$\phi'(t) = \sum_{j=1}^q a_j(t) Y_j(\phi(t))$$

with  $|a_j(t)| \leq \delta^{d_j}$ . Then define

$$\rho_L(X, Y) := \inf \{ \delta > 0 : \exists \phi \in C(\delta), \phi(0) = X, \phi(1) = Y \}.$$

In our case,  $q = 1 + 2d$  and  $\{Y_j, j = 1, \dots, q\}$  are an enumeration of

$$Y_1 := A_0, Y_{i+1} := A_i, Y_{i+d+1} := [A_0, A_i], \quad i = 1, \dots, d.$$

As explained in [20, pp.107] or [24, Section 18], for operators of the form (29) which are not just a sum-of-squares but with  $A_0$ , the degree of  $A_0$  counts 2 while the degree of each  $A_i$  counts 1, and likewise for their commutators. Hence, in our case we have that

$$d_1 = 2, \quad d_{i+1} = 1, \quad d_{i+d+1} = 3, \quad i = 1, \dots, d,$$

where  $d_j$  is the degree of  $Y_j$ . It is known that  $\rho_L$  is a metric and in our case, it compares with the Euclidean distance  $|\cdot|$  in the following way [20, Proposition 1.1]: for any compact set  $\mathbb{K} \subset \mathbb{R}^D$ , there are constants  $C_i = C_i(\mathbb{K}, d)$ ,  $i = 1, 2$ , such that for any  $X, Y \in \mathbb{K}$  we have that

$$C_1 |X - Y| \leq \rho_L(X, Y) \leq C_2 |X - Y|^{1/3}. \quad (31)$$

Given the control metric  $\rho_L$  we define balls in this metric of radius  $\delta > 0$  by

$$\mathbb{B}_L(X, \delta) := \{ Y \in \mathbb{R}^D : \rho_L(X, Y) \leq \delta \}.$$

These balls are locally volume doubling as proved in [20].

**Theorem 8 ([20, Theorem 1, pp. 110])** *For every compact set  $\mathbb{K} \subset \mathbb{R}^D$ , there exist constants  $C_1, C_2$  so that for all  $X \in \mathbb{K}$ ,*

$$0 < C_1 \leq \frac{|\mathbb{B}_L(X, \delta)|}{\Lambda(X, \delta)} \leq C_2 < \infty,$$

where

$$\Lambda(X, \delta) := \sum_{1 \leq i_1, \dots, i_D \leq q} \left| \det(Y_{i_1}, \dots, Y_{i_D})(X) \right| \delta^{d_{i_1} + \dots + d_{i_D}}.$$

In our case, it is straightforward to check that we have

$$\Lambda(X, \delta) = \sum_{i=1}^d |v_i| \delta^{4d-1} + \delta^{4d}, \quad X := (x_1, \dots, x_d, v_1, \dots, v_d). \quad (32)$$

It is then clear that

$$\frac{|\mathbb{B}_L(X, 2\delta)|}{|\mathbb{B}_L(X, \delta)|} \leq \frac{C_2 \Lambda(X, 2\delta)}{C_1 \Lambda(X, \delta)} \leq \frac{C_2}{C_1} 2^{4d}, \quad (33)$$

where the RHS is uniform for all  $X \in \mathbb{K}$ .

We can now state the pointwise estimates on Green's function and its partial derivatives in  $v$ , in the case of our operator  $L$ .

**Theorem 9** ([20, Corollary, pp. 114], see also [8, Theorem 1]) *Suppose that  $D = 2d \geq 2$ . For any compact set  $\mathbb{K} \subset \mathbb{R}^D$ , there exist some finite constants  $C_i = C_i(\mathbb{K}, d)$ ,  $i = 1, 2$ , such that for any  $X = (x, v), Y = (y, w) \in \mathbb{K}$ , we have that*

$$|G(X, Y)| \leq C_1 \frac{\rho_L(X, Y)^2}{|\mathbb{B}_L(X, \rho_L(X, Y))|}, \quad (34)$$

$$|\nabla_v G(X, Y)| \leq C_2 \frac{\rho_L(X, Y)}{|\mathbb{B}_L(X, \rho_L(X, Y))|}. \quad (35)$$

We check that (34) and (35) are sufficiently integrable near its singularity.

**Proposition 10** *Suppose that  $D = 2d \geq 2$ . There exists some  $\eta = \eta(d) > 0$  such that for any compact set  $\mathbb{K} \subset \mathbb{R}^D$ , and any  $X = (x, v) \in \mathbb{K}$ ,  $\delta \in (0, \infty)$ , there are finite constants  $C_i(\delta, \mathbb{K}, d)$ ,  $i = 1, 2$  such that*

$$\int_{\mathbb{B}_L(X, \delta)} |G(X, Y)|^{1+\eta} dY \leq C_1 \quad (36)$$

$$\int_{\mathbb{B}_L(X, \delta)} |\nabla_v G(X, Y)|^{1+\eta} dY \leq C_2. \quad (37)$$

**Proof.** We prove (37) while the proof of (36) is similar. For  $X \in \mathbb{K} \subset \mathbb{R}^D$  compact and  $Y \in \mathbb{B}_L(X, \delta)$  with  $\delta \in (0, \infty)$ , they both belong to some other compact set  $\mathbb{K}'$  that depends only on  $\mathbb{K}, \delta$ . This fact can be inferred from the meaning of the control metric Definition 7 in the case of our operator  $L$ . By (35), there exists some  $C = C(\mathbb{K}', d)$  such that for any  $\eta > 0$ ,

$$\begin{aligned} \int_{\mathbb{B}_L(X, \delta)} |\nabla_v G(X, Y)|^{1+\eta} dY &\leq C \int_{\mathbb{B}_L(X, \delta)} \frac{\rho_L(X, Y)^{1+\eta}}{|\mathbb{B}_L(X, \rho_L(X, Y))|^{1+\eta}} dY \\ &= C \sum_{k=0}^{\infty} \int_{\mathbb{B}_L(X, \delta 2^{-k}) \setminus \mathbb{B}_L(X, \delta 2^{-k-1})} \frac{\rho_L(X, Y)^{1+\eta}}{|\mathbb{B}_L(X, \rho_L(X, Y))|^{1+\eta}} dY \\ &\leq C \sum_{k=0}^{\infty} \frac{(\delta 2^{-k})^{1+\eta}}{|\mathbb{B}_L(X, \delta 2^{-k-1})|^{1+\eta}} |\mathbb{B}_L(X, \delta 2^{-k}) \setminus \mathbb{B}_L(X, \delta 2^{-k-1})| \\ &\leq C \sum_{k=0}^{\infty} \frac{(\delta 2^{-k})^{1+\eta}}{|\mathbb{B}_L(X, \delta 2^{-k-1})|^\eta} \frac{|\mathbb{B}_L(X, \delta 2^{-k})|}{|\mathbb{B}_L(X, \delta 2^{-k-1})|}. \end{aligned}$$

By (32),

$$|\mathbb{B}_L(X, \delta 2^{-k-1})| \geq C_1(\mathbb{K}) \Lambda(X, \delta 2^{-k-1}) \geq C_1(\delta 2^{-(k+1)})^{4d}.$$

Combined with the local volume doubling property (33) of  $\mathbb{B}_L(\cdot)$ , we conclude that

$$\int_{\mathbb{B}_L(X, \delta)} |\nabla_v G(X, Y)|^{1+\eta} dY \leq C' \sum_{k=0}^{\infty} 2^{-k(1+\eta-4d\eta)},$$

for some finite constant  $C' = C'(\mathbb{K}, d, \delta)$ . Choosing  $\eta > 0$  small enough such that  $1 + \eta - 4d\eta > 1/2$ , the sum on the RHS is convergent. ■

Another remark is that in the case of our operator  $-L$  (27), its Green's function is translation invariant in  $x$  (but not in  $v$ ). That is, we claim that

$$G((x, v), (y, w)) = G((0, v), (y - x, w)). \quad (38)$$

This can be seen as least starting from the explicit kernel  $P_t((x, v), (y, w))$  associated with the parabolic operator  $\partial_t - L$ . Since it is probabilistically linked to Gaussian processes, it can be explicitly calculated that, cf. [17], [7, Eq. (3.24)-(3.25)], [1, pp. 222],

$$\begin{aligned} P_t((x, v), (y, w)) &= \left( \frac{\sqrt{3}}{2\pi t^2} \right)^d \exp \left\{ -\frac{|v-w|^2}{t} - 3 \frac{(v-w) \cdot (x-y-tv)}{t^2} - 3 \frac{|x-y-tv|^2}{t^3} \right\} \\ &= \left( \frac{\sqrt{3}}{2\pi t^2} \right)^d \exp \left\{ -3 \left| \frac{v-w}{2t^{1/2}} + \frac{x-y-tv}{t^{3/2}} \right|^2 - \frac{|v-w|^2}{4t} \right\}. \end{aligned} \quad (39)$$

which depends on  $x, y$  through  $y - x$  only. Since the Green's function is the time-integral of the parabolic kernel (cf. [30, pp. 109], [2, pp. 220], [26, pp. 276]):

$$G((x, v), (y, w)) = \int_0^\infty P_t((x, v), (y, w)) dt,$$

the same translation invariance with respect to  $x$  transfers to  $G$  and thus we have (38).

## 5 The control of negligible terms

We proceed to bound all the terms coming out of Itô's formula applied to  $\mathcal{Y}_i$  in Section 3, that are expected to be negligible, thereby completing the proof of Proposition 5. By a slight abuse of notation, we define on the same probability space  $(\Omega, \mathcal{F}, \mathbb{P})$

$$X_i(t) := (x_i(t), v_i(t)) := (x_i(0) + v_i(0)t + \int_0^t B_i(s)ds, v_i(0) + B_i(t)), \quad i \in \mathbb{N}, t \geq 0, \quad (40)$$

a countable collection of independent particles, with the *same* Brownian motions  $\{B_i(t)\}_{i=1}^\infty$  as (1) and *same* initial condition for  $(x_i(0), v_i(0))_{i=1}^\infty$  with density  $f_0(x, v)$  satisfying Condition 1. The only difference with (1) is that the particles in (40) are non-interacting (i.e. they do not annihilate), hence they are independent, each with infinite lifespan. Further, there is an obvious a.s. domination between (1) and (40) such that whenever  $(x_i^N(t), v_i^N(t))$  is active, its trajectory coincides with that of  $(x_i(t), v_i(t))$ .

**Lemma 11** *Let  $d \geq 1$ . For any  $t \geq 0$ , the density of  $X_1(t) - X_2(t)$  is uniformly bounded by  $\Gamma$ , the uniform bound on  $f_0$  in Condition 1. Further, the density  $p(t, x, v)$  of one particle  $X_1(t)$  satisfies for every  $t \in [0, T]$ ,  $x, v \in \mathbb{R}^d$ , the bound*

$$p(t, x, v) \leq Ce^{-|v|/C} 1_{\{|v| \geq C\}} + \Gamma 1_{\{|v| < C\}}, \quad (41)$$

for some constant  $C = C(R, \Gamma, T, d)$ , where  $R$  is the maximal radius of the compact support of  $f_0$  in Condition 1.

**Proof.** First, we look at the density  $p(t, x, v)$  of one particle  $X_1(t)$ , which exists by hypoellipticity. Its evolution is governed by

$$\begin{aligned} \partial_t p(t, x, v) &= L^* p(t, x, v) \\ p|_{t=0} &= f_0(x, v), \end{aligned} \quad (42)$$

where  $L^*$  is the adjoint of  $L$ ,

$$L^* = -v \cdot \nabla_x + \frac{1}{2} \Delta_v.$$

One can write  $p(t, x, v)$  explicitly as  $\iint f_0(x_0, v_0) P_t^*((x_0, v_0), (x, v)) dx_0 dv_0$ , where the kernel  $P_t^*$  is given by a formula similar to (39) but with certain sign and constant changes:

$$P_t^*((x_0, v_0), (x, v)) := \left( \frac{\sqrt{3}}{\pi t^2} \right)^d \exp \left\{ -6 \left| \frac{v - v_0}{2t^{1/2}} + \frac{x - x_0 - tv_0}{t^{3/2}} \right|^2 - \frac{|v - v_0|^2}{2t} \right\}. \quad (43)$$

It then follows (e.g. from probabilistic representation of the solution of (42), cf. [23, Eq. (21)]) that

$$\|p(t, \cdot)\|_\infty \leq \|f_0\|_\infty \leq \Gamma.$$

Since the density of  $X_1(t) - X_2(t)$  is a convolution between the respective densities of  $X_1(t)$  and  $-X_2(t)$ , the two being independent, we conclude that it is also uniformly bounded by  $\Gamma$ .

Turning to a finer bound on  $p(t, x, v)$  when  $|v|$  is large compared to  $T$  and  $R$ , we note that by (43),

$$\begin{aligned} p(t, x, v) &= \iint_{\mathbb{R}^{2d}} f_0(x_0, v_0) P_t^*((x_0, v_0), (x, v)) dx_0 dv_0 \\ &\leq C(\Gamma) \iint_{|x_0|, |v_0| \leq R} \frac{1}{t^{2d}} \exp \left\{ -\frac{|v - v_0|^2}{4t} \right\} dx_0 dv_0. \end{aligned}$$

If  $|v_0| \leq R$  and

$$|v| \geq 2R + T \geq t,$$

then  $|v - v_0| \geq |v|/2$  and  $\lambda := |v|^2/t \geq |v|$ . Hence, for any such  $v$  we have that

$$\begin{aligned} \mathfrak{p}(t, x, v) &\leq C(\Gamma) R^{2d} \frac{|v|^{4d}}{t^{2d}} e^{-\frac{|v|^2}{16t}} |v|^{-4d} \\ &\leq C(\Gamma, R, T, d) \lambda^{2d} e^{-\lambda/16}. \end{aligned}$$

Since  $\lambda \mapsto \lambda^{2d} e^{-\lambda/16}$  is decreasing for  $\lambda \geq |v| \geq \lambda_0$  for some  $\lambda_0(d) < \infty$ , we get the bound

$$\mathfrak{p}(t, x, v) \leq C|v|^{2d} e^{-|v|/C},$$

for any  $|v| \geq C$ ,  $x \in \mathbb{R}^d$  and  $t \in [0, T]$ , where  $C = C(\Gamma, R, T, d)$  is a finite constant. Further adjusting the constant  $C$  to absorb the polynomial factor, we get (41). ■

The following proposition bounds (15). An analogous proof can treat (12) and (16), which we omit. We note at this stage that since the independent system (40) dominates our true system (1) in terms of lifespan, it suffices to prove all the following propositions for (40).

**Proposition 12** *For any finite  $T$  and  $d \geq 1$ , we have that*

$$\lim_{|z| \rightarrow 0} \limsup_{N \rightarrow \infty} \mathbb{E} \int_0^T dt \frac{1}{N^2} \sum_{i \neq j=1}^N \left| v_i(t) \cdot \nabla_x \phi(x_i(t), v_i(t)) \psi(x_j(t), v_j(t)) H^{\varepsilon, z}(x_i(t) - x_j(t), v_i(t) - v_j(t)) \right| = 0.$$

**Proof.** Recall that we have denoted by  $K = K^{\phi, \psi}$  the maximal radius of the compact supports of  $\phi, \psi$ , and defined a compactly-supported function  $\chi^K : \mathbb{R}^d \rightarrow [0, 1]$  (10). For the quantity in the thesis, it suffices to consider

$$|x_i(t)|, |x_j(t)|, |v_i(t)|, |v_j(t)| \leq K$$

(otherwise one of the arguments of  $\phi$  and  $\psi$  is zero). When this is the case, we have that

$$|x_i(t) - x_j(t)|, |v_i(t) - v_j(t)| \leq 2K.$$

We control each term of the double sum in the thesis. That is, consider particles indexed with 1, 2

$$\begin{aligned} &\mathbb{E} \int_0^T dt \left| v_1(t) \cdot \nabla_x \phi(x_1(t), v_1(t)) \psi(x_2(t), v_2(t)) H^{\varepsilon, z}(x_1(t) - x_2(t), v_1(t) - v_2(t)) \right| \\ &\leq K \|\phi\|_{C^1} \|\psi\|_{\infty} \mathbb{E} \int_0^T dt \left| H^{\varepsilon, z}(x_1(t) - x_2(t), v_1(t) - v_2(t)) \right| \chi^K(x_1(t) - x_2(t)) \chi^K(v_1(t) - v_2(t)). \end{aligned}$$

Denote by  $\mathfrak{p}^{12}(t, x, v)$  the density of  $(x_1(t) - x_2(t), v_1(t) - v_2(t))$ , which by Lemma 11 is uniformly bounded by  $\Gamma$ . We can write the expectation above explicitly as

$$\int_0^T dt \iint_{\mathbb{R}^{2d}} |H^{\varepsilon, z}(x, v)| \chi^K(x) \chi^K(v) \mathfrak{p}^{12}(t, x, v) dx dv.$$

By (30), (38), a change of variables  $\tilde{y} = y - x$  and the symmetry of  $\theta^\varepsilon$ , we have that

$$\begin{aligned} H^{\varepsilon, z}(x, v) &= \iint_{\mathbb{R}^{2d}} G((x, v), (y, w)) [\theta^\varepsilon(y) - \theta^\varepsilon(y + z)] \chi^K(w) dy dw \\ &= \iint_{\mathbb{R}^{2d}} G((0, v), (y - x, w)) [\theta^\varepsilon(y) - \theta^\varepsilon(y + z)] \chi^K(w) dy dw \\ &= \iint_{\mathbb{R}^{2d}} G((0, v), (\tilde{y}, w)) [\theta^\varepsilon(\tilde{y} + x) - \theta^\varepsilon(\tilde{y} + x + z)] \chi^K(w) d\tilde{y} dw \\ &= \iint_{\mathbb{R}^{2d}} G((0, v), (\tilde{y}, w)) [\theta^\varepsilon(-\tilde{y} - x) - \theta^\varepsilon(-\tilde{y} - x - z)] \chi^K(w) d\tilde{y} dw \\ &= \iint_{\mathbb{R}^{2d}} [G((0, v), (\tilde{y}, w)) - G((0, v), (\tilde{y} - z, w))] \theta^\varepsilon(-\tilde{y} - x) \chi^K(w) d\tilde{y} dw. \end{aligned} \tag{44}$$

It yields that

$$\begin{aligned}
& \iint_{\mathbb{R}^{2d}} |H^{\epsilon, z}(x, v)| \chi^K(x) \chi^K(v) p^{12}(t, x, v) dx dv \\
& \leq \iiint_{\mathbb{R}^{3d}} \left| G((0, v), (\tilde{y}, w)) - G((0, v), (\tilde{y} - z, w)) \right| \chi^K(w) \left( \int_{\mathbb{R}^d} \theta^\epsilon(-\tilde{y} - x) \chi^K(x) \chi^K(v) p^{12}(t, x, v) dx \right) d\tilde{y} dw dv \\
& \xrightarrow{\epsilon \rightarrow 0} \iiint_{\mathbb{R}^{3d}} \left| G((0, v), (\tilde{y}, w)) - G((0, v), (\tilde{y} - z, w)) \right| \chi^K(w) \chi^K(-\tilde{y}) \chi^K(v) p^{12}(t, -\tilde{y}, v) d\tilde{y} dw dv.
\end{aligned}$$

Since the RHS no longer depends on  $\epsilon$ , and also  $p^{12}$  is bounded uniformly by  $\Gamma$ , it suffices to prove that the following converges to 0 as  $|z| \rightarrow 0$ :

$$\iiint_{\mathbb{R}^{3d}} \left| G((0, v), (\tilde{y}, w)) - G((0, v), (\tilde{y} - z, w)) \right| \chi^K(w) \chi^K(-\tilde{y}) \chi^K(v) d\tilde{y} dw dv.$$

By the continuity of  $G$ , the integrand converges to 0 as  $|z| \rightarrow 0$  Lebesgue a.e. Further, due to the compact support of  $w, \tilde{y}, v$  we can apply the finite measure case of Vitali's convergence theorem, if we can check uniform integrability. By the de la Vallée-Poussin theorem it is enough to check that there exists some  $\eta = \eta(d) > 0$  and  $C = C(K, d)$  finite such that

$$\sup_{|z| \leq 1} \iiint_{\mathbb{R}^{3d}} \left| G((0, v), (\tilde{y} - z, w)) \right|^{1+\eta} \chi^K(w) \chi^K(-\tilde{y}) \chi^K(v) d\tilde{y} dw dv < C. \quad (45)$$

Note that the support of  $(0, v)$  is in a compact set, the Euclidean ball  $\mathbb{B}_E((0, 0), 2K)$  in  $\mathbb{R}^{2d}$ . By (31) there exists a ball in the metric  $\rho_L$  of some radius  $K'$  (depending only on  $K$  and  $d$ ) such that for any  $|z| \leq 1$  and  $-\tilde{y}, w$  in the compact support of  $\chi^K$ ,

$$(\tilde{y} - z, w) \in \mathbb{B}_E((0, v), 16K) \cap \mathbb{B}_E((0, 0), 32K) \subset \mathbb{B}_L((0, v), K').$$

Hence, we are in the setting of Proposition 10. By (36) there exists some  $\eta = \eta(d) > 0$  and constant  $C = C(K, K', d)$  finite such that for any fixed  $v$  with  $|v| \leq 2K$  we have that

$$\iint_{\mathbb{B}_L((0, v), K')} \left| G((0, v), (\tilde{y} - z, w)) \right|^{1+\eta} d\tilde{y} dw \leq C.$$

Integrating further in  $|v| \leq 2K$  yields a different constant which proves (45), hence the whole thesis. ■

To prepare for the next proposition, we require a crude pointwise bound on  $H^{\epsilon, z}$  and its partial derivative in  $v$ . It suffices to derive such for the function  $U^\epsilon$ .

**Lemma 13** *Suppose that  $D = 2d \geq 2$ . For any compact set  $\mathbb{K} \subset \mathbb{R}^D$ , there exists some finite constant  $C = C(\mathbb{K}, K, \theta, d)$  such that for any  $(x, v) \in \mathbb{K}$  we have that*

$$|U^\epsilon(x, v)|, |\nabla_v U^\epsilon(x, v)| \leq C \epsilon^{-d}. \quad (46)$$

**Proof.** We prove the thesis for the gradient, the other part being similar. Recall (30), we have that

$$\nabla_v U^\epsilon(x, v) = \int_{\mathbb{R}^{2d}} \nabla_v G((x, v), (y, w)) \theta^\epsilon(y) \chi^K(w) dy dw.$$

Note that, with the function  $\chi^1$  defined in (10), we can write

$$\theta^\epsilon(y) \leq \epsilon^{-d} \|\theta\|_\infty \chi^1(y).$$

Hence, by (37) (with  $\eta = 0$ ), there exists some finite constant  $C = C(\mathbb{K}, K, \theta, d)$  such that for any  $(x, v) \in \mathbb{K}$ , we have

$$|\nabla_v U^\epsilon(x, v)| \leq \epsilon^{-d} \|\theta\|_\infty \int_{\mathbb{R}^{2d}} |\nabla_v G((x, v), (y, w))| \chi^1(y) \chi^K(w) dy dw \leq \epsilon^{-d} C.$$

Indeed, the range of integration of  $(y, w)$  is in  $\mathbb{B}_E((0, 0), 8K)$ , which is inside  $\mathbb{B}_L((x, v), K')$  for some  $K' < \infty$  depending on  $\mathbb{K}, K$ , uniformly for all  $(x, v) \in \mathbb{K}$ . ■

The following proposition bounds (20). An analogous proof can treat (21), which we omit.

**Proposition 14** For any finite  $T$  and  $d \geq 1$ , we have that

$$\begin{aligned} & \lim_{|z| \rightarrow 0} \limsup_{N \rightarrow \infty} \mathbb{E} \int_0^T dt \frac{1}{N^4} \sum_{i=1}^N \left| \sum_{j \neq i} \nabla_v H^{\varepsilon, z}(x_i(t) - x_j(t), v_i(t) - v_j(t)) \phi(x_i(t), v_i(t)) \psi(x_j(t), v_j(t)) \right. \\ & \left. + \sum_{j \neq i} H^{\varepsilon, z}(x_i(t) - x_j(t), v_i(t) - v_j(t)) \psi(x_j(t), v_j(t)) \nabla_v \phi(x_i(t), v_i(t)) \right|^2 = 0. \end{aligned}$$

**Proof.** First, using the elementary inequality  $|\sum_{k=1}^M a_k|^2 \leq M \sum_{k=1}^M |a_k|^2$ , it suffices to bound separately

$$\mathbb{E} \int_0^T dt \frac{1}{N^3} \sum_{i,j=1}^N \left| \nabla_v H^{\varepsilon, z}(x_i(t) - x_j(t), v_i(t) - v_j(t)) \phi(x_i(t), v_i(t)) \psi(x_j(t), v_j(t)) \right|^2 \quad (47)$$

and

$$\mathbb{E} \int_0^T dt \frac{1}{N^3} \sum_{i,j=1}^N \left| H^{\varepsilon, z}(x_i(t) - x_j(t), v_i(t) - v_j(t)) \psi(x_j(t), v_j(t)) \nabla_v \phi(x_i(t), v_i(t)) \right|^2. \quad (48)$$

With the arguments of  $H^{\varepsilon, z}$  in a compact set, we can bound one term of the square in (48) by the crude bound (46) in Lemma 13, then using the relation  $\varepsilon^{-d} \leq N$  in Condition 2, we get that

$$(48) \leq C(K, d, \theta) \|\phi\|_{C^1} \|\psi\|_{\infty} \mathbb{E} \int_0^T dt \frac{1}{N^2} \sum_{i,j=1}^N \left| H^{\varepsilon, z}(x_i(t) - x_j(t), v_i(t) - v_j(t)) \psi(x_j(t), v_j(t)) \nabla_v \phi(x_i(t), v_i(t)) \right|.$$

Its negligibility follows by (a very similar statement and the same proof of) Proposition 12.

Turning to (47), again by (46) and Condition 2, it is sufficient to bound

$$\mathbb{E} \int_0^T dt \frac{1}{N^2} \sum_{i,j=1}^N \left| \nabla_v H^{\varepsilon, z}(x_i(t) - x_j(t), v_i(t) - v_j(t)) \psi(x_j(t), v_j(t)) \phi(x_i(t), v_i(t)) \right|.$$

The proof of the negligibility of the above quantity is analogous to the proof presented in Proposition 12, just using the bound (37) for the gradient instead of (36), hence we only sketch the idea. Indeed, as in (44), we have that

$$\nabla_v H^{\varepsilon, z}(x, v) = \iint_{\mathbb{R}^{2d}} \left[ \nabla_v G((0, v), (\tilde{y}, w)) - \nabla_v G((0, v), (\tilde{y} - z, w)) \right] \theta^\varepsilon(-\tilde{y} - x) \chi^K(w) d\tilde{y} dw.$$

One can now take  $\varepsilon \rightarrow 0$  to obtain

$$\begin{aligned} & \limsup_{\varepsilon \rightarrow 0} \iint_{\mathbb{R}^{2d}} |\nabla_v H^{\varepsilon, z}(x, v)| \chi^K(x) \chi^K(v) p^{12}(t, x, v) dx dv \\ & \leq \iiint_{\mathbb{R}^{3d}} \left| \nabla_v G((0, v), (\tilde{y}, w)) - \nabla_v G((0, v), (\tilde{y} - z, w)) \right| \chi^K(w) \chi^K(-\tilde{y}) \chi^K(v) p^{12}(t, -\tilde{y}, v) d\tilde{y} dw dv. \end{aligned}$$

In order to use Vitali convergence theorem to take  $|z| \rightarrow 0$ , we need to check uniform integrability similar to (45). In this case, the estimate (37) guarantees the existence of some  $\eta = \eta(d) > 0$  and  $C = C(K, d)$  finite such that

$$\sup_{|z| \leq 1} \iiint_{\mathbb{R}^{3d}} \left| \nabla_v G((0, v), (\tilde{y} - z, w)) \right|^{1+\eta} \chi^K(w) \chi^K(-\tilde{y}) \chi^K(v) d\tilde{y} dw dv < C.$$

■

The following proposition bounds (17); the same proof can treat (18), which we omit. For (19), note that upon bounding  $\|\theta^\varepsilon\|_{\infty} \leq \varepsilon^{-d} \|\theta\|_{\infty}$  and using  $\varepsilon^{-d} \leq N$ , it reduces to Proposition 12.

**Proposition 15** For any finite  $T$  and  $d \geq 1$ , we have that

$$\begin{aligned} & \lim_{|z| \rightarrow 0} \limsup_{N \rightarrow \infty} \mathbb{E} \int_0^T dt \frac{1}{N^3} \sum_{i \neq j, i \neq k=1}^N \theta^\varepsilon(x_i(t) - x_k(t)) \\ & \quad \cdot \left| H^{\varepsilon, z}(x_i(t) - x_j(t), v_i(t) - v_j(t)) \phi(x_i(t), v_i(t)) \psi(x_j(t), v_j(t)) \right| = 0. \end{aligned}$$

**Proof.** First, we note that for  $j = k$ , we can bound  $\|\theta^\epsilon\|_\infty \leq \epsilon^{-d}\|\theta\|_\infty$  and with the condition  $\epsilon^{-d} \leq N$ , we have a contribution

$$\begin{aligned} & \mathbb{E} \int_0^T dt \frac{1}{N^3} \sum_{i \neq j=1}^N \theta^\epsilon(x_i(t) - x_j(t)) \left| H^{\epsilon,z}(x_i(t) - x_j(t), v_i(t) - v_j(t)) \phi(x_i(t), v_i(t)) \psi(x_j(t), v_j(t)) \right| \\ & \leq \mathbb{E} \int_0^T dt \frac{1}{N^2} \sum_{i \neq j=1}^N \left| H^{\epsilon,z}(x_i(t) - x_j(t), v_i(t) - v_j(t)) \phi(x_i(t), v_i(t)) \psi(x_j(t), v_j(t)) \right|, \end{aligned}$$

whose negligibility follows from Proposition 12. Hence, it suffices to consider the contribution of  $i \neq j \neq k$  all distinct. By similar considerations as in the beginning of the proof of Proposition 12, it suffices to bound each term of the form

$$\mathbb{E} \int_0^T dt \theta^\epsilon(x_1(t) - x_3(t)) \left| H^{\epsilon,z}(x_1(t) - x_2(t), v_1(t) - v_2(t)) \right| \chi^K(x_1(t)) \chi^K(x_2(t)) \chi^K(v_1(t)) \chi^K(v_2(t)).$$

By independence, the joint density of  $(x_i(t), v_i(t))_{i=1}^3$  is given by  $\prod_{i=1}^3 p(t, x_i, v_i)$ . We write the above expectation explicitly

$$\int_0^T dt \int_{\mathbb{R}^{6d}} \theta^\epsilon(x_1 - x_3) \left| H^{\epsilon,z}(x_1 - x_2, v_1 - v_2) \right| \chi^K(x_1) \chi^K(x_2) \chi^K(v_1) \chi^K(v_2) \prod_{i=1}^3 p(t, x_i, v_i) dx_i dv_i.$$

By (41) estimate for the one-particle density, we integrate first in  $x_3, v_3$

$$\begin{aligned} & \int_{\mathbb{R}^{2d}} \theta^\epsilon(x_1 - x_3) p(t, x_3, v_3) dx_3 dv_3 \\ & \leq \int_{\mathbb{R}^{2d}} \theta^\epsilon(x_1 - x_3) \left[ C e^{-|v|/C} 1_{\{|v| \geq C\}} + \Gamma 1_{\{|v| < C\}} \right] dx_3 dv_3 \leq C'(\Gamma, R, T, d), \end{aligned}$$

where we used that  $\int \theta^\epsilon(x_1 - x_3) dx_3 = 1$  for any  $x_1$ . We are left to bound

$$\int_0^T dt \int_{\mathbb{R}^{4d}} \left| H^{\epsilon,z}(x_1 - x_2, v_1 - v_2) \right| \chi^K(x_1) \chi^K(x_2) \chi^K(v_1) \chi^K(v_2) \prod_{i=1}^2 p(t, x_i, v_i) dx_i dv_i,$$

which (after some rewriting) is the same type of quantity treated in the proof of Proposition 12, and its negligibility follows therefrom. ■

Finally, we remark that the control of (22), (23), (24) reduces to Propositions 15 and 12. Indeed, take (22) for instance, using the elementary inequality  $|\sum_{k=1}^M a_k|^2 \leq M \sum_{k=1}^M |a_k|^2$  and the bound (46), we have that

$$\begin{aligned} (22) &= \mathbb{E} \int_0^T dt \sum_{i \neq k=1}^N \frac{1}{N^5} \theta^\epsilon(x_i^N(t) - x_k^N(t)) \left| \sum_{j: j \neq i, k} H^{\epsilon,z}(x_i^N(t) - x_j^N(t), v_i^N(t) - v_j^N(t)) \phi \psi \right|^2 \\ &\leq \mathbb{E} \int_0^T dt \sum_{i \neq j \neq k=1}^N \frac{1}{N^4} \theta^\epsilon(x_i^N(t) - x_k^N(t)) \left| H^{\epsilon,z}(x_i^N(t) - x_j^N(t), v_i^N(t) - v_j^N(t)) \phi \psi \right|^2 \\ &\leq C \|\phi\|_\infty \|\psi\|_\infty \mathbb{E} \int_0^T dt \sum_{i \neq j \neq k=1}^N \frac{1}{N^3} \theta^\epsilon(x_i^N(t) - x_k^N(t)) \left| H^{\epsilon,z}(x_i^N(t) - x_j^N(t), v_i^N(t) - v_j^N(t)) \phi \psi \right|. \end{aligned}$$

It then reduces to Propositions 15.

We have controlled all the negligible terms, and by the discussions in Section 3, we have established Proposition 5.

## 6 Tightness and subsequential convergence

We show that for any finite  $T$ , the laws of the sequence  $\{\mu_t^N(dx, dv) : t \in [0, T]\}_{N \in \mathbb{N}}$  (2) of  $\mathcal{D}_T(\mathcal{M}_{+,1})$ -valued random variables is tight, hence weakly relatively compact by Prokhorov's theorem, where the Skorohod space  $\mathcal{D}_T(\mathcal{M}_{+,1})$  is defined in (3). Tightness criteria related to Skorohod spaces are well studied, cf. [6, 16] among many other sources. Here we recall a practical criterion to check (cf. [11, Section 6.1]).

**Proposition 16** *If (a). there exists constant finite  $C_1$  such that for any  $N \in \mathbb{N}$  and  $t \in [0, T]$  we have that*

$$\mathbb{E} \int_{\mathbb{R}^{2d}} (|x| + |v|) \mu_t^N(dx, dv) \leq C_1;$$

*and (b). there exists some  $\beta > 0$  such that for every  $\phi \in C_c^\infty(\mathbb{R}^{2d})$  there exists  $C_\phi < \infty$  such that for all  $0 \leq s < t \leq T$ ,*

$$\mathbb{E} \left| \langle \mu_t^N, \phi \rangle - \langle \mu_s^N, \phi \rangle \right| \leq C_\phi |t - s|^\beta;$$

*then the laws of the sequence  $\{\mu_t^N : t \in [0, T]\}_{N \in \mathbb{N}}$  is tight.*

Now we check the two conditions in our case. For (a), by the exchangeability of independent particles  $\{x_i(t), v_i(t)\}_{i=1}^N$  (40) whose trajectories coincide with the active particles in the true system  $\{x_i^N(t)\}_{i=1}^{N(t)}$  (1), we have that

$$\begin{aligned} \mathbb{E} \int_{\mathbb{R}^{2d}} (|x| + |v|) \mu_t^N(dx, dv) &= \frac{1}{N} \sum_{i=1}^{N(t)} \mathbb{E} (|x_i^N(t)| + |v_i^N(t)|) \\ &\leq \frac{1}{N} \sum_{i=1}^N \mathbb{E} (|x_i(t)| + |v_i(t)|) = \mathbb{E} (|x_1(t)| + |v_1(t)|). \end{aligned}$$

Since

$$(x_1(t), v_1(t)) = (x_1(0) + v_1(0)t + \int_0^t B_1(s)ds, v_1(0) + B_1(t))$$

for any  $t \in [0, T]$  and  $(x_1(0), v_1(0))$  is compactly supported in  $\mathbb{B}_E(0, R)$  by Condition 1, we have that

$$\begin{aligned} \mathbb{E}|v_1(t)| &\leq R + \mathbb{E}|B_1(t)| \leq R + c_d \sqrt{T} \\ \mathbb{E}|x_1(t)| &\leq R + RT + \mathbb{E} \int_0^t |B_1(s)|ds \leq R(1 + T) + c_d T^{3/2}. \end{aligned}$$

For (b), by (6) for any  $0 \leq s < t \leq T$  we have that

$$\begin{aligned} \mathbb{E} \left| \langle \mu_t^N, \phi \rangle - \langle \mu_s^N, \phi \rangle \right| &\leq \int_s^t \left| v \cdot \nabla_x \phi + \frac{1}{2} \Delta_v \phi, \mu_u^N \right| du \\ &+ \mathbb{E} \left| \int_s^t \frac{1}{N^2} \sum_{i \in \mathcal{N}(u)} \sum_{j: j \neq i} \theta^\epsilon (x_i^N(u) - x_j^N(u)) [\phi(x_i^N(u), v_i^N(u)) + \phi(x_j^N(u), v_j^N(u))] du \right| \\ &+ (\mathbb{E} |M_t^{\phi, D} - M_s^{\phi, D}|^2)^{1/2} + (\mathbb{E} |M_t^{\phi, J} - M_s^{\phi, J}|^2)^{1/2}, \end{aligned}$$

where, as in the analysis of Section 2 we can bound

$$\begin{aligned} \int_s^t \left| v \cdot \nabla_x \phi + \frac{1}{2} \Delta_v \phi, \mu_u^N \right| du &\leq (K^\phi \|\phi\|_{C^1} + \|\phi\|_{C^2}) |t - s|, \\ \mathbb{E} |M_t^{\phi, D} - M_s^{\phi, D}|^2 &\leq \frac{1}{N^2} \mathbb{E} \int_s^t \sum_{i \in \mathcal{N}(u)} |\nabla_{v_i} \phi(x_i^N(u), v_i^N(u))|^2 du \leq \frac{\|\phi\|_{C^1}^2}{N} |t - s|, \\ c_1 \mathbb{E} |M_t^{\phi, J} - M_s^{\phi, J}|^2 &\leq \frac{\|\phi\|_\infty^2}{N^3} \mathbb{E} \int_s^t \sum_{i \in \mathcal{N}(u)} \sum_{j: j \neq i} \theta^\epsilon (x_i^N(u) - x_j^N(u)) du \\ &\leq \frac{\epsilon^{-d} \|\theta\|_\infty \|\phi\|_\infty^2}{N} |t - s| \leq \|\theta\|_\infty \|\phi\|_\infty^2 |t - s|, \end{aligned}$$

where  $K^\phi$  denotes the maximal radius of the compact support of  $\phi$ , and  $c_1$  is a numerical constant. Further, upon dominating by the independent system, by (41) we have that

$$\begin{aligned} & \mathbb{E} \left| \int_s^t \frac{1}{N^2} \sum_{i \in \mathcal{N}(u)} \sum_{j \neq i} \theta^\epsilon(x_i^N(u) - x_j^N(u)) \phi(x_i^N(u), v_i^N(u)) du \right| \\ & \leq \mathbb{E} \int_s^t \theta^\epsilon(x_1(u) - x_2(u)) |\phi(x_1(u), v_1(u))| du \\ & = \Gamma |t - s| \int_{\mathbb{R}^{4d}} \theta^\epsilon(x_1 - x_2) |\phi(x_1, v_1)| [C e^{-|v_2|/C} 1_{\{|v_2| \geq C\}} + \Gamma 1_{\{|v_2| < C\}}] dx_1 dx_2 dv_1 dv_2 \\ & \leq C(\Gamma, R, \phi) |t - s|, \end{aligned}$$

due to  $\int \theta^\epsilon(x_1 - x_2) dx_2 = 1$  for any  $x_1$  and then  $\phi(x_1, v_1)$  has compact support. Thus, we verified (b) with  $\beta = 1/2$ .

We now show that any weak subsequential limit of  $\{\mu_t^N : t \in [0, T]\}$  is supported on the subset of  $\mathcal{D}_T(\mathcal{M}_{+,1})$  that are absolutely continuous with respect to Lebesgue measure of  $\mathbb{R}^{2d}$ , for every  $t \in [0, T]$ , and its density is dominated from above by  $p(t, x, v)$  of (41). To this end, we consider also the empirical measure of the independent system (40):

$$\mu_t^{N,f}(dx, dv) := \frac{1}{N} \sum_{i=1}^N \delta_{x_i(t)}(dx) \delta_{v_i(t)}(dv), \quad t \in [0, T].$$

Since the continuous paths  $(x_i(t), v_i(t), t \geq 0)_{i=1}^\infty$  are independent and identically distributed, by Varadarajan's theorem ([5, Theorem 11.4.1]), a.s. for every  $t \in [0, T]$

$$\mu_t^{N,f}(dx, dv) \xrightarrow{N \rightarrow \infty} \bar{\mu}_t^f(dx, dv) := p(t, x, v) dx dv,$$

where  $p(t, x, v)$  is the one-particle density (42), and  $\bar{\mu}_t^f$  is deterministic. Consider now the joint process of empirical measures

$$\{\mu_t^N, \mu_t^{N,f} : t \in [0, T]\}_{N \in \mathbb{N}}$$

which is also weakly relatively compact in  $\mathcal{D}_T(\mathcal{M}_{+,1})^2$ . Consider any weak subsequential limit

$$(\mu_t^{N_k}, \mu_t^{N_k,f} : t \in [0, T]) \xrightarrow{k \rightarrow \infty} (\bar{\mu}_t, \bar{\mu}_t^f : t \in [0, T]). \quad (49)$$

By Skorohod's representation theorem, there exists an auxiliary probability space and a sequence of random variables defined thereon with the same distributions as the ones in (49) (by an abuse of notation we denote hereafter by the same letters) such that the convergence in (49) is a.s. By the domination of paths between true and independent systems, a.s. for any open set  $A \subset \mathbb{R}^{2d}$ ,  $t \geq 0$  and  $k \in \mathbb{N}$ ,

$$0 \leq \mu_t^{N_k}(A) \leq \mu_t^{N_k,f}(A).$$

As  $k \rightarrow \infty$ , convergence in Skorohod topology implies that a.s. for any open set  $A$ ,  $t \in [0, T]$ ,

$$0 \leq \bar{\mu}_t(A) \leq \bar{\mu}_t^f(A).$$

Approximating a Borel set  $B \subset \mathbb{R}^{2d}$  by a sequence of open sets  $A_\ell \downarrow B$ , we have that a.s. for any Borel sets  $B$  and  $t \in [0, T]$ ,

$$0 \leq \bar{\mu}_t(B) \leq \bar{\mu}_t^f(B).$$

Since for every  $t$ ,  $\bar{\mu}_t^f$  is absolutely continuous with respect to Lebesgue measure, the same holds for  $\bar{\mu}_t$ , a.s. Further, the density of  $\bar{\mu}_t^f$  is a.s. dominated by  $p(t, x, v)$ , the density of  $\bar{\mu}_t^f$ . Hence, we have proven the following proposition.

**Proposition 17** *For any finite  $T$  and  $d \geq 1$ , the sequence  $\{\mu_t^N(dx, dv) : t \in [0, T]\}_{N \in \mathbb{N}}$  (2) of  $\mathcal{D}_T(\mathcal{M}_{+,1})$ -valued random variables is weakly relatively compact, and any subsequential limit law is concentrated on absolutely continuous paths, with nonnegative densities bounded above by  $p(t, x, v)$  of (41).*

Combining (6), (7), (8), Proposition 6 and Proposition 17, standard arguments can yield that any weak subsequential limit of  $\{\mu_i^N(dx, dv) : t \in [0, T]\}_{N \in \mathbb{N}}$  has a density  $\bar{f}(t, x, v)$  which is a (possibly random) weak solution of the PDE (4), in the sense of Definition 20, provided we can remove the extra test function  $\psi$  from the statement of Proposition 6, which is not present in the equation (6) satisfied by the empirical measure. We will also show in Section 7 that weak solutions to (4) are unique.

To this end, in place of  $\psi(x, v)$  in (9), consider a sequence  $\{\psi^\Lambda(x, v)\}_{\Lambda \in \mathbb{N}}$  of the form  $\psi^\Lambda(x, v) = \psi_1^\Lambda(x)\psi_2^\Lambda(v)$ , where for  $i = 1, 2$ ,

$$\psi_i^\Lambda : \mathbb{R}^d \rightarrow [0, 1]$$

is of class  $C_c^\infty$  that is identically 1 in  $\mathbb{B}_E(0, \Lambda)$  and identically 0 in  $\mathbb{B}_E(0, 2\Lambda)^c$ . For any  $i \neq j$ , we have that

$$\begin{aligned} & \int_0^T \theta^\epsilon(x_i^N(t) - x_j^N(t)) \phi(x_i^N(t), v_i^N(t)) dt - \int_0^T \theta^\epsilon(x_i^N(t) - x_j^N(t)) \phi(x_i^N(t), v_i^N(t)) \psi^\Lambda(x_j^N(t), v_j^N(t)) dt \\ &= \int_0^T \theta^\epsilon(x_i^N(t) - x_j^N(t)) \phi(x_i^N(t), v_i^N(t)) [1 - \psi_1^\Lambda(x_j^N(t)) \psi_2^\Lambda(v_j^N(t))] dt. \end{aligned} \quad (50)$$

Since  $|x_i^N(t) - x_j^N(t)| \leq \epsilon$  in the support of  $\theta^\epsilon$  and  $x_i^N(t)$  is in the compact support of  $\phi$ , for  $\Lambda$  large enough we have  $\psi_1^\Lambda(x_j^N(t)) = 1$  and therefore

$$(50) = \int_0^T \theta^\epsilon(x_i^N(t) - x_j^N(t)) \phi(x_i^N(t), v_i^N(t)) [1 - \psi_2^\Lambda(v_j^N(t))] dt.$$

Furthermore, we have that

**Lemma 18** *For any finite  $T$  and  $d \geq 1$ ,*

$$\lim_{\Lambda \rightarrow \infty} \sup_{N \in \mathbb{N}} \mathbb{E} \int_0^T \frac{1}{N^2} \sum_{i \in \mathcal{N}(t)} \sum_{j: j \neq i} \theta^\epsilon(x_i^N(t) - x_j^N(t)) \left| \phi(x_i^N(t), v_i^N(t)) [1 - \psi_2^\Lambda(v_j^N(t))] \right| dt = 0.$$

**Proof.** We can upper bound by the independent system (40) and consider each term of the form

$$\mathbb{E} \int_0^T \theta^\epsilon(x_1(t) - x_2(t)) \left| \phi(x_1(t), v_1(t)) [1 - \psi_2^\Lambda(v_2(t))] \right| dt.$$

Since  $1 - \psi_2^\Lambda(v_2(t))$  is identically 0 in  $\mathbb{B}_E(0, \Lambda)$  and bounded above by 1 elsewhere, utilizing (41) we have that

$$\begin{aligned} & \mathbb{E} \int_0^T \theta^\epsilon(x_1(t) - x_2(t)) \left| \phi(x_1(t), v_1(t)) [1 - \psi_2^\Lambda(v_2(t))] \right| dt \\ & \leq C(\Gamma, R, T) \int_{\mathbb{R}^{4d}} \theta^\epsilon(x_1 - x_2) 1_{\{|x_1|, |v_1| \leq K\phi\}} e^{-|v_2|/C} 1_{\{|v_2| \geq \Lambda\}} dx_1 dx_2 dv_1 dv_2 \\ & \leq C(\Gamma, R, T, \phi) \int_{\mathbb{R}^d} e^{-|v_2|/C} 1_{\{|v_2| \geq \Lambda\}} dv_2 \\ & \leq C e^{-\Lambda/C}, \end{aligned}$$

where  $K\phi$  denotes the maximal radius of the compact support of  $\phi$ , and  $\int \theta^\epsilon(x_1 - x_2) dx_2 = 1$  for every  $x_1$ . Note also  $1/N^2$  cancels with the cardinality of the double sum, and the limit as  $\Lambda \rightarrow \infty$  of the above expression is 0, uniformly in  $N$ . ■

Since Proposition 6 applies for  $\psi = \psi^\Lambda$  for each  $\Lambda > 0$ , (50) and Lemma 18 yield a version of Proposition 6 without the extra test function.

**Proposition 19** *For any finite  $T$  and  $d \geq 1$ ,*

$$\begin{aligned} & \lim_{\delta \rightarrow 0} \limsup_{N \rightarrow \infty} \mathbb{E} \left| \int_0^T \frac{1}{N^2} \sum_{i \in \mathcal{N}(t)} \sum_{j: j \neq i} \theta^\epsilon(x_i^N(t) - x_j^N(t)) \phi(x_i^N(t), v_i^N(t)) dt \right. \\ & \quad \left. - \int_0^T \int_{\mathbb{R}^d} dw \langle \eta^\delta(w - x_1) \eta^\delta(w - x_2) \phi(w, v_1), \mu_t^N(dx_1, dv_1) \mu_t^N(dx_2, dv_2) \rangle dt \right| = 0. \end{aligned}$$

This together with (6), Proposition 17 and standard arguments completes the proof of subsequential convergence, see e.g. [10, pp. 636-637] for a closely related context.

## 7 Uniqueness of weak solution

**Definition 20** Assume  $f_0 \in L^1_+(\mathbb{R}^{2d})$  (the space of nonnegative integrable functions) satisfies Condition 1, we say that

$$f \in L^\infty([0, T], L^1_+(\mathbb{R}^{2d})) \quad (51)$$

is a weak solution to

$$\begin{cases} \partial_t f(t, x, v) &= -v \cdot \nabla_x f + \frac{1}{2} \Delta_v f - 2 \int_{\mathbb{R}^d} f(t, x, v) f(t, x, v') dv' \\ f|_{t=0} &= f_0(x, v) \end{cases}, \quad (t, x, v) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d \quad (52)$$

if for any  $\phi \in C_c^\infty(\mathbb{R}^{2d})$  we have that

$$\begin{aligned} \int_{\mathbb{R}^{2d}} f(T, x, v) \phi(x, v) dx dv &= \int_{\mathbb{R}^{2d}} f_0(x, v) \phi(x, v) dx dv + \int_0^T \int_{\mathbb{R}^{2d}} (v \cdot \nabla_x f(t, x, v) + \frac{1}{2} \Delta_v f(t, x, v)) \phi(x, v) dx dv dt \\ &\quad - 2 \int_0^T \int_{\mathbb{R}^{3d}} f(t, x, v) f(t, x, v') \phi(x, v) dx dv dv' dt. \end{aligned}$$

**Proposition 21** For every finite  $T$  and  $d \geq 1$ , weak solutions to (52) are unique.

**Proof.** As in Lemma 11, we denote the operator

$$L^* = -v \cdot \nabla_x + \frac{1}{2} \Delta_v,$$

and by  $P(t)$  the diffusion Markov semigroup associated with  $L^*$ . Let  $f_1, f_2$  be two weak solutions of (52), and  $\bar{f} := f_1 - f_2$ . Then,  $\bar{f}$  satisfies in weak form

$$\begin{cases} (\partial_t - L^*) \bar{f}(t, x, v) &= -2 \int_{\mathbb{R}^d} f_1(t, x, v) \bar{f}(t, x, v') dv' - 2 \int_{\mathbb{R}^d} \bar{f}(t, x, v) f_2(t, x, v') dv' \\ \bar{f}|_{t=0} &= 0. \end{cases}$$

By the equivalence between weak and mild formulations we have that

$$\bar{f}(t, x, v) = -2 \int_0^t P(t-s) \left( \int_{\mathbb{R}^d} f_1(s, \cdot, \cdot) \bar{f}(s, \cdot, v') dv' + \int_{\mathbb{R}^d} \bar{f}(s, \cdot, \cdot) f_2(s, \cdot, v') dv' \right) ds. \quad (53)$$

Since  $P(t)$  is a contraction in  $L^1_+(\mathbb{R}^{2d})$ , cf. [23, page 225, last line], we have by taking  $L^1$ -norm of (53) that

$$\int_{\mathbb{R}^{2d}} |\bar{f}(t, x, v)| dx dv \leq 2 \int_0^t \int_{\mathbb{R}^{3d}} f_1(s, x, v) |\bar{f}(s, x, v')| dx dv dv' ds + 2 \int_0^t \int_{\mathbb{R}^{3d}} |\bar{f}(s, x, v)| f_2(s, x, v') dx dv dv' ds. \quad (54)$$

By a simple comparison, we have that

$$f_i(t, x, v) \leq g(t, x, v), \quad i = 1, 2,$$

where  $g$  is the unique weak solution to

$$\begin{cases} (\partial_t - L^*) g(t, x, v) = 0 \\ g|_{t=0} = f_0(x, v). \end{cases}$$

Further,  $g(t) = P(t)f_0$  is precisely the function  $p(t, x, v)$  in (41) hence, since  $f_0$  satisfies Condition 1, we have that for some finite constant  $C(T, \Gamma, R, d)$  and uniformly for any  $t \in [0, T]$ ,  $x \in \mathbb{R}^d$

$$\int_{\mathbb{R}^d} f_i(t, x, v) dv \leq \int_{\mathbb{R}^d} g(t, x, v) dv \leq C(T, \Gamma, R, d), \quad i = 1, 2. \quad (55)$$

Hence, we obtain by (54)-(55) that for any  $t \in [0, T]$ ,

$$\begin{aligned} \|\bar{f}(t)\|_{L^1_+(\mathbb{R}^{2d})} &\leq C \int_0^t \int_{\mathbb{R}^{2d}} |\bar{f}(s, x, v')| dx dv' ds + C \int_0^t \int_{\mathbb{R}^{2d}} |\bar{f}(s, x, v)| dx dv ds \\ &= C \int_0^t \|\bar{f}(s)\|_{L^1_+(\mathbb{R}^{2d})} ds. \end{aligned}$$

By Gronwall's lemma, we have that  $\bar{f} = f_1 - f_2 = 0$  in the class (51). ■

Since any weak subsequential limit obtained in Proposition 17 is supported in the class (51), by Proposition 21 we have uniqueness of subsequential limits hence convergence in law of the full sequence  $\{\mu_t^N\}_{N \in \mathbb{N}}$  to the unique weak solution of the PDE, as stated in Theorem 3. Since the limit is deterministic, we actually have convergence in probability.

**Acknowledgments.** The author thanks Prof. F. Flandoli for valuable discussions, corrections and pointers.

## References

- [1] F. Anceschi and S. Polidoro. A survey on the classical theory for Kolmogorov equation. *Le Matematiche* **LXXXV** (2020), 221-258.
- [2] G. Ben Arous and M. Gradinaru. Singularities of hypoelliptic Green functions. *Potential Anal.* **8** (1998), 217–258.
- [3] R.W.R. Darling and J.R. Norris. Differential equation approximations for Markov chains. *Probab. Surveys* **5** (2008), 37-79.
- [4] A. De Masi and E. Presutti. *Mathematical methods for hydrodynamic limits*, volume **1501** of Lecture Notes in Mathematics. Springer-Verlag, Berlin, 1991.
- [5] R.M. Dudley. *Real analysis and probability*. Cambridge studies in advanced mathematics, Cambridge University Press, 2002.
- [6] S.N. Ethier and T.G. Kurtz. *Markov processes: characterization and convergence*. John Wiley & Sons, 1986.
- [7] E. Fedrizzi, F. Flandoli, E. Priola and J. Vovelle. Regularity of stochastic kinetic equations. *Electron. J. Probab.* **48** (2017), 1-42.
- [8] C.L. Fefferman and A. Sánchez-Calle. Fundamental solutions for second order subelliptic operators. *Ann. of Math.* (2) **124** (1986) 247-272.
- [9] F. Flandoli, M. Gubinelli and E. Priola. Well-posedness of the transport equation by stochastic perturbation. *Invent. Math.* **180** (2010), 1-53.
- [10] F. Flandoli and R. Huang. The KPP equation as a scaling limit of locally interacting Brownian particles. *J. Differ. Equations* **303** (2021), 608-644.
- [11] F. Flandoli and R. Huang. Coagulation dynamics under environmental noise: Scaling limit to SPDE. *arXiv:2111.10859*.
- [12] F. Flandoli, M. Leimbach and C. Olivera. Uniform convergence of proliferating particles to the FKPP equation. *J. Math. Anal. Appl.*, **473** (2019), 27-52.
- [13] A. Hammond and F. Rezakhanlou. The kinetic limit of a system of coagulating Brownian particles. *Arch. Rat. Mech. Anal.* **185** (2007), 1-67.
- [14] L. Hörmander. Hypoelliptic second order differential equations. *Acta Math.* **119** (1967), 147-171.
- [15] P.-E. Jabin and Z. Wang. Mean field limit for stochastic particle systems. In: *Active Particles, Volume 1*, 379-402. Birkhäuser, Cham, 2017.
- [16] C. Kipnis and C. Landim. *Scaling limits of interacting particle systems*, volume **320** of Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, 1999.
- [17] A. Kolmogoroff. Zufällige Bewegungen (zur Theorie der Brownschen Bewegung). *Ann. of Math.* (2) **35** (1934), 116-117.
- [18] R. Lang and N.X. Xanh. Smoluchowski's theory of coagulation in colloids holds rigorously in the Boltzmann-Grad-limit. *Z. Wahrsch. Verw. Gebiete* **54** (1980), 227-280.

- [19] S. Méléard and S. Roelly-Coppoletta. A propagation of chaos result for a system of particles with moderate interaction. *Stochastic Process. Appl.* **26** (1987), 317-332.
- [20] A. Nagel, E.M. Stein and S. Wainger. Balls and metrics defined by vector fields I: Basic properties. *Acta Math.* **155** (1985), 103-147.
- [21] K. Oelschläger. A law of large numbers for moderately interacting diffusion processes. *Z. Wahrsch. Verw. Gebiete* **69** (1985), 279–322.
- [22] K. Oelschläger. On the derivation of reaction-diffusion equations as limit dynamics of systems of moderately interacting stochastic processes. *Probab. Theory Related Fields* **82** (1989), 565-586.
- [23] E. Priola. On weak uniqueness for some degenerate SDEs by global  $L^p$  estimates. *Potential Anal.* **42** (2015), 247-281.
- [24] L.P. Rothschild and E.M. Stein. Hypoelliptic differential operators and nilpotent groups. *Acta Math.* **137** (1977), 247-320.
- [25] A. Sánchez-Calle. Fundamental solutions and geometry of the sum of squares of vector fields. *Invent. Math.* **78** (1984), 143-160.
- [26] K.T. Sturm. Analysis on local Dirichlet spaces. II. Upper Gaussian estimates for the fundamental solutions of parabolic equations. *Osaka J. Math.* **32** (1995), 275-312.
- [27] A.-S. Sznitman. Propagation of chaos for a system of annihilating Brownian spheres. *Comm. Pure Appl. Math.* **XL** (1987), 663-690.
- [28] K. Uchiyama. Pressure in classical statistical mechanics and interacting Brownian particles in multi-dimensions. *Ann. Henri Poincaré* **1** (2000), 1159-1202
- [29] S.R.S. Varadhan. Scaling limits for interacting diffusions. *Comm. Math. Phys.* **135** (1991), 313-353.
- [30] N.Th. Varopoulos. Green's functions on positively curved manifolds. *J. Funct. Anal.* **45** (1982), 109-118.

Author address: Scuola Normale Superiore di Pisa. Piazza Dei Cavalieri 7. Pisa PI 56126. Italia.  
 E-mail: ruojun.huang@sns.it.