

Conservative Binary Dynamics with a Spinning Black Hole at $\mathcal{O}(G^3)$ from Scattering Amplitudes

Fernando Febres Cordero,^{1,*} Manfred Kraus,^{1,†} Guanda Lin,^{2,3,‡} Michael S. Ruf,^{4,§} and Mao Zeng^{3,¶}

¹*Physics Department, Florida State University, Tallahassee, FL 32306-4350, USA*

²*CAS Key Laboratory of Theoretical Physics, Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing, 100190, China*

³*Higgs Centre for Theoretical Physics, University of Edinburgh, James Clerk Maxwell Building, Peter Guthrie Tait Road, Edinburgh, EH9 3FD, United Kingdom*

⁴*Mani L. Bhaumik Institute for Theoretical Physics, University of California at Los Angeles, Los Angeles, CA 90095, USA*

We compute the conservative two-body Hamiltonian of a compact binary system with a spinning black hole through $\mathcal{O}(G^3)$ to all orders in velocity, including linear and quadratic spin terms. To obtain our results we calculate the classical limit of the two-loop amplitude for the scattering of a massive scalar particle with a massive spin-1 particle minimally coupled to gravity. We employ modern scattering amplitude and loop integration techniques, in particular numerical unitarity, integration-by-parts identities, and the method of regions. The conservative potential in terms of rest-frame spin vectors is extracted by matching to a non-relativistic effective field theory. We also apply the Kosower-Maybee-O’Connell (KMOC) formalism to calculate the impulse in the covariant spin formalism directly from the amplitude. We work systematically in conventional dimensional regularization and explicitly evaluate all divergent integrals that appear in full- and effective-theory amplitudes, as well as in the phase-space integrals that arise in the KMOC formalism.

INTRODUCTION

After the momentous observation of gravitational waves produced by the merging of a black hole pair [1], the LIGO-Virgo-KAGRA collaboration continues to explore gravitational waves generated by compact binary systems. The systems so far detected [2] are mainly formed by a pair of black holes with masses in the range of 5 to 100 solar masses, although a handful of events have been associated to systems involving one or two neutron stars. Extracting such details about the binary systems that produced the detected gravitational waves hinges on our ability to precisely predict the structure of such signals based on the parameters of the systems, like for example the masses of the coalescent compact objects, especially in the light of the next-generation gravitational wave detectors [3, 4]. A critical theoretical input to produce these predictions is the conservative Hamiltonian of the two-body system.

Great efforts have been devoted to improve the predictions of binary dynamics through perturbative means. We work in the post-Minkowskian (PM) framework [5–14] which expands in Newton’s constant while keeping the full dependence on velocity. Long considered difficult, high-order PM corrections to the conservative dynamics of spinless binary system enjoyed recent progress using scattering amplitudes and their classical limits [15–17], in combination with effective field theory and advanced integration techniques. The computation of the conservative two-body dynamics at $\mathcal{O}(G^3)$ [18–23] and $\mathcal{O}(G^4)$

[24, 25] for spinless binary systems has been completed.¹

Including spin effects is expected to be of high relevance in order to describe some of the detected binary systems (see e.g. Ref. [2]). This is particularly the case due to known degeneracies between the mass-ratio of the binary system and their spin, which leads to strongly biased parameter inference [40–43]. Therefore obtaining predictions for spin-dependent terms in the conservative potential is a necessity. Related results for up to fifth post-Newtonian (PN) order have appeared employing classical [44–63] and effective field theory methods [64–94]. In the PM approach, progress has been made at the first two PM orders [95–108]. At $\mathcal{O}(G^2)$, the scattering amplitudes approach have produced results up to quadratic-in-spin terms [109, 110], up to quartic-in-spin terms [111], fifth-power in spin [112], and even including all-orders-in-spin contributions [113, 114]. Furthermore, recently a study based on the worldline quantum field theory (QFT) formalism [115–118] has been presented on the impulse and spin kick in the scattering of two spinning compact objects including quadratic-in-spin contributions and up to order $\mathcal{O}(G^3)$ [119]. Proposals to express observables for spinning binaries through a generating function known as the eikonal exponent have appeared in Refs. [109, 121].

There is not a single definition of conservative dynamics in the literature, and here we adopt the definition used by Damour *et. al.* [14, 122] as well as Ref. [25],

¹ The $\mathcal{O}(G^3)$ and $\mathcal{O}(G^4)$ results have also been obtained [26–28] in the worldline approach. See also progress in post-Newtonian corrections in the last few years in Refs. [29–39].

based on a time-symmetric prescription for the graviton propagators, thereby excluding contributions from radiation-reaction and in particular contributions from zero-frequency graviton exchanges. These effects conserve the total energy of the system but in general not the orbital angular momentum. To determine the latter contributions it is efficient to employ complementary approaches based on linear response relations or soft theorems (see e.g. Refs.[119] and [120] for recent applications involving spin; and the latter reference for hidden relations with the spinless case).

In this letter we compute for the first time the conservative binary Hamiltonian with a spinning black hole up to $\mathcal{O}(G^3)$ including terms linear and quadratic in spin.

SCATTERING AMPLITUDES

We make use of the fact that $2 \rightarrow 2$ scattering amplitudes for processes involving spin- s massive particles can be used to fully characterize the conservative spinning dynamics through $2s$ powers of spin [123]. Therefore, we consider a theory of a massive scalar ϕ and a massive spin-1 (vector) field A_μ minimally coupled to gravity, which is described by the following Lagrangian

$$\mathcal{L} = \sqrt{-g} \left[-\frac{2R}{\kappa^2} + \frac{1}{2}g^{\mu\nu}\partial_\mu\phi\partial_\nu\phi - \frac{1}{2}m_\phi^2\phi^2 - \frac{1}{4}g^{\mu\alpha}g^{\nu\beta}F_{\alpha\beta}F_{\mu\nu} + \frac{1}{2}m_A^2g^{\mu\nu}A_\mu A_\nu \right], \quad (1)$$

where $\kappa = \sqrt{32\pi G}$, $g_{\mu\nu}$ is the metric tensor, $g = \det(g_{\mu\nu})$, R is the Ricci scalar associated to $g_{\mu\nu}$, and $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$. Furthermore we work in the weak field approximation, i.e. $g_{\mu\nu} = \eta_{\mu\nu} + \kappa h_{\mu\nu}$, where $h_{\mu\nu}$ is the graviton field. We consider the elastic scattering process $A(p_1, \epsilon_1) + \phi(p_2) \rightarrow \phi(p_3) + A(p_4, \epsilon_4)$, where $p_1^2 = p_4^2 = m_A^2$, $p_2^2 = p_3^2 = m_\phi^2$, and $\epsilon_{1,4}$ are the polarization vectors of the incoming and outgoing vector particles. We work in the center-of-mass (COM) frame in which the momentum transfer takes the form $q^\mu = p_2^\mu - p_3^\mu = (0, \mathbf{q})$. Momenta are decomposed [136, 137] as $p_1 = \bar{p}_1 - q/2$, $p_2 = \bar{p}_2 + q/2$, $p_3 = \bar{p}_2 - q/2$, $p_4 = \bar{p}_1 + q/2$, with $\bar{p}_i \cdot q = 0$. In the COM frame the spatial momenta read $\mathbf{p}_1 = -\mathbf{p}_2 = \bar{\mathbf{p}} - \mathbf{q}/2 \equiv \mathbf{p}$ and $E_{A,\phi} = \sqrt{m_{A,\phi}^2 + \mathbf{p}^2}$. We express the scattering amplitude in terms of the variables $\{m_A, m_\phi, \sigma, q^2\}$ with $\sigma = \frac{p_1 \cdot p_2}{m_A m_\phi}$.

We numerically compute the tree-level, one- and two-loop scattering amplitudes employing the multi-loop numerical unitarity method [129–133], as implemented in the CARAVEL framework [138], using finite field arithmetic to allow for the reconstruction of corresponding analytic expressions [139, 140]. This framework has already been used in the calculation of two-loop amplitudes in gravity [141] and we have extended it here to allow the handling of massive particles. We have also

included the interaction vertices needed in our calculation from the Lagrangian in Eq. (1) with the help of the xACT [142, 143] package.

We decompose the scattering amplitudes in terms of 5 form factors $\mathcal{M} = \sum_{n=1}^5 M_n T_n$, $T_n = \epsilon_{1,\mu} T_n^{\mu\nu} \epsilon_{4,\nu}^*$, with

$$\{T_1^{\mu\nu}, \dots, T_5^{\mu\nu}\} = \{\eta^{\mu\nu}, q^\mu q^\nu, q^2 \bar{p}_2^\mu \bar{p}_2^\nu, \bar{p}_2^\mu q^\nu, \bar{p}_2^\mu q^\nu\}, \quad (2)$$

where $a^{(\mu b^\nu)} = a^\mu b^\nu + a^\nu b^\mu$ and $a^{[\mu b^\nu]} = a^\mu b^\nu - a^\nu b^\mu$. The form factors M_n are obtained by computing 5 different helicity amplitudes and solving numerically for them. Parity invariance and crossing symmetry imply that only $M_{1,2,3,4}$ are present in the amplitude, though M_5 is present in the integrand before loop integration.

To perform the loop integration we employ integration-by-parts (IBP) identities [155], obtained with the FIRE 6 [156] program. This step is highly simplified by employing finite field values for the kinematic invariants and truncating the IBP relations by taking into account the power counting of the contributions relevant to the classical result. After doing an univariate rational reconstruction of the master-integral coefficients in q^2 , we expand in a Laurent series all master integrals and their coefficients for small momentum transfer q^2 and in ϵ , using the results of Ref. [137] and keeping terms up to $\mathcal{O}((q^2)^0)$. The coefficients in this double expansion are yet-unknown rational functions in the variables $\{m_A, m_\phi, \sigma\}$. We then employ multiple numerical evaluations to perform a multivariate rational reconstruction [140]. We are able to reconstruct the analytic expressions for the relevant classical terms of all required amplitudes by employing modest computational resources. The resulting amplitudes are given in the supplemental material, which includes computer-readable files [157].

EFFECTIVE FIELD THEORY

We employ effective field theory (EFT) techniques [16, 109, 134] to extract the two-body Hamiltonian for our system. We construct a non-relativistic EFT with non-local contact interactions described by the Lagrangian

$$L_{\text{EFT}} = \int_{\mathbf{k}} \hat{\phi}^\dagger(-\mathbf{k}) \left(i\partial_t - \sqrt{\mathbf{k}^2 + m_\phi^2} \right) \hat{\phi}(\mathbf{k}) + \int_{\mathbf{k}} \hat{A}^{\dagger,i}(-\mathbf{k}) \left(i\partial_t - \sqrt{\mathbf{k}^2 + m_A^2} \right) \hat{A}^i(\mathbf{k}) - \int_{\mathbf{k}, \mathbf{k}'} \tilde{V}_{ij}(\mathbf{k}, \mathbf{k}') \hat{A}^{\dagger,i}(\mathbf{k}') \hat{A}^j(\mathbf{k}) \hat{\phi}^\dagger(-\mathbf{k}') \hat{\phi}(-\mathbf{k}), \quad (3)$$

with $\int_{\mathbf{k}} \equiv \int \frac{d^3\mathbf{k}}{(2\pi)^3}$. The fields $\hat{\phi}$ and \hat{A} are defined in their respective rest frames. The potential \tilde{V}_{ij} can be decomposed into different spin operators according to

$$\tilde{V}_{ij}(\mathbf{k}, \mathbf{k}') = \sum_{n=1}^4 \tilde{V}^{(n)}(\mathbf{k}, \mathbf{k}') O_n^{ij} \left(\frac{\mathbf{k} + \mathbf{k}'}{2}, \mathbf{k}' - \mathbf{k} \right), \quad (4)$$

where the non-relativistic tensor structures $O_n^{ij}(\bar{\mathbf{p}}, \mathbf{q})$ are

$$\{O_1^{ij}, \dots, O_4^{ij}\} = \{\delta^{ij}, \bar{\mathbf{p}}^{[i} \mathbf{q}^{j]}, \mathbf{q}^i \mathbf{q}^j, \mathbf{q}^2 \bar{\mathbf{p}}^i \bar{\mathbf{p}}^j\}. \quad (5)$$

To connect to the classical spin we relate the non-relativistic tensor structures to spin operators using the representation $(\mathbf{S}_i)_{jk} = -i\epsilon_{ijk}$. We find

$$\begin{aligned} O_1 &= \mathbb{1}, & O_2 &= -i(\mathbf{q} \times \bar{\mathbf{p}}) \cdot \mathbf{S}, \\ O_3 &= \frac{1}{2} \mathbf{q}^2 \mathbf{S}^2 - (\mathbf{q} \cdot \mathbf{S})^2, \\ O_4 &= \mathbf{q}^2 \left(\frac{1}{2} \bar{\mathbf{p}}^2 \mathbf{S}^2 - (\bar{\mathbf{p}} \cdot \mathbf{S})^2 \right), \end{aligned} \quad (6)$$

where we defined $O_n \equiv O_n^{ij}(\bar{\mathbf{p}}, \mathbf{q}) \hat{A}^{\dagger, i}(\bar{\mathbf{p}} - \mathbf{q}/2) \hat{A}^j(\bar{\mathbf{p}} + \mathbf{q}/2)$. We ignore the Casimir operator $O_5 = \mathbf{q}^2 \mathbf{S}^2$ in the amplitude calculation for simplicity in this initial study, though we determine its coefficient in the potential by requiring consistency with the results of Ref. [119] in the aligned-spin limit². Note that the polarization tensor representation of O_n^{ij} in Eq. (5) can be used within conventional dimensional regularization for computing divergent EFT amplitudes, while the spin operators in Eq. (6) are intrinsically 3-dimensional and are used to represent the finite potential obtained by EFT matching. We normalize the coefficients of the operators in Eq. (4) including dimensional regularization factors as

$$\begin{aligned} \tilde{V}^{(n)}(\mathbf{k}, \mathbf{k}') &= \frac{(4\pi)^{3/2-\epsilon}}{\mathbf{q}^2} \sum_{L, Q=0}^{\infty} \left(\frac{\mu^2}{\mathbf{q}^2} \right)^{L\epsilon} \frac{G^{L+1} |\mathbf{q}|^{L+Q}}{2^{(L+1)(1-2\epsilon)}} \\ &\times \frac{\Gamma[1 - \frac{1}{2}(1-2\epsilon)L]}{\Gamma[\frac{1}{2}(1-2\epsilon)(L+1)]} \tilde{c}_{L+1, Q}^{(n)}(\mathbf{k}^2), \end{aligned} \quad (7)$$

where $\mathbf{q} = \mathbf{k}' - \mathbf{k}$ and μ is the dimensional regularization scale, while L denotes the loop order and Q the additional powers of \hbar beyond the classical limit. The unknown coefficients $\tilde{c}_{L, Q}^{(n)}(\mathbf{k}^2)$ are fixed by matching the EFT and full-theory amplitudes.




FIG. 1. The EFT amplitude is given by iterated bubble diagrams. Blue (red) lines refer to scalar (vector) particles. Each potential \tilde{V} has a perturbative expansion in terms of κ .

The EFT amplitude is given by iterated bubble diagrams [16], see Fig. 1. We start by decomposing the numerator of each diagram in terms of the operators O_n by

explicitly constructing projectors. Then, we follow the procedure outlined in Ref. [16], i.e. we integrate out energy components of the loop momenta using the residue theorem. Then the propagators are expanded for in the soft region $\ell \sim |\mathbf{q}|$, yielding the amplitude in terms of the *linearized* $(3 - 2\epsilon)$ -dimensional master integrals of Ref. [137]. We then obtain the L -loop EFT amplitude as an expansion in $|\mathbf{q}|$,

$$\mathcal{M}_{\text{EFT}}^{(L)} = \left(\frac{\kappa}{2} \right)^{2+2L} \left(\frac{\mu^2}{\mathbf{q}^2} \right)^{L\epsilon} \sum_{n=1}^4 \sum_{k=0}^{\infty} \frac{d_{L, k}^{(n)}}{|\mathbf{q}|^{2-k}} O_n. \quad (8)$$

As explained in Refs. [101, 109] to perform the matching to the relativistic loop amplitude the relativistic tensor structures T_n have to be converted into non-relativistic counterparts O_n . The necessary relations are given in the supplemental material [157].

Let us discuss the differences between the EFT matching procedure that we have employed and the one introduced in Ref. [16] and later used in Refs. [18, 19] at the two-loop order and in Ref. [109] for the case of spin. In these references, the integration of the relativistic amplitude is performed by an expansion and re-summation in the velocity. In this process IR-divergent integrals are kept unevaluated and explicitly matched to the corresponding EFT integrals, therefore strictly speaking no dimensional regularization is required. However, this strategy relies on subtle arguments for the cancellation of evanescent effects, i.e. terms subleading in ϵ or $|\mathbf{q}|$ at lower loops being promoted to physically-significant terms after multiplying divergent iteration integrals. Here we explicitly evaluate all such subleading terms in dimensional regularization in a transparent manner. This makes it natural to use dimensional regularization in an relativistic approach to conservative classical dynamics as introduced in Ref. [137] in conjunction with treating the EFT in dimensional regularization as described in this letter.

CLASSICAL HAMILTONIAN

The result of the EFT calculation described before are the matching coefficients $\tilde{c}_{L+1}^{(n)} = \tilde{c}_{L+1, 0}^{(n)}$. For the spin-orbit coupling we find

$$\tilde{c}_{L+1}^{(2)} = \tilde{c}_{L+1, \text{red}}^{(2)} + \tilde{c}_{L+1, \text{iter}}^{(2)} + \frac{\tilde{c}_{L+1, \text{red}}^{(1)}}{m_A^2 (\gamma_1 + 1)}, \quad (9)$$

with $\gamma_1 = E_A/m_A$. The coefficients $\tilde{c}_{L+1, \text{red}}^{(1)}$ are related to the scalar coupling [18, 19], while the $\tilde{c}_{L+1, \text{iter}}^{(2)}$ are given in terms of lower-order coefficients. We find

² Up to $O(G^2)$, this could be fixed by conjectured relations between different spin structures in Refs. [112] and [113].

$$\tilde{c}_{1,\text{red}}^{(1)}(\mathbf{k}^2) = \frac{m_A^2 m_\phi^2}{E^2 \xi} (1 - 2\sigma^2), \quad \tilde{c}_{2,\text{red}}^{(1)}(\mathbf{k}^2) = \frac{3(m_\phi + m_A) m_\phi^2 m_A^2}{4E^2 \xi} (1 - 5\sigma^2), \quad (10)$$

$$\begin{aligned} \tilde{c}_{3,\text{red}}^{(1)}(\mathbf{k}^2) = & \frac{m_A^2 m_\phi^2}{E^2 \xi} \left[-\frac{2}{3} m_A m_\phi \left(\frac{\text{arccosh}(\sigma)}{\sqrt{\sigma^2 - 1}} (-12\sigma^4 + 36\sigma^2 + 9) + 22\sigma^3 - 19\sigma \right) - 2(m_\phi^2 + m_A^2) (6\sigma^2 + 1) \right] \\ & + \frac{3E m_A^2 m_\phi^2}{4E^2 \xi} (m_A + m_\phi) \frac{(1 - 2\sigma^2)(1 - 5\sigma^2)}{(\sigma^2 - 1)} - \frac{3m_A^4 m_\phi^4}{E^2 \xi \mathbf{k}^2}, \end{aligned} \quad (11)$$

$$\tilde{c}_{1,\text{red}}^{(2)}(\mathbf{k}^2) = -\frac{2\sigma m_\phi}{E \xi}, \quad \tilde{c}_{2,\text{red}}^{(2)}(\mathbf{k}^2) = \frac{m_\phi (4m_A + 3m_\phi) \sigma (5\sigma^2 - 3)}{4E \xi (\sigma^2 - 1)}, \quad (12)$$

$$\begin{aligned} \tilde{c}_{3,\text{red}}^{(2)}(\mathbf{k}^2) = & \frac{m_\phi}{E \xi (\sigma^2 - 1)^2} \left[-2m_A^2 \sigma (3 - 12\sigma^2 + 10\sigma^4) - \left(\frac{83}{6} + 27\sigma^2 - 52\sigma^4 + \frac{44}{3}\sigma^6 \right) m_A m_\phi - m_\phi^2 \sigma \left(\frac{7}{2} - 14\sigma^2 + 12\sigma^4 \right) \right. \\ & \left. + \frac{(4m_A + 3m_\phi) E}{4} \sigma (2\sigma^2 - 1) (5\sigma^2 - 3) + 4m_A m_\phi \sigma (\sigma^2 - 6) (2\sigma^2 + 1) \sqrt{\sigma^2 - 1} \text{arccosh}(\sigma) \right], \end{aligned} \quad (13)$$

$$\tilde{c}_{1,\text{iter}}^{(2)}(\mathbf{k}^2) = 0, \quad \tilde{c}_{2,\text{iter}}^{(2)}(\mathbf{k}^2) = E \xi \tilde{c}_1^{(2)} \frac{\partial \tilde{c}_1^{(1)}}{\partial \mathbf{k}^2} + \tilde{c}_1^{(1)} \left(E \xi \frac{\partial \tilde{c}_1^{(2)}}{\partial \mathbf{k}^2} + \frac{\tilde{c}_1^{(2)} \left(\frac{2E^2 \xi}{\mathbf{k}^2} + \frac{1}{\xi} - 3 \right)}{2E} \right), \quad (14)$$

$$\begin{aligned} \tilde{c}_{3,\text{iter}}^{(2)}(\mathbf{k}^2) = & \left(\tilde{c}_1^{(1)} \right)^2 \left(-\frac{2}{3} E^2 \xi^2 \frac{\partial^2 \tilde{c}_1^{(2)}}{\partial (\mathbf{k}^2)^2} + \left(\xi \left(3 - \frac{E^2 \xi}{\mathbf{k}^2} \right) - 1 \right) \frac{\partial \tilde{c}_1^{(2)}}{\partial \mathbf{k}^2} + \tilde{c}_1^{(2)} \left(\frac{\frac{1}{2\xi} - 2}{E^2} + \frac{3\xi - 1}{\mathbf{k}^2} \right) \right) \\ & + \tilde{c}_1^{(1)} \left(\tilde{c}_1^{(2)} \left(\left(-\frac{3E^2 \xi^2}{\mathbf{k}^2} + 6\xi - 2 \right) \frac{\partial \tilde{c}_1^{(1)}}{\partial \mathbf{k}^2} - \frac{4}{3} E^2 \xi^2 \frac{\partial^2 \tilde{c}_1^{(1)}}{\partial (\mathbf{k}^2)^2} \right) \right. \\ & \left. + \frac{4}{3} E \xi \left(\frac{\partial \tilde{c}_2^{(2)}}{\partial \mathbf{k}^2} - 2E \xi \frac{\partial \tilde{c}_1^{(1)}}{\partial \mathbf{k}^2} \frac{\partial \tilde{c}_1^{(2)}}{\partial \mathbf{k}^2} \right) + \frac{E^2 \xi^2 \left(\tilde{c}_1^{(2)} \right)^2}{2\mathbf{k}^2} + \tilde{c}_2^{(2)} \left(\frac{\frac{2}{3\xi} - 2}{E} + \frac{E\xi}{\mathbf{k}^2} \right) \right) - \frac{1}{6} E^2 \xi^2 \left(\tilde{c}_1^{(2)} \right)^3 \\ & + \tilde{c}_1^{(2)} \left(\frac{2}{3} E \xi \left(\frac{\partial \tilde{c}_2^{(1)}}{\partial \mathbf{k}^2} - 2E \xi \left(\frac{\partial \tilde{c}_1^{(1)}}{\partial \mathbf{k}^2} \right)^2 \right) + \frac{\tilde{c}_2^{(1)} \left(\frac{3E^2 \xi}{\mathbf{k}^2} + \frac{1}{\xi} - 3 \right)}{3E} \right) + \frac{2}{3} E \xi \tilde{c}_2^{(1)} \frac{\partial \tilde{c}_1^{(2)}}{\partial \mathbf{k}^2} + \frac{4}{3} E \xi \tilde{c}_2^{(2)} \frac{\partial \tilde{c}_1^{(1)}}{\partial \mathbf{k}^2}, \end{aligned} \quad (15)$$

where $E = E_A + E_\phi$ and $\xi = E_A E_\phi / E^2$. These expressions represent our results for the spin-orbit term in the momentum-space potential in Eq. (7). The corresponding results for the quadratic-in-spin terms are given in the supplemental material [157]. Finally we convert to position space and obtain the Hamiltonian by computing the Fourier transform of the potential in Eq. (7). The resulting Hamiltonian reads

$$\begin{aligned} H = & \sqrt{\mathbf{p}^2 + m_A^2} + \sqrt{\mathbf{p}^2 + m_\phi^2} + V^{(1)}(\mathbf{r}^2, \mathbf{p}^2) \\ & + V^{(2)}(\mathbf{r}^2, \mathbf{p}^2) \frac{\mathbf{L} \cdot \mathbf{S}}{r^2} + V^{(3)}(\mathbf{r}^2, \mathbf{p}^2) \frac{(\mathbf{r} \cdot \mathbf{S})^2}{r^4} \\ & + V^{(4)}(\mathbf{r}^2, \mathbf{p}^2) \frac{(\mathbf{p} \cdot \mathbf{S})^2}{r^2} + V^{(5)}(\mathbf{r}^2, \mathbf{p}^2) \frac{\mathbf{S}^2}{r^2}, \end{aligned} \quad (16)$$

where each coefficient is expanded as

$$V^{(n)}(\mathbf{r}^2, \mathbf{p}^2) = \sum_{L=1}^3 \left(\frac{G}{r} \right)^L c_L^{(n)}(\mathbf{p}^2) + \mathcal{O}(G^4). \quad (17)$$

The scalar potential $V^{(1)}(\mathbf{r}^2, \mathbf{p}^2)$ coincides with the results of Ref. [18, 19]. The $c_L^{(n)}(\mathbf{p}^2)$ coefficients are pro-

vided in the supplemental material [157] as linear sums of the $\tilde{c}_{L,0}^{(n)}(\mathbf{p}^2)$ coefficients in Eq. (7).

Validation. We performed several checks of our computational framework in dimensional regularization. We checked that the EFT amplitudes reduce to their scalar counterparts given in Eq. (10.6) of Ref. [19] when setting $\tilde{c}_{L,Q}^{(n)} = 0$ for $n > 1$. Even more, we reproduced the two-loop matching coefficients of Refs. [18, 19]. For the EFT calculation including spin, we cross checked our tree-level and one-loop matching coefficients against Refs. [110, 158]. The scalar coefficients up to one-loop are also in agreement with Ref. [159] to all orders in ϵ . Beyond one-loop we checked explicitly the cancellation of the *super-classical* terms, i.e. unphysical terms behaving as powers of $1/\hbar \sim 1/|\mathbf{q}|$, where we observe a non-trivial interplay between the various matching coefficients. Furthermore the coefficients extracted in this way are finite in the non-relativistic limit $|\mathbf{p}| \ll 1$ (this can be seen explicitly from the post-Newtonian expanded coefficients provided in the supplemental material [157]).

We solved the equation of motion based on the com-

puted conservative potential to obtain the impulse in either the limit of an aligned spin or the limit of a spinless probe moving in a Kerr background. After translating to the covariant spin formalism using the procedure of Refs. [91, 97], the impulse agrees with the recent worldline QFT result of Ref. [119] in either of the two above limits, when specializing to the Kerr black hole case. The result is also consistent with the scattering angle in the probe limit obtained in Ref. [98]. Furthermore, the velocity expansion of our Hamiltonian is consistent with the next-to-next-to-leading order post-Newtonian results for spin-orbit and spin-squared coupling [79, 80] calculated by the EFTofPNG package [160], after finding a canonical transformation.³

OBSERVABLES FROM THE KMOC FORMALISM

We employ the full-theory amplitudes we obtained to compute classical gravitational observables using the KMOC formalism [17]. In this formalism, a change of classical variable O , associated with any operator \mathcal{O} , between *in* and *out* states is measured as $\Delta O = \langle \text{out} | \mathcal{O} | \text{out} \rangle - \langle \text{in} | \mathcal{O} | \text{in} \rangle$. Relating *in* and *out* states by the S-matrix allows to reformulate the problem in terms of scattering amplitudes. In the classical limit ΔO is given by [17]

$$\Delta O = \int \frac{d^D q}{(2\pi)^{D-2}} \delta(-2p_1 \cdot q) \delta(2p_2 \cdot q) e^{ib \cdot q} (\mathcal{I}_{O,v} + \mathcal{I}_{O,r}) \quad (18)$$

where $\mathcal{I}_{O,v}$ is the *virtual* kernel obtained from virtual amplitudes, while $\mathcal{I}_{O,r}$ is the *real* kernel depending on cut amplitudes with phase-space integration.

We begin with obtaining kernels \mathcal{I}_v and \mathcal{I}_r in order to compute the momentum impulse Δp_i (corresponding to the momentum operator \mathbb{P}_i). The spinless case has been discussed in Ref. [161] and we highlight new features introduced by polarizations and spins here. The *virtual* kernel $\mathcal{I}_{\Delta p,v}^\mu$ is simply $q^\mu \mathcal{M}$, which can be expanded with the tensors in Eq. (2). For the conservative dynamics we only need to consider two-particle cuts and the real kernel takes the form

$$\mathcal{I}_{\Delta p,r}^\mu = \int d\Phi_2 \sum_{\epsilon_c} \ell^\mu \mathcal{M}_L \mathcal{M}_R^*, \quad (19)$$

where $d\Phi_2$ is the two-body phase-space measure and \mathcal{M}_R and \mathcal{M}_L^* are amplitudes for the processes $A(p_1, \epsilon_1) + \phi(p_2) \rightarrow \phi(p_1 + \ell) + A(p_2 - \ell, \epsilon_c)$, and $\phi(p_1 + \ell) + A(p_2 - \ell, \epsilon_c) \rightarrow \phi(p_3) + A(p_4, \epsilon_4)$, respectively. The sum

runs over the polarizations of the vector particles in the cut. These one-loop integrals can be evaluated directly through sub-loop integration, for details see Ref. [21]. To expand the double-cut in terms of form factors, we need the additional tensor structures

$$\{T_6^{\mu\nu}, \dots, T_9^{\mu\nu}\} = \{\bar{p}_2^{[\mu} n^{\nu]}, \bar{p}_2^{(\mu} n^{\nu)}, q^{[\mu} n^{\nu]}, q^{(\mu} n^{\nu)}\}, \quad (20)$$

where $n^\mu = \epsilon^{\mu\nu\sigma\rho} q_\nu p_{1,\sigma} p_{2,\rho}$. After combining \mathcal{I}_r and \mathcal{I}_v , we find that all super-classical terms cancel.

The next step is to translate the polarization vectors into spin vectors. Following Ref. [135], we first boost the polarization vector ϵ_4^* into the frame of p_1 . With the Pauli-Lubanski operator $\mathbb{W}^\mu = \epsilon^{\mu\nu\sigma\rho} \mathbb{P}_\nu \mathbb{J}_{\rho\sigma}/2$, we define the spin vector s_1^μ as the expectation value of \mathbb{W}_1^μ/m_1 . It is straightforward to convert from ϵ_1 to s_1 given $s_1^\mu = i\epsilon^{\mu\nu\rho\sigma} p_{1,\nu} \epsilon_\rho(p_1) \epsilon_\sigma^*(p_1)/m_1$. Finally, we perform the Fourier transform from the momentum space to the impact parameter space as in Eq. (18). We compare our momentum impulse results up to $\mathcal{O}(G^3)$ with Ref. [91, 119] and find agreement in the Kerr black hole case. This serves as yet another check on the correctness of our amplitude and confirms the practical value of the KMOC formalism at nontrivial PM orders beyond the spinless case [21, 161]. There is no obstruction to computing other observables such as the spin kick using the KMOC formalism, which we leave to future work.

CONCLUSIONS

In this letter we have presented the conservative two-body Hamiltonian for a compact binary system with a spinning black hole at $\mathcal{O}(G^3)$ with exact velocity dependence and including terms linear and quadratic in spin. Though obtained from scattering, the Hamiltonian is directly applicable to bound orbits. This is especially valuable since analytic continuation for observables in these two kinds of orbits, through the boundary-to-bound map [162–164], has not been worked out for generic spin configurations. The computation employs powerful modern scattering amplitude techniques, including numerical unitarity, integration-by-parts identities, expansion-by-regions and functional reconstruction algorithms.

We have shown that the effective field theory approach to study the binary dynamics with spin [109, 123] can be applied in complex high-order calculations. Furthermore, we have performed matching calculations for the effective field theory of Ref. [16] at the two-loop level, for both the spinless case and spinning case, fully within dimensional regularization for the first time. We have also demonstrated the power of the KMOC formalism beyond the lowest order in G for spinning binary dynamics.

There are many directions to extend the applicability of our framework. For instance, one can study Hamiltonian terms with higher powers in spin by con-

³ We thank Justin Vines for carrying out the check, and thank Jan Steinhoff for sharing a private code used for the canonical transformation and comparison.

sidering minimally-coupled massive higher-spin particles [102, 123, 165]. Also including finite-size [166–173] and radiation [120, 150, 174–192] effects is an interesting possibility of relevance for phenomenological studies in gravitational wave detectors. Finally, our framework can be extended beyond the third-post-Minkowskian order following progress in the spinless case [24, 25, 27, 28].

Acknowledgements: We thank Harald Ita for collaboration during the initial phase of the project. We thank Jan Steinhoff and Justin Vines for their expertise and timely help in checking the consistency of our Hamiltonian with known post-Newtonian results. We thank Andres Luna, Chia-Hsien Shen, Dimitrios Kosmopoulos, Fei Teng, Justin Vines, Radu Roiban, and Zvi Bern for insightful discussions and/or comments on the manuscript. The work of F.F.C. and M.K. is supported in part by the

U.S. Department of Energy under grant DE-SC0010102. G.L. is supported in part by the National Natural Science Foundation of China Grants No. 11935013. G.L. thanks the Higgs Centre for Theoretical Physics at the University of Edinburgh for the Visiting Researcher Scheme. M.Z.’s work is supported in part by the U.K. Royal Society through Grant URF\R1\20109. M.S.R. thanks the Higgs Centre for Theoretical Physics at the University of Edinburgh for hospitality and the Mani L. Bhaumik Institute for generous support.

The computing for this project was performed on the HPC cluster at the Research Computing Center at the Florida State University (FSU). For the purpose of open access, the author have applied a Creative Commons Attribution (CC BY) license to any Author Accepted Manuscript version arising from this submission.

Supplementary Material

Full Theory Scattering Amplitudes

Here we summarize the form factor decomposition of the $2 \rightarrow 2$ scattering amplitudes in Eq. (2) of the main text in conventional dimensional regularization.

Tree Level

We decompose the tree-level scattering amplitude according to

$$\mathcal{M}^{(0)} = \left(\frac{\kappa}{2}\right)^2 \frac{1}{\mathbf{q}^2} \sum_{n=1}^4 M_n^{(0)} T_n + \mathcal{O}(|\mathbf{q}|^0), \quad (21)$$

with

$$M_1^{(0)} = 2m_A^2 m_\phi^2 \left(\frac{1}{1-\epsilon} - 2\sigma^2\right), \quad M_2^{(0)} = \frac{2m_\phi^2}{1-\epsilon}, \quad M_3^{(0)} = 2, \quad M_4^{(0)} = -4m_A m_\phi \sigma. \quad (22)$$

One Loop

The one-loop scattering amplitude is given in the form of

$$\mathcal{M}^{(1)} = \left(\frac{\kappa}{2}\right)^4 \frac{1}{\mathbf{q}^2} \sum_{n=1}^4 \sum_{k=0}^2 M_n^{(1,k)} |\mathbf{q}|^k T_n + \mathcal{O}(|\mathbf{q}|). \quad (23)$$

The super-classical terms are given by

$$M_1^{(1,0)} = -\frac{2m_A^3 m_\phi^3 f_2}{\sqrt{\sigma^2 - 1}} \left(\frac{1}{1-\epsilon} - 2\sigma^2\right)^2, \quad (24)$$

$$M_2^{(1,0)} = -\frac{m_A m_\phi^3 f_2}{(1-2\epsilon)\sqrt{\sigma^2 - 1}} \left[\frac{2-3\epsilon}{(1-\epsilon)^2} - 4\sigma^2(1-\epsilon)\right], \quad (25)$$

$$M_3^{(1,0)} = \frac{m_A m_\phi f_2}{\sqrt{\sigma^2 - 1}} \left[\frac{1-2\epsilon}{2(\sigma^2 - 1)(1-\epsilon)^2} + 4\sigma^2 - \frac{2\epsilon}{(1-\epsilon)(1-2\epsilon)}\right], \quad (26)$$

$$M_4^{(1,0)} = \frac{4m_A^2 m_\phi^2 \sigma f_2}{\sqrt{\sigma^2 - 1}} \left[\frac{1}{1-\epsilon} - 2\sigma^2\right]. \quad (27)$$

The classical contributions read

$$M_1^{(1,1)} = \frac{m_A^2 m_\phi^2 (m_A + m_\phi) f_1}{4(\sigma^2 - 1)} \frac{1 - 2\epsilon}{(1 - \epsilon)^2} \left(-3 + (3 - 4\epsilon)\sigma^2(6 - (5 - 4\epsilon)\sigma^2) \right), \quad (28)$$

$$M_2^{(1,1)} = \frac{m_\phi^2 f_1}{32(\sigma^2 - 1)} \left[\frac{(5 - 4\epsilon)(1 - 12\epsilon + 4\epsilon^2)}{(1 - \epsilon)^2} m_A \sigma^4 - \frac{4(5 - 2\epsilon)(1 - 2\epsilon)}{(1 - \epsilon)^2} m_\phi \right. \\ \left. + \frac{4(1 - \epsilon)(5 - 2\epsilon)(1 - 2\epsilon)(3 - 4\epsilon)}{(1 - \epsilon)^3} m_\phi \sigma^2 - \frac{27 - \epsilon(67 - 4\epsilon(14 - 3\epsilon))}{(1 - \epsilon)^3} m_A \right. \\ \left. + \frac{(78 - 2\epsilon(147 - 2\epsilon(113 - 2\epsilon(35 - 6\epsilon)))}{(1 - \epsilon)^3} m_A \sigma^2 \right], \quad (29)$$

$$M_3^{(1,1)} = \frac{f_1}{16(\sigma - 1)^2} \left[\frac{2 - 4\epsilon(3 - 4\epsilon)}{(1 - \epsilon)^2} m_\phi - \frac{8(1 - 2\epsilon)(3 - 4\epsilon)}{1 - \epsilon} m_\phi \sigma^2 + \frac{2(1 - 2\epsilon)(5 - 4\epsilon)(3 - 4\epsilon)}{(1 - \epsilon)^2} m_\phi \sigma^4 \right. \\ \left. + \frac{9 - 10\epsilon + 8\epsilon^2}{(1 - \epsilon)^2} m_A - \frac{2(33 - 2\epsilon(45 - 4\epsilon(11 - 4\epsilon)))}{(1 - \epsilon)^2} m_A \sigma^2 + \frac{(5 - 4\epsilon)(13 - 2\epsilon(15 - 8\epsilon))}{(1 - \epsilon)^2} m_A \sigma^4 \right], \quad (30)$$

$$M_4^{(1,1)} = \frac{m_A m_\phi \sigma f_1}{4(\sigma^2 - 1)} \frac{1 - 2\epsilon}{1 - \epsilon} \left[m_A(12 - 4(5 - 4\epsilon)\sigma^2) + m_\phi \left(\frac{3(3 - 4\epsilon)}{1 - \epsilon} - \frac{(5 - 4\epsilon)(3 - 4\epsilon)}{1 - \epsilon} \sigma^2 \right) \right]. \quad (31)$$

And finally, the quantum corrections correspond to

$$M_1^{(1,2)} = \frac{m_A m_\phi f_2}{\sqrt{\sigma^2 - 1}} \left[\frac{\epsilon}{2} \left(\frac{1}{1 - \epsilon} - 2\sigma^2 \right)^2 \frac{(m_A + m_\phi \sigma)^2}{\sigma^2 - 1} - m_\phi^2 \left(\frac{1 + 2\epsilon}{2(1 - \epsilon)(1 - 2\epsilon)} - \frac{4\epsilon}{1 - 2\epsilon} \sigma^2 + 2\epsilon \sigma^4 \right) \right], \quad (32)$$

$$M_2^{(1,2)} = \frac{m_\phi f_2}{8(\sigma^2 - 1)^{3/2}} \frac{1}{(1 - 2\epsilon)(1 - \epsilon)^2} \left[-2(1 - 6\epsilon + 3\epsilon^2) \frac{m_\phi^2}{m_A} + (8 - 44\epsilon + 64\epsilon^2 - 40\epsilon^3 + 8\epsilon^4) \frac{m_\phi^2 \sigma^2}{m_A} \right. \\ \left. - 4(1 - 7\epsilon + 7\epsilon^2) m_\phi \sigma + 16(1 - \epsilon)^2 (1 - 4\epsilon + \epsilon^2) m_\phi \sigma^3 - (1 - 4\epsilon + 6\epsilon^2) m_A \right. \\ \left. + 4\epsilon(1 + 3\epsilon - 6\epsilon^2 + 2\epsilon^3) m_A \sigma^2 + 8(1 - \epsilon)^2 (1 - 4\epsilon) m_A \sigma^4 \right], \quad (33)$$

$$M_3^{(1,2)} = \frac{f_2}{m_A m_\phi (\sigma^2 - 1)^{5/2}} \frac{1}{8(1 - 2\epsilon)} \left[\frac{3}{1 - \epsilon} m_A^2 - 4(6 - \epsilon(7 - 4\epsilon)) m_A^2 \sigma^2 + 8(1 - \epsilon)(3 - 2\epsilon) m_A^2 \sigma^4 \right. \\ \left. - \frac{1 + \epsilon}{(1 - \epsilon)^2} m_\phi^2 - \frac{4 - 2\epsilon(13 - 2\epsilon(10 - \epsilon(9 - 4\epsilon)))}{(1 - \epsilon)^2} m_\phi^2 \sigma^2 + \frac{8 - 16\epsilon(2 - (2 - \epsilon)\epsilon)}{1 - \epsilon} m_\phi^2 \sigma^4 \right. \\ \left. - \left(16 - \frac{6}{1 - \epsilon} \right) m_A m_\phi \sigma + 8(3 - 4\epsilon)\epsilon m_A m_\phi \sigma^3 + 16(1 - \epsilon)(1 - 2\epsilon) m_A m_\phi \sigma^5 \right], \quad (34)$$

$$M_4^{(1,2)} = \frac{f_2}{(\sigma^2 - 1)^{3/2}} \left[m_A^2 \sigma - 2(1 - \epsilon) m_A^2 \sigma^3 + \frac{1}{4(1 - \epsilon)^2} m_A m_\phi - \frac{\epsilon}{1 - \epsilon} m_A m_\phi \sigma^2 - (2 - 4\epsilon) m_A m_\phi \sigma^4 \right. \\ \left. + \left(1 + \frac{1}{2(1 - \epsilon)} + \frac{1}{2(1 - \epsilon)^2} - \frac{1}{1 - 2\epsilon} \right) m_\phi^2 \sigma - \left(1 - 2\epsilon - \frac{1}{1 - 2\epsilon} + \frac{2}{1 - \epsilon} \right) m_\phi^2 \sigma^3 \right]. \quad (35)$$

Here, we have defined

$$f_1 \equiv \frac{1}{4\pi} \left(\frac{4\pi\mu^2}{\mathbf{q}^2} \right)^\epsilon \frac{\Gamma^2(\frac{1}{2} - \epsilon)\Gamma(\frac{1}{2} + \epsilon)}{2\sqrt{\pi}\Gamma(1 - 2\epsilon)}, \quad f_2 \equiv \frac{1}{4\pi} \left(\frac{4\pi\mu^2}{\mathbf{q}^2} \right)^\epsilon \frac{i\Gamma^2(-\epsilon)\Gamma(1 + \epsilon)}{4\Gamma(-2\epsilon)}. \quad (36)$$

Two Loops

The two-loop scattering amplitude is given in the form of

$$\mathcal{M}^{(2)} = - \left(\frac{\kappa}{2} \right)^6 \frac{1}{\mathbf{q}^2} \mathcal{N} \sum_{n=1}^4 \sum_{k=0}^2 M_n^{(2,k)} |\mathbf{q}|^k T_n + \mathcal{O}(|\mathbf{q}|), \quad (37)$$

where we keep a normalization factor of

$$\mathcal{N} = \left(\frac{\mu^2}{\mathbf{q}^2}\right)^{2\epsilon} = \left(\frac{\mu^2}{\mathbf{q}^2}\right)^{2\epsilon} \left(\frac{e^{\gamma_E}}{4\pi}\right)^{-2\epsilon} \quad (38)$$

unexpanded. Note that, at the $\mathcal{O}(1/|\mathbf{q}|)$ order all master integrals are purely imaginary and therefore do not contribute to the classical limit. Furthermore, to extract the classical terms it is enough to know the $1/\epsilon$ poles of the scattering amplitudes which are equal to $-1/2$ times the coefficients of $\log(\mathbf{q}^2)$. With this, for the super-classical terms at $\mathcal{O}(1/\mathbf{q}^2)$ we obtain

$$M_1^{(2,0)} = \frac{m_A^4 m_\phi^4 (1-2\sigma^2)^2}{(8\pi)^2 (\sigma^2-1)} \left[\frac{1-2\sigma^2}{\epsilon^2} + \frac{3}{\epsilon} \right] + \mathcal{O}(\epsilon^0), \quad (39)$$

$$M_2^{(2,0)} = \frac{m_A^2 m_\phi^4 (1-2\sigma^2)}{(8\pi)^2 (\sigma^2-1)} \left[\frac{1-2\sigma^2}{\epsilon^2} + \frac{4-2\sigma^2}{\epsilon} \right] + \mathcal{O}(\epsilon^0), \quad (40)$$

$$M_3^{(2,0)} = \frac{m_A^2 m_\phi^2}{(16\pi)^2 (\sigma^2-1)^2} \left[\frac{(2\sigma^2-1)(1-8\sigma^2+8\sigma^4)}{\epsilon^2} + \frac{2(3-8\sigma^2)(\sigma^2-1)}{\epsilon} \right] + \mathcal{O}(\epsilon^0), \quad (41)$$

$$M_4^{(2,0)} = -\frac{m_A^3 m_\phi^3 \sigma (1-2\sigma^2)}{(4\pi)^2 (\sigma^2-1)} \left[\frac{1-2\sigma^2}{2\epsilon^2} + \frac{1}{\epsilon} \right] + \mathcal{O}(\epsilon^0). \quad (42)$$

The super-classical terms at $\mathcal{O}(1/|\mathbf{q}|)$ are given by

$$M_1^{(2,1)} = -i\pi \frac{3m_A^3 m_\phi^3 (m_A + m_\phi)}{(16\pi)^2 \sqrt{\sigma^2-1}} \left\{ (1-7\sigma^2+10\sigma^4) \left[\frac{1}{\epsilon} - 2\log(2) \right] - \frac{3+6\sigma^2-65\sigma^4+64\sigma^6}{3(\sigma^2-1)} \right\} + \mathcal{O}(\epsilon), \quad (43)$$

$$M_2^{(2,1)} = \frac{-i\pi m_A m_\phi^3}{2(32\pi)^2 (\sigma^2-1)^{3/2}} \left\{ (2\sigma^2-1) (m_\phi 20(1-3\sigma^2) + m_A(27-78\sigma^2-5\sigma^4)) \left[\frac{1}{\epsilon} - 2\log(2) \right] \right. \\ \left. + \frac{4}{3} m_\phi (-13-97\sigma^2+156\sigma^4) + m_A \frac{(-195+138\sigma^2+45\sigma^4+244\sigma^6)}{3} \right\} + \mathcal{O}(\epsilon), \quad (44)$$

$$M_3^{(2,1)} = \frac{i\pi m_A m_\phi}{(32\pi)^2 (\sigma^2-1)^{5/2}} \left\{ (m_A(2\sigma^2-1)(9-66\sigma^2+65\sigma^4) - m_\phi(2-28\sigma^2+78\sigma^4-60\sigma^6)) \left[\frac{1}{\epsilon} - 2\log(2) \right] \right. \\ \left. + \frac{2}{3} (m_\phi(7-134\sigma^2+327\sigma^4-192\sigma^6) + m_A(-66+249\sigma^2-264\sigma^4+89\sigma^6)) \right\} + \mathcal{O}(\epsilon), \quad (45)$$

$$M_4^{(2,1)} = \frac{-i\pi m_A^2 m_\phi^2 \sigma}{(16\pi)^2 (\sigma^2-1)^{3/2}} \left\{ (4m_A+3m_\phi)(3-11\sigma^2+10\sigma^4) \left[\frac{1}{\epsilon} - 2\log(2) \right] \right. \\ \left. + m_\phi(3+47\sigma^2-64\sigma^4) + 4m_A(3+6\sigma^2-13\sigma^4) \right\} + \mathcal{O}(\epsilon). \quad (46)$$

The classical terms are

$$M_1^{(2,2)} = \frac{1}{\epsilon^2} \frac{m_A^2 m_\phi^4 (1-2\sigma^2)}{(16\pi)^2 (\sigma^2-1)} + \frac{1}{\epsilon} \left[\frac{m_A^3 m_\phi^3 (3+12\sigma^2-4\sigma^4)}{(8\pi)^2 \sqrt{\sigma^2-1}} \operatorname{arccosh}(\sigma) + \frac{m_A^4 m_\phi^2 (-3+4\sigma^2)(1-8\sigma^2+8\sigma^4)}{(16\pi)^2 (\sigma^2-1)^2} \right. \\ \left. + \frac{m_A^2 m_\phi^4 (-9+48\sigma^2-78\sigma^4+40\sigma^6)}{(16\pi)^2 (\sigma^2-1)^2} + \frac{m_A^3 m_\phi^3 \sigma (-59+156\sigma^2-162\sigma^4+68\sigma^6)}{6(8\pi)^2 (\sigma^2-1)^2} \right] + \mathcal{O}(\epsilon^0), \quad (47)$$

$$M_2^{(2,2)} = \frac{1}{\epsilon^2} \left[\frac{m_\phi^4 (1-2\sigma^2)(1-4\sigma^2)}{2(16\pi)^2 (\sigma^2-1)^2} + \frac{m_A m_\phi^3 \sigma (1-2\sigma^2)(1-4\sigma^2)}{(16\pi)^2 (\sigma^2-1)^2} + \frac{m_A^2 m_\phi^2 (1-2\sigma^2)(1-8\sigma^4)}{(32\pi)^2 (\sigma^2-1)^2} \right] \\ + \frac{1}{\epsilon} \left[\frac{m_A m_\phi^3 (3+24\sigma^2-6\sigma^4+4\sigma^6)}{2(8\pi)^2 (\sigma^2-1)^{5/2}} \operatorname{arccosh}(\sigma) + \frac{m_\phi^4 (-9+62\sigma^2-72\sigma^4)}{(32\pi)^2 (\sigma^2-1)^2} \right. \\ \left. - \frac{m_A m_\phi^3 \sigma (1059-278\sigma^2+1064\sigma^4)}{30(16\pi)^2 (\sigma^2-1)^2} - \frac{m_A^2 m_\phi^2 (7-60\sigma^2+48\sigma^4+32\sigma^6)}{(32\pi)^2 (\sigma^2-1)^2} \right] + \mathcal{O}(\epsilon^0), \quad (48)$$

$$M_3^{(2,2)} = \frac{1}{\epsilon^2} \left[\frac{m_A m_\phi \sigma (1-2\sigma^2)(5-8\sigma^4)}{2(16\pi)^2 (\sigma^2-1)^3} - \frac{3m_A^2 (1-2\sigma^2)(1-8\sigma^2+8\sigma^4)}{(32\pi)^2 (\sigma^2-1)^3} + \frac{m_\phi^2 (1+2(\sigma^2-8\sigma^4+8\sigma^6))}{(32\pi)^2 (\sigma^2-1)^3} \right] \\ + \frac{1}{\epsilon} \left[\frac{m_A m_\phi (24+162\sigma^2+35\sigma^4-104\sigma^6+8\sigma^8)}{(16\pi)^2 (\sigma^2-1)^{7/2}} \operatorname{arccosh}(\sigma) + \frac{m_\phi^2 (19-98\sigma^2+200\sigma^4-128\sigma^6)}{2(32\pi)^2 (\sigma^2-1)^3} \right. \\ \left. - \frac{m_A m_\phi \sigma (7943+6986\sigma^2-8516\sigma^4+1192\sigma^6)}{15(32\pi)^2 (\sigma^2-1)^3} + \frac{m_A^2 (-21+2\sigma^2(7+64\sigma^2-64\sigma^4))}{2(32\pi)^2 (\sigma^2-1)^3} \right] + \mathcal{O}(\epsilon^0), \quad (49)$$

$$M_4^{(2,2)} = \frac{1}{\epsilon^2} \left[-\frac{(m_A^3 m_\phi + m_A m_\phi^3) \sigma (1-2\sigma^2)^2}{2(8\pi)^2 (\sigma^2-1)^2} - \frac{m_A^2 m_\phi^2 (1-2\sigma^2)(1-8\sigma^4)}{2(16\pi)^2 (\sigma^2-1)^2} \right] \\ + \frac{1}{\epsilon} \left[-\frac{m_A^2 m_\phi^2 \sigma (-6+\sigma^2)(1+2\sigma^2)}{2(4\pi)^2 (\sigma^2-1)^{3/2}} \operatorname{arccosh}(\sigma) + \frac{m_A m_\phi^3 \sigma (3-16\sigma^2+16\sigma^4)}{2(8\pi)^2 (\sigma^2-1)^2} \right. \\ \left. + \frac{m_A^3 m_\phi (\sigma-5\sigma^3+5\sigma^5)}{2(4\pi)^2 (\sigma^2-1)^2} + \frac{m_A^2 m_\phi^2 (77+189\sigma^2-360\sigma^4+136\sigma^6)}{3(16\pi)^2 (\sigma^2-1)^2} \right] + \mathcal{O}(\epsilon^0). \quad (50)$$

Polarization Tensors in Non-Relativistic EFT

In this section we give the expansion of the relativistic tensors $T_n = \epsilon_{1,\mu} T_n^{\mu\nu} \epsilon_{4,\nu}^*$ from Eq. (2) of the main text in terms of operators O_n defined by rest-frame polarization vectors (see Eq. (6) of the main text).

$$T_1 = -O_1 + \frac{O_2}{m_A^2 (\gamma_1 + 1)} \left(1 + \frac{\mathbf{q}^2}{4m_A^2 (\gamma_1 + 1)} \right) - \frac{O_3}{m_A^2 (\gamma_1 + 1)} \left(1 + \frac{\mathbf{q}^2}{8m_A^2 (\gamma_1 + 1)} \right) + \frac{O_4}{2m_A^4 (\gamma_1 + 1)^2}, \quad (51)$$

$$T_2 = -\frac{O_2 \mathbf{q}^2}{2m_A^2 (\gamma_1 + 1)} \left(1 + \frac{\mathbf{q}^2}{4m_A^2 (\gamma_1 + 1)} \right) + O_3 \left(1 + \frac{\mathbf{q}^2}{4m_A^2 (\gamma_1 + 1)} \right)^2 - \frac{O_4 \mathbf{q}^2}{4m_A^4 (\gamma_1 + 1)^2}, \quad (52)$$

$$T_3 = -\frac{O_2 \mathbf{q}^2}{2m_A^2} \left(E(E-m_A) - \frac{(2E-m_A)\mathbf{q}^2}{4m_A(\gamma_1+1)} + \frac{\mathbf{q}^2}{16m_A^2(\gamma_1+1)^2} \right) + \frac{O_3 \mathbf{q}^2}{4m_A^2} \left((E-m_A) - \frac{\mathbf{q}^2}{4m_A(\gamma_1+1)} \right)^2 \\ - \frac{O_4}{m_A^2} \left(E - \frac{\mathbf{q}^2}{4m_A(\gamma_1+1)} \right)^2, \quad (53)$$

$$T_4 = -\frac{O_2}{m_A} \left(E + \frac{(E-m_A)\mathbf{q}^2}{2m_A^2(\gamma_1+1)} - \frac{\mathbf{q}^2}{8m_A^3(\gamma_1+1)^2} \right) + O_3 \left(-1 + \frac{E}{m_A} + \frac{(E-2m_A)\mathbf{q}^2}{4m_A^3(\gamma_1+1)} - \frac{\mathbf{q}^2}{16m_A^4(\gamma_1+1)^2} \right) \\ - \frac{O_4}{m_A^3(\gamma_1+1)} \left(E - \frac{\mathbf{q}^2}{4m_A(\gamma_1+1)} \right), \quad (54)$$

where $E = E_A + E_\phi$, $\gamma_1 = E_A/m_A$ and, for brevity, $O_n \equiv \hat{\epsilon}_1^i O_n^{ij}(\vec{\mathbf{p}}, \mathbf{q}) \hat{\epsilon}_4^{*j}$ (where the $\hat{\epsilon}_i$ are the 3-dimensional rest frame polarization vectors). Note that these expressions are not truncated in $|\mathbf{q}|$.

Post-Newtonian expanded coefficients

The coefficients computed in the main text can directly be expanded in the non-relativistic limit $|\mathbf{p}| \ll 1$. Expressions including up to $\mathcal{O}(\mathbf{p}^{10})$ are given in the ancillary files to this manuscript⁴, here we only list the first two orders.

$$\tilde{c}_3^{(1)} = -\frac{1}{4}m_A m_\phi (m_A^2 + 8m_A m_\phi + m_\phi^2) - \frac{\mathbf{p}^2 \left(517m_A^2 m_\phi^2 + 230m_A^3 m_\phi + 25m_A^4 + 230m_A m_\phi^3 + 25m_\phi^4 \right)}{8m_A m_\phi} + \mathcal{O}(\mathbf{p}^4), \quad (55)$$

$$\tilde{c}_3^{(2)} = -\frac{1490m_A^3 m_\phi^2 + 1428m_A^2 m_\phi^3 + 636m_A^4 m_\phi + 84m_A^5 + 570m_A m_\phi^4 + 75m_\phi^5}{24m_A(m_A + m_\phi)^2} - \frac{\mathbf{p}^2 \left(40225m_A^4 m_\phi^2 + 34882m_A^3 m_\phi^3 + 12034m_A^2 m_\phi^4 + 19940m_A^5 m_\phi + 3384m_A^6 + 478m_A m_\phi^5 - 315m_\phi^6 \right)}{96m_A^3 m_\phi (m_A + m_\phi)^2} + \mathcal{O}(\mathbf{p}^4), \quad (56)$$

$$\tilde{c}_3^{(3)} = \frac{m_\phi \left(2025m_A^2 m_\phi^2 + 1000m_A^3 m_\phi + 132m_A^4 + 1381m_A m_\phi^3 + 264m_\phi^4 \right)}{240m_A(m_A + m_\phi)^2} + \frac{\mathbf{p}^2 \left(515832m_A^4 m_\phi^2 + 659844m_A^3 m_\phi^3 + 339726m_A^2 m_\phi^4 + 158568m_A^5 m_\phi + 14448m_A^6 + 45280m_A m_\phi^5 - 6027m_\phi^6 \right)}{6720m_A^3 m_\phi (m_A + m_\phi)^2} + \mathcal{O}(\mathbf{p}^4), \quad (57)$$

$$\tilde{c}_3^{(4)} = \frac{119982m_A^4 m_\phi^2 + 111716m_A^3 m_\phi^3 + 33749m_A^2 m_\phi^4 + 49536m_A^5 m_\phi + 5628m_A^6 - 4973m_A m_\phi^5 - 2891m_\phi^6}{6720m_A^3 m_\phi (m_A + m_\phi)^2} + \frac{\mathbf{p}^2 \left(965286m_A^6 m_\phi^2 + 2559750m_A^5 m_\phi^3 + 1043018m_A^4 m_\phi^4 - 1657386m_A^3 m_\phi^5 - 1397739m_A^2 m_\phi^6 \right)}{80640m_A^5 m_\phi^3 (m_A + m_\phi)^2} + \frac{\mathbf{p}^2 \left(-109202m_A^7 m_\phi - 45528m_A^8 - 164033m_A m_\phi^7 + 59997m_\phi^8 \right)}{80640m_A^5 m_\phi^3 (m_A + m_\phi)^2} + \mathcal{O}(\mathbf{p}^4), \quad (58)$$

$$\tilde{c}_3^{(5)} = \frac{m_\phi \left(1765m_A^2 m_\phi^2 + 1520m_A^3 m_\phi + 432m_A^4 + 901m_A m_\phi^3 + 214m_\phi^4 \right)}{960m_A(m_A + m_\phi)^2} + \frac{\mathbf{p}^2 \left(348602m_A^4 m_\phi^2 + 432624m_A^3 m_\phi^3 + 225346m_A^2 m_\phi^4 + 106768m_A^5 m_\phi + 4648m_A^6 + 33380m_A m_\phi^5 - 5537m_\phi^6 \right)}{26880m_A^3 m_\phi (m_A + m_\phi)^2} + \mathcal{O}(\mathbf{p}^4). \quad (59)$$

* ffebres@hep.fsu.edu

† mkraus@hep.fsu.edu

‡ linguandak@pku.edu.cn

§ mruf@physics.ucla.edu

¶ mao.zeng@ed.ac.uk

- [1] B. P. Abbott *et al.* (LIGO Scientific, Virgo), “Observation of Gravitational Waves from a Binary Black Hole Merger,” *Phys. Rev. Lett.* **116**, 061102 (2016), [arXiv:1602.03837 \[gr-qc\]](#).
- [2] R. Abbott *et al.* (LIGO Scientific, VIRGO, KAGRA), “GWTC-3: Compact Binary Coalescences Observed by LIGO and Virgo During the Second Part of the Third Observing Run,” (2021), [arXiv:2111.03606 \[gr-qc\]](#).
- [3] M Punturo *et al.*, “The einstein telescope: a third-generation gravitational wave observatory,” *Classical and Quantum Gravity* **27**, 194002 (2010).
- [4] David Reitze *et al.*, “Cosmic Explorer: The U.S. Contribution to Gravitational-Wave Astronomy beyond LIGO,” *Bull. Am. Astron. Soc.* **51**, 035 (2019), [arXiv:1907.04833 \[astro-ph.IM\]](#).

⁴ We include five computer files, including amplitudes computed in the full theory [FullTheory_Amplitudes.m], amplitudes computed from the EFT [EFT_Amplitudes.m], translations between

relativistic and non-relativistic tensors in Eqs. (3) and (7) [Tensor_expansion.m], and the matching coefficients defined in Eq. (8) [Coefficients.m] as well as non-relativistic expansions thereof [Coefficients_Expanded.m].

- [5] B. Bertotti, “On gravitational motion,” *Nuovo Cim.* **4**, 898–906 (1956).
- [6] R. P. Kerr, “The Lorentz-covariant approximation method in general relativity I,” *Nuovo Cim.* **13**, 469–491 (1959).
- [7] B. Bertotti and J. Plebanski, “Theory of gravitational perturbations in the fast motion approximation,” *Annals Phys.* **11**, 169–200 (1960).
- [8] M. Portilla, “Momentum and angular momentum of two gravitating particles,” *J. Phys. A* **12**, 1075–1090 (1979).
- [9] K. Westpfahl and M. Goller, “Gravitational scattering of two relativistic particles in post-linear approximation,” *Lett. Nuovo Cim.* **26**, 573–576 (1979).
- [10] M. Portilla, “Scattering of two gravitating particles: Classical approach,” *J. Phys. A* **13**, 3677–3683 (1980).
- [11] LLuis Bel, T. Damour, N. Deruelle, J. Ibanez, and J. Martin, “Poincaré-invariant gravitational field and equations of motion of two pointlike objects: The postlinear approximation of general relativity,” *Gen. Rel. Grav.* **13**, 963–1004 (1981).
- [12] Konradin Westpfahl, “High-Speed Scattering of Charged and Uncharged Particles in General Relativity,” *Fortsch. Phys.* **33**, 417–493 (1985).
- [13] Tomas Ledvinka, Gerhard Schafer, and Jiri Bicak, “Relativistic Closed-Form Hamiltonian for Many-Body Gravitating Systems in the Post-Minkowskian Approximation,” *Phys. Rev. Lett.* **100**, 251101 (2008), [arXiv:0807.0214 \[gr-qc\]](#).
- [14] Thibault Damour, “Gravitational scattering, post-Minkowskian approximation and Effective One-Body theory,” *Phys. Rev. D* **94**, 104015 (2016), [arXiv:1609.00354 \[gr-qc\]](#).
- [15] N. E. J. Bjerrum-Bohr, Poul H. Damgaard, Guido Festuccia, Ludovic Planté, and Pierre Vanhove, “General Relativity from Scattering Amplitudes,” *Phys. Rev. Lett.* **121**, 171601 (2018), [arXiv:1806.04920 \[hep-th\]](#).
- [16] Clifford Cheung, Ira Z. Rothstein, and Mikhail P. Solon, “From Scattering Amplitudes to Classical Potentials in the Post-Minkowskian Expansion,” *Phys. Rev. Lett.* **121**, 251101 (2018), [arXiv:1808.02489 \[hep-th\]](#).
- [17] David A. Kosower, Ben Maybee, and Donal O’Connell, “Amplitudes, Observables, and Classical Scattering,” *JHEP* **02**, 137 (2019), [arXiv:1811.10950 \[hep-th\]](#).
- [18] Zvi Bern, Clifford Cheung, Radu Roiban, Chia-Hsien Shen, Mikhail P. Solon, and Mao Zeng, “Scattering Amplitudes and the Conservative Hamiltonian for Binary Systems at Third Post-Minkowskian Order,” *Phys. Rev. Lett.* **122**, 201603 (2019), [arXiv:1901.04424 \[hep-th\]](#).
- [19] Zvi Bern, Clifford Cheung, Radu Roiban, Chia-Hsien Shen, Mikhail P. Solon, and Mao Zeng, “Black Hole Binary Dynamics from the Double Copy and Effective Theory,” *JHEP* **10**, 206 (2019), [arXiv:1908.01493 \[hep-th\]](#).
- [20] Paolo Di Vecchia, Carlo Heissenberg, Rodolfo Russo, and Gabriele Veneziano, “The eikonal approach to gravitational scattering and radiation at $\mathcal{O}(G^3)$,” *JHEP* **07**, 169 (2021), [arXiv:2104.03256 \[hep-th\]](#).
- [21] Enrico Herrmann, Julio Parra-Martinez, Michael S. Ruf, and Mao Zeng, “Radiative classical gravitational observables at $\mathcal{O}(G^3)$ from scattering amplitudes,” *JHEP* **10**, 148 (2021), [arXiv:2104.03957 \[hep-th\]](#).
- [22] N. Emil J. Bjerrum-Bohr, Poul H. Damgaard, Ludovic Planté, and Pierre Vanhove, “The amplitude for classical gravitational scattering at third Post-Minkowskian order,” *JHEP* **08**, 172 (2021), [arXiv:2105.05218 \[hep-th\]](#).
- [23] Andreas Brandhuber, Gang Chen, Gabriele Travaglini, and Congkao Wen, “Classical gravitational scattering from a gauge-invariant double copy,” *JHEP* **10**, 118 (2021), [arXiv:2108.04216 \[hep-th\]](#).
- [24] Zvi Bern, Julio Parra-Martinez, Radu Roiban, Michael S. Ruf, Chia-Hsien Shen, Mikhail P. Solon, and Mao Zeng, “Scattering Amplitudes and Conservative Binary Dynamics at $\mathcal{O}(G^4)$,” *Phys. Rev. Lett.* **126**, 171601 (2021), [arXiv:2101.07254 \[hep-th\]](#).
- [25] Zvi Bern, Julio Parra-Martinez, Radu Roiban, Michael S. Ruf, Chia-Hsien Shen, Mikhail P. Solon, and Mao Zeng, “Scattering Amplitudes, the Tail Effect, and Conservative Binary Dynamics at $\mathcal{O}(G^4)$,” *Phys. Rev. Lett.* **128**, 161103 (2022), [arXiv:2112.10750 \[hep-th\]](#).
- [26] Gregor Kälin, Zhengwen Liu, and Rafael A. Porto, “Conservative Dynamics of Binary Systems to Third Post-Minkowskian Order from the Effective Field Theory Approach,” *Phys. Rev. Lett.* **125**, 261103 (2020), [arXiv:2007.04977 \[hep-th\]](#).
- [27] Christoph Dlapa, Gregor Kälin, Zhengwen Liu, and Rafael A. Porto, “Dynamics of Binary Systems to Fourth Post-Minkowskian Order from the Effective Field Theory Approach,” (2021), [arXiv:2106.08276 \[hep-th\]](#).
- [28] Christoph Dlapa, Gregor Kälin, Zhengwen Liu, and Rafael A. Porto, “Conservative Dynamics of Binary Systems at Fourth Post-Minkowskian Order in the Large-Eccentricity Expansion,” *Phys. Rev. Lett.* **128**, 161104 (2022), [arXiv:2112.11296 \[hep-th\]](#).
- [29] Stefano Foffa, Pierpaolo Mastrolia, Riccardo Sturani, Christian Sturm, and William J. Torres Bobadilla, “Static two-body potential at fifth post-Newtonian order,” *Phys. Rev. Lett.* **122**, 241605 (2019), [arXiv:1902.10571 \[gr-qc\]](#).
- [30] J. Blümlein, A. Maier, and P. Marquard, “Five-Loop Static Contribution to the Gravitational Interaction Potential of Two Point Masses,” *Phys. Lett. B* **800**, 135100 (2020), [arXiv:1902.11180 \[gr-qc\]](#).
- [31] J. Blümlein, A. Maier, P. Marquard, and G. Schäfer, “The fifth-order post-Newtonian Hamiltonian dynamics of two-body systems from an effective field theory approach: potential contributions,” *Nucl. Phys. B* **965**, 115352 (2021), [arXiv:2010.13672 \[gr-qc\]](#).
- [32] J. Blümlein, A. Maier, P. Marquard, and G. Schäfer, “Fourth post-Newtonian Hamiltonian dynamics of two-body systems from an effective field theory approach,” *Nucl. Phys. B* **955**, 115041 (2020), [arXiv:2003.01692 \[gr-qc\]](#).
- [33] J. Blümlein, A. Maier, P. Marquard, and G. Schäfer, “The fifth-order post-Newtonian Hamiltonian dynamics of two-body systems from an effective field theory approach,” (2021), [arXiv:2110.13822 \[gr-qc\]](#).
- [34] J. Blümlein, A. Maier, P. Marquard, and G. Schäfer, “Testing binary dynamics in gravity at the sixth post-Newtonian level,” *Phys. Lett. B* **807**, 135496 (2020), [arXiv:2003.07145 \[gr-qc\]](#).
- [35] J. Blümlein, A. Maier, P. Marquard, and G. Schäfer, “The 6th post-Newtonian potential terms at $\mathcal{O}(G_N^4)$,” *Phys. Lett. B* **816**, 136260 (2021), [arXiv:2101.08630 \[gr-qc\]](#).

- [36] Donato Bini, Thibault Damour, and Andrea Geralico, “Binary dynamics at the fifth and fifth-and-a-half post-Newtonian orders,” *Phys. Rev. D* **102**, 024062 (2020), [arXiv:2003.11891 \[gr-qc\]](#).
- [37] Donato Bini, Thibault Damour, and Andrea Geralico, “Sixth post-Newtonian local-in-time dynamics of binary systems,” *Phys. Rev. D* **102**, 024061 (2020), [arXiv:2004.05407 \[gr-qc\]](#).
- [38] Donato Bini, Thibault Damour, Andrea Geralico, Stefano Laporta, and Pierpaolo Mastrolia, “Gravitational dynamics at $O(G^6)$: perturbative gravitational scattering meets experimental mathematics,” (2020), [arXiv:2008.09389 \[gr-qc\]](#).
- [39] Donato Bini, Thibault Damour, Andrea Geralico, Stefano Laporta, and Pierpaolo Mastrolia, “Gravitational scattering at the seventh order in G : nonlocal contribution at the sixth post-Newtonian accuracy,” *Phys. Rev. D* **103**, 044038 (2021), [arXiv:2012.12918 \[gr-qc\]](#).
- [40] Alessandra Buonanno, Bala Iyer, Evan Ochsner, Yi Pan, and B. S. Sathyaprakash, “Comparison of post-Newtonian templates for compact binary inspiral signals in gravitational-wave detectors,” *Phys. Rev. D* **80**, 084043 (2009), [arXiv:0907.0700 \[gr-qc\]](#).
- [41] Duncan A. Brown, Ian Harry, Andrew Lundgren, and Alexander H. Nitz, “Detecting binary neutron star systems with spin in advanced gravitational-wave detectors,” *Phys. Rev. D* **86**, 084017 (2012), [arXiv:1207.6406 \[gr-qc\]](#).
- [42] Emily Baird, Stephen Fairhurst, Mark Hannam, and Patricia Murphy, “Degeneracy between mass and spin in black-hole-binary waveforms,” *Phys. Rev. D* **87**, 024035 (2013), [arXiv:1211.0546 \[gr-qc\]](#).
- [43] N. V. Krishnendu and Frank Ohme, “Interplay of spin-precession and higher harmonics in the parameter estimation of binary black holes,” *Phys. Rev. D* **105**, 064012 (2022), [arXiv:2110.00766 \[gr-qc\]](#).
- [44] B. M. Barker and R. F. O’Connell, “Derivation of the equations of motion of a gyroscope from the quantum theory of gravitation,” *Phys. Rev. D* **2**, 1428–1435 (1970).
- [45] B. M. Barker and R. F. O’Connell, “Gravitational Two-Body Problem with Arbitrary Masses, Spins, and Quadrupole Moments,” *Phys. Rev. D* **12**, 329–335 (1975).
- [46] Lawrence E. Kidder, Clifford M. Will, and Alan G. Wiseman, “Spin effects in the inspiral of coalescing compact binaries,” *Phys. Rev. D* **47**, R4183–R4187 (1993), [arXiv:gr-qc/9211025](#).
- [47] Lawrence E. Kidder, “Coalescing binary systems of compact objects to postNewtonian 5/2 order. 5. Spin effects,” *Phys. Rev. D* **52**, 821–847 (1995), [arXiv:gr-qc/9506022](#).
- [48] Hideyuki Tagoshi, Akira Ohashi, and Benjamin J. Owen, “Gravitational field and equations of motion of spinning compact binaries to 2.5 postNewtonian order,” *Phys. Rev. D* **63**, 044006 (2001), [arXiv:gr-qc/0010014](#).
- [49] Guillaume Faye, Luc Blanchet, and Alessandra Buonanno, “Higher-order spin effects in the dynamics of compact binaries. I. Equations of motion,” *Phys. Rev. D* **74**, 104033 (2006), [arXiv:gr-qc/0605139](#).
- [50] Luc Blanchet, Alessandra Buonanno, and Guillaume Faye, “Higher-order spin effects in the dynamics of compact binaries. II. Radiation field,” *Phys. Rev. D* **74**, 104034 (2006), [Erratum: *Phys.Rev.D* 75, 049903 (2007), Erratum: *Phys.Rev.D* 81, 089901 (2010)], [arXiv:gr-qc/0605140](#).
- [51] Thibault Damour, Piotr Jaranowski, and Gerhard Schafer, “Hamiltonian of two spinning compact bodies with next-to-leading order gravitational spin-orbit coupling,” *Phys. Rev. D* **77**, 064032 (2008), [arXiv:0711.1048 \[gr-qc\]](#).
- [52] Jan Steinhoff, Steven Hergt, and Gerhard Schaefer, “On the next-to-leading order gravitational spin(1)-spin(2) dynamics,” *Phys. Rev. D* **77**, 081501 (2008), [arXiv:0712.1716 \[gr-qc\]](#).
- [53] Jan Steinhoff, Gerhard Schaefer, and Steven Hergt, “ADM canonical formalism for gravitating spinning objects,” *Phys. Rev. D* **77**, 104018 (2008), [arXiv:0805.3136 \[gr-qc\]](#).
- [54] Jan Steinhoff, Steven Hergt, and Gerhard Schaefer, “Spin-squared Hamiltonian of next-to-leading order gravitational interaction,” *Phys. Rev. D* **78**, 101503 (2008), [arXiv:0809.2200 \[gr-qc\]](#).
- [55] Sylvain Marsat, Alejandro Bohe, Guillaume Faye, and Luc Blanchet, “Next-to-next-to-leading order spin-orbit effects in the equations of motion of compact binary systems,” *Class. Quant. Grav.* **30**, 055007 (2013), [arXiv:1210.4143 \[gr-qc\]](#).
- [56] Steven Hergt, Jan Steinhoff, and Gerhard Schaefer, “Reduced Hamiltonian for next-to-leading order Spin-Squared Dynamics of General Compact Binaries,” *Class. Quant. Grav.* **27**, 135007 (2010), [arXiv:1002.2093 \[gr-qc\]](#).
- [57] Steven Hergt, Jan Steinhoff, and Gerhard Schaefer, “On the comparison of results regarding the post-Newtonian approximate treatment of the dynamics of extended spinning compact binaries,” *J. Phys. Conf. Ser.* **484**, 012018 (2014), [arXiv:1205.4530 \[gr-qc\]](#).
- [58] Alejandro Bohe, Sylvain Marsat, Guillaume Faye, and Luc Blanchet, “Next-to-next-to-leading order spin-orbit effects in the near-zone metric and precession equations of compact binaries,” *Class. Quant. Grav.* **30**, 075017 (2013), [arXiv:1212.5520 \[gr-qc\]](#).
- [59] Johannes Hartung, Jan Steinhoff, and Gerhard Schaefer, “Next-to-next-to-leading order post-Newtonian linear-in-spin binary Hamiltonians,” *Annalen Phys.* **525**, 359–394 (2013), [arXiv:1302.6723 \[gr-qc\]](#).
- [60] Sylvain Marsat, Luc Blanchet, Alejandro Bohe, and Guillaume Faye, “Gravitational waves from spinning compact object binaries: New post-Newtonian results,” (2013) [arXiv:1312.5375 \[gr-qc\]](#).
- [61] Alejandro Bohé, Guillaume Faye, Sylvain Marsat, and Edward K. Porter, “Quadratic-in-spin effects in the orbital dynamics and gravitational-wave energy flux of compact binaries at the 3PN order,” *Class. Quant. Grav.* **32**, 195010 (2015), [arXiv:1501.01529 \[gr-qc\]](#).
- [62] Donato Bini, Andrea Geralico, and Justin Vines, “Hyperbolic scattering of spinning particles by a Kerr black hole,” *Phys. Rev. D* **96**, 084044 (2017), [arXiv:1707.09814 \[gr-qc\]](#).
- [63] Nils Siemonsen, Jan Steinhoff, and Justin Vines, “Gravitational waves from spinning binary black holes at the leading post-Newtonian orders at all orders in spin,” *Phys. Rev. D* **97**, 124046 (2018), [arXiv:1712.08603 \[gr-qc\]](#).
- [64] Rafael A. Porto, “Post-Newtonian corrections to the motion of spinning bodies in NRGR,” *Phys. Rev. D* **73**, 104031 (2006), [arXiv:gr-qc/0511061](#).

- [65] Rafael A. Porto and Ira Z. Rothstein, “The Hyperfine Einstein-Infeld-Hoffmann potential,” *Phys. Rev. Lett.* **97**, 021101 (2006), [arXiv:gr-qc/0604099](#).
- [66] Rafael A. Porto and Ira Z. Rothstein, “Comment on ‘On the next-to-leading order gravitational spin(1) - spin(2) dynamics’ by J. Steinhoff et al,” (2007), [arXiv:0712.2032 \[gr-qc\]](#).
- [67] Rafael A. Porto and Ira Z. Rothstein, “Spin(1)Spin(2) Effects in the Motion of Inspiralling Compact Binaries at Third Order in the Post-Newtonian Expansion,” *Phys. Rev. D* **78**, 044012 (2008), [Erratum: *Phys.Rev.D* 81, 029904 (2010)], [arXiv:0802.0720 \[gr-qc\]](#).
- [68] Michele Levi, “Next to Leading Order gravitational Spin1-Spin2 coupling with Kaluza-Klein reduction,” *Phys. Rev. D* **82**, 064029 (2010), [arXiv:0802.1508 \[gr-qc\]](#).
- [69] Rafael A Porto and Ira Z. Rothstein, “Next to Leading Order Spin(1)Spin(1) Effects in the Motion of Inspiralling Compact Binaries,” *Phys. Rev. D* **78**, 044013 (2008), [Erratum: *Phys.Rev.D* 81, 029905 (2010)], [arXiv:0804.0260 \[gr-qc\]](#).
- [70] Rafael A. Porto, “Next to leading order spin-orbit effects in the motion of inspiralling compact binaries,” *Class. Quant. Grav.* **27**, 205001 (2010), [arXiv:1005.5730 \[gr-qc\]](#).
- [71] Michele Levi, “Next to Leading Order gravitational Spin-Orbit coupling in an Effective Field Theory approach,” *Phys. Rev. D* **82**, 104004 (2010), [arXiv:1006.4139 \[gr-qc\]](#).
- [72] Rafael A. Porto, Andreas Ross, and Ira Z. Rothstein, “Spin induced multipole moments for the gravitational wave flux from binary inspirals to third Post-Newtonian order,” *JCAP* **03**, 009 (2011), [arXiv:1007.1312 \[gr-qc\]](#).
- [73] Johannes Hartung and Jan Steinhoff, “Next-to-next-to-leading order post-Newtonian spin(1)-spin(2) Hamiltonian for self-gravitating binaries,” *Annalen Phys.* **523**, 919–924 (2011), [arXiv:1107.4294 \[gr-qc\]](#).
- [74] Michele Levi, “Binary dynamics from spin1-spin2 coupling at fourth post-Newtonian order,” *Phys. Rev. D* **85**, 064043 (2012), [arXiv:1107.4322 \[gr-qc\]](#).
- [75] Rafael A. Porto, Andreas Ross, and Ira Z. Rothstein, “Spin induced multipole moments for the gravitational wave amplitude from binary inspirals to 2.5 Post-Newtonian order,” *JCAP* **09**, 028 (2012), [arXiv:1203.2962 \[gr-qc\]](#).
- [76] Michele Levi and Jan Steinhoff, “Equivalence of ADM Hamiltonian and Effective Field Theory approaches at next-to-next-to-leading order spin1-spin2 coupling of binary inspirals,” *JCAP* **12**, 003 (2014), [arXiv:1408.5762 \[gr-qc\]](#).
- [77] Michele Levi and Jan Steinhoff, “Leading order finite size effects with spins for inspiralling compact binaries,” *JHEP* **06**, 059 (2015), [arXiv:1410.2601 \[gr-qc\]](#).
- [78] Michele Levi and Jan Steinhoff, “Spinning gravitating objects in the effective field theory in the post-Newtonian scheme,” *JHEP* **09**, 219 (2015), [arXiv:1501.04956 \[gr-qc\]](#).
- [79] Michele Levi and Jan Steinhoff, “Next-to-next-to-leading order gravitational spin-orbit coupling via the effective field theory for spinning objects in the post-Newtonian scheme,” *JCAP* **01**, 011 (2016), [arXiv:1506.05056 \[gr-qc\]](#).
- [80] Michele Levi and Jan Steinhoff, “Next-to-next-to-leading order gravitational spin-squared potential via the effective field theory for spinning objects in the post-Newtonian scheme,” *JCAP* **01**, 008 (2016), [arXiv:1506.05794 \[gr-qc\]](#).
- [81] Michele Levi and Jan Steinhoff, “Complete conservative dynamics for inspiralling compact binaries with spins at the fourth post-Newtonian order,” *JCAP* **09**, 029 (2021), [arXiv:1607.04252 \[gr-qc\]](#).
- [82] Natalia T. Maia, Chad R. Galley, Adam K. Leibovich, and Rafael A. Porto, “Radiation reaction for spinning bodies in effective field theory I: Spin-orbit effects,” *Phys. Rev. D* **96**, 084064 (2017), [arXiv:1705.07934 \[gr-qc\]](#).
- [83] Natalia T. Maia, Chad R. Galley, Adam K. Leibovich, and Rafael A. Porto, “Radiation reaction for spinning bodies in effective field theory II: Spin-spin effects,” *Phys. Rev. D* **96**, 084065 (2017), [arXiv:1705.07938 \[gr-qc\]](#).
- [84] Michèle Levi, Stavros Mougiakakos, and Mariana Vieira, “Gravitational cubic-in-spin interaction at the next-to-leading post-Newtonian order,” *JHEP* **01**, 036 (2021), [arXiv:1912.06276 \[hep-th\]](#).
- [85] Michèle Levi, Andrew J. McLeod, and Matthew Von Hippel, “N³LO gravitational spin-orbit coupling at order G⁴,” *JHEP* **07**, 115 (2021), [arXiv:2003.02827 \[hep-th\]](#).
- [86] Michèle Levi, Andrew J. McLeod, and Matthew Von Hippel, “N³LO gravitational quadratic-in-spin interactions at G⁴,” *JHEP* **07**, 116 (2021), [arXiv:2003.07890 \[hep-th\]](#).
- [87] Andrea Antonelli, Chris Kavanagh, Mohammed Khalil, Jan Steinhoff, and Justin Vines, “Gravitational spin-orbit coupling through third-subleading post-Newtonian order: from first-order self-force to arbitrary mass ratios,” *Phys. Rev. Lett.* **125**, 011103 (2020), [arXiv:2003.11391 \[gr-qc\]](#).
- [88] Michèle Levi and Fei Teng, “NLO gravitational quartic-in-spin interaction,” *JHEP* **01**, 066 (2021), [arXiv:2008.12280 \[hep-th\]](#).
- [89] Andrea Antonelli, Chris Kavanagh, Mohammed Khalil, Jan Steinhoff, and Justin Vines, “Gravitational spin-orbit and aligned spin₁-spin₂ couplings through third-subleading post-Newtonian orders,” *Phys. Rev. D* **102**, 124024 (2020), [arXiv:2010.02018 \[gr-qc\]](#).
- [90] Walter D. Goldberger, Jingping Li, and Ira Z. Rothstein, “Non-conservative effects on spinning black holes from world-line effective field theory,” *JHEP* **06**, 053 (2021), [arXiv:2012.14869 \[hep-th\]](#).
- [91] Zhengwen Liu, Rafael A. Porto, and Zixin Yang, “Spin Effects in the Effective Field Theory Approach to Post-Minkowskian Conservative Dynamics,” *JHEP* **06**, 012 (2021), [arXiv:2102.10059 \[hep-th\]](#).
- [92] Gihyuk Cho, Brian Pardo, and Rafael A. Porto, “Gravitational radiation from inspiralling compact objects: Spin-spin effects completed at the next-to-leading post-Newtonian order,” *Phys. Rev. D* **104**, 024037 (2021), [arXiv:2103.14612 \[gr-qc\]](#).
- [93] Jung-Wook Kim, Michèle Levi, and Zhewei Yin, “Quadratic-in-spin interactions at fifth post-Newtonian order probe new physics,” (2021), [arXiv:2112.01509 \[hep-th\]](#).
- [94] Gihyuk Cho, Rafael A. Porto, and Zixin Yang, “Gravitational radiation from inspiralling compact objects: Spin effects to fourth Post-Newtonian order,” (2022), [arXiv:2201.05138 \[gr-qc\]](#).

- [95] Donato Bini and Thibault Damour, “Gravitational spin-orbit coupling in binary systems, post-Minkowskian approximation and effective one-body theory,” *Phys. Rev. D* **96**, 104038 (2017), [arXiv:1709.00590 \[gr-qc\]](#).
- [96] Donato Bini and Thibault Damour, “Gravitational spin-orbit coupling in binary systems at the second post-Minkowskian approximation,” *Phys. Rev. D* **98**, 044036 (2018), [arXiv:1805.10809 \[gr-qc\]](#).
- [97] Justin Vines, “Scattering of two spinning black holes in post-Minkowskian gravity, to all orders in spin, and effective-one-body mappings,” *Class. Quant. Grav.* **35**, 084002 (2018), [arXiv:1709.06016 \[gr-qc\]](#).
- [98] Justin Vines, Jan Steinhoff, and Alessandra Buonanno, “Spinning-black-hole scattering and the test-black-hole limit at second post-Minkowskian order,” *Phys. Rev. D* **99**, 064054 (2019), [arXiv:1812.00956 \[gr-qc\]](#).
- [99] Alfredo Guevara, “Holomorphic Classical Limit for Spin Effects in Gravitational and Electromagnetic Scattering,” *JHEP* **04**, 033 (2019), [arXiv:1706.02314 \[hep-th\]](#).
- [100] Alfredo Guevara, Alexander Ochirov, and Justin Vines, “Scattering of Spinning Black Holes from Exponentiated Soft Factors,” *JHEP* **09**, 056 (2019), [arXiv:1812.06895 \[hep-th\]](#).
- [101] Ming-Zhi Chung, Yu-Tin Huang, Jung-Wook Kim, and Sangmin Lee, “The simplest massive S-matrix: from minimal coupling to Black Holes,” *JHEP* **04**, 156 (2019), [arXiv:1812.08752 \[hep-th\]](#).
- [102] Nima Arkani-Hamed, Yu-tin Huang, and Donal O’Connell, “Kerr black holes as elementary particles,” *JHEP* **01**, 046 (2020), [arXiv:1906.10100 \[hep-th\]](#).
- [103] Alfredo Guevara, Alexander Ochirov, and Justin Vines, “Black-hole scattering with general spin directions from minimal-coupling amplitudes,” *Phys. Rev. D* **100**, 104024 (2019), [arXiv:1906.10071 \[hep-th\]](#).
- [104] Ming-Zhi Chung, Yu-Tin Huang, and Jung-Wook Kim, “Classical potential for general spinning bodies,” *JHEP* **09**, 074 (2020), [arXiv:1908.08463 \[hep-th\]](#).
- [105] Poul H. Damgaard, Kays Haddad, and Andreas Helset, “Heavy Black Hole Effective Theory,” *JHEP* **11**, 070 (2019), [arXiv:1908.10308 \[hep-ph\]](#).
- [106] Rafael Aoude, Kays Haddad, and Andreas Helset, “On-shell heavy particle effective theories,” *JHEP* **05**, 051 (2020), [arXiv:2001.09164 \[hep-th\]](#).
- [107] Ming-Zhi Chung, Yu-tin Huang, Jung-Wook Kim, and Sangmin Lee, “Complete Hamiltonian for spinning binary systems at first post-Minkowskian order,” *JHEP* **05**, 105 (2020), [arXiv:2003.06600 \[hep-th\]](#).
- [108] Alfredo Guevara, Ben Maybee, Alexander Ochirov, Donal O’connell, and Justin Vines, “A worldsheet for Kerr,” *JHEP* **03**, 201 (2021), [arXiv:2012.11570 \[hep-th\]](#).
- [109] Zvi Bern, Andres Luna, Radu Roiban, Chia-Hsien Shen, and Mao Zeng, “Spinning black hole binary dynamics, scattering amplitudes, and effective field theory,” *Phys. Rev. D* **104**, 065014 (2021), [arXiv:2005.03071 \[hep-th\]](#).
- [110] Dimitrios Kosmopoulos and Andres Luna, “Quadratic-in-spin Hamiltonian at $\mathcal{O}(G^2)$ from scattering amplitudes,” *JHEP* **07**, 037 (2021), [arXiv:2102.10137 \[hep-th\]](#).
- [111] Wei-Ming Chen, Ming-Zhi Chung, Yu-tin Huang, and Jung-Wook Kim, “The 2PM Hamiltonian for binary Kerr to quartic in spin,” (2021), [arXiv:2111.13639 \[hep-th\]](#).
- [112] Zvi Bern, Dimitrios Kosmopoulos, Andrés Luna, Radu Roiban, and Fei Teng, “Binary Dynamics Through the Fifth Power of Spin at $\mathcal{O}(G^2)$,” (2022), [arXiv:2203.06202 \[hep-th\]](#).
- [113] Rafael Aoude, Kays Haddad, and Andreas Helset, “Searching for Kerr in the 2PM amplitude,” (2022), [arXiv:2203.06197 \[hep-th\]](#).
- [114] Rafael Aoude, Kays Haddad, and Andreas Helset, “Classical gravitational spinning-spinless scattering at $\mathcal{O}(G^2 S^\infty)$,” (2022), [arXiv:2205.02809 \[hep-th\]](#).
- [115] Gustav Mogull, Jan Plefka, and Jan Steinhoff, “Classical black hole scattering from a worldline quantum field theory,” *JHEP* **02**, 048 (2021), [arXiv:2010.02865 \[hep-th\]](#).
- [116] Gustav Uhre Jakobsen, Gustav Mogull, Jan Plefka, and Jan Steinhoff, “Classical Gravitational Bremsstrahlung from a Worldline Quantum Field Theory,” *Phys. Rev. Lett.* **126**, 201103 (2021), [arXiv:2101.12688 \[gr-qc\]](#).
- [117] Gustav Uhre Jakobsen, Gustav Mogull, Jan Plefka, and Jan Steinhoff, “Gravitational Bremsstrahlung and Hidden Supersymmetry of Spinning Bodies,” *Phys. Rev. Lett.* **128**, 011101 (2022), [arXiv:2106.10256 \[hep-th\]](#).
- [118] Gustav Uhre Jakobsen, Gustav Mogull, Jan Plefka, and Jan Steinhoff, “SUSY in the sky with gravitons,” *JHEP* **01**, 027 (2022), [arXiv:2109.04465 \[hep-th\]](#).
- [119] Gustav Uhre Jakobsen and Gustav Mogull, “Conservative and Radiative Dynamics of Spinning Bodies at Third Post-Minkowskian Order Using Worldline Quantum Field Theory,” *Phys. Rev. Lett.* **128**, 141102 (2022), [arXiv:2201.07778 \[hep-th\]](#).
- [120] Francesco Alessio and Paolo Di Vecchia, “Radiation reaction for spinning black-hole scattering,” (2022), [arXiv:2203.13272 \[hep-th\]](#).
- [121] Andrea Cristofoli, Riccardo Gonzo, Nathan Moynihan, Donal O’Connell, Alasdair Ross, Matteo Sergola, and Chris D. White, “The Uncertainty Principle and Classical Amplitudes,” (2021), [arXiv:2112.07556 \[hep-th\]](#).
- [122] Thibault Damour and Gilles Esposito-Farese, “Testing gravity to second postNewtonian order: A Field theory approach,” *Phys. Rev. D* **53**, 5541–5578 (1996), [arXiv:gr-qc/9506063](#).
- [123] Varun Vaidya, “Gravitational spin Hamiltonians from the S matrix,” *Phys. Rev. D* **91**, 024017 (2015), [arXiv:1410.5348 \[hep-th\]](#).
- [124] Zvi Bern, Lance J. Dixon, David C. Dunbar, and David A. Kosower, “One loop n point gauge theory amplitudes, unitarity and collinear limits,” *Nucl. Phys. B* **425**, 217–260 (1994), [arXiv:hep-ph/9403226](#).
- [125] Zvi Bern, Lance J. Dixon, David C. Dunbar, and David A. Kosower, “Fusing gauge theory tree amplitudes into loop amplitudes,” *Nucl. Phys. B* **435**, 59–101 (1995), [arXiv:hep-ph/9409265](#).

- [126] Z. Bern and A. G. Morgan, “Massive loop amplitudes from unitarity,” *Nucl. Phys. B* **467**, 479–509 (1996), [arXiv:hep-ph/9511336](#).
- [127] Zvi Bern, Lance J. Dixon, and David A. Kosower, “One loop amplitudes for $e^+ e^-$ to four partons,” *Nucl. Phys. B* **513**, 3–86 (1998), [arXiv:hep-ph/9708239](#).
- [128] Ruth Britto, Freddy Cachazo, and Bo Feng, “Generalized unitarity and one-loop amplitudes in $N=4$ super-Yang-Mills,” *Nucl. Phys. B* **725**, 275–305 (2005), [arXiv:hep-th/0412103](#).
- [129] Harald Ita, “Two-loop Integrand Decomposition into Master Integrals and Surface Terms,” *Phys. Rev. D* **94**, 116015 (2016), [arXiv:1510.05626 \[hep-th\]](#).
- [130] S. Abreu, F. Febres Cordero, H. Ita, M. Jaquier, and B. Page, “Subleading Poles in the Numerical Unitarity Method at Two Loops,” *Phys. Rev. D* **95**, 096011 (2017), [arXiv:1703.05255 \[hep-ph\]](#).
- [131] S. Abreu, F. Febres Cordero, H. Ita, M. Jaquier, B. Page, and M. Zeng, “Two-Loop Four-Gluon Amplitudes from Numerical Unitarity,” *Phys. Rev. Lett.* **119**, 142001 (2017), [arXiv:1703.05273 \[hep-ph\]](#).
- [132] Samuel Abreu, Fernando Febres Cordero, Harald Ita, Ben Page, and Mao Zeng, “Planar Two-Loop Five-Gluon Amplitudes from Numerical Unitarity,” *Phys. Rev. D* **97**, 116014 (2018), [arXiv:1712.03946 \[hep-ph\]](#).
- [133] S. Abreu, F. Febres Cordero, H. Ita, B. Page, and V. Sotnikov, “Planar Two-Loop Five-Parton Amplitudes from Numerical Unitarity,” *JHEP* **11**, 116 (2018), [arXiv:1809.09067 \[hep-ph\]](#).
- [134] Duff Neill and Ira Z. Rothstein, “Classical Space-Times from the S Matrix,” *Nucl. Phys. B* **877**, 177–189 (2013), [arXiv:1304.7263 \[hep-th\]](#).
- [135] Ben Maybee, Donal O’Connell, and Justin Vines, “Observables and amplitudes for spinning particles and black holes,” *JHEP* **12**, 156 (2019), [arXiv:1906.09260 \[hep-th\]](#).
- [136] P. V. Landshoff and J. C. Polkinghorne, “Iterations of regge cuts,” *Phys. Rev.* **181**, 1989–1995 (1969).
- [137] Julio Parra-Martinez, Michael S. Ruf, and Mao Zeng, “Extremal black hole scattering at $\mathcal{O}(G^3)$: graviton dominance, eikonal exponentiation, and differential equations,” *JHEP* **11**, 023 (2020), [arXiv:2005.04236 \[hep-th\]](#).
- [138] S. Abreu, J. Dormans, F. Febres Cordero, H. Ita, M. Kraus, B. Page, E. Pascual, M. S. Ruf, and V. Sotnikov, “Caravel: A C++ framework for the computation of multi-loop amplitudes with numerical unitarity,” *Comput. Phys. Commun.* **267**, 108069 (2021), [arXiv:2009.11957 \[hep-ph\]](#).
- [139] Andreas von Manteuffel and Robert M. Schabinger, “A novel approach to integration by parts reduction,” *Phys. Lett. B* **744**, 101–104 (2015), [arXiv:1406.4513 \[hep-ph\]](#).
- [140] Tiziano Peraro, “Scattering amplitudes over finite fields and multivariate functional reconstruction,” *JHEP* **12**, 030 (2016), [arXiv:1608.01902 \[hep-ph\]](#).
- [141] S. Abreu, F. Febres Cordero, H. Ita, M. Jaquier, B. Page, M. S. Ruf, and V. Sotnikov, “Two-Loop Four-Graviton Scattering Amplitudes,” *Phys. Rev. Lett.* **124**, 211601 (2020), [arXiv:2002.12374 \[hep-th\]](#).
- [142] David Brizuela, Jose M. Martin-Garcia, and Guillermo A. Mena Marugan, “xPert: Computer algebra for metric perturbation theory,” *Gen. Rel. Grav.* **41**, 2415–2431 (2009), [arXiv:0807.0824 \[gr-qc\]](#).
- [143] Teake Nutma, “xTras : A field-theory inspired xAct package for mathematica,” *Comput. Phys. Commun.* **185**, 1719–1738 (2014), [arXiv:1308.3493 \[cs.SC\]](#).
- [144] Gerard ’t Hooft and M. J. G. Veltman, “Regularization and Renormalization of Gauge Fields,” *Nucl. Phys. B* **44**, 189–213 (1972).
- [145] Frits A. Berends and W. T. Giele, “Recursive Calculations for Processes with n Gluons,” *Nucl. Phys. B* **306**, 759–808 (1988).
- [146] H. Kawai, D. C. Lewellen, and S. H. H. Tye, “A Relation Between Tree Amplitudes of Closed and Open Strings,” *Nucl. Phys. B* **269**, 1–23 (1986).
- [147] Z. Bern, J. J. M. Carrasco, and Henrik Johansson, “New Relations for Gauge-Theory Amplitudes,” *Phys. Rev. D* **78**, 085011 (2008), [arXiv:0805.3993 \[hep-ph\]](#).
- [148] Zvi Bern, John Joseph M. Carrasco, and Henrik Johansson, “Perturbative Quantum Gravity as a Double Copy of Gauge Theory,” *Phys. Rev. Lett.* **105**, 061602 (2010), [arXiv:1004.0476 \[hep-th\]](#).
- [149] Henrik Johansson and Alexander Ochirov, “Double copy for massive quantum particles with spin,” *JHEP* **09**, 040 (2019), [arXiv:1906.12292 \[hep-th\]](#).
- [150] Yilber Fabian Bautista and Alfredo Guevara, “On the double copy for spinning matter,” *JHEP* **11**, 184 (2021), [arXiv:1908.11349 \[hep-th\]](#).
- [151] Clifford Cheung and Grant N. Remmen, “Twofold Symmetries of the Pure Gravity Action,” *JHEP* **01**, 104 (2017), [arXiv:1612.03927 \[hep-th\]](#).
- [152] Clifford Cheung and Grant N. Remmen, “Hidden Simplicity of the Gravity Action,” *JHEP* **09**, 002 (2017), [arXiv:1705.00626 \[hep-th\]](#).
- [153] Gregor Kälin and Rafael A. Porto, “Post-Minkowskian Effective Field Theory for Conservative Binary Dynamics,” *JHEP* **11**, 106 (2020), [arXiv:2006.01184 \[hep-th\]](#).
- [154] Ruth Britto, Riccardo Gonzo, and Guy R. Jehu, “Graviton particle statistics and coherent states from classical scattering amplitudes,” *JHEP* **03**, 214 (2022), [arXiv:2112.07036 \[hep-th\]](#).
- [155] K. G. Chetyrkin and F. V. Tkachov, “Integration by Parts: The Algorithm to Calculate beta Functions in 4 Loops,” *Nucl. Phys. B* **192**, 159–204 (1981).
- [156] A. V. Smirnov and F. S. Chuharev, “FIRE6: Feynman Integral REduction with Modular Arithmetic,” *Comput. Phys. Commun.* **247**, 106877 (2020), [arXiv:1901.07808 \[hep-ph\]](#).
- [157] Supplemental material, includes three appendices on full-theory amplitudes, polarization tensors in non-relativistic EFT, and on the PN expansion of our coefficients. We also include five computer files, including amplitudes computed in the

- full theory [FullTheory_Amplitudes.m], amplitudes computed from the EFT [EFT_Amplitudes.m], translations between relativistic and non-relativistic tensors in Eqs. (2) and (6) [Tensor_expansion.m], and the matching coefficients defined in Eq. (7) [Coefficients.m] as well as non-relativistic expansions thereof [Coefficients_Expanded.m].
- [158] Barry R. Holstein and Andreas Ross, “Spin Effects in Long Range Gravitational Scattering,” (2008), [arXiv:0802.0716 \[hep-ph\]](#).
 - [159] Andrea Cristofoli, Poul H. Damgaard, Paolo Di Vecchia, and Carlo Heissenberg, “Second-order Post-Minkowskian scattering in arbitrary dimensions,” *JHEP* **07**, 122 (2020), [arXiv:2003.10274 \[hep-th\]](#).
 - [160] Michele Levi and Jan Steinhoff, “EFTofPNG: A package for high precision computation with the Effective Field Theory of Post-Newtonian Gravity,” *Class. Quant. Grav.* **34**, 244001 (2017), [arXiv:1705.06309 \[gr-qc\]](#).
 - [161] Enrico Herrmann, Julio Parra-Martinez, Michael S. Ruf, and Mao Zeng, “Gravitational Bremsstrahlung from Reverse Unitarity,” *Phys. Rev. Lett.* **126**, 201602 (2021), [arXiv:2101.07255 \[hep-th\]](#).
 - [162] Gregor Kälin and Rafael A. Porto, “From Boundary Data to Bound States,” *JHEP* **01**, 072 (2020), [arXiv:1910.03008 \[hep-th\]](#).
 - [163] Gregor Kälin and Rafael A. Porto, “From boundary data to bound states. Part II. Scattering angle to dynamical invariants (with twist),” *JHEP* **02**, 120 (2020), [arXiv:1911.09130 \[hep-th\]](#).
 - [164] Gihyuk Cho, Gregor Kälin, and Rafael A. Porto, “From boundary data to bound states. Part III. Radiative effects,” *JHEP* **04**, 154 (2022), [arXiv:2112.03976 \[hep-th\]](#).
 - [165] Marco Chiodaroli, Henrik Johansson, and Paolo Pichini, “Compton black-hole scattering for $s \leq 5/2$,” *JHEP* **02**, 156 (2022), [arXiv:2107.14779 \[hep-th\]](#).
 - [166] Clifford Cheung and Mikhail P. Solon, “Tidal Effects in the Post-Minkowskian Expansion,” *Phys. Rev. Lett.* **125**, 191601 (2020), [arXiv:2006.06665 \[hep-th\]](#).
 - [167] Gregor Kälin, Zhengwen Liu, and Rafael A. Porto, “Conservative Tidal Effects in Compact Binary Systems to Next-to-Leading Post-Minkowskian Order,” *Phys. Rev. D* **102**, 124025 (2020), [arXiv:2008.06047 \[hep-th\]](#).
 - [168] Zvi Bern, Julio Parra-Martinez, Radu Roiban, Eric Sawyer, and Chia-Hsien Shen, “Leading Nonlinear Tidal Effects and Scattering Amplitudes,” *JHEP* **05**, 188 (2021), [arXiv:2010.08559 \[hep-th\]](#).
 - [169] Clifford Cheung, Nabha Shah, and Mikhail P. Solon, “Mining the Geodesic Equation for Scattering Data,” *Phys. Rev. D* **103**, 024030 (2021), [arXiv:2010.08568 \[hep-th\]](#).
 - [170] Donato Bini, Thibault Damour, and Andrea Geralico, “Scattering of tidally interacting bodies in post-Minkowskian gravity,” *Phys. Rev. D* **101**, 044039 (2020), [arXiv:2001.00352 \[gr-qc\]](#).
 - [171] Kays Haddad and Andreas Helset, “Tidal effects in quantum field theory,” *JHEP* **12**, 024 (2020), [arXiv:2008.04920 \[hep-th\]](#).
 - [172] Rafael Aoude, Kays Haddad, and Andreas Helset, “Tidal effects for spinning particles,” *JHEP* **03**, 097 (2021), [arXiv:2012.05256 \[hep-th\]](#).
 - [173] Stavros Mougiakakos, Massimiliano Maria Riva, and Filippo Vernizzi, “Gravitational Bremsstrahlung with tidal effects in the post-Minkowskian expansion,” (2022), [arXiv:2204.06556 \[hep-th\]](#).
 - [174] Andrés Luna, Ricardo Monteiro, Isobel Nicholson, Donal O’Connell, and Chris D. White, “The double copy: Bremsstrahlung and accelerating black holes,” *JHEP* **06**, 023 (2016), [arXiv:1603.05737 \[hep-th\]](#).
 - [175] Walter D. Goldberger and Alexander K. Ridgway, “Radiation and the classical double copy for color charges,” *Phys. Rev. D* **95**, 125010 (2017), [arXiv:1611.03493 \[hep-th\]](#).
 - [176] Walter D. Goldberger, Siddharth G. Prabhu, and Jedidiah O. Thompson, “Classical gluon and graviton radiation from the bi-adjoint scalar double copy,” *Phys. Rev. D* **96**, 065009 (2017), [arXiv:1705.09263 \[hep-th\]](#).
 - [177] Walter D. Goldberger and Alexander K. Ridgway, “Bound states and the classical double copy,” *Phys. Rev. D* **97**, 085019 (2018), [arXiv:1711.09493 \[hep-th\]](#).
 - [178] David Chester, “Radiative double copy for Einstein-Yang-Mills theory,” *Phys. Rev. D* **97**, 084025 (2018), [arXiv:1712.08684 \[hep-th\]](#).
 - [179] Walter D. Goldberger, Jingping Li, and Siddharth G. Prabhu, “Spinning particles, axion radiation, and the classical double copy,” *Phys. Rev. D* **97**, 105018 (2018), [arXiv:1712.09250 \[hep-th\]](#).
 - [180] Chia-Hsien Shen, “Gravitational Radiation from Color-Kinematics Duality,” *JHEP* **11**, 162 (2018), [arXiv:1806.07388 \[hep-th\]](#).
 - [181] Walter D. Goldberger and Jingping Li, “Strings, extended objects, and the classical double copy,” *JHEP* **02**, 092 (2020), [arXiv:1912.01650 \[hep-th\]](#).
 - [182] Gabriel Luz Almeida, Stefano Foffa, and Riccardo Sturani, “Classical Gravitational Self-Energy from Double Copy,” *JHEP* **11**, 165 (2020), [arXiv:2008.06195 \[gr-qc\]](#).
 - [183] Alok Laddha and Ashoke Sen, “Gravity Waves from Soft Theorem in General Dimensions,” *JHEP* **09**, 105 (2018), [arXiv:1801.07719 \[hep-th\]](#).
 - [184] Alok Laddha and Ashoke Sen, “Logarithmic Terms in the Soft Expansion in Four Dimensions,” *JHEP* **10**, 056 (2018), [arXiv:1804.09193 \[hep-th\]](#).
 - [185] Alok Laddha and Ashoke Sen, “Observational Signature of the Logarithmic Terms in the Soft Graviton Theorem,” *Phys. Rev. D* **100**, 024009 (2019), [arXiv:1806.01872 \[hep-th\]](#).
 - [186] Biswajit Sahoo and Ashoke Sen, “Classical and Quantum Results on Logarithmic Terms in the Soft Theorem in Four Dimensions,” *JHEP* **02**, 086 (2019), [arXiv:1808.03288 \[hep-th\]](#).
 - [187] Marcello Ciafaloni, Dimitri Colferai, and Gabriele Veneziano, “Infrared features of gravitational scattering and radiation in the eikonal approach,” *Phys. Rev. D* **99**, 066008 (2019), [arXiv:1812.08137 \[hep-th\]](#).

- [188] Alok Laddha and Ashoke Sen, “Classical proof of the classical soft graviton theorem in $D > 4$,” *Phys. Rev. D* **101**, 084011 (2020), [arXiv:1906.08288 \[gr-qc\]](#).
- [189] Athira P. V. and A. Manu, “Classical double copy from Color Kinematics duality: A proof in the soft limit,” *Phys. Rev. D* **101**, 046014 (2020), [arXiv:1907.10021 \[hep-th\]](#).
- [190] Arnab Priya Saha, Biswajit Sahoo, and Ashoke Sen, “Proof of the classical soft graviton theorem in $D = 4$,” *JHEP* **06**, 153 (2020), [arXiv:1912.06413 \[hep-th\]](#).
- [191] A. Manu, Deboirna Ghosh, Alok Laddha, and P. V. Athira, “Soft radiation from scattering amplitudes revisited,” *JHEP* **05**, 056 (2021), [arXiv:2007.02077 \[hep-th\]](#).
- [192] Biswajit Sahoo, “Classical Sub-subleading Soft Photon and Soft Graviton Theorems in Four Spacetime Dimensions,” *JHEP* **12**, 070 (2020), [arXiv:2008.04376 \[hep-th\]](#).