

ON SOME GENERALIZED INVERSES AND PARTIAL ORDERS IN *-RINGS

JANKO MAROVT, DIJANA MOSIĆ, AND INSA CREMER

ABSTRACT. Let \mathcal{R} be a unital ring with involution. The notions of IMP-inverse and MP1-inverse are extended from $M_{m,n}(\mathbb{C})$, the set of all $m \times n$ matrices over \mathbb{C} , to the set \mathcal{R}^\dagger of all Moore-Penrose invertible elements in \mathcal{R} . We study partial orders on \mathcal{R}^\dagger that are induced by IMP-inverses and MP1-inverses. We also extend to the setting of Rickart *-rings the concept of another partial order, called the plus order, which has been recently introduced on the set of all bounded linear operators between Hilbert spaces. Properties of these relations are investigated and some known results are thus generalized.

1. INTRODUCTION

Let \mathcal{R} be a *-ring, i.e. a ring equipped with involution $*$. We call an element $a \in \mathcal{R}$ a **-regular* or *Moore-Penrose invertible* with respect to $*$ if there exists $x \in \mathcal{R}$ that satisfies the following four equations:

$$(1) \quad axa = a, \quad xax = x, \quad (ax)^* = ax, \quad (xa)^* = xa.$$

If such x exists, we write $x = a^\dagger$ and call it the *Moore-Penrose inverse* of a . It is known that a^\dagger is unique if it exists. We denote the set of all *-regular elements in \mathcal{R} by \mathcal{R}^\dagger . A ring \mathcal{R} where every element is *-regular is called a **-regular ring*. An example of a *-regular ring is the set $M_n(\mathbb{C})$ of all complex $n \times n$ matrices where A^* denotes the conjugate transpose of $A \in M_n(\mathbb{C})$. There are many other generalized inverses that can be introduced on \mathcal{R} and some of them may be defined with a subset of the set of equations (1). We say that an element $a \in \mathcal{R}$ is *regular* if there exists $x \in \mathcal{R}$ that satisfies the first equation in (1). Such x , if it exists, is called an *inner generalized inverse* or *{1}-inverse* of a , and we write $x = a^-$, i.e. $aa^-a = a$. The set of all regular elements in \mathcal{R} is denoted by $\mathcal{R}^{(1)}$. A ring \mathcal{R} is called *von Neumann regular* if every element in \mathcal{R} is regular. If $x \in \mathcal{R}$ satisfies the first three equations in (1), then it is called a *{1,2,3}-inverse* of a . We denote the sets of all {1}-inverses and {1,2,3}-inverses of $a \in \mathcal{R}$ by $a\{1\}$ and $a\{1,2,3\}$, respectively.

In a recent paper [7], Hernández et al. introduced two new types of hybrid generalized inverses on the set $M_{m,n}(\mathbb{C})$ of all complex $m \times n$ matrices. Note here that for $A \in M_{m,n}(\mathbb{C})$

2010 *Mathematics Subject Classification.* 15A09, 06F25, 06A06.

Key words and phrases. IMP-inverse; MP1-inverse; partial order; *-ring; Rickart *-ring.

This first author acknowledges the financial support from the Slovene Research Agency, ARRS (research core funding No. P1-0288). The second author is supported by the Ministry of Education, Science and Technological Development, Republic of Serbia (grant no. 174007 (451-03-9/2021-14/200124)). The authors also acknowledge the bilateral projects between Serbia and Slovenia (Generalized inverses, operator equations and applications), and Slovenia and Latvia (Ordered structures in Rickart rings) were financially supported by the Ministry of Education, Science and Technological Development, Republic of Serbia (grant no. 337-00-21/2020-09/32) and by the Slovene Research Agency, ARRS (grants BI-RS/20-21-039 and BI-LV/20-22-002).

the definitions of its inner generalized inverse $A^- \in M_{n,m}(\mathbb{C})$ and its unique Moore-Penrose inverse $A^\dagger \in M_{n,m}(\mathbb{C})$ are the same as on \mathcal{R} .

Definition 1. *Let $A \in M_{m,n}(\mathbb{C})$. For each inner generalized inverse A^- of A , the matrices*

$$A^{-\dagger} = A^-AA^\dagger \quad \text{and} \quad A^{\dagger-} = A^\dagger AA^-$$

are called a 1MP-inverse and a MP1-inverse of A , respectively.

An inner generalized inverse and the Moore-Penrose inverse of $A \in M_{m,n}(\mathbb{C})$ always exist and this guarantees the existence of a 1MP- and a MP1-inverse for every $A \in M_{m,n}(\mathbb{C})$. In general, 1MP- and MP1-inverses are not unique.

In [7], the 1MP- and MP1-inverses of $A \in M_{m,n}(\mathbb{C})$ were characterized. Also, a partial order induced by a 1MP-inverse and its dual case associated with a MP1-inverse were introduced. It is the aim of this paper to generalize the concept of a 1MP-inverse and a MP1-inverse to $*$ -regular rings or more generally to the set of all Moore-Penrose invertible elements in a $*$ -ring \mathcal{R} , study their properties, and the properties of partial orders on \mathcal{R} associated with these generalized inverses. We thus extend the results of [7]. The paper is structured as follows. In Section 2, we give some preliminaries and in Section 3 we introduce the concept of 1MP-inverses in the setting of $*$ -rings and present some characterizations of these generalized inverses. We introduce a relation associated with 1MP-inverses and study its properties in Section 4. The dual case (the MP1-inverses and a relation associated with these generalized inverses) is studied in Section 5. In the last section, we extend to the setting of Rickart $*$ -rings the concept of another partial order, called the plus order, which has been recently introduced in [1] on the set $B(\mathcal{H}, \mathcal{K})$ of all bounded linear operators between Hilbert spaces \mathcal{H} and \mathcal{K} .

2. PRELIMINARIES

Unless stated otherwise, let from now \mathcal{R} be a $*$ -ring with the (multiplicative) identity 1. If for $p \in \mathcal{R}$, $p^2 = p$, then p is said to be an idempotent. A projection $p \in \mathcal{R}$ is a self-adjoint idempotent, i.e. $p = p^2 = p^*$. The equality $1 = e_1 + e_2 + \cdots + e_n$, where e_1, e_2, \dots, e_n are idempotents in \mathcal{R} and $e_i e_j = 0$ for $i \neq j$, is called a decomposition of the identity of \mathcal{R} . Let $1 = e_1 + e_2 + \cdots + e_n$ and $1 = f_1 + f_2 + \cdots + f_n$ be two decompositions of the identity of \mathcal{R} . We have

$$x = 1 \cdot x \cdot 1 = (e_1 + e_2 + \cdots + e_n)x(f_1 + f_2 + \cdots + f_n) = \sum_{i,j=1}^n e_i x f_j.$$

Then any $x \in \mathcal{R}$ can be uniquely represented in the following matrix form:

$$(2) \quad x = \begin{bmatrix} x_{11} & \cdots & x_{1n} \\ \vdots & \ddots & \vdots \\ x_{n1} & \cdots & x_{nn} \end{bmatrix}_{e \times f},$$

where $x_{ij} = e_i x f_j \in e_i \mathcal{R} f_j$. With $e \times f$ we emphasize the use of the decompositions of the identity $1 = e_1 + e_2 + \cdots + e_n$ on the left side and $1 = f_1 + f_2 + \cdots + f_n$ on the right side of $x = 1 \cdot x \cdot 1$. If $x = (x_{ij})_{e \times f}$ and $y = (y_{ij})_{e \times f}$, then $x + y = (x_{ij} + y_{ij})_{e \times f}$. Moreover, if $1 = g_1 + \cdots + g_n$ is another decomposition of the identity of \mathcal{R} and $z = (z_{ij})_{f \times g}$, then, by the orthogonality of the idempotents involved, $xz = (\sum_{k=1}^n x_{ik} z_{kj})_{e \times g}$. Thus, if we

have decompositions of the identity of \mathcal{R} , then the usual algebraic operations in \mathcal{R} can be interpreted as simple operations between appropriate $n \times n$ matrices over \mathcal{R} . When $n = 2$ and $p, q \in \mathcal{R}$ are idempotents, we may write

$$x = pxq + px(1 - q) + (1 - p)xq + (1 - p)x(1 - q) = \begin{bmatrix} x_{11} & x_{12} \\ x_{21} & x_{22} \end{bmatrix}_{p \times q}.$$

Here $x_{11} = pxq$, $x_{12} = px(1 - q)$, $x_{21} = (1 - p)xq$, $x_{22} = (1 - p)x(1 - q)$.

By (2) we may write

$$x^* = \begin{bmatrix} x_{11}^* & \cdots & x_{n1}^* \\ \vdots & \ddots & \vdots \\ x_{1n}^* & \cdots & x_{nn}^* \end{bmatrix}_{f^* \times e^*},$$

where this matrix representation of x^* is given relative to the decompositions of the identity $1 = f_1^* + \cdots + f_n^*$ and $1 = e_1^* + \cdots + e_n^*$.

Let $a \in \mathcal{R}$ and a° denote the right annihilator of a , i.e. the set $a^\circ = \{x \in \mathcal{R} : ax = 0\}$. Similarly we denote the left annihilator ${}^\circ a$ of a , i.e. the set ${}^\circ a = \{x \in \mathcal{R} : xa = 0\}$. Suppose that $p, q \in \mathcal{R}$ are such idempotents that ${}^\circ a = {}^\circ p$ and $a^\circ = q^\circ$. Observe (or see [4, Lemma 2.2]) that ${}^\circ p = \mathcal{R}(1 - p)$ and $q^\circ = (1 - q)\mathcal{R}$. It follows that then $a = paq$, i.e.

$$(3) \quad a = \begin{bmatrix} a & 0 \\ 0 & 0 \end{bmatrix}_{p \times q}.$$

Suppose $a \in \mathcal{R}^{(1)}$, fix $h \in a\{1\}$, and let $p = ah$ and $q = ha$. Clearly, p and q are idempotents. Moreover, $pa = a$ and $aq = a$, and so ${}^\circ a = {}^\circ p$ and $a^\circ = q^\circ$. We may thus write a in the matrix form (3). Also, by [14, page 1044], $k \in a\{1\}$ if and only if

$$(4) \quad k = \begin{bmatrix} hah & k_{12} \\ k_{21} & k_{22} \end{bmatrix}_{q \times p}$$

where $k_{12} \in q\mathcal{R}(1 - p)$, $k_{21} \in (1 - q)\mathcal{R}p$, and $k_{22} \in (1 - q)\mathcal{R}(1 - p)$.

Suppose that $a \in \mathcal{R}^\dagger$ and let now $p = aa^\dagger$ and $q = a^\dagger a$. Then, p and q are projections and ${}^\circ a = {}^\circ p$ and $a^\circ = q^\circ$. Also, since $a^\dagger p = a^\dagger = qa^\dagger$, we may conclude that ${}^\circ a^\dagger = {}^\circ q$ and $a^{\dagger \circ} = p^\circ$. So,

$$a^\dagger = \begin{bmatrix} a^\dagger & 0 \\ 0 & 0 \end{bmatrix}_{q \times p}.$$

3. 1MP-INVERSES IN A *-RING

Let us now extend the concept of 1MP-inverses to the set of all Moore-Penrose invertible elements in a *-ring \mathcal{R} .

Definition 2. Let $a \in \mathcal{R}^\dagger$. For each $a^- \in a\{1\}$, the element

$$a^{-\dagger} = a^- aa^\dagger$$

is called a 1MP-inverse of a . The set of all 1MP-inverses of a is denoted by $a\{-\dagger\}$.

Since $a^\dagger = a^\dagger aa^\dagger$, we observe that $a^\dagger \in a\{-\dagger\}$. So, for every $a \in \mathcal{R}^\dagger$, the set $a\{-\dagger\}$ is nonempty. Let $a \in \mathcal{R}^\dagger$ and $a^- \in a\{1\}$. Denote $p = aa^\dagger$ and $q = a^\dagger a$. Then

$$a = \begin{bmatrix} a & 0 \\ 0 & 0 \end{bmatrix}_{p \times q} \quad \text{and} \quad a^\dagger = \begin{bmatrix} a^\dagger & 0 \\ 0 & 0 \end{bmatrix}_{q \times p}.$$

Observe that by (4)

$$a^- = \begin{bmatrix} a^\dagger a a^\dagger & d_{12} \\ d_{21} & d_{22} \end{bmatrix}_{q \times p} = \begin{bmatrix} a^\dagger & d_{12} \\ d_{21} & d_{22} \end{bmatrix}_{q \times p}$$

for some $d_{12} \in q\mathcal{R}(1-p)$, $d_{21} \in (1-q)\mathcal{R}p$, and $d_{22} \in (1-q)\mathcal{R}(1-p)$. Thus,

$$\begin{aligned} a^- a a^\dagger &= \begin{bmatrix} a^\dagger & d_{12} \\ d_{21} & d_{22} \end{bmatrix}_{q \times p} \begin{bmatrix} a & 0 \\ 0 & 0 \end{bmatrix}_{p \times q} \begin{bmatrix} a^\dagger & 0 \\ 0 & 0 \end{bmatrix}_{q \times p} \\ &= \begin{bmatrix} a^\dagger & d_{12} \\ d_{21} & d_{22} \end{bmatrix}_{q \times p} \begin{bmatrix} p & 0 \\ 0 & 0 \end{bmatrix}_{p \times p} \\ &= \begin{bmatrix} a^\dagger p & 0 \\ d_{21} p & 0 \end{bmatrix}_{q \times p} = \begin{bmatrix} a^\dagger & 0 \\ d_{21} & 0 \end{bmatrix}_{q \times p}. \end{aligned}$$

Since $\begin{bmatrix} a^\dagger & d_{12} \\ d_{21} & d_{22} \end{bmatrix}_{q \times p} \in a\{1\}$ for any $d_{12} \in q\mathcal{R}(1-p)$, $d_{21} \in (1-q)\mathcal{R}p$, and $d_{22} \in (1-q)\mathcal{R}(1-p)$, we may conclude that

$$(5) \quad a\{-\dagger\} = \left\{ \begin{bmatrix} a^\dagger & 0 \\ d_{21} & 0 \end{bmatrix}_{q \times p} : d_{21} \in (1-q)\mathcal{R}p \text{ is arbitrary} \right\}.$$

We will next give some characterizations of 1MP-inverses. First, let us prove a simple auxiliary result.

Lemma 3.1. *Let $a \in \mathcal{R}^\dagger$. Then $a\{-\dagger\} \subseteq a\{1, 2, 3\}$, $a^{-\dagger}a = a^-a$, and $aa^{-\dagger} = aa^\dagger$.*

Proof. Let $a^{-\dagger} \in a\{-\dagger\}$, i.e. $a^{-\dagger} = a^-aa^\dagger$ for some $a^- \in a\{1\}$. From

$$\begin{aligned} aa^{-\dagger}a &= aa^-aa^\dagger a = a, \\ a^{-\dagger}aa^{-\dagger} &= a^-aa^\dagger aa^-aa^\dagger = a^-aa^\dagger = a^{-\dagger}, \end{aligned}$$

and

$$(aa^{-\dagger})^* = (aa^-aa^\dagger)^* = (aa^\dagger)^* = aa^\dagger = aa^-aa^\dagger = aa^{-\dagger},$$

it follows that $a^{-\dagger} \in a\{1, 2, 3\}$ and thus $a\{-\dagger\} \subseteq a\{1, 2, 3\}$. Also, $aa^{-\dagger} = aa^\dagger$ and $a^{-\dagger}a = a^-aa^\dagger a = a^-a$. \square

Remark 3.2. *Let $a \in \mathcal{R}^\dagger$. Since aa^\dagger is a projection and a^-a is an idempotent, we may conclude, by Lemma 3.1, that $aa^{-\dagger}$ is a projection and $a^{-\dagger}a$ is an idempotent. Moreover, from $a = (aa^\dagger)a$ and since for $z \in \mathcal{R}$, $za = 0$ implies $zaa^\dagger = 0$, we have that $^\circ(aa^{-\dagger}) = ^\circ(aa^\dagger) = ^\circ a$. Similarly we obtain that $(a^{-\dagger}a)^\circ = a^\circ$.*

By Lemma 3.1 it follows that every $x = a^{-\dagger} \in a\{-\dagger\}$ satisfies the following system of equations:

$$(6) \quad xax = x \quad \text{and} \quad ax = aa^\dagger.$$

With the next theorem we extend [7, Theorem 3.1] from the set of all $m \times n$ complex matrices to the set \mathcal{R}^\dagger in a $*$ -ring \mathcal{R} with identity.

Theorem 3.3. *Let $a \in \mathcal{R}^\dagger$. Then the following statements are equivalent:*

- (i) $z \in a\{-\dagger\}$;
- (ii) $z \in \mathcal{R}$ is a solution of the system (6);

(iii) $z \in a\{1, 2, 3\}$.

Proof. (i) \Rightarrow (ii): It follows by Lemma 3.1.

(ii) \Rightarrow (i): Assume that for $z \in \mathcal{R}$ we have $zaz = z$ and $az = aa^\dagger$. Let $p = aa^\dagger$ and $q = a^\dagger a$. Then

$$a = \begin{bmatrix} a & 0 \\ 0 & 0 \end{bmatrix}_{p \times q} \quad \text{and} \quad a^\dagger = \begin{bmatrix} a^\dagger & 0 \\ 0 & 0 \end{bmatrix}_{q \times p}.$$

Let $z = \begin{bmatrix} z_{11} & z_{12} \\ z_{21} & z_{22} \end{bmatrix}_{q \times p}$. Then

$$az = \begin{bmatrix} a & 0 \\ 0 & 0 \end{bmatrix}_{p \times q} \begin{bmatrix} z_{11} & z_{12} \\ z_{21} & z_{22} \end{bmatrix}_{q \times p} = \begin{bmatrix} az_{11} & az_{12} \\ 0 & 0 \end{bmatrix}_{p \times p}$$

and

$$aa^\dagger = p = \begin{bmatrix} p & 0 \\ 0 & 0 \end{bmatrix}_{p \times p}.$$

So, $az_{12} = 0$ and since $z_{12} \in q\mathcal{R}(1-p)$ and $a^\circ = q^\circ$ this implies that $z_{12} = qz_{12} = 0$. Also, $az_{11} = p$, i.e. $az_{11} = aa^\dagger$. Since $z_{11} \in q\mathcal{R}p$ we therefore get

$$z_{11} = qz_{11} = a^\dagger az_{11} = a^\dagger aa^\dagger = a^\dagger.$$

From $z = zaz$ we obtain

$$\begin{aligned} z &= \begin{bmatrix} a^\dagger & 0 \\ z_{21} & z_{22} \end{bmatrix}_{q \times p} \begin{bmatrix} a & 0 \\ 0 & 0 \end{bmatrix}_{p \times q} \begin{bmatrix} a^\dagger & 0 \\ z_{21} & z_{22} \end{bmatrix}_{q \times p} \\ &= \begin{bmatrix} a^\dagger & 0 \\ z_{21} & z_{22} \end{bmatrix}_{q \times p} \begin{bmatrix} p & 0 \\ 0 & 0 \end{bmatrix}_{p \times p} \\ &= \begin{bmatrix} a^\dagger p & 0 \\ z_{21} p & 0 \end{bmatrix}_{q \times p} = \begin{bmatrix} a^\dagger & 0 \\ z_{21} & 0 \end{bmatrix}_{q \times p}. \end{aligned}$$

By (5) we may conclude that $z \in a\{-\dagger\}$.

(ii) \Rightarrow (iii): Let $z \in \mathcal{R}$ be a solution of the system (6). Since statement (ii) implies statement (i), $z \in a\{-\dagger\}$ and thus by Lemma 3.1 $z \in a\{1, 2, 3\}$.

(iii) \Rightarrow (ii): Let $z \in a\{1, 2, 3\}$, i.e. $aza = a$, $zaz = z$, and $(az)^* = az$. Let $p = aa^\dagger$ and $q = a^\dagger a$. To conclude the proof we must show that $az = aa^\dagger = p$. Since $z \in a\{1\}$, we may write by (4)

$$z = \begin{bmatrix} a^\dagger & z_{12} \\ z_{21} & z_{22} \end{bmatrix}_{q \times p}.$$

So,

$$az = \begin{bmatrix} a & 0 \\ 0 & 0 \end{bmatrix}_{p \times q} \begin{bmatrix} a^\dagger & z_{12} \\ z_{21} & z_{22} \end{bmatrix}_{q \times p} = \begin{bmatrix} p & az_{12} \\ 0 & 0 \end{bmatrix}_{p \times p}.$$

Recall that $p = p^*$. Thus

$$(az)^* = \begin{bmatrix} p & 0 \\ (az_{12})^* & 0 \end{bmatrix}_{p \times p}.$$

From $(az)^* = az$ we get $az_{12} = 0$. Since $a^\circ = q^\circ$, we get $0 = qz_{12} = z_{12}$. It follows that

$$az = \begin{bmatrix} p & 0 \\ 0 & 0 \end{bmatrix}_{p \times p} = p. \quad \square$$

Corollary 3.4. *Let $a \in \mathcal{R}^\dagger$ and fix $a^{-\dagger} \in a\{-\dagger\}$. Then*

$$a\{-\dagger\} = \left\{ a^{-\dagger} + \left(1 - a^{-\dagger}a\right) waa^{-\dagger} : w \in \mathcal{R} \text{ is arbitrary} \right\}.$$

Proof. Let

$$\mathcal{S} = \left\{ a^{-\dagger} + \left(1 - a^{-\dagger}a\right) waa^{-\dagger} : w \in \mathcal{R} \text{ is arbitrary} \right\}.$$

Let us first show that $\mathcal{S} \subseteq a\{-\dagger\}$. By Lemma 3.1, $aa^{-\dagger} = aa^\dagger$ and $aa^{-\dagger}a = a$, and thus

$$\begin{aligned} a \left(a^{-\dagger} + \left(1 - a^{-\dagger}a\right) waa^{-\dagger} \right) &= aa^{-\dagger} + \left(a - aa^{-\dagger}a \right) waa^{-\dagger} \\ &= aa^{-\dagger} + (a - a) waa^{-\dagger} \\ &= aa^{-\dagger} = aa^\dagger. \end{aligned}$$

Also, by Lemma 3.1, $a^{-\dagger}aa^{-\dagger} = a^{-\dagger}$ and hence

$$\left(a^{-\dagger} + \left(1 - a^{-\dagger}a\right) waa^{-\dagger} \right) a \left(a^{-\dagger} + \left(1 - a^{-\dagger}a\right) waa^{-\dagger} \right) = \left(a^{-\dagger} + \left(1 - a^{-\dagger}a\right) waa^{-\dagger} \right) aa^{-\dagger} = a^{-\dagger} + \left(1 - a^{-\dagger}a\right) waa^{-\dagger}. \text{ By Theorem 3.3 ((i) \Leftrightarrow (ii)) we have that } \mathcal{S} \subseteq a\{-\dagger\}.$$

To prove the reverse inclusion, let $z \in a\{-\dagger\}$. Then there exists $c \in a\{1\}$ such that $z = caa^\dagger$. Note that there exists $a^- \in a\{1\}$ such that $a^{-\dagger} = a^-aa^\dagger$, and let $p = aa^-$ and $q = a^-a$. Then $qcp = a^-(aca)a^- = a^-aa^-$ and we may thus write $c = a^-aa^- + c - qcp$. It follows that

$$\begin{aligned} z &= caa^\dagger = (a^-aa^- + c - qcp) aa^\dagger \\ &= a^-(aa^-a)a^\dagger + caa^\dagger - a^-ac(aa^-a)a^\dagger \\ &= a^{-\dagger} + (1 - a^-a) caa^\dagger. \end{aligned}$$

By Lemma 3.1, $a^{-\dagger}a = a^-a$ and $aa^{-\dagger} = aa^\dagger$. Thus, $z = a^{-\dagger} + (1 - a^{-\dagger}a) caa^{-\dagger}$ and so $z \in \mathcal{S}$, i.e. $a\{-\dagger\} \subseteq \mathcal{S}$. \square

We say that $a \in \mathcal{R}^\dagger$ is a partial isometry if $a^* = a^\dagger$ (see [13]). We now apply Theorem 3.3 and its corollary to the subset of all partial isometries in \mathcal{R} .

Proposition 3.5. *Let $a \in \mathcal{R}^\dagger$ be a partial isometry. Then the system $xax = x$, $ax = aa^*$ has a solution $x = c$ if and only if $c = a^-aa^* + (1 - a^-a)waa^*$ where $a^- \in a\{1\}$ and $w \in \mathcal{R}$.*

Proof. Suppose first the system $xax = x$, $ax = aa^*$ has a solution $x = c$. Since $a^* = a^\dagger$ we have $cac = c$ and $ac = aa^\dagger$. By Theorem 3.3, $c \in a\{-\dagger\}$, and thus by Corollary 3.4, $c = a^{-\dagger} + (1 - a^{-\dagger}a) waa^{-\dagger}$ for some $w \in \mathcal{R}$ and $a^- \in a\{1\}$ where $a^{-\dagger} = a^-aa^\dagger$. By Lemma 3.1, $a^{-\dagger}a = a^-a$, and $aa^{-\dagger} = aa^\dagger = aa^*$. So, $c = a^-aa^\dagger + (1 - a^-a) waa^*$.

Conversely, let $a^- \in a\{1\}$, $w \in \mathcal{R}$ and $c = a^-aa^* + (1 - a^-a)waa^*$. Then

$$ac = (aa^-a)a^* + (a - aa^-a)waa^* = aa^*$$

and

$$\begin{aligned} cac &= (a^-aa^* + (1 - a^-a)waa^*) (aa^-aa^* + (a - aa^-a) waa^*) \\ &= (a^-aa^* + (1 - a^-a)waa^*) aa^* \\ &= a^-aa^\dagger aa^\dagger + (1 - a^-a)waa^\dagger aa^\dagger \\ &= a^-aa^\dagger + (1 - a^-a)waa^\dagger \\ &= a^-aa^* + (1 - a^-a)waa^* = c, \end{aligned}$$

and hence $x = c$ is a solution of the system $xax = x$, $ax = aa^*$. \square

Several necessary and sufficient conditions for $x = a^{-\dagger}$ are next developed.

Lemma 3.6. *Let $a \in \mathcal{R}^\dagger$ and $a^- \in a\{1\}$. Then the following statements are equivalent:*

- (i) $x = a^{-\dagger}$;
- (ii) $ax = aa^\dagger$ and $x = a^-ax$;
- (iii) $a^*ax = a^*$ and $x = a^-ax$;
- (iv) $xa = a^-a$ and $x = xaa^\dagger$;
- (v) $xaa^- = a^-aa^-$ and $x = xaa^\dagger$;
- (vi) $axa = a$ and $a^-axaa^\dagger = x$;
- (vii) $xax = x$ and $a^*ax = a^*$.

Proof. (i) \Rightarrow (ii)–(vii): It can be verified by $x = a^-aa^\dagger$.

(ii) \Rightarrow (i): The assumptions $ax = aa^\dagger$ and $x = a^-ax$ give $x = a^-(ax) = a^-aa^\dagger$.

(iii) \Rightarrow (ii): Multiplying $a^*ax = a^*$ by $(a^\dagger)^*$ from the left hand side, we obtain $ax = aa^\dagger$.

(iv) \Rightarrow (i): From $xa = a^-a$ and $x = xaa^\dagger$, we have $x = (xa)a^\dagger = a^-aa^\dagger$.

The rest follows similarly. \square

We can observe that the existence of 1MP-inverses is closely related with the existence of adequate idempotents. If $x \in \mathcal{R}$ satisfies $(ax)^* = ax$, then it is called a $\{4\}$ -inverse of a . If $x \in \mathcal{R}$ satisfies the first two and the fourth equations in (1), then it is called a $\{1, 2, 4\}$ -inverse of a . We denote the sets of all $\{4\}$ -inverses and $\{1, 2, 4\}$ -inverses of $a \in \mathcal{R}$ by $a\{4\}$ and $a\{1, 2, 4\}$, respectively.

Theorem 3.7. *Let $a\{4\} \neq \emptyset$. The following statements are equivalent:*

- (i) $a\{-\dagger\} \neq \emptyset$;
- (ii) *there exist a projection $p \in \mathcal{R}$ and an idempotent $q \in \mathcal{R}$ such that $p\mathcal{R} = a\mathcal{R}$ and $\mathcal{R}q = \mathcal{R}a$.*

In addition, for arbitrary $a^- \in a\{1\}$, $qa^-p \in a\{-\dagger\}$, that is,

$$q \cdot a\{1\} \cdot p \subseteq a\{-\dagger\}.$$

Proof. (i) \Rightarrow (ii): For $x = a^-aa^\dagger$, let $p = ax$ and $q = xa$. Since $p = aa^\dagger$ and $q = a^-a$, we have $p = p^2 = p^*$, $q = q^2$, $p\mathcal{R} = aa^\dagger\mathcal{R} = a\mathcal{R}$ and $\mathcal{R}q = \mathcal{R}a^-a = \mathcal{R}a$.

(ii) \Rightarrow (i): The hypothesis $p\mathcal{R} = a\mathcal{R}$ implies $a = pa$ and $p = au$, for some $u \in \mathcal{R}$. Now, by $a = aua$, $p^* = p = au$ and $a\{4\} \neq \emptyset$, we deduce that a^\dagger exists and thus $a\{-\dagger\} \neq \emptyset$.

Notice that $p = au = aa^\dagger au = (auaa^\dagger)^* = aa^\dagger$. Let $x = qa^-p$, for $a^- \in a\{1\}$. From $\mathcal{R}q = \mathcal{R}a$, we conclude that $a = aq$. Therefore, $ax = (aq)a^-p = aa^-p = p = aa^\dagger$ and $xax = xp = x$, i.e. $x \in a\{-\dagger\}$ by Theorem 3.3. \square

For $x, y \in a\{-\dagger\}$, we prove that $xay \in a\{-\dagger\}$.

Theorem 3.8. *Let $a \in \mathcal{R}^\dagger$. Then*

$$a\{-\dagger\} \cdot a \cdot a\{-\dagger\} \subseteq a\{-\dagger\}.$$

Proof. Let $x, y \in a\{-\dagger\}$ and $x_1 = xay$. Then, by Theorem 3.3, $ax_1 = (axa)y = ay = aa^\dagger$ and $x_1ax_1 = x_1aa^\dagger = xay(aa^\dagger) = xa(yay) = xay = x_1$, i.e. $x_1 \in a\{-\dagger\}$. So, $a\{-\dagger\} \cdot a \cdot a\{-\dagger\} \subseteq a\{-\dagger\}$. \square

4. THE 1MP-PARTIAL ORDER

A partial order associated with 1MP-inverses was introduced on $M_{m,n}(\mathbb{C})$ in [7]. Let $A, B \in M_{m,n}(\mathbb{C})$. We say that A is below B with respect to the relation $\leq^{-\dagger}$ and write $A \leq^{-\dagger} B$ if there exists a 1MP-inverse $A^{-\dagger}$ of A such that

$$A^{-\dagger}A = A^{-\dagger}B \quad \text{and} \quad AA^{-\dagger} = BA^{-\dagger}.$$

It was proved in [7] that the relation $\leq^{-\dagger}$ is a partial order. We now extend this definition to $*$ -rings with identity.

Definition 3. Let $a \in \mathcal{R}^{\dagger}$ and $b \in \mathcal{R}$. We say that a is below b with respect to the 1MP-relation $\leq^{-\dagger}$ and write $a \leq^{-\dagger} b$ if there exists $a^{-\dagger} \in a\{-\dagger\}$ such that $a^{-\dagger}a = a^{-\dagger}b$ and $aa^{-\dagger} = ba^{-\dagger}$.

We will prove that the 1MP-relation $\leq^{-\dagger}$ is a partial order on \mathcal{R}^{\dagger} for any $*$ -ring \mathcal{R} with identity. First, let us present and prove two results that extend Theorem 4.2 in [7]. With the next theorem we describe the set of all elements $b \in \mathcal{R}$ that are above a given $a \in \mathcal{R}^{\dagger}$ with respect to the 1MP-relation.

Theorem 4.1. Let $a \in \mathcal{R}^{\dagger}$ and denote $p = aa^{\dagger}$ and $q = a^{\dagger}a$. Let $b \in \mathcal{R}$. Then the following statements are equivalent:

- (i) $a \leq^{-\dagger} b$;
- (ii)

$$(7) \quad b = \begin{bmatrix} a & 0 \\ -b_4da & b_4 \end{bmatrix}_{p \times q}$$

where $b_4 \in (1-p)\mathcal{R}(1-q)$ and $d \in (1-q)\mathcal{R}p$.

Proof. Suppose first $a \leq^{-\dagger} b$ for $a \in \mathcal{R}^{\dagger}$ and $b \in \mathcal{R}$. Then there exists $a^{-\dagger} \in a\{-\dagger\}$ such that $a^{-\dagger}a = a^{-\dagger}b$ and $aa^{-\dagger} = ba^{-\dagger}$. By (5) there is $d \in (1-q)\mathcal{R}p$ such that

$$a^{-\dagger} = \begin{bmatrix} a^{\dagger} & 0 \\ d & 0 \end{bmatrix}_{q \times p}.$$

Let $b = \begin{bmatrix} b_1 & b_2 \\ b_3 & b_4 \end{bmatrix}_{p \times q}$. Then

$$\begin{aligned} a^{-\dagger}b &= \begin{bmatrix} a^{\dagger} & 0 \\ d & 0 \end{bmatrix}_{q \times p} \begin{bmatrix} b_1 & b_2 \\ b_3 & b_4 \end{bmatrix}_{p \times q} \\ &= \begin{bmatrix} a^{\dagger}b_1 & a^{\dagger}b_2 \\ db_1 & db_2 \end{bmatrix}_{q \times q} \end{aligned}$$

which equals

$$a^{-\dagger}a = \begin{bmatrix} a^{\dagger} & 0 \\ d & 0 \end{bmatrix}_{q \times p} \begin{bmatrix} a & 0 \\ 0 & 0 \end{bmatrix}_{p \times q} = \begin{bmatrix} a^{\dagger}a & 0 \\ da & 0 \end{bmatrix}_{q \times q}.$$

So, $a^{\dagger}a = a^{\dagger}b_1$ and $a^{\dagger}b_2 = 0$, and thus $a = aa^{\dagger}b_1 = pb_1$ and $0 = aa^{\dagger}b_2 = pb_2$. Since $b_1, b_2 \in p\mathcal{R}$, we get $a = b_1$ and $b_2 = 0$. It follows that

$$ba^{-\dagger} = \begin{bmatrix} a & 0 \\ b_3 & b_4 \end{bmatrix}_{p \times q} \begin{bmatrix} a^{\dagger} & 0 \\ d & 0 \end{bmatrix}_{q \times p} = \begin{bmatrix} p & 0 \\ b_3a^{\dagger} + b_4d & 0 \end{bmatrix}_{p \times p}$$

which equals $aa^\dagger = \begin{bmatrix} p & 0 \\ 0 & 0 \end{bmatrix}_{p \times p}$. So, $b_3a^\dagger + b_4d = 0$, i.e. $b_3a^\dagger = -b_4d$, and thus $b_3q = -b_4da$. Since $b_3 \in \mathcal{R}q$, we have that $b_3 = -b_4da$ and therefore

$$b = \begin{bmatrix} a & 0 \\ -b_4da & b_4 \end{bmatrix}_{p \times q}.$$

To prove the converse implication, let $a \in \mathcal{R}^\dagger$ and let $b \in \mathcal{R}$ be of matrix form (7). Define

$$c = \begin{bmatrix} a^\dagger & 0 \\ d & 0 \end{bmatrix}_{q \times p}$$

where the element d is as in (7). By (5), $c \in a\{-\dagger\}$. We have

$$ca = \begin{bmatrix} a^\dagger & 0 \\ d & 0 \end{bmatrix}_{q \times p} \begin{bmatrix} a & 0 \\ 0 & 0 \end{bmatrix}_{p \times q} = \begin{bmatrix} q & 0 \\ da & 0 \end{bmatrix}_{q \times q}$$

and

$$cb = \begin{bmatrix} a^\dagger & 0 \\ d & 0 \end{bmatrix}_{q \times p} \begin{bmatrix} a & 0 \\ -b_4da & b_4 \end{bmatrix}_{p \times q} = \begin{bmatrix} q & 0 \\ da & 0 \end{bmatrix}_{q \times q}$$

and thus $ca = cb$. Also, $ac = aa^\dagger = p$ and since $d \in \mathcal{R}p$ we get

$$\begin{aligned} bc &= \begin{bmatrix} a & 0 \\ -b_4da & b_4 \end{bmatrix}_{p \times q} \begin{bmatrix} a^\dagger & 0 \\ d & 0 \end{bmatrix}_{q \times p} = \begin{bmatrix} p & 0 \\ -b_4dp + b_4d & 0 \end{bmatrix}_{p \times p} \\ &= \begin{bmatrix} p & 0 \\ -b_4d + b_4d & 0 \end{bmatrix}_{p \times p} = \begin{bmatrix} p & 0 \\ 0 & 0 \end{bmatrix}_{p \times p} = p. \end{aligned}$$

So, $ac = bc$ and therefore $a \leq^{-\dagger} b$. \square

For $a, b \in \mathcal{R}^\dagger$ where $a \leq^{-\dagger} b$ the next result describes the set $b\{-\dagger\}$ of all 1MP-inverses of b .

Theorem 4.2. *Let $a, b \in \mathcal{R}^\dagger$, suppose $a \leq^{-\dagger} b$, and represent b in the matrix form (7). Then $x \in b\{-\dagger\}$ if and only if there exist $x_3 \in (1-q)\mathcal{R}p$ and $x_4 \in (1-q)\mathcal{R}(1-p)$ such that*

$$x = \begin{bmatrix} a^\dagger & 0 \\ x_3 & x_4 \end{bmatrix}_{q \times p}$$

where (for b_4 and d from (7)) $b_4x_3 = b_4d$ and $x_4 \in b_4\{-\dagger\}$.

Proof. Suppose first $x \in b\{-\dagger\}$ and let

$$x = \begin{bmatrix} x_1 & x_2 \\ x_3 & x_4 \end{bmatrix}_{q \times p}$$

for $p = aa^\dagger$ and $q = a^\dagger a$. By Theorem 3.3, $b\{-\dagger\} = b\{1, 2, 3\}$ and so $bx b = b$, $xbx = x$, and $(bx)^* = bx$. We have

$$\begin{aligned} b &= bxb = \begin{bmatrix} a & 0 \\ -b_4da & b_4 \end{bmatrix}_{p \times q} \begin{bmatrix} x_1 & x_2 \\ x_3 & x_4 \end{bmatrix}_{q \times p} \begin{bmatrix} a & 0 \\ -b_4da & b_4 \end{bmatrix}_{p \times q} \\ &= \begin{bmatrix} ax_1a - ax_2b_4da & ax_2b_4 \\ (-b_4dax_1 + b_4x_3)a - (-b_4dax_2 + b_4x_4)b_4da & (-b_4dax_2 + b_4x_4)b_4 \end{bmatrix}_{p \times q} \end{aligned}$$

and therefore $ax_2b_4 = 0$. Also, $ax_1a - ax_2b_4da = a$ and thus $ax_1a = a$. Hence, $a^\dagger ax_1aa^\dagger = a^\dagger aa^\dagger = a^\dagger$ and so $qx_1p = a^\dagger$. Since $x_1 \in q\mathcal{R}p$, it follows that $x_1 = a^\dagger$. The element in the second row and second column yields $-b_4dax_2b_4 + b_4x_4b_4 = b_4$ and since $ax_2b_4 = 0$, we have

$$(8) \quad b_4x_4b_4 = b_4.$$

Also, $-b_4dax_1a + b_4x_3a + b_4dax_2b_4da - b_4x_4b_4da = -b_4da$. Since $ax_1a = a$, $ax_2b_4 = 0$, and $b_4x_4b_4 = b_4$, we get $b_4x_3a = b_4da$ and thus $b_4x_3p = b_4dp$. From $x_3, d \in \mathcal{R}p$ we obtain

$$b_4x_3 = b_4d.$$

It follows that

$$\begin{aligned} bx &= \begin{bmatrix} a & 0 \\ -b_4da & b_4 \end{bmatrix}_{p \times q} \begin{bmatrix} a^\dagger & x_2 \\ x_3 & x_4 \end{bmatrix}_{q \times p} \\ &= \begin{bmatrix} p & ax_2 \\ -b_4dp + b_4x_3 & -b_4dax_2 + b_4x_4 \end{bmatrix}_{p \times p} \\ &= \begin{bmatrix} p & ax_2 \\ 0 & -b_4dax_2 + b_4x_4 \end{bmatrix}_{p \times p}. \end{aligned}$$

Note that then

$$(bx)^* = \begin{bmatrix} p & 0 \\ (ax_2)^* & (-b_4dax_2 + b_4x_4)^* \end{bmatrix}_{p \times p}$$

and thus $ax_2 = 0$ which implies $qx_2 = 0$. Since $x_2 \in q\mathcal{R}$, we have $x_2 = 0$ and therefore

$$x = \begin{bmatrix} a^\dagger & 0 \\ x_3 & x_4 \end{bmatrix}_{q \times p}$$

and $-b_4dax_2 + b_4x_4 = b_4x_4$. So, $(bx)^* = bx$ implies

$$(9) \quad (b_4x_4)^* = b_4x_4.$$

Finally,

$$x = xbx = \begin{bmatrix} a^\dagger & 0 \\ x_3 & x_4 \end{bmatrix}_{q \times p} \begin{bmatrix} a & 0 \\ -b_4da & b_4 \end{bmatrix}_{p \times q} \begin{bmatrix} a^\dagger & 0 \\ x_3 & x_4 \end{bmatrix}_{q \times p} = \begin{bmatrix} a^\dagger & 0 \\ * & x_4b_4x_4 \end{bmatrix}_{q \times p}$$

and thus

$$(10) \quad x_4 = x_4b_4x_4.$$

By equations (8)-(10) and Theorem 3.3 we establish that $x_4 \in b_4\{-\dagger\}$.

Conversely, let

$$x = \begin{bmatrix} a^\dagger & 0 \\ x_3 & x_4 \end{bmatrix}_{q \times p}$$

with $b_4x_3 = b_4d$ and $x_4 \in b_4\{-\dagger\}$. Since $b_4x_4b_4 = b_4$, $b_4x_3 = b_4d$, and $d \in \mathcal{R}p$, we obtain

$$\begin{aligned} bxb &= \begin{bmatrix} a & 0 \\ -b_4da & b_4 \end{bmatrix}_{p \times q} \begin{bmatrix} a^\dagger & 0 \\ x_3 & x_4 \end{bmatrix}_{q \times p} \begin{bmatrix} a & 0 \\ -b_4da & b_4 \end{bmatrix}_{p \times q} \\ &= \begin{bmatrix} a & 0 \\ (-b_4dp + b_4x_3)a - b_4x_4b_4da & b_4x_4b_4 \end{bmatrix}_{p \times q} \\ &= \begin{bmatrix} a & 0 \\ -b_4da & b_4 \end{bmatrix}_{p \times q} = b. \end{aligned}$$

From $x_4b_4x_4 = x_4$, $b_4x_3 = b_4d$, and $x_3, d \in \mathcal{R}p$ we get

$$\begin{aligned} bx &= \begin{bmatrix} a^\dagger & 0 \\ x_3 & x_4 \end{bmatrix}_{q \times p} \begin{bmatrix} a & 0 \\ -b_4da & b_4 \end{bmatrix}_{p \times q} \begin{bmatrix} a^\dagger & 0 \\ x_3 & x_4 \end{bmatrix}_{q \times p} \\ &= \begin{bmatrix} a^\dagger & 0 \\ x_3p - x_4b_4dp + x_4b_4x_3 & x_4b_4x_4 \end{bmatrix}_{q \times p} = \begin{bmatrix} a^\dagger & 0 \\ x_3 & x_4 \end{bmatrix}_{q \times p} = x \end{aligned}$$

and

$$\begin{aligned} bx &= \begin{bmatrix} a & 0 \\ -b_4da & b_4 \end{bmatrix}_{p \times q} \begin{bmatrix} a^\dagger & 0 \\ x_3 & x_4 \end{bmatrix}_{q \times p} \\ &= \begin{bmatrix} p & 0 \\ -b_4dp + b_4x_3 & b_4x_4 \end{bmatrix}_{p \times p} = \begin{bmatrix} p & 0 \\ 0 & b_4x_4 \end{bmatrix}_{p \times p}. \end{aligned}$$

Since $p^* = p$ and $(b_4x_4)^* = b_4x_4$, we establish that $bx = (bx)^*$. Thus, $x \in b\{1, 2, 3\}$ and therefore by Theorem 3.3, $x \in b\{-\dagger\}$. \square

Let us now prove that the relation $\leq^{-\dagger}$ is a partial order on \mathcal{R}^\dagger .

Theorem 4.3. *Let \mathcal{R} be a *-ring with identity. The IMP-relation $\leq^{-\dagger}$ is a partial order on \mathcal{R}^\dagger .*

Proof. The set $a\{-\dagger\}$ is nonempty for every $a \in \mathcal{R}^\dagger$ which implies that $\leq^{-\dagger}$ is reflexive. To prove that it is antisymmetric, let for $a, b \in \mathcal{R}^\dagger$, $a \leq^{-\dagger} b$ and $b \leq^{-\dagger} a$. Thus, there exists $b^{-\dagger} \in b\{-\dagger\}$ such that $b^{-\dagger}b = b^{-\dagger}a$ and $bb^{-\dagger} = ab^{-\dagger}$. Let $p = aa^\dagger$ and $q = a^\dagger a$. Since $a \leq^{-\dagger} b$, it follows by Theorems 4.1 and 4.2 that

$$(11) \quad b = \begin{bmatrix} a & 0 \\ -b_4da & b_4 \end{bmatrix}_{p \times q} \quad \text{and} \quad b^{-\dagger} = \begin{bmatrix} a^\dagger & 0 \\ x_3 & x_4 \end{bmatrix}_{q \times p}$$

where $b_4 \in (1-p)\mathcal{R}(1-q)$, $d, x_3 \in (1-q)\mathcal{R}p$, $x_4 \in (1-q)\mathcal{R}(1-p)$, $b_4x_3 = b_4d$, and $x_4 \in b_4\{-\dagger\}$. We have

$$b^{-\dagger}b = \begin{bmatrix} q & 0 \\ x_3a - x_4b_4da & x_4b_4 \end{bmatrix}_{q \times q}$$

and

$$b^{-\dagger}a = \begin{bmatrix} a^\dagger & 0 \\ x_3 & x_4 \end{bmatrix}_{q \times p} \begin{bmatrix} a & 0 \\ 0 & 0 \end{bmatrix}_{p \times q} = \begin{bmatrix} q & 0 \\ x_3a & 0 \end{bmatrix}_{q \times q}.$$

It follows that $x_4b_4 = 0$ and thus $b_4x_4b_4 = 0$. Since $x_4 \in b_4\{-\dagger\} = b_4\{1, 2, 3\}$, we obtain $b_4 = b_4x_4b_4 = 0$ and therefore

$$b = \begin{bmatrix} a & 0 \\ 0 & 0 \end{bmatrix}_{p \times q} = a.$$

Let us now prove that $\leq^{-\dagger}$ is transitive. Let $a \leq^{-\dagger} b$ and $b \leq^{-\dagger} c$ for $a, b \in \mathcal{R}^\dagger$ and $c \in \mathcal{R}$. There exists $b^{-\dagger} \in b\{-\dagger\}$ such that $b^{-\dagger}b = b^{-\dagger}c$ and $bb^{-\dagger} = cb^{-\dagger}$. Let again $p = aa^\dagger$ and $q = a^\dagger a$. Since $a \leq^{-\dagger} b$, it follows as before by Theorems 4.1 and 4.2 that b and $b^{-\dagger}$ are of the form (11). Let

$$c = \begin{bmatrix} c_1 & c_2 \\ c_3 & c_4 \end{bmatrix}_{p \times q}.$$

From $b^{-\dagger}b = b^{-\dagger}c$ we have

$$\begin{bmatrix} q & 0 \\ x_3a - x_4b_4da & x_4b_4 \end{bmatrix}_{q \times q} = \begin{bmatrix} a^\dagger c_1 & a^\dagger c_2 \\ x_3c_1 + x_4c_3 & x_3c_2 + x_4c_4 \end{bmatrix}_{q \times q}$$

and therefore $a^\dagger c_2 = 0$. Thus, $pc_2 = 0$ and so $c_2 = 0$ since $c_2 \in p\mathcal{R}$. Also, $q = a^\dagger c_1$ and hence $aa^\dagger a = aa^\dagger c_1$, i.e. $a = pc_1 = c_1$. By $bb^{-\dagger} = cb^{-\dagger}$ we thus obtain

$$\begin{bmatrix} p & 0 \\ -b_4dp + b_4x_3 & b_4x_4 \end{bmatrix}_{p \times p} = \begin{bmatrix} p & 0 \\ c_3a^\dagger + c_4x_3 & c_4x_4 \end{bmatrix}_{p \times p}.$$

So, $-b_4dp + b_4x_3 = c_3a^\dagger + c_4x_3$. Since $d \in \mathcal{R}p$ and $b_4x_3 = b_4d$, we have $c_3a^\dagger = -c_4x_3$. From $c_3 \in \mathcal{R}q$, we establish that $c_3 = -c_4x_3a$. It follows that

$$c = \begin{bmatrix} a & 0 \\ -c_4x_3a & c_4 \end{bmatrix}_{p \times q}$$

and therefore by Theorem 4.1, $a \leq^{-\dagger} c$. \square

We give some equivalent conditions for $a \leq^{-\dagger} b$ to be satisfied.

Lemma 4.4. *Let $a \in \mathcal{R}^\dagger$ and $b \in \mathcal{R}$. Then the following statements are equivalent:*

- (i) $a \leq^{-\dagger} b$;
- (ii) *there exist $x, y \in a\{-\dagger\}$ such that $xa = xb$ and $ay = by$;*
- (iii) *there exists $a^- \in a\{1\}$ such that $aa^\dagger b = a = ba^-a$.*

Proof. (i) \Rightarrow (ii): It is clear.

(ii) \Rightarrow (iii): Since $x = zaa^\dagger$ and $y = a^-aa^\dagger$, for some $z, a^- \in a\{1\}$, we obtain $a = a(xa) = axb = (aza)a^\dagger b = aa^\dagger b$ and $a = (ay)a = bya = ba^-(aa^\dagger a) = ba^-a$.

(iii) \Rightarrow (i): Using $aa^\dagger b = a = ba^-a$, for $x = a^-aa^\dagger$, we have $xa = a^-aa^\dagger a = a^-a = a^-aa^\dagger b = xb$ and $ax = aa^-aa^\dagger = aa^\dagger = ba^-aa^\dagger = bx$. Hence, $a \leq^{-\dagger} b$. \square

Theorem 4.5. *Let $a\{4\} \neq \emptyset$. The following statements are equivalent:*

- (i) $a \leq^{-\dagger} b$;
- (ii) *there exist a projection $p \in \mathcal{R}$ and an idempotent $q \in \mathcal{R}$ such that $p\mathcal{R} = a\mathcal{R}$, $\mathcal{R}q = \mathcal{R}a$ and $pb = a = bq$.*

Proof. (i) \Rightarrow (ii): Since $ax = bx$ and $xa = xb$ for some $x = a^-aa^\dagger$, set $p = ax$ and $q = xa$. The rest is clear by the proof of Theorem 3.7 and $pb = axb = axa = a = bxa = bq$.

(ii) \Rightarrow (i): By Theorem 3.7 and $pb = a = bq$, we have $x = qa^-p \in a\{-\dagger\}$ and $ax = (aq)a^-p = aa^-p = bqa^-p = bx$ and $xa = qa^-(pa) = qa^-a = qa^-pb = xb$. \square

Theorem 4.6. *Let $a, b \in \mathcal{R}^\dagger$. If $a \leq^{-\dagger} b$, then*

$$b\{-\dagger\} \cdot a \cdot b\{-\dagger\} \subseteq a\{-\dagger\}.$$

Proof. Assume that $ax = bx$ and $xa = xb$ for some $x \in a\{-\dagger\}$. Let $y, z \in b\{-\dagger\}$ and $x_1 = zay$. Then $ay = a(xa)y = axby = (byax)^* = (bybx)^* = (bx)^* = (ax)^* = ax = aa^\dagger$ and similarly $az = aa^\dagger$, which give $ax_1 = azay = aa^\dagger aa^\dagger = aa^\dagger$. Since $x_1ax_1 = zayazay = zaa^\dagger aa^\dagger ay = zay = x_1$, we deduce that $x_1 \in b\{-\dagger\}$. \square

One of the best known relations on \mathcal{R} that may be defined with generalized inverses is the *minus order* [6]. We say that $a \in \mathcal{R}^{(1)}$ is below $b \in \mathcal{R}$ with respect to the minus order and write $a \leq^- b$ when there exists $a^- \in a\{1\}$ such that

$$(12) \quad a^- a = a^- b \quad \text{and} \quad a a^- = b a^-.$$

It is known (see [6]) that the relation \leq^- is a partial order on $\mathcal{R}^{(1)}$ for any ring \mathcal{R} .

Clearly, since $a\{-\dagger\} \subseteq a\{1\}$, $a \leq^{-\dagger} b$ implies $a \leq^- b$. We study additional conditions for the converse to hold.

Theorem 4.7. *Let $a \in \mathcal{R}^\dagger$ and $b \in \mathcal{R}$. Then the following statements are equivalent:*

- (i) $a \leq^{-\dagger} b$;
- (ii) $a \leq^- b$ and $a^\dagger b = a^\dagger a$;
- (iii) $a a^- = b a^-$ and $a^\dagger b = a^\dagger a$, for some $a^- \in a\{1\}$.

Proof. (i) \Rightarrow (ii): Applying (3) and (7), notice that

$$a^\dagger b = \begin{bmatrix} a^\dagger & 0 \\ 0 & 0 \end{bmatrix}_{q \times p} \begin{bmatrix} a & 0 \\ -b_4 d a & b_4 \end{bmatrix}_{p \times q} = \begin{bmatrix} q & 0 \\ 0 & 0 \end{bmatrix}_{q \times q} = a^\dagger a.$$

(ii) \Rightarrow (iii): It is evident.

(iii) \Rightarrow (i): Suppose that $a a^- = b a^-$ and $a^\dagger b = a^\dagger a$, for some $a^- \in a\{1\}$. Then $a a^\dagger b = a = b a^- a$ implies $a \leq^{-\dagger} b$, by Lemma 4.4. \square

5. THE DUAL CASE

A dual version of partial order $\leq^{-\dagger}$ was introduced in [7] on $M_{m,n}(\mathbb{C})$. Let $A, B \in M_{m,n}(\mathbb{C})$. We say that A is below B with respect to the relation $\leq^{\dagger-}$ and write $A \leq^{\dagger-} B$ if there exists a MP1-inverse $A^{\dagger-}$ of A such that

$$A^{\dagger-} A = A^{\dagger-} B \quad \text{and} \quad A A^{\dagger-} = B A^{\dagger-}.$$

We now extend the concept of MP1-inverses and the MP1-relation $\leq^{\dagger-}$ to the setting of *-rings. Recall that \mathcal{R} denotes a *-ring with identity.

Definition 4. *Let $a \in \mathcal{R}^\dagger$. For each $a^- \in a\{1\}$, the element*

$$a^{\dagger-} = a^\dagger a a^-$$

is called a MP1-inverse of a . The set of all MP1-inverses of a is denoted by $a\{\dagger-\}$.

A MP1-inverse of $a \in \mathcal{R}^\dagger$ always exists however in general it is not necessarily unique.

Definition 5. *Let $a \in \mathcal{R}^\dagger$ and $b \in \mathcal{R}$. We say that a is below b with respect to the MP1-relation $\leq^{\dagger-}$ and write $a \leq^{\dagger-} b$ if there exists $a^{\dagger-} \in a\{\dagger-\}$ such that $a^{\dagger-} a = a^{\dagger-} b$ and $a a^{\dagger-} = b a^{\dagger-}$.*

Consider now a new *-ring $\mathcal{R}_L = (\mathcal{R}, +, \cdot_L, *)$ where for $a, b \in \mathcal{R}$,

$$(13) \quad a \cdot_L b := b a.$$

Let $a, c \in \mathcal{R}$. Then $aca = a$ if and only if $a \cdot_L c \cdot_L a = aca = a$. So, $c = a^-$ is an inner generalized inverse of a in \mathcal{R} if and only if $c = a^-$ is an inner generalized inverse of a in \mathcal{R}_L . Similarly we observe that for $a \in \mathcal{R}^\dagger$, its unique Moore-Penrose inverse a^\dagger in \mathcal{R} is also its unique Moore-Penrose inverse in \mathcal{R}_L . Moreover, since $(a \cdot_L c)^* = a \cdot_L c$ if and only if

$(ca)^* = ca$, we establish that the set $a\{1, 2, 3\}$ of all $\{1, 2, 3\}$ -inverses of a in \mathcal{R}_L is the set $a\{1, 2, 4\}$ of all $\{1, 2, 4\}$ -inverses of a in \mathcal{R} . From

$$a^- \cdot_L a \cdot_L a^\dagger = a^\dagger a a^-$$

we have that $a^{-\dagger}$ is a 1MP-inverse of a in \mathcal{R}_L if and only if it is a MP1-inverse of a in \mathcal{R} , i.e. $z \in a\{-\dagger\}$ in \mathcal{R}_L if and only if $z \in a\{\dagger-\}$ in \mathcal{R} . The following theorem and its corollary are thus a direct corollary (13), Theorem 3.3 and Corollary 3.4.

Theorem 5.1. *Let $a \in \mathcal{R}^\dagger$. Then the following statements are equivalent:*

- (i) $z \in a\{\dagger-\}$;
- (ii) $z \in \mathcal{R}$ is a solution of the system $xax = x$ and $xa = a^\dagger a$;
- (iii) $z \in a\{1, 2, 4\}$.

Corollary 5.2. *Let $a \in \mathcal{R}^\dagger$ and fix $a^{\dagger-} \in a\{\dagger-\}$. Then*

$$a\{\dagger-\} = \left\{ a^{\dagger-} + a^{\dagger-} a w (1 - a a^{\dagger-}) : w \in \mathcal{R} \text{ is arbitrary} \right\}.$$

Remark 5.3. *Let $a \in \mathcal{R}^\dagger$. By Theorems 3.3 and 5.1 we have $a\{-\dagger\} = a\{1, 2, 3\}$ and $a\{\dagger-\} = a\{1, 2, 4\}$. It follows that*

$$a\{-\dagger\} \cap a\{\dagger-\} = \mathcal{R}^\dagger.$$

It is easy to see (compare Definitions 3 and 5) that for every $a \in \mathcal{R}^\dagger$ and $b \in \mathcal{R}$ we have

$$a \leq^{\dagger-} b \quad \text{if and only if} \quad a \leq_L^{-\dagger} b$$

where $\leq_L^{-\dagger}$ is the 1MP-relation in $(\mathcal{R}, +, \cdot_L, *)$. By Theorem 4.3 we may immediately establish that MP1-relation $\leq^{\dagger-}$ introduced with Definition 5 is a partial order on \mathcal{R}^\dagger . We conclude this section with two results that are direct corollaries of (13) and Theorems 4.1 and 4.2, respectively.

Theorem 5.4. *Let $a \in \mathcal{R}^\dagger$ and denote $p = a^\dagger a$ and $q = a a^\dagger$. Let $b \in \mathcal{R}$. Then the following statements are equivalent:*

- (i) $a \leq^{\dagger-} b$;
- (ii)

$$(14) \quad b = \begin{bmatrix} a & -adb_4 \\ 0 & b_4 \end{bmatrix}_{p \times q}$$

where $b_4 \in (1-p)\mathcal{R}(1-q)$ and $d \in q\mathcal{R}(1-p)$.

Theorem 5.5. *Let $a, b \in \mathcal{R}^\dagger$, suppose $a \leq^{\dagger-} b$, and represent b in the matrix form (14). Then $x \in b\{\dagger-\}$ if and only if there exist $x_3 \in q\mathcal{R}(1-p)$ and $x_4 \in (1-q)\mathcal{R}(1-p)$ such that*

$$x = \begin{bmatrix} a^\dagger & x_3 \\ 0 & x_4 \end{bmatrix}_{q \times p}$$

where $x_3 b_4 = d b_4$ and $x_4 \in b_4\{\dagger-\}$.

Let us end this section with a note that we may similarly obtain “the dual version” of Lemma 4.4 and Theorems 4.5–4.7.

6. THE PLUS PARTIAL ORDER IN RICKART *-RINGS

A ring \mathcal{R} is called a *Rickart ring* if for every $a \in \mathcal{R}$ there exist idempotent elements $p, q \in \mathcal{R}$ such that $a^\circ = p \cdot \mathcal{R}$ and ${}^\circ a = \mathcal{R} \cdot q$. A $*$ -ring \mathcal{R} is a *Rickart $*$ -ring* if the left annihilator ${}^\circ a$ of any element $a \in \mathcal{R}$ is generated by a projection $e \in \mathcal{R}$, i.e. ${}^\circ a = \mathcal{R} \cdot e$ where $e = e^* = e^2$. The projection e is unique and every Rickart ring has the (multiplicative) identity 1 (see [2] or [8]). Let us denote

$$LP(a) = \{p \in \mathcal{R} : p = p^2, {}^\circ a = {}^\circ p\} \quad \text{and} \quad RP(a) = \{q \in \mathcal{R} : q = q^2, a^\circ = q^\circ\}.$$

Note that the sets $LP(a)$ and $RP(a)$ are nonempty in case when \mathcal{R} is a Rickart ring. If \mathcal{R} is a Rickart $*$ -ring, there exists the unique projection in $LP(a)$. We denote it by $lp(a)$. Similarly, let $rp(a)$ denote the unique projection in $RP(a)$.

Let \mathcal{H}, \mathcal{K} be Hilbert spaces and let $\mathcal{B}(\mathcal{H}, \mathcal{K})$ be the set of all bounded linear operators from \mathcal{H} to \mathcal{K} . When $\mathcal{H} = \mathcal{K}$, we write $\mathcal{B}(\mathcal{H})$ instead of $\mathcal{B}(\mathcal{H}, \mathcal{H})$. Note that $\mathcal{B}(\mathcal{H})$ is an example of a Rickart $*$ -ring. Let $\text{Ker } A$, $\text{Im } A$, and $\overline{\text{Im } A}$ denote the kernel, the range, and the closure of the range of $A \in \mathcal{B}(\mathcal{H}, \mathcal{K})$, respectively. A new relation was introduced in [1] on $\mathcal{B}(\mathcal{H}, \mathcal{K})$. The definition follows.

Definition 6. *Let $A, B \in \mathcal{B}(\mathcal{H}, \mathcal{K})$. We say that A is below B with respect to the relation \leq^+ and write $A \leq^+ B$ if $\text{Im } A \subseteq \text{Im } B$, $\text{Im } A^* \subseteq \text{Im } B^*$, and there are idempotent operators $\tilde{Q} \in \mathcal{B}(\mathcal{H})$ and $Q \in \mathcal{B}(\mathcal{K})$ such that $A = \tilde{Q}BQ$. In such case, we can without loss of generality assume that $\text{Im } \tilde{Q} = \overline{\text{Im } A}$ and $\text{Im } Q^* = \overline{\text{Im } A^*}$.*

It was proved in [1] that \leq^+ is a partial order on $\mathcal{B}(\mathcal{H}, \mathcal{K})$. Authors of [1] named the relation \leq^+ the plus (partial) order. The plus partial order emerges on $\mathcal{B}(\mathcal{H}, \mathcal{K})$ as a generalization of the minus order and the *diamond order* [3]. It is known that for any von Neumann regular ring \mathcal{R} the definition of the minus (partial) order (12) is equivalent to

$$(15) \quad a = pb = bq$$

where $p, q \in \mathcal{R}$ are some idempotents. It turns out (see [10, Corollary 2.1] and [4, 5]) that the relation \leq^- defined with (15) (i.e. $a \leq^- b$ when $a = pb = bq$ for some $p = p^2$ and $q = q^2$) is a partial order when \mathcal{R} is a Rickart ring.

Let \mathcal{R} be a $*$ -ring. We say that a is below b with respect to the diamond relation \leq^\diamond and write $a \leq^\diamond b$ when

$$(16) \quad {}^\circ b \subseteq {}^\circ a, b^\circ \subseteq a^\circ, \quad \text{and} \quad ab^*a = aa^*a.$$

It turns out (see [9, 12]) that (at least) when \mathcal{R} is $*$ -regular ring with identity, the relation \leq^\diamond is a partial order.

For $A, B \in \mathcal{B}(\mathcal{H})$ we have (see [10, 12])

$$A^\circ \subseteq B^\circ \quad \text{if and only if} \quad \text{Ker } A \subseteq \text{Ker } B \quad \text{if and only if} \quad \overline{\text{Im } B^*} \subseteq \overline{\text{Im } A^*}$$

and

$${}^\circ A \subseteq {}^\circ B \quad \text{if and only if} \quad \overline{\text{Im } B} \subseteq \overline{\text{Im } A}.$$

Motivated by Definition 6 this observation leads us to the following definition.

Definition 7. *Let \mathcal{R} be a Rickart $*$ -ring and let $a, b \in \mathcal{R}$. We say that a is below b with respect to the plus order \leq^+ and write $a \leq^+ b$ if ${}^\circ b \subseteq {}^\circ a$, $b^\circ \subseteq a^\circ$, and there exist $q \in RP(a)$, $\tilde{q} \in LP(a)$ such that $a = \tilde{q}bq$.*

Theorem 6.1. *Let \mathcal{R} be a Rickart $*$ -ring. The relation \leq^+ introduced with Definition 7 is a partial order on \mathcal{R} .*

Proof. Since \mathcal{R} be a Rickart $*$ -ring, there exist $q \in \text{RP}(a)$ and $\tilde{q} \in \text{LP}(a)$. Note that for every idempotent p , $p(1-p) = (1-p)p = 0$. Since $q \in \text{RP}(a), \tilde{q} \in \text{LP}(a)$, we thus have $(1-\tilde{q})a = 0 = a(1-q)$ and hence $a = \tilde{q}a = aq$. So, $a = \tilde{q}aq$ and therefore \leq^+ is reflexive.

To show that \leq^+ is antisymmetric, let $a \leq^+ b$ and $b \leq^+ a$. Then there exist $q \in \text{RP}(a), q_1 \in \text{RP}(b), \tilde{q} \in \text{LP}(a)$, and $\tilde{q}_1 \in \text{LP}(b)$ such that $a = \tilde{q}bq$ and $b = \tilde{q}_1aq_1$. Also, ${}^\circ b = {}^\circ a$ and $b^\circ = a^\circ$, and therefore $(1-\tilde{q})\tilde{q}_1 = 0$ and $q_1(1-q) = 0$. So, $\tilde{q}_1 = \tilde{q}\tilde{q}_1$ and $q_1 = q_1q$. It follows that

$$a = \tilde{q}bq = \tilde{q}\tilde{q}_1aq_1q = \tilde{q}_1aq_1 = b.$$

Let us now prove that \leq^+ is transitive. Let for $a, b, c \in \mathcal{R}$, $a \leq^+ b$ and $b \leq^+ c$. So, ${}^\circ b \subseteq {}^\circ a, b^\circ \subseteq a^\circ, {}^\circ c \subseteq {}^\circ b, c^\circ \subseteq b^\circ$ and therefore ${}^\circ c \subseteq {}^\circ a$ and $c^\circ \subseteq a^\circ$. Also, there exist $q_1 \in \text{RP}(a), q_2 \in \text{RP}(b), \tilde{q}_1 \in \text{LP}(a)$, and $\tilde{q}_2 \in \text{LP}(b)$ with $a = \tilde{q}_1bq_1$ and $b = \tilde{q}_2cq_2$. It follows that $a = \tilde{q}_1\tilde{q}_2cq_2q_1$. Denote $q_3 = q_2q_1$ and $\tilde{q}_3 = \tilde{q}_1\tilde{q}_2$. So,

$$a = \tilde{q}_3cq_3.$$

To conclude the proof let us show that $q_3 \in \text{RP}(a)$ and $\tilde{q}_3 \in \text{LP}(a)$. From ${}^\circ b \subseteq {}^\circ a$ and $b^\circ \subseteq a^\circ$ we get ${}^\circ \tilde{q}_2 \subseteq {}^\circ \tilde{q}_1$ and $q_2^\circ \subseteq q_1^\circ$, and thus $(1-\tilde{q}_2)\tilde{q}_1 = 0$ and $q_1(1-q_2) = 0$. So, $\tilde{q}_1 = \tilde{q}_2\tilde{q}_1$ and $q_1 = q_1q_2$. It follows

$$\tilde{q}_3^2 = \tilde{q}_1(\tilde{q}_2\tilde{q}_1)\tilde{q}_2 = \tilde{q}_1^2\tilde{q}_2 = \tilde{q}_1\tilde{q}_2 = \tilde{q}_3$$

and

$$q_3^2 = q_2(q_1q_2)q_1 = q_2q_1^2 = q_2q_1 = q_3.$$

Observe that from $a = \tilde{q}_3cq_3$ we obtain ${}^\circ \tilde{q}_3 \subseteq {}^\circ a$ and $q_3^\circ \subseteq a^\circ$. Since $\tilde{q}_3 = \tilde{q}_1\tilde{q}_2$ and $q_3 = q_2q_1$, we have ${}^\circ a = {}^\circ \tilde{q}_1 \subseteq {}^\circ \tilde{q}_3$ and $a^\circ = q_1^\circ \subseteq q_3^\circ$. Thus ${}^\circ \tilde{q}_3 = {}^\circ a$ and $q_3^\circ = a^\circ$ and hence $q_3 \in \text{RP}(a)$ and $\tilde{q}_3 \in \text{LP}(a)$. Therefore, $a \leq^+ c$. \square

For the rest of the paper, let \mathcal{R} be a Rickart $*$ -ring. Let $a, b \in \mathcal{R}$ with $a \leq^\diamond b$. Then ${}^\circ b \subseteq {}^\circ a, b^\circ \subseteq a^\circ$, and $ab^*a = aa^*a$. It is easy to prove (see [11, proof of Lemma 5]) that

$$ab^*a = aa^*a \quad \text{if and only if} \quad a = \text{lp}(a)\text{brp}(a).$$

Since $\text{lp}(a) \in \text{LP}(a)$ and $\text{rp}(a) \in \text{RP}(a)$, it follows that $a \leq^+ b$. Let now $a \leq^- b$, i.e. $a = pb = bq$ for some idempotents $p, q \in \mathcal{R}$. Then $a = pbq, {}^\circ b \subseteq {}^\circ a$, and $b^\circ \subseteq a^\circ$. Also, by [10, Corollary 2.1] we may without loss of generality assume that ${}^\circ a = {}^\circ p$ and $a^\circ = q^\circ$, i.e. $p \in \text{LP}(a)$ and $q \in \text{RP}(a)$. It follows again that $a \leq^+ b$. We sum up these observations in the following proposition.

Proposition 6.2. *Let $a, b \in \mathcal{R}$. If $a \leq^\diamond b$, then $a \leq^+ b$, and if $a \leq^- b$, then $a \leq^+ b$.*

We end the paper with a characterization of the plus partial order in Rickart $*$ -rings. First, let us present an auxiliary result which was proved in [4].

Lemma 6.3. *Let $a \in \mathcal{R}$, $p \in \text{LP}(a)$, and $q \in \text{RP}(a)$. Then*

$$\text{LP}(a) = \left\{ \left[\begin{array}{cc} p & p_1 \\ 0 & 0 \end{array} \right]_{p \times p} : p_1 \in p\mathcal{R}(1-p) \right\} \quad \text{and} \quad \text{RP}(a) = \left\{ \left[\begin{array}{cc} q & 0 \\ q_1 & 0 \end{array} \right]_{q \times q} : q_1 \in (1-q)\mathcal{R}q \right\}.$$

Let $a \in \mathcal{R}$. For the purposes of the following characterization of the plus partial order we denote $l_a = \text{lp}(a)$ and $r_a = \text{rp}(a)$. The next theorem extends [1, Theorem 3.11] to the setting of Rickart $*$ -rings.

Theorem 6.4. *Let \mathcal{R} be a Rickart *-ring and let $a, b \in \mathcal{R}$. The following statements are equivalent:*

(i) $a \leq^+ b$;

(ii) $b = \begin{bmatrix} a + y(b_{22}x + w) + zx & yb_{22} + z \\ b_{22}x + w & b_{22} \end{bmatrix}_{l_a \times r_a}$

where $b_{22} \in (1 - l_a)\mathcal{R}(1 - r_a)$ is arbitrary, and $y \in l_a\mathcal{R}(1 - l_a)$, $x \in (1 - r_a)\mathcal{R}r_a$, $w \in (1 - l_a)\mathcal{R}r_a$, and $z \in l_a\mathcal{R}(1 - r_a)$ are such that ${}^\circ b \subseteq {}^\circ((y + 1)w)$ and $b^\circ \subseteq (z(x + 1))^\circ$.

Proof. Let $a, b \in \mathcal{R}$. Suppose $y \in l_a\mathcal{R}(1 - l_a)$, $x \in (1 - r_a)\mathcal{R}r_a$, and

$$\tilde{q} = \begin{bmatrix} l_a & -y \\ 0 & 0 \end{bmatrix}_{l_a \times l_a} \quad \text{and} \quad q = \begin{bmatrix} r_a & 0 \\ -x & 0 \end{bmatrix}_{r_a \times r_a}.$$

By Lemma 6.3, $\tilde{q} \in \text{LP}(a)$ and $q \in \text{RP}(a)$. We have $a = \tilde{q}bq$ if and only if

$$(17) \quad b = a + (1 - \tilde{q})b(1 - q) + (1 - \tilde{q})bq + \tilde{q}b(1 - q).$$

Let $b = \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix}_{l_a \times r_a}$ and denote $w = b_{21} - b_{22}x \in (1 - l_a)\mathcal{R}r_a$ and $z = b_{12} - yb_{22} \in l_a\mathcal{R}(1 - r_a)$. Then

$$\begin{aligned} (1 - \tilde{q})b(1 - q) &= \left(\begin{bmatrix} l_a & 0 \\ 0 & 1 - l_a \end{bmatrix}_{l_a \times l_a} - \begin{bmatrix} l_a & -y \\ 0 & 0 \end{bmatrix}_{l_a \times l_a} \right) \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix}_{l_a \times r_a} \\ &\quad \cdot \left(\begin{bmatrix} r_a & 0 \\ 0 & 1 - r_a \end{bmatrix}_{r_a \times r_a} - \begin{bmatrix} r_a & 0 \\ -x & 0 \end{bmatrix}_{r_a \times r_a} \right) \\ &= \begin{bmatrix} yb_{21} & yb_{22} \\ b_{21} & b_{22} \end{bmatrix}_{l_a \times r_a} \begin{bmatrix} 0 & 0 \\ x & 1 - r_a \end{bmatrix}_{r_a \times r_a} = \begin{bmatrix} yb_{22}x & yb_{22} \\ b_{22}x & b_{22} \end{bmatrix}_{l_a \times r_a}, \end{aligned}$$

$$\begin{aligned} (1 - \tilde{q})bq &= \begin{bmatrix} 0 & y \\ 0 & 1 - l_a \end{bmatrix}_{l_a \times l_a} \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix}_{l_a \times r_a} \begin{bmatrix} r_a & 0 \\ -x & 0 \end{bmatrix}_{r_a \times r_a} \\ &= \begin{bmatrix} yb_{21} & yb_{22} \\ b_{21} & b_{22} \end{bmatrix}_{l_a \times r_a} \begin{bmatrix} r_a & 0 \\ -x & 0 \end{bmatrix}_{r_a \times r_a} \\ &= \begin{bmatrix} y(b_{21} - b_{22}x) & 0 \\ b_{21} - b_{22}x & 0 \end{bmatrix}_{l_a \times r_a} = \begin{bmatrix} yw & 0 \\ w & 0 \end{bmatrix}_{l_a \times r_a}, \end{aligned}$$

and

$$\begin{aligned} \tilde{q}b(1 - q) &= \begin{bmatrix} l_a & -y \\ 0 & 0 \end{bmatrix}_{l_a \times l_a} \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix}_{l_a \times r_a} \begin{bmatrix} 0 & 0 \\ x & 1 - r_a \end{bmatrix}_{r_a \times r_a} \\ &= \begin{bmatrix} b_{11} - yb_{21} & b_{12} - yb_{22} \\ 0 & 0 \end{bmatrix}_{l_a \times r_a} \begin{bmatrix} 0 & 0 \\ x & 1 - r_a \end{bmatrix}_{r_a \times r_a} \\ &= \begin{bmatrix} (b_{12} - yb_{22})x & b_{12} - yb_{22} \\ 0 & 0 \end{bmatrix}_{l_a \times r_a} = \begin{bmatrix} zx & z \\ 0 & 0 \end{bmatrix}_{l_a \times r_a}. \end{aligned}$$

By (17) it follows that $a = \tilde{q}bq$ if and only if

$$\begin{aligned} b &= \begin{bmatrix} a & 0 \\ 0 & 0 \end{bmatrix}_{l_a \times r_a} + \begin{bmatrix} yb_{22}x & yb_{22} \\ b_{22}x & b_{22} \end{bmatrix}_{l_a \times r_a} + \begin{bmatrix} yw & 0 \\ w & 0 \end{bmatrix}_{l_a \times r_a} + \begin{bmatrix} zx & z \\ 0 & 0 \end{bmatrix}_{l_a \times r_a} \\ &= \begin{bmatrix} a + y(b_{22}x + w) + zx & yb_{22} + z \\ b_{22}x + w & b_{22} \end{bmatrix}_{l_a \times r_a}. \end{aligned}$$

From now on we assume that $a = \tilde{q}bq$. To finish the proof, let us show that ${}^\circ b \subseteq {}^\circ a$ if and only if ${}^\circ b \subseteq {}^\circ((y+1)w)$, and $b^\circ \subseteq a^\circ$ if and only if and $b^\circ \subseteq (z(x+1))^\circ$. We will prove only the first equivalence, the second equivalence can be proved similarly. Note first that on the one hand

$$(1 - \tilde{q})bq = \begin{bmatrix} yw & 0 \\ w & 0 \end{bmatrix}_{l_a \times r_a} = yw + w = (y+1)w$$

and on the other hand $(1 - \tilde{q})bq = bq - \tilde{q}bq = bq - a$. Let now ${}^\circ b \subseteq {}^\circ a$ and suppose $tb = 0$ for some $t \in \mathcal{R}$. Thus $ta = 0$ and therefore

$$0 = t(bq - a) = t(1 - \tilde{q})bq = t(y+1)w.$$

So, ${}^\circ b \subseteq {}^\circ((y+1)w)$. Conversely, let ${}^\circ b \subseteq {}^\circ((y+1)w)$ and $tb = 0$. Then $0 = t(y+1)w$ and thus $0 = t(bq - a) = -ta$. So, $ta = 0$ and hence ${}^\circ b \subseteq {}^\circ a$. \square

REFERENCES

- [1] M. L. Arias, A. Maestripieri, On partial orders of operators, preprint, available at [arXiv:2105.09098](https://arxiv.org/abs/2105.09098) [math.FA].
- [2] S. K. Berberian, Baer *-rings, Springer-Verlag, New York, 1972.
- [3] J.K. Baksalary, J. Hauke, A further algebraic version of Cochran's theorem and matrix partial orderings, *Linear Algebra Appl.* **127** (1990), 157-169.
- [4] D. S. Djordjević, D. S. Rakić, J. Marovt, Minus partial order in Rickart rings, *Publ. Math. Debrecen* **87** (2015), No. 3/4, 291-305.
- [5] G. Dolinar, B. Kuzma, J. Marovt, A note on partial orders of Hartwig, Mitsch, and Šemrl, *Appl. Math. Comput.* **270** (2015), 711-713.
- [6] R. E. Hartwig, How to partially order regular elements, *Math. Japon.* **25** (1980), 1-13.
- [7] M. V. Hernández, M. B. Lattanzi, N. Thome, From projectors to IMP and MP1 generalized inverses and their induced partial orders, *Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Math. RACSAM* **115** (2021), No. 3, article 148.
- [8] I. Kaplansky, *Rings of Operators*, Benjamin, New York, 1968.
- [9] L. Lebtahi, P. Patricio, N. Thome, The diamond partial order in rings, *Linear Multilinear Algebra* **62** (No. 3) (2014) 386-395.
- [10] J. Marovt, On partial orders in Rickart rings, *Linear Multilinear Algebra* **63** (2015), No. 9, 1707-1723.
- [11] J. Marovt, K. Mihelič, On sets of elements in Rickart rings induced by partial orders, *Appl. Math. Comput.* **315** (2017), 555-563.
- [12] J. Marovt, D. S. Rakić, D. S. Djordjević, Star, left-star, and right-star partial orders in Rickart *-rings, *Linear Multilinear Algebra* **63** (2015), No. 2, 343-365.
- [13] D. Mosić, D. S. Djordjević, Partial isometries and EP elements in rings with involution, *Electron. J. Linear Algebra* **18** (2009), 761-772.
- [14] D. S. Rakić, Decomposition of a ring induced by minus partial order, *Electron. J. Linear Algebra*, **23** (2012), 1040-1059.

(Janko Marovt) JANKO MAROVT, UNIVERSITY OF MARIBOR, FACULTY OF ECONOMICS AND BUSINESS,
RAZLAGOVA 14, SI-2000 MARIBOR, SLOVENIA, AND IMFM, JADRANSKA 19, SI-1000 LJUBLJANA, SLOVENIA
Email address: `janko.marovt@um.si`

(Dijana Mosić) FACULTY OF SCIENCES AND MATHEMATICS, UNIVERSITY OF NIŠ, P.O. BOX 224, 18000
NIŠ, SERBIA
Email address: `dijana@pmf.ni.ac.rs`

(Insa Cremer) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF LATVIA, JELGAVAS 3, RIGA, 1004, LATVIA
Email address: `insa.kremere@gmail.com`