

Optimal flat functions in Carleman-Roumieu ultraholomorphic classes in sectors

Javier Jiménez-Garrido

Ignacio Miguel-Cantero
Gerhard Schindl

Javier Sanz

May 17, 2022

Abstract

A general procedure is presented in order to obtain linear continuous extension operators, right inverses of the Borel map, whenever optimal flat functions are available in Carleman-Roumieu ultraholomorphic classes, defined on sectors and in terms of regular weight sequences in the sense of Dyn'kin. For a family of regular sequences, including the well-known q -Gevrey case, we construct such optimal flat functions in arbitrary sectors of the Riemann surface of the logarithm.

Key words: Carleman-Roumieu ultraholomorphic classes, asymptotic expansions, linear extension operators.

2020 MSC: Primary 47A57; secondary 44A10, 46E10.

1 Introduction

The asymptotic Borel map sends a function, admitting an asymptotic expansion in a sectorial region, into the formal power series providing such expansion. In many instances it is important to decide about the injectivity and surjectivity of this map when considered between so-called Carleman-Roumieu ultraholomorphic classes and the corresponding class of formal series, defined by restricting the growth of some of the characteristic data of their elements (the derivatives of the functions, the remainders in the expansion, or the coefficients of the series) in terms of a given weight sequence $\mathbb{M} = (M_p)_{p \in \mathbb{N}_0}$ of positive real numbers (see Subsection 2.3 for the definition of such classes). While the injectivity has been fully characterized for sectorial regions and general weight sequences [15, 23, 11], the surjectivity problem is still under study. The classical Borel-Ritt-Gevrey theorem of B. Malgrange and J.-P. Ramis [22], solving the case of Gevrey asymptotics (for which $\mathbb{M} = (p!^\alpha)_{p \in \mathbb{N}_0}$, $\alpha > 0$), was partially extended to different more general situations by J. Schmets and M. Valdivia [25], V. Thilliez [26, 27] and the authors [24, 11, 12]. Summing up, it is known now that the strong nonquasianalyticity condition (snq) for \mathbb{M} , equivalent to the fact that the index $\gamma(\mathbb{M})$ introduced by V. Thilliez is positive (see Subsection 2.2), is indeed necessary for surjectivity. Moreover, for an unbounded sector S_γ of opening $\pi\gamma$ ($\gamma > 0$) in the Riemann surface of the logarithm and for regular weight sequences in the sense of E. M. Dyn'kin [5] (see Subsection 2.1 for the definitions), the Borel map is surjective whenever $\gamma < \gamma(\mathbb{M})$, while it is not for $\gamma > \gamma(\mathbb{M})$ (the situation for $\gamma = \gamma(\mathbb{M})$ is still unclear in general). It is important to note that the proof in this more general situation is not constructive, but rests on the characterization, by abstract functional-analytic techniques, of the surjectivity of the Stieltjes moment mapping in Gelfand-Shilov spaces defined by regular sequences due to A. Debrouwere [4]. This information is transferred into the asymptotic framework in a halfplane

by means of the Fourier transform, and in [12] Laplace and Borel transforms of arbitrary order allow to conclude for general sectors. However, in the particular case of classes given by strongly regular sequences in the sense of V. Thilliez, the proof of surjectivity of the Borel map [27] rests on the construction of optimal flat functions in suitable sectors and a double application of Whitney extension results. Subsequently, A. Lastra, S. Malek and the third author [14] reproved surjectivity in a more explicit way by means of formal Borel- and truncated Laplace-like transforms, defined from suitable kernel functions obtained from those optimal flat functions.

The aim of this paper is to study the connection between the existence of such optimal flat functions in a sector and the surjectivity of the Borel map in it, as well as to provide a constructive technique for such extension results. For strongly regular sequences an equivalence is obtained between these two facts; however, for regular sequences we will get surjectivity whenever optimal flat functions are available, but no general procedure for their construction is currently available. Nonetheless, we will present a family of (non strongly) regular sequences for which such optimal flat functions can be provided in any sector of the Riemann surface of the logarithm, what agrees with the fact that the index $\gamma(\mathbb{M})$ is in this case equal to ∞ . We end by showing how optimal flat functions and extension results can be obtained for convolved sequences, in case the factor sequences admit such constructions separately. Some examples are commented on in regard with this technique.

The paper is organized as follows. Section 2 consists of all the preliminary information concerning weight sequences and some indices or auxiliary functions associated with them, and the main facts about ultraholomorphic classes and the (asymptotic) Borel map defined for them. In Section 3 we define optimal flat functions and carefully detail how their existence entails the surjectivity of the Borel map in ultraholomorphic classes defined by regular sequences. In the particular case of sequences of moderate growth, different statements are presented relating the property of strong non-quasianalyticity to the existence of such flat functions. In Section 4 we give a family of sequences (among which the classical q -Gevrey sequences are found) for which optimal flat functions can be constructed. We need to work first in $\mathbb{C} \setminus (-\infty, 0]$, and then apply a ramification in order to reason for arbitrary sectors in the Riemann surface of the logarithm. Finally, the last section is devoted to the work with convolved sequences.

2 Preliminaries

2.1 Weight sequences and their properties

We set $\mathbb{N} := \{1, 2, \dots\}$, $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$. In what follows, $\mathbb{M} = (M_p)_{p \in \mathbb{N}_0}$ will always stand for a sequence of positive real numbers with $M_0 = 1$. We define its *sequence of quotients* $\mathbf{m} = (m_p)_{p \in \mathbb{N}_0}$ by $m_p := M_{p+1}/M_p$, $p \in \mathbb{N}_0$; the knowledge of \mathbb{M} amounts to that of \mathbf{m} , since $M_p = m_0 \cdots m_{p-1}$, $p \in \mathbb{N}$. The following properties for a sequence will play a role in this paper:

- (i) \mathbb{M} is *logarithmically convex* (for short, (lc)) if $M_p^2 \leq M_{p-1}M_{p+1}$, $p \in \mathbb{N}$.
- (ii) \mathbb{M} is *stable under differential operators* or satisfies the *derivation closedness condition* (briefly, (dc)) if there exists $D > 0$ such that $M_{p+1} \leq D^{p+1}M_p$, $p \in \mathbb{N}_0$.
- (iii) \mathbb{M} is of, or has, *moderate growth* (for the sake of brevity, (mg)) if there exists $A > 0$ such that $M_{p+q} \leq A^{p+q}M_pM_q$, $p, q \in \mathbb{N}_0$.
- (iv) \mathbb{M} satisfies the condition (nq) if

$$\sum_{p=0}^{\infty} \frac{M_p}{(p+1)M_{p+1}} < +\infty.$$

(v) Finally, \mathbb{M} satisfies the condition (snq) if there exists $B > 0$ such that

$$\sum_{q=p}^{\infty} \frac{M_q}{(q+1)M_{q+1}} \leq B \frac{M_p}{M_{p+1}}, \quad p \in \mathbb{N}_0.$$

It is convenient to introduce the notation $\widehat{\mathbb{M}} := (p!M_p)_{p \in \mathbb{N}_0}$. All these properties are preserved when passing from \mathbb{M} to $\widehat{\mathbb{M}}$. In the classical work of H. Komatsu [13], the properties (lc), (dc) and (mg) are denoted by (M.1), (M.2)' and (M.2), respectively, while (snq) for \mathbb{M} is the same as property (M.3) for $\widehat{\mathbb{M}}$. Obviously, (mg) implies (dc).

The sequence of quotients \mathbf{m} is nondecreasing if and only if \mathbb{M} is (lc). In this case, it is well-known that $(M_p)^{1/p} \leq m_{p-1}$ for every $p \in \mathbb{N}$, the sequence $((M_p)^{1/p})_{p \in \mathbb{N}}$ is nondecreasing, and $\lim_{p \rightarrow \infty} (M_p)^{1/p} = \infty$ if and only if $\lim_{p \rightarrow \infty} m_p = \infty$. In order to avoid trivial situations, we will restrict from now on to (lc) sequences \mathbb{M} such that $\lim_{p \rightarrow \infty} m_p = \infty$, which will be called *weight sequences*.

Following E. M. Dyn'kin [5], if \mathbb{M} is a weight sequence and satisfies (dc), we say $\widehat{\mathbb{M}}$ is *regular*. According to V. Thilliez [27], if \mathbb{M} satisfies (lc), (mg) and (snq), we say \mathbb{M} is *strongly regular*; in this case \mathbb{M} is a weight sequence, and the corresponding $\widehat{\mathbb{M}}$ is regular.

We mention some interesting examples. In particular, those in (i) and (iii) appear in the applications of summability theory to the study of formal power series solutions for different kinds of equations.

- (i) The sequences $\mathbb{M}_{\alpha, \beta} := (p!^\alpha \prod_{m=0}^p \log^\beta(e+m))_{p \in \mathbb{N}_0}$, where $\alpha > 0$ and $\beta \in \mathbb{R}$, are strongly regular (in case $\beta < 0$, the first terms of the sequence have to be suitably modified in order to ensure (lc)). In case $\beta = 0$, we have the best known example of a strongly regular sequence, $\mathbb{M}_\alpha := \mathbb{M}_{\alpha, 0} = (p!^\alpha)_{p \in \mathbb{N}_0}$, called the *Gevrey sequence of order α* .
- (ii) The sequence $\mathbb{M}_{0, \beta} := (\prod_{m=0}^p \log^\beta(e+m))_{p \in \mathbb{N}_0}$, with $\beta > 0$, satisfies (lc) and (mg), and \mathbf{m} tends to infinity, but (snq) is not satisfied.
- (iii) For $q > 1$ and $1 < \sigma \leq 2$, $\mathbb{M}_{q, \sigma} := (q^{p^\sigma})_{p \in \mathbb{N}_0}$ satisfies (lc), (dc) and (snq), but not (mg). In case $\sigma = 2$, we get the well-known *q-Gevrey sequence*.

Two sequences $\mathbb{M} = (M_p)_{p \in \mathbb{N}_0}$ and $\mathbb{L} = (L_p)_{p \in \mathbb{N}_0}$ of positive real numbers, with $M_0 = L_0 = 1$ and with respective quotient sequences \mathbf{m} and $\boldsymbol{\ell}$, are said to be *equivalent*, and we write $\mathbb{M} \approx \mathbb{L}$, if there exist positive constants A, B such that $A^p M_p \leq L_p \leq B^p M_p$, $p \in \mathbb{N}_0$. They are said to be *strongly equivalent*, denoted by $\mathbf{m} \simeq \boldsymbol{\ell}$, if there exist positive constants a, b such that $am_p \leq \ell_p \leq bm_p$, $p \in \mathbb{N}_0$.

Whenever $\mathbf{m} \simeq \boldsymbol{\ell}$ we have $\mathbb{M} \approx \mathbb{L}$, but not conversely.

In case M_0 or L_0 is not equal to 1, the previous definitions of equivalence are meant to deal with the normalized sequences $(M_p/M_0)_{p \in \mathbb{N}_0}$ or $(L_p/L_0)_{p \in \mathbb{N}_0}$.

2.2 Index $\gamma(\mathbb{M})$ and auxiliary functions for weight sequences

The index $\gamma(\mathbb{M})$, introduced by V. Thilliez [27, Sect. 1.3] for strongly regular sequences \mathbb{M} , can be equally defined for (lc) sequences, and it may be equivalently expressed by different conditions:

- (i) A sequence $(c_p)_{p \in \mathbb{N}_0}$ is *almost increasing* if there exists $a > 0$ such that for every $p \in \mathbb{N}_0$ we have that $c_p \leq ac_q$ for every $q \geq p$. It was proved in [9, 10] that for any weight sequence \mathbb{M} one has

$$\gamma(\mathbb{M}) = \sup\{\gamma > 0 : (m_p/(p+1)^\gamma)_{p \in \mathbb{N}_0} \text{ is almost increasing}\} \in [0, \infty]. \quad (1)$$

(ii) For any $\beta > 0$ we say that \mathbf{m} satisfies the condition (γ_β) if there exists $A > 0$ such that

$$\sum_{\ell=p}^{\infty} \frac{1}{(m_\ell)^{1/\beta}} \leq \frac{A(p+1)}{(m_p)^{1/\beta}}, \quad p \in \mathbb{N}_0. \quad (\gamma_\beta)$$

In [7, 10] it is proved that for a weight sequence \mathbb{M} ,

$$\gamma(\mathbb{M}) = \sup\{\beta > 0; \mathbf{m} \text{ satisfies } (\gamma_\beta)\}; \quad \gamma(\mathbb{M}) > \beta \iff \mathbf{m} \text{ satisfies } (\gamma_\beta). \quad (2)$$

If we observe that the condition (snq) for \mathbb{M} is precisely (γ_1) for \widehat{m} , the sequence of quotients for $\widehat{\mathbb{M}}$, and that $\gamma(\widehat{\mathbb{M}}) = \gamma(\mathbb{M}) + 1$ (this is clear from (1)), we deduce from the second statement in (2) that

$$\mathbb{M} \text{ satisfies (snq) if, and only if, } \gamma(\mathbb{M}) > 0. \quad (3)$$

Given a weight sequence $\mathbb{M} = (M_p)_{p \in \mathbb{N}_0}$, we write

$$\omega_{\mathbb{M}}(t) := \sup_{p \in \mathbb{N}_0} \ln \left(\frac{t^p}{M_p} \right), \quad t > 0,$$

and $\omega_{\mathbb{M}}(0) = 0$. This is the classical function associated with the sequence \mathbb{M} , see [13].

Another associated function will play a key-role, namely

$$h_{\mathbb{M}}(t) := \inf_{p \in \mathbb{N}_0} M_p t^p, \quad t > 0.$$

The functions $h_{\mathbb{M}}$ and $\omega_{\mathbb{M}}$ are related by

$$h_{\mathbb{M}}(t) = \exp(-\omega_{\mathbb{M}}(1/t)), \quad t > 0. \quad (4)$$

In [13, Prop. 3.2] we find that, for a weight sequence \mathbb{M} ,

$$M_p = \sup_{t>0} t^p \exp(-\omega_{\mathbb{M}}(t)) = \sup_{t>0} t^p h_{\mathbb{M}}(1/t), \quad p \in \mathbb{N}_0. \quad (5)$$

By definition we immediately get the following.

Lemma 2.1. *Let $\mathbb{M} = (M_p)_{p \in \mathbb{N}_0}$ be a weight sequence, then:*

- (i) $t \in (0, \infty) \mapsto h_{\mathbb{M}}(t)$ is nondecreasing and continuous,
- (ii) $h_{\mathbb{M}}(t) \leq 1$ for all $t > 0$, $h_{\mathbb{M}}(t) = 1$ for all t sufficiently large and $\lim_{t \rightarrow 0} h_{\mathbb{M}}(t) = 0$.

Let us also introduce the counting function $\nu_{\mathbf{m}}$ by

$$\nu_{\mathbf{m}}(\lambda) := \#\{p \in \mathbb{N}_0 : m_p \leq \lambda\}. \quad (6)$$

If \mathbb{M} is a weight sequence, then the functions $\nu_{\mathbf{m}}$ and $\omega_{\mathbb{M}}$ are related by the following useful integral representation formula, e.g. see [13, (3.11)] and [15]:

$$\omega_{\mathbb{M}}(x) = \int_0^x \frac{\nu_{\mathbf{m}}(\lambda)}{\lambda} d\lambda = \int_{m_0}^x \frac{\nu_{\mathbf{m}}(\lambda)}{\lambda} d\lambda, \quad x > 0. \quad (7)$$

2.3 Asymptotic expansions, ultraholomorphic classes and the asymptotic Borel map

\mathcal{R} stands for the Riemann surface of the logarithm. $\mathbb{C}[[z]]$ is the space of formal power series in z with complex coefficients.

For $\gamma > 0$, we consider unbounded sectors bisected by direction 0,

$$S_\gamma := \{z \in \mathcal{R} : |\arg(z)| < \frac{\gamma\pi}{2}\}$$

or, in general, unbounded sectors with bisecting direction $d \in \mathbb{R}$ and opening $\gamma\pi$,

$$S(d, \gamma) := \{z \in \mathcal{R} : |\arg(z) - d| < \frac{\gamma\pi}{2}\}.$$

A sector T is said to be a *proper subsector* of a sector S if $\overline{T} \subset S$ (where the closure of T is taken in \mathcal{R} , and so the vertex of the sector is not under consideration).

In this paragraph S is an unbounded sector and \mathbb{M} a sequence. We start by recalling the concept of uniform asymptotic expansion.

We say a holomorphic function $f: S \rightarrow \mathbb{C}$ admits \widehat{f} as its *uniform \mathbb{M} -asymptotic expansion in S* (of type $1/A$ for some $A > 0$) if there exists $C > 0$ such that for every $p \in \mathbb{N}_0$, one has

$$\left| f(z) - \sum_{n=0}^{p-1} a_n z^n \right| \leq C A^p M_p |z|^p, \quad z \in S. \quad (8)$$

In this case we write $f \sim_{\mathbb{M}, A}^u \widehat{f}$ in S , and $\widetilde{\mathcal{A}}_{\mathbb{M}, A}^u(S)$ denotes the space of functions admitting uniform \mathbb{M} -asymptotic expansion of type $1/A$ in S , endowed with the norm

$$\|f\|_{\mathbb{M}, A, \widetilde{u}} := \sup_{z \in S, p \in \mathbb{N}_0} \frac{|f(z) - \sum_{k=0}^{p-1} a_k z^k|}{A^p M_p |z|^p}, \quad (9)$$

which makes it a Banach space. $\widetilde{\mathcal{A}}_{\{\mathbb{M}\}}^u(S)$ stands for the *(LB)* space of functions admitting a uniform $\{\mathbb{M}\}$ -asymptotic expansion in S , obtained as the union of the previous classes when A runs over $(0, \infty)$. When the type needs not be specified, we simply write $f \sim_{\{\mathbb{M}\}}^u \widehat{f}$ in S . Note that, taking $p = 0$ in (8), we deduce that every function in $\widetilde{\mathcal{A}}_{\{\mathbb{M}\}}^u(S)$ is a bounded function.

Finally, we define for every $A > 0$ the class $\mathcal{A}_{\mathbb{M}, A}(S)$ consisting of the holomorphic functions f in S such that

$$\|f\|_{\mathbb{M}, A} := \sup_{z \in S, p \in \mathbb{N}_0} \frac{|f^{(p)}(z)|}{A^p M_p} < \infty.$$

$(\mathcal{A}_{\mathbb{M}, A}(S), \|\cdot\|_{\mathbb{M}, A})$ is a Banach space, and $\mathcal{A}_{\{\mathbb{M}\}}(S) := \cup_{A > 0} \mathcal{A}_{\mathbb{M}, A}(S)$ is called a *Carleman-Roumieu ultraholomorphic class* in the sector S , whose natural inductive topology makes it an *(LB)* space.

We warn the reader that these notations, while the same as in the paper [12], do not agree with the ones used in [24, 11], where $\widetilde{\mathcal{A}}_{\{\mathbb{M}\}}^u(S)$ was denoted by $\widetilde{\mathcal{A}}_{\mathbb{M}}^u(S)$, $\mathcal{A}_{\mathbb{M}, A}(S)$ by $\mathcal{A}_{\mathbb{M}/\mathbb{L}_1, A}(S)$, and $\mathcal{A}_{\{\mathbb{M}\}}(S)$ by $\mathcal{A}_{\mathbb{M}/\mathbb{L}_1}(S)$.

If \mathbb{M} is (lc), the spaces $\mathcal{A}_{\{\mathbb{M}\}}(S)$ and $\widetilde{\mathcal{A}}_{\{\mathbb{M}\}}^u(S)$ are algebras, and if \mathbb{M} is (dc) they are stable under taking derivatives. Moreover, if $\mathbb{M} \approx \mathbb{L}$ the corresponding classes coincide.

Since the derivatives of $f \in \mathcal{A}_{\mathbb{M}, A}(S)$ are Lipschitz, for every $p \in \mathbb{N}_0$ one may define

$$f^{(p)}(0) := \lim_{z \in S, z \rightarrow 0} f^{(p)}(z) \in \mathbb{C}. \quad (10)$$

As a consequence of Taylor's formula and Cauchy's integral formula for the derivatives, there is a close relation between Carleman-Roumieu ultraholomorphic classes and the concept of asymptotic expansion (the proof may be easily adapted from [1]).

Proposition 2.2. *Let \mathbb{M} be a sequence and S be a sector. Then,*

- (i) *If $f \in \mathcal{A}_{\widehat{\mathbb{M}},A}(S)$ then f admits $\widehat{f} := \sum_{p \in \mathbb{N}_0} \frac{1}{p!} f^{(p)}(0) z^p$ as its uniform \mathbb{M} -asymptotic expansion in S of type $1/A$, where $(f^{(p)}(0))_{p \in \mathbb{N}_0}$ is given by (10). Moreover, $\|f\|_{\mathbb{M},A,\widetilde{u}} \leq \|f\|_{\widehat{\mathbb{M}},A}$, and so the identity map $\mathcal{A}_{\widehat{\mathbb{M}},A}(S) \hookrightarrow \widetilde{\mathcal{A}}_{\mathbb{M},A}^u(S)$ is continuous. Consequently, we also have that $\mathcal{A}_{\{\widehat{\mathbb{M}}\}}(S) \subseteq \widetilde{\mathcal{A}}_{\{\mathbb{M}\}}^u(S)$ and $\mathcal{A}_{\{\widehat{\mathbb{M}}\}}(S) \hookrightarrow \widetilde{\mathcal{A}}_{\{\mathbb{M}\}}^u(S)$ is continuous.*
- (ii) *If S is unbounded and T is a proper subsector of S , then there exists a constant $c = c(T, S) > 0$ such that the restriction to T , $f|_T$, of functions f defined on S and admitting a uniform \mathbb{M} -asymptotic expansion in S of type $1/A > 0$, belongs to $\mathcal{A}_{\widehat{\mathbb{M}},cA}(T)$, and $\|f|_T\|_{\widehat{\mathbb{M}},cA} \leq \|f\|_{\mathbb{M},A,\widetilde{u}}$. So, the restriction map from $\widetilde{\mathcal{A}}_{\mathbb{M},A}^u(S)$ to $\mathcal{A}_{\widehat{\mathbb{M}},cA}(T)$ is continuous, and it is also continuous from $\widetilde{\mathcal{A}}_{\{\mathbb{M}\}}^u(S)$ to $\mathcal{A}_{\{\widehat{\mathbb{M}}\}}(T)$.*

One may accordingly define classes of formal power series

$$\mathbb{C}[[z]]_{\mathbb{M},A} = \left\{ \widehat{f} = \sum_{p=0}^{\infty} a_p z^p \in \mathbb{C}[[z]] : \|\widehat{f}\|_{\mathbb{M},A} := \sup_{p \in \mathbb{N}_0} \frac{|a_p|}{A^p M_p} < \infty \right\}. \quad (11)$$

$(\mathbb{C}[[z]]_{\mathbb{M},A}, |\cdot|_{\mathbb{M},A})$ is a Banach space and we put $\mathbb{C}[[z]]_{\{\mathbb{M}\}} := \cup_{A>0} \mathbb{C}[[z]]_{\mathbb{M},A}$, again an (LB) space.

It is natural to consider the *asymptotic Borel map* $\widetilde{\mathcal{B}}$ sending a function $f \in \widetilde{\mathcal{A}}_{\mathbb{M},A}^u(S)$ into its \mathbb{M} -asymptotic expansion $\widehat{f} \in \mathbb{C}[[z]]_{\mathbb{M},A}$. By Proposition 2.2.(i) the asymptotic Borel map may be defined from $\widetilde{\mathcal{A}}_{\{\mathbb{M}\}}^u(S)$ or $\mathcal{A}_{\{\widehat{\mathbb{M}}\}}(S)$ into $\mathbb{C}[[z]]_{\{\mathbb{M}\}}$, and from $\mathcal{A}_{\widehat{\mathbb{M}},A}(S)$ into $\mathbb{C}[[z]]_{\mathbb{M},A}$.

If \mathbb{M} is (lc), $\widetilde{\mathcal{B}}$ is a homomorphism of algebras; if \mathbb{M} is also (dc), differentiation commutes with $\widetilde{\mathcal{B}}$. Moreover, it is continuous when considered between the corresponding Banach or (LB) spaces previously introduced. Finally, note that if $\mathbb{M} \approx \mathbb{L}$, then $\mathbb{C}[[z]]_{\{\mathbb{M}\}} = \mathbb{C}[[z]]_{\{\mathbb{L}\}}$, and the corresponding Borel maps are in all cases identical.

Since the problem under study is invariant under rotation, we will focus on the surjectivity of the Borel map in unbounded sectors S_γ . So, we define

$$\begin{aligned} S_{\{\widehat{\mathbb{M}}\}} &:= \{\gamma > 0; \quad \widetilde{\mathcal{B}} : \mathcal{A}_{\{\widehat{\mathbb{M}}\}}(S_\gamma) \longrightarrow \mathbb{C}[[z]]_{\{\mathbb{M}\}} \text{ is surjective}\}, \\ \widetilde{S}_{\{\mathbb{M}\}}^u &:= \{\gamma > 0; \quad \widetilde{\mathcal{B}} : \widetilde{\mathcal{A}}_{\{\mathbb{M}\}}^u(S_\gamma) \longrightarrow \mathbb{C}[[z]]_{\{\mathbb{M}\}} \text{ is surjective}\}. \end{aligned}$$

We again note that these intervals were respectively denoted by $S_{\mathbb{M}}$ and $\widetilde{S}_{\mathbb{M}}^u$ in [11].

It is clear that $S_{\{\widehat{\mathbb{M}}\}}$ and $\widetilde{S}_{\{\mathbb{M}\}}^u$ are either empty or left-open intervals having 0 as endpoint, called *surjectivity intervals*. Using Proposition 2.2, we easily see that

$$(\widetilde{S}_{\{\mathbb{M}\}}^u)^\circ \subseteq S_{\{\widehat{\mathbb{M}}\}} \subseteq \widetilde{S}_{\{\mathbb{M}\}}^u, \quad (12)$$

where I° stands for the interior of the interval I .

3 Optimal flat functions in ultraholomorphic classes

The following result appeared, in a slightly different form, in [11, Lemma 4.5]

Lemma 3.1. *Let \mathbb{M} be a weight sequence. If $\tilde{S}_{\{\mathbb{M}\}}^u \neq \emptyset$, then \mathbb{M} satisfies (snq) or, equivalently, $\gamma(\mathbb{M}) > 0$.*

Subsequently, a converse, more precise statement appeared in [12, Th. 3.7] under the additional hypothesis of condition (dc).

Theorem 3.2. *Let $\widehat{\mathbb{M}}$ be a regular sequence such that $\gamma(\mathbb{M}) > 0$. Then,*

$$(0, \gamma(\mathbb{M})) \subseteq S_{\{\widehat{\mathbb{M}}\}} \subseteq \tilde{S}_{\{\mathbb{M}\}}^u \subseteq (0, \gamma(\mathbb{M})).$$

In particular, if $\gamma(\mathbb{M}) = \infty$, we have that $S_{\{\widehat{\mathbb{M}}\}} = \tilde{S}_{\{\mathbb{M}\}}^u = (0, \infty)$.

So, the surjectivity of the Borel map for regular sequences is governed by the value of the index $\gamma(\mathbb{M})$. Our aim is to relate surjectivity in a sector to the existence of optimal flat functions in it, which we now define.

Definition 3.3. Let \mathbb{M} be a weight sequence, S an unbounded sector bisected by direction $d = 0$, i.e., by the positive real line $(0, +\infty) \subset \mathcal{R}$. A holomorphic function $G: S \rightarrow \mathbb{C}$ is called an *optimal $\{\mathbb{M}\}$ -flat function* in S if:

- (i) $G(x) > 0$ for all $x > 0$,
- (ii) There exist $K_1, K_2, K_3, K_4 > 0$ such that for all $z \in S$, one has

$$K_1 h_{\mathbb{M}}(K_2 |z|) \leq |G(z)| \leq K_3 h_{\mathbb{M}}(K_4 |z|). \quad (13)$$

Besides the symmetry imposed by condition (i), we note that the estimates in the right-hand inequality from (13) amount to the fact that $|G(z)| \leq K_3 K_4^p M_p |z|^p$ for every $p \in \mathbb{N}_0$ and every $z \in S$, what exactly means that $G \in \tilde{\mathcal{A}}_{\{\mathbb{M}\}}^u(S)$ and is $\{\mathbb{M}\}$ -flat, i.e., its uniform $\{\mathbb{M}\}$ -asymptotic expansion is given by the null series. The left-hand inequality imposed in (13) makes the function optimal in a sense, as its rate of decrease at 0 is accurately specified by the function $h_{\mathbb{M}}$.

If G is an optimal $\{\mathbb{M}\}$ -flat function in $\tilde{\mathcal{A}}_{\{\mathbb{M}\}}^u(S)$, the kernel function $e: S \rightarrow \mathbb{C}$ given by

$$e(z) := G\left(\frac{1}{z}\right), \quad z \in S,$$

is such that $e(x) > 0$ for all $x > 0$, and there exist $K_1, K_2, K_3, K_4 > 0$ such that

$$K_1 h_{\mathbb{M}}\left(\frac{K_2}{|z|}\right) \leq |e(z)| \leq K_3 h_{\mathbb{M}}\left(\frac{K_4}{|z|}\right), \quad z \in S. \quad (14)$$

For every $p \in \mathbb{N}_0$ we define the p -th moment of the function $e(z)$, given by

$$\mathbf{m}(p) := \int_0^\infty t^p e(t) dt.$$

Note that the positive value $\mathbf{m}(0)$ need not be equal to 1.

The following result is crucial for our aim.

Proposition 3.4. *Suppose $\widehat{\mathbb{M}}$ is a regular sequence and G is an optimal $\{\mathbb{M}\}$ -flat function in $\widetilde{\mathcal{A}}_{\{\mathbb{M}\}}^u(S)$. Consider the sequence of moments $\mathbf{m} := (\mathbf{m}(p))_{p \in \mathbb{N}_0}$ associated with the kernel function $e(z) = G(1/z)$. Then, there exist $B_1, B_2 > 0$ such that*

$$\mathbf{m}(0)B_1^p M_p \leq \mathbf{m}(p) \leq \mathbf{m}(0)B_2^p M_p, \quad p \in \mathbb{N}_0. \quad (15)$$

In other words, \mathbb{M} and \mathbf{m} are equivalent.

Proof. On the one hand, because of the right-hand inequalities in (14) and Lemma 2.1.(ii), for every $p \in \mathbb{N}_0$ and $s > 0$ we may write

$$\begin{aligned} \mathbf{m}(p) &= \int_0^s t^p e(t) dt + \int_s^\infty \frac{1}{t^2} t^{p+2} e(t) dt \\ &\leq K_3 \int_0^s t^p dt + K_3 \sup_{t>0} t^{p+2} h_{\mathbb{M}} \left(\frac{K_4}{t} \right) \int_s^\infty \frac{1}{t^2} dt \\ &= K_3 \frac{s^{p+1}}{p+1} + K_3 \frac{1}{s} K_4^{p+2} M_{p+2} \leq K_3 \left(\frac{s^{p+1}}{p+1} + \frac{(K_4 D)^{p+2} M_p}{s} \right). \end{aligned}$$

Note that in the last equality we have used (5), and then we have applied (dc) with a suitable constant $D > 0$. Since $s > 0$ was arbitrary, we finally get

$$\mathbf{m}(p) \leq \inf_{s>0} K_3 \left(\frac{s^{p+1}}{p+1} + \frac{(K_4 D)^{p+2} M_p}{s} \right) = K_3 \frac{p+2}{p+1} (K_4 D)^{p+1} (M_p)^{(p+1)/(p+2)} \leq \mathbf{m}(0) B_2^p M_p$$

for a suitably enlarged constant $B_2 > 0$ (observe that, eventually, $M_p \geq 1$).

On the other hand, by the left-hand inequalities in (14) and Lemma 2.1.(i), for every $p \in \mathbb{N}_0$ and $s > 0$ we may estimate

$$\mathbf{m}(p) \geq \int_0^s t^p e(t) dt \geq K_1 \int_0^s t^p h_{\mathbb{M}} \left(\frac{K_2}{t} \right) dt \geq K_1 h_{\mathbb{M}} \left(\frac{K_2}{s} \right) \frac{s^{p+1}}{p+1}.$$

Then, again by (5), we deduce that

$$\mathbf{m}(p) \geq \frac{K_1}{p+1} \sup_{s>0} h_{\mathbb{M}} \left(\frac{K_2}{s} \right) s^{p+1} = \frac{K_1}{p+1} K_2^{p+1} M_{p+1} \geq \mathbf{m}(0) B_1^p M_p$$

for a suitable constant $B_1 > 0$ (note that \mathbb{M} is eventually nondecreasing). \square

We can already state the following main result. The forthcoming implication (i) \Rightarrow (iv) for strongly regular sequences \mathbb{M} was first obtained by V. Thilliez [27, Th. 3.2.1], and the proof heavily rested on the moderate growth condition for \mathbb{M} because of the use of Whitney extension results in the ultradifferentiable setting. In [14] the implication (i) \Rightarrow (ii) was proved again for strongly regular sequences, but with a completely different technique, and it is this approach which allows here for the weakening of condition (mg) into (dc).

Theorem 3.5. *Let $\widehat{\mathbb{M}}$ be a regular sequence and $\gamma > 0$. Each of the following statements implies the next one:*

- (i) *There exists an optimal $\{\mathbb{M}\}$ -flat function in $\widetilde{\mathcal{A}}_{\{\mathbb{M}\}}^u(S_\gamma)$.*
- (ii) *There exists $c > 0$ such that for every $A > 0$ there exists a linear continuous map $T_{\mathbb{M},A}: \mathbb{C}[[z]]_{\mathbb{M},A} \rightarrow \widetilde{\mathcal{A}}_{\mathbb{M},cA}^u(S_\gamma)$ such that $\widetilde{\mathcal{B}} \circ T_{\mathbb{M},A}$ is the identity map in $\mathbb{C}[[z]]_{\mathbb{M},A}$ (i.e., $T_{\mathbb{M},A}$ is an extension operator, right inverse for $\widetilde{\mathcal{B}}$).*

(iii) The Borel map $\tilde{\mathcal{B}}: \tilde{\mathcal{A}}_{\{\mathbb{M}\}}^u(S_\gamma) \rightarrow \mathbb{C}[[z]]_{\{\mathbb{M}\}}$ is surjective. In other words, $(0, \gamma] \subset \tilde{S}_{\{\mathbb{M}\}}^u$.

(iv) $(0, \gamma) \subset S_{\{\widehat{\mathbb{M}}\}}$.

(v) $\gamma \leq \gamma(\mathbb{M})$.

Proof. (i) \Rightarrow (ii) Let $A > 0$ and $\hat{f} = \sum_{p=0}^{\infty} a_p z^p \in \mathbb{C}[[z]]_{\mathbb{M}, A}$ be given. Let $(\mathbf{m}(p))_{p \in \mathbb{N}_0}$ be the sequence of moments associated to the function $e(z) = G(1/z)$, where G is an optimal $\{\mathbb{M}\}$ -flat function in $\tilde{\mathcal{A}}_{\{\mathbb{M}\}}^u(S_\gamma)$. By the definition of the norm in $\mathbb{C}[[z]]_{\mathbb{M}, A}$ (see (11)), we have

$$|a_p| \leq |\hat{f}|_{\mathbb{M}, A} A^p M_p, \quad p \in \mathbb{N}_0.$$

From the left-hand inequalities in (15), we deduce that

$$\left| \frac{a_p}{\mathbf{m}(p)} \right| \leq \frac{|\hat{f}|_{\mathbb{M}, A}}{\mathbf{m}(0)} \left(\frac{A}{B_1} \right)^p, \quad p \in \mathbb{N}_0. \quad (16)$$

Hence, the formal Borel-like transform of \hat{f} ,

$$\hat{g} = \sum_{p=0}^{\infty} \frac{a_p}{\mathbf{m}(p)} z^p,$$

is convergent in the disc $D(0, R)$ for $R = B_1/A > 0$, and it defines a holomorphic function g there. Choose $R_0 := B_1/(2A) < R$, and define

$$T_{\mathbb{M}, A}(\hat{f})(z) := \frac{1}{z} \int_0^{R_0} e\left(\frac{u}{z}\right) g(u) du, \quad z \in S_\gamma,$$

which is a truncated Laplace-like transform of g with kernel e . By virtue of Leibniz's theorem for parametric integrals and the properties of e , we deduce that this function, denoted by f for the sake of brevity, is holomorphic in S_γ . We will prove that $f \sim_{\{\mathbb{M}\}}^u \hat{f}$ uniformly in S_γ , and that the map $\hat{f} \mapsto f$, which is obviously linear, is also continuous from $\mathbb{C}[[z]]_{\mathbb{M}, A}$ into $\tilde{\mathcal{A}}_{\mathbb{M}, cA}^u(S_\gamma)$ for suitable $c > 0$ independent from A .

Let $p \in \mathbb{N}_0$ and $z \in S_\gamma$. We have

$$\begin{aligned} f(z) - \sum_{n=0}^{p-1} a_n z^n &= f(z) - \sum_{n=0}^{p-1} \frac{a_n}{\mathbf{m}(n)} \mathbf{m}(n) z^n \\ &= \frac{1}{z} \int_0^{R_0} e\left(\frac{u}{z}\right) \sum_{n=0}^{\infty} \frac{a_n}{\mathbf{m}(n)} u^n du - \sum_{n=0}^{p-1} \frac{a_n}{\mathbf{m}(n)} \int_0^{\infty} v^n e(v) dv z^n. \end{aligned}$$

After a change of variable $u = zv$ in the last integral, one may use Cauchy's residue theorem and the right-hand estimates in (14) in order to rotate the path of integration and obtain

$$z^n \int_0^{\infty} v^n e(v) dv = \frac{1}{z} \int_0^{\infty} u^n e\left(\frac{u}{z}\right) du.$$

So, we can write the preceding difference as

$$\frac{1}{z} \left(\int_0^{R_0} e\left(\frac{u}{z}\right) \sum_{n=p}^{\infty} \frac{a_n}{\mathbf{m}(n)} u^n du - \int_{R_0}^{\infty} e\left(\frac{u}{z}\right) \sum_{n=0}^{p-1} \frac{a_n}{\mathbf{m}(n)} u^n du \right).$$

Then, we have

$$\left| f(z) - \sum_{n=0}^{p-1} a_n z^n \right| \leq \frac{1}{|z|} (f_1(z) + f_2(z)), \quad (17)$$

where

$$f_1(z) = \left| \int_0^{R_0} e\left(\frac{u}{z}\right) \sum_{n=p}^{\infty} \frac{a_n}{\mathfrak{m}(n)} u^n du \right|, \quad f_2(z) = \left| \int_{R_0}^{\infty} e\left(\frac{u}{z}\right) \sum_{n=0}^{p-1} \frac{a_n}{\mathfrak{m}(n)} u^n du \right|.$$

We now estimate $f_1(z)$ and $f_2(z)$. Observe that for every $u \in (0, R_0]$ we have $0 < Au/B_1 \leq 1/2$. So, from (16) we get

$$\sum_{n=p}^{\infty} \frac{|a_n|}{\mathfrak{m}(n)} u^n \leq \frac{|\widehat{f}|_{\mathbb{M},A}}{\mathfrak{m}(0)} \sum_{n=p}^{\infty} \left(\frac{Au}{B_1}\right)^n \leq \frac{2|\widehat{f}|_{\mathbb{M},A}}{\mathfrak{m}(0)} \left(\frac{A}{B_1}\right)^p u^p.$$

Hence,

$$f_1(z) \leq \frac{2|\widehat{f}|_{\mathbb{M},A}}{\mathfrak{m}(0)} \left(\frac{A}{B_1}\right)^p \int_0^{R_0} \left| e\left(\frac{u}{z}\right) \right| u^p du. \quad (18)$$

Regarding $f_2(z)$, for $u \geq R_0$ and $0 \leq n \leq p-1$ we have $(u/R_0)^n \leq (u/R_0)^p$, so $u^n \leq R_0^n u^p / R_0^p$. Again by (16), and taking into account the value of R_0 , we may write

$$\sum_{n=0}^{p-1} \frac{|a_n|}{\mathfrak{m}(n)} u^n \leq \frac{|\widehat{f}|_{\mathbb{M},A}}{\mathfrak{m}(0)} \frac{u^p}{R_0^p} \sum_{n=0}^{p-1} \left(\frac{AR_0}{B_1}\right)^n \leq \frac{|\widehat{f}|_{\mathbb{M},A}}{\mathfrak{m}(0)} \left(\frac{2A}{B_1}\right)^p u^p.$$

Then, we get

$$f_2(z) \leq \frac{|\widehat{f}|_{\mathbb{M},A}}{\mathfrak{m}(0)} \left(\frac{2A}{B_1}\right)^p \int_{R_0}^{\infty} \left| e\left(\frac{u}{z}\right) \right| u^p du. \quad (19)$$

In order to conclude, note that the second inequality in (14), followed by the first one, and the fact that $e(x) > 0$ for $x > 0$, together imply that for every $z \in S_\gamma$ and every $u > 0$ we have

$$|e(u/z)| \leq K_3 h_{\mathbb{M}} \left(K_4 \frac{|z|}{u} \right) \leq \frac{K_3}{K_1} e \left(\frac{K_2 u}{K_4 |z|} \right).$$

We use this fact, a simple change of variable and the right-hand estimates in (15), and obtain that

$$\begin{aligned} \int_0^{\infty} \left| e\left(\frac{u}{z}\right) \right| u^p du &\leq \int_0^{\infty} \frac{K_3}{K_1} e \left(\frac{K_2 u}{K_4 |z|} \right) u^p du \\ &= \frac{K_3}{K_1} \left(\frac{K_4 |z|}{K_2} \right)^{p+1} \mathfrak{m}(p) \leq \frac{\mathfrak{m}(0) K_3 K_4}{K_1 K_2} \left(\frac{K_4 B_2}{K_2} \right)^p M_p |z|^{p+1}. \end{aligned}$$

This estimate can be taken into both (18) and (19), and from (17) we easily get that for every $p \in \mathbb{N}_0$,

$$\left| f(z) - \sum_{n=0}^{p-1} a_n z^n \right| \leq \frac{3K_3 K_4}{K_1 K_2} |\widehat{f}|_{\mathbb{M},A} \left(\frac{2K_4 B_2 A}{K_2 B_1} \right)^p M_p |z|^p, \quad z \in S_\gamma,$$

and so f admits \widehat{f} as its uniform $\{\mathbb{M}\}$ -asymptotic expansion in S_γ . Moreover, recalling the definition (9) of the norm in these spaces with uniform asymptotics and fixed type, if we put $c := 2K_4 B_2 / (K_2 B_1) > 0$, we see that $f \in \widetilde{\mathcal{A}}_{\mathbb{M},cA}^u(S_\gamma)$ and

$$\|f\|_{\mathbb{M},cA,\widetilde{u}} \leq \frac{3K_3 K_4}{K_1 K_2} |\widehat{f}|_{\mathbb{M},A},$$

what proves the continuity of the linear map $T_{\mathbb{M},A}$.

(ii) \Rightarrow (iii) Immediate for any weight sequence \mathbb{M} .

(iii) \Rightarrow (iv) It follows from (12), again valid for any weight sequence.

(iv) \Rightarrow (v) This statement is a consequence of Theorem 3.2. \square

Remark 3.6. The crucial step taken by V. Thilliez [27, Th. 2.3.1] for strongly regular sequences was the explicit construction of optimal $\{\mathbb{M}\}$ -flat functions in sectors S_γ for every $\gamma > 0$ such that $\gamma < \gamma(\mathbb{M})$, what almost closes the circle of implications in Theorem 3.5.

The facts in Theorem 3.5.(ii) and Proposition 2.2.(ii) together guarantee that for every $\delta \in (0, \gamma)$ there exists $c' > 0$ such that for every $A > 0$ there exists a linear and continuous extension operator from $\mathbb{C}[[z]]_{\mathbb{M},A}$ into $\mathcal{A}_{\widehat{\mathbb{M}},c'A}(S_\delta)$. In fact, V. Thilliez stated his main result in this regard [27, Th. 3.2.1] in terms of the existence of such extension operators for every $\delta < \gamma(\mathbb{M})$.

The following three corollaries become now clear.

Corollary 3.7. *Let \mathbb{M} be a weight sequence satisfying (mg), and $\gamma > 0$. The following are equivalent:*

(i) $\gamma(\mathbb{M}) > \gamma$,

(ii) *There exists $\gamma_1 > \gamma$ such that the space $\widetilde{\mathcal{A}}_{\{\mathbb{M}\}}^u(S_{\gamma_1})$ contains optimal $\{\mathbb{M}\}$ -flat functions.*

(iii) *There exists $\gamma_1 > \gamma$ such that the Borel map $\widetilde{\mathcal{B}} : \widetilde{\mathcal{A}}_{\{\mathbb{M}\}}^u(S_{\gamma_1}) \rightarrow \mathbb{C}[[z]]_{\{\mathbb{M}\}}$ is surjective., i.e., $\gamma_1 \in \widetilde{S}_{\{\mathbb{M}\}}^u$.*

Proof. (ii) \Rightarrow (iii) and (iii) \Rightarrow (i) are respectively contained in Theorem 3.5 and Theorem 3.2, under weaker hypotheses. (i) \Rightarrow (ii) follows from [27, Th. 2.3.1] and Proposition 2.2. \square

As a consequence of (3) we deduce the following result.

Corollary 3.8. *Let \mathbb{M} be a weight sequence satisfying (mg). The following are equivalent:*

(i) \mathbb{M} is strongly regular.

(ii) *There exists $\gamma > 0$ such that the space $\widetilde{\mathcal{A}}_{\{\mathbb{M}\}}^u(S_\gamma)$ contains optimal $\{\mathbb{M}\}$ -flat functions.*

(iii) *There exists $\gamma > 0$ such that the Borel map $\widetilde{\mathcal{B}} : \widetilde{\mathcal{A}}_{\{\mathbb{M}\}}^u(S_\gamma) \rightarrow \mathbb{C}[[z]]_{\{\mathbb{M}\}}$ is surjective. In other words, $\widetilde{S}_{\{\mathbb{M}\}}^u \neq \emptyset$.*

Note that, according to Proposition 2.2, in the previous items (ii) and (iii) one could change $\widetilde{\mathcal{A}}_{\{\mathbb{M}\}}^u(S_\gamma)$ and $\widetilde{S}_{\{\mathbb{M}\}}^u$ into $\mathcal{A}_{\{\widehat{\mathbb{M}}\}}(S_\gamma)$ and $S_{\{\widehat{\mathbb{M}}\}}$, respectively.

Corollary 3.9. *Let \mathbb{M} be a weight sequence satisfying (mg), and $\gamma > 0$. The following are equivalent:*

(i) $\gamma(\mathbb{M}) > \gamma$,

(ii) *There exists $\gamma_1 > \gamma$ such that the space $\mathcal{A}_{\{\widehat{\mathbb{M}}\}}(S_{\gamma_1})$ contains optimal $\{\mathbb{M}\}$ -flat functions,*

(iii) *There exists $\gamma_1 > \gamma$ such that $\widetilde{\mathcal{B}} : \mathcal{A}_{\{\widehat{\mathbb{M}}\}}(S_{\gamma_1}) \rightarrow \mathbb{C}[[z]]_{\{\mathbb{M}\}}$ is surjective, i.e., $\gamma_1 \in S_{\{\widehat{\mathbb{M}}\}}$.*

The implication (ii) \Rightarrow (i) in the version of Corollary 3.8 for the space $\mathcal{A}_{\{\widehat{\mathbb{M}}\}}(S_\gamma)$ can be shown independently by using a result from J. Bruna [3], where a precise formula for nontrivial flat functions in Carleman-Roumieu ultradifferentiable classes, appearing in a work of T. Bang [2], is exploited. For the sake of completeness, we will present this proof below.

Theorem 3.10. *Let \mathbb{M} be a weight sequence satisfying (mg). If there exists $\gamma > 0$ such that $\mathcal{A}_{\{\widehat{\mathbb{M}}\}}(S_\gamma)$ contains optimal $\{\mathbb{M}\}$ -flat functions, then \mathbb{M} is strongly regular.*

The proof requires two auxiliary results which we state and prove now.

First, given a weight sequence \mathbb{M} , the sequence of quotients $\mathbf{m} = (m_p)_{p \in \mathbb{N}_0}$ is nondecreasing and tends to infinity, but it can happen that it remains constant on large intervals $[p_0, p_1]$ of indices, so that the counting function $\nu_{\mathbf{m}}$ defined in (6) yields $\nu_{\mathbf{m}}(m_{p_0}) = \nu_{\mathbf{m}}(m_{p_1}) = p_1 + 1$. However, in some applications or proofs it would be convenient to have $\nu_{\mathbf{m}}(m_p) = p + 1$ for all $p \geq 0$. This can be assumed without loss of generality by the following result.

Lemma 3.11. *Let $a = (a_p)_{p \geq 1}$ be a nondecreasing sequence of positive real numbers satisfying $\lim_{p \rightarrow +\infty} a_p = +\infty$ (it suffices that $a_{p-1} < a_p$ holds true for infinitely many indices p). Then there exists a sequence $b = (b_p)_{p \geq 1}$ of positive real numbers such that $p \mapsto b_p$ is strictly increasing and satisfies*

$$0 < \inf_{p \geq 1} \frac{b_p}{a_p} \leq \sup_{p \geq 1} \frac{b_p}{a_p} < +\infty.$$

So, in the language of weight sequences, we prove that for any weight sequence \mathbb{M} there exists a strongly equivalent weight sequence \mathbb{L} (and so $\mathbb{M} \approx \mathbb{L}$) such that $\nu_{\ell}(\ell_p) = p + 1$ for all $p \in \mathbb{N}_0$. Note that equivalent weight sequences define the same Carleman-Roumieu ultraholomorphic classes and associated classes of sequences.

Proof. Since a is nondecreasing and $\lim_{p \rightarrow +\infty} a_p = +\infty$ there exists a sequence $(p_j)_{j \geq 1}$ of indices such that $a_{p_{j-1}} < a_{p_j} = \dots = a_{p_{j+1}-1} < a_{p_{j+1}}$ for all $j \geq 1$ (and so $p_1 \geq 2$). For all $j \geq 1$ we have now $a_{p_j}/(a_{p_{j-1}}) > 1 + \varepsilon_j$ for a sequence $(\varepsilon_j)_{j \geq 1}$ with possibly small strictly positive numbers ε_j . Finally we put $p_0 := 1$.

We take some arbitrary $A > 1$ and choose $\delta_j > 0$ small enough so as to have $(1 + \delta_j)^{p_{j+1} - p_j - 1} \leq \min\{A, 1 + \varepsilon_{j+1}\}$. Then the sequence $(\delta_j)_{j \geq 0}$ satisfies

$$(1 + \delta_j)^{p_{j+1} - p_j - 1} \leq 1 + \varepsilon_{j+1} < \frac{a_{p_{j+1}}}{a_{p_{j+1}-1}}, \quad (1 + \delta_j)^{p_{j+1} - p_j - 1} \leq A, \quad j \geq 0. \quad (20)$$

We define now b as follows:

$$b_q := a_q \text{ if } q = p_j, j \geq 0, \quad b_q := (1 + \delta_j)b_{q-1} \text{ if } 1 + p_j \leq q \leq p_{j+1} - 1, j \geq 0. \quad (21)$$

So we have by iteration $b_q = (1 + \delta_j)^{q - p_j} b_{p_j} = (1 + \delta_j)^{q - p_j} a_{p_j} = (1 + \delta_j)^{q - p_j} a_q > a_q$ for all q with $1 + p_j \leq q \leq p_{j+1} - 1$, $j \geq 0$. On each such interval of indices the mapping $q \mapsto b_q$ is now clearly strictly increasing since $1 + \delta_j > 1$ for all j . Moreover, by the first half in (20), we have $b_{p_{j+1}-1} = (1 + \delta_j)^{p_{j+1} - p_j - 1} a_{p_j} < b_{p_{j+1}}$. Hence the sequence $q \mapsto b_q$ is strictly increasing.

By definition (21) we have $b_q = a_q$ for all $q = p_j$, $j \geq 0$, and $b_q > a_q$ otherwise. We conclude if we show that $b_q \leq A a_q$ for all q with $1 + p_j \leq q \leq p_{j+1} - 1$, $j \geq 0$. For this, since $q \mapsto b_q$ is strictly increasing, it suffices to observe that, thanks to the second half in (20), we have $b_{p_{j+1}-1} = (1 + \delta_j)^{p_{j+1} - p_j - 1} a_{p_j} \leq A a_{p_j} = A a_{p_{j+1}-1}$. \square

The second result is the following.

Lemma 3.12. *Let \mathbb{M} be a weight sequence. Then \mathbb{M} satisfies (mg) if and only if $\omega_{\mathbb{M}}(t) = O(\nu_{\mathbf{m}}(t))$ as $t \rightarrow +\infty$.*

Proof. The condition (mg) for \mathbb{M} is equivalent to $m_n \leq A(M_n)^{1/n}$ for some $A \geq 1$ and all $n \in \mathbb{N}$ (e.g., see [21, Lemma 2.2]). It is also known that $\omega_{\mathbb{M}}(m_n) = \log(m_n^n/M_n)$ for $n \in \mathbb{N}$ (see [15, Chapitre I]). So, if $m_{n-1} \leq t < m_n$ for some $n \geq 1$, we get

$$\omega_{\mathbb{M}}(t) \leq \omega_{\mathbb{M}}(m_n) = n \log \left(\frac{m_n}{M_n^{1/n}} \right) \leq n \log(A) = \log(A) \nu_{\mathbf{m}}(t),$$

that is, $\omega_{\mathbb{M}}(t) = O(\nu_{\mathbf{m}}(t))$ as $t \rightarrow +\infty$.

Conversely, suppose that there exists $A \geq 1$ such that $\omega_{\mathbb{M}}(t) \leq A \nu_{\mathbf{m}}(t)$ for all $t \geq m_0$. By [21, Lemma 2.2], (mg) for \mathbb{M} holds true if and only if there exists $H \geq 1$ such that for all t large enough one has $2\nu_{\mathbf{m}}(t) \leq \nu_{\mathbf{m}}(Ht) + H$, and this we will prove. Take $H \geq \exp(2A)$ and $t \geq m_0$. Using (7), and since $\nu_{\mathbf{m}}$ is nondecreasing, we estimate

$$\begin{aligned} \nu_{\mathbf{m}}(Ht) &\geq A^{-1} \omega_{\mathbb{M}}(Ht) = A^{-1} \int_{m_0}^{Ht} \frac{\nu_{\mathbf{m}}(\lambda)}{\lambda} d\lambda \geq A^{-1} \int_t^{Ht} \frac{\nu_{\mathbf{m}}(\lambda)}{\lambda} d\lambda \\ &\geq A^{-1} \nu_{\mathbf{m}}(t) \int_t^{Ht} \frac{1}{\lambda} d\lambda = A^{-1} \log(H) \nu_{\mathbf{m}}(t) \geq 2\nu_{\mathbf{m}}(t), \end{aligned}$$

as desired.

We mention that an alternative, more abstract proof can be based in the theory of O-regular variation and Matuszewska indices for functions. By [10, Th. 4.4] we have that the lower Matuszewska indices of $\nu_{\mathbf{m}}$ and $\omega_{\mathbb{M}}$ agree, that is, $\beta(\nu_{\mathbf{m}}) = \beta(\omega_{\mathbb{M}})$, and by [10, Cor.2.17 and Cor. 4.2] we know \mathbb{M} has (mg) if and only if $\beta(\nu_{\mathbf{m}}) > 0$. So, if $\beta(\nu_{\mathbf{m}}) > 0$, by [10, Th. 4.3] we have that $\liminf_{t \rightarrow \infty} \frac{\nu_{\mathbf{m}}(t)}{\omega_{\mathbb{M}}(t)} > 0$, and we deduce that $\omega_{\mathbb{M}}(t) = O(\nu_{\mathbf{m}}(t))$ as $t \rightarrow +\infty$. Conversely, if $\omega_{\mathbb{M}}(t) = O(\nu_{\mathbf{m}}(t))$ as $t \rightarrow +\infty$, then $\liminf_{t \rightarrow \infty} \frac{\nu_{\mathbf{m}}(t)}{\omega_{\mathbb{M}}(t)} > 0$, so by [10, Th. 4.3] we have that $\beta(\omega_{\mathbb{M}}) > 0$, and we are done. \square

Proof of Theorem 3.10. We follow the proof of necessity for [3, Th. 2.2]. By Lemma 3.11 and the remark following it, we can assume without loss of generality that \mathbf{m} is strictly increasing.

Let G be an optimal $\{\mathbb{M}\}$ -flat function in $\mathcal{A}_{\{\widehat{\mathbb{M}}\}}(S_\gamma)$ for some $\gamma > 0$. So, there exists some $A > 0$ such that

$$p_{\mathbb{M},A}(G) := \sup_{n \in \mathbb{N}_0, x \in (0, +\infty)} \frac{|G^{(n)}(x)|}{A^n n! M_n} < +\infty.$$

This shows that the Carleman-Roumieu ultradifferentiable class $\mathcal{E}_{\{\widehat{\mathbb{M}}\}}((-\varepsilon, +\infty))$, consisting of all smooth complex-valued functions g defined on the interval $(-\varepsilon, \infty)$ for some $\varepsilon > 0$, and such that

$$\sup_{n \in \mathbb{N}_0, x \in (-\varepsilon, +\infty)} \frac{|g^{(n)}(x)|}{D^n n! M_n} < +\infty$$

for suitable $D > 0$, contains nontrivial flat functions (it suffices to extend G by 0 for $x \in (-\varepsilon, 0]$). Then, the well-known Denjoy-Carleman theorem (e.g., see [6, Th. 1.3.8]) yields that \mathbb{M} satisfies (nq).

Let now

$$R_n := \sum_{k \geq n} \frac{1}{(k+1)m_k} < +\infty, \quad n \in \mathbb{N}_0,$$

and let the function \bar{h} be defined by $\bar{h}(t) := n$ if $R_{n+1} < t \leq R_n$, $n \in \mathbb{N}_0$.

By [2, (14), p. 142] we obtain that

$$G(x) = |G(x)| \leq p_{\mathbb{M},A}(G) \exp(-\bar{h}(Aex)), \quad x \in (0, +\infty).$$

Combining this with the left-hand side of (13), with (4) and setting $C := p_{\mathbb{M},A}(G)$, we get

$$\exp(\bar{h}(Aex)) \leq \frac{C}{|G(x)|} \leq CK_1^{-1} \exp(\omega_{\mathbb{M}}(1/(K_2x))), \quad x \in (0, +\infty).$$

If we put $t = Aex$ and $B := Ae/K_2$, we obtain that for every $t > 0$,

$$\bar{h}(t) \leq \log(CK_1^{-1}) + \omega_{\mathbb{M}}(B/t). \quad (22)$$

By Lemma 3.12, there exists $C_1 \geq 1$ such that $\omega_{\mathbb{M}}(s) \leq C_1\nu_{\mathbf{m}}(s) + C_1$ for $s > 0$. Choosing $t = B/m_n$ in (22), we see that

$$\bar{h}(B/m_n) \leq \log(CK_1^{-1}) + \omega_{\mathbb{M}}(m_n) \leq \log(CK_1^{-1}) + C_1\nu_{\mathbf{m}}(m_n) + C_1 = \log(CK_1^{-1}) + C_1(n+1) + C_1,$$

since \mathbf{m} is strictly increasing. Hence, $\bar{h}(B/m_n) \leq C_2(n+1)$ for some $C_2 \in \mathbb{N}$ and all $n \in \mathbb{N}_0$. By definition of \bar{h} , we get $R_{C_2(n+1)+1} \leq B/m_n$, i.e.,

$$m_n \sum_{k \geq C_2(n+1)+1} \frac{1}{(k+1)m_k} \leq B, \quad n \in \mathbb{N}_0.$$

Finally,

$$\begin{aligned} m_n \sum_{k \geq n} \frac{1}{(k+1)m_k} &= m_n \sum_{k \geq C_2(n+1)+1} \frac{1}{(k+1)m_k} + m_n \sum_{k=n}^{C_2(n+1)} \frac{1}{(k+1)m_k} \\ &\leq B + m_n \frac{n(C_2-1) + C_2 + 1}{(n+1)m_n} \leq B + 2C_2, \end{aligned}$$

which is (snq) for \mathbb{M} . □

4 Construction of optimal flat functions for a family of non strongly regular sequences

As deduced in Theorem 3.5, the construction of optimal $\{\mathbb{M}\}$ -flat functions in sectors within an ultraholomorphic class, given by a regular sequence $\widehat{\mathbb{M}}$, provides extension operators and surjectivity results. Although such construction is only available in general for strongly regular sequences, we wish to present here a family of (non strongly) regular sequences for which this technique works.

We recall that, for logarithmically convex sequences $(M_p)_{p \in \mathbb{N}_0}$, the condition (dc) is equivalent to the condition $\log(M_p) = O(p^2)$, $p \rightarrow \infty$ (see [15, Ch. 6]). On the other hand, the condition (mg) implies that the sequence is below some Gevrey order (there exists $\alpha > 0$ such that $M_p = O(p!^\alpha)$ as $p \rightarrow \infty$; see e.g. [16, 27]).

We will work, for $q > 1$ and $1 < \sigma \leq 2$, with the sequences $\mathbb{M}_{q,\sigma} := (q^{p^\sigma})_{p \in \mathbb{N}_0}$. They are clearly weight sequences and, by (1), it is immediate that $\gamma(\mathbb{M}_{q,\sigma}) = \infty$, so they satisfy (snq) (see (3)). According to the previous comments, they satisfy (dc) but not (mg). So, $\widehat{\mathbb{M}}_{q,\sigma}$ is regular, but $\mathbb{M}_{q,\sigma}$ is not strongly regular.

The case $\sigma = 2$ is well-known, as it corresponds to the so-called q -Gevrey sequences, appearing in the study of formal solutions for q -difference equations.

First, we will construct a holomorphic function on $\mathbb{C} \setminus (-\infty, 0]$ which will provide, by restriction, an optimal $\{\mathbb{M}_{q,\sigma}\}$ -flat function in any unbounded sector S_γ with $0 < \gamma < 2$. Subsequently, we will obtain such functions on general sectors of the Riemann surface \mathcal{R} of the logarithm by ramification. This, according to Theorems 3.2 and 3.5, agrees with the fact that $\gamma(\mathbb{M}_{q,\sigma}) = \infty$.

4.1 Flatness in the class given by $\mathbb{M}_{q,\sigma}$

It will be convenient to note that for a fixed $\sigma \in (1, 2]$, there exists a unique $s \geq 2$ such that $\sigma = s/(s-1)$.

We start by suitably estimating the function

$$\omega_{\mathbb{M}_{q,\sigma}}(t) = \sup_{p \in \mathbb{N}_0} \ln \left(\frac{t^p}{q^{p^\sigma}} \right) = \sup_{p \in \mathbb{N}_0} (p \ln(t) - p^{s/(s-1)} \ln(q)), \quad t > 0.$$

Due to the fact that $\omega_{\mathbb{M}_{q,\sigma}}(t) = 0$ for $t \leq 1$ (since $m_0 = M_1/M_0 = M_1 = q > 1$ and by (7)), we will restrict our attention to the case $t > 1$. Obviously, $\omega_{\mathbb{M}_{q,\sigma}}(t)$ is bounded above by the supremum of $x \ln(t) - x^{s/(s-1)} \ln(q)$ when x runs over $(0, \infty)$, which is easily obtained by elementary calculus and occurs at the point

$$x_0 = \left(\frac{(s-1) \ln(t)}{s \ln(q)} \right)^{s-1}.$$

If we put

$$b_{q,s} := \frac{1}{s} \left(\frac{s-1}{s \ln(q)} \right)^{s-1}, \quad (23)$$

then

$$\omega_{\mathbb{M}_{q,\sigma}}(t) \leq \left(\frac{(s-1) \ln(t)}{s \ln(q)} \right)^{s-1} \ln(t) - \left(\frac{(s-1) \ln(t)}{s \ln(q)} \right)^s \ln(q) = b_{q,s} \ln^s(t), \quad t > 1. \quad (24)$$

On the other hand, for $t > q^{s/(s-1)}$ (what amounts to $x_0 > 1$) we also have that $\omega_{\mathbb{M}_{q,\sigma}}(t)$ is at least the value of $x \ln(t) - x^{s/(s-1)} \ln(q)$ at $x = \lfloor x_0 \rfloor$, that is,

$$\begin{aligned} \omega_{\mathbb{M}_{q,\sigma}}(t) &\geq \left\lfloor \left(\frac{(s-1) \ln(t)}{s \ln(q)} \right)^{s-1} \right\rfloor \ln(t) - \left\lfloor \left(\frac{(s-1) \ln(t)}{s \ln(q)} \right)^{s-1} \right\rfloor^{s/(s-1)} \ln(q) \\ &\geq \left(\left(\frac{(s-1) \ln(t)}{s \ln(q)} \right)^{s-1} - 1 \right) \ln(t) - \left(\frac{(s-1) \ln(t)}{s \ln(q)} \right)^s \ln(q) \\ &= b_{q,s} \ln^s(t) - \ln(t). \end{aligned} \quad (25)$$

Lemma 4.1. *For every $t \geq q^{2s/(s-1)}$ it holds*

$$b_{q,s} \ln^s(t) - \ln(t) \geq b_{q,s} \ln^s \left(\frac{t}{q^{s/(s-1)}} \right) - \ln \left(q^{s/(s-1)} \right). \quad (26)$$

Proof. Observe that every $t \geq q^{2s/(s-1)}$ may be written as $t = q^{ys/(s-1)}$ for some $y \geq 2$. Then, we have that

$$b_{q,s} \ln^s(t) - b_{q,s} \ln^s \left(\frac{t}{q^{s/(s-1)}} \right) = b_{q,s} \left(\frac{s \ln(q)}{s-1} \right)^s (y^s - (y-1)^s) = \frac{\ln(q)}{s-1} (y^s - (y-1)^s).$$

By the mean value theorem, $y^s - (y-1)^s > s(y-1)^{s-1}$, and since $s \geq 2$ and $y \geq 2$, we have $(y-1)^{s-1} \geq y-1$. So we deduce that

$$\frac{\ln(q)}{s-1}(y^s - (y-1)^s) > \frac{s \ln(q)}{s-1}(y-1) = \ln(t) - \ln\left(q^{s/(s-1)}\right),$$

as desired. \square

Combining (24) with (25) and (26), and using (4), we get

$$\exp\left(-b_{q,s} \ln^s\left(\frac{1}{t}\right)\right) \leq h_{\mathbb{M}_{q,\sigma}}(t) \leq q^{s/(s-1)} \exp\left(-b_{q,s} \ln^s\left(\frac{1}{q^{s/(s-1)}t}\right)\right), \quad 0 < t \leq q^{-2s/(s-1)}. \quad (27)$$

We can say that these estimates express optimal $\{\mathbb{M}_{q,\sigma}\}$ -flatness.

4.2 Optimal $\{\mathbb{M}_{q,\sigma}\}$ -flat function in S_2

The estimates in (27) suggest considering the function $\exp(-b_{q,s} \log^s(1/z))$, with, say, principal branches, as a candidate for providing optimal flat functions. However, its analyticity in wide sectors is not guaranteed. Moreover, even in small sectors around the direction $d = 0$, its behaviour at ∞ might not be as desired: For example, when $s = 2$ it tends to 0 as $0 < x \rightarrow \infty$, what excludes the possibility of proving the first inequality in (13).

Because of these reasons, we will first define a suitably modified function in the sector $S_2 = \mathbb{C} \setminus (-\infty, 0]$, prove its flatness there, and then turn to general sectors by composing it with an appropriate ramification.

We define

$$G_2^{q,s}(z) := \exp\left(-b_{q,s} \log^s\left(1 + \frac{1}{z}\right)\right), \quad z \in S_2, \quad (28)$$

where the principal branch of the logarithm is chosen for both \log and the power $w \mapsto w^s = \exp(s \log(w))$ involved. Observe that if $z \in S_2$, then $1 + 1/z \in \mathbb{C} \setminus (-\infty, 1]$, and so $\log(1 + 1/z) = \ln(|1 + 1/z|) + i \arg(1 + 1/z) \notin (-\infty, 0]$. This ensures that the map

$$z \mapsto \log^s\left(1 + \frac{1}{z}\right) = \exp\left(s \log\left(\log\left(1 + \frac{1}{z}\right)\right)\right)$$

is also holomorphic in S_2 , and so is $G_2^{q,s}$.

Moreover, $G_2^{q,s}(x) > 0$ for $x > 0$. In order that $G_2^{q,s}$ is an optimal $\{\mathbb{M}_{q,\sigma}\}$ -flat function in S_2 , we are only left with proving the estimates (13). It turns out to be more convenient to work with the associated kernel

$$e_2(z) := G_2^{q,s}(1/z) = \exp(-b_{q,s} \log^s(1+z)), \quad z \in S_2,$$

and verify the following result.

Lemma 4.2. *There exist positive constants C_1, C_2, C_3, C_4 such that*

$$C_1 e_2(C_2|z|) \leq |e_2(z)| \leq C_3 e_2(C_4|z|), \quad z \in S_2.$$

Proof. We prove first the left inequality. Let $z \in S_2$ and $K > 0$. Since $K|z| + 1 > 1$, it is clear that $\log^s(K|z| + 1) = \ln^s(K|z| + 1) > 0$. On the other hand, we get

$$\begin{aligned} \Re(\log^s(z+1)) &\leq |\log^s(z+1)| = |\log(z+1)|^s \\ &= (\ln^2|z+1| + \arg^2(z+1))^{\frac{s}{2}} \leq (\ln^2|z+1| + \pi^2)^{\frac{s}{2}}. \end{aligned} \quad (29)$$

If we suppose that $|z| > 2$, then $\ln|z+1| > 0$ and $\ln^2|z+1| \leq \ln^2(|z+1|)$. Let us take $C_2 > e^\pi$, then $\lim_{x \rightarrow \infty} (\ln(C_2 x + 1) - \ln(x + 1)) = \ln(C_2) > \pi$, and there exists a constant $R > 2$ such that for $x \geq R$ we will have that $\ln(C_2 x + 1) - \ln(x + 1) \geq \pi$. So, let us take $z \in S_2$ with $|z| \geq R$, then

$$\ln^2(C_2|z| + 1) - \ln^2(|z| + 1) = \ln((C_2|z| + 1)(|z| + 1)) (\ln(C_2|z| + 1) - \ln(|z| + 1)) \geq \pi^2.$$

Consequently,

$$(\ln^2|z+1| + \pi^2)^{\frac{s}{2}} \leq (\ln^2(|z|+1) + \pi^2)^{\frac{s}{2}} \leq \ln^s(C_2|z| + 1).$$

Then, from (29), we obtain that $\Re(\log^s(z+1)) \leq \ln^s(C_2|z| + 1)$, and we can deduce that

$$|e_2(z)| = \exp(-b_{q,s} \Re(\log^s(z+1))) \geq \exp(-b_{q,s} \ln^s(C_2|z| + 1)) = e_2(C_2|z|), \quad z \in S_2, |z| \geq R.$$

Finally, since the function $|e_2(z)|$ stays bounded and bounded away from 0 for bounded $|z|$ (in particular, it tends to 1 when z tends to 0 in S_2), there exists a constant $C_1 > 0$ such that $C_1 e_2(C_2|z|) \leq |e_2(z)|$ for all $z \in S_2$.

We now turn to the right inequality. In the first place, we observe that for every $z \in S_2$,

$$\Re(\log^s(z+1)) = |\log^s(z+1)| \cos(\arg(\log^s(z+1))) = |\log(z+1)|^s \cos(s \arg(\log(z+1))). \quad (30)$$

Now,

$$s |\arg(\log(z+1))| = s \left| \arctan \left(\frac{\arg(z+1)}{\ln|z+1|} \right) \right| \leq s \left| \arctan \left(\frac{\pi}{\ln|z+1|} \right) \right|. \quad (31)$$

Hence, setting

$$R_0 := 1 + \exp \left(\frac{\pi}{\tan(\pi/(2s))} \right) \geq 2,$$

we get that $|z| > R_0$ implies that $|z+1| > R_0 - 1 \geq 1$, and therefore $\ln|z+1| > 0$ and

$$\frac{\pi}{\ln|z+1|} < \tan \left(\frac{\pi}{2s} \right).$$

From this and (31) we deduce that $\cos(s \arg(\log(z+1))) > 0$. Then, continuing with (30),

$$\begin{aligned} \Re(\log^s(z+1)) &\geq |\Re(\log(z+1))|^s \cos(s \arg(\log(z+1))) \\ &= \ln^s|z+1| - \ln^s|z+1| \frac{\sin^2(s \arg(\log(z+1)))}{1 + \cos(s \arg(\log(z+1)))}. \end{aligned} \quad (32)$$

Now, from the equality in (31) we see that $s \arg(\log(z+1)) \rightarrow 0$ as $z \rightarrow \infty$ in S_2 , and moreover

$$\lim_{\substack{z \rightarrow \infty \\ z \in S_2}} \left[\left(\frac{\sin^2(s \arg(\log(z+1)))}{1 + \cos(s \arg(\log(z+1)))} \right) / \left(\frac{s^2 \arg^2(z+1)}{2 \ln^2|z+1|} \right) \right] = 1.$$

Therefore, there exist $R_1 \geq R_0$ and $C > 0$ such that

$$\frac{\sin^2(s \arg(\log(z+1)))}{1 + \cos(s \arg(\log(z+1)))} \leq C \frac{1}{\ln^2|z+1|}, \quad |z| > R_1.$$

We deduce from (32) that for $z \in S_2$ with $|z| > R_1$,

$$\Re(\log^s(z+1)) \geq \ln^s|z+1| - C \ln^{s-2}|z+1| \geq \ln^s(|z|-1) - C \ln^{s-2}(|z|+1). \quad (33)$$

We would be almost done if we obtain, for the right-hand side in (33), a lower bound in terms of, say, $\ln^s(1 + |z|/2)$ for $|z|$ sufficiently large.

This is easy in case $s = 2$, for it suffices to take $|z| > 4$ in order to have $3 < 1 + |z|/2 < |z| - 1$, and so if $|z| \geq R_2 := \max\{R_1, 4\}$ we have

$$\Re(\log^s(z+1)) \geq \ln^s(|z| - 1) - C \geq \ln^s\left(1 + \frac{|z|}{2}\right) - C.$$

In case $s > 2$, it is not difficult to check that

$$\lim_{x \rightarrow +\infty} \left(\ln^s(x-1) - C \ln^{s-2}(x+1) - \ln^s\left(1 + \frac{x}{2}\right) \right) = +\infty,$$

so that, according to (33), there exists $R_2 \geq R_1$ such that for $z \in S_2$ with $|z| \geq R_2$ one has

$$\Re(\log^s(z+1)) \geq \ln^s\left(1 + \frac{|z|}{2}\right).$$

In any case, we can deduce an upper estimate of the form

$$\begin{aligned} |e_2(z)| &= \exp\left(-b_{q,s} \Re(\log^s(z+1))\right) \\ &\leq e^C \exp\left(-b_{q,s} \ln^s\left(1 + \frac{|z|}{2}\right)\right) = e^C e_2\left(\frac{|z|}{2}\right), \quad z \in S_2, |z| > R_2, \end{aligned}$$

which can be extended to the whole of S_2 by suitably enlarging the constant C , as before. \square

We are ready for the main objective of this section.

Theorem 4.3. *The function $G_2^{q,s}$ defined in (28) is an optimal $\{\mathbb{M}_{q,\sigma}\}$ -flat function in S_2 .*

Proof. The previous lemma ensures that there exist positive constants C_1, C_2, C_3, C_4 such that

$$C_1 \exp\left(-b_{q,s} \ln^s\left(1 + \frac{C_2}{|z|}\right)\right) \leq |G_2^{q,s}(z)| \leq C_3 \exp\left(-b_{q,s} \ln^s\left(1 + \frac{C_4}{|z|}\right)\right), \quad z \in S_2. \quad (34)$$

Observe that these inequalities guarantee that $|G_2^{q,s}(z)|$ is bounded and bounded away from 0 as soon as $|z| \geq r$ for any fixed $r > 0$, since then

$$C_1 \exp(-b_{q,s} \ln^s(1 + C_2/r)) \leq |G_2^{q,s}(z)| \leq C_3.$$

As the same is true for $h_{\mathbb{M}_{q,\sigma}}(t)$ for every $t \geq t_0$ and any fixed $t_0 > 0$ (see Lemma 2.1), we only need to check the estimates (13) for small enough $|z|$.

For $|z| \leq C_4$ it is clear that $\ln(1 + C_4/|z|) > \ln(C_4/|z|) \geq 0$. Then, from the first inequality in (27) we have that

$$\begin{aligned} |G_2^{q,s}(z)| &\leq C_3 \exp\left(-b_{q,s} \ln^s\left(1 + \frac{C_4}{|z|}\right)\right) \\ &\leq C_3 \exp\left(-b_{q,s} \ln^s\left(\frac{C_4}{|z|}\right)\right) \leq C_3 h_{\mathbb{M}_{q,\sigma}}\left(\frac{|z|}{C_4}\right), \quad |z| \leq C_4 q^{-2s/(s-1)}, \end{aligned}$$

and we have proved the right-hand part of (13).

By using the left-hand side of (34), we have for $z \in S_2$ that

$$|G_2^{q,s}(z)| \geq C_1 \exp\left(-b_{q,s} \ln^s\left(\frac{C_2}{|z|}\right)\right) \exp\left(-b_{q,s} \left[\ln^s\left(1 + \frac{C_2}{|z|}\right) - \ln^s\left(\frac{C_2}{|z|}\right) \right]\right).$$

The mean value theorem gives that $\ln^s(1 + C_2/|z|) - \ln^s(C_2/|z|)$ tends to zero if $|z| \rightarrow 0$, and we deduce that there exists L such that

$$|G_2^{q,s}(z)| \geq L \exp\left(-b_{q,s} \ln^s\left(\frac{C_2}{|z|}\right)\right), \quad |z| \leq 1.$$

The second inequality in (27) implies now that, as long as $|z| \leq \min\{1, C_2 q^{-s/(s-1)}\}$, we have

$$|G_2^{q,s}(z)| \geq L q^{-s/(s-1)} h_{\mathbb{M}_{q,\sigma}}\left(\frac{|z|}{C_2 q^{s/(s-1)}}\right),$$

and so the left-hand side of (13) also holds. \square

4.3 Optimal $\{\mathbb{M}_{q,\sigma}\}$ -flat function in arbitrary sectors

Let us consider a sector $S_\gamma \subset \mathcal{R}$ with $\gamma > 2$, and define the function

$$G_\gamma^{q,s}(z) := \exp\left(-b_{q,s} \left(\frac{\gamma}{2}\right)^s \log^s\left(1 + z^{-2/\gamma}\right)\right) = \left(G_2^{q,s}(z^{2/\gamma})\right)^{(\gamma/2)^s}, \quad z \in S_\gamma. \quad (35)$$

The map $z \mapsto z^{2/\gamma}$ is holomorphic from S_γ into S_2 , and so $G_\gamma^{q,s}$ is holomorphic in S_γ . We will prove that this function is an optimal $\{\mathbb{M}_{q,\sigma}\}$ -flat function in this sector. It is clear that $G_\gamma^{q,s}(x) > 0$ for every $x > 0$, so only the estimates in (13) are pending.

As before, we consider the kernel

$$e_\gamma(z) := G_\gamma^{q,s}(1/z) = \exp\left(-b_{q,s} \left(\frac{\gamma}{2}\right)^s \log^s\left(1 + z^{2/\gamma}\right)\right) = \left(e_2(z^{2/\gamma})\right)^{(\gamma/2)^s}, \quad z \in S_\gamma.$$

Lemma 4.4. *There exist constants $B_1, B_2, B_3, B_4 > 0$ such that*

$$B_1 e_2(B_2|z|) \leq |e_\gamma(z)| \leq B_3 e_2(B_4|z|), \quad z \in S_\gamma. \quad (36)$$

Proof. According to the definition of e_γ and by applying Lemma 4.2, there exist constants $C_1, C_2, C_3, C_4 > 0$ such that for every $z \in S_\gamma$ one has

$$\left(C_1 e_2(C_2|z|^{2/\gamma})\right)^{(\gamma/2)^s} \leq |e_\gamma(z)| = \left|e_2(z^{2/\gamma})\right|^{(\gamma/2)^s} \leq \left(C_3 e_2(C_4|z|^{2/\gamma})\right)^{(\gamma/2)^s}.$$

We recall that the function $|e_2(z)|$ stays bounded and bounded away from 0 for bounded $|z|$; from the previous estimates, the same can be said about $|e_\gamma(z)|$, and so we can prove (36) by restricting our considerations to large enough values of $|z|$ and well chosen $B_2, B_4 > 0$, and then suitably enlarging the constants $B_1, B_3 > 0$ involved. Let us observe that

$$\begin{aligned} \left(e_2(C_4|z|^{2/\gamma})\right)^{(\gamma/2)^s} &= \exp\left(-b_{q,s} \ln^s\left[(1 + C_4|z|^{2/\gamma})^{\gamma/2}\right]\right), \\ e_2(B_4|z|) &= \exp\left(-b_{q,s} \ln^s(1 + B_4|z|)\right). \end{aligned}$$

So, we will be done with the upper estimates in (36) if we see that

$$\ln^s(1 + B_4|z|) - \ln^s\left[(1 + C_4|z|^{2/\gamma})^{\gamma/2}\right]$$

admits an upper bound for large enough $|z|$ and suitably chosen $B_4 > 0$. But this follows from the clear fact that

$$\ln^s(1 + B_4|z|) - \ln^s\left[\left(1 + C_4|z|^{2/\gamma}\right)^{\gamma/2}\right] \sim -s \ln\left(\frac{C_4^{\gamma/2}}{B_4}\right) \ln^{s-1}(1 + B_4|z|), \quad |z| \rightarrow \infty,$$

where \sim means that the quotient of both expressions tends to 1. Indeed, in view of this equivalence it suffices to choose any $B_4 < C_4^{\gamma/2}$ in order to have the desired estimation for suitably large B_3 and $|z|$.

In the same way, one can obtain the first estimation in (36) by choosing any $B_2 > C_2^{\gamma/2}$ and considering large enough $B_1 > 0$ and $|z|$. \square

Corollary 4.5. *The function $G_\gamma^{q,s}$ defined in (35) is an optimal $\{\mathbb{M}_{q,\sigma}\}$ -flat function in S_γ .*

Proof. By the previous lemma, there exist $B_1, B_2, B_3, B_4 > 0$ such that

$$B_1 \exp\left(-b_{q,s} \ln^s\left(1 + \frac{B_2}{|z|}\right)\right) \leq |G_\gamma^{q,s}(z)| \leq B_3 \exp\left(-b_{q,s} \ln^s\left(1 + \frac{B_4}{|z|}\right)\right), \quad z \in S_\gamma.$$

Note that these estimates are essentially those in (34), and so the conclusion follows in exactly the same way as in the proof of Theorem 4.3. \square

Remark 4.6. We mention that a similar approach has been followed in the preprint [8], by A. Lastra and the first and third authors, in order to construct extension operators for the ultraholomorphic classes associated with the sequences $\mathbb{M}^{\tau,\sigma} = (p^\tau p^\sigma)_{p \in \mathbb{N}_0}$, for $\tau > 0$ and $\sigma \in (1, 2)$. These sequences have appeared in a series of papers by S. Pilipović, N. Teofanov and F. Tomić [17, 18, 19, 20], inducing ultradifferentiable spaces of so-called extended Gevrey regularity. However, in that case the construction of suitable kernels for our technique involves the Lambert function, whose handling is not so convenient. This fact has caused our results to be available only in sectors strictly contained in S_2 , in spite of the fact that $\gamma(\mathbb{M}_{\tau,\sigma}) = \infty$, what would in principle allow for such extension operators to exist in sectors of arbitrary opening.

5 Convolved sequences, flat functions and extension results

We show in this section that whenever two weight sequences are given and there exist optimal flat functions in the respectively associated classes, then optimal flat functions exist in the class defined by the so-called convolved sequence as well. Moreover, the extension technique works if one of the convolved sequences satisfies (dc).

5.1 Convolved sequences

Let $\mathbb{M}^1 = (M_p^1)_{p \in \mathbb{N}_0}$, $\mathbb{M}^2 = (M_p^2)_{p \in \mathbb{N}_0}$ be two sequences of positive real numbers, then the *convolved sequence* $\mathbb{L} := \mathbb{M}^1 \star \mathbb{M}^2$ is $(L_p)_{p \in \mathbb{N}_0}$ given by

$$L_p := \min_{0 \leq q \leq p} M_q^1 M_{p-q}^2, \quad p \in \mathbb{N}_0,$$

see [13, (3.15)]. Hence, obviously $\mathbb{M}^1 \star \mathbb{M}^2 = \mathbb{M}^2 \star \mathbb{M}^1$.

For all $p \in \mathbb{N}_0$ we have $L_p \leq \min\{M_0^1 M_p^2, M_0^2 M_p^1\}$. So, if in addition $M_0^1 = M_0^2 = 1$, then we get $L_0 = 1$ and

$$L_p \leq \min\{M_p^1, M_p^2\}, \quad p \in \mathbb{N}_0. \quad (37)$$

Given $\mathbb{M} = (M_p)_{p \in \mathbb{N}_0}$ with $M_0 = 1$, put $\mathbb{L} = (L_p)_{p \in \mathbb{N}_0} = \mathbb{M} \star \mathbb{M}$. The condition (mg) states precisely that there exists $A > 0$ such that $M_p \leq A^p L_p$ for every $p \in \mathbb{N}_0$; according to (37), \mathbb{M} satisfies (mg) if and only if \mathbb{M} and $\mathbb{M} \star \mathbb{M}$ are equivalent.

Remark 5.1. Let $\mathbb{M}, \mathbb{M}^1, \mathbb{M}^2$ be weight sequences.

- (i) In [13, Lemma 3.5] the following facts are shown: $\mathbb{M}^1 \star \mathbb{M}^2$ is again a weight sequence. The corresponding quotient sequence $\mathbf{m}^1 \star \mathbf{m}^2$ is obtained when rearranging resp. ordering the sequences \mathbf{m}^1 and \mathbf{m}^2 in the order of growth. This yields, by definition of the counting function (see (6)), that for all $t \geq 0$ one has

$$\nu_{\mathbf{m}^1 \star \mathbf{m}^2}(t) = \nu_{\mathbf{m}^1}(t) + \nu_{\mathbf{m}^2}(t);$$

so, by (7) we get

$$\omega_{\mathbb{M}^1 \star \mathbb{M}^2}(t) = \omega_{\mathbb{M}^1}(t) + \omega_{\mathbb{M}^2}(t), \quad t \geq 0,$$

and by (4) we obtain

$$h_{\mathbb{M}^1 \star \mathbb{M}^2}(t) = h_{\mathbb{M}^1}(t)h_{\mathbb{M}^2}(t), \quad t > 0. \quad (38)$$

- (ii) If either \mathbb{M}^1 or \mathbb{M}^2 has (dc), then $\mathbb{M}^1 \star \mathbb{M}^2$ as well: As said before, for sequences $(M_p)_{p \in \mathbb{N}_0}$ satisfying (lc), the condition (dc) amounts to the condition $\log(M_p) = O(p^2)$, $p \rightarrow \infty$. Then, it suffices to apply (37).

- (iv) As seen in item (i), for every $t \geq 0$ we have

$$2\omega_{\mathbb{M}}(t) = \omega_{\mathbb{M} \star \mathbb{M}}(t).$$

Since \mathbb{M} satisfies (mg) if and only if there exists $H \geq 1$ such that

$$2\omega_{\mathbb{M}}(t) \leq \omega_{\mathbb{M}}(Ht) + H, \quad t \geq 0$$

(see [13, Prop. 3.6]), it turns out that (mg) amounts to the fact that

$$\omega_{\mathbb{M} \star \mathbb{M}}(t) \leq \omega_{\mathbb{M}}(Ht) + H, \quad t \geq 0,$$

for some $H \geq 1$, or in other words,

$$h_{\mathbb{M}}(t) \leq e^H h_{\mathbb{M} \star \mathbb{M}}(Ht), \quad t > 0.$$

5.2 Optimal flat functions and extension procedure

Let \mathbb{M}^1 and \mathbb{M}^2 be weight sequences such that optimal flat functions $G_{\mathbb{M}^1}$ and $G_{\mathbb{M}^2}$ exist in the corresponding classes with uniform asymptotic expansion in a given sector S . Then, we claim that $G_{\mathbb{M}^1 \star \mathbb{M}^2} := G_{\mathbb{M}^1} \cdot G_{\mathbb{M}^2}$ is an optimal flat function (on the same sector S) in the class associated with the sequence $\mathbb{M}^1 \star \mathbb{M}^2$. Obviously, $G_{\mathbb{M}^1 \star \mathbb{M}^2}(x) > 0$ for all $x > 0$. Suppose K_m and J_m , $m = 1, 2, 3, 4$, are the constants appearing in (13) for $G_{\mathbb{M}^1}$ and $G_{\mathbb{M}^2}$, respectively. By (38) we get that, for all $z \in S$,

$$|G_{\mathbb{M}^1}(z) \cdot G_{\mathbb{M}^2}(z)| \leq K_3 h_{\mathbb{M}^1}(K_4|z|) J_3 h_{\mathbb{M}^2}(J_4|z|) \leq K_3 J_3 h_{\mathbb{M}^1}(D|z|) h_{\mathbb{M}^2}(D|z|) = C h_{\mathbb{M}^1 \star \mathbb{M}^2}(D|z|),$$

with $C := K_3 J_3$ and $D := \max\{K_4, J_4\}$, since each function $h_{\mathbb{M}}$ is nondecreasing. Similarly, we can estimate

$$\begin{aligned} |G_{\mathbb{M}^1}(z) \cdot G_{\mathbb{M}^2}(z)| &\geq K_1 h_{\mathbb{M}^1}(K_2|z|) J_1 h_{\mathbb{M}^2}(J_2|z|) \\ &\geq K_1 J_1 h_{\mathbb{M}^1}(D_1|z|) h_{\mathbb{M}^2}(D_1|z|) = C_1 h_{\mathbb{M}^1 \star \mathbb{M}^2}(D_1|z|), \end{aligned}$$

with $C_1 := K_1 J_1$ and $D_1 := \min\{K_2, J_2\}$, and the conclusion follows.

In case at least one of the sequences \mathbb{M}^1 and \mathbb{M}^2 satisfies (dc), $\mathbb{M}^1 \star \mathbb{M}^2$ does so, and the extension operators from Theorem 3.5 will be available for the convolved sequence.

5.3 Some examples

Fix $q > 1$ and $\sigma \in (1, 2]$. Let us put $\mathbb{L}_{q,\sigma} := \mathbb{M}_{q,\sigma} \star \mathbb{M}_{q,\sigma}$, $\mathbb{L}_{q,\sigma} = (L_p)_{p \in \mathbb{N}_0}$. It is not difficult to check that

$$L_{2p} = q^{2p^\sigma}, \quad L_{2p+1} = q^{p^\sigma + (p+1)^\sigma}, \quad p \in \mathbb{N}_0.$$

Observe that $2p^\sigma = 2^{1-\sigma}(2p)^\sigma$, so that L_{2p} equals the $2p$ -th term of the sequence $\mathbb{M}_{q^{2^{1-\sigma}}, \sigma}$. Regarding the odd terms, it is a consequence of Taylor's formula at $x = 0$ for the functions of the form $x \mapsto (1+x)^\alpha$, $\alpha > 0$, that

$$p^\sigma + (p+1)^\sigma - 2^{1-\sigma}(2p+1)^\sigma = O(p^{\sigma-2}), \quad p \rightarrow \infty.$$

Since $\sigma \in (1, 2]$, we deduce that $\mathbb{L}_{q,\sigma}$ is equivalent to $\mathbb{M}_{q^{2^{1-\sigma}}, \sigma}$.

According to Subsection 5.2, an optimal flat function in the class associated with $\mathbb{L}_{q,\sigma}$ in, say, the sector S_2 is the function

$$G(z) := G_2^{q,s}(z)G_2^{q,s}(z) = \exp\left(-2b_{q,s} \log^s\left(1 + \frac{1}{z}\right)\right), \quad z \in S_2.$$

It is not a surprise that, from the definition (23) of $b_{q,s}$ and the relation between σ and s , one obtains $b_{q^{2^{1-\sigma}}, s} = 2b_{q,s}$, and so G is precisely $G_2^{q^{2^{1-\sigma}}, s}$, what agrees with the aforementioned equivalence of sequences.

If we consider instead $1 < \sigma < 2$ and $\mathbb{J} := \mathbb{M}_{q,\sigma} \star \mathbb{M}_{q,2}$, $\mathbb{J} = (J_p)_{p \in \mathbb{N}_0}$, the computation of the terms J_p is no longer possible in closed form, since their values depend for general p on the position of σ within the interval $(1, 2)$. However, the previous subsection shows that, for s associated with σ as usual, the function

$$G(z) := G_2^{q,s}(z)G_2^{q,2}(z) = \exp\left(-b_{q,s} \log^s\left(1 + \frac{1}{z}\right) - b_{q,2} \log^2\left(1 + \frac{1}{z}\right)\right), \quad z \in S_2,$$

is a flat function in the class associated with \mathbb{J} in S_2 . Note that s is not equal to 2, hence the very aspect of the exponent in this function, and the fact that the restriction $G|_{(0,\infty)}$ is closely related to the function $h_{\mathbb{J}}$ (see Definition 3.3), shows that \mathbb{J} is not equivalent to any of the sequences $\mathbb{M}_{q,\sigma}$. Since the sequence \mathbb{J} does satisfy (dc), the extension procedure described in this paper is available for the classes associated with \mathbb{J} .

Observe that these examples of optimal flat functions can also be provided in general sectors S_γ , $\gamma > 2$, by using the functions $G_\gamma^{q,s}$ introduced in (35).

Acknowledgements: The first three authors are partially supported by the Spanish Ministry of Science and Innovation under the project PID2019-105621GB-I00. The fourth author is supported by FWF-Project P33417-N.

References

- [1] W. Balser, *Formal power series and linear systems of meromorphic ordinary differential equations*, Springer, Berlin, 2000.
- [2] T. Bang, *The theory of metric spaces applied to infinitely differentiable functions*, Math. Scand., 1:137–152, 1953.
- [3] J. Bruna, *An extension theorem of Whitney type for non quasi-analytic classes of functions*. J. London Math. Soc., 22(2):495–505, 1980.

- [4] A. Debrouwere, Solution to the Stieltjes moment problem in Gelfand-Shilov spaces, *Studia Math.* 254 (2020), 295–323, doi: 10.4064/sm190627-8-10.
- [5] E. M. Dyn'kin, Pseudoanalytic extension of smooth functions. The uniform scale, *Amer. Math. Soc. Transl.* (2) 115 (1980), 33–58.
- [6] L. Hörmander. *The analysis of linear partial differential operators I, Distribution theory and Fourier analysis.* Springer-Verlag, 2003.
- [7] J. Jiménez-Garrido, *Applications of regular variation and proximate orders to ultraholomorphic classes, asymptotic expansions and multisummability*, PhD dissertation, University of Valladolid, 2018. Available at <http://uvadoc.uva.es/handle/10324/29501> (last accessed July 13th, 2020).
- [8] J. Jiménez-Garrido, A. Lastra, J. Sanz, Extension operators for some ultraholomorphic classes defined by sequences of rapid growth, submitted, available at <https://arxiv.org/abs/2204.01316>.
- [9] J. Jiménez-Garrido, J. Sanz, Strongly regular sequences and proximate orders, *J. Math. Anal. Appl.* 438 (2016), no. 2, 920–945.
- [10] J. Jiménez-Garrido, J. Sanz, G. Schindl, Indices of O-regular variation for weight functions and weight sequences, *Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM* 113 (4) (2019), 3659–3697.
- [11] J. Jiménez-Garrido, J. Sanz, G. Schindl, Injectivity and surjectivity of the asymptotic Borel map in Carleman ultraholomorphic classes, *J. Math. Anal. Appl.* 469 (2019), no. 1, 136–168.
- [12] J. Jiménez-Garrido, J. Sanz, G. Schindl, Surjectivity of the asymptotic Borel map in Carleman-Roumieu ultraholomorphic classes defined by regular sequences, *Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM* (2021) 115:181, <https://doi.org/10.1007/s13398-021-01119-y>.
- [13] H. Komatsu, Ultradistributions, I: Structure theorems and a characterization, *J. Fac. Sci. Univ. Tokyo Sect. IA Math.* 20 (1973), 25–105.
- [14] A. Lastra, S. Malek, and J. Sanz. Continuous right inverses for the asymptotic Borel map in ultraholomorphic classes via a Laplace-type transform. *J. Math. Anal. Appl.*, 396:724–740, 2012.
- [15] S. Mandelbrojt, *Séries adhérentes, régularisation des suites, applications*, Collection de monographies sur la théorie des fonctions, Gauthier-Villars, Paris, 1952.
- [16] W. Matsumoto, Characterization of the separativity of ultradifferentiable classes, *J. Math. Kyoto Univ.* 24-4 (1984), 667–678.
- [17] S. Pilipović, N. Teofanov, F. Tomić, On a class of ultradifferentiable functions. *Novi Sad J. Math.* 45, no. 1, 125–142 (2015).
- [18] S. Pilipović, N. Teofanov, F. Tomić, Beyond Gevrey regularity, *J. Pseudo-Differ. Oper. Appl.* 7 (2016), 113–140. doi: 10.1007/s11868-016-0145-0
- [19] S. Pilipović, N. Teofanov, F. Tomić, A Paley-Wiener theorem in extended Gevrey regularity. *J. Pseudo-Differ. Oper. Appl.* 11, 593–612 (2020). doi: <https://doi.org/10.1007/s11868-019-00298-y>
- [20] S. Pilipović, N. Teofanov, F. Tomić, Boundary values in ultradistribution spaces related to extended Gevrey regularity, *Mathematics* 2021, 9, 7. doi: <https://dx.doi.org/10.3390/math9010007>

- [21] A. Rainer and G. Schindl. Extension of Whitney jets of controlled growth. *Math. Nachr.*, (290):2356–2374, 2017.
- [22] J. P. Ramis, Dévissage Gevrey, *Asterisque* 59–60 (1978), 173–204.
- [23] B. Rodríguez-Salinas, Funciones con momentos nulos, *Rev. Acad. Ci. Madrid* 49 (1955), 331–368.
- [24] J. Sanz, Flat functions in Carleman ultraholomorphic classes via proximate orders, *J. Math. Anal. Appl.* 415 (2014), 623–643.
- [25] J. Schmets, M. Valdivia, Extension maps in ultradifferentiable and ultraholomorphic function spaces, *Studia Math.* 143 (3) (2000), 221–250.
- [26] V. Thilliez, Extension Gevrey et rigidité dans un secteur, *Studia Math.* 117 (1995), 29–41.
- [27] V. Thilliez, Division by flat ultradifferentiable functions and sectorial extensions, *Results Math.* 44 (2003), 169–188.

Affiliation:

Javier Jiménez-Garrido:
 Departamento de Matemáticas, Estadística y Computación
 Universidad de Cantabria
 Avda. de los Castros, s/n, 39005 Santander, Spain
 Instituto de Investigación en Matemáticas IMUVA, Universidad de Valladolid
 ORCID: 0000-0003-3579-486X
 E-mail: jesusjavier.jimenez@unican.es

Ignacio Miguel-Cantero:
 Departamento de Álgebra, Análisis Matemático, Geometría y Topología
 Universidad de Valladolid
 Facultad de Ciencias, Paseo de Belén 7, 47011 Valladolid, Spain.
 Instituto de Investigación en Matemáticas IMUVA
 ORCID: 0000-0001-5270-0971
 E-mail: ignacio.miguel@uva.es

Javier Sanz:
 Departamento de Álgebra, Análisis Matemático, Geometría y Topología
 Universidad de Valladolid
 Facultad de Ciencias, Paseo de Belén 7, 47011 Valladolid, Spain.
 Instituto de Investigación en Matemáticas IMUVA
 ORCID: 0000-0001-7338-4971
 E-mail: javier.sanz.gil@uva.es

Gerhard Schindl:
 Fakultät für Mathematik, Universität Wien, Oskar-Morgenstern-Platz 1, A-1090 Wien, Austria.
 ORCID: 0000-0003-2192-9110
 E-mail: gerhard.schindl@univie.ac.at