

# STUDY OF THE OPERATOR $\partial^k \bar{\partial}^k + c$ IN THE WEIGHTED HILBERT SPACE $L^2(\mathbb{C}, e^{-|z|^2})$ .

ERAMANE BODIAN & SOUHAIBOU SAMBOU & PAPA BADIANE & WINNIE OSSETE INGOBA & SALOMON SAMBOU

**ABSTRACT.** By the Hörmander's  $L^2$ -method, we study the operator  $\partial^k \bar{\partial}^k + c$  for any order  $k$  in the weighted Hilbert space  $L^2(\mathbb{C}, e^{-|z|^2})$ . We prove the existence of its right inverse which is also a bounded operator.

**Keywords :** The operator  $\partial^k \bar{\partial}^k + c$ , Weighted Hilbert space  $L^2(\mathbb{C}, e^{-|z|^2})$ , Hörmander's  $L^2$ -method.

**Mathematics Subject Classification (2010) .** 32F32.

## 1. INTRODUCTION

In [1], Shoayu DAI and Yifei PAN are interested to the operator's study  $\frac{d^k}{dx^k} + a$ . This work has been extended in [2] to the operator  $\bar{\partial}^k + a$ . Inspired by the  $\partial \bar{\partial}$  problem, we study the operator  $\partial^k \bar{\partial}^k + c$  in the weighted Hilbert space  $L^2(\mathbb{C}, e^{-|z|^2})$ .

We prove the existence of a weak solution of the equation

$$\partial^k \bar{\partial}^k u + cu = f$$

where  $u, f \in L^2(\mathbb{C}, e^{-\varphi})$ ,  $c$  is a complex constant,  $\bar{\partial}^k = \frac{\partial^k}{\partial \bar{z}^k}$  and  $\partial^k = \frac{\partial^k}{\partial z^k}$ . We thus have the following result:

**Theorem 1.1.** *For any  $f \in L^2(\mathbb{C}, e^{-|z|^2})$ , there exists a weak solution  $u \in L^2(\mathbb{C}, e^{-|z|^2})$  of the equation*

$$\partial^k \bar{\partial}^k u + cu = f$$

with the norm estimate

$$\int_{\mathbb{C}} |u|^2 e^{-|z|^2} d\sigma \leq \frac{1}{(k!)^2} \int_{\mathbb{C}} |f|^2 e^{-|z|^2} d\sigma.$$

Moreover for  $k = 1$ , we have for any smooth and nonnegative function  $\varphi$  define on  $\mathbb{C}$  such that  $\Delta(e^\varphi \Delta e^{-\varphi}) > 0$ . For all  $f \in L^2(\mathbb{C}, e^{-\varphi})$  such that

$$\frac{f}{\sqrt{\Delta(e^\varphi \Delta e^{-\varphi})}} \in L^2(\mathbb{C}, e^{-\varphi}),$$

there exists a weak solution  $u \in L^2(\mathbb{C}, e^{-\varphi})$  of the equation

$$\partial^1 \bar{\partial}^1 u + cu = f$$

with the norm estimate

$$\int_{\mathbb{C}} |u|^2 e^{-\varphi} d\sigma \leq 16 \int_{\mathbb{C}} \frac{|f|^2}{\Delta(e^\varphi \Delta e^{-\varphi})} e^{-\varphi} d\sigma.$$

As a consequence of theorem 1.1, we show that the operator  $\partial^k \bar{\partial}^k + c$  has a right inverse and that its inverse is bounded in  $L^2(\mathbb{C}, e^{-|z|^2})$ . For  $k = 1$  then  $4\partial^1 \bar{\partial}^1 = \Delta$  is the complex Laplacian and  $\Delta$  is the Laplacian in  $\mathbb{R}^2$ .

## 2. SOME LEMMAS

Consider the weighted Hilbert space

$$L^2(\mathbb{C}, e^{-\varphi}) = \{f \in L^2_{loc}(\mathbb{C}) : \int_{\mathbb{C}} |f|^2 e^{-\varphi} d\sigma < +\infty\}$$

where  $\varphi$  is a nonnegative function on  $\mathbb{C}$ .

We define the weighted inner product on  $L^2(\mathbb{C}, e^{-\varphi})$  by:  $\forall f, g \in L^2(\mathbb{C}, e^{-\varphi})$ , we have

$$\langle f, g \rangle_{\varphi} = \int_{\mathbb{C}} \bar{f} g e^{-\varphi} d\sigma$$

and the norm on  $L^2(\mathbb{C}, e^{-\varphi})$  by

$$\|f\|_{\varphi} = \sqrt{\langle f, f \rangle_{\varphi}} \quad \forall f \in L^2(\mathbb{C}, e^{-\varphi}).$$

We denote by  $C_0^\infty(\mathbb{C})$  the space of the functions  $\phi : \mathbb{C} \rightarrow \mathbb{C}$  of class  $C^\infty$  with compact support.

Let  $u, f \in L^2_{loc}(\mathbb{C})$ , we say that  $u$  is a weak solution of the equation  $\partial^k \bar{\partial}^k u = f$  if

$$\int_{\mathbb{C}} u \partial^k \bar{\partial}^k \phi d\sigma = \int_{\mathbb{C}} f \phi d\sigma.$$

Let  $\varphi$  be a nonnegative function of class  $C^\infty$  on  $\mathbb{C}$  and  $\phi \in C_0^\infty(\mathbb{C})$ , we define the formal adjoint of  $\partial^k \bar{\partial}^k$  with respect to the definition of weighted inner product in  $L^2(\mathbb{C}, e^{-\varphi})$  and with respect to the definition of the weak solution as :

$$\begin{aligned} \langle \phi, \partial^k \bar{\partial}^k u \rangle &= \int_{\mathbb{C}} \bar{\phi} \partial^k \bar{\partial}^k u e^{-\varphi} d\sigma \\ &= \int_{\mathbb{C}} \bar{\partial}^k \partial^k (\bar{\phi} e^{-\varphi}) u d\sigma \\ &= \int_{\mathbb{C}} e^\varphi \bar{\partial}^k \partial^k (\bar{\phi} e^{-\varphi}) u e^{-\varphi} d\sigma \\ &= \int_{\mathbb{C}} \overline{e^\varphi \partial^k \bar{\partial}^k (\phi e^{-\varphi})} u e^{-\varphi} d\sigma \\ &= \langle e^\varphi \partial^k \bar{\partial}^k (\phi e^{-\varphi}), u \rangle_{\varphi} \\ &=: \langle \bar{\partial}_\varphi^{k*} \partial_\varphi^{k*}(\phi), u \rangle_{\varphi} \end{aligned}$$

where  $\bar{\partial}_\varphi^{k*} \partial_\varphi^{k*}(\phi) = e^\varphi \partial^k \bar{\partial}^k(\phi e^{-\varphi})$  is the formal adjoint of  $\partial^k \bar{\partial}^k$  in  $C_0^\infty(\mathbb{C})$ . Let  $(\partial^k \bar{\partial}^k + c)_\varphi^*$  be the formal adjoint of  $\partial^k \bar{\partial}^k + c$  in  $C_0^\infty(\mathbb{C})$ . Let  $I_\varphi^* = I$  where  $I$  the identity operator, then

$$(\partial^k \bar{\partial}^k + c)_\varphi^* = \bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} + c.$$

We start by giving some lemmas in the general case of weighted spaces based on the functional analysis which will be very useful to us in the proof of the Theorem 1.1.

**Lemma 2.1.** *For each  $f \in L^2(\mathbb{C}, e^{-\varphi})$ , there exists a weak solution  $u \in L^2(\mathbb{C}, e^{-\varphi})$  of the equation*

$$\partial^k \bar{\partial}^k u + cu = f$$

with the norm estimate

$$\|u\|_\varphi^2 \leq a$$

if and only if

$$|\langle f, \phi \rangle_\varphi|^2 \leq a \|(\partial^k \bar{\partial}^k + c)_\varphi^* \phi\|_\varphi^2$$

$\forall \phi \in C_0^\infty(\mathbb{C})$  where  $a$  is a constant.

**Proof:** Let  $H = \partial^k \bar{\partial}^k + c$  and  $H_\varphi^* = (\partial^k \bar{\partial}^k + c)_\varphi^*$ . We suppose that there exists  $u \in L^2(\mathbb{C}, e^{-\varphi})$  such that

$$\partial^k \bar{\partial}^k u + cu = f$$

with the norm estimate

$$\|u\|_\varphi^2 \leq a,$$

then we have

$$\begin{aligned} |\langle f, \phi \rangle_\varphi|^2 &= |\langle Hu, \phi \rangle_\varphi|^2 = |\langle u, H_\varphi^* \phi \rangle_\varphi|^2 \\ &\leq \|u\|_\varphi^2 \|H_\varphi^* \phi\|_\varphi^2 \\ &\leq a \|H_\varphi^* \phi\|_\varphi^2 \end{aligned}$$

therefore

$$|\langle f, \phi \rangle_\varphi|^2 \leq a \|(\partial^k \bar{\partial}^k + c)_\varphi^* \phi\|_\varphi^2.$$

We suppose that

$$|\langle f, \phi \rangle_\varphi|^2 \leq a \|(\partial^k \bar{\partial}^k + c)_\varphi^* \phi\|_\varphi^2.$$

Consider the subspace

$$E = \{H_\varphi^* \phi : \phi \in C_0^\infty(\mathbb{C})\} \subset L^2(\mathbb{C}, e^{-\varphi}).$$

We define a linear map by

$$L_f : E \rightarrow \mathbb{C}$$

with

$$L_f(H_\varphi^* \phi) = \langle f, \phi \rangle_\varphi = \int_{\mathbb{C}} \bar{f} \phi e^{-\varphi} d\sigma$$

however

$$|L_f(H_\varphi^*\phi)| = | \langle f, \phi \rangle_\varphi | \leq \sqrt{a} \|H_\varphi^*\phi\|_\varphi$$

therefore  $L_f$  is bounded in  $E$ .

According to the Hahn-Banach extension theorem,  $L_f$  can be extended to an operator  $\tilde{L}_f$  on  $L^2(\mathbb{C}, e^{-\varphi})$  with

$$|\tilde{L}_f(g)| \leq \sqrt{a} \|g\|_\varphi \quad \forall g \in L^2(\mathbb{C}, e^{-\varphi}).$$

According to Riesz's representation theorem, there exists a unique  $u_0 \in L^2(\mathbb{C}, e^{-\varphi})$  such that

$$\tilde{L}_f(g) = \langle u_0, g \rangle_\varphi \quad \forall g \in L^2(\mathbb{C}, e^{-\varphi})$$

therefore

$$\tilde{L}_f(H_\varphi^*\phi) = \langle u_0, H_\varphi^*\phi \rangle_\varphi = \langle Hu_0, \phi \rangle_\varphi$$

however

$$\tilde{L}_f(H_\varphi^*\phi) = L_f(H_\varphi^*\phi) = \langle f, \phi \rangle_\varphi$$

in  $E$  therefore

$$\langle Hu_0, \phi \rangle_\varphi = \langle f, \phi \rangle_\varphi$$

so

$$Hu_0 = f$$

which implies

$$\partial^k \bar{\partial}^k u_0 + cu_0 = f$$

and

$$\|u_0\|_\varphi^2 = | \langle u_0, u_0 \rangle_\varphi | = |\tilde{L}_f(u_0)| \leq \sqrt{a} \|u_0\|_\varphi$$

therefore

$$\|u_0\|_\varphi^2 \leq a$$

however  $u_0 \in L^2(\mathbb{C}, e^{-\varphi})$  then let  $u_0 = u$ , so there exists a  $u \in L^2(\mathbb{C}, e^{-\varphi})$  such that

$$\partial^k \bar{\partial}^k u + cu = f$$

with

$$\|u\|_\varphi^2 \leq a.$$

□

**Lemma 2.2.**

$$\|(\partial^k \bar{\partial}^k + c)_\varphi^* \phi\|_\varphi^2 = \|(\partial^k \bar{\partial}^k + c)\phi\|_\varphi^2 + \langle \phi, \partial^k \bar{\partial}^k (\bar{\partial}^{k*} \partial^{k*} \phi) - \bar{\partial}^{k*} \partial^{k*} (\partial^k \bar{\partial}^k \phi) \rangle_\varphi$$

$\forall \phi \in C_0^\infty(\mathbb{C})$ .

**Proof:** Let  $H = \partial^k \bar{\partial}^k + c$  and  $H_\varphi^* = (\partial^k \bar{\partial}^k + c)_\varphi^*$ .  
For any  $\phi \in C_0^\infty(\mathbb{C})$

$$\begin{aligned} \|H_\varphi^* \phi\|_\varphi^2 &= \langle H_\varphi^* \phi, H_\varphi^* \phi \rangle_\varphi \\ &= \langle \phi, HH_\varphi^* \phi \rangle_\varphi \\ &= \langle \phi, H_\varphi^* H \phi + HH_\varphi^* \phi - H_\varphi^* H \phi \rangle_\varphi \\ &= \langle H \phi, H \phi \rangle + \langle \phi, HH_\varphi^* \phi - H_\varphi^* H \phi \rangle_\varphi \\ &= \|H \phi\|_\varphi^2 + \langle \phi, HH_\varphi^* \phi - H_\varphi^* H \phi \rangle_\varphi \end{aligned}$$

we have

$$\begin{aligned} HH_\varphi^* \phi &= (\partial^k \bar{\partial}^k + c)(\partial^k \bar{\partial}^k + c)_\varphi^* \phi \\ &= (\partial^k \bar{\partial}^k + c)(\bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} + c) \phi \\ &= \partial^k \bar{\partial}^k (\bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} \phi) + c \partial^k \bar{\partial}^k \phi + c \bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} \phi + c^2 \phi \end{aligned}$$

We have the same,

$$H_\varphi^* H \phi = \bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} (\partial^k \bar{\partial}^k \phi) + c \partial^k \bar{\partial}^k \phi + c \bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} \phi + c^2 \phi$$

therefore

$$HH_\varphi^* \phi - H_\varphi^* H \phi = \partial^k \bar{\partial}^k (\bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} \phi) - \bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} (\partial^k \bar{\partial}^k \phi)$$

so

$$\|H_\varphi^* \phi\|_\varphi^2 = \|H \phi\|_\varphi^2 + \langle \phi, \partial^k \bar{\partial}^k (\bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} \phi) - \bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} (\partial^k \bar{\partial}^k \phi) \rangle_\varphi$$

therefore

$$\|(\partial^k \bar{\partial}^k + c)_\varphi^* \phi\|_\varphi^2 = \|(\partial^k \bar{\partial}^k + c) \phi\|_\varphi^2 + \langle \phi, \partial^k \bar{\partial}^k (\bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} \phi) - \bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} (\partial^k \bar{\partial}^k \phi) \rangle_\varphi .$$

□

**Lemma 2.3.**

$$\begin{aligned} \langle \phi, \partial^k \bar{\partial}^k (\bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} \phi) - \bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} (\partial^k \bar{\partial}^k \phi) \rangle_\varphi &= \langle \phi, \sum_{i,j=1}^k \sum_{l,m=1}^k C_k^i C_k^j C_k^l C_k^m \times \\ &\quad (\partial^{k-m} \bar{\partial}^{k-l} \partial^{k-j} \bar{\partial}^{k-i} \phi) \partial^m \bar{\partial}^l (e^\varphi \partial^j \bar{\partial}^i e^{-\varphi}) \rangle \end{aligned}$$

For any  $\phi \in C_0^\infty(\mathbb{C})$ .

**Proof:**

$$\begin{aligned}
\bar{\partial}_\varphi^{k*} \partial_\varphi^{k*}(\phi) &= e^\varphi \partial^k \bar{\partial}^k (\phi e^{-\varphi}) \\
&= e^\varphi \partial^k \left( \sum_{i=0}^k C_k^i \bar{\partial}^{k-i} \phi \bar{\partial}^i e^{-\varphi} \right) \\
&= e^\varphi \sum_{i=0}^k C_k^i \partial^k (\bar{\partial}^{k-i} \phi \bar{\partial}^i e^{-\varphi}) \\
&= e^\varphi \sum_{i=0}^k C_k^i \sum_{j=0}^k C_k^j \partial^{k-j} \bar{\partial}^{k-i} \phi \bar{\partial}^j \bar{\partial}^i e^{-\varphi} \\
&= \sum_{i,j=0}^k C_k^i C_k^j \partial^{k-j} \bar{\partial}^{k-i} \phi (e^\varphi \bar{\partial}^j \bar{\partial}^i e^{-\varphi})
\end{aligned}$$

$$\begin{aligned}
\partial^k \bar{\partial}^k (\bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} \phi) &= \sum_{i,j=0}^k C_k^i C_k^j \partial^k \bar{\partial}^k [\partial^{k-j} \bar{\partial}^{k-i} \phi (e^\varphi \bar{\partial}^j \bar{\partial}^i e^{-\varphi})] \\
&= \sum_{i,j=0}^k C_k^i C_k^j \partial^k \sum_{l=0}^k C_k^l (\bar{\partial}^{k-l} \partial^{k-j} \bar{\partial}^{k-i} \phi) \bar{\partial}^l (e^\varphi \bar{\partial}^j \bar{\partial}^i e^{-\varphi}) \\
&= \sum_{i,j,l=0}^k C_k^i C_k^j C_k^l \partial^k [(\bar{\partial}^{k-l} \partial^{k-j} \bar{\partial}^{k-i} \phi) \bar{\partial}^l (e^\varphi \bar{\partial}^j \bar{\partial}^i e^{-\varphi})] \\
&= \sum_{i,j,l=0}^k C_k^i C_k^j C_k^l \sum_{m=0}^k C_k^m (\partial^{k-m} \bar{\partial}^{k-l} \partial^{k-j} \bar{\partial}^{k-i} \phi) \partial^m \bar{\partial}^l (e^\varphi \bar{\partial}^j \bar{\partial}^i e^{-\varphi}) \\
&= \sum_{i,j,l,m=0}^k C_k^i C_k^j C_k^l C_k^m (\partial^{k-m} \bar{\partial}^{k-l} \partial^{k-j} \bar{\partial}^{k-i} \phi) \partial^m \bar{\partial}^l (e^\varphi \bar{\partial}^j \bar{\partial}^i e^{-\varphi}).
\end{aligned}$$

We have the same

$$\bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} (\partial^k \bar{\partial}^k \phi) = \sum_{i,j=0}^k C_k^i C_k^j (\partial^{k-j} \bar{\partial}^{k-i} \partial^k \bar{\partial}^k \phi) (e^\varphi \bar{\partial}^j \bar{\partial}^i e^{-\varphi}).$$

then

$$\begin{aligned}
\partial^k \bar{\partial}^k (\bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} \phi) - \bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} (\partial^k \bar{\partial}^k \phi) &= \sum_{i,j=1}^k \sum_{l,m=1}^k C_k^i C_k^j C_k^l C_k^m \times \\
&\quad (\partial^{k-m} \bar{\partial}^{k-l} \partial^{k-j} \bar{\partial}^{k-i} \phi) \partial^m \bar{\partial}^l (e^\varphi \bar{\partial}^j \bar{\partial}^i e^{-\varphi})
\end{aligned}$$

therefore

$$\begin{aligned} \langle \phi, \partial^k \bar{\partial}^k (\bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} \phi) - \bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} (\partial^k \bar{\partial}^k \phi) \rangle &= \langle \phi, \sum_{i,j=1}^k \sum_{l,m=1}^k C_k^i C_k^j C_k^l C_k^m \times \\ &\quad (\partial^{k-m} \bar{\partial}^{k-l} \partial^{k-j} \bar{\partial}^{k-i} \phi) \partial^m \bar{\partial}^l (e^\varphi \partial^j \bar{\partial}^i e^{-\varphi}) \rangle. \end{aligned}$$

□

**Lemma 2.4.** *Let  $\varphi = |z|^2$  then*

$$\begin{aligned} \langle \phi, \partial^k \bar{\partial}^k (\bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} \phi) - \bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} (\partial^k \bar{\partial}^k \phi) \rangle_\varphi &= \langle \phi, \sum_{j=1}^k \sum_{l=1}^k \sum_{i=j-n}^k \sum_{n=l}^j \sum_{m=1}^{i-j+n} C_k^i C_k^j C_k^l C_k^m C_j^n \times \\ &\quad (\partial^{k-m} \bar{\partial}^{k-l} \partial^{k-j} \bar{\partial}^{k-i} \phi) \times \\ &\quad (-1)^{i+n} \frac{i!}{(i-j+n-m)!} \\ &\quad \frac{(n)!}{(n-l)!} (z)^{i-j+n-m} (\bar{z})^{n-l} \rangle \end{aligned}$$

For all  $\phi \in C_0^\infty(\mathbb{C})$ .

**Proof:** Let  $h(g) = e^g$  with  $g = -\varphi$ , according to the formula of Faà di Bruno [3], we have

$$\begin{aligned} \bar{\partial}^i e^g &= \bar{\partial}^i (h(g)) \\ &= \sum \frac{i!}{m_1! m_2! \cdots m_i!} h^{(m_1 + \cdots + m_i)}(g) \prod_{\gamma=1}^i \left( \frac{\bar{\partial}^\gamma g}{\gamma!} \right)^{m_\gamma} \\ &= \left( \sum \frac{i!}{m_1! m_2! \cdots m_i!} \prod_{\gamma=1}^i \left( \frac{-\bar{\partial}^\gamma \varphi}{\gamma!} \right)^{m_\gamma} \right) e^{-\varphi} \\ &=: P_i e^{-\varphi} \end{aligned}$$

Where the sum runs through all the  $i$ -tuples of nonnegative integers  $(m_1, m_2, \dots, m_i)$  satisfying the constraint :  $1m_1 + 2m_2 + \cdots + im_i = i$ . Let  $\varphi = |z|^2$ , we have

$$\begin{aligned} \bar{\partial}^\gamma \varphi &= \begin{cases} z & \text{if } \gamma = 1 \\ 0 & \text{if } \gamma \geq 2 \end{cases} \\ P_i &= (-\bar{\partial}^\gamma \varphi)^i = (-z)^i = (-1)^i (z)^i \\ \bar{\partial}^i e^{-|z|^2} &= (-1)^i (z)^i e^{-|z|^2} \end{aligned}$$

So

$$(2.1) \quad \partial^j \bar{\partial}^i e^{-|z|^2} = (-1)^i \partial^j (z)^i e^{-|z|^2}$$

$$(2.2) \quad \partial^j \bar{\partial}^i e^{-|z|^2} = (-1)^i \sum_{n=0}^j C_j^n \partial^{j-n} (z)^i \partial^n e^{-|z|^2}$$

By still applying Faà di Bruno's formula, we have

$$P_j = (-1)^j (z)^j$$

$$\partial^n e^{-|z|^2} = (-1)^n (\bar{z})^n e^{-|z|^2}$$

By replacing in 2.2, we have

$$(2.3) \quad \partial^j \bar{\partial}^i e^{-|z|^2} = (-1)^i \sum_{n=0}^j (-1)^n C_j^n \partial^{j-n} (z)^i (\bar{z})^n e^{-|z|^2}$$

however

$$\partial^{j-n} (z)^i = \begin{cases} 0 & \text{if } i < j - n \\ \frac{i!}{(i-j+n)!} (z)^{i-j+n} & \text{if } i \geq j - n \end{cases}$$

By replacing in 2.3, we have

$$\partial^j \bar{\partial}^i e^{-|z|^2} = \begin{cases} 0 & \text{if } i < j - n \\ \sum_{n=0}^j (-1)^{n+i} C_j^n \frac{i!}{(i-j+n)!} (z)^{i-j+n} (\bar{z})^n e^{-|z|^2} & \text{if } i \geq j - n \end{cases}$$

$$e^{|z|^2} \partial^j \bar{\partial}^i e^{-|z|^2} = \begin{cases} 0 & \text{if } i < j - n \\ \sum_{n=0}^j (-1)^{n+i} C_j^n \frac{i!}{(i-j+n)!} (z)^{i-j+n} (\bar{z})^n & \text{if } i \geq j - n \end{cases}$$

we have

$$\bar{\partial}^l (e^{|z|^2} \partial^j \bar{\partial}^i e^{-|z|^2}) = \sum_{n=0}^j (-1)^{n+i} C_j^n \frac{i!}{(i-j+n)!} (z)^{i-j+n} \bar{\partial}^l (\bar{z})^n \text{ if } i \geq j - n$$

$$\bar{\partial}^l (e^{|z|^2} \partial^j \bar{\partial}^i e^{-|z|^2}) = \begin{cases} 0 & \text{if } n < l \text{ and } i < j - n \\ \sum_{n=0}^j (-1)^{n+i} C_j^n \frac{i!}{(i-j+n)!} \frac{(n)!}{(n-l)!} (z)^{i-j+n} (\bar{z})^{n-l} & \text{if } i \geq j - n \text{ and } n \geq l \end{cases}$$

so

$$\partial^m \bar{\partial}^l (e^{|z|^2} \partial^j \bar{\partial}^i e^{-|z|^2}) = \begin{cases} 0 & \text{if } n < l, i - j + n < m \text{ and } i < j - n \\ \sum_{n=0}^j (-1)^{n+i} C_j^n \frac{i!}{(i-j+n)!} \frac{(n)!}{(n-l)!} \frac{(i-j+n)!}{(i-j+n-m)!} (z)^{i-j+n-m} (\bar{z})^{n-l} & \text{if } i \geq j - n, i - j + n \geq m \text{ and } n \geq l \end{cases}$$



$$\begin{aligned} & \langle \phi, \partial^k \bar{\partial}^k (\bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} \phi) - \bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} (\partial^k \bar{\partial}^k \phi) \rangle = \langle \phi, \\ & \sum_{j=1}^k \sum_{l=1}^k \sum_{i=j-n}^k \sum_{n=l}^j \sum_{m=1}^{i-j+n} C_k^i C_k^j C_k^l C_k^m C_j^m (\partial^{k-m} \bar{\partial}^{k-l} \partial^{k-j} \bar{\partial}^{k-i} \phi) (-1)^{i+n} \times \\ & \frac{i!}{(i-j+n-m)!} \frac{(n)!}{(n-l)!} (z)^{i-j+n-m} (\bar{z})^{n-l} \rangle. \end{aligned}$$

□

**Lemma 2.5.** Let  $\varphi = |z|^2$  then

$$\langle \phi, \partial^k \bar{\partial}^k (\bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} \phi) - \bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} (\partial^k \bar{\partial}^k \phi) \rangle_\varphi = \sum_{\alpha=0}^{k-1} \sum_{\beta=k-i+j-n}^{k-1} \frac{(k!)^4}{(\alpha!)^2 (\beta!)^2 ((k-\beta)! ((k-\alpha)!))} \times \|\bar{\partial}^\beta \partial^\alpha \phi\|_\varphi^2$$

**Proof:** According to the lemma 2.4, we have

$$\begin{aligned} & \langle \phi, \partial^k \bar{\partial}^k (\bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} \phi) - \bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} (\partial^k \bar{\partial}^k \phi) \rangle = \langle \phi, \\ & \sum_{j=1}^k \sum_{l=1}^k \sum_{i=j-n}^k \sum_{n=l}^j \sum_{m=1}^{i-j+n} C_k^i C_k^j C_k^l C_k^m C_j^m (\partial^{k-m} \bar{\partial}^{k-l} \partial^{k-j} \bar{\partial}^{k-i} \phi) (-1)^{i+n} \times \\ & \frac{i!}{(i-j+n-m)!} \frac{(n)!}{(n-l)!} (z)^{i-j+n-m} (\bar{z})^{n-l} \rangle \end{aligned}$$

Let  $s = n - l$

$$\begin{aligned} & \langle \phi, \partial^k \bar{\partial}^k (\bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} \phi) - \bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} (\partial^k \bar{\partial}^k \phi) \rangle = \langle \phi, \\ & \sum_{j=1}^k \sum_{l=1}^k \sum_{i=j-n}^k \sum_{m=1}^{i-j+n} A (\partial^{k-m} \bar{\partial}^{k-l} \partial^{k-j} \bar{\partial}^{k-i} \phi) \sum_{s=0}^{j-l} (-1)^{i+l-m+m+s} C_{j-l}^s \times \\ & \frac{(i-m)!}{(i-j+l+s-m)!} (z)^{i-j-m+l+s} (\bar{z})^s \rangle \end{aligned}$$

with

$$A = A(k, j, l, m, i) := \frac{(k!)^4}{m! l! (k-l)! (k-m)! (k-j)! (k-i)! (j-l)! (i-m)!}$$

$$\begin{aligned} & \langle \phi, \partial^k \bar{\partial}^k (\bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} \phi) - \bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} (\partial^k \bar{\partial}^k \phi) \rangle = \langle \phi, \\ & \sum_{j=1}^k \sum_{l=1}^k \sum_{i=j-n}^k \sum_{m=1}^{i-j+n} A (\partial^{k-m} \bar{\partial}^{k-l} \partial^{k-j} \bar{\partial}^{k-i} \phi) (-1)^{m+l} e^{|z|^2} \partial^{j-l} \bar{\partial}^{i-m} e^{-|z|^2} \rangle \end{aligned}$$

$$[\partial^k \bar{\partial}^k (\bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} \phi) - \bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} (\partial^k \bar{\partial}^k \phi)] e^{-|z|^2} =$$

$$\sum_{j=1}^k \sum_{l=1}^k \sum_{i=j-n}^k \sum_{m=1}^{i-j+n} A(\partial^{k-m} \bar{\partial}^{k-l} \partial^{k-j} \bar{\partial}^{k-i} \phi) (-1)^{m+l} \partial^{j-l} \bar{\partial}^{i-m} e^{-|z|^2}$$

**Remark 2.1.** *Since  $i \geq j - n$ ,  $i - j + n \geq m$  and  $n \geq l$  then  $j \geq l$  and  $i \geq m$ . If  $j < l$  or  $i < m$  then*

$$\partial^k \bar{\partial}^k (\bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} \phi) - \bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} (\partial^k \bar{\partial}^k \phi) e^{-|z|^2} = 0.$$

Let  $s = j - l$  and  $t = i - m$  then

$$[\partial^k \bar{\partial}^k (\bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} \phi) - \bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} (\partial^k \bar{\partial}^k \phi)] e^{-|z|^2} =$$

$$\sum_{l=1}^k \sum_{m=1}^{i-j+n} A' \sum_{s=0}^{k-l} \sum_{t=0}^{k-m} C_{k-m}^s C_{k-l}^t (\partial^{k-m} \bar{\partial}^{k-l} \partial^{k-l-s} \bar{\partial}^{k-m-t} \phi) (-1)^{m+l} \partial^s \bar{\partial}^t e^{-|z|^2}$$

with

$$A' = A'(k, l, m) := \frac{(k!)^4}{m! l! [(k-l)!]^2 [(k-m)!]^2}$$

$$[\partial^k \bar{\partial}^k (\bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} \phi) - \bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} (\partial^k \bar{\partial}^k \phi)] e^{-|z|^2} =$$

$$\sum_{l=1}^k \sum_{m=1}^{i-j+n} A' (-1)^{m+l} \partial^{k-m} \bar{\partial}^{k-l} (\partial^{k-l} \bar{\partial}^{k-m} \phi e^{-|z|^2})$$

By returning to the computation with respect of weighted inner product we have,

$$\begin{aligned}
\langle \phi, \partial^k \bar{\partial}^k (\bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} \phi) - \bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} (\partial^k \bar{\partial}^k \phi) \rangle_\varphi &= \int_{\mathbb{C}} \bar{\phi} \partial^k \bar{\partial}^k (\bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} \phi) e^{-|z|^2} d\sigma - \\
&\int_{\mathbb{C}} \bar{\phi} \bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} (\partial^k \bar{\partial}^k \phi) e^{-|z|^2} d\sigma \\
&= \sum_{m=1}^k \sum_{l=1}^{i-j+n} A' (-1)^{m+l} \times \\
&\int_{\mathbb{C}} \bar{\phi} \partial^{k-m} \bar{\partial}^{k-l} [(\partial^{k-l} \bar{\partial}^{k-m} \phi) e^{-|z|^2}] d\sigma \\
&= \sum_{m=1}^k \sum_{l=1}^{i-j+n} A' (-1)^{m+l} \times \\
&(-1)^{-m-l} \int_{\mathbb{C}} \bar{\partial}^{k-l} \partial^{k-m} \bar{\phi} (\partial^{k-l} \bar{\partial}^{k-m} \phi) e^{-|z|^2} d\sigma \\
&= \sum_{m=1}^k \sum_{l=1}^{i-j+n} A' \times \\
&\int_{\mathbb{C}} \overline{\partial^{k-l} \bar{\partial}^{k-m} \phi} (\partial^{k-l} \bar{\partial}^{k-m} \phi) e^{-|z|^2} d\sigma \\
&= \sum_{m=1}^k \sum_{l=1}^{i-j+n} A' \times \\
&\langle \partial^{k-l} \bar{\partial}^{k-m} \phi, \partial^{k-l} \bar{\partial}^{k-m} \phi \rangle_\varphi \\
&= \sum_{m=1}^k \sum_{l=1}^{i-j+n} A' \|\partial^{k-l} \bar{\partial}^{k-m} \phi\|_\varphi^2 \\
&= \sum_{m=1}^k \sum_{l=1}^{i-j+n} \frac{(k!)^4}{m! l! ((k-l)! ((k-m)!))^2} \times \\
&\|\partial^{k-l} \bar{\partial}^{k-m} \phi\|_\varphi^2
\end{aligned}$$

Let  $\alpha = k - l$  and  $\beta = k - m$

$$\langle \phi, \partial^k \bar{\partial}^k (\bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} \phi) - \bar{\partial}_\varphi^{k*} \partial_\varphi^{k*} (\partial^k \bar{\partial}^k \phi) \rangle_\varphi = \sum_{\alpha=0}^{k-1} \sum_{\beta=0}^{k-1} \frac{(k!)^4}{(\alpha!)^2 (\beta!)^2 ((k-\beta)! ((k-\alpha)!))} \|\partial^\alpha \bar{\partial}^\beta \phi\|_\varphi^2.$$

□

## 3. PROOF OF THE MAIN THEOREM

**Proof:**(Theorem 1.1) Let  $\varphi = |z|^2$  and  $\phi \in C_0^\infty(\mathbb{C})$ . According to the lemma 2.4

$$\|(\partial^k \bar{\partial}^k + c)_\varphi^* \phi\|_\varphi^2 \geq \langle \phi, \partial^k \bar{\partial}^k (\bar{\partial}^{k*} \partial^{k*} \phi) - \bar{\partial}^{k*} \partial^{k*} (\partial^k \bar{\partial}^k \phi) \rangle_\varphi.$$

According to the lemma 2.3

$$\langle \phi, \partial^k \bar{\partial}^k (\bar{\partial}^{k*} \partial^{k*} \phi) - \bar{\partial}^{k*} \partial^{k*} (\partial^k \bar{\partial}^k \phi) \rangle_\varphi = \sum_{\alpha=0}^{k-1} \sum_{\beta=0}^{k-j} \frac{(k!)^4}{(\alpha!)^2 (\beta!)^2 ((k-\beta)!) ((k-\alpha)!) \| \partial^\alpha \bar{\partial}^\beta \phi \|_\varphi^2}$$

therefore

$$\|(\partial^k \bar{\partial}^k + c)_\varphi^* \phi\|_\varphi^2 \geq (k!)^2 \|\phi\|_\varphi^2.$$

According to the Cauchy-Schwarz inequality, we have

$$\begin{aligned} |\langle f, \phi \rangle_\varphi|^2 &\leq \|f\|_\varphi^2 \|\phi\|_\varphi^2 \\ &= \left(\frac{1}{(k!)^2} \|f\|_\varphi^2\right) ((k!)^2 \|\phi\|_\varphi^2) \\ &\leq \left(\frac{1}{(k!)^2} \|f\|_\varphi^2\right) \|(\partial^k \bar{\partial}^k + c)_\varphi^* \phi\|_\varphi^2. \end{aligned}$$

According to the lemma 2.1, there exists a  $u \in L^2(\mathbb{C}, e^{-|z|^2})$  such that

$$\partial^k \bar{\partial}^k u + cu = f$$

with the norm estimate

$$\|u\|_\varphi^2 \leq \frac{1}{(k!)^2} \|f\|_\varphi^2$$

that is to say

$$\int_{\mathbb{C}} |u|^2 e^{-|z|^2} d\sigma \leq \frac{1}{(k!)^2} \int_{\mathbb{C}} |f|^2 e^{-|z|^2} d\sigma.$$

Moreover for  $k = 1$ , for all  $\phi \in C_0^\infty(\mathbb{C})$ , we have

$$\partial^1 \bar{\partial}^1 (\bar{\partial}_\varphi^{1*} \partial_\varphi^{1*} \phi) - \bar{\partial}_\varphi^{1*} \partial_\varphi^{1*} (\partial^1 \bar{\partial}^1 \phi) = \phi \partial^1 \bar{\partial}^1 (e^\varphi \partial^1 \bar{\partial}^1 e^{-\varphi}).$$

According to the lemma 2.4, we have

$$\begin{aligned} \|(\partial^1 \bar{\partial}^1 + c)_\varphi^* \phi\|_\varphi^2 &\geq \langle \phi, \partial^1 \bar{\partial}^1 (\bar{\partial}_\varphi^{1*} \partial_\varphi^{1*} \phi) - \bar{\partial}_\varphi^{1*} \partial_\varphi^{1*} (\partial^1 \bar{\partial}^1 \phi) \rangle_\varphi \\ &= \langle \phi, \phi \partial^1 \bar{\partial}^1 (e^\varphi \partial^1 \bar{\partial}^1 e^{-\varphi}) \rangle_\varphi \\ &= \langle \phi \sqrt{\partial^1 \bar{\partial}^1 (e^\varphi \partial^1 \bar{\partial}^1 e^{-\varphi})}, \phi \sqrt{\partial^1 \bar{\partial}^1 (e^\varphi \partial^1 \bar{\partial}^1 e^{-\varphi})} \rangle_\varphi \\ &= \left\| \phi \sqrt{\partial^1 \bar{\partial}^1 (e^\varphi \partial^1 \bar{\partial}^1 e^{-\varphi})} \right\|_\varphi^2 \end{aligned}$$

By applying the Cauchy-Schwarz inequality, we have

$$\begin{aligned} |\langle f, \phi \rangle_\varphi|^2 &= \left| \left\langle \frac{f}{\sqrt{\partial^1 \bar{\partial}^1 (e^\varphi \partial^1 \bar{\partial}^1 e^{-\varphi})}}, \phi \sqrt{\partial^1 \bar{\partial}^1 (e^\varphi \partial^1 \bar{\partial}^1 e^{-\varphi})} \right\rangle_\varphi \right|^2 \\ &\leq \left\| \frac{f}{\sqrt{\partial^1 \bar{\partial}^1 (e^\varphi \partial^1 \bar{\partial}^1 e^{-\varphi})}} \right\|_\varphi^2 \left\| \phi \sqrt{\partial^1 \bar{\partial}^1 (e^\varphi \partial^1 \bar{\partial}^1 e^{-\varphi})} \right\|_\varphi^2 \\ &\leq \left\| \frac{f}{\sqrt{\partial^1 \bar{\partial}^1 (e^\varphi \partial^1 \bar{\partial}^1 e^{-\varphi})}} \right\|_\varphi^2 \left\| (\partial^1 \bar{\partial}^1 + c)_\varphi^* \phi \right\|_\varphi^2 \end{aligned}$$

According to the lemma 2.1, there exists a weak solution  $u \in L^2(\mathbb{C}, e^{-\varphi})$  of the equation

$$\partial^1 \bar{\partial}^1 u + cu = f$$

with

$$\|u\|_\varphi^2 \leq \left\| \frac{f}{\sqrt{\partial^1 \bar{\partial}^1 (e^\varphi \partial^1 \bar{\partial}^1 e^{-\varphi})}} \right\|_\varphi^2$$

that is to say

$$\int_{\mathbb{C}} |u|^2 e^{-\varphi} d\sigma \leq 16 \int_{\mathbb{C}} \frac{|f|^2}{\Delta(e^\varphi \Delta e^{-\varphi})} e^{-\varphi} d\sigma.$$

□

**Theorem 3.1.** *There is a liner operator and bounded*

$$T_k : L^2(\mathbb{C}, e^{-|z|^2}) \longrightarrow L^2(\mathbb{C}, e^{-|z|^2})$$

such that

$$(\partial^k \bar{\partial}^k + c)T_k = I$$

with the norm estimate

$$\|T_k\|_\varphi \leq \frac{1}{(k!)^2}$$

where  $\|T_k\|_\varphi$  is the norm of  $T_k$  in  $L^2(\mathbb{C}, e^{-|z|^2})$ .

**Proof:** Let  $f \in L^2(\mathbb{C}, e^{-|z|^2})$ . According to the theorem 1.1, there is a weak solution  $u \in L^2(\mathbb{C}, e^{-|z|^2})$  such that

$$\partial^k \bar{\partial}^k u + cu = f$$

with the norm estimate

$$\|u\|_\varphi \leq \frac{1}{(k!)^2} \|f\|_\varphi$$

therefore

$$(\partial^k \bar{\partial}^k + c)T_k(f) = f$$

with

$$\|T_k(f)\|_\varphi \leq \frac{1}{(k!)^2} \|f\|_\varphi.$$

So

$$(\partial^k \bar{\partial}^k + c)T_k = I$$

with

$$\|T_k\|_\varphi \leq \frac{1}{(k!)^2}.$$

□

#### 4. SOME CONSEQUENCES

Let  $\lambda > 0$  and  $z_0 \in \mathbb{C}$ . Let  $\varphi = \lambda^2|z - z_0|^2$ , we obtain the corollary of the following Theorem 1.1 :

**Corollary 4.1.** *Let  $f \in L^2(\mathbb{C}, e^{-\lambda^2|z-z_0|^2})$ , there exists a weak solution  $u \in L^2(\mathbb{C}, e^{-\lambda^2|z-z_0|^2})$  of the equation*

$$\partial^k \bar{\partial}^k u + cu = f$$

with the norm estimate

$$\int_{\mathbb{C}} |u|^2 e^{-\lambda^2|z-z_0|^2} d\sigma \leq \frac{1}{(\lambda^k k!)^2} \int_{\mathbb{C}} |f|^2 e^{-\lambda^2|z-z_0|^2} d\sigma.$$

**Proof:** Let  $f \in L^2(\mathbb{C}, e^{-\lambda^2|z-z_0|^2})$ , we have

$$\int_{\mathbb{C}} |f|^2 e^{-\lambda^2|z-z_0|^2} d\sigma < +\infty.$$

Let  $z = \frac{w}{\lambda} + z_0$  and  $g(w) = f(z) = f(\frac{w}{\lambda} + z_0)$  then

$$\frac{1}{\lambda^2} \int_{\mathbb{C}} |g|^2 e^{-|w|^2} d\sigma(w) < +\infty$$

therefore  $g \in L^2(\mathbb{C}, e^{-|w|^2})$ . According to the Theorem 1.1, there exists a weak solution  $v \in L^2(\mathbb{C}, e^{-|w|^2})$  of the equation

$$\partial^k \bar{\partial}^k v + \frac{c}{\lambda^k} v = g$$

with the norm estimate

$$\int_{\mathbb{C}} |v|^2 e^{-|w|^2} d\sigma(w) \leq \frac{1}{(k!)^2} \int_{\mathbb{C}} |g|^2 e^{-|w|^2} d\sigma(w).$$

Let  $u(z) = \frac{1}{\lambda^k} v(w) = \frac{1}{\lambda^k} v(\lambda(z - z_0))$ .

Then

$$\partial^k \bar{\partial}^k u + cu = f$$

with the norm estimate

$$\int_{\mathbb{C}} |u|^2 e^{-\lambda^2|z-z_0|^2} d\sigma \leq \frac{1}{(\lambda^k k!)^2} \int_{\mathbb{C}} |f|^2 e^{-\lambda^2|z-z_0|^2} d\sigma.$$

□

As a consequence of the Corollary 2 we have

**Corollary 4.2.** *Let  $U \in \mathbb{C}$  be a bounded open set. Let  $f \in L^2(U)$ , there exists a weak solution  $u \in L^2(U)$  of the equation*

$$\partial^k \bar{\partial}^k u + cu = f$$

with

$$\|u\|_{L^2(U)} \leq \left( \frac{e^{|U|^2}}{(k!)^2} \right) \|f\|_{L^2(U)}$$

where  $|U|$  is the diameter of  $U$ .

**Proof:** Let  $z_0 \in U$  and  $f \in L^2(U)$ , we have

$$\tilde{f}(z) = \begin{cases} f(z) & \text{si } z \in U \\ 0 & \text{si } z \in \mathbb{C} \setminus U \end{cases}$$

Therefore  $\tilde{f} \in L^2(\mathbb{C}) \subset L^2(\mathbb{C}, e^{-|z-z_0|^2})$ . According to the Corollary 2, there exists a weak solution  $\tilde{u} \in L^2(\mathbb{C}, e^{-|z-z_0|^2})$  of the equation

$$\partial^k \bar{\partial}^k \tilde{u} + c\tilde{u} = \tilde{f}$$

with

$$\int_{\mathbb{C}} |\tilde{u}|^2 e^{-|z-z_0|^2} d\sigma \leq \frac{1}{(k!)^2} \int_{\mathbb{C}} |\tilde{f}|^2 e^{-|z-z_0|^2} d\sigma$$

therefore

$$\int_{\mathbb{C}} |\tilde{u}|^2 e^{-|z-z_0|^2} d\sigma \leq \frac{1}{(k!)^2} \int_U |f|^2 d\sigma$$

however

$$\int_{\mathbb{C}} |\tilde{u}|^2 e^{-|z-z_0|^2} d\sigma \geq e^{-|U|^2} \int_U |\tilde{u}|^2 d\sigma$$

therefore

$$\int_U |\tilde{u}|^2 d\sigma \leq \left( \frac{e^{|U|^2}}{(k!)^2} \right) \int_U |f|^2 d\sigma.$$

Let  $\tilde{u}|_U = u$  then we have

$$\partial^k \bar{\partial}^k u + cu = f$$

with

$$\|u\|_{L^2(U)} \leq \left( \frac{e^{|U|^2}}{(k!)^2} \right) \|f\|_{L^2(U)}.$$

□

#### ACKNOWLEDGMENTS

The authors would like to thank the director of Ziguinchor Polytechnic Institute (ZIP) and all the members of the SAGELAB laboratory at ZIP where we stayed to do this work.

REFERENCES

- [1] S. Dai, Y. Pan, A right inverse of differential operator  $\frac{d^k}{dx^k} + a$  in Weighted Hilbert space  $L^2(\mathbb{R}, e^{-x^2})$ , J. Math. Anal. Appl. 486 (2020) 123890.
- [2] S. Dai, Y. Pan, A right inverse of Cauchy-Riemann operator  $\bar{\partial}^k + a$  in Weighted Hilbert space  $L^2(\mathbb{C}, e^{-|z|^2})$ , a paraitre aux Annales de la Faculté des Sciences de Toulouse.
- [3] W. P. Johnson, The curious history of Faà di Bruno's formula, Am. Math. Man. 109 (3) (2002), 217 – 237.

MATHEMATICS DEPARTMENT, UFR OF SCIENCE AND TECHNOLOGY, ASSANE SECK UNIVERSITY OF ZIGUINCHOR, BP: 523 (SENEGAL)  
*Email address:* m.bodian@univ-zig.sn

MATHEMATICS DEPARTMENT, UFR OF SCIENCE AND TECHNOLOGY, ASSANE SECK UNIVERSITY OF ZIGUINCHOR, BP: 523 (SENEGAL)  
*Email address:* s.sambou1440@zig.univ.sn

MATHEMATICS DEPARTMENT, UFR OF SCIENCE AND TECHNOLOGY, ASSANE SECK UNIVERSITY OF ZIGUINCHOR, BP: 523 (SENEGAL)  
*Email address:* p.badiane4963@zig.univ.sn

MATHEMATICS DEPARTMENT, FACULTY OF SCIENCE AND TECHNOLOGY, MARIEN NGOUABI UNIVERSITY, (CONGO)  
*Email address:* wnossete@gmail.com

MATHEMATICS DEPARTMENT, UFR OF SCIENCE AND TECHNOLOGY, ASSANE SECK UNIVERSITY OF ZIGUINCHOR, BP: 523 (SENEGAL)  
*Email address:* ssambou@univ-zig.sn