

Multiple solutions of a nonlinear biharmonic equation on graphs

Songbo Hou*

Department of Applied Mathematics, College of Science, China Agricultural University, Beijing, 100083, P.R. China

Abstract

In this paper, we consider a biharmonic equation with respect to the Dirichlet problem on a domain of a locally finite graph. Using the variation method, we prove that the equation has two distinct solutions under certain conditions.

Key words: Locally finite graph, Biharmonic equation, Distinct solutions.

2010 MSC: 35A15, 35G30

1. Introduction

The existence and non-existence of solutions of boundary value problems for biharmonic equations has been studied by many authors. A lot of results are devoted to the following problem in $H_0^2(\Omega)$,

$$\begin{cases} \Delta^2 u = \lambda u + |u|^{p-2}u, & x \in \Omega \\ u|_{\partial\Omega} = 0, \quad \frac{\partial u}{\partial n}|_{\partial\Omega} = 0, \end{cases} \quad (1.1)$$

where Ω is a bounded smooth domain in \mathbb{R}^N ; λ is a constant; $p = \frac{2N}{N-4}$. If $\lambda < 0$ and Ω is star-shaped, then the problem (1.1) has only trivial solution. Denote by λ_1 the first eigenvalue of the problem

$$\begin{cases} \Delta^2 u = \lambda u, & x \in \Omega \\ u|_{\partial\Omega} = 0, \quad \frac{\partial u}{\partial n}|_{\partial\Omega} = 0. \end{cases}$$

If $0 < \lambda < \lambda_1$ and $N \geq 8$, then Problem (1.1) has at least one nontrivial solution [4]. For $5 \leq n \leq 7$, there exists a constant $\bar{\lambda} > 0$ such that Problem (1.1) has a non-trivial solution for all $\lambda \in (\bar{\lambda}, \lambda_1)$. Deng and Wang [3] studied the existence and non-existence of multiple solutions of biharmonic equations boundary value problem,

$$\begin{cases} \Delta^2 u = \lambda u + |u|^{p-2}u + f(x), & x \in \Omega, \\ u|_{\partial\Omega} = 0, \quad \frac{\partial u}{\partial n}|_{\partial\Omega} = 0. \end{cases}$$

*Corresponding author: Songbo Hou

Email address: housb@cau.edu.cn (Songbo Hou)

Preprint submitted to ***

May 17, 2022

We refer the reader to [1, 2, 13, 14] for more related results.

In this paper, we will study the similar problem on a graph. A discrete graph is denoted by $G = (V, E)$, where V is the vertex set and E is the edge set. We say that G is locally finite if for any $x \in V$, there are only finite $y \in V$ such that $xy \in E$. A graph is called connected if for any $x, y \in V$, they can be connected via finite edges. Throughout this paper, we assume that G is locally finite and connected. Let ω_{xy} be the weight of an edge $xy \in E$ such that $\omega_{xy} > 0$ and $\omega_{xy} = \omega_{yx}$. We use a positive function $\mu : V \rightarrow \mathbb{R}^+$ to define a measure on G . For a bounded domain $\Omega \subset V$, the boundary of Ω is defined as

$$\partial\Omega := \{y \notin \Omega : \exists x \in \Omega \text{ such that } xy \in E\}.$$

We introduce some notations about the partial differential equations on a graph. The μ -Laplacian of a function $u : V \rightarrow \mathbb{R}$ is defined by

$$\Delta u(x) := \frac{1}{\mu(x)} \sum_{y \sim x} \omega_{xy}(u(y) - u(x)),$$

where $y \sim x$ stands for $xy \in E$. For any two functions u and v on the graph, the gradient form is defined by

$$\Gamma(u, v) = \frac{1}{2\mu(x)} \sum_{y \sim x} \omega_{xy}(u(y) - u(x))(v(y) - v(x)).$$

If $u = v$, we write $\Gamma(u) = \Gamma(u, u)$, which is used to define the length of the gradient for u

$$|\nabla u|(x) = \sqrt{\Gamma(u)(x)} = \left(\frac{1}{2\mu(x)} \sum_{y \sim x} \omega_{xy}(u(y) - u(x))^2 \right)^{1/2}.$$

For a function u over V , the integral is defined by

$$\int_V u d\mu = \sum_{x \in V} \mu(x)u(x).$$

Recently, Grigor'yan, Lin and Yang [6, 7] applied the mountain-pass theorem to establish existence results for some nonlinear equations. Zhang and Zhao [15] studied the convergence of ground state solutions for nonlinear Schrödinger equations via the Nehari method on graphs. Using the similar method, Han, Shao and Zhao [9] studied existence and convergence of solutions for nonlinear biharmonic equations on graphs. One may refer to [10] for the results about the heat equation and refer to [5, 8, 11] for the results about the Kazdan-Warner equation on a graph.

In this paper, we consider the following equation

$$\begin{cases} \Delta^2 u = \lambda u + |u|^{p-2}u + \epsilon f, & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega, \end{cases} \quad (1.2)$$

where Ω is a bounded domain of V ; λ , p and ϵ are positive constants, $p > 2$; f is a given function on V .

Let $W^{2,2}(\Omega)$ be the space of functions $u : V \rightarrow \mathbb{R}$ under the norm

$$\|u\|_{W^{2,2}(\Omega)} = \left(\int_{\Omega \cup \partial\Omega} (|\Delta u|^2 + |\nabla u|^2) d\mu + \int_{\Omega} u^2 d\mu \right)^{\frac{1}{2}}. \quad (1.3)$$

Let $H(\Omega) = W^{2,2}(\Omega) \cap W_0^{1,2}(\Omega)$, where $W_0^{1,2}(\Omega)$ is the completion of $C_c(\Omega)$ under the norm

$$\|u\|_{W_0^{1,2}(\Omega)} = \left(\int_{\Omega \cup \partial\Omega} |\nabla u|^2 d\mu + \int_{\Omega} u^2 d\mu \right)^{\frac{1}{2}}.$$

Define

$$\|u\|_H = \left(\int_{\Omega \cup \partial\Omega} |\Delta u|^2 d\mu \right)^{\frac{1}{2}}$$

for any $u \in H$. It is easy to see that $\|\cdot\|_H$ is a norm on H . Noting that the dimension of $W^{2,2}(\Omega)$ is finite, we have that $\|u\|_H = \left(\int_{\Omega \cup \partial\Omega} |\Delta u|^2 d\mu \right)^{\frac{1}{2}}$ is a norm equivalent to (1.3) on $H(\Omega)$. Lemma 2.6 in [9] implies that $H(\Omega)$ is embedded in $L^q(\Omega)$ for all $1 \leq q < +\infty$ and there is a constant depending only on q and Ω such that

$$\left(\int_{\Omega} |u|^q d\mu \right)^{1/q} \leq C \left(\int_{\Omega \cup \partial\Omega} |\Delta u|^2 d\mu \right)^{\frac{1}{2}}. \quad (1.4)$$

Definition 1.1. For a $u \in H$, if for any $\varphi \in C_c(\Omega)$, there holds that

$$\int_{\Omega \cup \partial\Omega} \Delta u \Delta \varphi d\mu = \lambda \int_{\Omega} u \varphi d\mu + \int_{\Omega} |u|^{p-2} u \varphi d\mu + \epsilon \int_{\Omega} f \varphi d\mu,$$

then u is called a weak solution of (1.2).

Define

$$J_{\epsilon}(u) = \frac{1}{2} \int_{\Omega \cup \partial\Omega} |\Delta u|^2 d\mu - \frac{\lambda}{2} \int_{\Omega} \lambda u^2 d\mu - \frac{1}{p} \int_{\Omega} |u|^p d\mu - \epsilon \int_{\Omega} f(x) u d\mu,$$

which will be used in the later variation procedure. Define

$$\lambda_1(\Omega) = \inf_{u \neq 0, u|_{\partial\Omega} = 0} \frac{\int_{\Omega \cup \partial\Omega} |\Delta u|^2 d\mu}{\int_{\Omega} u^2 d\mu}.$$

Using the variation method similar to that in [6, 7], we prove the following theorem.

Theorem 1.1. Let $G = (V, E)$ be a locally finite graph. Suppose that $0 < \lambda < \lambda_1(\Omega)$, $f \in H'(\Omega)$ where H' is the dual space of H . Then there exists $\epsilon_1 > 0$ such that (1.2) has two distinct solutions if $0 < \epsilon < \epsilon_1$.

2. Proof of the main results

Lemma 2.1. There exist positive constants r_{ϵ} and δ_{ϵ} such that $J_{\epsilon} \geq \delta_{\epsilon}$ for all $u \in H$ with $r_{\epsilon} \leq \|u\|_H \leq 2r_{\epsilon}$ if $0 < \epsilon < \epsilon_1$ for a sufficiently small ϵ_1 .

Proof.

$$\begin{aligned} J_{\epsilon}(u) &\geq \frac{1}{2} \|u\|_H^2 - \frac{\lambda}{2} \int_{\Omega} |u|^2 d\mu - \frac{C}{p} \|u\|_H^p - \epsilon \|f\|_{H'} \|u\|_H \\ &\geq \frac{\tau}{2} \|u\|_H^2 - \frac{C}{p} \|u\|_H^p - \epsilon \|f\|_{H'} \|u\|_H \\ &\geq \|u\|_H \left(\frac{\tau}{2} \|u\|_H - \frac{C}{2} \|u\|_H^{p-1} - \epsilon \|f\|_{H'} \right), \end{aligned} \quad (2.1)$$

where $\tau = \frac{\lambda_1(\Omega) - \lambda}{\lambda_1(\Omega)}$. Take $r_\epsilon = \sqrt{\epsilon}$. Since

$$\lim_{\epsilon \rightarrow 0^+} \frac{\frac{\tau}{2} \sqrt{\epsilon} - 2^{p-2} C \epsilon^{(p-1)/2} - \epsilon \|f\|_{H'}}{\frac{\tau}{2} \sqrt{\epsilon}} = 1,$$

there exists some ϵ_1 such that

$$\frac{\tau}{2} \sqrt{\epsilon} - 2^{p-2} C \epsilon^{(p-1)/2} - \epsilon \|f\|_{H'} \geq \frac{\tau}{4} \sqrt{\epsilon}$$

if $0 < \epsilon < \epsilon_1$.

Setting $\delta_\epsilon = \frac{\tau \epsilon}{4}$, we obtain $J_\epsilon(u) \geq \delta_\epsilon$ if $0 < \epsilon < \epsilon_1$. \square

Lemma 2.2. *For any $c \in \mathbb{R}$, J_ϵ satisfies the $(PS)_c$ condition. If $(u_k) \subset H$ is a sequence such that $J_\epsilon(u_k) \rightarrow c$ and $J'_\epsilon(u_k) \rightarrow 0$, then up to a subsequence, u_k converges to some u in H .*

Proof. If $J_\epsilon(u_k) \rightarrow c$ and $J'_\epsilon(u_k) \rightarrow 0$, then we have

$$\frac{1}{2} \int_{\Omega \cup \partial\Omega} |\Delta u_k|^2 d\mu - \frac{1}{2} \int_{\Omega} \lambda u_k^2 d\mu - \frac{1}{p} \int_{\Omega} |u_k|^p d\mu - \epsilon \int_{\Omega} f(x) u_k d\mu = c + o_k(1), \quad (2.2)$$

$$\left| \int_{\Omega \cup \partial\Omega} |\Delta u_k|^2 d\mu - \int_{\Omega} \lambda u_k^2 d\mu - \int_{\Omega} |u_k|^p d\mu - \epsilon \int_{\Omega} f(x) u_k d\mu \right| = o_k(1) \|u_k\|_H, \quad (2.3)$$

where $o_k(1) \rightarrow 0$ as $k \rightarrow +\infty$.

From (2.2) and (2.3), we have

$$\left(\frac{1}{2} - \frac{1}{p} \right) \int_{\Omega} |u_k|^p d\mu = c + \frac{\epsilon}{2} \int_{\Omega} f u_k d\mu + o_k(1) \|u_k\|_H + o_k(1). \quad (2.4)$$

Hence

$$\begin{aligned} \tau \|u_k\|_H^2 &\leq \int_{\Omega \cup \partial\Omega} |\Delta u_k|^2 d\mu - \int_{\Omega} \lambda u_k^2 d\mu \\ &\leq \frac{2pc}{p-2} + \frac{(2p-2)\epsilon}{p-2} \|f\|_{H'} \|u_k\|_H + o_k(1) \|u_k\|_H + o_k(1) \\ &\leq \frac{2pc}{p-2} + \frac{4(p-1)^2 \epsilon^2}{(p-2)^2 \tau} \|f\|_{H'}^2 + \frac{\tau}{4} \|u_k\|_H^2 + \frac{\tau}{4} \|u_k\|_H^2 + o_k(1). \end{aligned} \quad (2.5)$$

Thus we have (u_k) is bounded in H . Since H is pre-compact, it follows that up to a subsequence, u_k converges to some u in H . \square

Now we arrive at a position to prove the main theorem. For any $u^* \in H$, passing to the limit $t \rightarrow +\infty$, we get

$$J_\epsilon(tu^*) = \frac{t^2}{2} \int_{\Omega \cup \partial\Omega} |\Delta u^*|^2 d\mu - \frac{t^2}{2} \int_{\Omega} \lambda u^{*2} d\mu - \frac{t^p}{p} \int_{\Omega} |u^*|^p d\mu - t \epsilon \int_{\Omega} f(x) u^* d\mu \rightarrow -\infty$$

as $t \rightarrow \infty$. Hence there exists some $\tilde{u} \in H$ such that $J_\epsilon(\tilde{u}) < 0$ with $\|\tilde{u}\|_H > r_\epsilon$. Combining Lemma 2.1, we see that J_ϵ satisfies all the hypotheses of the mountain-pass theorem: $J_\epsilon \in C^1(H, \mathbb{R})$; $J_\epsilon(0) = 0$; when $\|u\|_H = r_\epsilon$, $J_\epsilon(u) \geq \delta_\epsilon$; $J_\epsilon(\tilde{u}) < 0$ for some \tilde{u} with $\|\tilde{u}\|_H > r_\epsilon$.

Then we have

$$c = \min_{\gamma \in \Gamma} \max_{u \in \gamma} J_\epsilon(u)$$

is the critical point of J_ϵ , where

$$\Gamma = \{\gamma \in C([0, 1], H) : \gamma(0) = 0, \gamma(1) = \bar{u}\}.$$

Thus there exists a weak solution $u_c \in H$ with $J_\epsilon(u_c) \geq \delta_\epsilon$.

Lemma 2.3. *There exists τ_0 and $u^* \in H$ with $\|u^*\|_H = 1$ such that $J_\epsilon(tu^*) < 0$ if $0 < t < \tau_0$.*

Proof. We study the equation

$$\Delta^2 u = \lambda u + f \tag{2.6}$$

in $H(\Omega)$. Define the functional

$$J_f(u) = \frac{1}{2} \int_{\Omega \cup \partial\Omega} |\Delta u|^2 d\mu - \frac{1}{2} \int_{\Omega} \lambda u^2 d\mu - \int_{\Omega} f(x) u d\mu.$$

Noting that

$$\frac{1}{2} \int_{\Omega \cup \partial\Omega} |\Delta u|^2 d\mu - \frac{1}{2} \int_{\Omega} \lambda u^2 d\mu \geq \frac{\tau}{2} \|u\|_H^2 \tag{2.7}$$

and

$$\left| \int_{\Omega} f(x) u d\mu \right| \leq \|f\|_{H'} \|u\|_H \leq \frac{\tau}{4} \|u\|_H^2 + \frac{1}{\tau} \|f\|_{H'}^2, \tag{2.8}$$

we have that J_f have a lower bound on H .

Set

$$m_f = \inf_{u \in H} J_f(u).$$

There exists $u_k \in H$ such that $J_f(u_k) \rightarrow m_f$. From (2.7) and (2.8), we know u_k is bounded in H .

Hence there exists $u^* \in H$ such that $u_k \rightharpoonup u^*$ weakly in H . Then

$$J_f(\bar{u}) \leq \liminf_{k \rightarrow \infty} J_f(u_k) = m_f$$

and \bar{u} is the weak solution of (2.6). It follows that

$$\int_{\Omega} f \bar{u} d\mu = \int_{\Omega \cup \partial\Omega} |\Delta \bar{u}|^2 d\mu - \lambda \int_{\Omega} (\bar{u})^2 d\mu > 0 \tag{2.9}$$

Now we compute the derivative of $J_\epsilon(t\bar{u})$:

$$\frac{d}{dt} J_\epsilon(t\bar{u}) = t \int_{\Omega \cup \partial\Omega} |\Delta \bar{u}|^2 d\mu - t \int_{\Omega} \lambda (\bar{u})^2 d\mu - t^{p-1} \int_{\Omega} |\bar{u}|^p d\mu - \epsilon \int_{\Omega} f(x) \bar{u} d\mu.$$

By (2.9), we obtain

$$\left. \frac{d}{dt} J_\epsilon(t\bar{u}) \right|_{t=0} < 0.$$

Letting $u^* = \frac{\bar{u}}{\|\bar{u}\|_H}$, we finish the proof. \square

Lemma 2.4. Choose ϵ such that $0 < \epsilon < \epsilon_1$ where ϵ_1 is the same as in Lemma 2.1. Then there exists a function $u_0 \in H$ with $\|u_0\|_H \leq 2r_\epsilon$ such that

$$J_\epsilon(u_0) = c_\epsilon = \inf_{\|u\| \leq 2r_\epsilon} J_\epsilon(u),$$

where $r_\epsilon = \sqrt{\epsilon}$, $c_\epsilon < 0$.

Proof. By (2.1), we see that J_ϵ has a lower bound on

$$B_{2r_\epsilon} = \{u \in H : \|u\|_H \leq 2r_\epsilon\}.$$

Combining Lemma 2.3, we get $c_\epsilon < 0$.

Let $(u_k) \subset H$ be a sequence satisfying $\|u_k\|_H \leq 2r_\epsilon$ and $J_\epsilon(u_k) \rightarrow c_\epsilon$. Then up to a subsequence, u_k converges weakly to u_0 in H and converges strongly to u_0 in $L^q(V)$ for any $1 \leq q \leq +\infty$. It follows that

$$\begin{aligned} \lim_{k \rightarrow +\infty} \int_{\Omega} f u_k d\mu &= \int_{\Omega} f u_0 d\mu, \\ \|u_0\|_H &\leq \limsup_{k \rightarrow +\infty} \|u_k\|_H \leq 2r_\epsilon, \end{aligned}$$

$$\lim_{k \rightarrow +\infty} \int_{\Omega} \lambda u_k^2 d\mu = \int_{\Omega} \lambda u_0^2 d\mu,$$

$$\lim_{k \rightarrow +\infty} \int_{\Omega} u_k^p d\mu = \int_{\Omega} u_0^p d\mu.$$

Hence

$$J_\epsilon(u_0) \leq \limsup_{k \rightarrow +\infty} J_\epsilon(u_k) = c_\epsilon$$

and u_0 is the minimizer of J_ϵ on B_{2r_ϵ} . Lemma 2.1 implies that $\|u_0\|_H < r_\epsilon$. For any $\varphi \in C_c(V)$, let $\psi(t) = J_\epsilon(u_0 + t\varphi)$. Then ψ is a smooth function in t . It is easy to see that there is $\eta > 0$ such that $u_0 + t\varphi \in B_{2r_\epsilon}$ if $|t| < \eta$. This means that $\psi(0)$ is the minimum of $\psi(t)$ on $(-\eta, \eta)$. By $\psi'(0) = 0$, we get

$$\int_{\Omega \cup \partial\Omega} \Delta u_0 \Delta \varphi d\mu - \lambda \int_{\Omega} u_0 \varphi d\mu - \int_{\Omega} |u_0|^{p-1} u_0 \varphi d\mu - \epsilon \int_{\Omega} f \varphi d\mu = 0.$$

We conclude that u_0 is a weak solution of (1.2) and complete the proof. \square

Clearly, u_c and u_0 are two distinct solutions of (1.2) since $J_\epsilon(u_c) > 0$ and $J_\epsilon(u_0) < 0$.

Acknowledgement

This work is partially supported by the National Natural Science Foundation of China (Grant No. 11721101), and by National Key Research and Development Project SQ2020YFA070080.

References

- [1] Claudianor O. Alves, Giovany M. Figueiredo, Multiplicity of nontrivial solutions to a biharmonic equation via Lusternik-Schnirelman theory, *Math. Methods Appl. Sci.* 36(6) (2013) 683-694.
- [2] Daomin Cao, Wei Dai, Classification of nonnegative solutions to a bi-harmonic equation with Hartree type nonlinearity, *Proc. Roy. Soc. Edinburgh Sect. A* 149 (4) (2019) 979-994.
- [3] Yinbin Deng, Cengsheng Wang, On inhomogeneous biharmonic equations involving critical exponents, *Proc. Roy. Soc. Edin.* 129 A (1999) 925-946.
- [4] D. E. Edmunds, D. Fortunato, E. Jannelli, Critical exponents, critical dimensions and the biharmonic operator, *Arch. Rational Mech. Anal.* 112(3) (1990) 269-289.
- [5] Huabin Ge, Wenfeng Jiang, Kazdan-Warner equation on infinite graphs, *J. Korean Math. Soc.* 55(5) (2018) 1091-1101.
- [6] Alexander Grigor'yan, Yong Lin, Yunyan Yang, Yamabe type equations on graphs, *J. Differential Equations* 261 (2016) 4924-4943.
- [7] Alexander Grigor'yan, Yong Lin, Yunyan Yang, Existence of positive solutions to some nonlinear equations on locally finite graphs, *Science China Mathematics* 60(7) (2017) 1311-1324.
- [8] Alexander Grigor'yan, Yong Lin, Yunyan Yang, Kazdan-Warner equation on graph, *Calc. Var. Partial Differential Equations* 55(4) (2016) Art. 92 13 pp.
- [9] Xiaoli Han, Mengqiu Shao, Liang Zhao, Existence and convergence of solutions for nonlinear biharmonic equations on graphs, *J. Differential Equations* 268 (2020) 3936-3961.
- [10] Yong Lin, Yiting Wu, The existence and nonexistence of global solutions for a semilinear heat equation on graphs, *Calc. Var. Partial Differential Equations* 56(4) (2017) Paper No.102 22 pp.
- [11] Shuang Liu, Yunyan Yang, Multiple solutions of Kazdan-Warner equation on graphs in the negative case, *Calc. Var. Partial Differential Equations* 59(5) (2020) 164.
- [12] Patrizia Pucci, James Serrin, A general variational identity, *Indiana Univ. Math. J.* 35(3) (1986) 681-703.
- [13] Youjun Wang, Yaotian She, Multiple and sign-changing solutions for a class of semilinear biharmonic equation, *J. Differential Equations* 246 (2009) 3109-3125.
- [14] Weihua Wang, Peihao Zhao, Nonuniformly nonlinear elliptic equations of p-biharmonic type, *J. Math. Anal. Appl.* 348 (2008) 730-738.
- [15] Ning Zhang, Liang Zhao, Convergence of ground state solutions for nonlinear Schrödinger equations on graphs, *Sci. China Math.* 61(8) (2018) 1481-1494.