

Sharp Hessian estimates for fully nonlinear elliptic equations under relaxed convexity assumptions, oblique boundary conditions and applications

by

Junior da S. Bessa*, João Vitor da Silva†, Maria N.B. Frederico‡

&

Gleydson C. Ricarte §

May 17, 2022

Abstract

In this work we derive global $W^{2,p}$ estimates (with $n \leq p < \infty$) for viscosity solutions to fully nonlinear elliptic equations under relaxed structural assumptions on the governing operator which are weaker than convexity and oblique boundary conditions as follows

$$\begin{cases} F(D^2u, Du, u, x) = f(x) & \text{in } \Omega \\ \beta(x) \cdot Du + \gamma(x)u(x) = g(x) & \text{on } \partial\Omega, \end{cases}$$

for $f \in L^p(\Omega)$ and under suitable assumptions on the data β, γ, g and $\Omega \subset \mathbb{R}^n$. Our approach makes use of geometric tangential methods, which consists of importing “fine regularity estimates” from a limiting profile, i.e., the *Recession* operator, associated with the original second order one via compactness and stability procedures. As a result, we devote a special attention to the borderline scenario, i.e., $f \in \text{BMO}_p \supseteq L^\infty$. In such a setting, we prove that solutions enjoy BMO_p type estimates for their second derivatives. In the end, as another application of our findings, we obtain Hessian estimates to obstacle type problems under oblique boundary conditions and no convexity assumptions, which may have their own mathematical interest. A density result in a suitable class of viscosity solutions will be also addressed.

Keywords: Hessian estimates, fully nonlinear elliptic equations, oblique boundary conditions, relaxed convexity assumptions, obstacle type problems.

AMS Subject Classification: 35J25, 35J60, 35B65, 35R35.

*Universidade Federal do Ceará. Department of Mathematics. Fortaleza - CE, Brazil. E-mail address: junior.bessa@alu.ufc.br

†Universidade Estadual de Campinas - UNICAMP. Department of Mathematics. Campinas - SP, Brazil. E-mail address: jdasilva@unicamp.br

‡Universidade Federal do Ceará. Department of Mathematics. Fortaleza - CE, Brazil. E-mail address: nildebarreto@gmail.com

§Universidade Federal Ceará. Department of Mathematics. Fortaleza, CE-Brazil 60455-760. E-mail address: ricarte@mat.ufc.br

1 Introduction

In this manuscript, we study global $W^{2,p}$ estimates (with $n \leq p < \infty$) for viscosity solutions to fully nonlinear elliptic equations of the form

$$(1.1) \quad \begin{cases} F(D^2u, Du, u, x) = f(x) & \text{in } \Omega \\ \mathcal{B}(x, u, Du) = g(x) & \text{on } \partial\Omega, \end{cases}$$

where

$$\mathcal{B}(x, u, Du) = \beta(x) \cdot Du + \gamma(x)u(x)$$

is the oblique boundary condition, $\Omega \subset \mathbb{R}^n$ a bounded domain with regular boundary and data f , g , β and γ under appropriate conditions to be clarified soon.

Throughout this work $F : \text{Sym}(n) \times \mathbb{R}^n \times \mathbb{R} \times \Omega \rightarrow \mathbb{R}$ is assumed to be a second order operator with uniformly elliptic structure, i.e., there exist *ellipticity constants* $0 < \lambda \leq \Lambda < \infty$ such that

$$(1.2) \quad \lambda \|Y\| \leq F(X + Y, \varsigma, s, x) - F(X, \varsigma, s, x) \leq \Lambda \|Y\|$$

for every $X, Y \in \text{Sym}(n)$ with $Y \geq 0$ (in the usual partial ordering on symmetric matrices) and $(\varsigma, s, x) \in \mathbb{R}^n \times \mathbb{R} \times \Omega$. Moreover, our approach does not impose any extra regularity assumption on nonlinearity F , e.g. concavity, convexity or smoothness hypothesis (cf. [8], [9], [33] and [41] for classical and modern references). As a matter of fact, we obtain global $W^{2,p}$ -estimates, i.e.,

$$\|u\|_{W^{2,p}(\Omega)} \leq C(\text{universal}) (\|u\|_{L^\infty(\Omega)} + \|f\|_{L^p(\Omega)} + \|g\|_{C^{1,\alpha}(\partial\Omega)})$$

(with $n \leq p < \infty$ and some $\alpha \in (0, 1)$) for viscosity solutions to (1.1) under relaxed convexity assumptions on governing operator (cf. [26], [35] and [44] for modern results) – a sort of asymptotic convexity condition at infinity on F .

To do this concept more precise, we introduce the notion of *Recession operator*, whose terminology comes from Giga-Sato's work [21] on theory of Hamilton-Jacobi PDEs.

Definition 1.1 (Recession profile). We say that $F(X, \varsigma, s, x)$ is asymptotically fully nonlinear elliptic operator if there exists a uniformly elliptic operator $F^\sharp(X, \varsigma, s, x)$, designated the *Recession operator*, such that

$$(R.O.) \quad F^\sharp(X, \varsigma, s, x) := \lim_{\tau \rightarrow 0^+} \tau \cdot F\left(\frac{1}{\tau}X, \varsigma, s, x\right)$$

for all $X \in \text{Sym}(n)$, $\varsigma \in \mathbb{R}^n$, $s \in \mathbb{R}$ and $x \in \Omega$.

For instance, by way of illustration such a limiting profile (R.O.) appears naturally in singularly perturbed free boundary problems ruled by fully nonlinear equations, whose Hessian of solutions blows-up through the phase transition of the model, i.e., $\partial\{u^\varepsilon > \varepsilon\}$, where u^ε satisfies in the viscosity sense

$$F(D^2u^\varepsilon, Du^\varepsilon, x) = \mathcal{Q}_0(x) \frac{1}{\varepsilon} \zeta \left(\frac{u^\varepsilon}{\varepsilon} \right).$$

For such approximations, we have $0 < \mathcal{Q}_0 \in C^0(\bar{\Omega})$, $0 \leq \zeta \in C^\infty(\mathbb{R})$ with $\text{supp } \zeta = [0, 1]$. For this reason, in the above model the limiting free boundary condition is governed by the F^\sharp rather than F , i.e.,

$$F^\sharp(z_0, \nabla u(z_0) \otimes \nabla u(z_0)) = 2T, \quad z_0 \in \partial\{u_0 > 0\}$$

in some appropriate viscosity sense, for a certain total mass $T > 0$ (cf. [37, Section 6] for some enlightening example and details).

Additionally, limit profiles like (R.O.) also appears in higher order convergence rates in periodic homogenization of fully nonlinear uniformly parabolic Cauchy problems with rapidly oscillating initial data as follows

$$\begin{cases} \frac{d}{dt} u^\varepsilon(x, t) = \frac{1}{\varepsilon^2} F(\varepsilon^2 D^2 u^\varepsilon, x, t, \frac{x}{\varepsilon}, \frac{t}{\varepsilon}) & \text{in } \mathbb{R}^n \times (0, T) \\ u^\varepsilon(x, 0) = g(x, \frac{x}{\varepsilon}) & \text{on } \mathbb{R}^n \end{cases}$$

In such a context

$$\frac{1}{\varepsilon^2} F(\varepsilon^2 P, x, t, y, s) \rightarrow F^\sharp(P, x, t, y, s) \quad \text{as } \varepsilon \rightarrow 0^+,$$

uniformly for all $(P, x, t, y, s) \in (\text{Sym}(n) \setminus \{\mathcal{O}_{n \times n}\}) \times \mathbb{R}^n \times [0, T] \times \mathbb{T}^n \times \mathbb{T}$ (see [23]). As a result, there exists a unique function $v : \mathbb{R}^n \times [0, T] \times \mathbb{T}^n \times [0, \infty) \rightarrow \mathbb{R}$ such that $v(x, t, \cdot, \cdot)$ is a viscosity solution to

$$\begin{cases} \frac{d}{ds} v(y, s) = F^\sharp(D_y^2 v, x, t, y, s) & \text{in } \mathbb{T}^n \times (0, \infty) \\ v(x, t, y, 0) = g(y, x) & \text{on } \mathbb{T}^n \end{cases}$$

From now on, the crucial idea will be imposed F^\sharp enjoys a sort of good structural properties (e.g convexity/concavity or suitable *a priori* estimates). At this point, by making use of a geometric tangential analysis, we may to access such a good regularity properties, and in a suitable manner, transfer such estimates to solutions of the original operator via compactness and stability processes.

It is worth to stress that convexity assumptions with respect to the second order derivatives play an essential role in obtaining a number of regularity issues in fully nonlinear elliptic models, e.g. Calderón-Zygmung estimates [8], Schauder type estimates [9] and higher estimates in some nonlinear models [18], [24], [45] and [46], and free boundary problems [13], [14], [19] and [38] just to mention a few. Nevertheless, it is not a trivial task to assure that such regularity issues hold true to problems like (1.1) driven by operators satisfying the structure **(R.O.)**. For instance, consider perturbations of Bellman operators of the form

$$\begin{cases} F(X, \varsigma, s, x) := \inf_{\iota \in \hat{\mathcal{A}}} \left(- \sum_{i,j=1}^n a_{ij}^\iota(x) X_{ij} + \sum_{i=1}^n b_i^\iota(x) \cdot \varsigma_i + c^\iota(x) s \right) + \sum_{i=1}^n \arctg(1 + \lambda_i(X)) = f(x) & \text{in } \Omega \\ \mathcal{B}(x, s, \varsigma) = g(x) & \text{on } \partial\Omega, \end{cases}$$

where $(\lambda_i(X))_{i=1}^n$ are eigenvalues of the matrix $X \in \text{Sym}(n)$, $b_i^\iota, c^\iota : \Omega \rightarrow \mathbb{R}$ are real functions for each $\iota \in \hat{\mathcal{A}}$ (for $\hat{\mathcal{A}}$ any set of index), and $(a_{ij}^\iota(x))_{i,j=1}^n \subset C^{0,\alpha}(\Omega)$ have eigenvalues in $[\lambda, \Lambda]$ for each $x \in \Omega$ and $\iota \in \hat{\mathcal{A}}$. Thus, it is easy to check that such operators are neither concave nor convex. However, its Recession profile **(R.O.)** inherits good structure properties, i.e.,

$$\begin{cases} F^\sharp(D^2 \mathfrak{h}, D\mathfrak{h}, \mathfrak{h}, x) = \inf_{\iota \in \hat{\mathcal{A}}} \left(- \sum_{i,j=1}^n a_{ij}^\iota(x) \mathfrak{h}_{ij} \right) = 0 & \text{in } \Omega \\ \mathcal{B}(x, \mathfrak{h}, D\mathfrak{h}) = g(x) & \text{on } \partial\Omega, \end{cases}$$

which is a convex operator, thereby enjoying higher regularity estimates in face of Evans-Krylov-Trudinger's type theory (with oblique boundary conditions) addressed in [16, Theorem 2.4.], [29, Theorem 1.3], [31, Theorem 5.4], [39, §8], [40, Theorem 3.3] and [34, Theorem 8.3] (see also [9, Chapter 6], [18], [24], [45] and [46] for local estimates), i.e., for some $\alpha \in (0, 1)$ and $\Omega' \subset\subset \Omega \cup \Gamma$ - with $C^{1,\alpha} \ni \Gamma \subset \partial\Omega$, $\beta, \gamma, g \in C^{1,\alpha}(\bar{\Gamma})$, there holds

$$(1.3) \quad \|\mathfrak{h}\|_{C^{2,\alpha}(\bar{\Omega}')} \leq C \left(n, \lambda, \Lambda, \alpha, \|a_{ij}^\iota\|_{C^{0,\alpha}(\Omega)}, \|\beta\|_{C^{1,\alpha}(\bar{\Gamma})}, \|\gamma\|_{C^{1,\alpha}(\bar{\Gamma})}, \Omega', \Omega \right) \left(\|\mathfrak{h}\|_{L^\infty(\Omega)} + \|g\|_{C^{1,\alpha}(\bar{\Gamma})} \right).$$

Heuristically, if the “limiting profile” in **(R.O.)** there exists point-wisely and fulfils a suitable *a priori* estimate like (1.3), then a $W^{2,p}$ -regularity theory to (1.1) holds true up to the boundary via geometric tangential methods (cf. [6], [12] and [35]).

Therefore, one of the main purposes in this work is establishing sharp Hessian estimates for problems enjoying a regular asymptotically elliptic property as **(R.O.)** on the governing operator in (1.1), as well as present a number of applications in related analysis themes and free boundary problems of obstacle type.

Historically, the notion the asymptotically regular problem dates back to Chipot-Evans' work [11] in the context of certain problems in the calculus of variations. Since then there have been an increasing research interest in such class of problems and their related topics, see [7], [20], [36], [42], [43] for an incomplete list of contributions. However, to the best of our knowledge, there is no investigation on such boundary asymptotic qualitative estimates of general problems like (1.1) and their intrinsic connections and applications in Analysis of PDEs and Free Boundary Problems.

1.1 Assumptions and main results

For a given $r > 0$, we denote by $B_r(x)$ the ball of radius r centered at $x = (x', x_n) \in \mathbb{R}^n$, where $x' = (x_1, x_2, \dots, x_{n-1}) \in \mathbb{R}^{n-1}$. We write $B_r = B_r(0)$ and $B_r^+ = B_r \cap \mathbb{R}_+^n$. Also, we write $T_r := \{(x', 0) \in \mathbb{R}^{n-1} : |x'| < r\}$ and $T_r(x_0) := T_r + x'_0$ where $x'_0 \in \mathbb{R}^{n-1}$. Finally, $\text{Sym}(n)$ will denote the set of all $n \times n$ real symmetric matrices.

Throughout this manuscript, we will assume the following assumptions:

- (A1) **(Structural conditions)** We assume that $F \in C^0(\text{Sym}(n), \mathbb{R}^n, \mathbb{R}, \Omega)$. Moreover, there are constants $0 < \lambda \leq \Lambda$, $\sigma \geq 0$ and $\xi \geq 0$ such that

$$(1.4) \quad \begin{aligned} \mathcal{P}_{\lambda, \Lambda}^-(X - Y) - \sigma|q_1 - q_2| - \xi|r_1 - r_2| &\leq F(X, q_1, r_1, x) - F(Y, q_2, r_2, x) \\ &\leq \mathcal{P}_{\lambda, \Lambda}^+(X - Y) + \sigma|q_1 - q_2| + \xi|r_1 - r_2| \end{aligned}$$

for all $X, Y \in \text{Sym}(n)$, $q_1, q_2 \in \mathbb{R}^n$, $r_1, r_2 \in \mathbb{R}$, $x \in \Omega$, where

$$(1.5) \quad \mathcal{P}_{\lambda, \Lambda}^+(X) := \Lambda \sum_{e_i > 0} e_i + \lambda \sum_{e_i < 0} e_i \quad \text{and} \quad \mathcal{P}_{\lambda, \Lambda}^-(X) := \Lambda \sum_{e_i < 0} e_i + \lambda \sum_{e_i > 0} e_i,$$

are the *Pucci's extremal operators* and $e_i = e_i(X)$ ($1 \leq i \leq n$) denote the eigenvalues of X .

- (A2) **(Regularity of the data)** The data satisfy $f \in C^0(\Omega) \cap L^p(\Omega)$ for $n \leq p < \infty$, $g, \gamma \in C^0(\partial\Omega)$ with $\gamma \leq 0$ and $\beta \in C^0(\partial\Omega; \mathbb{R}^n)$ with $\|\beta\|_{L^\infty(\partial\Omega)} \leq 1$ and there exists a positive constant μ_0 such that $\beta \cdot \vec{\mathbf{n}} \geq \mu_0$, where $\vec{\mathbf{n}}$ is the outward normal vector of Ω .

- (A3) **(Continuous coefficients in the L^p -average sense)** According to [10] for a fixed $x_0 \in \Omega$, we define the quantity:

$$\psi_{F^\sharp}(x; x_0) := \sup_{X \in \text{Sym}(n)} \frac{|F^\sharp(X, 0, 0, x) - F^\sharp(X, 0, 0, x_0)|}{\|X\| + 1},$$

which measures the oscillation of the coefficients of F around x_0 . Furthermore, for notation purposes, we shall often write $\psi_F(x, 0) = \psi_F(x)$. Finally, the mapping $\Omega \ni x \mapsto F^\sharp(X, 0, 0, x)$ is assumed to be Hölder continuous function (in the L^p -average sense) for every $X \in \text{Sym}(n)$. This means that, there exist universal constants¹ $\hat{\alpha} \in (0, 1)$, $\theta_0 > 0$ and $0 < r_0 \leq 1$ such that

$$\left(\int_{B_r(x_0) \cap \Omega} \psi_{F^\sharp}(x, x_0)^p dx \right)^{1/p} \leq \theta_0 r^{\hat{\alpha}}$$

for $x_0 \in \bar{\Omega}$ and $0 < r \leq r_0$.

- (A4) **($C^{1,1}$ a priori estimates)** We shall assume that the recession operator F^\sharp there exists and fulfils a up-to-the boundary $C^{1,1}$ a priori estimates, i.e., for $x_0 \in B_1^+$ and $g_0 \in C^{1,\alpha}(T_1)$ (for some $\alpha \in (0, 1)$), there exists a solution $\mathfrak{h} \in C^{1,1}(B_1^+) \cap C^0(\bar{B}_1^+)$ of

$$\begin{cases} F^\sharp(D^2\mathfrak{h}, x_0) &= 0 & \text{in } B_1^+ \\ \mathcal{B}(x, w, D\mathfrak{h}) &= g_0(x) & \text{on } T_1 \end{cases}$$

such that

$$\|\mathfrak{h}\|_{C^{1,1}(\bar{B}_{\frac{1}{2}}^+)} \leq C_1(\text{universal}) \left(\|\mathfrak{h}\|_{L^\infty(B_1^+)} + \|g_0\|_{C^{1,\alpha}(\bar{T}_1)} \right)$$

for some constant $C_1 > 0$.

¹Throughout this work, a constant is said to be *universal* if it depends only on $n, \lambda, \Lambda, p, \mu_0, \|\gamma\|_{C^{1,\alpha}(\partial\Omega)}$ and $\|\beta\|_{C^{1,\alpha}(\partial\Omega)}$

From now on, an operator fulfilling (A1) will be called $(\lambda, \Lambda, \sigma, \xi)$ -elliptic operator. Furthermore, we shall always assume the normalization condition: $F(\mathcal{O}_{n \times n}, \vec{0}, 0, x) = 0$ for all $x \in \Omega$, which is not restrictive, because one can reduce the problem in order to check it.

We now state our first main result, which endures global $W^{2,p}$ estimates for viscosity solutions to asymptotically fully nonlinear equations.

Theorem 1.2 ($W^{2,p}$ estimates under oblique boundary conditions). *Let $n \leq p < \infty$, $\Omega \subset \mathbb{R}^n$, $\partial\Omega \in C^{2,\alpha}$, $\beta, \gamma, g \in C^{1,\alpha}(\partial\Omega)$ (for some $\alpha \in (0, 1)$) and u be an L^p -viscosity solution of (1.1). Further, assume that structural assumptions (A1)-(A4) are in force. Then, there exists a constant $C = C(n, \lambda, \Lambda, p, \mu_0, \|\gamma\|_{C^{1,\alpha}(\partial\Omega)}, \|\beta\|_{C^{1,\alpha}(\partial\Omega)}, \|\partial\Omega\|_{C^{1,1}}) > 0$ such that $u \in W^{2,p}(\Omega)$. Furthermore, the following estimate holds true*

$$(1.6) \quad \|u\|_{W^{2,p}(\Omega)} \leq C \cdot (\|u\|_{L^\infty(\Omega)} + \|f\|_{L^p(\Omega)} + \|g\|_{C^{1,\alpha}(\partial\Omega)}).$$

We also supply a sharp estimate in the scenery that f exceeds the scope of L^∞ functions. Precisely, our second result concerns BMO type *a priori* estimates for second derivatives provided the RHS of (1.1) is a BMO function and F^\sharp enjoys classical *a priori* estimates.

In contrast to the assumptions on Section 2, we will assume, for the next Theorem, the following hypothesis on F^\sharp :

(A4)* (**Higher a priori estimates**) F^\sharp fulfils a $C^{2,\psi}$ *a priori* estimate up to the boundary, for some $\psi \in (0, 1)$, i.e., for any $g_0 \in C^0(\mathbb{B}_1^+) \cap C^{1,\psi}(\mathbb{T}_1)$, $\beta \in C^{1,\psi}(\mathbb{T}_1)$, there exists $C_* > 0$, depending only on universal parameters, such that any viscosity solution of

$$\begin{cases} F^\sharp(D^2\mathfrak{h}, 0, 0, x) = 0 & \text{in } \mathbb{B}_1^+, \\ \beta \cdot D\mathfrak{h} = g_0(x) & \text{on } \mathbb{T}_1. \end{cases}$$

belong to $C^{2,\psi}(\mathbb{B}_1^+) \cap C^0(\overline{\mathbb{B}_1^+})$ and fulfils

$$\|\mathfrak{h}\|_{C^{2,\psi}(\overline{\mathbb{B}_\rho^+})} \leq \frac{C_*}{\rho^{2+\psi}} \left(\|\mathfrak{h}\|_{L^\infty(\mathbb{B}_1^+)} + \|g_0\|_{C^{1,\psi}(\mathbb{T}_1)} \right) \quad \forall 0 < \rho \ll 1.$$

Definition 1.3. For a function $f \in L^1_{\text{loc}}(\mathbb{R}^n)$ $x_0 \in \Omega$ and $\rho > 0$ we define $(f)_{x_0, \rho}$ by

$$(f)_{x_0, \rho} := \int_{\mathbb{B}_\rho(x_0) \cap \Omega} f(x) dx.$$

Moreover, when $x_0 = 0$ we will denote $(f)_\rho$ by simplicity.

An $f \in L^1_{\text{loc}}(\Omega)$ is a function of p -bounded mean oscillation, in short $f \in p\text{-BMO}(\Omega)$, if

$$(1.7) \quad \|f\|_{p\text{-BMO}(\Omega)} := \sup_{\substack{\mathbb{B}_\rho \subseteq \Omega \\ x_0 \in \Omega}} \left(\int_{\mathbb{B}_\rho(x_0) \cap \Omega} |f(x) - (f)_{x_0, \rho}|^p dx \right)^{\frac{1}{p}} \leq C(\text{universal}),$$

for a constant $C > 0$. Such functions equipped with the norm (1.7) define a Banach space.

Theorem 1.4 (Boundary p -BMO type estimates). *Let u be an L^p -viscosity solution to*

$$\begin{cases} F(D^2u, Du, u, x) = f(x) & \text{in } \mathbb{B}_1^+ \\ \beta \cdot Du = g(x) & \text{on } \mathbb{T}_1, \end{cases}$$

where $f \in p\text{-BMO}(\mathbb{B}_1^+) \cap L^p(\mathbb{B}_1^+)$, for some $n \leq p \leq \infty$. Assume further that assumptions (A1)-(A3) and (A4)* are in force. Then, $D^2u \in p\text{-BMO}(\mathbb{B}_1^+)$, Moreover, the following estimate holds

$$(1.8) \quad \|D^2u\|_{p\text{-BMO}(\overline{\mathbb{B}_{\frac{1}{2}}^+})} \leq C(n, \lambda, \Lambda, p, C_*, \mu_0, \|\beta\|_{C^{1,\alpha}(\mathbb{T}_1)}) \left(\|u\|_{L^\infty(\mathbb{B}_1^+)} + \|g\|_{C^{1,\alpha}(\mathbb{T}_1)} + \|f\|_{p\text{-BMO}(\mathbb{B}_1^+)} \right).$$

Remark 1.5 ($C^{1,\text{Log-Lip}}$ estimates). Since viscosity solutions to (1.1) enjoy a Hessian control in the L^p -average sense, then by invoking the embedding result from [1, Lemma 1], we may establish $C^{1,\text{Log-Lip}}$ type estimates: Let $h : B_1^+ \rightarrow \mathbb{R}$ be a function such that locally $D_{ij}h \in BMO_p(B_1^+)$ for all $1 \leq i, j \leq n$. Then, $h \in C_{loc}^{1,\text{Log-Lip}}(B_1^+)$. In other words,

$$\sup_{\rho \in (0,1)} \sup_{z \in \partial\Omega} \sup_{B_\rho(z) \cap \Omega} \frac{|u(x) - [u(z) + Du(z) \cdot (x-z)]|}{\rho^2 \ln \rho^{-1}} \leq \hat{C} \cdot (\|u\|_{L^\infty(\Omega)} + \|g\|_{C^{1,\alpha}(\partial\Omega)} + \|f\|_{p\text{-BMO}(\Omega)}),$$

for some constant $\hat{C} = \hat{C}(n, \lambda, \Lambda, p, C_*, \mu_0, \|\beta\|_{C^{1,\alpha}(\overline{T_1})}, \|\partial\Omega\|_{C^{1,1}})$ (cf. [12, Theorem 5.2] for related boundary estimates).

We would like to mention that although our manuscript has been strongly motivated by recent works [2], [12] and [35], our approach has required a sort of non-trivial adaptations due to the presence of a non-homogeneous oblique boundary condition and the asymptotic nature of limiting operator. Furthermore, different from [48], our findings also provide further quantitative applications, such as BMO type estimates (see Sect. 4), $W^{2,p}$ -regularity to obstacle type problems under oblique boundary conditions (see Sect. 5) and density results for oblique type problems (see Section 6) via tangential methods. Additionally, our recession profiles **R.O.** encode more general operators than linear ones as addressed in [48] (see also [6]). Particularly, our results extend regarding oblique boundary scenario, former results from [6], [35] and [44] (see also [12]), and to some extent, those from [6], [26], [27] and [48] by making use of techniques adapted to the general framework of the fully nonlinear models under relaxed convexity assumptions and oblique boundary datum.

Finally, we also highlight that Theorems 1.2 and 1.4 can be extended to a more general class of models, including fully nonlinear parabolic PDEs. For sake of simplicity and clarity, we have decided to consider just the elliptic scenario in this paper. We pretend study such issues in a forthcoming project.

1.2 Applications to obstacle type problems

In closing, we would like to highlight that boundary Hessian estimates are also useful in the context of obstacle type problems. Physically the classical obstacle problem refers to the equilibrium position of an elastic membrane (i.e $u : \Omega \rightarrow \mathbb{R}$), whose boundary is held fixed (i.e $u|_{\partial\Omega} = g$), lying above a given obstacle and subject to the action of external forces e.g. friction, tension and/or gravity ($f : \Omega \rightarrow \mathbb{R}$). More precisely, given an elastic membrane u attached to a fixed boundary $\partial\Omega$ and an obstacle ϕ , we seek for the equilibrium position of the membrane when we move it downwards toward the obstacle satisfying:

$$\begin{cases} \Delta u \leq f & \text{in the weak sense in } \Omega \\ \Delta u = f & \text{in the weak sense in } \{u > \phi\} \\ u \geq \phi & \text{in } \Omega \end{cases}$$

with $u - g \in H_0^1(\Omega)$. We refer [4] for Calderón-Zygmund estimates for elliptic and parabolic obstacle problems in non-divergence form (under convex structure) with discontinuous coefficients and irregular obstacles. We must quote [30] concerning gradient estimates for obstacle problems of elliptic models (under convex/convex structure) with oblique boundary conditions.

Now, we would also like to focus our attention to the recent work [3, Theorem 1.1.], where $W^{2,p}$ estimates for the obstacle problem with oblique boundary conditions

$$(1.9) \quad \begin{cases} F(D^2u, Du, u, x) \leq f(x) & \text{in } \Omega \\ (F(D^2u, Du, u, x) - f)(u - \phi) = 0 & \text{in } \Omega \\ u(x) \geq \phi(x) & \text{in } \Omega \\ \beta \cdot Du + \gamma u = g(x) & \text{on } \partial\Omega \end{cases}$$

for a given obstacle $\phi \in W^{2,p}(\Omega)$ satisfying $\beta(x) \cdot D\phi \geq g$ a.e. on $\partial\Omega$, have been obtained in the case when F is a convex operator and $\gamma = g \equiv 0$.

Motivated by above problem, we will investigate $W^{2,p}$ regularity estimates for obstacle type problems (1.9) under an asymptotic convexity assumption weaker than convexity and non-homogeneous oblique conditions.

For our purpose, it is necessary to assume the following assumptions:

(A5) There exists a modulus of continuity $\omega : [0, +\infty) \rightarrow [0, +\infty)$ with $\omega(0) = 0$, such that

$$F(X_1, q, r, x_1) - F(X_2, q, r, x_2) \leq \omega(|x_1 - x_2|) [(|q| + 1) + \alpha_0|x_1 - x_2|^2]$$

holds for any $x_1, x_2 \in \Omega$, $q \in \mathbb{R}^n$, $r \in \mathbb{R}$, $\alpha_0 > 0$ and $X_1, X_2 \in \text{Sym}(n)$ satisfying

$$-3\alpha_0 \begin{pmatrix} \text{Id}_n & 0 \\ 0 & \text{Id}_n \end{pmatrix} \leq \begin{pmatrix} X_2 & 0 \\ 0 & -X_1 \end{pmatrix} \leq 3\alpha_0 \begin{pmatrix} \text{Id}_n & -\text{Id}_n \\ -\text{Id}_n & \text{Id}_n \end{pmatrix},$$

where Id_n is the identity matrix.

(A6) F is a proper operator in the following sense:

$$d \cdot (r_2 - r_1) \leq F(X, q, r_1, x) - F(X, q, r_2, x),$$

for any $X \in \text{Sym}(n)$, $r_1, r_2 \in \mathbb{R}$, with $r_1 \leq r_2$, $x \in \Omega$, $q \in \mathbb{R}^n$, and some $d > 0$.

The previous assumptions are invoked to assure the validity of the Comparison Principle for oblique derivative problems like (1.1) (see [10, Theorem 2.10] and [32, Theorem 7.17]), thereby ensuring the possibility of employing the Perron's Method for viscosity solutions (see Lieberman's Book [32, Chapter 7.4 and 7.6]).

Finally, our third result addresses a $W^{2,p}$ -regularity theory for (1.9), whose approach does not require any extra regularity assumption on nonlinearity F .

Theorem 1.6 (Obstacle problems with oblique boundary conditions). *Let u be an L^p -viscosity solution of (1.9) with $n < p < \infty$, where F satisfies the structural assumption (A1)-(A4) and (A5)-(A6), $\partial\Omega \in C^3$, $f \in L^p(\Omega)$, $\beta \in C^2(\partial\Omega)$ with $\beta \cdot \vec{\mathbf{n}} \geq \mu_0$ for some $\mu_0 > 0$, $\phi \in W^{2,p}(\Omega)$ and $g \in C^{1,\alpha}(\partial\Omega)$ (for some $\alpha \in (0, 1)$). Then, $u \in W^{2,p}(\Omega)$ with the following estimate*

$$(1.10) \quad \|u\|_{W^{2,p}(\Omega)} \leq C_0 \cdot (\|f\|_{L^p(\Omega)} + \|\phi\|_{W^{2,p}(\Omega)} + \|g\|_{C^{1,\alpha}(\partial\Omega)}).$$

where $C_0 = C_0(n, \lambda, \Lambda, p, \mu_0, \sigma, \omega, \|\beta\|_{C^2(\partial\Omega)}, \|\gamma\|_{C^2(\partial\Omega)}, \partial\Omega, \text{diam}(\Omega), \theta_0)$.

In conclusion, we must stress that similar Hessian estimates hold true for obstacle problems with Dirichlet boundary conditions

$$\begin{cases} F(D^2u, Du, u, x) \leq f(x) & \text{in } \Omega \\ (F(D^2u, Du, u, x) - f)(u - \phi) = 0 & \text{in } \Omega \\ u(x) \geq \phi(x) & \text{in } \Omega \\ u(x) = g(x) & \text{on } \partial\Omega \end{cases}$$

by invoking the related global $W^{2,p}$ regularity estimates addressed in [6, Theorem 2.7] and [12, Theorems 1.1] with following estimate

$$\|u\|_{W^{2,p}(\Omega)} \leq C(\text{universal}) \cdot (\|f\|_{L^p(\Omega)} + \|\phi\|_{W^{2,p}(\Omega)} + \|g\|_{W^{2,p}(\partial\Omega)}) \quad \text{for } n < p < \infty.$$

1.3 Hessian estimates to models in non-divergence form: State-of-Art

Regularity estimates in Sobolev spaces for nonlinear models have instituted an important chapter in the modern history of elliptic theory of viscosity solutions. Moreover, because of its non-variational nature, obtaining sharp integrability properties of solutions of such elliptic operators have been a demanding work throughout the last decades. One of the most important

issues was on whether $W^{2,p}$ a priori estimates could be addressed for any fully nonlinear operator, thereby attracting notable efforts by the academic community during the last three decades or more, one remained open up to the advance of modern techniques.

In the seminal work [8, Theorem 7.1], Caffarelli proved that C^0 -viscosity solutions to

$$(1.11) \quad F(D^2u, x) = f(x) \quad \text{in } B_1$$

satisfy an interior $W^{2,p}$ -regularity for $n < p < \infty$, with the following estimate

$$\|u\|_{W^{2,p}(B_{\frac{1}{2}})} \leq C(n, p, \lambda, \Lambda) (\|u\|_{L^\infty(B_1)} + \|f\|_{L^p(B_1)}),$$

provided a small oscillation of $F(X, x)$ in the variable x and $C^{1,1}$ a priori estimates for homogeneous equation with “frozen coefficients” $F(D^2w, x_0) = 0$ are in force. Furthermore, Caffarelli shown that for any $p < n$ there exists a fully nonlinear operator fulfilling the previous hypothesis for which $W^{2,p}$ -estimates fail to hold. In this direction, we must also mention the Dong-Kim’ work [15] for notable examples of linear elliptic operators in non-divergence form, with piecewise constant coefficients, whose solutions have a lacking of $W^{2,p}$ estimates.

A few years later, Caffarelli’s $W^{2,p}$ -estimates were extended by Escauriaza in [17, Theorem 1] in the context of L^n -viscosity solutions to the range of exponents $n - \varepsilon_0 < p < \infty$ with $\varepsilon_0 = \varepsilon_0(\frac{\Lambda}{\lambda}, n) \in (\frac{n}{2}, n)$ (the Escauriaza’s constant), which provides the minimal range of integrability for which the Harnack inequality (resp. Hölder regularity) holds for viscosity solutions to (1.11). Additionally, when the boundary of the domain is smooth enough, namely $C^{1,1}$, a global $W^{2,p}$ estimate is obtained for $n - \varepsilon_0 < p < \infty$ by Winter in [47, Theorem 4.5].

In the aftermath, Byun-Lee-Palagachev in [5] extend the Escauriaza-Winter’s results by studying global regularity of viscosity solutions to the following Dirichlet problem for a fully nonlinear uniformly elliptic equation

$$(1.12) \quad \begin{cases} F(D^2u, Du, u, x) = f(x) & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega \end{cases}$$

in weighted Lebesgue spaces. Under appropriate structural conditions, if $f \in L_w^p(\Omega)$ for $p > n - \varepsilon_0$ and $w \in \mathcal{A}_{\frac{p}{n-\varepsilon_0}}$ (Muckenhoupt classes), then the problem (1.12) admits a unique solution $u \in W_w^{2,p}(\Omega)$ such that

$$\|u\|_{W_w^{2,p}(\Omega)} \leq C(\text{universal})\|f\|_{L_w^p(\Omega)}$$

Recently, in a new landmark for this theory, Caffarelli’s local $W^{2,p}$ regularity estimates were extended by Pimentel-Teixeira in [35]. Precisely, the novelty with respect to the Caffarelli’s results is the concept of *Recession* operator (a sort of tangent profile for F at “infinity”, see **(R.O.)**). In such a context, the authors relaxed the hypothesis of $C^{1,1}$ a priori estimates for solutions of equations without dependence on x , by assuming that F must be “convex or concave” only at the ends of $Sym(n)$. In this setting, the authors proved that any viscosity solution to

$$(1.13) \quad F(D^2u) = f(x) \quad \text{in } B_1,$$

where $f \in L^p(B_1) \cap C^0(B_1)$, fulfils $u \in W_{loc}^{2,p}(B_1)$ with an a priori estimate

$$\|u\|_{W^{2,p}(B_{\frac{1}{2}})} \leq C(n, p, \lambda, \Lambda) (\|u\|_{L^\infty(B_1)} + \|f\|_{L^p(B_1)}).$$

We must quote similar global $W^{2,p}$ -estimates were proved independently by Byan *et al* in [6, Theorems 2.5 and 2.7] for solutions to asymptotically linear operators like (1.13) with zero Dirichlet boundary condition. Their approach is based on a proper transformation which converts a given asymptotically elliptic PDE to a suitable uniformly elliptic one. We also cite Silvestre-Teixeira’s

work [44, Theorems 1 and 4] for an interesting survey on this theme in the setting of sharp gradient estimates.

In the sequel, global $W^{2,p}$, BMO_p and $C^{1,\text{Log-Lip}}$ regularity estimates, in the context of asymptotic convex operators (see Definition (1.1)), i.e.,

$$\begin{cases} F(D^2u, Du, u, x) = f(x) & \text{in } \Omega \\ u(x) = g(x) & \text{on } \partial\Omega \end{cases}$$

were addressed by da Silva-Ricarte in [12, Theorems 1.1, 1.2 and 5.2]. Precisely, they obtain

$$\|u\|_{W^{2,p}(\Omega)} \leq C(n, p, \lambda, \Lambda, \|\partial\Omega\|_{C^{1,1}}) (\|u\|_{L^\infty(\Omega)} + \|f\|_{L^p(\Omega)} + \|g\|_{W^{2,p}(\partial\Omega)}).$$

We highlight that the proof of the such estimates in [12] is based on successful adaptation of compactness arguments inspired by the ideas from Pimentel-Teixeira [35], Silvestre-Teixeira [44] and Winter [47]. We also refer the reader to Lee's work [28] for Hessian estimates for solutions of (1.13) in the framework of weighted Orlicz spaces.

Among other works on this research topic, we must cite the sequence of Krylov's fundamental manuscripts [25], [26] and [27], which are associated with the existence of L^p -viscosity solutions under either relaxed or no convexity assumptions on F .

Now, concerning problems with oblique boundary conditions, Hessian estimates for viscosity solutions of elliptic equations like (1.1) has been regarded as an important issue in the study of PDEs for a long time. The theory of this subject has also been improved over the last several decades. For general fully nonlinear problems like (1.1), the existence and uniqueness for viscosity solutions of oblique boundary problems was proved in [22] and [32].

Inspired by the ideas from Caffarelli's fundamental work [8], Byan and Han in [2] establish global $W^{2,p}$ -estimates for viscosity solutions of convex fully nonlinear elliptic models of the form

$$\begin{cases} F(D^2u, Du, u, x) = f(x) & \text{in } \Omega \\ \beta(x) \cdot Du = 0 & \text{on } \partial\Omega \end{cases}$$

Moreover, the following estimate holds true

$$\|u\|_{W^{2,p}(\Omega)} \leq C(\text{universal}) (\|u\|_{L^\infty(\Omega)} + \|f\|_{L^p(\Omega)}),$$

It is also worth highlighting that $C^{0,\alpha}$, $C^{1,\alpha}$ and $C^{2,\alpha}$ boundary regularity were addressed for the Neumann problem on flat boundary by Milakis-Silvestre in [34]. Moreover, such results were recently extended to the general oblique scenario by Li-Zhang in [29].

Finally, we must quote Zhang *et al.*'s work [48], which establish $W^{2,p}$ -regularity for viscosity solutions of fully nonlinear elliptic equations with a homogeneous oblique derivative boundary condition. In this context, the nonlinearity is assumed to fulfil a certain asymptotic regularity condition.

In conclusion, taking into account such historical breakthroughs, in this work, we aim to obtain a global $W^{2,p}$ estimates for a large class of nonlinear elliptic PDEs. To this end, we only assume that $F(X, \varsigma, s, x)$ is asymptotically elliptic, that is, under a relaxed convexity assumption (a suitable asymptotic structure of F just at "infinity" of $Sym(n)$ according to the Definition 1.1) we establish global Sobolev *a priori* estimates for solutions of (1.1).

This paper is organized as follows: Section 2 contains the main notations and preliminary results which we will work throughout this manuscript. In Section 2.1, we present the *Modus Operandi*, i.e., the tangential approximation mechanism used in order to relate the estimates coming from *Recession* operator F^\sharp with its original counterpart F . Section 3 is devoted to proving Theorem 1.2. In Section 4 we establish the regularity estimates in the borderline setting, namely for p-BMO spaces. Finally, in the last sections we establish the global $W^{2,p}$ estimate for the obstacle problems with oblique boundary conditions and density results in a proper class of viscosity solutions.

2 Preliminaries and auxiliary results

We introduce the appropriate notions of viscosity solutions to (1.1).

Definition 2.1 (C^0 -viscosity solutions). Let F be a $(\lambda, \Lambda, \sigma, \xi)$ -elliptic operator and $\Gamma \subset \Omega$ a relatively open set. A function $u \in C(\Omega \cup \Gamma)$ is said a C^0 -viscosity solution if the following condition hold:

a) for all $\forall \varphi \in C^2(\Omega \cup \Gamma)$ touching u by above at $x_0 \in \Omega \cup \Gamma$,

$$\begin{cases} F(D^2\varphi(x), D\varphi(x), \varphi(x), x) \geq f(x_0) & \text{when } x_0 \in \Omega \\ \text{and } \mathcal{B}(x_0, u(x_0), D\varphi(x_0)) \geq g(x_0) & \text{at } x_0 \in \Gamma \end{cases}$$

b) for all $\forall \varphi \in C^2(\Omega \cup \Gamma)$ touching u by below at $x_0 \in \Omega \cup \Gamma$,

$$\begin{cases} F(D^2\varphi(x), D\varphi(x), \varphi(x), x) \leq f(x_0) & \text{when } x_0 \in \Omega \\ \text{and } \mathcal{B}(x_0, u(x_0), D\varphi(x_0)) \leq g(x_0) & \text{at } x_0 \in \Gamma \end{cases}$$

Definition 2.2 (L^p -viscosity solution). Let F be a $(\lambda, \Lambda, \sigma, \xi)$ -elliptic operator, $p \geq n$ and $f \in L^p(\Omega)$. Assume that F is continuous in X , q, r and measurable in x . A function $u \in C(\bar{\Omega})$ is said an L^p -viscosity solution for 1.1 if the following assertions hold:

a) For all $\varphi \in W^{2,p}(\Omega)$ whenever $\varepsilon > 0$, \mathcal{O} is a relatively open in $\bar{\Omega}$ and

$$\begin{cases} F(D^2\varphi(x), D\varphi(x), \varphi(x), x) \leq f(x) - \varepsilon, & \text{a.e. in } \mathcal{O} \cap \Omega \\ \text{and } \mathcal{B}(x, \varphi(x), D\varphi(x)) \leq g(x) - \varepsilon & \text{on } \mathcal{O} \cap \partial\Omega, \end{cases}$$

$u - \varphi$ cannot attain a local maximum in \mathcal{O} .

b) For all $\varphi \in W^{2,p}(\Omega)$ whenever $\varepsilon > 0$, \mathcal{O} is a relatively open in $\bar{\Omega}$ and

$$\begin{cases} F(D^2\varphi(x), D\varphi(x), \varphi(x), x) \geq f(x) - \varepsilon, & \text{a.e. in } \mathcal{O} \cap \Omega \\ \text{and } \mathcal{B}(x, \varphi(x), D\varphi(x)) \geq g(x) - \varepsilon & \text{on } \mathcal{O} \cap \partial\Omega, \end{cases}$$

$u - \varphi$ cannot attain a local minimum in \mathcal{O} .

For convenience of notation we define

$$\mathcal{L}^\pm(u) := \mathcal{P}_{\lambda, \Lambda}^\pm(D^2u) \pm \sigma|Du|.$$

Definition 2.3. We define the class $\bar{\mathcal{S}}(\lambda, \Lambda, \sigma, f)$ and $\underline{\mathcal{S}}(\lambda, \Lambda, \sigma, f)$ to be the set of all continuous functions u that satisfy $\mathcal{L}^+(u) \geq f$, respectively $\mathcal{L}^-(u) \leq f$ in the viscosity sense (see Definition 2.1). We define

$$\mathcal{S}(\lambda, \Lambda, \sigma, f) := \bar{\mathcal{S}}(\lambda, \Lambda, \sigma, f) \cap \underline{\mathcal{S}}(\lambda, \Lambda, \sigma, f) \text{ and } \mathcal{S}^*(\lambda, \Lambda, \sigma, f) := \bar{\mathcal{S}}(\lambda, \Lambda, \sigma, |f|) \cap \underline{\mathcal{S}}(\lambda, \Lambda, \sigma, -|f|).$$

Moreover, when $\sigma = 0$, we denote $\mathcal{S}^*(\lambda, \Lambda, 0, f)$ just by $\mathcal{S}^*(\lambda, \Lambda, f)$ (resp. by $\underline{\mathcal{S}}, \bar{\mathcal{S}}, \mathcal{S}$).

Now, we present a compactness result, whose proof can be found in [29, Theorem 1.1].

Theorem 2.4 ($C^{0, \alpha'}$ regularity). Let $u \in C^0(\Omega) \cap C^0(\Gamma)$ be satisfying

$$\begin{cases} u \in \mathcal{S}(\lambda, \Lambda, f) & \text{in } \Omega \\ \mathcal{B}(x, u, Du) = g(x) & \text{on } \Gamma \end{cases}$$

Then, for any $\Omega' \subset\subset \Omega \cup \Gamma$, $u \in C^{0, \alpha'}(\bar{\Omega}')$ and

$$\|u\|_{C^{0, \alpha'}(\bar{\Omega}')} \leq C(n, \lambda, \Lambda, \mu_0, \|\gamma\|_{L^\infty(\Gamma)}, \Omega', \Omega) (\|u\|_{L^\infty(\Omega)} + \|f\|_{L^n(\Omega)} + \|g\|_{L^\infty(\Gamma)})$$

where $\alpha' \in (0, 1)$ depends on n, λ, Λ and μ_0 .

Next, we present the following Stability result (see [10, Theorem 3.8] for a proof).

Lemma 2.5 (Stability Lemma). *For $k \in \mathbb{N}$ let $\Omega_k \subset \Omega_{k+1}$ be an increasing sequence of domains and $\Omega := \bigcup_{k=1}^{\infty} \Omega_k$. Let $p \geq n$ and F, F_k be $(\lambda, \Lambda, \sigma, \xi)$ -elliptic operators. Assume $f \in L^p(\Omega)$, $f_k \in L^p(\Omega_k)$ and that $u_k \in C^0(\Omega_k)$ are L^p -viscosity sub-solutions (resp. super-solutions) of*

$$F_k(D^2 u_k, Du_k, u_k, x) = f_k(x) \quad \text{in } \Omega_k.$$

Suppose that $u_k \rightarrow u_\infty$ locally uniformly in Ω and that for $B_r(x_0) \subset \Omega$ and $\varphi \in W^{2,p}(B_r(x_0))$ we have

$$(2.1) \quad \|(\hat{g} - \hat{g}_k)^+\|_{L^p(B_r(x_0))} \rightarrow 0 \quad (\text{resp. } \|(\hat{g} - \hat{g}_k)^-\|_{L^p(B_r(x_0))} \rightarrow 0),$$

where $\hat{g}(x) := F(D^2 \varphi, D\varphi, u, x) - f(x)$ and $\hat{g}_k(x) = F_k(D^2 \varphi, D\varphi, u_k, x) - f_k(x)$. Then u is an L^p -viscosity sub-solution (resp. super-solution) of

$$F(D^2 u, Du, u, x) = f(x) \quad \text{in } \Omega.$$

Moreover, if F, f are continuous, then u is a C^0 -viscosity sub-solution (resp. super-solution) provided that (2.1) holds for all $\varphi \in C^2(B_r(x_0))$ test function.

The next result is an ABP Maximum Principle (see [29, Theorem 2.1] for a proof).

Lemma 2.6 (ABP Maximum Principle). *Let $\Omega \subset B_1$ and u satisfying*

$$\begin{cases} u \in \mathcal{S}(\lambda, \Lambda, f) & \text{in } \Omega \\ \mathcal{B}(x, u, Du) = g(x) & \text{on } \Gamma. \end{cases}$$

Suppose that exists $\varsigma \in \partial B_1$ such that $\beta \cdot \varsigma \geq \mu_0$ and $\gamma \leq 0$ in Γ . Then,

$$\|u\|_{L^\infty(\Omega)} \leq \|u\|_{L^\infty(\partial\Omega \setminus \Gamma)} + C(n, \lambda, \Lambda, \mu_0)(\|g\|_{L^\infty(\Gamma)} + \|f\|_{L^n(\Omega)})$$

In the sequel, we comment on the existence and uniqueness of viscosity solutions with oblique boundary conditions. For that purpose, we will assume the following condition on F :

(SC) There exists a modulus of continuity $\tilde{\omega}$, i.e., $\tilde{\omega}$ is nondecreasing with $\lim_{t \rightarrow 0} \tilde{\omega}(t) = 0$ and

$$\psi_F(x, y) \leq \tilde{\omega}(|x - y|).$$

Now, we will ensure existence and uniqueness of viscosity solutions to following problem:

$$\begin{cases} F(D^2 u, x) = f(x) & \text{in } \Omega, \\ \mathcal{B}(x, u, Du) = g(x) & \text{on } \Gamma \\ u(x) = \varphi(x) & \text{on } \partial\Omega \setminus \Gamma, \end{cases}$$

where Γ is relatively open of $\partial\Omega$. The next proof follows the ideas from [29, Theorem 3.1] with minor modifications. For this reason, we will omit it here.

Theorem 2.7. *Suppose that $\Gamma \in C^2$ and $\beta \in C^2(\bar{\Gamma})$. Assume that F satisfies (A1) and (SC). Let u and v be satisfying*

$$\begin{cases} F(D^2 u, x) \geq f_1(x) & \text{in } \Omega, \\ \mathcal{B}(x, u, Du) \geq g_1(x) & \text{on } \Gamma \end{cases}$$

and

$$\begin{cases} F(D^2 v, x) \leq f_2(x) & \text{in } \Omega, \\ \mathcal{B}(x, v, Dv) \leq g_2(x) & \text{on } \Gamma \end{cases}$$

Then,

$$\begin{cases} u - v \in \mathcal{S}(\frac{\lambda}{n}, \Lambda, f_1 - f_2) & \text{in } \Omega, \\ \mathcal{B}(x, u - v, D(u - v)) \geq (g_1 - g_2)(x) & \text{on } \Gamma \end{cases}$$

Theorem 2.8 (Uniqueness). *Let $\Gamma \in C^2$, $\beta \in C^2(\bar{\Gamma})$, $\gamma \leq 0$ and $\varphi \in C^0(\partial\Omega \setminus \Gamma)$. Suppose that there exists $\varsigma \in \partial B_1$ such that $\beta \cdot \varsigma \geq \mu_0$ on Γ . Assume that F satisfies (A1). Then, there exists at most one viscosity solution of*

$$\begin{cases} F(D^2u, x) = f(x) & \text{in } \Omega, \\ \mathcal{B}(x, u, Du) = g(x) & \text{on } \Gamma \\ u(x) = \varphi(x) & \text{on } \partial\Omega \setminus \Gamma. \end{cases}$$

Proof. The proof is direct consequence of Theorem 2.7 with ABP estimate (Lemma 2.6). \square

Theorem 2.9 (Existence). *Let $\Gamma \in C^2$, $\beta \in C^2(\bar{\Gamma})$, $\gamma \leq 0$ and $\varphi \in C^0(\partial\Omega \setminus \Gamma)$. Suppose that there exists $\varsigma \in \partial B_1$ such that $\beta \cdot \varsigma \geq \mu_0$ on Γ and assume (SC). In addition, suppose that Ω satisfies an exterior cone condition at any $x \in \partial\Omega \setminus \bar{\Gamma}$ and satisfies an exterior sphere condition at any $x \in \bar{\Gamma} \cap (\partial\Omega \setminus \Gamma)$. Then, there exists a unique viscosity solution $u \in C^0(\bar{\Omega})$ of*

$$\begin{cases} F(D^2u, x) = f(x) & \text{in } \Omega, \\ \mathcal{B}(x, u, Du) = g(x) & \text{on } \Gamma \\ u(x) = \varphi(x) & \text{on } \partial\Omega \setminus \Gamma. \end{cases}$$

Proof. The proof follows the same lines as the one in [29, Theorem 3.3] with minor modifications. For instance, we must invoke Theorem 2.8 instead of [29, Theorem 3.2]. Other necessary modifications are the same made in the proof of Theorem 2.7. \square

Next result is a key tool in our tangential approach. Precisely, it describes how the point-wise convergence of F_τ to F^\sharp takes place.

Lemma 2.10. *Let F be a uniformly elliptic operator and assume F^\sharp there exists. Then, given $\epsilon > 0$ there exists $\tau_0(\lambda, \Lambda, \epsilon, \psi_{F^*}) > 0$ such that, for every $\tau \in (0, \tau_0)$ there holds*

$$\frac{|\tau F(\tau^{-1}X, 0, 0, x) - F^\sharp(X, 0, 0, x)|}{1 + \|X\|} \leq \epsilon,$$

for every $X \in \text{Sym}(n)$.

Proof. The proof follows the same lines of the one in [44] (see also [35]). For this reason we omit it here. \square

Remark 2.11. It is worth highlighting that the $W^{2,p}$ estimates from Theorem 1.2 depends not only on universal constants, but in effect on the “modulus of convergence” $F_\tau \rightarrow F^\sharp$. Precisely, by defining $\varsigma : (0, \infty) \rightarrow (0, \infty)$ as follows

$$\varsigma(\epsilon) := \sup_{\substack{X \in \text{Sym}(n) \\ \tau \in (0, \tau_0)}} \left\{ \frac{|\tau F(\tau^{-1}X, 0, 0, x) - F^\sharp(X, 0, 0, x)|}{1 + \|X\|} \leq \epsilon \right\},$$

then the constant $C > 0$ appearing in up-to-the boundary $W^{2,p}$ a priori estimates (1.6) also depends on ς .

2.1 Approximation device in geometric tangential analysis

The main result of this section provides a compactness method that will be used as a key point throughout the whole paper. As a matter of fact, we will show that if our equation is close enough to the homogeneous equation with constant coefficients, then our solution is sufficiently close to a solution of the homogeneous equation with frozen coefficients. At the core of our techniques, is the notion of recession operator (see Definition 1.1). The appropriate way to formalize this intuition is by an approximation lemma.

Lemma 2.12 (Approximation Lemma). *Let $n \leq p < \infty$, $0 \leq \nu \leq 1$ and assume that (A1) – (A4) are in force. Then, for every $\delta > 0$, $\varphi \in C(\partial B_1(0', \nu))$ with $\|\varphi\|_{L^\infty(\partial B_1(0', \nu))} \leq \mathbf{c}_1$ and $g \in C^{1,\alpha}(\overline{T}_2)$ with $0 < \alpha < 1$ and $\|g\|_{C^{1,\alpha}(\overline{T}_2)} \leq \mathbf{c}_2$ for some $\mathbf{c}_2 > 0$ there exist positive constants $\epsilon = \epsilon(\delta, n, \mu_0, p, \lambda, \Lambda, \gamma, \mathbf{c}_1, \mathbf{c}_2) < 1$ and $\tau_0 = \tau_0(\delta, n, \lambda, \Lambda, \mu_0, \mathbf{c}_1, \mathbf{c}_2) > 0$ such that, if*

$$\max \left\{ |F_\tau(X, x) - F^\sharp(X, x)|, \|\psi_{F^\sharp}(\cdot, 0)\|_{L^p(B_r^+)}, \|f\|_{L^p(B_r^+)} \right\} \leq \epsilon \quad \text{and} \quad \tau \leq \tau_0$$

then, any two L^p -viscosity solutions u (normalized, i.e., $\|u\|_{L^\infty(B_r^+(0', \nu))} \leq 1$) and \mathfrak{h} of

$$\begin{cases} F_\tau(D^2u, x) = f(x) & \text{in } B_r^+(0', \nu) \cap \mathbb{R}_+^n \\ \mathcal{B}(x, u, Du) = g(x) & \text{on } B_r(0', \nu) \cap T_r \\ u(x) = \varphi(x) & \text{on } \partial B_r(0', \nu) \cap \mathbb{R}_+^n \end{cases}$$

and

$$\begin{cases} F^\sharp(D^2\mathfrak{h}, 0) = 0 & \text{in } B_{\frac{7}{8}r}^+(0', \nu) \cap \mathbb{R}_+^n \\ \mathcal{B}(x, \mathfrak{h}, D\mathfrak{h}) = g(x) & \text{on } B_{\frac{7}{8}r}(0', \nu) \cap T_r \\ \mathfrak{h}(x) = u(x) & \text{on } \partial B_{\frac{7}{8}r}(0', \nu) \cap \mathbb{R}_+^n \end{cases}$$

satisfy

$$\|u - \mathfrak{h}\|_{L^\infty(B_{\frac{7}{8}r}^+(0', \nu))} \leq \delta.$$

Proof. Without loss of generality, we may assume that $r = 1$. We will proceed by a *reductio at absurdum* argument. Thus, let us assume that the claim is not satisfied. Then, there exists $\delta_0 > 0$ and a sequence of functions $(F_{\tau_j})_{j \in \mathbb{N}}$, $(F_j^\sharp)_{j \in \mathbb{N}}$, $(u_j)_{j \in \mathbb{N}}$, $(f_j)_{j \in \mathbb{N}}$, $(\varphi_j)_{j \in \mathbb{N}}$, $\{g_j\}_{j \in \mathbb{N}}$ and $(\mathfrak{h}_j)_{j \in \mathbb{N}}$ linked through

$$\begin{cases} F_{\tau_j}(D^2u_j, x) = f_j(x) & \text{in } B_1(0', \nu_j) \cap \mathbb{R}_+^n \\ \mathcal{B}(x, u_j, Du_j) = g_j(x) & \text{on } B_1(0', \nu_j) \cap T_1 \\ u_j(x) = \varphi_j(x) & \text{on } \partial B_1(0', \nu) \cap \mathbb{R}_+^n \end{cases}$$

and

$$\begin{cases} F_j^\sharp(D^2\mathfrak{h}_j, 0) = 0 & \text{in } B_{\frac{7}{8}}(0', \nu_j) \cap \mathbb{R}_+^n \\ \mathcal{B}(x, \mathfrak{h}_j, D\mathfrak{h}_j) = g_j(x) & \text{on } B_{\frac{7}{8}}(0', \nu_j) \cap T_1 \\ \mathfrak{h}_j(x) = u_j(x) & \text{on } \partial B_{\frac{7}{8}}(0', \nu) \cap \mathbb{R}_+^n. \end{cases}$$

where τ_j , $\|\psi_{F_{\tau_j}^\sharp}(\cdot, 0)\|_{L^p(B_1^+)}$, $\|f_j\|_{L^p(B_1^+)}$ go to zero as $j \rightarrow \infty$ and such that

$$(2.2) \quad \|u_j - \mathfrak{h}_j\|_{L^\infty(B_{\frac{7}{8}}(0', \nu) \cap \mathbb{R}_+^n)} > \delta_0$$

Here, $\varphi_j \in C(\partial B_1(0', \nu_j))$ and $g_j \in C^{1,\alpha}(\overline{T}_2)$ satisfy $\|\varphi_j\|_{L^\infty(\partial B_1(0', \nu_j))} \leq \mathbf{c}_1$ and $\|g_j\|_{C^{1,\alpha}(\overline{T}_2)} \leq \mathbf{c}_2$, respectively. From Theorem 2.4, we have for all $0 < \rho < 1$

$$(2.3) \quad \|u_j\|_{C^{0,\alpha'}(\overline{B_{1-\rho}(0', \nu_j)} \cap \mathbb{R}_+^n)} \leq C(n, \lambda, \Lambda, \mathbf{c}_1, \mathbf{c}_2, \mu_0) \rho^{-\alpha'}$$

for some $\alpha' = \alpha'(n, \lambda, \Lambda, \mu_0)$ and sufficiently large j . Suppose that there exists a number ν_∞ and a subsequence $\{\nu_{j_k}\}$ such that $\nu_{j_k} \rightarrow \nu_\infty$ as $k \rightarrow +\infty$. We can assume that such a subsequence is monotone. If ν_{j_k} is decreasing, we can check that

$$B_1(0', \nu_\infty) \cap \mathbb{R}_+^n \subset B_1(0', \nu_{j_k}) \cap \mathbb{R}_+^n$$

for any k . Thus, we observe that

$$(2.4) \quad \|u_{j_k}\|_{C^{0,\alpha'}(\overline{B_{15/16}(0', \nu_\infty)} \cap \mathbb{R}_+^n)} \leq C(\mathbf{c}_1, \mathbf{c}_2, n, \lambda, \Lambda, \mu_0)$$

by making use (2.3). On the other hand, if ν_{j_k} is increasing, there exists a number k_0 such that

$$B_{31/32}(0', \nu_{k_j}) \cap \mathbb{R}_+^n \supset B_{15/16}(0', \nu_\infty) \cap \mathbb{R}_+^n \quad \text{for } k \geq k_0.$$

Then, we can deduce once again (2.4) for some proper subsequence u_{j_k} . Thus, by Arzelà-Ascoli's compactness criterium, there exist functions $u_\infty \in C^{0,\alpha}(\overline{B_{15/16}(0', \nu_\infty) \cap \mathbb{R}_+^n})$, $g_\infty \in C^0(\partial B_1^+)$ and subsequences such that $u_{j_k} \rightarrow u_\infty$ in $C^{0,\alpha'}(B_1^+)$ and $u_\infty = g_\infty$ on $B_{15/16}(0', \nu_\infty) \cap T_1$.

Since the functions $F_{j_k}^\sharp(\cdot, 0) \rightarrow F_\infty^\sharp(\cdot, 0)$ uniformly in compact sets of $Sym(n)$ and for every $\varphi \in C^2(\overline{B_2^+})$,

$$\begin{aligned} |F_{\tau_{j_k}}(D^2\varphi, x) - f_{j_k}(x) - F_\infty^\sharp(D^2\varphi, 0)| &\leq |F_{\tau_{j_k}}(D^2\varphi, x) - F_{j_k}^\sharp(D^2\varphi, x)| + |f_{j_k}| + \\ &\quad + |F_{j_k}^\sharp(D^2\varphi, x) - F_{j_k}^\sharp(D^2\varphi, 0)| + \\ &\quad + |F_{j_k}^\sharp(D^2\varphi, 0) - F_\infty^\sharp(D^2\varphi, 0)| \\ &\leq |F_{\tau_{j_k}}(D^2\varphi, x) - F_{j_k}^\sharp(D^2\varphi, x)| + |f_{j_k}| + \\ &\quad + \psi_{F_{j_k}^\sharp}(x, 0)(1 + |D^2\varphi|) \end{aligned}$$

then

$$\lim_{k \rightarrow +\infty} \|F_{\tau_{j_k}}(D^2\varphi, x) - f_{j_k}(x) - F_\infty^\sharp(D^2\varphi, 0)\|_{L^p(B_r(x_0))} = 0,$$

for any ball $B_r(x_0) \subset B_{15/16}(0, \nu_\infty) \cap \mathbb{R}_+^n$. Therefore, the Stability Lemma 2.5 ensures that u_∞ satisfies

$$\begin{cases} F_\infty^\sharp(D^2u_\infty, 0) = 0 & \text{in } B_{15/16}(0', \nu_\infty) \cap \mathbb{R}_+^n \\ \mathcal{B}(x, u_\infty, Du_\infty) = g_\infty(x) & \text{on } B_{15/16}(0', \nu_\infty) \cap T_1 \end{cases}$$

in the viscosity sense. Now, we consider $w_{j_k} := u_\infty - \mathfrak{h}_{j_k}$ for each k . Then, by [2, Lemma 4.3], w_{j_k} satisfies

$$\begin{cases} w_{j_k} \in \mathcal{S}(\frac{\lambda}{n}, \Lambda, 0) & \text{in } B_{7/8}(0', \nu_\infty) \cap \mathbb{R}_+^n \\ \mathcal{B}(x, w_{j_k}, Dw_{j_k}) = g_\infty(x) - g_{j_k}(x) & \text{on } B_{7/8}(0', \nu_\infty) \cap T_1 \\ w_{j_k}(x) = u_\infty(x) - u_{j_k}(x) & \text{on } \overline{\partial B_{7/8}(0', \nu_\infty) \cap \mathbb{R}_+^n}. \end{cases}$$

Using Lemma 2.6, we observe that

$$\begin{aligned} \|w_{j_k}\|_{L^\infty(B_{7/8}(0', \nu_\infty) \cap \mathbb{R}_+^n)} &\leq \|u_\infty - u_{j_k}\|_{L^\infty(\partial B_{7/8}(0', \nu_\infty))} + \\ &\quad + C(n, \lambda, \Lambda, \mu_0) \|g_\infty - g_{j_k}\|_{L^\infty(B_{7/8}(0', \nu_\infty) \cap T_1)} \rightarrow 0 \quad \text{as } k \rightarrow +\infty. \end{aligned}$$

This is, w_{j_k} converges uniformly to zero. This implies that \mathfrak{h}_{j_k} converges uniformly to u_∞ in $\overline{B_{7/8}(0', \nu_\infty) \cap \mathbb{R}_+^n}$, which contradicts (2.2) for $k \gg 1$. \square

3 $W^{2,p}$ estimates: Proof of Theorem 1.2

Before starting the proof of Theorem 1.2 we must introduce some useful terminologies: let

$$\mathcal{Q}_r^d(x_0) := \left(x_0 - \frac{r}{2}, x_0 + \frac{r}{2}\right)^d$$

be the d -dimension cube of side-length r and center x_0 . In the case $x_0 = 0$, we will write \mathcal{Q}_r^d . Furthermore, if $d = n$ we will write $\mathcal{Q}_r(x_0)$.

We will need later on the Calderón-Zygmund decomposition Lemma for cubes: Let $\mathcal{Q}_1^{n-1} \times (0, 1)$ unit cube. We split it into 2^n cubes of half side. We do the same splitting procedure with each one of these 2^n cubes, and we iterate such a process. Each cube in such a procedure is called a dyadic cube. Given two $\mathcal{Q}, \tilde{\mathcal{Q}} \neq \mathcal{Q}_1^{n-1} \times (0, 1)$ dyadic cubes, we say that \mathcal{Q} is a predecessor cube of $\tilde{\mathcal{Q}}$ if \mathcal{Q} is one of the 2^n cubes obtained in the partition of $\tilde{\mathcal{Q}}$. In the sequel, we may enunciate a key result whose proof can be found in Caffarelli-Cabr e's Book [9, Lemma 4.2].

Lemma 3.1 (Calderón-Zygmund cube decomposition). *Let $\mathcal{A} \subset \mathcal{B} \subset \mathcal{Q}_1^{n-1} \times (0, 1)$ be measurable sets and $\delta \in (0, 1)$ such that*

(a) $\mathcal{L}^n(\mathcal{A}) \leq \delta$;

(b) *If \mathcal{Q} is a dyadic cube such that $\mathcal{L}^n(\mathcal{A} \cap \mathcal{Q}) > \delta \mathcal{L}^n(\mathcal{Q})$, then $\tilde{\mathcal{Q}} \subset \mathcal{B}$.*

Then, $\mathcal{L}^n(\mathcal{A}) \leq \delta \mathcal{L}^n(\mathcal{B})$.

For $f \in L^1_{\text{loc}}(\mathbb{R}^n)$ recall that the *Hardy-Littlewood maximal operator* is defined as follows

$$\mathcal{M}(f)(x) := \sup_{\rho > 0} \int_{B_\rho(x)} |f(y)| dy.$$

We say that P_M is a paraboloid with *opening* $M > 0$ whenever $P_M(x) = \pm \frac{M}{2}|x|^2 + p_1 \cdot x + p_0$. Observe that such a paraboloid is a convex function in the case of “plus” sign and a concave function otherwise. Now, for $u \in C^0(\Omega)$, $\Omega' \subset \bar{\Omega}$ and $M > 0$ we define

$$\underline{\mathcal{G}}_M(u, \Omega') := \{x_0 \in \Omega' : \exists P_M \text{ concave paraboloid s. t. } P_M(x_0) = u(x_0), P_M(x) \leq u(x) \forall x \in \Omega'\}$$

and

$$\underline{\mathcal{A}}_M(u, \Omega') := \Omega' \setminus \underline{\mathcal{G}}_M(u, \Omega').$$

In a similar way, by using convex paraboloids we define $\bar{\mathcal{G}}_M(u, \Omega')$ and $\bar{\mathcal{A}}_M(u, \Omega')$ and the sets

$$\mathcal{G}_M(u, \Omega') := \underline{\mathcal{G}}_M(u, \Omega') \cap \bar{\mathcal{G}}_M(u, \Omega') \text{ and } \mathcal{A}_M(u, \Omega') := \underline{\mathcal{A}}_M(u, \Omega') \cap \bar{\mathcal{A}}_M(u, \Omega').$$

In addition, we define:

$$\bar{\Theta}(u, \Omega', x) := \inf \{M > 0; x \in \bar{\mathcal{G}}_M(u, \Omega')\}.$$

We also can define $\underline{\Theta}(u, \Omega', x)$. Finally, we define:

$$\Theta(u, \Omega', x) := \sup \{\underline{\Theta}(u, \Omega', x), \bar{\Theta}(u, \Omega', x)\}.$$

The first step towards $W^{2,p}$ -estimates up to the boundary are estimates for paraboloids on the boundary. Hence, in this section we will prove an appropriated power decay on the boundary for $\mathcal{L}^n(\mathcal{G}_t(u, \Omega))$. For this end, we will need the following standard technical result (cf. [9]):

Proposition 3.2. *Let $0 \leq h : \Omega \rightarrow \mathbb{R}$ be a measurable function, $\mu_h(t) := \mathcal{L}^n(\{x \in \Omega : h(x) \geq t\})$ its distribution function and $\eta > 0$, $M > 1$ constants. Then,*

$$h \in L^p(\Omega) \iff \sum_{j=1}^{\infty} M^{pj} \mu_h(\eta M^j) < \infty$$

for every $p \in (0, +\infty)$. Particularly, there exists a constant $C = C(n, \eta, M)$ such that

$$C^{-1} \cdot \left(\sum_{j=1}^{\infty} M^{pj} \cdot \mu_h(\eta M^j) \right)^{\frac{1}{p}} \leq \|h\|_{L^p(\Omega)} \leq C \left(\mathcal{L}^n(\Omega) + \sum_{j=1}^{\infty} M^{pj} \mu_h(\eta M^j) \right)^{\frac{1}{p}}.$$

Once we have such estimates for paraboloids at the boundary, the ideas will follow as the ones of Caffarelli’s original proof for the interior estimates (cf. [9]). In effect, consider the distribution function of Θ , i.e. $\mu_{\Theta, \Omega'}(t) := \mathcal{L}^n(\{x \in \Omega' : \Theta(x) > t\})$. Then, one check that $\mu_{\Theta, \Omega'}(t) = \mathcal{L}^n(\mathcal{A}_t(u, \Omega'))$. Moreover, as an application of Proposition 3.2 we have

$$\Theta(u, \Omega', \cdot) \in L^p(\Omega') \iff \sum_{j=1}^{\infty} M^{pj} \mathcal{L}^n(\mathcal{A}_{\eta M^j}(u, \Omega')) < \infty.$$

Finally, from [9, Proposition 1.1] we conclude

$$\|D^2u\|_{L^p(\Omega')} \leq C(\eta, M, p) \left(\mathcal{L}^n(\Omega) + \sum_{j=1}^{\infty} M^{pj} \mathcal{L}^n(\mathcal{A}_{\eta M^j}(u, \Omega')) \right).$$

Therefore, to obtain the desired $W^{2,p}$ -estimates in Ω' it is enough to prove the corresponding summability for $\sum_{j=1}^{\infty} M^{pj} \mathcal{L}^n(\mathcal{A}_{\eta M^j}(u, \Omega'))$.

We gather few elements involved in the proof of Theorem 1.2, although such results are well-known facts in the literature. We observe that such *a priori* estimate is independent of further assumptions on the operator F , and it follows merely from uniform ellipticity and the integrability of the source term. Thus, the proof is omitted in what follows. See [9, Lemma 7.8] and [47, Lemma 2.7] for details.

Proposition 3.3 (Power Decay on the boundary). *Let $u \in \mathcal{S}(\lambda, \Lambda, f)$ in $B_{12\sqrt{n}}^+ \subset \Omega \subset \mathbb{R}_+^n$, $u \in C^0(\Omega)$ and $\|u\|_{L^\infty(\Omega)} \leq 1$. Then, there exist universal constants $C > 0$ and $\delta > 0$ such that $\|f\|_{L^n(B_{12\sqrt{n}}^+)} \leq 1$ implies*

$$\mathcal{L}^n(\mathcal{A}_t(u, \Omega) \cap (\mathcal{Q}_1^{n-1} \times (0, 1) + x_0)) \leq C.t^{-\delta}$$

for any $x_0 \in B_{9\sqrt{n}} \cap \overline{\mathbb{R}_+^n}$ and $t > 1$.

Proposition 3.3 yields key information on the measure of $\mathcal{G}_M(u, \Omega) \cap ((\mathcal{Q}_1^{n-1} \times (0, 1)) + x_0)$, provided

$$\mathcal{G}_1(u, \Omega) \cap ((\mathcal{Q}_2^{n-1} \times (0, 2)) + x_0) \neq \emptyset.$$

The proof of Theorem 1.2 will be split into two main steps: Firstly, we will investigate equations governed by operators without dependence on the function and gradient entry. In the sequel, we reduce the analysis for operators with full dependence. The first step towards such estimates is the following Proposition.

Proposition 3.4. *Let u be a normalized viscosity solution of*

$$\begin{cases} F(D^2u, x) = f(x) & \text{in } B_1^+, \\ \mathcal{B}(x, u, Du) = g(x) & \text{on } T_1, \end{cases}$$

where $\beta, \gamma, g \in C^{1,\alpha}(T_1)$ with $\beta \cdot \vec{n} \geq \mu_0$ for some $\mu_0 > 0$, $\gamma \leq 0$ and $f \in L^p(B_1^+) \cap C^0(B_1^+)$, for $n \leq p < \infty$. Further, assume that assumptions (A1)-(A4) are in force. Then, there exist positive constants ψ_0 and r_0 depending on $n, \lambda, \Lambda, p, \mu_0, \alpha, \|\beta\|_{C^{1,\alpha}(\overline{T_1})}$ and $\|\gamma\|_{C^{1,\alpha}(\overline{T_1})}$ such that if

$$\left(\int_{B_r(x_0) \cap B_1^+} \psi_{F^\sharp}(x, x_0)^p dx \right)^{\frac{1}{p}} \leq \psi_0$$

for any $x_0 \in B_1^+$ and $r \in (0, r_0)$, then $u \in W^{2,p}(\overline{B_{\frac{1}{2}}^+})$ and

$$\|u\|_{W^{2,p}(\overline{B_{\frac{1}{2}}^+})} \leq C \cdot \left(\|u\|_{L^\infty(B_1^+)} + \|f\|_{L^p(B_1^+)} + \|g\|_{C^{1,\alpha}(\overline{T_1})} \right),$$

where $C = C(n, \lambda, \Lambda, \mu_0, p, \|\gamma\|_{C^{1,\alpha}(\overline{T_1})}, \alpha, r_0) > 0$.

In order to prove Proposition 3.4 we need a number of auxiliary results:

Proposition 3.5. *Assume that assumptions (A1)-(A4) there hold. Let $B_{14\sqrt{n}}^+ \subset \Omega \subset \mathbb{R}_+^n$ and u be a viscosity solution of*

$$\begin{cases} F_\tau(D^2u, x) &= f(x) & \text{in } B_{14\sqrt{n}}^+, \\ \mathcal{B}(x, u, Du) &= g(x) & \text{on } T_{14\sqrt{n}}. \end{cases}$$

Assume further that $\max \left\{ \|f\|_{L^n(B_{14\sqrt{n}}^+)}, \tau \right\} \leq \epsilon$. Finally, suppose for some $\tilde{x}_0 \in B_{9\sqrt{n}} \cap \{x_n \geq 0\}$ the following

$$\mathcal{G}_1(u, \Omega) \cap ((\mathcal{Q}_2^{n-1} \times (0, 2)) + \tilde{x}_0) \neq \emptyset.$$

Then,

$$\mathcal{L}^n(\mathcal{G}_M(u; \Omega) \cap ((\mathcal{Q}_1^{n-1} \times (0, 1)) + x_0)) \geq 1 - \epsilon_0,$$

where $x_0 \in B_{9\sqrt{n}} \cap \{x_n \geq 0\}$, $M > 1$ depends only on $n, \lambda, \Lambda, \mu_0, \alpha, \|\gamma\|_{C^{1,\alpha}(\overline{T}_{14\sqrt{n}})}, C_1$ (from Assumption A4) and $\|g\|_{C^{1,\alpha}(T_{14\sqrt{n}})}$, and $\epsilon_0 \in (0, 1)$.

Proof. Consider $x_1 \in \mathcal{G}_1(u, \Omega) \cap (\mathcal{Q}_2^{n-1} \times (0, 2) + \tilde{x}_0)$. Then, there exist paraboloids with opening $t = 1$ touching u at x_1 from above and below, this is,

$$-\frac{1}{2}|x - x_1|^2 \leq u(x) - \ell(x) \leq \frac{1}{2}|x - x_1|^2$$

for $x \in \Omega$ and an affine function ℓ .

Now, we set

$$v(x) = \frac{u(x) - \ell(x)}{C_*},$$

where $C_* > 0$ is a dimensional constant chosen (large enough) such that $\|v\|_{L^\infty(B_{12\sqrt{n}}^+)} \leq 1$ and

$$-|x|^2 \leq v(x) \leq |x|^2 \quad \text{in } \Omega \setminus B_{12\sqrt{n}}^+.$$

Next, observe that v is a viscosity solution to

$$\begin{cases} \tilde{F}_\tau(D^2v, x) &= \tilde{f}(x) & \text{in } B_{14\sqrt{n}}^+, \\ \mathcal{B}(x, v, Dv) &= \frac{1}{C_*}[g(x) - \beta \cdot D\ell - \gamma\ell] & \text{on } T_{14\sqrt{n}}. \end{cases}$$

where

$$\tilde{F}_\tau(X, x) := \frac{1}{C_*}F_\tau(C_*X, x) \quad \text{and} \quad \tilde{f}(x) := \frac{1}{C_*}f(x)$$

Now, consider \mathfrak{h} the function ϵ -close to u , coming from Lemma 2.12, this is, let $\mathfrak{h} \in C^{1,1}(B_{13\sqrt{n}}^+) \cap C^0(\overline{B_{13\sqrt{n}}^+})$ be solution of

$$\begin{cases} \tilde{F}^\sharp(D^2\mathfrak{h}, 0) &= 0 & \text{in } B_{13\sqrt{n}}^+, \\ \mathcal{B}(x, \mathfrak{h}, D\mathfrak{h}) &= \frac{1}{C_*}[g(x) - \beta(x) \cdot D\ell(x) - \gamma(x)\ell(x)] & \text{on } T_{13\sqrt{n}}. \end{cases}$$

such that

$$\|v - \mathfrak{h}\|_{L^\infty(B_{13\sqrt{n}}^+)} \leq 2$$

. Notice that $\beta \cdot D\ell \in C^{1,\alpha}(\overline{T}_{14\sqrt{n}})$ since $\beta \in C^{1,\alpha}(\overline{T}_{14\sqrt{n}})$ and $D\ell$ is a constant vector. Hence, ABP Maximum Principle (Lemma 2.6) assures that

$$\begin{aligned} \|\mathfrak{h}\|_{L^\infty(B_{13\sqrt{n}}^+)} &\leq \|v\|_{L^\infty(\partial B_{13\sqrt{n}}^+ \setminus T_{13\sqrt{n}})} + \frac{C}{C_*} \left[\|g\|_{L^\infty(\overline{T}_{13\sqrt{n}})} + |D\ell| \|\beta\|_{L^\infty(\overline{T}_{14\sqrt{n}})} + \|\gamma\ell\|_{L^\infty(\overline{T}_{13\sqrt{n}})} \right] \\ &\leq C(n, \|\ell\|_{L^\infty(\overline{T}_{14\sqrt{n}})}, \|\gamma\|_{C^{1,\alpha}(T_{14\sqrt{n}})}, \|g\|_{C^{1,\alpha}(\overline{T}_{14\sqrt{n}})}) \\ &:= \tilde{C} \end{aligned}$$

In consequence, condition (A4) ensures that

$$\|\mathfrak{h}\|_{C^{1,1}(\overline{B_{12\sqrt{n}}^+})} \leq C(C_1, \tilde{C}) \implies \mathcal{A}_N(\mathfrak{h}, B_{12\sqrt{n}}^+) \cap ((\mathcal{Q}_1^{n-1} \times (0, 1)) + x_0) = \emptyset$$

for some $N = N(C_1, \tilde{C}) > 1$, and we extended $\mathfrak{h}|_{B_{12\sqrt{n}}^+}$ (continuously) outside $B_{12\sqrt{n}}^+$ such that $\mathfrak{h} = v$ outside $B_{13\sqrt{n}}^+$ and $\|v - \mathfrak{h}\|_{L^\infty(\Omega)} = \|v - \mathfrak{h}\|_{L^\infty(B_{12\sqrt{n}}^+)}$. For this reason,

$$\|v - \mathfrak{h}\|_{L^\infty(\Omega)} \leq \overline{C}(C_1, \tilde{C})$$

and

$$-(\overline{C}(C_1, \tilde{C}) + |x|^2) \leq \mathfrak{h}(x) \leq \overline{C}(C_1, \tilde{C}) + |x|^2 \quad \text{in } \Omega \setminus B_{12\sqrt{n}}^+.$$

Therefore, there exists $M_0 = M_0(C_1, \tilde{C}) \geq N > 1$ for which

$$\mathcal{A}_{M_0}(\mathfrak{h}, \Omega) \cap ((\mathcal{Q}_1^{n-1} \times (0, 1)) + x_0) = \emptyset.$$

Summarizing,

$$(3.1) \quad (\mathcal{Q}_1^{n-1} \times (0, 1) + x_0) \subset \mathcal{G}_{M_0}(\mathfrak{h}, \Omega).$$

Now, we define

$$w(x) := \frac{\delta}{2C\epsilon}(v - \mathfrak{h})(x).$$

Therefore, w fulfills the assumptions of Proposition 3.3. Thus, we obtain for $t > 1$ the following

$$\mathcal{L}^n(\mathcal{A}_t(w, \Omega) \cap ((\mathcal{Q}_1^{n-1} \times (0, 1)) + x_0)) \leq Ct^{-\kappa}.$$

By using $\mathcal{A}_{2M_0}(u) \subset \mathcal{A}_{M_0}(w) \cup \mathcal{A}_{M_0}(\mathfrak{h})$ and (3.1) we conclude that

$$\mathcal{L}^n(\mathcal{G}_{2M_0}(v - \mathfrak{h}, \Omega) \cap ((\mathcal{Q}_1^{n-1} \times (0, 1)) + x_0)) \geq 1 - C\epsilon^{-\kappa}.$$

Finally, we conclude that

$$\mathcal{L}^n(\mathcal{G}_{2M_0}(v, \Omega) \cap ((\mathcal{Q}_1^{n-1} \times (0, 1)) + x_0)) \geq 1 - C\epsilon^{-\kappa}.$$

The proof finishes by choosing $\epsilon \ll 1$ in an appropriate manner and by setting $M \equiv 2M_0$. \square

Lemma 3.6. *Given $\epsilon_0 \in (0, 1)$ and let u be a normalized viscosity solution to*

$$\begin{cases} F_\tau(D^2u, x) = f(x) & \text{in } B_{14\sqrt{n}}^+, \\ \mathcal{B}(x, u, Du) = g(x) & \text{on } T_{14\sqrt{n}}. \end{cases}$$

Assume that (A1)-(A4) hold and extend f by zero outside $B_{14\sqrt{n}}^+$. For $x \in B_{14\sqrt{n}}$, let

$$\max \left\{ \tau, \|f\|_{L^n(B_{14\sqrt{n}})} \right\} \leq \epsilon$$

for some $\epsilon > 0$ depending only $n, \epsilon_0, \lambda, \Lambda, \mu_0, \alpha$. Then, for $k \in \mathbb{N} \setminus \{0\}$ we define

$$\begin{aligned} \mathcal{A} &:= \mathcal{A}_{M^{k+1}}(u, B_{14\sqrt{n}}^+) \cap (\mathcal{Q}_1^{n-1} \times (0, 1)) \\ \mathcal{B} &:= \left(\mathcal{A}_{M^k}(u, B_{14\sqrt{n}}^+) \cap (\mathcal{Q}_1^{n-1} \times (0, 1)) \right) \cup \{x \in \mathcal{Q}_1^{n-1} \times (0, 1); \mathcal{M}(f^n) \geq (C_0 M^k)^n\}, \end{aligned}$$

where $M = M(n, C_0) > 1$. Then,

$$\mathcal{L}^n(\mathcal{A}) \leq \epsilon_0(n, \epsilon, \lambda, \Lambda) \mathcal{L}^n(\mathcal{B}).$$

Proof. We will use Lemma 3.1. Observe that $\mathcal{A} \subset \mathcal{B} \subset (\mathcal{Q}_1^{n-1} \times (0, 1))$ and of the Proposition 3.5 we conclude that $\mathcal{L}^n(\mathcal{A}) \leq \delta < 1$ provided we choice $\delta = \epsilon_0$. Thus, it remains to show the following implication: for dyadic cubes \mathcal{Q}

$$\mathcal{L}^n(\mathcal{A} \cap \mathcal{Q}) > \epsilon_0 \mathcal{L}^n(\mathcal{Q}) \quad \Rightarrow \quad \tilde{\mathcal{Q}} \subset \mathcal{B}.$$

For that purpose, assume that for some $i \geq 1$, $\mathcal{Q} = \left(\mathcal{Q}_1^{n-1} \times (0, \frac{1}{2^i}) \right) + x_0$ is a dyadic cube with predecessor $\tilde{\mathcal{Q}} = \left(\mathcal{Q}_1^{n-1} \times (0, \frac{1}{2^{i-1}}) \right) + \tilde{x}_0$. Now, we assume that \mathcal{Q} satisfies

$$(3.2) \quad \mathcal{L}^n(\mathcal{A} \cap \mathcal{Q}) = \mathcal{L}^n\left(\mathcal{A}_{M^{k+1}}(u, B_{14\sqrt{n}}^+) \cap \mathcal{Q}\right) > \epsilon_0 \mathcal{L}^n(\mathcal{Q}),$$

however the inclusion $\tilde{\mathcal{Q}} \subseteq \mathcal{B}$ does not hold. In consequence, there must be $x_1 \in \tilde{\mathcal{Q}} \setminus \mathcal{B}$, i.e.,

$$(3.3) \quad x_1 \in \tilde{\mathcal{Q}} \cap \mathcal{G}_{M^k}(u, B_{14\sqrt{n}}^+) \quad \text{and} \quad \mathcal{M}(f^n)(x_1) < (C_0 M^k)^n.$$

In the sequel, we must split the analysis into two cases:

Case 1: If $|x_0 - (x'_0, 0)| < \frac{1}{2^{i-3}} \sqrt{n}$.

In this case, we define: $T(y) := (x'_0, 0) + \frac{1}{2^i} \cdot y$ and $\tilde{u} : \tilde{\Omega} \rightarrow \mathbb{R}$, where $\tilde{\Omega} = T^{-1}(\Omega)$, given by $\tilde{u}(y) := \frac{2^{2i}}{M^k} u(T(y))$. Now, observe that $\mathcal{Q} \subset (\mathcal{Q}_1^{n-1} \times (0, 1))$ implies that $B_{14\sqrt{n}/2^i}^+(x'_0, 0) \subset B_{14\sqrt{n}}^+$. Furthermore, such a \tilde{u} is a viscosity solution to

$$\begin{cases} \tilde{F}_\tau(D^2 \tilde{u}, y) &= \tilde{f}(x) \quad \text{in } B_{14\sqrt{n}}^+, \\ \tilde{B}(y, \tilde{u}, D\tilde{u}) &= \tilde{g}(x) \quad \text{on } T_{14\sqrt{n}}, \end{cases}$$

where

$$\begin{cases} \tilde{F}_\tau(X, q, r, y) &:= \frac{\tau}{M^k} F\left(\frac{M^k}{\tau} X, T(y)\right), \\ \tilde{f}(y) &:= \frac{1}{M^k} f(T(y)), \\ \tilde{B}(y, s, \vec{v}) &:= \beta(y) \cdot \vec{v} + \tilde{\gamma}(y)s, \\ \tilde{\beta}(y) &:= \beta(T(y)), \\ \tilde{\gamma}(y) &:= \frac{1}{2^i} \gamma(T(y)), \\ \tilde{g}(y) &:= \frac{2^i}{M^k} g(T(y)) \end{cases}$$

Now, notice that \tilde{F}^\sharp fulfills $C^{1,1}$ -estimates with the same constant as F^\sharp . Moreover, from (3.3) we obtain

$$\|\tilde{f}\|_{L^n(B_{14\sqrt{n}}^+)} \leq \frac{2^i}{M^k} \left(\int_{\mathcal{Q}_{\frac{28\sqrt{n}}{2^i}}(x_1)} |f(x)|^n dx \right)^{\frac{1}{n}} \leq 2^n C_0;$$

As a result, $\|\tilde{f}\|_{L^n(B_{14\sqrt{n}}^+)} \leq \epsilon$ provided we select C_0 small enough in (3.3). In addition, from (3.3) we conclude that

$$\mathcal{G}_1(\tilde{u}, T^{-1}(B_{14\sqrt{n}}^+)) \cap (\mathcal{Q}_2^{n-2} \times (0, 2) + 2^i(\tilde{x}_0 - (x'_0, 0))) \neq \emptyset.$$

Furthermore, $|x_0 - \tilde{x}_0| \leq \frac{1}{2^i} \sqrt{n}$ implies $|2^i(\tilde{x}_0 - (x'_0, 0))| < 9\sqrt{n}$.

Therefore, we have shown that the assumptions of Proposition 3.5 are in force. Hence, it follows

$$\mathcal{L}^n\left(\mathcal{G}_M(\tilde{u}, T^{-1}(B_{14\sqrt{n}}^+)) \cap ((\mathcal{Q}_1^{n-1} \times (0, 1)) + 2^i(x_0 - (x'_0, 0)))\right) \geq 1 - \epsilon_0.$$

Thus,

$$\mathcal{L}^n \left(\mathcal{G}_{M^{k+1}}(u; \mathbb{B}_{14\sqrt{n}}^+ \cap \mathcal{Q} \right) \geq (1 - \epsilon_0) \mathcal{L}^n(\mathcal{Q}),$$

which contradicts (3.2).

Case 2: If $|x_0 - (x'_0, 0)| \geq \frac{1}{2^{i-3}} \sqrt{n}$.

In this scenario, we conclude that $\mathbb{B}_{\frac{\sqrt{n}}{2^{i-3}}}(x_0 + \frac{1}{2^{i+1}} e_n) \subset \mathbb{B}_{8\sqrt{n}}^+$, where e_n is the n^{th} unit vector of canonical base. Now, by defining the transformation: $\mathbb{T}(y) := (x_0 + \frac{1}{2^{i+1}} e_n) + \frac{1}{2^i} y$, we proceed similarly to the first part of the proof. Finally, applying [35, Lemma 5.2] instead of Proposition 3.5 we obtain a contradiction to (3.2). This completes the proof of the Lemma. \square

In the sequel, we will establish the proof of Proposition 3.4.

Proof of Proposition 3.4. Fix $x_0 \in \mathbb{B}_{1/2} \cap \{x_n \geq 0\}$. When $x_0 \in \mathbb{T}_{\frac{1}{2}}$, $0 < r < \frac{1-|x_0|}{14\sqrt{n}}$ define

$$\kappa := \frac{\epsilon r}{\epsilon r^{-1} \|u\|_{L^\infty(\mathbb{B}_{14r\sqrt{n}}^+(x_0))} + \|f\|_{L^n(\mathbb{B}_{14r\sqrt{n}}^+(x_0))} + \epsilon r^{-1} \|g\|_{C^{1,\alpha}(\mathbb{T}_{14r\sqrt{n}}(x_0))}}$$

where the constant $\epsilon = \epsilon(n, \epsilon_0, \lambda, \Lambda, p, \mu_0, \alpha, \|\beta\|_{C^{1,\alpha}(\mathbb{T}_1)})$ is as the one in Proposition 3.5 and $\epsilon_0 \in (0, 1)$ will be determined *a posteriori*. Now, we define: $\tilde{u}(y) := \frac{\kappa}{r^2} u(x_0 + ry)$. Then, \tilde{u} is a normalized viscosity solution to

$$\begin{cases} \tilde{F}(D^2 \tilde{u}, y) &= \tilde{f}(x) \quad \text{in } \mathbb{B}_{14\sqrt{n}}^+, \\ \tilde{\mathcal{B}}(y, \tilde{u}, D\tilde{u}) &= \tilde{g}(x) \quad \text{on } \mathbb{T}_{14\sqrt{n}}. \end{cases}$$

where

$$\begin{cases} \tilde{F}(X, y) &:= \kappa F\left(\frac{1}{\kappa} X, ry + x_0\right) \\ \tilde{f}(y) &:= \kappa f(x_0 + ry) \\ \tilde{\mathcal{B}}(y, s, \vec{v}) &:= \tilde{\beta}(y) \cdot \vec{v} + \tilde{\gamma}(y) s \\ \tilde{\beta}(y) &:= \beta(x_0 + ry) \\ \tilde{\gamma}(y) &:= r\gamma(x_0 + ry) \\ \tilde{g}(y) &:= \frac{\kappa}{r} g(x_0 + ry). \end{cases}$$

Hence, \tilde{F} fulfills (A1)-(A4). Moreover,

$$(3.4) \quad \|\tilde{f}\|_{L^n(\mathbb{B}_{14\sqrt{n}}^+)} = \frac{\kappa}{r} \|f\|_{L^n(\mathbb{B}_{14r\sqrt{n}}^+)} \leq \epsilon \quad \text{and} \quad \|\tilde{\beta}\|_{C^{1,\alpha}(\mathbb{T}_{14\sqrt{n}})} \leq 1$$

which ensures that hypotheses of Lemma 3.6 are in force. Now, let $M > 0$ and $C_0 > 0$ be as in the Lemma 3.6 and choose $\epsilon_0 := \frac{1}{2M^p}$. Now, for $k \geq 0$ we define

$$\begin{aligned} \alpha_k &:= \mathcal{L}^n \left(\mathcal{A}_{M^k}(\tilde{u}, \mathbb{B}_{14\sqrt{n}}^+) \cap (\mathcal{Q}_1^{n-1} \times (0, 1)) \right) \\ \beta_k &:= \mathcal{L}^n \left(\left\{ x \in (\mathcal{Q}_1^{n-1} \times (0, 1)); \mathcal{M}(\tilde{f}^n)(x) \geq (C_0 M^k)^n \right\} \right). \end{aligned}$$

Thus, Lemma 3.6 implies that $\alpha_{k+1} \leq \epsilon_0 (\alpha_k + \beta_k)$ and consequently

$$(3.5) \quad \alpha_k \leq \epsilon_0^k + \sum_{i=0}^{k-1} \epsilon_0^{k-i} \beta_i.$$

On the other hand, from assumption (A2), we have $\tilde{f}^n \in L^{\frac{n}{n-1}}(\mathbb{B}_{14\sqrt{n}}^+)$, consequently $\mathcal{M}(\tilde{f}^n) \in L^{\frac{n}{n-1}}(\mathbb{B}_{14\sqrt{n}}^+)$. Thus, by (3.4)

$$\|\mathcal{M}(\tilde{f}^n)\|_{L^{\frac{n}{n-1}}(\mathbb{B}_{14\sqrt{n}}^+)} \leq C(n, p) \|\tilde{f}^n\|_{L^{\frac{n}{n-1}}(\mathbb{B}_{14\sqrt{n}}^+)} \leq C(n, p) \|\tilde{f}\|_{L^p(\mathbb{B}_{14\sqrt{n}}^+)}^n \leq C.$$

Therefore, by Proposition 3.2 we obtain

$$(3.6) \quad \sum_{k=1}^{\infty} M^{pk} \beta_k \leq C(n, p).$$

Finally, taking into account the choice of ϵ_0 , (3.5) and (3.6) we conclude that

$$\sum_{k=1}^{\infty} M^{pk} \alpha_k \leq \sum_{k=1}^{\infty} 2^{-k} + \left(\sum_{k=0}^{\infty} M^{pk} \beta_k \right) \cdot \left(\sum_{k=1}^{\infty} M^{pk} \epsilon_0^k \right) \leq C(n, p),$$

which implies that $\|D^2 \tilde{u}\|_{L^p(\overline{\mathbb{B}_{1/2}^+})} \leq C(n, p, M)$ and consequently

$$\|D^2 u\|_{L^p(\overline{\mathbb{B}_{\frac{1}{2}}^+(x_0)})} \leq C(n, \lambda, \Lambda, p, r) \left(\|u\|_{L^\infty(\mathbb{B}_1^+)} + \|f\|_{L^p(\mathbb{B}_1^+)} + \|g\|_{C^{1,\alpha}(\overline{\mathbb{T}_1})} \right).$$

On the other hand, if $x_0 \in \mathbb{B}_{1/2}^+$, we can apply the result of interior estimates (cf. [35, Theorem 6.1], see also [12]). Finally, by combining interior and boundary estimates, we obtain the desired results by using a standard covering argument. This finishes proof of the Proposition. \square

Corollary 3.7. *Let u be a bounded viscosity solution of*

$$\begin{cases} F(D^2 u, Du, u, x) = f(x) & \text{in } \mathbb{B}_1^+, \\ \mathcal{B}(x, u, Du) = g(x) & \text{on } \mathbb{T}_1, \end{cases}$$

where $\beta, \gamma, g \in C^{1,\alpha}(\mathbb{T}_1)$ with $\beta \cdot \vec{n} \geq \mu_0$ for some $\mu_0 > 0$, $\gamma \leq 0$. Further, assume that (A1)-(A4) are in force. Then, there exists a constant $C > 0$ depending on $n, \lambda, \Lambda, p, C_1$, such that $u \in W^{2,p}(\mathbb{B}_{\frac{1}{2}}^+)$ and

$$\|u\|_{W^{2,p}(\mathbb{B}_{\frac{1}{2}}^+)} \leq C \cdot \left(\|u\|_{L^\infty(\mathbb{B}_1^+)} + \|f\|_{L^p(\mathbb{B}_1^+)} + \|g\|_{C^{1,\alpha}(\overline{\mathbb{T}_1})} \right).$$

Proof. Notice that u is also a viscosity solution of

$$\begin{cases} \tilde{F}(D^2 u, x) = \tilde{f}(x) & \text{in } \mathbb{B}_1^+, \\ \mathcal{B}(x, u, Du) = g(x) & \text{on } \mathbb{T}_1. \end{cases}$$

where $\tilde{F}(X, x) := F(X, 0, 0, x)$ and \tilde{f} is a function satisfying

$$|\tilde{f}| \leq \sigma |Du| + \xi |u| + |f|.$$

Thus, we can apply Proposition 3.4 and conclude that

$$(3.7) \quad \|u\|_{W^{2,p}(\mathbb{B}_{\frac{1}{2}}^+)} \leq C \cdot \left(\|u\|_{L^\infty(\mathbb{B}_1^+)} + \|\tilde{f}\|_{L^p(\mathbb{B}_1^+)} + \|g\|_{C^{1,\alpha}(\overline{\mathbb{T}_1})} \right).$$

Now, by proceeding similarly as in [2, Corollary 4.5] we conclude that $u \in W^{1,p}(\mathbb{B}_1^+)$ and

$$(3.8) \quad \|u\|_{W^{1,p}(\mathbb{B}_1^+)} \leq C \cdot \left(\|u\|_{L^\infty(\mathbb{B}_1^+)} + \|f\|_{L^p(\mathbb{B}_1^+)} \right).$$

Finally, combining (3.7) and (3.8) we complete the desired estimate. \square

Corollary 3.8. *Let u be a bounded L^p -viscosity solution of*

$$\begin{cases} F(D^2u, Du, u, x) = f(x) & \text{in } B_1^+, \\ \mathcal{B}(x, u, Du) = g(x) & \text{on } T_1, \end{cases}$$

where $\beta, \gamma, g \in C^{1,\alpha}(T_1)$ with $\beta \cdot \vec{n} \geq \mu_0$ for some $\mu_0 > 0$, $\gamma \leq 0$ and $f \in L^p(B_1^+)$, for $n < p < \infty$. Further, assume that F^\sharp satisfies (A4) and F fulfills the condition (SC). Then, there exist positive constants $\beta_0 = \beta_0(n, \lambda, \Lambda, p)$, $r_0 = r_0(n, \lambda, \Lambda, p)$ and $C = C(n, \lambda, \Lambda, p, r_0) > 0$, such that if

$$\left(\int_{B_r(x_0) \cap B_1^+} \psi_{F^\sharp}(x, x_0)^p dx \right)^{\frac{1}{p}} \leq \psi_0$$

for any $x_0 \in B_1^+$ and $r \in (0, r_0)$, then $u \in W^{2,p}(\overline{B_{\frac{r}{2}}^+})$ and

$$\|u\|_{W^{2,p}(\overline{B_{\frac{r}{2}}^+})} \leq C \cdot \left(\|u\|_{L^\infty(B_1^+)} + \|f\|_{L^p(B_1^+)} + \|g\|_{C^{1,\alpha}(\overline{T_1})} \right).$$

Proof. It is enough to prove the result for equations without dependence on Du and u (see [47, Theorem 4.3] for details). We will approximate f in L^p by functions $f_j \in C^\infty(\overline{B_1^+}) \cap L^p(B_1^+)$ such that $f_j \rightarrow f$ in $L^p(B_1^+)$ and also approximate g by a sequence (g_j) in $C^{1,\alpha}(T_1)$ with the property that $g_j \rightarrow g$ in $C^{1,\alpha}(T_1)$. By Theorems of Uniqueness and Existence (Theorem 2.8 and 2.9), there exists a sequence of functions $u_j \in C^0(\overline{B_1^+})$, which they are viscosity solutions of following family of PDEs:

$$\begin{cases} F(D^2u_j, x) = f_j(x) & \text{in } B_1^+, \\ \mathcal{B}(x, u_j, Du_j) = g_j(x) & \text{on } T_1 \\ u_j(x) = u(x) & \text{on } \partial B_1^+ \setminus T_1. \end{cases}$$

Therefore, the assumptions of the Theorem 3.4 are in force. As a result,

$$\|u_j\|_{W^{2,p}(\overline{B_{\frac{r}{2}}^+})} \leq C(\text{universal}) \cdot \left(\|u_j\|_{L^\infty(B_1^+)} + \|f_j\|_{L^p(B_1^+)} + \|g_j\|_{C^{1,\alpha}(\overline{T_1})} \right),$$

for a constant $C > 0$. Furthermore, a standard covering argument yields $u_j \in W^{2,p}(B_1^+)$. From Lemma 2.6, $(u_j)_{j \in \mathbb{N}}$ is uniformly bounded in $W^{2,p}(\overline{B_\rho^+})$ for $\rho \in (0, 1)$.

Once again, we can apply Lemma 2.6 and obtain

$$\|u_j - u_k\|_{L^\infty(B_1^+)} \leq C(n, \lambda, \Lambda, \mu_0) (\|f_j - f_k\|_{L^p(B_1^+)} + \|g_j - g_k\|_{C^{1,\alpha}(\overline{T_1})}).$$

Thus, $u_j \rightarrow u_\infty$ in $C^0(\overline{B_1^+})$. Moreover, since $(u_j)_{j \in \mathbb{N}}$ is bounded in $W^{2,p}(B_{\frac{1}{2}}^+)$ we obtain $u_j \rightarrow u_\infty$ weakly in $W^{2,p}(B_{\frac{1}{2}}^+)$. Thus,

$$\|u_\infty\|_{W^{2,p}(\overline{B_{1/2}^+})} \leq C \cdot \left(\|u_\infty\|_{L^\infty(B_1^+)} + \|f\|_{L^p(B_1^+)} + \|g\|_{C^{1,\alpha}(\overline{T_1})} \right).$$

Finally, stability results (see Lemma 2.5) ensure that u_∞ is an L^p -viscosity solution to

$$\begin{cases} F(D^2u_\infty, x) = f(x) & \text{in } B_1^+, \\ \mathcal{B}(x, u_\infty, Du_\infty) = g(x) & \text{on } T_1 \\ u_\infty(x) = u(x) & \text{on } \partial B_1^+ \setminus T_1. \end{cases}$$

Thus, $w := u_\infty - u$ fulfills

$$\begin{cases} w \in S(\frac{\lambda}{n}, \Lambda, 0) & \text{in } B_1^+, \\ \mathcal{B}(x, w, Dw) = 0 & \text{on } T_1 \\ w = 0 & \text{on } \partial B_1^+ \setminus T_1. \end{cases}$$

which by Lemma 2.6 we conclude that $w = 0$ in $\overline{B_1^+} \setminus T_1$. By continuity, $w = 0$ in $\overline{B_1^+}$ which finishes the proof. \square

Finally, we are now in position to prove the Theorem 1.2.

Proof of Theorem 1.2. Based on standard reasoning (cf. [47]) it is important to highlight that is always possible to perform a change of variables in order to flatten the boundary. For this end, consider $x_0 \in \partial\Omega$. Since $\partial\Omega \in C^{2,\alpha}$ there exists a neighborhood of x_0 , which we will label $\mathcal{V}(x_0)$ and a $C^{2,\alpha}$ -diffeomorphism $\Phi : \mathcal{V}(x_0) \rightarrow \mathbb{B}_1(0)$ such that $\Phi(x_0) = 0$ and $\Phi(\Omega \cap \mathcal{V}(x_0)) = \mathbb{B}_1^+$. Now, for $\tilde{\varphi} \in W^{2,p}(\mathbb{B}_1^+)$ we set $\varphi = \tilde{\varphi} \circ \Phi \in W^{2,p}(\mathcal{V}(x_0))$. Thus, we obtain that

$$D\varphi = (D\tilde{\varphi} \circ \Phi)D\Phi \quad \text{and} \quad D^2\varphi = D\Phi^t \cdot (D^2\tilde{\varphi} \circ \Phi) \cdot D\Phi + ((D\tilde{\varphi} \circ \Phi)\partial_{ij}\Phi)_{1 \leq i,j \leq n}.$$

Moreover, $\tilde{u} := u \circ \Phi^{-1} \in C^0(\mathbb{B}_1^+)$. Next, observe that \tilde{u} is an L^p -viscosity solution to

$$\begin{cases} \tilde{F}(D^2\tilde{u}, D\tilde{u}, \tilde{u}, y) &= \tilde{f}(x) \quad \text{in } \mathbb{B}_1^+, \\ \tilde{B}(y, \tilde{u}, D\tilde{u}) &= \tilde{g}(x) \quad \text{on } \mathbb{T}_1. \end{cases}$$

where for $y = \Phi^{-1}(x)$ we have

$$\begin{cases} \tilde{F}(D^2\tilde{\varphi}, D\tilde{\varphi}, x) &:= F(D\Phi^t(y)D^2\tilde{\varphi}D\Phi(y) + D\tilde{\varphi}D^2\Phi(y), D\tilde{\varphi}D\Phi(y), \tilde{u}, y) \\ \tilde{f}(x) &:= f \circ \Phi^{-1}(x) \\ \tilde{B}(x, s, \vec{v}) &:= \tilde{\beta}(x) \cdot \vec{v} + \tilde{\gamma}(x)s \\ \tilde{\beta}(x) &:= (\beta \circ \Phi^{-1}) \cdot (D\Phi \circ \Phi^{-1})^t \\ \tilde{\gamma}(x) &:= (\gamma \circ \Phi^{-1}) \cdot (D\Phi \circ \Phi^{-1})^t \\ \tilde{g}(x) &:= g \circ \Phi^{-1} \end{cases}$$

Furthermore, note that $\tilde{F}(X, \varsigma, \eta, y) = F(D\Phi^t(y) \cdot X \cdot D\Phi(y) + \varsigma D^2\Phi, \varsigma D\Phi(y), \eta, y)$ is a uniformly elliptic operator with ellipticity constants $\lambda C(\Phi)$, $\Lambda C(\Phi)$. Thus,

$$\tilde{F}^\sharp(X, \varsigma, \eta, x) = F^\sharp(D\Phi^t(\Phi^{-1}(x)) \cdot X \cdot D\Phi(\Phi^{-1}(x)) + \varsigma D^2\Phi(\Phi^{-1}(x)), 0, 0, \Phi^{-1}(x)).$$

In consequence, we conclude that $\psi_{\tilde{F}^\sharp}(x, x_0) \leq C(\Phi)\psi_{F^\sharp}(x, x_0)$, which ensures that \tilde{F} falls into the assumptions of Corollary 3.8. This finishes the proof, as well as establishes the proof of the Theorem 1.2. \square

4 BMO type estimates: Proof of Theorem 1.4

This section will be devoted to establishing boundary BMO type estimates. Before proving Theorem 1.4 we will present some key results which play a crucial role in our strategy.

Lemma 4.1 (Approximation Lemma II). *Let $g \in C^{1,\psi}(\mathbb{T}_1)$ such that $\|g\|_{C^{1,\psi}(\mathbb{T}_1)} \leq C_g$ and $\beta \in C^{1,\psi}(\overline{\mathbb{T}_1})$. Let F and F^∞ be $(\frac{\lambda}{n}, \Lambda)$ -elliptic operators. Given $\delta > 0$, there exists $\epsilon_0 = \epsilon_0(\delta, n, \lambda, \Lambda, C_g) < 1$ such that, if*

$$\max \left\{ \frac{|F(X, x) - F^\infty(X, x_0)|}{\|X\| + 1}, \psi_{F^\infty}(x), \|f\|_{L^p(\mathbb{B}_1^+) \cap p\text{-BMO}(\mathbb{B}_1^+)} \right\} \leq \epsilon_0,$$

then any two L^p -viscosity solutions u and v of

$$\begin{cases} F(D^2u, x) &= f(x) \quad \text{in } \mathbb{B}_1^+ \\ \beta \cdot Du &= g(x) \quad \text{on } \mathbb{T}_1. \end{cases} \quad \text{and} \quad \begin{cases} F^\infty(D^2\mathfrak{h}, x_0) &= 0 \quad \text{in } \mathbb{B}_1^+ \\ \beta \cdot D\mathfrak{h} &= g(x) \quad \text{on } \mathbb{T}_1. \end{cases}$$

satisfy

$$\|u - \mathfrak{h}\|_{L^\infty(\mathbb{B}_{\frac{1}{8}}^+)} \leq \delta.$$

Proof. The proof holds the same lines from Lemma 2.12 along with minor modifications. \square

Lemma 4.2 (A quadratic approximation). *Under the assumptions from Lemma 4.1, assume that F^∞ satisfies $(A_4)^*$. Let u be an L^p -viscosity solution to*

$$\begin{cases} F(D^2u, x) = f(x) & \text{in } B_1^+, \\ \beta \cdot Du = g(x) & \text{on } T_1. \end{cases}$$

Then, there exist universal constants $C_\# > 0$ and $r_0 \in (0, 1/2)$ and a quadratic polynomial \mathfrak{P} , with $\|\mathfrak{P}\|_{L^\infty(B_1^+)} \leq C_\#$ such that

$$\sup_{B_r^+} |u(x) - \mathfrak{P}(x)| \leq r^2 \quad \text{for every } r \leq r_0$$

Proof. Firstly, take $\delta \in (0, 1)$, to be chosen *a posteriori*, and apply Lemma 4.1 to obtain $\epsilon_0 > 0$ and an L^p -viscosity solution to

$$\begin{cases} F^\infty(D^2\mathfrak{h}, x) = 0 & \text{in } B_1^+, \\ \beta \cdot D\mathfrak{h} = g(x) & \text{on } T_1. \end{cases}$$

such that

$$\sup_{B_1^+} |u - \mathfrak{h}| \leq \delta.$$

Remember that F^∞ fulfills assumption $(A_4)^*$, then \mathfrak{h} enjoys classical estimates. Hence, its Taylor's expansion of second order, namely \mathfrak{P} , is well-defined. Furthermore, we have

$$\sup_{B_\rho^+} |\mathfrak{h} - \mathfrak{P}| \leq C_* \rho^{2+\psi} \quad \forall \rho \leq r_0.$$

Now, we select $0 < r \ll 1$ (small enough) in such a way that

$$r \leq \min \left\{ r_0, \left(\frac{1}{10C_*} \right)^{\frac{1}{\psi}} \right\}.$$

Thus,

$$\sup_{B_r^+} |\mathfrak{h} - \mathfrak{P}| \leq \frac{1}{10} r^2.$$

On the other hand, we choose $\delta := \frac{1}{10} r^2$. As a result, we obtain

$$\sup_{B_r^+} |u - \mathfrak{h}| \leq \frac{1}{10} r^2.$$

Finally, by combining the previous inequalities, we conclude that

$$\sup_{B_r^+} |u - \mathfrak{P}| \leq r^2.$$

□

As a byproduct of the above result, we can to yield a quadratic approximation at the recession level. The proof holds along straightforward modifications in the Lemma 4.2. For this reason, we will omit it here.

Corollary 4.3. *Assume the assumptions from Lemma 4.1. Then, there exist universal constants $C_\# > 0$, $\tau_0 > 0$ and $r > 0$, such that if u is a (normalized) viscosity solution of*

$$\begin{cases} F_\tau(D^2u, x) = f(x) & \text{in } B_1^+, \\ \beta \cdot Du = g(x) & \text{on } T_1. \end{cases}$$

with $\max \left\{ \tau, \|f\|_{p\text{-BMO}(\mathbb{B}_1^+)} \right\} \leq \tau_0$, there exists a quadratic polynomial \mathfrak{P} , with $\|\mathfrak{P}\|_\infty \leq C_\sharp$ satisfying

$$\sup_{\mathbb{B}_1^+} |u(x) - \mathfrak{P}(x)| \leq r^2.$$

In face of such auxiliary results, we are already ready to prove Theorem 1.4.

Proof of Theorem 1.4. In fact, for $\kappa \in (0, 1)$ to be determined *a posteriori*, we define $v(x) := \kappa u(x)$ such that v is a (normalized) L^p -viscosity solution to

$$\begin{cases} F_\tau(D^2v, x) = \tilde{f}(x) & \text{in } \mathbb{B}_1^+, \\ \beta \cdot Dv = \tilde{g}(x) & \text{on } \mathbb{T}_1. \end{cases}$$

where $\tau := \kappa$, $\tilde{f}(x) := \kappa f(x)$ and $\tilde{g}(x) := \kappa g(x)$. Now, we are going to determine κ . This will be performed in such a way that $\max \left\{ \tau, \|\tilde{f}\|_{p\text{-BMO}(\mathbb{B}_1^+)} \right\} \leq \tau_0$, where τ_0 comes from Corollary 4.3. Finally, we will establish the result for v , which will lead to the statement of the Theorem.

From now on, we will show that there exists a sequence of second order polynomials $(\mathfrak{P}_k)_{k \in \mathbb{N}}$ given by $\mathfrak{P}_k(x) = \frac{1}{2}x^T \cdot \mathcal{A}_k \cdot x + \mathcal{B}_k \cdot x + \mathcal{C}_k$ such that

$$(4.1) \quad F^\sharp(\mathcal{A}_k, x) = (\tilde{f})_1,$$

$$(4.2) \quad r^{2(k-1)}|\mathcal{A}_k - \mathcal{A}_{k-1}| + r^{k-1}|\mathcal{B}_{k-1} - \mathcal{B}_k| + |\mathcal{C}_{k-1} - \mathcal{C}_k| \leq C_\sharp \cdot r^{2(k-1)},$$

where $C_\sharp > 0$ is a universal constant, $0 < r \ll 1$ comes from Lemma 4.2 and

$$\sup_{\mathbb{B}_{r,k}^+} |v(x) - \mathfrak{P}_k(x)| \leq r^{2k}.$$

The proof will follow by an induction process. Let us put \mathfrak{P}_0 and \mathfrak{P}_{-1} to be $\mathfrak{P}_0(x) = \mathfrak{P}_{-1}(x) = \frac{1}{2}x^T \cdot X_0 \cdot x$, where $X_0 \in \text{Sym}(n)$ fulfills $F^\sharp(X_0, x) = (\tilde{f})_1$.

The first step of the argument, i.e., the case $k = 0$, is naturally verified. Now, suppose we have established the existence of such polynomials for $k = 0, 1, \dots, j$. Then, defining the following auxiliary function $v_j : \mathbb{B}_1^+ \rightarrow \mathbb{R}$ given by

$$v_j(x) := \frac{(v - \mathfrak{P}_j)(r^j x)}{r^{2j}},$$

where we have by the induction assumption that v_j is a (normalized) L^p -viscosity solution to

$$\begin{cases} F_j(D^2v_j, x) = \tilde{f}_j(x) & \text{in } \mathbb{B}_1^+ \\ \tilde{\beta}(x) \cdot Dv_j(x) = \tilde{g}_j(x) & \text{on } \mathbb{T}_1. \end{cases}$$

where

$$\begin{cases} F_j(X, x) & := \tau F\left(\frac{1}{\tau}(X + \mathcal{A}_j), r^j x\right) \\ \tilde{f}_j(x) & := \tilde{f}(r^j x) \\ \tilde{g}_j(x) & := \tilde{g}(r^j x) - \frac{1}{r^j} \tilde{\beta}(x) \cdot D\mathfrak{P}_j(r^j x) \\ \tilde{\beta}(x) & := r^j \beta(r^j x) \end{cases}$$

with

$$\begin{aligned} \|f_j\|_{p\text{-BMO}(\mathbb{B}_1^+)} &= \sup_{0 < s \leq 1} \left(\int_{\mathbb{B}_s^+} |f_j(x) - (f_j)_s|^p dx \right)^{\frac{1}{p}} \\ &= \sup_{0 < s \leq 1} \left(\int_{\mathbb{B}_{sr}^+} |f(z) - (f)_{sr}|^p dz \right)^{\frac{1}{p}} \\ &\leq \|\tilde{f}\|_{p\text{-BMO}(\mathbb{B}_1^+)} \\ &\leq \tau_0. \end{aligned}$$

Furthermore, since $F^\sharp(\mathcal{A}_j, x) = (\tilde{f})_1$, the PDE with oblique boundary conditions

$$\begin{cases} F_j^\sharp(D^2\mathfrak{h}, x) = (\tilde{f})_1 & \text{in } \mathbb{B}_1^+ \\ \beta \cdot D\mathfrak{h}(x) = \tilde{g}(x) & \text{on } \mathbb{T}_1 \end{cases}$$

satisfies the same (up to the boundary) $C^{2,\psi}$ a priori estimates from the problem driven by F^\sharp , and it is under the assumption of Corollary 4.3. Thus, there exists a quadratic polynomial $\tilde{\mathfrak{P}}$ with $\|\tilde{\mathfrak{P}}\|_\infty \leq C_\sharp$ such that

$$(4.3) \quad \sup_{\mathbb{B}_r^+} |v_j - \tilde{\mathfrak{P}}| \leq r^2.$$

Rewriting (4.3) back to the original domain yields

$$\sup_{\mathbb{B}_{r^{k+1}}^+} \left| v(x) - \left[\mathfrak{P}_k(x) + r^{2k} \tilde{\mathfrak{P}}\left(\frac{x}{r^k}\right) \right] \right| \leq r^{2(k+1)}.$$

Finally, by defining $\mathfrak{P}_{j+1}(x) := \mathfrak{P}_j(x) + r^{2j} \tilde{\mathfrak{P}}(r^{-j}x)$ we check the $(k+1)$ th step of induction. Furthermore, the required conditions (4.1) and (4.2) are satisfied. In conclusion, for $\rho > 0$, choose k an integer such that $r^{k+1} < \rho \leq r^k$. Then, by applying Theorem 1.2 to v_k , we obtain

$$\begin{aligned} \sup_{\rho \in (0,1/2)} \left(\int_{\mathbb{B}_\rho^+} |D^2v(z) - \mathcal{A}_k|^p dz \right)^{\frac{1}{p}} &\leq \sup_{\rho \in (0,1/2)} \left(\frac{1}{r^n} \int_{\mathbb{B}_{r^k}^+} |D^2v(z) - \mathcal{A}_k|^p dz \right)^{\frac{1}{p}} \\ &= \sup_{\rho \in (0,1/2)} \left(\frac{1}{r^n} \int_{\mathbb{B}_1^+} |D^2v_k|^p dx \right)^{\frac{1}{p}} \\ &\leq C(\text{universal}). \end{aligned}$$

Now, recall the general inequality:

$$\int_{\mathbb{B}_\rho^+} \left| D^2v - \int_{\mathbb{B}_\rho^+} D^2v \, dy \right|^p dx \leq 2^p \int_{\mathbb{B}_\rho^+} |D^2v - \mathcal{A}_k|^p dx.$$

In effect, by triangular inequality,

$$\begin{aligned} \left(\int_{\mathbb{B}_\rho^+} \left| D^2v - \int_{\mathbb{B}_\rho^+} D^2v \, dy \right|^p dx \right)^{\frac{1}{p}} &\leq \left(\int_{\mathbb{B}_\rho^+} |D^2v - \mathcal{A}_k|^p dx \right)^{\frac{1}{p}} + \left| \int_{\mathbb{B}_\rho^+} D^2v \, dy - \mathcal{A}_k \right| \\ &\leq \left(\int_{\mathbb{B}_\rho^+} |D^2v - \mathcal{A}_k|^p dx \right)^{\frac{1}{p}} + \int_{\mathbb{B}_\rho^+} |D^2v - \mathcal{A}_k| dx \\ &\leq 2 \left(\int_{\mathbb{B}_\rho^+} |D^2v - \mathcal{A}_k|^p dx \right)^{\frac{1}{p}}, \end{aligned}$$

Therefore,

$$\|Dv\|_{p\text{-BMO}\left(\overline{\mathbb{B}_{\frac{1}{2}}^+}\right)} := \sup_{\rho \in (0,1/2)} \left(\int_{\mathbb{B}_\rho^+} \left| D^2v - \int_{\mathbb{B}_\rho^+} D^2v \, dy \right|^p dx \right)^{\frac{1}{p}} \leq C(\text{universal}),$$

thereby finishing the proof of the estimate (1.8). \square

5 Obstacle type problems: Proof of Theorem 1.6

In this section we will study an important class of free boundary problems, namely the obstacle problem with oblique boundary datum (1.9) for a given obstacle $\phi \in W^{2,p}(\Omega)$ satisfying $\mathcal{B}(x, u, D\phi) \geq g$ a.e. on $\partial\Omega$, and F a uniformly elliptic operator fulfilling $F(\mathcal{O}_{n \times n}, \vec{0}, 0, x) = 0$. The main purpose will be establish a $W^{2,p}$ -regularity theory for (1.9), whose strategy does not require any extra regularity assumption on nonlinearity F .

In the sequel, similarly to [4, Theorem 3.3] and [13, Appendix] we are going to reduce the analysis of the obstacle problem (1.9) in studying a family of penalized problems as follows

$$(5.1) \quad \begin{cases} F(D^2u_\varepsilon, Du_\varepsilon, u_\varepsilon, x) &= h^+(x)\Psi_\varepsilon(u_\varepsilon - \phi) + f(x) - h^+(x) & \text{in } \Omega \\ \mathcal{B}(x, u_\varepsilon, Du_\varepsilon) &= g(x) & \text{on } \partial\Omega, \end{cases}$$

for $\varepsilon \in (0, 1)$, where $\Psi_\varepsilon(s)$ is a smooth function such that

$$\Psi_\varepsilon(s) \equiv 0 \quad \text{if } s \leq 0; \quad \Psi_\varepsilon(s) \equiv 1 \quad \text{if } s \geq \varepsilon,$$

$$0 \leq \Psi_\varepsilon(s) \leq 1 \quad \text{for any } s \in \mathbb{R}.$$

and

$$h(x) := f(x) - F(D^2\phi, D\phi, \phi, x).$$

Now, we define $\hat{f}_{u_\varepsilon}(x) := h^+(x)\Psi_\varepsilon(u_\varepsilon - \phi) + f(x) - h^+(x)$. Then, note that, $h \in L^p(\Omega)$ with the estimate

$$(5.2) \quad \begin{aligned} \|h\|_{L^p(\Omega)} &\leq \|f\|_{L^p(\Omega)} + \|F(D^2\phi, D\phi, \phi, x)\|_{L^p(\Omega)} \\ &\leq C \cdot (\|f\|_{L^p(\Omega)} + \|\phi\|_{W^{2,p}(\Omega)}) \end{aligned}$$

for some $C = C(n, \lambda, \Lambda, \sigma, \xi, p) > 0$, since F fulfills (A1). In consequence, we have

$$(5.3) \quad \|\hat{f}_{u_\varepsilon}\|_{L^p(\Omega)} \leq C(n, \lambda, \Lambda, \sigma, \xi, p) (\|f\|_{L^p(\Omega)} + \|\phi\|_{W^{2,p}(\Omega)})$$

Next, we claim that the problem (5.1) admits a viscosity solution that enjoys *a priori* estimates. In effect, according to Perron's Method (see Lieberman's Book [32, Theorem 7.19]) it follows that for each $v_0 \in L^p(\Omega)$ fixed, there exists a unique viscosity solution $u_\varepsilon \in W^{2,p}(\Omega)$ satisfying in the viscosity sense

$$\begin{cases} F(D^2u_\varepsilon, Du_\varepsilon, u_\varepsilon, x) &= h^+(x)\Psi_\varepsilon(v_0 - \phi) + f(x) - h^+(x) & \text{in } \Omega \\ \mathcal{B}(x, u_\varepsilon, Du_\varepsilon) &= g(x) & \text{on } \partial\Omega, \end{cases}$$

fulfilling (due to Theorem 1.2) the estimate

$$(5.4) \quad \|u_\varepsilon\|_{W^{2,p}(\Omega)} \leq C \cdot \left(\|u_\varepsilon\|_{L^\infty(\Omega)} + \|\hat{f}_{v_0}\|_{L^p(\Omega)} + \|g\|_{C^{1,\alpha}(\partial\Omega)} \right)$$

for some $C = C(n, \lambda, \Lambda, p, \mu_0, \sigma, \omega, \|\beta\|_{C^2(\partial\Omega)}, \|\gamma\|_{C^2(\partial\Omega)}, \theta_0, \text{diam}(\Omega)) > 0$. Moreover, as in (5.3), we have $\hat{f}_{v_0} \in L^p(\Omega)$.

Now, from ABP estimate (Lemma 2.6) we obtain

$$(5.5) \quad \begin{aligned} \|u_\varepsilon\|_{W^{2,p}(\Omega)} &\leq C \cdot \left(\|\hat{f}_{v_0}\|_{L^p(\Omega)} + \|g\|_{C^{1,\alpha}(\partial\Omega)} \right) \\ &\leq C \cdot \left(\|f\|_{L^p(\Omega)} + \|h\|_{L^p(\Omega)} + \|g\|_{C^{1,\alpha}(\partial\Omega)} \right) \\ &\leq \hat{C} \cdot \left(\|f\|_{L^p(\Omega)} + \|g\|_{C^{1,\alpha}(\partial\Omega)} + \|\phi\|_{W^{2,p}(\Omega)} \right) \end{aligned}$$

Thus,

$$\|u_\varepsilon\|_{W^{2,p}(\Omega)} \leq C_0(\hat{C}, n, \lambda, \Lambda, p, \mu_0, \sigma, \omega, \|\beta\|_{C^2(\partial\Omega)}, \theta_0, \text{diam}(\Omega), \|f\|_{L^p(\Omega)}, \|g\|_{C^{1,\alpha}(\partial\Omega)}, \|\phi\|_{W^{2,p}(\Omega)}),$$

where $C_0 > 0$ is independent on v_0 .

In this point, by defining the operator $\mathcal{T} : L^p(\Omega) \rightarrow W^{2,p}(\Omega) \subset L^p(\Omega)$ given by $\mathcal{T}(v_0) = u_\varepsilon$, we conclude that \mathcal{T} maps the C_0 -ball (in $L^p(\Omega)$) into itself. Hence, \mathcal{T} is a compact operator. Therefore, by Schauder's fixed point theorem, there exists u_ε such that $\mathcal{T}(u_\varepsilon) = u_\varepsilon$, which is a viscosity solution to (5.1).

Finally, we are in a position to establish our main result.

Proof of Theorem 1.6. From (5.5) we observe that

$$\|u_\varepsilon\|_{W^{2,p}(\Omega)} \leq C \cdot (\|f\|_{L^p(\Omega)} + \|g\|_{C^{1,\alpha}(\partial\Omega)} + \|\phi\|_{W^{2,p}(\Omega)})$$

for some $C = C(n, \lambda, \Lambda, p, \mu_0, \sigma, \xi, \|\beta\|_{C^2(\partial\Omega)}, \text{diam}(\Omega), \theta_0)$. This ensures that $\{u_\varepsilon\}_{\varepsilon>0}$ is uniformly bounded in $W^{2,p}(\Omega)$. Hence, by standard compactness arguments, we can find a subsequence $\{u_{\varepsilon_j}\}_{j \in \mathbb{N}}$ with $\varepsilon_j \rightarrow 0$ and a function $u_\infty \in W^{2,p}(\Omega)$ such that

$$\begin{cases} u_{\varepsilon_j} \rightharpoonup u_\infty & \text{in } W^{2,p}(\Omega) \\ u_{\varepsilon_j} \rightarrow u_\infty & \text{in } C^{0,\alpha_0}(\overline{\Omega}) \\ u_{\varepsilon_j} \rightarrow u_\infty & \text{in } C^{1,1-\frac{n}{p}}(\overline{\Omega}) \end{cases}$$

for some universal $\alpha_0 \in (0, 1)$.

Now, we claim u_∞ is indeed a viscosity solution of (1.9). In effect:

- ✓ Since $\beta(x) \cdot Du_{\varepsilon_j}(x) + \gamma(x)u_{\varepsilon_j}(x) = g(x)$ and $\{u_{\varepsilon_j}\}_{j \in \mathbb{N}}$ are uniformly bounded and equi-continuous on $\partial\Omega$, then from (5.5) and Morrey embedding $W^{2,p}(\Omega) \hookrightarrow C^{1,1-\frac{n}{p}}$, we have

$$\beta(x) \cdot Du_\infty(x) + \gamma(x)u_\infty(x) = g(x) \quad \text{on } \partial\Omega$$

in the viscosity sense (and point-wisely).

- ✓ On the other hand, from (5.1), we have that

$$\begin{aligned} F(D^2u_{\varepsilon_j}, Du_{\varepsilon_j}, u_{\varepsilon_j}, x) &= h^+(x)\Psi_{\varepsilon_j}(u_{\varepsilon_j} - \phi) + f(x) - h^+(x) \\ &\leq f(x) \quad \text{in } \Omega \quad \text{for each } j \in \mathbb{N}. \end{aligned}$$

Thus, by Stability results (Lemma 2.5), by passing to the limit $j \rightarrow +\infty$, we conclude

$$F(D^2u_\infty, Du_\infty, u_\infty, x) \leq f \quad \text{in } \Omega.$$

- ✓ Next, we are going to prove that

$$u_\infty \geq \phi \quad \text{in } \overline{\Omega}.$$

Firstly, we see that $\Psi_{\varepsilon_j}(u_{\varepsilon_j} - \phi) \equiv 0$ on the set $\mathcal{O}_j := \{x \in \overline{\Omega} : u_{\varepsilon_j}(x) < \phi(x)\}$. Note that, if $\mathcal{O}_j = \emptyset$, we have nothing to prove. Thus, we suppose that $\mathcal{O}_j \neq \emptyset$. Then,

$$F(D^2u_{\varepsilon_j}, Du_{\varepsilon_j}, u_{\varepsilon_j}, x) = f(x) - h^+(x) \quad \text{for } x \in \mathcal{O}_j.$$

Now, note that \mathcal{O}_j is relatively open in $\overline{\Omega}$ for each $j \in \mathbb{N}$ since $u_{\varepsilon_j} \in C(\overline{\Omega})$. Moreover, from definition of h we have

$$F(D^2\phi, D\phi, \phi, x) = f(x) - h(x) \geq F(D^2u_{\varepsilon_j}, Du_{\varepsilon_j}, u_{\varepsilon_j}, x) \quad \text{in } \mathcal{O}_j$$

Moreover, we have $u_{\varepsilon_j} = \phi$ on $\partial\mathcal{O}_j \setminus \partial\Omega$. Thus, from the Comparison Principle (see [10, Theorem 2.10] and [32, Theorem 7.17]) we conclude that $u_{\varepsilon_j} \geq \phi$ in \mathcal{O}_j , which yields a contradiction. Therefore, $\mathcal{O}_j = \emptyset$ and $u_\infty \geq \phi$ in $\overline{\Omega}$.

✓ Finally, it remains to show that

$$F(D^2u_\infty, Du_\infty, u_\infty, x) = f(x) \text{ in } \{u_\infty > \phi\}$$

in the viscosity sense. For this end, notice that for each $k \in \mathbb{N}$ we have

$$h^+(x)\Psi_{\varepsilon_j}(u_{\varepsilon_j} - \phi) + f(x) - h^+(x) \rightarrow f(x) \text{ a.e. in } \left\{x \in \Omega : u_\infty(x) > \phi(x) + \frac{1}{k}\right\}.$$

Therefore, via Stability results (Lemma 2.5) we conclude (in the viscosity sense) that

$$F(D^2u_\infty, Du_\infty, u_\infty, x) = f(x) \text{ in } \{u_\infty > \phi\} = \bigcup_{k=1}^{\infty} \left\{u_\infty > \phi + \frac{1}{k}\right\} \text{ as } j \rightarrow +\infty,$$

thereby proving the claim.

In conclusion, from (5.5) u_∞ satisfies the following regularity estimate

$$\begin{aligned} \|u_\infty\|_{W^{2,p}(\Omega)} &\leq \liminf_{j \rightarrow +\infty} \|u_{\varepsilon_j}\|_{W^{2,p}(\Omega)} \\ &\leq C(n, \lambda, \Lambda, p, \mu_0, \sigma, \omega, \|\beta\|_{C^2(\partial\Omega)}, \|\gamma\|_{C^2(\partial\Omega)}, \text{diam}(\Omega), \theta_0) \cdot (\|f\|_{L^p(\Omega)} + \|g\|_{C^{1,\alpha}(\partial\Omega)} + \|\phi\|_{W^{2,p}(\Omega)}) \end{aligned}$$

thereby finishing the proof of the Theorem. \square

As a consequence, we obtain the uniqueness of existing solutions.

Corollary 5.1 (Uniqueness). *The viscosity solution found in the Theorem 1.6 is unique.*

Proof. Indeed, let u and v be two viscosity solutions of (1.9). Assume that $u \neq v$. Then, we may suppose without loss of generality that

$$\mathcal{O}_\sharp = \{v > u\} \neq \emptyset.$$

Since $v > u \geq \phi$ in \mathcal{O}_\sharp , we obtain in the viscosity sense

$$F(D^2v, Dv, v, x) = f(x) \text{ in } \mathcal{O}_\sharp$$

Hence, we conclude that

$$\begin{cases} F(D^2u, Du, u, x) \leq f(x) \leq F(D^2v, Dv, v, x) & \text{in } \mathcal{O}_\sharp \\ u(x) = v(x) & \text{on } \partial\mathcal{O}_\sharp \setminus \partial\Omega, \\ \mathcal{B}(x, u, Du) = \mathcal{B}(x, v, Dv) & \text{on } \partial\mathcal{O}_\sharp \cap \partial\Omega \end{cases}$$

Therefore, by the Comparison Principle for problems with oblique boundary conditions [10, Theorem 2.10] and [32, Theorem 7.17], we conclude that $u \geq v$ in \mathcal{O}_\sharp whether $\partial\mathcal{O}_\sharp \cap \partial\Omega = \emptyset$ or not. However, this contradicts the definition of the set \mathcal{O}_\sharp , thereby proving that $u = v$. \square

6 Final conclusions: Density of viscosity solutions

In this final part we will deliver another application to $W^{2,p}$ regularity estimates.

Theorem 6.1 ($W^{2,p}$ density on the class C^0 -viscosity solutions). *Let u be a C^0 -viscosity solution of*

$$\begin{cases} F(D^2u, x) = f(x) & \text{in } B_1^+ \\ \mathcal{B}(x, u, Du) = g(x) & \text{on } T_1, \end{cases}$$

where $f \in L^p(B_1^+) \cap C^0(B_1^+)$ (for $n \leq p < \infty$), $\beta, \gamma, g \in C^{1,\alpha}(T_1)$ with $\gamma \leq 0$ and $\beta \cdot \vec{n} \geq \mu_0$ on T_1 , for some $\mu_0 > 0$. Then, for any $\delta > 0$, there exists a sequence $(u_j)_{j \in \mathbb{N}} \subset W_{loc}^{2,p}(B_1^+) \cap \mathcal{S}(\lambda - \delta, \Lambda + \delta, f)$ converging local uniformly to u .

Proof. The proof follows some ideas from [35, Theorem 8.1] adapted to oblique boundary scenery. We will present the details for completeness to the reader. Firstly, we are going to build up such a desired sequence of operators $F_j : \text{Sym}(n) \times \mathbb{B}_1^+ \rightarrow \mathbb{R}$ as follow: Given $\delta > 0$ we consider the Pucci Maximal operator

$$\mathcal{L}_\delta(X) := \mathcal{P}_{(\lambda-\delta), (\Lambda+\delta)}^+(X) = (\Lambda + \delta) \sum_{e_i > 0} e_i(X) + (\lambda - \delta) \sum_{e_i < 0} e_i(X),$$

where $e_i(X)$ are the eigenvalues of matrix $X \in \text{Sym}(n)$. Now, we define

$$F_j : \text{Sym}(n) \times \mathbb{B}_1^+ \rightarrow \mathbb{R} \\ (X, x) \mapsto \max\{F(X, x), \mathcal{L}_\delta(X) - C_j\},$$

for $(C_j)_{j \in \mathbb{N}}$ a sequence of divergence positive number, which will be chosen precisely *a posteriori*. Firstly, notice that F_j is continuous (since it is the maximum between two continuous functions) and uniformly elliptic with ellipticity constants $\lambda - \delta \leq \Lambda + \delta$, because F and \mathcal{L}_δ are too. Now, taking into account the (λ, Λ) -ellipticity of F there holds

$$\begin{aligned} F(X, x) &\geq \lambda \sum_{e_i > 0} e_i(X) + \Lambda \sum_{e_i < 0} e_i \\ &\geq \lambda \sum_{e_i > 0} e_i(X) - \Lambda \|X\| \\ &= \mathcal{L}_\delta(X) - (\Lambda + \delta - \lambda) \sum_{e_i > 0} e_i - (\lambda - \delta) \sum_{e_i < 0} e_i - \Lambda \|X\| \\ &\geq \mathcal{L}_\delta(X) - (2\Lambda - \lambda + \delta) \|X\| \\ &\geq \mathcal{L}_\delta(X) - C_j, \quad \forall B_j \subset \text{Sym}(n), \quad \forall x \in \mathbb{B}_1^+, \end{aligned}$$

where we select $C_j := j(2\Lambda - \lambda + \delta)$. Hence, $F \equiv F_j$ in $B_j \times \mathbb{B}_1^+ \subset \text{Sym}(n) \times \mathbb{B}_1^+$. On the other hand, we analyse the recession operator associated to F_j . For that purpose, for each $\mu > 0$ we have

$$(F_j)_\mu(X, x) = \mu F_j(\mu^{-1}X, x) = \max\{F_\mu(X, x), L_\delta(X) - \mu C_j\}.$$

Since F_μ is (λ, Λ) -elliptic for any $\mu > 0$ we get

$$\begin{aligned} F_\mu(X, x) &\leq \Lambda \sum_{e_i > 0} e_i(X) + \lambda \sum_{e_i < 0} e_i(X) \\ &= \mathcal{L}_\delta(X) - \delta \sum_{e_i > 0} e_i(X) + \delta \sum_{e_i < 0} e_i(X) \\ &\leq \mathcal{L}_\delta(X) - \delta \|X\| \\ &\leq \mathcal{L}_\delta(X) - \mu C_j, \end{aligned}$$

for all $X \in B_{\frac{\mu C_j}{\delta}} \subset \text{Sym}(n)$ and $x \in \mathbb{B}_1^+$. Therefore, we conclude that $F_j(X, x) = \mathcal{L}_\delta(X) - C_j$ outside a ball of radius $\sim C_j$, then $F_j^\sharp = \mathcal{L}_\delta$. Now, notice that F_j^\sharp fulfills $C^{1,1}$ *a priori* estimates, i.e., given $g_0 \in C^{1,\alpha}(\mathbb{T}_1)$ any viscosity solution to

$$\begin{cases} F_j^\sharp(D^2\mathfrak{h}, x_0) &= 0 & \text{in } \mathbb{B}_1^+ \\ \mathcal{B}(x, \mathfrak{h}, D\mathfrak{h}) &= g_0(x) & \text{on } \mathbb{T}_1 \end{cases}$$

satisfies $\mathfrak{h} \in C^{1,1}(\overline{\mathbb{B}_{1/2}^+})$ (see e.g. [29, Teorema 1.3]) with the following estimate

$$\|\mathfrak{h}\|_{C^{1,1}(\overline{\mathbb{B}_{1/2}^+})} \leq C \cdot \left(\|\mathfrak{h}\|_{L^\infty(\mathbb{B}_1^+)} + \|g\|_{C^{1,\alpha}(\mathbb{T}_1)} \right).$$

Therefore, statements of Proposition 3.4 are in force. Then, for each $j \in \mathbb{N}$ fixed, any viscosity solution of

$$\begin{cases} F_j(D^2v, x) = f(x) & \text{in } B_1^+ \\ \mathcal{B}(x, v, Dv) = g(x) & \text{on } T_1 \end{cases}$$

enjoys $W^{2,p}$ regularity estimates. Precisely, for each $j \in \mathbb{N}$ there exists a constant $\kappa_j > 0$ such that

$$\|v\|_{W^{2,p}(B_{1/2}^+)} \leq \kappa_j \cdot (\|v\|_{L^\infty(B_1^+)} + \|f\|_{L^p(B_1^+)} + \|g\|_{C^{1,\alpha}(\overline{T_1})}).$$

Finally, we build up the desired sequence $(u_j)_{j \in \mathbb{N}}$ to be a viscosity solution to

$$\begin{cases} F_j(D^2u_j, x) = f(x) & \text{in } B_1^+ \\ \mathcal{B}(x, u_j, Du_j) = g(x) & \text{on } T_1 \\ u(x) = u_j(x) & \text{on } \partial B_1^+ \setminus T_1, \end{cases}$$

whose existence does hold from Theorem 2.9. Furthermore, each $u_j \in W^{2,p}(B_1^+)$ and since operators F_j are uniformly elliptic, it follow that $F_j(\cdot, x) \rightarrow F_0(\cdot, x)$ local uniformly in $\text{Sym}(n)$ for each $x \in B_1^+$. Moreover, since $F_j \equiv F$ in $B_j \times B_1^+$ then $F_0 \equiv F$. In conclusion, by local $C^{0,\alpha}$ regularity and ABP estimates (see Theorem 2.4 and Lemma 2.6 respectively) the sequence $(u_j)_{j \in \mathbb{N}}$, converge, up to a subsequence, local uniformly to u_0 in the $C^{0,\alpha}$ -topology. Furthermore, by Stability results (Lemma 2.5) u_0 is a viscosity solution of

$$\begin{cases} F(D^2u_0, x) = f(x) & \text{in } B_1^+ \\ \mathcal{B}(x, u_0, Du_0) = g(x) & \text{on } T_1 \\ u(x) = u_0(x) & \text{on } \partial B_1^+ \setminus T_1. \end{cases}$$

Finally, by taking $w = u_0 - u$ we obtain that w satisfies in the viscosity sense

$$\begin{cases} w \in \mathcal{S}(\lambda/n, \Lambda, 0) & \text{in } B_1^+ \\ \mathcal{B}(x, w, Dw) = 0 & \text{on } T_1 \\ w = 0 & \text{on } \partial B_1^+ \setminus T_1, \end{cases}$$

and one more time, by ABP estimates (Lemma 2.6) we conclude that $w = 0$ in $\overline{B_1^+} \setminus T_1$. Therefore, $w \equiv 0$, i.e., $u = u_0$, thereby finishing the proof. \square

Acknowledgments

J.S. Bessa was partially supported by CAPES-Brazil under Grant No. 88887.482068/2020-00. J.V. da Silva, M. N. Barreto Frederico and G.C. Ricarte have been partially supported by CNPq-Brazil under Grants No. 310303/2019-2, No 165746/2020-3 and No. 304239/2021-6. Part of this work was developed during the *Fortaleza Conference on Analysis and PDEs (2022)* at Universidade Federal do Ceará (UFC-Brazil). J.V. da Silva would like to thanks to UFC's Department of Mathematics by fostering a pleasant scientific and research atmosphere during his visit in the Summer of 2022.

References

- [1] Azzam, J. and Bedrossian, J. *Bounded mean oscillation and the uniqueness of active scalar equations*. Trans. Amer. Math. Soc. 367 (2015), no. 5, 3095-3118.
- [2] Byun, S.S. and Han, J. *$W^{2,p}$ -estimates for fully nonlinear elliptic equations with oblique boundary conditions*. J. Differential Equations 268 (2020), no. 5, 2125-2150.

- [3] Byun, S.S., Han J. and Oh, J., *On $W^{2,p}$ -estimates for solutions of obstacle problems for fully nonlinear elliptic equations with oblique boundary conditions*. Arxiv Preprint. <https://arxiv.org/pdf/2012.07442.pdf>.
- [4] Byun, S.S., Lee, K.-A., Oh, J. and Park, J. *Nondivergence elliptic and parabolic problems with irregular obstacles*. Math. Z. 290 (2018), no. 3-4, 973-990.
- [5] Byun, S.-S.; Lee, M. and Palagachev, D.K. *Hessian estimates in weighted Lebesgue spaces for fully nonlinear elliptic equations*. J. Differential Equations 260 (2016), no. 5, 4550-4571.
- [6] Byun, S.-S.; Oh, J. and Wang, L. *$W^{2,p}$ estimates for solutions to asymptotically elliptic equations in nondivergence form*. J. Differential Equations 260 (2016), no. 11, 7965-7981.
- [7] Byun, S.S. and Wang, L. *Global Calderón-Zygmund theory for asymptotically regular nonlinear elliptic and parabolic equations*. Int. Math. Res. Not. IMRN 2015, no. 17, 8289-8308.
- [8] Caffarelli, L.A. *Interior a priori estimates for solutions of fully nonlinear equations*. Ann. of Math.(2) 130 (1989), no. 1, 189–213.
- [9] Caffarelli, L.A. and Cabré, X. *Fully Nonlinear Elliptic Equations*. American Mathematical Society Colloquium Publications, 43. American Mathematical Society, Providence, RI, 1995. vi+104 pp. ISBN: 0-8218-0437-5.
- [10] Caffarelli, L.A., Crandall, M.G., Kocan, M. and Świąch, A. *On viscosity solutions of fully nonlinear equations with measurable ingredients*. Comm, Pure Appl. Math. 49 (1996) (4), 365–397.
- [11] Chipot, M. and Evans, L.C. *Linearisation at infinity and Lipschitz estimates for certain problems in the calculus of variations*. Proc. Roy. Soc. Edinburgh Sect. A. 102 (3-4) (1986).
- [12] da Silva, J.V. and Ricarte, G.C. *An asymptotic treatment for non-convex fully nonlinear elliptic equations: Global Sobolev and BMO type estimates*. Commun. Contemp. Math. 21 (2019), no. 7, 1850053, 28 pp.
- [13] da Silva, J.V. and Vivas, H. *Sharp regularity for degenerate obstacle type problems: a geometric approach*. Discrete Contin. Dyn. Syst. 41 (2021), no. 3, 1359-1385.
- [14] da Silva, J.V. and Vivas, H. *The obstacle problem for a class of degenerate fully nonlinear operators*. Rev. Mat. Iberoam. 37 (2021), no. 5, 1991-2020.
- [15] Dong, H. and Kim, D. *On the impossibility of $W^{2,p}$ estimates for elliptic equations with piecewise constant coefficients*. J. Funct. Anal. 267 (10) (2014) 3963–3974.
- [16] Dong, H. and Li, Z. *Classical solutions of oblique derivative problem in nonsmooth domains with mean Dini coefficients*. Trans. Amer. Math. Soc. 373 (2020), no. 7, 4975-4997.
- [17] Escauriaza, L. *$W^{2,n}$ a priori estimates for solutions to fully non-linear elliptic equation*. Indiana Univ. Math. J. 42, no. 2 (1993), 413-423.
- [18] Evans, L. C., *Classical solutions of fully nonlinear, convex, second-order elliptic equations*. Comm. Pure Appl. Math. 35(3), 333-363, 1982.
- [19] Figalli, A. and Shahgholian, H., *A general class of free boundary problems for fully nonlinear elliptic equations*. Arch. Ration. Mech. Anal. 213 (2014), no. 1, 269–286.
- [20] Foss, M. *Global regularity for almost minimizers of nonconvex variational problems*. Ann. Mat. Pura Appl. (4) 187 (2) (2008), 263-321.

- [21] Giga, Y. and Sato, M.-H. *On semicontinuous solutions for general Hamilton-Jacobi equations.* Comm. Partial Differential Equations 26 (2001), 813-839.
- [22] Ishii, H. and Lions, P.L. *Fully nonlinear oblique derivative problems for nonlinear second-order elliptic PDEs.* Duke math. J. 62 (3) (1991) 633-661.
- [23] Kim, S. and Lee, K-A. *Higher order convergence rates in theory of homogenization II: Oscillatory initial data.* Adv. Math. 362 (2020), 106960, 52 pp.
- [24] Krylov, N.V. *Boundedly nonhomogeneous elliptic and parabolic equations.* Izv. Akad. Nauk. SSSR Ser. Mat. 46 (1982), 487-523; English transl. in Math USSR Izv. 20 (1983), 459-492.
- [25] Krylov, N.V. *On the existence of smooth solutions for fully nonlinear elliptic equations with measurable "coefficients" without convexity assumptions.* Methods Appl. Anal., 19 (2012), pp. 119–146.
- [26] Krylov, N.V. *On the existence of $W^{2,p}$ solutions for fully nonlinear elliptic equations under relaxed convexity assumptions.* Comm. Partial Diff. Eqs., 38 (2013), pp. 687–710.
- [27] Krylov, N.V. *On the existence of $W^{2,p}$ solutions for fully nonlinear elliptic equations under either relaxed or no convexity assumptions.* Commun. Contemp. Math. 19 (2017), no. 6, 1750009, 39 pp.
- [28] Lee, M. *Weighted Orlicz regularity estimates for fully nonlinear elliptic equations with asymptotic convexity.* Commun. Contemp. Math. 21 (2019), no. 4, 1850024, 29 pp.
- [29] Li, D. and Zhang, K. *Regularity for fully nonlinear elliptic equations with oblique boundary conditions.* Arch. Ration. Mech. Anal. 228(3)(2018) 923-967.
- [30] Lieberman, G.M. *Regularity of solutions of obstacle problems for elliptic equations with oblique boundary conditions.* Pacific J. Math. 201 (2001), no. 2, 389–419.
- [31] Lieberman, G.M. *Higher regularity for nonlinear oblique derivative problems in Lipschitz domains.* Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) 1 (2002), no. 1, 111-151.
- [32] Lieberman, G.M. *Oblique derivative problems for elliptic equations.* World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2013. xvi+509 pp. ISBN: 978-981-4452-32-8.
- [33] Lieberman, G.M. and Trudinger, N.S. *Nonlinear oblique boundary value problems for nonlinear elliptic equations.* Trans. Amer. Math. Soc. 295 (1986), no. 2, 509-546.
- [34] Milakis, E. and Silvestre, L. *Regularity for fully nonlinear elliptic equations with Neumann boundary data.* Commun. Partial Differ. Equ. 31 (7-9) (2006) 1227-1252.
- [35] Pimentel, E. and Teixeira, E.V. *Sharp hessian integrability estimates for nonlinear elliptic equations: an asymptotic approach.* J. Math. Pures Appl. 106 (2016), pp. 744–767.
- [36] Raymond, J.P. *Lipschitz regularity of solutions of some asymptotically convex problems.* Proc. Roy. Soc. Edinburgh Sect. A. 117 (1-2) (1991), 59-73.
- [37] Ricarte G.C. and Teixeira, E.V. *Fully nonlinear singularly perturbed equations and asymptotic free boundary.* J. Funct. Anal., vol. 261, Issue 6, 2011, 1624-1673.
- [38] Ros-Oton, X. and Serra, J. *The structure of the free boundary in the fully nonlinear thin obstacle problem.* Adv. Math. 316 (2017), 710-747.
- [39] Safonov, M.V. *Nonlinear elliptic equations of second order*, Lecture notes. Dipartimento di Matematica Applicata "G. Sansone", Universita degli studi di Firenze, 1991, 52 p.

- [40] Safonov, M.V. *On the boundary value problems for fully nonlinear elliptic equations of second order*. CMA Research Report MRR 049-94, 1994, 34 p.
- [41] Savin, O. *Small perturbation solutions for elliptic equations*. Comm. Partial Differential Equations 32 (2007), no. 4-6, 557-578.
- [42] Scheven, C. and Schmidt, T. *Asymptotically regular problems I: Higher integrability*. J. Differential Equations 248 (4) (2010), 745-791.
- [43] Scheven, C. and Schmidt, T. *Asymptotically regular problems II: Partial Lipschitz continuity and a singular set of positive measure*. Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) 8 (3) (2009), 469-507.
- [44] Silvestre, L. and Teixeira, E.V. *Regularity estimates for fully non linear elliptic equations which are asymptotically convex*. Contributions to nonlinear elliptic equations and systems, 425-438, Progr. Nonlinear Differential Equations Appl., 86, Birkhäuser/Springer, Cham, 2015.
- [45] Trudinger, N.S. *Fully nonlinear, uniformly elliptic equations under natural structure conditions*. Trans. Amer. Math. Soc. 278 (1983), no. 2, 751-769.
- [46] Trudinger, N.S. *Regularity of solutions of fully nonlinear elliptic equations*. Boll. Un. Mat. Ital. A (6) 3 (1984), no. 3, 421-430.
- [47] Winter, N. *$W^{2,p}$ and $W^{1,p}$ -Estimates at the Boundary for Solutions of Fully Nonlinear, Uniformly Elliptic Equations*. Z. Anal. Adwend. (J. Anal. Appl.) 28 (2009), 129-164.
- [48] Zhang, J., Zheng, S. and Zuo, C. *$W^{2,p}$ -regularity for asymptotically regular fully nonlinear elliptic and parabolic equations with oblique boundary values*. Discrete Contin. Dyn. Syst. Ser. S 14 (2021), no. 9, 3305-3318.