

# A Fisher matrix for gravitational-wave population inference

Jonathan R. Gair<sup>1\*</sup>, Andrea Antonelli<sup>2†</sup>, Riccardo Barbieri<sup>1‡</sup>

<sup>1</sup>Max Planck Institute for Gravitational Physics (Albert Einstein Institute), Am Mühlenberg 1, Potsdam-Golm, 14476, Germany

<sup>2</sup>Department of Physics and Astronomy, Johns Hopkins University, 3400 N. Charles Street, Baltimore, Maryland, 21218, USA

18 May 2022

## ABSTRACT

We derive a Fisher matrix for the parameters characterising a population of gravitational-wave events. This provides a guide to the precision with which population parameters can be estimated with multiple observations, which becomes increasingly accurate as the number of events and the signal-to-noise ratio of the sampled events increases. The formalism takes into account individual event measurement uncertainties and selection effects, and can be applied to arbitrary population models. We illustrate the framework with two examples: an analytical calculation of the Fisher matrix for the mean and variance of a Gaussian model describing a population affected by selection effects, and an estimation of the precision with which the slope of a power law distribution of supermassive black-hole masses can be measured using extreme-mass-ratio inspiral observations. We compare the Fisher predictions to results from Monte Carlo analyses, finding very good agreement.

**Key words:** gravitational waves.

## 1 INTRODUCTION

Population analyses aim at inferring the parameters that describe the distribution of the properties of a set of observed events drawn from a common population. In the context of gravitational-wave (GW) astrophysics, such analyses have been carried out for the 90 coalescing compact-object binaries that have so far been observed by ground-based gravitational wave detectors, and reported in the third gravitational wave transient catalogue, GWTC-3 (Abbott et al. 2021b,c,d). These population analyses aimed at understanding the astrophysical processes that lead to the formation of the binaries (Abbott et al. 2019), or at using them to constrain cosmic expansion history by estimating parameters including the Hubble constant (Abbott et al. 2021a).

Given a set of observed events, the usual approach to estimate distribution parameters is to complete a Bayesian hierarchical analysis using techniques such as Markov Chain Monte Carlo (MCMC). While these are the most reliable way to obtain posterior samples from actual data, they are typically computationally expensive and so it can become impractical to use these approaches to make forecasts for future observations that include surveys over param-

eter space. However, such surveys are crucial for scoping out the science cases of future detectors, such as the Einstein Telescope (Punturo et al. 2010), Cosmic Explorer (Reitze et al. 2019) and the spaceborne LISA mission (Amaro-Seoane et al. 2017), all of which are expected to detect thousands of sources from multiple populations. For explorations of this nature, one can trade off accuracy in the estimates of parameter-measurement precision for computational speed by using approximations that are valid in the limit of high signal-to-noise ratio (SNR). In the context of *source* parameters for *individual* signals, the Fisher matrix is commonly used to cheaply assess the measurement precision of a parameter (Vallisneri 2008). Within the linear-signal approximation, valid for high SNR sources, the inverse of the Fisher matrix is an approximation to the covariance matrix and therefore the width of the likelihood function. Under the assumption of flat priors, it also approximates the shape of the Bayesian posterior probability distribution we would expect to obtain in an MCMC analyses. For a parameter set  $\vec{\lambda}$ , the Fisher matrix can be written in general terms as the expectation value over the data generating process of derivatives of the log-likelihood  $p(\mathbf{d}|\vec{\lambda})$ ,

$$(\Gamma_{\lambda})_{ij} = \mathbb{E} \left[ -\frac{\partial^2 \ln p(\mathbf{d}|\vec{\lambda})}{\partial \lambda^i \partial \lambda^j} \right]. \quad (1)$$

While this provides a guide to measurement uncertainties for individual events, the Fisher matrix does not directly

\* jonathan.gair@aei.mpg.de  
 † aantone3@jh.edu  
 ‡ riccardo.barbieri@aei.mpg.de

provide an indication of how well the properties of the population can be inferred when those events are subsequently combined in a hierarchical model. In this paper we address this shortcoming by deriving a Fisher Matrix for the population parameters assuming Gaussian noise and using the likelihood for population inference in the presence of selection effects from (Mandel, Farr & Gair 2019). The expression we obtain is valid for high SNRs and small biases in the individual events' parameters. We illustrate our formalism with two examples. First, we consider a ‘‘Gaussian-Gaussian’’ case, in which both noise and the data generation process are normally distributed, and check our expressions against the direct calculation of the Fisher matrix as an expectation value over data realizations. We also perform MCMC analyses with and without selection effects, as a cross-check to verify our results. Secondly, we consider the more astrophysically relevant case of a power-law distributed population (while still assuming Gaussian noise) and again validate our results against MCMC analyses with and without selection effects. We generally find an excellent agreement between the Fisher and MCMC estimates, while confirming results in (Gair, Tang & Volonteri 2010) for the latter scenario.

The paper is organized as follows. In Section 2 we describe our population Fisher Matrix formalism, highlighting the main assumptions and steps to obtain the result. A derivation of corrections to this formula and their scalings is found in Appendix 2. In Section 3, we consider the Gaussian population model, checking our formula against a direct calculation of the Fisher Matrix and an MCMC analysis. In section 4, we consider the case of inference of a power-law massive black hole mass distribution using extreme-mass-ratio inspiral (EMRI) observations, once again comparing the result against MCMC. Finally, in section 5 we discuss our results and prospects for future work. The framework we develop here could be applied in a wide variety of contexts. The focus on gravitational wave detectors and the choice of the examples provided here are driven purely by the authors' areas of expertise. The results can be fully reproduced with codes made publicly available at <https://github.com/aantonelli94/PopFisher>.

## 2 THE FISHER MATRIX FOR POPULATION DISTRIBUTIONS

The standard model used to represent the data stream,  $\mathbf{d}$ , of a gravitational wave detector is as a linear combination of a signal,  $\mathbf{h}(\vec{\theta})$ , dependent on some parameters  $\vec{\theta}$ , and noise,  $\mathbf{n}$ , that is usually assumed to be a realisation of a stationary and Gaussian stochastic process described by a power spectral density  $S_h(f)$ ,

$$\mathbf{d} = \mathbf{h}(\vec{\theta}) + \mathbf{n}, \quad \langle \tilde{n}^*(f) \tilde{n}(f') \rangle = S_h(f) \delta(f - f'). \quad (2)$$

In this model the likelihood is

$$p(\mathbf{d}|\vec{\theta}) \propto \exp \left[ -\frac{1}{2} \left( \mathbf{d} - \mathbf{h}(\vec{\theta}) | \mathbf{d} - \mathbf{h}(\vec{\theta}) \right) \right],$$

$$\text{where } \langle \mathbf{a} | \mathbf{b} \rangle = 4 \text{Re} \int_0^\infty \frac{\tilde{a}^*(f) \tilde{b}(f)}{S_h(f)} df. \quad (3)$$

To understand the precision with which gravitational wave observations can determine the parameters of a source, it is

common to compute the Fisher information matrix, defined by

$$(\Gamma_\theta)_{ij} = \mathbb{E} \left[ \frac{\partial \ln p(\mathbf{d}|\vec{\theta})}{\partial \theta^i} \frac{\partial \ln p(\mathbf{d}|\vec{\theta})}{\partial \theta^j} \right], \quad (4)$$

where the expectation value is taken over realizations of the data drawn from the data generating process,  $\mathbf{d}$ . For the gravitational wave detector likelihood in Eq. (3), the Fisher information matrix can be seen to reduce to

$$(\Gamma_\theta)_{ij} = \left( \frac{\partial \mathbf{h}}{\partial \theta^i} \middle| \frac{\partial \mathbf{h}}{\partial \theta^j} \right), \quad (5)$$

where we are using the inner product introduced in Eq. (3). The Fisher matrix provides a leading order approximation to the shape of the likelihood and hence also the Bayesian posterior when using priors that are approximately flat over the support of the likelihood. It becomes an increasingly good guide to the precision of parameter estimation as the SNR with which the source is observed increases.

In population inference, we are no longer primarily interested in the parameters of the individual events, but in the parameters that characterise the population from which the individual events are drawn. We assume that we have some population model,  $p(\vec{\theta}|\vec{\lambda})$ , that describes the probability distribution of the parameters,  $\vec{\theta}$ , of individual events drawn randomly from a population characterised by parameters,  $\vec{\lambda}$ . We want to infer the parameters of the population by combining the information from many observed events. For a given choice of population parameters, the distribution of observed datasets is characterised by

$$p(\mathbf{d}|\vec{\lambda}) = \frac{p_{\text{full}}(\mathbf{d}|\vec{\lambda})}{p_{\text{det}}(\vec{\lambda})} \quad (6)$$

$$\text{where } p_{\text{full}}(\mathbf{d}|\vec{\lambda}) = \int p(\mathbf{d}|\vec{\theta}) p(\vec{\theta}|\vec{\lambda}) d\vec{\theta}$$

$$p_{\text{det}}(\vec{\lambda}) = \int p_{\text{det}}(\vec{\theta}) p(\vec{\theta}|\vec{\lambda}) d\vec{\theta}$$

$$p_{\text{det}}(\vec{\theta}) = \int_{\mathbf{d} > \text{thresh}} p(\mathbf{d}|\vec{\theta}) d\mathbf{d}. \quad (7)$$

This expression accounts for the fact that not all events that occur in the Universe are detected. Detection is a property of the observed data,  $\mathbf{d}$ , and the last integral is over all data sets that would pass the threshold to be counted as a detected event and hence included in the population inference. The normalisation term,  $p_{\text{det}}(\vec{\lambda})$ , depends only on the population parameters and represents the fraction of events in the Universe that are detectable. We refer the reader to (Mandel, Farr & Gair 2019) for further details. This form of the likelihood assumes that the number of events observed in a fixed time period does not convey any information about the population parameters. However, the precision with which the population parameters are estimated asymptotically is independent of that assumption. This is discussed in more detail in Appendix 1.

Equation (1) is the equivalent of Eq. (4) for this population likelihood, and so it should give a guide to the precision with which the population parameters can be measured. Note that the two forms of the expression are slightly different, but it is straightforward to show that the two results are equivalent by integrating by parts and using conservation of

probability. This will be a good guide for a “high signal-to-noise ratio”, which for populations means a large number of observed events. The fact that Eq. (1) is a good approximation to the precision of population inference can be seen as follows. In a general population inference problem, we have observed a set of events, indexed by  $i$ , with corresponding datasets  $\{\mathbf{d}_i\}$ . The posterior distribution on the population parameters from this set of events can be found from Bayes’ theorem and takes the form

$$p(\vec{\lambda}|\{\mathbf{d}_i\}) \propto \pi(\vec{\lambda}) \prod_{i=1}^n p(\mathbf{d}_i|\vec{\lambda}) \quad (8)$$

where  $n$  is the total number of events observed,  $\pi(\vec{\lambda})$  is the prior on the population parameters and  $p(\mathbf{d}_i|\vec{\lambda})$  is the likelihood of the population parameters  $\vec{\lambda}$  for dataset  $\mathbf{d}_i$ . The log-posterior is

$$\ln p(\vec{\lambda}|\{\mathbf{d}_i\}) \propto \ln \pi(\vec{\lambda}) + \sum_{i=1}^n \ln p(\mathbf{d}_i|\vec{\lambda}). \quad (9)$$

The latter quantity is a sum of independent random variables (assuming that all observations are independent). In the limit that  $n \rightarrow \infty$  we can use the central limit theorem to deduce

$$\frac{1}{n} \sum_{i=1}^n \ln p(\mathbf{d}_i|\vec{\lambda}) \sim N\left(\mu(\vec{\lambda}|\vec{\lambda}_t), \frac{\sigma^2(\vec{\lambda}|\vec{\lambda}_t)}{n}\right) \quad (10)$$

where

$$\mu(\vec{\lambda}|\vec{\lambda}_t) = \mathbb{E} \left[ \ln p(\mathbf{d}|\vec{\lambda}) \right], \quad \sigma^2(\vec{\lambda}|\vec{\lambda}_t) = \mathbb{E} \left[ (\ln p(\mathbf{d}|\vec{\lambda}) - \mu)^2 \right] \quad (11)$$

and the expectation value is taken over the data generating process, which we assume to be consistent with the likelihood we are using, evaluated for the true values of the population parameters  $\vec{\lambda}_t$ . Since

$$\begin{aligned} \frac{\partial \mu}{\partial \lambda^i} &= \int p(\mathbf{d}|\vec{\lambda}_t) \frac{1}{p(\mathbf{d}|\vec{\lambda})} \frac{\partial p(\mathbf{d}|\vec{\lambda})}{\partial \lambda^i} d\mathbf{d} \\ \Rightarrow \frac{\partial \mu}{\partial \lambda^i} \Big|_{\vec{\lambda}=\vec{\lambda}_t} &= \int \frac{\partial p(\mathbf{d}|\vec{\lambda})}{\partial \lambda^i} d\mathbf{d} = \frac{\partial}{\partial \lambda^i} \int p(\mathbf{d}|\vec{\lambda}) d\mathbf{d} \\ &= \frac{\partial}{\partial \lambda^i} (1) = 0, \end{aligned} \quad (12)$$

we deduce that the population likelihood is peaked at the true parameters asymptotically (this is not true on average for a finite number of observations, as discussed in Appendix 2). Note that this happens by virtue of the assumed consistency between the likelihood and the data-generating process, and would not be the case if the likelihood was only an approximation to the true population. As  $n \rightarrow \infty$  the log-posterior converges to the function  $n\mu(\vec{\lambda})$ , and so the posterior becomes increasingly concentrated around  $\vec{\lambda}_t$ . Expanding the function  $\mu(\vec{\lambda}|\vec{\lambda}_t)$  near  $\vec{\lambda}_t$  we have

$$\begin{aligned} \mu(\vec{\lambda}|\vec{\lambda}_t) &= \mu(\vec{\lambda}_t|\vec{\lambda}_t) \\ &+ \frac{1}{2} \left( \frac{d^2 \mu}{d\lambda^i d\lambda^j} \right)_{\vec{\lambda}_t} (\lambda^i - \lambda_t^i)(\lambda^j - \lambda_t^j) + \dots \end{aligned} \quad (13)$$

We deduce that the asymptotic covariance matrix is  $\Gamma_\lambda^{-1}/n$ , where

$$(\Gamma_\lambda)_{ij} = - \left( \frac{d^2 \mu}{d\lambda^i d\lambda^j} \right)_{\vec{\lambda}_t} = \mathbb{E} \left[ - \frac{\partial^2 \ln p(\mathbf{d}|\vec{\lambda})}{\partial \lambda^i \partial \lambda^j} \right]. \quad (14)$$

In the last equality, we use Eq. (12). This justifies the use of Eq. (1) to characterise the precision of parameter estimation in the limit  $n \rightarrow \infty$ . It becomes increasingly reliable as  $n \rightarrow \infty$ , as corrections to this formula scale like  $n^{-\frac{1}{2}}$  relative to leading order. This is justified in more detail in Appendix 2.

This result can be evaluated at various levels of approximation. The full asymptotic posterior is described by the function  $\mu(\vec{\lambda}|\vec{\lambda}_t)$ , which can be evaluated through Monte Carlo integration. This is computationally expensive as it requires evaluation over different choices of  $\vec{\lambda}$  and  $\vec{\lambda}_t$ . The next level of approximation is to evaluate Eq. (1) directly. This makes a linear signal approximation in the population parameters, but no approximation to the evaluation of  $p(\mathbf{d}|\vec{\lambda})$ . This is less complex because evaluation is only needed in the vicinity of  $\vec{\lambda}_t$ . A final level of approximation is to simplify  $p(\mathbf{d}|\vec{\lambda})$  by using the linear signal approximation for the individual event parameters as well. This is the approach we will now describe.

We consider a single observation of a source with parameters  $\vec{\theta}_0$ , and data  $\mathbf{d} = \mathbf{h}(\vec{\theta}_0) + \mathbf{n}$ . Taking the expectation value over the true data distribution then reduces to taking the expectation value over the distribution of the noise  $\mathbf{n}$  and the distribution of the parameters  $\vec{\theta}_0$ , which is  $p(\vec{\theta}_0|\vec{\lambda}_t)$ . Under the linear signal approximation we expand

$$\mathbf{h}(\vec{\theta}) = \mathbf{h}(\vec{\theta}_0) + \frac{\partial \mathbf{h}}{\partial \theta^i} \Delta \theta^i \quad (15)$$

where  $\Delta \theta^i = \theta^i - \theta_0^i$ . The gravitational wave likelihood can then be written

$$\begin{aligned} \tilde{p}(\{\mathbf{d}\}|\vec{\theta}) &\propto \exp \left[ -\frac{1}{2} (\mathbf{d} - \mathbf{h}(\vec{\theta})) | \mathbf{d} - \mathbf{h}(\vec{\theta}) \right] \\ &\approx \exp \left[ -\frac{1}{2} (\mathbf{n}|\mathbf{n}) + N_i \Delta \theta^i - \frac{1}{2} (\Gamma_\theta)_{ij} \Delta \theta^i \Delta \theta^j \right] \end{aligned} \quad (16)$$

where  $N_i = \left( \frac{\partial \mathbf{h}}{\partial \theta^i} | \mathbf{n} \right)$

and  $(\Gamma_\theta)_{ij}$  is the single source Fisher matrix defined in Eq. (5). We similarly expand the source prior term

$$\begin{aligned} \ln p(\vec{\theta}|\vec{\lambda}) &= \ln p(\vec{\theta}_0|\vec{\lambda}) + P_i \Delta \theta^i - \frac{1}{2} H_{ij} \Delta \theta^i \Delta \theta^j + \dots \\ \text{where } P_i &= \frac{\partial \ln p(\vec{\theta}|\vec{\lambda})}{\partial \theta^i}, \quad H_{ij} = - \frac{\partial^2 \ln p(\vec{\theta}|\vec{\lambda})}{\partial \theta^i \partial \theta^j}, \end{aligned} \quad (17)$$

in which the derivatives are evaluated at the parameter space point  $\theta_0$ . Substituting the preceding two expressions into Eq. 7 and integrating over  $\vec{\theta}$ , which is equivalent to integrating over  $\Delta \vec{\theta}$  in the linear signal approximation, we obtain

$$\begin{aligned} \tilde{p}(\{\mathbf{d}\}|\vec{\lambda}) &\approx \frac{\exp[-(\mathbf{n}|\mathbf{n})/2]}{p_{\det}(\vec{\lambda})} \int d\Delta \vec{\theta} \left[ p(\vec{\theta}_0|\vec{\lambda}) \right. \\ &\left. \exp \left\{ -\frac{1}{2} (\Gamma_{ij} + H_{ij})(\Delta \theta^i - \Delta \theta_{\text{bf}}^i)(\Delta \theta^j - \Delta \theta_{\text{bf}}^j) \right. \right. \\ &\left. \left. + \frac{1}{2} (N_i + P_i)(\Gamma + H)_{ij}^{-1} (N_j + P_j) \right\} \right] \\ &= (2\pi)^{N/2} \frac{p(\vec{\theta}_0|\vec{\lambda}) \exp[-(\mathbf{n}|\mathbf{n})/2]}{p_{\det}(\vec{\lambda}) \sqrt{\det(\Gamma + H)}} \\ &\times \exp \left[ \frac{1}{2} (N_i + P_i)(\Gamma + H)_{ij}^{-1} (N_j + P_j) \right], \end{aligned} \quad (18)$$

where we have written

$$\Delta \theta_{\text{bf}}^i = (\Gamma + H)_{ik}^{-1} (N_k + P_k).$$

This is the point at which the likelihood is maximized and hence is the “best-fit” point in parameter space. We can now evaluate the population Fisher matrix using the expression

$$\begin{aligned} -(\Gamma_\lambda)_{ij} &= \int \left( \frac{\partial^2 \ln p(\mathbf{d}|\vec{\lambda})}{\partial \lambda^i \partial \lambda^j} \right)_{\vec{\lambda}_t} p(\mathbf{d}|\vec{\lambda}_t) d\mathbf{d} \\ &= \iint \left( \frac{\partial^2 \ln p(\mathbf{d}|\vec{\lambda})}{\partial \lambda^i \partial \lambda^j} \right)_{\vec{\lambda}_t} p(\vec{\theta}_0|\vec{\lambda}_t) p(\mathbf{n}|\vec{\theta}_0) d\mathbf{n} d\vec{\theta}_0. \end{aligned} \quad (19)$$

In the above the integral over the noise distribution is conditioned on  $\vec{\theta}_0$  because of selection effects. This integral is over all noise realisations that ensure  $\mathbf{d} = \mathbf{h}(\vec{\theta}_0) + \mathbf{n}$  is above the detection threshold. Substituting Eq. (18) into the above we obtain a sequence of terms. To simplify these we carry out the integral over the noise,  $\mathbf{n}$ . The only terms in Eq. (18) that depend on  $\mathbf{n}$  are  $N_i$  and the prefactor  $\exp[-(\mathbf{n}|\mathbf{n})/2]$ . The latter enters  $\ln \bar{p}$  additively and has no dependence on the population parameters, so it does not contribute to the final result. The former term also has no explicit dependence on the population parameters, but it appears multiplied by terms that do. There are thus three distinct types of term that appear in the argument of the integral - terms that have no explicit dependence on  $\mathbf{n}$ , terms that are linear in  $N_i$  and terms that are quadratic in  $N_i$ . We define these integrals as follows

$$\begin{aligned} p_{\text{det}}(\vec{\theta}_0) &= \int p(\mathbf{n}|\vec{\theta}_0) d\mathbf{n} \\ D_i &\equiv \int \left( \mathbf{n} \left| \frac{\partial \mathbf{h}}{\partial \theta^i} \right. \right) p(\mathbf{n}) d\mathbf{n} = \frac{\partial p_{\text{det}}(\vec{\theta}_0)}{\partial \theta^i}, \\ D_{ij} &\equiv \int \left( \mathbf{n} \left| \frac{\partial \mathbf{h}}{\partial \theta^i} \right. \right) \left( \mathbf{n} \left| \frac{\partial \mathbf{h}}{\partial \theta^j} \right. \right) p(\mathbf{n}) d\mathbf{n}. \end{aligned} \quad (20)$$

Using these expressions to carry out the integrals over  $\mathbf{n}$ , we obtain the final result

$$-(\Gamma_\lambda)_{ij} \equiv (\Gamma_{\text{I}})_{ij} + (\Gamma_{\text{II}})_{ij} + (\Gamma_{\text{III}})_{ij} + (\Gamma_{\text{IV}})_{ij} + (\Gamma_{\text{V}})_{ij}, \quad (21)$$

with

$$\begin{aligned} (\Gamma_{\text{I}})_{ij} &= \int \frac{\partial^2 \ln(p(\vec{\theta}_0|\vec{\lambda})/p_{\text{det}}(\vec{\lambda}))}{\partial \lambda^i \partial \lambda^j} \frac{p_{\text{det}}(\vec{\theta}_0)}{p_{\text{det}}(\vec{\lambda})} p(\vec{\theta}_0|\vec{\lambda}) d\vec{\theta}_0, \\ (\Gamma_{\text{II}})_{ij} &= -\frac{1}{2} \int \frac{\partial^2 \ln \det(\Gamma + H)}{\partial \lambda^i \partial \lambda^j} \frac{p_{\text{det}}(\vec{\theta}_0)}{p_{\text{det}}(\vec{\lambda})} p(\vec{\theta}_0|\vec{\lambda}) d\vec{\theta}_0, \\ (\Gamma_{\text{III}})_{ij} &= \frac{1}{2} \int \frac{\partial^2}{\partial \lambda^i \partial \lambda^j} [(\Gamma + H)^{-1}]_{kl} D_{kl} \frac{p(\vec{\theta}_0|\vec{\lambda})}{p_{\text{det}}(\vec{\lambda})} d\vec{\theta}_0, \\ (\Gamma_{\text{IV}})_{ij} &= \int \frac{\partial^2}{\partial \lambda^i \partial \lambda^j} [P_k(\Gamma + H)^{-1}]_{kl} D_l \frac{p(\vec{\theta}_0|\vec{\lambda})}{p_{\text{det}}(\vec{\lambda})} d\vec{\theta}_0, \\ (\Gamma_{\text{V}})_{ij} &= \frac{1}{2} \int \frac{\partial^2}{\partial \lambda^i \partial \lambda^j} [P_k(\Gamma + H)^{-1} P_l] \frac{p_{\text{det}}(\vec{\theta}_0)}{p_{\text{det}}(\vec{\lambda})} p(\vec{\theta}_0|\vec{\lambda}) d\vec{\theta}_0. \end{aligned}$$

This is an approximate expression for the population Fisher matrix which can be used to estimate the precision with which observations will be able to determine the population parameters. In the next sections we will illustrate the use of this result for a few sample problems.

### 3 ILLUSTRATION I: A GAUSSIAN-GAUSSIAN MODEL

The first application of Eq. (21) we will consider is to a “Gaussian-Gaussian” model in which both observations and noise are normally distributed. We simplify the setting by assuming a waveform dependent on a single parameter  $\theta$ . The distribution of the parameter is

$$p(\theta|\vec{\lambda}) = \mathcal{N}(\mu, \Sigma^2) = \frac{1}{\sqrt{2\pi\Sigma^2}} \exp\left[-\frac{(\theta - \mu)^2}{2\Sigma^2}\right], \quad (22)$$

with population parameters (henceforth, hyperparameters)  $\vec{\lambda} = \{\mu, \Sigma^2\}$ . Noise is also a Gaussian with zero mean and variance  $\sigma$ . Since the data stream is a sum of Gaussians, it is modelled by  $\mathcal{N}(\mu, \sigma^2 + \Sigma^2)$ , with mean and variance given by the sums of individual means and variances,

$$p(\mathbf{d}|\vec{\lambda}) = \frac{1}{\sqrt{2\pi(\sigma^2 + \Sigma^2)}} \exp\left[-\frac{(\mathbf{d} - \mu)^2}{2(\sigma^2 + \Sigma^2)}\right]. \quad (23)$$

Given the implicit simple choice for the signals, the Fisher matrix of source parameters reduces to

$$\Gamma_\theta = \left( \frac{\partial \mathbf{h}}{\partial \theta} \middle| \frac{\partial \mathbf{h}}{\partial \theta} \right) = \frac{1}{\sigma^2} \left| \frac{\partial \mathbf{h}}{\partial \theta} \right|^2 = \frac{1}{\sigma^2}, \quad (24)$$

while from Eq. (22) we have that  $P$  and  $H$  in (17) are

$$P = -\frac{(\theta - \mu)}{\Sigma^2} \quad \text{and} \quad H = \frac{1}{\Sigma^2}. \quad (25)$$

The example reported in this section does not have an immediate analogy in GW astrophysics, but it can be thought of as a more general application of the population Fisher matrix. The advantage of choosing such a simple setting is that the matrix entries can be directly integrated as expectation values over data realizations. In the presence of selection effects, the integrals to be solved are

$$\begin{aligned} (\Gamma_\lambda)_{ij} &= \mathbb{E} \left( -\frac{\partial^2}{\partial \lambda^i \partial \lambda^j} \left[ \ln \left( \frac{p(\mathbf{d}|\theta, \vec{\lambda})}{p_{\text{det}}(\vec{\lambda})} \right) \right] \right) \\ &= -\int_{\mathbf{d}_{\text{th}}}^{\infty} \frac{\partial^2}{\partial \lambda^i \partial \lambda^j} \left[ \ln \left( \frac{p(\mathbf{d}|\vec{\lambda})}{p_{\text{det}}(\vec{\lambda})} \right) \right] \frac{p(\mathbf{d}|\vec{\lambda})}{p_{\text{det}}(\vec{\lambda})} d\mathbf{d}. \end{aligned} \quad (26)$$

We only select realizations of the data  $\mathbf{d} > \mathbf{d}_{\text{th}}$  that are above a certain threshold. The predictions for the various components of the population Fisher matrix are<sup>1</sup>

$$\begin{aligned} (\Gamma_\lambda)_{\mu\mu} &= \frac{\partial^2 \ln p_{\text{det}}(\vec{\lambda})}{\partial \mu^2} + \frac{1}{(\sigma^2 + \Sigma^2)}, \\ (\Gamma_\lambda)_{\mu\Sigma^2} &= \frac{\partial^2 \ln p_{\text{det}}(\vec{\lambda})}{\partial \mu \partial \Sigma^2} + \frac{1}{(\sigma^2 + \Sigma^2)} \frac{p(\mathbf{d}_{\text{th}}|\vec{\lambda})}{p_{\text{det}}(\vec{\lambda})}, \\ (\Gamma_\lambda)_{\Sigma^2\Sigma^2} &= \frac{\partial^2 \ln p_{\text{det}}(\vec{\lambda})}{(\partial \Sigma^2)^2} + \frac{1}{2(\sigma^2 + \Sigma^2)^2} \\ &\quad + \frac{(\mathbf{d}_{\text{th}} - \mu)}{(\sigma^2 + \Sigma^2)^2} \frac{p(\mathbf{d}_{\text{th}}|\vec{\lambda})}{p_{\text{det}}(\vec{\lambda})}, \end{aligned} \quad (27)$$

<sup>1</sup> For the last integral, it is useful to notice that

$$\int_{\mathbf{d}_{\text{th}}}^{\infty} (\mathbf{d} - \mu)^2 p(\mathbf{d}|\vec{\lambda}) d\mathbf{d} = (\sigma^2 + \Sigma^2) \left[ 1 + (\mathbf{d}_{\text{th}} - \mu) \frac{p(\mathbf{d}_{\text{th}}|\vec{\lambda})}{p_{\text{det}}(\vec{\lambda})} \right],$$

which follows from  $(\mathbf{d} - \mu)p(\mathbf{d}|\vec{\lambda}) = -(\sigma^2 + \Sigma^2)\partial p(\mathbf{d}|\vec{\lambda})/\partial \mathbf{d}$ , integrating by parts, and using Eq.(28).

where we have used Eqs.(23), (26), the fact that the integral is normalized through

$$p_{\text{det}}(\vec{\lambda}) = \int_{\mathbf{d}_{\text{th}}}^{\infty} p(\mathbf{d}|\vec{\lambda})d\mathbf{d} = \frac{1}{2} \operatorname{erfc} \left( \frac{(\mathbf{d}_{\text{th}} - \mu)}{\sqrt{2(\sigma^2 + \Sigma^2)}} \right), \quad (28)$$

and the definition

$$p(\mathbf{d}_{\text{th}}|\vec{\lambda}) \equiv \frac{1}{\sqrt{2\pi(\sigma^2 + \Sigma^2)}} \exp \left[ -\frac{(\mathbf{d}_{\text{th}} - \mu)^2}{2(\sigma^2 + \Sigma^2)} \right]. \quad (29)$$

Finally, from Eq. (28) it follows that

$$\begin{aligned} \frac{\partial^2 \ln p_{\text{det}}(\lambda)}{(\partial \mu)^2} &= \left( \frac{\mathbf{d}_{\text{th}} - \mu}{\sigma^2 + \Sigma^2} \right) \frac{p(\mathbf{d}_{\text{th}}|\lambda)}{p_{\text{det}}(\lambda)} - \frac{p(\mathbf{d}_{\text{th}}|\lambda)^2}{p_{\text{det}}(\lambda)^2}, \\ \frac{\partial^2 \ln p_{\text{det}}(\lambda)}{\partial \mu \partial \Sigma^2} &= \frac{p(\mathbf{d}_{\text{th}}|\lambda)}{p_{\text{det}}(\lambda)} \left[ \frac{(\mathbf{d}_{\text{th}} - \mu)^2}{2(\sigma^2 + \Sigma^2)^2} - \frac{1}{2(\sigma^2 + \Sigma^2)} \right] \\ &\quad - \frac{p(\mathbf{d}_{\text{th}}|\lambda)^2}{2p_{\text{det}}(\lambda)^2} \left( \frac{\mathbf{d}_{\text{th}} - \mu}{\sigma^2 + \Sigma^2} \right), \\ \frac{\partial^2 \ln p_{\text{det}}(\lambda)}{(\partial \Sigma^2)^2} &= \frac{p(\mathbf{d}_{\text{th}}|\lambda)}{p_{\text{det}}(\lambda)} \left[ \frac{(\mathbf{d}_{\text{th}} - \mu)^3}{4(\sigma^2 + \Sigma^2)^3} - \frac{3(\mathbf{d}_{\text{th}} - \mu)}{4(\sigma^2 + \Sigma^2)^2} \right] \\ &\quad - \frac{p(\mathbf{d}_{\text{th}}|\lambda)^2}{4p_{\text{det}}(\lambda)^2} \left( \frac{\mathbf{d}_{\text{th}} - \mu}{\sigma^2 + \Sigma^2} \right)^2. \end{aligned} \quad (30)$$

### 3.1 Solution from the population Fisher matrix

While in this simple setting the direct evaluation of the Fisher Matrix as expectation value is much simpler, we wish to now evaluate it using Eq. (21) as an important sanity check of that general formula. For  $(\Gamma_{\lambda})_{\mu\mu}$ , we notice that the  $(\Gamma_{\text{II}})_{\mu\mu}$ ,  $(\Gamma_{\text{III}})_{\mu\mu}$  and  $(\Gamma_{\text{IV}})_{\mu\mu}$  vanish. The second and third terms vanish because  $\Gamma + H$  does not depend on  $\mu$ , while the fourth term vanishes because  $P$  is only linear in  $\mu$  and two derivatives with respect to it are needed for  $(\Gamma_{\lambda})_{\mu\mu}$ . The only terms contributing are therefore  $(\Gamma_{\text{I}})_{\mu\mu}$  and  $(\Gamma_{\text{V}})_{\mu\mu}$ , which leads to

$$\begin{aligned} (\Gamma_{\lambda})_{\mu\mu} &= - \int \frac{\partial^2 \ln(p(\theta|\vec{\lambda})/p_{\text{det}}(\vec{\lambda}))}{\partial \mu^2} \frac{p_{\text{det}}(\theta)}{p_{\text{det}}(\vec{\lambda})} p(\theta|\vec{\lambda})d\theta \\ &\quad - \frac{1}{2} \int \frac{\partial^2}{\partial \mu^2} [P^2(\Gamma + H)^{-1}] \frac{p_{\text{det}}(\theta)}{p_{\text{det}}(\vec{\lambda})} p(\theta|\vec{\lambda})d\theta \\ &= \frac{\partial^2 \ln p_{\text{det}}(\vec{\lambda})}{\partial \mu^2} + \frac{1}{(\sigma^2 + \Sigma^2)}, \end{aligned} \quad (31)$$

where we have used  $P$ ,  $\Gamma$  and  $H$  given in Eqs. (22), (24) and (25), as well as the normalisation

$$\int \frac{p_{\text{det}}(\theta)}{p_{\text{det}}(\vec{\lambda})} p(\theta|\vec{\lambda})d\theta = 1. \quad (32)$$

The result matches the prediction of Eq.(27) as expected.

In the case of  $(\Gamma_{\lambda})_{\mu\Sigma^2}$ , the second and third terms,  $(\Gamma_{\text{II}})_{\mu\Sigma^2}$  and  $(\Gamma_{\text{III}})_{\mu\Sigma^2}$ , vanish for the same reason as above, but now the fourth term  $(\Gamma_{\text{IV}})_{\mu\Sigma^2}$  does contribute. Overall

we have that

$$\begin{aligned} (\Gamma_{\lambda})_{\mu\Sigma^2} &= - \int \frac{\partial^2 \ln(p(\theta|\vec{\lambda})/p_{\text{det}}(\vec{\lambda}))}{\partial \mu \partial \Sigma^2} \frac{p_{\text{det}}(\theta)}{p_{\text{det}}(\vec{\lambda})} p(\theta|\vec{\lambda})d\theta \\ &\quad - \int \frac{\partial^2}{\partial \mu \partial \Sigma^2} [P(\Gamma + H)^{-1}] D_i \frac{p_{\text{det}}(\theta)}{p_{\text{det}}(\vec{\lambda})} p(\theta|\vec{\lambda})d\theta \\ &\quad - \frac{1}{2} \int \frac{\partial^2}{\partial \mu \partial \Sigma^2} [P^2(\Gamma + H)^{-1}] \frac{p_{\text{det}}(\theta)}{p_{\text{det}}(\vec{\lambda})} p(\theta|\vec{\lambda})d\theta \\ &= \frac{\partial^2 \ln p_{\text{det}}(\vec{\lambda})}{\partial \mu \partial \Sigma^2} + \int \frac{(\theta - \mu)}{\Sigma^4} \frac{p_{\text{det}}(\theta)}{p_{\text{det}}(\vec{\lambda})} p(\theta|\vec{\lambda})d\theta \\ &\quad + \int \left[ \frac{\sigma^2}{(\sigma^2 + \Sigma^2)^2} \right] \frac{\partial p_{\text{det}}(\theta)}{\partial \theta} \frac{p(\theta|\vec{\lambda})}{p_{\text{det}}(\vec{\lambda})} d\theta \\ &\quad - \frac{\sigma^2(\sigma^2 + 2\Sigma^2)}{\Sigma^4(\sigma^2 + \Sigma^2)^2} \frac{1}{p_{\text{det}}(\vec{\lambda})} \int (\theta - \mu) p_{\text{det}}(\theta) p(\theta|\vec{\lambda})d\theta, \end{aligned}$$

where we have used the definition of  $D_i$  in Eq. (20) and (32). We also notice that, from the definition

$$\begin{aligned} p_{\text{det}}(\theta) &= \int_{\mathbf{d}_{\text{th}}}^{\infty} p(\mathbf{d}|\theta)d\mathbf{d} \\ &= \int_{\mathbf{d}_{\text{th}}}^{\infty} \frac{1}{\sqrt{2\pi\sigma^2}} \exp \left[ -\frac{(\mathbf{d} - \theta)^2}{2\sigma^2} \right] d\mathbf{d} = \frac{1}{2} \operatorname{erfc} \left( \frac{\mathbf{d}_{\text{th}} - \theta}{\sqrt{2\sigma^2}} \right), \end{aligned} \quad (33)$$

it follows that

$$\begin{aligned} \frac{\partial p_{\text{det}}(\theta)}{\partial \theta} &= \int_{\mathbf{d}_{\text{th}}}^{\infty} \frac{(\mathbf{d} - \theta)}{\sigma^2} p(\mathbf{d}|\theta)d\mathbf{d} \\ &= \frac{1}{\sqrt{2\pi\sigma^2}} \exp \left[ -\frac{(\mathbf{d}_{\text{th}} - \theta)^2}{2\sigma^2} \right] \equiv p(\mathbf{d}_{\text{th}}|\theta), \end{aligned} \quad (34)$$

which can be rearranged to give

$$\begin{aligned} \int (\theta - \mu) \frac{\partial p_{\text{det}}(\theta)}{\partial \theta} p(\theta|\vec{\lambda})d\theta \\ &= \Sigma^2 \frac{\partial}{\partial \mu} \int p(\mathbf{d}_{\text{th}}|\theta) p(\theta|\vec{\lambda})d\theta \\ &= \Sigma^2 \frac{\partial p(\mathbf{d}_{\text{th}}|\vec{\lambda})}{\partial \mu} = \frac{\Sigma^2}{(\sigma^2 + \Sigma^2)} (\mathbf{d}_{\text{th}} - \mu) p(\mathbf{d}_{\text{th}}|\vec{\lambda}). \end{aligned} \quad (35)$$

From Eq. (28) it also follows that  $\partial p_{\text{det}}(\vec{\lambda})/\partial \mu = p(\mathbf{d}_{\text{th}}|\vec{\lambda})$ . Using this constraint, and the fact that

$$\begin{aligned} \frac{\partial p(\mathbf{d}_{\text{th}}|\vec{\lambda})}{\partial \mu} &= \int \frac{\partial}{\partial \mu} [p_{\text{det}}(\theta) p(\theta|\vec{\lambda})] d\theta \\ &= \frac{1}{\Sigma^2} \int (\theta - \mu) p_{\text{det}}(\theta) p(\theta|\vec{\lambda}) d\theta, \end{aligned} \quad (36)$$

we immediately get

$$\int (\theta - \mu) p_{\text{det}}(\theta) p(\theta|\vec{\lambda}) d\theta = \Sigma^2 p(\mathbf{d}_{\text{th}}|\vec{\lambda}), \quad (37)$$

and, together with (32) and (35), that

$$(\Gamma_{\lambda})_{\mu\Sigma^2} = \frac{\partial^2 \ln p_{\text{det}}(\vec{\lambda})}{\partial \mu \partial \Sigma^2} + \frac{1}{(\sigma^2 + \Sigma^2)} \frac{p(\mathbf{d}_{\text{th}}|\vec{\lambda})}{p_{\text{det}}(\vec{\lambda})}. \quad (38)$$

This agrees with Eq. (27) as expected.

Finally, we consider the case of  $(\Gamma_{\lambda})_{\Sigma^2\Sigma^2}$ , in which no term in (21) vanishes. The first term can be rearranged to

give<sup>2</sup>

$$\begin{aligned}
 (\Gamma_I)_{\Sigma^2\Sigma^2} &= \frac{\partial^2 \ln p_{\text{det}}(\vec{\lambda})}{(\partial\Sigma^2)^2} - \frac{1}{2\Sigma^4} \\
 &+ \frac{1}{\Sigma^6} \int (\theta - \mu)^2 \frac{p_{\text{det}}(\theta)}{p_{\text{det}}(\vec{\lambda})} p(\theta|\vec{\lambda}) d\theta \\
 &= \frac{\partial^2 \ln p_{\text{det}}(\vec{\lambda})}{(\partial\Sigma^2)^2} + \frac{1}{2\Sigma^4} + \frac{(\mathbf{d}_{\text{th}} - \mu)}{\Sigma^2(\sigma^2 + \Sigma^2)} \frac{p(\mathbf{d}_{\text{th}}|\vec{\lambda})}{p_{\text{det}}(\vec{\lambda})}.
 \end{aligned} \quad (40)$$

Using Eq. (32), the second term  $(\Gamma_{II})_{\Sigma^2\Sigma^2}$  is easily found to be

$$(\Gamma_{II})_{\Sigma^2\Sigma^2} = \frac{\sigma^2}{2\Sigma^4} \frac{(\sigma^2 + 2\Sigma^2)}{(\sigma^2 + \Sigma^2)^2}. \quad (41)$$

The third term  $(\Gamma_{III})_{\Sigma^2\Sigma^2}$  contains  $D_{ij}$  from Eq.(20), which can be rearranged to give

$$\begin{aligned}
 D_{\theta\theta} &= \frac{1}{\sqrt{2\pi}\sigma^3} \int_{\mathbf{d}_{\text{th}}}^{\infty} (\mathbf{d} - \theta)^2 \exp\left[-\frac{(\mathbf{d} - \theta)^2}{2\sigma^2}\right] d\mathbf{d} \\
 &= \frac{(\mathbf{d}_{\text{th}} - \theta)}{\sqrt{2\pi}\sigma^3} \exp\left[-\frac{(\mathbf{d}_{\text{th}} - \theta)^2}{2\sigma^2}\right] + \frac{1}{2\sigma^2} \text{erfc}\left(\frac{\mathbf{d}_{\text{th}} - \theta}{\sqrt{2}\sigma}\right).
 \end{aligned}$$

The complementary error function can be replaced by  $p_{\text{det}}(\theta)$  using (33). With this in mind, we have

$$(\Gamma_{III})_{\Sigma^2\Sigma^2} = \frac{\sigma^4}{(\sigma^2 + \Sigma^2)^4} (\mathbf{d}_{\text{th}} - \mu) \frac{p(\mathbf{d}_{\text{th}}|\vec{\lambda})}{p_{\text{det}}(\vec{\lambda})} + \frac{\sigma^2}{(\sigma^2 + \Sigma^2)^3}. \quad (42)$$

The results follows from rearranging Eqs. (34) and (29) into

$$\begin{aligned}
 &\int (\mathbf{d}_{\text{th}} - \theta) \exp\left[-\frac{(\mathbf{d}_{\text{th}} - \mu)^2}{2\sigma^2}\right] \frac{p(\theta|\vec{\lambda})}{p_{\text{det}}(\vec{\lambda})} d\theta \\
 &= -\sigma^3 \sqrt{2\pi} \int \frac{\partial p(\mathbf{d}_{\text{th}}|\theta)}{\partial \mathbf{d}_{\text{th}}} \frac{p(\theta|\vec{\lambda})}{p_{\text{det}}(\vec{\lambda})} d\theta \\
 &= -\frac{\sigma^3 \sqrt{2\pi}}{p_{\text{det}}(\vec{\lambda})} \int \frac{\partial}{\partial \mathbf{d}_{\text{th}}} [p(\mathbf{d}_{\text{th}}|\theta) p(\theta|\vec{\lambda})] d\theta \\
 &= \frac{\sigma^3 \sqrt{2\pi}}{(\sigma^2 + \Sigma^2)} (\mathbf{d}_{\text{th}} - \mu) \frac{p(\mathbf{d}_{\text{th}}|\vec{\lambda})}{p_{\text{det}}(\vec{\lambda})}.
 \end{aligned}$$

The fourth term  $(\Gamma_{IV})_{\Sigma^2\Sigma^2}$  is found using (37), and reads

$$\begin{aligned}
 (\Gamma_{IV})_{\Sigma^2\Sigma^2} &= \int \frac{\partial^2}{(\partial\Sigma^2)^2} [P(\Gamma + H)^{-1}] D_{\theta} \frac{p_{\text{det}}(\theta)}{p_{\text{det}}(\vec{\lambda})} p(\theta|\vec{\lambda}) d\theta \\
 &= \frac{2\sigma^2\Sigma^2}{(\sigma^2 + \Sigma^2)^4} (\mathbf{d}_{\text{th}} - \mu) \frac{p(\mathbf{d}_{\text{th}}|\vec{\lambda})}{p_{\text{det}}(\vec{\lambda})}.
 \end{aligned} \quad (43)$$


Finally, the final term  $(\Gamma_V)_{\Sigma^2\Sigma^2}$  is found through (39) to be

$$\begin{aligned}
 (\Gamma_V)_{\Sigma^2\Sigma^2} &= -\frac{1}{2} \int \frac{\partial^2}{(\partial\Sigma^2)^2} [P^2(\Gamma + H)^{-1}] \frac{p_{\text{det}}(\theta)}{p_{\text{det}}(\vec{\lambda})} p(\theta|\vec{\lambda}) d\theta \\
 &= -\frac{\sigma^2}{\Sigma^4} \frac{(\sigma^4 + 3\sigma^2 + \Sigma^2 + 3\Sigma^4)}{(\sigma^2 + \Sigma^2)^3} \\
 &\quad \left[ 1 + \frac{\Sigma^2}{(\sigma^2 + \Sigma^2)} (\mathbf{d}_{\text{th}} - \mu) \frac{p(\mathbf{d}_{\text{th}}|\vec{\lambda})}{p_{\text{det}}(\vec{\lambda})} \right].
 \end{aligned} \quad (44)$$

<sup>2</sup> The integral over  $\theta$  can be evaluated by parts to obtain

$$\int (\theta - \mu)^2 \frac{p_{\text{det}}(\theta)}{p_{\text{det}}(\vec{\lambda})} p(\theta|\vec{\lambda}) d\theta = \Sigma^2 + \frac{(\mathbf{d}_{\text{th}} - \mu)\Sigma^4}{(\sigma^2 + \Sigma^2)} \frac{p(\mathbf{d}_{\text{th}}|\vec{\lambda})}{p_{\text{det}}(\vec{\lambda})}, \quad (39)$$

using (28) and (37) with vanishing boundary terms.



**Figure 1.** Normally-distributed data with (red) and without (black) selection effects.

Adding up (40) to (44), we find

$$\Gamma_{\Sigma^2\Sigma^2} = \frac{\partial^2 \ln p_{\text{det}}(\vec{\lambda})}{(\partial\Sigma^2)^2} + \frac{1}{2(\sigma^2 + \Sigma^2)^2} + \frac{(\mathbf{d}_{\text{th}} - \mu)}{(\sigma^2 + \Sigma^2)^2} \frac{p(\mathbf{d}_{\text{th}}|\vec{\lambda})}{p_{\text{det}}(\vec{\lambda})}, \quad (45)$$

which matches (27) as expected. This concludes the analytic check. Equation (21) can be used to reproduce the predictions (27) obtained from the more general definition of the Fisher matrix as an expectation value over data realizations.


### 3.2 MCMC analysis

The Fisher predictions for the Gaussian-Gaussian model can be compared with MCMC simulations as a further check of the formalism. While this example is arguably textbook material, see for instance Sec. (6) in (Vitale et al. 2020), we report a few details below for completeness. The results of the present section can be fully reproduced with the codes accompanying the present paper. We simulate synthetic data including  $N_{\text{tot}} = 10^5$  observations from the observation model (23), choosing true mean  $\mu_{\text{tr}} = 0.5$ , true variance  $\Sigma_{\text{tr}}^2 = 1.0$  and noise variance  $\sigma = 0.1$ . The latter two indicate that each event is taken with a high SNR. We then apply an arbitrary cutoff, imposing that only positive data are observed. That is,  $\mathbf{d}_{\text{th}} = 0$ , resulting in around  $N_{\text{det}} \sim 70000$  detected events. Figure 1 shows the total (black) and detected (red) populations under our specified assumptions. We perform the MCMC analyses in both cases in which we do and do not have selection effects using `emcee` (Foreman-Mackey et al. 2013). As log-likelihood, we take the sum of individual log-likelihoods,

$$\log p(d|\lambda) = -N \log p_{\text{det}}(\lambda) + \sum_i^N \log p(d_i|\lambda), \quad (46)$$

where  $N = N_{\text{tot}}$  in the case in which we do not include selection effects, and  $N = N_{\text{det}}$  in the case in which we do. We choose flat hyperpriors over a very broad range that includes the true values. The selection function is defined and integrated as in Eq. (28), and is nontrivial only in the latter case. The MCMC posteriors for  $\mu$  and  $\Sigma^2$ , in both the cases considered, can be found in Fig. 2.

We can then compare the predictions from the population Fisher matrix with what we obtain numerically. To this



**Figure 2.** Fisher and MCMC predictions for the toy Gaussian case with (red) and without (black) selection effects.

end, we invert the matrix

$$(\Gamma\lambda)_{ij} = \begin{pmatrix} \Gamma_{\mu\mu} & \Gamma_{\mu\Sigma^2} \\ \Gamma_{\mu\Sigma^2} & \Gamma_{\Sigma^2\Sigma^2} \end{pmatrix} \quad (47)$$

with entries given in Eq. (27) and below. The errors are normalized by the number of events,

$$\Delta\mu = \sqrt{N(\Gamma_\lambda^{-1})_{\mu\mu}}, \quad \Delta\Sigma^2 = \sqrt{N(\Gamma_\lambda^{-1})_{\Sigma^2\Sigma^2}}.$$

In Fig. 2, the Fisher predictions are shown (in black) to reproduce the widths from the MCMC runs.

The same widths can be well approximated by the inverse of the matrix

$$(\Gamma\lambda)_{ij} = \begin{pmatrix} (\Gamma_1)_{\mu\mu} & (\Gamma_1)_{\mu\Sigma^2} \\ (\Gamma_1)_{\mu\Sigma^2} & (\Gamma_1)_{\Sigma^2\Sigma^2} \end{pmatrix}, \quad (48)$$

in which only the first terms in Eq. (21) are retained. The predictions are shown in red in Fig. 2, and they overlap well with the full Fisher matrix predictions. Interestingly, retaining the first terms in Eq. (48) is equivalent to setting  $\sigma = 0$  in Eq. (47), which is the limit in which the parameters of the individual events are measured perfectly. This suggests that the reason that the approximation in Eq. (48) works well in this case is because the individual measurements are much more precise than the scale over which the population is varying. We would expect the other terms contributing to Eq. (21) to become increasingly important as the measurement errors become larger. This hypothesis holds exactly in

the simplified Gaussian-Gaussian case considered here, but we will revisit it in the GW-like illustration in the next section.

#### 4 ILLUSTRATION II: AN EXAMPLE FROM GRAVITATIONAL-WAVE ASTROPHYSICS

The spaceborne LISA mission is expected to detect extreme mass ratio inspirals (EMRIs), namely binary systems in which one compact object, typically a stellar remnant, has a mass that is much smaller than the companion, typically a supermassive black hole (SMBH) in the centre of a galaxy (Amaro-Seoane et al. 2007; Barack 2009; Babak et al. 2017). Inference of the parameters that characterise EMRI systems is expected to provide accurate constraints on the theory of gravity (Gair et al. 2013), as well as an insight into the astrophysical population of and stellar environments surrounding SMBHs (Barausse, Cardoso & Pani 2014). LISA might also detect a foreground generated by individually unresolved EMRIs, which would provide information about the properties of the population of these systems (Gair et al. 2011; Gair, Tang & Volonteri 2010; Sesana et al. 2011; Bonetti & Sesana 2020). The use of LISA observations of EMRIs to provide measurements of the BH mass function in the range probed by LISA has previously been investigated in (Gair, Tang & Volonteri 2010), henceforth ‘‘GTV’’. GTV assumed that the mass function was described by a power law  $p(\theta|\lambda) \equiv p(M|\lambda) \propto M^{\alpha-1}$ , with a true value that is close to flat in the log of the masses, i.e.,  $\alpha \approx 0$ . GTV explored the ability of LISA to constrain the parameters of this mass function, using MCMC techniques to carry out hierarchical analyses on an extensive set of populations of simulated events. They found that with 10(1000) events, the spectral index  $\alpha$  could be constrained at a level of precision  $\Delta\alpha = 0.3(0.03)$ . Here, we will use the population Fisher Matrix formalism described above to predict the precision with which a set of EMRI observations might be able to constrain a power-law mass function. We do not expect to get exactly the same answer here, as the two analyses make a few different simplifying assumptions. In GTV the raw data was taken to be counts of events in a binned analysis, provided by point estimates of the parameters. Selection effects were included in the rate of events in each bin, by accounting for the length of time a source with the given parameters would be observable. This ignores the fact that the time remaining to plunge is constrained by the gravitational wave data. In this analysis we again approximate the observation process, assuming that the data can be reduced to a measurement of a single parameter, but we handle selection effects more carefully. GTV’s results was computed ignoring measurement uncertainties in the model used in the analysis, although they did demonstrate consistency between results obtained on simulated data with and without measurement uncertainties. Here we will include measurement uncertainties, but we will approximate these as Gaussian, with equal variance for all observations. We will see that despite these differences in assumptions, the population Fisher matrix is able to predict qualitatively the results observed in GTV without the need for costly computational sampling of many posterior distributions. For a more direct comparison with numerical results we also perform our own MCMC analysis,

under identical assumptions to those used to compute the population Fisher matrix, and find even better agreement. All the results in this section can be reproduced with the codes made available with this publication.

We draw  $N = 100$  masses from a power law distribution,

$$p(M|\alpha) = \frac{\alpha}{M_{\max}^{\alpha} - M_{\min}^{\alpha}} M^{\alpha-1}, \quad (49)$$

with maximal and minimal observable masses  $M_{\min} = 10^4 M_{\odot}$  and  $M_{\max} = 10^7 M_{\odot}$  and a true value for the spectral index that is exactly flat in the log of the masses,  $\alpha = 0$ . The observed data is assumed to be a point estimate of the log of the mass, which is equal to the true value plus a normally-distributed uncertainty that has a variance  $\sigma = 0.1$ . An arbitrary hard cutoff  $\mathbf{d}_{\text{th}}$  corresponding to masses  $M \sim 5 \times 10^5 M_{\odot}$  is imposed, and only events with observed values above this threshold are included in the analysis. This introduces a large selection effect, and leads to only  $N_{\text{det}} = 39$  of the original 100 sources being observed. The true (underlying) and observed populations of events are represented in the top panel of Fig. 3.

The Fisher-matrix prediction can again be obtained from Eq. (21), this time with  $\theta = M$  and  $\lambda = \alpha$ . For sufficiently simple models, the integrals are analytically tractable. With the power-law distribution considered here, and in the presence of selection effects, this is already not possible. Because of this, we obtain the Fisher prediction in a semi-analytical fashion by solving the integrals with Monte-Carlo methods, i.e., by generating a sufficiently large set of  $N_s$  samples,  $\{M_i\}$ , from a distribution  $p(M|\alpha)$ , we can approximate the integral of an arbitrary function  $X(M)$  via

$$\int X(M)p(M|\alpha)dM \approx \frac{1}{N_s} \sum_i X(M_i). \quad (50)$$

In this case we can draw samples from the distribution (49) directly using the method of inversion. The various terms entering the arguments of Eq. (21) —  $D_i$ ,  $D_{ij}$ ,  $P_i$  and  $H_{ij}$  — can be computed analytically. A *Mathematica* notebook that solves for the arguments of the Fisher-matrix integrals can be found in the accompanying codes. Following this procedure, we find that the Fisher matrix predicts an error  $\Delta\alpha = \sqrt{(N_{\text{det}}\Gamma_{\alpha})^{-1}} \approx 0.19$ . When rescaled to 10 observations by multiplying by  $\sqrt{39/10}$ , the inferred error is  $\Delta\alpha \approx 0.37$ , which is in good agreement with what was obtained in GTV, despite the differences in the assumptions used in each case.

As in the Gaussian-Gaussian example, we find that the dominant contribution comes from the first term of the Fisher matrix,  $\Gamma_I$ . Once again, this behaviour appears to be driven by the assumed precision of measurement on the individual events. If the noise in the individual measurements,  $\sigma$ , is increased, the other terms make a larger contribution, although always a sub-dominant contribution for the range of values we have tried. This fact suggests that, when individual events are expected to be characterised with a precision better than the typical lengthscale over which the population prior varies, keeping only the first term in the sum will provide a good estimate for the expected precision of population inference, regardless of the population model chosen. This observation could help to reduce the complexity of using our formalism.

We validate the Fisher predictions with an MCMC anal-

ysis for the same data set. The MCMC setup is similar to the one used in the first illustrative example. The likelihood is modified, and the selection function is no longer known analytically, but is approximated by a Monte Carlo integral,  $p_{\text{det}}(\alpha) = (1/N_s) \sum \text{erfc}[(\mathbf{d}_{\text{th}} - M_i)/\sqrt{2\sigma^2}]/2$ , with  $\{M_i\}$  drawn from the power law distribution  $p(M|\alpha)$ . The posterior, KDE and  $2\sigma$  percentile for the estimate of  $\alpha$  are shown in the bottom panel of Fig. (3). These are compared with the Fisher predictions above, and once again show very good agreement. We have also repeated the calculation in the absence of selection effects,  $\mathbf{d}_{\text{th}} \rightarrow -\infty$ , and find a similar level of agreement between the Fisher predictions and the MCMC analysis. In this case, the first (dominant) term of the Fisher matrix can be integrated to get the following estimate of precision

$$\Gamma_{\alpha} \approx N_{\text{tot}} \left( \frac{1}{\alpha^2} - \frac{M_{\max}^{\alpha} M_{\min}^{\alpha} (\ln M_{\max} - \ln M_{\min})^2}{(M_{\max}^{\alpha} - M_{\min}^{\alpha})^2} \right). \quad (51)$$


The relative ease in obtaining and evaluating the above expression and the fact that it leads to results that are consistent with independent MCMC analyses showcase how useful our Fisher formalism could be in forecast studies for future detectors, which could nicely complement the source parameter precision estimated with state-of-the-art Fisher codes (Borhanian 2021; Harms et al. 2022).

We expect that the accuracy of the population Fisher matrix prediction should improve as the number of observations included in the analysis,  $N_{\text{tot}}$ , increases. We assess this by comparing the result of MCMC analyses of data sets with increasing numbers of observations to the population Fisher matrix. We do this first for the case without selection effects, so that we can make use of the analytical prediction given above. These results are shown in Fig. 4 as  $N_{\text{tot}}$  is varied from 2 to 30 events. For each  $N_{\text{tot}}$ , we repeat the MCMC analysis several times to allow us to estimate the variance in the posterior width between different runs. This variance is larger when there are fewer events, as expected, and so more MCMC analyses were performed for lower  $N_{\text{tot}}$ 's to ensure the variance was accurately characterised. We find that for  $N_{\text{tot}} \gtrsim 10$  the (simplified) Fisher widths agree well with numerical simulations. For  $N_{\text{tot}} < 10$ , the differences are progressively more pronounced, but the variance in the MCMC widths also increases, and the Fisher matrix prediction is usually within the range spanned by the MCMC runs. The agreement becomes worse for very small numbers of observations, consistent with the expectation that this is an approximation valid in the limit of large  $N_{\text{tot}}$ . Finally, we check whether a similar level of agreement is seen in the case when the observations are subject to selection effects. For this, we obtain the population Fisher matrix prediction (dashed orange line) by rescaling the  $\Delta\alpha$  prediction in Fig. 3 by a factor  $N_{\text{det}}^{-1/2}$ . These results are also shown in Fig. 3. We see a similar trend — the population Fisher matrix is very accurate for  $N_{\text{tot}} > 10$ , but the accuracy diminishes for very small numbers of observations, as expected.


## 5 CONCLUSIONS

The Fisher information matrix is a valuable tool for estimating the precision attainable in parameter inference, especially in contexts where the cost of doing full posterior





**Figure 3.** *Top panel:* distribution of masses for the underlying true population of Sec. 4 (red) and for the observed population (dark gray). The true population is composed of  $N = 100$  events drawn from a power-law model that is flat in the logarithm of the masses. The threshold has been arbitrarily set at  $\sim 5 \times 10^5 M_\odot$ , leading to 39 events actually being observed. *Bottom panel:* MCMC posterior distribution for the spectral index describing the mass distribution. The histogram and KDE are compared against the Fisher estimate obtained as described in the text, demonstrating very good agreement between the two.



**Figure 4.** Fisher predictions for the width of the spectral index  $\alpha$  as a function of the number of observed events  $N$ , compared to the range of measured uncertainties obtained over a set of MCMC runs (red and black points with error bars). The Fisher matrix prediction is approximated with Eq. (51). The agreement between the Fisher matrix predictions and the MCMC results is very good, especially if the number of events is increased beyond  $N \sim 10$ .

estimation via Bayesian sampling is highly expensive (Valisneri 2008). The Fisher matrix has been widely used in GW analyses to make forecasts for the precision with which the *source* parameters describing individual GW signals can be estimated by current and future detectors. In this paper we have extended the Fisher matrix concept to the estimation of the parameters characterising the *population* from

which a set of observed sources is drawn. Our result was based on Eq. (1), which is the most general definition for the population Fisher matrix. However, we have obtained a simpler to evaluate expression, Eq. (21), which is valid when the signal-to-noise ratio of the individual events making up the population is high. We have tested this result both analytically and against numerical Monte-Carlo results for a reference Gaussian model (Sec. 3) and for a more GW-like scenario (Sec. 4), in which we are using GW events to estimate the slope of a power-law population. We find that Eq. (21) is generally in very good agreement with the numerical results, for a sufficiently large number of observations. In this case sufficiently large was only  $O(10)$ . Results for the power-law population case can be compared to previous results in the literature (Gair, Tang & Volonteri 2010), and are found to be in very good agreement. We conclude that we can reproduce the results of extensive sets of computationally expensive MCMC simulations much more cheaply, while also including a more careful treatment of selection effects. In addition, we have found that the first term in the sum that gives the population Fisher matrix,  $(\Gamma_I)_{ij}$  in Eq. (21), is enough to reproduce well the precision attainable by the full Fisher matrix in both of the examples considered. This is most likely due to the fact that the noise-induced uncertainty in the parameter measurements of each individual event,  $\sigma$ , is much smaller than the scale on which the population model varies, and so measurement errors are essentially ignorable. This result could be useful to further reduce the computational cost of computing the population Fisher matrix in many contexts.

The results presented in this paper represent the first attempt at describing population inference within a Fisher formalism for generic population models, and with a likelihood that takes into account selection effects in the way of (Mandel, Farr & Gair 2019). The formalism herein developed can be used to obtain forecasts for the precision of population analyses with future ground-based and spaceborne detectors, which are expected to detect many thousands (or even millions) of signals. Obtain such population inference forecasts in specific contexts of relevance to current and future observations is one possible future direction for the present project. Finally, it would be interesting to generalize the results of (Cutler & Vallisneri 2007) and (Antonelli, Burke & Gair 2021) to assess inference biases on population parameters from waveform modelling errors or confusion noise.

*Data Availability Statement.* The results in this paper use `numpy` (Harris et al. 2020), `matplotlib` (Hunter 2007), `seaborn` (Waskom 2021), `emcee` (Foreman-Mackey et al. 2013), `arviz` (Kumar et al. 2019). The results can be fully reproduced with codes made publicly available at <https://github.com/aantonelli94/PopFisher>.

## REFERENCES

- Abbott B. P., et al., 2019, *Phys. Rev. X*, 9, 031040  
 Abbott R., et al., 2021a  
 Abbott R., et al., 2021b  
 Abbott R., et al., 2021c  
 Abbott R., et al., 2021d  
 Amaro-Seoane P., Gair J. R., Freitag M., Coleman Miller M., Mandel I., Cutler C. J., Babak S., 2007, *Class. Quant. Grav.*, 24, R113  
 Amaro-Seoane P., et al., 2017  
 Antonelli A., Burke O., Gair J. R., 2021, *Mon. Not. Roy. Astron. Soc.*, 507, 5069  
 Babak S. et al., 2017, *Phys. Rev. D*, 95, 103012  
 Barack L., 2009, *Class. Quant. Grav.*, 26, 213001  
 Barausse E., Cardoso V., Pani P., 2014, *Phys. Rev. D*, 89, 104059  
 Bonetti M., Sesana A., 2020, *Phys. Rev. D*, 102, 103023  
 Borhanian S., 2021, *Class. Quant. Grav.*, 38, 175014  
 Cutler C., Vallisneri M., 2007, *Phys. Rev. D*, 76, 104018  
 Foreman-Mackey D., Hogg D. W., Lang D., Goodman J., 2013, *Publ. Astron. Soc. Pac.*, 125, 306  
 Gair J. R., Sesana A., Berti E., Volonteri M., 2011, *Class. Quant. Grav.*, 28, 094018  
 Gair J. R., Tang C., Volonteri M., 2010, *Phys. Rev. D*, 81, 104014  
 Gair J. R., Vallisneri M., Larson S. L., Baker J. G., 2013, *Living Rev. Rel.*, 16, 7  
 Harms J. et al., 2022  
 Harris C. R. et al., 2020, *Nature*, 585, 357  
 Hunter J. D., 2007, *Computing in Science & Engineering*, 9, 90  
 Kumar R., Carroll C., Hartikainen A., Martin O., 2019, *Journal of Open Source Software*, 4, 1143  
 Mandel I., Farr W. M., Gair J. R., 2019, *Mon. Not. Roy. Astron. Soc.*, 486, 1086  
 Punturo M., et al., 2010, *Class. Quant. Grav.*, 27, 194002  
 Reitze D., et al., 2019, *Bull. Am. Astron. Soc.*, 51, 035  
 Sesana A., Gair J., Berti E., Volonteri M., 2011, *Phys. Rev. D*, 83, 044036  
 Vallisneri M., 2008, *Phys. Rev. D*, 77, 042001  
 Vitale S., Gerosa D., Farr W. M., Taylor S. R., 2020  
 Waskom M. L., 2021, *Journal of Open Source Software*, 6, 3021

## 1 Dealing with rates

The likelihood in Eq. (6) assumes that the number of events that are detected conveys no information about the population. Relaxing this assumption the joint likelihood takes the alternative form

$$p(\{\mathbf{d}_i\}|\vec{\lambda}, R) \propto \left[ \prod_{i=1}^n p_{\text{full}}(\mathbf{d}_i|\vec{\lambda}) \right] R^n \exp[-Rp_{\text{det}}(\vec{\lambda})], \quad (1)$$

in which  $R$  is the rate of events occurring in the Universe over the total time data has been collected, and all other terms are as before. The derivation of this expression can be found in (Mandel, Farr & Gair 2019). Imposing an (improper) scale-invariant prior on the total rate,  $p(R) \propto 1/R$ , and marginalising over  $R$  we obtain the form of the joint likelihood used in Eq. (6) and (8).

We denote the rate-dependent terms by

$$p_{\text{rate}}(n|R, \vec{\lambda}) \equiv R^n \exp[-Rp_{\text{det}}(\vec{\lambda})]. \quad (2)$$

The contribution of these terms to the joint log-likelihood is

$$\ln p_{\text{rate}}(n|R, \vec{\lambda}) = n \ln R - Rp_{\text{det}}(\vec{\lambda}), \quad (3)$$

which is maximized when  $R = n/p_{\text{det}}(\vec{\lambda})$ . Expanding about this maximum-likelihood point we can write  $R = n/p_{\text{det}} + \delta R$  and obtain

$$\begin{aligned} \ln p_{\text{rate}}(n|R, \vec{\lambda}) &= n \ln n - n \ln p_{\text{det}}(\vec{\lambda}) + n \ln(1 + \delta R p_{\text{det}}/n) \\ &\quad - n - \delta R p_{\text{det}} \\ &= (n \ln n - n) - n \ln p_{\text{det}}(\vec{\lambda}) - \frac{\delta R^2 p_{\text{det}}^2}{2n} + \dots \end{aligned} \quad (4)$$

The second term here,  $-n \ln p_{\text{det}}(\vec{\lambda})$ , is what is needed to change  $p_{\text{full}}(\mathbf{d}_i|\vec{\lambda})$  into  $p(\mathbf{d}_i|\vec{\lambda})$  in the product term in Eq. (1), reducing that to the form analysed in the main body of the paper. We deduce that the asymptotic Fisher matrix for the joint estimation of  $R$  and  $\vec{\lambda}$  is block diagonal

$$\Gamma = \begin{pmatrix} \Gamma_{\lambda} & 0 \\ 0 & \Gamma_R \end{pmatrix} \quad (5)$$

with the shape parameter block,  $\Gamma_{\lambda}$ , as before and the rate precision given by the inverse of  $\Gamma_R = p_{\text{det}}^2/2n$ , where this result can be obtained directly from the coefficient of  $\delta R^2$  in equation (4). We conclude that the precision with which the shape parameters can be determined does not depend on which particular form of the likelihood is being used. This makes sense since we know that the two forms are equivalent for a particular choice of rate prior, and we expect results to be asymptotically independent of the initial prior choice. We note also that the precision with which the rate of observed events,  $R p_{\text{det}}$ , can be measured is  $\sqrt{n}$ , consistent with the expected uncertainty in the estimation of the rate of a Poisson process.

To conclude this section, we note that in the above we have been assuming that the rate parameter  $R$  is an additional parameter of the model, separate to the parameters  $\vec{\lambda}$  that characterise the shape of the population distribution. If instead both  $R$  and  $\vec{\lambda}$  are functions of another set of population parameters,  $\vec{\mu}$ , we can use the linear signal approximation to change variables and obtain the usual result that the Fisher matrix for the  $\vec{\mu}$  parameters is

$$(\Gamma_\mu)_{cd} = (\Gamma_\lambda)_{ab} \frac{\partial \lambda^a}{\partial \mu^c} \frac{\partial \lambda^b}{\partial \mu^d} + \Gamma_R \frac{\partial R}{\partial \mu^c} \frac{\partial R}{\partial \mu^d}. \quad (6)$$

In this case, the result is different to what would be obtained by transforming the Fisher matrix that ignores the rate, which would be the first term only. This reflects the fact that if  $R$  also depends on  $\vec{\mu}$ , the measurement of  $R$  provides additional information that can help to improve the estimation of the  $\vec{\mu}$  parameters.

## 2 The asymptotic behaviour of the Fisher matrix

In this section we will rederive the expression for the Fisher matrix given in the main body of the paper by directly expanding the posterior distribution. In doing so we will derive the form and scaling of the leading corrections to the Fisher matrix approximation. We will proceed by computing the posterior mode, mean and variance in the limit  $n \gg 1$ . These are all random variables, since they depend on the particular realisation of the data that is being analysed, and so we can characterise them by their expectation value and variance. We will show how to compute the first two terms in a large- $n$  expansion of both the mean and variance for all three posterior summary statistics, and give the result explicitly for the posterior mode. We write

$$\hat{\mu}(n, \{\mathbf{d}_i\}, \vec{\lambda}) = -\frac{1}{n} \sum_{i=1}^n \ln p(\mathbf{d}_i | \vec{\lambda}). \quad (7)$$

such that the posterior distribution is proportional to  $\exp[-n\hat{\mu}(n, \{\mathbf{d}_i\}, \vec{\lambda}) + \ln \pi(\vec{\lambda})]$ . We use  $\vec{\beta}(n, \{\mathbf{d}_i\}, \vec{\lambda}_t)$  to denote the solution to

$$U_i + U_i^\pi + (V_{ij} + V_{ij}^\pi) \beta^j = 0 \quad (8)$$

where

$$U_i = \left( \frac{\partial \hat{\mu}}{\partial \lambda^i} \right)_{|\vec{\lambda}_t}, \quad U_i^\pi = -\frac{1}{n} \left( \frac{\partial \ln \pi}{\partial \lambda^i} \right)_{|\vec{\lambda}_t}$$

$$V_{ij} = \left( \frac{\partial^2 \hat{\mu}}{\partial \lambda^i \partial \lambda^j} \right)_{|\vec{\lambda}_t}, \quad V_{ij}^\pi = -\frac{1}{n} \left( \frac{\partial^2 \ln \pi}{\partial \lambda^i \partial \lambda^j} \right)_{|\vec{\lambda}_t}. \quad (9)$$

The quantity  $\hat{\mu}$ , and its derivatives, are averages of a set of independent identically distributed (IID) random variables and so have predictable scalings. The expectation value is  $O(1)$ , covariances are  $O(1/n)$ , three and four point functions are  $O(1/n^2)$ , and so on. In this case  $\mathbb{E}[U] = 0$ , as shown in Eq. (12) in the main body of the paper. This facilitates obtaining a solution for  $\vec{\beta}$  perturbatively

$$\vec{\beta} = \vec{\beta}_{\frac{1}{2}} + \vec{\beta}_1 + \vec{\beta}_{\frac{3}{2}} + \dots$$

$$\text{where } \vec{\beta}_{\frac{1}{2}} = (\mathbb{E}[\mathbf{V}])^{-1} \mathbf{U}$$

$$\vec{\beta}_1 = (\mathbb{E}[\mathbf{V}])^{-1} \left[ (\mathbf{V} - \mathbb{E}[\mathbf{V}]) \vec{\beta}_{\frac{1}{2}} + \mathbf{U}^\pi \right]$$

$$\vec{\beta}_{\frac{3}{2}} = (\mathbb{E}[\mathbf{V}])^{-1} \left[ (\mathbf{V} - \mathbb{E}[\mathbf{V}]) \vec{\beta}_1 + \mathbf{V}^\pi \vec{\beta}_{\frac{1}{2}} \right], \quad (10)$$

in which  $\vec{\beta}_k \sim n^{-k}$ .

The  $\vec{\beta}_k$ 's are random variables, but we can compute their means

$$\mathbb{E} \left[ \vec{\beta}_{\frac{1}{2}} \right] = 0$$

$$\mathbb{E} \left[ \beta_1^i \right] = (\mathbb{E}[\mathbf{V}])_{ij}^{-1} \left[ (\mathbb{E}[\mathbf{V}])_{kl}^{-1} \Sigma_{jkl}^{VU} + \mathbf{U}_j^\pi \right]$$

$$\mathbb{E} \left[ \beta_{\frac{3}{2}}^i \right] = (\mathbb{E}[\mathbf{V}])_{ij}^{-1} (\mathbb{E}[\mathbf{V}])_{kl}^{-1} (\mathbb{E}[\mathbf{V}])_{mn}^{-1} \Sigma_{jklmn}^{VUU}$$

$$\Sigma_{ijk}^{VU} = \mathbb{E}[(V_{ij} - \mathbb{E}[V_{ij}])U_k]$$

$$= \frac{1}{n} \int l_{,ij}(\mathbf{d}) l_{,k}(\mathbf{d}) p(\mathbf{d} | \vec{\lambda}_t) d\mathbf{d}$$

$$\Sigma_{ijklm}^{VUU} = \mathbb{E}[(V_{ij} - \mathbb{E}[V_{ij}]) (V_{kl} - \mathbb{E}[V_{kl}]) U_m]$$

$$= \frac{1}{n^2} \int l_{,ij}(\mathbf{d}) l_{,kl}(\mathbf{d}) l_{,m}(\mathbf{d}) p(\mathbf{d} | \vec{\lambda}_t) d\mathbf{d}$$

$$- \mathbb{E}[V_{ij}] \Sigma_{klm}^{VU} - \mathbb{E}[V_{kl}] \Sigma_{ijm}^{VU} \quad (11)$$

where we are using the notation  $l_{,i}(\mathbf{d})$  to denote the derivative  $\partial \ln p(\mathbf{d} | \vec{\lambda}) / \partial \lambda^i$  evaluated at  $\vec{\lambda} = \vec{\lambda}_t$ . Additional indices after the comma indicate further partial derivatives as usual. The fact that the first expectation value vanishes is why we have continued the expansion to three terms, allowing us to obtain the first two terms in an expansion of the mean.

We can also compute their covariances, using the usual notation  $\text{cov}(a^i, b^j) = \mathbb{E}[(a^i - \mathbb{E}[a^i])(b^j - \mathbb{E}[b^j])]$ .

$$\text{cov}(\beta_{\frac{1}{2}}^i, \beta_{\frac{1}{2}}^j) = (\mathbb{E}[\mathbf{V}])_{ik}^{-1} (\mathbb{E}[\mathbf{V}])_{jl}^{-1} \Sigma_{kl}^{UU}$$

$$\text{cov}(\beta_{\frac{1}{2}}^i, \beta_1^j) = (\mathbb{E}[\mathbf{V}])_{im}^{-1} (\mathbb{E}[\mathbf{V}])_{jk}^{-1} (\mathbb{E}[\mathbf{V}])_{ln}^{-1} \Sigma_{klmn}^{VUU}$$

$$\text{cov}(\beta_1^i, \beta_1^j) = (\mathbb{E}[\mathbf{V}])_{ik}^{-1} (\mathbb{E}[\mathbf{V}])_{lm}^{-1} (\mathbb{E}[\mathbf{V}])_{jp}^{-1} (\mathbb{E}[\mathbf{V}])_{qr}^{-1} \Sigma_{klpqmr}^{VUUU}$$

$$- (\mathbb{E}[\mathbf{V}])_{ik}^{-1} (\mathbb{E}[\mathbf{V}])_{lm}^{-1} \Sigma_{klm}^{VU} \Sigma_{pq}^{VU}$$

$$\text{cov}(\beta_{\frac{3}{2}}^i, \beta_{\frac{3}{2}}^j) = (\mathbb{E}[\mathbf{V}])_{ik}^{-1} (\mathbb{E}[\mathbf{V}])_{jq}^{-1} (\mathbb{E}[\mathbf{V}])_{lm}^{-1}$$

$$\times \left[ (\mathbb{E}[\mathbf{V}])_{np}^{-1} \Sigma_{klmnpq}^{VUUU} + \mathbf{U}_m^\pi \Sigma_{klq}^{VU} + \mathbf{V}_{kl}^\pi \Sigma_{klmq}^{VUUU} \right]$$

$$\Sigma_{ij}^{UU} = \mathbb{E}[U_i U_j]$$

$$= \frac{1}{n} \int l_{,i}(\mathbf{d}) l_{,j}(\mathbf{d}) p(\mathbf{d} | \vec{\lambda}_t) d\mathbf{d}$$

$$\Sigma_{ijkl}^{VUU} = \mathbb{E}[(V_{ij} - \mathbb{E}[V_{ij}]) U_k U_l]$$

$$= \frac{1}{n^2} \int l_{,ij}(\mathbf{d}) l_{,k}(\mathbf{d}) l_{,l}(\mathbf{d}) p(\mathbf{d} | \vec{\lambda}_t) d\mathbf{d}$$

$$- \mathbb{E}[V_{ij}] \Sigma_{kl}^{UU}$$

$$\Sigma_{ijklmn}^{VUUU} = \mathbb{E}[(V_{ij} - \mathbb{E}[V_{ij}]) (V_{kl} - \mathbb{E}[V_{kl}]) U_m U_n]$$

$$= \frac{n(n-1)}{n^4} \left[ \int l_{,ij}(\mathbf{d}) l_{,kl}(\mathbf{d}) p(\mathbf{d} | \vec{\lambda}_t) d\mathbf{d} \right.$$

$$\times \int l_{,m}(\mathbf{d}) l_{,n}(\mathbf{d}) p(\mathbf{d} | \vec{\lambda}_t) d\mathbf{d}$$

$$+ \int l_{,ij}(\mathbf{d}) l_{,m}(\mathbf{d}) p(\mathbf{d} | \vec{\lambda}_t) d\mathbf{d}$$

$$\times \int l_{,kl}(\mathbf{d}) l_{,n}(\mathbf{d}) p(\mathbf{d} | \vec{\lambda}_t) d\mathbf{d}$$

$$+ m \leftrightarrow n \left. \right]$$

$$+ \frac{1}{n^3} \int l_{,ij}(\mathbf{d}) l_{,kl}(\mathbf{d}) l_{,m}(\mathbf{d}) l_{,n}(\mathbf{d}) p(\mathbf{d} | \vec{\lambda}_t) d\mathbf{d}$$

$$- \mathbb{E}[V_{ij}] \Sigma_{klmn}^{VUU} - \mathbb{E}[V_{kl}] \Sigma_{ijmn}^{VUU}$$

$$- \mathbb{E}[V_{ij}] \mathbb{E}[V_{kl}] \Sigma_{mn}^{UU} \quad (12)$$

All four of these terms are needed to obtain the first two terms in the covariance of  $\vec{\beta}$ . We need only retain terms of  $O(1/n^2)$  in the final expression, which means the last term in  $\Sigma_{ijklmn}^{VUUU}$  can be ignored and  $n(n-1)/n^4$  replaced by  $n^{-2}$ . The final result, keeping the first two orders is

$$\begin{aligned} \text{cov}(\beta^i, \beta^j) &= \text{cov}\left(\beta_{\frac{1}{2}}^i, \beta_{\frac{1}{2}}^j\right) + \text{cov}\left(\beta_{\frac{1}{2}}^i, \beta_1^j\right) \\ &+ \text{cov}\left(\beta_{\frac{1}{2}}^j, \beta_1^i\right) + \text{cov}\left(\beta_1^i, \beta_1^j\right) \\ &+ \text{cov}\left(\beta_{\frac{3}{2}}^i, \beta_{\frac{3}{2}}^j\right) + \text{cov}\left(\beta_{\frac{3}{2}}^j, \beta_{\frac{3}{2}}^i\right) \end{aligned} \quad (13)$$

The posterior mode is at  $\vec{\lambda} = \vec{\lambda}_t + \vec{\beta} + \vec{\delta}$ . The random variable  $\vec{\delta}$  obeys the equation

$$\begin{aligned} 0 &= \delta^j V_{ij} + \frac{1}{2}(\beta^j + \delta^j)(\beta^k + \delta^k)W_{ijk} + V_{ij}^\pi \delta^j \\ &+ \frac{1}{2}(\beta^j + \delta^j)(\beta^k + \delta^k)W_{ijk}^\pi \\ W_{ijk} &= \left. \frac{\partial^3 \hat{\mu}}{\partial \lambda^i \partial \lambda^j \partial \lambda^k} \right|_{\vec{\lambda}_t} \\ W_{ijk}^\pi &= -\frac{1}{n} \left. \frac{\partial^3 \ln \pi}{\partial \lambda^i \partial \lambda^j \partial \lambda^k} \right|_{\vec{\lambda}_t}. \end{aligned} \quad (14)$$

We can find a perturbative solution as we did for  $\vec{\beta}$

$$\vec{\delta} = \vec{\delta}_1 + \vec{\delta}_{\frac{3}{2}} + \dots$$

$$\begin{aligned} \text{where } \delta_1^i &= \frac{1}{2} (\mathbb{E}[\mathbf{V}])_{ij}^{-1} \mathbb{E}[W_{jkl}] \beta_{\frac{1}{2}}^k \beta_{\frac{1}{2}}^l \\ \delta_{\frac{3}{2}}^i &= (\mathbb{E}[\mathbf{V}])_{ij}^{-1} \left[ \beta_{\frac{1}{2}}^k \beta_1^l \mathbb{E}[W_{jkl}] + \beta_{\frac{1}{2}}^k \delta_1^l \mathbb{E}[W_{jkl}] \right. \\ &+ (V_{jk} - \mathbb{E}[V_{jk}]) \delta_1^k \\ &\left. + \frac{1}{2} (W_{jkl} - \mathbb{E}[W_{jkl}]) \beta_{\frac{1}{2}}^k \beta_{\frac{1}{2}}^l \right] \end{aligned} \quad (15)$$

The means and relevant covariances are

$$\begin{aligned} \mathbb{E}[\delta_1^i] &= \frac{1}{2} (\mathbb{E}[\mathbf{V}])_{ij}^{-1} \mathbb{E}[W_{jkl}] \text{cov}\left(\beta_{\frac{1}{2}}^k, \beta_{\frac{1}{2}}^l\right) \\ \mathbb{E}[\delta_{\frac{3}{2}}^i] &= (\mathbb{E}[\mathbf{V}])_{ij}^{-1} \left[ \mathbb{E}[W_{jkl}] \text{cov}\left(\beta_{\frac{1}{2}}^k, \beta_1^l\right) \right. \\ &+ \mathbb{E}[W_{jkl}] \text{cov}\left(\beta_{\frac{1}{2}}^k, \delta_1^l\right) \\ &+ \frac{1}{2} (\mathbb{E}[\mathbf{V}])_{kn}^{-1} \mathbb{E}[W_{npq}] (\mathbb{E}[\mathbf{V}])_{pl}^{-1} (\mathbb{E}[\mathbf{V}])_{qm}^{-1} \Sigma_{ijklmn}^{VUUU} \\ &\left. + \frac{1}{2} (\mathbb{E}[\mathbf{V}])_{km}^{-1} (\mathbb{E}[\mathbf{V}])_{ln}^{-1} \Sigma_{ijklmn}^{WUUU} \right] \\ \text{cov}(\delta_1^i, \delta_1^j) &= \frac{1}{2} (\mathbb{E}[\mathbf{V}])_{ik}^{-1} (\mathbb{E}[\mathbf{V}])_{jl}^{-1} (\mathbb{E}[\mathbf{V}])_{mx}^{-1} (\mathbb{E}[\mathbf{V}])_{ny}^{-1} \\ &(\mathbb{E}[\mathbf{V}])_{pr}^{-1} (\mathbb{E}[\mathbf{V}])_{qs}^{-1} \mathbb{E}[W_{kmn}] \mathbb{E}[W_{lpq}] \Sigma_{xyrs}^{UUUU} \\ &- \mathbb{E}[\delta_1^i] \mathbb{E}[\delta_1^j] \\ \text{cov}(\delta_1^i, \beta_{\frac{1}{2}}^j) &= \frac{1}{2} (\mathbb{E}[\mathbf{V}])_{ik}^{-1} (\mathbb{E}[\mathbf{V}])_{jp}^{-1} (\mathbb{E}[\mathbf{V}])_{lq}^{-1} (\mathbb{E}[\mathbf{V}])_{mr}^{-1} \\ &\mathbb{E}[W_{klm}] \Sigma_{pqr}^{UUU} \\ \text{cov}(\delta_1^i, \beta_1^j) &= \frac{1}{2} (\mathbb{E}[\mathbf{V}])_{ik}^{-1} (\mathbb{E}[\mathbf{V}])_{lp}^{-1} (\mathbb{E}[\mathbf{V}])_{mq}^{-1} (\mathbb{E}[\mathbf{V}])_{jr}^{-1} \\ &\mathbb{E}[W_{klm}] \left( (\mathbb{E}[\mathbf{V}])_{sx}^{-1} \Sigma_{rsxpq}^{UUUU} + U_r^\pi \Sigma_{pq}^{UU} \right) \\ &- \mathbb{E}[\delta_1^i] \mathbb{E}[\beta_1^j] \end{aligned}$$

$$\begin{aligned} \text{cov}\left(\delta_{\frac{3}{2}}^i, \beta_{\frac{3}{2}}^j\right) &= (\mathbb{E}[\mathbf{V}])_{ik}^{-1} (\mathbb{E}[\mathbf{V}])_{jp}^{-1} \left[ \mathbb{E}[W_{klm}] (\mathbb{E}[\mathbf{V}])_{lq}^{-1} \right. \\ &\left( (\mathbb{E}[\mathbf{V}])_{mx}^{-1} (\mathbb{E}[\mathbf{V}])_{yz}^{-1} \Sigma_{xyzpq}^{UUUU} + U_x^\pi \Sigma_{pq}^{UU} \right) \\ &+ \frac{1}{2} \mathbb{E}[W_{klm}] \mathbb{E}[W_{xyz}] (\mathbb{E}[\mathbf{V}])_{lq}^{-1} (\mathbb{E}[\mathbf{V}])_{mx}^{-1} \\ &(\mathbb{E}[\mathbf{V}])_{yu}^{-1} (\mathbb{E}[\mathbf{V}])_{zv}^{-1} \Sigma_{pquv}^{UUUU} \\ &+ \frac{1}{2} (\mathbb{E}[\mathbf{V}])_{lq}^{-1} (\mathbb{E}[\mathbf{V}])_{xr}^{-1} (\mathbb{E}[\mathbf{V}])_{ys}^{-1} \\ &\mathbb{E}[W_{qxy}] \Sigma_{klrsp}^{UUUU} \\ &\left. + \frac{1}{2} (\mathbb{E}[\mathbf{V}])_{ln}^{-1} (\mathbb{E}[\mathbf{V}])_{mq}^{-1} \Sigma_{klmnpq}^{WUUU} \right] \end{aligned} \quad (16)$$

where

$$\begin{aligned} \Sigma_{ijklm}^{WUU} &= \mathbb{E}[(W_{ijk} - \mathbb{E}[W_{ijk}])U_l U_m] \\ &= \frac{1}{n^2} \int l_{,ijk}(\mathbf{d}) l_{,l}(\mathbf{d}) l_{,m}(\mathbf{d}) p(\mathbf{d}|\vec{\lambda}_t) \mathbf{d}\mathbf{d} \\ &\quad - \mathbb{E}[W_{ijk}] \Sigma_{lm}^{UU} \\ \Sigma_{ijk}^{UUU} &= \mathbb{E}[U_i U_j U_k] \\ &= \frac{1}{n^2} \int l_{,i}(\mathbf{d}) l_{,j}(\mathbf{d}) l_{,k}(\mathbf{d}) p(\mathbf{d}|\vec{\lambda}_t) \mathbf{d}\mathbf{d} \\ \Sigma_{ijkl}^{UUUU} &= \mathbb{E}[U_i U_j U_k U_l] \\ &= \frac{n(n-1)}{n^4} \left[ \int l_{,i}(\mathbf{d}) l_{,j}(\mathbf{d}) p(\mathbf{d}|\vec{\lambda}_t) \mathbf{d}\mathbf{d} \right. \\ &\quad \left. \times \int l_{,k}(\mathbf{d}) l_{,l}(\mathbf{d}) p(\mathbf{d}|\vec{\lambda}_t) \mathbf{d}\mathbf{d} \right] \\ &+ \frac{1}{n^3} \int l_{,i}(\mathbf{d}) l_{,j}(\mathbf{d}) l_{,k}(\mathbf{d}) l_{,l}(\mathbf{d}) p(\mathbf{d}|\vec{\lambda}_t) \mathbf{d}\mathbf{d} \\ \Sigma_{ijklm}^{VUUU} &= \mathbb{E}[(V_{ij} - \mathbb{E}[V_{ij}])U_k U_l U_m] \\ &= \frac{n(n-1)}{n^4} \left[ \int l_{,ij}(\mathbf{d}) l_{,k}(\mathbf{d}) p(\mathbf{d}|\vec{\lambda}_t) \mathbf{d}\mathbf{d} \right. \\ &\quad \times \int l_{,l}(\mathbf{d}) l_{,m}(\mathbf{d}) p(\mathbf{d}|\vec{\lambda}_t) \mathbf{d}\mathbf{d} \\ &\quad \left. + k \leftrightarrow l + k \leftrightarrow m \right] \\ &+ \frac{1}{n^3} \int l_{,ij}(\mathbf{d}) l_{,k}(\mathbf{d}) l_{,l}(\mathbf{d}) l_{,m}(\mathbf{d}) p(\mathbf{d}|\vec{\lambda}_t) \mathbf{d}\mathbf{d} \\ &\quad - \mathbb{E}[V_{ij}] \Sigma_{klm}^{UUU} \\ \Sigma_{ijklmn}^{WUUU} &= \mathbb{E}[(W_{ijk} - \mathbb{E}[W_{ijk}])U_l U_m U_n] \\ &= \frac{n(n-1)}{n^4} \left[ \int l_{,ijk}(\mathbf{d}) l_{,l}(\mathbf{d}) p(\mathbf{d}|\vec{\lambda}_t) \mathbf{d}\mathbf{d} \right. \\ &\quad \times \int l_{,m}(\mathbf{d}) l_{,n}(\mathbf{d}) p(\mathbf{d}|\vec{\lambda}_t) \mathbf{d}\mathbf{d} \\ &\quad \left. + l \leftrightarrow m + l \leftrightarrow n \right] \\ &+ \frac{1}{n^3} \int l_{,ijk}(\mathbf{d}) l_{,l}(\mathbf{d}) l_{,m}(\mathbf{d}) l_{,n}(\mathbf{d}) p(\mathbf{d}|\vec{\lambda}_t) \mathbf{d}\mathbf{d} \\ &\quad - \mathbb{E}[W_{ijk}] \Sigma_{lmn}^{UUU}. \end{aligned} \quad (17)$$

From these we can construct the leading order covariances

$$\begin{aligned} \text{cov}(\beta^i, \delta^j) &= \text{cov}\left(\beta_{\frac{1}{2}}^i, \delta_1^j\right) + \text{cov}\left(\beta_1^i, \delta_1^j\right) + \text{cov}\left(\beta_{\frac{3}{2}}^i, \delta_{\frac{3}{2}}^j\right) \\ \text{cov}(\delta^i, \delta^j) &= \text{cov}\left(\delta_1^i, \delta_1^j\right). \end{aligned} \quad (18)$$

Putting these together we can obtain the mean and covariance of the posterior mode,  $\hat{\lambda}^i - \lambda_t^i = \beta^i + \delta^i$ , expressed as

a deviation from the true population parameters.

$$\begin{aligned} \mathbb{E}[\hat{\lambda}^i - \lambda_t^i] &= \mathbb{E}[\beta^i] + \mathbb{E}[\delta^i] \\ \text{cov}(\hat{\lambda}^i - \lambda_t^i, \hat{\lambda}^j - \lambda_t^j) &= \text{cov}(\beta^i, \beta^j) + \text{cov}(\delta^i, \delta^j) \\ &\quad + \text{cov}(\beta^i, \delta^j) + \text{cov}(\beta^j, \delta^i). \end{aligned} \quad (19)$$

We now turn our attention to the posterior mean and variance. These are averages over the posterior. The definition of  $\vec{\beta}$  was motivated to ensure the leading terms of the log-posterior can be written as a quadratic in  $\vec{\lambda} - \vec{\beta} - \vec{\lambda}_t$ . Denoting  $\vec{B} = \vec{\lambda}_t + \vec{\beta}$ , averages of a function  $f(\vec{\lambda})$  over the posterior then take the form

$$\begin{aligned} \langle f \rangle &= \frac{\int f(\vec{\lambda}) g(\vec{\lambda}) \exp\left[-\frac{n}{2}(\lambda^i - B^i)(V_{ij} + V_{ij}^\pi)(\lambda^j - B^j)\right] d\vec{\lambda}}{\int g(\vec{\lambda}) \exp\left[-\frac{n}{2}(\lambda^i - B^i)(V_{ij} + V_{ij}^\pi)(\lambda^j - B^j)\right] d\vec{\lambda}} \\ \ln g(\vec{\lambda}) &= -\frac{n}{6} W_{ijk} (\lambda^i - \lambda_t^i)(\lambda^j - \lambda_t^j)(\lambda^k - \lambda_t^k) \\ &\quad - \frac{n}{24} X_{ijkl} (\lambda^i - \lambda_t^i)(\lambda^j - \lambda_t^j)(\lambda^k - \lambda_t^k)(\lambda^l - \lambda_t^l) \\ X_{ijkl} &= \left( \frac{\partial^4 \hat{\mu}}{\partial \lambda^i \partial \lambda^j \partial \lambda^k \partial \lambda^l} \right)_{|\vec{\lambda}_t} \end{aligned} \quad (20)$$

There are further corrections in  $g(\vec{\lambda})$  from higher derivatives in the expansion, and from the prior terms,  $W_{ijk}^\pi$ ,  $X_{ijkl}^\pi$  etc. However, the contributions from the included terms can be seen to be  $1/n$  down relative to the leading terms in the integral, and these other corrections are at least  $1/n^{\frac{3}{2}}$  down from leading and are hence sub-dominant.

Integrals of these form are standard and we will make use of the following results

$$I_0(\Gamma) = \int \exp\left[-\frac{1}{2} \mathbf{x}^T \Gamma^{-1} \mathbf{x}\right] d\mathbf{x} = (2\pi)^{\frac{N}{2}} \sqrt{|\Gamma|} \quad (21)$$

$$\begin{aligned} I_{ij}(\Gamma) &= \int x_i x_j \exp\left[-\frac{1}{2} \mathbf{x}^T \Gamma^{-1} \mathbf{x}\right] d\mathbf{x} \\ &= (2\pi)^{\frac{N}{2}} \sqrt{|\Gamma|} \Gamma_{ij} \end{aligned} \quad (22)$$

$$\begin{aligned} I_{ijkl}(\Gamma) &= \int x_i x_j x_k x_l \exp\left[-\frac{1}{2} \mathbf{x}^T \Gamma^{-1} \mathbf{x}\right] d\mathbf{x} \\ &= (2\pi)^{\frac{N}{2}} \sqrt{|\Gamma|} (\Gamma_{ij} \Gamma_{kl} + \Gamma_{ik} \Gamma_{jl} + \Gamma_{il} \Gamma_{jk}) \end{aligned} \quad (23)$$

$$\begin{aligned} I_{ijklmn}(\Gamma) &= \int x_i x_j x_k x_l x_m x_n \exp\left[-\frac{1}{2} \mathbf{x}^T \Gamma^{-1} \mathbf{x}\right] d\mathbf{x} \\ &= (2\pi)^{\frac{N}{2}} \sqrt{|\Gamma|} (\Gamma_{ij} \Gamma_{kl} \Gamma_{mn} + \Gamma_{ij} \Gamma_{km} \Gamma_{ln} \\ &\quad + \Gamma_{ij} \Gamma_{kn} \Gamma_{lm} + \Gamma_{ik} \Gamma_{jl} \Gamma_{mn} + \Gamma_{ik} \Gamma_{jm} \Gamma_{ln} \\ &\quad + \Gamma_{ik} \Gamma_{jn} \Gamma_{lm} + \Gamma_{il} \Gamma_{jk} \Gamma_{mn} + \Gamma_{il} \Gamma_{jm} \Gamma_{kn} \\ &\quad + \Gamma_{il} \Gamma_{jn} \Gamma_{km} + \Gamma_{im} \Gamma_{jk} \Gamma_{ln} + \Gamma_{im} \Gamma_{jl} \Gamma_{kn} \\ &\quad + \Gamma_{im} \Gamma_{jn} \Gamma_{kl} + \Gamma_{in} \Gamma_{jk} \Gamma_{lm} + \Gamma_{in} \Gamma_{jl} \Gamma_{km} \\ &\quad + \Gamma_{in} \Gamma_{jm} \Gamma_{kl}) \end{aligned} \quad (24)$$

where  $|\Gamma|$  denotes the determinant of  $\Gamma$ . We will also use the notation  $\tilde{I}_{ij}(\Gamma) \equiv I_{ij}(\Gamma)/I_0(\Gamma)$  and similarly for other terms. In this case, the covariance matrix  $\Gamma = (\mathbf{V} + \mathbf{V}^\pi)^{-1}/n$ . Every additional factor of  $\Gamma$  therefore introduces an extra negative power of  $n$ . This allows us to identify the dominant terms. To evaluate the above expressions we need to be able to compute  $\Gamma$ , which can be done perturbatively by noting

$$n\Gamma(\mathbb{E}[\mathbf{V}] + (\mathbf{V} - \mathbb{E}[\mathbf{V}]) + \mathbf{V}^\pi) = \mathbf{I} \quad (25)$$

from which

$$\begin{aligned} \Gamma &= \frac{1}{n} (\Gamma_0 + \Gamma_{\frac{1}{2}} + \Gamma_1 + \dots) \\ \Gamma_0 &= (\mathbb{E}[\mathbf{V}])^{-1} \\ \Gamma_{\frac{1}{2}} &= -(\mathbb{E}[\mathbf{V}])^{-1} (\mathbf{V} - \mathbb{E}[\mathbf{V}]) \\ \Gamma_1 &= -(\mathbb{E}[\mathbf{V}])^{-1} \left[ \Gamma_{\frac{1}{2}} (\mathbf{V} - \mathbb{E}[\mathbf{V}]) + \mathbf{V}^\pi \right]. \end{aligned} \quad (26)$$

To obtain the posterior mean, we first compute

$$\begin{aligned} \langle g \rangle &= \int g(\vec{\lambda}) \exp\left[-\frac{n}{2}(\lambda^i - B^i)(V_{ij} + V_{ij}^\pi)(\lambda^j - B^j)\right] d\vec{\lambda} \\ &= I_0(\Gamma) - \frac{n}{6} W_{ijk} \left[ \beta^i I_{jk}(\Gamma) + \beta^j I_{ki}(\Gamma) + \beta^l I_{ij}(\Gamma) \right] \\ &\quad - \frac{n}{6} W_{ijk} \beta^i \beta^j \beta^k I_0(\Gamma) + \frac{n^2}{72} W_{ijk} W_{lmn} I_{ijklmn}(\Gamma) \\ &\quad - \frac{n}{24} X_{ijkl} I_{ijkl}(\Gamma). \end{aligned} \quad (27)$$

Using similar notation to before we can write

$$\begin{aligned} \langle g \rangle &= I_0(\Gamma) \left( 1 + g_{\frac{1}{2}} + g_1 + \dots \right) \\ g_{\frac{1}{2}} &= -\frac{1}{6} \mathbb{E}[W_{ijk}] \left( \beta_{\frac{1}{2}}^i (\Gamma_0)_{jk} + \beta_{\frac{1}{2}}^j (\Gamma_0)_{kl} + \beta_{\frac{1}{2}}^l (\Gamma_0)_{ij} \right) \\ &\quad - \frac{n}{6} \mathbb{E}[W_{ijk}] \beta_{\frac{1}{2}}^i \beta_{\frac{1}{2}}^j \beta_{\frac{1}{2}}^k \\ g_1 &= -\frac{1}{6} (W_{ijk} - \mathbb{E}[W_{ijk}]) \\ &\quad \times \left( \beta_{\frac{1}{2}}^i (\Gamma_0)_{jk} + \beta_{\frac{1}{2}}^j (\Gamma_0)_{kl} + \beta_{\frac{1}{2}}^l (\Gamma_0)_{ij} \right) \\ &\quad - \frac{1}{6} \mathbb{E}[W_{ijk}] \left( \beta_{\frac{1}{2}}^i (\Gamma_{\frac{1}{2}})_{jk} + \beta_{\frac{1}{2}}^j (\Gamma_{\frac{1}{2}})_{kl} + \beta_{\frac{1}{2}}^l (\Gamma_{\frac{1}{2}})_{ij} \right) \\ &\quad + \beta_1^i (\Gamma_0)_{jk} + \beta_1^j (\Gamma_0)_{kl} + \beta_1^l (\Gamma_0)_{ij} \\ &\quad - \frac{n}{6} (W_{ijk} - \mathbb{E}[W_{ijk}]) \beta_{\frac{1}{2}}^i \beta_{\frac{1}{2}}^j \beta_{\frac{1}{2}}^k \\ &\quad - \frac{n}{6} \mathbb{E}[W_{ijk}] \left( \beta_1^i \beta_{\frac{1}{2}}^j \beta_{\frac{1}{2}}^k + \beta_{\frac{1}{2}}^i \beta_1^j \beta_{\frac{1}{2}}^k + \beta_{\frac{1}{2}}^i \beta_{\frac{1}{2}}^j \beta_1^k \right) \\ &\quad + \frac{1}{72} \mathbb{E}[W_{ijk}] \mathbb{E}[W_{lmn}] \frac{1}{n} \tilde{I}_{ijklmn}(\Gamma_0) \\ &\quad - \frac{1}{24} \frac{1}{n} \mathbb{E}[X_{ijkl}] \tilde{I}_{ijkl}(\Gamma_0) \end{aligned} \quad (28)$$

Now we compute the posterior mean, expressed as a distance from the true population parameters

$$\begin{aligned} \bar{\lambda}^i - \lambda_t^i &\equiv \langle (\lambda^i - \lambda_t^i) \rangle = \beta^i + \langle (\lambda^i - B^i) \rangle \\ &= \beta^i + \frac{1}{\langle g \rangle} \left[ -\frac{n}{6} W_{jkl} \left( I_{ijkl}(\Gamma) + 3\beta^j \beta^k I_{il}(\Gamma) \right) \right. \\ &\quad - \frac{n}{6} X_{jklm} \left( \beta^j I_{iklm}(\Gamma) + \beta^i \beta^k \beta^l I_{im}(\Gamma) \right) \\ &\quad + \frac{n^2}{36} W_{jkl} W_{mnp} \left( 3\beta^j I_{iklmnp}(\Gamma) \right. \\ &\quad \left. + \beta^j \beta^k \beta^l I_{imnp}(\Gamma) + 3\beta^j \beta^k \beta^m I_{ilnp}(\Gamma) \right. \\ &\quad \left. + 3\beta^j \beta^k \beta^l \beta^m \beta^n I_{ip}(\Gamma) \right) \left. \right]. \end{aligned} \quad (29)$$

We note that the leading order correction in the bracketed term is  $1/n$ . To obtain the posterior mean to the same order as  $\beta$  we therefore only need to retain terms up to  $g_{\frac{1}{2}}$  in  $\langle g \rangle$ .

Specifically we can write

$$\begin{aligned}
\bar{\lambda}^i - \lambda_t^i &= \bar{\lambda}_{\frac{1}{2}}^i + \bar{\lambda}_1^i + \bar{\lambda}_{\frac{3}{2}}^i + \dots \\
\bar{\lambda}_{\frac{1}{2}}^i &= \beta_{\frac{1}{2}}^i \\
\bar{\lambda}_1^i &= \beta_1^i - \frac{1}{6}\mathbb{E}[W_{jkl}] \left( \frac{1}{n}\tilde{I}_{ijkl}(\Gamma_0) + 3\beta_{\frac{1}{2}}^j\beta_{\frac{1}{2}}^k(\Gamma_0)_{il} \right) \\
\bar{\lambda}_{\frac{3}{2}}^i &= \beta_{\frac{3}{2}}^i + \left( \frac{g_{\frac{1}{2}}}{6}\mathbb{E}[W_{jkl}] - \frac{1}{6}(W_{jkl} - \mathbb{E}[W_{jkl}]) \right) \\
&\quad \times \left( \frac{1}{n}\tilde{I}_{ijkl}(\Gamma_0) + 3\beta_{\frac{1}{2}}^j\beta_{\frac{1}{2}}^k(\Gamma_0)_{il} \right) \\
&\quad - \frac{1}{6}\mathbb{E}[W_{jkl}] \left( \frac{2}{n} \left( (\Gamma_0)_{ij}(\Gamma_{\frac{1}{2}})_{kl} + (\Gamma_0)_{ik}(\Gamma_{\frac{1}{2}})_{jl} \right. \right. \\
&\quad \quad \left. \left. + (\Gamma_0)_{il}(\Gamma_{\frac{1}{2}})_{jk} \right) \right. \\
&\quad \left. + 3(\beta_1^j\beta_{\frac{1}{2}}^k + \beta_{\frac{1}{2}}^j\beta_1^k)(\Gamma_0)_{il} \right. \\
&\quad \left. + 3\beta_{\frac{1}{2}}^j\beta_{\frac{1}{2}}^k(\Gamma_{\frac{1}{2}})_{il} \right) \\
&\quad - \frac{1}{6}\mathbb{E}[X_{jklm}] \left( \beta_{\frac{1}{2}}^j\frac{1}{n}\tilde{I}_{iklm} + \beta_{\frac{1}{2}}^j\beta_{\frac{1}{2}}^k\beta_{\frac{1}{2}}^l(\Gamma_0)_{im} \right) \\
&\quad + \frac{1}{36}\mathbb{E}[W_{jkl}]\mathbb{E}[W_{mnp}] \left( \frac{3}{n}\beta_{\frac{1}{2}}^j\tilde{I}_{iklmnp}(\Gamma_0) \right. \\
&\quad \left. + \beta_{\frac{1}{2}}^j\beta_{\frac{1}{2}}^k\beta_{\frac{1}{2}}^l\tilde{I}_{imnp}(\Gamma_0) + 3\beta_{\frac{1}{2}}^j\beta_{\frac{1}{2}}^k\beta_{\frac{1}{2}}^m\tilde{I}_{ilnp}(\Gamma_0) \right. \\
&\quad \left. + 3n\beta_{\frac{1}{2}}^j\beta_{\frac{1}{2}}^k\beta_{\frac{1}{2}}^l\beta_{\frac{1}{2}}^m\beta_{\frac{1}{2}}^n(\Gamma_0)_{ip} \right) \quad (30)
\end{aligned}$$

Finally, we consider the posterior covariance

$$\begin{aligned}
\hat{\Gamma}_{ij} &\equiv \langle (\lambda^i - \bar{\lambda}^i)(\lambda^j - \bar{\lambda}^j) \rangle \\
&= \langle (\lambda^i - B^i)(\lambda^j - B^j) \rangle \\
&\quad + (B^i - \bar{\lambda}^i)\langle (\lambda^j - B^j) \rangle + i \leftrightarrow j \\
&\quad + (B^i - \bar{\lambda}^i)(B^j - \bar{\lambda}^j). \quad (31)
\end{aligned}$$

The term on the first line has a leading order dependence of  $1/n$ , plus corrections of  $1/n^{\frac{3}{2}}$ . The terms on the second and third lines are  $O(1/n^2)$  and so are sub-dominant. We deduce

$$\hat{\Gamma}_{ij} = \frac{1}{\langle g \rangle} \left[ I_{ij}(\Gamma) - \frac{n}{6}W_{klm} \left( 3\beta^k I_{ijlm}(\Gamma) + \beta^k\beta^l\beta^m I_{ij}(\Gamma) \right) \right] \quad (32)$$

and expand

$$\begin{aligned}
\hat{\Gamma}_{ij} &= (\hat{\Gamma}_1)_{ij} + \left( \hat{\Gamma}_{\frac{3}{2}} \right)_{ij} + \dots \\
(\hat{\Gamma}_1)_{ij} &= \frac{1}{n}(\Gamma_0)_{ij} \\
\left( \hat{\Gamma}_{\frac{3}{2}} \right)_{ij} &= \frac{1}{n} \left( \Gamma_{\frac{1}{2}} \right)_{ij} - \frac{g_{\frac{1}{2}}}{n}(\Gamma_0)_{ij} \\
&\quad - \frac{1}{6}\mathbb{E}[W_{klm}] \left( \frac{3}{n}\beta_{\frac{1}{2}}^k\tilde{I}_{ijlm}(\Gamma_0) + \beta_{\frac{1}{2}}^k\beta_{\frac{1}{2}}^l\beta_{\frac{1}{2}}^m(\Gamma_0)_{ij} \right) \quad (33)
\end{aligned}$$

Using the preceding expressions, we could now compute the mean and variance of the posterior mean and covariance as we did for the shift in the posterior mode. However, this calculation is very similar to the calculations carried out above and is tedious so we leave it out. Instead we note a number of features.

- The leading order posterior covariance is constant and equal to  $(\mathbb{E}[\mathbf{V}])^{-1}/n$ , which is the expression we used to derive the population Fisher matrix in the body of the paper.

- The leading order difference between either the posterior mode or mean and the true parameter value has expectation value that scales like  $1/n$  and variance that also scales like  $1/n$ . Thus, the posterior bias is noise dominated, i.e., fluctuations due to the particular random realisation of the population that was observed dominate over the fixed bias. If multiple sets of observations of  $n$  events were repeated and averaged, then this bias would eventually be significant. In practice we would never do this since it weakens the precision of inference. This means that computing corrections to the posterior mean is unnecessary.

- The leading order correction to the expected value of the posterior covariance scales like  $n^{-\frac{3}{2}}$ , while the variance in the posterior covariance scales like  $n^{-3}$ . So, fluctuations in the posterior covariance due to randomness in the observed population are comparable to the size of corrections from the finite number of observations.