Enhanced EDMs from Small Instantons

Ravneet S. Bedi^a, Tony Gherghetta^a and Maxim Pospelov^{a,b}

 ^a School of Physics and Astronomy, University of Minnesota, Minneapolis, Minnesota 55455, USA
 ^b William I. Fine Theoretical Physics Institute, University of Minnesota, Minneapolis, Minnesota 55455 USA

Abstract

We show that models in which the strong CP problem is solved by introducing an axion field with a mass enhanced by non-QCD UV dynamics at a scale Λ_{SI} , exhibit enhanced sensitivity to external sources of CP-violation. In the presence of higher-dimensional CP-odd sources at a scale Λ_{CP} , the same mechanisms that enhance the axion mass, also modify the axion potential, shifting the potential minimum by a factor $\propto \Lambda_{SI}^2/\Lambda_{CP}^2$. This phenomenon of CP-violation enhancement, which puts stringent constraints on the scale of new physics, is explicitly demonstrated within a broad class of "small instanton" models with CP-odd sources arising from the dimension-six Weinberg gluonic and four-fermion operators. We find that for heavy axion masses $\gtrsim 100 \text{ MeV}$, arising from new dynamics at $\Lambda_{SI} \lesssim 10^{10} \text{ GeV}$, CP-violation generated up to the Planck scale can be probed by future EDM experiments.

1 Introduction

The Standard Model (SM) of particle physics has two sources of CP violation. The well established and measured source of CP-violation in the quark mixing sector, the Kobayashi-Maskawa phase [1], is responsible for a multitude of CP violating phenomena observed in the quark flavorchanging transitions. At the same time, this phase induces electric dipole moments (EDMs) of neutrons and heavy atoms well below current experimental limits. The other source of CP violation, the non-perturbative parameter θ of quantum chromodynamics (QCD), is largely irrelevant for flavor physics, but tends to induce large EDMs. The non-observation of EDMs that imply the smallness of theta, $|\theta| \leq 10^{-10}$ [2, 3], contrasted with the naive expectation of $\theta \sim O(1)$, poses a naturalness problem for the Standard Model, the strong CP problem.

There are two generic approaches to resolve the strong CP problem. The first approach involves promoting the θ parameter to a new dynamical field, the QCD axion [4–10], which symbolically can be represented as

$$\frac{\theta}{32\pi^2}G^c_{\mu\nu}\widetilde{G}^{c\mu\nu} \rightarrow \frac{1}{2}(\partial_\mu a)^2 + \frac{a}{32\pi^2 f_a}G^c_{\mu\nu}\widetilde{G}^{c\mu\nu},\tag{1}$$

where $G^c_{\mu\nu}$ is the gluon field strength, $\tilde{G}^{c\mu\nu} \equiv \frac{1}{2} \varepsilon^{\mu\nu\rho\sigma} G^c_{\rho\sigma}$ with *c* the adjoint index and f_a is the decay constant of the axion field *a*. The QCD vacuum energy, which for small θ can be parametrically expressed as

$$E(\theta) \propto \theta^2 m_q \Lambda_{\rm QCD}^3 \rightarrow V(a) = \frac{1}{2} m_a^2 a^2,$$
 (2)

can be made to dynamically relax to the minimum of the potential V(a). In this expression, Λ_{QCD} is the non-perturbative scale of the strong interactions, and m_q is the light quark mass. As a result, any initial value of $\theta = a/f_a$ will relax to the minimum of the axion potential. In the absence of additional sources of CP violation, this minimum is exactly at $\theta = 0$, as in Eq.(2). Therefore, the neutron EDM that scales as

$$d_n \propto \frac{m_q \theta}{\Lambda_{\rm QCD}^2} ,$$
 (3)

is also relaxed to zero.

Consider now additional sources of CP violation placed at some new physics scale Λ_{CP} that we will assume to be larger than the electroweak scale (for example, this could be due to supersymmetric theories with large CP-violating phases). Integrating out the new physics at this scale will, in general, result in a number of generic consequences:

- 1. The theta parameter may receive additive corrections to its value, $\theta \to \theta + \theta_{rad}$. Since GG is a dimension four operator, θ_{rad} can depend only on the ratio of scales, and therefore has $\Lambda^0_{\rm CP}$ scaling. Potentially, this can be a large correction, but the axion mechanism will remove the theta term together with θ_{rad} .
- 2. *CP*-violating new physics will generically induce higher-dimensional *CP*-odd operators, of which the most relevant are dimension six operators, \mathcal{O}_6 that are suppressed by the square of the new physics scale, and the resulting EDMs will have scaling $d_n(\mathcal{O}_6) \propto \Lambda_{\rm QCD}/\Lambda_{\rm CP}^2$ (or $m_q/\Lambda_{\rm CP}^2$, depending on the chiral properties of \mathcal{O}_6).

3. In the presence of higher-dimensional *CP*-odd new physics operators, the axion potential minimum shifts away from zero inducing a low-energy value of theta, $\theta_{\rm ind} \propto \Lambda_{\rm QCD}^2/\Lambda_{\rm CP}^2$. This leads to an additional θ -induced contribution to d_n that has, for example, a comparable $m_q/\Lambda_{\rm CP}^2$ scaling [11–13].

An important conclusion can be drawn from these observations: the QCD axion mechanism ensures that for sufficiently large $\Lambda_{\rm CP}$, the observable EDMs can be made small and indeed within current bounds for $\Lambda_{\rm CP} \gtrsim 100 \,\text{TeV}$, one can allow for an arbitrarily large amount of (strong) *CP* violation above these scales. In this sense, the axion mechanism allows for a proper decoupling of new physics contributions to EDMs.

The second class of models do not introduce an axion, and instead appeal to symmetry arguments that help to argue why θ is zero or small. Historically, models with an exact CPsymmetry or exact parity that is spontaneously broken at some UV scale, have been argued to give a viable solution to the strong CP problem (see Refs. [14–21] for a representative set of ideas). Models based on mirror symmetries have also been used to implement this approach [22, 23]. The most important feature of these models is the absence of a dynamical axion and the sensitivity of EDM observables to the value of θ generated at a UV scale. For example, the spontaneous breaking of CP symmetry may also result in complex quark Yukawa couplings that feed into θ_{rad} (a representative set of calculations can be found in Refs. [24–28]). Since the θ term has Λ_{CP}^0 scaling, this non-decoupling means that all possible sources of CP breaking have to be "controlled" to very high scales.

Recently, there has been renewed interest in models that solve the strong CP problem which occupy an intermediate niche between the QCD axion solution and solutions based on discrete symmetries. In this class of models there is still a dynamical axion field and Peccei-Quinn symmetry at a high scale, but the axion mass is now enhanced compared to (2) by additional dynamical mechanisms at the small-instanton scale Λ_{SI} . By small instantons we refer to instantons whose size $1/\Lambda_{\rm SI}$ is smaller than the inverse electroweak scale. For example, extending the strong gauge interactions and the corresponding axion to a larger group where the non-QCD partners confine at a much larger scale $\Lambda'_{\rm QCD}$ (identified with $\Lambda_{\rm SI}$) can lead to a significant parametric increase in the axion mass provided $\Lambda'_{\rm QCD} \gg \Lambda_{\rm QCD}$ [29–33]. Similarly, an axion "portal" between QCD and a mirror QCD with the alignment of θ and θ' can also result in a heavier axion for $\Lambda'_{\rm QCD} \gg \Lambda_{\rm QCD}$ [23, 34]. Alternatively, if the QCD coupling running is modified to become strong above the TeV scale, the QCD axion mass would receive new contributions from "small"-size instantons [35–39]. This naturally occurs in models where at some UV scale, QCD propagates in five dimensions [40, 41]. These models which significantly enhance the axion mass compared to the minimal QCD axion models have a distinctively different phenomenology. Indeed, given the conventional axion mass range $10^{-6} - 10^{-3}$ eV, the enhancement mechanisms imply heavy axions could be in the 100 MeV range or above. These heavier axions avoid most of the astrophysical bounds, and makes the axion amenable to searches at beam dump and collider experiments [34, 38]. Moreover, such heavy axions will be less susceptible to possible distortions of the axion potential by the imperfections of the Peccei-Quinn global symmetry.

Besides the enhanced axion mass it is therefore also interesting to consider whether EDM observables could be enhanced in these models. In this paper we investigate heavy axion models in the presence of additional sources of CP-violation, which are parametrized as higher-dimensional operators that arise from SM fields. The central question we would like to address is whether there is a similar decoupling as for the standard QCD axion, where all observables from, for



Figure 1: Schematic diagram of the different scales referred to in the text. The scale of CP-violation, $\Lambda_{\rm CP}$ due to dimension six operators is a UV scale near the Planck scale, M_P , and $\Lambda_{\rm SI}$ is the small-instanton scale (assumed to be above the electroweak scale, v and QCD strong coupling scale, $\Lambda_{\rm QCD}$) where new dynamics enhances the axion mass. The PQ symmetry breaking scale f_a is assumed to be an independent parameter that can either be above (as shown in the figure) or below the scale $\Lambda_{\rm SI}$.

example, dimension six operators, scale as $\Lambda^2_{\text{QCD}}/\Lambda^2_{\text{CP}}$, or if there is an enhancement of CP violation mediated by the induced θ which is similar to models attempting to solve the strong CP problem using exact parity or CP symmetries.

To answer this question we compute the topological susceptibility and mixed correlators in heavy axion models that arise from two sources of CP violation: the dimension-six Weinberg gluonic operator and a CP-odd four-fermion operator. Such CP-odd operators induce a linear term in θ (or equivalently a) in the axion potential leading to a shift θ_{ind} in the potential minimum. While our computation employs a simple, non-interacting instanton (or anti-instanton) background that ignores strong coupling effects, we are able to extract qualitative results which show that the induced theta, $\theta_{ind} \propto \Lambda_{SI}^2/\Lambda_{CP}^2$.

This induced shift is qualitatively different from the usual QCD axion scenario and solutions based on exact discrete symmetries due to the presence of the new scale Λ_{SI} . While there is still decoupling in the $\Lambda_{CP} \rightarrow \infty$ limit, our results show that the induced θ can enhance the magnitude of observable EDMs, even to the point that if $\Lambda_{SI}^2/\Lambda_{CP}^2$ is too large, the strong CPproblem will re-appear. Thus, models with a dynamically enhanced axion mass are subject to bounds depending on the amount of CP violation that is present at energy scales that may significantly exceed 100 TeV. Interestingly, the enhanced EDMs are potentially observable in future EDM experiments.

This paper is organized as follows: in Section 2 we investigate vacuum correlators in an instanton (or anti-instanton) background with different sources of CP violation that shift the axion potential minimum. In section 3, we consider different heavy QCD axion models with small instantons, deriving the resulting size of the induced θ and subsequent constraints on the CP-violating scale, Λ_{CP} . We reach our conclusions in Section 4.

2 Instanton Correlation Functions

We begin with briefly reviewing QCD dynamics and the instanton solution that will be used to compute various instanton correlation functions. The pure Yang-Mills part of the QCD Lagrangian is given by

$$\mathcal{L}_{\text{QCD}} = -\frac{1}{4g^2} G^a_{\mu\nu} G^{a\mu\nu} + \frac{\theta}{32\pi^2} G^a_{\mu\nu} \widetilde{G}^{a\mu\nu} , \qquad (4)$$

where g is the QCD gauge coupling, θ is the QCD vacuum angle and $a = 1, \ldots, 8$ labels the gauge adjoint representation. The BPST instanton solution [42] is given by

$$A^{a}_{\mu}(x) = \frac{2\eta^{a}_{\mu\nu}(x-x_{0})_{\nu}}{(x-x_{0})^{2}+\rho^{2}},$$
(5)

where the instanton is located at x_0 and has a size ρ . The $\eta^a_{\mu\nu}$ denote the group-theoretic 't Hooft η -symbols [43]. The topological charge is defined to be

$$Q = \frac{1}{32\pi^2} \int d^4x \, G^a_{\mu\nu} \widetilde{G}^{a\mu\nu} \,, \tag{6}$$

where Q = 1 for the one instanton solution (5). We will next compute correlation functions in the instanton (or anti-instanton) background (5) that will be useful in obtaining contributions to EDM observables such as the neutron EDM.

2.1 Topological susceptibility

The vacuum-to-vacuum amplitude in QCD can be written as:

$$\langle 0|0\rangle = \sum_{Q} \int \mathcal{D}A_{\mu}^{(Q)} e^{-S_{E}} , \qquad (7)$$

where the Euclidean action for (4) in an instanton background of charge Q [44] is given by

$$S_E = \frac{8\pi^2}{g^2} |Q| + iQ\theta .$$
(8)

The topological susceptibility is then introduced as [8, 11, 45]

$$\chi(0) = -i \lim_{k \to 0} \int d^4 x \, e^{ikx} \left\langle 0 \left| T \left\{ \frac{1}{32\pi^2} G \widetilde{G}(x), \frac{1}{32\pi^2} G \widetilde{G}(0) \right\} \right| 0 \right\rangle , \tag{9}$$

where $G\widetilde{G}$ is shorthand notation for $G^a_{\mu\nu}\widetilde{G}^{a\mu\nu}$.

Since the amplitude in the |Q| > 1 instanton background becomes more exponentially suppressed, only the $Q = \pm 1$ configurations dominate the path integral. Henceforth, we refer to S_E in (8) only for |Q| = 1. In the instanton background (5) we then obtain the two-point correlator

$$\left\langle 0 \left| T \left\{ G \widetilde{G}(x), G \widetilde{G}(0) \right\} \right| 0 \right\rangle_{Q=+1}$$

= $\int \mathcal{D}A_{\mu} \ G \widetilde{G}(x) G \widetilde{G}(0) \ e^{-\frac{8\pi^2}{g_0^2}},$ (10)

$$= \int d^4x_0 \frac{d\rho}{\rho^5} C[N] \left(\frac{8\pi^2}{g^2(1/\rho)}\right)^{2N} e^{-\frac{8\pi^2}{g^2(1/\rho)}} \frac{192\rho^4}{((x-x_0)^2+\rho^2)^4} \frac{192\rho^4}{(x_0^2+\rho^2)^4} , \qquad (11)$$

where the running coupling $g(1/\rho)$ encodes corrections from the quantum fluctuations. In (10) we have replaced the path integral over the fluctuation A_{μ} with an integration in (11) over the collective coordinates (see Ref. [43]) where, assuming an SU(N) gauge group,¹ the coefficient

$$C[N] = \frac{C_1 e^{-C_2 N}}{(N-1)!(N-2)!} , \qquad (12)$$

and C_1, C_2 are order one constants ($C_1 = 0.466, C_2 = 1.679$ using Pauli-Villars regularization [47]). The gauge coupling running is given by

$$\frac{8\pi^2}{g^2(1/\rho)} = \frac{8\pi^2}{g_0^2} - b_0 \log(M_{UV}\rho) , \qquad (13)$$

where $b_0 = 4N - N/3 = 11N/3$ is the pure SU(N) Yang-Mills β -function coefficient and $g_0 = g(M_{UV})$ with UV cutoff M_{UV} .

In principle, we could consider an ensemble of instantons and anti-instantons [48–50] to compute correlation functions. However, the qualitative aspects of such an ensemble can be simply captured by one instanton and one anti-instanton [51, 52], where the (anti-)instantons are assumed to be non-interacting with each other. Thus, we will compute correlation functions by adding the contribution from an instanton background to that in an anti-instanton background. The total contribution to the topological susceptibility, obtained by performing the x integration first that arises from (9), followed by the x_0 integration in (11), is then given by

$$\chi(0) = -2i \int \frac{d\rho}{\rho^5} C[N] \left(\frac{8\pi^2}{g^2(1/\rho)}\right)^{2N} e^{-\frac{8\pi^2}{g^2(1/\rho)}} .$$
(14)

Assuming an asymptotically free theory, the integral in (14) is divergent for large instantons but can be evaluated with a IR cutoff ρ_{IR} on the instanton size. Assuming N = 3 with $\rho_{IR} = 1/\Lambda_{\rm QCD}$ we obtain $\chi(0) \propto \Lambda_{\rm QCD}^4$.

2.1.1 Fermion contributions

The introduction of fermions modifies the path integral and the collective coordinate integration. In the massless fermion limit, the pure vacuum-to-vacuum transition amplitude is zero. Instead, the instanton now causes transitions from left-handed to right-handed fermions violating the U(1) chiral symmetry so that, for example, $\langle 0|\bar{\psi}_{Ri}\psi_{Li}|0\rangle \neq 0$. Thus, instantons only contribute to correlation functions in which each fermion flavor and chirality appears at least once.

The effect of massless fermions is usually formulated as an "effective" Lagrangian [43, 47]

$$\mathcal{L}_f = \int d^4 x_0 \, \frac{d\rho}{\rho^5} \, C[N] e^{0.292N_f} \left(\frac{8\pi^2}{g^2(1/\rho)}\right)^{2N} e^{-\frac{8\pi^2}{g^2(1/\rho)} - i\theta} \rho^{3N_f} \det\left[\bar{\psi}_R(x_0)\psi_L(x_0)\right] + h.c. \,, \quad (15)$$

where the determinant is taken over the N_f fermion flavors, and $\psi^{\alpha}_{L,R}(x_0)$ are the fermion zero modes. The constant $e^{0.292N_f}$ assumes Pauli-Villars regularization and the gauge coupling running (13) now includes the fermion contributions $b_0 \rightarrow b_0 - 2/3N_f$.

¹In principle, we should also include the normalized Haar measure of the group, as computed in [39, 46]. We will omit this measure since its value is simply one for (11) (or an $\mathcal{O}(1)$ number in more generic cases), and therefore our qualitative results remain unchanged.

Note that because of the explicit appearance of the fermion zero modes $\psi_{L,R}(x_0)$ in (15), there is only a contribution to the axion potential if the external fermion zero mode legs are closed. There are two ways this can occur. The first way is to assume that the fermions have an explicit mass m_f (corresponding to a nonzero Higgs vacuum expectation value (VEV), $v \approx 246$ GeV) that connects left and right-handed fermion fields. The determinant in the effective action then gives a contribution $\propto (\rho m_f)^{N_f}$ for N_f fermion flavors. This is the case for the usual contributions from "large" instantons with $\rho \sim \rho_{IR} = 1/\Lambda_{\rm QCD}$ and $m_f \lesssim \Lambda_{\rm QCD}$. However, since we are interested in "small" instantons corresponding to instanton sizes ($\sim 1/\Lambda_{\rm SI}$) much smaller than the inverse of the electroweak scale, a second possibility is to close the external fermion zero-mode legs in (15) with $N_f/2$ Higgs bosons. This contribution will be proportional to the product of Yukawa couplings (times a loop factor) and is larger than the Higgs VEV contribution that now scales as $\sim (m_f/\Lambda_{\rm SI})^{N_f}$ (assuming $\Lambda_{\rm SI} \gg v$). Instead of proceeding with the 't Hooft determinant operator in the effective Lagrangian (15) we will follow the approach taken in Ref. [37, 39] and directly compute the vacuum-to-vacuum amplitude by including the Higgs-fermion Yukawa interaction in the path integral.

Consider a Higgs field H which couples to N_f flavors of massless fermions with the following Euclidean action

$$S_H = S_H^{(0)} - i \int d^4x \sum_{i=1}^{N_f} \frac{y_i}{\sqrt{2}} H(x) \bar{\psi}_i(x) \psi_i(x) , \qquad (16)$$

where $S_H^{(0)}$ is the quadratic (free) part of the Higgs action and y_i are the Yukawa couplings. The Yukawa couplings, or equivalently the fermion masses, have been redefined to be real with their phase included in $\bar{\theta} = \theta + \text{Arg Det} M_q$, where M_q is the quark mass matrix. The vacuum-to-vacuum amplitude now takes the form

$$\langle 0|0\rangle_{\Delta Q=1} = \int d^4 x_0 \frac{d\rho}{\rho^5} C[N] \left(\frac{8\pi^2}{g^2(1/\rho)}\right)^{2N} e^{-S_E} \times \int \mathcal{D}\mathcal{H} e^{-S_H^{(0)}} \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{-S_{\psi}^{(0)} + i\int d^4 x \sum_{i=1}^{N_f} \frac{y_i}{\sqrt{2}} H(x)\bar{\psi}_i(x)\psi_i(x)} = \int d^4 x_0 \frac{d\rho}{\rho^5} C[N] e^{0.292N_f} \left(\frac{8\pi^2}{g^2(1/\rho)}\right)^{2N} e^{-S_E} (N_f - 1)!! \left(\prod_{i=1}^{N_f} \frac{y_i\rho}{\sqrt{2}}\right) \mathcal{I}^{N_f/2}(17)$$

where the action S_E is defined in (8) with $\theta \to \overline{\theta}$. The first line in (17) shows the collective coordinate integration arising from the gauge field part of the path integral and the second line contains the Higgs and massless fermion contributions to the path integral with $S_{\psi}^{(0)}$ the quadratic (free) part of the fermion action. Integrating over the fermionic fields introduces the factor $e^{0.292N_f}$ and the running gauge coupling now contains fermionic contributions via $b_0 \to b_0 - 2/3N_f$. Finally, the path integral over the Higgs field gives a nonzero contribution to the amplitude provided all Higgs fields are contracted where $(N_f - 1)!!$ is the number of Higgs contractions and the quantity \mathcal{I} is given by [37, 39]

$$\begin{aligned} \mathcal{I} &= -\int d^4 x_1 \int d^4 x_2 \, \bar{\psi}_i^{(0)}(x_1) \psi_i^{(0)}(x_1) \bar{\psi}_j^{(0)}(x_2) \psi_j^{(0)}(x_2) \Delta_H(x_1 - x_2) , \\ &= \frac{\rho^4}{4\pi^8} \int d^4 x_1 \int d^4 x_2 \int d^4 k \, \frac{1}{k^2 + m_H^2} \frac{e^{-ik(x_1 - x_0)}}{((x_1 - x_0)^2 + \rho^2)^3} \frac{e^{ik(x_2 - x_0)}}{((x_2 - x_0)^2 + \rho^2)^3} , \end{aligned}$$

$$\approx \begin{cases} \frac{1}{12\pi^{2}\rho^{2}} & m_{H}\rho \ll 1 ,\\ \frac{1}{5\pi^{2}m_{H}^{2}\rho^{4}} & m_{H}\rho \gg 1 . \end{cases}$$
(18)

In the second line of (18) we have substituted for the scalar Feynman propagator $\Delta_H(x_1 - x_2)$ and the fermions have been replaced with their respective zero mode expressions given in [47]. Note that for an instanton background we have two zero modes $\bar{\psi}_{i,L}^{(0)}, \psi_{j,R}^{(0)}$ (and $\bar{\psi}_{i,R}^{(0)}, \psi_{j,L}^{(0)}$ in an anti-instanton background) where the subscripts L, R, which are suppressed hereon, denote leftand right- handed fields, respectively. Thus, combining (18) and (17) gives the final expression (assuming $m_H \rho \ll 1$)

$$\langle 0|0\rangle_{\Delta Q=1} = \int d^4 x_0 \, \frac{d\rho}{\rho^5} \, C_f[N] \left(\frac{8\pi^2}{g^2(1/\rho)}\right)^{2N} e^{-S_E} \,, \tag{19}$$

with S_E defined in (8) (assuming $\theta \to \overline{\theta}$), and

$$C_f[N] \equiv (N_f - 1)!! \left(\frac{2}{3}\right)^{N_f/2} \left(\prod_{i=1}^{N_f} \frac{y_i}{4\pi}\right) e^{0.292N_f} C[N] .$$
(20)

The expression (19) shows how the instanton density in the vacuum-to-vacuum amplitude is modified in the presence of massless fermions and a Higgs-fermion Yukawa interaction. As expected, the amplitude vanishes if any Yukawa coupling is zero. Thus, the topological susceptibility (14) in the presence of massless fermions is obtained by the substitutions $C[N] \rightarrow C_f[N]$, $\theta \rightarrow \bar{\theta}$ and $b_0 \rightarrow b_0 - 2/3N_f$.

In the case of "large" instantons associated with the scale $1/\Lambda_{\rm QCD}$, the expression for the vacuum-to-vacuum amplitude differs from (19). As already mentioned, each light fermion ($m_f \leq \Lambda_{\rm QCD}$) introduces a $e^{0.292}\rho m_f$ factor ². This can be seen via the first line in (18) where Δ_H can be replaced by v^2 , which just gives $\mathcal{I} = v^2$, and hence:

$$\langle 0|0\rangle_{\Delta Q=1} = \int d^4 x_0 \, \frac{d\rho}{\rho^5} \left(\prod_{i=1}^{N_L} \rho \, m_i\right) \, e^{0.292N_L} C[N] \left(\frac{8\pi^2}{g^2(1/\rho)}\right)^{2N} e^{-S_E} \,, \tag{21}$$

where the product runs only over N_L light fermions and $m_i = y_i v / \sqrt{2}$.

2.2 Weinberg gluonic operator

The Weinberg operator is a purely gluonic, CP odd, dimension six term given by $\mathcal{O}_W = GG\widetilde{G}$ [54] that leads to the Lagrangian term

$$\mathcal{L} \supset \frac{1}{\Lambda_W^2} G G \widetilde{G} , \qquad (22)$$

²In QCD, $\chi(0) \propto m_f$, whereas the $\chi(0)$ resulting from (21) $\propto m_f^{N_L}$. The difference can be understood in terms of instanton-(anti-)instanton interactions- either via mixing between the fermion zero modes of the instanton with those of the anti-instanton [53], or using 't Hooft vertices with fermion legs joined between an instanton and anti-instanton [49].

where Λ_W is an effective UV scale. The operator (22) can induce a shift in the axion potential minimum, which can be computed by considering the mixed correlator [13, 55]

$$\chi_W(0) = -i \lim_{k \to 0} \int d^4 x \, e^{ikx} \left\langle 0 \left| T \left\{ \frac{1}{32\pi^2} G\widetilde{G}(x), \frac{1}{\Lambda_W^2} G G\widetilde{G}(0) \right\} \right| 0 \right\rangle \,. \tag{23}$$

In the instanton background (5) we obtain

$$\mathcal{O}_W = f_{abc} G^a_{\mu\kappa} G^b_{\kappa\nu} \widetilde{G}^{c\nu\mu}(x) = -\frac{1536\rho^6}{((x-x_0)^2 + \rho^2)^6} , \qquad (24)$$

where f_{abc} are the structure constants. Note that for an SU(N) gauge group, the SU(2) instanton solution is embedded in the top left corner of the $N \times N$ matrix of SU(N) generators. Thus, the sum in (24) only gives nonzero contributions for a, b, c = 1, 2, 3. Furthermore,

$$\left\langle 0 \left| T \left\{ G \widetilde{G}(x), G G \widetilde{G}(0) \right\} \right| 0 \right\rangle_{Q=+1}$$

$$= \int \mathcal{D}A_{\mu} \ G \widetilde{G}(x) G G \widetilde{G}(0) \ e^{-\frac{8\pi^2}{g_0^2}} ,$$

$$= \int d^4 x_0 \frac{d\rho}{\rho^5} C[N] \left(\frac{8\pi^2}{g^2(1/\rho)} \right)^{2N} e^{-\frac{8\pi^2}{g^2(1/\rho)}} \frac{192\rho^4}{((x-x_0)^2+\rho^2)^4} \frac{-1536\rho^6}{(x_0^2+\rho^2)^6} .$$

$$(25)$$

Again performing the integrals first over x and then x_0 gives

$$\chi_W(0) = 2i \frac{384\pi^2}{5\Lambda_W^2} \int \frac{d\rho}{\rho^7} C[N] \left(\frac{8\pi^2}{g^2(1/\rho)}\right)^{2N} e^{-\frac{8\pi^2}{g^2(1/\rho)}} , \qquad (26)$$

where we have also included the anti-instanton contribution.

In the presence of fermions, $\chi_W(0)$ is obtained by making the substitutions $C[N] \to C_f[N]$ for small instantons (or by introducing the factor $(\rho m_f)^{N_L}$, as in (21) for large instantons), $\theta \to \bar{\theta}$ and $b_0 \to b_0 - 2/3N_f$ in the running gauge coupling $g(1/\rho)$.

2.3 Four-fermion operators

Another class of dimension six operators which can affect the axion solution are the four-fermion operators. Such operators are suppressed by an effective mass scale Λ_F and given by

$$\mathcal{L} \supset \sum_{ijkl} \frac{\lambda_{ijkl}}{\Lambda_F^2} \bar{\psi}_i \psi_j \bar{\psi}_k \psi_l , \qquad (27)$$

where λ_{ijkl} are complex coefficients with flavor indices i, j, k, l. Note that the spinor and electroweak structure has been suppressed in (27), although it is straightforward to incorporate these details. Of particular interest is the spinor structure of (27) resulting in CP violation. These are operators of the type $\mathcal{O}_{F,ijkl} = \bar{\psi}_i i \gamma_5 \psi_j \bar{\psi}_k \psi_l$ which are anti-Hermitian with the corresponding λ_{ijkl} purely imaginary.

The CP-violating effect arising from (27) can be obtained by including the four-fermion interactions in the path integral (17). These operators allow for new ways to close the fermion legs in the 't Hooft vertex, as depicted in Figure 2. The largest contribution arises from just

one insertion of \mathcal{O}_F , as shown in Figure 2(a), while more insertions of the four-fermion operator, such as in Figure 2(b) are suppressed by powers of Λ_F . Similar to the definition (23) for $\chi_W(0)$ we can define a fermion mixed correlator

$$\chi_{F,ijkl}(0) = -i \lim_{k \to 0} \int d^4 x \, e^{ikx} \left\langle 0 \left| T \left\{ \frac{1}{32\pi^2} G\widetilde{G}(x), \frac{\lambda_{ijkl}}{\Lambda_F^2} \mathcal{O}_{F,ijkl}(0) \right\} \right| 0 \right\rangle .$$
(28)



Figure 2: The t'Hooft vertex that includes the insertion of four-fermion operators. Fermion legs are closed with one four-fermion operator \mathcal{O}_F and two Higgs-fermion Yukawa interactions in (a) and three four-fermion operators \mathcal{O}_F in (b).

The only operators contributing to the fermion path integral are those with two pairs of flavor indices $(i = j \neq k = l \text{ or } i = l \neq k = j)$, i.e. $\mathcal{O}_{F,iijj}$, and $\mathcal{O}_{F,ijji}$, both of which are hereon generically referred to as $\mathcal{O}_{F,ij}$ with the corresponding coupling constant $\lambda_{ij} \equiv \lambda_{iijj}$ (or λ_{ijij}). The explicit expression for such a generic operator $\mathcal{O}_{F,ij}$ can be computed as:

$$\chi_{F,ij}(0) = -2i \int d^4 x_0 \frac{d\rho}{\rho^5} C[N] e^{0.292N_f} \left(\frac{8\pi^2}{g^2(1/\rho)}\right)^{2N} e^{-\frac{8\pi^2}{g^2(1/\rho)}} \\ \times \frac{2\lambda_{ij}}{y_i y_j} (N_f - 3)!! \left(\prod_{k=1}^{N_f} \frac{y_k \rho}{\sqrt{2}}\right) \mathcal{I}^{N_f/2 - 1} \frac{1}{\Lambda_F^2} \bar{\psi}_i^{(0)} i\gamma_5 \psi_i^{(0)} \bar{\psi}_j^{(0)} \psi_j^{(0)}(0) \frac{1}{32\pi^2} \int d^4 x \, G \widetilde{G}(x) , \\ = 2i \, \frac{2(-i\lambda_{ij})}{y_i y_j} \int \frac{d\rho}{\rho^5} \frac{C_f[N]}{N_f - 1} \left(\frac{8\pi^2}{g^2(1/\rho)}\right)^{2N} \frac{12}{5\rho^2 \Lambda_F^2} e^{-\frac{8\pi^2}{g^2(1/\rho)}} ,$$
(29)

where we have also included the effect of the anti-instanton. The part of $\mathcal{O}_{F,ij}$ contributing to the path integral in the instanton background is $i\psi^{\dagger}_{L,i}\psi_{R,i}\psi^{\dagger}_{L,j}\psi_{R,j}$, while in the anti-instanton background (where $G\tilde{G} \rightarrow -G\tilde{G}$) it is $-i\psi^{\dagger}_{R,i}\psi_{L,i}\psi^{\dagger}_{R,j}\psi_{L,j}$. These two contributions add up³ to give the factor of 2*i* in (29).

The result (29) can also be understood in terms of the results (17) and (18) from the fermionic path integral, up to the overall ratio of couplings. If we assume that \mathcal{O}_F is generated by a heavy scalar of mass Λ_F , interacting with Standard Model quarks via Yukawa interactions, (18)

³Instead, for *CP* even operators of the type $\bar{\psi}_i \psi_i \bar{\psi}_j \psi_j$ there is a cancellation between the two contributions since $\psi_{L,i}^{\dagger} \psi_{R,i} \psi_{L,j}^{\dagger} \psi_{R,j}$ and $\psi_{R,i}^{\dagger} \psi_{L,i} \psi_{R,j}^{\dagger} \psi_{L,j}$ both appear with the same sign.

implies a factor of $12\pi^2 \rho^2 / 5\pi^2 \Lambda_F^2 \rho^4 = 12/5\rho^2 \Lambda_F^2$ relative to the expression (19), which matches the factor inside the integral. The factor $1/(N_f - 1)$ arises from having a fewer number of contractions- $(N_f - 3)!!$ compared to (17), assuming only one insertion of the operator $\mathcal{O}_{F,ij}$.

Furthermore, notice that y_i and y_j have been explicitly factored out of (29) to write the result in terms of $C_f[N]$ defined in (20). For $-i\lambda_{ij} \sim 1$, this shows that the effect of the fourfermion operator, being $\propto 1/y_i y_j$, is most enhanced for the up and down quarks compared to that from the Weinberg gluonic operator or the second and third generation quarks. However, the four-fermion operator coefficient λ_{ij} can be chirally suppressed by Yukawa couplings [56]. For example, such four-fermion operators with a chiral suppression can arise from the overlap of fermion profiles in extra dimension models [57]. Thus, we will henceforth assume that $-i\lambda_{ij} \propto$ $y_i y_j$ so that the effect of the four-fermion operator is similar to that of the Weinberg gluonic operator as well as the contributions from the other generations of quarks.

Assuming $-i\lambda_{ij} = y_i y_j/2$, we then have $N_f(N_f-1)$ contributions of the fermion susceptibility (29) for both types of operators $\mathcal{O}_{F,iijj}$, and $\mathcal{O}_{F,ijji}$, each. Thus, for $N_f = 6$ we obtain

$$\chi_F(0) \equiv 2N_f(N_f - 1)\chi_{F,ij}(0) = 2i\frac{144}{5\Lambda_F^2} \int \frac{d\rho}{\rho^7} C_f[N] \left(\frac{8\pi^2}{g^2(1/\rho)}\right)^{2N} e^{-\frac{8\pi^2}{g^2(1/\rho)}} .$$
 (30)

Using (30) we will place limits on a generic scale Λ_F that represents all these fermion effects.

Finally, note that in supersymmetric theories the operator \mathcal{O}_F can arise from a dimensionfour term in the superpotential [58]. After integrating out the scalar superpartners this leads to a four-fermion term with

$$\frac{1}{\Lambda_F^2} \sim \frac{g^2}{16\pi^2} \frac{1}{\Lambda_{\rm UV} m_{\rm SUSY}} , \qquad (31)$$

where $\Lambda_{\rm UV}$ is the UV scale of the superpotential term and $m_{\rm SUSY}$ is the supersymmetry-breaking scale of the scalar superpartners. The bounds on Λ_F can thus be interpreted as bounds on the scalar superpartner masses.

3 Induced θ

Using the results in Section 2 we can now obtain an estimate for the shift in the axion potential minimum due to CP odd operators. In the presence of the Weinberg operator the axion potential is modified by a linear term in the axion field

$$V(a) = \chi_W(0) \left(\frac{a}{f_a}\right) + \frac{1}{2}\chi(0) \left(\frac{a}{f_a}\right)^2 , \qquad (32)$$

where we have promoted the theta angle to the axion field, $\bar{\theta} \to a/f_a$. This leads to a shift in the potential minimum by an amount

$$\left\langle \frac{a}{f_a} \right\rangle \equiv \theta_{\rm ind} = -\frac{\chi_W(0)}{\chi(0)} \ .$$
 (33)

In the case of four-fermion operators the linear potential term again causes a shift in the potential minimum given by (33), with $\chi_W(0)$ replaced by $\chi_F(0)$.

The induced θ then directly contributes to EDM observables such as the neutron EDM where

$$d_n \propto \frac{m_q}{\Lambda_{\rm QCD}^2} |\theta_{\rm ind}| = \frac{m_q}{\Lambda_{\rm QCD}^2} \frac{\chi_{W,F}(0)}{\chi(0)} .$$
(34)

The experimental limit arising from the neutron EDM gives the constraint

$$|\theta_{\rm ind}| \lesssim 10^{-10} , \qquad (35)$$

which can now be used to obtain constraints on various heavy axion scenarios⁴.

3.1 QCD

We first consider the effect of dimension six operators in QCD with N_L light fermions (i.e. $m_f \leq \Lambda_{\rm QCD}$). The induced θ (33) that arises from including the Weinberg operator is given by

$$\theta_{\rm ind}^{\rm QCD} \approx \xi_W \frac{b_0 - 4 + N_L}{b_0 - 6 + N_L} \frac{\Lambda_{\rm QCD}^2}{\Lambda_W^2} , \qquad (36)$$

where $\xi_W = 384\pi^2/5$, b_0 is the β -function coefficient and the *CP*-violation scale $\Lambda_{\rm CP}$ is identified with Λ_W . Note that in (36) the product of all light quark masses cancel and the induced θ becomes small (or decouples) as $\Lambda_W \to \infty$. Imposing the constraint (35) for QCD ($b_0^{\rm QCD} = 9$, $N_L = 3$ and $\Lambda_{\rm QCD} \approx 300$ MeV), gives the limit $\Lambda_W \gtrsim 10^6$ GeV on the effective scale of the Weinberg operator.

For the case of the CP- odd four-fermion operator, the 't Hooft vertex now has two fewer factors of ρm_f compared to the topological susceptibility resulting from (21). This gives a bound similar to Λ_W when there is no chirality suppression in the four-fermion operator, otherwise the Λ_F bound is much weaker. A calculation for θ_{ind} using the chiral anomaly can be found in [59], which agrees with our estimate of the bound on Λ_F within an order of magnitude.

As such, current constraints on the neutron EDM correspond to new CP-violating physics at ~ 10⁶ GeV. Thus, future neutron EDM experiments can probe new CP violating sources at scales ranging from ~ 10⁶ - 10⁹ GeV, beyond which the SM contribution due to the CKM phase becomes comparable in size.

3.2 4D Small Instantons

3.2.1 Product gauge group

A heavy axion can be generated by extending the QCD gauge group into a product gauge group $SU(3)^k = SU(3)_1 \times SU(3)_2 \times \cdots \times SU(3)_k$ which is spontaneously broken at a scale Λ_{SI} [38, 39]. Small instantons at the scale Λ_{SI} associated with the product gauge groups lead to this enhancement. The SM quarks are assumed to be charged under only $SU(3)_1$. In addition, there are k axions, labelled by i, which couple to the k SU(3) $G\tilde{G}$ terms with decay constants f_{a_i} , eliminating the k theta terms.

At the scale Λ_{SI} the QCD gauge coupling, α is matched to the $SU(3)_k$ gauge couplings, α_i via the relation

$$\frac{1}{\alpha(\Lambda_{\rm SI})} = \sum_{i=1}^{k} \frac{1}{\alpha_i(\Lambda_{\rm SI})} .$$
(37)

This relation implies that each individual coupling α_i must be larger than the QCD coupling at the scale Λ_{SI} . Therefore, the larger couplings $\alpha_i(\Lambda_{SI})$ can make the small instanton effects

⁴For simplicity, we will present limits that arise from the individual operators \mathcal{O}_W and \mathcal{O}_F separately. Our results can be straightforwardly generalized by summing the contributions in (33) if both operators are present.

dominate over the usual QCD large instantons. This effect is most dominant in the limit $k \gg 1$, where the axion masses scale as $m_{a_1} \sim \sqrt{\prod_f y_f} \Lambda_{SI}^2 / f_{a_1}$ (with y_f the quark Yukawa couplings) and $m_{a_i} \sim \Lambda_{SI}^2 / f_{a_i}$ for i = 2..., k, showing that the lightest axion mass (m_{a_1}) can remain much heavier that the QCD axion mass for $\Lambda_{SI} \gg \Lambda_{QCD}$.

For concreteness, let us consider the case with small k, where there is some perturbative control and the instanton (or anti-instanton) background still gives us qualitatively accurate results. Assuming the product gauge group is broken by scalars with a VEV, v_{ϕ} , the effective cutoff for the instanton size then becomes $2\pi v_{\phi}$, in contrast to the naive expectation, $\Lambda_{\rm SI}$ [39]. The constraint (35) can then be used to obtain limits on the scales associated with the sources of CP violation from the Weinberg and four-fermion operators. Since the QCD instanton contribution to $\chi_{W,F}(0)$ is suppressed by at least $\Lambda^2_{\rm QCD}/\Lambda^2_{W,F}$, the small instanton contribution from the UV gauge group dominates and results in

$$\theta_{\rm ind} \approx \xi_{W,F} \frac{2}{b_{0,i} - 6} \frac{(2\pi v_{\phi})^2}{\Lambda_{W,F}^2} \approx \xi_{W,F} \frac{8\pi^2}{b_{0,i} - 6} \frac{\Lambda_{\rm SI}^2}{\Lambda_{W,F}^2} , \qquad (38)$$

where $\xi_F = 24N_f/5$, ξ_W is defined under (36) and we have assumed $\Lambda_{\rm SI} \approx v_{\phi}$ in the second expression in (38). The constraint (35) then implies $\Lambda_{\rm SI}/\Lambda_W \lesssim 10^{-8}$ and $\Lambda_{\rm SI}/\Lambda_F \lesssim 10^{-7}$ or $\Lambda_{\rm SI} \lesssim 10^{10}(10^{11})$ GeV for $\Lambda_W(\Lambda_F) = M_P$ where $M_P = 2.4 \times 10^{18}$ GeV is the (reduced) Planck mass⁵, $b_{0,1} = 13/2$ and $b_{0,k} = 21/2$. For $i = 2, \ldots, k - 1$, the same expression (38) holds with $b_{0,i} = 10$, and $v_{\phi} \to \sqrt{2}v_{\phi}$, which does not change the bounds significantly⁶. Note that if UV couplings are included in (22) then the effective scale Λ_W can be larger than M_P . Assuming $f_a > \Lambda_{\rm SI}$, the limits on $\Lambda_{W,F}$ correspond to a maximum possible axion mass enhancement of $\sim 10^7$ for k = 3 relative to the QCD axion [38, 39]. As such, axion masses $m_a \gtrsim 100$ MeV with $f_a \lesssim 10^7$ GeV [60, 61] can be explored in future experimental searches.

However, when $f_a < \Lambda_{\rm SI}$, we need to UV complete the dimension five axion-GG coupling and explain the PQ breaking. This can be done in a minimal KSVZ-type scenario [7, 8], by introducing a single heavy Dirac fermion Ψ , with mass m_{Ψ} , charged under the $U(1)_{PQ}$ symmetry, which changes the instanton measure by a factor of $e^{0.292}\rho m_{\Psi}$. Combining this with the contribution arising from the running of the gauge coupling between m_{Ψ} and $\Lambda_{\rm SI}$, the topological susceptibility (or any similar correlator) is modified to

$$\chi(0) \to \chi(0) \frac{b_0 - 4}{b_0 - 11/3} \left(\frac{m_\Psi}{\Lambda_{\rm SI}}\right)^{-2/3} e^{0.292} \left(\frac{m_\Psi}{\Lambda_{\rm SI}}\right) \approx \chi(0) \left(\frac{f_a}{\Lambda_{\rm SI}}\right)^{1/3} , \qquad (39)$$

where the Yukawa coupling between Ψ and the PQ scalar is assumed to be order one, i.e. $m_{\Psi} \approx f_a$. Since $m_a^2 \propto \chi(0)$ this suppresses the axion mass enhancement by an amount $(f_a/\Lambda_{\rm SI})^{1/6}$ [62]. Thus for the experimentally interesting region of $m_a \gtrsim 100$ MeV and $f_a \lesssim 10^7$ GeV, the axion mass enhancement is reduced by up to a factor of 10 when $f_a < \Lambda_{\rm SI}$.

A similar result is also obtained for an enlarged color group [32, 35, 36] where $\Lambda_{\rm SI}$ is identified with the scale where the enlarged symmetry group is broken and the appropriate b_0 is used. In all these cases, there is again a non-decoupling effect that depends on the ratio $\Lambda_{\rm SI}/\Lambda_{W,F}$.

⁵The difference in these two bounds results from the size of the different prefactors $\xi_{W,F}$, where ξ_W results from the large number of color contractions in (23), while ξ_F arises from the smaller flavor multiplicity of the four-fermion operator (27).

⁶It is possible that the axion mass could instead be dominated by QCD large instantons. But in this case the *CP* violation arising from small instantons of the product gauge group gives the much weaker constraint that $\Lambda_{\rm SI}/\Lambda_W \lesssim 10^{-8} \times m_{a,\rm QCD}/m_{a_1}$. For instance, assuming $m_{a,\rm QCD}/m_{a_1} = 10^3$ implies that $\Lambda_{\rm SI} \lesssim 10^{13}$ GeV for $\Lambda_W = M_P$.

3.2.2 Mirror QCD

A heavy axion can also be obtained by assuming that there exists a \mathbb{Z}_2 mirror copy of QCD that becomes strong at a scale $\Lambda'_{\text{QCD}} (\equiv \Lambda_{\text{SI}}) \gg \Lambda_{\text{QCD}}$ [22, 23, 30, 31, 34]. The axion is \mathbb{Z}_2 neutral and couples to both QCD and mirror QCD, via the interaction

$$\frac{1}{32\pi^2} \frac{a}{f_a} \varepsilon^{\mu\nu\rho\sigma} \left(G^c_{\mu\nu} G^c_{\rho\sigma} + G^{\prime c}_{\mu\nu} G^{\prime c}_{\rho\sigma} \right), \tag{40}$$

where $G'_{\mu\nu}$ is the mirror QCD field strength. The axion now receives contributions from the mirror QCD instantons (which are small relative to those from QCD).

The mirror QCD expression for the induced θ due to the Weinberg operator can be obtained by substituting $\Lambda_{\rm SI}$ in (36). This leads to the bounds $\Lambda_{\rm SI}/\Lambda_W \lesssim 10^{-7}$ or $\Lambda_{\rm SI} \lesssim 10^{11}$ GeV for $\Lambda_W = M_P$, assuming the mirror Higgs VEV, $v' \gg \Lambda_{\rm SI}$ such that QCD' is a pure Yang-Mills theory at $\Lambda_{\rm SI}$ with $b_0^{\rm QCD'} = 11$. These bounds for the Weinberg operator do not change appreciably if this assumption is relaxed.

The induced θ from the four-fermion operator can be obtained by considering $N_L \geq 2$ light flavors in QCD'. Applying the QCD result (21) for QCD' then gives

$$\theta_{\rm ind,F} \approx \frac{2N_L(N_L-1)}{5\pi^2} \frac{b_0 - 4 + N_L}{b_0 - 8 + N_L} \frac{\Lambda_{\rm SI}^4}{v'^2 \Lambda_F^2} \approx \frac{2N_L(N_L-1)}{5\pi^2} \frac{b_0 - 4 + N_L}{b_0 - 8 + N_L} \frac{\Lambda_{\rm SI}^2}{\Lambda_F^2} , \qquad (41)$$

where we have taken $v' \approx \Lambda_{\rm SI}$ in the last expression in (41). Assuming $b_0 = 9$ and $N_L = 3$, implies $\Lambda_{\rm SI}/\Lambda_F \lesssim 10^{-5}$, or $\Lambda_{\rm SI} \lesssim 10^{13}$ GeV for $\Lambda_F = M_P$. Again, the difference in the $\Lambda_{W,F}$ bounds arises from the different color and flavor multiplicity factors.

3.3 5D Small Instantons

Another way for the QCD coupling to become large at a UV scale and increase the effect of small instantons is to consider a 5D model where QCD gluons propagate in a 5th dimension of size R. The axion can be identified with a UV boundary localized field that couples to QCD via a coupling proportional to $1/f_a$, with f_a an independent parameter of the theory. Above the scale 1/R the QCD coupling increases in strength until the coupling becomes strong at the cutoff scale Λ_5 which is defined by the relation [40]

$$\Lambda_5 R = \frac{6\pi\epsilon}{\alpha(1/R)} , \qquad (42)$$

where $\alpha = g^2/(4\pi)$ and $\epsilon \leq 1$ is a perturbativity parameter⁷. The small instanton scale can be identified as $\Lambda_{\rm SI} \equiv \Lambda_5$. The 4D effective action is approximately given by [40]

$$S_{\text{eff}} \approx \frac{2\pi}{\alpha_s(1/R)} - \frac{R}{\rho} + b_0 \ln \frac{R}{\rho} , \qquad (43)$$

where the power-law term R/ρ arises from summing over the 5D Kaluza-Klein gluons. Thus, small instantons of size $1/\Lambda_{\rm SI} \lesssim \rho \lesssim R$ can now reduce the effective action and contribute greatly to the path integral.

⁷Note that in the 5D model, small instantons can be made to dominate when perturbativity still holds. This implies that our instanton (or anti-instanton) approximation used for the correlators will give more accurate quantitative results relative to QCD.



Figure 3: Lower limit on the effective scale of the dimension-six Weinberg (four-fermion) operator, depicted in purple (orange) as a function of the extra dimension scale 1/R, assuming $\epsilon = 0.30$. The Planck scale is shown as a dotted line for reference. The dashed lines represent the limit from the approximation (44). The deviation from (44) arises since for small 1/R, large QCD instantons begin to dominate the instanton integral $\chi(0)$.

Using an approximate expression for the integrals in (14) and (26) with the effective action (43), the induced θ from 5D small instantons is

$$\theta_{\rm ind} \approx \xi_{W,F} \frac{\Lambda_{\rm SI}^2}{\Lambda_{WF}^2} ,$$
(44)

where ξ_W and ξ_F are defined under (36) and (38), respectively. The induced θ no longer necessarily decouples in the limit $\Lambda_{\rm SI}, \Lambda_{W,F} \to \infty$. Imposing the constraint (35) leads to the limit $\Lambda_{\rm SI}/\Lambda_W(\Lambda_F) \leq 10^{-7}(10^{-6})$. For $\Lambda_W(\Lambda_F) = M_P$ this implies an upper bound $\Lambda_{\rm SI} \leq 10^{11}(10^{12})$ GeV on the 5D strong coupling scale. The limit on $\Lambda_{W,F}$ from an exact numerical evaluation of $\theta_{\rm ind}$ is shown in Figure 3. We see that the limit on $\Lambda_{W,F}$ deviates from (44) for small 1/R (and hence small Λ_5). The limits on the ratio $\Lambda_{\rm SI}/\Lambda_{W,F}$ imply that for the case when $\Lambda_{W,F} \sim \Lambda_5(=\Lambda_{\rm SI})$, the dimension six terms would need to be generated in the UV completion of the 5D model with an additional suppression in the otherwise order-one coefficients.

The corresponding range of axion mass enhancement is depicted in Figure 4. Note that both effects of small instantons - the enhancement of the axion mass and the shift in the axion potential minimum due to CP violating operators - are dominant only for large 1/R, since eventually large (QCD) instantons dominate the susceptibility at small values of 1/R.

Furthermore, when $f_a < \Lambda_{\rm SI}$ the axion mass enhancement is reduced by the factor $(f_a/\Lambda_{\rm SI})^{1/6}$ as obtained from (39). This means that in the experimentally viable region of $m_a \gtrsim 100$ MeV and $f_a \lesssim 10^7$ GeV [60, 61], the axion mass enhancement is reduced by up to an order of magnitude, as can be seen in Figure 4, where we have taken $f_a = 10^6$ GeV as a representative value.



Figure 4: The ratio of the enhanced axion mass to the QCD axion mass as a function of the extra dimension scale, 1/R. The dotted contour lines assume $f_a > \Lambda_{\rm SI}$ and depict the ratio for different values of the perturbitivity parameter ϵ , up to the maximum possible enhancement in the red shaded region. The dashed contours assume $f_a = 10^6$ GeV and include the suppression (39) when $f_a < \Lambda_{\rm SI}$. The blue shaded region to the right shows the excluded 1/R range due to the Weinberg gluonic operator.

3.4 Enhanced EDMs

Compared to QCD, the small instanton contributions provide an enhancement to the EDMs due to CP violating sources. In particular, using (38), (41) and (44) we see that the neutron EDM (34) is enhanced by a factor of $\Lambda_{\rm SI}^2/\Lambda_{\rm QCD}^2$ compared to the θ induced from new CP- odd sources in QCD (see (36)). Therefore, measuring the neutron EDM can be interpreted as a probe of the small instanton scale, $\Lambda_{\rm SI}$. For example, if $\Lambda_{W,F} = M_P$, this corresponds to modified strong dynamics at scales of order $\Lambda_{\rm SI} \sim 10^8 - 10^{11}$ GeV, where the lower limit represents a neutron EDM value equivalent to the Standard Model CKM contribution. Furthermore, if the CP violating sources appear at scales lower than the Planck scale, then any new contribution due to small instantons will appear at even lower scales 10^4 GeV $\leq \Lambda_{\rm SI} \leq 10^8$ GeV, where the model-dependent lower limit corresponds to the scale of axion mass enhancement.

Finally note that when $f_a \lesssim \Lambda_{\rm SI}$, the UV completion of the dimension five axion-gluon coupling does not affect the predictions for the induced θ . Since the neutron EDM (34) depends only on the ratio of the mixed correlators with the topological susceptibility, the suppression factor in (39) cancels, leaving the results for the induced θ unchanged.

4 Conclusion

The QCD axion solution provides an elegant mechanism for solving the strong CP problem in such a way that an arbitrarily large amount of CP violation at UV scales, Λ_{CP} can be sufficiently

decoupled as $\Lambda_{CP} \to \infty$. This is in contrast with solutions to the strong CP problem that invoke exact discrete symmetries. For these solutions there is a non-decoupling of the additional sources of CP violation, which means that arbitrarily large amounts of CP violation cannot be tolerated at UV scales in models with exact parity or CP symmetry.

Heavy axion models represent a qualitatively different class of solution to the strong CP problem in which new dynamics at some UV scale, Λ_{SI} magnifies the effect of small instantons (which are normally exponentially suppressed), giving rise to a new contribution and enhancement of the axion mass. This has led to renewed interest in axion searches outside the usual QCD axion mass window. However, in the presence of additional sources of CP- violation, the enhanced effect of small instantons could also lead to enhanced EDM observables such as the neutron EDM as well as possible non-decoupling effects.

We have estimated these effects by calculating the topological susceptibility and mixed correlators in the presence of two *CP*-violating dimension-six operators: the Weinberg gluonic operator and a *CP*-odd four-fermion operator. The calculation is performed using an instanton (or anti-instanton) background where Standard Model fermion chiral zero modes in the t'Hooft vertex are closed with the Higgs boson. Identifying the scale of the additional sources of *CP* violation with Λ_{CP} we find that the axion potential minimum shifts by an amount $\theta_{ind} \propto \Lambda_{SI}^2/\Lambda_{CP}^2$ in several heavy axion models, where Λ_{SI} is the scale where small instanton effects dominate. This result reveals that unlike the minimal QCD axion models, the amount of decoupling is limited, although not as restrictive as models with exact discrete symmetries. Imposing the neutron EDM derived limit $|\bar{\theta}| \leq 10^{-10}$, we obtain the constraint $\Lambda_{SI}/\Lambda_{CP} \leq 10^{-8}$, which is stronger than the naive estimate of 10^{-5} due to sizeable prefactors that depend on the particular heavy axion model. In particular, for a benchmark value of $\Lambda_{CP} \simeq M_P$ requires $\Lambda_{SI} \leq 10^{10}$ GeV (as can be seen in Figure 4 for the 5D small instanton model).

The modification of the decoupling behaviour is a direct consequence of the new dynamical scale $\Lambda_{\rm SI}$. Our results therefore imply that EDM observables such as the neutron EDM can be enhanced in heavy axion models up to the current experimental limit $d_n \leq 10^{-26}$ e·cm. This compares with the SM CKM prediction ($\sim 10^{-32} - 10^{-31}$ e·cm). Thus, besides axion searches, EDM observables provide another probe of UV scales in heavy axion models associated with new dynamics, assuming that this class of models plays any role in solving the strong *CP* problem.

Acknowledgments

T.G. thanks Raymond Co and Keisuke Harigaya for useful discussions. This work is supported in part by the Department of Energy under Grant DE-SC0011842 at the University of Minnesota. T.G. is also supported by the Simons Foundation.

References

- M. Kobayashi and T. Maskawa, CP Violation in the Renormalizable Theory of Weak Interaction, Prog. Theor. Phys. 49 (1973) 652.
- [2] C. Abel et al., Measurement of the Permanent Electric Dipole Moment of the Neutron, Phys. Rev. Lett. 124 (2020) 081803 [2001.11966].

- [3] B. Graner, Y. Chen, E.G. Lindahl and B.R. Heckel, Reduced Limit on the Permanent Electric Dipole Moment of Hg199, Phys. Rev. Lett. 116 (2016) 161601 [1601.04339].
- [4] R.D. Peccei and H.R. Quinn, CP Conservation in the Presence of Instantons, Phys. Rev. Lett. 38 (1977) 1440.
- [5] S. Weinberg, A New Light Boson?, Phys. Rev. Lett. 40 (1978) 223.
- [6] F. Wilczek, Problem of Strong P and T Invariance in the Presence of Instantons, Phys. Rev. Lett. 40 (1978) 279.
- [7] J.E. Kim, Weak Interaction Singlet and Strong CP Invariance, Phys. Rev. Lett. 43 (1979) 103.
- [8] M.A. Shifman, A.I. Vainshtein and V.I. Zakharov, Can Confinement Ensure Natural CP Invariance of Strong Interactions?, Nucl. Phys. B 166 (1980) 493.
- [9] M. Dine, W. Fischler and M. Srednicki, A Simple Solution to the Strong CP Problem with a Harmless Axion, Phys. Lett. 104B (1981) 199.
- [10] A.R. Zhitnitsky, On Possible Suppression of the Axion Hadron Interactions. (In Russian), Sov. J. Nucl. Phys. 31 (1980) 260.
- [11] I.I.Y. Bigi and N.G. Uraltsev, Induced Multi Gluon Couplings and the Neutron Electric Dipole Moment, Nucl. Phys. B 353 (1991) 321.
- [12] M. Pospelov and A. Ritz, Neutron EDM from electric and chromoelectric dipole moments of quarks, Phys. Rev. D 63 (2001) 073015 [hep-ph/0010037].
- [13] M. Pospelov and A. Ritz, *Electric dipole moments as probes of new physics*, Annals Phys. 318 (2005) 119 [hep-ph/0504231].
- [14] A.E. Nelson, Naturally Weak CP Violation, Phys. Lett. B 136 (1984) 387.
- [15] S.M. Barr, Solving the Strong CP Problem Without the Peccei-Quinn Symmetry, Phys. Rev. Lett. 53 (1984) 329.
- [16] K.S. Babu and R.N. Mohapatra, CP Violation in Seesaw Models of Quark Masses, Phys. Rev. Lett. 62 (1989) 1079.
- [17] K.S. Babu and R.N. Mohapatra, A Solution to the Strong CP Problem Without an Axion, Phys. Rev. D 41 (1990) 1286.
- [18] R.N. Mohapatra and A. Rasin, Simple supersymmetric solution to the strong CP problem, Phys. Rev. Lett. 76 (1996) 3490 [hep-ph/9511391].
- [19] R. Kuchimanchi, Solution to the strong CP problem: Supersymmetry with parity, Phys. Rev. Lett. 76 (1996) 3486 [hep-ph/9511376].
- [20] B. Holdom, Nonstandard order parameters and the origin of CP violation, Phys. Rev. D 61 (2000) 011702 [hep-ph/9907361].

- [21] G. Hiller and M. Schmaltz, Solving the Strong CP Problem with Supersymmetry, Phys. Lett. B 514 (2001) 263 [hep-ph/0105254].
- [22] Z. Berezhiani, L. Gianfagna and M. Giannotti, Strong CP problem and mirror world: The Weinberg-Wilczek axion revisited, Phys. Lett. B500 (2001) 286 [hep-ph/0009290].
- [23] A. Hook, Anomalous solutions to the strong CP problem, Phys. Rev. Lett. 114 (2015) 141801 [1411.3325].
- [24] J.R. Ellis and M.K. Gaillard, Strong and Weak CP Violation, Nucl. Phys. B 150 (1979) 141.
- [25] I.B. Khriplovich, Quark Electric Dipole Moment and Induced θ Term in the Kobayashi-Maskawa Model, Phys. Lett. B 173 (1986) 193.
- [26] M.E. Pospelov, Radiative corrections to theta term in the left-right supersymmetric models, Phys. Lett. B 391 (1997) 324 [hep-ph/9609458].
- [27] P.H. Frampton and O.C.W. Kong, Strong CP and low-energy supersymmetry, Phys. Lett. B 402 (1997) 297 [hep-ph/9612452].
- [28] J. de Vries, P. Draper and H.H. Patel, Do Minimal Parity Solutions to the Strong CP Problem Work?, 2109.01630.
- [29] S. Dimopoulos, A Solution of the Strong CP Problem in Models With Scalars, Phys. Lett. B 84 (1979) 435.
- [30] V.A. Rubakov, Grand unification and heavy axion, JETP Lett. 65 (1997) 621 [hep-ph/9703409].
- [31] H. Fukuda, K. Harigaya, M. Ibe and T.T. Yanagida, Model of visible QCD axion, Phys. Rev. D92 (2015) 015021 [1504.06084].
- [32] T. Gherghetta, N. Nagata and M. Shifman, A Visible QCD Axion from an Enlarged Color Group, Phys. Rev. D93 (2016) 115010 [1604.01127].
- [33] T. Gherghetta and M.D. Nguyen, A Composite Higgs with a Heavy Composite Axion, JHEP 12 (2020) 094 [2007.10875].
- [34] A. Hook, S. Kumar, Z. Liu and R. Sundrum, *High Quality QCD Axion and the LHC*, *Phys. Rev. Lett.* **124** (2020) 221801 [1911.12364].
- [35] B. Holdom and M.E. Peskin, Raising the Axion Mass, Nucl. Phys. B208 (1982) 397.
- [36] B. Holdom, Strong QCD at High-energies and a Heavy Axion, Phys. Lett. 154B (1985) 316.
- [37] J.M. Flynn and L. Randall, A Computation of the Small Instanton Contribution to the Axion Potential, Nucl. Phys. B293 (1987) 731.
- [38] P. Agrawal and K. Howe, Factoring the Strong CP Problem, JHEP 12 (2018) 029 [1710.04213].

- [39] C. Csáki, M. Ruhdorfer and Y. Shirman, UV Sensitivity of the Axion Mass from Instantons in Partially Broken Gauge Groups, JHEP 04 (2020) 031 [1912.02197].
- [40] T. Gherghetta, V.V. Khoze, A. Pomarol and Y. Shirman, The Axion Mass from 5D Small Instantons, JHEP 03 (2020) 063 [2001.05610].
- [41] T. Gherghetta and A. Pomarol, Small instantons in weakly-gauged holographic models, JHEP 11 (2021) 136 [2110.01762].
- [42] A.A. Belavin, A.M. Polyakov, A.S. Schwartz and Y.S. Tyupkin, Pseudoparticle Solutions of the Yang-Mills Equations, Phys. Lett. B 59 (1975) 85.
- [43] G. 't Hooft, Computation of the Quantum Effects Due to a Four-Dimensional Pseudoparticle, Phys. Rev. D14 (1976) 3432.
- [44] R. Jackiw, C. Nohl and C. Rebbi, Conformal properties of pseudoparticle configurations, Physical Review D 15 (1977) 1642.
- [45] E. Witten, Current Algebra Theorems for the U(1) Goldstone Boson, Nucl. Phys. B 156 (1979) 269.
- [46] S.F. Cordes, The Instanton Induced Superpotential in Supersymmetric QCD, Nucl. Phys. B 273 (1986) 629.
- [47] A.I. Vainshtein, V.I. Zakharov, V.A. Novikov and M.A. Shifman, ABC's of Instantons, Sov. Phys. Usp. 25 (1982) 195.
- [48] C.G. Callan, Jr., R.F. Dashen and D.J. Gross, Toward a Theory of the Strong Interactions, Phys. Rev. D17 (1978) 2717.
- [49] D. Diakonov and V.Y. Petrov, Instanton Based Vacuum from Feynman Variational Principle, Nucl. Phys. B 245 (1984) 259.
- [50] E.V. Shuryak, Instantons in QCD. 1. Properties of the 'Instanton Liquid', Nucl. Phys. B 319 (1989) 521.
- [51] D. Diakonov and V.Y. Petrov, A Theory of Light Quarks in the Instanton Vacuum, Nucl. Phys. B 272 (1986) 457.
- [52] D. Diakonov, M.V. Polyakov and C. Weiss, Hadronic matrix elements of gluon operators in the instanton vacuum, Nucl. Phys. B 461 (1996) 539 [hep-ph/9510232].
- [53] D. Diakonov and V.Y. Petrov, Chiral Condensate in the Instanton Vacuum, Phys. Lett. B 147 (1984) 351.
- [54] S. Weinberg, Larger Higgs Exchange Terms in the Neutron Electric Dipole Moment, Phys. Rev. Lett. 63 (1989) 2333.
- [55] C. Weiss, Nucleon matrix element of Weinberg's CP-odd gluon operator from the instanton vacuum, Phys. Lett. B 819 (2021) 136447 [2103.13471].

- [56] R. Kitano and W. Yin, Strong CP problem and axion dark matter with small instantons, JHEP 07 (2021) 078 [2103.08598].
- [57] Q. Bonnefoy, P. Cox, E. Dudas, T. Gherghetta and M.D. Nguyen, Flavoured Warped Axion, JHEP 04 (2021) 084 [2012.09728].
- [58] M. Pospelov, A. Ritz and Y. Santoso, Flavor and CP violating physics from new supersymmetric thresholds, Phys. Rev. Lett. 96 (2006) 091801 [hep-ph/0510254].
- [59] H. An, X. Ji and F. Xu, P-odd and CP-odd Four-Quark Contributions to Neutron EDM, JHEP 02 (2010) 043 [0908.2420].
- [60] D. Cadamuro and J. Redondo, Cosmological bounds on pseudo Nambu-Goldstone bosons, JCAP 02 (2012) 032 [1110.2895].
- [61] J. Jaeckel and M. Spannowsky, Probing MeV to 90 GeV axion-like particles with LEP and LHC, Phys. Lett. B 753 (2016) 482 [1509.00476].
- [62] R.T. Co, T. Gherghetta and K. Harigaya, to appear, .