

Local Completeness Logic on Kleene Algebra with Tests

Marco Milanese and Francesco Ranzato

Dipartimento di Matematica, University of Padova, Italy

Abstract. Local Completeness Logic (LCL) has been put forward as a program logic for proving both the correctness and incorrectness of program specifications. LCL is an abstract logic, parameterized by an abstract domain that allows combining over- and under-approximations of program behaviors. It turns out that LCL instantiated to the trivial singleton abstraction boils down to O’Hearn incorrectness logic, which allows us to prove the presence of program bugs. It has been recently proved that suitable extensions of Kleene algebra with tests (KAT) allow representing both O’Hearn incorrectness and Hoare correctness program logics within the same equational framework. In this work, we generalize this result by showing how KATs extended either with a modal diamond operator or with a top element are able to represent the local completeness logic LCL. This is achieved by studying how these extended KATs can be endowed with an abstract domain so as to define the validity of correctness/incorrectness LCL triples and to show that the LCL proof system is logically sound and, under some hypotheses, complete.

Keywords: Local Completeness Logic · Incorrectness Logic · Complete Abstract Interpretation · Kleene Algebra with Tests.

1 Introduction

Kleene algebra [6] with tests (KAT) [16] allows an equational reasoning on programs and their properties. Programs are modeled as elements of a KAT, so that their properties can be algebraically derived through the general equational theory of KATs. KATs feature sound, complete, and decidable equational theories and have found successful applications in several different contexts, most notably in network programming [1, 2, 11, 29, 30]. The foundational study of Kozen [17] has shown that the reasoning of Hoare correctness logic [14] can be encoded and formulated equationally within a KAT. Later work by Desharnais, Möller and Struth [9, 23] extended KAT with a domain (KAD) to express the modal operators of propositional dynamic logic [10], thus enabling a more natural way of reasoning through a map from actions to propositions. The expressive power of KAD has been recently substantiated by Möller, O’Hearn and Hoare [22], who have shown how to encode both Hoare correctness and O’Hearn [24] incorrectness program logics in a unique class of KAD where a backward diamond modality is exploited to encode strongest postconditions. Furthermore, very recently, Zhang, De Amorim and Gaboardi [32, Theorem 1] have shown that O’Hearn incorrectness logic cannot be formulated within a conventional KAT, but, at the same time, a full fledged modal KAT is not needed. In fact, [32] proves that a KAT including a

greatest element, called TopKAT, is capable to encode both Hoare and O’Hearn logic in a purely equational fashion. Moreover, [32] provides a PSPACE algorithm to decide TopKAT equality, based on a reduction to Cohen et al. [5]’s algorithm for KAT.

This stream of works made it possible to reason equationally on both program correctness and incorrectness in the same algebraic framework. For example, in the KAD framework where a backward diamond modality $\langle a|p$ plays the role of strongest postcondition of a KAT element a (viz., a program) for a KAT test p (viz., a precondition), the validity of a Hoare correctness triple $\{p\} a \{q\}$ is determined by the inequality $\langle a|p \leq q$, while the validity of an O’Hearn incorrectness triple $[p] a [q]$ boils down to $q \leq \langle a|p$. Moreover, if a KAT test s plays the role of specification for a program a and a Hoare triple $\{p\} a \{q\}$ is provable, then a can be proved correct through the inequality $q \leq s$. Vice versa, if $[p] a [q]$ is a provable incorrectness triple, then incorrectness of a can be verified as $q \leq \neg s$.

Recently, Bruni et al. [4] put forward a novel program logic, called local completeness logic LCL, which is parameterized by an abstract domain [7, 8] of program stores and simultaneously combines over- and under-approximations of program behaviours. This program logic leverages the notion of *locally complete* abstract interpretation, meaning that the abstract interpretation of atomic program commands, such as variable assignments and Boolean guards, is complete, i.e. with no false alarm, locally on the preconditions, as opposed to standard completeness [12, 28] which must be satisfied *globally* for all the preconditions. While a global completeness program logic was proposed in [13], Bruni et al. [4] design a proof system for inferring that a program analysis is locally complete. It turns out [4, Section VI] that the instantiation of this LCL program logic to the trivial store abstraction with a unique “don’t know” value abstracting any concrete store property, boils down to O’Hearn incorrectness logic [24].

In this work, we show that the local completeness logic LCL can be made fully algebraic in a suitable KAT, yet preserving all its noteworthy logical properties. In particular, we study two different formulations of LCL given: (1) in a KAD, the KAT model used in [22], and (2) in a TopKAT, the KAT model employed in [32]. In both frameworks, we introduce a suitable notion of abstract domain of KAT that, correspondingly, induces a sound abstract semantics for KAT programs (i.e., KAT terms). Our local completeness logic on KAT, called LCK, turns out to be logically sound w.r.t. this abstract semantics, meaning that a provable LCK triple $[p] a [q]$ for an abstract domain A on a KAT K satisfies:

- (i) q is below the strongest postcondition in K of the program term a for the precondition p ;
- (ii) the program term a is locally complete for the precondition p in the abstract domain A ;
- (iii) the approximations in A of q and of the strongest postcondition of a for p coincide.

Moreover, by generalizing analogous results proved in [4], we show that:

- Our proof systems are logically complete under some additional hypotheses.
- By instantiating the LCK logic to the trivial domain abstracting any concrete value to “don’t know” we exactly obtain O’Hearn incorrectness program logic [24], thus retrieving its logical soundness and completeness as consequences of our results.

Triples of O’Hearn incorrectness logic carry two postconditions, corresponding to normal and erroneous program termination. While the original local completeness logic LCL in [4] only considers normal termination, we propose a generalization that also supports erroneous termination. Moreover, we use the KAD construction of [22] to generalize our logical soundness and completeness results to incorrectness triples.

Due to lack of space, all the proofs have been placed in Appendix ??.

2 Background on Kleene Algebra with Tests

Kleene algebra with tests (KAT) is a purely algebraic structure that provides an elegant equational framework for program reasoning. A KAT consists of actions, playing the role of programs, and tests, interpreted as pre/postconditions and Boolean guards. KAT elements can be combined with three basic operations: nondeterministic choice $a_1 + a_2$, sequential composition $a_1; a_2$, and Kleene iteration a^* . A standard model of KAT used to represent computations is the relational model, in which KAT elements are binary relations on some set, thus modeling programs as a relation between input and output states. Further models of KAT include regular languages over a finite alphabet and square matrices over another Kleene algebra. In the following, we briefly recall some basics of KAT. For more details, the reader is referred to [6, 9, 16].

An *idempotent-semiring* (*i-semiring*) is a tuple $(A, +, \cdot, 0, 1)$ where: (1) $(A, +, 0)$ is a commutative monoid with an idempotent addition, i.e., for all $a \in A$, $a + a = a$; (2) $(A, \cdot, 1)$ is a monoid, where the multiplication symbol \cdot is often omitted, such that, for any $a \in A$, $0 \cdot a = a \cdot 0$; (3) multiplication distributes over addition (in both arguments). In an *i-semiring* A , the relation $a \leq b \triangleq a + b = b$ is a partial order, referred to as the natural ordering, that we will implicitly use throughout the paper. Note that the addition $+$ is the join w.r.t. the natural ordering.

A *test-semiring* is a tuple $(A, \text{test}(A), +, \cdot, \neg, 0, 1)$ where: (1) $(A, +, \cdot, 0, 1)$ is an *i-semiring*; (2) $\text{test}(A) \subseteq A$, and $(\text{test}(A), \vee, \wedge, \neg, 0, 1)$ is a Boolean subalgebra of A with greatest element 1 and least element 0. It turns out that the meet \wedge and join \vee of the Boolean algebra $\text{test}(A)$ coincide, resp., with multiplication \cdot and addition $+$.

A *Kleene algebra* is a tuple $(K, +, \cdot, *, 0, 1)$ where: (1) $(K, +, \cdot, 0, 1)$ is an *i-semiring*; (2) $(\cdot)^* : K \rightarrow K$ is a unary operation, called Kleene star or iteration, satisfying the following conditions:

$$\begin{array}{lll} 1 + aa^* \leq a^* & 1 + a^*a \leq a^* & (*\text{-unfold}) \\ b + ac \leq c \Rightarrow a^*b \leq c & b + ca \leq c \Rightarrow ba^* \leq c & (*\text{-induction}) \end{array}$$

Definition 2.1 (KAT [16]). A *Kleene algebra with tests (KAT)* is a two-sorted algebra $(K, \text{test}(K), +, \cdot, *, \neg, 0, 1)$ such that $(K, \text{test}(K), +, \cdot, \neg, 0, 1)$ is a test-semiring and $(K, +, \cdot, *, 0, 1)$ is a Kleene algebra.

A KAT K is *countably-test-complete (CTC)* if any countable subset of $\text{test}(K)$ admits least upper bound (lub).

A KAT is **-continuous*, referred to as KAT^* , if it satisfies the following condition: for all $a, b, c \in K$, $ab^*c = \bigvee_{n \in \mathbb{N}} ab^n c$ (this equation implicitly assumes that the lub $\bigvee_{n \in \mathbb{N}} ab^n c$, w.r.t. the natural order of K , exists). \square

A *relational KAT* [18] on a carrier set X is given by a set $K \subseteq \wp(X \times X)$ of binary relations on X with tests $\text{test}(K) \subseteq \wp(\{(x, x) \mid x \in X\})$, where addition is union, multiplication is composition of relations, the additive identity is the empty relation, the multiplicative identity is $\{(x, x) \mid x \in X\}$, the Kleene star is the reflexive-transitive closure, and test complement is set complementation w.r.t. the multiplicative identity.

Informally, a backward diamond $\langle \cdot | \cdot$ on a KAT allows us to compute strongest postconditions of programs, that is, $\langle a | p$ can be interpreted as $\text{post}[a]p$.

Definition 2.2 (bdKAT [22]). A *backward-diamond KAT (bdKAT)* is a two-sorted algebra $(K, \text{test}(K), +, \cdot, *, \neg, 0, 1, \langle | \rangle)$ such that:

- (1) $(K, \text{test}(K), +, \cdot, *, \neg, 0, 1)$ is a KAT;
- (2) $\langle \cdot | \cdot : K \rightarrow (\text{test}(K) \rightarrow \text{test}(K))$ is a *backward-diamond* operator satisfying the following conditions: for all $a, b \in K$ and $p, q \in \text{test}(K)$,

$$\langle a | p \leq q \Leftrightarrow pa \leq aq \quad (\text{bd1})$$

$$\langle ab | p = \langle b | (\langle a | p) \quad (\text{bd2})$$

□

The following proposition summarizes few well-known operative properties of the $\langle \cdot | \cdot$ operator. Such operator can be defined using the codomain $\rho(\cdot)$ as $\langle a | p = \rho(pa)$ (e.g., as in [23, Section 5]) and those properties are simply derived as a consequence of [9, Lemma 4.11]. On the other hand, in this paper we consider a (equivalent) direct axiomatization of the backward-diamond; therefore for the sake of completeness we provide a direct proof.

Proposition 2.3. *Let $(K, \text{test}(K), +, \cdot, *, \neg, 0, 1, \langle | \rangle)$ a backward-diamond KAT. For any $a, b \in K$ and $p, q \in \text{test}(K)$, the following properties hold*

$$\langle a + b | p = \langle a | p + \langle b | p \quad (1)$$

$$\langle a | (p + q) = \langle a | p + \langle a | q \quad (2)$$

$$a \leq b \Rightarrow \langle a | p \leq \langle b | p \quad (3)$$

$$p \leq q \Rightarrow \langle a | p \leq \langle a | q \quad (4)$$

$$\langle s | p = p \cdot s \quad (5)$$

$$p + \langle a | \langle a^* | p \leq \langle a^* | p \quad (6)$$

If in addition, K satisfies the CTC condition,

$$\langle a^* | p = \bigvee_{n \in \mathbb{N}} \langle a^n | p. \quad (7)$$

Proof.

(1): For any $p, q \in \text{test}(K)$ and $a \in K$

$$pa \leq aq \Rightarrow paq' \leq aqq' \Rightarrow paq' = 0$$

$$paq' = 0 \Rightarrow pa = pa \cdot 1 = pa(q + q') = paq + paq' = paq \Rightarrow pa \leq aq$$

thus

$$pa \leq aq \Leftrightarrow paq' = 0 \quad (8)$$

where $\neg q$ is denoted with q' . We have

$$\begin{aligned} \langle a + b | p \leq q &\Leftrightarrow [\text{By (bd1)}] \\ p(a + b) \leq (a + b)q &\Leftrightarrow [\text{By (8)}] \\ p(a + b)q' = 0 &\Leftrightarrow \\ paq' + pbq' = 0 &\Leftrightarrow \\ paq' = 0 \wedge pbq' = 0 &\Leftrightarrow [\text{By (8)}] \\ pa \leq aq \wedge pb \leq bq &\Leftrightarrow [\text{By (bd1)}] \\ \langle a | p \leq q \wedge \langle b | p \leq q &\Leftrightarrow \\ \langle a | p + \langle b | p \leq q & \end{aligned}$$

Finally, since q is arbitrary:

$$\begin{aligned} \langle a + b | p \leq \langle a + b | p &\Rightarrow \langle a | p + \langle b | p \leq \langle a + b | p \\ \langle a | p + \langle b | p \leq \langle a + b | p &\Rightarrow \langle a + b | p \leq \langle a | p + \langle b | p \end{aligned}$$

implying the claim $\langle a + b | p = \langle a | p + \langle b | p$.

(2): Same as (1).

(3): By definition $a \leq b \Leftrightarrow a + b = b$, thus

$$\langle b | p = \langle a + b | p \stackrel{(1)}{=} \langle a | p + \langle b | p$$

implying $\langle a | p \leq \langle b | p$.

(4): Same as (3).

(5): Since $p, s \in \text{test}(K)$ and $\text{test}(K)$ forms a boolean algebra $ps = sps$ and by (bd1) $\langle s | p \leq ps$. Still by (bd1), $ps \leq s \langle s | p \leq \langle s | p$, implying $\langle s | p = ps$.

(6): The axiom ($*$ -unfold) implies $1 + a^*a \leq a^*$ so that

$$\begin{aligned} 1 + a^*a \leq a^* &\Rightarrow [\text{By (3)}] \\ \langle 1 + a^*a | p \leq \langle a^* | p &\Rightarrow [\text{By (1)}] \\ \langle 1 | p + \langle a^*a | p \leq \langle a^* | p &\Rightarrow [\text{By (5)}] \\ p + \langle a^*a | p \leq \langle a^* | p &\Rightarrow [\text{By (bd2)}] \\ p + \langle a | \langle a^* | p \leq \langle a^* | p. & \end{aligned}$$

(7): The proof of this property can be found in [22, Lemma 3.3]. \square

The axiom (bd1) is equivalent to require that $\langle a | p$ is the least test in K satisfying $pa \leq aq$ (the original definition of Kleene algebra with domain in [9] is of this form). Moreover, $pa \leq aq$ in (bd1) is equivalent to $pa = paq$ (see [9, Lemma 3.4]). For the sake of completeness we provide a direct proof of this fact.

Lemma 2.4. *Let $(K, \text{test}(K), +, \cdot, *, \neg, 0, 1, \langle | \rangle)$ a bdKAT. For all $p \in \text{test}(K)$ and $a \in K$, (bd1) holds iff $\langle a|p$ is the least (w.r.t. \leq) $q \in \text{test}(K)$ such that $pa = paq$.*

Proof. First we show the equivalence between $pa \leq aq$ and $pa = paq$. Indeed: $pa \leq aq$ implies $pa = ppa \leq paq$ and $1 \geq q$ implies $pa \geq paq$, meaning that $pa = paq$. Viceversa if $pa = paq$, then $p \leq 1$ implies $pa = paq \leq aq$.

We can show that (bd1) is equivalent to say that $\langle a|p$ is the least $q \in \text{test}(K)$ satisfying $pa \leq aq$ (which is equivalent to $pa = paq$). In formulas the latter is

$$pa \leq a\langle a|p \tag{9}$$

$$pa \leq aq \Rightarrow \langle a|p \leq q \tag{10}$$

Assume (bd1). (10) is the direction \Leftarrow of (bd1). (9) holds by instantiating $q = \langle a|p$ in (bd1). Viceversa, assume (9) and (10). (10) is the direction \Leftarrow of (bd1). The other direction holds because $pa \leq a\langle a|p \leq aq$. \square

Definition 2.5 (TopKAT [20]). A KAT with top (TopKAT) is a KAT K that contains a largest element $\top \in K$, that is, for all $a \in K$, $a \leq \top$. \square

3 Local Completeness Logic in KAT

We investigate how the local completeness program logic LCL [4] can be interpreted on a KAT. To achieve this, we need to address the following tasks:

- To define a notion of abstract domain of a KAT, with the aim of abstracting the set of program predicates, namely tests of a KAT;
- To establish a concrete semantics and a corresponding sound abstract semantics of programs on KATs;
- To adapt the local completeness proof system to attain valid triples on a KAT;
- To prove logical soundness and completeness w.r.t. a KAT of this new proof system.

3.1 Program Properties in KAT

Program properties can be broadly classified as intensional and extensional. The former relate to how programs are written, while the latter concern the input-output relation of the program, i.e., its strongest postcondition denotational semantics. Local completeness logic LCL relies on an abstract interpretation of programs which crucially depends on intensional properties of programs, meaning that even if two programs share the same denotation, they could well have different abstract semantics. Thus, we expect that a proper definition of abstract semantics based on a KAT model should also be intensional. Given two elements a and b of a modal bdKAT playing the role of programs, we therefore expect that their backward diamond functions might coincide, i.e. $\langle a| = \langle b|$, even if a and b encode different programs, i.e. $a \neq b$. However, as shown by the following result for the fundamental relational model of KAT, it might happen that for certain classes of KAT models the backward diamond interpretation is injective.

In order to prove such result, we require an auxiliary Lemma to provide an explicit formula for the $\langle \cdot |$ operator in the special case of full set of tests (i.e., $\text{test}(K) = \wp(\{(x, x) \mid x \in X\})$).

Lemma 3.1. *Let K be a relational KAT on a set X where $\text{test}(K) = \wp(\{(x, x) \mid x \in X\})$. Then, for all $a \in K$ and $p \in \text{test}(K)$,*

$$\langle a \mid p = \{(y, y) \mid \exists x \in X. (x, x) \in p, (x, y) \in a\}$$

Proof. We want to show that $\langle a \mid p$ exists and is equal to

$$s = \{(y, y) \mid \exists x \in X. (x, x) \in p, (x, y) \in a\}.$$

Using the characterization of $\langle a \mid p$ of Lemma 2.4, we need to show that s is the least $q \in \text{test}(K)$ satisfying $pa = paq$. In this context

$$\begin{aligned} paq &= \{(x, y) \mid (x, x) \in p, (x, y) \in a, (y, y) \in q\} \\ pa &= \{(x, y) \mid (x, x) \in p, (x, y) \in a\} \end{aligned}$$

First we show that $pa = pas$. Assume $(x, y) \in pa$, then by definition of s it holds $(y, y) \in s$, meaning that $(x, y) \in pas$. This shows $pa \leq pas$, the other inequality is immediate since $s \leq 1$ so that $pa \geq pas$.

Secondly we show that s is the least such element. Assume by contradiction that $\exists t \in \text{test}(K)$ satisfying $pa = pat$ and $t < s$. Let $(y, y) \in s \setminus t$. By definition of s exists $x \in X$ such that $(x, x) \in p$ and $(x, y) \in a$, therefore $(x, y) \notin pat$ because t is a subidentity and $(y, y) \notin t$. This contradicts the hypothesis $pa = pat$. \square

Proposition 3.2. *Let K be a relational KAT on a set X where $\text{test}(K) = \wp(\{(x, x) \mid x \in X\})$. Then, for all $a, b \in K$, $\langle a \mid = \langle b \mid \Leftrightarrow a = b$.*

Proof. The direction \Leftarrow is trivial, because $\langle \cdot \mid$ is a function on the first argument. Instead, we focus on the \Rightarrow direction.

Assume $\langle a \mid = \langle b \mid$. If $(x, y) \in a$, then by Lemma 3.1

$$\langle a \mid \{(x, x)\} = \langle b \mid \{(x, x)\} \supseteq \{(y, y)\}$$

This implies the existence of $z \in X$ such that $(z, z) \in \{(x, x)\}$ and $(z, y) \in b$, but the only possibility is $z = x$ so that $(x, y) \in b$. By symmetry we can obtain $(x, y) \in b \Rightarrow (x, y) \in a$.

Notice that we can be sure that $\{(x, x)\} \in \text{test}(K)$ because of the hypothesis $\text{test}(K) = \wp(\{(x, x) \mid x \in X\})$. \square

This means that, at least for some major KAT models, a backward diamond-based semantic interpretation of KAT elements could not preserve intensional properties, as it is the case, instead, of abstract interpretation-based semantics.

3.2 KAT Language

As a consequence of the discussion in Section 3.1, the concrete semantics cannot be directly defined on KAT elements. A solution is to define it on an inductive language. Actually, in a language two elements are equal iff they are syntactically equal, or, in

equivalent words, if the corresponding programs are written in the same way. This property makes a language an ideal basis upon which a concrete semantics can be defined, because all the intensional properties are retained.

A natural choice for defining this language of programs is the so-called *KAT language*, as originally defined by Kozen and Smith [18, Section 2.3], because it contains all and only the operators of a KAT, so that the interpretation of language terms as KAT elements is the most natural one. This language is inductively defined from two disjoint sets of primitive actions and tests through the basic elements/operations $0, 1, +, \cdot, *$ of KATs. More precisely, given a set Σ of *primitive actions* and a set B of *primitive tests* such that $\Sigma \cap B = \emptyset$, the corresponding *KAT language* $T_{\Sigma, B}$ of terms is defined as follows:

$$\begin{aligned} \text{Atom} \ni \mathbf{a} &::= a \in \Sigma \mid p \in B \\ T_{\Sigma, B} \ni \mathbf{t} &::= \mathbf{a} \mid 0 \mid 1 \mid \mathbf{t}_1 + \mathbf{t}_2 \mid \mathbf{t}_1 \cdot \mathbf{t}_2 \mid \mathbf{t}^* \end{aligned}$$

For simplicity, we assume that 0 and 1 are primitive tests in B , so that 0 and $1 \in \text{Atom}$. The notation $\text{Atom}(\mathbf{t}) \subseteq \text{Atom}$ will denote the set of atoms occurring in a term $\mathbf{t} \in T_{\Sigma, B}$. Notice that a KAT language $T_{\Sigma, B}$ is an equivalent presentation of the language of regular commands used in [4, 24] for their program logics.

Given a KAT K , an evaluation of atoms in K is a mapping $u : \text{Atom} \rightarrow K$ such that $\mathbf{p} \in B \Rightarrow u(\mathbf{p}) \in \text{test}(K)$. An evaluation u induces an interpretation of terms $\llbracket \cdot \rrbracket_u : T_{\Sigma, B} \rightarrow K$, which is inductively defined as expected:

$$\begin{aligned} \llbracket \mathbf{a} \rrbracket_u &\triangleq u(\mathbf{a}) & \llbracket \mathbf{t}_1 + \mathbf{t}_2 \rrbracket_u &\triangleq \llbracket \mathbf{t}_1 \rrbracket_u + \llbracket \mathbf{t}_2 \rrbracket_u \\ \llbracket \mathbf{t}_1 \cdot \mathbf{t}_2 \rrbracket_u &\triangleq \llbracket \mathbf{t}_1 \rrbracket_u \cdot \llbracket \mathbf{t}_2 \rrbracket_u & \llbracket \mathbf{t}^* \rrbracket_u &\triangleq \llbracket \mathbf{t} \rrbracket_u^* \end{aligned}$$

In turn, the concrete semantic function

$$\llbracket \cdot \rrbracket^K : T_{\Sigma, B} \rightarrow (\text{test}(K) \rightarrow \text{test}(K))$$

models the strongest postcondition of a program, i.e. a language term, for a given precondition, i.e. a KAT test. This is therefore defined in terms of the backward diamond of a bdKAT as follows:

$$\llbracket \mathbf{t} \rrbracket^K p \triangleq \langle \llbracket \mathbf{t} \rrbracket_u \mid p \rangle. \quad (11)$$

We will often use $\llbracket \mathbf{t} \rrbracket$ to denote a concrete semantics, by omitting the superscript K when it is clear from the context.

The construction of the concrete semantic leverages the backward-diamond (11) and the interpretation $\llbracket \cdot \rrbracket_u$, hence the properties of Proposition 2.3 can be transferred to

$\llbracket \cdot \rrbracket^K$. For completeness we summarize them in the following:

$$p \leq q \Rightarrow \llbracket \mathfrak{t} \rrbracket p \leq \llbracket \mathfrak{t} \rrbracket q \quad (12)$$

$$\llbracket \mathfrak{t}_1 + \mathfrak{t}_2 \rrbracket p = \llbracket \mathfrak{t}_1 \rrbracket p + \llbracket \mathfrak{t}_2 \rrbracket p \quad (13)$$

$$\llbracket \mathfrak{t} \rrbracket (p + q) = \llbracket \mathfrak{t} \rrbracket p + \llbracket \mathfrak{t} \rrbracket q \quad (14)$$

$$\llbracket \mathfrak{t}_1 \cdot \mathfrak{t}_2 \rrbracket p = \llbracket \mathfrak{t}_2 \rrbracket \llbracket \mathfrak{t}_1 \rrbracket p \quad (15)$$

$$\llbracket \mathfrak{t}^* \rrbracket p = \bigvee_{n \in \mathbb{N}} (\llbracket \mathfrak{t} \rrbracket)^n p \quad (16)$$

$$p + \llbracket \mathfrak{t} \rrbracket \llbracket \mathfrak{t}^* \rrbracket p \leq \llbracket \mathfrak{t}^* \rrbracket p \quad (17)$$

3.3 Kleene Abstractions

As well as an abstract domain is used in abstract interpretation for approximating store properties, i.e., sets of program stores, in our KAT model, the role of concrete domain is played by the set of tests $\text{test}(K)$ of a KAT K , ordered by the natural ordering induced by K .

Definition 3.3 (Kleene abstract domain). A poset (A, \leq_A) is a *Kleene abstract domain* of a bdKAT K if: (i) There exists a Galois Insertion $\gamma : A \rightarrow \text{test}(K)$ and $\alpha : \text{test}(K) \rightarrow A$ of the poset (A, \leq_A) into the poset $(\text{test}(K), \leq_K)$; (ii) A is countably-complete, i.e., any countable subset of A admits a lub. \square

The abstract semantic function $\llbracket \cdot \rrbracket_A^\sharp : T_{\Sigma, B} \rightarrow (A \rightarrow A)$ defines how abstract preconditions are transformed into abstract postconditions. Likewise store-based abstract interpretation, this abstract semantics is inductively defined as follows:

$$\begin{aligned} \llbracket \mathfrak{a} \rrbracket_A^\sharp p^\sharp &\triangleq \alpha(\llbracket \mathfrak{a} \rrbracket^K \gamma(p^\sharp)) & \llbracket \mathfrak{t}_1 + \mathfrak{t}_2 \rrbracket_A^\sharp p^\sharp &\triangleq \llbracket \mathfrak{t}_1 \rrbracket_A^\sharp p^\sharp + \llbracket \mathfrak{t}_2 \rrbracket_A^\sharp p^\sharp \\ \llbracket \mathfrak{t}_1 \cdot \mathfrak{t}_2 \rrbracket_A^\sharp p^\sharp &\triangleq \llbracket \mathfrak{t}_2 \rrbracket_A^\sharp (\llbracket \mathfrak{t}_1 \rrbracket_A^\sharp p^\sharp) & \llbracket \mathfrak{t}^* \rrbracket_A^\sharp p^\sharp &\triangleq \bigvee_{n \in \mathbb{N}} (\llbracket \mathfrak{t} \rrbracket_A^\sharp)^n p^\sharp \end{aligned} \quad (18)$$

Let us point that condition (ii) of Definition 3.3 ensures that the Kleene star clause in (18) is well defined.

We recall in the following Proposition few properties of Galois Insertion. Proofs and a more in depth-presentation are available in [21].

Proposition 3.4. Let $(C, \leq_C) \xleftrightarrow[\alpha]{\gamma} (A, \leq_A)$ a Galois Insertion.

$$\forall c_1, c_2 \in C. c_1 \leq_C c_2 \Rightarrow \alpha(c_1) \leq_A \alpha(c_2) \quad (19)$$

$$\forall a_1, a_2 \in A. a_1 \leq_A a_2 \Rightarrow \gamma(a_1) \leq_C \gamma(a_2) \quad (20)$$

$$\forall a_1, a_2 \in A. a_1 \neq a_2 \Rightarrow \gamma(a_1) \neq \gamma(a_2) \quad (21)$$

$$\forall c \in C. c \leq_C \gamma(\alpha(c)) \quad (22)$$

$$\gamma \circ \alpha \circ \gamma \circ \alpha = \gamma \circ \alpha \quad (23)$$

It turns out that $\llbracket \cdot \rrbracket_A^\sharp$ is a sound abstract semantics.

Theorem 3.5 (Soundness of bdKAT Abstract Semantics). *Let A be a Kleene abstraction of a CTC bdKAT K . For all $p^\sharp, q^\sharp \in A$ and $\mathfrak{t} \in T_{\Sigma, B}$:*

$$\begin{aligned} p^\sharp \leq_A q^\sharp &\Rightarrow \llbracket \mathfrak{t} \rrbracket_A^\sharp p^\sharp \leq_A \llbracket \mathfrak{t} \rrbracket_A^\sharp q^\sharp && \text{(monotonicity)} \\ \alpha(\llbracket \mathfrak{t} \rrbracket^K p) &\leq_A \llbracket \mathfrak{t} \rrbracket_A^\sharp \alpha(p) && \text{(soundness)} \end{aligned}$$

Proof. Let us prove the first implication. The proof is by induction on the structure of $\mathfrak{t} \in T_{\Sigma, B}$.

($\mathfrak{a} \in \text{Atom}$):

$$\begin{aligned} \llbracket \mathfrak{a} \rrbracket_A^\sharp p^\sharp &= \\ \alpha(\llbracket \mathfrak{a} \rrbracket \gamma(p^\sharp)) &\leq [\text{isotony of } \alpha(\cdot), \gamma(\cdot), \llbracket \cdot \rrbracket] \\ \alpha(\llbracket \mathfrak{a} \rrbracket \gamma(q^\sharp)) &= \\ \llbracket \mathfrak{a} \rrbracket_A^\sharp q^\sharp & \end{aligned}$$

($\mathfrak{t}_1 + \mathfrak{t}_2$): By induction it holds $\llbracket \mathfrak{t}_i \rrbracket_A^\sharp p^\sharp \leq \llbracket \mathfrak{t}_i \rrbracket_A^\sharp q^\sharp$ for $i \in \{1, 2\}$

$$\llbracket \mathfrak{t}_1 + \mathfrak{t}_2 \rrbracket_A^\sharp p^\sharp = \llbracket \mathfrak{t}_1 \rrbracket_A^\sharp p^\sharp + \llbracket \mathfrak{t}_2 \rrbracket_A^\sharp p^\sharp \leq \llbracket \mathfrak{t}_1 \rrbracket_A^\sharp q^\sharp + \llbracket \mathfrak{t}_2 \rrbracket_A^\sharp q^\sharp = \llbracket \mathfrak{t}_1 + \mathfrak{t}_2 \rrbracket_A^\sharp q^\sharp$$

($\mathfrak{t}_1 \cdot \mathfrak{t}_2$): By induction it holds $\llbracket \mathfrak{t}_1 \rrbracket_A^\sharp p^\sharp \leq \llbracket \mathfrak{t}_1 \rrbracket_A^\sharp q^\sharp$ and also $\llbracket \mathfrak{t}_2 \rrbracket_A^\sharp (\llbracket \mathfrak{t}_1 \rrbracket_A^\sharp p^\sharp) \leq \llbracket \mathfrak{t}_2 \rrbracket_A^\sharp (\llbracket \mathfrak{t}_1 \rrbracket_A^\sharp q^\sharp)$

$$\llbracket \mathfrak{t}_1 \cdot \mathfrak{t}_2 \rrbracket_A^\sharp p^\sharp = \llbracket \mathfrak{t}_2 \rrbracket_A^\sharp \llbracket \mathfrak{t}_1 \rrbracket_A^\sharp p^\sharp \leq \llbracket \mathfrak{t}_2 \rrbracket_A^\sharp \llbracket \mathfrak{t}_1 \rrbracket_A^\sharp q^\sharp = \llbracket \mathfrak{t}_1 \cdot \mathfrak{t}_2 \rrbracket_A^\sharp q^\sharp$$

(\mathfrak{t}_0^*): The result is a consequence of the following claim

$$\forall n \in \mathbb{N}. p^\sharp \leq q^\sharp \Rightarrow (\llbracket \mathfrak{t}_0 \rrbracket_A^\sharp)^n p^\sharp \leq (\llbracket \mathfrak{t}_0 \rrbracket_A^\sharp)^n q^\sharp \quad (24)$$

which can be shown by induction on n . $(\llbracket \mathfrak{t}_0 \rrbracket_A^\sharp)^0 p^\sharp = p^\sharp \leq q^\sharp = (\llbracket \mathfrak{t}_0 \rrbracket_A^\sharp)^0 q^\sharp$ and for the inductive case: $(\llbracket \mathfrak{t}_0 \rrbracket_A^\sharp)^{n+1} p^\sharp = \llbracket \mathfrak{t}_0 \rrbracket_A^\sharp ((\llbracket \mathfrak{t}_0 \rrbracket_A^\sharp)^n p^\sharp) \leq \llbracket \mathfrak{t}_0 \rrbracket_A^\sharp ((\llbracket \mathfrak{t}_0 \rrbracket_A^\sharp)^n q^\sharp) \leq (\llbracket \mathfrak{t}_0 \rrbracket_A^\sharp)^{n+1} q^\sharp$. Therefore

$$\llbracket \mathfrak{t}_0^* \rrbracket_A^\sharp p^\sharp = \bigvee_{n \in \mathbb{N}} (\llbracket \mathfrak{t}_0 \rrbracket_A^\sharp)^n p^\sharp \stackrel{(24)}{\leq} \bigvee_{n \in \mathbb{N}} (\llbracket \mathfrak{t}_0 \rrbracket_A^\sharp)^n q^\sharp = \llbracket \mathfrak{t}_0^* \rrbracket_A^\sharp q^\sharp$$

Similarly, the soundness is proven by induction on the structure of $\mathfrak{t} \in T_{\Sigma, B}$.

($\mathfrak{a} \in \text{Atom}$): By isotony of $\alpha(\cdot)$, $\llbracket \cdot \rrbracket$ and (22) we have

$$\alpha(\llbracket \mathfrak{a} \rrbracket p) \leq \alpha(\llbracket \mathfrak{a} \rrbracket \gamma(\alpha(p))) = \llbracket \mathfrak{a} \rrbracket_A^\sharp \alpha(p)$$

($\mathfrak{t}_1 + \mathfrak{t}_2$): By additivity of $\llbracket \cdot \rrbracket$ and $\alpha(\cdot)$ we have

$$\begin{aligned} \alpha(\llbracket \mathfrak{t}_1 + \mathfrak{t}_2 \rrbracket p) &= \alpha(\llbracket \mathfrak{t}_1 \rrbracket p + \llbracket \mathfrak{t}_2 \rrbracket p) \\ &= \alpha(\llbracket \mathfrak{t}_1 \rrbracket p) + \alpha(\llbracket \mathfrak{t}_2 \rrbracket p) \\ &\leq \llbracket \mathfrak{t}_1 \rrbracket_A^\sharp \alpha(p) + \llbracket \mathfrak{t}_2 \rrbracket_A^\sharp \alpha(p) \\ &= \llbracket \mathfrak{t}_1 + \mathfrak{t}_2 \rrbracket_A^\sharp \alpha(p) \end{aligned}$$

$(\mathfrak{t}_1 \cdot \mathfrak{t}_2)$: By isotony of $\llbracket \cdot \rrbracket_A^\sharp$ we have

$$\begin{aligned} \alpha(\llbracket \mathfrak{t}_1 \cdot \mathfrak{t}_2 \rrbracket p) &= \alpha(\llbracket \mathfrak{t}_2 \rrbracket \llbracket \mathfrak{t}_1 \rrbracket p) \\ &\leq \llbracket \mathfrak{t}_2 \rrbracket_A^\sharp \alpha(\llbracket \mathfrak{t}_1 \rrbracket p) \\ &\leq \llbracket \mathfrak{t}_2 \rrbracket_A^\sharp \llbracket \mathfrak{t}_1 \rrbracket_A^\sharp \alpha(p) \\ &= \llbracket \mathfrak{t}_1 \cdot \mathfrak{t}_2 \rrbracket_A^\sharp \alpha(p) \end{aligned}$$

(\mathfrak{t}_0^*) : In order to show the result we need an auxiliary claim:

$$\alpha(\llbracket \mathfrak{t}_0 \rrbracket^n p) \leq (\llbracket \mathfrak{t}_0 \rrbracket_A^\sharp)^n \alpha(p) \quad (25)$$

The proof is by induction on n . The base case holds as $\alpha(\llbracket \mathfrak{t}_0 \rrbracket^0 p) = \alpha(p) = (\llbracket \mathfrak{t}_0 \rrbracket_A^\sharp)^0 \alpha(p)$ and similarly the inductive case as $\alpha(\llbracket \mathfrak{t}_0 \rrbracket^{n+1} p) = \alpha(\llbracket \mathfrak{t}_0 \rrbracket^n \llbracket \mathfrak{t}_0 \rrbracket p) \leq (\llbracket \mathfrak{t}_0 \rrbracket_A^\sharp)^n \alpha(\llbracket \mathfrak{t}_0 \rrbracket p) \leq (\llbracket \mathfrak{t}_0 \rrbracket_A^\sharp)^n \llbracket \mathfrak{t}_0 \rrbracket_A^\sharp \alpha(p) = (\llbracket \mathfrak{t}_0 \rrbracket_A^\sharp)^{n+1} \alpha(p)$. Therefore we have

$$\begin{aligned} \alpha(\llbracket \mathfrak{t}_0^* \rrbracket p) &= \text{[By (16)]} \\ \alpha\left(\bigvee_{n \in \mathbb{N}} (\llbracket \mathfrak{t}_0 \rrbracket^n p)\right) &= \text{[By additivity of } \alpha(\cdot)\text{]} \\ \bigvee_{n \in \mathbb{N}} \alpha(\llbracket \mathfrak{t}_0 \rrbracket^n p) &\leq \text{[By (25)]} \\ \bigvee_{n \in \mathbb{N}} (\llbracket \mathfrak{t}_0 \rrbracket_A^\sharp)^n \alpha(p) &= \\ &= \llbracket \mathfrak{t}_0^* \rrbracket_A^\sharp \alpha(p) \end{aligned}$$

Notice that the CTC condition on K must hold in order to apply (16). \square

3.4 Local Completeness Logic on bdKAT

Given a Kleene abstraction A , we will slightly abuse notation by using $A \triangleq \gamma \circ \alpha : \text{test}(K) \rightarrow \text{test}(K)$ as a function (indeed, this is the upper closure operator on tests induced by the Galois insertion defining A). Let us recall the notions of global vs. local completeness. If $f : \text{test}(K) \rightarrow \text{test}(K)$ is any test transformer then:

- A is *globally complete* for f , denoted $\mathbb{C}^A(f)$, iff $A \circ f = A \circ f \circ A$;
- A is *locally complete* for f on a concrete test $p \in \text{test}(K)$, denoted $\mathbb{C}_p^A(f)$, iff $A \circ f(p) = A \circ f \circ A(p)$.

It is known [13] that global completeness is hard to achieve in practice, even for simple programs. Moreover, a complete and compositional (i.e., inductively defined on program structure) abstract interpretation is even harder to design [3]. This motivated to study local completeness in abstract interpretation [4] as a pragmatic and more attainable weakening of standard global completeness.

In our local completeness logic on Kleene algebra, a triple $[p] \mathfrak{t} [q]$, where p and q are tests and \mathfrak{t} is a language term, will be valid when:

$\frac{\mathbf{a} \in \Sigma \cup B \quad \mathbb{C}_p^A(\llbracket \mathbf{a} \rrbracket)}{\vdash_A^K [p] \mathbf{a} \llbracket \mathbf{a} \rrbracket p} \text{ (transfer)} \quad \frac{p' \leq p \leq A(p') \quad \vdash_A^K [p'] \mathbf{t} [q'] \quad q \leq q' \leq A(q')}{\vdash_A^K [p] \mathbf{t} [q]} \text{ (relax)}$
$\frac{\vdash_A^K [p] \mathbf{t}_1 [r] \quad \vdash_A^K [r] \mathbf{t}_2 [q]}{\vdash_A^K [p] \mathbf{t}_1 \cdot \mathbf{t}_2 [q]} \text{ (seq)} \quad \frac{\vdash_A^K [p] \mathbf{t}_1 [q_1] \quad \vdash_A^K [p] \mathbf{t}_2 [q_2]}{\vdash_A^K [p] \mathbf{t}_1 + \mathbf{t}_2 [q_1 + q_2]} \text{ (join)}$
$\frac{\vdash_A^K [p] \mathbf{t} [r] \quad \vdash_A^K [p+r] \mathbf{t}^* [q]}{\vdash_A^K [p] \mathbf{t}^* [q]} \text{ (rec)} \quad \frac{\vdash_A^K [p] \mathbf{t} [q] \quad q \leq A(p)}{\vdash_A^K [p] \mathbf{t}^* [p+q]} \text{ (iterate)}$

Fig. 1. Proof system LCK_A .

- (1) q is an under-approximation of the concrete semantics of \mathbf{t} from a precondition p ;
- (2) A is locally complete for $\llbracket \mathbf{t} \rrbracket$ on the precondition p ;
- (3) q and $\llbracket \mathbf{t} \rrbracket p$ share the same over-approximation in A .

Definition 3.6 (Triple Validity). Let K a CTC bdKAT, A be a Kleene abstraction of K , and $T_{\Sigma, B}$ be a KAT language interpreted on K . For all $p, q \in \text{test}(K)$ and $\mathbf{t} \in T_{\Sigma, B}$, a triple $[p] \mathbf{t} [q]$ is valid in A , denoted by $\models_A^K [p] \mathbf{t} [q]$, if

- (a) $q \leq_K \llbracket \mathbf{t} \rrbracket^K p$;
- (b) $\llbracket \mathbf{t} \rrbracket_A^\# \alpha(p) = \alpha(q) = \alpha(\llbracket \mathbf{t} \rrbracket^K p)$. □

The local completeness proof system in [4] can be adapted to our algebraic framework, yielding the set of rules denoted by LCK_A in Figure 1. The only syntactic difference concerns the usage of elements of $\text{test}(K)$ as pre/postconditions and the language of terms $T_{\Sigma, B}$ playing the role of programs.

It turns out that the logic LCK_A is logically sound (we use “logical” soundness to avoid overloading the soundness of abstract semantics).

Theorem 3.7 (Logical Soundness of \vdash_A^K). *If $\vdash_A^K [p] \mathbf{t} [q]$ then*

- (i) $q \leq_K \llbracket \mathbf{t} \rrbracket p$;
- (ii) $\llbracket \mathbf{t} \rrbracket_A^\# \alpha(p) = \alpha(q) = \alpha(\llbracket \mathbf{t} \rrbracket p)$.

Proof. The proof is adapted from [4, Theorem 5.5]. The first equality of (ii) is a consequence of the first one, (i) and soundness:

$$\alpha(q) \leq \alpha(\llbracket \mathbf{t} \rrbracket p) \leq \llbracket \mathbf{t} \rrbracket_A^\# \alpha(p) = \alpha(q)$$

that implies $\alpha(q) = \alpha(\llbracket \mathbf{t} \rrbracket p)$. For this reason we only need to prove $\llbracket \mathbf{t} \rrbracket_A^\# \alpha(p) = \alpha(q)$.

The proof is on the structure of the derivation tree of $\vdash_A^K [p] \mathbf{t} [q]$.

(transfer): (i) is trivial, while (ii) is a consequence of the local completeness axiom $\mathbb{C}_p^A(\llbracket \mathbf{a} \rrbracket)$, $\llbracket \mathbf{a} \rrbracket_A^\# \alpha(p) = \alpha(\llbracket \mathbf{a} \rrbracket \gamma(\alpha(p))) \stackrel{\text{LC}}{=} \alpha(\llbracket \mathbf{a} \rrbracket p) = \alpha(q)$.

(relax): By induction $q' \leq \llbracket \mathbf{t} \rrbracket p'$ so that $q \leq q' \leq \llbracket \mathbf{t} \rrbracket p' \leq \llbracket \mathbf{t} \rrbracket p$, implying (i). By hypothesis we have $p' \leq p \leq A(p')$. By isotony of $\alpha(\cdot)$ and $\gamma(\cdot)$, and (23) we have that

$A(p') \leq A(p) \leq A(A(p')) = A(p')$, which means $A(p') = A(p)$. By the same reasoning $A(q) = A(q')$. By injectivity of $\gamma(\cdot)$ we have $\alpha(p') = \alpha(p)$ and $\alpha(q') = \alpha(q)$. This proves (ii).

(seq): (i) holds by isotony of $\llbracket \cdot \rrbracket$ and by induction $q \leq \llbracket \mathfrak{t}_2 \rrbracket r \leq \llbracket \mathfrak{t}_2 \rrbracket \llbracket \mathfrak{t}_1 \rrbracket p = \llbracket \mathfrak{t}_1 \cdot \mathfrak{t}_2 \rrbracket p$, while (ii) holds by induction as $\llbracket \mathfrak{t}_1 \cdot \mathfrak{t}_2 \rrbracket_A^\sharp \alpha(p) = \llbracket \mathfrak{t}_2 \rrbracket_A^\sharp \llbracket \mathfrak{t}_1 \rrbracket_A^\sharp \alpha(p) = \llbracket \mathfrak{t}_2 \rrbracket_A^\sharp \alpha(r) = \alpha(q)$.

(join): The inductive hypothesis yields $q_1 \leq \llbracket \mathfrak{t}_1 \rrbracket p$ and $q_2 \leq \llbracket \mathfrak{t}_2 \rrbracket p$, thus by additivity of $\llbracket \cdot \rrbracket$, $q_1 + q_2 \leq \llbracket \mathfrak{t}_1 \rrbracket p + \llbracket \mathfrak{t}_2 \rrbracket p = \llbracket \mathfrak{t}_1 + \mathfrak{t}_2 \rrbracket p$, implying (i). (ii) instead can be obtained by induction and additivity of $\alpha(\cdot)$:

$$\llbracket \mathfrak{t}_1 + \mathfrak{t}_2 \rrbracket_A^\sharp \alpha(p) = \llbracket \mathfrak{t}_1 \rrbracket_A^\sharp \alpha(p) + \llbracket \mathfrak{t}_2 \rrbracket_A^\sharp \alpha(p) = \alpha(q_1) + \alpha(q_2) = \alpha(q_1 + q_2)$$

(rec): By induction we have that $r \leq \llbracket \mathfrak{t} \rrbracket p$. For any $a \in K$ it holds $a \leq a^*$ [15, Section 2.1], and by isotony of $\langle \cdot | \cdot \rangle$,

$$r \leq \llbracket \mathfrak{t} \rrbracket p = \langle \llbracket \mathfrak{t} \rrbracket_u | p \rangle \leq \langle (\llbracket \mathfrak{t} \rrbracket_u)^* | p \rangle = \langle \llbracket \mathfrak{t}^* \rrbracket_u | p \rangle = \llbracket \mathfrak{t}^* \rrbracket p$$

Moreover, for any $a \in K$ it holds $a^* a^* = a^*$ [15, Section 2.1], so that

$$\begin{aligned} \llbracket \mathfrak{t}^* \rrbracket \llbracket \mathfrak{t}^* \rrbracket p &= [\text{By (15)}] \\ \llbracket \mathfrak{t}^* \cdot \mathfrak{t}^* \rrbracket p &= \\ \langle \llbracket \mathfrak{t}^* \cdot \mathfrak{t}^* \rrbracket_u | p \rangle &= \\ \langle (\llbracket \mathfrak{t} \rrbracket_u)^* (\llbracket \mathfrak{t} \rrbracket_u)^* | p \rangle &= \\ \llbracket \mathfrak{t}^* \rrbracket p & \end{aligned}$$

By induction, it holds $q \leq \llbracket \mathfrak{t}^* \rrbracket (p+r) = \llbracket \mathfrak{t}^* \rrbracket p + \llbracket \mathfrak{t}^* \rrbracket r$, but $\llbracket \mathfrak{t}^* \rrbracket r \leq \llbracket \mathfrak{t}^* \rrbracket \llbracket \mathfrak{t}^* \rrbracket p = \llbracket \mathfrak{t}^* \rrbracket p$, thus implying (i). (ii) can be retrieved by (i), isotony of $\llbracket \cdot \rrbracket_A^\sharp$, $\alpha(\cdot)$ and soundness

$$\alpha(q) \leq \alpha(\llbracket \mathfrak{t}^* \rrbracket p) \leq \llbracket \mathfrak{t}^* \rrbracket_A^\sharp \alpha(p) \leq \llbracket \mathfrak{t}^* \rrbracket_A^\sharp \alpha(p+r) = \alpha(q)$$

(iterate): For any $a \in K$ it holds $1 \leq a^*$ and $a \leq a^*$ [15, Section 2.1], thus by isotony of $\langle \cdot | \cdot \rangle$, $q \leq \llbracket \mathfrak{t} \rrbracket p = \langle \llbracket \mathfrak{t} \rrbracket_u | p \rangle \leq \langle (\llbracket \mathfrak{t} \rrbracket_u)^* | p \rangle = \langle \llbracket \mathfrak{t}^* \rrbracket_u | p \rangle \leq \llbracket \mathfrak{t}^* \rrbracket p$ and by (5) and isotony of $\langle \cdot | \cdot \rangle$, $p = p \cdot 1 = \langle 1 | p \rangle \leq \langle (\llbracket \mathfrak{t} \rrbracket_u)^* | p \rangle = \llbracket \mathfrak{t}^* \rrbracket p$, thus implying (i) $p + q \leq \llbracket \mathfrak{t}^* \rrbracket p$. To show (ii) we need the following preliminary fact:

$$\llbracket \mathfrak{t} \rrbracket_A^\sharp \alpha(p) \leq \alpha(p) \Rightarrow \bigvee_{n \in \mathbb{N}} (\llbracket \mathfrak{t} \rrbracket_A^\sharp)^n \alpha(p) = \alpha(p) \quad (26)$$

First, let us prove by induction on n that $\alpha(p)$ is an upper-bound of the elements of the disjunction, i.e., $\forall n \in \mathbb{N}. (\llbracket \mathfrak{t} \rrbracket_A^\sharp)^n \alpha(p) \leq \alpha(p)$. The base case $n = 0$ is true since $(\llbracket \mathfrak{t} \rrbracket_A^\sharp)^0 \alpha(p) = \alpha(p) \leq \alpha(p)$, while the inductive case holds by isotony of $\llbracket \cdot \rrbracket_A^\sharp$, $(\llbracket \mathfrak{t} \rrbracket_A^\sharp)^{n+1} \alpha(p) = (\llbracket \mathfrak{t} \rrbracket_A^\sharp)^n \llbracket \mathfrak{t} \rrbracket_A^\sharp \alpha(p) \leq (\llbracket \mathfrak{t} \rrbracket_A^\sharp)^n \alpha(p) \leq \alpha(p)$. Notice that $\alpha(p)$ is also the least upper bound because $(\llbracket \mathfrak{t} \rrbracket_A^\sharp)^0 \alpha(p) = \alpha(p)$. The hypothesis $q \leq A(p)$ implies $\alpha(q) \leq \alpha(p)$ by isotony of $\alpha(\cdot)$ and $\gamma(\cdot)$, (21) and (23). Notice that $\llbracket \mathfrak{t} \rrbracket_A^\sharp \alpha(p) = \alpha(q) \leq \alpha(p)$, meaning that we can apply (26):

$$\llbracket \mathfrak{t}^* \rrbracket_A^\sharp \alpha(p) = \bigvee_{n \in \mathbb{N}} (\llbracket \mathfrak{t} \rrbracket_A^\sharp)^n \alpha(p) = \alpha(p) = \alpha(p) + \alpha(q) = \alpha(p+q). \quad \square$$

Analogously to what happens for LCL, we can prove that LCK_A is logically complete under these two additional hypotheses:

(A) The following infinitary rule is added to LCK_A :

$$\frac{\forall n \in \mathbb{N}. \vdash_A^K [p_n] \mathfrak{t} [p_{n+1}]}{\vdash_A^K [p_0] \mathfrak{t}^* [\bigvee_{n \in \mathbb{N}} p_n]} \text{ (limit)}$$

Let us point out that the lub $\bigvee_{n \in \mathbb{N}} p_n$ always exists in K , as a consequence of the CTC requirement on K .

(B) The concrete semantics of the primitive actions and tests occurring in the program are globally complete.

It turns out that the rule (limit) preserves logical soundness:

Lemma 3.8. *With the hypothesis of Theorem 3.7, the rule (limit) preserves logical soundness.*

Proof. Let's assume that (limit) proves a triple $[p_0] \mathfrak{t}^* [\bigvee_{n \in \mathbb{N}} p_n]$. The conditions (i) is $\bigvee_{n \in \mathbb{N}} p_n \leq \llbracket \mathfrak{t}^* \rrbracket p_0$, which is equivalent to $\forall n \in \mathbb{N}. p_n \leq \llbracket \mathfrak{t}^* \rrbracket p_0$. The latter can be shown by induction on n . By (17), $p_0 \leq p_0 + \llbracket \mathfrak{t} \rrbracket \llbracket \mathfrak{t}^* \rrbracket p_0 \leq \llbracket \mathfrak{t}^* \rrbracket p_0$. Similarly for the inductive case $p_{n+1} \leq \llbracket \mathfrak{t} \rrbracket p_n \leq \llbracket \mathfrak{t} \rrbracket \llbracket \mathfrak{t}^* \rrbracket p_0 \leq p_0 + \llbracket \mathfrak{t} \rrbracket \llbracket \mathfrak{t}^* \rrbracket p_0 \leq \llbracket \mathfrak{t}^* \rrbracket p_0$, where the first inequality holds by structural induction, i.e., $\vdash_A^K [p_n] \mathfrak{t} [p_{n+1}]$ implies $\models_A^K [p_n] \mathfrak{t} [p_{n+1}]$. To show (ii) we need the auxiliary claim $\forall n \in \mathbb{N}. (\llbracket \mathfrak{t} \rrbracket_A^\sharp)^n \alpha(p_0) = \alpha(p_n)$. That can be proved by induction on n . The base case is trivially true as $(\llbracket \mathfrak{t} \rrbracket_A^\sharp)^0 \alpha(p_0) = \alpha(p_0)$. While the inductive case holds by structural induction, i.e., $\llbracket \mathfrak{t} \rrbracket_A^\sharp \alpha(p_n) = \alpha(p_{n+1})$, as $(\llbracket \mathfrak{t} \rrbracket_A^\sharp)^{n+1} \alpha(p_0) = \llbracket \mathfrak{t} \rrbracket_A^\sharp (\llbracket \mathfrak{t} \rrbracket_A^\sharp)^n \alpha(p_0) = \llbracket \mathfrak{t} \rrbracket_A^\sharp \alpha(p_n) = \alpha(p_{n+1})$. Therefore by additivity of $\alpha(\cdot)$,

$$\llbracket \mathfrak{t}^* \rrbracket_A^\sharp \alpha(p_0) = \bigvee_{n \in \mathbb{N}} (\llbracket \mathfrak{t} \rrbracket_A^\sharp)^n \alpha(p_0) = \bigvee_{n \in \mathbb{N}} \alpha(p_n) = \alpha(\bigvee_{n \in \mathbb{N}} p_n). \quad \square$$

The condition (B) entails global completeness:

Lemma 3.9. *Let K a backward-diamond CTC Kleene algebra, A a Kleene abstract domain on K and $T_{\Sigma, B}$ a KAT language on K . For any $\mathfrak{t} \in T_{\Sigma, B}$ and $p \in \text{test}(K)$ we have*

$$(\forall \mathfrak{a} \in \text{Atom}(\mathfrak{t}). \mathbb{C}^A(\llbracket \mathfrak{a} \rrbracket)) \Rightarrow \llbracket \mathfrak{t} \rrbracket_A^\sharp \alpha(p) = \alpha(\llbracket \mathfrak{t} \rrbracket p)$$

Proof. The proof is on the structure of \mathfrak{t} :

($\mathfrak{a} \in \text{Atom}$):

$$\begin{aligned} \llbracket \mathfrak{a} \rrbracket_A^\sharp \circ \alpha &= \\ \alpha \circ \llbracket \mathfrak{a} \rrbracket \circ \gamma \circ \alpha &= [\text{By } \mathbb{C}^A(\llbracket \mathfrak{a} \rrbracket), (21)] \\ \alpha \circ \llbracket \mathfrak{a} \rrbracket & \end{aligned}$$

$(\mathfrak{t}_1 + \mathfrak{t}_2)$: Let us denote with p a generic test.

$$\begin{aligned} \llbracket \mathfrak{t}_1 + \mathfrak{t}_2 \rrbracket_A^\sharp \alpha(p) &= \\ \llbracket \mathfrak{t}_1 \rrbracket_A^\sharp \alpha(p) + \llbracket \mathfrak{t}_2 \rrbracket_A^\sharp \alpha(p) &= \text{[By induction]} \\ \alpha(\llbracket \mathfrak{t}_1 \rrbracket p) + \alpha(\llbracket \mathfrak{t}_2 \rrbracket p) &= \text{[By additivity of } \alpha(\cdot), \llbracket \cdot \rrbracket \text{]} \\ \alpha(\llbracket \mathfrak{t}_1 + \mathfrak{t}_2 \rrbracket p) & \end{aligned}$$

$(\mathfrak{t}_1 \cdot \mathfrak{t}_2)$: Let us denote with p a generic test.

$$\begin{aligned} \llbracket \mathfrak{t}_1 \cdot \mathfrak{t}_2 \rrbracket_A^\sharp \alpha(p) &= \\ \llbracket \mathfrak{t}_2 \rrbracket_A^\sharp \llbracket \mathfrak{t}_1 \rrbracket_A^\sharp \alpha(p) &= \text{[By induction]} \\ \llbracket \mathfrak{t}_2 \rrbracket_A^\sharp \alpha(\llbracket \mathfrak{t}_1 \rrbracket p) &= \text{[By induction]} \\ \alpha(\llbracket \mathfrak{t}_2 \rrbracket \llbracket \mathfrak{t}_1 \rrbracket p) &= \text{[By (15)]} \\ \alpha(\llbracket \mathfrak{t}_1 \cdot \mathfrak{t}_2 \rrbracket p) & \end{aligned}$$

(\mathfrak{t}_0^*) : The result is a consequence of the following claim

$$\forall n \in \mathbb{N}. (\llbracket \mathfrak{t}_0 \rrbracket_A^\sharp)^n \alpha(p) = \alpha((\llbracket \mathfrak{t}_0 \rrbracket)^n p) \quad (27)$$

that can be proved by induction on n . The base case is $(\llbracket \mathfrak{t}_0 \rrbracket_A^\sharp)^0 \alpha(p) = \alpha(p) = \alpha((\llbracket \mathfrak{t}_0 \rrbracket)^0 p)$. While the inductive case can be proved as follows

$$(\llbracket \mathfrak{t}_0 \rrbracket_A^\sharp)^{n+1} \alpha(p) = \llbracket \mathfrak{t}_0 \rrbracket_A^\sharp (\llbracket \mathfrak{t}_0 \rrbracket_A^\sharp)^n \alpha(p) = \llbracket \mathfrak{t}_0 \rrbracket_A^\sharp \alpha((\llbracket \mathfrak{t}_0 \rrbracket)^n p) = \alpha((\llbracket \mathfrak{t}_0 \rrbracket)^{n+1} p)$$

Therefore,

$$\begin{aligned} \llbracket \mathfrak{t}_0^* \rrbracket_A^\sharp \alpha(p) &= \\ \bigvee_{n \in \mathbb{N}} (\llbracket \mathfrak{t}_0 \rrbracket_A^\sharp)^n \alpha(p) &= \text{[By (27)]} \\ \bigvee_{n \in \mathbb{N}} \alpha((\llbracket \mathfrak{t}_0 \rrbracket)^n p) &= \text{[By additivity of } \alpha(\cdot) \text{]} \\ \alpha\left(\bigvee_{n \in \mathbb{N}} (\llbracket \mathfrak{t}_0 \rrbracket)^n p\right) &= \text{[By (16)]} \\ \alpha(\llbracket \mathfrak{t}_0^* \rrbracket p) & \quad \square \end{aligned}$$

Theorem 3.10 (Logical Completeness of \vdash_A^K). *Assume that conditions (A) and (B) hold. If $\models_A^K [p] \vDash [q]$ then $\vdash_A^K [p] \vDash [q]$.*

Proof. The proof is organized in two parts. In the first part we prove that $\vdash_A^K [p] \vDash \llbracket \llbracket \mathfrak{t} \rrbracket p \rrbracket$. In the second one instead we prove

$$\frac{\vdash_A^K [p] \vDash \llbracket \llbracket \mathfrak{t} \rrbracket p \rrbracket}{\vdash_A^K [p] \vDash [q]}$$

1) The proof is by induction on the structure of $\mathfrak{t} \in T_{\Sigma, B}$:

($\mathfrak{a} \in \text{Atom}$): By the global completeness hypothesis we have $\mathbb{C}_p^A(\llbracket \mathfrak{a} \rrbracket)$, hence (transfer) yields $\vdash_A^{\kappa} [p] \mathfrak{a} \llbracket \llbracket \mathfrak{a} \rrbracket p \rrbracket$.

($\mathfrak{t}_1 + \mathfrak{t}_2$): Let $q_i = \llbracket \mathfrak{t}_i \rrbracket p$, and by Lemma 3.9 $\llbracket \mathfrak{t}_i \rrbracket_A^{\#} \alpha(p) = \alpha(q_i)$, which means $\models_A^{\kappa} [p] \mathfrak{t}_i [q_i]$, thus by induction $\vdash_A^{\kappa} [p] \mathfrak{t}_i [q_i]$. Notice that by additivity of $\llbracket \cdot \rrbracket$, $q_1 + q_2 = \llbracket \mathfrak{t}_1 \rrbracket p + \llbracket \mathfrak{t}_2 \rrbracket p = \llbracket \mathfrak{t}_1 + \mathfrak{t}_2 \rrbracket p$. Finally, (join) yields the result $\vdash_A^{\kappa} [p] \mathfrak{t}_1 + \mathfrak{t}_2 \llbracket \llbracket \mathfrak{t}_1 + \mathfrak{t}_2 \rrbracket p \rrbracket$

($\mathfrak{t}_1 \cdot \mathfrak{t}_2$): Let $q_1 = \llbracket \mathfrak{t}_1 \rrbracket p$ and $q_2 = \llbracket \mathfrak{t}_2 \rrbracket q_1$. By Lemma 3.9 $\llbracket \mathfrak{t}_1 \rrbracket_A^{\#} \alpha(p) = \alpha(\llbracket \mathfrak{t}_1 \rrbracket p)$ and $\llbracket \mathfrak{t}_2 \rrbracket_A^{\#} \alpha(q_1) = \alpha(\llbracket \mathfrak{t}_2 \rrbracket q_1)$, which means $\models_A^{\kappa} [p] \mathfrak{t}_1 [q_1]$ and $\models_A^{\kappa} [q_1] \mathfrak{t}_2 [q_2]$, and by induction $\vdash_A^{\kappa} [p] \mathfrak{t}_1 [q_1]$ and $\vdash_A^{\kappa} [q_1] \mathfrak{t}_2 [q_2]$. Finally, (seq) yields the result $\vdash_A^{\kappa} [p] \mathfrak{t}_1 \cdot \mathfrak{t}_2 \llbracket \llbracket \mathfrak{t}_1 \cdot \mathfrak{t}_2 \rrbracket p \rrbracket$, where by (15) $q_2 = \llbracket \mathfrak{t}_2 \rrbracket \llbracket \mathfrak{t}_1 \rrbracket p = \llbracket \mathfrak{t}_1 \cdot \mathfrak{t}_2 \rrbracket p$.

(\mathfrak{t}_0^*): Let $p_n = (\llbracket \mathfrak{t}_0 \rrbracket)^n p$. As preliminary step let us prove that $\models_A^{\kappa} [p_n] \mathfrak{t}_0 [p_{n+1}]$ for any n . (i) holds because $p_{n+1} = \llbracket \mathfrak{t}_0 \rrbracket p_n$, while (ii) is immediate by Lemma 3.9, $\llbracket \mathfrak{t}_0 \rrbracket_A^{\#} \alpha(p_n) = \alpha(\llbracket \mathfrak{t}_0 \rrbracket p_n) = \alpha(p_{n+1})$. By induction we have $\forall n \in \mathbb{N}. \vdash_A^{\kappa} [p_n] \mathfrak{t}_0 [p_{n+1}]$, thus (limit) yields the result $\vdash_A^{\kappa} [p] \mathfrak{t}_0^* \llbracket \llbracket \mathfrak{t}_0^* \rrbracket p \rrbracket$ (notice that $p = (\llbracket \mathfrak{t}_0 \rrbracket)^0 p = p_0$ and by (16) $\llbracket \mathfrak{t}_0^* \rrbracket p = \bigvee_{n \in \mathbb{N}} (\llbracket \mathfrak{t}_0 \rrbracket)^n p = \bigvee_{n \in \mathbb{N}} p_n$).

2) By hypothesis $\models_A^{\kappa} [p] \mathfrak{t} [q]$, thus $q \leq \llbracket \mathfrak{t} \rrbracket p$ and also $A(q) = A(\llbracket \mathfrak{t} \rrbracket p)$ so that we can use (relax) to obtain the result: $\vdash_A^{\kappa} [p] \mathfrak{t} [q]$. \square

Summing up, this shows that the local completeness logic LCL introduced in [4] can be made *fully algebraic* by means of a natural interpretation on modal KATs with a backward diamond operator, still preserving its logical soundness and completeness, which are proved by using just the algebraic axioms of this class of KATs, thus proving that there is no need to leverage particular semantic properties of programs to reason on their local completeness.

3.5 An Example of a Language-Theoretic KAT

To give an example digressing from programs and showing the generality of the KAT-based approach, we describe a language-theoretic model of Kleene algebra, introduced by Kozen and Smith [18, Section 3].

Let $\Sigma = \{\mathfrak{u}\}$ and $B = \{\mathfrak{b}_1, \mathfrak{b}_2\}$ be, respectively, the sets of primitive actions and tests. An *atom* is a string $c_1 c_2$, where $c_i \in \{\mathfrak{b}_i, \overline{\mathfrak{b}_i}\}$. If $c_i = \mathfrak{b}_i$, where $i \in \{1, 2\}$, then \mathfrak{b}_i appears positively in the atom $c_1 c_2$, while if $c_i = \overline{\mathfrak{b}_i}$ it appears negatively. A *guarded string* is either a single atom or a string $\alpha_0 \mathfrak{a}_1 \alpha_1 \dots \mathfrak{a}_n \alpha_n$, where α_i are atoms and $\mathfrak{a}_i \in \Sigma$. If we are only interested in the first (last) atom of a guarded string $\alpha \mathfrak{a}_1 \alpha_1 \dots \mathfrak{a}_n \beta$, we may refer to it through the syntax $\alpha x (\beta)$. Concatenation of guarded strings is given by a coalesced product operation \diamond , which is partially defined as follows:

$$x\alpha \diamond \beta y \triangleq \begin{cases} x\alpha y & \text{if } \alpha = \beta \\ \text{undefined} & \text{otherwise} \end{cases}$$

The elements of this KAT \mathcal{G} are sets of guarded strings. Thus, $+$ is set union, the product is defined as pointwise coalesced product:

$$A \cdot B \triangleq \{x \diamond y \mid x \in A, y \in B\},$$

while the Kleene iteration is: $A^* \triangleq \bigcup_{n \in \mathbb{N}} A^n$. The product identity $1_{\mathcal{G}}$ corresponds to the whole set of atoms $\{\mathbf{b}_1 \mathbf{b}_2, \mathbf{b}_1 \bar{\mathbf{b}}_2, \bar{\mathbf{b}}_1 \mathbf{b}_2, \bar{\mathbf{b}}_1 \bar{\mathbf{b}}_2\}$, while $0_{\mathcal{G}}$ is the empty set. The set of tests is $\text{test}(\mathcal{G}) \triangleq \wp(1_{\mathcal{G}})$.

It turns out that \mathcal{G} is a bdKAT, whose backward diamond is as follows: for all $a \in \mathcal{G}$ and $p \in \text{test}(\mathcal{G})$,

$$\langle a | p \triangleq \{\beta \mid x\beta \in pa\}. \quad (28)$$

Proof. Let $r = \{\beta \mid x\beta \in pa\}$. By Lemma 2.4, it is enough to show that r is the least $q \in \text{test}(\mathcal{G})$ satisfying $pa = paq$.

Let $x\beta \in pa$. By definition of r , $\beta \in r$ means that $x\beta \diamond \beta = x\beta$ is contained in par . This therefore means that $pa \leq par$. The opposite inequality is trivial since r is a test, hence $r \leq 1_{\mathcal{G}}$, and by monotonicity of \cdot , we have that $pa \geq par$, thus implying $pa = par$.

Assume now, by contradiction, that there exists $t \in \text{test}(\mathcal{G})$ such that $pa = pat$ and $t < r$. This means that there is at least an atom β in r which is not in t . By definition of r , there is a guarded string $x\beta \in pa$. Since $pa = pat$, the last atom of all the guarded strings in pa must be in t , but this does not hold for $x\beta$ as $\beta \notin t$. \square

Let us consider the evaluation function $G : \text{Atom} \rightarrow \mathcal{G}$ as defined in [18]:

$$\begin{aligned} G(\mathbf{a}) &\triangleq \{\alpha \mathbf{a} \beta \mid \alpha, \beta \in 1_{\mathcal{G}}\}, \\ G(\mathbf{b}) &\triangleq \{\alpha \in 1_{\mathcal{G}} \mid \mathbf{b} \text{ appears positively in } \alpha\}. \end{aligned}$$

We consider the abstract domain $A \triangleq \{\top, e, o, \perp\}$ determined by the following abstraction $\alpha : \text{test}(\mathcal{G}) \rightarrow A$ and concretization $\gamma : A \rightarrow \text{test}(\mathcal{G})$ maps:

$$\alpha(p) \triangleq \begin{cases} \perp & \text{if } p = \emptyset \\ e & \text{if } \emptyset \subsetneq p \subseteq \{\mathbf{b}_1 \mathbf{b}_2, \bar{\mathbf{b}}_1 \bar{\mathbf{b}}_2\} \\ o & \text{if } \emptyset \subsetneq p \subseteq \{\bar{\mathbf{b}}_1 \mathbf{b}_2, \mathbf{b}_1 \bar{\mathbf{b}}_2\} \\ \top & \text{otherwise} \end{cases} \quad \gamma(p^\#) \triangleq \begin{cases} \emptyset & \text{if } p^\# = \perp \\ \{\mathbf{b}_1 \mathbf{b}_2, \bar{\mathbf{b}}_1 \bar{\mathbf{b}}_2\} & \text{if } p^\# = e \\ \{\bar{\mathbf{b}}_1 \mathbf{b}_2, \mathbf{b}_1 \bar{\mathbf{b}}_2\} & \text{if } p^\# = o \\ 1_{\mathcal{G}} & \text{if } p^\# = \top \end{cases}$$

By counting, in an atom, the number of primitive tests that appear positively we obtain an integer that may be even or odd. Hence, this abstract domain A represents the property of being even e or odd o of all the atoms in a test $p \in \text{test}(\mathcal{G})$.

By using our logic LCK, we study the correctness of the program $\mathbf{r} \triangleq (\mathbf{u} \cdot \mathbf{b}_1)^* \in \mathcal{G}$, assuming a precondition $p \triangleq \{\mathbf{b}_1 \mathbf{b}_2, \bar{\mathbf{b}}_1 \bar{\mathbf{b}}_2\} \in \text{test}(\mathcal{G})$ and a specification $\text{Spec} \triangleq p = \gamma(e)$. Let us define two auxiliary tests: $q \triangleq \{\mathbf{b}_1 \mathbf{b}_2, \mathbf{b}_1 \bar{\mathbf{b}}_2\}$, $s \triangleq \{\mathbf{b}_1 \mathbf{b}_2, \mathbf{b}_1 \bar{\mathbf{b}}_2, \bar{\mathbf{b}}_1 \bar{\mathbf{b}}_2\}$. Using the formula (28) we can easily check the following local completeness equations:

$$\begin{aligned} \alpha(\llbracket \mathbf{u} \rrbracket A(s)) &= \alpha(\llbracket \mathbf{u} \rrbracket 1_{\mathcal{G}}) = \alpha(1_{\mathcal{G}}) = \top = \alpha(1_{\mathcal{G}}) = \alpha(\llbracket \mathbf{u} \rrbracket s) &\Rightarrow \mathbb{C}_s^A(\llbracket \mathbf{u} \rrbracket) \\ \alpha(\llbracket \mathbf{u} \rrbracket A(p)) &= \alpha(\llbracket \mathbf{u} \rrbracket p) = \alpha(1_{\mathcal{G}}) = \top = \alpha(1_{\mathcal{G}}) = \alpha(\llbracket \mathbf{u} \rrbracket p) &\Rightarrow \mathbb{C}_p^A(\llbracket \mathbf{u} \rrbracket) \\ \alpha(\llbracket \mathbf{b}_1 \rrbracket A(1_{\mathcal{G}})) &= \alpha(\llbracket \mathbf{b}_1 \rrbracket 1_{\mathcal{G}}) = \alpha(q) = \top = \alpha(q) = \alpha(\llbracket \mathbf{b}_1 \rrbracket 1_{\mathcal{G}}) &\Rightarrow \mathbb{C}_{1_{\mathcal{G}}}^A(\llbracket \mathbf{b}_1 \rrbracket) \end{aligned}$$

Therefore, we have the following derivation in LCK_A of the triple $[p] \mathbf{r} [s]$:

$$\frac{\frac{\frac{\mathbb{C}_p^A(\llbracket \mathbf{u} \rrbracket)}{\vdash_A^{\mathbf{K}} [p] \mathbf{u} [1_{\mathcal{G}}]} \text{ (transfer)}}{\vdash_A^{\mathbf{K}} [p] \mathbf{u} \cdot \mathbf{b}_1 [q]} \text{ (seq)}}{\vdash_A^{\mathbf{K}} [p] (\mathbf{u} \cdot \mathbf{b}_1)^* [s]} \text{ (rec)} \quad \frac{\frac{\frac{\frac{\mathbb{C}_s^A(\llbracket \mathbf{u} \rrbracket)}{\vdash_A^{\mathbf{K}} [s] \mathbf{u} [1_{\mathcal{G}}]} \text{ (transfer)}}{\vdash_A^{\mathbf{K}} [s] \mathbf{u} \cdot \mathbf{b}_1 [q]} \text{ (seq)}}{\vdash_A^{\mathbf{K}} [s] (\mathbf{u} \cdot \mathbf{b}_1)^* [s]} \text{ (rec)}}{\vdash_A^{\mathbf{K}} [s] \mathbf{u} \cdot \mathbf{b}_1 [q]} \text{ (seq)} \quad \frac{\frac{\mathbb{C}_{1_{\mathcal{G}}}^A(\llbracket \mathbf{b}_1 \rrbracket)}{\vdash_A^{\mathbf{K}} [1_{\mathcal{G}}] \mathbf{b}_1 [q]} \text{ (transfer)}}{\vdash_A^{\mathbf{K}} [s] \mathbf{u} \cdot \mathbf{b}_1 [q]} \text{ (seq)} \quad q \leq A(s) \text{ (iterate)}$$

$\frac{a \in \text{Atom}}{\vdash_{\text{UL}} [p] a \llbracket [a] p \rrbracket} \text{ (transfer)}$	
$\frac{}{\vdash_{\text{UL}} [p] \mathbf{t} [0]} \text{ (empty)}$	$\frac{p' \leq p \quad \vdash_{\text{UL}} [p'] \mathbf{t} [q'] \quad q \leq q'}{\vdash_{\text{UL}} [p] \mathbf{t} [q]} \text{ (consequence)}$
$\frac{\vdash_{\text{UL}} [p_1] \mathbf{t} [q_1] \quad \vdash_{\text{UL}} [p_2] \mathbf{t} [q_2]}{\vdash_{\text{UL}} [p_1 + p_2] \mathbf{t} [q_1 + q_2]} \text{ (disj)}$	$\frac{\vdash_{\text{UL}} [p] \mathbf{t}_1 [r] \quad \vdash_{\text{UL}} [r] \mathbf{t}_2 [q]}{\vdash_{\text{UL}} [p] \mathbf{t}_1 \cdot \mathbf{t}_2 [q]} \text{ (seq)}$
$\frac{}{\vdash_{\text{UL}} [p] \mathbf{t}^* [p]} \text{ (iterate zero)}$	$\frac{\vdash_{\text{UL}} [p] \mathbf{t}^* \cdot \mathbf{t} [q]}{\vdash_{\text{UL}} [p] \mathbf{t}^* [q]} \text{ (iterate non-zero)}$
$\frac{\forall n \in \mathbb{N}. \vdash_{\text{UL}} [p_n] \mathbf{t} [p_{n+1}]}{\vdash_{\text{UL}} [p_0] \mathbf{t}^* [\bigvee_{n \in \mathbb{N}} p_n]} \text{ (back-v)}$	$\frac{\vdash_{\text{UL}} [p] \mathbf{t}_i [q], \text{ with } i = 1 \text{ or } i = 2}{\vdash_{\text{UL}} [p] \mathbf{t}_1 + \mathbf{t}_2 [q]} \text{ (choice)}$

Fig. 2. Proof System UL.

Here, in accordance with the soundness Theorem 3.7, we have that $s \subseteq \llbracket \mathbf{r} \rrbracket p \subseteq A(s)$. Observe that $A(s) \not\subseteq \text{Spec}$, meaning that an abstract interpretation-based analysis fails to prove that the program \mathbf{r} is correct for Spec . However, unlike conventional abstract interpretation, LCK_A is capable to show that $s \setminus \text{Spec} = \{\mathbf{b}_1 \bar{\mathbf{b}}_2\}$ is indeed a true alert, meaning that the program \mathbf{r} is really incorrect and the failure to prove its correctness was not due to a false alarm.

3.6 Under-Approximation Logic

O’Hearn [24] incorrectness logic (IL) establishes two main novelties w.r.t. the seminal Hoare logic of program correctness [14]: (1) a valid postcondition of an incorrectness triple for a program P is an *under-approximation* of the strongest postcondition of P , rather than an over-approximation of Hoare logic; (2) incorrectness triples feature two postconditions: one corresponding to a “normal” program termination and one corresponding to an erroneous termination. Even if IL was originally defined with both those features, we first neglect the second one — i.e., we consider “normal” termination only — and we refer to the resulting program logic as Under-approximation Logic (UL). For the sake of clarity, Figure 2 recalls the UL proof system, adapted to our algebraic framework. We only focus on the “propositional” fragment of this logic, meaning that the roles of all the special program commands (i.e., `error`, `assume`, `skip`, `nondet` used in [24]) and variable manipulations commands of incorrectness logic are played by some corresponding elements in Atom . Hence, for all of them, the single rule (transfer) is unifying and enough for our purposes.

Analogously to what has been proved in [4, Section 6] for LCL, it turns out that the trivial abstraction, i.e., the abstract domain $A_{tr} \triangleq \{\top\}$ that approximates all the concrete elements to a single abstract element \top , allows us to show that the instantiation $\text{LCK}_{A_{tr}}$, with the additional rule (limit), boils down to UL, namely, our LCK logic generalizes UL, even when both are interpreted on KATs.

LCK _{A_{tr}}	UL
transfer	transfer
relax	consequence
seq	seq
limit	back-v

Table 1. Correspondence between the rules of LCK_{A_{tr}} and UL.

Theorem 3.11 (LCK_{A_{tr}} \equiv UL). *Let K be a CTC bdkAT. Assume that LCK_{A_{tr}} includes the rule (limit). For any $p, q \in \text{test}(K)$ and $\mathfrak{t} \in T_{\Sigma, B}$:*

$$\vdash_{A_{tr}}^K [p] \mathfrak{t} [q] \Leftrightarrow \vdash_{UL} [p] \mathfrak{t} [q].$$

Proof. Notice that in the LCK_{A_{tr}} proof system, the premise $\mathbb{C}_p^{A_{tr}}(\llbracket \mathbf{a} \rrbracket)$ of (transfer) is always true because $A_{tr} \circ \llbracket \mathbf{a} \rrbracket p = 1 = A_{tr} \circ \llbracket \mathbf{a} \rrbracket \circ A_{tr}(p)$ and similarly the conditions $p \leq A_{tr}(p')$ and $q' \leq A_{tr}(q')$ of (relax) are always true because $A_{tr}(p) = 1$ which is the greatest element of $\text{test}(K)$.

Moreover, there is a strong correspondence between the two proof systems: in particular the rules of Table 1 coincide.

\Rightarrow) As preliminary step, notice that derivation trees built by the proof of Theorem 3.10 do not require the rules (rec) and (iterate), thus by soundness and completeness we can build a derivation tree of $\vdash_{A_{tr}}^K [p] \mathfrak{t} [q]$ which does not contain such rules.

The proof is by induction on the derivation tree of $\vdash_{A_{tr}}^K [p] \mathfrak{t} [q]$. The rules of Table 1 are immediate. The only rule left to prove is (join). By induction we have $\vdash_{UL} [p] \mathfrak{t}_1 [q_1]$ and $\vdash_{UL} [p] \mathfrak{t}_2 [q_2]$ and we need to show $\vdash_{UL} [p] \mathfrak{t}_1 + \mathfrak{t}_2 [q_1 + q_2]$:

$$\frac{\frac{\vdash_{UL} [p] \mathfrak{t}_1 [q_1]}{\vdash_{UL} [p] \mathfrak{t}_1 + \mathfrak{t}_2 [q_1]} \text{ (choice)} \quad \frac{\vdash_{UL} [p] \mathfrak{t}_2 [q_2]}{\vdash_{UL} [p] \mathfrak{t}_1 + \mathfrak{t}_2 [q_2]} \text{ (choice)}}{\vdash_{UL} [p] \mathfrak{t}_1 + \mathfrak{t}_2 [q_1 + q_2]} \text{ (disj)}$$

\Leftarrow) The proof is by induction on the derivation tree of $\vdash_{UL} [p] \mathfrak{t} [q]$. As in the previous case, the rules of Table 1 are immediate. The rules that are left to prove are (empty), (choice), (disj), (iterate zero) and (iterate non-zero).

In order to carry out the derivation in those cases we exploit the completeness of LCK. To do so we only need to verify condition (i), because (ii) is trivially true: $\llbracket \mathfrak{t} \rrbracket_A^\# \alpha(p) = \top = \alpha(q) = \top = \alpha(\llbracket \mathfrak{t} \rrbracket p)$ for any $p, q \in \text{test}(K)$ and $\mathfrak{t} \in T_{\Sigma, B}$.

(empty): $\vdash_{A_{tr}}^K [p] \mathfrak{t} [0]$ is trivial since $0 \leq \llbracket \mathfrak{t} \rrbracket p$ is always true, therefore by completeness of LCK_{A_{tr}}, $\vdash_{A_{tr}}^K [p] \mathfrak{t} [0]$.

(choice): By induction we have, without loss of generality, $\vdash_{A_{tr}}^K [p] \mathfrak{t}_1 [q]$ and we need to show $\vdash_{A_{tr}}^K [p] \mathfrak{t}_1 + \mathfrak{t}_2 [q]$. Notice that for any $\mathfrak{t}_2 \in T_{\Sigma, B}$, $\vdash_{A_{tr}}^K [p] \mathfrak{t}_2 [0]$ because (i) is trivially true: $0 \leq \llbracket \mathfrak{t}_2 \rrbracket p$, therefore by completeness of LCK_{A_{tr}} $\vdash_{A_{tr}}^K [p] \mathfrak{t}_2 [0]$. Finally, (join) yields the result $\vdash_{A_{tr}}^K [p] \mathfrak{t}_1 + \mathfrak{t}_2 [q]$.

(disj): By induction we have $\vdash_{A_{tr}}^K [p_1] \mathfrak{t} [q_1]$ and $\vdash_{A_{tr}}^K [p_2] \mathfrak{t} [q_2]$ and we need to show $\vdash_{A_{tr}}^K [p_1 + p_2] \mathfrak{t} [q_1 + q_2]$. By soundness we have $\models_{A_{tr}}^K [p_1] \mathfrak{t} [q_1]$ and $\models_{A_{tr}}^K [p_2] \mathfrak{t} [q_2]$,

meaning $q_1 \leq \llbracket \mathfrak{t} \rrbracket p_1$ and $q_2 \leq \llbracket \mathfrak{t} \rrbracket p_2$, so that $q_1 + q_2 \leq \llbracket \mathfrak{t} \rrbracket (p_1 + p_2)$. Finally, by completeness we retrieve the result: $\vdash_{A_{tr}}^K [p_1 + p_2] \mathfrak{t} [q_1 + q_2]$.

(iterate zero): By (17) it holds $\llbracket \mathfrak{t}^* \rrbracket p \geq p$ which means $\models_{A_{tr}}^K [p] \mathfrak{t}^* [p]$, and by completeness we obtain the result $\vdash_{A_{tr}}^K [p] \mathfrak{t}^* [p]$.

(iterate non-zero): By (17) it holds $\llbracket \mathfrak{t}^* \rrbracket p \geq \llbracket \mathfrak{t}^* \cdot \mathfrak{t} \rrbracket p$. By hypothesis $\vdash_{A_{tr}}^K [p] \mathfrak{t}^* \cdot \mathfrak{t} [q]$, hence by soundness $q \leq \llbracket \mathfrak{t}^* \cdot \mathfrak{t} \rrbracket p$. Combining the two we yields $q \leq \llbracket \mathfrak{t}^* \rrbracket p$, which means $\models_{A_{tr}}^K [p] \mathfrak{t}^* [q]$ and by completeness $\vdash_{A_{tr}}^K [p] \mathfrak{t}^* [q]$. \square

Moreover, since the abstraction map defining A_{tr} is $\alpha_{A_{tr}} = \lambda x. \top$, we have that condition (b) of Definition 3.6 for the validity of a LCK triple trivially holds, that is, $\llbracket \mathfrak{t} \rrbracket_A^d \alpha(p) = \top = \alpha(q) = \top = \alpha(\llbracket \mathfrak{t} \rrbracket^K p)$. This therefore entails that

$$\models_{A_{tr}}^K [p] \mathfrak{t} [q] \Leftrightarrow \models_{UL} [p] \mathfrak{t} [q]. \quad (29)$$

This allows us to retrieve the soundness and completeness results of incorrectness logic [24] as a consequence of those for LCK.

Corollary 3.12. *Under the same hypotheses of Theorem 3.11, the proof system UL is sound and complete, that is, $\vdash_{UL} [p] \mathfrak{t} [q] \Leftrightarrow \models_{UL} [p] \mathfrak{t} [q]$.*

Proof.

$$\begin{aligned} \vdash_{UL} [p] \mathfrak{t} [q] &\Leftrightarrow \text{[by Theorem 3.11]} \\ \vdash_{A_{tr}}^K [p] \mathfrak{t} [q] &\Leftrightarrow \text{[by Theorems 3.7 and 3.10]} \\ \models_{A_{tr}}^K [p] \mathfrak{t} [q] &\Leftrightarrow \text{[by (29)]} \\ \models_{UL} [p] \mathfrak{t} [q] & \quad \square \end{aligned}$$

4 Incorrectness Logic in KAT

Incorrectness logic IL has been introduced by O’Hearn [24] as a natural under-approximating counterpart of the pivotal Hoare correctness logic [14], and quickly attracted a lot of research interest [19,25,26,27,31]. Incorrectness logic distinguishes two postconditions corresponding to normal and erroneous/abnormal program termination. Here, we generalize the algebraic formulation of our LCK logic to support abnormal termination. We follow the approach of Möller, O’Hearn and Hoare [22], namely, each language term is interpreted as a pair of KAT elements which model the normal and abnormal execution. The evaluation function has type $u : \text{Atom} \rightarrow (K \times K)$, while the interpretation function has type $\llbracket \cdot \rrbracket_u : T_{\Sigma, B} \rightarrow (K \times K)$. As a shorthand $\llbracket \cdot \rrbracket_u$ can be subscripted with **ok** or **err** to denote, resp., its first normal and second erroneous component. The definition is as follows:

$$\begin{aligned} \llbracket \mathfrak{a} \rrbracket_u &\triangleq u(\mathfrak{a}) \\ \llbracket \mathfrak{t}_1 + \mathfrak{t}_2 \rrbracket_u &\triangleq (\llbracket \mathfrak{t}_1 \rrbracket_{u_{\text{ok}}} + \llbracket \mathfrak{t}_2 \rrbracket_{u_{\text{ok}}}, \llbracket \mathfrak{t}_1 \rrbracket_{u_{\text{err}}} + \llbracket \mathfrak{t}_2 \rrbracket_{u_{\text{err}}}) \\ \llbracket \mathfrak{t}_1 \cdot \mathfrak{t}_2 \rrbracket_u &\triangleq (\llbracket \mathfrak{t}_1 \rrbracket_{u_{\text{ok}}} \cdot \llbracket \mathfrak{t}_2 \rrbracket_{u_{\text{ok}}}, \llbracket \mathfrak{t}_1 \rrbracket_{u_{\text{err}}} + \llbracket \mathfrak{t}_1 \rrbracket_{u_{\text{ok}}} \cdot \llbracket \mathfrak{t}_2 \rrbracket_{u_{\text{err}}}) \\ \llbracket \mathfrak{t}^* \rrbracket_u &\triangleq (\llbracket \mathfrak{t} \rrbracket_{u_{\text{ok}}}^*, \llbracket \mathfrak{t} \rrbracket_{u_{\text{ok}}}^* \cdot \llbracket \mathfrak{t} \rrbracket_{u_{\text{err}}}) \end{aligned} \quad (30)$$

Following the original definition of IL, the precondition encodes an **ok** condition only, while the postcondition contains both an **ok** and an **err** component. Hence, the latter is given by a pair $(p, q) \in \text{test}(K) \times \text{test}(K)$, typically denoted by **ok** : p , **err** : q . The concrete semantics $\llbracket \cdot \rrbracket : T_{\Sigma, B} \rightarrow (\text{test}(K) \rightarrow (\text{test}(K) \times \text{test}(K)))$ is defined as

$$\llbracket \mathfrak{t} \rrbracket p \triangleq \mathbf{ok} : \langle \llbracket \mathfrak{t} \rrbracket_{u_{\mathbf{ok}}} | p, \mathbf{err} : \langle \llbracket \mathfrak{t} \rrbracket_{u_{\mathbf{err}}} | p$$

To refer to one of its components, $\llbracket \cdot \rrbracket$ can be subscripted with **ok** or **err**, e.g., $\llbracket \mathfrak{t} \rrbracket_{\mathbf{ok}} p$.

Like for the LCK formulation, the concrete semantic of LCIL leverages the backward-diamond operator, therefore it inherits its properties (e.g., isotony, additivity). Moreover, by (1), (bd2) we have that

$$\llbracket \mathfrak{t}_1 \cdot \mathfrak{t}_2 \rrbracket_{\mathbf{err}} p = \llbracket \mathfrak{t}_1 \rrbracket_{\mathbf{err}} p + \llbracket \mathfrak{t}_2 \rrbracket_{\mathbf{err}} \llbracket \mathfrak{t}_1 \rrbracket_{\mathbf{ok}} p \quad (31)$$

$$\llbracket \mathfrak{t}^* \rrbracket_{\mathbf{err}} p = \llbracket \mathfrak{t} \rrbracket_{\mathbf{err}} \llbracket \mathfrak{t}^* \rrbracket_{\mathbf{ok}} p \quad (32)$$

Given a Kleene abstract domain A on K , the corresponding abstract semantics $\llbracket \cdot \rrbracket_A^\sharp : T_{\Sigma, B} \rightarrow (A \rightarrow (A \times A))$ is defined as follows:

$$\begin{aligned} \llbracket \mathfrak{a} \rrbracket_A^\sharp p^\sharp &\triangleq \mathbf{ok} : \alpha(\llbracket \mathfrak{a} \rrbracket_{\mathbf{ok}} \gamma(p^\sharp)), \mathbf{err} : \alpha(\llbracket \mathfrak{a} \rrbracket_{\mathbf{err}} \gamma(p^\sharp)) \\ \llbracket \mathfrak{t}_1 + \mathfrak{t}_2 \rrbracket_A^\sharp p^\sharp &\triangleq \mathbf{ok} : \llbracket \mathfrak{t}_1 \rrbracket_{A_{\mathbf{ok}}}^\sharp p^\sharp + \llbracket \mathfrak{t}_2 \rrbracket_{A_{\mathbf{ok}}}^\sharp p^\sharp, \mathbf{err} : \llbracket \mathfrak{t}_1 \rrbracket_{A_{\mathbf{err}}}^\sharp p^\sharp + \llbracket \mathfrak{t}_2 \rrbracket_{A_{\mathbf{err}}}^\sharp p^\sharp \\ \llbracket \mathfrak{t}_1 \cdot \mathfrak{t}_2 \rrbracket_A^\sharp p^\sharp &\triangleq \mathbf{ok} : \llbracket \mathfrak{t}_2 \rrbracket_{A_{\mathbf{ok}}}^\sharp (\llbracket \mathfrak{t}_1 \rrbracket_{A_{\mathbf{ok}}}^\sharp p^\sharp), \mathbf{err} : \llbracket \mathfrak{t}_1 \rrbracket_{A_{\mathbf{err}}}^\sharp p^\sharp + \llbracket \mathfrak{t}_2 \rrbracket_{A_{\mathbf{err}}}^\sharp (\llbracket \mathfrak{t}_1 \rrbracket_{A_{\mathbf{ok}}}^\sharp p^\sharp) \\ \llbracket \mathfrak{t}^* \rrbracket_A^\sharp p^\sharp &\triangleq \mathbf{ok} : \bigvee_{n \in \mathbb{N}} (\llbracket \mathfrak{t} \rrbracket_{A_{\mathbf{ok}}}^\sharp)^n p^\sharp, \mathbf{err} : \llbracket \mathfrak{t} \rrbracket_{A_{\mathbf{err}}}^\sharp \bigvee_{n \in \mathbb{N}} (\llbracket \mathfrak{t} \rrbracket_{A_{\mathbf{ok}}}^\sharp)^n p^\sharp \end{aligned} \quad (33)$$

The **ok** part coincides with the semantics of LCK, while the **err** component puts in place some differences. In particular, the composition exhibits a short-circuiting behavior, meaning that an error in the first command aborts the execution without executing the second one, while the Kleene star allows an error to occur after some error-free iterations. It is straightforward to check that this definition of abstract semantics is monotonic and sound.

The proof system LCK can be extended with incorrectness triples. In particular, a triple $[p] \mathfrak{t} [\mathbf{ok} : q][\mathbf{err} : r]$ is valid if the standard validity conditions hold for both **ok** and **err**.

Definition 4.1 (Incorrectness Triple). Let K be a CTC bdKAT K and $T_{\Sigma, B}$ be a language interpreted on K . An *incorrectness triple* is either $[p] \mathfrak{t} [\mathbf{ok} : q]$ or $[p] \mathfrak{t} [\mathbf{err} : r]$, where $p, q, r \in \text{test}(K)$ and $\mathfrak{t} \in T_{\Sigma, B}$.

Let A be a Kleene abstract domain on K with abstraction map $\alpha : \text{test}(K) \rightarrow A$.

- The triple $[p] \mathfrak{t} [\mathbf{ok} : q]$ is valid if: (1) $q \leq \llbracket \mathfrak{t} \rrbracket_{\mathbf{ok}} p$, and (2) $\llbracket \mathfrak{t} \rrbracket_{A_{\mathbf{ok}}}^\sharp \alpha(p) = \alpha(q) = \alpha(\llbracket \mathfrak{t} \rrbracket_{\mathbf{ok}} p)$.
- The triple $[p] \mathfrak{t} [\mathbf{err} : r]$ is valid if: (1) $r \leq \llbracket \mathfrak{t} \rrbracket_{\mathbf{err}} p$, and (2) $\llbracket \mathfrak{t} \rrbracket_{A_{\mathbf{err}}}^\sharp \alpha(p) = \alpha(r) = \alpha(\llbracket \mathfrak{t} \rrbracket_{\mathbf{err}} p)$.
- A triple $[p] \mathfrak{t} [\mathbf{ok} : q][\mathbf{err} : r]$ is valid when both $[p] \mathfrak{t} [\mathbf{ok} : q]$ and $[p] \mathfrak{t} [\mathbf{err} : r]$ are valid. In particular, if $q = r$ then the triple $[p] \mathfrak{t} [\mathbf{e} : q]$ is valid. \square

The proof system LCIL_A defining the local completeness incorrectness logic is given in Figure 3.

$$\begin{array}{c}
\frac{\mathbf{a} \in \text{Atom} \quad \mathbb{C}_p^A(\llbracket \mathbf{a} \rrbracket_{\text{ok}}) \quad \mathbb{C}_p^A(\llbracket \mathbf{a} \rrbracket_{\text{err}})}{\vdash_A^K [p] \mathbf{a} \text{ [ok : } \llbracket \mathbf{a} \rrbracket_{\text{ok}} p \text{ [err : } \llbracket \mathbf{a} \rrbracket_{\text{err}} p]} \text{ (transfer)} \\
\\
\frac{p' \leq p \leq A(p') \quad \vdash_A^K [p'] \mathbf{t} \text{ [}\epsilon : q'] \quad q \leq q' \leq A(q)}{\vdash_A^K [p] \mathbf{t} \text{ [}\epsilon : q]} \text{ (relax)} \\
\\
\frac{\vdash_A^K [p] \mathbf{t}_1 \text{ [ok : } r] \quad \vdash_A^K [r] \mathbf{t}_2 \text{ [ok : } q]}{\vdash_A^K [p] \mathbf{t}_1 \cdot \mathbf{t}_2 \text{ [ok : } q]} \text{ (seq-ok)} \\
\\
\frac{\vdash_A^K [p] \mathbf{t}_1 \text{ [ok : } q \text{ [err : } r] \quad \vdash_A^K [q] \mathbf{t}_2 \text{ [err : } s]}{\vdash_A^K [p] \mathbf{t}_1 \cdot \mathbf{t}_2 \text{ [err : } r + s]} \text{ (seq-err)} \\
\\
\frac{\vdash_A^K [p] \mathbf{t}^* \text{ [ok : } q] \quad \vdash_A^K [q] \mathbf{t} \text{ [err : } r]}{\vdash_A^K [p] \mathbf{t}^* \text{ [err : } r]} \text{ (rec-err)} \\
\\
\frac{\vdash_A^K [p] \mathbf{t}_1 \text{ [}\epsilon : q_1] \quad \vdash_A^K [p] \mathbf{t}_2 \text{ [}\epsilon : q_2]}{\vdash_A^K [p] \mathbf{t}_1 + \mathbf{t}_2 \text{ [}\epsilon : q_1 + q_2]} \text{ (join)} \quad \frac{\forall n \in \mathbb{N}. \vdash_A^K [p_n] \mathbf{t} \text{ [ok : } p_{n+1}]}{\vdash_A^K [p_0] \mathbf{t}^* \text{ [ok : } \bigvee_{n \in \mathbb{N}} p_n]} \text{ (limit)}
\end{array}$$

Fig. 3. Proof system LCIL_A.

Theorem 4.2 (Logical Soundness of LCIL_A). *The triples provable in LCIL_A are valid.*

Proof. Definitions of $\llbracket \cdot \rrbracket_{u_{\text{ok}}}$ and $\llbracket \cdot \rrbracket_{A_{\text{ok}}}^\sharp$ coincide respectively with $\llbracket \cdot \rrbracket_u$ and $\llbracket \cdot \rrbracket_A^\sharp$ of LCK, and the **ok** part of LCIL is a subset of LCK¹. For this reason the soundness of the **ok** part of the triple follows by Theorem 3.7. Moreover, the base case $\mathbf{t} \equiv \mathbf{a} \in \text{Atom}$ and the inductive case $\mathbf{t} \equiv \mathbf{t}_1 + \mathbf{t}_2$ of $\llbracket \cdot \rrbracket_{u_{\text{err}}}$ and $\llbracket \cdot \rrbracket_{A_{\text{err}}}^\sharp$ are completely symmetric to their **ok** counterparts. For this reason the **err** part of (transfer),(join) can be shown in the same way as the **ok** one. Likewise, the soundness of the **err** part of (relax) follows as for the **ok** one.

The only cases that are left to prove are (seq-err) and (rec-err). As in Theorem 3.7, notice that the first equality of (2) entails the second one, i.e., if $\llbracket \mathbf{t} \rrbracket_{A_{\text{err}}}^\sharp \alpha(p) = \alpha(r)$ is true, then $\llbracket \mathbf{t} \rrbracket_{A_{\text{err}}}^\sharp \alpha(p) = \alpha(\llbracket \mathbf{t} \rrbracket_{\text{err}} p)$. For this reason it is enough to show only the former.

(seq-err): By induction we have $q \leq \llbracket \mathbf{t}_1 \rrbracket_{\text{ok}} p$, $r \leq \llbracket \mathbf{t}_1 \rrbracket_{\text{err}} p$ and $s \leq \llbracket \mathbf{t}_2 \rrbracket_{\text{err}} q$, thus $s \leq \llbracket \mathbf{t}_2 \rrbracket_{\text{err}} q \leq \llbracket \mathbf{t}_2 \rrbracket_{\text{err}} \llbracket \mathbf{t}_1 \rrbracket_{\text{ok}} p$. By (31) we have that $\llbracket \mathbf{t}_1 \cdot \mathbf{t}_2 \rrbracket_{\text{err}} p = \llbracket \mathbf{t}_1 \rrbracket_{\text{err}} p + \llbracket \mathbf{t}_2 \rrbracket_{\text{err}} \llbracket \mathbf{t}_1 \rrbracket_{\text{ok}} p \geq r + s$, thus implying (1). By induction we have $\llbracket \mathbf{t}_1 \rrbracket_{A_{\text{ok}}}^\sharp \alpha(p) = \alpha(q)$, $\llbracket \mathbf{t}_1 \rrbracket_{A_{\text{err}}}^\sharp \alpha(p) = \alpha(r)$ and $\llbracket \mathbf{t}_2 \rrbracket_{A_{\text{err}}}^\sharp \alpha(q) = \alpha(s)$, so that $\llbracket \mathbf{t}_1 \rrbracket_{A_{\text{err}}}^\sharp \llbracket \mathbf{t}_1 \rrbracket_{A_{\text{ok}}}^\sharp \alpha(p) =$

¹ Here, we refer to the proof system LCK with the additional rule (limit).

$\llbracket \mathfrak{t}_2 \rrbracket_{A_{\text{err}}}^\# \alpha(q) = \alpha(s)$. By definition of $\llbracket \mathfrak{t}_1 \cdot \mathfrak{t}_2 \rrbracket_{A_{\text{err}}}^\#$, we can prove (2) as follows

$$\begin{aligned} & \llbracket \mathfrak{t}_1 \cdot \mathfrak{t}_2 \rrbracket_{A_{\text{err}}}^\# \alpha(p) = \\ & \llbracket \mathfrak{t}_1 \rrbracket_{A_{\text{err}}}^\# \alpha(p) + \llbracket \mathfrak{t}_2 \rrbracket_{A_{\text{err}}}^\# \llbracket \mathfrak{t}_1 \rrbracket_{A_{\text{ok}}}^\# \alpha(p) = \\ & \alpha(r) + \alpha(s) = [\text{Additivity of } \alpha(\cdot)] \\ & \alpha(r + s). \end{aligned}$$

(rec-err): By induction we have $q \leq \llbracket \mathfrak{t}^* \rrbracket_{\text{ok}} p$, $r \leq \llbracket \mathfrak{t} \rrbracket_{\text{err}} q$. By (32) we have that $\llbracket \mathfrak{t}^* \rrbracket_{\text{err}} p = \llbracket \mathfrak{t} \rrbracket_{\text{err}} \llbracket \mathfrak{t}^* \rrbracket_{\text{ok}} p \geq \llbracket \mathfrak{t} \rrbracket_{\text{err}} q \geq r$, thus implying (1). By induction we have that $\llbracket \mathfrak{t}^* \rrbracket_{A_{\text{ok}}}^\# \alpha(p) = \alpha(q)$ and $\llbracket \mathfrak{t} \rrbracket_{A_{\text{err}}}^\# \alpha(q) = \alpha(r)$. Using the definition of $\llbracket \mathfrak{t}^* \rrbracket_{A_{\text{err}}}^\#$ we can prove (2) as $\llbracket \mathfrak{t}^* \rrbracket_{A_{\text{err}}}^\# \alpha(p) = \llbracket \mathfrak{t} \rrbracket_{A_{\text{err}}}^\# \llbracket \mathfrak{t}^* \rrbracket_{A_{\text{ok}}}^\# \alpha(p) = \llbracket \mathfrak{t} \rrbracket_{A_{\text{err}}}^\# \alpha(q) = \alpha(r)$. \square

Furthermore, it turns out that LCIL_A is logically complete. To prove this, we need a preliminary result.

Lemma 4.3. *Let A be a Kleene abstract domain on a CTC bdKAT K and $T_{\Sigma, B}$ be a language interpreted on K . If the atoms in $\mathfrak{t} \in T_{\Sigma, B}$ are globally complete, i.e., for all $\mathfrak{a} \in \text{Atom}(\mathfrak{t})$, $\mathbb{C}^A(\llbracket \mathfrak{a} \rrbracket_{\text{ok}})$ and $\mathbb{C}^A(\llbracket \mathfrak{a} \rrbracket_{\text{err}})$, then, for any $p \in \text{test}(K)$, $\llbracket \mathfrak{t} \rrbracket_{A_{\text{ok}}}^\# \alpha(p) = \alpha(\llbracket \mathfrak{t} \rrbracket_{\text{ok}} p)$ and $\llbracket \mathfrak{t} \rrbracket_{A_{\text{err}}}^\# \alpha(p) = \alpha(\llbracket \mathfrak{t} \rrbracket_{\text{err}} p)$.*

Proof. Analogous to the proof of Lemma 3.9. \square

Theorem 4.4 (Logical Completeness of LCIL_A). *Let A be a Kleene abstract domain on a CTC bdKAT K and $T_{\Sigma, B}$ be a language interpreted on K . Assume that the atoms in $\mathfrak{t} \in T_{\Sigma, B}$ are globally complete, i.e., for all $\mathfrak{a} \in \text{Atom}(\mathfrak{t})$, $\mathbb{C}^A(\llbracket \mathfrak{a} \rrbracket_{\text{ok}})$ and $\mathbb{C}^A(\llbracket \mathfrak{a} \rrbracket_{\text{err}})$. If $[p] \mathfrak{t} [\text{ok} : q] [\text{err} : r]$ is valid, then it is provable in LCIL_A .*

Proof. As discussed in the proof of Theorem 4.2, the **ok** part of the semantic coincides with the LCK logic, so that $[p] \mathfrak{t} [\text{ok} : q]$ is provable by the completeness theorem of LCK_A . For this reason only $[p] \mathfrak{t} [\text{err} : r]$ must be shown. The proof is by induction on the structure of $\mathfrak{t} \in T_{\Sigma, B}$. By construction, the **err** base case $\mathfrak{t} \equiv \mathfrak{a} \in \text{Atom}$ and the inductive case $\mathfrak{t} \equiv \mathfrak{t}_1 + \mathfrak{t}_2$ are symmetric to their **ok** counterparts, so that the proof of those cases is equivalent to Theorem 3.10. Therefore we only need to prove \cdot and $*$.

($\mathfrak{t}_1 \cdot \mathfrak{t}_2$): Consider the following assignments (the equality after the \Rightarrow sign is true by Lemma 4.3):

$$\begin{aligned} s &= \llbracket \mathfrak{t}_1 \rrbracket_{\text{ok}} p \Rightarrow \llbracket \mathfrak{t}_1 \rrbracket_{A_{\text{ok}}}^\# \alpha(p) = \alpha(\llbracket \mathfrak{t}_1 \rrbracket_{\text{ok}} p) = \alpha(s) \\ t &= \llbracket \mathfrak{t}_1 \rrbracket_{\text{err}} p \Rightarrow \llbracket \mathfrak{t}_1 \rrbracket_{A_{\text{err}}}^\# \alpha(p) = \alpha(\llbracket \mathfrak{t}_1 \rrbracket_{\text{err}} p) = \alpha(t) \\ u &= \llbracket \mathfrak{t}_2 \rrbracket_{\text{err}} s \Rightarrow \llbracket \mathfrak{t}_2 \rrbracket_{A_{\text{err}}}^\# \alpha(s) = \alpha(\llbracket \mathfrak{t}_2 \rrbracket_{\text{err}} s) = \alpha(u) \end{aligned}$$

and by induction we have that

$$\begin{aligned} & \models_A^K [p] \mathfrak{t}_1 [\text{ok} : s] \Rightarrow \vdash_A^K [p] \mathfrak{t}_1 [\text{ok} : s] \\ & \models_A^K [p] \mathfrak{t}_1 [\text{err} : t] \Rightarrow \vdash_A^K [p] \mathfrak{t}_1 [\text{err} : t] \\ & \models_A^K [s] \mathfrak{t}_2 [\text{err} : u] \Rightarrow \vdash_A^K [s] \mathfrak{t}_2 [\text{err} : u] \end{aligned}$$

Then by (seq-err) the triple $[p] \mathbf{t}_1 \cdot \mathbf{t}_2 [\mathbf{err} : t + u]$ is provable. By hypothesis we have

$$r \leq \llbracket \mathbf{t}_1 \cdot \mathbf{t}_2 \rrbracket_{\mathbf{err}} p \stackrel{(31)}{=} \llbracket \mathbf{t}_1 \rrbracket_{\mathbf{err}} p + \llbracket \mathbf{t}_2 \rrbracket_{\mathbf{err}} \llbracket \mathbf{t}_1 \rrbracket_{\mathbf{ok}} p = t + u$$

and

$$\alpha(r) = \alpha(\llbracket \mathbf{t}_1 \cdot \mathbf{t}_2 \rrbracket_{\mathbf{err}} p) = \alpha(t + u)$$

so that we can apply (relax) to obtain the result $\vdash_A^K [p] \mathbf{t}_1 \cdot \mathbf{t}_2 [\mathbf{err} : r]$.

(\mathbf{t}_0^*): Let $q = \llbracket \mathbf{t}_0^* \rrbracket_{\mathbf{ok}} p$ and $s = \llbracket \mathbf{t}_0 \rrbracket_{\mathbf{err}} \llbracket \mathbf{t}_0^* \rrbracket_{\mathbf{ok}} p$. By Lemma 4.3 $\llbracket \mathbf{t}_0^* \rrbracket_{A_{\mathbf{ok}}}^\# \alpha(p) = \alpha(q)$ so that $\vdash_A^K [p] \mathbf{t}_0^* [\mathbf{ok} : q]$ and the completeness of the **ok** part yields $\vdash_A^K [p] \mathbf{t}_0^* [\mathbf{ok} : q]$. By Lemma 4.3 $\llbracket \mathbf{t}_0 \rrbracket_{A_{\mathbf{err}}}^\# \alpha(q) = \alpha(s)$, meaning that $\vdash_A^K [q] \mathbf{t}_0 [\mathbf{err} : s]$ and by induction $\vdash_A^K [q] \mathbf{t}_0 [\mathbf{err} : s]$. By (rec-err) we obtain $\vdash_A^K [p] \mathbf{t}_0^* [\mathbf{err} : s]$.

In order to use (relax), we need to ensure that the additional conditions $r \leq s$ and $\alpha(r) = \alpha(s)$ are satisfied. By $\vdash_A^K [p] \mathbf{t}_0^* [\mathbf{err} : r]$ and (32), $r \leq \llbracket \mathbf{t}_0^* \rrbracket_{\mathbf{err}} p = s$ and $\alpha(r) = \alpha(\llbracket \mathbf{t}_0^* \rrbracket_{\mathbf{err}} p) = \alpha(\llbracket \mathbf{t}_0 \rrbracket_{\mathbf{err}} \llbracket \mathbf{t}_0^* \rrbracket_{\mathbf{ok}} p) = \alpha(s)$, so that (relax) yields the result $\vdash_A^K [p] \mathbf{t}_0^* [\mathbf{err} : r]$. \square

Example 4.5. Consider a relational bdKAT $K = \wp(\mathbb{Z} \times \mathbb{Z})$ on the set of integers \mathbb{Z} , where $\mathbf{1}_K \triangleq \{\langle z, z \rangle \mid z \in \mathbb{Z}\}$ and $\mathbf{0}_K \triangleq \emptyset$, and the standard integer interval abstraction Int [7, 8]. Let us consider a language with primitive actions $\Sigma = \{\text{skip}, \mathbf{err}\}$ and primitive tests $B = \{x = 0, x \neq 0\}$. The evaluation function $u : \Sigma \cup B \rightarrow K_{\mathbf{ok}} \times K_{\mathbf{err}}$ is defined as expected:

$$\begin{aligned} u(\text{skip}) &= (\mathbf{1}_K, \mathbf{0}_K) & u(\mathbf{err}) &= (\mathbf{0}_K, \mathbf{1}_K) \\ u(x = 0) &= (\{\langle 0, 0 \rangle\}, \mathbf{0}_K) & u(x \neq 0) &= (\mathbf{1}_K \setminus \{\langle 0, 0 \rangle\}, \mathbf{0}_K) \end{aligned}$$

We study the correctness of the program $\mathbf{r} \triangleq ((x \neq 0) \cdot \text{skip}) + ((x = 0) \cdot \mathbf{err})$, for precondition $p \triangleq \{\langle z, z \rangle \mid z \in [-10, 10]\}$ and specification $\text{Spec} \triangleq (\mathbf{ok} : p, \mathbf{err} : \mathbf{0}_K)$. Let us define two auxiliary tests $q \triangleq \{\langle 0, 0 \rangle\}$ and $s \triangleq p \setminus q$.

The atoms are locally complete because $\text{Int}(p) = p$ holds and $\text{skip}, \mathbf{err}$ are globally complete. We can compute the concrete semantics by Lemma 3.1 characterizing the backward diamond operator in relational KATs. We therefore have the following derivation in LCIL_{Int} for \mathbf{r} :

$$\begin{aligned} & \frac{\frac{\mathbb{C}_p^{\text{Int}}(\llbracket [x \neq 0]_{\mathbf{ok}} \rrbracket)}{\vdash_{\text{Int}}^K [p] x \neq 0 [\mathbf{ok} : s] [\mathbf{err} : 0]} \quad \mathbb{C}_p^{\text{Int}}(\llbracket [x \neq 0]_{\mathbf{err}} \rrbracket)}{\text{(transfer)}} \quad \frac{\mathbb{C}_s^{\text{Int}}(\llbracket [\text{skip}]_{\mathbf{ok}} \rrbracket)}{\vdash_{\text{Int}}^K [s] \text{skip} [\mathbf{ok} : s] [\mathbf{err} : 0]} \quad \mathbb{C}_s^{\text{Int}}(\llbracket [\text{skip}]_{\mathbf{err}} \rrbracket)}{\text{(transfer)}}}{\text{(seq-ok), (seq-err)}} \\ & \frac{\frac{\mathbb{C}_p^{\text{Int}}(\llbracket [x = 0]_{\mathbf{ok}} \rrbracket)}{\vdash_{\text{Int}}^K [p] x = 0 [\mathbf{ok} : q] [\mathbf{err} : 0]} \quad \mathbb{C}_p^{\text{Int}}(\llbracket [x = 0]_{\mathbf{err}} \rrbracket)}{\text{(transfer)}} \quad \frac{\mathbb{C}_q^{\text{Int}}(\llbracket [\mathbf{err}]_{\mathbf{ok}} \rrbracket)}{\vdash_{\text{Int}}^K [q] \mathbf{err} [\mathbf{ok} : 0] [\mathbf{err} : q]} \quad \mathbb{C}_q^{\text{Int}}(\llbracket [\mathbf{err}]_{\mathbf{err}} \rrbracket)}{\text{(transfer)}}}{\text{(seq-ok), (seq-err)}} \\ & \frac{\frac{\vdash_{\text{Int}}^K [p] (x \neq 0) \cdot \text{skip} [\mathbf{ok} : s] [\mathbf{err} : 0]}{\text{(seq-ok), (seq-err)}} \quad \frac{\vdash_{\text{Int}}^K [p] (x = 0) \cdot \mathbf{err} [\mathbf{ok} : 0] [\mathbf{err} : q]}{\text{(seq-ok), (seq-err)}}}{\text{(join)}} \\ & \frac{\vdash_{\text{Int}}^K [p] ((x \neq 0) \cdot \text{skip}) + ((x = 0) \cdot \mathbf{err}) [\mathbf{ok} : s] [\mathbf{err} : q]}{\text{(join)}} \end{aligned}$$

By soundness of LCIL_{Int} in Theorem 4.2, the program \mathbf{r} satisfies the **ok** part of Spec because

$$\llbracket \mathbf{r} \rrbracket_{\mathbf{ok}} p \subseteq \text{Int}(s) = p \subseteq p = \text{Spec}_{\mathbf{ok}}.$$

However, the **err** part is not satisfied as $\text{Int}(q) = q \not\subseteq \emptyset = \mathbf{0}_K = \text{Spec}_{\mathbf{err}}$. Moreover, LCIL_{Int} also catches a true alert as $q \setminus \text{Spec}_{\mathbf{err}} = q = \{\langle 0, 0 \rangle\}$. \square

$\frac{\mathbf{a} \in \text{Atom}}{\vdash_{\text{IL}} [p] \mathbf{a} [\mathbf{ok} : \llbracket \mathbf{a} \rrbracket_{\mathbf{ok}} p] [\mathbf{err} : \llbracket \mathbf{a} \rrbracket_{\mathbf{err}} p]} \text{ (transfer)}$	$\frac{}{\vdash_{\text{IL}} [p] \mathbf{t} [\mathbf{\epsilon} : 0]} \text{ (empty)}$
$\frac{p' \leq p \quad \vdash_{\text{IL}} [p'] \mathbf{t} [\mathbf{\epsilon} : q'] \quad q \leq q'}{\vdash_{\text{IL}} [p] \mathbf{t} [\mathbf{\epsilon} : q]} \text{ (consequence)}$	
$\frac{\vdash_{\text{IL}} [p_1] \mathbf{t} [\mathbf{\epsilon} : q_1] \quad \vdash_{\text{IL}} [p_2] \mathbf{t} [\mathbf{\epsilon} : q_2]}{\vdash_{\text{IL}} [p_1 + p_2] \mathbf{t} [\mathbf{\epsilon} : q_1 + q_2]} \text{ (disj)}$	$\frac{\vdash_{\text{IL}} [p] \mathbf{t}_1 [\mathbf{err} : q]}{\vdash_{\text{IL}} [p] \mathbf{t}_1 \cdot \mathbf{t}_2 [\mathbf{err} : q]} \text{ (short-circuit)}$
$\frac{\vdash_{\text{IL}} [p] \mathbf{t}_1 [\mathbf{ok} : r] \quad \vdash_{\text{IL}} [r] \mathbf{t}_2 [\mathbf{\epsilon} : q]}{\vdash_{\text{IL}} [p] \mathbf{t}_1 \cdot \mathbf{t}_2 [\mathbf{\epsilon} : q]} \text{ (seq-normal)}$	
$\frac{\forall n \in \mathbb{N}. \vdash_{\text{IL}} [p_n] \mathbf{t} [\mathbf{ok} : p_{n+1}]}{\vdash_{\text{IL}} [p_0] \mathbf{t}^* [\mathbf{ok} : \bigvee_{n \in \mathbb{N}} p_n]} \text{ (back-v)}$	$\frac{\vdash_{\text{IL}} [p] \mathbf{t}^* \cdot \mathbf{t} [\mathbf{\epsilon} : q]}{\vdash_{\text{IL}} [p] \mathbf{t}^* [\mathbf{\epsilon} : q]} \text{ (iterate non-zero)}$
$\frac{\vdash_{\text{IL}} [p] \mathbf{t}_i [\mathbf{\epsilon} : q], \text{ with } i \in \{1, 2\}}{\vdash_{\text{IL}} [p] \mathbf{t}_1 + \mathbf{t}_2 [\mathbf{\epsilon} : q]} \text{ (choice)}$	

Fig. 4. Proof system IL.

4.1 Relationship with Incorrectness logic

Section 3.6 has shown that LCK is a generalization of UL. The same can be done for IL, i.e., we prove that LCIL_A with incorrectness triples generalizes the incorrectness logic of [24]. For the sake of clarity, we recall in Figure 4 an algebraic version of IL. Analogously to the reduction of Theorem 3.11, this generalization is obtained by letting $A = A_{tr}$, where A_{tr} is the trivial abstract domain.

Theorem 4.6. *Let K be a CTC bdKAT and $T_{\Sigma, B}$ a language interpreted on K . For any $p, q \in \text{test}(K)$, $\mathbf{t} \in T_{\Sigma, B}$,*

$$\vdash_{A_{tr}}^K [p] \mathbf{t} [\mathbf{ok} : q] [\mathbf{err} : r] \Leftrightarrow \vdash_{\text{IL}} [p] \mathbf{t} [\mathbf{ok} : q] [\mathbf{err} : r].$$

Proof. Notice that the **ok** part of LCIL coincides with LCK^2 and likewise IL coincides with UL. Hence this part of the equivalence follows by Theorem 3.11. We only need to show the equivalence of the **err** part.

\Rightarrow) Assuming $\vdash_{A_{tr}}^K [p] \mathbf{t} [\mathbf{err} : r]$ we need to show $\vdash_{\text{IL}} [p] \mathbf{t} [\mathbf{err} : r]$. The proof is by induction on the derivation tree.

(transfer): Coincides with (transfer).

(relax): Coincides with (consequence).

(seq-ok): Is the **ok** part of (seq-normal).

(seq-err): Can be derived as follows

$$\frac{\frac{\vdash_{\text{IL}} [p] \mathbf{t}_1 [\mathbf{ok} : q] \quad \vdash_{\text{IL}} [q] \mathbf{t}_2 [\mathbf{err} : s]}{\vdash_{\text{IL}} [p] \mathbf{t}_1 \cdot \mathbf{t}_2 [\mathbf{err} : s]} \text{ (seq-normal)} \quad \frac{\vdash_{\text{IL}} [p] \mathbf{t}_1 [\mathbf{err} : r]}{\vdash_{\text{IL}} [p] \mathbf{t}_1 \cdot \mathbf{t}_2 [\mathbf{err} : r]} \text{ (short-circuit)}}{\vdash_{\text{IL}} [p] \mathbf{t}_1 \cdot \mathbf{t}_2 [\mathbf{err} : s + r]} \text{ (disj)}$$

² With the additional rule (limit) and without the rules (rec), (iterate).

(rec-err): Can be derived as follows

$$\frac{\frac{\vdash_{\text{IL}} [p] \mathfrak{t}^* [\mathbf{ok} : q] \quad \vdash_{\text{IL}} [q] \mathfrak{t} [\mathbf{err} : r]}{\vdash_{\text{IL}} [p] \mathfrak{t}^* \cdot \mathfrak{t} [\mathbf{err} : r]} \text{ (seq-normal)}}{\vdash_{\text{IL}} [p] \mathfrak{t}^* [\mathbf{err} : r]} \text{ (iterate non-zero)}$$

(join): As in Theorem 3.11.

(limit): Coincides with (back-v).

\Leftarrow) Assuming $\vdash_{\text{IL}} [p] \mathfrak{t} [\mathbf{err} : r]$ we need to show $\vdash_{A_{tr}}^{\text{K}} [p] \mathfrak{t} [\mathbf{err} : r]$. As in Theorem 3.11, we exploit the completeness of LCIL and likewise we observe that the condition (2) is trivially true as $\llbracket \mathfrak{t} \rrbracket_{A_{tr, \mathbf{err}}}^{\sharp} \alpha(p) = \top = \alpha(q) = \top = \alpha(\llbracket \mathfrak{t} \rrbracket_{\mathbf{err}} p)$, meaning that the only condition to check is (1). The proof is by structural induction on the derivation tree of $\vdash_{\text{IL}} [p] \mathfrak{t} [\mathbf{err} : r]$.

(transfer): The premises $\mathbb{C}_p^A(\llbracket \mathbf{a} \rrbracket_{\mathbf{err}})$ and $\mathbb{C}_p^A(\llbracket \mathbf{a} \rrbracket_{\mathbf{ok}})$ hold for any \mathbf{a} and p because $A_{tr}(\cdot) = 1$, so that (transfer) yields $\vdash_{A_{tr}}^{\text{K}} [p] \mathbf{a} [\mathbf{err} : \llbracket \mathbf{a} \rrbracket_{\mathbf{err}} p]$.

(empty): For any $\mathfrak{t} \in T_{\Sigma, B}$ it holds $0 \leq \llbracket \mathfrak{t} \rrbracket_{\mathbf{err}} p$, meaning $\models_{A_{tr}}^{\text{K}} [p] \mathfrak{t} [\mathbf{err} : 0]$ and by completeness of LCIL we have $\vdash_{A_{tr}}^{\text{K}} [p] \mathfrak{t} [\mathbf{err} : 0]$.

(consequence): The inequalities $p \leq A_{tr}(p')$ and $q' \leq A_{tr}(q)$ are always true because $A_{tr}(\cdot) = 1$ which is the greatest element of $\text{test}(K)$ meaning that the rule coincides with (relax).

(disj) and (choice): Same as in Theorem 3.11.

(short-circuit): By induction $\vdash_{A_{tr}}^{\text{K}} [p] \mathfrak{t}_1 [\mathbf{err} : r]$ and by soundness $\models_{A_{tr}}^{\text{K}} [p] \mathfrak{t}_1 [\mathbf{err} : r]$. By (31) $\llbracket \mathfrak{t}_1 \cdot \mathfrak{t}_2 \rrbracket_{\mathbf{err}} p = \llbracket \mathfrak{t}_1 \rrbracket_{\mathbf{err}} p + \llbracket \mathfrak{t}_2 \rrbracket_{\mathbf{err}} \llbracket \mathfrak{t}_1 \rrbracket_{\mathbf{ok}} p \geq \llbracket \mathfrak{t}_1 \rrbracket_{\mathbf{err}} p \geq r$ which means $\models_{A_{tr}}^{\text{K}} [p] \mathfrak{t}_1 \cdot \mathfrak{t}_2 [\mathbf{err} : r]$ and by completeness $\vdash_{A_{tr}}^{\text{K}} [p] \mathfrak{t}_1 \cdot \mathfrak{t}_2 [\mathbf{err} : r]$.

(seq-normal): By induction and soundness we have $r \leq \llbracket \mathfrak{t}_1 \rrbracket_{\mathbf{ok}} p$ and $q \leq \llbracket \mathfrak{t}_2 \rrbracket_{\mathbf{err}} r$. By (31) $\llbracket \mathfrak{t}_1 \cdot \mathfrak{t}_2 \rrbracket_{\mathbf{err}} p = \llbracket \mathfrak{t}_1 \rrbracket_{\mathbf{err}} p + \llbracket \mathfrak{t}_2 \rrbracket_{\mathbf{err}} \llbracket \mathfrak{t}_1 \rrbracket_{\mathbf{ok}} p \geq \llbracket \mathfrak{t}_2 \rrbracket_{\mathbf{err}} \llbracket \mathfrak{t}_1 \rrbracket_{\mathbf{ok}} p \geq \llbracket \mathfrak{t}_2 \rrbracket_{\mathbf{err}} r \geq q$, which means $\models_{A_{tr}}^{\text{K}} [p] \mathfrak{t}_1 \cdot \mathfrak{t}_2 [\mathbf{err} : q]$ and by completeness $\vdash_{A_{tr}}^{\text{K}} [p] \mathfrak{t}_1 \cdot \mathfrak{t}_2 [\mathbf{err} : q]$.

(iterate non-zero): By induction and soundness

$$q \leq \llbracket \mathfrak{t}^* \cdot \mathfrak{t} \rrbracket_{\mathbf{err}} p \stackrel{(31)}{=} \llbracket \mathfrak{t}^* \rrbracket_{\mathbf{err}} p + \llbracket \mathfrak{t} \rrbracket_{\mathbf{err}} \llbracket \mathfrak{t}^* \rrbracket_{\mathbf{ok}} p \stackrel{(32)}{=} \llbracket \mathfrak{t}^* \rrbracket_{\mathbf{err}} p$$

which means $\models_{A_{tr}}^{\text{K}} [p] \mathfrak{t}^* [\mathbf{err} : q]$ and by completeness $\vdash_{A_{tr}}^{\text{K}} [p] \mathfrak{t}^* [\mathbf{err} : q]$. \square

The abstraction map $\alpha = \lambda x. \top$ of A_{tr} makes the validity of a triple trivially true. In particular, $\llbracket \mathfrak{t} \rrbracket_{A_{tr, \mathbf{ok}}}^{\sharp} \alpha(p) = \top = \alpha(q) = \top = \alpha(\llbracket \mathfrak{t} \rrbracket_{\mathbf{ok}} p)$ and $\llbracket \mathfrak{t} \rrbracket_{A_{tr, \mathbf{err}}}^{\sharp} \alpha(p) = \top = \alpha(q) = \top = \alpha(\llbracket \mathfrak{t} \rrbracket_{\mathbf{err}} p)$ hold. As a consequence, we obtain that

$$\models_{A_{tr}}^{\text{K}} [p] \mathfrak{t} [\mathbf{ok} : q] [\mathbf{err} : r] \Leftrightarrow \vdash_{\text{IL}} [p] \mathfrak{t} [\mathbf{ok} : q] [\mathbf{err} : r] \quad (34)$$

By this equivalence (34) and Theorems 4.2 and 4.4, we can thus retrieve the logical soundness and completeness of IL as a consequence of the one of LCIL $_{A_{tr}}$.

Corollary 4.7. *Let K be a CTC bdKAT and $T_{\Sigma, B}$ a language interpreted on K . For any $p, q \in \text{test}(K)$, $\mathfrak{t} \in T_{\Sigma, B}$, $\vdash_{\text{IL}} [p] \mathfrak{t} [\mathbf{ok} : q] [\mathbf{err} : r] \Leftrightarrow \models_{\text{IL}} [p] \mathfrak{t} [\mathbf{ok} : q] [\mathbf{err} : r]$.*

5 Local Completeness Logic in TopKAT

We have shown in Section 3 how KAT extended with a modal backward-diamond operator allows us to interpret and represent the local completeness program logic. This result follows the approach used by Moller, O’Hearn and Hoare [22], who leverage a backward-diamond operator in their KAT interpretation of correctness/incorrectness logics. On the other hand, very recently, Zhang et al. [32] have shown that incorrectness logic can be formulated for a standard KAT, provided that it contains a top element, so-called TopKAT. In particular, [32] observed that TopKAT is enough to express the codomain of relational KATs. In this section we take a similar path in studying an alternative formulation of local completeness logic based on a TopKAT.

5.1 Abstracting TopKATs

We expect that the base case of abstract semantics $\llbracket a \rrbracket_A^\sharp p^\sharp$ of a basic action $a \in \text{Atom}$, is defined as best correct approximation in A of the concrete semantics of a on the concretization of p^\sharp . In a bdKAT this is achieved through its backward-diamond operator, which is crucially used in (11) to define the strongest postcondition as $\llbracket a \rrbracket^K \gamma(p^\sharp) = \langle \llbracket a \rrbracket_u | \gamma(p^\sharp) \rangle$. Zhang et al. [32] observed that in a relational model of KAT, the codomain inclusion $\text{cod}(q) \subseteq \text{cod}(pa)$ defining the meaning of an under-approximation triple $[p] a [q]$ can be expressed in a TopKAT as the inequality $\top q \leq \top pa$, thus hinting that this condition could be taken as definition of validity of incorrectness triples in a TopKAT. We therefore follow a similar approach by considering the element $\top p \llbracket a \rrbracket_u$ as a proxy for strongest postconditions in a TopKAT. Notice that while in a bdKAT a strongest postcondition $\langle \llbracket a \rrbracket_u | p \rangle$ is always a test, in a TopKAT K , given $p \in \text{test}(K)$ and $\mathbf{a} \in T_{\Sigma, B}$, it is not guaranteed that there exists a test $q \in \text{test}(K)$ such that $\top p \llbracket \mathbf{a} \rrbracket_u = \top q$, as shown by the following example.

Example 5.1 (Strongest Postconditions in TopKAT). Consider the Kleene algebra $A_3 = \{0, 1, a\}$ consisting of 3 elements and characterized in [6, Chapter 12]. This algebra can be lifted to a KAT by letting $\text{test}(A_3) \triangleq \{0, 1\}$ and defining the KAT operators as follows:

$$\begin{array}{c} + \mid 0 \ 1 \ a \\ \hline 0 \mid 0 \ 1 \ a \\ 1 \mid 1 \ 1 \ 1 \\ a \mid a \ 1 \ a \end{array} \quad \begin{array}{c} \cdot \mid 0 \ 1 \ a \\ \hline 0 \mid 0 \ 0 \ 0 \\ 1 \mid 0 \ 1 \ a \\ a \mid 0 \ a \ 0 \end{array} \quad \begin{array}{c} * \mid 0 \ 1 \ a \\ \hline 1 \mid 1 \ 1 \end{array}$$

We have that $1 \geq a$ and $1 \geq 0$, because $1 + a = 1$ and $1 + 0 = 1$, so that A_3 is a TopKAT with $\top = 1$. Moreover, $\top \cdot 1 \cdot a = 1 \cdot 1 \cdot a = a$, whereas there exists no $q \in \text{test}(A_3)$ satisfying $\top \cdot q = a$. Indeed, $\top \cdot 1 = 1 \cdot 1 = 1 \neq a$ and $\top \cdot 0 = 0 \neq a$. \square

In general, the lack of such a $q \in \text{test}(K)$ implies that the abstract domain cannot be defined as an abstraction of the set of topped-tests $\{\top p \mid p \in \text{test}(K)\}$, because in such a case we could miss the abstraction $\alpha(\top p \llbracket \mathbf{a} \rrbracket_u)$. To settle this issue, an abstract domain must provide an abstraction of the larger set $\text{top}(K) \triangleq \{\top a \mid a \in K\}$ which contains all the multiplicative elements $\top a$.

Definition 5.2 (Top Kleene abstract domain). A poset (A, \leq) is a *top Kleene abstract domain* of a TopKAT K if: (1) There exists a Galois Insertion $\gamma : A \rightarrow \text{top}(K)$ and $\alpha : \text{top}(K) \rightarrow A$ of the poset (A, \leq_A) into the poset $(\text{top}(K), \leq_K)$; (2) A is countably-complete. \square

The abstract semantic function $\llbracket \cdot \rrbracket_A^\sharp : T_{\Sigma, B} \rightarrow (A \rightarrow A)$ on a top Kleene abstraction A can be therefore defined for the base cases $\mathbf{a} \in \text{Atom}$ as

$$\llbracket \mathbf{a} \rrbracket_A^\sharp p^\sharp \triangleq \alpha(\gamma(p^\sharp) \llbracket \mathbf{a} \rrbracket_u),$$

while the remaining inductive cases are defined as in (18) for Kleene abstractions. The monotonicity and soundness properties of this abstract semantics hold, provided that the TopKAT is \ast -continuous³, which is referred to as TopKAT \ast .

Theorem 5.3 (Soundness of TopKAT Abstract Semantics). *Let A be a Kleene abstraction of a TopKAT \ast K and $T_{\Sigma, B}$ a language interpreted on K . For all $p^\sharp, q^\sharp \in A$, $a \in K$ and $\mathbf{t} \in T_{\Sigma, B}$:*

$$\begin{aligned} p^\sharp \leq_A q^\sharp &\Rightarrow \llbracket \mathbf{t} \rrbracket_A^\sharp p^\sharp \leq_A \llbracket \mathbf{t} \rrbracket_A^\sharp q^\sharp && \text{(monotonicity)} \\ \alpha(\top a \llbracket \mathbf{t} \rrbracket_u) &\leq_A \llbracket \mathbf{t} \rrbracket_A^\sharp \alpha(\top a) && \text{(soundness)} \end{aligned}$$

Proof. The first property can be proved by induction on the structure of $\mathbf{t} \in T_{\Sigma, B}$. Notice that the inductive cases are the same as Theorem 3.5, therefore we only need to verify the base case.

$$\llbracket \mathbf{a} \rrbracket_A^\sharp p^\sharp = \alpha(\gamma(p^\sharp) \llbracket \mathbf{a} \rrbracket_u) \leq \alpha(\gamma(q^\sharp) \llbracket \mathbf{a} \rrbracket_u) = \llbracket \mathbf{a} \rrbracket_A^\sharp q^\sharp$$

where we used the isotony of $\alpha(\cdot)$, $\gamma(\cdot)$ and \cdot .

Likewise we prove the second property by induction on the structure of $\mathbf{t} \in T_{\Sigma, B}$. The computation can be easily adapted from Theorem 3.5. As an example we explicitly prove the \mathbf{t}_0^\ast case.

(\mathbf{t}_0^\ast): In order to prove the result we need a preliminary fact

$$\alpha(\top a (\llbracket \mathbf{t}_0 \rrbracket_u)^n) \leq (\llbracket \mathbf{t}_0 \rrbracket_A^\sharp)^n \alpha(\top a) \tag{35}$$

which can be shown by induction on n . The base case holds as $\alpha(\top a (\llbracket \mathbf{t}_0 \rrbracket_u)^0) = \alpha(\top a) = (\llbracket \mathbf{t}_0 \rrbracket_A^\sharp)^0 \alpha(\top a)$. The inductive case can be proved as follows

$$\alpha(\top a (\llbracket \mathbf{t}_0 \rrbracket_u)^{n+1}) = \alpha(\top a (\llbracket \mathbf{t}_0 \rrbracket_u)^n \llbracket \mathbf{t}_0 \rrbracket_u) \leq \llbracket \mathbf{t}_0 \rrbracket_A^\sharp \alpha(\top a (\llbracket \mathbf{t}_0 \rrbracket_u)^n) \leq (\llbracket \mathbf{t}_0 \rrbracket_A^\sharp)^{n+1} \alpha(\top a)$$

³ This condition plays a role similar to the CTC condition for bdKATs.

The previous fact and the $*$ -continuity of K yield the soundness

$$\begin{aligned}
& \alpha(\top a \llbracket \mathfrak{t}_0^* \rrbracket_u) = \\
& \alpha(\top a (\llbracket \mathfrak{t}_0 \rrbracket_u)^*) = [\text{By } *\text{-continuity}] \\
& \alpha\left(\bigvee_{n \in \mathbb{N}} \top a (\llbracket \mathfrak{t}_0 \rrbracket_u)^n\right) = [\text{By additivity of } \alpha(\cdot)] \\
& \bigvee_{n \in \mathbb{N}} \alpha(\top a (\llbracket \mathfrak{t}_0 \rrbracket_u)^n) \leq [\text{By (35)}] \\
& \bigvee_{n \in \mathbb{N}} (\llbracket \mathfrak{t}_0 \rrbracket_A^\sharp)^n \alpha(\top a) = \\
& \llbracket \mathfrak{t}_0^* \rrbracket_A^\sharp \alpha(\top a). \quad \square
\end{aligned}$$

5.2 Local Completeness Logic on TopKAT

Completeness and triple validity are adapted to the TopKAT framework as follows. Given a Top Kleene abstract domain A on a TopKAT* K , A is defined to be locally complete for $a \in K$ on an element $b \in K$, denoted by $\mathbb{C}_b^A(a)$, when

$$A(\top ba) = A(A(\top b)a).$$

Moreover, A is globally complete for a , denoted $\mathbb{C}^A(a)$, iff it is locally complete for any $b \in K$.

Likewise, a triple $[a] \mathfrak{t} [b]$, with $a, b \in K$ and $\mathfrak{t} \in T_{\Sigma, B}$, is valid, denoted by $\models_A^{\text{TK}} [a] \mathfrak{t} [b]$, when:

$$(1) \top b \leq \top a \llbracket \mathfrak{t} \rrbracket_u; \quad (2) \llbracket \mathfrak{t} \rrbracket_A^\sharp \alpha(\top a) = \alpha(\top b) = \alpha(\top a \llbracket \mathfrak{t} \rrbracket_u).$$

The corresponding proof system, denoted by LCTK_A , has the same rules of LCK_A in Figure 1 except (transfer), (relax) and (iterate) which are modified as follows:

$$\begin{aligned}
& \frac{c \in \text{Atom} \quad \mathbb{C}_a^A(\llbracket c \rrbracket_u)}{\vdash_A^{\text{TK}} [a] c [a \llbracket c \rrbracket_u]} \text{ (transfer)} \\
& \frac{\top a' \leq \top a \leq A(\top a') \quad \vdash_A^{\text{TK}} [a'] \mathfrak{t} [b'] \quad \top b \leq \top b' \leq A(\top b)}{\vdash_A^{\text{TK}} [a] \mathfrak{t} [b]} \text{ (relax)} \\
& \frac{\vdash_A^{\text{TK}} [a] \mathfrak{t} [b] \quad \top b \leq A(\top a)}{\vdash_A^{\text{TK}} [a] \mathfrak{t}^* [a + b]} \text{ (iterate)}
\end{aligned}$$

This incarnation LCTK_A of local completeness logic for TopKAT* turns out to be logically sound and, with additional hypotheses, complete.

Theorem 5.4 (Logical Soundness of \vdash_A^{TK}). *If $\vdash_A^{\text{TK}} [a] \mathfrak{t} [b]$ then*

- (i) $\top b \leq \top a \llbracket \mathfrak{t} \rrbracket_u$;
- (ii) $\llbracket \mathfrak{t} \rrbracket_A^\sharp \alpha(\top a) = \alpha(\top b) = \alpha(\top a \llbracket \mathfrak{t} \rrbracket_u)$.

Proof. The proof is as in Theorem 3.7. As an example we explicitly prove the cases (transfer) and (iterate).

(transfer): (i) is immediate because $\top b = \top a \llbracket c \rrbracket_u \leq \top a \llbracket c \rrbracket_u$. (ii) instead is a consequence of local completeness $\llbracket c \rrbracket_A^\sharp \alpha(\top a) = \alpha(\gamma(\alpha(\top a)) \llbracket c \rrbracket_u) \stackrel{\text{LC}}{=} \alpha(\top a \llbracket c \rrbracket_u) = \alpha(\top b)$.

(iterate): For any $k \in K$ it holds $1 \leq k^*$ and $k \leq k^*$ [15, Section 2.1]. In particular $\top b \leq \top a \llbracket t \rrbracket_u \leq \top a (\llbracket t \rrbracket_u)^*$ and $\top a = \top a \cdot 1 \leq \top a (\llbracket t \rrbracket_u)^*$, thus implying (i) $\top(a + b) \leq \top a (\llbracket t \rrbracket_u)^* = \top a \llbracket t^* \rrbracket_u$. To show the (ii) we need a preliminary fact

$$\llbracket t \rrbracket_A^\sharp \alpha(\top a) \leq \alpha(\top a) \Rightarrow \bigvee_{n \in \mathbb{N}} (\llbracket t \rrbracket_A^\sharp)^n \alpha(\top a) = \alpha(\top a) \quad (36)$$

As a first step, let us prove by induction on n that $\alpha(\top a)$ is an upper-bound of the elements of the disjunction, i.e., $\forall n \in \mathbb{N}. (\llbracket t \rrbracket_A^\sharp)^n \alpha(\top a) \leq \alpha(\top a)$. The base case is trivially true $(\llbracket t \rrbracket_A^\sharp)^0 \alpha(\top a) = \alpha(\top a) \leq \alpha(\top a)$, while the inductive case can be shown as follows: $(\llbracket t \rrbracket_A^\sharp)^{n+1} \alpha(\top a) = (\llbracket t \rrbracket_A^\sharp)^n \llbracket t \rrbracket_A^\sharp \alpha(\top a) \leq (\llbracket t \rrbracket_A^\sharp)^n \alpha(\top a) \leq \alpha(\top a)$. Notice that $\alpha(\top a)$ is also the least upper bound because it is part of the disjunction as $(\llbracket t \rrbracket_A^\sharp)^0 \alpha(\top a) = \alpha(\top a)$. The hypothesis $\top b \leq A(\top a)$ can be rewritten as $\alpha(\top b) \leq \alpha(\top a)$ by isotony of $\alpha(\cdot), \gamma(\cdot)$, (21), (23). Finally, $\llbracket t \rrbracket_A^\sharp \alpha(\top a) = \alpha(\top b) \leq \alpha(\top a)$, meaning that the premise of (36) holds, so that we can conclude

$$\llbracket t^* \rrbracket_A^\sharp \alpha(\top a) = \bigvee_{n \in \mathbb{N}} (\llbracket t \rrbracket_A^\sharp)^n \alpha(\top a) = \alpha(\top a) = \alpha(\top a) + \alpha(\top b) = \alpha(\top(a + b))$$

□

Logical completeness needs the following additional conditions:

(a) Likewise LCK_A , the same infinitary rule for Kleene star:

$$\frac{\forall n \in \mathbb{N}. \vdash_A^{\text{TK}} [a_n] \mathfrak{t} [a_{n+1}]}{\vdash_A^{\text{TK}} [a_0] \mathfrak{t}^* [\bigvee_{n \in \mathbb{N}} a_n]} \text{ (limit)}$$

where we assume that:

- $\bigvee_{n \in \mathbb{N}} a_n$ always exists. Let us remark that for bdKAT , such explicit condition was not needed, as it was entailed by the CTC requirement on the KAT.
- \top distributes over $\bigvee_{n \in \mathbb{N}} a_n$, i.e., $\top \bigvee_{n \in \mathbb{N}} a_n = \bigvee_{n \in \mathbb{N}} \top a_n$.

(b) Global completeness of all the primitive actions and tests occurring in the program.

As in LCK_A , (limit) is sound:

Lemma 5.5. *With the hypothesis of Theorem 5.4, the rule (limit) is logically sound.*

Proof. Instantiating (i) we obtain $\top \bigvee_{n \in \mathbb{N}} a_n \leq \top a_0 \llbracket t^* \rrbracket_u$. The distributivity condition of (limit) yields $\top \bigvee_{n \in \mathbb{N}} a_n = \bigvee_{n \in \mathbb{N}} \top a_n$. For this reason (i) is equivalent to $\forall n \in \mathbb{N}. \top a_0 \llbracket t^* \rrbracket_u \geq \top a_n$. This fact can be shown by induction on n .

For $n = 0$: $\top a_0 \llbracket t^* \rrbracket_u = \top a_0 (\llbracket t \rrbracket_u)^* \geq \top a_0$, as for any $k \in K, k^* \geq 1$ by $(^*-\text{unfold})$.

For $n+1$: The rule has a premise $\vdash_A^{\text{TK}} [a_n] \mathfrak{t} [a_{n+1}]$, and by induction (on the derivation tree) we have that $\models_A^{\text{TK}} [a_n] \mathfrak{t} [a_{n+1}]$.

$$\begin{aligned}
& \top a_0 \llbracket \mathfrak{t}^* \rrbracket_u = \\
& \top a_0 (\llbracket \mathfrak{t} \rrbracket_u)^* \geq [\text{By } (*\text{-unfold})] \\
& \top a_0 (1 + (\llbracket \mathfrak{t} \rrbracket_u)^* \llbracket \mathfrak{t} \rrbracket_u) \geq \\
& \top a_0 (\llbracket \mathfrak{t} \rrbracket_u)^* \llbracket \mathfrak{t} \rrbracket_u = \\
& \top a_0 \llbracket \mathfrak{t}^* \rrbracket_u \llbracket \mathfrak{t} \rrbracket_u \geq \\
& \top a_n \llbracket \mathfrak{t} \rrbracket_u \geq [\text{By } \models_A^{\text{TK}} [a_n] \mathfrak{t} [a_{n+1}]] \\
& \top a_{n+1}
\end{aligned}$$

For condition (ii) instead we need to show $\llbracket \mathfrak{t}^* \rrbracket_A^\# \alpha(\top a_0) = \alpha(\top \bigvee_{n \in \mathbb{N}} a_n)$, provided that $\llbracket \mathfrak{t} \rrbracket_A^\# \alpha(\top a_n) = \alpha(\top a_{n+1})$. By definition, $\llbracket \mathfrak{t}^* \rrbracket_A^\# \alpha(\top a_0) = \bigvee_{n \in \mathbb{N}} (\llbracket \mathfrak{t} \rrbracket_A^\#)^n \alpha(\top a_0)$ and we have also that $\alpha(\top \bigvee_{n \in \mathbb{N}} a_n) = \alpha(\bigvee_{n \in \mathbb{N}} \top a_n) = \bigvee_{n \in \mathbb{N}} \alpha(\top a_n)$. Therefore (ii) is implied by

$$\forall n \in \mathbb{N}. (\llbracket \mathfrak{t} \rrbracket_A^\#)^n \alpha(\top a_0) = \alpha(\top a_n)$$

That can be show by induction on n . The base case is trivial ($(\llbracket \mathfrak{t} \rrbracket_A^\#)^0 \alpha(\top a_0) = \alpha(\top a_0)$), while the inductive can be shown as follows

$$(\llbracket \mathfrak{t} \rrbracket_A^\#)^{n+1} \alpha(\top a_0) = \llbracket \mathfrak{t} \rrbracket_A^\# (\llbracket \mathfrak{t} \rrbracket_A^\#)^n \alpha(\top a_0) = \llbracket \mathfrak{t} \rrbracket_A^\# \alpha(\top a_n) = \alpha(\top a_{n+1}). \quad \square$$

Condition (b) entails global completeness:

Lemma 5.6. *Let K a TopKAT*, A a top Kleene abstract domain on K and $T_{\Sigma, B}$ a KAT language on K . For any $\mathfrak{t} \in T_{\Sigma, B}$ and $a \in K$ we have*

$$(\forall \mathfrak{b} \in \text{Atom}(\mathfrak{t}). \mathbb{C}^A(\llbracket \mathfrak{b} \rrbracket_u)) \Rightarrow \llbracket \mathfrak{t} \rrbracket_A^\# \alpha(\top a) = \alpha(\top a \llbracket \mathfrak{t} \rrbracket_u)$$

Proof. The proof is as in Lemma 3.9. We only prove explicitly the \mathfrak{t}_0^* case.

(\mathfrak{t}_0^*): The result is a consequence of the following claim

$$\forall n \in \mathbb{N}. (\llbracket \mathfrak{t}_0 \rrbracket_A^\#)^n \alpha(\top a) = \alpha(\top a (\llbracket \mathfrak{t}_0 \rrbracket_u)^n) \quad (37)$$

that can be proved by induction. The base case is as follows: $(\llbracket \mathfrak{t}_0 \rrbracket_A^\#)^0 \alpha(\top a) = \alpha(\top a) = \alpha(\top a (\llbracket \mathfrak{t}_0 \rrbracket_u)^0)$. The inductive case can be proved as follows

$$\begin{aligned}
& (\llbracket \mathfrak{t}_0 \rrbracket_A^\#)^{n+1} \alpha(\top a) = \\
& \llbracket \mathfrak{t}_0 \rrbracket_A^\# (\llbracket \mathfrak{t}_0 \rrbracket_A^\#)^n \alpha(\top a) = \\
& \llbracket \mathfrak{t}_0 \rrbracket_A^\# \alpha(\top a (\llbracket \mathfrak{t}_0 \rrbracket_u)^n) = \\
& \alpha(\top a (\llbracket \mathfrak{t}_0 \rrbracket_u)^n \llbracket \mathfrak{t}_0 \rrbracket_u) = \\
& \alpha(\top a (\llbracket \mathfrak{t}_0 \rrbracket_u)^{n+1})
\end{aligned}$$

Moreover by $*$ -continuity we have that $\top a \llbracket \mathfrak{t}_0^* \rrbracket_u = \top a(\llbracket \mathfrak{t}_0 \rrbracket_u)^* = \bigvee_{n \in \mathbb{N}} \top a(\llbracket \mathfrak{t}_0 \rrbracket_u)^n$, meaning that

$$\begin{aligned}
& \llbracket \mathfrak{t}_0^* \rrbracket_A^\# \alpha(\top a) = \\
& \bigvee_{n \in \mathbb{N}} (\llbracket \mathfrak{t}_0 \rrbracket_A^\#)^n \alpha(\top a) = [\text{By (37)}] \\
& \bigvee_{n \in \mathbb{N}} \alpha(\top a(\llbracket \mathfrak{t}_0 \rrbracket_u)^n) = [\text{Additivity of } \alpha(\cdot)] \\
& \alpha\left(\bigvee_{n \in \mathbb{N}} \top a(\llbracket \mathfrak{t}_0 \rrbracket_u)^n\right) = \\
& \alpha(\top a \llbracket \mathfrak{t}_0^* \rrbracket_u) \quad \square
\end{aligned}$$

Theorem 5.7 (Logical Completeness of \vdash_A^{TK}). *Assume that conditions (a) and (b) hold. If $\models_A^{\text{TK}} [a] \mathfrak{t} [b]$ then $\vdash_A^{\text{TK}} [a] \mathfrak{t} [b]$.*

Proof. The proof is as in Theorem 3.10. The case (\mathfrak{t}_0^*) requires special attention, and is proved explicitly.

(\mathfrak{t}_0^*) : Let $a_n = a_0(\llbracket \mathfrak{t}_0 \rrbracket_u)^n$. As a first step we prove that $\forall n \in \mathbb{N}. \models_A^{\text{TK}} [a_n] \mathfrak{t}_0 [a_{n+1}]$: condition (i) holds because $\top a_{n+1} = \top a_0(\llbracket \mathfrak{t}_0 \rrbracket_u)^{n+1} = \top a_0(\llbracket \mathfrak{t}_0 \rrbracket_u)^n \llbracket \mathfrak{t}_0 \rrbracket_u = \top a_n \llbracket \mathfrak{t}_0 \rrbracket_u$, while (ii) holds by Lemma 5.6, $\llbracket \mathfrak{t}_0 \rrbracket_A^\# \alpha(\top a_n) = \alpha(\top a_n \llbracket \mathfrak{t}_0 \rrbracket_u) = \alpha(\top a_{n+1})$. Finally, the inductive hypothesis yields $\forall n \in \mathbb{N}. \vdash_A^{\text{TK}} [a_n] \mathfrak{t}_0 [a_{n+1}]$.

We want to use the rule (limit) with postcondition $\bigvee_{n \in \mathbb{N}} a_n$. To do so, we need to show two additional facts: 1) the disjunction exists 2) \top distributes over the disjunction. 1) holds as $a_0(\llbracket \mathfrak{t}_0 \rrbracket_u)^* = \bigvee_{n \in \mathbb{N}} a_0(\llbracket \mathfrak{t}_0 \rrbracket_u)^n = \bigvee_{n \in \mathbb{N}} a_n$, where the first equality holds by $*$ -continuity. By $*$ -continuity we have that $\top a_0(\llbracket \mathfrak{t}_0 \rrbracket_u)^* = \bigvee_{n \in \mathbb{N}} \top a_0(\llbracket \mathfrak{t}_0 \rrbracket_u)^n = \bigvee_{n \in \mathbb{N}} \top a_n$ and $\top a_0(\llbracket \mathfrak{t}_0 \rrbracket_u)^* = \top \bigvee_{n \in \mathbb{N}} a_0(\llbracket \mathfrak{t}_0 \rrbracket_u)^n = \top \bigvee_{n \in \mathbb{N}} a_n$, thus implying $\bigvee_{n \in \mathbb{N}} \top a_n = \top \bigvee_{n \in \mathbb{N}} a_n$. Finally, by (limit) we obtain the result $\vdash_A^{\text{TK}} [a] \mathfrak{t}_0^* [a \llbracket \mathfrak{t}_0^* \rrbracket_u]$. \square

Example 5.8. Consider a relational KAT $K = \wp(\mathbb{Z} \times \mathbb{Z})$ on the set of integers \mathbb{Z} , where $\mathbf{1}_K \triangleq \{(z, z) \mid z \in \mathbb{Z}\}$ and $\mathbf{0}_K \triangleq \emptyset$. Notice that $\mathbb{Z} \times \mathbb{Z} \in K$ is the top element \top of K , meaning that K is a TopKAT. Let us consider a language with primitive actions $\Sigma = \{x := x + 1\}$ and primitive tests $B = \{x \geq 0, x < 0\}$. The evaluation function $u : \Sigma \cup B \rightarrow K$ is defined as expected by the following relations:

$$\begin{aligned}
u(x := x + 1) &\triangleq \{(z, z + 1) \mid z \in \mathbb{Z}\}, & u(x \geq 0) &\triangleq \{(z, z) \mid z \in \mathbb{Z}, z \geq 0\}, \\
u(x < 0) &\triangleq \{(z, z) \mid z \in \mathbb{Z}, z < 0\}.
\end{aligned}$$

Consider the following sign abstraction $\text{Sign} \triangleq \{\mathbb{Z}, \mathbb{Z}_{<0}, \mathbb{Z}_{\neq 0}, \mathbb{Z}_{\geq 0}, \mathbb{Z}_{<0}, \mathbb{Z}_{=0}, \mathbb{Z}_{>0}, \emptyset\}$. We verify that the program

$$r \equiv ((x \geq 0) \cdot (x := x + 1))^* \cdot (x < 0)$$

does not terminate with precondition $p \triangleq \{(0, 0), (10, 10)\}$, i.e., we prove the specification $\text{Spec} \triangleq \emptyset$. Let us define the following auxiliary elements: $q \triangleq \{(1, 1), (11, 11)\}$, $s \triangleq p + q$, $t_{\geq 0} \triangleq \{(x, z) \mid x \in \mathbb{Z}, z \in \mathbb{Z}_{\geq 0}\}$, and observe that $\text{Sign}(t_{\geq 0}) = t_{\geq 0}$.

The following local completeness conditions for the atoms hold:

$$\begin{aligned}\alpha(\text{Sign}(\top p)\llbracket \mathbf{x} \geq 0 \rrbracket_u) &= \alpha(t_{\geq 0}\llbracket \mathbf{x} \geq 0 \rrbracket_u) = \mathbb{Z}_{\geq 0} = \alpha(\top p\llbracket \mathbf{x} \geq 0 \rrbracket_u) \\ \alpha(\text{Sign}(\top p)\llbracket \mathbf{x} := \mathbf{x} + 1 \rrbracket_u) &= \alpha(t_{\geq 0}\llbracket \mathbf{x} := \mathbf{x} + 1 \rrbracket_u) = \mathbb{Z}_{> 0} = \alpha(\top p\llbracket \mathbf{x} := \mathbf{x} + 1 \rrbracket_u) \\ \alpha(\text{Sign}(\top s)\llbracket \mathbf{x} < 0 \rrbracket_u) &= \alpha(t_{\geq 0}\llbracket \mathbf{x} < 0 \rrbracket_u) = \emptyset = \alpha(\top s\llbracket \mathbf{x} < 0 \rrbracket_u)\end{aligned}$$

Moreover, we also have that:

$$\top q = \{(x, z) \mid x \in \mathbb{Z}, z \in \{1, 11\}\} \leq t_{\geq 0} = \text{Sign}(\top p).$$

The following derivation shows that the triple $[p] \text{ r } [\mathbf{0}_K]$ is provable:

$$\frac{\frac{\frac{\mathbb{C}_p^{\text{Sign}}(\llbracket \mathbf{x} \geq 0 \rrbracket_u)}{\vdash_{\text{Sign}}^{\text{TK}} [p] \mathbf{x} \geq 0} \text{ (transfer)}}{\vdash_{\text{Sign}}^{\text{TK}} [p] (\mathbf{x} \geq 0) \cdot (\mathbf{x} := \mathbf{x} + 1) [q]} \text{ (seq)}}{\frac{\frac{\frac{\mathbb{C}_p^{\text{Sign}}(\llbracket \mathbf{x} := \mathbf{x} + 1 \rrbracket_u)}{\vdash_{\text{Sign}}^{\text{TK}} [p] \mathbf{x} := \mathbf{x} + 1} \text{ (transfer)}}{\vdash_{\text{Sign}}^{\text{TK}} [p] ((\mathbf{x} \geq 0) \cdot (\mathbf{x} := \mathbf{x} + 1))^* [s]} \text{ (iterate)}}{\vdash_{\text{Sign}}^{\text{TK}} [p] ((\mathbf{x} \geq 0) \cdot (\mathbf{x} := \mathbf{x} + 1))^* \cdot (\mathbf{x} < 0) [\mathbf{0}_K]} \text{ (seq)}} \text{ (seq)}$$

By Theorem 5.4, we have that $\top \mathbf{0}_K \subseteq \top p\llbracket \mathbf{x} \rrbracket_u \subseteq \text{Sign}(\top \mathbf{0}_K) = \emptyset$, meaning that the program does not terminate and Spec is satisfied as $\top p\llbracket \mathbf{x} \rrbracket_u = \emptyset = \top \text{Spec}$. \square

5.3 Relationship with Under-Approximation Logic

We have shown in Section 3.6 that the backward-diamond formulation of LCK generalizes UL. The same can be done for the TopKAT formulation. A TopKAT version of the UL proof system was already proposed in [32, Figure 6]. The reduction here considered refers to such system, with the following minor differences:

- We consider only propositional fragments of the logic, meaning that the rules (assume) and (identity) are replaced by the following single rule (transfer):

$$\frac{c \in \text{Atom}}{\vdash_{\text{UL}} [a] c [a\llbracket c \rrbracket_u]} \text{ (transfer)}$$

- The premises of the rule (consequence) in [32, Figure 6], $b \leq b'$ and $c' \leq c$, are relaxed to $\top b \leq \top b'$ and $\top c' \leq \top c$. Notice that the former implies the latter. Furthermore, the soundness proof of [32, Theorem 4] is not affected by this change, because $(\top b' \geq \top b \wedge \top c \geq \top c' \wedge \top bp \geq c) \Rightarrow \top b'p \geq \top bp \geq \top c \geq \top c'$, and, by [32, Theorem 3], it holds that $\top b'p \geq \top c'$ entails $\top b'p \geq c'$.
- The (limit) rules of LCTK_A and UL differ on the distributivity condition. We assume that distributivity also holds in UL.

By instantiating to the trivial abstract domain A_{tr} , it turns out that the two proof systems become equivalent.

Theorem 5.9 ($\text{LCTK}_{A_{tr}} \equiv \text{UL}$). *Let K be a TopKAT*. For any $a, b \in K$, $\mathfrak{t} \in T_{\Sigma, B}$:*

$$\vdash_{A_{tr}}^{\text{TK}} [a] \mathfrak{t} [b] \Leftrightarrow \vdash_{\text{UL}} [a] \mathfrak{t} [b].$$

Proof. The \Rightarrow direction can be shown as in Theorem 3.11. We focus instead on the opposite direction.

\Leftarrow) The proof is by induction on the derivation tree of $\vdash_{\text{UL}} [a] \mathfrak{t} [b]$. The rules of Table 1 are immediate, meaning that only (empty),(choice),(disj),(iterate zero) and (iterate non-zero) are left to prove. As in Theorem 3.11, we leverage the completeness of the proof system and likewise, in order to prove that a triple is valid, it is enough to show that condition (1) holds because (2) is trivially true.

(empty): $\models_{A_{tr}}^{\text{TK}} [a] \mathfrak{t} [0]$ is trivial since $\top \cdot 0 = 0 \leq \top a \llbracket \mathfrak{t} \rrbracket_u$ is always true, therefore by completeness of $\text{LCTK}_{A_{tr}}$, $\vdash_{A_{tr}}^{\text{TK}} [a] \mathfrak{t} [0]$.

(choice): By induction we have, without loss of generality, $\vdash_{A_{tr}}^{\text{TK}} [a] \mathfrak{t}_1 [b]$ and we need to show $\vdash_{A_{tr}}^{\text{TK}} [a] \mathfrak{t}_1 + \mathfrak{t}_2 [b]$. Notice that for any $\mathfrak{t}_2 \in T_{\Sigma, B}$, $\models_{A_{tr}}^{\text{TK}} [a] \mathfrak{t}_2 [0]$ because $\top \cdot 0 = 0 \leq \top a \llbracket \mathfrak{t}_2 \rrbracket_u$, therefore by completeness of $\text{LCTK}_{A_{tr}}$ $\vdash_{A_{tr}}^{\text{TK}} [a] \mathfrak{t}_2 [0]$. Finally, (join) yields the result $\vdash_{A_{tr}}^{\text{TK}} [a] \mathfrak{t}_1 + \mathfrak{t}_2 [b]$.

(disj): By induction we have $\vdash_{A_{tr}}^{\text{TK}} [a_1] \mathfrak{t} [b_1]$ and $\vdash_{A_{tr}}^{\text{TK}} [a_2] \mathfrak{t} [b_2]$ and we need to show $\vdash_{A_{tr}}^{\text{TK}} [a_1 + a_2] \mathfrak{t} [b_1 + b_2]$. By soundness $\top b_1 \leq \top a_1 \llbracket \mathfrak{t} \rrbracket_u$ and $\top b_2 \leq \top a_2 \llbracket \mathfrak{t} \rrbracket_u$, so that $\top (b_1 + b_2) \leq \top (a_1 + a_2) \llbracket \mathfrak{t} \rrbracket_u$. Finally, by completeness we retrieve the result: $\vdash_{A_{tr}}^{\text{TK}} [a_1 + a_2] \mathfrak{t} [b_1 + b_2]$.

(iterate zero): By (*-unfold) $\llbracket \mathfrak{t}^* \rrbracket_u = (\llbracket \mathfrak{t} \rrbracket_u)^* \geq 1$, so that $\top a \llbracket \mathfrak{t}^* \rrbracket_u \geq \top a$ and by completeness $\vdash_{A_{tr}}^{\text{TK}} [a] \mathfrak{t}^* [a]$.

(iterate non-zero): By (*-unfold) $\llbracket \mathfrak{t}^* \rrbracket_u = (\llbracket \mathfrak{t} \rrbracket_u)^* \geq (\llbracket \mathfrak{t} \rrbracket_u)^* \llbracket \mathfrak{t} \rrbracket_u = \llbracket \mathfrak{t}^* \rrbracket_u \llbracket \mathfrak{t} \rrbracket_u = \llbracket \mathfrak{t}^* \cdot \mathfrak{t} \rrbracket_u$. Thus, by hypothesis $\top b \leq \top a \llbracket \mathfrak{t}^* \cdot \mathfrak{t} \rrbracket_u \leq \top a \llbracket \mathfrak{t}^* \rrbracket_u$ and by completeness $\vdash_{A_{tr}}^{\text{TK}} [a] \mathfrak{t}^* [b]$. \square

Finally, the logical soundness and completeness of UL can be retrieved as a consequence of those of LCTK.

Corollary 5.10. *Under the same hypotheses of Theorem 5.9, the proof system UL is sound and complete, that is, $\vdash_{\text{UL}} [p] \mathfrak{t} [q] \Leftrightarrow \models_{\text{UL}} [p] \mathfrak{t} [q]$.*

Proof.

$$\begin{aligned} \vdash_{\text{UL}} [p] \mathfrak{t} [q] &\Leftrightarrow \text{[by Theorem 5.9]} \\ \vdash_{A_{tr}}^{\text{TK}} [p] \mathfrak{t} [q] &\Leftrightarrow \text{[by Theorems 5.4, 5.7]} \\ \models_{A_{tr}}^{\text{TK}} [p] \mathfrak{t} [q] &\Leftrightarrow \\ \models_{\text{UL}} [p] \mathfrak{t} [q] &\quad \square \end{aligned}$$

6 Incorrectness Logic in TopKAT

In Section 5 we proposed an alternative formulation of local completeness logic based on a TopKAT. Here, we take a similar path as in Section 4 in adding support for abnormal termination. Like in the modal formulation, each language term is interpreted as a pair of KAT elements, meaning that the interpretation function has signature $\llbracket \cdot \rrbracket_u : T_{\Sigma, B} \rightarrow (K \times K)$ and is defined inductively as in (30). Following the approach

of Section 5 we consider an element $\top pa$ as a proxy for the strongest postcondition and in particular we have two such elements corresponding to respectively normal and abnormal termination: $\top p[[\mathfrak{t}]]_{u_{ok}}$ and $\top p[[\mathfrak{t}]]_{u_{err}}$.

The abstract semantic is a function of type $[[\cdot]]_A^\sharp : T_{\Sigma, B} \rightarrow (A \rightarrow (A \times A))$, where A is a top Kleene abstract domain. As usual it is defined inductively, with base case $a \in \text{Atom}$ defined as

$$[[a]]_A^\sharp p^\sharp \triangleq \mathbf{ok} : \alpha(\gamma(p^\sharp)[[a]]_{u_{ok}}), \mathbf{err} : \alpha(\gamma(p^\sharp)[[a]]_{u_{err}})$$

and the inductive cases as in (33). It turns out that if K is a TopKAT* then the abstract semantic is sound and monotone.

6.1 Local completeness incorrectness logic on TopKAT

The proof system LCIL can be adapted to the TopKAT framework as follows. Let A be a top Kleene abstract domain on a TopKAT* K and a language $T_{\Sigma, B}$. If $a, b \in K$ and $\mathfrak{t} \in T_{\Sigma, B}$, then

- The triple $[a] \mathfrak{t} [\mathbf{ok} : b]$ is valid (1) $\top b \leq \top a[[\mathfrak{t}]]_{u_{ok}}$, and (2) $[[\mathfrak{t}]]_{A_{ok}}^\sharp \alpha(\top a) = \alpha(\top b) = \alpha(\top a[[\mathfrak{t}]]_{u_{ok}})$.
- The triple $[a] \mathfrak{t} [\mathbf{err} : c]$ is valid (1) $\top c \leq \top a[[\mathfrak{t}]]_{u_{err}}$, and (2) $[[\mathfrak{t}]]_{A_{err}}^\sharp \alpha(\top a) = \alpha(\top c) = \alpha(\top a[[\mathfrak{t}]]_{u_{err}})$.
- A triple $[a] \mathfrak{t} [\mathbf{ok} : b][\mathbf{err} : c]$ is valid if both $[a] \mathfrak{t} [\mathbf{ok} : b]$ and $[a] \mathfrak{t} [\mathbf{err} : c]$ are valid. In particular, if $b = c$ then the triple $[a] \mathfrak{t} [\epsilon : b]$ is valid.

The proof system, denoted LCTIL_A , has the same rules as LCIL_A in Figure 3 except for the rules (transfer) and (relax), modified as follows:

$$\frac{c \in \text{Atom} \quad \mathbb{C}_a^A([\mathfrak{c}]_{u_{ok}}) \quad \mathbb{C}_a^A([\mathfrak{c}]_{u_{err}})}{\vdash_A^{\text{TK}} [a] \mathfrak{c} [\mathbf{ok} : a[[\mathfrak{c}]]_{u_{ok}}][\mathbf{err} : a[[\mathfrak{c}]]_{u_{err}}]} \text{ (transfer)}$$

$$\frac{\top a' \leq \top a \leq A(\top a') \quad \vdash_A^{\text{TK}} [a'] \mathfrak{t} [\epsilon : b'] \quad \top b \leq \top b' \leq A(\top b)}{\vdash_A^{\text{TK}} [a] \mathfrak{t} [\epsilon : b]} \text{ (relax)}$$

Moreover, for the rule (limit) we assume that the infinitary disjunction exists and that \top distributes over $\bigvee_{n \in \mathbb{N}} a_n$. Observe that these additional conditions are the same as (a) for the TopKAT formulation of the local completeness logic.

It turn out that LCTIL_A is logically sound and complete.

Theorem 6.1 (Logical Soundness of LCTIL_A). *The triples provable in LCTIL_A are valid.*

Proof. Analogous to the proof of Theorem 4.2. □

Theorem 6.2 (Logical completeness of LCTIL_A). *Assume that the atoms in $\mathfrak{t} \in T_{\Sigma, B}$ are globally complete, i.e., for all $a \in \text{Atom}$, $\mathbb{C}^A([\mathfrak{a}]_{u_{ok}})$ and $\mathbb{C}^A([\mathfrak{a}]_{u_{err}})$. If $[a] \mathfrak{t} [\mathbf{ok} : b][\mathbf{err} : c]$ is valid, then it is provable in LCTIL_A .*

Proof. It is easy to check that $[[\mathfrak{t}]]_{A_{ok}}^\sharp \alpha(\top a) = \alpha(\top a[[\mathfrak{t}]]_{u_{ok}})$ and $[[\mathfrak{t}]]_{A_{err}}^\sharp \alpha(\top a) = \alpha(\top a[[\mathfrak{t}]]_{u_{err}})$, as in Lemmas 4.3 and 5.6. Then we can proceed as in Theorem 4.4 to obtain the result. □

6.2 Relationship with Incorrectness logic

In Section 4.1 and 5.3 we have shown that the modal formulation of LCIL generalizes IL and LCTK generalizes UL. Likewise, it is possible to show that LCTIL generalizes IL. A TopKAT formulation of the IL proof system was already proposed in [32, Figure 4]. We refer to such system for the proposed reduction, with the same minor changes that we discussed in Section 5.3, namely: 1) we consider only propositional fragments of the logic, 2) the premises of (consequence) are relaxed to $\top b \leq \top b'$ and $\top c' \leq \top c$, 3) we assume that the distributivity condition is satisfied also in IL.

Similarly to the reductions already proposed in this paper, we let $A = A_{tr}$, where A_{tr} is the trivial abstract domain.

Theorem 6.3 (LCTIL \equiv IL). *Let K be a TopKAT*. For any $a, b \in K$, $\mathfrak{t} \in T_{\Sigma, B}$,*

$$\vdash_A^{\text{TK}} [a] \mathfrak{t} [\text{ok} : b][\text{err} : c] \Leftrightarrow \vdash_{\text{IL}} [a] \mathfrak{t} [\text{ok} : b][\text{err} : c]$$

Proof. Analogous to the proof of Theorem 4.6. □

Finally, we can retrieve the logical soundness and completeness of IL as a consequence of those of LCTIL.

Corollary 6.4. *Under the hypothesis of Theorem 6.3 the proof system IL is logically sound and complete, that is, $\vdash_{\text{IL}} [a] \mathfrak{t} [\text{ok} : b][\text{err} : c] \Leftrightarrow \models_{\text{IL}} [a] \mathfrak{t} [\text{ok} : b][\text{err} : c]$.*

7 Conclusion

This work has shown that the abstract interpretation-based local completeness logic introduced in [4] can be generalized to and interpreted in Kleene algebra with tests. In particular, we proved that this can be achieved both for KATs extended with a modal backward diamond operator playing the role of strongest postcondition, and for KATs endowed with a top element. Our results generalize both the modal [22] and top [32] KAT approaches that encode Hoare correctness and O'Hearn incorrectness logic using different classes of KATs. In particular, our KAT-based logic leverages an abstract interpretation of KAT, a problem that was not studied so far.

Our plan for future work includes, but is not limited to, the following questions.

- For a KAT with top \top , following the technical idea underlying the approach by Zhang et al. [32], we defined an abstract domain as an approximation of all the algebraic elements of type $\top \cdot a$, where a is any element of the KAT (cf. Definition 5.2). Although this definition technically works, it is somehow artificial, in the sense that elements $\top \cdot a$ do not carry a clear intuitive meaning. As an interesting future task, we would like to characterize under which conditions an element $\top \cdot a$ coincides with $\top \cdot p$ for some test $p \in \text{test}(K)$, and if such test p is unique.
- This work is a first step towards an *algebraic and equational* approach to abstract interpretation. We envisage that the reasoning made by an abstract interpreter of programs could be made purely equational within a KAT equipped with a suitable collection of axioms. The ambition would be to conceive a notion of *abstract Kleene algebra* (AKA) making this slogan true: *AKA is for the abstract interpretation of programs what KAT is for concrete interpretation of programs.*

Acknowledgements. This research has been funded by the *Italian MIUR*, under the PRIN2017 project no. 201784YSZ5 “AnalysiS of PRogram Analyses (ASPRA)”, by *Facebook Research*, under a “Probability and Programming Research Award”, and by a *Meta Research* gift.

References

1. Anderson, C.J., Foster, N., Guha, A., Jeannin, J.B., Kozen, D., Schlesinger, C., Walker, D.: NetKAT: Semantic foundations for networks. In: Proceedings of the 41st ACM SIGPLAN-SIGACT Symposium on Principles of Programming Languages. p. 113–126. POPL ’14, ACM (2014). <https://doi.org/10.1145/2535838.2535862>
2. Beckett, R., Greenberg, M., Walker, D.: Temporal NetKAT. In: Proceedings of the 37th ACM SIGPLAN Conference on Programming Language Design and Implementation, PLDI 2016. pp. 386–401. ACM (2016). <https://doi.org/10.1145/2908080.2908108>
3. Bruni, R., Giacobazzi, R., Gori, R., Garcia-Contreras, I., Pavlovic, D.: Abstract extensionality: on the properties of incomplete abstract interpretations. Proc. ACM Program. Lang. **4**(POPL), 28:1–28:28 (2020). <https://doi.org/10.1145/3371096>, <https://doi.org/10.1145/3371096>
4. Bruni, R., Giacobazzi, R., Gori, R., Ranzato, F.: A Logic for Locally Complete Abstract Interpretations. In: Proceedings 36th ACM/IEEE Symposium on Logic in Computer Science (LICS 2021). pp. 1–13. IEEE (2021). <https://doi.org/10.1109/LICS52264.2021.9470608>
5. Cohen, E., Kozen, D., Smith, F.: The complexity of Kleene algebra with tests. Tech. rep., Cornell University, USA (1996), <https://www.cs.cornell.edu/~kozen/Papers/ckat>
6. Conway, J.: Regular Algebra and Finite Machines. Chapman and Hall mathematics series, Dover Publications (2012)
7. Cousot, P., Cousot, R.: Abstract interpretation: a unified lattice model for static analysis of programs by construction or approximation of fixpoints. In: Proceedings of the 4th ACM SIGACT-SIGPLAN Symposium on Principles of Programming Languages (POPL 1977). pp. 238–252. ACM (1977). <https://doi.org/10.1145/512950.512973>
8. Cousot, P., Cousot, R.: Systematic design of program analysis frameworks. In: Proceedings of the 6th ACM SIGACT-SIGPLAN Symposium on Principles of Programming Languages (POPL 1979). pp. 269–282. ACM (1979). <https://doi.org/10.1145/567752.567778>
9. Desharnais, J., Möller, B., Struth, G.: Kleene algebra with domain. ACM Trans. Comput. Logic **7**(4), 798–833 (oct 2006). <https://doi.org/10.1145/1183278.1183285>
10. Fischer, M.J., Ladner, R.E.: Propositional dynamic logic of regular programs. Journal of Computer and System Sciences **18**(2), 194–211 (Apr 1979). [https://doi.org/10.1016/0022-0000\(79\)90046-1](https://doi.org/10.1016/0022-0000(79)90046-1)
11. Foster, N., Kozen, D., Milano, M., Silva, A., Thompson, L.: A coalgebraic decision procedure for NetKAT. In: Proceedings of the 42nd Annual ACM SIGPLAN-SIGACT Symposium on Principles of Programming Languages, POPL 2015. pp. 343–355. ACM (2015). <https://doi.org/10.1145/2676726.2677011>
12. Giacobazzi, R., Ranzato, F., Scozzari, F.: Making abstract interpretation complete. Journal of the ACM **47**(2), 361–416 (March 2000). <https://doi.org/10.1145/333979.333989>, <https://doi.org/10.1145/333979.333989>
13. Giacobazzi, R., Logozzo, F., Ranzato, F.: Analyzing program analyses. In: Proceedings of the 42nd Annual ACM SIGPLAN-SIGACT Symposium on Principles of Programming Languages, POPL 2015. pp. 261–273 (2015). <https://doi.org/10.1145/2676726.2676987>, <http://doi.acm.org/10.1145/2676726.2676987>

14. Hoare, C.A.R.: An axiomatic basis for computer programming. *Commun. ACM* **12**(10), 576–580 (oct 1969). <https://doi.org/10.1145/363235.363259>
15. Kozen, D.: A completeness theorem for Kleene algebras and the algebra of regular events. *Information and Computation* **110**(2), 366–390 (1994). <https://doi.org/https://doi.org/10.1006/inco.1994.1037>, <https://www.sciencedirect.com/science/article/pii/S0890540184710376>
16. Kozen, D.: Kleene algebra with tests. *ACM Transactions on Programming Languages and Systems* **19**(3), 427–443 (May 1997). <https://doi.org/10.1145/256167.256195>
17. Kozen, D.: On Hoare logic and Kleene algebra with tests. *ACM Trans. Comput. Logic* **1**(1), 60–76 (jul 2000). <https://doi.org/10.1145/343369.343378>
18. Kozen, D., Smith, F.: Kleene algebra with tests: Completeness and decidability. In: *Proceedings 10th International Workshop on Computer Science Logic, CSL 1996, Annual Conference of the EACSL. Lecture Notes in Computer Science*, vol. 1258, pp. 244–259. Springer (1996). https://doi.org/10.1007/3-540-63172-0_43
19. Le, Q.L., Raad, A., Villard, J., Berdine, J., Dreyer, D., O’Hearn, P.W.: Finding real bugs in big programs with incorrectness logic. *Proc. ACM Program. Lang.* **6**(OOPSLA1) (apr 2022). <https://doi.org/10.1145/3527325>
20. Mamouras, K.: Equational theories of abnormal termination based on kleene algebra. In: *Proceedings 20th International Conference on Foundations of Software Science and Computation Structures, FOSSACS 2017. Lecture Notes in Computer Science*, vol. 10203, pp. 88–105 (2017). https://doi.org/10.1007/978-3-662-54458-7_6
21. Miné, A.: Tutorial on static inference of numeric invariants by abstract interpretation. *Found. Trends Program. Lang.* **4**(3–4), 120–372 (dec 2017). <https://doi.org/10.1561/25000000034>, <https://doi.org/10.1561/25000000034>
22. Möller, B., O’Hearn, P.W., Hoare, T.: On algebra of program correctness and incorrectness. In: *Proceedings of the 19th International Conference on Relational and Algebraic Methods in Computer Science, RAMiCS 2021. Lecture Notes in Computer Science*, vol. 13027, pp. 325–343. Springer (2021). https://doi.org/10.1007/978-3-030-88701-8_20
23. Möller, B., Struth, G.: Algebras of modal operators and partial correctness. *Theoretical Computer Science* **351**(2), 221–239 (Feb 2006). <https://doi.org/10.1016/j.tcs.2005.09.069>
24. O’Hearn, P.W.: Incorrectness logic. *Proceedings of the ACM on Programming Languages* **4**(POPL), 1–32 (Jan 2020). <https://doi.org/10.1145/3371078>
25. Poskitt, C.M.: Incorrectness logic for graph programs. In: *Proceedings of the 14th International Conference on Graph Transformation, ICGT 2021. Lecture Notes in Computer Science*, vol. 12741, pp. 81–101. Springer (2021). https://doi.org/10.1007/978-3-030-78946-6_5
26. Raad, A., Berdine, J., Dang, H., Dreyer, D., O’Hearn, P.W., Villard, J.: Local reasoning about the presence of bugs: Incorrectness separation logic. In: *Proceedings 32nd International Conference on Computer Aided Verification, CAV 2020. Lecture Notes in Computer Science*, vol. 12225, pp. 225–252. Springer (2020). https://doi.org/10.1007/978-3-030-53291-8_14
27. Raad, A., Berdine, J., Dreyer, D., O’Hearn, P.W.: Concurrent incorrectness separation logic. *Proc. ACM Program. Lang.* **6**(POPL), 1–29 (2022). <https://doi.org/10.1145/3498695>
28. Ranzato, F.: Complete abstractions everywhere. In: *Proceedings of the 14th International Conference on Verification, Model Checking, and Abstract Interpretation, VMCAI 2013. Lecture Notes in Computer Science*, vol. 7737, pp. 15–26. Springer (2013). https://doi.org/10.1007/978-3-642-35873-9_3
29. Smolka, S., Eliopoulos, S.A., Foster, N., Guha, A.: A fast compiler for NetKAT. In: *Proceedings of the 20th ACM SIGPLAN International Conference on Functional Programming, ICFP 2015*. pp. 328–341. ACM (2015). <https://doi.org/10.1145/2784731.2784761>

30. Smolka, S., Kumar, P., Foster, N., Kozen, D., Silva, A.: Cantor meets Scott: Semantic foundations for probabilistic networks. In: Proceedings of the 44th ACM SIGPLAN Symposium on Principles of Programming Languages. p. 557–571. POPL 2017, ACM (2017). <https://doi.org/10.1145/3009837.3009843>
31. Yan, P., Jiang, H., Yu, N.: On incorrectness logic for quantum programs. Proc. ACM Program. Lang. **6**(OOPSLA1) (apr 2022). <https://doi.org/10.1145/3527316>, <https://doi.org/10.1145/3527316>
32. Zhang, C., de Amorim, A.A., Gaboardi, M.: On Incorrectness Logic and Kleene Algebra with Top and Tests. Proceedings of the ACM on Programming Languages **6**(POPL), 1–30 (Jan 2022). <https://doi.org/10.1145/3498690>