

# Center of mass motion in bag model

Chia-Wei Liu\* and Chao-Qiang Geng

*School of Fundamental Physics and Mathematical Sciences,  
Hangzhou Institute for Advanced Study, UCAS, Hangzhou 310024, China  
University of Chinese Academy of Sciences, 100190 Beijing, China*

## Abstract

Despite the great success on the mass spectra, the reputation of the bag model has been closely followed by the embarrassment from the center of mass motion. It leads to severe theoretical inconsistencies. For instance, the masses and the decay constants would no longer be independent of the momentum. In this work, we provide a systematical approach to resolve this problem. The meson decay constants as well as the baryon transition form factors can be computed consistently in our framework. Notably, the form factors in the neutron  $\beta$  decays do not depend on any free parameters, and are determined to be  $F_1^V = 1$  and  $F_1^A = 1.31$  or  $F_1^A/F_1^V = 1.31$ , which is close the experimental value of  $F_1^A/F_1^V = 1.27$ . In addition, we find that  $\mathcal{B}(\Lambda_b \rightarrow \Lambda\gamma) = (7.1 \pm 0.3) \times 10^{-6}$ , which is well consistent to the experimental value of  $(7.1 \pm 1.7) \times 10^{-5}$ . These results show that our approach for the center of mass motion in the bag model is suitable for both light and heavy quark systems.

---

\* chiaweiliu@ucas.ac.cn

## I. INTRODUCTION

The Massachusetts Institute of Technology (MIT) bag model describes a hadron as a bag which contains various quarks, antiquarks and perhaps gluons [1]. It was proposed to reconcile two very different ideas, behind the quantum chromodynamics (QCD) with a single parameter of the bag radius ( $R$ ), as follows:

- Due to the asymptotic freedom, quarks move freely inside the bag;
- Quarks are not allowed to penetrate the bag because of the QCD confinement.

Expressing the ideas mathematically, we start with the free Dirac equations, given by

$$(i\cancel{\partial} - M_q)\psi_q(x) = 0, \quad \text{in } r < R, \quad (1)$$

along with the boundary condition,

$$\vec{x} \cdot (\bar{\psi}_q(x)\vec{\gamma}\psi_q(x)) = 0, \quad \text{at } r = R, \quad (2)$$

where  $\psi_q$  stands for the quark wave function,  $M_q$  represents the quark mass with  $q$  being the quark flavor,  $r = |\vec{x}|$ ,  $\vec{\gamma} = (\gamma_1, \gamma_2, \gamma_3)$ , and  $R$  is the bag radius. The description is similar to the infinite square well in quantum physics. In practice, “bag” can be understood as the abbreviation for “infinite spherical well”. The higher order mass corrections can be done by taking  $\psi_q$  as an unperturbed state.

Thanks to its simplicity, the bag model can easily cooperate with various QCD systems. The applications range from atoms to burning stars. In nuclear physics, the bag model offers an intuitive understandable ground for studying the mass spectra [2] and the hypothetical objects, such as glue balls and pentaquarks [3]. See Ref. [4] for a historical review. In astrophysics, under the framework of the bag model, the strange stars, neutron stars and quark cores are viewed as large bags containing countless quarks [5]. In spite of these great progresses, the bag model has little application in the particle decay system, mainly due to the center of mass motion (CMM) [6]. This study is devoted to introduce the homogeneous bag model to provide a consistent framework to deal with the CMM.

The paper is organized as follows. We give a brief review of the MIT bag model in Sec. II, where we concentrate on the problem of the CMM rather than computational details. In Sec. III, we display some of the methods tackling with the CMM given in the literature. In Sec. IV, we present a systematic framework to deal with the CMM, and show the numerical results of the neutron  $\beta$  decay and  $\Lambda_b \rightarrow \Lambda\gamma$ . We conclude the study in Sec. V.

## II. THE MIT BAG MODEL

Since we only consider the low-lying hadrons, the quark wave function of  $\psi_q$  can be safely taken to be isotropic. Consequently, Eq. (1) can be easily solved as

$$\psi_q(x) = \phi_q(\vec{x})e^{-iE_q t} = N \left( \begin{array}{c} \omega_{q+} j_0(p_q r) \chi \\ i\omega_{q-} j_1(p_q r) \hat{r} \cdot \vec{\sigma} \chi \end{array} \right) e^{-iE_q t}, \quad (3)$$

where  $\phi_q(\vec{x})$  corresponds to the spatial part of the quark wave function with  $E_q^k = \sqrt{p_q^2 + M_q^2}$ ,  $\omega_{q\pm} = \sqrt{E_q^k \pm M_q}$ ,  $p_q$  and  $E_q$  are the magnitudes of the quark 3-momentum and energy, respectively,  $N$  is the normalization constant,  $\chi$  corresponds to the two-component spinor, and  $j_{0,1}$  are the first and the second spherical Bessel functions, which are basically the cosine and sine functions in the spherical coordinate. For the absence of energy corrections, we have that  $E_q = E_q^k$ . In analogy to the case of the infinite square well, imposing the boundary condition would quantize  $p_q$ . At the massless and heavy quark limits, we obtain

$$\lim_{M_q R \rightarrow 0} p_q R = 2.043, 5.396 \dots, \quad \lim_{M_q R \rightarrow \infty} p_q R = \pi, 2\pi \dots, \quad (4)$$

respectively. These results are handy for quick estimations as  $M_{u,d}$  can be taken as massless, whereas  $M_{c,b}$  as infinite in practice. It states that the magnitude of the 3-momentum grows along with the quark mass, approaching  $\pi/R$ .

At this stage, one can readily give the estimations of the hadron masses by adding up the quark energies. In this work, we take the proton as an example, and the formalism can be easily generalized to other hadrons. The proton wave function is given by superposing the quark states

$$\Psi(x_1, x_2, x_3) = \langle 0 | \hat{u}(x_1) \hat{u}(x_2) \hat{d}(x_3) | p_B(0) \rangle = \psi_u(x_1) \psi_u(x_2) \psi_d(x_3), \quad (5)$$

where  $\hat{u}$  and  $\hat{d}$  are the quark field operators, without writing down the spin-flavor and color indices explicitly, and  $|p_B(0)\rangle$  represents the proton state from the bag model, centered in the coordinate. The experimental proton charge radius of 0.840 fm [7] corresponds to  $R = 5.85 \text{ GeV}^{-1}$  in the bag model, leading to the proton and  $\Delta$  masses as

$$M_{p,\Delta} = \frac{2.043}{5.85} \times 3 = 1.047 \text{ GeV}, \quad (6)$$

which are close to the experimental value,

$$\frac{1}{2} (M_p + M_\Delta)_{\text{EXP}} = 1.085 \text{ GeV}. \quad (7)$$

However, the neutron charge radius is essentially zero in the bag model, which is inconsistent with the experiments.

There are several mass corrections, which can be summarized as follows [3]:

- The energy of the bag is proportional to its volume, given by  $E_V = 4\pi BR^3/3$  with  $B$  the bag energy density.
- The zero-point energy is negative, arising from the bag boundary, given by  $E_0 = -Z_0/R$ . It can be understood as the Casimir effect of QCD, proportional to the inverse of the bag radius.
- The quark-gluon interaction introduces the strong coupling constant of  $\alpha_s$ .

Here,  $B$ ,  $Z_0$  and  $\alpha_s$  are treated as free parameters in the model. Surprisingly, the bag radius itself is *not* a free parameter. It is chosen to minimize the hadron masses, resulting in different hadron bag radii. Typically, the baryon and meson radii are found to be around 5 and 3  $\text{GeV}^{-1}$ , respectively.

Although the mass corrections from the bag and the zero-point energies are intuitively satisfactory, they are problematic under the spacetime symmetry. For instance, particles at rest are invariant under the space translations. How can their masses related to a finite volume? How do the zero-point energies transform under the Lorentz boost? These problems partly come from that the description of the bag model is a semi-classical one. Quarks are quantum objects as they satisfy the Dirac equation, whereas the bag itself, having a definite position as well as a concrete boundary, is a classical object. These are the essential problems that have haunted the bag model ever since it was proposed, closely related to the CMM. As a pioneer of quark models, it provides an excellent framework to understand the hadrons in relativistic systems. However, it can hardly be applied to the particle decay processes.

One of the minor issues after considering the mass corrections is that we lost the ability to track the  $t$ -dependencies of the individual quarks. The best we can get is the  $t$ -dependency of the proton wave function, read as

$$\Psi(t, \vec{x}_1, \vec{x}_2, \vec{x}_3) = e^{-iM_p t} \phi_u(\vec{x}_1) \phi_u(\vec{x}_2) \phi_d(\vec{x}_3). \quad (8)$$

Here, we can no longer take  $t_{1,2,3}$  to be different as done in Eq. (5) for  $2E_u^k + E_d^k \neq M_p$  after including the bag and zero-point energies. Due to this reason, we can not apply a Lorentz boost on Eq. (8), which would mix up the spacetime coordinates. Moreover, except for  $u$

and  $d$ , the quark masses are taken to be free parameters in the model, fitted to be [3]

$$M_s = 0.279 \text{ GeV}, \quad M_c = 1.641 \text{ GeV}, \quad M_b = 5.093 \text{ GeV}. \quad (9)$$

It can be viewed as the effective masses when the quarks swim inside the bags, but it is contradicted to the idea of the asymptotic freedom.

### III. THE WAVE PACKET AND ELLIPSOIDAL BAG APPROACHES

We are interested in the applications of the bag model in the decay system, in which momentum eigenstates are required. In the literature, there are two methods concerning decays with the bag model. These methods have their own advantages but fail to achieve a consistent framework with the Poincaré symmetry. Nonetheless, they shed light on the CMM problem and we discuss them briefly.

#### A bag as a wave packet

The most naïve solution to explain the CMM is the wave packet approach, which treats the bag wave function as a localized wave packet. We *extract* the momentum states based on the Fourier analysis, read as

$$|p(\vec{p})\rangle = N(\vec{p}) \int d^3x e^{-i\vec{p}\cdot\vec{x}} |p_B(\vec{x})\rangle, \quad (10)$$

where the left hand side is a proton state with  $\vec{p}$  the 3-momentum, leading to the quark state decomposition as

$$\Psi(t, \vec{x}_1, \vec{x}_2, \vec{x}_3) = N(\vec{p}) \int d^3x e^{-i\vec{p}\cdot\vec{x} - iM_p t} \phi_u(\vec{x}_1 - \vec{x}) \phi_u(\vec{x}_2 - \vec{x}) \phi_d(\vec{x}_3 - \vec{x}). \quad (11)$$

One of the advantages of this method is that the  $t$ -dependencies of the quark states are not required, so it can easily cooperate with the bag and zero-point energies<sup>1</sup>. However, the  $t$ -dependencies of different  $|p\rangle$  are the same since  $|p_B(\vec{x})\rangle$  have the same energy. It would result in a severe inconsistency as the mass depends on  $\vec{p}$ , which is a fatal mistake in the decay system. In addition, it has also been shown that the meson decay constants also depend on  $\vec{p}$  [8].

---

<sup>1</sup> See the discussion following Eq (8).

## The ellipsoidal bag approach

The contradiction occurred in the wave packet approach is attributed to that we can not have a wave packet with a definite energy. To solve the problem, the Lorentz boost for the bag state is needed. Boosting Eq. (5), the wave function is read as

$$\Psi^v(x_1, x_2, x_3) = \psi_u^v(x_1)\psi_u^v(x_2)\psi_d^v(x_3) \quad (12)$$

with  $v$  the velocity and

$$\psi_q^v(x) = S_v \psi_q(\Lambda^{-1}x), \quad (13)$$

where  $S_v$  and  $\Lambda$  are the Lorentz boost matrices toward  $\hat{z}$  direction for Dirac spinors and spacetime coordinates, respectively. In contrast to Eq. (5), the spherical bag deforms to an ellipsoid due to the Lorentz contraction, and the bag itself is moving.

Clearly, the  $t$ -dependencies ( $E_q$ ) of the quark states are needed by Eq. (13). There are two schemes for this ambiguity. In the effective quark mass scheme, we stick to the quark masses given by the MIT bag model in Eq. (9), and we take the quark energies as  $E_q^k$ . The computation in the decay system does not require any new parameters, but it would yield wrong hadron masses as it does not include the bag and zero-point energies.

On the other hand, in the pole mass scheme we take the quark masses as the pole ones as the quarks move freely in the bags, given as [9]

$$M_s = 0.1 \text{ GeV}, \quad M_c = 1.655 \text{ GeV}, \quad M_b = 4.78 \text{ GeV}. \quad (14)$$

Note that  $M_s$  is taken as the current quark mass since the pole mass for the strange quark is ill-defined. On the other hand,  $E_q$  are fitted roughly with the mass spectra, found to be

$$E_{u,d} = 0.32 \text{ GeV}, \quad E_s = 0.44 \text{ GeV}, \quad E_c = 1.56 \text{ GeV}, \quad E_b = 4.97 \text{ GeV}, \quad (15)$$

from the ground state baryons, vector mesons and heavy mesons. The final results depend little on which scheme we use as we will see in Eq. (29).

If we take Eq. (12) as a momentum eigenstate, there are couple requirements to be added by hand before the calculations :

- To have the same bag volume, the initial and final hadrons must be opposite in velocities.
- As the bag is moving, a specific timing for the computation has to be chosen.

The studies of the ellipsoidal bag approach are firstly carried out in the decays with the  $b \rightarrow c$  transition [10], where the overlapping between the heavy quarks is treated by the heavy quark symmetry. On the other hand, the overlapping of the light quarks would lead to the  $v$ -dependency of the Isgur-Wise function, read as

$$\langle l_v | l_{-v} \rangle = \int d^3\vec{x} \psi_l^{v\dagger}(0, \vec{x}) \psi_l^{-v}(0, \vec{x}) = D_l^v(0), \quad (16)$$

with

$$D_l^v(\vec{x}_\Delta) = \sqrt{1-v^2} \int d^3\vec{x} \phi_l^\dagger \left( \vec{x} + \frac{1}{2}\vec{x}_\Delta \right) \phi_l \left( \vec{x} - \frac{1}{2}\vec{x}_\Delta \right) e^{-2ivE_l z}, \quad (17)$$

where  $l$  is the spectator quark in the heavy hadron transition, and  $D_l^v(\vec{x}_\Delta)$  is a function of both velocity and position, which is defined for the latter convenience. In Ref. [10], it was found that  $\mathcal{B}(B^0 \rightarrow D^- l^+ \nu_l) = 2.12\%$ , which is close to  $(2.31 \pm 0.10)\%$  given by the experiments [9]. Remarkably, Eq. (16) can be understood intuitively, in which  $\sqrt{1-v^2}$  comes from the Lorentz contraction of the bag volume, while the exponential in the integral causes the damping, which is a punishment for not being at the same speed.

However, it is unarguable that both the Lorentz and translational symmetries are broken in the ellipsoidal bag approach, since a specific initial frame and timing are demanded in the computation. Furthermore, Eq (16) tell us that the inner products between  $|l_{\pm v}\rangle$  are nonzero, violating the energy momentum conservation.

#### IV. THE HOMOGENEOUS BAG MODEL

The homogeneous bag model, first proposed in Ref. [11], has been widely been applied in various decay systems [12]. It is meant to reconcile the Poincaré symmetry and the bag model, without necessarily introducing a new parameter.

In the following discussion, we will construct hadron states by combining the features of the wave packet and the ellipsoidal bag approaches. We will show that the inconsistencies with the Poincaré symmetry are resolved in our framework. We also take the neutron  $\beta$  decay for an instance of the computation, as it is independent of free parameters.

##### Hadron wave functions

We start with a hadron state at rest. As a zero momentum state has to distribute homogeneously over the space, we linearly superpose infinite bags at different locations. A

proton state at rest is *constructed* as

$$|p(\vec{p} = 0)\rangle = \mathcal{N} \int d^3\vec{x} |p_B(\vec{x})\rangle, \quad (18)$$

where  $\mathcal{N}$  is the normalization constant. Notice that the formula is identical to Eq. (10) when  $\vec{p} = 0$  but the idea is totally different. Eq. (10) is meant to extract the specific momentum component from the wave packet, whereas we are trying to build up a momentum eigenstate from infinite static bags here.

The hadron wave function can be decomposed to the quark states as

$$\Psi(x_1, x_2, x_3) = \mathcal{N} \int d^3\vec{x} \prod_{i=1,2,3} \phi_{q_i}(\vec{x}_i - \vec{x}) e^{-iE_{q_i} t_i}, \quad (19)$$

where  $q_i$  are the quark flavors of the hadron. After the Lorentz boost in  $\hat{z}$  direction, we have

$$\Psi^v(x_1, x_2, x_3) = \mathcal{N} \int d^3\vec{x} \prod_{i=1,2,3} S_v \phi_{q_i}(\vec{x}_i^v - \vec{x}) e^{-i\gamma E_{q_i} (t_i - v z_i)}, \quad (20)$$

with

$$\vec{x}_i^v = (x_i, y_i, \gamma z_i - \gamma v t_i), \quad \gamma = \sqrt{\frac{1}{1 - v^2}}. \quad (21)$$

To obtain  $\mathcal{N}$ , we calculate the overlapping

$$\begin{aligned} \langle \vec{p}' | \vec{p} \rangle &= \int \Psi_v^\dagger(\vec{x}_1, \vec{x}_2, \vec{x}_3, t) \Psi_v(\vec{x}_1, \vec{x}_2, \vec{x}_3, t) \prod_{i=1,2,3} d^3\vec{x}_i \\ &= \mathcal{N}^2 \int d^3\vec{x} d^3\vec{x}' \prod_{i=1,2,3} d^3\vec{x}_i \phi_{q_i}^\dagger(\vec{x}_i^v - \vec{x}') S_v^2 \phi_{q_i}(\vec{x}_i^v - \vec{x}) e^{iE_{q_i}(\gamma v - \gamma' v') z_i}. \end{aligned} \quad (22)$$

Here, we have taken  $S_{v'}^\dagger = S_v$  by anticipating that  $v = v'$  or else the integral vanishes. To simplify the integral, we adopt the following variables:

$$\begin{aligned} \vec{x}_i^r &= \vec{x}_i^v - \frac{1}{2}(\vec{x} + \vec{x}'), \\ \vec{x}_\Delta &= \vec{x} - \vec{x}', \quad \vec{x}_A = \frac{1}{2}(\vec{x} + \vec{x}'). \end{aligned} \quad (23)$$

Now, the integral is read as

$$\begin{aligned} &\frac{\mathcal{N}^2}{\gamma^3} \int d^3\vec{x}_\Delta d^3\vec{x}_A \prod_{i=1,2,3} d^3\vec{x}_i \phi_{q_i}^\dagger\left(\vec{x}_i^r + \frac{1}{2}\vec{x}_\Delta\right) S_v^2 \phi_{q_i}\left(\vec{x}_i^r - \frac{1}{2}\vec{x}_\Delta\right) e^{iE_{q_i}(v-v')z_i^r} e^{iE_{q_i}(v-v')z_A} \\ &= \mathcal{N}^2 \gamma (2\pi)^3 \delta(\vec{p} - \vec{p}') \int d^3\vec{x}_\Delta \prod_{i=1,2,3} D_{q_i}^0(\vec{x}_\Delta) \end{aligned} \quad (24)$$

where  $1/\gamma^3$  comes from the Jacobian in Eq. (23),  $D_q^0(\vec{x}_\Delta)$  is define in Eq. (17) with  $v = 0$ , and

$$M_H = \sum_{i=1,2,3} E_{q_i}, \quad (25)$$

with  $M_H$  the hadron mass. The  $\delta$  function arises from that the overlapping of  $D_q^0(\vec{x}_\Delta)$  for the bags with a distance  $\vec{x}_\Delta$  occurs infinite times in the integral, which is essentially a result of the translational symmetry.

By taking the normalization for a momentum state as

$$\langle \vec{p} | \vec{p}' \rangle = 2\gamma M_H (2\pi)^3 \delta^3(\vec{p} - \vec{p}'), \quad (26)$$

we find

$$\frac{1}{\mathcal{N}^2} = \frac{1}{2M_H} \int d^3\vec{x}_\Delta \prod_{i=1,2,3} D_{q_i}^0(\vec{x}_\Delta). \quad (27)$$

It is important that  $\mathcal{N}$  must be independent of the velocity because the Lorentz boosts are unitary for physical states. Here, our result shows that it is indeed the case in contrast to Eq. (10).

### Transition matrix elements

After the hadron wave functions are constructed, the calculations of the transition matrix elements are straightforward. Here, we choose the neutron-proton transition as an example with  $d \rightarrow u$  at quark level. By sandwiching the quark transition operators with the hadron states, we arrive

$$\int \langle p(\vec{v}) | u^\dagger \Gamma d(x) e^{iqx} | n(-\vec{v}) \rangle d^4x = \mathcal{Z} \int d^3\vec{x}_\Delta \Gamma_{ud}(\vec{x}_\Delta) \prod_{q_j=u,d} D_{q_j}^v(\vec{x}_\Delta), \quad (28)$$

along with

$$\begin{aligned} \mathcal{Z} &\equiv (2\pi)^4 \delta^4(p_n - p_p - q) \mathcal{N}_n \mathcal{N}_p, \\ \Gamma_{ud}(\vec{x}_\Delta) &= \int d^3\vec{x} \phi_u^\dagger \left( \vec{x} + \frac{1}{2}\vec{x}_\Delta \right) S_{\vec{v}} \Gamma S_{-\vec{v}} \phi_d \left( \vec{x} - \frac{1}{2}\vec{x}_\Delta \right) e^{i(M_p + M_n - E_u - E_d)\vec{v}\cdot\vec{x}}, \end{aligned} \quad (29)$$

where  $\mathcal{N}_{n,p}$  are the normalization constants and  $\Gamma$  is an arbitrary Dirac matrix.

Each term in Eq (28) has a concrete physical meaning, which can be summarized as follows:

- $\mathcal{Z}$  represents the overall constant along with the energy momentum conservation;
- $D_{q_j}^v(\vec{x}_\Delta)$  are the overlapping coefficients of the spectator quarks ( $u$  and  $d$ ) in the initial and final states as found in the ellipsoidal bag approach. Note that the centers of the quark wave functions are separated at a distance of  $\vec{x}_\Delta$ ;
- $\Gamma_{ud}$  is the overlapping coefficient of  $d \rightarrow u$  at quark level. Again, the centers of the bags are separated at a distance of  $\vec{x}_\Delta$ .

Here, we have found that the overlapping integrals of the spectator quarks with different velocities ( $D_q^v$ ) do not vanish, which is a feature inherited from the ellipsoidal bag approach. However, we have the energy momentum conservation when we consider the whole wave functions. It can be viewed as the spectator quarks are kicked by the bag, which are in turn kicked by the quark transition operators.

The main ambiguity of the homogeneous bag model comes from  $E_q$  in the exponential as discussed in Eq. (15). However, the deviations are insensible at low velocities, which make the calculation for the neutron-proton transition unaffected. In addition, from the formula,  $D_q^v$  and  $\Gamma_{ud}$  have opposite dependencies on  $E_q$ . Explicitly,  $D_q^v$  goes down, whereas  $\Gamma_{ud}$  goes up, along with the increase of  $E_q$ . It turns out that the physical quantities depend little on  $E_q$  as long as the values of  $E_q$  are reasonable.

For the neutron  $\beta$  decay, the dimensionless form factors  $F_1^{V,A}$  are defined by [13]

$$\begin{aligned} \langle p | \bar{u} \gamma^\mu d | n \rangle &= \bar{u}_p \left( F_1^V(q^2) \gamma^\mu - F_2^V(q^2) i \sigma_{\mu\nu} \frac{q^\nu}{M_n} + F_3^V(q^2) \frac{q^\mu}{M_n} \right) u_n, \\ \langle p | \bar{u} \gamma^\mu \gamma^5 d | n \rangle &= \bar{u}_p \left( F_1^A(q^2) \gamma^\mu - F_2^A(q^2) i \sigma_{\mu\nu} \frac{q^\nu}{M_n} + F_3^A(q^2) \frac{q^\mu}{M_n} \right) \gamma^5 u_n, \end{aligned} \quad (30)$$

where  $u_{p(n)}$  represents the proton (neutron) Dirac spinor, and  $q$  corresponds to the 4-momentum difference of the neutron and proton, which can be safely taken as zero in practice.  $F_1^V$  and  $F_1^A$  can be extracted straightforwardly after computing the transition matrix elements with  $\Gamma = 1$  and  $\Gamma = \gamma^0 \gamma^1 \gamma_5$ , respectively. In the model,  $u$  and  $d$  are taken as massless, and thus there is only one free parameter  $R$ , having the length dimension. The twist is that  $F_1^{V,A}$  do *not* rely on the bag radius since the length dimension can not be canceled. At the  $v \rightarrow 0$  limit, we find that

$$F_1^V = 1, \quad F_1^A = 1.31, \quad (31)$$

which are close to the experimental value of  $F_1^A/F_1^V = 1.27$ . Compared to the result of

$F_1^A/F_1^V = 1.09$ , given previously in the bag model [3, 6], the ratio improves significantly after considering the correction from the CMM.

For the heavy quark transitions, we use the decay of  $\Lambda_b \rightarrow \Lambda\gamma$  for an example, which is governed by the tensor operator of  $b \rightarrow s$ . The bag radii of  $\Lambda_b$  and  $\Lambda$  are found to be around 5 and 4.6 GeV<sup>-1</sup>, respectively [3]. In the calculation, we take both of them as 4.8 GeV<sup>-1</sup> to simplify the numerical calculations. The corresponding tensor form factor and the branching ratio are

$$f_2^{TV}(q^2 = 0) = 0.120 \pm 0.003, \quad (32)$$

and

$$\mathcal{B}(\Lambda_b \rightarrow \Lambda\gamma) = (7.1 \pm 0.3) \times 10^{-6}, \quad (33)$$

respectively, where the definition of  $f_2^{TV}$  and the formalism of the branching ratio can be referred to Ref. [14]. The uncertainties come from the deviation between the effective quark mass and the pole mass schemes. Our result of the branching ratio is well consistent with  $(7.1 \pm 1.7) \times 10^{-6}$  given by the experiment [9].

## V. SUMMARY

We have reviewed the attempts tackling with the CMM of the bag model in the literature. We have discussed the advantages of the wave packet and the ellipsoidal bag approaches as well as their inconsistencies with the Poincaré symmetry. By combing their merits, we have proposed the framework of the homogeneous bags, which is consistent with the Poincaré symmetry. Notably, we have shown that in our framework, the dominated form factors of the neutron  $\beta$  decay do not depend on any free parameters, given as  $F_1^A/F_1^V = 1.31$ , which is close to the experimental value of 1.27. For the heavy quark transition, we have taken the decay of  $\Lambda_b \rightarrow \Lambda\gamma$  as an example, and obtained that  $\mathcal{B}(\Lambda_b \rightarrow \Lambda\gamma) = (7.1 \pm 0.3) \times 10^{-6}$ , which is well consistent with the experimental measurement of  $(7.1 \pm 1.7) \times 10^{-6}$ . As a conclusion, we have found that the homogeneous bag model is useful in both light and heavy quark systems. It is clear that the homogeneous bag model can provide a consistent framework for the computations concerning the hadron transitions, including the form factors as well

as the decay constants.

- 
- [1] A. Chodos, R. L. Jaffe, K. Johnson, C. B. Thorn and V. F. Weisskopf, Phys. Rev. D **9**, 3471 (1974)
- [2] A. W. Thomas, Czech. J. Phys. B **32**, 239 (1982); A. Hosaka and H. Toki, Phys. Rept. **277**, 65 (1996).
- [3] A. Chodos, R. L. Jaffe, K. Johnson and C. B. Thorn, Phys. Rev. D **10**, 2599 (1974); T. A. DeGrand, R. L. Jaffe, K. Johnson and J. E. Kiskis, Phys. Rev. D **12**, 2060 (1975); S. Fleck and J. M. Richard, Prog. Theor. Phys. **82**, 760 (1989); D. H. He, K. Qian, Y. B. Ding, X. Q. Li and P. N. Shen, Phys. Rev. D **70**, 094004 (2004); M. Karliner and J. L. Rosner, Phys. Rev. D **90**, 094007 (2014); W. X. Zhang, H. Xu and D. Jia, Phys. Rev. D **104**, 114011 (2021).
- [4] F. Fernández and J. Segovia, Symmetry **13**, 252 (2021).
- [5] G. Baym and S. A. Chin, Phys. Lett. B **62**, 241 (1976); H. Li, X. L. Luo and H. S. Zong, Phys. Rev. D **82**, 065017 (2010); R. Mallick, Phys. Rev. C **87**, 025804 (2013) A. Aziz, S. Ray, F. Rahaman, M. Khlopov and B. K. Guha, Int. J. Mod. Phys. D **28**, 1941006 (2019); S. Joshi, S. Sau and S. Sanyal, JHEAp **30**, 16 (2021).
- [6] K. Ushio and H. Konashi, Phys. Lett. B **135**, 468 (1984); S. D. Bass and A. W. Thomas, Phys. Lett. B **684**, 216 (2010).
- [7] Y. H. Lin, H. W. Hammer and U. G. Meißner, Phys. Rev. Lett. **128**, 052002 (2022).
- [8] I. Duck, Phys. Lett. B **64**, 163 (1976); J. F. Donoghue and K. Johnson, Phys. Rev. D **21**, 1975 (1980); C. W. Wong, Phys. Rev. D **24**, 1416 (1981).
- [9] P. A. Zyla *et al.* [Particle Data Group], PTEP **2020**, 083C01 (2020); A. Vafaei, K. Javidan and S. Atashbar Tehrani, arXiv:1806.07995 [hep-ph].
- [10] M. Sadzikowski and K. Zalewski, Z. Phys. C **59**, 677 (1993).
- [11] C. Q. Geng, C. W. Liu and T. H. Tsai, Phys. Rev. D **102**, 034033 (2020).
- [12] C. Q. Geng and C. W. Liu, JHEP **11**, 104 (2021); C. W. Liu and C. Q. Geng, JHEP **01**, 128 (2022); X. N. Jin, C. W. Liu and C. Q. Geng, Phys. Rev. D **105**, 053005 (2022); C. W. Liu and B. D. Wan, arXiv:2204.08207 [hep-ph].
- [13] S. Groote, J. G. Körner and B. Melić, Eur. Phys. J. C **79**, 948 (2019).
- [14] C. Q. Geng, C. W. Liu, Z. Y. Wei and J. Zhang, Phys. Rev. D **05**, 073007 (2022).