

BLOWUP OF CYLINDRICALLY SYMMETRIC SOLUTIONS FOR BIHARMONIC NLS

TIANXIANG GOU

*School of Mathematics and Statistics, Xi'an Jiaotong University,
710049, Xi'an, Shaanxi, China*

ABSTRACT. In this paper, we consider blowup of solutions to the Cauchy problem for the following biharmonic NLS,

$$i \partial_t u = \Delta^2 u - \mu \Delta u - |u|^{2\sigma} u \quad \text{in } \mathbb{R} \times \mathbb{R}^d,$$

where $d \geq 1$, $\mu \in \mathbb{R}$ and $0 < \sigma < \infty$ if $1 \leq d \leq 4$ and $0 < \sigma < 4/(d-4)$ if $d \geq 5$. In the mass critical and supercritical cases, we establish the existence of blowup solutions to the problem for cylindrically symmetric data. Our result extends the one obtained in [7], where blowup of solutions to the problem for radially symmetric data was considered.

1. INTRODUCTION

In this paper, we are concerned with blowup of cylindrically symmetric solutions to the Cauchy problem for the following biharmonic NLS,

$$i \partial_t u = \Delta^2 u - \mu \Delta u - |u|^{2\sigma} u \quad \text{in } \mathbb{R} \times \mathbb{R}^d, \quad (1.1)$$

where $d \geq 1$, $\mu \in \mathbb{R}$ and $0 < \sigma < \infty$ if $1 \leq d \leq 4$ and $0 < \sigma < 4/(d-4)$ if $d \geq 5$. The first study of biharmonic NLS traces back to Karpman [12] and Karpman-Shagalov [13], where the authors investigated the regularization and stabilization effect of the fourth-order dispersion. Later, Fibich et al. [10] carried out a rigorous survey to biharmonic NLS from mathematical point of views and proved global existence in time of solutions to the Cauchy problem for (1.1). During recent years, there are a large number of literature devoted to the study of well-posedness and scattering of solutions to the Cauchy problem for (1.1), see for example [8, 11, 15, 16, 17, 18, 19] and references therein. In [7], Boulenger and Lenzmann rigorously and completely discussed the existence of blowup solutions to the Cauchy problem for (1.1) with radially symmetric initial data, which in turn confirms a series of numerical studies conducted in [1, 2, 3]. We also refer to the readers to the papers due to Bonheure et al. [5, 6] with respect to orbital instability of radially symmetric standing waves to (1.1). Inspired by the aforementioned works, the aim of the present paper is to investigate blowup of solutions to the Cauchy problem for (1.1) with cylindrically symmetric initial data, i.e. initial data belong to Σ_d defined by

$$\Sigma_d := \{u \in H^2(\mathbb{R}^d) : u(y, x_d) = u(|y|, x_d), x_d u \in L^2(\mathbb{R}^d)\},$$

where $x = (y, x_d) \in \mathbb{R}^d$ and $x_d = (x_1, \dots, x_{d-1}) \in \mathbb{R}^{d-1}$.

For further clarifications, we shall fix some notations. Let us define

$$s_c := \frac{d}{2} - \frac{2}{\sigma}.$$

We refer to the cases $s_c < 0$, $s_c = 0$ and $s_c > 0$ as mass subcritical, critical and supercritical, respectively. The end case $s_c = 2$ is energy critical. Note that the cases $s_c = 0$ and $s_c = 2$ correspond to the exponents $\sigma = 4/d$

E-mail address: tianxiang.gou@xjtu.edu.cn.

2010 *Mathematics Subject Classification.* Primary: 35Q55; Secondary: 35B35, 35B44.

Key words and phrases. Blowup; Biharmonic NLS; Cylindrical symmetry solutions.

The author was supported by the National Natural Science Foundation of China (No. 12101483) and the Postdoctoral Science Foundation of China.

and $\sigma = 4/(d-4)$, respectively. For $1 \leq p < \infty$, we denote by $L^q(\mathbb{R}^d)$ the usual Lebesgue space with the norm

$$\|u\|_p := \left(\int_{\mathbb{R}^d} |u|^p dx \right)^{\frac{1}{p}}.$$

The Sobolev space $H^2(\mathbb{R}^d)$ is equipped with the standard norm

$$\|u\| := \|\Delta u\|_2 + \|\nabla u\|_2 + \|u\|_2.$$

In addition, we denote by $Q \in H^2(\mathbb{R}^d)$ a ground state to the following nonlinear elliptic equation,

$$\Delta^2 Q + Q - |Q|^{2\sigma} Q = 0 \quad \text{in } \mathbb{R}^d.$$

The main results of the present paper read as follows, which gives blowup criteria for solutions to the Cauchy problem for (1.1) with cylindrically symmetric data.

Theorem 1.1. (Blowup for Mass-Supercritical Case) *Let $d \geq 5$, $\mu \in \mathbb{R}$ and $0 < s_c < 2$ with $0 < \sigma \leq 1$. Suppose that $u_0 \in \Sigma_d$ satisfies one of the following condition.*

(i) *If $\mu \neq 0$, we assume that*

$$E[u_0] < \begin{cases} 0, & \text{for } \mu > 0, \\ -\chi\mu^2 M[u_0], & \text{for } \mu < 0, \end{cases}$$

with some constant $\chi = \chi(d, \sigma) > 0$.

(ii) *If $\mu = 0$, we assume that either $E[u_0] < 0$ or if $E[u_0] \geq 0$, we suppose that*

$$E[u_0]^{s_c} M[u_0]^{2-s_c} < E[Q]^{s_c} M[Q]^{2-s_c}$$

and

$$\|\Delta u_0\|_2^2 \|u_0\|_2^{2-s_c} > \|\Delta Q\|_2^2 \|Q\|_2^{2-s_c}.$$

Then the solution $u \in C([0, T), H^2(\mathbb{R}^d))$ to the Cauchy problem for (1.1) with initial datum u_0 blows up in finite time, i.e. $0 < T < +\infty$ and $\lim_{t \rightarrow T^-} \|\Delta u\|_2 = +\infty$.

Remark 1.1. *The extra restriction on σ comes from the use of the well-known radial Sobolev inequality in \mathbb{R}^{d-1} . Note that if $4/d < \sigma \leq 1$, then $d \geq 5$. This is the reason that we need to assume that $d \geq 5$. Such restrictions are reminiscent of analogous ones for the study of blowup of solutions to NLS with cylindrically symmetric data, see [4, 9, 14].*

Theorem 1.2. (Blowup for Mass-Critical Case) *Let $d \geq 4$, $\mu \geq 0$ and $s_c = 0$. Let $u_0 \in \Sigma_d$ be such that $E(u_0) < 0$. Then the solution $u \in C([0, T), H^2(\mathbb{R}^d))$ to the Cauchy problem for (1.1) satisfies the following.*

(i) *If $\mu > 0$, then $u(t)$ blows up in finite time.*

(ii) *If $\mu = 0$, then $u(t)$ either blows up in finite time or $u(t)$ blows up in infinite time.*

To prove Theorems 1.1 and 1.2, the essential argument is to deduce the evolution of a localized virial quantity $M_{\varphi_R}[u(t)]$ defined by (2.2) along time, see Lemma 2.2. To this end, we shall make use of some ideas from [7, 14]. For cylindrically symmetric solutions, the radial Sobolev inequality is only applicable in \mathbb{R}^{d-1} , then we need to utilize different arguments to estimate error terms due to the nonlinearity in the process of discussion of time evolution of the localized virial quantity.

2. PROOFS OF MAIN RESULTS

In this section, we are going to prove Theorems 1.1 and 1.2. To do this, we first need to introduce a localized virial quantity, which is inspired by [7] and [14]. For $d \geq 2$, let $\psi : \mathbb{R}^{d-1} \rightarrow \mathbb{R}$ be a radially symmetric and smooth function such that $|\nabla \psi^j| \in L^\infty(\mathbb{R}^{d-1})$ for $1 \leq j \leq 6$ and

$$\psi(r) := \begin{cases} \frac{r^2}{2} & \text{for } r \leq 1, \\ \text{const.} & \text{for } r \geq 10, \end{cases} \quad \psi''(r) \leq 1 \text{ for } r \geq 0.$$

For $R > 0$ given, we define a radial function $\psi_R : \mathbb{R}^{d-1} \rightarrow \mathbb{R}$ by

$$\psi_R(r) := R^2 \varphi\left(\frac{r}{R}\right).$$

It follows from (3.3) in [7] that

$$1 - \psi_R''(r) \geq 0, \quad 1 - \frac{\psi_R'(r)}{r} \geq 0, \quad d - 1 - \Delta \psi_R(r) \geq 0, \quad r \geq 0. \quad (2.1)$$

Let

$$\varphi_R(x) := \psi_R(r) + \frac{x_d^2}{2}, \quad x = (y, x_d) \in \mathbb{R}^{d-1} \times \mathbb{R}, \quad r = |y|.$$

Define

$$\mathcal{M}_{\varphi_R}[u] := 2\text{Im} \int_{\mathbb{R}^d} \bar{u} (\nabla \varphi_R \cdot \nabla u) \, dx. \quad (2.2)$$

It is simple see that $\mathcal{M}_{\varphi_R}[u]$ is well-defined for any $u \in \Sigma_d$. For later use, we shall give the well-known radial Sobolve inequality in [20]. For every radial function $f \in H^1(\mathbb{R}^{d-1})$ with $d \geq 3$, then

$$|y|^{\frac{d-2}{2}} |f(y)| \leq 2 \|f\|_2^{\frac{1}{2}} \|\nabla f\|_2^{\frac{1}{2}}, \quad y \neq 0. \quad (2.3)$$

We also present the well-known Gagliardo-Nirenberge's inequality in one dimension. For any $f \in H^1(\mathbb{R})$ and $p > 2$, then

$$\|f\|_p \leq C_p \|f'\|_2^\alpha \|f\|_2^{1-\alpha}, \quad \alpha = \frac{p-2}{2p}. \quad (2.4)$$

Let $f : \mathbb{R}^{d-1} \rightarrow \mathbb{C}$ be a radial and smooth function, then

$$\partial_{kl}^2 f = \left(\delta_{kl} - \frac{x_k x_l}{r^2} \right) \frac{\partial_r f}{r} + \frac{x_k x_l}{r^2} \partial_r^2 f. \quad (2.5)$$

Next we present the well-posedness of solutions to the Cauchy problem for (1.1) in $H^2(\mathbb{R}^d)$, which was established by Pausader [16].

Lemma 2.1. [16, Proposition 4.1] *Let $d \geq 1$, $\mu \in \mathbb{R}$ and $s_c < 2$. Then, for any $u_0 \in H^2(\mathbb{R}^d)$, there exist a constant $T > 0$ and a unique solution $u \in C([0, T], H^2(\mathbb{R}^d))$ to the Cauchy problem for (1.1) with initial datum u_0 . The solution has conserved mass and energy in the sense that*

$$M[u(t)] = M[u_0], \quad E[u(t)] = E[u_0], \quad t \in [0, T],$$

where

$$M[u] := \int_{\mathbb{R}^d} |u|^2 \, dx$$

and

$$E[u] = \frac{1}{2} \int_{\mathbb{R}^d} |\Delta u|^2 \, dx + \frac{\mu}{2} \int_{\mathbb{R}^d} |\nabla u|^2 \, dx - \frac{1}{2\sigma + 2} \int_{\mathbb{R}^d} |u|^{2\sigma+2} \, dx.$$

Moreover, blowup alternative holds, i.e. either $T = +\infty$ or $\|u(t)\|_{H^2} = +\infty$ as $t \rightarrow T^-$. The solution map

$$u_0 \in H^2(\mathbb{R}^d) \mapsto u \in C([0, T], H^2(\mathbb{R}^d))$$

is continuous.

In the following, we give the evolution of $\mathcal{M}[u(t)]$ along time, which is the key argument to prove Theorems 1.1 and 1.2.

Lemma 2.2. *Let $d \geq 3$, $R > 0$ and $0 < \sigma \leq 1$. Suppose that $u \in C([0, T]; H^2(\mathbb{R}^d))$ is the solution to the Cauchy problem for (1.1) with initial datum $u_0 \in \Sigma_d$. Then, for any $t \in [0, T]$, there holds that*

$$\begin{aligned} \frac{d}{dt} \mathcal{M}_R[u(t)] &\leq 8 \int_{\mathbb{R}^d} |\Delta u|^2 \, dx + 4\mu \int_{\mathbb{R}^d} |\nabla u|^2 \, dx - \frac{2\sigma d}{\sigma + 1} \int_{\mathbb{R}^d} |u|^{2\sigma+2} \, dx + X_\mu[u] \\ &\quad + \mathcal{O}\left(R^{-4} + R^{-2} \|\nabla u\|_2^2 + R^{-\sigma(d-2)} \|\nabla u\|_2^{2\sigma} + |\mu| R^{-2}\right) \\ &= 4d\sigma E(u_0) - (2d\sigma - 8) \|\Delta u\|_2^2 - \mu(2d\sigma - 4) \|\nabla u\|_2^2 + X_\mu[u] \\ &\quad + \mathcal{O}\left(R^{-4} + R^{-2} \|\nabla u\|_2^2 + R^{-\sigma(d-2)} \|\nabla u\|_2^{2\sigma} + |\mu| R^{-2}\right), \end{aligned}$$

where

$$X_\mu[u] \lesssim \begin{cases} 0, & \text{for } \mu \geq 0, \\ |\mu| \|\nabla u\|_2^2, & \text{for } \mu < 0. \end{cases}$$

Proof. To achieve this, we shall adapt some elements from [7] and [14]. In view of Step 1 of the proof of [7, Lemma 3.1], we first have that

$$\begin{aligned} \frac{d}{dt} \mathcal{M}_R[u(t)] &= 8 \sum_{k,l,m=1}^d \langle u, \partial_{kl}^2 (\partial_{lm}^2 \varphi_R) \partial_{mk}^2 u \rangle + 4 \sum_{k,l=1}^d \langle u, \partial_k (\partial_{kl} \Delta \varphi_R) \partial_l u \rangle \\ &\quad + 2 \sum_{k=1}^d \langle u, \partial_k (\Delta^2 \varphi_R) \partial_l u \rangle + \langle u, (\Delta^3 \varphi_R) u \rangle \\ &\quad - 4\mu \sum_{k,l=1}^d \langle u, \partial_k (\partial_{kl}^2 \varphi_R) \partial_l u \rangle - \mu \langle u, (\Delta^2 \varphi_R) u \rangle - \langle u, [|u|^{2\sigma}, \nabla \varphi_R \cdot \nabla + \nabla \cdot \nabla \varphi_R] u \rangle, \\ &=: \mathcal{A}_R^{(1)}[u] + \mathcal{A}_R^{(2)}[u] + \mathcal{B}_R[u]. \end{aligned}$$

In what follows, we are going to estimate the terms $\mathcal{A}_R^{(1)}[u]$, $\mathcal{A}_R^{(2)}[u]$ and $\mathcal{B}_R[u]$. The estimates of dispersive terms $\mathcal{A}_R^{(1)}[u]$ and $\mathcal{A}_R^{(2)}[u]$ are inspired by the proof of [7, Lemma 3.1]. Let us begin with treating the term $\mathcal{A}_R^{(1)}[u]$. Using integration by parts and the definition of φ_R , we are able to derive that

$$\begin{aligned} \sum_{k,l,m=1}^d \langle u, \partial_{kl}^2 (\partial_{lm}^2 \varphi_R) \partial_{mk}^2 u \rangle &= \sum_{k,l,m=1}^d \int_{\mathbb{R}^d} (\partial_{kl}^2 \bar{u}) (\partial_{lm}^2 \varphi_R) (\partial_{mk}^2 u) \, dx \\ &= \sum_{k,l,m=1}^{d-1} \int_{\mathbb{R}^d} (\partial_{kl}^2 \bar{u}) (\partial_{lm}^2 \psi_R) (\partial_{mk}^2 u) \, dx \\ &\quad + \sum_{l,m=1}^{d-1} \int_{\mathbb{R}^d} (\partial_{dl}^2 \bar{u}) (\partial_{lm}^2 \psi_R) (\partial_{md}^2 u) \, dx \\ &\quad + \sum_{k=1}^{d-1} \int_{\mathbb{R}^d} (\partial_{kd}^2 \bar{u}) (\partial_{dk}^2 u) \, dx + \int_{\mathbb{R}^d} |\partial_{dd}^2 u|^2 \, dx. \end{aligned} \tag{2.6}$$

We now compute each term in the right hand side of (2.6). Utilizing (2.5), we can derive that

$$\begin{aligned} &\sum_{k,l,m=1}^{d-1} \int_{\mathbb{R}^d} (\partial_{kl}^2 \bar{u}) (\partial_{lm}^2 \psi_R) (\partial_{mk}^2 u) \, dx \\ &= \sum_{k,l,m=1}^{d-1} \int_{\mathbb{R}} \int_{\mathbb{R}^{d-1}} (\partial_{kl}^2 \bar{u}) (\partial_{lm}^2 \psi_R) (\partial_{mk}^2 u) \, dy dx_d \\ &= \int_{\mathbb{R}} \int_{\mathbb{R}^{d-1}} \left(\partial_r^2 \psi_R |\partial_r^2 u|^2 + \frac{d-2}{r^2} \frac{\partial_r \psi_R}{r} |\partial_r u|^2 \right) \, dy dx_d \\ &= \int_{\mathbb{R}^d} |\Delta_y u|^2 - (1 - \partial_r^2 \psi_R) |\partial_r^2 u|^2 - \left(1 - \frac{\partial_r \psi_R}{r} \right) \frac{d-2}{r^2} |\partial_r u|^2 \, dx \end{aligned} \tag{2.7}$$

and

$$\begin{aligned}
\sum_{l,m=1}^{d-1} \int_{\mathbb{R}^d} (\partial_{dl}^2 \bar{u}) (\partial_{lm}^2 \psi_R) (\partial_{md}^2 u) dx &= \sum_{k=1}^{d-1} \int_{\mathbb{R}^d} (\partial_r^2 \psi_R) |\partial_{kd}^2 u|^2 dx \\
&= \sum_{k=1}^{d-1} \int_{\mathbb{R}^d} |\partial_{kd}^2 u|^2 - (1 - \partial_r^2 \psi_R) |\partial_{kd}^2 u|^2 dx \\
&= \sum_{k=1}^{d-1} \int_{\mathbb{R}^d} (\partial_{kd}^2 \bar{u}) (\partial_{dk}^2 u) - (1 - \partial_r^2 \psi_R) |\partial_{kd}^2 u|^2 dx.
\end{aligned} \tag{2.8}$$

In addition, applying integration by parts, we have that

$$\int_{\mathbb{R}^d} (\partial_{kd}^2 \bar{u}) (\partial_{dk}^2 u) dx = \int_{\mathbb{R}^d} (\partial_{kk}^2 \bar{u}) (\partial_{dd}^2 u) dx. \tag{2.9}$$

As a consequence, coming back to (2.6) and using (2.1), (2.6), (2.7), (2.8) and (2.9), we now conclude that

$$\sum_{k,l,m=1}^d \langle u, \partial_{kl}^2 (\partial_{lm}^2 \varphi_R) \partial_{mk}^2 u \rangle \leq \int_{\mathbb{R}^d} |\Delta u|^2 dx. \tag{2.10}$$

Furthermore, by the definitions of φ_R and ψ_R , there holds that

$$\begin{aligned}
\sum_{k,l=1}^d |\langle u, \partial_k (\partial_{kl} \Delta \varphi_R) \partial_l u \rangle| &= \sum_{k,l=1}^{d-1} |\langle u, \partial_k (\partial_{kl} \Delta \psi_R) \partial_l u \rangle| \lesssim R^{-2} \|\nabla u\|_2^2, \\
\sum_{k,l=1}^d |\langle u, \partial_k (\Delta^2 \varphi_R) \partial_l u \rangle| &= \sum_{k,l=1}^{d-1} |\langle u, \partial_k (\Delta^2 \psi_R) \partial_l u \rangle| \lesssim R^{-2} \|\nabla u\|_2^2,
\end{aligned}$$

and

$$|\langle u, (\Delta^3 \varphi_R) u \rangle| = |\langle u, (\Delta^3 \psi_R) u \rangle| \lesssim R^{-4} \|u\|_2^2.$$

This along with (2.10) and the conservation of mass implies that

$$\mathcal{A}_R^{(1)}[u] \leq 8 \int_{\mathbb{R}^d} |\Delta u|^2 dx + \mathcal{O}(R^{-4} + R^{-2} \|\nabla u\|_2^2).$$

We next deal with the term $\mathcal{A}_R^{(2)}[u]$. In virtue of integration by parts, the definition of φ_R and (2.5), we can show that

$$\begin{aligned}
\mathcal{A}_R^{(2)}[u(t)] &= 4\mu \sum_{k,l=1}^d \int_{\mathbb{R}^d} (\partial_k \bar{u}) (\partial_{kl}^2 \varphi_R) (\partial_l u) dx - \mu \int_{\mathbb{R}^d} (\Delta^2 \varphi_R) |u|^2 dx \\
&= 4\mu \sum_{k,l=1}^{d-1} \int_{\mathbb{R}} \int_{\mathbb{R}^{d-1}} (\partial_k \bar{u}) (\partial_{kl}^2 \psi_R) (\partial_l u) dy dx_d + 4\mu \int_{\mathbb{R}^d} |\partial_d u|^2 dx - \mu \int_{\mathbb{R}^d} (\Delta^2 \psi_R) |u|^2 dx \\
&= 4\mu \int_{\mathbb{R}} \int_{\mathbb{R}^{d-1}} (\partial_r^2 \psi_R) |\partial_r u|^2 dy dx_d + 4\mu \int_{\mathbb{R}^d} |\partial_d u|^2 dx - \mu \int_{\mathbb{R}^d} (\Delta^2 \psi_R) |u|^2 dx \\
&= 4\mu \int_{\mathbb{R}^d} |\nabla_y u|^2 dx + X_\mu[u] + 4\mu \int_{\mathbb{R}^d} |\partial_d u|^2 dx - \mu \int_{\mathbb{R}^d} (\Delta^2 \psi_R) |u|^2 dx \\
&= 4\mu \int_{\mathbb{R}^d} |\nabla u|^2 dx + X_\mu[u] - \mu \int_{\mathbb{R}^d} (\Delta^2 \psi_R) |u|^2 dx
\end{aligned}$$

where

$$X_\mu[u] := -4\mu \int_{\mathbb{R}^d} (1 - \partial_r^2 \psi_R) |\partial_r u|^2 dx.$$

Due to $\|\Delta^2 \psi_R\|_\infty \lesssim R^{-2}$, then

$$\mathcal{A}_R^{(2)}[u(t)] = 4\mu \int_{\mathbb{R}^d} |\nabla u|^2 dx + X_\mu[u] + \mathcal{O}(|\mu| R^{-2}).$$

We now turn to handle the term $\mathcal{B}_R[u]$. Here we need some special treatments. Applying integration by parts and the definition of φ_R , we first derive that

$$\begin{aligned}\mathcal{B}_R[u] &= 2 \int_{\mathbb{R}^d} |u|^2 \nabla \varphi_R \cdot \nabla (|u|^{2\sigma}) = -\frac{2\sigma}{\sigma+1} \int_{\mathbb{R}^d} (\Delta \varphi_R) |u|^{2\sigma+2} dx \\ &= -\frac{2\sigma}{\sigma+1} \int_{\mathbb{R}^d} (\Delta \psi_R) |u|^{2\sigma+2} dx - \frac{2\sigma}{\sigma+1} \int_{\mathbb{R}^d} |u|^{2\sigma+2} dx \\ &= -\frac{2\sigma d}{\sigma+1} \int_{\mathbb{R}^d} |u|^{2\sigma+2} dx - \frac{2\sigma}{\sigma+1} \int_{\mathbb{R}^d} (\Delta \psi_R - d + 1) |u|^{2\sigma+2} dx.\end{aligned}$$

In virtue of the definition of ψ_R and (2.5), then there holds that $\Delta \psi_R(r) - d + 1 = 0$ for $0 \leq r \leq R$. This further implies that

$$\mathcal{B}_R[u] = -\frac{2\sigma d}{\sigma+1} \int_{\mathbb{R}^d} |u|^{2\sigma+2} dx - \frac{2\sigma}{\sigma+1} \int_{\mathbb{R}} \int_{|y| \geq R} (\Delta \psi_R - d + 1) |u|^{2\sigma+2} dy dx_d. \quad (2.11)$$

In the following, we shall estimate the second term in the right hand side of (2.11). Observe first that

$$\int_{\mathbb{R}} \int_{|y| \geq R} |u|^{2\sigma+2} dy dx_d \leq \int_{\mathbb{R}} \|u\|_{L^\infty(|y| \geq R)}^{2\sigma} \|u\|_{L_y^2}^2 dx_d. \quad (2.12)$$

To proceed the proof, we first consider the case that $\sigma = 1$. In this case, by (2.3), Hölder's inequality and the conservation of mass, then

$$\begin{aligned}\int_{\mathbb{R}} \|u\|_{L^\infty(|y| \geq R)}^2 dx_d &\lesssim R^{-(d-2)} \int_{\mathbb{R}} \|u\|_{L_y^2} \|\nabla_y u\|_{L_y^2} dx_d \\ &\leq R^{-(d-2)} \|u\|_2 \|\nabla_y u\|_2 \lesssim R^{-(d-2)} \|\nabla_y u\|_2.\end{aligned} \quad (2.13)$$

On the other hand, by Hölder's inequality and the conservation of mass, we know that

$$\begin{aligned}\|u\|_{L^\infty(\mathbb{R}, L_y^2(\mathbb{R}^{d-1}))}^2 &= \sup_{x_d \in \mathbb{R}} \int_{\mathbb{R}^{d-1}} |u|^2 dy = \sup_{x_d \in \mathbb{R}} \int_{\mathbb{R}^{d-1}} \int_{-\infty}^{x_d} \partial_d (|u|^2) dy dx_d \\ &= 2 \operatorname{Re} \sup_{x_d \in \mathbb{R}} \int_{\mathbb{R}^{d-1}} \int_{-\infty}^{x_d} \bar{u} (\partial_d u) dy dx_d \leq 2 \|u\|_2 \|\partial_d u\|_2 \lesssim \|\partial_d u\|_2.\end{aligned} \quad (2.14)$$

Consequently, going back to (2.12) and using (2.13) and (2.14), we derive that

$$\int_{\mathbb{R}} \int_{|y| \geq R} |u|^4 dy dx_d \lesssim R^{-(d-2)} \|\nabla_y u\|_2 \|\partial_d u\|_2 \lesssim R^{-(d-2)} \|\nabla u\|_2^2. \quad (2.15)$$

We next consider the case that $0 < \sigma < 1$. In this case, from (2.12) and Hölder's inequality, it follows that

$$\int_{\mathbb{R}} \int_{|y| \geq R} |u|^{2\sigma+2} dx \leq \left(\int_{\mathbb{R}} \|u\|_{L^\infty(|y| \geq R)}^2 dx_d \right)^\sigma \left(\int_{\mathbb{R}} \|u\|_{L_y^2}^{\frac{2}{1-\sigma}} dx_d \right)^{1-\sigma}. \quad (2.16)$$

In view of (2.4) and the conservation of mass, we get that

$$\begin{aligned}\int_{\mathbb{R}} \|u\|_{L_y^2}^{\frac{2}{1-\sigma}} dx_d &\leq \left(\int_{\mathbb{R}} \left| \partial_d (\|u\|_{L_y^2}^2) \right|^2 dx_d \right)^{\frac{\sigma}{2(1-\sigma)}} \left(\int_{\mathbb{R}} \|u\|_{L_y^2}^2 dx \right)^{\frac{2-\sigma}{2(1-\sigma)}} \\ &\lesssim \left(\int_{\mathbb{R}} \left| \partial_d (\|u\|_{L_y^2}^2) \right|^2 dx_d \right)^{\frac{\sigma}{2(1-\sigma)}}.\end{aligned} \quad (2.17)$$

Furthermore, notice that

$$\left| \partial_d (\|u\|_{L_y^2}^2) \right| \|u\|_{L_y^2} = \frac{1}{2} \left| \partial_d (\|u\|_{L_y^2}^2) \right| = \frac{1}{2} \left| \operatorname{Re} \int_{\mathbb{R}^{d-1}} \bar{u} (\partial_d u) dy \right| \leq \frac{1}{2} \|\partial_d u\|_{L_y^2} \|u\|_{L_y^2}.$$

This means that

$$\left| \partial_d (\|u\|_{L_y^2}^2) \right| \leq \frac{1}{2} \|\partial_d u\|_{L_y^2}. \quad (2.18)$$

As a result, via (2.17), we obtain that

$$\int_{\mathbb{R}} \|u\|_{L_y^2}^{\frac{2}{1-\sigma}} dx_d \lesssim \left(\int_{\mathbb{R}} \|\partial_d u\|_{L_y^2}^2 dx_d \right)^{\frac{\sigma}{2(1-\sigma)}} = \|\partial_d u\|_2^{\frac{\sigma}{1-\sigma}}. \quad (2.19)$$

Using (2.13) and (2.19), we then obtain from (2.16) that

$$\int_{|y| \geq R} |u|^{2\sigma+2} dx \lesssim R^{-\sigma(d-2)} \|\nabla_y u\|_2^\sigma \|\partial_d u\|_2^\sigma \lesssim R^{-\sigma(d-2)} \|\nabla u\|_2^{2\sigma}. \quad (2.20)$$

To sum up, it then follows from (2.11), (2.15) and (2.20) that

$$\mathcal{B}_R[u] \lesssim -\frac{2\sigma d}{\sigma+1} \int_{\mathbb{R}^d} |u|^{2\sigma+2} dx + R^{-\sigma(d-2)} \|\nabla u\|_2^{2\sigma}.$$

Accordingly, applying the estimates to $\mathcal{A}_R^{(1)}[u]$, $\mathcal{A}_R^{(2)}[u]$ and $\mathcal{B}_R[u]$ and the conservation of energy, we finally derive that

$$\begin{aligned} \frac{d}{dt} \mathcal{M}_R[u(t)] &\leq 8 \int_{\mathbb{R}^d} |\Delta u|^2 dx + 4\mu \int_{\mathbb{R}^d} |\nabla u|^2 dx - \frac{2\sigma d}{\sigma+1} \int_{\mathbb{R}^d} |u|^{2\sigma+2} dx + X_\mu[u] \\ &\quad + \mathcal{O}\left(R^{-4} + R^{-2} \|\nabla u\|_2^2 + R^{-\sigma(d-2)} \|\nabla u\|_2^{2\sigma} + |\mu|R^{-2}\right) \\ &= 4d\sigma E(u_0) - (2d\sigma - 8) \|\Delta u\|_2^2 - \mu(2d\sigma - 4) \|\nabla u\|_2^2 + X_\mu[u] \\ &\quad + \mathcal{O}\left(R^{-4} + R^{-2} \|\nabla u\|_2^2 + R^{-\sigma(d-2)} \|\nabla u\|_2^{2\sigma} + |\mu|R^{-2}\right), \end{aligned}$$

This completes the proof. \square

Proof of Theorem 1.1. Noting that $0 < \sigma \leq 1$ and applying Lemma 2.2, then the proof can be completed by closely following the one of [7, Theorem 1]. \square

Proof of Theorem 1.2. If $\mu > 0$, using Lemma 2.2 and arguing as the proof of [7, Theorem 3], we can get the desired result. To complete the proof, we only need to consider the case that $\mu = 0$. In this case, we need to conduct a more refined analysis to the evolution of $\mathcal{M}[u(t)]$ along time. From the proof of Lemma 2.2, (2.1) and (2.5), we first have that

$$\begin{aligned} \frac{d}{dt} \mathcal{M}_R[u(t)] &\leq 16E(u_0) - 8 \int_{\mathbb{R}^d} (1 - \partial_r^2 \psi_R) |\partial_r^2 u|^2 dx + \int_{\mathbb{R}^d} (\Delta^3 \psi_R) |u|^2 dx \\ &\quad - \int_{\mathbb{R}^d} A_R |\partial_r u|^2 dx + \int_{\mathbb{R}^d} B_R |u|^{2+\frac{8}{d}} dx, \end{aligned} \quad (2.21)$$

where

$$A_R := 4\partial_r^2 \Delta \psi_R + 2\Delta^2 \psi_R(r), \quad B_R := \frac{8}{d+4} (d-1 - \Delta \psi_R).$$

Thereinafter, we shall estimate the last two terms in the right hand side of (2.21). Using (7.7) in [7] and the conservation of mass, we can derive that

$$\begin{aligned} \left| \int_{\mathbb{R}^d} A_R |\partial_r u|^2 dx \right| &= \left| \int_{\mathbb{R}} \int_{\mathbb{R}^{d-1}} A_R |\partial_r u|^2 dy dx_d \right| \lesssim 8\eta R^4 \int_{\mathbb{R}} \|A_R \partial_r^2 u\|_{L_y^2}^2 dx_d + \eta^{-1} R^{-4} \int_{\mathbb{R}} \|u\|_{L_y^2}^2 dx_d \\ &= 8\eta R^4 \|A_R \partial_r^2 u\|_2^2 + \eta^{-1} R^{-4}, \end{aligned}$$

where $\eta > 0$ is an arbitrary constant. We now treat the other term. To do this, we first consider the case that $d = 4$. In this case, we have that

$$\begin{aligned} \left| \int_{\mathbb{R}^d} B_R |u|^4 dx \right| &= \left| \int_{\mathbb{R}} \int_{|y| \geq R} B_R |u|^4 dy dx_d \right| \leq \int_{\mathbb{R}} \|u\|_{L_y^2}^2 \|B_R^{\frac{1}{2}} u\|_{L^\infty(|y| \geq R)}^2 dx_d \\ &\leq \|u\|_{L^\infty(R, L_y^2(\mathbb{R}^{N-1}))}^2 \int_{\mathbb{R}} \|B_R^{\frac{1}{2}} u\|_{L^\infty(|y| \geq R)}^2 dx_d, \end{aligned} \quad (2.22)$$

because of $B_R = 0$ for $0 \leq r \leq R$. In view of (2.3) with $d = 4$ and the definition of B_R , we are able to infer that

$$\begin{aligned} \int_{\mathbb{R}} \|B_R^{\frac{1}{2}} u\|_{L^\infty(|y| \geq R)}^2 dx_d &\lesssim R^{-2} \int_{\mathbb{R}} \|B_R^{\frac{1}{2}} u\|_{L_y^2} \|\nabla_y (B_R^{\frac{1}{2}} u)\|_{L_y^2} dx_d \\ &\lesssim R^{-2} \int_{\mathbb{R}} \|u\|_{L_y^2} \|\nabla_y (B_R^{\frac{1}{2}} u)\|_{L_y^2} dx_d. \end{aligned} \quad (2.23)$$

It follows from (7.9) with $d = 4$ in [7] that

$$\|\nabla_y(B_R^{\frac{1}{2}}u)\|_{L_y^2}^2 \lesssim \left(\eta^{-\frac{1}{4}} + R^{-2}\right) \|u\|_{L_y^2}^2 + 8\eta^{\frac{1}{4}} \|B_R \partial_r^2 u\|_{L_y^2}^2.$$

By means of (2.23), the conservation of mass and Hölder's inequality, we then get that

$$\begin{aligned} \int_{\mathbb{R}} \|B_R^{\frac{1}{2}}u\|_{L^\infty(|y|\geq R)}^2 dx_d &\lesssim R^{-2} \left(\eta^{-\frac{1}{4}} + R^{-2}\right) + 8R^{-2}\eta^{\frac{1}{4}} \int_{\mathbb{R}^d} \|u\|_{L_y^2} \|B_R \partial_r^2 u\|_{L_y^2} dx_d \\ &\lesssim R^{-2} \left(\eta^{-\frac{1}{4}} + R^{-2}\right) + 8R^{-2}\eta^{\frac{1}{4}} \|B_R \partial_r^2 u\|_2. \end{aligned} \quad (2.24)$$

Going back to (2.22) and using (2.14) and (2.24), we then get that

$$\begin{aligned} \left| \int_{\mathbb{R}^N} B_R |u|^4 dx \right| &\lesssim R^{-2} \left(\eta^{-\frac{1}{4}} + R^{-2}\right) \|\partial_d u\|_2 + 8R^{-2}\eta^{\frac{1}{4}} \|B_R \partial_r^2 u\|_2 \|\partial_d u\|_2 \\ &\leq R^{-2} \left(\eta^{-\frac{1}{4}} + R^{-2}\right) \|\partial_d u\|_2 + R^{-4}\eta^{-\frac{1}{2}} \|\partial_d u\|_2^2 + 8\eta \|B_R \partial_r^2 u\|_2^2. \end{aligned}$$

Taking into account (2.21) and noting that $\|\Delta^2 \varphi_R\|_\infty \lesssim R^{-2}$, we then obtain that

$$\begin{aligned} \frac{d}{dt} \mathcal{M}_R[u(t)] &\leq 16E(u_0) - 8 \int_{\mathbb{R}^d} (1 - \partial_r^2 \psi_R - \eta(R^4 A_R^2 + B_R^2)) |\partial_r^2 u|^2 dx \\ &\quad + R^{-2} \left(\eta^{-\frac{1}{4}} + R^{-2}\right) \|\partial_d u\|_2 + R^{-4}\eta^{-\frac{1}{2}} \|\partial_d u\|_2^2 + \eta^{-1} R^{-4} + R^{-2}. \end{aligned} \quad (2.25)$$

Next we consider the case that $d \geq 5$. In this case, by Hölder's inequality and the definition of B_R , we can obtain that

$$\begin{aligned} \left| \int_{\mathbb{R}^N} B_R |u|^{2+\frac{8}{d}} dx \right| &= \left| \int_{\mathbb{R}} \int_{|y|\geq R} B_R |u|^{2+\frac{8}{d}} dy dx_d \right| \leq \int_{\mathbb{R}} \|B_R^{\frac{1}{2}}u\|_{L_y^\infty(|y|\geq R)}^{\frac{8}{d}} \|B_R^{\frac{1}{2}-\frac{2}{d}}u\|_{L_y^2}^2 dx_d \\ &\leq \left(\int_{\mathbb{R}} \|B_R^{\frac{1}{2}}u\|_{L_y^\infty(|y|\geq R)}^2 dx_d \right)^{\frac{4}{d}} \left(\int_{\mathbb{R}} \|u\|_{L_y^2}^{\frac{2d}{d-4}} dx_d \right)^{\frac{d-4}{d}}. \end{aligned} \quad (2.26)$$

As an application of (2.4) leads to

$$\int_{\mathbb{R}} \|u\|_{L_y^2}^{\frac{2d}{d-4}} dx_d \lesssim \left(\int_{\mathbb{R}^d} |\partial_d (\|u\|_{L_y^2})|^2 dx_d \right)^{\frac{2}{d-4}} \left(\int_{\mathbb{R}} \|u\|_{L_y^2}^2 dx_d \right)^{\frac{d-2}{d-4}}. \quad (2.27)$$

It then follows from (2.18), (2.27) and the conservation of mass that

$$\int_{\mathbb{R}} \|u\|_{L_y^2}^{\frac{2d}{d-4}} dx_d \lesssim \left(\int_{\mathbb{R}^d} \|\partial_d u\|_{L_y^2}^2 dx_d \right)^{\frac{2}{d-4}} = \|\partial_d u\|_2^{\frac{4}{d-4}}. \quad (2.28)$$

Therefore, coming back to (2.26) and using (2.24) and (2.28), we derive that

$$\begin{aligned} \left| \int_{\mathbb{R}^N} B_R |u|^{2+\frac{8}{d}} dx \right| &\lesssim \left(\int_{\mathbb{R}} \|B_R^{\frac{1}{2}}u\|_{L_y^\infty(|y|\geq R)}^2 dx_d \right)^{\frac{4}{d}} \|\partial_d u\|_2^{\frac{4}{d}} \\ &\lesssim R \int_{\mathbb{R}} \|B_R^{\frac{1}{2}}u\|_{L_y^\infty(|y|\geq R)}^2 dx_d + R^{-\frac{4}{d-4}} \|\partial_d u\|_2^{\frac{4}{d-4}} \\ &\leq R^{-1} \left(\eta^{-\frac{1}{4}} + R^{-2}\right) \|\partial_d u\|_2 + 8R^{-1}\eta^{\frac{1}{4}} \|B_R \partial_r^2 u\|_2 + R^{-\frac{4}{d-4}} \|\partial_d u\|_2^{\frac{4}{d-4}} \\ &\lesssim R^{-1} \left(\eta^{-\frac{1}{4}} + R^{-2}\right) \|\partial_d u\|_2 + 8\eta \|B_R \partial_r^2 u\|_2^2 + R^{-\frac{4}{d-4}} \|\partial_d u\|_2^{\frac{4}{d-4}} + R^{-2}\eta^{-\frac{1}{2}}. \end{aligned}$$

As a consequence, invoking (2.21), we finally derive that

$$\begin{aligned} \frac{d}{dt} \mathcal{M}_R[u(t)] &\leq 16E(u_0) - 8 \int_{\mathbb{R}^d} (1 - \partial_r^2 \psi_R - \eta(R^4 A_R^2 + B_R^2)) |\partial_r^2 u|^2 dx \\ &\quad + R^{-1} \left(\eta^{-\frac{1}{4}} + R^{-2}\right) \|\partial_d u\|_2 + R^{-4}\eta^{-\frac{1}{2}} \|\partial_d u\|_2^2 \\ &\quad + R^{-\frac{4}{d-4}} \|\partial_d u\|_2^{\frac{4}{d-4}} + R^{-2}\eta^{-\frac{1}{2}} + \eta^{-1} R^{-4} + R^{-2}. \end{aligned} \quad (2.29)$$

At this point, using (2.25) and (2.29) and reasoning as the proof of [7, Theorem 3], we are able to finish the proof. This completes the proof. \square

REFERENCES

- [1] G. Baruch, G. Fibich: *Singular solutions of the L^2 -supercritical biharmonic nonlinear Schrödinger equation*, Nonlinearity 24 (6) (2011) 1843-1859.
- [2] G. Baruch, G. Fibich, E. Mandelbaum: *Ring-type singular solutions of the biharmonic nonlinear Schrödinger equation*, Nonlinearity 23 (11) (2010) 2867-2887.
- [3] G. Baruch, G. Fibich, E. Mandelbaum: *Singular solutions of the biharmonic nonlinear Schrödinger equation*, SIAM J. Appl. Math. 70 (8) (2010) 3319-3341.
- [4] J. Bellazzini, L. Forcella, V. Georgiev: *Ground state energy threshold and blow-up for NLS with competing nonlinearities*, arXiv:2012.10977.
- [5] D. Bonheure, J.-B. Castéras, T. Gou, L. Jeanjean: *Strong instability of ground states to a fourth order Schrödinger equation*, Int. Math. Res. Not. IMRN (17) 2019 5299-5315.
- [6] D. Bonheure, J.-B. Castéras, T. Gou, L. Jeanjean: *Normalized solutions to the mixed dispersion nonlinear Schrödinger equation in the mass critical and supercritical regime*, Trans. Amer. Math. Soc. 372 (3) (2019) 2167-2212.
- [7] T. Boulenger, E. Lenzmann: *Blowup for Biharmonic NLS*, Ann. Sci. Éc. Norm. Supér. (4) 50 (3) (2017) 503-44.
- [8] V.D. Dinh: *Dynamics of radial solutions for the focusing fourth-order nonlinear Schrödinger equations*, Nonlinearity 34 (2) (2021) 776-821.
- [9] V.D. Dinh, S. Keraani: *Long time dynamics of nonradial solutions to inhomogeneous nonlinear Schrödinger equations*, SIAM J. Math. Anal. 53 (4) (2021) 4765-4811.
- [10] G. Fibich, B. Ilan, G. Papanicolaou: *Self-focusing with fourth-order dispersion*, SIAM J. Appl. Math. 62 (4) (2002) 1437-1462.
- [11] Q. Guo: *Scattering for the focusing L^2 -supercritical and H^2 -subcritical biharmonic NLS equations*, Comm. Partial Differential Equations 41 (2) (2016) 185-207.
- [12] V.I. Karpman: *Stabilization of soliton instabilities by higher-order dispersion: Fourth-order non-linear schrödinger-type equations*, Phys. Rev. E 53 (1996) R1336-R1339.
- [13] V.I. Karpman, A.G. Shagalov: *Stability of solitons described by nonlinear Schrödinger-type equations with higher-order dispersion*, Phys. D 144 (1-2) (2000) 194-210.
- [14] Y. Martel: *Blow-up for the nonlinear Schrödinger equation in nonisotropic spaces*, Nonlinear Anal. 28 (12) (1997) 1903-1908.
- [15] C. Miao, G. Xu, L. Zhao: *Global well-posedness and scattering for the focusing energy-critical nonlinear Schrödinger equations of fourth order in the radial case*, J. Differential Equations 246 (9) (2009) 3715-3749.
- [16] B. Pausader: *Global well-posedness for energy critical fourth-order Schrödinger equations in the radial case*, Dyn. Partial Differ. Equ. 4 (3) (2007) 197-225.
- [17] B. Pausader: *The cubic fourth-order Schrödinger equation*, J. Funct. Anal. 256 (8) (2009) 2473-2517.
- [18] B. Pausader: *The focusing energy-critical fourth-order Schrödinger equation with radial data*, Discrete Contin. Dyn. Syst. 24 (4) (2009) 1275-1292.
- [19] B. Pausader, S. Xia, *Scattering theory for the fourth-order Schrödinger equation in low dimensions*, Nonlinearity 26 (8) (2013) 2175-2191.
- [20] W.A. Strauss: *Existence of solitary waves in higher dimensions*, Comm. Math. Phys. 55 (2) (1977) 149-162.