

QUASI-CONICAL DOMAINS WITH EMBEDDED EIGENVALUES

DAVID KREJČÍŘÍK AND VLADIMIR LOTOREICHIK

ABSTRACT. The spectrum of the Dirichlet Laplacian on any quasi-conical open set coincides with the non-negative semi-axis. We show that there is a connected quasi-conical open set such that the respective Dirichlet Laplacian has a positive (embedded) eigenvalue. This open set is constructed as the tower of cubes of growing size connected by windows of vanishing size. Moreover, we show that the sizes of the windows in this construction can be chosen so that the absolutely continuous spectrum of the Dirichlet Laplacian is empty.

1. INTRODUCTION

An open set $\Omega \subset \mathbb{R}^d$ with any $d \geq 2$ is said to be *quasi-conical* if it contains an arbitrarily large ball. We consider the self-adjoint Dirichlet Laplacian $-\Delta_D^\Omega$ on Ω acting in the Hilbert space $L^2(\Omega)$. It is easy to see that

$$(1) \quad \sigma(-\Delta_D^\Omega) = [0, \infty)$$

for any quasi-conical open set.

Sufficient conditions to ensure an absence of eigenvalues of the Dirichlet Laplacian on quasi-conical sets are available in the literature. In particular, Rellich proved in [17] that there are no embedded eigenvalues for the Dirichlet Laplacian on the exterior of any compact set. Jones [13] excluded embedded eigenvalues for certain other classes of quasi-conical open sets such as domains conical at infinity. More recently, Bonnet-Ben Dhia *et al.* [2] excluded the existence of eigenvalues in non-convex conical sectors. (The critical case of the Robin Laplacian in a half-space is considered in [3].)

On the other hand, it is easy to construct examples of disconnected quasi-conical open sets for which there are (embedded) eigenvalues in $[0, \infty)$ (*e.g.*, take the union of any bounded set and its exterior). The question whether there are examples of quasi-conical *domains* (*i.e.* *connected* open sets) for which there are eigenvalues seems not to have been considered in the literature. Our goal is to fill in this gap. The main result of the present paper reads as follows.

Theorem 1.

- (i) *Given any positive number λ , there exists a connected quasi-conical open set such that*

$$\lambda \in \sigma_p(-\Delta_D^\Omega).$$

- (ii) *The connected quasi-conical open set in (i) can be constructed so that*

$$\sigma_{ac}(-\Delta_D^\Omega) = \emptyset.$$

Our proof relies on an explicit construction, which we believe is of independent interest. The desired connected quasi-conical open set is constructed as the tower of cubes of growing size connected by windows of vanishing size (see Figure 1). We show that the sizes of the windows connecting the cubes can be chosen so small that we get an embedded eigenvalue and that the absolutely continuous spectrum turns out to be empty. Our result is partially of qualitative nature, because we have no explicit control on the sizes of the windows, which ensure the desired property.

In order to show the existence of an embedded eigenvalue, we start from the tower made of disjoint cubes. The respective Dirichlet Laplacian clearly has an embedded eigenvalue. Then we start to




FIGURE 1. The disconnected quasi-conical open set Ω with places for the small holes to be dug indicated by circles.

subsequently open the windows choosing their sizes so small that this embedded eigenvalue and the respective eigenfunction do not change much (in a proper sense). In the course of this construction we get a family of eigenfunctions weakly convergent to a function, which turns out to be an eigenfunction corresponding to an embedded eigenvalue for the tower of cubes with all the windows opened. In this argument we rely on perturbation results for Dirichlet eigenvalues and eigenfunctions under variation of the open set [4].

The absence of the absolutely continuous spectrum is shown via a perturbation type argument. Clearly, for the disjoint cubes the absolutely continuous of the Dirichlet Laplacian spectrum is empty. Adapting the technique of [12], we show that provided the sizes of the windows are sufficiently small, a suitable resolvent power difference between the Dirichlet Laplacian on a tower of cubes with opened windows and the Dirichlet Laplacian on the disjoint cubes belongs to the trace class. The claim follows from the stability of the absolutely continuous under such a perturbation.

In the classification of Glazman [10] (see also [7, Section X.6.1]) there are three types of open sets: *quasi-conical* (*i.e.* presently considered), *quasi-cylindrical* (*i.e.* not quasi-conical but containing infinitely many pairwise disjoint balls of the same radius), and *quasi-bounded* (*i.e.* neither quasi-conical nor quasi-cylindrical). Quasi-cylindrical open sets with embedded eigenvalues are constructed in a large number of papers. Behind these constructions there is typically a symmetry based argument. For example, it follows from the considerations in [8, 1] (see also [6]) that there is an embedded eigenvalue in the essential spectrum of the Dirichlet Laplacian in an infinite Swiss-cross waveguide. Other examples are given in [22, 21]. The construction in the present paper for quasi-conical domains is significantly different and not based on symmetry. The Dirichlet Laplacian on a quasi-bounded open set typically has purely discrete spectrum, but not necessarily if the set is not bounded.*

To the best of our knowledge, as in the present situation of quasi-conical domains, embedded eigenvalues for unbounded quasi-bounded open sets are not studied in the literature.

The structure of the paper is as follows. First, in Section 2 we provide preliminary material that is used throughout the paper. In Subsection 2.1 we rigorously introduce the Dirichlet Laplacian on a quasi-conical open set and evoke its spectral property (1). Next, in Subsection 2.2 we recall general results on perturbation of eigenvalues, eigenfunctions, and the resolvent of the Dirichlet Laplacian under variation of the open set. Further, in Subsection 2.3 we recall the definitions of the singular values for a compact operator and of the trace class. In Section 3 we prove Theorem 1 (i) on the existence of embedded eigenvalues. Theorem 1 (ii) on the absence of the absolutely continuous spectrum is proved in Section 4. The paper is complemented by Section 5 with a discussion of a related open question.

*In passing, our present construction can be considered as a dual geometry to the celebrated domain “rooms and passages” due to Fraenkel [9], which exhibits an essential spectrum for the Neumann Laplacian on a bounded domain.

2. PRELIMINARIES

2.1. Dirichlet Laplacian on a (quasi-conical) open set. Let $\Omega \subset \mathbb{R}^d$ ($d \geq 2$) be an open set. The self-adjoint Dirichlet Laplacian $-\Delta_D^\Omega$ in $L^2(\Omega)$ can be rigorously introduced using the first representation theorem [18, Thm. 10.7] via the quadratic form

$$\mathfrak{h}_D^\Omega[u] := \int_{\Omega} |\nabla u|^2 dx, \quad \text{dom } \mathfrak{h}_D^\Omega := H_0^1(\Omega).$$

The operator $-\Delta_D^\Omega$ can be characterised as

$$-\Delta_D u = -\Delta u, \quad \text{dom}(-\Delta_D) = \{u \in H_0^1(\Omega) : \Delta u \in L^2(\Omega)\}.$$

Since the Dirichlet Laplacian is a non-negative operator, we have $\sigma(-\Delta_D^\Omega) \subseteq [0, \infty)$ for any open set Ω . Now, assume additionally that Ω is a quasi-conical open set (recall that an open set is called quasi-conical if it contains a ball of arbitrarily large radius). Then, using quasi-modes based on plane waves modulated by suitable cut-off functions in the enlarging balls (see, e.g., [7, Thm. X.6.5]), the opposite inclusion of (1) follows. In particular, the spectrum of the Dirichlet Laplacian on any quasi-conical open set is purely essential.

2.2. Dirichlet Laplacian on varying domains. In this subsection we recall some convergence results for the Dirichlet Laplacian on varying domains contained in [4] (see also [5, 15, 20]). We begin by introducing a type of convergence for a sequence of open sets.

Definition 2.1 ([4, Def. 1.1]). If $\Omega_n, \Omega \subset \mathbb{R}^d$ are such that

- (i) the weak limit points of every sequence $u_n \in H_0^1(\Omega_n)$, $n \in \mathbb{N}$, in $H^1(\mathbb{R}^d)$ are in $H_0^1(\Omega)$,
- (ii) for every $\varphi \in H_0^1(\Omega)$ there exist $\varphi_n \in H_0^1(\Omega_n)$ such that $\varphi_n \rightarrow \varphi$ in $H^1(\mathbb{R}^d)$,

then we write $\Omega_n \rightarrow \Omega$ and say that Ω_n converges to Ω *in the sense of Mosco*.

In general, verifying the conditions of Definition 2.1 can be a difficult problem. However, in the case that the sequence Ω_n approximates Ω monotonously from inside or outside the convergence in the sense of Mosco can be easily checked. In the present paper we only use the approximation from outside. (For any set $S \subset \mathbb{R}^d$ we denote by $\text{int}(S)$ its interior.)

Proposition 2.2 ([4, Prop. 7.4 and Thm. 7.5], [5, Sec. 5.4]). Suppose that $\Omega_n \supset \Omega_{n+1} \supset \Omega$ for all $n \in \mathbb{N}$, and that $\text{int}(\bigcap_{n \in \mathbb{N}} \Omega_n) = \Omega$. If $H_0^1(\Omega) = H_0^1(\Omega \cup \Gamma)$, where

$$\Gamma := \bigcap_{n \in \mathbb{N}} \left(\overline{\bigcup_{k \geq n} (\Omega_k \cap \partial\Omega)} \right) \subset \partial\Omega,$$

then $\Omega_n \rightarrow \Omega$.

The convergence of the open sets in the sense of Mosco is convenient in the study of convergence of the respective Dirichlet Laplacians.

Proposition 2.3 ([4, Cor. 4.7]). Suppose that Ω_n, Ω are contained in a fixed bounded set and that $\Omega_n \rightarrow \Omega$. Then

$$\left\| (-\Delta_D^{\Omega_n} + 1)^{-1} - (-\Delta_D^\Omega + 1)^{-1} \right\| \rightarrow 0, \quad n \rightarrow \infty.$$

This norm-resolvent convergence particularly implies the convergence of the respective spectral projections.

2.3. Singular values and the trace class. In this subsection we recall the concepts of singular values and the trace class. Further details can be found in the monographs [11, 19].

Definition 2.4. For a compact operator K in a Hilbert space the eigenvalues $\{s_k(K)\}_{k \geq 1}$ of the self-adjoint non-negative compact operator $(K^*K)^{1/2}$ enumerated in the non-increasing order and repeated with multiplicities taken into account are called the *singular values* of K .

For a compact operator K and a bounded closed operator B acting in a Hilbert space the products BK and KB are compact and by [19, Thm. 1.6]

$$(2) \quad s_k(BK) \leq \|B\| s_k(K) \quad \text{and} \quad s_k(KB) \leq \|B\| s_k(K).$$

Definition 2.5. A compact operator K is said to belong to the *trace class* if $\sum_{k=1}^{\infty} s_k(K) < \infty$.

According to [19, Sec. 2], the trace class is a Banach space with the norm given by

$$(3) \quad \|K\|_1 := \sum_{k=1}^{\infty} s_k(K).$$

3. EMBEDDED EIGENVALUES

In this section we prove part (i) of Theorem 1 on the existence of a connected quasi-conical open set with an embedded eigenvalue. The argument is performed via an explicit construction. For clarity, we divide the proof into several steps.

Disconnected set. Let $Q_a(x) \subset \mathbb{R}^d$ denote an open cube with sides of half-width $a > 0$ centred at a point $x \in \mathbb{R}^d$. With a slight abuse of notation we will occasionally interpret $t \in \mathbb{R}$ as $(t, 0, 0, \dots, 0) \in \mathbb{R}^d$. We aim at constructing a tower of cubes of growing size, in which each next cube is placed “on the top” of the previous one. To this aim, let us introduce the sequence of centres $0 < x_1 < x_2 < \dots < x_n < \dots$ such that the associated sequence of half-widths $a_1 := x_1$, $a_n := x_n - x_{n-1} - a_{n-1}$ ($n \geq 2$) satisfies $a_n > 0$ for all $n \in \mathbb{N}$ and $a_n \rightarrow \infty$ as $n \rightarrow \infty$. At this stage we fix only $x_1 > 0$ while the sequence $\{x_n\}_{n \geq 2}$ is not fixed and we will specify it in the course of the construction. Let λ_1 be the lowest (simple) eigenvalue of $-\Delta_D^{Q_{a_1}(x_1)}$. Let ψ_1 denote a corresponding eigenfunction, normalised to 1 in $L^2(Q_{a_1}(x_1))$.

Define a (disconnected) quasi-conical open set Ω as a disjoint union of enlarging cubes:

$$(4) \quad \Omega := \bigcup_{n=1}^{\infty} \omega_n \quad \text{with} \quad \omega_n := Q_{a_n}(x_n).$$

Then (extending ψ_1 by zero to Ω), it is easy to see that ψ_1 is an eigenfunction of $-\Delta_D^\Omega$ corresponding to λ . Thus, we have constructed a disconnected quasi-conical open set with an embedded eigenvalue.

To construct a connected open set based on Ω , we dig small holes at the places where the disjoint cubes ω_n touch (see Figure 1). A careful attention is needed in order to ensure that this construction does not eliminate the embedded eigenvalue.

1st interconnection. Let us set $\Omega_1 := \omega_1$ and let us choose $x_2 > x_1$ so that $a_2 = x_2 - x_1 - a_1 \geq a_1 + 1$ and that λ_1 is a simple eigenvalue of $-\Delta_D^{\Omega_1 \cup \omega_2}$. For any positive $\delta_1 < a_1$, let us connect the first two cubes $\Omega_1 = \omega_1$ and ω_2 by defining

$$(5) \quad \Omega_2 := \Omega_1 \cup \omega_2 \cup Q_{\delta_1}(x_1 + a_1).$$

It follows from Proposition 2.2 (with $\Gamma = \emptyset$) that Ω_2 converges to $\Omega_1 \cup \omega_2$ as $\delta_1 \rightarrow 0$ in the sense of Definition 2.1. Hence, Proposition 2.3 combined with [14, Sec. IV.3.5] yields that given any (small) positive

number ε_1 , there exist a positive half-width δ_1 , a simple eigenvalue λ_2 of $-\Delta_D^{\Omega_2}$ and a corresponding eigenfunction ψ_2 , normalised to 1 in $L^2(\Omega_2)$, such that

$$(6) \quad |\lambda_2 - \lambda_1| \leq \varepsilon_1 \quad \text{and} \quad \|\psi_2 - \psi_1\|_{L^2(\Omega_2)} \leq \varepsilon_1.$$

Using additionally the monotonicity of Dirichlet eigenvalues with respect to inclusion of open sets, the first inequality in (6) implies that we can choose δ_1 so small that

$$(7) \quad \lambda_1 - \varepsilon_1 \leq \lambda_2 \leq \lambda_1.$$

Using the normalisations of ψ_1 and ψ_2 , together with the fact that $\psi_1 = 0$ in ω_2 , the second inequality in (6) yields

$$(8) \quad \|\psi_2\|_{L^2(\Omega_2 \setminus \Omega_1)} \leq \varepsilon_1 \quad \text{and} \quad \|\psi_2\|_{L^2(\Omega_1)} \geq 1 - \varepsilon_1.$$

2nd interconnection. Similarly, let us choose $x_3 > x_2$ so that $a_3 = x_3 - x_2 - a_2 \geq a_2 + 1$ and that λ_2 (constructed in the previous step) is a simple eigenvalue of $-\Delta_D^{\Omega_2 \cup \omega_3}$. For any $\delta_2 < a_2$, let us connect Ω_2 with the third cube ω_3 by defining

$$(9) \quad \Omega_3 := \Omega_2 \cup \omega_3 \cup Q_{\delta_2}(x_2 + a_2).$$

Arguing as before, given any positive number ε_2 , there exist a positive half-width δ_2 , a simple eigenvalue λ_3 of $-\Delta_D^{\Omega_3}$, and a corresponding eigenfunction ψ_3 , normalised to 1 in $L^2(\Omega_3)$, such that

$$(10) \quad |\lambda_3 - \lambda_2| \leq \varepsilon_2 \quad \text{and} \quad \|\psi_3 - \psi_2\|_{L^2(\Omega_3)} \leq \varepsilon_2.$$

Using the monotonicity of Dirichlet eigenvalues and (7), the first inequality in (10) implies that we can choose δ_2 so small that

$$(11) \quad \lambda_1 - \varepsilon_1 - \varepsilon_2 \leq \lambda_3 \leq \lambda_2 \leq \lambda_1.$$

From the second inequality in (10) and the first inequality in (8), we deduce

$$(12) \quad \|\psi_3\|_{L^2(\Omega_3 \setminus \Omega_1)} \leq \|\psi_3 - \psi_2\|_{L^2(\Omega_3 \setminus \Omega_1)} + \|\psi_2\|_{L^2(\Omega_3 \setminus \Omega_1)} \leq \varepsilon_2 + \varepsilon_1.$$

Consequently,

$$(13) \quad 1 = \|\psi_3\|_{L^2(\Omega_3)} \leq \|\psi_3\|_{L^2(\Omega_1)} + \|\psi_3\|_{L^2(\Omega_3 \setminus \Omega_1)} \leq \|\psi_3\|_{L^2(\Omega_1)} + \varepsilon_2 + \varepsilon_1.$$

That is,

$$(14) \quad \|\psi_3\|_{L^2(\Omega_1)} \geq 1 - \varepsilon_1 - \varepsilon_2.$$

nth interconnection. Continuing by induction, given any natural number $n \geq 1$, we choose $x_{n+1} > x_n$ so that $a_{n+1} = x_{n+1} - x_n - a_n \geq a_n + 1$ and λ_n (constructed in the $(n-1)$ st step) is a simple eigenvalue of $-\Delta_D^{\Omega_n \cup \omega_{n+1}}$. For any $\delta_n < a_n$, let us connect Ω_n with the $(n+1)$ th cube ω_{n+1} by defining

$$(15) \quad \Omega_{n+1} := \Omega_n \cup \omega_{n+1} \cup Q_{\delta_n}(x_n + a_n).$$

As before given any positive number ε_n , there exist a positive half-width δ_n , a simple eigenvalue λ_{n+1} of $-\Delta_D^{\Omega_{n+1}}$ and a corresponding eigenfunction ψ_{n+1} , normalised to 1 in $L^2(\Omega_{n+1})$, such that

$$(16) \quad \lambda_1 - (\varepsilon_1 + \dots + \varepsilon_n) \leq \lambda_{n+1} \leq \lambda_n \leq \lambda_1 \quad \text{and} \quad \|\psi_{n+1}\|_{L^2(\Omega_1)} \geq 1 - (\varepsilon_1 + \dots + \varepsilon_n).$$

Given any positive number ε , and choosing $\varepsilon_k := 2^{-k}\varepsilon$ for every $k \geq 1$, we get uniform estimates

$$(17) \quad \lambda_1 - \varepsilon \leq \lambda_{n+1} \leq \lambda_1 \quad \text{and} \quad \|\psi_{n+1}\|_{L^2(\Omega_1)} \geq 1 - \varepsilon.$$

Connected set. By using the procedure of connecting the individual cubes described above, we define a connected open set Ω' by the limit

$$(18) \quad \Omega' := \lim_{n \rightarrow \infty} \Omega_n.$$

It remains to show that the embedded eigenvalue “survives” this limit. After the n th interconnection, we have the eigenvalue equation

$$(19) \quad \forall \phi \in H_0^1(\Omega_{n+1}), \quad (\nabla \phi, \nabla \psi_{n+1})_{L^2(\Omega_{n+1})} = \lambda_{n+1} (\phi, \psi_{n+1})_{L^2(\Omega_{n+1})}.$$

As usual, extending ψ_{n+1} and ϕ by zero to the whole Ω' , the inner product can be replaced by that of $L^2(\Omega')$. Choosing $\phi := \psi_{n+1}$ for the test function in (19) and using the normalisation of ψ_{n+1} , we obtain

$$(20) \quad \|\nabla \psi_{n+1}\|_{L^2(\Omega')}^2 \leq \lambda_{n+1} \leq \lambda_1 \quad \text{and} \quad \|\psi_{n+1}\|_{L^2(\Omega')}^2 = 1.$$

Here the uniform bound to λ_{n+1} follows from (17). It follows that $\{\psi_{n+1}\}_{n=0}^\infty$ is a bounded sequence in $H_0^1(\Omega')$. Consequently, there exists a subsequence $\{\psi_{n_j+1}\}_{j=0}^\infty$ which is weakly converging to some ψ_∞ in $H_0^1(\Omega')$. In particular, $\{\psi_{n_j+1}\}_{j=0}^\infty$ is also weakly converging to ψ_∞ in $L^2(\Omega')$. Inserting this subsequence into (19) and sending j to infinity, we obtain

$$(21) \quad \forall \phi \in H_0^1(\Omega'), \quad (\nabla \phi, \nabla \psi_\infty)_{L^2(\Omega')} = \lambda_\infty (\phi, \psi_\infty)_{L^2(\Omega')},$$

where the limit

$$(22) \quad \lambda_\infty := \lim_{n \rightarrow \infty} \lambda_n$$

is well defined because $\{\lambda_n\}_{n=1}^\infty$ is monotonously decreasing by construction. Consequently, ψ_∞ belongs to the domain of the operator $-\Delta_D^{\Omega'}$ and solves the equation $-\Delta_D^{\Omega'} \psi_\infty = \lambda_\infty \psi_\infty$ with $\lambda_1 - \varepsilon \leq \lambda_\infty \leq \lambda_1$. It remains to ensure that ψ_∞ is not identically equal to zero. However, this follows from the uniform estimate (17) on the norm of ψ_{n+1} in the n -independent bounded set Ω_1 , provided that $\varepsilon < 1$.

We remark that by rescaling the domain Ω' by a constant factor $\alpha > 0$ we get the domain $\alpha\Omega'$ and the respective Dirichlet Laplacian $-\Delta_D^{\alpha\Omega'}$ has an embedded eigenvalue $\alpha^{-2}\lambda_\infty$. Hence, we can construct a connected quasi-conical open set with an embedded eigenvalue at any prescribed position.

Remark 3.1. The same type of examples of quasi-conical domains with embedded eigenvalues can be given for other cell domains than cubes. Moreover, the interconnections obtained by digging the holes for flat boundaries can be replaced for general cell domains by using thin tubes as connections.

4. ABSENCE OF THE ABSOLUTELY CONTINUOUS SPECTRUM

In this section we prove part (ii) of Theorem 1 by showing that by choosing the sizes of the windows $\{\delta_n\}_{n \geq 1}$ sufficiently small the absolutely continuous spectrum of the operator $-\Delta_D^{\Omega'}$ becomes empty. In order to prove this result we adapt the technique developed in [12]. As in Section 3, we subsequently open the windows between the cubes. The half-widths of the windows $\{\delta_n\}_{n \geq 1}$ will be determined so as to satisfy simultaneously the conditions in Section 3 and some new additional conditions.

Let the sequence of the bounded domains $\{\Omega_n\}_{n \in \mathbb{N}}$ be constructed as in Section 3. We intend to modify the sizes of the windows $\{\delta_n\}_{n \geq 1}$ in order to satisfy some additional assumptions. The choice of the sequence $\{x_n\}_{n \in \mathbb{N}}$ in Section 3 easily adjusts to possibly different choices of the sizes of the holes. It is not hard to see that in the construction in Section 3 on the n th interconnection we are allowed to take any $\delta_n \in (0, \delta_n^*)$ for some sufficiently small $\delta_n^* > 0$, which depends on the values of the parameters chosen on the previous steps of the construction. For the convenience of the reader we split the remaining argument into two steps.

Step 1: modified sizes of the windows. Without loss of generality we can assume that $\delta_n \leq a_1$ for all $n \in \mathbb{N}$. Since $\Omega_{n+1} \rightarrow \Omega_n \cup \omega_{n+1}$ as $\delta_n \rightarrow 0$ in the sense of Definition 2.1, applying Proposition 2.3 we conclude that for any $\varepsilon > 0$ there exists $\delta_n \in (0, \delta_n^*)$ such that

$$(23) \quad \left\| (-\Delta_D^{\Omega_{n+1}} + 1)^{-1} - (-\Delta_D^{\Omega_n \cup \omega_{n+1}} + 1)^{-1} \right\| \leq \varepsilon.$$

Let the power $\ell \in \mathbb{N}$ be fixed. By purely algebraic manipulations the resolvent power difference of the Dirichlet Laplacians on Ω_{n+1} and $\Omega_n \cup \omega_{n+1}$ acting in the same Hilbert space $L^2(\Omega_n \cup \omega_{n+1})$ can be represented by

$$(24) \quad \begin{aligned} & (-\Delta_D^{\Omega_{n+1}} + 1)^{-\ell} - (-\Delta_D^{\Omega_n \cup \omega_{n+1}} + 1)^{-\ell} = \\ & = \sum_{k=1}^{\ell} (-\Delta_D^{\Omega_{n+1}} + 1)^{-(\ell-k)} \left[(-\Delta_D^{\Omega_{n+1}} + 1)^{-1} - (-\Delta_D^{\Omega_n \cup \omega_{n+1}} + 1)^{-1} \right] (-\Delta_D^{\Omega_n \cup \omega_{n+1}} + 1)^{-(k-1)}. \end{aligned}$$

We conclude from the Weyl spectral asymptotics for the Dirichlet Laplacian on a bounded open set (see *e.g.* [18, Thm. 12.14]) that for any $k \in \{1, 2, \dots, \ell\}$ at least one of the two operators $(-\Delta_D^{\Omega_{n+1}} + 1)^{-(\ell-k)}$ and $(-\Delta_D^{\Omega_n \cup \omega_{n+1}} + 1)^{-k-1}$ belongs to the trace class provided that ℓ is chosen to be sufficiently large. The trace norm of this trace class resolvent power can be estimated from above by a constant $C_n > 0$ independent of k and δ_n while the operator norm of the other resolvent power can be estimated from above by 1 thanks to non-negativity of the Dirichlet Laplacian. Let us comment on the possibility to get an upper bound on this trace norm independent of δ_n . Indeed the operator $-\Delta_D^{\Omega_n \cup \omega_{n+1}}$ is independent of δ_n and hence the trace norm of the corresponding resolvent power (provided it belongs to the trace class) is also independent of δ_n . The trace norm of the resolvent power corresponding to $-\Delta_D^{\Omega_{n+1}}$ (provided it belongs to the trace class) can be expressed as

$$(25) \quad \left\| (-\Delta_D^{\Omega_{n+1}} + 1)^{-(\ell-k)} \right\|_1 = \sum_{m=1}^{\infty} \frac{1}{[1 + \lambda_m(-\Delta_D^{\Omega_{n+1}})]^{\ell-k}}$$

where $\{\lambda_m(-\Delta_D^{\Omega_{n+1}})\}_{m \geq 1}$ are the eigenvalues $-\Delta_D^{\Omega_{n+1}}$ enumerated in the non-decreasing way and repeated with multiplicities taken into account. It is easy to see by the min-max principle that the eigenvalues of $-\Delta_D^{\Omega_{n+1}}$ are non-increasing functions of δ_n . Hence, the trace norm in (25) is an increasing function of δ_n and therefore it can be estimated from above by the respective trace norm for $\delta_n = a_1$.

Combining the estimate (23) with the property of singular values in (2) and the definition of the trace norm in (3) we derive from (24) the bound

$$(26) \quad \left\| (-\Delta_D^{\Omega_{n+1}} + 1)^{-\ell} - (-\Delta_D^{\Omega_n \cup \omega_{n+1}} + 1)^{-\ell} \right\|_1 \leq C_n \ell \varepsilon, \quad n \in \mathbb{N}.$$

It follows from (23) and (26) that by choosing $\varepsilon > 0$ sufficiently small we can find a sufficiently small $\delta_n \in (0, \delta_n^*)$ so that

$$(27a) \quad \left\| (-\Delta_D^{\Omega_{n+1}} + 1)^{-1} - (-\Delta_D^{\Omega_n \cup \omega_{n+1}} + 1)^{-1} \right\| \leq 2^{-n}, \quad n \in \mathbb{N},$$

$$(27b) \quad \left\| (-\Delta_D^{\Omega_{n+1}} + 1)^{-\ell} - (-\Delta_D^{\Omega_n \cup \omega_{n+1}} + 1)^{-\ell} \right\|_1 \leq 2^{-n}, \quad n \in \mathbb{N}.$$

Thus, we have selected the sequence $\{\delta_n\}_{n \geq 1}$ which fits the construction in Section 3 and simultaneously the above two estimates hold.

Step 2: trace class resolvent power difference. Consider the sequence of quasi-conical unbounded domains

$$\begin{aligned}\Omega'_1 &:= \Omega, \\ \Omega'_2 &:= \Omega \cup Q_{\delta_1}(x_1 + a_1), \\ \Omega'_3 &:= \Omega \cup Q_{\delta_1}(x_1 + a_1) \cup Q_{\delta_2}(x_2 + a_2), \\ &\vdots \\ \Omega'_n &:= \Omega \cup \left(\bigcup_{k=1}^{n-1} Q_{\delta_k}(x_k + a_k) \right), \\ &\vdots\end{aligned}$$

The Dirichlet Laplacian on Ω'_n can be decomposed into the orthogonal sum

$$-\Delta_{\mathbb{D}}^{\Omega'_n} = (-\Delta_{\mathbb{D}}^{\Omega_n}) \oplus (-\Delta_{\mathbb{D}}^{\Omega \setminus \overline{\Omega_n}})$$

with respect to the decomposition $L^2(\Omega) = L^2(\Omega_n) \oplus L^2(\Omega \setminus \overline{\Omega_n})$. In view of this decomposition we obtain the following representation for the resolvent difference of the Dirichlet Laplacians on Ω'_n and Ω'_{n+1} :

$$(-\Delta_{\mathbb{D}}^{\Omega'_{n+1}} + 1)^{-1} - (-\Delta_{\mathbb{D}}^{\Omega'_n} + 1)^{-1} = \left((-\Delta_{\mathbb{D}}^{\Omega_{n+1}} + 1)^{-1} - (-\Delta_{\mathbb{D}}^{\Omega_n \cup \omega_{n+1}} + 1)^{-1} \right) \oplus 0.$$

Hence, it follows from (27a) that the sequence $\mathbb{N} \ni n \mapsto (-\Delta_{\mathbb{D}}^{\Omega'_n} + 1)^{-1}$ is a Cauchy sequence in the operator norm. Thus, there exists an operator \mathbb{T} in the Hilbert space $L^2(\Omega)$ such that

$$(28) \quad \left\| (-\Delta_{\mathbb{D}}^{\Omega'_n} + 1)^{-1} - \mathbb{T} \right\| \rightarrow 0, \quad n \rightarrow \infty.$$

On the other hand the sequence of operators $\{-\Delta_{\mathbb{D}}^{\Omega'_n}\}_{n \geq 1}$ is decreasing in the form sense. In view of the identity $\cup_{n \geq 1} H_0^1(\Omega'_n) = H_0^1(\Omega')$ ($\Omega' = \cup_{n \geq 1} \Omega'_n$) we get by [16, Thm. S.16] that $-\Delta_{\mathbb{D}}^{\Omega'_n}$ converges to $-\Delta_{\mathbb{D}}^{\Omega'}$ in the strong resolvent sense. Thanks to (28) we obtain that $\mathbb{T} = (-\Delta_{\mathbb{D}}^{\Omega'} + 1)^{-1}$ and that $-\Delta_{\mathbb{D}}^{\Omega'_n}$ converges to $-\Delta_{\mathbb{D}}^{\Omega'}$ in the norm resolvent sense.

Analogously it follows from (27b) that the sequence of resolvent powers $\mathbb{N} \ni n \mapsto (-\Delta_{\mathbb{D}}^{\Omega'_n} + 1)^{-\ell}$ is a Cauchy sequence in the trace norm. Hence, it has a limit \mathbb{T}' in the trace norm. Since the trace norm is stronger than the operator norm, we obtain that $(-\Delta_{\mathbb{D}}^{\Omega'_n} + 1)^{-\ell}$ converges to \mathbb{T}' also in the operator norm as $n \rightarrow \infty$. By the uniqueness of the limit we get that $\mathbb{T}' = (-\Delta_{\mathbb{D}}^{\Omega'} + 1)^{-\ell}$. Therefore, we obtain

$$\begin{aligned}\left\| (-\Delta_{\mathbb{D}}^{\Omega} + 1)^{-\ell} - (-\Delta_{\mathbb{D}}^{\Omega'} + 1)^{-\ell} \right\|_1 &\leq \sum_{n=1}^{\infty} \left\| (-\Delta_{\mathbb{D}}^{\Omega'_n} + 1)^{-\ell} - (-\Delta_{\mathbb{D}}^{\Omega'_{n+1}} + 1)^{-\ell} \right\|_1 \\ &= \sum_{n=1}^{\infty} \left\| (-\Delta_{\mathbb{D}}^{\Omega_{n+1}} + 1)^{-\ell} - (-\Delta_{\mathbb{D}}^{\Omega_n \cup \omega_{n+1}} + 1)^{-\ell} \right\|_1 \leq \sum_{n=1}^{\infty} 2^{-n} < \infty.\end{aligned}$$

Thus, the resolvent power difference $(-\Delta_{\mathbb{D}}^{\Omega} + 1)^{-\ell} - (-\Delta_{\mathbb{D}}^{\Omega'} + 1)^{-\ell}$ is a trace class operator. Hence, by [14, Thm. X.4.12, Rem. X.4.13] we have

$$\sigma_{\text{ac}}(-\Delta_{\mathbb{D}}^{\Omega'}) = \sigma_{\text{ac}}(-\Delta_{\mathbb{D}}^{\Omega}) = \emptyset.$$

In the course of this construction we have also satisfied all the assumption in Section 3. Hence, the operator $-\Delta_{\mathbb{D}}^{\Omega'}$ possesses also a positive embedded eigenvalue λ_{∞} .

5. DISCUSSION

It remains an open question whether it is possible to select the sizes of the windows so small that the singular continuous spectrum of the operator $-\Delta_D^{\Omega'}$ is empty. If it is possible to do, then we would get the Dirichlet Laplacian on a connected quasi-conical open set having only purely point spectrum, which densely fills the non-negative semi-axis.

Acknowledgement. D.K. was supported by the EXPRO grant No. 20-17749X of the Czech Science Foundation. V.L. was supported by the grant No. 21-07129S of the Czech Science Foundation.

REFERENCES

- [1] Y. Avishai, D. Bessis, B. G. Giraud, and G. Mantica, *Quantum bound states in open geometries*, Phys. Rev. B **44** (1991), 8028–8034.
- [2] A.-S. Bonnet-Ben Dhia, S. Fliss, C. Hazard, and A. Tonnoir, *A Rellich type theorem for the Helmholtz equation in a conical domain*, C. R. Acad. Sci. Paris **354** (2016), 27–32.
- [3] L. Cossetti and D. Krejčířík, *Absence of eigenvalues of non-self-adjoint Robin Laplacians on the half-space*, Proc. London. Math. Soc. **121** (2020), 584–616.
- [4] D. Daners, *Dirichlet problems on varying domains*, J. Differential Equations **188** (2003), 591–624.
- [5] ———, *Domain perturbation for linear and semi-linear boundary value problems*, Handbook of differential equations: Stationary partial differential equations. Vol. VI, Amsterdam: Elsevier/North Holland, 2008, pp. 1–81.
- [6] M. Dauge, Y. Lafranche, and N. Raymond, *Quantum waveguides with corners*, ESAIM, Proc. **35** (2012), 14–45.
- [7] D. E. Edmunds and W. D. Evans, *Spectral theory and differential operators*, Oxford University Press, Oxford, 1987.
- [8] P. Exner, P. Šeba, and P. Šťovíček, *On existence of a bound state in an L-shaped waveguide*, Czech. J. Phys. B **39** (1989), 1181–1191.
- [9] L. E. Fraenkel, *On regularity of the boundary in the theory of Sobolev spaces*, Proc. Lond. Math. Soc. **39** (1979), 385–427.
- [10] I. M. Glazman, *Direct methods of qualitative spectral analysis of singular differential operators*, Israel Program for Scientific Translations, 1965.
- [11] I. C. Gohberg and M. G. Kreĭn, *Introduction to the theory of linear nonselfadjoint operators in Hilbert space*, Amer. Math. Soc., 1969.
- [12] R. Hempel, L. A. Seco, and B. Simon, *The essential spectrum of Neumann Laplacians on some bounded singular domains*, J. Funct. Anal. **102** (1991), 448–483.
- [13] D. S. Jones, *The eigenvalues of $\nabla^2 u + \lambda u = 0$ when the boundary conditions are given on semi-infinite domains*, Mathematical Proceedings of the Cambridge Philosophical Society **49** (1953), no. 4, 668–684.
- [14] T. Kato, *Perturbation theory for linear operators*, Springer-Verlag, Berlin, 1966.
- [15] J. Rauch and M. Taylor, *Potential and scattering theory on wildly perturbed domains*, J. Funct. Anal. **18** (1975), 27–59.
- [16] M. Reed and B. Simon, *Methods of modern mathematical physics, I. Functional analysis*, Academic Press, New York, 1972.
- [17] F. Rellich, *Über das asymptotische Verhalten der Lösungen von $\Delta u + \lambda u = 0$ in unendlichen Gebieten*, Jber. dtsh. MatVer. **53** (1943), 57–65.
- [18] K. Schmüdgen, *Unbounded self-adjoint operators on Hilbert space*, vol. 265, Dordrecht: Springer, 2012.
- [19] B. Simon, *Trace ideals and their applications*, vol. 120, Providence, RI: American Mathematical Society, 2005.
- [20] P. Stollmann, *A convergence theorem for Dirichlet forms with applications to boundary value problems with varying domains*, Math. Z. **219** (1995), 275–287.
- [21] K. J. Witsch, *Examples of embedded eigenvalues for the Dirichlet-Laplacian in domains with infinite boundaries*, Math. Methods Appl. Sci. **12** (1990), 177–182.
- [22] ———, *Examples of embedded eigenvalues for the Dirichlet Laplacian in perturbed waveguides*, Math. Methods Appl. Sci. **12** (1990), 91–93.

(D. KREJČÍŘÍK) DEPARTMENT OF MATHEMATICS, FACULTY OF NUCLEAR SCIENCES AND PHYSICAL ENGINEERING, CZECH TECHNICAL UNIVERSITY IN PRAGUE, TROJANOVA 13, 120 00, PRAGUE, CZECH REPUBLIC
E-MAIL: DAVID.KREJCIRIK@FJFI.CVUT.CZ

(V. LOTOREICHIK) DEPARTMENT OF THEORETICAL PHYSICS, NUCLEAR PHYSICS INSTITUTE, CZECH ACADEMY OF SCIENCES, 25068 ŘEŽ, CZECH REPUBLIC
E-MAIL: LOTOREICHIK@UJF.CAS.CZ