

# UNIFORM DENSITY ESTIMATES AND $\Gamma$ -CONVERGENCE FOR THE ALT-PHILLIPS FUNCTIONAL OF NEGATIVE POWERS

D. DE SILVA AND O. SAVIN

ABSTRACT. We obtain density estimates for the free boundaries of minimizers  $u \geq 0$  of the Alt-Phillips functional involving negative power potentials

$$\int_{\Omega} (|\nabla u|^2 + u^{-\gamma} \chi_{\{u>0\}}) dx, \quad \gamma \in (0, 2).$$

These estimates remain uniform as the parameter  $\gamma \rightarrow 2$ . As a consequence we establish the uniform convergence of the corresponding free boundaries to a minimal surface as  $\gamma \rightarrow 2$ . The results are based on the  $\Gamma$ -convergence of these energies (properly rescaled) to the Dirichlet-perimeter functional

$$\int_{\Omega} |\nabla u|^2 dx + Per_{\Omega}(\{u = 0\}),$$

considered by Athanasopoulous, Caffarelli, Kenig, and Salsa.

## 1. INTRODUCTION

Energy functionals involving the Dirichlet integral of a density  $u$  and a potential term  $W(u)$

$$\int_{\Omega} |\nabla u|^2 + W(u) dx,$$

appear in various models in the calculus of variations. A classical example is the Allen-Cahn [AC] energy given by the double-well potential

$$W(t) = (1 - t^2)^2,$$

which is relevant in the theory of phase-transitions and minimal surfaces. In their celebrated result, Modica and Mortola [MM] showed that 0-homogenous rescalings of bounded minimizers  $|u| \leq 1$ , converge up to subsequences to a  $\pm 1$  configuration separated by a minimal surface, i.e.

$$(1.1) \quad u_{\epsilon}(x) = u\left(\frac{x}{\epsilon}\right) \rightarrow \chi_E - \chi_{E^c} \quad \text{in } L^1_{loc}, \quad \text{as } \epsilon \rightarrow 0,$$

with  $E$  a set of minimal perimeter. At the level of the energy, this result is expressed in terms of the Gamma-convergence of the rescaled energies

$$\int_{\Omega} \epsilon |\nabla u|^2 + \frac{1}{\epsilon} W(u) dx,$$

to a multiple of the perimeter functional  $c_0 Per_{\Omega}(E)$ .

Other examples of energies appear in the theory of free boundary problems. When the potential  $W(t)$  is not of class  $C^{1,1}$  near a minimum point, say  $t = 0$ , minimizers can develop patches where they take this value. The boundary of such a patch  $\partial\{u = 0\}$  is the free boundary. Two particular potentials of interest are given by

$$W(t) = t^+,$$

which corresponds to the obstacle problem (for a comprehensive survey see [PSU]), and by

$$W(t) = \chi_{\{t>0\}},$$

which corresponds to the Bernoulli free boundary problem (see for example [AC, ACF, CS]). These can be viewed as part of the family of power-potentials

$$W(t) = (t^+)^{\beta}, \quad \beta \in [0, 2),$$

which were considered by Alt and Phillips [AP] in the early 80's.

Recently in [DS], we investigated properties of non-negative minimizers and their free boundaries for Alt-Phillips potentials of negative powers

$$W(t) = t^{-\gamma} \chi_{\{t>0\}}, \quad \gamma \in (0, 2).$$

These potentials are relevant in the applications, for example in liquid models with large cohesive internal forces in regions of low density. The upper bound  $\gamma < 2$  is necessary for the finiteness of the energy.

In [DS] we showed that minimizers  $u \geq 0$  of the Alt-Phillips functional involving negative power potentials

$$(1.2) \quad \mathcal{E}_{\gamma}(u) := \int_{\Omega} (|\nabla u|^2 + u^{-\gamma} \chi_{\{u>0\}}) dx, \quad \gamma \in (0, 2),$$

have optimal  $C^{\alpha}$  Hölder continuity. The free boundary

$$F(u) := \partial\{u > 0\}$$

is characterized by an expansion of the type

$$u = c_{\alpha} d^{\alpha} + o(d^{2-\alpha}), \quad \alpha := \frac{2}{2+\gamma} \in \left(\frac{1}{2}, 1\right),$$

where  $d$  denotes the distance to  $F(u)$  and  $c_{\alpha} d^{\alpha}$  represents the explicit 1D homogeneous solution. Furthermore, we showed that  $F(u)$  is a hypersurface of class  $C^{1,\beta}$  up to a closed singular set of dimension at most  $n - k(\gamma)$ , where  $k(\gamma) \geq 3$  is the first dimension in which a nontrivial  $\alpha$ -homogeneous minimizer exists. We also established the Gamma-convergence of a suitable multiple of the  $\mathcal{E}_{\gamma}$  to the perimeter of the positivity set  $Per_{\Omega}(\{u > 0\})$  as  $\gamma \rightarrow 2$ .

In this work we investigate in more detail the properties of minimizers as the parameter  $\gamma$  tends to the critical value 2, and make precise the connection between their free boundaries and the theory of minimal surfaces. In particular we establish density estimates and the uniform convergence (up to subsequences) of the free boundaries  $F(u_k)$  to a minimal surface, for a sequence of bounded minimizers  $u_k$  corresponding to parameters  $\gamma_k \rightarrow 2$ , see Corollary 2.6. Uniform convergence results in different settings were obtained by Caffarelli and Cordoba [CC] for the Allen-Cahn energy and the convergence in (1.1), and by Caffarelli and Valdinoci [CV] for the  $s$ -nonlocal minimal surfaces with  $s \rightarrow 1$ . We also refer the reader to other related works in similar contexts [ADM, DFPV, PV, SV, V].

The constants in the Hölder and density estimates obtained in [DS] degenerate as  $\gamma \rightarrow 2$ . However, here we develop uniform estimates in  $\gamma$ , and for this it is convenient to rescale the potential term in the functional  $\mathcal{E}_{\gamma}$  in a suitable way (see (2.1)). We further establish the Gamma-convergence to the Dirichlet-perimeter functional

$$\mathcal{F}(u) := \int_{\Omega} |\nabla u|^2 dx + Per_{\Omega}(\{u = 0\}),$$

which was studied by Athanassopoulos, Caffarelli, Kenig, Salsa in [ACKS]. Heuristically, this shows that the cohesive term  $W$  has the effect of surface tension as  $\gamma \rightarrow 2$ .

## 2. MAIN RESULTS

Let  $\Omega$  be a bounded domain in  $\mathbb{R}^n$  with Lipschitz boundary. We consider  $J_\gamma$ , a rescaling of  $\mathcal{E}_\gamma$ , which acts on functions

$$u : \Omega \rightarrow \mathbb{R}, \quad u \in H^1(\Omega), \quad u \geq 0,$$

and it is defined as

$$(2.1) \quad J_\gamma(u, \Omega) := \int_{\Omega} |\nabla u|^2 + W_\gamma(u) \, dx,$$

where

$$(2.2) \quad W_\gamma(u) := c_\gamma u^{-\gamma} \chi_{\{u>0\}}, \quad \text{with } c_\gamma := \frac{1}{16} \cdot (2 - \gamma)^2, \quad \gamma \in (0, 2).$$

We study uniform properties of the minimizers of  $J_\gamma$  as  $\gamma \rightarrow 2^-$ . We often drop the dependence on  $\gamma$  from  $J$  and  $W$  when there is no possibility of confusion.

Notice that  $u$  is a minimizer of  $\mathcal{E}_\gamma$  defined in (1.2), if and only if  $c(\gamma)u$  is a minimizer of  $J_\gamma$ , with  $c(\gamma) = c_\gamma^{\frac{1}{\gamma+2}}$  an appropriate constant depending only on  $\gamma$ , and  $c(\gamma) \rightarrow 0$  as  $\gamma \rightarrow 2$ .

The constant  $c_\gamma$  in (2.2) is chosen such that

$$(2.3) \quad \int_0^1 2\sqrt{W_\gamma(s)} \, ds = 1.$$

The homogenous 1D solution  $\varphi$  plays an important role in the analysis. It is given by

$$(2.4) \quad \varphi(t) := c_\gamma^* (t^+)^{\alpha},$$

with

$$\alpha := \frac{2}{2 + \gamma}, \quad c_\gamma^* := \left( \left(1 + \frac{\gamma}{2}\right)^2 c_\gamma \right)^{\frac{1}{\gamma+2}},$$

and satisfies

$$(2.5) \quad \varphi' = (W_\gamma(\varphi))^{1/2}, \quad \text{in } \{\varphi > 0\}.$$

We differentiate the last equality and obtain that  $\varphi$  solves the Euler-Lagrange equation

$$(2.6) \quad 2\varphi'' = W'_\gamma(\varphi) \quad \text{in } \{\varphi > 0\}.$$

*Positive constants depending only on the dimension  $n$  are denoted by  $c$ ,  $C$ , and referred to as universal constants.*

The first result is an optimal uniform growth estimate.

**Theorem 2.1.** *Let  $u$  be a minimizer of  $J_\gamma$  in  $B_1$  and assume  $u(0) = 0$ . Then, there exists a universal constant  $C$  such that*

$$u(x) \leq C|x|^\alpha, \quad \alpha := \frac{2}{2 + \gamma}, \quad \forall x \in B_{1/2}.$$

The second theorem gives the uniform density estimate of the free boundary.

**Theorem 2.2** (Density estimates). *There exists a universal constant  $c_0$  such that if  $u$  is a nonnegative minimizer of  $J_\gamma$  in  $B_1$  and  $0 \in F(u)$  then*

$$1 - c_0 \geq \frac{|\{u > 0\} \cap B_r|}{|B_r|} \geq c_0, \quad \forall r \leq \frac{1}{2}.$$

The following result is a direct consequence of Theorems 2.1 and 2.2.

**Corollary 2.3.** *Let  $u$  be a nonnegative minimizer of  $J_\gamma$  in  $B_1$ . If  $0 \in F(u)$  then for all  $r \in (0, 1/2)$  each of the sets  $\{u = 0\} \cap B_r$  and  $\{u > 0\} \cap B_r$  contains an interior ball of radius  $cr$ . Moreover*

$$cr^{n-\alpha\gamma} \leq J(u, B_r) \leq Cr^{n-\alpha\gamma}.$$

Next we introduce the Dirichlet-perimeter functional  $\mathcal{F}$  introduced by Athanassopoulous, Caffarelli, Kenig, Salsa in [ACKS]. It acts on the space of admissible pairs  $(u, E)$  consisting of functions  $u \geq 0$  and measurable sets  $E \subset \Omega$  which have the property that  $u = 0$  a.e. on  $E$ ,

$$\mathcal{A}(\Omega) := \{(u, E) \mid u \in H^1(\Omega), \quad E \text{ Caccioppoli set, } u \geq 0 \text{ in } \Omega, u = 0 \text{ a.e. in } E\}.$$

The functional  $\mathcal{F}$  is given by the Dirichlet - perimeter energy

$$\mathcal{F}_\Omega(u, E) = \int_\Omega |\nabla u|^2 dx + P_\Omega(E),$$

where  $P_\Omega(E)$  represents the perimeter of  $E$  in  $\Omega$

$$\begin{aligned} P_\Omega(E) &= \int_\Omega |\nabla \chi_E| \\ &= \sup \int_\Omega \chi_E \operatorname{div} g \, dx \quad \text{with } g \in C_0^\infty(\Omega), \quad |g| \leq 1. \end{aligned}$$

The next theorem establishes the  $\Gamma$ -convergence of the  $J_\gamma$ 's.

**Theorem 2.4.** *As  $\gamma \rightarrow 2$ , the functionals  $J_\gamma$   $\Gamma$ -converge to  $\mathcal{F}$ .*

*More precisely we have:*

a) *(lower semicontinuity) if  $\gamma_k \rightarrow 2$  and  $u_k$  satisfy*

$$u_k^{1-\gamma_k/2} \rightarrow \chi_{E^c} \quad \text{in } L^1(\Omega), \quad u_k \rightarrow u \quad \text{in } L^2(\Omega),$$

*then*

$$\liminf J_{\gamma_k}(u_k, \Omega) \geq \mathcal{F}_\Omega(u, E).$$

b) *(approximation) given  $(u, E) \in \mathcal{A}(\Omega)$  with  $u$  a continuous in  $\overline{\Omega}$ , there exists  $\gamma_k \rightarrow 2$  and  $u_k$  such that*

$$\begin{aligned} u_k^{1-\gamma_k/2} &\rightarrow \chi_{E^c} \quad \text{in } L^1(\Omega), \quad u_k \rightarrow u \quad \text{in } L^2(\Omega), \\ J_{\gamma_k}(u_k, \Omega) &\rightarrow \mathcal{F}_\Omega(u, E). \end{aligned}$$

Our main result gives the strong convergence of the minimizers of  $J_\gamma$  and their zero set to the minimizing pairs  $(u, E)$  of  $\mathcal{F}$ .

**Theorem 2.5.** *Let  $\Omega$  be a bounded domain with Lipschitz boundary,  $\gamma_k \rightarrow 2^-$ , and  $u_k$  a sequence of functions with uniform bounded energies*

$$\|u_k\|_{L^2(\Omega)} + J_{\gamma_k}(u_k, \Omega) \leq M,$$

*for some  $M > 0$ . Then, after passing to a subsequence, we can find  $(u, E) \in \mathcal{A}(\Omega)$  such that*

$$u_k^{1-\gamma_k/2} \rightarrow \chi_{E^c} \quad \text{in } L^1(\Omega), \quad u_k \rightarrow u \quad \text{in } L^2(\Omega),$$

and

$$\chi_{\{u_k > 0\}} \rightarrow \chi_{E^c} \quad \text{in } L^1(\Omega).$$

Moreover, if  $u_k$  are minimizers of  $J_{\gamma_k}$  then the limit  $(u, E)$  is a minimizer of  $\mathcal{F}$ . The convergence of  $u_k$  to  $u$  and respectively of the free boundaries  $\partial\{u_k > 0\}$  to  $\partial E$  is uniform on compact sets (in the Hausdorff distance sense).

As a consequence we obtain the connection between bounded minimizers of  $\mathcal{E}_\gamma$  with  $\gamma \rightarrow 2$  and minimal surfaces, as stated in the Introduction. The uniform boundedness of minimizers can be deduced for example from a uniform bound of the boundary data on  $\partial\Omega$ .

**Corollary 2.6.** *Assume that  $u_k$  are uniformly bounded minimizers of  $\mathcal{E}_{\gamma_k}$  defined in (1.2), and  $\gamma_k \rightarrow 2$ . Then, up to subsequences,  $F(u_k)$  converge uniformly on compact sets to a minimal surface  $\partial E$ .*

Indeed,  $c(\gamma_k)u_k$  is a minimizer for  $J_{\gamma_k}$  and, since  $c(\gamma_k) \rightarrow 0$ , the limiting function  $u$  of Theorem 2.5 is identically 0. This means that the limiting set  $E$  must be a set of minimal perimeter in  $\Omega$ .

The paper is organized as follows. In Section 3 we prove the uniform growth estimate Theorem 2.1 and in Section 4 we obtain the uniform density estimates. In the last section we prove the main result Theorem 2.5.

### 3. PROOF OF THEOREM 2.1

In this section we prove Theorem 2.1. We state it here again for the reader convenience. We remark that this statement was proved in [DS] with a constant  $C$  depending on  $\gamma$ . The purpose of this section is to show that in fact the statement holds with a universal constant  $C$ . In the proof, we use that minimizers are viscosity solution in the sense of Definition 4.1 of [DS], as showed in Proposition 4.4 of [DS].

**Theorem 3.1.** *Let  $u$  be a minimizer of  $J_\gamma$  in  $B_1$ , and assume  $u(0) = 0$ . Then*

$$u(x) \leq C|x|^\alpha, \quad \forall x \in B_{1/2},$$

with  $C$  universal.

*Proof.* Minimizers of  $J$  are invariant under  $\alpha$ -homogenous rescalings

$$\tilde{u}(x) = \frac{u(y_0 + \lambda x)}{\lambda^\alpha}.$$

After such a rescaling, we may assume that we are in the situation  $B_1 \subset \{u > 0\}$  and  $u$  vanishes at some point  $x_0 \in \partial B_1$ . We need to prove that  $u(0)$  is bounded above by a large universal constant.

Notice that in  $B_1$  we satisfy

$$\Delta u \leq 0, \quad \Delta(u-1)^+ \geq -1.$$

Thus, if

$$u(0) \geq M \gg 1,$$

then by the weak Harnack inequality we find

$$u \geq cM \quad \text{in } B_{1/2}, \quad \text{with } c > 0 \text{ universal.}$$

**Lemma 3.2.** *There exists a one dimensional increasing function  $\psi$ ,*

$$\psi : [0, t_0] \rightarrow \mathbb{R}, \quad \psi(0) = 0, \quad t_0 \leq \frac{1}{4},$$

such that (see (2.4) for the definition of  $\varphi$ )

1)

$$\psi(t) = \varphi(t) + \epsilon t^{2-\alpha} + O(t^{2-\alpha+\delta}) \quad \text{near } 0, \quad \delta > 0,$$

2)

$$2\psi'' \geq 4n\psi' + W'(\psi),$$

3)

$$\psi(t_0) \leq 1, \quad \psi'(t_0) \leq C_0 \quad \text{universal.}$$

Using Lemma 3.2 we construct a barrier  $\Psi : B_1 \setminus B_{1/2} \rightarrow \mathbb{R}$ , as

$$\Psi(x) = \psi(1 - |x|) \quad \text{in } B_1 \setminus B_{1-t_0}$$

and

$$\Delta \Psi = 0 \quad \text{in } B_{1-t_0} \setminus B_{1/2},$$

with boundary conditions

$$\Psi = cM \quad \text{on } \partial B_{1/2}, \quad \Psi = \psi(t_0) \quad \text{on } \partial B_{1-t_0}.$$

Since  $\psi(t_0) \leq 1$ , it follows that

$$|\nabla \Psi| > C_0 \quad \text{in the annulus } B_{1-t_0} \setminus B_{1/2},$$

provided that  $M$  is large universal.

We claim that

$$(3.1) \quad 2\Delta \Psi \geq W'(\Psi), \quad \text{in } B_1 \setminus B_{1/2}.$$

The inequality is satisfied in the outer annulus  $B_1 \setminus B_{1-t_0}$  by property 2) above, and in the inner annulus  $B_{1-t_0} \setminus B_{1/2}$  since  $0 > W'$ .

Moreover, the inequalities between the normal derivatives on either side of  $\partial B_{1-t_0}$  guarantee that (3.1) holds in the whole domain.

Since  $W'(t)$  is increasing for  $t > 0$ , we can apply the maximum principle and conclude that

$$u \geq \Psi \quad \text{in } B_1 \setminus B_{1/2}.$$

We contradict the free boundary condition at the point  $x_0 \in F(u)$  for a minimizer, see Proposition 4.4 in [DS]. Indeed, property 1) above shows that  $\Psi - \varphi(d)$  has a positive correction term  $\epsilon d^{2-\alpha}$  in the expansion near its free boundary and therefore it is a strict viscosity subsolution on  $\partial B_1$ , see Definition 4.1 in [DS].  $\square$

It remains to prove the lemma above.

*Proof of Lemma 3.2:* We reduce the second order ODE to a 1st order ODE by taking  $\psi$  as an independent variable. More precisely, with a strictly increasing function  $\psi$  we associate the function  $g > 0$  defined on the range of  $\psi$  as

$$(3.2) \quad g(\psi) := (\psi')^2.$$

After differentiation we obtain

$$2\psi'' = g'(\psi).$$

The function  $\psi$  can be recovered from  $g$  by the formula

$$(3.3) \quad \psi(t) = G^{-1}(t), \quad G(r) := \int_0^r \frac{1}{\sqrt{g(s)}} ds.$$

In the case when  $\psi$  coincides with the 1D solution  $\varphi$  given in (2.4), then the associated function  $g$  equals  $W$ , see (2.5).

In our setting we define  $g$  explicitly as

$$g(s) := W(s) + \bar{\varepsilon} + C_1 s^{1-\frac{\gamma}{2}}, \quad s \in (0, s_0],$$

with  $C_1 = 8n$  universal, and  $s_0$  given by the solution to

$$C_1 s^{1-\frac{\gamma}{2}} = c_\gamma s^{-\gamma} = W(s) \quad \text{when } s = s_0,$$

and  $\bar{\varepsilon} > 0$  arbitrarily small. Notice that  $s_0 \rightarrow 0$  as  $\gamma \rightarrow 2^-$ , and from the formula for  $c_\gamma$  in (2.3) it follows

$$(3.4) \quad s_0 \sim 2 - \gamma.$$

Notice that

$$g(s_0) \leq 3C_1 s_0^{1-\frac{\gamma}{2}} \leq 3C_1 =: C_0^2.$$

This gives property 3) since

$$\psi(t_0) = s_0, \quad \psi'(t_0) = (g(s_0))^{1/2}.$$

By construction  $g \geq W$  which by (3.3) implies  $\psi \geq \varphi$ . Thus

$$s_0 = \psi(t_0) \geq \varphi(t_0),$$

and by (2.4), (3.4), it follows that also  $t_0 \rightarrow 0$  as  $\gamma \rightarrow 2^-$ .

We compute

$$g' = W' + C_1 \left(1 - \frac{\gamma}{2}\right) s^{-\frac{\gamma}{2}}$$

and use the inequality

$$C_1 \left(1 - \frac{\gamma}{2}\right) \geq 8n\sqrt{c_\gamma},$$

and that  $g \leq 3W$  in the interval  $[0, s_0]$  to obtain

$$g' \geq W' + 4n(3W)^{1/2} \geq W' + 4ng^{1/2},$$

which gives 2).

Finally, we obtain property 1) from (3.3) and the expansion

$$\frac{1}{\sqrt{g(s)}} = \frac{1}{\sqrt{W(s)}} \left(1 - c(\gamma)\bar{\varepsilon}s^\gamma + O(s^{1+\frac{\gamma}{2}})\right).$$

□

#### 4. DENSITY ESTIMATES

In this technical section we prove Theorem 2.2 and Corollary 2.3. We follow the classical ideas from the minimal surface theory by constructing appropriate competitors for the minimizer  $u$ , and then make use of the isoperimetric inequality. They allow us to obtain discrete differential inequalities for the measure of the sets  $\{u > 0\}$  in  $B_r$ , which give the desired conclusion after iteration.

We start with the lower bound.

**Lemma 4.1.** *Let  $u$  be a minimizer of  $J_\gamma$  in  $B_1$  and assume  $0 \in F(u)$ . Then*

$$|\{u > 0 \cap B_r\}| \geq c_0 |B_r|.$$

*Proof.* After a dilation, assume  $u$  minimizes  $J$  in  $B_3$ . Since  $0 \in F(u)$ ,

$$u \leq C_0 \quad \text{in } B_2,$$

by Theorem 2.1. Define

$$A_r := \{u > 0\} \cap B_r, \quad a(r) := |A_r|.$$

It suffices to show that

$$a(1) \leq c_0 \implies a(r) = 0 \quad \text{for all } r \text{ sufficiently small,}$$

which is not possible since  $0 \in F(u)$ . We consider the case when  $\gamma$  is close to 2.

Define  $s_0, t_0$ , as

$$(4.1) \quad W(s_0) = 1, \quad \varphi(t_0) = s_0 \implies \varphi'(t_0) = \sqrt{W(s_0)} = 1$$

and notice that  $s_0, t_0 \rightarrow 0$  as  $\gamma \rightarrow 2^-$ .

*Step 1:* We show that the densities of the sets  $A_r$  in  $B_r$  decay geometrically as we rescale by a factor of  $1 - 2t_0$ , i.e if  $a(1) \leq c_0$  then

$$(4.2) \quad r_0^{-n} a(r_0) \leq r_0 a(1) \quad \text{with } r_0 := 1 - 2t_0.$$

First we construct a 1D function.

**Lemma 4.2.** *There exists a piecewise  $C^1$  function  $\psi$  in  $[0, 1]$  such that*

1)

$$\psi(t) = \varphi(t) \quad \text{if } t \leq t_0,$$

2)

$$2\psi'' + 4n\psi' \leq W'(\psi) \quad \text{if } t \geq t_0,$$

3)

$$\psi(1) \geq 2C_0, \quad \psi'(t_0) \leq C_1 \quad \text{for some } C_1 \text{ large universal.}$$

*Proof.* Indeed, we may take

$$\psi := \varphi + Kg(t) \chi_{\{t \geq t_0\}}$$

with  $g$  an increasing  $C^2$  function in  $[t_0, 1]$  such that

$$g(t_0) = 0, \quad g'' + 2ng' \leq -c \text{ in } [t_0, 1],$$

and  $K$  a sufficiently large universal constant. Properties 1), 3) follow immediately from the definition of  $\psi$ . For 2) we use that in  $[t_0, 1]$

$$2\varphi'' = W'(\varphi), \quad \varphi' \leq \varphi'(t_0) = 1,$$

hence

$$\begin{aligned} 2\psi'' + 4n\psi' &\leq 2\varphi'' + 4n\varphi' - 2Kc \\ &\leq W'(\varphi) + 4n - 2Kc \\ &\leq W'(\varphi) \\ &\leq W'(\psi), \end{aligned}$$

where in the last inequality we used that  $W'$  is an increasing function. □



*Proof of Step 1:* We use Lemma 4.2 to define

$$\Psi(x) := \psi(|x| - (1 - t_0)),$$

and let

$$D := \{u > \Psi\} \subset B_{2-t_0} \cap \{u > 0\}.$$

Notice that  $u = \Psi$  on  $\partial D$ , hence the minimality of  $J$  implies

$$(4.3) \quad J(u, D) \leq J(\Psi, D).$$

We decompose  $D$  as the disjoint union

$$D = D_1 \cup D_2, \quad D_1 := D \cap B_1, \quad D_2 := D \setminus B_1,$$

and notice that

$$(4.4) \quad \begin{aligned} J(\Psi, D_2) - J(u, D_2) &= \\ &= \int_{D_2} -2\nabla(u - \Psi) \cdot \nabla\Psi - |\nabla(u - \Psi)|^2 + W(\Psi) - W(u) dx \\ &\leq \int_{D_2} (u - \Psi)2\Delta\Psi + W(\Psi) - W(u) dx + \int_{\partial D_2} 2(u - \Psi)|\nabla\Psi| d\sigma \\ &\leq \int_{D_2} (u - \Psi)W'(\Psi) + W(\Psi) - W(u) dx + \int_{\partial D_2 \cap \partial B_1} 2(u - \Psi)|\nabla\Psi| d\sigma \\ &\leq C\mathcal{H}^{n-1}(\{u > 0\} \cap \partial B_1), \end{aligned}$$

where we have used that

$$0 \leq u - \Psi \leq C, \quad 2\Delta\Psi \leq W'(\Psi), \quad |\nabla\Psi| \leq C \quad \text{on } \partial B_1,$$

and that  $W$  is convex on its positivity set.

Combining (4.3) and (4.4) we find

$$(4.5) \quad J(u, D_1) \leq J(\Psi, D_1) + C\mathcal{H}^{n-1}(\{u > 0\} \cap \partial B_1).$$

In  $D_1$  we use the Cauchy-Schwartz inequality and the coarea formula to obtain

$$(4.6) \quad \begin{aligned} J(u, D_1) &\geq \int_{D_1 \cap \{u < \varphi(t_0)\}} 2|\nabla u| \sqrt{W(u)} dx \\ &= \int_0^{s_0} \mathcal{H}^{n-1}(\{u = s\} \cap D_1) 2\sqrt{W(s)} ds. \end{aligned}$$

On the other hand  $|\nabla\Psi| = \sqrt{W'(\Psi)}$  in  $D_1$  by construction (see 1) in Lemma 4.2 and (2.5)) and the inequality above becomes an equality for  $\Psi$ :

$$(4.7) \quad J(\Psi, D_1) = \int_0^{s_0} \mathcal{H}^{n-1}(\{\Psi = s\} \cap D_1) 2\sqrt{W(s)} ds.$$

Next we use that

$$\{u > s\} \cap B_{1-2t_0} \subset D_1 \cap \{u > s > \Psi\}, \quad s > 0,$$

and the isoperimetric inequality implies

$$c_n |\{u > s\} \cap B_{1-2t_0}|^{\frac{n-1}{n}} \leq \mathcal{H}^{n-1}(\{u = s\} \cap D_1) + \mathcal{H}^{n-1}(\{\Psi = s\} \cap D_1),$$

hence

$$(4.8) \quad \int_0^{s_0} c_n |\{u > s\} \cap B_{1-2t_0}|^{\frac{n-1}{n}} 2\sqrt{W(s)} ds \leq J(u, D_1) + J(\Psi, D_1).$$

We combine this with (4.5), (4.7) and use that

$$(4.9) \quad \int_{B_{1-t_0}} W(u) dx \leq J(u, D_1),$$

and obtain

$$(4.10) \quad \begin{aligned} \int_{B_{1-2t_0}} W(u) dx + \int_0^{s_0} c_n |\{u > s\} \cap B_{1-2t_0}|^{\frac{n-1}{n}} 2\sqrt{W(s)} ds &\leq \\ &\leq C \mathcal{H}^{n-1}(\{\Psi = s_0\} \cap \{u > 0\}) + \\ &+ C \int_0^{s_0} \mathcal{H}^{n-1}(\{\Psi = s\} \cap \{u > 0\}) 2\sqrt{W(s)} ds. \end{aligned}$$

The inequality holds also when we replace  $\Psi$  by  $\Psi_t$  defined as

$$\Psi_s(x) := \psi(|x| - (1 - t_0 - t)), \quad t \in [0, t_0].$$

Notice that  $\Psi_0 = \Psi$  and  $\{\Psi_t = s\}$  is the sphere at distance  $t$  from the sphere  $\{\Psi = s\}$ . Thus, if we write the inequality above for  $t \in [0, t_0]$  and average it over this interval we obtain

$$(4.11) \quad \begin{aligned} \int_{B_{1-2t_0}} W(u) dx + \int_0^{s_0} c_n |\{u > s\} \cap B_{1-2t_0}|^{\frac{n-1}{n}} 2\sqrt{W(s)} ds &\leq \\ &\leq C t_0^{-1} |\{u > 0\} \cap (B_1 \setminus B_{1-2t_0})| \int_0^{s_0} 2\sqrt{W(s)} ds. \end{aligned}$$

Let  $s_1 \in [0, s_0]$  and denote by

$$b := |\{0 < u \leq s_1\} \cap B_{1-2t_0}|,$$

hence if  $s \leq s_1$  then

$$|\{u > s\} \cap B_{1-2t_0}| \leq |\{u > s_1\} \cap B_{1-2t_0}| = a(1 - 2t_0) - b.$$

Notice that by the choice of  $c_\gamma$  we have

$$\int_0^{s_1} 2\sqrt{W(s)} ds = s_1^{1-\frac{\gamma}{2}}.$$

Since  $W(u) \geq W(s_1)$  in the set  $\{0 < u \leq s_1\}$ , we can bound below the left hand side in (4.11) by

$$(4.12) \quad W(s_1)b + c_1 s_1^{1-\frac{\gamma}{2}} (a(1 - 2t_0) - b)^{\frac{n-1}{n}},$$

while the right hand side in (4.11) is bounded above by

$$C_2 \frac{a(1) - a(1 - 2t_0)}{2t_0},$$

with  $C_2, c_1$  universal constants.

We choose  $s_1$  such that

$$W(s_1) = C_3 \gg C_2, \quad \text{i.e.} \quad s_1 = C_3^{-\frac{1}{\gamma}} s_0 = (c_\gamma / C_3)^{\frac{1}{\gamma}}.$$

Using that  $c_\gamma \sim (2 - \gamma)^2$  we find that the coefficient

$$c_1 s_1^{1-\frac{\gamma}{2}}$$

which appears in (4.12) remains bounded below as  $\gamma \rightarrow 2^-$ . This means that if  $a(1-2t_0) \leq a(1) \leq c_0$  small, universal, then the expression in (4.12) is decreasing in the variable  $b \in [0, a(1-2t_0)]$  and is bounded below by  $C_3 a(1-2t_0)$ . In conclusion

$$C_3 a(1-2t_0) \leq C_2 \frac{a(1) - a(1-2t_0)}{2t_0}$$

which gives (4.2):

$$a(1-2t_0)(1-2t_0)^{-(n+1)} \leq a(1),$$

and Step 1 is proved.  $\square$

As we iterate Step 1 we find that the densities of the positivity set in  $B_r$ ,  $a(r)r^{-n}$ , tend to 0 as  $r = r_0^m \rightarrow 0$ . After rescaling, it remains to show that if  $a(1)$  is sufficiently small, depending on  $\gamma$ , then  $a(1/2) = 0$ .

*Step 2:* If  $a(1) \leq c(\gamma)$  small then for all  $r \in [1/2, 1]$ ,

$$(4.13) \quad a(r-2t)^\delta \leq \frac{a(r) - a(r-2t)}{2t}, \quad t = a(r)^\mu,$$

with  $\delta, \mu$  universal constants.

*Proof of Step 2:* Assume for simplicity that  $r = 1$ . Notice that by Theorem 2.1 it follows that

$$u \leq Ca(1)^{\frac{\alpha}{n}} \quad \text{in } B_1.$$

We argue as in Step 1 and improve the last part of the argument. Take

$$\Psi = \psi(|x| - (1-t_1))$$

with  $t_1 \in (0, t_0]$  such that

$$\varphi(t_1) = a(1)^\mu \gg \|u\|_{L^\infty(B_1)}.$$

This means that  $\{u < \Psi\}$  on  $\partial B_1$  and now we may take  $D = \{u > \Psi\} \cap B_1$ . We obtain as above the corresponding inequality (4.10) with  $t_0$  replaced by  $t_1$ . After averaging over the family of translates  $\Psi_t$  with  $t \in [0, t_1]$  we establish the inequality (4.11) with  $t_0$  replaced by  $t_1$ . We bound the left hand side as before by taking

$$s_1 = \varphi(t_1) = a(1)^\mu,$$

and obtain

$$W(s_1)s_1^{\frac{\gamma}{2}-1}b + (a(1-2t_1) - b)^{\frac{n-1}{n}} \leq C \frac{a(1) - a(1-2t_1)}{2t_1}.$$

Using that

$$s_1 = a(1)^\mu \geq a(1-2t_1)^\mu,$$

the coefficient of  $b$  in the left hand side is bounded below by a negative power of  $a(1-2t_1)$  (provided that  $a(1)$  is sufficiently small, depending on  $\gamma$ ). Then, by arguing that

$$\text{either } b \leq \frac{a(1-2t_1)}{2} \quad \text{or } b \geq \frac{a(1-2t_1)}{2},$$

we obtain that the left hand side is bounded below by

$$a(1-2t_1)^{1-\delta},$$

for some  $\delta$  universal. After relabeling  $\delta$  if necessary we reach the desired discrete differential inequality claimed in Step 2.

$$a(1 - 2s_1)^{1-\delta} \leq \frac{a(1) - a(1 - 2s_1)}{2s_1}, \quad s_1 = a(1)^\mu.$$

□

*End of the proof:* Now it is straightforward to check that a nondecreasing function  $a(r)$  that satisfies (4.13) must vanish when  $r = 1/2$  if  $a(1)$  is sufficiently small. In the continuous setting we obtain  $a' \geq a^{1-\delta}$  which implies

$$a(r) \leq (r - 1/2)^M,$$

for some large  $M$ , provided that the inequality is satisfied at  $r = 1$ . In the discrete setting it follows by induction that the inequality above holds for  $r = r_k$  where  $r_k$  is the sequence

$$r_{k+1} = r_k - 2a(r_k)^\mu, \quad r_0 = 1.$$

□

*Remark 4.3.* From (4.5) and (4.7) it follows that

$$J(u, B_{1/2}) \leq J(u, D_1) \leq C,$$

with  $C$  universal.

Next we prove the other side of the density bound using a similar analysis.

**Lemma 4.4.** *Let  $u$  be a minimizer of  $J_\gamma$  in  $B_1$  and assume  $0 \in F(u)$ . Then*

$$|\{u = 0\} \cap B_r| \geq c_0 |B_r|.$$

*Proof.* Let  $s_0$ ,  $s_1$ , and  $t_1$  be defined as

$$W(s_0) = 1, \quad W(s_1) = M, \quad \varphi(t_1) = s_1,$$

with  $M$  a large universal constant to be made precise later. Let

$$A_r := \{u \leq s_1\} \cap B_r, \quad a(r) := |A_r|.$$

*Step 1:* We prove that if  $a(1) \leq c_0$  universal,  $M \geq C_0$  and  $\gamma$  sufficiently close to 2 (depending on  $M$ ) then

$$(4.14) \quad a(r_0)r_0^{-n} \leq r_0 a(1) \quad \text{for some fixed } r_0 < 1.$$

We first construct a 1D profile.

**Lemma 4.5.** *There exists a nondecreasing Lipschitz function  $\psi : [0, 1] \rightarrow \mathbb{R}$ , with  $\psi(0) = 0$ , which is  $C^1$  in the intervals  $\{\psi < s_1\}$ ,  $\{\psi > s_1\}$  such that*

$$1) \ \psi = \varphi \text{ in } [0, t_1] = \{\psi \leq s_1\},$$

$$2) \text{ in } (t_1, 1] = \{\psi > s_1\},$$

$$2\psi'' - 8n\psi' \geq W'(\psi)$$

and  $\psi$  is constant in  $[1/4, 1]$ ,

3)

$$\frac{1}{2}W(\psi) \leq (\psi')^2 \leq W(\psi) \quad \text{in } [0, t_0] := \{\psi \leq s_0\}.$$

Here  $t_0$  is defined such that

$$\psi(t_0) = s_0, \quad \text{thus } W(\psi(t_0)) = 1.$$

*Proof of Step 1:* Define in  $\overline{B_1}$  the function

$$\Psi(x) = \psi(1 - |x|),$$

and denote by

$$D := \{u < \Psi\}.$$

Notice that  $\Psi$  vanishes on  $\partial B_1$  and coincides with  $\varphi(1 - |x|)$  near  $\partial B_1$ , hence

$$(4.15) \quad |\nabla \Psi| = \sqrt{W(\Psi)} \quad \text{in } B_1 \setminus B_{1-t_1} = \{\Psi \leq s_1\}.$$

Also by 2)

$$2\Delta \Psi \geq W'(\Psi) \quad \text{in } \{\Psi > s_1\},$$

and 3) implies

$$(4.16) \quad \frac{1}{2}W(\Psi) \leq |\nabla \Psi|^2 \leq W(\Psi) \quad \text{in } B_1 \setminus B_{1-t_0},$$

and

$$(4.17) \quad W(\Psi) \leq 1 \quad \text{in } B_{1-t_0}.$$

Denote by

$$\begin{aligned} D_1 &:= \{u > s_1\} \cap D, & D_2 &:= D \setminus D_1, \\ F_1 &:= \{\Psi > s_1\} \cap D, & F_2 &:= D \setminus F_1. \end{aligned}$$

Then  $J(u, D) \leq J(\Psi, D)$  implies

$$J(u, D_2) \leq J(\Psi, F_2) + J(\Psi, F_1) - J(u, D_1).$$

In

$$F_1 = D_1 \cup A_{1-t_1}$$

we write

$$\max\{u, \sigma\} = \Psi - w, \quad \text{with } \Psi \geq w \geq 0,$$

and notice that  $w$  vanishes on  $\partial F_1$  hence

$$(4.18) \quad \begin{aligned} \int_{D_1} |\nabla u|^2 &= \int_{F_1} |\nabla(\Psi - w)|^2 dx \\ &\geq \int_{F_1} |\nabla \Psi|^2 + 2w\Delta \Psi dx \\ &\geq \int_{F_1} |\nabla \Psi|^2 + wW'(\Psi) dx \\ &\geq \int_{F_1} |\nabla \Psi|^2 + (W(\Psi) - W(\Psi - w))\chi_{D_1} - CW(\Psi)\chi_{A_{1-t_1}} dx, \end{aligned}$$

where in the last inequality we used the convexity of  $W$  in  $D_1$  and the fact that  $W'(\Psi) < 0$  in  $A_{1-t_1}$  thus

$$wW'(\Psi) \geq \Psi W'(\Psi) = -\gamma W(\Psi).$$

Since  $\Psi - w = u$  in  $D_1$  we find

$$J(u, D_1) \geq J(\Psi, F_1) - C \int_{A_{1-t_1}} W(\Psi) dx,$$

hence

$$J(u, D_2) \leq J(\Psi, F_2) + C \int_{A_1} W(\Psi) dx.$$

By Cauchy-Schwartz and co-area formula we obtain

$$J(u, D_2) \geq \int_0^{s_1} \mathcal{H}^{n-1}(\{u = s\} \cap D) \sqrt{W(s)} ds,$$

while, by (4.15),

$$J(\Psi, F_2) = \int_0^{s_1} \mathcal{H}^{n-1}(\{\Psi = s\} \cap D) \sqrt{W(s)} ds.$$

Hence

$$J(\Psi, F_2) \leq \int_0^{s_1} \mathcal{H}^{n-1}(\{\Psi = s\} \cap A_1) \sqrt{W(s)} ds,$$

and we also write

$$\int_{A_1} W(\Psi) dx = \int_{A_1 \cap B_{1-t_0}} W(\Psi) dx + \int_{A_1 \setminus B_{1-t_0}} W(\Psi) dx.$$

By (4.17) the second term is bounded by  $|A_1|$ , while by (4.16) and the co-area formula as above, the first integral is bounded by

$$C \int_0^{s_0} \mathcal{H}^{n-1}(\{\Psi = s\} \cap A_1) \sqrt{W(s)} ds.$$

Using that

$$E := \{u = 0\} \cap B_{1-t_1} \subset \{u \leq s \leq \Psi\}, \quad s \in [0, s_1],$$

we find by the isoperimetric inequality that

$$|E|^{\frac{n-1}{n}} \int_0^{s_1} \sqrt{W(s)} ds \leq J(u, D_2) + J(\Psi, F_2).$$

Notice that as  $\gamma \rightarrow 2$  (and fixed  $M$ ), the integral converges to

$$\int_0^1 \sqrt{W(s)} ds = \frac{1}{2}.$$

Also

$$W(s_1)|A_{1-t_1} \setminus E| \leq \int_{A_{1-t_1}} W(u) dx \leq J(u, D_2)$$

In conclusion

$$\begin{aligned} & \frac{1}{4}|E|^{\frac{n-1}{n}} + M|A_{1-t_1} \setminus E| \leq \\ (4.19) \quad & \leq C \int_0^{s_0} \mathcal{H}^{n-1}(\{\Psi = s\} \cap A_1) \sqrt{W(s)} ds + C|A_1|. \end{aligned}$$

Since  $|E| \leq a(1) \leq c_0$  is sufficiently small, and  $M \geq C_0$ , the left hand side is bounded below by

$$\frac{C_0}{2}|A_{1-t_1}| \geq \frac{C_0}{2}a(1-2t_0).$$

We average the right hand side by taking as test functions

$$\Psi_t(x) = \psi(1-t-|x|), \quad t \in [0, t_0],$$

and obtain

$$\frac{C_0}{2}a(1-2t_0) \leq C \frac{a(1) - a(1-2t_0)}{2t_0} + Ca(1)$$

which implies the desired conclusion (4.14) with  $r_0 = 1 - 2t_0$ ,

$$a(1-2t_0)(1-2t_0)^{-(n+1)} \leq a(1),$$

provided  $C_0$  is chosen sufficiently large. □

Next we prove the lemma when  $\gamma$  is close to 2.

*Step 2:* If  $\gamma$  is sufficiently close to 2 then  $|\{u = 0\} \cap B_1| \geq c_0/2$ .

*Proof of Step 2:* If the conclusion does not hold then

$$(4.20) \quad |\{u = 0\} \cap B_1| \leq c_0/2 \implies a(1) \leq c_0.$$

Indeed, otherwise

$$|\{0 < u \leq s_1\} \cap B_1| \geq c_0/2,$$

and we can apply inequality (4.19) (with  $A_{1-t_1}$ ,  $A_1$  replaced by  $A_1$ , respectively  $A_{1+t_1}$ ) and obtain

$$M \frac{c_0}{2} \leq C \int_0^{s_0} \mathcal{H}^{n-1}(\{\Psi = s\} \cap A_{1+t_1}) \sqrt{W(s)} ds + C|A_{1+t_1}| \leq C.$$

We get a contradiction by choosing  $M$  universal, sufficiently large, and (4.20) is proved. Now we may apply Step 1 and obtain

$$|\{u \leq r_0^\alpha s_1\} \cap B_{r_0}| r_0^{-n} \leq a(r_0) r_0^{-n} \leq r_0 a(1),$$

which can be rescaled and iterated indefinitely. Thus, after a rescaling of  $u$  of factor  $r_0^m$  with  $m$  large we find that  $a(1)$  can be made arbitrarily small.

We reached a contradiction to  $0 \in F(u)$  since, by Theorem 2.1,

$$a(1) \geq c(s_1) > 0. \quad \square$$

Finally, we prove the conclusion also when  $\gamma$  stays away from 2.

*Step 3:* If  $\gamma \leq 2 - \delta$  then  $|\{u > 0\} \cap B_1| \geq c(\delta)$ .

*Proof of Step 3:* This follows easily by compactness. However, here we sketch a direct proof that follows from an argument in Step 1.

First we claim that

$$\max_{\partial B_1} u \geq c(\delta),$$

for some  $c(\delta) > 0$  small. Otherwise, the energy of  $u$  in  $B_{1/2}$  is sufficiently small, which implies that  $\{u > 0\}$  has small measure in  $B_{1/2}$  and contradicts Lemma 4.1.

Next, let  $v$  be the solution to the Euler-Lagrange equation  $2\Delta\Psi = W'(\Psi)$  in  $B_1$ ,  $v = u$  on  $\partial B_1$ . Since  $v$  is superharmonic,  $v(0) > c(\delta)$ . Moreover,  $W(v)$  is bounded by an integrable function in  $B_1$ . As in Step 1, the inequality

$$J(u, B_1) \leq J(v, B_1)$$

implies (see (4.18) with  $s_1 = 0$ ,  $D_1 = F_1 = B_1$ ),

$$\int_{B_1} |\nabla(v-u)|^2 dx \leq C \int_{\{u=0\}} W(v) dx.$$

The left hand side is bounded below by a  $c_1(\delta)$  which follows from Theorem 2.1 and  $(v-u)(0) = v(0) \geq c(\delta)$ . This shows that  $\{u=0\}$  cannot have arbitrarily small measure.  $\square$

It remains to prove the existence of the 1D profile of Lemma 4.5.

*Proof of Lemma 4.5:* We construct  $\psi$  by defining its corresponding function  $g$  as in (3.2), (3.3). Let  $g$  be the perturbation of  $W$

$$g(s) = W(s) + \left( -\frac{1}{2} + C_n(s^{1-\frac{\gamma}{2}} - s_1^{1-\frac{\gamma}{2}}) \right) \chi_{[s_1, 1]},$$

with  $C_n = 8n$ . Let  $s_2$  be defined as

$$W(s_2) = \frac{1}{4},$$

hence  $s_2 = 4^{1/\gamma} s_0 \sim s_0$ , and notice that  $s_2 \rightarrow 0$  as  $\gamma \rightarrow 2$ . Moreover

$$W' = -\gamma W/s \leq -C \quad \text{in } [0, s_2]$$

which implies that  $g' \leq -C$  in the same interval. Furthermore, for  $\gamma$  sufficiently close to 2 (depending on  $M$ ), then  $s_1^{1-\gamma/2}$  is close to 1 hence the error  $g(s) - W(s)$  is uniformly close to the constant  $-1/2$  in the interval  $[s_1, 1]$ .

These facts imply that  $g \leq W$ , and  $g$  crosses 0 at some point  $\sigma \in [s_0, s_2]$ , and

$$g \geq \frac{1}{2}W \quad \text{in } [s_1, s_0],$$

which gives property 3). Property 1) follows directly from the definition. Finally, property 2) holds since in  $(s_1, 1] \cap \{g > 0\}$

$$g' - W' = C_n(1 - \frac{\gamma}{2})s^{-\gamma/2} \geq 8n\sqrt{W} \geq 8n\sqrt{g}.$$

Moreover,

$$\begin{aligned} \int_{\{g>0\}} (2g)^{-1/2} ds &= \int_0^\sigma (2g)^{-1/2} ds \\ &\leq \int_0^{s_0} W^{-1/2} ds + C \int_{s_0}^\sigma s_0^{1/2} (\sigma - s)^{-1/2} ds \\ &\leq 2^{1/2} t_0 + C s_0 \\ &\leq 1/4 \end{aligned}$$

which shows that  $\psi$  is constant outside an interval of length  $1/4$ .  $\square$

We conclude this section with a proof of Corollary 2.3.

*Proof of Corollary 2.3.* Assume that  $u$  is a minimizer of  $J$  in  $B_2$  and  $0 \in F(u)$ . First we prove that

$$(4.21) \quad c \leq J(u, B_1) \leq C,$$

with  $c, C$  universal constants.



The upper bound follows from Remark 4.3. For the lower bound, we use that

$$(1 - c_0)|B_1| \geq |\{u > 0\} \cap B_1| \geq c_0|B_1|.$$

Let  $s_0$  be defined as in the proof of Lemma 4.1, see (4.1). If

$$(4.22) \quad |\{u > s_0\} \cap B_1| \leq \frac{c_0}{2}|B_1|,$$

then

$$|\{0 < u \leq s_0\} \cap B_1| \geq \frac{c_0}{2}|B_1|.$$

In this last set  $W(u) \geq W(s_0) = 1$ , and the lower bound is obtained from the potential term.

On the other hand, if the opposite inequality in (4.22) holds, then for all  $s \in (0, s_0)$  the density of  $\{u > s\}$  in  $B_1$  is bounded both above and below by universal constants. Now the lower bound follows from (4.6) and the Poincaré inequality for  $\chi_{\{u > s\}}$  in  $B_1$ .

The existence of a full ball of radius  $c'$  included in  $\{u > 0\} \cap B_1$  (or  $\{u = 0\} \cap B_1$ ) follows by a standard covering argument. We sketch it below.

We take a collection of  $m$  disjoint balls  $B_\rho(x_i)$ ,  $x_i \in \{u > 0\} \cap B_1$  such that  $\cup B_{5\rho}(x_i)$  covers  $\{u > 0\} \cap B_1$ . It follows that  $m \sim \rho^{-n}$ . If we assume that each  $B_{\rho/2}(x_i)$  intersects the free boundary then, by the rescaled version of (4.21),

$$J(u, B_\rho(x_i)) \geq c\rho^{n-\alpha\gamma}.$$

We obtain

$$J(u, B_1) \geq m c \rho^{n-\alpha\gamma},$$

and we contradict the upper bound if  $\rho$  is chosen small, universal.  $\square$

## 5. THE GAMMA CONVERGENCE

In this section we prove our main result Theorem 2.5. We start by constructing an interpolation between two functions which are close to each other in a ring.

**Proposition 5.1.** *Let  $u_k, v_k$  be sequences in  $H^1(B_1)$  and  $\gamma_k \rightarrow 2^-$ . Assume that for some  $\rho \in (\frac{1}{2}, 1)$  and  $\delta > 0$  small,*

$$J_{\gamma_k}(u_k, B_{\rho+\delta}), \quad J_{\gamma_k}(v_k, B_{\rho+\delta})$$

are uniformly bounded, and

$$\|u_k - v_k\|_{L^2} + \|u_k^{1-\frac{\gamma_k}{2}} - v_k^{1-\frac{\gamma_k}{2}}\|_{L^1} \rightarrow 0 \quad \text{in } B_{\rho+\delta} \setminus \bar{B}_\rho, \quad \text{as } k \rightarrow \infty.$$

Then, there exists  $w_k \in H^1(B_1)$  with

$$w_k := \begin{cases} v_k & \text{in } B_\rho \\ u_k & \text{in } B_1 \setminus \bar{B}_{\rho+\delta} \end{cases}$$

such that

$$J_{\gamma_k}(w_k, B_1) \leq J_{\gamma_k}(v_k, B_{\rho+\delta}) + J_{\gamma_k}(u_k, B_1 \setminus \bar{B}_\rho) + o(1),$$

with  $o(1) \rightarrow 0$  as  $k \rightarrow \infty$ .

*Proof.* Fix  $\epsilon > 0$  small. We prove the conclusion with  $o(1)$  replaced by  $C\epsilon$  for some  $C$  universal. Since the energies of  $u_k$  and  $v_k$  are uniformly bounded, we can decompose the annulus  $B_{\rho+\delta} \setminus B_\rho$  into a disjoint union of  $\sim \epsilon^{-1}$  annuli, and after relabeling  $\rho$  and  $\delta$  we may assume that

$$J_{\gamma_k}(u_k, B_{\rho+\delta} \setminus B_\rho) \leq \epsilon, \quad J_{\gamma_k}(v_k, B_{\rho+\delta} \setminus B_\rho) \leq \epsilon.$$

For simplicity of notation we drop the subindex  $k$ .

First we prove the result under the additional assumption

$$(5.1) \quad u \geq v \text{ in } B_{\rho+\delta} \setminus B_\rho.$$

Denote by

$$\psi_r(x) = \varphi(|x| - r), r \in [\rho, \rho + \frac{\delta}{4}],$$

and let

$$\Psi_r = \min\{u, \max\{\psi_r, v\}\}.$$

Notice that

$$u \geq \Psi_r \geq v \text{ in } B_1, \text{ and } \Psi_r = v \text{ in } B_\rho.$$

Let

$$D_r := \{u > \Psi_r > v\} \cap B_{\rho+\delta},$$

then, by the property (2.5) of the one-dimensional solution  $\varphi$ , we find

$$(5.2) \quad J(\Psi_r, D_r) = J(\psi_r, D_r) = \int_0^1 \mathcal{H}^{n-1}(\{\Psi_r = s\} \cap D_r) 2\sqrt{W(s)} ds.$$

Notice that

$$\{\Psi_r = s\} \cap D_r = \{u > s > v\} \cap \partial B_{r+\varphi^{-1}(s)} \cap B_{\rho+\delta}.$$

Thus, we average (5.2) for  $r \in [\rho, \rho + \delta/4]$ , and obtain

$$(5.3) \quad \int_\rho^{\rho+\delta/4} J(\Psi_r, D_r) dr \leq \frac{C}{\delta} \int_0^1 \mathcal{H}^n(\{u > s > v\} \cap (B_{\rho+\delta} \setminus B_\rho)) 2\sqrt{W(s)} ds.$$

We use (2.3) and the change of coordinates

$$s^{1-\gamma/2} = \sigma \quad \text{and obtain} \quad 2\sqrt{W(s)} ds = d\sigma.$$

The right hand side in (5.3) equals

$$\begin{aligned} & \frac{C}{\delta} \int_0^1 \mathcal{H}^n(\{u^{1-\gamma/2} > \sigma > v^{1-\gamma/2}\} \cap (B_{\rho+\delta} \setminus B_\rho)) d\sigma \\ & \leq \frac{C}{\delta} \left\| u^{1-\gamma/2} - v^{1-\gamma/2} \right\|_{L^1(B_{\rho+\delta} \setminus B_\rho)}. \end{aligned}$$

Thus, for all  $k$  sufficiently large, we can find an  $r = r_k \in [\rho, \rho + \delta/4]$ , such that

$$J(\Psi_r, D_r) \leq \epsilon.$$

Since in the annulus  $B_{\rho+\delta} \setminus B_\rho$  the function  $\Psi_r$  coincides with  $u$  or  $v$  outside  $D_r$  we find

$$(5.4) \quad J(\Psi_r, B_{\rho+\delta} \setminus B_\rho) \leq 3\epsilon.$$

Finally we define

$$w = \eta \Psi_r + (1 - \eta)u,$$

with  $\eta \in C_0^\infty(B_{\rho+\delta})$  a cutoff function with  $\eta = 1$  in  $B_{\rho+\delta/2}$ . Clearly,  $w = u$  outside  $B_{\rho+\delta}$  and  $w = \Psi_r$  in  $B_{\rho+\delta/2}$ , hence  $w = v$  in  $B_\rho$ . Moreover,

$$u \geq w \geq \Psi_r > 0 \implies W(w) \leq W(\Psi_r) \quad \text{in } B_{\rho+\delta} \setminus B_{\rho+\delta/2}.$$

Since

$$|\nabla w|^2 \leq 3 (|\nabla \Psi_r|^2 + |\nabla u|^2 + |\nabla \eta|^2 (u - \Psi_r)^2),$$

we find

$$J(w, B_{\rho+\delta} \setminus B_\rho) \leq 3 (J(\Psi_r, B_{\rho+\delta} \setminus B_\rho) + J(u, B_{\rho+\delta} \setminus B_\rho) + C(\delta) \|\Psi_r - u\|_{L^2}^2).$$

Using that,

$$|u - \Psi_r| \leq |u - v|,$$

we obtain

$$C(\delta) \|\Psi_r - u\|_{L^2}^2 \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

We find

$$J(w, B_{\rho+\delta} \setminus B_\rho) \leq 15\epsilon,$$

for all large  $k$ , which gives the desired conclusion under the assumption (5.1).

The general case follows easily from the interpolation procedure between the two ordered functions described above. We apply it two times, first in the annulus  $B_{\rho+\delta} \setminus B_{\rho+\delta/2}$  where we interpolate between  $u$  and  $\min\{u, v\}$  and then in the annulus  $B_{\rho+\delta/2} \setminus B_\rho$  where we interpolate between  $\min\{u, v\}$  and  $v$ .  $\square$

We recall now the functional  $\mathcal{F}$  introduced in Section 2, which is defined on the space of pairs  $(u, E) \in \mathcal{A}(\Omega)$

$$\mathcal{A}(\Omega) := \{(u, E) \mid u \in H^1(\Omega), \quad E \text{ Caccioppoli set}, \quad u \geq 0 \text{ in } \Omega, \quad u = 0 \text{ a.e. in } E\},$$

given by the Dirichlet - perimeter energy

$$\mathcal{F}_\Omega(u, E) = \int_\Omega |\nabla u|^2 dx + P_\Omega(E).$$

Here  $P_\Omega(E)$  represents the perimeter of  $E$  in  $\Omega$

$$P_\Omega(E) = [\nabla \chi_E]_{BV(\Omega)} = \int_\Omega |\nabla \chi_E|.$$

In the next two lemmas we establish the  $\Gamma$ -convergence of the  $J_\gamma$  to  $\mathcal{F}$ .

**Lemma 5.2** (Lower semicontinuity). *Let  $\gamma_k \rightarrow 2^-$  and  $u_k$  satisfy*

$$u_k^{1-\gamma_k/2} \rightarrow \chi_{E^c} \quad \text{in } L^1(\Omega), \quad u_k \rightarrow u \quad \text{in } L^2(\Omega).$$

*Then*

$$\liminf J_{\gamma_k}(u_k, \Omega) \geq \mathcal{F}_\Omega(u, E).$$

*Proof.* After passing to a subsequence we may assume that the two convergences above hold pointwise a.e. in  $\Omega$ . This implies that  $\{u > 0\} \setminus E^c$  is a set of measure zero, hence  $u = 0$  a.e. on  $E$ , and  $(u, E)$  is an admissible pair.

We write

$$J_{\gamma_k}(u_k, \Omega) = J_{\gamma_k}(u_k, \Omega \cap \{u_k \leq \epsilon\}) + J_{\gamma_k}(u_k, \Omega \cap \{u_k > \epsilon\}).$$

By the coarea formula and the definition of  $W$  (see (2.1))

$$\begin{aligned}
J_{\gamma_k}(u_k, \Omega \cap \{u_k \leq \epsilon\}) &\geq \int_{\{u_k \leq \epsilon\}} |\nabla u_k| 2\sqrt{W(u_k)} dx \\
&= \int_{\{u_k \leq \epsilon\}} |\nabla u_k^{1-\gamma_k/2}| dx \\
(5.5) \qquad \qquad \qquad &= \int_{\Omega} |\nabla \bar{u}_k^{1-\gamma_k/2}| dx, \quad \text{with } \bar{u}_k := \min\{u_k, \epsilon\}.
\end{aligned}$$

Moreover,  $\bar{u}_k^{1-\gamma_k/2}$  converges in  $L^1$  to  $\chi_E$ , hence

$$\liminf J_{\gamma_k}(u_k, \Omega \cap \{u_k \leq \epsilon\}) \geq \int_{\Omega} |\nabla \chi_E| dx,$$

by the lower semicontinuity of the BV norm. On the other hand

$$J_{\gamma_k}(u_k, \Omega \cap \{u_k > \epsilon\}) \geq \int_{\Omega} |\nabla (u_k - \epsilon)^+|^2 dx,$$

and since  $(u_k - \epsilon)^+ \rightarrow (u - \epsilon)^+$  in  $L^2$ , we obtain

$$\liminf J_{\gamma_k}(u_k, \Omega \cap \{u_k > \epsilon\}) \geq \int_{\Omega} |\nabla (u - \epsilon)^+|^2 dx.$$

By adding the inequalities we find

$$\liminf J_{\gamma_k}(u_k, \Omega) \geq \int_{\Omega} |\nabla (u - \epsilon)^+|^2 dx + P_{\Omega}(E),$$

and the conclusion is proved by letting  $\epsilon \rightarrow 0$ .  $\square$

**Lemma 5.3.** *Let  $(u, E) \in \mathcal{A}(\Omega)$  with  $u$  a continuous function in a Lipschitz domain  $\bar{\Omega}$ . Then, given a sequence  $\gamma_k \rightarrow 2^-$  we can construct a sequence  $u_k$  such that*

$$\begin{aligned}
u_k^{1-\gamma_k/2} &\rightarrow \chi_{E^c} \quad \text{in } L^1(\Omega), \quad u_k \rightarrow u \quad \text{in } L^2(\Omega), \\
J_{\gamma_k}(u_k, \Omega) &\rightarrow \mathcal{F}_{\Omega}(u, E).
\end{aligned}$$

In view of the lower semicontinuity property in  $\Omega \setminus \bar{D}$ , where  $D \subset \Omega$  is a subdomain, we obtain that

$$\int_{\bar{D}} |\nabla u|^2 dx + \int_{\bar{D}} |\nabla \chi_E| \geq \limsup J_{\gamma_k}(u_k, D).$$

*Proof.* For the convergence of the energies it suffices to show that

$$\limsup J_{\gamma_k}(u_k, \Omega) \leq \mathcal{F}_{\Omega}(u, E).$$

Fix  $\epsilon > 0$  small. First we approximate  $E$  in  $\Omega$  by a smooth set  $F \subset \mathbb{R}^n$  which is included in the open set  $\{u < \epsilon\}$  in  $\Omega$  (which contains a neighborhood of  $E$ ). Precisely, by the results of Modica [M], there exists a smooth set  $F$  which approximates  $E$  in  $\Omega$  in the sense that

$$\begin{aligned}
(5.6) \qquad \qquad \qquad &F \cap \Omega \subset \{u < \epsilon\}, \\
&\|\chi_{F \cap \Omega} - \chi_E\|_{L^1} \leq \epsilon, \quad P_{B_1}(F \cap \Omega) \leq P_{\Omega}(E) + \epsilon, \\
&\mathcal{H}^{n-1}(\partial F \cap \partial \Omega) = 0.
\end{aligned}$$

In view of this, it suffices to prove the lemma with  $E$  replaced by  $\tilde{E} := F$  and  $u$  replaced by  $\tilde{u} := (u - 2\epsilon)^+$  which is an approximation of  $u$  in  $H^1(\Omega)$ . Notice that

by construction  $\tilde{u}$  vanishes in a  $\delta$ -neighborhood of  $\tilde{E}$  for some small  $\delta$ . We define  $u_k$  in  $B_1$  as

$$u_k := \max\{\varphi_k(d), \tilde{u}\},$$

where  $d$  represents the distance in  $\mathbb{R}^n$  to  $\tilde{E}$ . Next we check that  $u_k$  satisfies the desired conclusions.

Clearly  $u_k = 0$  on  $\tilde{E}$ , and using that

$$C \geq u_k \geq \varphi_k(d) \quad \text{on } \Omega \setminus \tilde{E},$$

and  $1 - \gamma_k/2 \rightarrow 0^+$  we have

$$u_k^{1-\gamma_k/2} \rightarrow 1 \quad \text{in } \Omega \setminus \tilde{E},$$

hence

$$u_k^{1-\gamma_k/2} \rightarrow \chi_{\tilde{E}} \quad \text{in } L^1(\Omega).$$

Here we used that  $\varphi_k(d) = c_\gamma^* d^\alpha$ , with  $c_\gamma^*$  defined in (2.4), and we have

$$(c_\gamma^*)^{1-\gamma/2} \rightarrow 1 \quad \text{as } \gamma \rightarrow 2.$$

Since  $\varphi_k(d)$  converges uniformly to 0 as  $k \rightarrow \infty$  we also obtain

$$u_k \rightarrow \tilde{u} \quad \text{in } L^2(\Omega).$$

Using property (2.5) we obtain that

$$\begin{aligned} J(\varphi(d), \{a < d < b\} \cap \Omega) &= \int_{\varphi(a)}^{\varphi(b)} \mathcal{H}^{n-1}(\{\varphi(d) = s\} \cap \Omega) 2\sqrt{W(s)} ds \\ &= \int_a^b \mathcal{H}^{n-1}(\{d = t\} \cap \Omega) \omega_\gamma(t) dt, \end{aligned}$$

with

$$\omega_\gamma(t) := 2\sqrt{W(\varphi(t))} \varphi'(t).$$

Notice that

$$\omega_\gamma(t) dt = \varphi'(t)^2 + W(\varphi(t)) dt,$$

represents the measure of the one-dimensional solution which, as  $k \rightarrow \infty$ , converges weakly in any bounded interval  $[-a, a]$  to the Dirac delta measure at 0. On the other hand (5.6) implies that

$$\mathcal{H}^{n-1}(\{\varphi(d) = t\} \cap \Omega) \rightarrow P_\Omega(\tilde{E} \cap \Omega) \quad \text{as } t \rightarrow 0.$$

In conclusion, we find that as  $k \rightarrow \infty$

$$J(\varphi(d), \{0 < d < \delta\} \cap \Omega) \rightarrow P_\Omega(\tilde{E} \cap \Omega)$$

and

$$J(\varphi(d), \{d > \delta\} \cap \Omega) \rightarrow 0.$$

Using that

$$u_k = \varphi(d) \quad \text{if } d < \delta,$$

and

$$u_k \geq \varphi(d) \implies W(u_k) \leq W(\varphi(d)) \quad \text{if } d > \delta,$$

we find

$$\limsup J(u_k, \Omega) \leq P_\Omega(\tilde{E} \cap \Omega) + \int_\Omega |\nabla \tilde{u}|^2 dx.$$

□

We are finally ready to prove our main theorem.

*Proof of Theorem 2.5.* The  $L^2$  convergence follows from the uniform bound of the  $u_k$  in  $H^1(\Omega)$ .

By the coarea formula (see (5.5)) we find that

$$[u_k^{1-\gamma_k/2}]_{BV(\Omega)} \leq M$$

and using the inequality

$$u_k^{1-\gamma_k/2} \leq 1 + u_k^2$$

we find that  $u_k^{1-\gamma_k/2}$  are uniformly bounded in  $BV(\Omega)$ . Thus, after passing to a subsequence, we have

$$(5.7) \quad u_k^{1-\gamma_k/2} \rightarrow g \quad \text{in } L^1(\Omega),$$

for some non-negative  $g \in BV(\Omega)$ . We claim that

$$(5.8) \quad g = \chi_{E^c} \text{ for some set } E.$$

First we show that for all  $\delta > 0$  small

$$\{\delta \leq g \leq 1 - \delta\} \quad \text{has measure zero.}$$

Otherwise, for all large  $k$ , the set

$$\{\delta/2 \leq u_k^{1-\gamma_k/2} \leq 1 - \delta/2\}$$

has measure bounded below by a fixed positive constant. On this set

$$(5.9) \quad W(u_k) \geq W((1 - \delta/2)^{\frac{2}{2-\gamma_k}}) = c_{\gamma_k} (1 - \delta/2)^{-\frac{2\gamma_k}{2-\gamma_k}} \rightarrow \infty,$$

as  $k \rightarrow \infty$  and we contradict the uniform upper bound for the energy of  $u_k$  in  $\Omega$ .

Similarly we find that the set

$$\{g \geq 1 + \delta\} \quad \text{has measure zero.}$$

Indeed, otherwise

$$\{u_k^{1-\gamma_k/2} \geq 1 + \delta/2\}$$

has measure bounded below by a fixed positive constant. Then we contradict the uniform upper bound for the  $L^2$  norm of  $u_k$  since on the set above

$$u_k^2 \geq (1 + \delta/2)^{\frac{4}{2-\gamma_k}} \rightarrow \infty$$

as  $k \rightarrow \infty$ , and the claim (5.8) is proved.

The argument above implies also that

$$(5.10) \quad \chi_{\{u_k > 0\}} \rightarrow \chi_{E^c} \quad \text{in } L^1(\Omega).$$

For example if

$$|\{u_k > 0\} \setminus E^c| \geq \mu > 0$$

for some positive constant  $\mu$  independent of  $k$ , then (5.7)-(5.8) imply

$$|\{0 < u_k^{1-\gamma_k/2} \leq \frac{1}{2}\}| \geq \mu/2,$$

and we get a contradiction as in (5.9). Also

$$|E^c \setminus \{u_k > 0\}| = |E^c \cap \{u_k = 0\}| \rightarrow 0,$$

as  $k \rightarrow \infty$ , follows from the convergence (5.7)-(5.8).

Next we assume that  $u_k$  are minimizers for  $J_{\gamma_k}$  and prove the minimality of  $(u, E)$  for  $\mathcal{F}$ . The argument is standard and follows from Proposition 5.1. We sketch it for completeness.

For simplicity let  $\Omega = B_1$ . Since the functions  $u_k$  are uniformly Hölder continuous on compact sets of  $B_1$  we find that the limiting function  $u$  is Hölder continuous in  $B_1$  and the convergence  $u_k \rightarrow u$  is uniform on compact subsets.

Let  $(v, F)$  be an admissible pair which coincides with  $(u, E)$  near  $\partial B_1$  and let

$$\mathcal{R} := B_{\rho+\delta} \setminus B_\rho,$$

be an annulus near  $\partial B_1$  where the two pairs coincide.

Denote by  $v_k$  be the functions constructed in Lemma 5.3 corresponding to the pair  $(v, F)$  in  $B_{\rho+\delta}$ . Since  $u_k$  and  $v_k$  satisfy the hypotheses of Proposition 5.1 we can construct  $w_k$  as the interpolation between  $u_k$  and  $v_k$ . By the minimality of  $u_k$  in  $B_1$  and the conclusion of Proposition 5.1 we have

$$J(u_k, B_1) \leq J(w_k, B_1) \leq J(u_k, B_1 \setminus B_\rho) + J(v_k, B_{\rho+\delta}) + o(1).$$

This gives

$$J(u_k, B_\rho) \leq J(v_k, B_{\rho+\delta}) + o(1),$$

and by taking  $k \rightarrow \infty$ , we find from Lemmas 5.2 and 5.3

$$\mathcal{F}_{B_\rho}(u, E) \leq \mathcal{F}_{B_{\rho+\delta}}(v, F).$$

We let  $\rho \rightarrow 1$  and obtain the desired conclusion

$$\mathcal{F}_{B_1}(u, E) \leq \mathcal{F}_{B_1}(v, F).$$

Finally, the uniform convergence of the free boundaries follows from the uniform density estimates and the  $L^1$  convergence (5.10).  $\square$

#### REFERENCES

- [AIC] Allen S. M., Cahn J. W., *Ground State Structures in Ordered Binary Alloys with Second Neighbor Interactions*. Acta Metall. 20, (1972) 423–433.
- [AC] Alt H. W., Caffarelli L. A., *Existence and regularity for a minimum problem with free boundary*. J. Reine Angew. Math 325, 105–144 (1981)
- [ACF] Alt H., Caffarelli L., Friedman A., *Variational problems with two phases and their free boundaries*. Trans. Amer. Math. Soc. 282 (1984), no. 2, 43–461.
- [AP] Alt H.W., Phillips D., *A free boundary problem for semilinear elliptic equations*. J. Reine Angew. Math., 368 (1986), pp. 63–107.
- [ADM] Ambrosio L., De Philippis G., Martinazzi L., *Gamma-convergence of nonlocal perimeter functionals*. Manuscripta Math. 134 (2011), no. 3-4, 377–403.
- [ACKS] Athanopoulos I., L. A. Caffarelli L., Kenig C. and Salsa S., *An area-Dirichlet integral minimization problem*. Comm. Pure Appl. Math. 54 (2001), no. 4, 479–499.
- [CC] Caffarelli L., Cordoba A., *Uniform convergence of a singular perturbation problem*. Comm. Pure Appl. Math. 48 (1995), no. 1, 1–12.
- [CS] Caffarelli L., Salsa S. *A geometric approach to free boundary problems*. Graduate Studies in Mathematics, 68. American Mathematical Society, Providence, RI, 2005. x+270
- [CV] Caffarelli L., Valdinoci E., *Uniform estimates and limiting arguments for non- local minimal surfaces*. Calc. Var. Partial Differential Equations 41 (2011), no. 1-2, 203–240.
- [DS] De Silva D., Savin O., *Regularity of Lipschitz free boundaries for the thin one-phase problem*. J. Eur. Math. Soc. (JEMS) 17 (2015), no. 6, 1293–1326.
- [DFPV] Dipierro S., Figalli A., Palatucci G., Valdinoci E., *Asymptotics of the s perimeter as  $s \rightarrow 0$* . Discrete Contin. Dyn. Syst. 33 (2013), no. 7, 2777–2790.
- [M] Modica L., *The gradient theory of phase transitions and the minimal interface criterion*. Arch. Rational Mech. Anal. 98 (1987), no. 2, 123–142.

- [MM] Modica L., Mortola S. *Un esempio di  $\Gamma$ -convergenza.* (Italian) Boll. Un. Mat. Ital. B (5) **14** (1977), no. 1, 285–299.
- [PSU] Petrosyan A., Shahgholian H., Uraltseva N., *Regularity of free boundaries in obstacle-type problems.* Graduate Studies in Mathematics, 136. American Mathematical Society, Providence, RI, 2012. x+221 pp.
- [PV] Petrosyan A., Valdinoci E., *Density estimates for a degenerate/singular phase-transition model.* SIAM J. Math. Anal. 36 (2005), no. 4, 1057–1079.
- [SV] Savin O., Valdinoci E., *Density estimates for a nonlocal variational model via the Sobolev inequality.* SIAM J. Math. Anal. 43 (2011), no. 6, 2675–2687.
- [V] Valdinoci E., *Plane-like minimizers in periodic media: jet flows and Ginzburg-Landau-type functionals.* J. Reine Angew. Math. 574 (2004), 147–185.

DEPARTMENT OF MATHEMATICS, BARNARD COLLEGE, COLUMBIA UNIVERSITY, NEW YORK, NY 10027

*Email address:* `desilva@math.columbia.edu`

DEPARTMENT OF MATHEMATICS, COLUMBIA UNIVERSITY, NEW YORK, NY 10027

*Email address:* `savin@math.columbia.edu`