

A trail of dark matter-free galaxies from a bullet dwarf collision

Pieter van Dokkum¹, Zili Shen¹, Michael A. Keim¹, Sebastian Trujillo-Gomez², Shany Danieli³, Dhruva Dutta Chowdhury¹, Roberto Abraham⁴, Charlie Conroy⁵, J. M. Diederik Kruijssen², Daisuke Nagai⁶, Aaron Romanowsky^{7,8}

¹*Department of Astronomy, Yale University, New Haven, CT 06511, USA*

²*Astronomisches Rechen-Institut, Zentrum für Astronomie der Universität Heidelberg, Mönchhofstraße 12-14, D-69120 Heidelberg, Germany*

³*Department of Astrophysical Sciences, 4 Ivy Lane, Princeton University, Princeton, NJ 08544, USA*

⁴*Department of Astronomy & Astrophysics, University of Toronto, 50 St. George Street, Toronto, ON M5S 3H4, Canada*

⁵*Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA, USA*

⁶*Department of Physics, Yale University, New Haven, CT 06511, USA*

⁷*Department of Physics and Astronomy, San José State University, San Jose, CA 95192, USA*

⁸*University of California Observatories, 1156 High Street, Santa Cruz, CA 95064, USA*

The ultra-diffuse galaxies DF2 and DF4 in the NGC1052 group share several unusual properties: they both have large sizes,¹ rich populations of overluminous and large globular clusters,^{2–6} and very low velocity dispersions indicating little or no dark matter.^{7–10} It has been suggested that these galaxies were formed in the aftermath of high velocity collisions of gas rich galaxies,^{11–13} events that resemble the collision that created the bullet cluster¹⁴ but on much smaller scales. The gas separates from the dark matter in the collision and subsequent star formation leads to the formation of one or more dark matter-free galaxies.¹² Here we show that the present-day line-of-sight distances and radial velocities of DF2 and DF4 are consistent with their joint formation in the aftermath of a single bullet-dwarf collision, around eight billion years ago. Moreover, we find that DF2 and DF4 are part of an apparent linear substructure of seven to eleven large, low-luminosity objects. We propose that these all originated in the same event, forming a trail of dark matter-free galaxies that is roughly more than two megaparsecs long and angled $7^\circ \pm 2^\circ$ from the line of sight. We also tentatively identify the highly dark matter-dominated remnants of the two progenitor galaxies that are expected¹¹ at the leading edges of the trail.

We begin with the assumption that it is not a coincidence that DF2 and DF4 have the same set of otherwise-unique properties and that they were in close proximity to one another at the time of their formation. With that assumption, collisional formation is implied by their present-day ra-

dial velocities and three-dimensional locations. The geometry is shown in Fig. 1a. The relative radial velocity of the galaxies is high, 358 km s^{-1} , which is three times the velocity dispersion of the NGC1052 group (about 115 km s^{-1}). Furthermore, while the two galaxies are separated by only 0.24 Mpc in the plane of the sky³ a differential tip of the red giant branch (TRGB) analysis has shown that they are 2.1 ± 0.5 Mpc apart along the line of sight,^{15, 16} which is five times the virial radius of NGC1052 (about 400 kpc^{16, 17}). In the context of a shared origin of both galaxies, their radial velocities are consistent with their line-of-sight distances, that is, the closest galaxy (DF4) is moving toward us (with respect to the mean velocity) and the farthest galaxy (DF2) is moving away from us. Tracing their line-of-sight positions back in time, we infer that they must have formed in a high-velocity encounter. The minimum time since that encounter is about 6 Gyr, for constant motion along the line of sight.

The situation at that point, a $\gtrsim 300 \text{ km s}^{-1}$ encounter between two galaxies in the gas-rich environment of a young group, is a close match to the initial conditions of mini-bullet-cluster¹⁴ scenarios that have previously been proposed for the formation of DF2 and DF4.^{11–13} In a near head-on collision between two gas-rich galaxies, the collisional gas can be shocked and separated from the collisionless dark matter and pre-existing stars.^{11, 12} The accompanying star formation favours massive clumps in highly compressed gas,^{11, 13} producing the unusual globular clusters as well as the lack of dark matter. Galaxies that form this way are initially compact,¹³ in apparent conflict with the large half-light radii of DF2 and DF4. However, intense feedback accompanying the formation of the globular clusters is expected to increase the sizes of the newly formed galaxies, an effect that is particularly efficient when the stars are not bound by a dark matter halo.^{18, 19} These previous studies focused on the formation of a single new galaxy at the collision site, but we propose that both DF2 and DF4 were formed in a single ‘bullet dwarf’ event. An illustration of the proposed scenario is shown in the Methods section.

We have sufficient information to construct a plausible model for the geometry and timing of the collision. It is likely that the event took place near the central elliptical galaxy, NGC1052, as it is roughly halfway between DF2 and DF4 in projection and its deep potential well is conducive to high-speed interactions. It is also likely that at least one of the progenitor galaxies was a satellite of NGC1052 as the probability of a collision of two unbound galaxies is ex-




Figure 1: **Geometry of the DF2 and DF4 system.** **a.** The radial velocity difference between DF2 and DF4 is 358 km s^{-1} and this large velocity difference is accompanied by a large line-of-sight separation of $2.1 \pm 0.5 \text{ Mpc}$.^{15,16} The geometry implies that the galaxies are moving away from each other. Tracing their positions back in time, we infer that they were formed in a high-speed encounter $\geq 6 \text{ Gyr}$ ago. **b.** Example of a collisional scenario involving NGC1052. Velocities are given with respect to that galaxy ($cz = 1,488 \text{ km s}^{-1}$). An infalling gas-rich galaxy on an unbound orbit collided with a satellite of NGC1052 about 8 Gyr ago, leading to two dark remnants (possibly RCP32 and DF7), DF2 and DF4, and three to seven other dark matter-free galaxies.

tremely small.¹² We assume that the second progenitor was not bound to NGC1052, and either on first infall or a satellite of another massive galaxy in the group.²⁰ DF4 has a velocity difference of -43 km s^{-1} with respect to NGC1052, whereas the velocity of DF2 is $+315 \text{ km s}^{-1}$. In Fig. 1b we show a configuration for DF2, DF4, and NGC1052 that satisfies these constraints. Progenitor 1 arrived in the vicinity of NGC1052 with a high ($> 300 \text{ km s}^{-1}$) relative velocity and collided with progenitor 2, which was on a bound orbit. The collision produced DF2 and DF4, with DF2's velocity and orbit similar to progenitor 1 and DF4's similar to progenitor 2. DF2 is currently unbound and about 3 Mpc behind the group, whereas DF4 remained bound and has begun falling back. This geometry is not unique but it is similar to examples that have been explored in simulations.¹²

It is also consistent with the near-identical tidal distortions of the two galaxies:²¹ in the geometry of Fig. 1b, DF2 and DF4 were at the same distance from NGC1052 when they were formed, and as neither galaxy experienced a stronger tidal field afterwards, their morphologies have remained the same. In this model, DF2 has travelled about 3 Mpc since the collision with an average velocity with respect to NGC1052 that is probably higher than its present-day value of 315 km s^{-1} . Assuming an average post-collisional velocity of $\langle v \rangle \sim 350 \text{ km s}^{-1}$ dates the collision to about 8 Gyr ago, in excellent agreement with the ages of the globular clusters and the diffuse light in DF2 ($9 \pm 2 \text{ Gyr}$, as measured from optical spectra^{2,22}).

We further investigate the possible joint formation of




Figure 2: **A linear feature in the spatial distribution of faint galaxies in the NGC1052 field.** **a.** Distribution of galaxies with $g > 16.5$ (black circles) from a recent compilation of low surface brightness objects in the NGC1052 field.²³ The positions are with respect to the coordinates of NGC1052, the central bright elliptical galaxy in the group. **b.** Hough transform²⁴ of the spatial distribution. The peak corresponds to a line that has 11 galaxies located within ± 30 kpc. The significance of the feature, as determined from randomized realizations of the data, is 97%. **c.** Zoom of **a**, shown with the four brightest members of the NGC1052 group. Yellow circles correspond to the positions of galaxies in the box shown in panel **a**. The orientation and offset of the line corresponds to the location of the peak of the Hough transform. Both DF2 and DF4 are part of the linear feature.

DF2 and DF4 by examining the spatial distribution of galaxies along the DF2–DF4 axis. Given the complex gas distribution during and after the event, it may be that more than two dark matter-deficient objects were formed in the wake of the collision.¹² Furthermore, the bullet-dwarf scenario predicts the existence of two dark matter-dominated objects that are the remnants of the progenitor galaxies. These should precede DF2, DF4 and any other dark matter-deficient galaxies along their path, as they have the highest velocities relative to the barycentre.¹¹

The spatial distribution of galaxies with magnitude $g > 16.5$ in a recently compiled catalog of the NGC1052 field²³ is shown in Fig. 2a. An objective search for linear features in this distribution is performed using a discrete implementation of the Hough transform.²⁴ The transform is shown in Fig. 2b. There is a clear peak with 11 galaxies on a line, corresponding to the relation

$$\Delta \text{DEC} = 0.45 \Delta \text{RA} - 2.8, \quad (1)$$

in units of arcminutes north and west of NGC1052. This relation is shown by the line in Fig. 2c. Both DF2 and DF4 are in the sample of 11 galaxies. The probability that the peak arose by chance is 3%, and the probability that the peak arose by chance and that both DF2 and DF4 are part of it is 0.6% (see Methods). Before turning to the properties of the galaxies in the trail, we note that the Hough transform provides post hoc validation of our initial assumption that DF2 and DF4 are related to each other.

Images of the 11 galaxies that are part of the trail are shown in Fig. 3. The average galaxy density in the central $R < 30'$ implies that 2 ± 2 of the 11 are chance projections, and we infer that there are 7–11 galaxies in the struc-

ture. Besides DF2 and DF4 other galaxies in the trail are also uncommonly large for their luminosity. The relation between size and apparent magnitude for faint galaxies in the NGC1052 group is shown in Fig. 4a. After subtracting a simple linear least squares fit to the running median (broken line), we find that galaxies in the trail are on average 26% larger than the rest of the sample. The Wilcoxon probability that the trail galaxies and the rest of the sample are drawn from the same size distribution is $< 1\%$. The spatial distribution of galaxies color-coded by their (magnitude-dependent) relative size is shown in Fig. 4b. The unusual prevalence of large, low surface brightness galaxies in the central regions of the NGC1052 group had been noticed previously;^{1,23} here we propose that the bullet dwarf event was responsible for it.

We highlight three large and unusual galaxies that are part of the sample of eleven. DF5 looks very similar to DF2 and DF4 but has a much lower surface brightness.^{1,21} It is so close in projection to DF4 (only $1.6'$ away) that both galaxies were observed in the same HST/ACS pointing.^{1,9} In the bullet dwarf scenario this puzzling object is readily explained as another dark matter-free fragment that formed in the aftermath of the collision. RCP32 and DF7 are the furthest away from the center of the structure, and are “ahead” of DF2 and DF4 respectively. RCP32 is the faintest ultra-diffuse galaxy (UDG) that has so far been discovered in integrated light and may have a globular cluster system.²³ DF7 is an elongated UDG that was previously observed with HST¹ (see Methods section). We tentatively identify RCP32 and DF7 as the candidate remnants of the original, pre-collision galaxies. In the geometry of Fig. 1b RCP32 could be the remnant of progenitor 1, the gas-rich object that was not bound to NGC1052, and the brighter galaxy DF7 could be the remnant of progen-




Figure 3: **Galaxies on the DF2–DF4 axis.** Legacy survey image of the central region of the NGC1052 group, highlighting the 11 galaxies that are part of the trail according to the Hough transform. Several (2 ± 2) of these are expected to be chance projections. LEDA4014647 is a candidate for an interloper (that is, an unrelated group member) given its brightness and relative compactness. Its radial velocity was listed as 1680 ± 60 in earlier SDSS releases (DR3) but was later erroneously revised to a $z = 0.7$ QSO (DR16). Judging from morphology alone, RCP26 and Ta21-12000 may also be chance projections. Besides DF2 and DF4, RCP32, DF5, and DF7 all satisfy or nearly satisfy the UDG criteria. RCP32 and DF7 are candidates for the two dark matter-dominated remnants that have been predicted to precede dark matter-deficient galaxies along the post-collision trajectory.¹¹ We also highlight DF9 (SDSS J024007.01–081344.4), a galaxy with a bright star cluster that falls on the trail but is not part of the objectively-selected sample.²³

itor 2, the galaxy that was a NGC1052 satellite at the time of collision. In this context the elongation can be understood as the effect of the strong tidal forces at the time of collision.²⁵

The scenario that is proposed here makes predictions for the properties of the collision products, and further observations can test and refine this explanation. As the formation of DF2 and DF4 was triggered by a single event the ages of the globular clusters of DF4, which have not yet been measured, should be identical to those of the clusters in DF2 (9 ± 2 Gyr.^{2,22}). A stringent and model-independent test is to directly compare the (averaged) spectra of clusters in the two galaxies; any clear differences could falsify our model, particularly differences in age-sensitive spectral features. Turning to other galaxies on the trail, their kinematics are predicted to be consistent with baryon-only models — except RCP32 and DF7, which could show evidence for an unusually *low* baryon fraction. Furthermore, the radial velocities of trail galaxies should follow the approximate re-

lation $cz \approx 1700 - 10\Delta RA \text{ km s}^{-1}$, with ΔRA defined as in Eq. 1. Similarly, their line-of-sight distances are predicted to be $D \approx 21.5 - 0.06\Delta RA \text{ Mpc}$, for TRGB measurements that are calibrated to $D = 22.1 \text{ Mpc}$ for DF2.¹⁶ We note that these relations likely have considerable scatter and are not expected to be linear for most geometries. Also, 0–4 galaxies of the 11 are expected to be interlopers (see Fig. 3), and some galaxies that seem to be off the trail may in fact be part of it due to the foreshortening (see Fig. 1a). This may be the case for DF1, a very large and diffuse galaxy (marked in Fig. 4) that is only $14'$ off the DF2–DF4 axis and is elongated toward the center of the trail.

Bullet dwarf collisions hold the potential to constrain the self-interaction cross section of dark matter. Modeling of the bullet cluster has provided an upper limit,²⁶ but as self-interacting dark matter was introduced to explain the “cored” dark matter density profiles of low mass galaxies²⁷ it is important to measure the cross section on small scales.²⁸ Quan-




Figure 4: **The sizes of galaxies on the DF2–DF4 axis.** **a.** The apparent size – apparent magnitude relation for low surface brightness objects in NGC1052, using recent measurements from Legacy data.²³ Galaxies that are part of the trail are labeled with solid symbols. The thin solid line is a running median, with $N = 7$. The broken line is a least squares fit to the running median and has the form $\log r_e = -0.09(g - 20) + 0.87$ kpc. Galaxies in the trail are typically larger than other galaxies of the same magnitude. **b.** Distribution of galaxies color-coded by their location with respect to the least squares fit. The largest galaxies in the group are preferentially located along the DF2–DF4 axis.

titative constraints will likely require more than a single example of a bullet dwarf. Encouragingly, these events are probably more common than the collision that produced the bullet cluster;²⁹ a search for plausible DF2/DF4 progenitors in the Illustris TNG cosmological simulation³⁰ produced 248 head-on high velocity collisions in a 100^3 Mpc^3 volume,¹² corresponding to ~ 8 within $D < 20 \text{ Mpc}$.

1. Cohen, Y. *et al.* The Dragonfly Nearby Galaxies Survey. V. HST/ACS Observations of 23 Low Surface Brightness Objects in the Fields of NGC 1052, NGC 1084, M96, and NGC 4258. *ApJ* **868**, 96 (2018).
2. van Dokkum, P. *et al.* An Enigmatic Population of Luminous Globular Clusters in a Galaxy Lacking Dark Matter. *ApJ* **856**, L30 (2018).
3. van Dokkum, P., Danieli, S., Abraham, R., Conroy, C. & Romanowsky, A. J. A Second Galaxy Missing Dark Matter in the NGC 1052 Group. *ApJ* **874**, L5 (2019).
4. Dutta Chowdhury, D., van den Bosch, F. C. & van Dokkum, P. On the Orbital Decay of Globular Clusters in NGC 1052-DF2: Testing a Baryon-only Mass Model. *ApJ* **877**, 133 (2019).
5. Dutta Chowdhury, D., van den Bosch, F. C. & van Dokkum, P. On the Evolution of the Globular Cluster System in NGC 1052-DF2: Dynamical Friction, Globular-Globular Interactions, and Galactic Tides. *ApJ* **903**, 149 (2020).
6. Shen, Z., van Dokkum, P. & Danieli, S. A Complex Luminosity Function for the Anomalous Globular Clusters in NGC 1052-DF2 and NGC 1052-DF4. *ApJ* **909**, 179 (2021).

7. van Dokkum, P. *et al.* A galaxy lacking dark matter. *Nature* **555**, 629–632 (2018).
8. van Dokkum, P. *et al.* A Revised Velocity for the Globular Cluster GC-98 in the Ultra Diffuse Galaxy NGC 1052-DF2. *Research Notes of the American Astronomical Society* **2**, 54 (2018).
9. Danieli, S., van Dokkum, P., Conroy, C., Abraham, R. & Romanowsky, A. J. Still Missing Dark Matter: KCWI High-resolution Stellar Kinematics of NGC1052-DF2. *ApJ* **874**, L12 (2019).
10. Emsellem, E. *et al.* The ultra-diffuse galaxy NGC 1052-DF2 with MUSE. I. Kinematics of the stellar body. *A&A* **625**, A76 (2019).
11. Silk, J. Ultra-diffuse galaxies without dark matter. *MNRAS* **488**, L24–L28 (2019).
12. Shin, E.-j. *et al.* Dark Matter Deficient Galaxies Produced via High-velocity Galaxy Collisions in High-resolution Numerical Simulations. *ApJ* **899**, 25 (2020).
13. Lee, J., Shin, E.-j. & Kim, J.-h. Dark Matter Deficient Galaxies and Their Member Star Clusters Form Simultaneously during High-velocity Galaxy Collisions in 1.25 pc Resolution Simulations. *ApJ* **917**, L15 (2021).
14. Clowe, D. *et al.* A Direct Empirical Proof of the Existence of Dark Matter. *ApJ* **648**, L109–L113 (2006).
15. Danieli, S. *et al.* A Tip of the Red Giant Branch Distance to the Dark Matter Deficient Galaxy NGC 1052-DF4 from Deep Hubble Space Telescope Data. *ApJ* **895**, L4 (2020).

16. Shen, Z. *et al.* A Tip of the Red Giant Branch Distance of 22.1 ± 1.2 Mpc to the Dark Matter Deficient Galaxy NGC 1052-DF2 from 40 Orbits of Hubble Space Telescope Imaging. *ApJ* **914**, L12 (2021).
17. Forbes, D. A., Alabi, A., Brodie, J. P. & Romanowsky, A. J. Dark matter and no dark matter: on the halo mass of NGC 1052. *MNRAS* **489**, 3665–3669 (2019).
18. Trujillo-Gomez, S., Kruijssen, J. M. D., Keller, B. W. & Reina-Campos, M. Constraining the formation of NGC 1052-DF2 from its unusual globular cluster population. *MNRAS* **506**, 4841–4854 (2021).
19. Trujillo-Gomez, S., Kruijssen, J. M. D. & Reina-Campos, M. The emergence of dark matter-deficient ultra-diffuse galaxies driven by scatter in the stellar mass-halo mass relation and feedback from globular clusters. *MNRAS*, in press arXiv:2103.08610 (2021).
20. van Dokkum, P., Danieli, S., Romanowsky, A., Abraham, R. & Conroy, C. The Distance to NGC 1042 in the Context of its Proposed Association with the Dark Matter-deficient Galaxies NGC 1052-DF2 and NGC 1052-DF4. *Research Notes of the American Astronomical Society* **3**, 29 (2019).
21. Keim, M. A. *et al.* Tidal Distortions in NGC1052-DF2 and NGC1052-DF4: Independent Evidence for a Lack of Dark Matter. *arXiv e-prints* arXiv:2109.09778 (2021).
22. Fensch, J. *et al.* The ultra-diffuse galaxy NGC 1052-DF2 with MUSE. II. The population of DF2: stars, clusters, and planetary nebulae. *A&A* **625**, A77 (2019).
23. Román, J., Castilla, A. & Pascual-Granado, J. Discovery and analysis of low-surface-brightness galaxies in the environment of NGC 1052. *A&A* **656**, A44 (2021).
24. Duda, R. O. & Hart, P. E. Use of the Hough transformation to detect lines and curves in pictures. *Communications of the ACM* **15**, 11–15 (1972).
25. Rich, R. M. *et al.* A tidally distorted dwarf galaxy near NGC 4449. *Nature* **482**, 192–194 (2012).
26. Randall, S. W., Markevitch, M., Clowe, D., Gonzalez, A. H. & Bradač, M. Constraints on the Self-Interaction Cross Section of Dark Matter from Numerical Simulations of the Merging Galaxy Cluster 1E 0657-56. *ApJ* **679**, 1173–1180 (2008).
27. Spergel, D. N. & Steinhardt, P. J. Observational Evidence for Self-Interacting Cold Dark Matter. *Phys. Rev. Lett.* **84**, 3760–3763 (2000).
28. Tulin, S. & Yu, H.-B. Dark matter self-interactions and small scale structure. *Phys. Rep.* **730**, 1–57 (2018).
29. Bouillot, V. R., Alimi, J.-M., Corasaniti, P.-S. & Rasera, Y. Probing dark energy models with extreme pairwise velocities of galaxy clusters from the DEUS-FUR simulations. *MNRAS* **450**, 145–159 (2015).
30. Pillepich, A. *et al.* Simulating galaxy formation with the IllustrisTNG model. *MNRAS* **473**, 4077–4106 (2018).
31. Dark Energy Survey Collaboration. The Dark Energy Survey: more than dark energy - an overview. *MNRAS* **460**, 1270–1299 (2016).
32. Peng, C. Y., Ho, L. C., Impey, C. D. & Rix, H.-W. Detailed Structural Decomposition of Galaxy Images. *AJ* **124**, 266–293 (2002).
33. Beers, T. C., Flynn, K. & Gebhardt, K. Measures of location and scale for velocities in clusters of galaxies - A robust approach. *AJ* **100**, 32–46 (1990).
34. Bailin, J., Power, C., Norberg, P., Zaritsky, D. & Gibson, B. K. The anisotropic distribution of satellite galaxies. *MNRAS* **390**, 1133–1156 (2008).
35. Tempel, E., Guo, Q., Kipper, R. & Libeskind, N. I. The alignment of satellite galaxies and cosmic filaments: observations and simulations. *MNRAS* **450**, 2727–2738 (2015).
36. Wang, Y. *et al.* Probing the intrinsic shape and alignment of dark matter haloes using SDSS galaxy groups. *MNRAS* **385**, 1511–1522 (2008).
37. Silverman, B. W. *Density Estimation for Statistics and Data Analysis* (Chapman & Hall, London, 1986).
38. Trujillo, I. *et al.* Introducing the LBT Imaging of Galactic Halos and Tidal Structures (LIGHTS) survey. A preview of the low surface brightness Universe to be unveiled by LSST. *A&A* **654**, A40 (2021).
39. Martin, N. F., Collins, M. L. M., Longeard, N. & Tollerud, E. Current Velocity Data on Dwarf Galaxy NGC 1052-DF2 do not Constrain it to Lack Dark Matter. *ApJ* **859**, L5 (2018).
40. Trujillo, I. *et al.* A distance of 13 Mpc resolves the claimed anomalies of the galaxy lacking dark matter. *MNRAS* **486**, 1192–1219 (2019).
41. Monelli, M. & Trujillo, I. The TRGB Distance to the Second Galaxy “Missing Dark Matter”: Evidence for Two Groups of Galaxies at 13.5 and 19 Mpc in the Line of Sight of NGC 1052. *ApJ* **880**, L11 (2019).
42. van Dokkum, P. *et al.* A Revised Velocity for the Globular Cluster GC-98 in the Ultra Diffuse Galaxy NGC 1052-DF2. *Research Notes of the American Astronomical Society* **2**, 54 (2018).
43. Ogiya, G. Tidal stripping as a possible origin of the ultra diffuse galaxy lacking dark matter. *MNRAS* **480**, L106–L110 (2018).
44. Macciò, A. V. *et al.* Creating a galaxy lacking dark matter in a dark matter-dominated universe. *MNRAS* **501**, 693–700 (2021).
45. Jackson, R. A. *et al.* Dark matter-deficient dwarf galaxies form via tidal stripping of dark matter in interactions with massive companions. *MNRAS* **502**, 1785–1796 (2021).
46. Moreno, J. *et al.* Galaxies lacking dark matter produced by close encounters in a cosmological simulation. *Nature Astronomy* (2022).

47. Montes, M. *et al.* The Galaxy “Missing Dark Matter” NGC 1052-DF4 is Undergoing Tidal Disruption. *ApJ* **904**, 114 (2020).
48. Montes, M., Trujillo, I., Infante-Sainz, R., Monelli, M. & Borlaff, A. S. A Disk and No Signatures of Tidal Distortion in the Galaxy “Lacking” Dark Matter NGC 1052-DF2. *ApJ* **919**, 56 (2021).

Methods

Illustration of the collision scenario. The proposed scenario for the formation of DF2, DF4, and the other trail galaxies is shown in Fig. Extended Data Figure 1. As discussed in the main text, the scenario is a combination of the original idea that a bullet dwarf collision might have formed DF2 and/or DF4;¹¹ the results from subsequent hydrodynamical simulations, showing that multiple dark matter-free clumps can form in such a collision¹² and that the formation of massive star clusters is indeed promoted;¹³ and the independent finding that feedback from massive cluster formation in these conditions leads to a rapid expansion of the galaxies.¹⁸

Faint galaxy sample. We make use of a recently compiled catalog of low surface brightness objects in the NGC1052 field,²³ augmented by a catalog of all brighter galaxies with redshifts in the range $1000 \text{ km s}^{-1} < cz < 2000 \text{ km s}^{-1}$ that is provided in the same study. Ref 23 makes use of the publicly available DECaLS dataset.³¹ The galaxies were initially identified with a combination of automated techniques and visual inspection, with the majority coming from visual inspection. Their structural parameters were measured with GALFIT.³² We caution that the DECaLS dataset suffers from sky subtraction errors around low surface brightness galaxies, and that this may bias the size measurements. The main point of Fig. 4 is a relative comparison of the sizes of galaxies on and off the trail and this should be more robust than the absolute size measurements.

Velocity dispersion of the NGC1052 group. We use the latest compilation of radial velocities in the NGC1052 field²³ for an updated value of the velocity dispersion of the group. Table 2 of ref 23 contains 30 galaxies with redshifts $cz < 2000 \text{ km s}^{-1}$. Two were removed: DF2, as it is almost certainly not bound to the group, and LEDA4014647. LEDA4014647 was assigned a radial velocity of 1680 ± 60 in earlier SDSS releases (DR3) but its redshift was later revised to $z = 0.7$ (DR16). A visual inspection of the SDSS spectrum shows no clear features. Using the biweight estimator³³ we find a central velocity for the remaining 28 galaxies of $\langle cz \rangle = 1435 \pm 20 \text{ km s}^{-1}$ and a line of sight velocity dispersion of $\sigma = 115 \pm 15 \text{ km s}^{-1}$.

The Hough transform. We use the Hough transform to look for linear features in the galaxy distribution, a standard




Figure Extended Data Figure 1: **Illustration of the proposed formation scenario of DF2 and DF4.** Two gas rich dwarf galaxies experience a high speed encounter with a small impact parameter (top). Following previous studies^{11–13} the collisional gas gets stripped and shocked at closest approach, and forms stars at a prodigious rate with a bias toward massive clumps (second from top). The dark matter and previously formed stars are tidally distorted but continue ahead of the newly forming galaxies (third from top). Feedback in the absence of a dark matter halo leads to expansion of the newly formed galaxies.¹⁸ Most of the newly formed stellar mass is in two clumps, but several lower mass galaxies have also formed in the wake (bottom).

method for detecting lines in images.²⁴ The transform provides the number of galaxies along all possible directions, characterized by an angle and a distance from the center. A width and maximum linear extent have to be chosen; we use $\pm 30 \text{ kpc}$ ($\pm 5.2'$) for the width and $< 400 \text{ kpc}$ ($69'$) for the linear extent. Although the exact number of galaxies that the Hough transform associates with the linear feature depends on the precise limits that are chosen, the qualitative results are not sensitive to them. In Fig. 2b the orientation of the line is on the vertical axis and the offset with respect to NGC1052 on the horizontal axis.

Statistical significance of the trail. We use simulations to assess the probability that the alignment of the 11 galaxies arose by chance. We generate $N = 1000$ realizations of the (x, y) pairs by maintaining the angular distance from

NGC1052 for each pair and randomizing the angle. This procedure ensures that the density profile of the sample is maintained for all realizations. We then create Hough transforms for all realizations and determine how often the strongest linear feature contains ≥ 11 galaxies. We find that the probability of a chance alignment of ≥ 11 galaxies is 3 %.

This calculation assumes that galaxies are oriented randomly with respect to NGC1052, and does not take into account anisotropy associated with the filamentary structure of the cosmic web.^{34,35} Galaxy groups are generally not spherical but have an average projected axis ratio of 0.77.³⁶ We examined the large scale structure in the NGC1052 field using a recently compiled catalog of galaxies in this general area.²³ Selecting all low surface brightness galaxies that were identified in that study plus all bright galaxies with $cz < 2000 \text{ km s}^{-1}$ gives a sample of 72 probable group members. Their distribution is shown in Fig. Extended Data Figure 2. The smooth density field was calculated with the non-parametric kernel density estimator.³⁷ There is no evidence for large scale structure associated with the trail. In fact, there are no galaxies at all in the trail direction in the outskirts of the group, and the overall orientation of the group is perpendicular to the trail. The assumption of isotropy is therefore slightly conservative, in the sense that more galaxies will be scattered toward the line than away from it.




Figure Extended Data Figure 2: **Morphology of the NGC1052 group.** Distribution of 72 probable group members from a recent compilation of galaxies in the NGC1052 field.²³ Contours were derived with the non-parametric kernel density method.³⁷ The dashed line indicates the trail. There is no evidence that the trail is in the general direction of large scale structure in this field.

Finally, we note that the probability that there is a chance alignment *and* that it is a coincidence that both DF2 and DF4 are part of it is very low. This joint probability can be calculated directly for the isotropic case: of the 31 simulations that have ≥ 11 aligned galaxies only 6 have both DF2 and DF4 in the sample, corresponding to a combined probability of the observed arrangement of 0.6 %.

A twelfth low surface brightness dwarf galaxy on the trail. Visual inspection of the DECaLS imaging³¹ readily shows that there is a fairly prominent 12th galaxy that is part of the apparent trail. The object is SDSS J024007.01–081344.4; it was previously pointed out as a likely low luminosity group member with a central star cluster.³⁸ It is not in the objective catalog that we use for the main analysis.²³ This may be because of its redshift in the SDSS database (it is erroneously listed as a $z = 0.933$ active galactic nucleus) or because the light from the central cluster moved the object outside of the size and surface brightness criteria. We refer to the galaxy as DF9 as that was the catalog number in our initial Dragonfly catalog.¹ We do not use the galaxy in the objective analysis but we show its DECaLS image in Fig. 3. For convenience we provide the coordinates of all trail galaxies in Table 1.




Figure Extended Data Figure 3: **A candidate dark matter-dominated galaxy.** HST/ACS images¹ of DF7, which is located at the Western end of the galaxy trail, beyond DF4. At left is a color image generated from the F606W and F814W data. At right is a median-filtered version of the F606W image, with the arrow depicting the direction toward DF4. DF7 is a candidate for the dark matter-dominated remnant of one of the original galaxies,¹¹ given its location and its elongation in the direction of DF4.

Hubble Space Telescope imaging of the candidate dark galaxy DF7. DF7 is at one of the leading edges of the trail, “ahead” of DF4. The galaxy was observed with HST/ACS as part of an exploratory survey of Dragonfly-identified low surface brightness galaxies in several groups.¹ The observations constituted two orbits, one orbit in F606W and one orbit in F814W. In Fig. Extended Data Figure 3 we show the HST imaging at two different contrast levels. The galaxy is elongated and appears distorted, with the elongation in the direction of DF4. DF7’s apparent distortion, combined with its location at the leading edge of the trail, lead us to speculate that the galaxy is the highly dark matter-dominated rem-

Id	RA [deg]	DEC [deg]
RCP 32	40.6202	−8.3768
NGC 1052–DF2	40.4451	−8.4028
RCP 28	40.4215	−8.3475
RCP 26	40.2897	−8.2968
RCP 21	40.1200	−8.2434
NGC 1052–DF9	40.0292	−8.2290
RCP 17	39.9696	−8.2121
TA 21–12000	39.9139	−8.2285
NGC 1052–DF4	39.8128	−8.1160
NGC 1052–DF5	39.8028	−8.1408
LEDA 4014647	39.7021	−8.0494
NGC 1052–DF7	39.6241	−7.9257

Table 1: Coordinates (J2000) of the trail galaxies.

nant of one of the two progenitor galaxies. We note that DF7 may be largely disrupted in this interpretation: the observed¹ axis ratio is $b/a = 0.42$, but given the extreme foreshortening of the geometry the intrinsic axis ratio could be a stream-like $\sim 1 : 20$.

Other proposed scenarios. The joint formation of DF2 and DF4 in a single bullet dwarf event explains their lack of dark matter, large sizes, luminous and large globular clusters, striking similarity, large distance between them, large radial velocity difference, and the presence of a trail of other galaxies on the DF2 – DF4 axis. Here we briefly discuss other scenarios that have been proposed to explain the properties of DF2 and DF4.

Initially follow-up studies focused on possible errors in the measurements, either in the masses³⁹ or in the distances of the galaxies.^{40,41} However, with four independent velocity dispersion measurements^{3,9,10,42} (three for DF2 and one for DF4) and TRGB distances from extremely deep HST data^{15,16} these issues have now largely been settled.

Most astrophysical explanations center on the absence of dark matter only, and invoke some form of extreme tidal interaction (with NGC1052 or other galaxies) to strip the dark matter (along with a large fraction of the initial stellar population).^{43–46} These models do not explain the low metallicity of the galaxies, why there are two nearly identical objects in the same group, the newly discovered trail, or their overluminous and too-large globular clusters. The globular clusters, which have the same age (within the errors) as the diffuse light,²² show that the galaxies were *formed* in an unusual way and did not merely *evolve* in an unusual way. Besides the bullet scenario, the only model that explains the globular clusters is a study of star formation in galaxies that are in the tails of the scatter in the halo mass – stellar mass relation.^{18,19} This model has ad hoc initial conditions and does not account for the presence of two near-identical

galaxies, but the key aspects of it (the formation of luminous globular clusters in a compact configuration and the subsequent puffing up of the galaxies due to feedback) likely apply to the collision products in the bullet scenario (see main text).

It has recently been suggested that DF2 and DF4 are entirely unrelated, with DF4 being stripped of its dark matter by NGC1035, which is near it in projection, and DF2 a face-on disk galaxy with a normal dark matter content.^{47,48} The association of DF4 with NGC1035 is not seen in all datasets,²¹ and there is no compelling evidence that DF2 is a disk.⁷ Furthermore, the globular clusters and the trail remain unexplained, and there is the question of the likelihood that DF2 and DF4 have entirely different explanations but coincidentally share several otherwise-unique properties.

Data availability. The HST data for DF7 are available in the Mikulski Archive for Space Telescopes (MAST; <http://archive.stsci.edu>), under program ID 14644. The Legacy Survey data shown in Fig. 3 are available at <https://www.legacysurvey.org/>. All other data that support the findings of this study are available in published studies that are referenced in the text.

Code availability. We have made use of standard data analysis tools in the Python environment.

Acknowledgements A.J.R. was supported by National Science Foundation grant AST-1616710 and as a Research Corporation for Science Advancement Cottrell Scholar. J.M.D.K. acknowledges funding from the German Research Foundation (DFG) in the form of an Emmy Noether Research Group grant no. KR4801/1-1. J.M.D.K. and S.T.-G. acknowledge funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme via the ERC Starting Grant MUSTANG (grant agreement no. 714907). Support from Space Telescope Science Institute grants HST-GO-14644, HST-GO-15695, and HST-GO-15851 is gratefully acknowledged. Celeste van Dokkum created the illustration in Fig. Extended Data Figure 1.

Author Contributions P.v.D. led the analysis and wrote the manuscript. Z.S. performed the relative distance measurement (published in Shen et al. 2021) that is at the basis of the study. M.K. created Fig. 3 and measured structural parameters for DF5. S.T.-G. created an early version of Fig. 1b. All authors commented on the manuscript and aided in the interpretation.

Author Information The authors declare that they have no competing financial interests. Correspondence and requests for materials should be addressed to P.v.D. (email: pieter.vandokkum@yale.edu).