

Observational constraints on the fractal cosmology

Mahnaz Asghari ^{*1,2} and Ahmad Sheykhi ^{†1,2}

¹*Department of Physics, College of Sciences, Shiraz University, Shiraz 71454, Iran*

²*Biruni Observatory, College of Sciences, Shiraz University, Shiraz 71454, Iran*

Abstract

In this paper, we explore a fractal model of the universe proposed by Calcagni [JHEP03(2010)120] for a power-counting renormalizable field theory living in a fractal spacetime. Considering a timelike fractal profile, we constrain the fractal cosmology by observational data, including Planck cosmic microwave background (CMB), weak lensing, supernovae, baryon acoustic oscillations (BAO), and redshift-space distortions (RSD) data. Our results indicate that fractal cosmology is not capable to alleviate the tensions between local and global observational probes. However, according to our MCMC results, the proposed fractal model of the universe is in a reasonable consistency with standard model of cosmology.

1 Introduction

Recent observational data from type Ia supernovae (SNeIa) [1, 2], the cosmic microwave background (CMB) anisotropies [3, 4, 5], large scale structures [6, 7, 8], and baryon acoustic oscillations (BAO) [9, 10, 11], confirm Λ CDM as the concordance model of cosmology. On the other hand, Λ CDM model which is based on the theory of general relativity (GR), encounters some observational discrepancies, principally σ_8 and H_0 tensions. To be more specific, there is a disagreement between low-redshift determinations and Planck CMB measurements of matter perturbation amplitude, σ_8 [5, 12]. Moreover, direct measurements of Hubble constant are in significant tension with Planck observations [5, 13, 14, 15, 16]. So, inconsistencies between local and global data, motivate cosmologists to prob beyond Λ CDM model. Accordingly, it is suggested to consider some corrections on GR, describing as modified theory of gravity [17, 18, 19].

On the other hand, Calcagni proposed an effective quantum field theory which is power counting renormalizable and Lorentz invariant, living in a fractal universe [20, 21]. The fractal nature firstly introduced by Mandelbrot in 1983 [22] suggests conditional cosmological principle in fractal universe, where the universe appears the same from every galaxy. Thereafter, in 1986, Linde [23] propounded a model of an eternally existing chaotic inflationary universe, explaining a fractal cosmology. Accordingly, there are several investigations on the theory of fractal cosmology in literatures. Rasseem and Ahmed [24] in 1996 considered a nonhomogeneous cosmological model with a fractal distribution of matter which evolves to a homogeneous universe as time passes. The conditional cosmological principle in fractal cosmology is discussed in [25, 26]. [27] studies Multi-fractal geometry. Thermodynamics of the apparent horizon in a fractal universe is explored in [28]. In order to find more theoretical studies on fractal cosmology refer to e.g. [29, 30, 31, 32, 33].

*mahnaz.asghari@shirazu.ac.ir

†asheykhi@shirazu.ac.ir

Furthermore, it is interesting to explore fractal models with cosmological data as discussed in [34, 35, 36, 37, 38, 39] (also see [40] for a review). Correspondingly, in the present work we are going to investigate the fractal universe in background and perturbation levels, as well as studying observational constraints on parameters of fractal cosmology.

The paper is organized as follows. Section 2 is dedicated to field equations in a fractal universe. In section 3, we study the fractal model numerically, and further we constrain the model with current observational data in section 4. We summarize our results in section 5.

2 Field equations in a fractal universe

The total action in a fractal spacetime is given by [20]

$$S = \frac{1}{16\pi G} \int d\rho(x) \sqrt{-g} (R - 2\Lambda - \omega \partial_\mu v \partial^\mu v) + S_m, \quad (1)$$

where ω is the fractal parameter, v is the fractional function, and $d\rho(x)$ is a Lebesgue-Stieltjes measure. It is possible to derive field equations from action (1) similar to scalar-tensor theories. Thus, in a fractal universe we obtain [20]

$$R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} (R - 2\Lambda) + g_{\mu\nu} \frac{\square v}{v} - \frac{\nabla_\mu \nabla_\nu v}{v} + \omega \left(\frac{1}{2} g_{\mu\nu} \partial_\sigma v \partial^\sigma v - \partial_\mu v \partial_\nu v \right) = 8\pi G T_{\mu\nu}. \quad (2)$$

Furthermore, continuity equation in a fractal spacetime takes the form [20]

$$\nabla_\mu (v T_\nu^\mu) - \partial_\nu v \mathcal{L}_m = 0. \quad (3)$$

It should be noted that for $v = 1$, standard equations in GR will be recovered. We consider a fractal spacetime with the following flat Friedmann-Lemaître-Robertson-Walker (FLRW) metric in the synchronous gauge

$$ds^2 = a^2(\tau) \left(-d\tau^2 + (\delta_{ij} + h_{ij}) dx^i dx^j \right), \quad (4)$$

where $h_{ij} = \text{diag}(-2\eta, -2\eta, h + 4\eta)$, with scalar perturbations h and η . Here, we focus on a timelike fractal, then v is only time dependent given by

$$v = H_0^\beta \left(\frac{a}{a_0} \right)^\beta, \quad (5)$$

in which $\beta = 4(1 - \alpha)$ is the fractal dimension, and the parameter α ranges as $0 < \alpha \leq 1$. Then, considering the energy content of the universe as a perfect fluid with $T_{\mu\nu} = (\rho + p)u_\mu u_\nu + g_{\mu\nu}p$, field equations in background level take the form

$$H^2 \left(1 + \beta - \frac{1}{6} \omega H_0^{2\beta} \beta^2 \left(\frac{a}{a_0} \right)^{2\beta} \right) = \frac{8\pi G}{3} \sum_i \bar{\rho}_i, \quad (6)$$

$$(\beta + 2) \frac{H'}{a} + H^2 \left(3 + 2\beta + \beta^2 + \frac{1}{2} \omega H_0^{2\beta} \beta^2 \left(\frac{a}{a_0} \right)^{2\beta} \right) = -8\pi G \sum_i \bar{p}_i, \quad (7)$$

where a prime indicates a deviation with respect to the conformal time. It can be easily seen that, $\beta = 0$ restores field equations in standard cosmology. According to equation (6), total density parameter can be find as

$$\Omega_{\text{tot}} = 1 + \beta - \frac{1}{6} \omega H_0^{2\beta} \beta^2 \left(\frac{a}{a_0} \right)^{2\beta}, \quad (8)$$

where we have considered a universe filled with radiation (R), baryons (B), dark matter (DM) and cosmological constant (Λ). Also field equations to linear order of perturbations can be written as

$$\frac{a'}{a} \left(1 + \frac{1}{2}\beta\right) h' - 2k^2\eta = 8\pi G a^2 \sum_i \delta\rho_i, \quad (9)$$

$$k^2\eta' = 4\pi G a^2 \sum_i (\bar{\rho}_i + \bar{p}_i)\theta_i, \quad (10)$$

$$\frac{1}{2}h'' + 3\eta'' + \frac{a'}{a}(h' + 6\eta') \left(1 + \frac{1}{2}\beta\right) - k^2\eta = 0, \quad (11)$$

$$\frac{a'}{a} (2 + \beta) h' + h'' - 2k^2\eta = -24\pi G a^2 \sum_i \delta p_i. \quad (12)$$

In addition, regarding equation (3), conservation equations of fractal cosmology for i th component of the universe in background and perturbation levels become

$$\bar{\rho}'_i + (3 + \beta) \frac{a'}{a} (\bar{\rho}_i + \bar{p}_i) = 0, \quad (13)$$

$$\delta'_i = -\frac{a'}{a} (3 + \beta) \left[\delta_i (c_{si}^2 - w_i) + (c_{si}^2 - c_{ai}^2) \frac{a'}{a} (3 + \beta) (1 + w_i) \frac{\theta_i}{k^2} \right] - (1 + w_i)\theta_i - \frac{1}{2}(1 + w_i)h', \quad (14)$$

$$\theta'_i = \theta_i \left[-\frac{a'}{a} (4 + \beta) + \frac{a'}{a} (3 + \beta) (1 + w_i + c_{si}^2 - c_{ai}^2) \right] + \frac{k^2 c_{si}^2}{1 + w_i} \delta_i. \quad (15)$$

Then, choosing $\beta = 0$ would recover equations in concordance Λ CDM model.

Now that we have described main equations in fractal cosmology, it is possible to derive observational constraints on fractal model, using a modified version of the CLASS¹ code [41], and also applying an MCMC² approach via the MONTE PYTHON code [42, 43].

3 Numerical results

In this section, we modify the CLASS cosmological Boltzmann code according to the field equations of fractal model outlined in section 2. Correspondingly, we take into account the Planck 2018 data [5] (i.e. $\Omega_{B,0}h^2 = 0.02242$, $\Omega_{DM,0}h^2 = 0.11933$, $H_0 = 67.66 \frac{\text{km}}{\text{sMpc}}$, $A_s = 2.105 \times 10^{-9}$, and $\tau_{\text{reio}} = 0.0561$) in our numerical investigation.

In figure (1) we depict the CMB temperature power spectra in fractal cosmology compared to Λ CDM model. Upper panels display the CMB power spectra for different values of β , where we have considered $\omega = 0$. In lower panels the CMB temperature anisotropy diagrams are illustrated for different values of ω , while regarding $\beta = 0.04$. Furthermore, for a better comparison we have shown the relative ratio with respect to $\omega = 0$ in the bottom-right panel, which indicates that the effect of ω is not significant in CMB power spectra.

Figure (2) demonstrates the matter power spectra of fractal model in comparison with standard cosmological model. In left panel we see matter power spectra for different values of β , where ω is considered to be zero. Right panel shows the power spectra for different values of ω , in which we have fixed β to be 0.04. Considering this figure, fractal cosmology predicts a higher growth of structure which is in conflict with local probes of large scale structures. In addition, this plot explains that the fractal parameter ω has no remarkable influence on structure formation.

¹Cosmic Linear Anisotropy Solving System

²Markov Chain Monte Carlo




Figure 1: Upper panels show the CMB power spectra diagrams (left) and their relative ratio with respect to standard cosmological model (right) for different values of β , considering $\omega = 0$. Lower panels show the the CMB power spectra diagrams for different values of ω , considering $\beta = 0.04$, compared to Λ CDM model (left) and their relative ratio with respect to the case $\omega = 0$ (right).




Figure 2: Matter power spectra diagrams for different values of β compared to Λ CDM model, considering $\omega = 0$ (left), and analogous diagrams for different values of ω , where $\beta = 0.04$ (right).

Moreover, the evolution of Hubble parameter in fractal cosmology is depicted in figure (3). Accordingly, we can conclude that Hubble tension is aggravated in fractal model, and also the expansion history would not be affected by the fractal parameter ω .

4 Constraints from observational data

In this part, we employ the MCMC sampling package MONTE PYTHON to investigate observational constraints on fractal model. In this respect, we use the following set of cosmological parameters: $\{ 100 \Omega_{B,0} h^2, \Omega_{DM,0} h^2, 100 \theta_s, \ln(10^{10} A_s), n_s, \tau_{\text{reio}}, \beta \}$, where $\Omega_{B,0} h^2$ and $\Omega_{DM,0} h^2$ stand for the baryon and cold dark matter densities respectively, θ_s represents the ratio of the sound horizon to the angular diameter distance at decoupling, A_s in-




Figure 3: Hubble parameter in term of conformal time for different values of β compared to Λ CDM model, where $\omega = 0$ (left), and analogous diagrams for different values of ω , regarding $\beta = 0.04$ (right).

dicates the amplitude of the primordial scalar perturbation spectrum, n_s stands for the scalar spectral index, τ_{reio} is the optical depth to reionization, and β is the fractal dimension. We have also put constraints on four derived parameters which are reionization redshift (z_{reio}), the matter density parameter ($\Omega_{M,0}$), the Hubble constant (H_0), and the root-mean-square mass fluctuations on scales of $8 h^{-1}$ Mpc (σ_8). Additionally, preliminary numerical explorations consider the prior range $[0, 0.04]$ for the fractal dimension. It is worth mentioning that, since the fractal parameter ω is irrelevant to cosmological observable (based on numerical results in section 3), we fix ω to be zero in MCMC analysis.

The dataset we use to constrain fractal model includes the Planck likelihood with Planck 2018 data (containing high- l TT, TE, EE, low- l EE, low- l TT, and lensing) [5], the Planck-SZ likelihood for the Sunyaev-Zeldovich effect measured by Planck [44, 45], the CFHTLenS likelihood with the weak lensing data [46, 47], the Pantheon likelihood with the supernovae data [48], the BAO likelihood with the baryon acoustic oscillations data [49, 50], and the BAORS D likelihood for BAO and redshift-space distortions (RSD) measurements [51, 52].

The best fit and 1σ and 2σ confidence intervals for cosmological parameters of fractal model and also Λ CDM as the reference model, from "Planck + Planck-SZ + CFHTLenS + Pantheon + BAO + BAORS D" dataset are summarized in table (1). Correspondingly, contour plots for some selected parameters of fractal model compared to Λ CDM are displayed in figure (4). Considering the obtained constraints on fractal dimension, observational data detect no significant deviation from Λ CDM model. Therefore, according to MCMC analysis, it can be understood that a fractal model of the universe is in excellent agreement with standard model of cosmology.

Moreover, the Akaike information criterion (AIC) defined as $AIC = -2 \ln \mathcal{L}_{\text{max}} + 2K$ where \mathcal{L}_{max} is the maximum likelihood function and K indicates the number of free parameters [53, 54], results in $AIC_{(\Lambda\text{CDM})} = 3847.12$ and $AIC_{(\text{fractal})} = 3849.70$, then consequently $\Delta AIC = 2.58$. Thus, Λ CDM provides a better fit to observations, while fractal model is still allowed.

5 Conclusions

We have considered a fractal model of the universe introduced by Calcagni [20, 21], which is a power counting renormalizable and also Lorentz invariant model. Taking into account a timelike fractal, we have investigated the influence of fractal model on observables, mainly power spectra and Hubble constant, by using a modified version of the publicly available CLASS code. Considering our results, there is an enhancement in structure growth for non-zero values of fractal dimension β , being incompatible with low redshift structure formation. Moreover, the current value of Hubble

Table 1: Best fit values and 68% and 95% confidence limits for cosmological parameters from "Planck + Planck-SZ + CFHTLenS + Pantheon + BAO + BAORSD" data set for Λ CDM and fractal model.

parameter	Λ CDM		fractal model	
	best fit	68% & 95% limits	best fit	68% & 95% limits
$100 \Omega_{B,0} h^2$	2.261	$2.263^{+0.012+0.026}_{-0.013-0.025}$	2.274	$2.265^{+0.014+0.028}_{-0.015-0.027}$
$\Omega_{DM,0} h^2$	0.1163	$0.1164^{+0.00078+0.0015}_{-0.00079-0.0015}$	0.1164	$0.1168^{+0.00086+0.0017}_{-0.00093-0.0018}$
$100 \theta_s$	1.042	$1.042^{+0.00029+0.00055}_{-0.00026-0.00053}$	1.042	$1.042^{+0.00030+0.00055}_{-0.00028-0.00054}$
$\ln(10^{10} A_s)$	3.034	$3.024^{+0.010+0.023}_{-0.014-0.021}$	3.023	$3.025^{+0.0091+0.022}_{-0.014-0.021}$
n_s	0.9712	$0.9719^{+0.0036+0.0072}_{-0.0039-0.0074}$	0.9724	$0.9720^{+0.0036+0.0074}_{-0.0038-0.0072}$
τ_{reio}	0.05358	$0.04963^{+0.0041+0.010}_{-0.0074-0.0096}$	0.04706	$0.04919^{+0.0039+0.010}_{-0.0077-0.0092}$
β	—	—	0.0003854	$0.0005421^{+0.00012+0.00092}_{-0.00054-0.00054}$
z_{reio}	7.502	$7.084^{+0.50+1.0}_{-0.69-1.0}$	6.811	$7.046^{+0.49+1.0}_{-0.70-1.0}$
$\Omega_{M,0}$	0.2871	$0.2876^{+0.0043+0.0086}_{-0.0044-0.0086}$	0.2828	$0.2841^{+0.0056+0.010}_{-0.0050-0.012}$
$H_0 \left[\frac{\text{km}}{\text{s Mpc}} \right]$	69.56	$69.54^{+0.37+0.73}_{-0.36-0.71}$	70.16	$70.08^{+0.46+1.3}_{-0.68-1.1}$
σ_8	0.8079	$0.8044^{+0.0045+0.0096}_{-0.0051-0.0091}$	0.8063	$0.8092^{+0.0049+0.012}_{-0.0067-0.011}$




Figure 4: The one-dimensional posterior distribution and two-dimensional posterior contours with 68% and 95% confidence limits for the selected cosmological parameters of fractal model (orange) compared to Λ CDM (blue).

parameter decreases in fractal cosmology, which is inconsistent with local determinations of H_0 . Consequently, numerical results indicate that fractal cosmology has no positive impact on relieving

cosmological tensions.

Furthermore, we put constraints on the parameters of the fractal universe by utilizing current observations, chiefly Planck CMB, weak lensing, supernovae, BAO, and RSD data. According to our MCMC results, one can conclude that fractal model is in a reasonable consistency with the concordance Λ CDM model.

Acknowledgments

We thank Shiraz University Research Council.

References

- [1] Adam G. Riess et al. Observational Evidence from Supernovae for an Accelerating Universe and a Cosmological Constant. *The Astronomical Journal*, 116(3):1009–1038, September 1998.
- [2] S. Perlmutter et al. Measurements of Ω and Λ from 42 High-Redshift Supernovae. *The Astrophysical Journal*, 517(2):565–586, June 1999.
- [3] P. A. R. Ade et al. Planck 2013 results. XVI. Cosmological parameters. *Astronomy & Astrophysics*, 571:A16, November 2014.
- [4] P. A. R. Ade et al. Planck 2015 results. XIII. Cosmological parameters. *Astronomy & Astrophysics*, 594:A13, September 2016.
- [5] N. Aghanim et al. Planck 2018 results. VI. Cosmological parameters. *Astronomy & Astrophysics*, 641:A6, September 2020.
- [6] Max Tegmark et al. Cosmological parameters from SDSS and WMAP. *Phys. Rev. D*, 69:103501, May 2004.
- [7] Kevork Abazajian et al. The Second Data Release of the Sloan Digital Sky Survey. *The Astronomical Journal*, 128(1):502–512, July 2004.
- [8] Kevork Abazajian et al. The Third Data Release of the Sloan Digital Sky Survey. *The Astronomical Journal*, 129(3):1755–1759, March 2005.
- [9] H. Lampeitl et al. First-year Sloan Digital Sky Survey-II supernova results: consistency and constraints with other intermediate-redshift data sets. *Monthly Notices of the Royal Astronomical Society*, 401(4):2331–2342, 2010.
- [10] V. Sahni, A. Shafieloo, and A. A. Starobinsky. Model-independent Evidence for Dark Energy Evolution from Baryon Acoustic Oscillations. *The Astrophysical Journal Letters*, 793(2):L40, October 2014.
- [11] Will J. Percival et al. Baryon acoustic oscillations in the Sloan Digital Sky Survey Data Release 7 galaxy sample. *Monthly Notices of the Royal Astronomical Society*, 401(4):2148–2168, February 2010.
- [12] S. W. Allen, R. W. Schmidt, A. C. Fabian, and H. Ebeling. Cosmological constraints from the local X-ray luminosity function of the most X-ray-luminous galaxy clusters. *Monthly Notices of the Royal Astronomical Society*, 342(1):287, 2003.
- [13] Adam G. Riess et al. A 2.4% DETERMINATION OF THE LOCAL VALUE OF THE HUBBLE CONSTANT. *The Astrophysical Journal*, 826(1):56, 2016.
- [14] Adam G. Riess et al. New parallaxes of galactic cepheids from spatially scanning the Hubble space telescope: Implications for the Hubble constant. *The Astrophysical Journal*, 855(2):136, 2018.
- [15] Adam G. Riess, Stefano Casertano, Wenlong Yuan, Lucas M. Macri, and Dan Scolnic. Large magellanic cloud cepheid standards provide a 1% foundation for the determination of the Hubble constant and stronger evidence for physics beyond Λ CDM. *The Astrophysical Journal*, 876(1):85, 2019.

- [16] Adam G. Riess et al. Cosmic distances calibrated to 1% precision with Gaia EDR3 parallaxes and Hubble space telescope photometry of 75 milky way cepheids confirm tension with Λ CDM. *The Astrophysical Journal*, 908(1):L6, 2021.
- [17] S. Tsujikawa. *Modified Gravity Models of Dark Energy*, pages 99–145. Springer Berlin Heidelberg, Berlin, Heidelberg, 2010.
- [18] Shin’ichi Nojiri and Sergei D. Odintsov. Unified cosmic history in modified gravity: From F(R) theory to Lorentz non-invariant models. *Physics Reports*, 505(2):59–144, 2011.
- [19] Timothy Clifton, Pedro G. Ferreira, Antonio Padilla, and Constantinos Skordis. Modified gravity and cosmology. *Physics Reports*, 513(1):1–189, 2012. Modified Gravity and Cosmology.
- [20] Gianluca Calcagni. Quantum field theory, gravity and cosmology in a fractal universe. *Journal of High Energy Physics*, 03:120, 2010.
- [21] Gianluca Calcagni. Fractal universe and quantum gravity. *Phys. Rev. Lett.*, 104:251301, Jun 2010.
- [22] B. B. Mandelbrot. *The fractal geometry of nature*. W. H. Freeman and Comp., New York, 3 edition, 1983.
- [23] A.D. Linde. Eternally existing self-reproducing chaotic inflationary universe. *Physics Letters B*, 175(4):395–400, 1986.
- [24] M. Rassem and E. Ahmed. On Fractal Cosmology. *Astrophysical Letters and Communications*, 35:311, January 1996.
- [25] A. K. Mittal and T. R. Seshadri. Fractals and the large-scale structure in the universe. *Resonance*, 7:39, Apr 2002.
- [26] A. K. Mittal and D. Lohiya. Fractal dust model of the universe based on Mandelbrot’s conditional cosmological principle and general theory of relativity. arXiv:astro-ph/0206300, 2002.
- [27] Gianluca Calcagni. Geometry and field theory in multi-fractional spacetime. *Journal of High Energy Physics*, 2012:65, Jan 2012.
- [28] Ahmad Sheykhi, Zeinab Teimoori, and Bin Wang. Thermodynamics of fractal universe. *Physics Letters B*, 718(4):1203–1207, Jan 2013.
- [29] K. Karami, Mubasher Jamil, S. Ghaffari, and K. Fahimi. Holographic, new agegraphic, and ghost dark energy models in fractal cosmology. *Canadian Journal of Physics*, 91(10):770–776, Oct 2013.
- [30] Surajit Chattopadhyay, Antonio Pasqua, and Sudipto Roy. A study on some special forms of Holographic Ricci Dark Energy in fractal universe. *ISRN High Energy Physics*, 2013:251498, Oct 2013.
- [31] Gianluca Calcagni. Multi-scale gravity and cosmology. *Journal of Cosmology and Astroparticle Physics*, 2013(12):041–041, Dec 2013.
- [32] Sayani Maity and Ujjal Debnath. Co-existence of modified Chaplygin gas and other dark energies in the framework of fractal universe. *International Journal of Theoretical Physics*, 55:2668, May 2016.
- [33] Gianluca Calcagni. Multiscale spacetimes from first principles. *Phys. Rev. D*, 95:064057, Mar 2017.
- [34] Gianluca Calcagni, Sachiko Kuroyanagi, and Shinji Tsujikawa. Cosmic microwave background and inflation in multi-fractional spacetimes. *Journal of Cosmology and Astroparticle Physics*, 2016(08):039–039, Aug 2016.
- [35] Dipanjana Das, Sourav Dutta, Abdulla Al Mamon, and Subenoy Chakraborty. Does fractal universe describe a complete cosmic scenario ? *The European Physical Journal C*, 78:849, Oct 2018.
- [36] Ehsan Sadri, Martiros Khurshudyan, and Surajit Chattopadhyay. An interacting new holographic dark energy in the framework of fractal cosmology. *Astrophysics and Space Science*, 363:230, Oct 2018.
- [37] Ujjal Debnath and Kazuharu Bamba. Parametrizations of dark energy models in the background of general non-canonical scalar field in D-dimensional fractal universe. *The European Physical Journal C*, 79:722, Aug 2019.

- [38] S. Ghaffari, E. Sadri, and A. H. Ziaie. Tsallis holographic dark energy in fractal universe. *Modern Physics Letters A*, 35(14):2050107, Mar 2020.
- [39] Gianluca Calcagni and Antonio De Felice. Dark energy in multifractional spacetimes. *Phys. Rev. D*, 102:103529, Nov 2020.
- [40] Gianluca Calcagni. Multifractional theories: An updated review. *Modern Physics Letters A*, 36(14):2140006, 2021.
- [41] Diego Blas, Julien Lesgourgues, and Thomas Tram. The Cosmic Linear Anisotropy Solving System (CLASS). Part II: Approximation schemes. *Journal of Cosmology and Astroparticle Physics*, 2011(07):034, 2011.
- [42] Benjamin Audren, Julien Lesgourgues, Karim Benabed, and Simon Prunet. Conservative Constraints on Early Cosmology: an illustration of the MONTE PYTHON cosmological parameter inference code. *JCAP*, 1302:001, 2013.
- [43] Thejs Brinckmann and Julien Lesgourgues. MontePython 3: boosted MCMC sampler and other features. arXiv:1804.07261, 2018.
- [44] P. A. R. Ade et al. Planck 2015 results. XXIV. Cosmology from Sunyaev-Zeldovich cluster counts. *Astron. Astrophys.*, 594:A24, 2016.
- [45] P. A. R. Ade et al. Planck 2013 results. XX. Cosmology from Sunyaev-Zeldovich cluster counts. *Astron. Astrophys.*, 571:A20, 2014.
- [46] Martin Kilbinger et al. CFHTLenS: Combined probe cosmological model comparison using 2D weak gravitational lensing. *Monthly Notices of the Royal Astronomical Society*, 430(3):2200, 2013.
- [47] Catherine Heymans et al. CFHTLenS tomographic weak lensing cosmological parameter constraints: Mitigating the impact of intrinsic galaxy alignments. *Monthly Notices of the Royal Astronomical Society*, 432(3):2433, 2013.
- [48] D. M. Scolnic et al. The complete light-curve sample of spectroscopically confirmed SNe Ia from pan-STARRS1 and cosmological constraints from the combined pantheon sample. *The Astrophysical Journal*, 859(2):101, 2018.
- [49] Florian Beutler et al. The 6DF Galaxy Survey: Baryon Acoustic Oscillations and the Local Hubble Constant. *Monthly Notices of the Royal Astronomical Society*, 416(4):3017–3032, 2011.
- [50] Lauren Anderson et al. The clustering of galaxies in the SDSS-III Baryon Oscillation Spectroscopic Survey: Baryon Acoustic Oscillations in the Data Releases 10 and 11 Galaxy Samples. *Monthly Notices of the Royal Astronomical Society*, 441(1):24–62, 2014.
- [51] Shadab Alam et al. The clustering of galaxies in the completed SDSS-III baryon oscillation spectroscopic survey: cosmological analysis of the DR12 galaxy sample. *Monthly Notices of the Royal Astronomical Society*, 470(3):2617–2652, 2017.
- [52] Manuel A. Buen-Abad, Martin Schmaltz, Julien Lesgourgues, and Thejs Brinckmann. Interacting Dark Sector and Precision Cosmology. *JCAP*, 1801(01):008, 2018.
- [53] Hirotugu Akaike. A new look at the statistical model identification. *IEEE Transactions on Automatic Control*, 19(6):716–723, 1974.
- [54] K.P. Burnham and D.R. Anderson. *Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach*. Springer Verlag, 2002.