

NEW EXAMPLES OF G_2 -STRUCTURES WITH DIVERGENCE-FREE TORSION

AGUSTÍN GARRONE

ABSTRACT. Interest in Riemannian manifolds with holonomy equal to the exceptional Lie group G_2 have spurred extensive research in geometric flows of G_2 -structures defined on seven-dimensional manifolds in recent years. Among many possible geometric flows, the so-called *isometric flow* has the distinctive feature of preserving the underlying metric induced by that G_2 -structure, so it can be used to evolve a G_2 -structure to one with the smallest possible torsion in a given metric class. This flow is built upon the divergence of the full torsion tensor of the flowing G_2 -structures in such a way that its critical points are precisely G_2 -structures with divergence-free torsion. In this article we study three large families of pairwise non-equivalent non-closed left-invariant G_2 -structures defined on simply connected solvable Lie groups previously studied in [16] and compute the divergence of their full torsion tensor, obtaining that it is identically zero in all cases.

CONTENTS

1. Introduction	1
2. G_2 -structures and their geometry	2
2.1. Definitions	2
2.2. Left-invariant structures	4
3. G_2 -structure over the Lie algebras $\mathfrak{g}_{A,B,C}$	4
3.1. Definitions and remarks	4
3.2. Formulas for the torsion forms	5
3.3. Auxiliary formulas	7
3.4. Formulas for the Levi-Civita connection and the Ricci operator	9
3.5. Formula for the divergence of the torsion	10
4. Skew-symmetric case	12
5. Diagonal case	12
6. Antidiagonal case	13
References	14

1. INTRODUCTION

A G_2 -structure on a 7-dimensional smooth manifold is a *positive* smooth 3-form $\varphi \in \Omega^3(M)$, by which we mean that for all $p \in M$ it can be written as

$$(1) \quad \varphi_p = e^{127} + e^{347} + e^{567} + e^{135} - e^{146} - e^{236} - e^{245}$$

with respect to an ordered basis $\{e_1, \dots, e_7\}$ of $T_p M$. The existence of a G_2 -structure imposes several topological restrictions on M beyond dimension, for example that M be orientable and spin ([17], [14], [3], or [11, Proposition 10.1.6]). Notably, they are non-trivially related to the exceptional Lie group G_2 , the normed algebra of the octonions, generalizations of vector cross products, and even theoretical physics (most notably, M -theory); for a non-exhaustive review of these features, see [7], [6], [17, Section 2], [1], and [8, Section 1].

A striking result of G_2 -geometry is that a G_2 -structure naturally determines a Riemannian metric g and a volume form vol in M ; see Section 2. In particular, they induce a Hodge star operator $\star : \Omega^k(M) \rightarrow \Omega^{7-k}(M)$, $0 \leq k \leq 7$. This induced Riemannian structure allows us to consider the Levi-Civita derivative $\nabla \varphi$ of φ , which is an interesting object in its own right. Of special interest is the *torsion-free* case, which is

characterized by the condition $\nabla\varphi = 0$. Torsion-free G_2 -structures are important for a variety of reasons, which include that the holonomy of M with respect to g is contained in G_2 *precisely* when $\nabla\varphi = 0$ (see [11, Proposition 10.1.3]), which in turn is relevant because of the famous Berger's classification theorem (see [2]). Historically, torsion-free G_2 -structures with holonomy equal to G_2 were thought to be non-existent for decades until the first non-compact and compact examples were found respectively by R. Bryant in 1987 (see [3]) and D. Joyce in 1996 (see [11]). Since then, geometric flows techniques have become a feature in the pursuit of new examples. Different geometric flows of G_2 -structures have been studied, notably the *Laplacian flow of closed G_2 -structures* (i.e., such that $d\varphi = 0$) (see [4]) and the *Laplacian coflow of coclosed G_2 -structures* (i.e., such that $d\star\varphi = 0$) (see [15]). The most relevant flow to us is the *isometric flow*, first defined by S. Grigorian (see [9] and [10]) and independently by S. Dwivedi, P. Giannotis, and S. Karigiannis (see [5]), which is governed by the equation

$$(2) \quad \frac{\partial\varphi(t)}{\partial t} = \iota_{\operatorname{div} T(t)}(\star\varphi(t)).$$

Several notions are introduced in equation (2):

- The map ι , called *contraction*, simply given by $\iota_X(\eta)(Y, Z, W) = \eta(X, Y, Z, W)$ for a 4-form $\eta \in \Omega^4(M)$ and vector fields $X, Y, Z, W \in \mathfrak{X}(M)$.
- The *full torsion tensor* T of φ , which is the only smooth tensor field of type $(1, 1)$ on M such that (see Section 2).

$$(3) \quad \nabla_X\varphi = \iota_{T(X)}(\star\varphi), \quad X \in \mathfrak{X}(M).$$

- The *divergence* of T , which is the smooth vector field $\operatorname{div} T$ given by

$$(4) \quad g(\operatorname{div} T, E_j) = \sum_{i=1}^7 (\nabla_{E_i} T)(E_i, E_j),$$

where $\{E_i \mid 1 \leq i \leq 7\}$ is an arbitrary local orthonormal frame of (M, g) .

It is clear that that G_2 -structures such that $\operatorname{div} T = 0$ are critical points of (2); such structures are said to have *divergence-free full torsion tensor*, or *divergence-free torsion* for short.

The main motivation behind this article is to find new examples of G_2 -structures with divergence-free full torsion tensor. Our research led us to left-invariant G_2 -structures defined on simply connected solvable Lie groups $G_{A,B,C}$, introduced in Section 3, whose definition depends on the choice of a triple $A, B, C \in \mathfrak{sl}(4, \mathbb{R})$ of pairwise-commuting 4×4 matrices. Recall from [16] that these groups admit lattices when A, B, C are chosen to be linearly independent, so there are compact versions of the examples. Three broad families of triples are considered: the *skew-symmetric case*, in which $A, B, C \in \mathfrak{sl}(4, \mathbb{R})$ are skew-symmetric; the *diagonal case*, in which $A, B, C \in \mathfrak{sl}(4, \mathbb{R})$ are diagonal (which is proven to be equivalent to the *symmetric case*, see Proposition 5.1); and the *antidiagonal case*, in which $A, B, C \in \mathfrak{sl}(4, \mathbb{R})$ are antidiagonal (see equation (70)). These three cases are non-equivalent under mild restrictions on A, B, C , as computations regarding the Ricci operator of the underlying Riemannian structure of $(\mathfrak{g}_{A,B,C}, \varphi)$ readily show (see Proposition 3.12). The main result can be stated as follows:

Theorem 1.1. *Left-invariant G_2 -structures given by (1) defined on Lie groups $G_{A,B,C}$ with $A, B, C \in \mathfrak{sl}(4, \mathbb{R})$ pairwise-commuting 4×4 real matrices that are all skew-symmetric, all symmetric, or all antidiagonal have a divergence-free full torsion tensor T ; furthermore, those G_2 -structures are non-closed and pairwise non-equivalent under mild restrictions on A, B , and C .*

2. G_2 -STRUCTURES AND THEIR GEOMETRY

2.1. Definitions. A smooth 3-form φ defined on a 7-dimensional differentiable manifold M is called a G_2 -structure if it is *positive*; that is, if at every point $p \in M$ it can be written as

$$(5) \quad \varphi_p = e^{127} + e^{347} + e^{567} + e^{135} - e^{146} - e^{236} - e^{245}$$

with respect to an ordered basis $\{e_1, \dots, e_7\}$ of $T_p M$. Note that this definition is slightly different but equivalent to other more frequently used alternatives (e.g., [3], [4], [13], [14], among others). Two G_2 -structures (M, φ) and (M', φ') are said to be *equivalent* if there is a diffeomorphism $f : M \rightarrow M'$ such that $\varphi = f^*\varphi'$; such a map is called an *equivalence*.

One of the most astounding and well-known facts about G_2 -structures is that they canonically induce a Riemannian metric g and a volume form vol by

$$(6) \quad g(X, Y) \text{vol} = \frac{1}{6} \iota_X(\varphi) \wedge \iota_Y(\varphi) \wedge \varphi, \quad X, Y \in \mathfrak{X}(M);$$

here, the map ι is simply defined as $\iota_X(\varphi)(W, Z) := \varphi(X, W, Z)$ for $W, Z \in \mathfrak{X}(M)$. A proof of this can be found in [12, Proposition 2.3.1]. This induces a Hodge star operator $\star : \Omega^k(M) \rightarrow \Omega^{7-k}(M)$, $0 \leq k \leq 7$, on M ; in particular, it allows us to define the dual 4-form of φ , $\psi := \star\varphi$, which can readily be shown to satisfy

$$(7) \quad \psi_p = e^{3456} + e^{1256} + e^{1234} - e^{2467} + e^{2357} + e^{1457} + e^{1367}$$

with respect to the ordered basis $\{e_1, \dots, e_7\}$.

Some of the most important aspects of the geometry that can be done in presence of G_2 -structures are encoded in the *torsion forms*, which are the unique differential forms $\tau_i \in \Omega^i(M)$ with $i = 0, 1, 2, 3$ such that

$$(8) \quad d\varphi = \tau_0\psi + 3\tau_1 \wedge \varphi + \star\tau_3, \quad d\psi = 4\tau_1 \wedge \psi + \star\tau_2;$$

this is proven in [13, Theorem 2.3] and in [4, Proposition 1]. Equivalently, we have

$$(9) \quad \begin{aligned} \tau_0 &= \frac{1}{7} \star(d\varphi \wedge \varphi), & \tau_1 &= -\frac{1}{12} \star(\star d\varphi \wedge \varphi), \\ \tau_2 &= -\star d\psi + 4\star(\tau_1 \wedge \psi), & \tau_3 &= \star d\varphi - \tau_0\varphi - 3\star(\tau_1 \wedge \varphi); \end{aligned}$$

see [20] or [19] for a careful derivation of this equivalence. The vanishing of some torsion forms determines different kinds of G_2 -structures, for example:

- $d\varphi = 0$ if and only if $\tau_0 = \tau_1 = \tau_3 = 0$, in which case we say φ is *closed*.
- $d\psi = 0$ if and only if $\tau_1 = \tau_2 = 0$, in which case we say φ is *coclosed*.
- $d\varphi = 0$ and $d\psi = 0$ if and only if $\tau_i = 0$ for all $i = 0, 1, 2, 3$, in which case we say φ is *torsion-free*.

Recall that the *torsion* of a G_2 -structure is $\nabla\varphi$, the covariant derivative of φ with respect to the metric g induced by φ . It can be shown that there is a tensor field $T \in \mathfrak{T}^{(1,1)}(M)$ such that

$$(10) \quad \nabla_X\varphi = \iota_{T(X)}(\psi), \quad X \in \mathfrak{X}(M);$$

we refer to [14, Theorem 4.44] for a proof. We call T the *full torsion tensor* of φ . As the name suggests, T is directly related to the torsion forms $\tau_i \in \Omega^i(M)$, $i = 0, 1, 2, 3$, as it can be shown that

$$(11) \quad T(X, Y) = \frac{1}{4}\tau_0g(X, Y) - \iota_{\tau_1}(\varphi)(X, Y) - \frac{1}{2}\tau_2(X, Y) - \tau_{27}(X, Y), \quad X, Y \in \mathfrak{X}(M),$$

where $\iota_{\tau_1}(\varphi)$ is the contraction of φ by the smooth vector field $\tilde{\tau}_1$ uniquely determined by the condition

$$(12) \quad g(\tilde{\tau}_1, X) = \tau_1(X), \quad X \in \mathfrak{X}(M),$$

which is called τ_1 by abuse of language, and τ_{27} is the symmetric 2-tensor given by

$$(13) \quad \tau_{27}(X, Y) := \star(\iota_X(\varphi) \wedge \iota_Y(\varphi) \wedge \tau_3), \quad X, Y \in \mathfrak{X}(M);$$

a proof can be found in [13, Theorem 2.27]. We see that a G_2 -structure is torsion-free if and only if $T = 0$, which is also equivalent to $\nabla\varphi = 0$. This is often called Fernández-Gray theorem; see [6].

Interest in torsion-free G_2 -structures is widespread, as these structures have many desirable properties, among them having holonomy contained in the exceptional Lie group G_2 and being Ricci-flat (see [11, Propositions 10.1.3 and 10.1.5]), and are of historical relevance, as the famous Berger's classification theorem points out (see [2]). Geometric flows techniques have been introduced in this area of research, most notably in [3] and in [15]. Of special relevance to us is the *isometric flow*, first defined by S. Grigorian (see [9] and [10]) and independently by S. Dwivedi, P. Giannotis, and S. Karigiannis (see [5]), which is governed by the equation

$$(14) \quad \frac{\partial\varphi(t)}{\partial t} = \iota_{\text{div } T(t)}(\psi(t));$$

here the *divergence* $\text{div } T$ of the full torsion tensor T is the vector field defined as

$$(15) \quad g(\text{div } T, E_j) = \sum_{i=1}^7 (\nabla_{E_i} T)(E_i, E_j),$$

where $\{E_i \mid 1 \leq i \leq 7\}$ is an arbitrary local orthonormal frame with respect to the induced metric g . It is clear that G_2 -structures such that $\text{div } T = 0$ are critical points of (14); such structures are said to have

divergence-free full torsion tensor, or *divergence-free torsion* for short. It is known from [9, Theorem 4.3] that $\operatorname{div} T = 0$ if $d\varphi = 0$. The G_2 -structures considered in this article are non-closed in general.

2.2. Left-invariant structures. We will restrict our attention to left-invariant G_2 -structures defined on Lie groups G , for it is usual to come across simpler formulas in this setting. An important observation in this regard is that $\nabla_X Y \in \operatorname{Lie}(G)$ for all $X, Y \in \operatorname{Lie}(G)$, which in turn allows us to write the Koszul formula for the Levi-Civita connection as

$$(16) \quad g(\nabla_X Y, Z) = \frac{1}{2} (g([X, Y], Z) - g([Y, Z], X) + g([Z, X], Y)), \quad X, Y, Z \in \operatorname{Lie}(G).$$

It is natural then to consider the map $U : \operatorname{Lie}(G) \times \operatorname{Lie}(G) \rightarrow \operatorname{Lie}(G)$ determined by

$$(17) \quad g(U(X, Y), Z) = \frac{1}{2} (g([Z, X], Y) - g([Y, Z], X)), \quad X, Y, Z \in \operatorname{Lie}(G).$$

Note in particular that $U(X, Y) = U(Y, X)$ for all $X, Y \in \operatorname{Lie}(G)$. In terms of this map, equation (16) can be written more simply as

$$(18) \quad \nabla_X Y = \frac{[X, Y]}{2} + U(X, Y), \quad X, Y \in \operatorname{Lie}(G);$$

equation (18) will come in handy. It is worth mentioning that U is identically zero when the metric g is bi-invariant; that is, g is both left and right-invariant.

Turning again to G_2 -structures, we first recall that two left-invariant G_2 -structures (G, φ) and (G', φ') defined on Lie groups G and G' are called *equivariantly equivalent* if there is a Lie group isomorphism $f : G \rightarrow G'$ such that $f^* \varphi' = \varphi$; such a map is called an *equivariant equivalence*. We also recall that any 3-form defined on the Lie algebra $\operatorname{Lie}(G)$ of a Lie group G by

$$(19) \quad \varphi = e^{127} + e^{347} + e^{567} + e^{135} - e^{146} - e^{236} - e^{245} \in \Lambda^3(\operatorname{Lie}(G)^*),$$

with respect to some ordered basis $\{e_1, \dots, e_7\}$ on $\operatorname{Lie}(G)$ gives rise to a left-invariant G_2 -structure on G , typically denoted by the same name; moreover, φ induces on $\operatorname{Lie}(G)$ an inner product $\langle \cdot, \cdot \rangle$ and a volume form vol on $\operatorname{Lie}(G)$ given by a formula akin to (6), which can in turn be extended to corresponding left-invariant structures on the Lie group G . Similar considerations apply to other G_2 -related paraphernalia, most importantly the full torsion tensor T , whose divergence satisfies

$$(20) \quad \langle \operatorname{div}(T), e_j \rangle = - \sum_{1 \leq i \leq 7} T(\nabla_{e_i} e_i, e_j) - \sum_{1 \leq i \leq 7} T(e_i, \nabla_{e_i} e_j)$$

when restricted to $\operatorname{Lie}(G)$, where $\{e_1, \dots, e_7\}$ is an ordered orthonormal basis of $\operatorname{Lie}(G)$. This prompts us to work entirely at the Lie algebra level.

3. G_2 -STRUCTURE OVER THE LIE ALGEBRAS $\mathfrak{g}_{A,B,C}$

3.1. Definitions and remarks. Let $A, B, C \in \mathfrak{sl}(4, \mathbb{R})$ be pairwise-commuting 4×4 real matrices. Over a real seven-dimensional vector space with basis $\{e_1, \dots, e_7\}$, which we denote $\mathfrak{g}_{A,B,C}$, we put

$$(21) \quad \mathfrak{a} := \operatorname{span}\{e_1, e_2, e_7\}, \quad \mathfrak{n} := \operatorname{span}\{e_3, e_4, e_5, e_6\},$$

and define a bilinear, anticommutative product $[\cdot, \cdot] : \mathfrak{g}_{A,B,C} \times \mathfrak{g}_{A,B,C} \rightarrow \mathfrak{g}_{A,B,C}$ by setting $[\mathfrak{a}, \mathfrak{a}] = 0$, $[\mathfrak{n}, \mathfrak{n}] = 0$, and

$$(22) \quad [e_7, v] = Av, \quad [e_1, v] = Bv, \quad [e_2, v] = Cv$$

for all $v \in \mathfrak{n}$, so as to make $(\mathfrak{g}_{A,B,C}, [\cdot, \cdot])$ a real 7-dimensional Lie algebra.

These algebras were first defined in [16], and they constitute a generalization of almost abelian Lie algebras. Note that the Jacobi condition is actually equivalent to the fact that A, B, C are pairwise-commuting. By definition, $\mathfrak{a} = \operatorname{span}\{e_1, e_2, e_7\}$ is an abelian subalgebra and $\mathfrak{n} = \operatorname{span}\{e_3, e_4, e_5, e_6\}$ is an abelian ideal of $\mathfrak{g}_{A,B,C}$; furthermore, it is readily verified that $\mathfrak{n}$ is the nilradical of $\mathfrak{g}_{A,B,C}$ only if no linear combination of A, B, C is a nilpotent matrix. For all choices of A, B, C we have that $\mathfrak{g}_{A,B,C}$ is a solvable Lie algebra and, as a consequence of tracelessness, that $\mathfrak{g}_{A,B,C}$ is unimodular. These last two facts are exploited in [16] to determine equivalence classes among left-invariant G_2 -structures defined on Lie groups whose Lie algebra is isomorphic to $\mathfrak{g}_{A,B,C}$, and we will use this result to establish that there are many non-equivalent examples of G_2 -structures with divergence-free full torsion tensor defined on $\mathfrak{g}_{A,B,C}$, or equivalently, left-invariant G_2 -structures with divergence-free full torsion tensor defined on the simply connected Lie group $G_{A,B,C}$ such that its Lie algebra equals $\mathfrak{g}_{A,B,C}$.

Under the further restriction that $A, B, C \in \mathfrak{sl}(4, \mathbb{R})$ are linearly independent and that they simultaneously diagonalize over $\mathbb{R}$, in which case we call the triple *compatible*, all Lie algebras $\mathfrak{g}_{A,B,C}$ are isomorphic to each other, and the corresponding Lie groups $G_{A,B,C}$ are isomorphic to G_J , the only completely solvable unimodular Lie group in the classification appearing in [18]. More on this topic as well as their relation to G_2 -structures can be found in [16]. We will not be dealing with compatible triples in the rest of this article.

We consider over $\mathfrak{g}_{A,B,C}$ the G_2 -structure defined with respect to the basis $\{e_1, \dots, e_7\}$ as

$$(23) \quad \varphi := e^{127} + e^{347} + e^{567} + e^{135} - e^{146} - e^{236} - e^{245} \in \Lambda^3(\mathfrak{g}_{A,B,C}^*),$$

which, as we know, induces an inner product $\langle \cdot, \cdot \rangle$ and a volume form vol on $\mathfrak{g}_{A,B,C}$ according to

$$(24) \quad g(X, Y) \text{vol} = \frac{1}{6} \iota_X(\varphi) \wedge \iota_Y(\varphi) \wedge \varphi, \quad X, Y \in \mathfrak{g}_{A,B,C}.$$

These definitions ensure that $\{e_1, \dots, e_7\}$ is orthonormal with respect to $\langle \cdot, \cdot \rangle$, as well as $\mathfrak{a}$ and $\mathfrak{n}$ are orthogonal complements to each other. Accordingly, this structure induces a Hodge star operator $\star : \Lambda^k(\mathfrak{g}_{A,B,C}^*) \rightarrow \Lambda^{7-k}(\mathfrak{g}_{A,B,C}^*)$, which in turn allows us to define the dual 4-form $\psi := \star \varphi$ as

$$(25) \quad \psi := e^{3456} + e^{1256} + e^{1234} - e^{2467} + e^{2357} + e^{1457} + e^{1367}.$$

3.2. Formulas for the torsion forms. In order to simplify future formulas and calculations, we introduce the 2-forms on $\mathfrak{g}_{A,B,C}$ given by

$$(26) \quad \omega_7 := e^{34} + e^{56}, \quad \omega_1 := e^{35} - e^{46}, \quad \omega_2 := -e^{36} - e^{45},$$

$$(27) \quad \bar{\omega}_7 := e^{34} - e^{56}, \quad \bar{\omega}_1 := e^{35} + e^{46}, \quad \bar{\omega}_2 := -e^{36} + e^{45}.$$

We interpret the 2-forms given by equations (26) and (27) as elements of $\Lambda^2(\mathfrak{n}^*)$. We now list some immediate properties of these 2-forms in the following result for future reference. We call $g_{\mathfrak{n}}$, $\text{vol}_{\mathfrak{n}}$, and $\star_{\mathfrak{n}}$ the metric, volume form, and Hodge star operator induced on $\mathfrak{n}$ by the corresponding structures in $\mathfrak{g}_{A,B,C}$.

Lemma 3.1. *Let $\mathfrak{n} = \text{span}\{e_3, e_4, e_5, e_6\}$ be the nilpotent ideal of $\mathfrak{g}_{A,B,C}$, and let $\omega_7, \omega_1, \omega_2, \bar{\omega}_7, \bar{\omega}_1$, and $\bar{\omega}_2 \in \Lambda^2(\mathfrak{n}^*)$ be given as in equations (26) and (27). Then*

- (i) $\varphi = e^{127} + \omega_7 \wedge e^7 + \omega_1 \wedge e^1 + \omega_2 \wedge e^2$.
- (ii) $\psi = e^{3456} + \omega_7 \wedge e^{12} + \omega_1 \wedge e^{27} - \omega_2 \wedge e^{17}$.
- (iii) $\star_{\mathfrak{n}} \omega_i = \omega_i$ and $\star_{\mathfrak{n}} \bar{\omega}_j = -\bar{\omega}_j$ for $i, j = 1, 2, 7$.
- (iv) $\omega_i \wedge \omega_j = \omega_i \wedge \bar{\omega}_j = \bar{\omega}_i \wedge \bar{\omega}_j = 0$ for all $i \neq j$.
- (v) the following equations hold:

$$(28) \quad \omega_7 \wedge \omega_7 = \omega_1 \wedge \omega_1 = \omega_2 \wedge \omega_2 = 2e^{3456},$$

$$(29) \quad \bar{\omega}_7 \wedge \bar{\omega}_7 = \bar{\omega}_1 \wedge \bar{\omega}_1 = \bar{\omega}_2 \wedge \bar{\omega}_2 = -2e^{3456}.$$

- (vi) the following equations hold:

$$(30) \quad e^{34} = \frac{\bar{\omega}_7 + \omega_7}{2}, \quad e^{35} = \frac{\bar{\omega}_1 + \omega_1}{2}, \quad e^{36} = -\frac{\bar{\omega}_2 + \omega_2}{2},$$

$$(31) \quad e^{45} = \frac{\bar{\omega}_2 - \omega_2}{2}, \quad e^{46} = \frac{\bar{\omega}_1 - \omega_1}{2}, \quad e^{56} = -\frac{\bar{\omega}_7 - \omega_7}{2}.$$

- (vii) $\mathfrak{B} := \{\bar{\omega}_7, \bar{\omega}_1, \bar{\omega}_2, \omega_7, \omega_1, \omega_2\}$ is an orthogonal basis of $\Lambda^2(\mathfrak{n}^*)$ with respect to the metric induced by $g_{\mathfrak{n}}$, with every element having norm equal to $\sqrt{2}$.

Let $\theta : \mathfrak{sl}(4, \mathbb{R}) \rightarrow \text{End}(\Lambda^2(\mathfrak{n}^*))$ denote the natural representation of $\mathfrak{sl}(4, \mathbb{R})$ on $\Lambda^2(\mathfrak{n}^*)$, given by

$$(32) \quad (\theta(D)\eta)(\cdot, \cdot) := -\eta(D\cdot, \cdot) - \eta(\cdot, D\cdot).$$

It is clear from the definition that $\theta(M)e^i = -\sum_{j=3}^6 m_{ij}e^j$, where $M \in \mathfrak{sl}(4, \mathbb{R})$ is a matrix with coefficients $[m_{ij}]$, $i, j = 3, 4, 5, 6$. Moreover,

$$(33) \quad \begin{aligned} \theta(M)\omega_7 := & -(m_{33} + m_{44})e^{34} + (m_{63} - m_{45})e^{35} - (m_{46} + m_{53})e^{36} \\ & + (m_{64} + m_{35})e^{45} + (m_{36} - m_{54})e^{46} - (m_{55} + m_{66})e^{56}, \end{aligned}$$

$$(34) \quad \begin{aligned} \theta(M)\omega_1 := & -(m_{54} + m_{63})e^{34} - (m_{33} - m_{55})e^{35} + (m_{43} - m_{56})e^{36} \\ & + (m_{65} - m_{34})e^{45} + (m_{44} + m_{66})e^{46} + (m_{45} + m_{36})e^{56}, \end{aligned}$$

$$(35) \quad \begin{aligned} \theta(M)\omega_2 := & (m_{64} - m_{53})e^{34} + (m_{43} + m_{65})e^{35} + (m_{33} + m_{66})e^{36} \\ & + (m_{44} + m_{55})e^{45} + (m_{56} + m_{54})e^{46} + (m_{35} - m_{46})e^{56}; \end{aligned}$$

see [22, Remark 2.8]. We recall the following result.

Proposition 3.2. [22, Theorems 2.3 and 2.4] *For each set of pairwise-commuting matrices $A, B, C \in \mathfrak{sl}(4, \mathbb{R})$, the following formulas hold in $(\mathfrak{g}_{A,B,C}, \varphi)$:*

$$(36) \quad d\varphi = (\theta(B)\omega_7 - \theta(A)\omega_1) \wedge e^{17} + (\theta(C)\omega_7 - \theta(A)\omega_2) \wedge e^{27} + (\theta(B)\omega_2 - \theta(C)\omega_1) \wedge e^{12},$$

$$(37) \quad \star d\varphi = (\theta(B^\top)\omega_7 - \theta(A^\top)\omega_1) \wedge e^2 - (\theta(C^\top)\omega_7 - \theta(A^\top)\omega_2) \wedge e^1 - (\theta(B^\top)\omega_2 - \theta(C^\top)\omega_1) \wedge e^7,$$

$$(38) \quad d\psi = (\theta(A)\omega_7 + \theta(B)\omega_1 + \theta(C)\omega_2) \wedge e^{127},$$

$$(39) \quad \star d\psi = -(\theta(A^\top)\omega_7 + \theta(B^\top)\omega_1 + \theta(C^\top)\omega_2).$$

Remark 3.3. The formulas in Proposition 3.2 show that the G_2 -structures $(\mathfrak{g}_{A,B,C}, \varphi)$ are neither closed nor coclosed in general. In fact, $(\mathfrak{g}_{A,B,C}, \varphi)$ is closed if and only if

$$(40) \quad \theta(B)\omega_7 = \theta(A)\omega_1, \quad \theta(C)\omega_7 = \theta(A)\omega_2, \quad \theta(B)\omega_2 = \theta(C)\omega_1,$$

and $(\mathfrak{g}_{A,B,C}, \varphi)$ is coclosed if and only if

$$(41) \quad \theta(A)\omega_7 + \theta(B)\omega_1 + \theta(C)\omega_2 = 0.$$

Proposition 3.2 can be used to give explicit formulas for the torsion forms τ_0, τ_1, τ_2 , and τ_3 of $(\mathfrak{g}_{A,B,C}, \varphi)$ in terms of the action of θ via equations (9). This has been done in [22, Proposition 2.7] with the help of equations (33) to (35); we recall the ugly-looking result¹.

Proposition 3.4. [22, Proposition 2.7] *For each set of pairwise-commuting matrices $A = [a_{ij}]$, $B = [b_{ij}]$, $C = [c_{ij}] \in \mathfrak{sl}(4, \mathbb{R})$ for $i, j \in \{3, 4, 5, 6\}$, the following formulas hold in $(\mathfrak{g}_{A,B,C}, \varphi)$:*

$$(42) \quad \tau_0 = \frac{2}{7}(a_{46} - a_{64} + a_{53} - a_{35} + b_{35} - b_{53} + b_{64} - b_{46} + c_{54} - c_{45} + c_{63} - c_{36}),$$

$$(43) \quad \begin{aligned} \tau_1 = & -\frac{1}{12}(a_{36} - a_{63} + a_{45} - a_{54} + c_{56} - c_{65} + c_{34} - c_{43})e^1 \\ & -\frac{1}{12}(a_{64} - a_{46} + a_{35} - a_{53} + b_{43} - b_{34} + b_{65} - b_{56})e^2 \\ & -\frac{1}{12}(b_{63} - b_{36} + b_{54} - b_{45} + c_{46} - c_{64} + c_{53} - c_{35})e^7, \end{aligned}$$

$$(44) \quad \begin{aligned} \tau_2 = & \frac{1}{3}(b_{45} - b_{54} + b_{36} - b_{63} + c_{35} - c_{53} + c_{64} - c_{46})e^{12} \\ & + \frac{1}{3}(a_{64} - a_{46} + a_{35} - a_{53} + b_{65} - b_{56} + b_{43} - b_{34})e^{17} \\ & + \frac{1}{3}(a_{54} - a_{45} + a_{63} - a_{36} + c_{65} - c_{56} + c_{43} - c_{34})e^{27} \\ & + \frac{1}{3}(-3a_{33} - 3a_{44} + 2c_{46} - 2c_{35} - 2b_{45} - 2b_{36} - c_{53} + c_{64} - b_{63} - b_{54})e^{34} \\ & + \frac{1}{3}(-2a_{54} + 2a_{36} + 2c_{56} + 2c_{34} + a_{63} - a_{45} + c_{65} + c_{43} - 3b_{55} - 3b_{33})e^{35} \\ & + \frac{1}{3}(-2a_{64} - 2a_{35} - 2b_{65} + 2b_{34} - a_{46} - a_{53} - b_{56} + b_{43} - 3c_{44} - 3c_{55})e^{36} \\ & + \frac{1}{3}(a_{64} + a_{35} + b_{65} - b_{34} + 2a_{46} + 2a_{53} + 2b_{56} - 2b_{43} + 3c_{55} + 3c_{44})e^{45} \\ & + \frac{1}{3}(-a_{54} + a_{36} + c_{56} + c_{34} + 2a_{63} - 2a_{45} + 2c_{65} + 2c_{43} - 3b_{33} - 3b_{55})e^{46} \\ & + \frac{1}{3}(3a_{33} + 3a_{44} - c_{46} + c_{35} + b_{45} + b_{36} + 2c_{53} - 2c_{64} + 2b_{63} + 2b_{54})e^{56}, \end{aligned}$$

¹There is no explicit formula for the torsion form τ_3 in [22, Proposition 2.7], although clear guidelines for obtaining it are given; we have τ_3 following those with the help of symbolic and numeric computing software.

$$\begin{aligned}
(45) \quad \tau_3 = & -\frac{2}{7}(a_{56} + c_{54} + a_{34} - c_{36} - a_{65} + c_{63} - a_{43} - c_{45} + b_{64} + b_{35} - b_{53} - b_{46})e^{127} \\
& -\frac{1}{4}(-b_{65} - b_{43} + b_{56} + b_{34} - a_{64} + a_{53} - 3a_{46} + 3a_{35} - c_{33} - c_{44})e^{134} \\
& +\frac{1}{7}(5a_{56} + 5c_{54} + 5a_{34} - 5c_{36} + 2a_{65} - 2c_{63} + 2a_{43} + 2c_{45} - 2b_{64} - 2b_{35} + 2b_{53} + 2b_{46})e^{135} \\
& -\frac{1}{4}(-b_{63} + b_{45} - b_{54} + b_{36} - c_{53} - c_{46} - 3c_{64} - 3c_{35} + a_{55} + a_{44})e^{136} \\
& +\frac{1}{4}(b_{63} - b_{45} + b_{54} - b_{36} - 3c_{53} - 3c_{46} - c_{64} - c_{35} + 4a_{44} + 4a_{55})e^{145} \\
& +\frac{1}{7}(2a_{56} + 2c_{54} + 2a_{34} - 2c_{36} + 5a_{65} - 5c_{63} + 5a_{43} + 5c_{45} + 2b_{64} + 2b_{35} - 2b_{53} - 2b_{46})e^{146} \\
& +\frac{1}{4}(b_{65} + b_{43} - b_{56} - b_{34} - 3a_{64} + 3a_{53} - a_{46} + a_{35} - 4c_{44} - 4c_{33})e^{156} \\
& +\frac{1}{4}(-c_{56} + c_{43} + c_{65} - c_{34} + a_{63} + a_{54} + 3a_{45} + 3a_{36} - 4b_{33} - 4b_{44})e^{234} \\
& +\frac{1}{4}(b_{63} - b_{45} - 3b_{54} + 3b_{36} + c_{53} + c_{46} - c_{64} - c_{35} + 4a_{33} + 4a_{55})e^{235} \\
& -\frac{1}{7}(-2a_{56} - 2c_{54} + 5a_{34} + 2c_{36} - 5a_{65} - 2c_{63} + 2a_{43} + 2c_{45} + 5b_{64} + 5b_{35} + 2b_{53} + 2b_{46})e^{236} \\
& +\frac{1}{7}(-5a_{56} + 2c_{54} + 2a_{34} - 2c_{36} - 2a_{65} + 2c_{63} + 5a_{43} - 2c_{45} + 2b_{64} + 2b_{35} + 5b_{53} + 5b_{46})e^{245} \\
& +\frac{1}{4}(3b_{63} - 3b_{45} - b_{54} + b_{36} - c_{53} - c_{46} + c_{64} + c_{35} + 4a_{55} + 4a_{33})e^{246} \\
& -\frac{1}{4}(c_{56} - c_{43} - c_{65} + c_{34} + 3a_{63} + 3a_{54} + a_{45} + a_{36} - 4b_{44} - 4b_{33})e^{256} \\
& -\frac{1}{7}(2a_{56} + 2c_{54} + 2a_{34} + 5c_{36} - 2a_{65} + 2c_{63} - 2a_{43} + 5c_{45} + 2b_{64} - 5b_{35} - 2b_{53} + 5b_{46})e^{347} \\
& -\frac{1}{4}(b_{65} + b_{43} + 3b_{56} + 3b_{34} + a_{64} - a_{53} - a_{46} + a_{35} + 4c_{33} + 4c_{55})e^{357} \\
& -\frac{1}{4}(c_{56} - c_{43} + 3c_{65} - 3c_{34} - a_{63} - a_{54} + a_{45} + a_{36} - 4b_{55} - 4b_{44})e^{367} \\
& -\frac{1}{4}(-3c_{56} + 3c_{43} - c_{65} + c_{34} - a_{63} - a_{54} + a_{45} + a_{36} + 4b_{44} + 4b_{55})e^{457} \\
& +\frac{1}{4}(-3b_{65} - 3b_{43} - b_{56} - b_{34} + a_{64} - a_{53} - a_{46} + a_{35} - 4c_{55} - 4c_{33})e^{467} \\
& -\frac{1}{7}(2a_{56} - 5c_{54} + 2a_{34} - 2c_{36} - 2a_{65} - 5c_{63} - 2a_{43} - 2c_{45} - 5b_{64} + 2b_{35} + 5b_{53} - 2b_{46})e^{567}.
\end{aligned}$$

The next result is a straightforward application of (i) of Lemma 3.1, where we put

$$(46) \quad k_1 := -\frac{1}{12}(a_{36} - a_{63} + a_{45} - a_{54} + c_{56} - c_{65} + c_{34} - c_{43}),$$

$$(47) \quad k_2 := -\frac{1}{12}(a_{64} - a_{46} + a_{35} - a_{53} + b_{43} - b_{34} + b_{65} - b_{56}),$$

$$(48) \quad k_7 := -\frac{1}{12}(b_{63} - b_{36} + b_{54} - b_{45} + c_{46} - c_{64} + c_{53} - c_{35}).$$

so that $\tau_1 = k_1 e^1 + k_2 e^2 + k_7 e^7$.

Corollary 3.5. *For each set of pairwise-commuting matrices $A, B, C \in \mathfrak{sl}(4, \mathbb{R})$, the following formula holds in $(\mathfrak{g}_{A,B,C}, \varphi)$:*

$$(49) \quad \iota_{\tau_1}(\varphi) = k_1(e^{27} + \omega_1) + k_2(-e^{17} + \omega_2) + k_7(e^{12} + \omega_7).$$

Remark 3.6. We note for future reference that Proposition 3.4 and Corollary 3.5 imply that

$$\iota_{\tau_1}(\varphi), \tau_2 \in \text{span}\{e^{12}, e^{17}, e^{27}, e^{34}, e^{35}, e^{36}, e^{45}, e^{46}, e^{56}\},$$

as well as

$$\tau_3 \in \text{span}\{e^{127}, e^{134}, e^{135}, e^{136}, e^{145}, e^{146}, e^{156}, e^{234}, e^{235}, e^{236}, e^{245}, e^{246}, e^{256}, e^{347}, e^{357}, e^{367}, e^{457}, e^{467}, e^{567}\}.$$

3.3. Auxiliary formulas. The following technical results aim to provide a better understanding of the symmetric 2-tensor τ_{27} . All of them can be established by straightforward computations. We assume only that $A, B, C \in \mathfrak{sl}(4, \mathbb{R})$ are pairwise-commuting and that φ is given as in equation (23).

Lemma 3.7. *The following formulas hold in $(\mathfrak{g}_{A,B,C}, \varphi)$:*

$$(50) \quad \iota_{e_j}(\varphi) = \begin{cases} e^{27} + e^{35} - e^{46}, & j = 1. \\ -e^{17} - e^{36} - e^{45}, & j = 2. \\ e^{47} - e^{15} + e^{26}, & j = 3. \\ -e^{37} + e^{16} + e^{25}, & j = 4. \\ e^{67} + e^{13} - e^{24}, & j = 5. \\ -e^{57} - e^{14} - e^{23}, & j = 6. \\ e^{12} + e^{34} + e^{56}, & j = 7. \end{cases}$$

$$(51) \quad \iota_{e_1}(\varphi) \wedge \iota_{e_j}(\varphi) = \begin{cases} +e^{1257} - e^{3457} + e^{2356} - e^{1456}, & j = 3. \\ -e^{1267} + e^{1356} + e^{3467} + e^{2456}, & j = 4. \\ -e^{1237} + e^{3567} + e^{2345} - e^{1346}, & j = 5. \\ +e^{1247} + e^{1345} - e^{4567} + e^{2346}, & j = 6. \end{cases}$$

$$(52) \quad \iota_{e_2}(\varphi) \wedge \iota_{e_j}(\varphi) = \begin{cases} -e^{1267} + e^{3467} - e^{1356} - e^{2456}, & j = 3. \\ -e^{1257} + e^{2356} + e^{3457} - e^{1456}, & j = 4. \\ +e^{1247} - e^{2346} - e^{4567} - e^{1345}, & j = 5. \\ +e^{1237} - e^{3567} - e^{1346} + e^{2345}, & j = 6. \end{cases}$$

$$(53) \quad \iota_{e_7}(\varphi) \wedge \iota_{e_j}(\varphi) = \begin{cases} +e^{1247} - e^{1345} + e^{2346} + e^{4567}, & j = 3. \\ -e^{1237} + e^{1346} + e^{2345} - e^{3567}, & j = 4. \\ +e^{1267} + e^{3467} + e^{1356} - e^{2456}, & j = 5. \\ -e^{1257} - e^{3457} - e^{1456} - e^{2356}, & j = 6. \end{cases}$$

$$(54) \quad \frac{1}{2}\iota_{e_j}(\varphi) \wedge \iota_{e_j}(\varphi) = \begin{cases} +e^{1457} - e^{2467} + e^{1256}, & j = 3. \\ +e^{1367} + e^{2357} + e^{1256}, & j = 4. \\ +e^{1367} - e^{2467} + e^{1234}, & j = 5. \\ +e^{1457} + e^{2357} + e^{1234}, & j = 6. \end{cases}$$

$$(55) \quad \iota_{e_j}(\varphi) \wedge \iota_{e_{(9-j)}}(\varphi) = \begin{cases} -e^{2347} + e^{1246} + e^{2567} + e^{1235}, & j = 3, 6. \\ -e^{2347} - e^{1246} + e^{2567} - e^{1235}, & j = 4, 5. \end{cases}$$

Corollary 3.8. *Define*

$$\mathfrak{S}_1 := \text{span}\{e^{127}, e^{134}, e^{135}, e^{136}, e^{145}, e^{146}, e^{156}, e^{234}, e^{235}, e^{236}, e^{245}, e^{246}, e^{256}, e^{347}, e^{357}, e^{367}, e^{457}, e^{467}, e^{567}\},$$

$$\mathfrak{S}_2 := \text{span}\{e^{134}, e^{136}, e^{145}, e^{156}, e^{234}, e^{235}, e^{246}, e^{256}, e^{357}, e^{367}, e^{457}, e^{467}\},$$

$$\mathfrak{S}_3 := \text{span}\{e^{127}, e^{135}, e^{136}, e^{145}, e^{146}, e^{234}, e^{235}, e^{236}, e^{245}, e^{246}, e^{256}, e^{347}, e^{367}, e^{457}, e^{567}\}.$$

The following formulas hold in $(\mathfrak{g}_{A,B,C}, \varphi)$ for all $\chi_1 \in \mathfrak{S}_1$, $\chi_2 \in \mathfrak{S}_2$, and $\chi_3 \in \mathfrak{S}_3$:

$$(56) \quad \iota_{e_k}(\varphi) \wedge \iota_{e_i}(\varphi) \wedge \chi_1 = 0, \quad k = 1, 2, 7, \quad 3 \leq i \leq 6,$$

$$(57) \quad \iota_{e_n}(\varphi) \wedge \iota_{e_n}(\varphi) \wedge \chi_2 = 0, \quad 3 \leq n \leq 6,$$

$$(58) \quad \iota_{e_m}(\varphi) \wedge \iota_{e_{(9-m)}}(\varphi) \wedge \chi_3 = 0, \quad m \leq l \leq 6,$$

Following the definition of τ_{27} given in equation (13), the next result is an immediate consequence of Remark 3.6 and equation (56) from Corollary 3.8.

Corollary 3.9. *The following formulas hold in $(\mathfrak{g}_{A,B,C}, \varphi)$:*

$$(59) \quad \tau_{27}(e_k, e_i) = 0 \quad \text{for all } k = 1, 2, 7 \text{ and } 3 \leq i \leq 6,$$

3.4. Formulas for the Levi-Civita connection and the Ricci operator. We now turn to the goal of finding the Levi-Civita connection of the underlying Riemannian structure in $(\mathfrak{g}_{A,B,C}, \varphi)$. Special notation is to be introduced in order to state the result more briefly:

- Given $Z \in \mathfrak{a}$ of the form $Z = \lambda_1 e_1 + \lambda_2 e_2 + \lambda_7 e_7$ for some $\lambda_1, \lambda_2, \lambda_7 \in \mathbb{R}$, we define

$$(60) \quad M_Z := \lambda_1 B + \lambda_2 C + \lambda_7 A.$$

Note that if $Z \in \mathfrak{a}$ and $W \in \mathfrak{n}$ then $[Z, W] = M_Z W$.

- We call $S(M)$ and $A(M)$ the symmetric and antisymmetric parts of a matrix $M \in \mathfrak{gl}(4, \mathbb{R})$ respectively; that is,

$$(61) \quad S(M) := \frac{M+M^\top}{2}, \quad A(M) := \frac{M-M^\top}{2}.$$

- We define

$$(62) \quad D^l = B\delta_{l1} + C\delta_{l2} + A\delta_{l7},$$

where δ_{ij} is the Kronecker delta and $A, B, C \in \mathfrak{sl}(4, \mathbb{R})$ are the matrices appearing in the definition of $\mathfrak{g}_{A,B,C}$. Note that we are **not** performing a summation over the index l .

Proposition 3.10. *Let $A, B, C \in \mathfrak{sl}(4, \mathbb{R})$ be pairwise-commuting. The Levi-Civita connection ∇ of the underlying Riemannian structure in $(\mathfrak{g}_{A,B,C}, \varphi)$ is given by*

$$(63) \quad \nabla_X Y = \begin{cases} 0 & X \in \mathfrak{a}, Y \in \mathfrak{a}. \\ A(M_X)Y & X \in \mathfrak{a}, Y \in \mathfrak{n}. \\ -S(M_Y)X & X \in \mathfrak{n}, Y \in \mathfrak{a}. \\ \sum_{l=1,2,7} \langle S(D^l)X, Y \rangle e_l & X \in \mathfrak{n}, Y \in \mathfrak{n}. \end{cases}$$

Proof. We consider four separate cases.

Let $X, Y \in \mathfrak{a}$. Then $[X, Y] = 0$, as $\mathfrak{a}$ is an abelian subalgebra. Therefore, equation (17) implies that $U(X, Y)$ has no components in $\mathfrak{a}$. In addition, $U(X, Y)$ has no components in $\mathfrak{n}$, for if $Z \in \mathfrak{n}$ then $[Z, X] \in \mathfrak{n}$ and $[Z, Y] \in \mathfrak{n}$ because $\mathfrak{n}$ is an ideal, and then

$$2\langle U(X, Y), Z \rangle = \langle [Z, X], Y \rangle + \langle [Z, Y], X \rangle = 0,$$

where the last equality holds because $\mathfrak{a}$ and $\mathfrak{n}$ are orthogonal complements in $\mathfrak{g}_{A,B,C}$. Thus, equation (18) implies that $\nabla_X Y = 0$ if $X, Y \in \mathfrak{a}$.

Let $X \in \mathfrak{a}$ and $Y \in \mathfrak{n}$. As equation (18) implies that $\nabla_X Y = \frac{M_X Y}{2} + U(X, Y)$, it suffices to show that $U(X, Y) = -\frac{M_X^\top Y}{2}$. To such end, let $W \in \mathfrak{a}$ and $Z \in \mathfrak{n}$ be arbitrary. Note that $[W, X] = 0$ and $[Z, Y] = 0$ because both $\mathfrak{a}$ and $\mathfrak{n}$ are abelian. In addition, as $[W, Y] \in \mathfrak{n}$ for $\mathfrak{n}$ is an ideal, it follows that $\langle [W, Y], X \rangle = 0$ due to $\mathfrak{a}$ and $\mathfrak{n}$ being orthogonal complements in $\mathfrak{g}_{A,B,C}$. Thus, equation (17) implies that

$$2\langle U(X, Y), W \rangle = \langle [W, X], Y \rangle + \langle [W, Y], X \rangle = 0;$$

this establishes that $U(X, Y) \in \mathfrak{n}$. Furthermore, as $[Z, X] = -M_X Z$, the same equations show that

$$2\langle U(X, Y), Z \rangle = \langle [Z, X], Y \rangle + \langle [Z, Y], X \rangle = -\langle M_X Z, Y \rangle = -\langle M_X^\top Y, Z \rangle.$$

We can write the last equality $\langle U(X, Y), Z \rangle = -\frac{\langle M_X^\top Y, Z \rangle}{2}$. Combining the last two results, we get $U(X, Y) = -\frac{M_X^\top Y}{2}$.

Let $X \in \mathfrak{n}$ and $Y \in \mathfrak{a}$. From the torsion-free property of ∇ and the previous paragraph, we see that

$$\nabla_X Y = \nabla_Y X + [X, Y] = A(M_Y)X - M_Y X = -S(M_Y)X.$$

Let $X, Y \in \mathfrak{n}$. Then $[X, Y] = 0$, as $\mathfrak{n}$ is abelian. Therefore, equation (17) implies that $U(X, Y)$ has no components in $\mathfrak{n}$. Let $Z \in \mathfrak{a}$, e.g. $Z = e_7$. We see from equation (17) that

$$2\langle U(X, Y), e_7 \rangle = \langle [e_7, X], Y \rangle + \langle [e_7, Y], X \rangle = \langle AX, Y \rangle + \langle AY, X \rangle = \langle (A + A^\top)X, Y \rangle;$$

i.e., that $\langle U(X, Y), e_7 \rangle = \langle S(A)X, Y \rangle$. Doing the same for $Z = e_1$ y $Z = e_2$, we find that

$$U(X, Y) = \langle S(B)X, Y \rangle e_1 + \langle S(C)X, Y \rangle e_2 + \langle S(A)X, Y \rangle e_7. \quad \square$$

Remark 3.11. A similar result to Proposition 3.10 is valid in any Lie algebra $\mathfrak{g}$ that splits as an orthogonal direct sum $\mathfrak{a} \oplus \mathfrak{n}$ with respect to an inner product $\langle \cdot, \cdot \rangle$, where $\mathfrak{a}$ is an abelian subalgebra and $\mathfrak{n}$ is an abelian ideal.

We recall the following result for further reference.

Proposition 3.12. [22, Section 2.5] *Let $A, B, C \in \mathfrak{sl}(4, \mathbb{R})$ be pairwise-commuting. The Ricci operator $\text{Ric} : \mathfrak{g}_{A,B,C} \rightarrow \mathfrak{g}_{A,B,C}$ of the underlying Riemannian structure in $(\mathfrak{g}_{A,B,C}, \varphi)$ is given by*

$$(64) \quad \text{Ric}|_{\mathfrak{a} \times \mathfrak{n}} = 0,$$

$$(65) \quad \text{Ric}|_{\mathfrak{n} \times \mathfrak{n}} = \frac{1}{2}([A, A^\top] + [B, B^\top] + [C, C^\top]),$$

$$(66) \quad \text{Ric}|_{\mathfrak{a} \times \mathfrak{a}} = - \begin{pmatrix} \text{tr}(S(A)^2) & \text{tr}(S(A)B) & \text{tr}(S(A)C) \\ \text{tr}(S(A)B) & \text{tr}(S(B)^2) & \text{tr}(S(B)C) \\ \text{tr}(S(A)C) & \text{tr}(S(B)C) & \text{tr}(S(C)^2) \end{pmatrix}.$$

3.5. Formula for the divergence of the torsion. Remark 3.6, Corollary 3.9, and Proposition 3.10 allow us to give a simple expression for the divergence of the full torsion tensor of $(\mathfrak{g}_{A,B,C}, \varphi)$, given by equation (20).

Theorem 3.13. *Let $A, B, C \in \mathfrak{sl}(4, \mathbb{R})$ be pairwise-commuting. The divergence $\text{div}(T)$ of the full torsion tensor T of $(\mathfrak{g}_{A,B,C}, \varphi)$ satisfies*

$$(67) \quad \langle \text{div}(T), e_j \rangle = \begin{cases} -\sum_{3 \leq n \leq 6} (D^j)_{nn} \tau_{27}(e_n, e_n) + \sum_{3 \leq i \neq l \leq 6} S(D^j)_{il} \tau_{27}(e_i, e_l) & j = 1, 2, 7, \\ 0 & 3 \leq j \leq 6, \end{cases}$$

with D^j as in equation (62).

Proof. Recall from equation (20) that

$$\langle \text{div}(T), e_j \rangle = - \sum_{1 \leq i \leq 7} T(\nabla_{e_i} e_i, e_j) - \sum_{1 \leq i \leq 7} T(e_i, \nabla_{e_i} e_j).$$

We claim that Proposition 3.10 and A, B, C being traceless ensure that the first term is identically zero. Indeed, for $1 \leq j \leq 7$, we have

$$\begin{aligned} \sum_{1 \leq i \leq 7} T(\nabla_{e_i} e_i, e_j) &= \sum_{3 \leq i \leq 6} T(\nabla_{e_i} e_i, e_j) = \sum_{3 \leq i \leq 6} T \left(\sum_{k=1,2,7} \langle S(D^k) e_i, e_i \rangle e_k, e_j \right) \\ &= \sum_{3 \leq i \leq 6} \sum_{k=1,2,7} S(D^k)_{ii} T(e_k, e_j) = \sum_{k=1,2,7} \text{tr}(S(D^k)) T(e_k, e_j) \\ &= \sum_{k=1,2,7} \text{tr}(D^k) T(e_k, e_j) = \sum_{k=1,2,7} 0 \cdot T(e_k, e_j) = 0. \end{aligned}$$

On the other hand, Proposition 3.10 ensures that for $j = 1, 2, 7$ we have

$$\begin{aligned} \sum_{1 \leq i \leq 7} T(e_i, \nabla_{e_i} e_j) &= \sum_{3 \leq i \leq 6} T(e_i, \nabla_{e_i} e_j) = - \sum_{3 \leq i \leq 6} T(e_i, S(D^j) e_i) = - \sum_{3 \leq i, l \leq 6} S(D^j)_{il} T(e_i, e_l) \\ &= \sum_{3 \leq n \leq 6} (D^j)_{nn} T(e_n, e_n) + \sum_{3 \leq i \neq l \leq 6} S(D^j)_{il} T(e_i, e_l), \end{aligned}$$

whereas for $3 \leq j \leq 6$ we have

$$\begin{aligned} \sum_{1 \leq i \leq 7} T(e_i, \nabla_{e_i} e_j) &= \sum_{i=1,2,7} T(e_i, A(D^i) e_j) + \sum_{3 \leq i \leq 6} T \left(e_i, \sum_{k=1,2,7} \langle S(D^k) e_i, e_j \rangle e_k \right) \\ &= \sum_{i=1,2,7} T(e_i, A(D^i) e_j) + \sum_{3 \leq i \leq 6} \sum_{k=1,2,7} S(D^k)_{ij} T(e_i, e_k) \\ &= \sum_{k=1,2,7} \left(T(e_k, A(D^k) e_j) + \sum_{3 \leq i \leq 6} S(D^k)_{ij} T(e_i, e_k) \right). \end{aligned}$$

Let us recall now equation (11), which states that

$$(68) \quad T(u, v) = \frac{1}{4} \tau_0 g(u, v) - \iota_{\tau_1}(\varphi)(u, v) - \frac{1}{2} \tau_2(u, v) - \tau_{27}(u, v), \quad u, v \in \mathfrak{g}_{A,B,C},$$

and equation (13) defining the symmetric 2-tensor τ_{27} . When $u = v = e_n$ for $3 \leq n \leq 6$ we find that the first term in equation (68) is $\frac{\tau_0}{4}$ because $\{e_j \mid 1 \leq j \leq 7\}$ is chosen to be an orthonormal basis, and that

the second and third terms are zero because they are skew-symmetric, but the last term is non-zero in general; moreover, given that $\sum_{3 \leq n \leq 6} (D^j)_{nn} \frac{\tau_0}{4} = 0$ because of tracelessness of $D = A, B, C$, we have that

$$\sum_{3 \leq n \leq 6} (D^j)_{nn} T(e_n, e_n) = - \sum_{3 \leq n \leq 6} (D^j)_{nn} \tau_{27}(e_n, e_n).$$

Similarly, when $u = e_i$ and $v = e_l$ for $3 \leq i \neq l \leq 6$, we find that the first term in equation (68) is zero because $\{e_j \mid 1 \leq j \leq 7\}$ is chosen to be an orthonormal basis; moreover, the fact that $S(D^j)_{il}$ is symmetric in the indices i and l while the second and third terms in equation (68) are skew-symmetric entails that

$$\sum_{3 \leq i \neq l \leq 6} S(D^j)_{il} T(e_i, e_l) = - \sum_{3 \leq i \neq l \leq 6} S(D^j)_{il} \tau_{27}(e_i, e_l).$$

We can then summarize the findings of the last paragraph as

$$\langle \operatorname{div}(T), e_j \rangle = - \sum_{3 \leq n \leq 6} (D^j)_{nn} \tau_{27}(e_n, e_n) + \sum_{3 \leq i \neq l \leq 6} S(D^j)_{il} \tau_{27}(e_i, e_l) \quad \text{for } j = 1, 2, 7.$$

On the other hand, when $u = e_i$ and $v = e_k$ for $3 \leq i \leq 6$ and $k = 1, 2, 7$ we find that the first term in equation (68) is zero because $\{e_j \mid 1 \leq j \leq 7\}$ is chosen to be an orthonormal basis, the second and third terms are zero because of Remark 3.6, and the last term is zero because of Remark 3.6 and Corollary 3.9. This means that $T(e_i, e_k) = 0$ for $3 \leq i \leq 6$ and $k = 1, 2, 7$; moreover, it also implies that $T(e_k, A(D^k)e_j) = 0$ for $k = 1, 2, 7$ and $3 \leq j \leq 6$, for $A(D^k)e_j \in \mathfrak{n} = \operatorname{span}\{e_3, \dots, e_6\}$ and T is bilinear. Therefore, for $3 \leq j \leq 6$ we have

$$\sum_{1 \leq i \leq 7} T(e_i, \nabla_{e_i} e_j) = \sum_{k=1,2,7} \left(T(e_k, A(D^k)e_j) + \sum_{3 \leq i \leq 6} S(D^k)_{ij} T(e_i, e_k) \right) = 0,$$

and thus,

$$\langle \operatorname{div}(T), e_j \rangle = 0 \quad \text{for } 3 \leq j \leq 6. \quad \square$$

Equation (67) reveals that in order to have an explicit formula for $\operatorname{div}(T)$ we only need to compute

$$(69) \quad \tau_{27}(e_n, e_n) \quad \text{for } 3 \leq n \leq 6, \quad \tau_{27}(e_i, e_l) \quad \text{for } 3 \leq i \neq l \leq 6,$$

a task rendered possible by equation (45). It appears that little gain is made in performing such amount of computations in this general context, so we omit it. It remains unclear how to characterize when the full torsion tensor T is divergence-free in terms of conditions on the coefficients of A, B , and C , and even more so which of those conditions lead to non-equivalent G_2 -structures. Going forward, we will restrict ourselves to the three broad cases that proved tractable: The case in which A, B, C are all skew-symmetric, the case in which A, B, C are all diagonal (which turns out to be equivalent to the symmetric case, see Proposition 5.1), and the case in which A, B, C are all antidiagonal. Recall that a matrix $D = [d_{ij}]$ for $3 \leq i, j \leq 6$ is called *antidiagonal* if it is of the form

$$(70) \quad D = \begin{pmatrix} 0 & 0 & 0 & d_{36} \\ 0 & 0 & d_{45} & 0 \\ 0 & d_{54} & 0 & 0 \\ d_{63} & 0 & 0 & 0 \end{pmatrix}.$$

Generally speaking, these cases are indeed non-equivalent as a consequence of Proposition 3.12, for we have the following:

- In the skew symmetric case, Ric is identically zero (in fact, ∇ is flat in this case; see Proposition 4.1).
- In the diagonal (or symmetric) case, both $\operatorname{Ric}|_{\mathfrak{a} \times \mathfrak{n}}$ and $\operatorname{Ric}|_{\mathfrak{n} \times \mathfrak{n}}$ are identically zero, whereas $\operatorname{Ric}|_{\mathfrak{a} \times \mathfrak{a}}$ is generally not.
- In the antidiagonal case, both $\operatorname{Ric}|_{\mathfrak{a} \times \mathfrak{n}}$ and $\operatorname{Ric}|_{\mathfrak{a} \times \mathfrak{a}}$ are identically zero, whereas $\operatorname{Ric}|_{\mathfrak{n} \times \mathfrak{n}}$ is generally not at least when A, B, C are not all skew-symmetric or diagonal/symmetric.

4. SKEW-SYMMETRIC CASE

This case is of particular interest due in part to the following well-known result.

Proposition 4.1. [21, Theorem 1.5] *If $A, B, C \in \mathfrak{sl}(4, \mathbb{R})$ are skew-symmetric pairwise-commuting then the metric $\langle \cdot, \cdot \rangle$ induced in $\mathfrak{g}_{A,B,C}$ by φ is flat.*

(Ricci)-flatness is a relevant assumption (though not crucial) in the context of the isometric flow of G_2 -structures, for torsion-free G_2 -structures are known to be Ricci-flat (see [11, Propositions 10.1.3 and 10.1.5]) and the isometric flow preserves the underlying metric. It is worth mentioning that three pairwise-commuting skew-symmetric matrices are always a linearly-dependent set, for the dimension of the Cartan subalgebra² of $\mathfrak{so}(3)$ is 2.

According to equation (67), this case is trivial. We state the result for the sake of formality.

Theorem 4.2. *Let $A, B, C \in \mathfrak{sl}(4, \mathbb{R})$ be skew-symmetric. Then $(\mathfrak{g}_{A,B,C}, \varphi)$ is a G_2 -structure with divergence-free full torsion tensor.*

Remark 4.3. Theorem 4.2 is consistent with [23, Theorem 4.8], in which a much larger class of 7-dimensional solvable Lie groups is found to admit a G_2 -structure with divergence-free full torsion tensor that is compatible with a given flat left-invariant metric.

We calculate the torsion forms for this case for the sake of completeness.

Proposition 4.4. *The torsion forms τ_0, τ_1, τ_2 , and τ_3 of $(\mathfrak{g}_{A,B,C}, \varphi)$ in the skew-symmetric case are given by the following formulas:*

$$\begin{aligned}
 (71) \quad \tau_0 &= \frac{4}{7}(a_{46} + a_{53} + b_{35} + b_{64} + c_{54} + c_{63}), \\
 (72) \quad \tau_1 &= -\frac{1}{6}(a_{36} + a_{45} + c_{56} + c_{34})e^1 - \frac{1}{6}(a_{64} + a_{35} + b_{43} + b_{65})e^2 - \frac{1}{6}(b_{63} + b_{54} + c_{46} + c_{53})e^7, \\
 (73) \quad \tau_2 &= \frac{2}{3}(b_{45} + b_{36} + c_{35} + c_{64})e^{12} + \frac{2}{3}(a_{64} + a_{35} + b_{65} + b_{43})e^{17} + \frac{2}{3}(a_{54} + a_{63} + c_{65} + c_{43})e^{27} \\
 &\quad + \frac{1}{3}(c_{46} - c_{35} - b_{45} - b_{36})e^{34} + \frac{1}{3}(-a_{54} + a_{36} + c_{56} + c_{34})e^{35} + \frac{1}{3}(-a_{64} - a_{35} - b_{65} + b_{34})e^{36} \\
 &\quad + \frac{1}{3}(a_{46} + a_{53} + b_{56} - b_{43})e^{45} + \frac{1}{3}(a_{63} - a_{45} + c_{65} + c_{43})e^{46} + \frac{1}{3}(c_{53} - c_{64} + b_{63} + b_{54})e^{56}, \\
 (74) \quad \tau_3 &= \frac{4}{7}(a_{65} - c_{63} + a_{43} - c_{54} + b_{53} - b_{64})e^{127} + \frac{1}{2}(b_{65} + b_{43} - a_{64} + a_{53})e^{134} \\
 &\quad + \frac{1}{7}(-3a_{65} + 3c_{63} - 3a_{43} + 3c_{54} + 4b_{53} - 4b_{64})e^{135} + \frac{1}{2}(b_{63} + b_{54} - c_{53} + c_{64})e^{136} \\
 &\quad + \frac{1}{2}(b_{63} + b_{54} - c_{53} + c_{64})e^{145} + \frac{1}{7}(3a_{65} - 3c_{63} + 3a_{43} - 3c_{54} - 4b_{53} + 4b_{64})e^{146} \\
 &\quad + \frac{1}{2}(b_{65} + b_{43} - a_{64} + a_{53})e^{156} + \frac{1}{2}(c_{65} + c_{43} - a_{54} - a_{63})e^{234} \\
 &\quad + \frac{1}{2}(-b_{63} - b_{54} + c_{53} - c_{64})e^{235} + \frac{1}{7}(3a_{65} + 4c_{63} + 3a_{43} + 4c_{54} + 3b_{53} - 3b_{64})e^{236} \\
 &\quad + \frac{1}{7}(3a_{65} + 4c_{63} + 3a_{43} + 4c_{54} + 3b_{53} - 3b_{64})e^{245} + \frac{1}{2}(b_{63} + b_{54} - c_{53} + c_{64})e^{246} \\
 &\quad + \frac{1}{2}(c_{65} + c_{43} - a_{54} - a_{63})e^{256} + \frac{1}{7}(4a_{65} + 3c_{63} + 4a_{43} + 3c_{54} - 3b_{53} + 3b_{64})e^{347} \\
 &\quad + \frac{1}{2}(b_{65} + b_{43} - a_{64} + a_{53})e^{357} + \frac{1}{2}(-c_{65} - c_{43} + a_{54} + a_{63})e^{367} \\
 &\quad + \frac{1}{2}(-c_{65} - c_{43} + a_{54} + a_{63})e^{457} + \frac{1}{2}(-b_{65} - b_{43} + a_{64} - a_{53})e^{467} \\
 &\quad + \frac{1}{7}(4a_{65} + 3c_{63} + 4a_{43} + 3c_{54} - 3b_{53} + 3b_{64})e^{567}.
 \end{aligned}$$

5. DIAGONAL CASE

This case is of particular interest due in part to the following result.

Proposition 5.1. [16, Lemma 4.5] *If $A, B, C \in \mathfrak{sl}(4, \mathbb{R})$ are symmetric pairwise-commuting then there exist diagonal $A_0, B_0, C_0 \in \mathfrak{sl}(4, \mathbb{R})$ such that $(G_{A,B,C}, \varphi)$ is equivariantly equivalent to $(G_{A_0,B_0,C_0}, \varphi)$, where φ is as given in equation (5).*

Proposition 5.1 shows that the diagonal case is in some sense the “opposite” of the skew-symmetric case; moreover, it ensures that it is enough to consider only diagonal triples $A, B, C \in \mathfrak{sl}(4, \mathbb{R})$ to fully understand the case in which all three $A, B, C \in \mathfrak{sl}(4, \mathbb{R})$ are symmetric and pairwise-commuting. Note

²By *Cartan subalgebra* of $\mathfrak{so}(3)$ we mean a maximal abelian Lie subalgebra consisting of diagonalizable elements of the complexification of $\mathfrak{so}(3)$; it is known that any two Cartan subalgebras are isomorphic.

that any triple of diagonal matrices is trivially pairwise-commuting, so tracelessness is the only extra condition we need to impose.

According to Theorem 3.13, if all three matrices A, B, C are diagonal then the divergence of the full torsion tensor T of $(\mathfrak{g}_{A,B,C}, \varphi)$ is

$$(75) \quad \langle \operatorname{div}(T), e_j \rangle = \begin{cases} -\sum_{3 \leq n \leq 6} (D^j)_{nn} \tau_{27}(e_n, e_n) & j = 1, 2, 7, \\ 0 & 3 \leq j \leq 6. \end{cases}$$

To proceed, we first adapt Proposition 3.4 to the diagonal case.

Proposition 5.2. *The torsion forms τ_0, τ_1, τ_2 , and τ_3 of $(\mathfrak{g}_{A,B,C}, \varphi)$ in the diagonal case are given by the following formulas:*

$$(76) \quad \tau_0 = 0,$$

$$(77) \quad \tau_1 = 0,$$

$$(78) \quad \tau_2 = -(a_{33} + a_{44})e^{34} - (b_{33} + b_{55})e^{35} - (c_{44} + c_{55})e^{36} \\ + (c_{44} + c_{55})e^{45} - (b_{33} - b_{55})e^{46} + (a_{33} + a_{44})e^{56},$$

$$(79) \quad \tau_3 = (c_{33} + c_{44})e^{134} - (a_{44} + a_{55})e^{136} + (a_{44} + a_{55})e^{145} \\ - (c_{33} + c_{44})e^{156} - (b_{33} + b_{44})e^{234} + (a_{33} + a_{55})e^{235} \\ + (a_{33} + a_{55})e^{246} + (b_{33} + b_{44})e^{256} - (c_{33} + c_{55})e^{357} \\ + (b_{44} + b_{55})e^{367} - (b_{44} + b_{55})e^{457} - (c_{33} + c_{55})e^{467},$$

Remark 5.3. Upon further inspection, we note that $\tau_2 = -\star d\psi$ and $\tau_3 = \star d\varphi$ in the diagonal case. This is consistent with [16, Corollary 4.8].

Given that $\tau_3 \in \operatorname{span}\{e^{134}, e^{136}, e^{145}, e^{156}, e^{234}, e^{235}, e^{246}, e^{256}, e^{357}, e^{367}, e^{457}, e^{467}\}$, equation (57) from Corollary (3.8) entails the following result.

Corollary 5.4. *The symmetric 2-tensor τ_{27} of $(\mathfrak{g}_{A,B,C}, \varphi)$ in the diagonal case satisfies*

$$(80) \quad \tau_{27}(e_n, e_n) = 0 \quad \text{for all } 3 \leq n \leq 6.$$

We apply Corollary 5.4 to equation (75) to establish the main result of this Section.

Theorem 5.5. *Let $A, B, C \in \mathfrak{sl}(4, \mathbb{R})$ be diagonal. Then $(\mathfrak{g}_{A,B,C}, \varphi)$ is a G_2 -structure with divergence-free full torsion tensor.*

Theorem 5.5 can be enhanced as a result of Proposition 5.1.

Corollary 5.6. *Let $A, B, C \in \mathfrak{sl}(4, \mathbb{R})$ be symmetric and pairwise-commuting. Then $(\mathfrak{g}_{A,B,C}, \varphi)$ is a G_2 -structure with divergence-free full torsion tensor.*

Remark 5.7. It is worth mentioning that equations (76) and (77) imply all by their own that $\operatorname{div} T = 0$ as a consequence of [9, Theorem 4.3]. The discussion above is an alternative proof of this fact.

6. ANTIDIAGONAL CASE

It is not known whether an analogous result to Proposition 5.1 of Section 5 holds in the antidiagonal case. Note that in order to produce non-equivalent examples of G_2 -structures with divergence-free full torsion tensor to those found in the previous sections we need to consider only triples $A, B, C \in \mathfrak{sl}(4, \mathbb{R})$ such that not all three matrices are skew-symmetric or symmetric. We also note that it is not guaranteed that any election of a triple $A, B, C \in \mathfrak{sl}(4, \mathbb{R})$ of antidiagonal matrices pairwise commute. On the contrary, this imposes a series of restrictions on such matrices. Those restrictions, however, play no role in what follows, and therefore any discussion regarding them is omitted.

According to Theorem 3.13, if all three matrices A, B, C are antidiagonal then the divergence of the full torsion tensor T of $(\mathfrak{g}_{A,B,C}, \varphi)$ is

$$(81) \quad \langle \operatorname{div}(T), e_j \rangle = \begin{cases} \sum_{3 \leq m \leq 6} S(D^j)_{m,9-m} \tau_{27}(e_m, e_{9-m}) & j = 1, 2, 7, \\ 0 & 3 \leq j \leq 6. \end{cases}$$

As in the previous Section, we first adapt Proposition 3.4 to the antidiagonal case to proceed.

Proposition 6.1. *The torsion forms τ_0 , τ_1 , τ_2 , and τ_3 of $(\mathfrak{g}_{A,B,C}, \varphi)$ in the antidiagonal case are given by the following formulas:*

$$(82) \quad \tau_0 = \frac{2}{7}(c_{54} - c_{45} + c_{63} - c_{36}),$$

$$(83) \quad \tau_1 = \frac{1}{12}(a_{63} - a_{36} + a_{54} - a_{45})e^1 + \frac{1}{12}(b_{36} - b_{63} + b_{45} - b_{54})e^7,$$

$$(84) \quad \tau_2 = \frac{1}{3}(b_{45} - b_{54} + b_{36} - b_{63})e^{12} + \frac{1}{3}(a_{54} - a_{45} + a_{63} - a_{36})e^{27} \\ + \frac{1}{3}(2b_{36} - b_{63} + b_{54} - 2b_{45})e^{34} + \frac{1}{3}(2a_{36} + a_{63} - 2a_{54} - a_{45})e^{35} \\ + \frac{1}{3}(2a_{63} + a_{36} - 2a_{45} - a_{54})e^{46} + \frac{1}{3}(2b_{63} + b_{36} + 2b_{54} - b_{45})e^{56},$$

$$(85) \quad \tau_3 = \frac{2}{7}(c_{45} - c_{54} + c_{36} - c_{63})e^{127} + \frac{1}{7}(5c_{54} + 2c_{45} - 5c_{36} - 2c_{63})e^{135} \\ + \frac{1}{4}(b_{63} - b_{36} + b_{54} - b_{45})e^{136} + \frac{1}{4}(b_{63} - b_{36} + b_{54} - b_{45})e^{145} \\ + \frac{1}{7}(5c_{45} + 2c_{54} - 5c_{63} - 2c_{36})e^{146} + \frac{1}{4}(3a_{36} + 3a_{45} + a_{63} + a_{54})e^{234} \\ + \frac{1}{4}(3b_{36} + b_{63} - 3b_{54} - b_{45})e^{235} + \frac{2}{7}(c_{54} - c_{45} + c_{63} - c_{36})e^{236} \\ + \frac{2}{7}(c_{54} - c_{45} + c_{63} - c_{36})e^{245} + \frac{1}{4}(3b_{63} + b_{36} - 3b_{45} - b_{54})e^{246} \\ - \frac{1}{4}(3a_{63} + a_{36} + 3a_{54} + a_{45})e^{256} - \frac{1}{7}(5c_{45} + 2c_{54} + 5c_{36} + 2c_{63})e^{347} \\ + \frac{1}{4}(a_{54} - a_{45} + a_{63} - a_{36})e^{367} + \frac{1}{4}(a_{54} - a_{45} + a_{63} - a_{36})e^{457} \\ + \frac{1}{7}(5c_{54} + 2c_{45} + 5c_{63} + 2c_{36})e^{567},$$

Given that $\tau_3 \in \text{span}\{e^{127}, e^{135}, e^{136}, e^{145}, e^{146}, e^{234}, e^{235}, e^{236}, e^{245}, e^{246}, e^{256}, e^{347}, e^{367}, e^{457}, e^{567}\}$, equation (58) from Corollary 3.8 entails the following result.

Corollary 6.2. *The symmetric 2-tensor τ_{27} of $(\mathfrak{g}_{A,B,C}, \varphi)$ in the antidiagonal case satisfies*

$$(86) \quad \tau_{27}(e_m, e_{9-m}) = 0 \quad \text{for all } 3 \leq m \leq 6.$$

We apply Corollary 6.2 to equation (81) to establish the main result of this Section.

Theorem 6.3. *Let $A, B, C \in \mathfrak{sl}(4, \mathbb{R})$ be pairwise-commuting and antidiagonal. Then $(\mathfrak{g}_{A,B,C}, \varphi)$ is a G_2 -structure with divergence-free full torsion tensor.*

Acknowledgements. I am very grateful with Dr. Adrián Andrada for his continued guidance during the preparation of this paper. I would also like to thank Dr. Jorge Lauret, for a suggestion of his originated this article. Special thanks go to Alejandro Tolcachier for thoughtful conversations and encouragement, and to Facundo Gelatti for his patient and constant assistance in technical matters.

REFERENCES

- [1] J. Baez, The Octonions, Bull. Amer. Math. Soc. **39** (2002), 145–205, <https://doi.org/10.1090/S0273-0979-01-00934-X>
- [2] M. Berger, Sur les groupes d’holonomie homogènes de variétés à connexion affine et des variétés Riemanniennes, Bull. Soc. Math. France **83** (1955), 279–330, <https://doi.org/10.24033/bsmf.1464>
- [3] R. Bryant, Metrics with exceptional holonomy, Ann. Math. **126** (1987), 525–576, <https://doi.org/10.2307/1971360>
- [4] M. Bryant, Some remarks on G_2 -structures, Proc. Gökova Geom.-Topol. Conf. **126** (2005), 75–109, <https://doi.org/10.48550/arXiv.math/0305124>
- [5] S. Dwivedi, P. Gianniotis, S. Karigiannis, A gradient flow of isometric G_2 -structures, J. Geom. Anal. **31** (2021), 1855–1933, <http://doi.org/10.14288/1.0385122>
- [6] M. Fernández, A. Gray, Riemannian manifolds with structure group G_2 , Ann. Mat. Pura Appl. **132** (1982), 19–45, <https://doi.org/10.1007/BF01760975>
- [7] A. Gray, Vector cross products on manifolds, Trans. Amer. Math. Soc. **141** (1969), 465–504, <http://doi.org/10.1090/S0002-9947-1969-0243469-5>
- [8] S. Grigorian, Moduli spaces of G_2 -manifolds, Rev. Math. Phys. **22** (2010), 1061–1097, <http://doi.org/10.1142/S0129055X10004132>
- [9] S. Grigorian, Estimates and monotonicity for a heat flow of isometric G_2 -structures, Calc. Var. Partial Differential Equations **5** (2019), 1–37, <https://doi.org/10.1007/s00526-019-1630-0>
- [10] S. Grigorian, Isometric flows of G_2 -structures, Current Trends in Analysis, its Applications and Computation, Trends in Mathematics, **12** (2022), 545–553, <https://doi.org/10.1007/978-3-030-87502-2>
- [11] D. Joyce, Compact Manifolds with Special Holonomy, Oxford Mathematical Monographs, Oxford University Press Inc, New York, 1st edition (2000).

- [12] S. Karigiannis, Deformations of G_2 and $\text{Spin}(7)$ -structures, *Can. J. Math.* **57** (2005), 1012–1055, <https://doi.org/10.4153/CJM-2005-039-x>
- [13] S. Karigiannis, Flows of G_2 -structures I, *Q. J. Math.* **60** (2008), 487–522, <https://doi.org/10.1093/qmath/han020>
- [14] S. Karigiannis, Introduction to G_2 -geometry, *Lectures and Surveys on G_2 -Manifolds and Related Topics*, Fields Institute Communications, Springer, **84** (2020), 3–50.
- [15] S. Karigiannis, B. McKay, T. Mao-Pei, Soliton solutions for the Laplacian co-flow of some G_2 -structures with symmetry, *Diff. Geom. Appl.* **30** (2012), 318–333, <http://dx.doi.org/10.1016/j.difgeo.2012.05.003>
- [16] I. Kath, J. Lauret, A new example of a compact ERP G_2 -structure, *Bull. London Math. Soc.* **53** (2021), 1692–1710, <https://doi.org/10.1112/blms.12520>
- [17] J. Lauret, Laplacian flow of homogeneous G_2 -structures and its solitons, *Proc. London Math. Soc.* **114** (2017), 527–560, <http://doi.org/10.1112/plms.12014>
- [18] J. Lauret, M. Nicolini, The classification of ERP G_2 -structures on Lie groups, *Ann. Mat. Pura App.* **199** (2020), 2489–2510, <http://doi.org/10.1007/s10231-020-00977-4>
- [19] E. Loubeau, A. Moreno, H. Sá Earp, J. Saavedra, Harmonic $\text{Sp}(2)$ -invariant G_2 -structures on the 7-sphere, *J. Geom. Anal.* **32** (2022), article number 240, <https://doi.org/10.1007/s12220-022-00953-9>
- [20] V. Manero, A. Otal, R. Villacampa, Laplacian coflow for warped G_2 -structures, *Diff. Geom. Appl.* **69** (2020), 93–101, <https://doi.org/10.1016/j.difgeo.2020.101593>
- [21] J. Milnor, Curvatures of left invariant metrics on Lie groups, *Advances in Mathematics* **21** (1976), 293–329.
- [22] M. Nicolini, New examples of shrinking Laplacian solitons, *Q. J. Math.* **73** (2021), 239–259, <https://doi.org/10.1093/qmath/haab029>
- [23] A. Tolcachier, G_2 -structures on flat solvmanifolds, *Abh. Math. Sem. Univ. Hamburg.*, **92** (2022), 179–207 <https://doi.org/10.1007/s12188-022-00261-7>

FAMAF, UNIVERSIDAD NACIONAL DE CÓRDOBA, MEDINA ALLENDE S/N, CIUDAD UNIVERSITARIA (X5000HUA), CÓRDOBA, ARGENTINA.

Email address: agustin.garrone@mi.unc.edu.ar