

Full Counting Statistics and Fluctuation Theorem for the Currents in the Discrete Model of Feynman's Ratchet

Yu-Xin Wu¹, Jiayin Gu^{1,*} and H. T. Quan^{1,2,3†}

¹*School of Physics, Peking University, Beijing 100871, China;*

²*Collaborative Innovation Center of Quantum Matter, Beijing 100871, China; and*

³*Frontiers Science Center for Nano-Optoelectronics, Peking University, Beijing 100871, China*

We provide a detailed investigation on the fluctuations of two currents in the discrete model of Feynman's ratchet proposed by Jarzynski and Mazonka in 1999. Two macroscopic currents are identified, with the corresponding affinities determined using Schnakenberg's graph analysis. We also compute full counting statistics of the two currents and show that a fluctuation theorem holds for their joint probability distribution. Moreover, fluctuation-dissipation relation, Onsager reciprocal relation and their nonlinear generalizations are numerically shown to be satisfied in this model.

I. INTRODUCTION

Feynman's ratchet and pawl system – a device capable of converting thermal random motion into useful work – acts as a microscopic mechanical Maxwell's demon' [1]. As a paradigm for rectifying thermal noise, it has inspired extensive theoretical studies [2–11] as well as experimental efforts [12–15]. Its basic setup includes a ratchet and a pawl (see Fig. 1). From time to time, the pawl switches between being engaged and disengaged. The ratchet is attached to a windmill immersed in a heat reservoir at temperature T_B where the gas molecules collide against the vanes of the windmill, causing the ratchet to rotate. The thermal motion of the microscopic pawl, which is immersed in another reservoir at temperature T_A , will cause the pawl itself to rise up or fall down from time to time. When it rises up, the pawl and the ratchet become disengaged, allowing the free rotation of the ratchet in both directions. But when the pawl falls down, it will press against the teeth of the ratchet, enforcing the ratchet to move in one direction only but never in the other. Feynman found the rates of rotation in two directions to be equal when the temperatures of the two reservoirs are equal. This is in accordance with the second law of thermodynamics. But when there is temperature difference between the two, the device can work as a heat engine or a refrigerator. In the heat engine "phase" the ratchet can rectify the thermal random motion of gas molecules to do useful work while in the refrigerator "phase" it can pump heat from the cold reservoir to the hot reservoir at the expenditure of work.

In 1999, Jarzynski and Mazonka proposed a very elegant discrete model that precisely captures the essential features of Feynman's ratchet [16], and it was recently realized experimentally [17]. Their model consists of only six states whose underlying dynamics is Markovian jump process described by a master equation. They calculated the mean values of mass velocity, energy flow and entropy production rate in nonequilibrium steady state.

They also discussed the linear response relations for two currents—mass displacement and energy flow. However, they didn't explore the fluctuation properties of the currents (but see Ref. [18, 19]). As we know, Feynman proposed this automatic mechanical version of Maxwell's demon to demonstrate that on average the second law can never be violated, but occasionally, in a individual realization, the second law can be probabilistically "violated". Since the original Feynman's setting is too complex to be analyzed from the first principle, he himself was unable to quantitatively characterize the probability of "violating" the second law.

Over the last three decades, fluctuation theorems for systems in nonequilibrium steady state have been studied extensively, both theoretically [20–30] and experimentally [31–36]. In particular, Gaspard, Andrieux, and coworkers extended the fluctuation theorem to coupled currents [37–40]. They provided a unified framework for deducing fluctuation-dissipation theorem, Onsager reciprocal relations, and relations for higher order response coefficients from fluctuation theorem [37, 41–43].

In this article, we focus on the fluctuation of currents, in purpose of showing fluctuation theorem and its implications for the response properties of Feynman's ratchet model. By utilizing the method of graph decomposition proposed by Schnakenberg [44], we decompose the currents in Feynman's ratchet model and calculate the full counting statistics of the currents. With these results, we find the Gallavotti-Cohen symmetry of the currents. Thus we are able to quantitatively characterize the fluctuations and the probabilistic "violation" of the second law in Feynman's ratchet model. Our results confirm what Feynman has envisioned and enhance our understanding about the fluctuating properties of this famous model from the qualitative way to the quantitative way.

The rest of the paper is organized as follows. Sec. II gives a detailed description of the discrete Feynman's ratchet model. Sec. III concerns the currents and the corresponding affinities. Sec. IV is devoted to the full counting statistics of the currents. The response relations are numerically verified in Sec V. Conclusions are given in Sec. VI.

* gujiayin@pku.edu.cn

† htquan@pku.edu.cn




Figure 1. Schematic illustration of the Feynman's ratchet.




Figure 2. The landscape of potential energy $U_i^{(m)}$ for two modes, $m = 1, 2$, in the presence of an external load $f = \alpha/2d$. Alternatively, we denote the potential energy as $U_n, n = 1, 2, \dots, 6$. See Table I for one to one correspondence.

II. DISCRETE MODEL OF FEYNMAN'S RATCHET

The discrete model of Feynman's ratchet proposed by Jarzynski and Mazonka can be envisaged as a particle hopping between neighboring sites on a 1D regular lattice. As shown in Figure 2, the potential energy of each site i has two modes $m = 1, 2$.

$$U_i^{(m)} = \begin{cases} -ifd, & m = 1 \\ \alpha \cdot [(i \bmod 3) - 1] - ifd, & m = 2. \end{cases} \quad (1)$$

Here, f denotes the load, α is the energy unit and d is the lattice spacing. Due to periodicity, we only need to consider one tooth on the ratchet: the system consists of six possible states, $n = 1, \dots, 6$. The energies are listed in Table I. Only those transitions between neighboring sites in the same mode and those between modes on the same site are allowed.

We now denote by $\mathcal{P}_n(t)$ the probability of finding the system in state n at time t . Its evolution is governed by the master equation

$$\frac{d}{dt} \mathcal{P}_n(t) = \sum_{n'(\neq n)} [W_{nn'} \mathcal{P}_{n'}(t) - W_{n'n} \mathcal{P}_n(t)]. \quad (2)$$

Transition rates are given by [45]

$$W_{nn'} = \frac{\beta \Delta U_{n \leftarrow n'}}{\exp(\beta \Delta U_{n \leftarrow n'}) - 1} \quad (3)$$

Table I. The energies of the six states.

state n	mode m	site i	potential energy U_n
1	1	0	0
2	1	1	$-fd$
3	1	2	$-2fd$
4	2	0	$-\alpha$
5	2	1	$-fd$
6	2	2	$\alpha - 2fd$

for allowed transitions and $W_{nn'} = 0$ for forbidden ones. Here $\Delta U_{n \leftarrow n'}$ is the potential energy change associated with the transition from state n' to n . For transition $n' \rightarrow n$ within one period, the energy difference is $\Delta U_{n \leftarrow n'} = U_n - U_{n'}$ with energies U_n listed in Table I. However, for those transitions across the boundaries, the potential energy changes are given by

$$\Delta U_{3 \leftarrow 1} = fd, \quad (4)$$

$$\Delta U_{1 \leftarrow 3} = -fd, \quad (5)$$

$$\Delta U_{6 \leftarrow 4} = 2\alpha + fd, \quad (6)$$

$$\Delta U_{4 \leftarrow 6} = -2\alpha - fd. \quad (7)$$

$\beta \equiv 1/(k_B T)$ is the inverse temperature and k_B is Boltzmann's constant. Depending on the type of transition, T may take different values: $T = T_A$ for switches between two modes and $T = T_B$ for jumps between neighbouring sites. The rate function (3) satisfies the detailed balance condition,

$$W_{nn'} = W_{n'n} \exp(-\beta \Delta U_{n \leftarrow n'}) \quad (8)$$

originating from the microscopic reversibility.

The master equation (2) with explicit transition rates can be simulated with the Gillespie's algorithm [46], which is an explicit method to generate random trajectories for Markov stochastic processes. If we define the diagonal elements to be

$$W_{nn} \equiv - \sum_{n'(\neq n)} W_{n'n}, \quad (9)$$

the master equation (2) can be rewritten into a matrix form as

$$\frac{d}{dt} \mathcal{P} = \mathbf{L} \cdot \mathcal{P}, \quad (10)$$

where $\mathcal{P} = (\mathcal{P}_1, \dots, \mathcal{P}_6)^T$ and $\mathbf{L} = [W_{nn'}]$. Each column of \mathbf{L} sums to zero so that it guarantees the conservation of probability.

III. THE CURRENTS AND THEIR CORRESPONDING AFFINITIES

Intuitively we can infer that there are two coupled currents in the ratchet and pawl system, namely the energy flow between two heat baths and the displacement of the load. When the system switches the mode, reservoir A exchanges an energy α with the system. Therefore the instantaneous energy current from reservoir A to the system is defined as

$$j_E(t) \equiv \alpha \sum_{k=1}^{+\infty} \left[\delta(t - t_k^{1 \leftarrow 4}) - \delta(t - t_k^{4 \leftarrow 1}) + \delta(t - t_k^{6 \leftarrow 3}) - \delta(t - t_k^{3 \leftarrow 6}) \right], \quad (11)$$

$$j_X(t) \equiv d \sum_{k=1}^{+\infty} \left[\delta(t - t_k^{2 \leftarrow 1}) - \delta(t - t_k^{1 \leftarrow 2}) + \delta(t - t_k^{3 \leftarrow 2}) - \delta(t - t_k^{2 \leftarrow 3}) + \delta(t - t_k^{1 \leftarrow 3}) - \delta(t - t_k^{3 \leftarrow 1}) + \delta(t - t_k^{5 \leftarrow 4}) - \delta(t - t_k^{4 \leftarrow 5}) + \delta(t - t_k^{6 \leftarrow 5}) - \delta(t - t_k^{5 \leftarrow 6}) + \delta(t - t_k^{4 \leftarrow 6}) - \delta(t - t_k^{6 \leftarrow 4}) \right]. \quad (12)$$

Correspondingly, there are two driving forces or affinities, which can be determined by analyzing the master equation (2) with Schnakenberg's network theory [44, 47]. In this theory, a graph is associated with the Markov jump process. Vertices represent the mesoscopic states while the edges stand for the allowed transitions between states. The graph for the ratchet system is depicted in Fig. 3. From the so-constructed graph, the affinities can be calculated from the transition rates along cyclic paths and their reversals. For example, the cyclic path associated with an energy transfer α from reservoir A to B and its reversed path could be

$$\mathcal{C}_E : 1 \rightarrow 2 \rightarrow 5 \rightarrow 4 \rightarrow 1, \quad (13)$$

$$\tilde{\mathcal{C}}_E : 1 \rightarrow 4 \rightarrow 5 \rightarrow 2 \rightarrow 1, \quad (14)$$

and the affinity is given by

$$A_E = \frac{1}{\alpha} \ln \frac{W_{14}W_{45}W_{52}W_{21}}{W_{12}W_{25}W_{54}W_{41}} = \beta_B - \beta_A. \quad (15)$$

The prefactor is aimed to set the affinity in the unit of energy. Similarly, for a load displacement of $3d$, the cyclic path and its reversal could be

$$\mathcal{C}_X : 1 \rightarrow 2 \rightarrow 3 \rightarrow 1, \quad (16)$$

$$\tilde{\mathcal{C}}_X : 1 \rightarrow 3 \rightarrow 2 \rightarrow 1. \quad (17)$$

The corresponding affinity is

$$A_X = \frac{1}{3d} \ln \frac{W_{13}W_{32}W_{21}}{W_{12}W_{23}W_{31}} = f\beta_B. \quad (18)$$

where $t_k^{n' \leftarrow n}$ is the k_{th} random transition from state n to n' . Similarly, the instantaneous current of displacement (velocity) of the load is defined by considering transitions between neighboring sites in both modes, yielding




Figure 3. The graph of the discrete model with vertices representing the distinct states, edges denoting the allowed transitions. The subgraph consists of thick yellow line is chosen as the maximal tree. The corresponding chords are numbered with ①, ②, ③, ④.

We notice that although the transition rates usually depend on the mesoscopic states, the so-obtained affinities only depend on the macroscopic thermodynamic forces which are of physical importance. Since the graph only contains six vertices, it can be easily decomposed into four independent cycles which form the fundamental set. Detailed analysis is presented in Appendix A.




Figure 4. The contour map of the cumulant generating function $Q(\lambda_X, \lambda_E)$ centering at $(A_X/2 = 1/2, A_E/2 = 1/3)$ with $T_A = 3, T_B = 1, f = 1$.

IV. FLUCTUATION THEOREM FOR THE CURRENTS

A. Full Counting Statistics

Giving access to all cumulants of the current, full counting statistics is a powerful theoretical method to study fluctuations. In order to calculate the full counting statistics, we first define the accumulated energy flowing across the ratchet system and the accumulated displacement of the load over the time interval $[0, t]$

$$E(t) \equiv \int_0^t j_E(t') dt', \quad X(t) \equiv \int_0^t j_X(t') dt'. \quad (19)$$

Then it is easy to obtain the extended master equation for the joint probability $\mathcal{P}_n(E, X, t)$ of finding the system in state n at time t while having absorbed energy E from reservoir A and moved by a distance X (see Appendix B). By summing over all states, we can also define the marginal probability $\mathcal{P}(E, X, t) \equiv \sum_n \mathcal{P}_n(E, X, t)$. Note that the energy and displacement take discrete values and change according to $E \rightarrow E \pm \alpha$ and $X \rightarrow X \pm d$. We further define the cumulant generating function as

$$Q(\lambda_E, \lambda_X) \equiv \lim_{t \rightarrow \infty} -\frac{1}{t} \ln \sum_{E, X} \mathcal{P}(E, X, t) e^{-\lambda_E E - \lambda_X X}, \quad (20)$$

where λ_E and λ_X are the counting parameters. The cumulant generating function (CGF) $Q(\lambda_E, \lambda_X)$ can be obtained by solving for the leading eigenvalue of operator $\mathbf{L}_{\lambda_E, \lambda_X}$

$$\mathbf{L}_{\lambda_E, \lambda_X} \cdot \Psi_{\lambda_E, \lambda_X} = -Q(\lambda_E, \lambda_X) \Psi_{\lambda_E, \lambda_X}, \quad (21)$$

with $\mathbf{L}_{\lambda_E, \lambda_X}$ being

$$\mathbf{L}_{\lambda_E, \lambda_X} \equiv \begin{pmatrix} W_{11} & W_{12} e^{d\lambda_X} & W_{13} e^{-d\lambda_X} & W_{14} e^{-\alpha\lambda_E} & W_{15} & W_{16} \\ W_{21} e^{-d\lambda_X} & W_{22} & W_{23} e^{d\lambda_X} & W_{24} & W_{25} & W_{26} \\ W_{31} e^{d\lambda_X} & W_{32} e^{-d\lambda_X} & W_{33} & W_{34} & W_{35} & W_{36} e^{\alpha\lambda_E} \\ W_{41} e^{\alpha\lambda_E} & W_{42} & W_{43} & W_{44} & W_{45} e^{d\lambda_X} & W_{46} e^{-d\lambda_X} \\ W_{51} & W_{52} & W_{53} & W_{54} e^{-d\lambda_X} & W_{55} & W_{56} e^{d\lambda_X} \\ W_{61} & W_{62} & W_{63} e^{-\alpha\lambda_E} & W_{64} e^{d\lambda_X} & W_{65} e^{-d\lambda_X} & W_{66} \end{pmatrix}. \quad (22)$$

Since the matrix exponential $\exp(\mathbf{L}_{\lambda_E, \lambda_X} t) > 0$, the Perron-Frobenius theorem applies and the leading eigenvalue $-Q(\lambda_E, \lambda_X)$ of $\mathbf{L}_{\lambda_E, \lambda_X}$ corresponds to the real maximum eigenvalue $\exp[-Q(\lambda_E, \lambda_X)]$ of $\exp(\mathbf{L}_{\lambda_E, \lambda_X} t)$ in magnitude ($t > 0$). So the $\Psi_{\lambda_E, \lambda_X}$ can be asymptoti-

cally evaluated as

$$\Psi_{\lambda_E, \lambda_X} \sim \lim_{t \rightarrow \infty} \exp(\mathbf{L}_{\lambda_E, \lambda_X} t) \cdot \psi, \quad (23)$$

which is then normalized

$$\Psi_{\lambda_E, \lambda_X} \leftarrow \frac{\Psi_{\lambda_E, \lambda_X}}{\sqrt{\Psi_{\lambda_E, \lambda_X}^T \cdot \Psi_{\lambda_E, \lambda_X}}}. \quad (24)$$

Here ψ is the randomly chosen distribution supposed to include the desired component. The matrix exponential can be computed using Padé approximation. The cumulant generating function is calculated as

$$Q(\lambda_E, \lambda_X) = -\Psi_{\lambda_E, \lambda_X}^T \cdot \mathbf{L}_{\lambda_E, \lambda_X} \cdot \Psi_{\lambda_E, \lambda_X}. \quad (25)$$

The stationary distribution is given by $\mathcal{P}_{\text{st}} = \Psi_{0,0}$. Fig. 4 shows in contour map the cumulant generating function in the plane of the counting parameters λ_E and λ_X . Two cross sections are plotted in Fig. 5.

B. Fluctuation Theorem

We find that matrix $\mathbf{L}_{\lambda_E, \lambda_X}$ (22) exhibits the symmetry relation

$$\mathbf{M}^{-1} \cdot \mathbf{L}_{\lambda_E, \lambda_X} \cdot \mathbf{M} = \mathbf{L}_{A_E - \lambda_E, A_X - \lambda_X}^T \quad (26)$$

under similarity transformation with matrix \mathbf{M} [48–50]

$$\mathbf{M} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & e^{\alpha/dT_B} & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & e^{-\alpha/dT_B} \end{pmatrix}. \quad (27)$$

As similarity transformation of the matrix does not change the eigenvalues, the relation (26) implies

$$Q(\lambda_E, \lambda_X) = Q(A_E - \lambda_E, A_X - \lambda_X), \quad (28)$$

which is known as Gallavotti-Cohen symmetry [23, 47]. This symmetry for the CGF can be seen in Fig. 4 and Fig. 5. The CGF is directly related to the large deviation function through Legendre-Fenchel transform

$$I(e, x) = \max_{\lambda_E, \lambda_X} \left[Q(\lambda_E, \lambda_X) - (\lambda_E e + \lambda_X x) \right]. \quad (29)$$

$I(e, x)$ is the large deviation rate function of the probability $\mathcal{P}(E = et, X = xt, t)$ defined as follows

$$\mathcal{P}(E = et, X = xt, t) \sim \lim_{t \rightarrow \infty} e^{-tI(e, x)}. \quad (30)$$

The Gallavotti-Cohen symmetry relation (28) immediately implies

$$I(-e, -x) - I(e, x) = A_E e + A_X x, \quad (31)$$

leading all the way to

$$\lim_{t \rightarrow \infty} \frac{\mathcal{P}(E, X, t)}{\mathcal{P}(-E, -X, t)} \simeq e^{A_E E + A_X X}, \quad (32)$$

which is the usual form of fluctuation theorem. This relation can also be derived based on the cycle theory (See Appendix. A).

What's more, the similarity between $\mathbf{L}_{\lambda_E, \lambda_X}$ and $\mathbf{L}_{A_E - \lambda_E, A_X - \lambda_X}^T$ implies that all eigenvalues are the same, leading to a finite-time fluctuation theorem

$$\langle e^{-\lambda_E E - \lambda_X X} \rangle = \langle e^{-(A_E - \lambda_E)E - (A_X - \lambda_X)X} \rangle. \quad (33)$$

This coincides with the fluctuation theorem of entropy production [51].

C. Thermodynamic Entropy Production

For the ratchet system in nonequilibrium steady state, the mean currents of energy and displacement can be evaluated as

$$J_E = \lim_{t \rightarrow \infty} \frac{1}{t} \sum_{E, X} \mathcal{P}(E, X, t) E, \quad (34)$$

$$J_X = \lim_{t \rightarrow \infty} \frac{1}{t} \sum_{E, X} \mathcal{P}(E, X, t) X. \quad (35)$$

The thermodynamic entropy production rate [52, 53] in units of Boltzmann's constant is identified as the sum of products of the affinities and the corresponding mean currents. According to the fluctuation theorem (32), it can be expressed as

$$\begin{aligned} \frac{1}{k_B} \frac{d_i S}{dt} &= A_E J_E + A_X J_X \\ &= \lim_{t \rightarrow \infty} \frac{1}{t} \sum_{E, X} \mathcal{P}(E, X, t) \ln \frac{\mathcal{P}(E, X, t)}{\mathcal{P}(-E, -X, t)}, \end{aligned} \quad (36)$$

in terms of the Kullback-Leibler divergence of $\mathcal{P}(E, X, t)$ and $\mathcal{P}(-E, -X, t)$. The term on the r.h.s. of Eq. (36) is always non-negative, which is in accordance with the 2nd law of thermodynamics. Also, Eqs. (32,33) indicate that there is a nonzero probability of observing the "violation" of the second law, i.e., $A_E E + A_X X < 0$. But in the Feynman's ratchet, the probability of observing such a "violation" becomes negligibly small in the long time limit since both E and X are proportional to t . These results obviously agree with our intuition about Feynman's ratchet.

V. SYMMETRY RELATIONS FOR THE RESPONSE COEFFICIENTS

In previous sections we fix the affinities A_X, A_E . In this section, we will evaluate the response properties when the affinities are varied. Hence, the CGF




Figure 5. The cross sections of the cumulant generating function $Q(\lambda_X, \lambda_E)$. In the left panel, the counting parameter $\lambda_E = \frac{A_E}{2}$ while in the right panel $\lambda_X = \frac{A_X}{2}$. In both panels, $T_A = 3$

$Q(\lambda_X, \lambda_E)$ should be rewritten as $Q(\boldsymbol{\lambda}, \mathbf{A})$. Here $\boldsymbol{\lambda} = \{\lambda_1, \lambda_2\}$, $\mathbf{A} = \{A_1, A_2\}$, $1 = X, 2 = E$. The cumulants can be evaluated by taking successive derivatives of the CGF with respect to the counting parameters. The average currents and diffusivities are given by [40, 43, 54–57]

$$J_i(\mathbf{A}) = \left. \frac{\partial Q(\boldsymbol{\lambda}; \mathbf{A})}{\partial \lambda_i} \right|_{\boldsymbol{\lambda}=\mathbf{0}}, \quad (37)$$

$$D_{ij}(\mathbf{A}) = -\frac{1}{2} \left. \frac{\partial^2 Q(\boldsymbol{\lambda}; \mathbf{A})}{\partial \lambda_i \partial \lambda_j} \right|_{\boldsymbol{\lambda}=\mathbf{0}}. \quad (38)$$

In the mean time, we can expand the mean currents in the power series of affinities as

$$J_i(\mathbf{A}) = \sum_j L_{i,j} A_j + \frac{1}{2} \sum_{j,k} M_{i,jk} A_j A_k + \dots \quad (39)$$

in terms of the linear and nonlinear response coefficients defined by [40]

$$L_{i,j} \equiv \left. \frac{\partial^2 Q(\boldsymbol{\lambda}; \mathbf{A})}{\partial \lambda_i \partial A_j} \right|_{\boldsymbol{\lambda}=\mathbf{A}=\mathbf{0}}, \quad (40)$$

$$M_{i,jk} \equiv \left. \frac{\partial^3 Q(\boldsymbol{\lambda}; \mathbf{A})}{\partial \lambda_i \partial A_j \partial A_k} \right|_{\boldsymbol{\lambda}=\mathbf{A}=\mathbf{0}}, \quad (41)$$

where we notice $M_{i,jk} = M_{i,kj}$ by definition. From the Gallavotti-Cohen symmetry relation (28), we can easily deduce the fluctuation-dissipation theorem [58]

$$L_{i,j} = D_{ij}(\mathbf{A} = \mathbf{0}), \quad (42)$$

and Onsager reciprocal relations [16, 40, 59, 60]

$$L_{i,j} = L_{j,i}. \quad (43)$$

We can also prove the relations characterizing the second order response properties [40]

$$M_{i,jk} = R_{ij,k} + R_{ik,j}, \quad (44)$$




Figure 6. The comparison of linear response coefficients $L_{i,j}$ (orange triangles) and diffusivities in equilibrium $D_{ij}(\mathbf{0})$ (blue circles). The parameters are taken to be the same as those in Fig. 4.

where $R_{ij,k}$ is defined as the linear response coefficient of the diffusivity around equilibrium,

$$R_{ij,k} \equiv \left. \frac{\partial D_{ij}}{\partial A_k} \right|_{\mathbf{A}=\mathbf{0}} = -\frac{1}{2} \left. \frac{\partial^3 Q(\boldsymbol{\lambda}; \mathbf{A})}{\partial \lambda_i \partial \lambda_j \partial A_k} \right|_{\boldsymbol{\lambda}=\mathbf{A}=\mathbf{0}}. \quad (45)$$

We notice that Onsager reciprocal relation (43) is the direct consequence of the fluctuation-dissipation relations (42) together with the symmetry of the diffusivities $D_{ij} = D_{ji}$. Higher order relations can be obtained likewise.

We do a numerical test of the relations (42)-(44) in the ratchet system. For this purpose, we first calculate the CGF $Q(\lambda_X, \lambda_E; A_X, A_E)$ for several points around $\lambda_X = \lambda_E = A_X = A_E = 0$. Then we perform the Lagrange interpolation to obtain the multivariate polynomial approximating the CGF. Finally, the values of $L_{i,j}$, D_{ij} , $M_{i,jk}$ and $R_{ij,k}$ for $i, j, k = 1, 2$ (1 for X, 2 for E) are calculated by taking derivatives according to their definitions. The agreement between the related quantities is clearly demonstrated in Fig. 6 and 7. Thus for the ratchet system, the validity of predictions of fluctuation theorems is tested in both linear and nonlinear regimes.




Figure 7. The comparison between the second-order response coefficients $M_{i,jk}$ and $R_{i,jk} + R_{i,kj}$, where $R_{i,jk}$ is defined by Eq. (45). Same parameters as in Fig. 4.

VI. CONCLUSION AND PERSPECTIVES

In this article, we calculate the full counting statistics of Feynman's ratchet and show that Gallavotti-Cohen fluctuation theorem holds for the currents in the discrete model of Feynman's ratchet. The Markovian stochastic jump process is described by the master equation, with transition rates determined by energy difference. Detailed balance condition is guaranteed so that the dynamics is compatible with the underlying law of microreversibility. Feynman's ratchet and pawl system is driven out of equilibrium by the temperature difference between two heat baths and also by the external load. Correspondingly, there are two physical currents coupled to each other. By performing full counting statistics of the currents, we obtain the CGF which manifests a symmetry relation. This symmetry quantitatively characterizes the fluctuation and the probabilistic "violation" of the second law by the Feynman's ratchet and hence promotes our understanding about fluctuating properties of the famous model from the qualitative way to the quantitative way. In addition, we point out that the currents exhibit linear and nonlinear response properties in the ratchet system, as predicted by the fluctuation theorem. We numerically verified in detail that the second-order response coefficients are related to the linear response of the diffusivities.

As the six-state model of Feynman's ratchet is simple enough, it can be regarded as a basis for studying more sophisticated systems, e.g., spatially extended models. Considering that the pawl stochastically switches between two modes, a feedback control can be applied to determine whether or not the external load is to be attached. In this way, a fluctuation theorem involving information flow could possibly be explored.

ACKNOWLEDGMENTS

Financial support from National Science Foundation of China under the Grant No. 12147162, 11775001 and 11825501 is acknowledged.

Appendix A: Cycle Decomposition

Schnakenberg found that [44] stochastic processes described by the master equation can be investigated by carrying out graph analysis and that the nonequilibrium constraints exerted on a system are related to affinities of the cycles in the graph. This mathematical tool is applied here to study the fluctuation theorem for the currents – energy flow and load displacement.

In the basic graph G , the vertices represent the distinct states while the edges stand for the allowed transitions between states. Then we choose a maximal tree $T(G)$ which is a covering subgraph of G . The requirements for a maximal tree are listed as follows:

- $T(G)$ covers all the vertices of G and all the edges of $T(G)$ belong to G ;
- $T(G)$ must be connected;
- $T(G)$ contains no circuit.

The edges l of graph G that does not belong to $T(G)$ are referred to as chords of maximal tree $T(G)$. By respectively adding one chord to the maximal tree, we get exactly one closed circuit at a time in the resulting subgraph $T(G) + l$. The set of circuits is called a fundamental set $\{C_l\}$. As marked in Fig. 3, by choosing the maximal tree $T(G)$ in yellow, the remaining four edges automatically become its chords. It has been identified that there are four independent cycles in the fundamental set $\{C_l\} = \{C_I, C_{II}, C_{III}, C_{IV}\}$

$$C_I = (e_{1 \rightarrow 2}, e_{2 \rightarrow 3}, e_{3 \rightarrow 1}), \quad (A1)$$

$$C_{II} = (e_{2 \rightarrow 3}, e_{3 \rightarrow 6}, e_{6 \rightarrow 5}, e_{5 \rightarrow 2}), \quad (A2)$$

$$C_{III} = (e_{4 \rightarrow 6}, e_{6 \rightarrow 5}, e_{5 \rightarrow 4}), \quad (A3)$$

$$C_{IV} = (e_{1 \rightarrow 2}, e_{2 \rightarrow 3}, e_{3 \rightarrow 6}, e_{6 \rightarrow 5}, e_{5 \rightarrow 4}, e_{4 \rightarrow 1}), \quad (A4)$$

where $e_{i \rightarrow j}$ denotes the directional edge representing the transition from state i to j . Schnakenberg's graph analysis tells us that each cycle C in the basic graph can be expressed as linear combination of cycles in the fundamental set. In the long-time limit, the initial state becomes irrelevant and the whole trajectory forms a closed cycle $C(t)$. So we have

$$C(t) = a_I(t)C_I + a_{II}(t)C_{II} + a_{III}(t)C_{III} + a_{IV}(t)C_{IV}. \quad (A5)$$

Now we apply Schnakenberg's network theory to prove that Gallavotti-Cohen symmetry exists in the CGF of the energy flow and load displacement. The load displacement can be evaluated by counting the jump times between states in both modes

$$X(t) = d \left[S_{1 \rightarrow 2} + S_{2 \rightarrow 3} + S_{3 \rightarrow 1} + S_{4 \rightarrow 5} + S_{5 \rightarrow 6} + S_{6 \rightarrow 1} \right] C(t), \quad (A6)$$

where the operator $S_{i \rightarrow j}$ counts the jump times across the edge $e_{i \rightarrow j}$. Such linear operators $S_{i \rightarrow j}$ separately act on independent cycles

$$S_{1 \rightarrow 2}C(t) = a_I(t) + a_{IV}(t), \quad (\text{A7})$$

$$S_{2 \rightarrow 3}C(t) = a_I(t) + a_{II}(t) + a_{IV}(t), \quad (\text{A8})$$

$$S_{3 \rightarrow 1}C(t) = a_I(t), \quad (\text{A9})$$

$$S_{4 \rightarrow 5}C(t) = -a_{III}(t) - a_{IV}(t), \quad (\text{A10})$$

$$S_{5 \rightarrow 6}C(t) = -a_{II}(t) - a_{III}(t) - a_{IV}(t), \quad (\text{A11})$$

$$S_{6 \rightarrow 4}C(t) = -a_{III}(t). \quad (\text{A12})$$

Substituting (A7-A12) into A6 we have

$$X(t) = d \left[3a_I(t) - 3a_{III}(t) \right]. \quad (\text{A13})$$

The energy transferred from the reservoir A to the system can be similarly evaluated by considering only the transitions between two modes

$$E(t) = \alpha \left[S_{4 \rightarrow 1} + S_{3 \rightarrow 6} \right] C(t) = \alpha \left[a_{II}(t) + 2a_{IV}(t) \right]. \quad (\text{A14})$$

The CGF for the energy flow and load displacement can be written as

$$\begin{aligned} Q(\lambda_E, \lambda_X) &= \lim_{t \rightarrow \infty} -\frac{1}{t} \ln \left\langle e^{-\lambda_E E(t) - \lambda_X X(t)} \right\rangle \\ &= \lim_{t \rightarrow \infty} -\frac{1}{t} \ln \left\langle e^{-3\lambda_X a_I(t) - \lambda_E a_{II}(t) + 3\lambda_X a_{III}(t) - 2\lambda_E a_{IV}(t)} \right\rangle. \end{aligned} \quad (\text{A15})$$

Recalling the fluctuation theorem for the currents crossing the chords proved by Andrieux and Gaspard [39], the CGF of independent currents along the chords

$$q(\{\lambda_l\}) = \lim_{t \rightarrow \infty} -\frac{1}{t} \ln \left\langle e^{\sum_l \lambda_l a_l(t)} \right\rangle \quad (\text{A16})$$

exhibits the following symmetry

$$q(\{\lambda_l\}) = q(\{A_l - \lambda_l\}), \quad (\text{A17})$$

where A_l is the affinity of the independent cycle in the fundamental set and the notation $\{\lambda_l\}$ is short for $\{\lambda_I, \dots, \lambda_{IV}\}$. In this case, the affinities $\{A_l\}$ can be calculated by taking the log of ratio of transition rate products in both directions

$$A_I = \frac{1}{d} \ln \frac{W_{13}W_{32}W_{21}}{W_{12}W_{23}W_{31}} = 3f\beta_B, \quad (\text{A18})$$

and similarly

$$A_{II} = \beta_B - \beta_A, \quad (\text{A19})$$

$$A_{III} = -3f\beta_B, \quad (\text{A20})$$

$$A_{IV} = 2(\beta_B - \beta_A). \quad (\text{A21})$$

When $\{C_l\}$ is a fundamental set, the coefficients $\{a_l(t)\}$ in cycle decomposition equal the integrated current along the chords, i.e., the number of times crossing each chord. Then the CGF for joint energy and displacement flow $Q(\lambda_E, \lambda_X)$ can be mapped onto $q(\{\lambda_l\})$ as

$$\begin{aligned} Q(\lambda_E, \lambda_X) &= q(3\lambda_X, \lambda_E, -3\lambda_X, 2\lambda_E) \\ &= q(A_I - 3\lambda_X, A_{II} - \lambda_E, A_{III} + 3\lambda_X, A_{IV} - 2\lambda_E) \\ &= q(3f\beta_B - 3\lambda_X, \beta_B - \beta_A - \lambda_E, -3f\beta_B + 3\lambda_X, 2\beta_B - 2\beta_A - 2\lambda_E) \\ &= Q(A_E - \lambda_E, A_X - \lambda_X), \end{aligned} \quad (\text{A22})$$

which gives the Gallavotti-Cohen symmetry (28).

Appendix B: Derivation of Eq. (22)

The generating function for the joint energy flow (from reservoir A to the system) and the displacement is defined as

$$\Psi_{\lambda_E, \lambda_X}(t) = \sum_n (\Psi_{\lambda_E, \lambda_X})_n(t), \quad (\text{B1})$$

$$(\Psi_{\lambda_E, \lambda_X})_n(t) = \int dE dX e^{-\lambda_E E - \lambda_X X} \mathcal{P}_n(E, X, t), \quad (\text{B2})$$

where $\mathcal{P}_n(E, X, t)$ represents the probability of finding the system in state n at time t while having absorbed energy $E(t)$ from reservoir A and moved by a displacement $X(t)$. The displacement of the particle and the heat absorbed from reservoir A during a transition from state m to state n is given by ΔX_{nm} and ΔE_{nm} .

In a small time interval τ the variation of $\mathcal{P}_n(E, X, t)$ is given by

$$\mathcal{P}_n(E, X, t + \tau) \approx \mathcal{P}_n(E, X, t) + \tau \sum_{m(\neq n)} [W_{nm} \mathcal{P}_m(E - \Delta E_{nm}, X - \Delta X_{nm}, t) - W_{mn} \mathcal{P}_n(E, X, t)] \quad (\text{B3})$$

$$= \mathcal{P}_n(E, X, t) + \tau \sum_{m(\neq n)} \{W_{nm} \sum_{k, l=0} \left[\frac{(-\Delta E_{nm})^k}{k!} \frac{(-\Delta X_{nm})^l}{l!} \frac{\partial^{k+l} \mathcal{P}_m(E, X, t)}{\partial E^k \partial X^l} \right] - W_{mn} \mathcal{P}_n(E, X, t)\}. \quad (\text{B4})$$

The joint distribution function $\mathcal{P}_n(E, X, t)$ evolves as

$$\frac{\partial \mathcal{P}_n(E, X, t)}{\partial t} = \sum_{m(\neq n)} \{W_{nm} \sum_{k, l} \left[\frac{(-\Delta E_{nm})^k}{k!} \frac{(-\Delta X_{nm})^l}{l!} \frac{\partial^{k+l} \mathcal{P}_m(E, X, t)}{\partial E^k \partial X^l} \right] - W_{mn} \mathcal{P}_n(E, X, t)\} \quad (\text{B5})$$

with $(\Delta E)_{nm}$ and $(\Delta X)_{nm}$ respectively given by

$$(\Delta X)_{nm} = d \begin{pmatrix} 0 & -1 & 1 & 0 & 0 & 0 \\ 1 & 0 & -1 & 0 & 0 & 0 \\ -1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 1 \\ 0 & 0 & 0 & 1 & 0 & -1 \\ 0 & 0 & 0 & -1 & 1 & 0 \end{pmatrix}, \quad (\Delta E)_{nm} = \alpha \begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 \\ -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}. \quad (\text{B6})$$

From the differential equation (2), we get the time evolution for $(\Psi_{\lambda_E, \lambda_X})_n(t)$

$$\frac{(\Psi_{\lambda_E, \lambda_X})_n(t)}{\partial t} = \sum_m \{W_{nm} e^{-\lambda_E \Delta E_{nm} - \lambda_X \Delta X_{nm}} - \delta_{nm} [\sum_{l(\neq n)} W_{lm}]\} (\Psi_{\lambda_E, \lambda_X})_m(t) \quad (\text{B7})$$

$$= \sum_m (\mathbf{L}_{\lambda_E, \lambda_X})_{nm} (\Psi_{\lambda_E, \lambda_X})_m(t). \quad (\text{B8})$$

This way we get the evolution matrix $\mathbf{L}_{\lambda_E, \lambda_X}$ (22).

- [1] R. P. Feynman, R. B. Leighton, M. Sands, and E. M. Hafner, *American Journal of Physics* **33**, 750 (1965).
 [2] M. O. Magnasco, *Physical Review Letters* **71**, 1477 (1993).
 [3] J. M. R. Parrondo and P. Español, *American Journal of*

- Physics* **64**, 1125 (1996).
 [4] K. Sekimoto, *Journal of the Physical Society of Japan* **66**, 1234 (1997).
 [5] M. O. Magnasco and G. Stolovitzky, *Journal of Statistical Physics* **93**, 615 (1998).

- [6] R. D. Astumian and I. Derényi, *European Biophysics Journal* **27**, 474 (1998).
- [7] C. V. den Broeck, R. Kawai, and P. Meurs, *Physical Review Letters* **93**, 090601 (2004).
- [8] A. Ryabov, V. Holubec, M. H. Yaghoubi, M. Varga, M. E. Foulaadvand, and P. Chvosta, *Journal of Statistical Mechanics: Theory and Experiment* **2016**, 093202 (2016).
- [9] A. Gomez-Marin and J. M. Sancho, *Physical Review E* **73**, 045101 (2006).
- [10] N. Nakagawa and T. S. Komatsu, *Europhysics Letters (EPL)* **75**, 22 (2006).
- [11] Z. C. Tu, *Journal of Physics A: Mathematical and Theoretical* **41**, 312003 (2008).
- [12] I. A. Martínez, É. Roldán, L. Dinis, and R. A. Rica, *Soft Matter* **13**, 22 (2017).
- [13] J. A. C. Albay, G. Paneru, H. K. Pak, and Y. Jun, *Optics Express* **26**, 29906 (2018).
- [14] J. A. C. Albay, P.-Y. Lai, and Y. Jun, *Applied Physics Letters* **116**, 103706 (2020).
- [15] P. Eshuis, K. van der Weele, D. Lohse, and D. van der Meer, *Physical Review Letters* **104**, 248001 (2010).
- [16] C. Jarzynski and O. Mazonka, *Physical Review E* **59**, 6448 (1999).
- [17] J. Bang, R. Pan, T. M. Hoang, J. Ahn, C. Jarzynski, H. T. Quan, and T. Li, *New Journal of Physics* **20**, 103032 (2018).
- [18] H. Sakaguchi, *Journal of the Physical Society of Japan* **69**, 104 (2000).
- [19] W. D. Roeck and C. Maes, *Physical Review E* **76**, 051117 (2007).
- [20] G. Nicolis and I. Prigogine, *Proceedings of the National Academy of Sciences* **68**, 2102 (1971).
- [21] I. Prigogine and G. Nicolis, in *Bifurcation Analysis* (Springer Netherlands, 1985) pp. 3–12.
- [22] D. J. Evans, E. G. D. Cohen, and G. P. Morriss, *Physical Review Letters* **71**, 2401 (1993).
- [23] G. Gallavotti and E. G. D. Cohen, *Physical Review Letters* **74**, 2694 (1995).
- [24] G. Gallavotti and E. G. D. Cohen, *Journal of Statistical Physics* **80**, 931 (1995).
- [25] J. Kurchan, *Journal of Physics A: Mathematical and General* **31**, 3719 (1998).
- [26] J. L. Lebowitz and H. Spohn, *Journal of Statistical Physics* **95**, 333 (1999).
- [27] C. Maes, *Journal of Statistical Physics* **95**, 367 (1999).
- [28] C. Jarzynski, *Physical Review Letters* **78**, 2690 (1997).
- [29] C. Jarzynski, *Physical Review E* **56**, 5018 (1997).
- [30] G. E. Crooks, *Journal of Statistical Physics* **90**, 1481 (1998).
- [31] G. M. Wang, E. M. Sevick, E. Mittag, D. J. Searles, and D. J. Evans, *Physical Review Letters* **89**, 050601 (2002).
- [32] E. Seitaridou, M. M. Inamdar, R. Phillips, K. Ghosh, and K. Dill, *The Journal of Physical Chemistry B* **111**, 2288 (2007).
- [33] Y. Utsumi, D. S. Golubev, M. Marthaler, K. Saito, T. Fujisawa, and G. Schön, *Physical Review B* **81**, 125331 (2010).
- [34] B. Küng, C. Rössler, M. Beck, M. Marthaler, D. S. Golubev, Y. Utsumi, T. Ihn, and K. Ensslin, *Physical Review X* **2**, 011001 (2012).
- [35] N. Garnier and S. Ciliberto, *Physical Review E* **71**, 060101 (2005).
- [36] S. Ciliberto, A. Imparato, A. Naert, and M. Tanase, *Physical Review Letters* **110**, 180601 (2013).
- [37] D. Andrieux and P. Gaspard, *The Journal of Chemical Physics* **121**, 6167 (2004).
- [38] D. Andrieux and P. Gaspard, *Journal of Statistical Mechanics: Theory and Experiment* **2006**, P01011 (2006).
- [39] D. Andrieux and P. Gaspard, *Journal of statistical physics* **127**, 107 (2007).
- [40] D. Andrieux, P. Gaspard, T. Monnai, and S. Tasaki, *New Journal of Physics* **11**, 043014 (2009).
- [41] D. Andrieux and P. Gaspard, *Journal of Statistical Mechanics: Theory and Experiment* **2007**, P02006 (2007).
- [42] P. Gaspard, *New Journal of Physics* **15**, 115014 (2013).
- [43] M. Barbier and P. Gaspard, *Journal of Physics A: Mathematical and Theoretical* **51**, 355001 (2018).
- [44] J. Schnakenberg, *Reviews of Modern Physics* **48**, 571 (1976).
- [45] In their paper, Jarzynski and Mazonka only gave the ratio of the transition rates and adopt the Metropolis algorithm to simulate the dynamics.
- [46] D. T. Gillespie, *Journal of Computational Physics* **22**, 403 (1976).
- [47] A. Faggionato and D. D. Pietro, *Journal of Statistical Physics* **143**, 11 (2011).
- [48] A. W. C. Lau, D. Lacoste, and K. Mallick, *Physical Review Letters* **99**, 158102 (2007).
- [49] D. Lacoste and K. Mallick, *Physical Review E* **80**, 021923 (2009).
- [50] D. Lacoste, A. W. Lau, and K. Mallick, *Physical Review E* **78**, 011915 (2008).
- [51] U. Seifert, *Physical Review Letters* **95**, 040602 (2005).
- [52] I. Prigogine and P. V. Rysselberghe, *Journal of The Electrochemical Society* **110**, 97C (1963).
- [53] H. B. Callen and H. L. Scott, *American Journal of Physics* **66**, 164 (1998).
- [54] M. Barbier and P. Gaspard, *Physical Review E* **102**, 022141 (2020).
- [55] M. Barbier and P. Gaspard, *Journal of Physics A: Mathematical and Theoretical* **53**, 145002 (2020).
- [56] J. Gu and P. Gaspard, *Journal of Statistical Mechanics: Theory and Experiment* **2020**, 103206 (2020).
- [57] J. Gu and P. Gaspard, *Physical Review E* **99**, 012137 (2019).
- [58] R. Kubo, *Reports on Progress in Physics* **29**, 255 (1966).
- [59] L. Onsager, *Physical Review* **38**, 2265 (1931).
- [60] H. B. G. Casimir, *Reviews of Modern Physics* **17**, 343 (1945).