

Relations among $B_c \rightarrow J/\psi, \eta_c$ form factors

P. Colangelo^a, F. De Fazio^a, F. Loparco^a, N. Losacco^{a,b}, M. Novoa-Brunet^a

^aIstituto Nazionale di Fisica Nucleare, Sezione di Bari, Via Orabona 4, I-70126 Bari, Italy

^bDipartimento Interateneo di Fisica “M. Merlin”, Università e Politecnico di Bari,
via Orabona 4, 70126 Bari, Italy

Abstract

We analyze the form factors parametrizing the $B_c \rightarrow J/\psi, \eta_c$ matrix elements of the operators in a generalized low-energy $b \rightarrow c$ semileptonic Hamiltonian. We consider an expansion in nonrelativistic QCD, classifying the heavy quark spin symmetry breaking terms and expressing the form factors in terms of universal functions in a selected kinematical range. Using as an input the lattice QCD results for the $B_c \rightarrow J/\psi$ matrix element of the SM operator, we obtain information on other form factors. The extrapolation to the full kinematical range is also presented.

1 Introduction

The hadronic uncertainty affecting the description of several weak processes represents an important limitation for the Standard Model (SM) predictions. Such a theoretical error needs to be reduced or (at least) controlled in view of the ongoing and planned high precision measurements. Moreover, the error assessment is a preliminary step before interpreting the deviations of experimental results from the SM expectations, an important issue considering the recently observed tensions, the so-called flavour anomalies. In the time in which the search for phenomena beyond the Standard Model (BSM) through the direct production of new particles at the colliders has not produced compelling evidence, the search for BSM signals relies on the precision analysis of virtual effects distorting the SM predictions, as well as on the improved measurements of the fundamental parameters.

In the heavy flavour sector, anomalies have been detected in charged current $b \rightarrow c\ell\nu$ induced transition of $B_{d,u}$ mesons, comparing the $\ell = e, \mu$ with the $\ell = \tau$ modes [1, 2]. If such deviations are due to genuine new physics (NP) phenomena producing violation of lepton flavour universality (LFU), related deviations should be found in B_s , B_c and b -baryon decay modes, both exclusive and inclusive [3, 4]. However, the various processes are affected by hadronic uncertainties in different ways, hence it is necessary to analyze each mode separately.

The present study is devoted to the exclusive semileptonic $b \rightarrow c\ell\nu$ decays of the B_c meson, in particular $B_c \rightarrow J/\psi\ell\bar{\nu}_\ell$ and $B_c \rightarrow \eta_c\ell\bar{\nu}_\ell$ which are under experimental scrutiny [5]. For such processes possible BSM effects can be analyzed in terms of a low-energy Hamiltonian generalizing the Standard Model one with the inclusion of the full set of dimension-6 operators,

as done for other modes [3,6–12]. The matrix elements of the various operators in the effective Hamiltonian, parametrized in terms of form factors, introduce the hadronic uncertainty. The peculiarity of the processes is that they involve mesons each one comprising two heavy quarks, strengthening the interest in B_c , a meson with quarkonium structure and only weak decays.

The form factors for the $B_c \rightarrow J/\psi$ matrix elements of SM operators have been computed in the full kinematical range of dilepton invariant mass q^2 by lattice nonrelativistic QCD, by the HPQCD Collaboration [13]. Other QCD-based computations employ QCD sum rules in the low q^2 range [14,15]. In the same kinematical range, using standard NRQCD methods the $\langle J/\psi | \bar{c} \Gamma_i b | B_c \rangle$ matrix elements are expressed in the form $\langle J/\psi | \bar{c} \Gamma_i b | B_c \rangle \simeq \psi_{B_c}(0) \psi_{J/\psi}(0) T_i$, in terms of meson wave functions at the origin $\psi_{B_c}(0)$, $\psi_{J/\psi}(0)$, and of perturbatively calculable hard-scattering kernels T_i describing spectator interaction corrections and vertex corrections to factorization [16–20]. Perturbative QCD calculations use an analogous principle [21–24]. Light-Cone sum rules have been applied for all q^2 [25]. These approaches are affected by their own theoretical uncertainties. As for the determinations based on quark models, beyond the variation of the input parameters the uncertainty attached to the model can hardly be quantified [26,27].

In Ref. [28] it has been observed that the semileptonic B_c form factors can be expressed in terms of universal functions in selected kinematical regions, on the basis of the heavy quark spin symmetry for large heavy quark masses. This remark has prompted a number of phenomenological analyses of B_c decays [29–31]. The relations to the universal functions have been provided at the leading order in the heavy quark mass expansion. Here we extend the analysis at the next-to-leading order in the expansion, to establish relations among the form factors and a set of universal functions based on the heavy quark spin symmetry and the power counting rules of nonrelativistic QCD (NRQCD). The relations obtained in such a systematic expansion have two applications. They can be used to test the form factors obtained by different methods, for a quantitative assessment of their theoretical uncertainty. Moreover, information can be gained on form factors that have not been computed yet, which are needed for analyses based on the generalized low-energy Hamiltonian. For this purpose, the available lattice QCD results can be employed as input information.

We shall use the power counting of NRQCD, the effective QCD theory relevant for mesons comprising two heavy quarks [32,33]. A classification of the NRQCD operators is in [34]. Using this effective theory and the potential-NRQCD effective theory (pNRQCD) formulated at a lower scale, fundamental QCD parameters have been precisely computed, namely the beauty and charm quark mass and the coupling constant α_s [35]. Precise determinations of the B_c mass [36,37] and lifetime have also been obtained [38,39]. Here we focus on semileptonic form factors based on a systematic expansion.

The plan of the paper is as follows. In Sec.2 we introduce the $b \rightarrow c \ell \nu$ generalized low-energy Hamiltonian in terms of $D = 6$ operators. In Sec.3 we describe the formalism of the NRQCD expansion and in Sec.4 we define the universal functions. In Sec.5 we present a set of numerical results. In App. A we define the parametrization used for the hadronic matrix elements. In App. B we provide the form factors in terms of universal functions. In App. C we give the results of the fit of some form factors obtained through the universal functions.

2 Generalized $b \rightarrow c$ semileptonic effective Hamiltonian and $B_c \rightarrow J/\psi, \eta_c$ form factors

The low-energy Hamiltonian comprising the full set of $D = 6$ semileptonic $b \rightarrow c$ operators with left-handed neutrinos can be written in the form:

$$H_{\text{eff}}^{b \rightarrow c \ell \bar{\nu}} = \frac{G_F}{\sqrt{2}} V_{cb} \left[(1 + \epsilon_V^\ell) (\bar{c} \gamma_\mu (1 - \gamma_5) b) (\bar{\ell} \gamma^\mu (1 - \gamma_5) \nu_\ell) + \epsilon_R^\ell (\bar{c} \gamma_\mu (1 + \gamma_5) b) (\bar{\ell} \gamma^\mu (1 - \gamma_5) \nu_\ell) \right. \\ \left. + \epsilon_S^\ell (\bar{c} b) (\bar{\ell} (1 - \gamma_5) \nu_\ell) + \epsilon_P^\ell (\bar{c} \gamma_5 b) (\bar{\ell} (1 - \gamma_5) \nu_\ell) + \epsilon_T^\ell (\bar{c} \sigma_{\mu\nu} (1 - \gamma_5) b) (\bar{\ell} \sigma^{\mu\nu} (1 - \gamma_5) \nu_\ell) \right]. \quad (1)$$

G_F is the Fermi constant and V_{cb} an element of the Cabibbo-Kobayashi-Maskawa (CKM) mixing matrix. $\mathcal{O}_{SM} = 4(\bar{c}_L \gamma^\mu b_L) (\bar{\ell}_L \gamma_\mu \nu_{\ell L})$ is the SM operator. The Hamiltonian Eq. (1) also comprises the operator $\mathcal{O}_R = 4(\bar{c}_R \gamma^\mu b_R) (\bar{\ell}_L \gamma_\mu \nu_{\ell L})$, the scalar $\mathcal{O}_S = (\bar{c} b) (\bar{\ell} (1 - \gamma_5) \nu_\ell)$, pseudoscalar $\mathcal{O}_P = (\bar{c} \gamma_5 b) (\bar{\ell} (1 - \gamma_5) \nu_\ell)$ and tensor $\mathcal{O}_T = (\bar{c} \sigma_{\mu\nu} (1 - \gamma_5) b) (\bar{\ell} \sigma^{\mu\nu} (1 - \gamma_5) \nu_\ell)$ operators. The Wilson coefficients $\epsilon_{V,R,S,P,T}^\ell$ are complex and lepton-flavour dependent, in general. Eq. (1) reduces to the SM for $\epsilon_i^\ell = 0$.¹ The Hamiltonian Eq. (1) has been considered in connection with the anomalies in $B \rightarrow D^{(*)} \tau \nu_\tau$ vs $B \rightarrow D^{(*)} \ell \nu_\ell$ decays, obtaining information on the various operators and bounding the parameter space of the Wilson coefficients [3, 6–12]. Exclusive B semileptonic modes induced by the $b \rightarrow u$ transition and inclusive b -baryon modes have been studied analogously [4, 43].

The B_c and $J/\psi, \eta_c$ matrix elements of the operators in Eq. (1) require hadronic form factors, for which different parametrizations can be used. The $B_c \rightarrow \eta_c$ matrix elements of the vector $\bar{Q}' \gamma_\mu Q$, scalar $\bar{Q}' Q$, and tensor $\bar{Q}' \sigma_{\mu\nu} Q$ and $\bar{Q}' \sigma_{\mu\nu} \gamma_5 Q$ currents can be written in terms of form factors f_i as

$$\langle P(p') | \bar{Q}' \gamma_\mu Q | B_c(p) \rangle = f_+^{B_c \rightarrow P}(q^2) \left(p_\mu + p'_\mu - \frac{m_{B_c}^2 - m_P^2}{q^2} q_\mu \right) + f_0^{B_c \rightarrow P}(q^2) \frac{m_{B_c}^2 - m_P^2}{q^2} q_\mu, \\ \langle P(p') | \bar{Q}' Q | B_c(p) \rangle = f_S^{B_c \rightarrow P}(q^2), \\ \langle P(p') | \bar{Q}' \sigma_{\mu\nu} Q | B_c(p) \rangle = -i \frac{2f_T^{B_c \rightarrow P}(q^2)}{m_{B_c} + m_P} (p_\mu p'_\nu - p_\nu p'_\mu), \\ \langle P(p') | \bar{Q}' \sigma_{\mu\nu} \gamma_5 Q | B_c(p) \rangle = -\frac{2f_T^{B_c \rightarrow P}(q^2)}{m_{B_c} + m_P} \epsilon_{\mu\nu\alpha\beta} p^\alpha p'^\beta, \quad (2)$$

where $P = \eta_c$, $q = p - p'$ is the momentum transfer to the lepton pair, and the condition $f_+^{B_c \rightarrow P}(0) = f_0^{B_c \rightarrow P}(0)$ holds. We use $\epsilon^{0123} = +1$ and the relation $\sigma_{\mu\nu} \gamma_5 = \frac{i}{2} \epsilon_{\mu\nu\alpha\beta} \sigma^{\alpha\beta}$. $f_S^{B_c \rightarrow P}$ is related to $f_0^{B_c \rightarrow P}$: $f_S^{B_c \rightarrow P}(q^2) = \frac{m_{B_c}^2 - m_P^2}{m_Q - m_{Q'}} f_0^{B_c \rightarrow P}(q^2)$ with quark masses m_Q and $m_{Q'}$. The

¹The operator \mathcal{O}_R is included in the set of dimension 6 operators with left-handed neutrinos. In the Standard Model Effective Field Theory the only dimension-6 operator with right-handed quark current is nonlinear in the Higgs field [40–42].

$B_c \rightarrow J/\psi$ matrix elements can be parametrized as

$$\begin{aligned}
\langle V(p', \epsilon) | \bar{Q}' \gamma_\mu Q | B_c(p) \rangle &= -\frac{2V^{B_c \rightarrow V}(q^2)}{m_{B_c} + m_V} i \epsilon_{\mu\nu\alpha\beta} \epsilon^{*\nu} p^\alpha p'^\beta, \\
\langle V(p', \epsilon) | \bar{Q}' \gamma_\mu \gamma_5 Q | B_c(p) \rangle &= (m_{B_c} + m_V) \left(\epsilon_\mu^* - \frac{(\epsilon^* \cdot q)}{q^2} q_\mu \right) A_1^{B_c \rightarrow V}(q^2) \\
&\quad - \frac{(\epsilon^* \cdot q)}{m_{B_c} + m_V} \left((p + p')_\mu - \frac{m_{B_c}^2 - m_V^2}{q^2} q_\mu \right) A_2^{B_c \rightarrow V}(q^2) \\
&\quad + (\epsilon^* \cdot q) \frac{2m_V}{q^2} q_\mu A_0^{B_c \rightarrow V}(q^2), \\
\langle V(p', \epsilon) | \bar{Q}' \gamma_5 Q | B_c(p) \rangle &= -\frac{2m_V}{m_Q + m_{Q'}} (\epsilon^* \cdot q) A_0^{B_c \rightarrow V}(q^2), \\
\langle V(p', \epsilon) | \bar{Q}' \sigma_{\mu\nu} Q | B_c(p) \rangle &= T_0^{B_c \rightarrow V}(q^2) \frac{\epsilon^* \cdot q}{(m_{B_c} + m_V)^2} \epsilon_{\mu\nu\alpha\beta} p^\alpha p'^\beta \\
&\quad + T_1^{B_c \rightarrow V}(q^2) \epsilon_{\mu\nu\alpha\beta} p^\alpha \epsilon^{*\beta} + T_2^{B_c \rightarrow V}(q^2) \epsilon_{\mu\nu\alpha\beta} p'^\alpha \epsilon^{*\beta}, \\
\langle V(p', \epsilon) | \bar{Q}' \sigma_{\mu\nu} \gamma_5 Q | B_c(p) \rangle &= i T_0^{B_c \rightarrow V}(q^2) \frac{\epsilon^* \cdot q}{(m_{B_c} + m_V)^2} (p_\mu p'_\nu - p_\nu p'_\mu) \\
&\quad + i T_1^{B_c \rightarrow V}(q^2) (p_\mu \epsilon_\nu^* - \epsilon_\mu^* p_\nu) + i T_2^{B_c \rightarrow V}(q^2) (p'_\mu \epsilon_\nu^* - \epsilon_\mu^* p'_\nu),
\end{aligned} \tag{3}$$

with $V = J/\psi$ and ϵ the J/ψ polarization vector. The condition holds:

$$A_0^{B_c \rightarrow V}(0) = \frac{m_{B_c} + m_V}{2m_V} A_1^{B_c \rightarrow V}(0) - \frac{m_{B_c} - m_V}{2m_V} A_2^{B_c \rightarrow V}(0). \tag{4}$$

The results for V and $A_{1,2,0}$ computed in [13] will be exploited in our numerical analysis.

In our study it is convenient to use a different basis of form factors defined in Appendix A. For $B_c \rightarrow \eta_c$ the relations between the two basis are:

$$\begin{aligned}
\frac{m_{B_c}^2 - m_P^2}{q^2} \left(f_0(q^2) - f_+(q^2) \right) &= \frac{1}{2\sqrt{m_{B_c} m_P}} \left((m_{B_c} + m_P) h_-(w) - (m_{B_c} - m_P) h_+(w) \right) \\
f_+(q^2) &= \frac{1}{2\sqrt{m_{B_c} m_P}} \left((m_{B_c} + m_P) h_+(w) - (m_{B_c} - m_P) h_-(w) \right) \tag{5} \\
f_T(q^2) &= -\frac{m_{B_c} + m_P}{\sqrt{m_{B_c} m_P}} h_T(w),
\end{aligned}$$

where $v = \frac{p}{m_{B_c}}$, $v' = \frac{p'}{m_P}$ and $w = v \cdot v'$, hence $q^2 = m_{B_c}^2 + m_P^2 - 2m_{B_c} m_P w$. For $B_c \rightarrow J/\psi$

the relations are:

$$\begin{aligned}
V(q^2) &= \frac{m_{B_c} + m_V}{2\sqrt{m_{B_c}m_V}} h_V(w) \\
A_1(q^2) &= \sqrt{m_{B_c}m_V} \frac{w+1}{m_{B_c} + m_V} h_{A_1}(w) \\
A_2(q^2) &= \frac{m_{B_c} + m_V}{2\sqrt{m_{B_c}m_V}} \left(h_{A_3}(w) + \frac{m_V}{m_B} h_{A_2}(w) \right) \\
A_0(q^2) &= \frac{1}{2\sqrt{m_{B_c}m_V}} \left(m_{B_c}(w+1)h_{A_1}(w) - (m_{B_c} - m_V w)h_{A_2}(w) - (m_{B_c} w - m_V)h_{A_3}(w) \right) \\
T_0(q^2) &= -\frac{(m_{B_c} + m_V)^2}{m_{B_c}\sqrt{m_{B_c}m_V}} h_{T_3}(w) \\
T_1(q^2) &= \frac{m_V}{\sqrt{m_{B_c}m_V}} \left(h_{T_1}(w) + h_{T_2}(w) \right) \\
T_2(q^2) &= \frac{m_{B_c}}{\sqrt{m_{B_c}m_V}} \left(h_{T_1}(w) - h_{T_2}(w) \right) ,
\end{aligned} \tag{6}$$

with $q^2 = m_{B_c}^2 + m_V^2 - 2m_{B_c}m_V w$. The form factors h_i can be related to a set of universal functions in a kinematical range close to $w = 1$. For hadrons comprising a single heavy quark, this has been done in [44–46]. The modifications for the heavy quarkonium are discussed in the next sections.

3 Expansion of the heavy quark field and the QCD Lagrangian

To construct the heavy quark expansion, the heavy quark QCD field $Q(x)$ with mass m_Q is written factorizing a fast oscillation mass term:

$$Q(x) = e^{-im_Q v \cdot x} \psi(x) = e^{-im_Q v \cdot x} (\psi_+(x) + \psi_-(x)) \tag{7}$$

with $\psi_{\pm} = P_{\pm} \psi(x)$ and $P_{\pm} = \frac{1 \pm \not{v}}{2}$. ψ_+ is the positive energy component of the field (we use the notation adopted in [38]). v is identified with the heavy meson (quarkonium) 4-velocity with $v^2 = 1$. The equation of motion allows us to relate ψ_- to ψ_+ ,

$$\psi_-(x) = \frac{1}{2m_Q + iv \cdot D} i \not{D}_{\perp} \psi_+(x) \tag{8}$$

where $D_{\perp\mu} = D_{\mu} - (v \cdot D)v_{\mu}$. In the rest frame $v = (1, 0, 0, 0)$ we have $v \cdot D = D_t$ and $D_{\perp\mu} = (0, D_i)$.

Using (7) and (8) $Q(x)$ can be expressed in terms of $\psi_+(x)$,

$$Q(x) = e^{-im_Q v \cdot x} \left(1 + \frac{1}{2m_Q + iv \cdot D} i \not{D}_{\perp} \right) \psi_+(x), \tag{9}$$

a nonlocal expression which can be expanded:

$$Q(x) = e^{-im_Q v \cdot x} \left(1 + \frac{i \not{D}_\perp}{2m_Q} + \frac{(-iv \cdot D)}{2m_Q} \frac{i \not{D}_\perp}{2m_Q} + \dots \right) \psi_+(x). \quad (10)$$

The power counting of the various operators is set within NRQCD: $D_t \sim \tilde{v}^2$, $D_\perp \sim \tilde{v}$ and $\psi_+ \sim \tilde{v}^{3/2}$, where $\tilde{v} = |\vec{v}| \ll 1$ is the relative heavy quark 3-velocity in the hadron rest frame [33]. Therefore, the second term in Eq. (10) is $\mathcal{O}(\tilde{v} \times \tilde{v}^{3/2})$, the third one is $\mathcal{O}(\tilde{v}^3 \times \tilde{v}^{3/2})$. From now on, the power $\tilde{v}^{3/2}$ for each quark field will be omitted in the power counting of the operators.

The QCD Lagrangian expressed in terms of ψ_+

$$\mathcal{L}_{QCD} = \bar{\psi}_+(x) \left(iv \cdot D + i \not{D}_\perp \frac{1}{2m_Q + iv \cdot D} i \not{D}_\perp \right) \psi_+(x) \quad (11)$$

can be expanded:

$$\begin{aligned} \mathcal{L}_{QCD} &= \bar{\psi}_+(x) \left(iv \cdot D + \frac{(iD_\perp)^2}{2m_Q} + \frac{g}{4m_Q} \sigma \cdot G_\perp + \frac{i \not{D}_\perp}{2m_Q} \frac{(-iv \cdot D)}{2m_Q} (i \not{D}_\perp) + \dots \right) \psi_+(x) \\ &= \mathcal{L}_0 + \mathcal{L}_1 + \dots \end{aligned} \quad (12)$$

In this expression $G_{\perp\mu\nu}$ is $G_{\perp\mu\nu} = (g_{\mu\alpha} - v_\mu v_\alpha)(g_{\nu\beta} - v_\nu v_\beta)G^{\alpha\beta}$. In the rest frame $G_{\perp\mu\nu} = G_{ij}$ for $\mu = i = 1, 2, 3$ and $\nu = j = 1, 2, 3$, while the other components vanish. The power counting of the chromoelectric field components $E_i = G_{0i}$ and of the chromomagnetic ones $B_i = \frac{1}{2}\epsilon_{ijk}G^{jk}$ is \tilde{v}^3 and \tilde{v}^4 , respectively [32].

The first and second term in Eq. (12) are $\mathcal{O}(\tilde{v}^2)$ and provide the leading order Lagrangian

$$\mathcal{L}_0 = \bar{\psi}_+(x) \left(iv \cdot D + \frac{(iD_\perp)^2}{2m_Q} \right) \psi_+(x) \quad (13)$$

giving the equation of motion for $\psi_+(x)$

$$\left(iv \cdot D + \frac{(iD_\perp)^2}{2m_Q} \right) \psi_+(x) = 0. \quad (14)$$

The third and fourth term in Eq. (12) are $\mathcal{O}(\tilde{v}^4)$ and give the NLO Lagrangian

$$\mathcal{L}_1 = \mathcal{L}_{1,1} + \mathcal{L}_{1,2}$$

where

$$\begin{aligned} \mathcal{L}_{1,1} &= \bar{\psi}_+(x) \frac{g\sigma \cdot G_\perp}{4m_Q} \psi_+(x) \\ \mathcal{L}_{1,2} &= \bar{\psi}_+(x) \frac{i \not{D}_\perp}{2m_Q} \frac{(-iv \cdot D)}{2m_Q} (i \not{D}_\perp) \psi_+(x). \end{aligned} \quad (15)$$

Using the equation of motion together with $[i\vec{D}_\perp, iv \cdot D] = ig\gamma^\mu v^\nu G_{\mu\nu}$ and $\vec{D}_\perp \vec{D}_\perp = D_\perp^2 - \frac{1}{2}g\sigma \cdot G_\perp$, $\mathcal{L}_{1,2}$ can be expressed in the form

$$\begin{aligned}\mathcal{L}_{1,2} &= -\frac{1}{4m_Q^2} \left(\bar{\psi}_+(x) \left(-\frac{(iD_\perp)^4}{2m_Q} \right) \psi_+(x) + \bar{\psi}_+(x) \frac{g}{2} \sigma \cdot G_\perp \left(-\frac{(iD_\perp)^2}{2m_Q} \right) \psi_+(x) \right. \\ &\quad \left. + igv^\alpha \bar{\psi}_+(x) iD_\perp^\sigma G_{\alpha\sigma} \psi_+(x) + gv^\alpha \bar{\psi}_+(x) iD_{\perp\tau} \sigma^{\tau\sigma} G_{\alpha\sigma} \psi_+(x) \right) \\ &= \mathcal{L}_{1,2}^{(1)} + \mathcal{L}_{1,2}^{(2)} + \mathcal{L}_{1,2}^{(3)} + \mathcal{L}_{1,2}^{(4)}.\end{aligned}\tag{16}$$

$\mathcal{L}_{1,2}^{(2)}$ is of higher order in the \tilde{v} expansion. \mathcal{L}_1 can be arranged in the form

$$\mathcal{L}_1 = \mathcal{L}_1^A + \mathcal{L}_1^B\tag{17}$$

with

$$\mathcal{L}_1^A = \mathcal{L}_{1,1} + \mathcal{L}_{1,2}^{(4)} = \frac{1}{4m_Q} \bar{\psi}_+(x) A_{\tau\sigma} \sigma^{\tau\sigma} \psi_+(x)\tag{18}$$

$$\mathcal{L}_1^B = \mathcal{L}_{1,2}^{(1)} + \mathcal{L}_{1,2}^{(3)} = \frac{1}{4m_Q^2} \bar{\psi}_+(x) B \psi_+(x),\tag{19}$$

where in (18) we have factorized the leading $1/m_Q$ power. To deal with the antiquark, the QCD field $Q(x)$ is written as

$$Q(x) = e^{im_Q v \cdot x} X(x) = e^{im_Q v \cdot x} \left(X_+(x) + X_-(x) \right) = e^{im_Q v \cdot x} \left(1 + \frac{1}{2m_Q - iv \cdot D} i\vec{D}_\perp \right) X_-(x),\tag{20}$$

with X_- containing the negative energy component. The QCD Lagrangian written in terms of X_-

$$\mathcal{L}_{QCD} = \bar{X}_-(x) \left(-iv \cdot D + i\vec{D}_\perp \frac{1}{2m_Q - iv \cdot D} i\vec{D}_\perp \right) X_-(x)\tag{21}$$

is expanded as

$$\mathcal{L}_{QCD} = \bar{X}_-(x) \left(-iv \cdot D + \frac{(iD_\perp)^2}{2m_Q} + \dots \right) X_-(x).\tag{22}$$

The above expressions define the effective theory in which to work out the meson form factors.

4 Meson form factors in the effective theory

To obtain the meson form factors in the effective theory, we expand the weak current involving two heavy quarks $\bar{Q}'\Gamma Q$, with Γ a generic Dirac matrix:

$$\begin{aligned}\bar{Q}'(x)\Gamma Q(x) &= \bar{\psi}'_+(x) \left(1 - \frac{i\overleftarrow{\vec{D}}'_\perp}{2m_{Q'}} - \frac{1}{4m_{Q'}^2} (i\overleftarrow{\vec{D}}'_\perp)(iv' \cdot \overleftarrow{D}') + \dots \right) \\ &\quad \Gamma \left(1 + \frac{i\vec{D}_\perp}{2m_Q} + \frac{1}{4m_Q^2} (-iv \cdot \vec{D}) i\vec{D}_\perp + \dots \right) \psi_+(x)\end{aligned}\tag{23}$$

where $D'_{\perp\mu} = D_\mu - (v' \cdot D)v'_\mu$. Keeping terms up to $\mathcal{O}(1/m_Q^2)$, the current can be written as

$$\bar{Q}'(x)\Gamma Q(x) = J_0 + \left(\frac{J_{1,0}}{2m_Q} + \frac{J_{0,1}}{2m_{Q'}}\right) + \left(-\frac{J_{2,0}}{4m_Q^2} - \frac{J_{0,2}}{4m_{Q'}^2} + \frac{J_{1,1}}{4m_Q m_{Q'}}\right), \quad (24)$$

with J_i terms

$$\begin{aligned} J_0 &= \bar{\psi}'_+ \Gamma \psi_+ \\ J_{1,0} &= \bar{\psi}'_+ \Gamma i \vec{D}_\perp \psi_+ \\ J_{0,1} &= \bar{\psi}'_+ \left(-i \overleftarrow{D}'_\perp\right) \Gamma \psi_+ \\ J_{2,0} &= \bar{\psi}'_+ \Gamma \left(iv \cdot \vec{D}\right) i \vec{D}_\perp \psi_+ \\ J_{0,2} &= \bar{\psi}'_+ i \overleftarrow{D}'_\perp \left(iv' \cdot \overleftarrow{D}\right) \Gamma \psi_+ \\ J_{1,1} &= \bar{\psi}'_+ \left(-i \overleftarrow{D}'_\perp\right) \Gamma \left(i \vec{D}_\perp\right) \psi_+. \end{aligned} \quad (25)$$

Considering the power counting in \tilde{v} , Eq. (24) comprises terms up to $\mathcal{O}(\tilde{v}^3)$. The $\mathcal{O}(1/m_Q^3)$ terms involving three derivatives have not been included in (24), even though they can be of the same order in \tilde{v} of some terms appearing in the (nonlocal) corrections discussed in the following: we assume that they provide numerically suppressed effects.

We have neglected the perturbative α_s corrections. Considering such corrections the short distance expansion of the current in (24) would contain more operators and a set of matching coefficients would appear:

$$\bar{Q}'(x)\Gamma Q(x) = \sum_i C_i(\mu, w)(J_0)_i + \sum_j \left[\frac{B_j(\mu, w)}{2m_b}(J_{1,0})_j + \frac{B'_j(\mu, w)}{2m_c}(J_{0,1})_j \right] + \dots \quad (26)$$

The various structures in Eq. (24) can be identified with the first terms in each of the sums in (26). The coefficients C_i and $B_j^{(\prime)}$ are perturbatively expanded in α_s . Only the coefficients of the operators in (24) contribute at leading order in α_s , the others start at $\mathcal{O}(\alpha_s)$. At leading order in the inverse HQ mass expansion and at $\mathcal{O}(\alpha_s)$ one finds for the various currents:

$$\begin{aligned} \bar{Q}'Q &= \bar{\psi}'_+ \left(1 + \frac{\alpha_s}{\pi} C_S\right) \psi_+ + \dots \\ \bar{Q}'\gamma_5 Q &= \bar{\psi}'_+ \left(1 + \frac{\alpha_s}{\pi} C_P\right) \gamma_5 \psi_+ + \dots \\ \bar{Q}'\gamma_\mu Q &= \bar{\psi}'_+ \left[\left(1 + \frac{\alpha_s}{\pi} C_{V_1}\right) \gamma_\mu + \frac{\alpha_s}{\pi} C_{V_2} v_\mu + \frac{\alpha_s}{\pi} C_{V_3} v'_\mu\right] \psi_+ + \dots \\ \bar{Q}'\gamma_\mu \gamma_5 Q &= \bar{\psi}'_+ \left[\left(1 + \frac{\alpha_s}{\pi} C_{A_1}\right) \gamma_\mu + \frac{\alpha_s}{\pi} C_{A_2} v_\mu + \frac{\alpha_s}{\pi} C_{A_3} v'_\mu\right] \gamma_5 \psi_+ + \dots \\ \bar{Q}'\sigma^{\mu\nu} Q &= \bar{\psi}'_+ \left[\left(1 + \frac{\alpha_s}{\pi} C_{T_1}\right) \sigma^{\mu\nu} + \frac{\alpha_s}{\pi} C_{T_2} i(v^\mu \gamma^\nu - v^\nu \gamma^\mu) + \frac{\alpha_s}{\pi} C_{T_3} i(v'^\mu \gamma^\nu - v'^\nu \gamma^\mu)\right] \psi_+ + \dots \end{aligned} \quad (27)$$

The coefficients C_i have been computed in [47–49], the results for $B_j^{(\prime)}$ for the vector and axial vector currents are in [50]. In Sec. 5 we comment on the accuracy of using (24) instead of (26).

The B_c and $J/\psi, \eta_c$ matrix elements of the various terms in the expansion (24) can be expressed using the trace formalism [51]. In this formalism, the lowest-lying S-wave $\bar{b}c$ and $\bar{c}c$ bound states are described by 4×4 matrices [28]

$$H^{c\bar{b}}(v) = \frac{1 + \not{v}}{2} [B_c^{*\mu} \gamma_\mu - B_c \gamma_5] \frac{1 - \not{v}}{2} \quad (28)$$

$$H^{c\bar{c}}(v') = \frac{1 + \not{v}'}{2} [\Psi^{*\mu} \gamma_\mu - \eta_c \gamma_5] \frac{1 - \not{v}'}{2} \quad (29)$$

satisfying the relations $\not{v}H(v) = H(v) = -H(v)\not{v}$ and $H(v')\not{v}' = H(v') = -\not{v}'H(v')$. $B_c^{*\mu}$, B_c and $\Psi^{*\mu}$, η_c annihilate vector and pseudoscalar $\bar{b}c$ and $\bar{c}c$ mesons of velocity v and v' , respectively. They are normalized to $\sqrt{m_M}$ with M one of the mesons in the spin doublet. The trace formalism has been used to write the effective Lagrangians governing strong and radiative heavy quarkonium transitions in the soft-exchange approximation [52–54].

The x -dependence of the matrix element $\mathcal{M}_0(x) = \langle M'(v') | \bar{\psi}'_+(x) \Gamma \psi_+(x) | M(v) \rangle$ can be obtained exploiting the dependence in the effective theory [55, 56]:

$$|M(x)\rangle = e^{-i\tilde{\Lambda}v \cdot x} |M(0)\rangle \quad (30)$$

where, for a $Q\bar{Q}'$ meson,

$$\tilde{\Lambda} = m_H - m_Q - m_{Q'}. \quad (31)$$

Eq. (31) means that for the heavy quarkonium under scrutiny the heavy quark binding is generated by nonperturbative effects. B_c and $J/\psi, \eta_c$ are not considered as purely Coulombic states. The small binding energy scale $\tilde{\Lambda}$ can display a residual heavy quark mass dependence. From (30) and (31) we have

$$\mathcal{M}_0(x) = e^{-i\phi \cdot x} \mathcal{M}_0(0) \quad (32)$$

with $\phi = \tilde{\Lambda}v - \tilde{\Lambda}'v'$. For $B_c \rightarrow J/\psi(\eta_c)$, the binding energies $\tilde{\Lambda}^{(\prime)}$ are given by $\tilde{\Lambda} = m_{B_c} - m_b - m_c$ and $\tilde{\Lambda}' = m_{J/\psi(\eta_c)} - 2m_c$.

Using the trace formalism, we define

$$\langle M'(v') | \bar{\psi}'_+ \Gamma (iv \cdot \overrightarrow{D}) \psi_+ | M(v) \rangle = -\phi_K(w) \text{Tr} [\bar{H}'(v') \Gamma H(v)] \quad (33)$$

$$\langle M'(v') | \bar{\psi}'_+ (-iv' \cdot \overleftarrow{D}) \Gamma \psi_+ | M(v) \rangle = -\phi'_K(w) \text{Tr} [\bar{H}'(v') \Gamma H(v)] \quad (34)$$

with $w = v \cdot v'$. The same formalism allows us to parametrize the matrix elements of the various terms in (24). The matrix element of J_0 is written as

$$\langle M'(v') | J_0 | M(v) \rangle = -\Delta(w) \text{Tr} [\bar{H}'(v') \Gamma H(v)] \quad (35)$$

and involves the form factor $\Delta(w)$ obtained in [28]. The $1/m_{Q,Q'}$ terms involve the functions Δ_α and $\bar{\Delta}_\alpha$,

$$\langle M'(v') | \bar{\psi}'_+ \Gamma^\alpha i D_\alpha \psi_+ | M(v) \rangle = -\text{Tr} [\Delta_\alpha(v, v') \bar{H}'(v') \Gamma^\alpha H(v)] \quad (36)$$

$$\langle M'(v') | \bar{\psi}'_+ (-i \overleftarrow{D}_\alpha) \Gamma^\alpha \psi_+ | M(v) \rangle = -\text{Tr} [\bar{\Delta}_\alpha(v, v') \bar{H}'(v') \Gamma^\alpha H(v)] , \quad (37)$$

which are expressed in general as

$$\Delta_\alpha(v, v') = \Delta_+(w)(v + v')_\alpha + \Delta_-(w)(v - v')_\alpha - \Delta_3(w)\gamma_\alpha \quad (38)$$

$$\bar{\Delta}_\alpha(v, v') = \bar{\Delta}_+(w)(v + v')_\alpha + \bar{\Delta}_-(w)(v' - v)_\alpha - \bar{\Delta}_3(w)\gamma_\alpha. \quad (39)$$

Exploiting the relation

$$i\partial_\alpha(\bar{\psi}'_+\Gamma\psi_+) = \bar{\psi}'_+(i\overleftarrow{D}_\alpha)\Gamma\psi_+ + \bar{\psi}'_+\Gamma(i\overrightarrow{D}_\alpha)\psi_+ \quad (40)$$

and using (32), the functions in (38)-(39) can be connected to Δ in (35):

$$(\tilde{\Lambda} - \tilde{\Lambda}'w)\Delta = (\Delta_+ - \bar{\Delta}_+)(1 + w) + (\Delta_- + \bar{\Delta}_-)(1 - w) + (\Delta_3 - \bar{\Delta}_3) \quad (41)$$

$$(\tilde{\Lambda}w - \tilde{\Lambda}')\Delta = (\Delta_+ - \bar{\Delta}_+)(1 + w) + (\Delta_- + \bar{\Delta}_-)(w - 1) + (\Delta_3 - \bar{\Delta}_3). \quad (42)$$

Δ_i satisfy the equations

$$\Delta_+(1 + w) + \Delta_-(1 - w) + \Delta_3 = \phi_K \quad (43)$$

$$\bar{\Delta}_+(1 + w) + \bar{\Delta}_-(1 - w) + \bar{\Delta}_3 = \phi'_K \quad (44)$$

$$\Delta_3 = \bar{\Delta}_3 \quad (45)$$

obtained using Eqs. (14) and (32), with solutions

$$\Delta_+(w) = -\frac{\Delta_3(w)}{(1 + w)} + \frac{\Delta(w)}{2(1 + w)} \left(\tilde{\Lambda}w - \tilde{\Lambda}' \right) + \frac{\phi_K(w) + \phi'_K(w)}{2(1 + w)} \quad (46)$$

$$\bar{\Delta}_+(w) = -\frac{\Delta_3(w)}{(1 + w)} + \frac{\Delta(w)}{2(1 + w)} \left(\tilde{\Lambda}'w - \tilde{\Lambda} \right) + \frac{\phi_K(w) + \phi'_K(w)}{2(1 + w)} \quad (47)$$

$$\Delta_-(w) = \frac{\Delta(w)}{2(w - 1)} \left(\tilde{\Lambda}w - \tilde{\Lambda}' \right) - \frac{\phi_K(w) - \phi'_K(w)}{2(w - 1)} \quad (48)$$

$$\bar{\Delta}_-(w) = \frac{\Delta(w)}{2(w - 1)} \left(\tilde{\Lambda}'w - \tilde{\Lambda} \right) + \frac{\phi_K(w) - \phi'_K(w)}{2(w - 1)}. \quad (49)$$

The functions Δ_- and $\bar{\Delta}_-$ are finite at $w = 1$ if $\phi_K(w)$ and $\phi'_K(w)$ are related by the condition

$$\phi_K(w) - \phi'_K(w) = \left(\tilde{\Lambda} - \tilde{\Lambda}' \right) \Delta(w). \quad (50)$$

Solving Eqs. (41)-(44) after an expansion of the universal functions close to $w = 1$ we find that Eq. (50) holds for $w = 1$. There is an interesting analogy to the case of heavy-light mesons, where $\phi_K(w) = \phi'_K(w)$ and $\phi_K(1)$ is proportional to the heavy quark kinetic energy [45]. If the relation holds for all values of w , the heavy-light case is recovered for $\tilde{\Lambda} = \tilde{\Lambda}'$. The condition

(50) allows us to obtain:

$$\Delta_+(w) = -\frac{\Delta_3(w)}{(w+1)} + \Delta(w) \frac{\tilde{\Lambda}(w-1)}{2(1+w)} + \frac{\phi_K(w)}{w+1} \quad (51)$$

$$\bar{\Delta}_+(w) = -\frac{\Delta_3(w)}{(w+1)} + \Delta(w) \frac{\tilde{\Lambda}'(w-1)}{2(1+w)} + \frac{\phi'_K(w)}{w+1} \quad (52)$$

$$\Delta_-(w) = \frac{\Delta(w)}{2} \tilde{\Lambda} \quad (53)$$

$$\bar{\Delta}_-(w) = \frac{\Delta(w)}{2} \tilde{\Lambda}' . \quad (54)$$

The matrix elements of the operators with $1/m_{Q,Q'}$ in (24) are expressed in terms of Δ_i :

$$\begin{aligned} \frac{1}{2m_Q} \langle M'(v') | J_{1,0} | M(v) \rangle = & -\frac{1}{2m_Q} \left\{ -\text{Tr} [\bar{H}'(v') \Gamma H(v)] \left[\frac{\Delta_3(w)}{1+w} + \frac{w}{1+w} (\phi_K(w) - \Delta(w) \tilde{\Lambda}) \right] \right. \\ & + \text{Tr} [\bar{H}'(v') \Gamma \psi' H(v)] \left[-\frac{\Delta_3(w)}{1+w} + \frac{1}{1+w} (\phi_K(w) - \Delta(w) \tilde{\Lambda}) \right] \\ & \left. - [\gamma^\beta \bar{H}'(v') \Gamma \gamma_\beta H(v)] \Delta_3(w) \right\} \end{aligned} \quad (55)$$

$$\begin{aligned} \frac{1}{2m_{Q'}} \langle M'(v') | J_{0,1} | M(v) \rangle = & -\frac{1}{2m_{Q'}} \left\{ -\text{Tr} [\bar{H}'(v') \Gamma H(v)] \left[\frac{\Delta_3(w)}{1+w} + \frac{w}{1+w} (\phi_K(w) - \Delta(w) \tilde{\Lambda}) \right] \right. \\ & + \text{Tr} [\bar{H}'(v') \psi \Gamma H(v)] \left[-\frac{\Delta_3(w)}{1+w} + \frac{1}{1+w} (\phi_K(w) - \Delta(w) \tilde{\Lambda}) \right] \\ & \left. - [\gamma^\beta \bar{H}'(v') \gamma_\beta \Gamma H(v)] \Delta_3(w) \right\}. \end{aligned} \quad (56)$$

To consider the $1/m_Q^2$ terms we define

$$\langle M'(v') | \bar{\psi}'_+ (-i \overleftarrow{D}_\alpha) \Gamma^{\alpha\beta} (i \overrightarrow{D}_\beta) \psi_+ | M(v) \rangle = -\text{Tr} [\psi_{\alpha\beta}(v, v') \bar{H}'(v') \Gamma^{\alpha\beta} H(v)] . \quad (57)$$

The function $\psi_{\alpha\beta}(v, v')$ is written in terms of its symmetric ψ^S and antisymmetric ψ^A parts

$$\psi_{\alpha\beta} = \frac{1}{2} [\psi_{\alpha\beta}^S + \psi_{\alpha\beta}^A] \quad (58)$$

which can be parametrized as

$$\begin{aligned} \psi_{\alpha\beta}^S = & \psi_1^S(w) g_{\alpha\beta} + \psi_2^S(w) (v+v')_\alpha (v+v')_\beta + \psi_3^S(w) (v-v')_\alpha (v-v')_\beta \\ & + \psi_4^S(w) [(v+v')_\alpha \gamma_\beta + (v+v')_\beta \gamma_\alpha] + \psi_5^S(w) [(v-v')_\alpha \gamma_\beta + (v-v')_\beta \gamma_\alpha] \\ & + \psi_6^S(w) [(v+v')_\alpha (v-v')_\beta + (v+v')_\beta (v-v')_\alpha] \end{aligned} \quad (59)$$

$$\begin{aligned} \psi_{\alpha\beta}^A = & \psi_1^A(w) [v_\alpha v'_\beta - v_\beta v'_\alpha] + \psi_2^A(w) [(v-v')_\alpha \gamma_\beta - (v-v')_\beta \gamma_\alpha] \\ & + \psi_3^A(w) i\sigma_{\alpha\beta} + \psi_4^A(w) [(v+v')_\alpha \gamma_\beta - (v+v')_\beta \gamma_\alpha]. \end{aligned} \quad (60)$$

The hadronic matrix element of $J_{1,1}$ in (24) can be expressed in terms of $\psi_{\alpha\beta}$. For $J_{2,0}$ and $J_{0,2}$ integration by parts is also needed. We obtain:

$$\begin{aligned}
\langle M'(v') | J_{2,0} | M(v) \rangle = & -(\tilde{\Lambda} - w\tilde{\Lambda}') \left\{ \Delta_+ \left(\text{Tr} [\bar{H}'(v') \Gamma H(v)] + \text{Tr} [\bar{H}'(v') \Gamma \not{p}' H(v)] \right) \right. \\
& + \Delta_- \left(\text{Tr} [\bar{H}'(v') \Gamma H(v)] - \text{Tr} [\bar{H}'(v') \Gamma \not{p}' H(v)] \right) \\
& \left. - \Delta_3 \text{Tr} [\bar{H}'(v') \Gamma \gamma^\beta H(v) \gamma_\beta] - \phi_K \text{Tr} [\bar{H}'(v') \Gamma H(v)] \right\} \\
& - \frac{1}{2} \left((1+w)\psi_2^S - (1-w)\psi_3^S - 2w\psi_6^S + \psi_1^A - (\psi_4^S - \psi_5^S + \psi_2^A - \psi_4^A) \right) \\
& \times \left(-w \text{Tr} [\bar{H}'(v') \Gamma H(v)] + \text{Tr} [\bar{H}'(v') \Gamma \not{p}' H(v)] \right) \\
& - \frac{1}{2} \left((1+w)(\psi_4^S + \psi_4^A) + (1-w)(\psi_5^S + \psi_2^A) + \psi_3^A \right) \\
& \times \left(\text{Tr} [\bar{H}'(v') \Gamma H(v)] + \text{Tr} [\bar{H}'(v') \Gamma \gamma^\beta H(v) \gamma_\beta] \right), \tag{61}
\end{aligned}$$

$$\begin{aligned}
\langle M'(v') | J_{0,2} | M(v) \rangle = & (\tilde{\Lambda}w - \tilde{\Lambda}') \left\{ \bar{\Delta}_+ \left(\text{Tr} [\bar{H}'(v') \Gamma H(v)] + \text{Tr} [\bar{H}'(v') \not{p} \Gamma H(v)] \right) \right. \\
& + \bar{\Delta}_- \left(\text{Tr} [\bar{H}'(v') \Gamma H(v)] - \text{Tr} [\bar{H}'(v') \not{p} \Gamma H(v)] \right) \\
& \left. - \bar{\Delta}_3 \text{Tr} [\gamma_\beta \bar{H}'(v') \gamma^\beta \Gamma H(v)] - \phi'_K \text{Tr} [\bar{H}'(v') \Gamma H(v)] \right\} \\
& - \frac{1}{2} \left((1+w)\psi_2^S - (1-w)\psi_3^S + 2w\psi_6^S + \psi_1^A - \psi_4^S - \psi_5^S - \psi_2^A - \psi_4^A \right) \\
& \times \left(-w \text{Tr} [\bar{H}'(v') \Gamma H(v)] + \text{Tr} [\bar{H}'(v') \not{p} \Gamma H(v)] \right) \\
& - \frac{1}{2} \left((1+w)(\psi_4^S - \psi_4^A) - (1-w)(\psi_5^S - \psi_2^A) + \psi_3^A \right) \\
& \times \left(\text{Tr} [\bar{H}'(v') \Gamma H(v)] + \text{Tr} [\gamma_\beta \bar{H}'(v') \gamma^\beta \Gamma H(v)] \right), \tag{62}
\end{aligned}$$

$$\begin{aligned}
\langle M'(v') | J_{1,1} | M(v) \rangle = & -\frac{1}{2}(\psi_1^S - \psi_3^A) \left(w \text{Tr} [\bar{H}'(v') \Gamma H(v)] - \text{Tr} [\bar{H}'(v') \not{p} \Gamma H(v)] \right. \\
& - \text{Tr} [\bar{H}'(v') \Gamma \not{p}' H(v)] + \text{Tr} [\bar{H}'(v') \gamma^\beta \Gamma \gamma_\beta H(v)] \left. \right) \\
& - \frac{1}{2}(\psi_2^S - \psi_3^S + \psi_1^A) \left(w^2 \text{Tr} [\bar{H}'(v') \Gamma H(v)] \right. \\
& - w (\text{Tr} [\bar{H}'(v') \not{p} \Gamma H(v)] + \text{Tr} [\bar{H}'(v') \Gamma \not{p}' H(v)]) + \text{Tr} [\bar{H}'(v') \not{p} \Gamma \not{p}' H(v)] \left. \right) \\
& - \frac{1}{2}(\psi_4^S + \psi_2^A + \psi_5^S + \psi_4^A) \left(\text{Tr} [\bar{H}'(v') (\not{p} - w) \Gamma H(v)] \right. \\
& + \text{Tr} [\bar{H}'(v') (\not{p} - w) \Gamma \gamma^\beta H(v) \gamma_\beta] \left. \right) \\
& - \frac{1}{2}(\psi_4^S + \psi_2^A - \psi_5^S - \psi_4^A) \left(\text{Tr} [\bar{H}'(v') \Gamma (\not{p}' - w) H(v)] \right. \\
& + \text{Tr} [\gamma^\beta \bar{H}'(v') \gamma^\beta \Gamma (\not{p}' - w) H(v)] \left. \right) \\
& - \frac{1}{2}\psi_3^A \left(\text{Tr} [\bar{H}'(v') \Gamma H(v)] + \text{Tr} [\gamma^\beta \bar{H}'(v') \gamma^\beta \Gamma \gamma_\alpha H(v) \gamma_\alpha] \right. \\
& + \text{Tr} [\gamma_\beta \bar{H}'(v') \gamma^\beta \Gamma H(v)] + \text{Tr} [\bar{H}'(v') \Gamma \gamma^\beta H(v) \gamma_\beta] \left. \right). \tag{63}
\end{aligned}$$

Using Eqs. (14), (33) and (34) two relations follow:

$$\begin{aligned}
2m_Q \phi_K = & \tilde{\Lambda}'(\tilde{\Lambda} \Delta - \phi_K)(w - 1) \\
& - \frac{1}{2}(3\psi_1^S + (1 - w^2)(\psi_2^S + \psi_3^S - 2\psi_6^S) + 2(w - 1)(\psi_4^S - \psi_5^S)) \tag{64}
\end{aligned}$$

$$\begin{aligned}
2m_Q \phi'_K = & \tilde{\Lambda}(\tilde{\Lambda}' \Delta - \phi'_K)(w - 1) \\
& - \frac{1}{2}(3\psi_1^S + (1 - w^2)(\psi_2^S + \psi_3^S + 2\psi_6^S) + 2(w - 1)(\psi_4^S + \psi_5^S)). \tag{65}
\end{aligned}$$

In addition to the corrections obtained by the expansion of the weak currents, we must consider the corrections to the states. Using Eqs. (18),(19), they can be written as

$$\begin{aligned}
\langle M'(v') | i \int d^4x T[J_0(0) \mathcal{L}_1(x)] | M(v) \rangle = & -\frac{1}{2m_Q^2} \chi_1(w) \text{Tr} [\bar{H}'(v') \Gamma H(v)] \\
& - \frac{1}{4m_Q} \text{Tr} \left[\chi_{2\mu\nu}(w) \bar{H}'(v') \Gamma P_+ \left(-\frac{i}{2} \right) \sigma^{\mu\nu} H(v) \right] \tag{66} \\
= & \mathcal{M}_3 + \mathcal{M}_4
\end{aligned}$$

and, considering the term analogous to \mathcal{L}_1 for the antiquark in Eq.(22),

$$\begin{aligned} \langle M'(v') | i \int d^4x T[J_0(0)\mathcal{L}'_1(x)] | M(v) \rangle &= -\frac{1}{2m_{Q'}^2} \bar{\chi}_1(w) \text{Tr} [\bar{H}'(v') \Gamma H(v)] \\ &\quad - \frac{1}{4m_{Q'}} \text{Tr} \left[\bar{\chi}_{2\mu\nu}(w) \bar{H}'(v') \left(-\frac{i}{2} \right) \sigma^{\mu\nu} P'_+ \Gamma H(v) \right] \\ &= \bar{\mathcal{M}}_3 + \bar{\mathcal{M}}_4 . \end{aligned} \quad (67)$$

We deal with such corrections parametrizing the functions χ_i , $\bar{\chi}_i$ as

$$\chi_{2\mu\nu} = \chi_2^A i\sigma_{\mu\nu} + \chi_2^B (v_\mu \gamma_\nu - v_\nu \gamma_\mu) + \chi_2^C (v'_\mu \gamma_\nu - v'_\nu \gamma_\mu) \quad (68)$$

$$\bar{\chi}_{2\mu\nu} = \bar{\chi}_2^A i\sigma_{\mu\nu} + \bar{\chi}_2^B (v_\mu \gamma_\nu - v_\nu \gamma_\mu) + \bar{\chi}_2^C (v'_\mu \gamma_\nu - v'_\nu \gamma_\mu). \quad (69)$$

Using $\not{v}H(v) = H(v)$ and $\bar{H}'(v')\not{v}' = \bar{H}'(v')$ one sees that χ_2^B and $\bar{\chi}_2^C$ do not contribute to the matrix elements. Hence, \mathcal{M}_4 and $\bar{\mathcal{M}}_4$ are given by:

$$\begin{aligned} \mathcal{M}_4 &= -\frac{1}{4m_Q} \chi_2^A(w) d_M \text{Tr} [\bar{H}'(v') \Gamma H(v)] \\ &\quad - \frac{1}{4m_Q} \chi_2^C(w) \left[\text{Tr} [\bar{H}'(v') \Gamma H(v)] + \text{Tr} [\gamma^\nu \bar{H}'(v') \Gamma P_+ \not{v}' \gamma_\nu H(v)] \right] \end{aligned} \quad (70)$$

$$\begin{aligned} \bar{\mathcal{M}}_4 &= -\frac{1}{4m_{Q'}} \bar{\chi}_2^A(w) d'_M \text{Tr} [\bar{H}'(v') \Gamma H(v)] \\ &\quad + \frac{1}{4m_{Q'}} \bar{\chi}_2^B(w) \left[\text{Tr} [\bar{H}'(v') \Gamma H(v)] + \text{Tr} [\gamma^\nu \bar{H}'(v') \gamma_\nu \not{v}' P'_+ \Gamma H(v)] \right] , \end{aligned} \quad (71)$$

with $M = P, V$ and $d_P^{(\prime)} = 3$, $d_V^{(\prime)} = -1$.

The above relations can be used to write the various form factors in terms of universal functions. The resulting expressions are collected in Appendix B.

5 Numerical results

We present a numerical analysis based on the relations in App. B. Since the number of universal functions increases including the various terms, we shall limit the study neglecting $\mathcal{O}(1/m^2)$ terms (with m generically m_b , m_c). This already produces interesting relations. For example, the combinations

$$F_1(w) = \frac{\phi_K(w) - \Delta(w)\tilde{\Lambda}}{2} + \Delta_3(w) \quad (72)$$

$$F_2(w) = \frac{\phi_K(w) - \Delta(w)\tilde{\Lambda}}{2} - \Delta_3(w) \quad (73)$$

$$K(w) = 3\chi_2^A(w) + 2(w-1)\chi_2^C(w) \quad (74)$$

$$\bar{K}(w) = 3\bar{\chi}_2^A(w) - 2(w-1)\bar{\chi}_2^B(w) \quad (75)$$

allow us to simplify the expressions for the form factors and to establish relations among them. For the $B_c \rightarrow \eta_c$ form factors we find:

$$\begin{aligned}
h_+(w) &= \Delta(w) + \frac{1}{4m_b}K(w) + \frac{1}{4m_c}\bar{K}(w) \\
h_-(w) &= \left(-\frac{1}{m_b} + \frac{1}{m_c}\right)F_1(w) \\
h_T(w) &= \Delta(w) - \left(\frac{1}{m_b} + \frac{1}{m_c}\right)F_1(w) + \frac{1}{4m_b}K(w) + \frac{1}{4m_c}\bar{K}(w) \\
h_S(w) &= \Delta(w) - \left(\frac{1}{m_b} + \frac{1}{m_c}\right)\frac{w-1}{w+1}F_1(w) + \frac{1}{4m_b}K(w) + \frac{1}{4m_c}\bar{K}(w) .
\end{aligned} \tag{76}$$

For $B_c \rightarrow J/\psi$ form factors we obtain:

$$\begin{aligned}
h_V(w) &= \Delta(w) + \frac{1}{4m_b}[-4F_1(w) + K(w)] - \frac{1}{4m_c}[\bar{\chi}_2^A(w) + 2(F_1(w) + F_2(w))] \\
h_{A_1}(w) &= \Delta(w) - \frac{1}{m_b}\frac{w-1}{w+1}F_1(w) - \frac{1}{2m_c}\frac{w-1}{w+1}(F_1(w) + F_2(w)) + \frac{1}{4m_b}K(w) - \frac{1}{4m_c}\bar{\chi}_2^A(w) \\
h_{A_2}(w) &= \frac{1}{m_c}\left[\frac{1}{2(1+w)}(F_1(w) + 3F_2(w)) + \frac{1}{2}\bar{\chi}_2^B(w)\right] \\
h_{A_3}(w) &= \Delta(w) - \frac{1}{m_b}F_1(w) + \frac{1}{4m_b}K(w) - \frac{1}{2m_c(1+w)}[wF_1(w) + (w-2)F_2(w)] \\
&\quad - \frac{1}{4m_c}[\bar{\chi}_2^A(w) + 2\bar{\chi}_2^B(w)] \\
h_{T_1}(w) &= \Delta(w) + \frac{1}{4m_b}K(w) - \frac{1}{4m_c}\bar{\chi}_2^A(w) \\
h_{T_2}(w) &= -\frac{1}{m_b}F_1(w) + \frac{1}{2m_c}(F_1(w) + F_2(w)) \\
h_{T_3}(w) &= -\frac{1}{2m_c}\left[\bar{\chi}_2^B(w) - \frac{1}{1+w}(F_1(w) + 3F_2(w))\right] \\
h_P(w) &= \Delta(w) + \frac{1}{4m_b}[-4F_1(w) + K(w)] - \frac{1}{4m_c}[4F_2(w) + \bar{K}(w) - 2\bar{\chi}_2^A(w)] .
\end{aligned} \tag{77}$$

The lattice results for $V(q^2)$ and $A_{1,2,0}(q^2)$ [13] are translated into $h_V(w)$ and $h_{A_{1,2,3}}(w)$ in Fig. 1. Keeping the leading term in the heavy quark mass expansion leads to $h_V = h_{A_1} = h_{A_3} = \Delta$ and $h_{A_2} = 0$, relations badly violated by the results obtained by lattice QCD, as shown in Fig. 1. Therefore, subleading terms must be considered. In this respect, the same lattice QCD results can be used to predict other form factors, exploiting Eqs. (76) and (77). A number of universal functions can be determined. In particular, we find:




Figure 1: $B_c \rightarrow J/\psi$ form factors h_V , h_{A_1} , h_{A_2} and h_{A_3} obtained using the lattice QCD results for V , A_1 , A_2 , A_0 .

$$\begin{aligned} \frac{\phi_K(w) - \Delta(w)\tilde{\Lambda}}{2} &= \frac{m_c}{2(m_b + 3m_c)}(1 + w) \left(m_b h_{A_1}(w) \right. \\ &\quad \left. + m_c (h_{A_2}(w) + h_{A_3}(w)) - (m_b + m_c) h_V(w) \right) \end{aligned} \quad (78)$$

$$\begin{aligned} \Delta_3(w) &= -\frac{m_c}{2(m_b + 3m_c)}(1 + w) \left(-2m_b h_{A_1}(w) \right. \\ &\quad \left. + (m_b + m_c) (h_{A_2}(w) + h_{A_3}(w)) + (m_b - m_c) h_V(w) \right) \end{aligned} \quad (79)$$

$$\bar{\chi}_B^2(w) = m_c (h_{A_2}(w) - h_{A_3}(w) + h_V(w)). \quad (80)$$

Such functions are displayed in Fig. 2. In the figure the relations are applied to the full kinematical range, a useful extrapolation for comparing with other calculations. This allows




Figure 2: Functions in Eqs. (78), (79) and (80) extrapolated to the full kinematical range.

us to derive other form factors near the zero recoil point. For $B_c \rightarrow J/\psi$ the relations hold:

$$h_{T_1}(w) = \frac{1}{2} \left((1+w)h_{A_1}(w) - (w-1)h_V(w) \right) \quad (81)$$

$$h_{T_2}(w) = \frac{1+w}{2(m_b+3m_c)} \left((m_b-3m_c)h_{A_1}(w) + 2m_c(h_{A_2}(w) + h_{A_3}(w)) - (m_b-m_c)h_V(w) \right) \quad (82)$$

$$h_{T_3}(w) = h_{A_3}(w) - h_V(w) \quad (83)$$

$$h_P(w) = \frac{1}{m_b+3m_c} \left((1+w)(m_b h_{A_1}(w) + 2m_c h_V(w)) + (-m_b + (w-2)m_c)h_{A_2}(w) - (w m_b + (2w-1)m_c)h_{A_3}(w) \right). \quad (84)$$

These functions are depicted in Fig. 3 extrapolated to the full kinematical range. For $B_c \rightarrow \eta_c$, $h_-(w)$ and two form factor differences can be derived:

$$h_-(w) = \frac{m_b-m_c}{2(m_b+3m_c)}(1+w) \left(3h_{A_1}(w) - h_{A_2}(w) - h_{A_3}(w) - 2h_V(w) \right) \quad (85)$$

$$h_T(w) - h_+(w) = -\frac{m_b+m_c}{2(m_b+3m_c)}(1+w) \left(3h_{A_1}(w) - h_{A_2}(w) - h_{A_3}(w) - 2h_V(w) \right) \quad (86)$$

$$h_T(w) - h_S(w) = -\frac{m_b+m_c}{(m_b+3m_c)} \left(3h_{A_1}(w) - h_{A_2}(w) - h_{A_3}(w) - 2h_V(w) \right). \quad (87)$$




Figure 3: Tensor $B_c \rightarrow J/\psi$ form factors obtained applying Eqs. (81), (82), (83) and (84) in the full kinematical range and using lattice QCD results for V and $A_{1,2,0}$.

Their extrapolations are displayed in Fig. 4. The value of the universal functions at $w = 1$ is not predicted, however from Eqs. (81) and (76) the relations $h_{T_1}(w = 1) = h_{A_1}(w = 1)$ and $h_S(w = 1) = h_+(w = 1)$ are obtained, respectively. It is interesting to observe that if we consider the limit $m_b \rightarrow \infty$ keeping the $1/m_c$ terms, some results remain unaffected. This is the case of Eqs. (81) and (83), and of the relations $h_{T_1}(w = 1) = h_{A_1}(w = 1)$ and $h_S(w = 1) = h_+(w = 1)$. In the η_c case, the relations (85)-(87) are rescaled replacing $\frac{m_b \pm m_c}{m_b + 3m_c} \rightarrow 1$.

The relations among the various form factors are modified using the expansion (27). In




Figure 4: $B_c \rightarrow \eta_c$ form factors in Eqs. (85), (86) and (87) extended to the full kinematical range, obtained using lattice QCD results for V and $A_{1,2,0}$.

particular, the right-hand sides of Eqs. (81)-(87) acquire the extra terms:

$$\begin{aligned}
\delta h_{T_1}(w) &= -\frac{1}{2} \frac{\alpha_s}{\pi} [2C_{T_1} + (w-1)(C_{T_2} - C_{T_3} + C_{V_1}) - (w+1)C_{A_1}] \Delta(w) \\
\delta h_{T_2}(w) &= -\frac{1+w}{2} \frac{\alpha_s}{\pi} [(C_{T_2} + C_{T_3}) + \frac{m_b - m_c}{m_b + 3m_c}(C_{V_1} - C_{A_1}) - \frac{2m_c}{(m_b + 3m_c)}(C_{A_2} + C_{A_3})] \Delta(w) \\
\delta h_{T_3}(w) &= -\frac{\alpha_s}{\pi} [C_{T_2} + C_{V_1} - C_{A_1} - C_{A_3}] \Delta(w) \\
\delta h_P(w) &= -\frac{\alpha_s}{\pi} [C_P - \frac{2m_c}{(m_b + 3m_c)}(1+w)C_{V_1} - \frac{m_b - (2w-1)m_c}{(m_b + 3m_c)}C_{A_1} \\
&\quad + \frac{m_b - (w-2)m_c}{m_b + 3m_c}C_{A_2} + \frac{wm_b + (2w-1)m_c}{(m_b + 3m_c)}C_{A_3}] \Delta(w) \\
\delta h_-(w) &= -\frac{1+w}{2} \frac{\alpha_s}{\pi} [\frac{m_b - m_c}{(m_b + 3m_c)}(2C_{V_1} - 2C_{A_1} + C_{A_2} + C_{A_3}) + C_{V_2} - C_{V_3}] \Delta(w) \\
\delta h_{T+}(w) &= -\frac{1}{2} \frac{\alpha_s}{\pi} [2C_{T_1} - 2C_{T_2} + 2C_{T_3} - \frac{2[m_b(w+2) + m_c(w+4)]}{m_b + 3m_c}C_{V_1} \\
&\quad - (1+w)(C_{V_2} + C_{V_3}) + \frac{m_b + m_c}{m_b + 3m_c}(1+w)(2C_{A_1} - C_{A_2} - C_{A_3})] \Delta(w) \\
\delta h_{TS}(w) &= -\frac{\alpha_s}{\pi} [C_{T_1} - C_{T_2} + C_{T_3} - C_S + \frac{m_b + m_c}{m_b + 3m_c}(-2C_{V_1} + 2C_{A_1} - C_{A_2} - C_{A_3})] \Delta(w) .
\end{aligned}$$

To assess the size of the new contributions we choose $w = 1$ and $\Delta(1) \simeq 1$ [29]. Setting $\mu = \sqrt{m_c m_b}$ and $\alpha_s(\mu) = 0.27$, we find $\delta h_{T_1}(1) \simeq -0.065$, $\delta h_{T_2}(1) \simeq -0.02$, $\delta h_{T_3}(1) \sim \mathcal{O}(10^{-4})$, $\delta h_P(1) \simeq 0.025$, $\delta h_-(1) \simeq -0.04$, $\delta h_{T_+}(1) \simeq -0.01$, $\delta h_{TS}(1) \simeq -0.09$. Such results compared to the values of the form factors at $w = 1$ in Figs. 3, 4 show that the impact of the matching coefficients is small.

The difference of our approach with other calculations based on NRQCD must be noticed. We have used the heavy quark expansion and the NRQCD power counting to relate the various form factors close to the zero recoil point and exploited lattice QCD to determine, e.g., the tensor form factors. A systematic control of the error can be achieved by the expansion. Then the results are extrapolated to small q^2 . Different analyses are carried out at $q^2 \simeq 0$, where a perturbative approach to the form factor calculation can be attempted, and extrapolated to higher values of q^2 after a normalization to the lattice QCD results [20]. Numerically, the results match for the form factors T_1 and T_2 in the basis (3), while T_0 is affected by a larger uncertainty.

A final remark is in order about the extrapolation from the kinematical range close to the zero-recoil point $w \sim 1$ to the full kinematical range. There are methods to constrain the extrapolation, starting from the form factors evaluated in few points and using unitarity constraints and the dispersion matrix [57–64]. The application of such methods to the form factors discussed here is deferred to a dedicated analysis. Their availability strengthens the significance of our study, allowing us to foresee the control of the hadronic uncertainties not only near the zero-recoil point but in the full kinematical range.

6 Conclusions

Using the heavy quark expansion, the heavy quark spin symmetry and NRQCD power counting we have expressed the form factors parametrizing the matrix elements $\langle J/\psi(\eta_c) | \bar{c} \Gamma_i b | B_c \rangle$ in terms of universal functions near the zero-recoil point. This can be done for the various operators in the generalized low-energy Hamiltonian Eq. (1), establishing relations among form factors in a kinematical range around the zero-recoil point. Lattice QCD results for the matrix element of the Standard Model operator between B_c and J/ψ allow us to predict the pseudoscalar and tensor form factors. $B_c \rightarrow \eta_c$ form factors are also related to the previous ones, obtaining h_- and the differences between the remaining form factors. We have also presented the results of the extrapolation to the full kinematical range. The relations worked out in our study can be checked if further information from lattice QCD is available. The effort is to efficiently control the hadronic uncertainties affecting the predictions for semileptonic $B_c \rightarrow J/\psi, \eta_c$ decays in the Standard Model and beyond.

Acknowledgements. This study has been carried out within the INFN project (Iniziativa Specifica) QFT-HEP.

A Form factors basis h_i for $B_c \rightarrow \eta_c, J/\psi$

The basis of form factors h_i is defined below.

$B_c \rightarrow \eta_c$:

$$\begin{aligned}\langle P(v') | \bar{Q}' \gamma_\mu Q | B_c(v) \rangle &= \sqrt{m_P m_{B_c}} [h_+(w) (v + v')_\mu + h_-(w) (v - v')_\mu] \\ \langle P(v') | \bar{Q}' Q | B_c(v) \rangle &= \sqrt{m_P m_{B_c}} h_S(w) (1 + w) \\ \langle P(v') | \bar{Q}' \sigma_{\mu\nu} Q | B_c(v) \rangle &= -i \sqrt{m_P m_{B_c}} h_T(w) (v_\mu v'_\nu - v_\nu v'_\mu)\end{aligned}\tag{A.1}$$

with $P = \eta_c$, $v = \frac{p}{m_{B_c}}$, $v' = \frac{p'}{m_P}$ and $w = v \cdot v'$.

$B_c \rightarrow J/\psi$:

$$\begin{aligned}\langle V(v', \epsilon) | \bar{Q}' \gamma_\mu Q | B_c(v) \rangle &= i \sqrt{m_V m_{B_c}} h_V(w) \epsilon_{\mu\nu\alpha\beta} \epsilon^{*\nu} v'^\alpha v^\beta \\ \langle V(v', \epsilon) | \bar{Q}' \gamma_\mu \gamma_5 Q | B_c(v) \rangle &= \sqrt{m_V m_{B_c}} [h_{A_1}(w) (1 + w) \epsilon_\mu^* - h_{A_2}(w) (\epsilon^* \cdot v) v_\mu - h_{A_3}(w) (\epsilon^* \cdot v) v'_\mu] \\ \langle V(v', \epsilon) | \bar{Q}' \gamma_5 Q | B_c(v) \rangle &= -\sqrt{m_V m_{B_c}} h_P(w) (\epsilon^* \cdot v) \\ \langle V(v', \epsilon) | \bar{Q}' \sigma_{\mu\nu} Q | B_c(v) \rangle &= -\sqrt{m_V m_{B_c}} \epsilon^{\mu\nu\alpha\beta} [h_{T_1}(w) \epsilon_\alpha^* (v + v')_\beta + h_{T_2}(w) \epsilon_\alpha^* (v - v')_\beta \\ &\quad + h_{T_3}(w) (\epsilon^* \cdot v) v_\alpha v'_\beta]\end{aligned}\tag{A.2}$$

with $V = J/\psi$, ϵ the J/ψ polarization vector and $v' = \frac{p'}{m_V}$.

B Form factors in terms of universal functions

The expressions of the form factors h_i in terms of universal functions are given in the following.

B.1 $B_c \rightarrow \eta_c$

$$\begin{aligned}h_+(w) &= \Delta(w) + \frac{1}{4m_b} (3\chi_2^A(w) + 2(w-1)\chi_2^C(w)) + \frac{1}{4m_c} (3\bar{\chi}_2^A(w) - 2(w-1)\bar{\chi}_2^B(w)) \\ &\quad + \frac{1}{2m_b^2} \chi_1(w) + \frac{1}{2m_c^2} \bar{\chi}_1(w) \\ &\quad + \frac{1}{8m_b m_c} \left(-(2+w)\psi_1^S(w) - (w^2-1)(\psi_2^S(w) - \psi_3^S(w) + \psi_1^A(w)) \right. \\ &\quad \left. + 6(w-1)(\psi_4^S(w) + \psi_2^A(w)) + (w-7)\psi_3^A(w) \right)\end{aligned}\tag{B.1}$$

$$\begin{aligned}
h_-(w) = & \left(-\frac{1}{m_b} + \frac{1}{m_c} \right) \left(\frac{\phi_K(w) - \Delta(w)\tilde{\Lambda}}{2} + \Delta_3(w) \right) \\
& + \frac{1}{2m_b^2} \left[(\tilde{\Lambda} - w\tilde{\Lambda}') \left(\frac{\phi_K(w) - \Delta(w)\tilde{\Lambda}}{2} + \Delta_3(w) \right) + \frac{1}{4}(1+w)^2\psi_2^S(w) + \frac{1}{4}(w^2-1)\psi_3^S(w) \right. \\
& + (1+w) \left(-\psi_4^S(w) - \frac{w}{2}\psi_6^S(w) + \frac{1}{4}\psi_1^A(w) - \frac{1}{2}\psi_4^A(w) \right) \\
& \left. + \frac{2w-1}{2}\psi_5^S(w) + \frac{w-2}{2}\psi_2^A(w) - \frac{3}{4}\psi_3^A(w) \right] \\
& + \frac{1}{2m_c^2} \left[(\tilde{\Lambda}w - \tilde{\Lambda}') \left(\frac{\phi_K(w) - \Delta(w)\tilde{\Lambda}}{2} + \Delta_3(w) \right) \right. \\
& - \frac{1}{4}(1+w)^2\psi_2^S(w) - \frac{1}{4}(w^2-1)\psi_3^S(w) \\
& + (1+w) \left(\psi_4^S(w) - \frac{w}{2}\psi_6^S(w) - \frac{1}{4}\psi_1^A(w) - \frac{1}{2}\psi_4^A(w) \right) \\
& \left. + \frac{2w-1}{2}\psi_5^S(w) - \frac{w-2}{2}\psi_2^A(w) + \frac{3}{4}\psi_3^A(w) \right]
\end{aligned} \tag{B.2}$$

$$\begin{aligned}
h_T(w) = & \Delta(w) - \left(\frac{1}{m_b} + \frac{1}{m_c} \right) \left(\frac{\phi_K(w) - \Delta(w)\tilde{\Lambda}}{2} + \Delta_3(w) \right) \\
& + \frac{1}{4m_b} \left(3\chi_2^A(w) + 2(w-1)\chi_2^C(w) \right) + \frac{1}{4m_c} \left(3\bar{\chi}_2^A(w) - 2(w-1)\bar{\chi}_2^B(w) \right) \\
& + \frac{1}{2m_b^2} \left[(\tilde{\Lambda} - w\tilde{\Lambda}') \left(\frac{\phi_K(w) - \Delta(w)\tilde{\Lambda}}{2} + \Delta_3(w) \right) + \chi_1(w) \right. \\
& + \frac{1}{4}(1+w)^2\psi_2^S(w) + \frac{1}{4}(w^2-1)\psi_3^S(w) \\
& + (1+w) \left(-\psi_4^S(w) - \frac{w}{2}\psi_6^S(w) + \frac{1}{4}\psi_1^A(w) - \frac{1}{2}\psi_4^A(w) \right) \\
& \left. + \frac{2w-1}{2}\psi_5^S(w) + \frac{w-2}{2}\psi_2^A(w) - \frac{3}{4}\psi_3^A(w) \right] \\
& - \frac{1}{2m_c^2} \left[(\tilde{\Lambda}w - \tilde{\Lambda}') \left(\frac{\phi_K(w) - \Delta(w)\tilde{\Lambda}}{2} + \Delta_3(w) \right) - \bar{\chi}_1(w) \right. \\
& - \frac{1}{4}(1+w)^2\psi_2^S(w) - \frac{1}{4}(w^2-1)\psi_3^S(w) \\
& + (1+w) \left(\psi_4^S(w) - \frac{w}{2}\psi_6^S(w) - \frac{1}{4}\psi_1^A(w) - \frac{1}{2}\psi_4^A(w) \right) \\
& \left. + \frac{2w-1}{2}\psi_5^S(w) - \frac{w-2}{2}\psi_2^A(w) + \frac{3}{4}\psi_3^A(w) \right] \\
& + \frac{1}{2m_b m_c} \left[\frac{w+2}{4}\psi_1^S(w) + \frac{w^2-1}{4}(\psi_2^S(w) - \psi_3^S(w) + \psi_1^A(w)) \right. \\
& \left. - \frac{3}{2}(w-1)(\psi_4^S(w) + \psi_2^A(w)) - \frac{w-7}{4}\psi_3^A(w) \right]
\end{aligned} \tag{B.3}$$

$$\begin{aligned}
h_S(w) = & \Delta(w) - \left(\frac{1}{m_b} + \frac{1}{m_c} \right) \frac{w-1}{w+1} \left(\frac{\phi_K(w) - \Delta(w)\tilde{\Lambda}}{2} + \Delta_3(w) \right) \\
& + \frac{1}{4m_b} \left(3\chi_2^A(w) + 2(w-1)\chi_2^C(w) \right) + \frac{1}{4m_c} \left(3\bar{\chi}_2^A(w) - 2(w-1)\bar{\chi}_2^B(w) \right) \\
& + \frac{1}{2m_b^2} \left[(\tilde{\Lambda} - w\tilde{\Lambda}') \frac{w-1}{w+1} \left(\frac{\phi_K(w) - \Delta(w)\tilde{\Lambda}}{2} + \Delta_3(w) \right) + \chi_1(w) \right] \\
& - \frac{1}{2m_c^2} \left[(\tilde{\Lambda}w - \tilde{\Lambda}') \frac{w-1}{w+1} \left(\frac{\phi_K(w) - \Delta(w)\tilde{\Lambda}}{2} + \Delta_3(w) \right) - \bar{\chi}_1(w) \right] \\
& + \frac{1}{8m_b^2} (w-1) \left[(w+1)\psi_2^S(w) + (w-1)\psi_3^S(w) - 4\psi_4^S(w) + 2\frac{2w-1}{w+1}\psi_5^S(w) \right. \\
& \left. - 2w\psi_6^S(w) + \psi_1^A(w) + 2\frac{w-2}{w+1}\psi_2^A(w) - \frac{3}{w+1}\psi_3^A(w) - 2\psi_4^A(w) \right] \quad (\text{B.4}) \\
& + \frac{1}{8m_c^2} (w-1) \left[(w+1)\psi_2^S(w) + (w-1)\psi_3^S(w) - 4\psi_4^S(w) - 2\frac{2w-1}{w+1}\psi_5^S(w) \right. \\
& \left. + 2w\psi_6^S(w) + \psi_1^A(w) + 2\frac{w-2}{w+1}\psi_2^A(w) - \frac{3}{w+1}\psi_3^A(w) + 2\psi_4^A(w) \right] \\
& + \frac{1}{8m_b m_c} \left[(2+w)\psi_1^S(w) - (w-7)\psi_3^A(w) - 6(w-1)[\psi_4^S(w) + \psi_2^A(w)] \right. \\
& \left. + (w^2-1)[\psi_2^S(w) - \psi_3^S(w) + \psi_1^A(w)] \right]
\end{aligned}$$

B.2 $B_c \rightarrow J/\psi$

$$\begin{aligned}
h_V(w) = & \Delta(w) + \frac{1}{4m_b} \left[3\chi_2^A(w) + 2(w-1)\chi_2^C(w) - 4 \left(\Delta_3(w) + \frac{\phi_K(w) - \Delta(w)\tilde{\Lambda}}{2} \right) \right] \\
& - \frac{1}{4m_c} \left[\bar{\chi}_2^A(w) + 2 \left(\phi_K(w) - \Delta(w)\tilde{\Lambda} \right) \right] \\
& + \frac{1}{2m_b^2} \left[\left(\tilde{\Lambda} - w\tilde{\Lambda}' \right) \left(\frac{\phi_K(w) - \Delta(w)\tilde{\Lambda}}{2} + \Delta_3(w) \right) + \chi_1(w) - \frac{3}{4}\psi_3^A(w) \right. \\
& + (1+w) \left(\frac{1}{4}\psi_1^A(w) - \psi_4^S(w) - \frac{1}{2}\psi_4^A(w) - \frac{1}{2}w\psi_6^S(w) \right) + \frac{w-2}{2}\psi_2^A(w) \\
& \left. + \frac{2w-1}{2}\psi_5^S(w) + \frac{w^2-1}{4}\psi_3^S(w) + \frac{(w+1)^2}{4}\psi_2^S(w) \right] \tag{B.5} \\
& + \frac{1}{2m_c^2} \left[\left(\tilde{\Lambda}' - w\tilde{\Lambda} \right) \frac{\phi_K(w) - \Delta(w)\tilde{\Lambda}}{2} + \bar{\chi}_1(w) - \frac{1}{4}\psi_3^A(w) - \frac{1}{2}\psi_2^A(w) \right. \\
& + (1+w) \left(\frac{1}{4}\psi_1^A(w) - \frac{1}{2}\psi_4^S(w) + \frac{1}{2}w\psi_6^S(w) \right) - \frac{1}{2}w\psi_5^S(w) + \frac{w^2-1}{4}\psi_3^S(w) + \frac{(1+w)^2}{4}\psi_2^S(w) \left. \right] \\
& + \frac{1}{8m_b m_c} \left[w\psi_1^S(w) + (w^2-1) \left(\psi_2^S(w) - \psi_3^S(w) + \psi_1^A(w) \right) \right. \\
& \left. + 2(1-w) \left(2\psi_4^S(w) + 2\psi_2^A(w) + \psi_4^A + \psi_5^S \right) + (3-w)\psi_3^A(w) \right]
\end{aligned}$$

$$\begin{aligned}
h_P(w) = & \Delta(w) + \frac{1}{4m_b} \left[3\chi_2^A(w) + 2(w-1)\chi_2^C(w) - 4 \left(\Delta_3(w) + \frac{\phi_K(w) - \Delta(w)\tilde{\Lambda}}{2} \right) \right] \\
& - \frac{1}{4m_c} \left[\bar{\chi}_2^A(w) + 4 \left(\frac{\phi_K(w) - \Delta(w)\tilde{\Lambda}}{2} - \Delta_3(w) \right) - 2(w-1)\bar{\chi}_2^B \right] \\
& + \frac{1}{2m_b^2} \left[(\tilde{\Lambda} - w\tilde{\Lambda}') \left(\frac{\phi_K(w) - \Delta(w)\tilde{\Lambda}}{2} + \Delta_3(w) \right) + \chi_1(w) - \frac{3}{4}\psi_3^A(w) \right. \\
& + (1+w) \left(\frac{1}{4}\psi_1^A(w) - \psi_4^S(w) - \frac{1}{2}\psi_4^A(w) - \frac{1}{2}w\psi_6^S(w) \right) + \frac{w-2}{2}\psi_2^A(w) \\
& \left. + \frac{2w-1}{2}\psi_5^S(w) + \frac{w^2-1}{4}\psi_3^S(w) + \frac{(w+1)^2}{4}\psi_2^S(w) \right] \\
& - \frac{1}{2m_c^2} \left[(\tilde{\Lambda}' - w\tilde{\Lambda}) \left(-\frac{\phi_K(w) - \Delta(w)\tilde{\Lambda}}{2} + \Delta_3(w) \right) \right. \\
& - \bar{\chi}_1(w) - \frac{1}{4}\psi_3^A(w) + \frac{1}{2}\psi_5^S(w) + \frac{1}{2}w\psi_2^A(w) \\
& - (1+w) \left(\frac{1}{4}\psi_1^A(w) - \frac{1}{2}\psi_4^A(w) + \frac{1}{2}w\psi_6^S(w) \right) + \frac{1-w^2}{4}\psi_3^S(w) - \frac{(1+w)^2}{4}\psi_2^S(w) \left. \right] \\
& + \frac{1}{8m_b m_c} \left[(w-2)\psi_1^S(w) + (w^2-1)(\psi_2^S(w) - \psi_3^S(w) + \psi_1^A(w)) \right. \\
& \left. + 2(1-w)(2\psi_4^A(w) + \psi_2^A(w) + \psi_4^S(w) + 2\psi_5^S(w)) - (1+w)\psi_3^A(w) \right]
\end{aligned} \tag{B.6}$$

$$\begin{aligned}
h_{A_1} = & \Delta(w) - \frac{1}{m_b} \frac{w-1}{w+1} \left(\frac{\phi_K(w) - \Delta(w)\tilde{\Lambda}}{2} + \Delta_3(w) \right) - \frac{1}{m_c} (w-1) \frac{\phi_K(w) - \Delta(w)\tilde{\Lambda}}{2(w+1)} \\
& + \frac{1}{4m_b} \left(3\chi_2^A(w) + 2(w-1)\chi_2^C(w) \right) - \frac{1}{4m_c} \bar{\chi}_2^A(w) \\
& + \frac{1}{2m_b^2} \left[\left(\tilde{\Lambda} - w\tilde{\Lambda}' \right) \frac{w-1}{(w+1)} \left(\frac{\phi_K(w) - \Delta(w)\tilde{\Lambda}}{2} + \Delta_3(w) \right) + \chi_1(w) \right] \\
& + \frac{1}{8m_b^2} \left[(w^2-1)\psi_2^S(w) + (w-1)^2\psi_3^S(w) + 2\frac{w^2-3w+2}{1+w}\psi_2^A(w) \right. \\
& \left. + (w-1) \left[-4\psi_4^S(w) + 2\frac{2w-1}{w+1}\psi_5^S(w) - 2w\psi_6^S(w) + \psi_1^A - \frac{3}{w+1}\psi_3^A(w) - 2\psi_4^A(w) \right] \right] \\
& + \frac{1}{2m_c^2} \left[- \left(\tilde{\Lambda}w - \tilde{\Lambda}' \right) \frac{w-1}{2(w+1)} (\phi_K(w) - \Delta(w)\tilde{\Lambda}) + \bar{\chi}_1(w) \right] \\
& + \frac{1}{8m_c^2} \left[(w^2-1)\psi_2^S(w) + (w-1)^2\psi_3^S(w) \right. \\
& \left. + (w-1) \left[-2\psi_4^S(w) - 2\frac{w}{w+1}\psi_5^S(w) + 2w\psi_6^S(w) + \psi_1^A - \frac{2}{w+1}\psi_2^A - \frac{1}{w+1}\psi_3^A(w) \right] \right] \\
& + \frac{1}{8m_b m_c} \left[w\psi_1^S(w) + (w^2-1)(\psi_2^S(w) - \psi_3^S(w) + \psi_1^A(w)) \right. \\
& \left. + (w-1) \left[-4\psi_4^S(w) - 2\psi_5^S(w) - 4\psi_2^A(w) - 2\psi_4^A(w) \right] - (w-3)\psi_3^A(w) \right]
\end{aligned} \tag{B.7}$$

$$\begin{aligned}
h_{A_2} = & \frac{1}{m_c} \left(\frac{1}{1+w} [\phi_K(w) - \Delta(w)\tilde{\Lambda} - \Delta_3(w)] + \frac{1}{2}\bar{\chi}_2^B(w) \right) \\
& + \frac{1}{m_c^2} \frac{1}{2(w+1)} \left(\tilde{\Lambda}w - \tilde{\Lambda}' \right) [\phi_K(w) - \Delta(w)\tilde{\Lambda} - \Delta_3(w)] \\
& + \frac{1}{4m_c^2} \left[- (1+w)\psi_2^S(w) - (w-1)\psi_3^S(w) \right. \\
& \left. + \psi_4^S(w) + \psi_5^S(w) - 2w\psi_6^S(w) - \psi_1^A(w) + \psi_2^A(w) + \psi_4^A(w) \right] \\
& + \frac{1}{4m_b m_c} \left[- \psi_1^S(w) - (1+w)\psi_2^S(w) + (w+1)\psi_3^S(w) \right. \\
& \left. + 3\psi_4^S(w) + 3\psi_5^S(w) - (1+w)\psi_1^A(w) + 3\psi_2^A(w) + \psi_3^A(w) + 3\psi_4^A(w) \right]
\end{aligned} \tag{B.8}$$

$$\begin{aligned}
h_{A_3} = & \Delta(w) - \frac{1}{m_b} \left(\frac{\phi_K(w) - \Delta(w)\tilde{\Lambda}}{2} + \Delta_3(w) \right) + \frac{1}{4m_b} \left(3\chi_2^A(w) + 2(w-1)\chi_2^C(w) \right) \\
& - \frac{1}{m_c(1+w)} \left[(w-1) \frac{\phi_K(w) - \Delta(w)\tilde{\Lambda}}{2} + \Delta_3(w) \right] - \frac{1}{4m_c} [\bar{\chi}_2^A(w) + 2\bar{\chi}_2^B(w)] \\
& + \frac{1}{2m_b^2} \left[\left(\tilde{\Lambda} - w\tilde{\Lambda}' \right) \left[\frac{\phi_K(w) - \Delta(w)\tilde{\Lambda}}{2} + \Delta_3(w) \right] + \chi_1(w) \right] \\
& - \frac{1}{2m_c^2} \left[\frac{1}{1+w} \left(\tilde{\Lambda}w - \tilde{\Lambda}' \right) \left[(w-1) \frac{\phi_K(w) - \Delta(w)\tilde{\Lambda}}{2} + \Delta_3(w) \right] - \bar{\chi}_1(w) \right] \\
& + \frac{1}{8m_b^2} \left[(1+w)^2\psi_2^S(w) + (w^2-1)\psi_3^S(w) + 2(2w-1)\psi_5^S(w) + 2(w-2)\psi_2^A(w) \right. \\
& \left. - 3\psi_3^A(w) + (1+w)(-4\psi_4^S(w) - 2w\psi_6^S(w) + \psi_1^A(w) - 2\psi_4^A(w)) \right] \\
& + \frac{1}{8m_c^2} \left[(w^2-1)\psi_2^S(w) + (w-1)^2\psi_3^S(w) - 2w\psi_4^S(w) - 2(w-1)\psi_5^S(w) \right. \\
& \left. - \psi_3^A(w) + 2\psi_4^A(w) + (w-1)(2w\psi_6^S(w) + \psi_1^A(w)) \right] \\
& + \frac{1}{8m_b m_c} \left[(2+w)(\psi_1^S(w) - 2\psi_5^S(w) - 2\psi_4^A(w)) - 2(2w+1)(\psi_4^S(w) + \psi_2^A(w)) \right. \\
& \left. + (1+w)^2(\psi_2^S(w) - \psi_3^S(w) + \psi_1^A(w)) - (w-1)\psi_3^A(w) \right]
\end{aligned} \tag{B.9}$$

$$\begin{aligned}
h_{T_1} = & \Delta(w) + \frac{1}{4m_b} \left(3\chi_2^A(w) + 2(w-1)\chi_2^C(w) \right) - \frac{1}{4m_c} \bar{\chi}_2^A(w) \\
& + \frac{1}{2m_b^2} \chi_1(w) + \frac{1}{2m_c^2} \bar{\chi}_1(w) \\
& + \frac{1}{8m_b m_c} \left[-w\psi_1^S(w) - (w^2-1)[\psi_2^S(w) - \psi_3^S(w) + \psi_1^A(w)] \right. \\
& \left. + (w-1)[4\psi_4^S(w) + 2\psi_5^S(w) + 4\psi_2^A(w) + 2\psi_4^A(w)] + (w-3)\psi_3^A(w) \right]
\end{aligned} \tag{B.10}$$

$$\begin{aligned}
h_{T_2} = & -\frac{1}{m_b} \left(\frac{\phi_K(w) - \Delta(w)\tilde{\Lambda}}{2} + \Delta_3(w) \right) + \frac{1}{2m_c} (\phi_K(w) - \Delta(w)\tilde{\Lambda}) \\
& + \frac{1}{2m_b^2} (\tilde{\Lambda} - w\tilde{\Lambda}') \left[\frac{\phi_K(w) - \Delta(w)\tilde{\Lambda}}{2} + \Delta_3(w) \right] + \frac{1}{2m_c^2} (\tilde{\Lambda}w - \tilde{\Lambda}') \frac{\phi_K(w) - \Delta(w)\tilde{\Lambda}}{2} \\
& + \frac{1}{8m_b^2} \left[(w+1)^2 \psi_2^S(w) + (w^2-1) \psi_3^S(w) + (1+w) [-4\psi_4^S(w) - 2w\psi_6^S(w) + \psi_1^A(w) - 2\psi_4^A(w)] \right. \\
& + 2(2w-1) \psi_5^S(w) + 2(w-2) \psi_2^A(w) - 3\psi_3^A(w) \left. \right] \\
& + \frac{1}{8m_c^2} \left[-(w+1)^2 \psi_2^S(w) - (w^2-1) \psi_3^S(w) + (1+w) [2\psi_4^S(w) - 2w\psi_6^S(w) - \psi_1^A(w)] \right. \\
& + 2w\psi_5^S(w) + 2\psi_2^A(w) + \psi_3^A(w) \left. \right]
\end{aligned} \tag{B.11}$$

$$\begin{aligned}
h_{T_3}(w) = & -\frac{1}{2m_c} \left[\bar{\chi}_2^B(w) + \frac{2}{1+w} \left(\Delta_3(w) - (\phi_K(w) - \Delta(w)\tilde{\Lambda}) \right) \right] \\
& - \frac{1}{4m_c^2} \left[\frac{2}{w+1} (\tilde{\Lambda}' - w\tilde{\Lambda}) \left((\phi_K(w) - \Delta(w)\tilde{\Lambda}) - \Delta_3(w) \right) \right. \\
& + \psi_1^A(w) - \psi_2^A(w) - \psi_4^S(w) - \psi_4^A(w) \\
& + (w-1) \psi_3^S(w) - \psi_5^S(w) + (1+w) \psi_2^S(w) + 2w\psi_6^S(w) \left. \right] \\
& - \frac{1}{4m_b m_c} \left[\psi_1^S(w) - \psi_3^A(w) - 3(\psi_2^A(w) + \psi_4^A(w) + \psi_4^S(w) + \psi_5^S(w)) \right. \\
& + (1+w) (\psi_1^A(w) + \psi_2^S(w) - \psi_3^S(w)) \left. \right].
\end{aligned} \tag{B.12}$$

C Fit of the form factors h_{T_i} and h_P

The form factors h_{T_i} and h_P depicted in Fig. 3 are fitted using the parametrization adopted by the HPQCD Collaboration [65]. The variable $z(q^2, t_0)$ is defined:

$$z(q^2, t_0) = \frac{\sqrt{t_+ - q^2} - \sqrt{t_+ - t_0}}{\sqrt{t_+ - q^2} + \sqrt{t_+ - t_0}} \tag{C.1}$$

with $t_+ = (m_B + m_{D^*})^2$ and $t_0 = t_- = (m_{B_c} - m_{J/\psi})^2$. The form factors $h_i(q^2)$ are fitted using a truncated expansion in powers of z :

$$h_i(q^2) = \frac{1}{P(q^2)} \sum_{n=0}^N a_n z^n \tag{C.2}$$

Table 1: Mass M_{P_i} (in GeV) of the $\bar{b}c$ resonances P_i included in the Blaschke factors (C.3).

| J^P/i | 1^- | 1^+ | 0^- |
|---------|-------|-------|-------|
| 1 | 6.336 | 6.745 | 6.275 |
| 2 | 6.926 | 6.75 | 6.872 |
| 3 | 7.02 | 7.15 | 7.25 |
| 4 | 7.28 | 7.15 | |

where the Blaschke factors $P(t)$ account for the resonances in the t -channel:

$$P(q^2) = \prod_{P_i} z(q^2, M_{P_i}^2). \quad (\text{C.3})$$

h_{T_3} involves poles with $J^P = 1^+$, $h_{T_{1,2}}$ with $J^P = 1^-$, h_P with $J^P = 0^-$, with masses in Table 1. The values of a_n resulting from the fit are collected in Table 2.

Table 2: Parameters a_n in Eq. (C.2)

| | a_0 | a_1 | a_2 | a_3 |
|-----------|--------------------|--------------------|--------------------|--------------------|
| h_{T_1} | 0.058 ± 0.002 | -0.476 ± 0.001 | 1.228 ± 0.131 | 3.419 ± 3.977 |
| h_{T_2} | -0.021 ± 0.003 | 0.096 ± 0.014 | 0.030 ± 0.665 | 0.473 ± 0.670 |
| h_{T_3} | -0.047 ± 0.014 | 0.675 ± 0.165 | -0.701 ± 1.693 | 0.894 ± 7.468 |
| h_P | 0.142 ± 0.008 | -0.944 ± 0.053 | 0.157 ± 1.629 | -1.495 ± 1.131 |

References

- [1] **HFLAV** Collaboration, Y. S. Amhis et al., *Averages of b -hadron, c -hadron, and τ -lepton properties as of 2018*, *Eur. Phys. J. C* **81** (2021) 226, [[arXiv:1909.12524](#)].
- [2] P. Gambino et al., *Challenges in semileptonic B decays*, *Eur. Phys. J. C* **80** (2020) 966, [[arXiv:2006.07287](#)].
- [3] P. Colangelo and F. De Fazio, *Tension in the inclusive versus exclusive determinations of $|V_{cb}|$: a possible role of new physics*, *Phys. Rev. D* **95** (2017) 011701, [[arXiv:1611.07387](#)].
- [4] P. Colangelo, F. De Fazio, and F. Loparco, *Inclusive semileptonic Λ_b decays in the Standard Model and beyond*, *JHEP* **11** (2020) 032, [[arXiv:2006.13759](#)].
- [5] **LHCb** Collaboration, R. Aaij et al., *Measurement of the ratio of branching fractions $\mathcal{B}(B_c^+ \rightarrow J/\psi \tau^+ \nu_\tau)/\mathcal{B}(B_c^+ \rightarrow J/\psi \mu^+ \nu_\mu)$* , *Phys. Rev. Lett.* **120** (2018) 121801, [[arXiv:1711.05623](#)].
- [6] P. Biancofiore, P. Colangelo, and F. De Fazio, *Anomalous enhancement observed in $B \rightarrow D^{(*)} \tau \bar{\nu}_\tau$ decays*, *Phys. Rev. D* **87** (2013) 074010, [[arXiv:1302.1042](#)].

- [7] D. Becirevic, S. Fajfer, I. Nisandzic, and A. Tayduganov, *Angular distributions of $\bar{B} \rightarrow D^{(*)} \ell \bar{\nu}_\ell$ decays and search of New Physics*, *Nucl. Phys. B* **946** (2019) 114707, [[arXiv:1602.03030](#)].
- [8] R. Alonso, B. Grinstein, and J. Martin Camalich, *Lifetime of B_c^- Constrains Explanations for Anomalies in $B \rightarrow D^{(*)} \tau \nu$* , *Phys. Rev. Lett.* **118** (2017) 081802, [[arXiv:1611.06676](#)].
- [9] M. Jung and D. M. Straub, *Constraining new physics in $b \rightarrow c \ell \nu$ transitions*, *JHEP* **01** (2019) 009, [[arXiv:1801.01112](#)].
- [10] P. Colangelo and F. De Fazio, *Scrutinizing $\bar{B} \rightarrow D^* (D\pi) \ell^- \bar{\nu}_\ell$ and $\bar{B} \rightarrow D^* (D\gamma) \ell^- \bar{\nu}_\ell$ in search of new physics footprints*, *JHEP* **06** (2018) 082, [[arXiv:1801.10468](#)].
- [11] C. Murgui, A. Peñuelas, M. Jung, and A. Pich, *Global fit to $b \rightarrow c \tau \nu$ transitions*, *JHEP* **09** (2019) 103, [[arXiv:1904.09311](#)].
- [12] M. Algueró, S. Descotes-Genon, J. Matias, and M. Novoa-Brunet, *Symmetries in $B \rightarrow D^* \ell \nu$ angular observables*, *JHEP* **06** (2020) 156, [[arXiv:2003.02533](#)].
- [13] **HPQCD** Collaboration, J. Harrison, C. T. H. Davies, and A. Lytle, *$B_c \rightarrow J/\psi$ form factors for the full q^2 range from lattice QCD*, *Phys. Rev. D* **102** (2020) 094518, [[arXiv:2007.06957](#)].
- [14] P. Colangelo, G. Nardulli, and N. Paver, *QCD sum rules calculation of B_c decays*, *Z. Phys. C* **57** (1993) 43.
- [15] V. Kiselev, A. Likhoded, and A. Onishchenko, *Semileptonic B_c meson decays in sum rules of QCD and NRQCD*, *Nucl. Phys. B* **569** (2000) 473, [[hep-ph/9905359](#)].
- [16] C.-F. Qiao and R.-L. Zhu, *Estimation of semileptonic decays of B_c meson to S -wave charmonia with nonrelativistic QCD*, *Phys. Rev. D* **87** (2013) 014009, [[arXiv:1208.5916](#)].
- [17] R. Zhu, Y. Ma, X.-L. Han, and Z.-J. Xiao, *Relativistic corrections to the form factors of B_c into S -wave Charmonium*, *Phys. Rev. D* **95** (2017) 094012, [[arXiv:1703.03875](#)].
- [18] D. Shen, H. Ren, F. Wu, and R. Zhu, *$B_c \rightarrow J/\psi$ tensor form factors at large momentum recoil*, *Int. J. Mod. Phys. A* **36** (2021) 2150135.
- [19] R.-Y. Tang, Z.-R. Huang, C.-D. Lü, and R. Zhu, *Scrutinizing New Physics in Semi-leptonic $B_c \rightarrow J/\psi \tau \nu$ Decay*, [arXiv:2204.04357](#).
- [20] W. Tao, Z.-J. Xiao, and R. Zhu, *Next-to-leading order QCD calculation of B_c to charmonium tensor form factors*, *Phys. Rev. D* **105** (2022) 114026, [[arXiv:2204.06385](#)].
- [21] W.-F. Wang, Y.-Y. Fan, and Z.-J. Xiao, *Semileptonic decays $B_c \rightarrow (\eta_c, J/\Psi) l \nu$ in the perturbative QCD approach*, *Chin. Phys. C* **37** (2013) 093102, [[arXiv:1212.5903](#)].

- [22] J.-M. Shen, X.-G. Wu, H.-H. Ma, and S.-Q. Wang, *QCD corrections to the B_c to charmonia semileptonic decays*, *Phys. Rev. D* **90** (2014) 034025, [[arXiv:1407.7309](#)].
- [23] Z. Rui, H. Li, G.-x. Wang, and Y. Xiao, *Semileptonic decays of B_c meson to S -wave charmonium states in the perturbative QCD approach*, *Eur. Phys. J. C* **76** (2016) 564, [[arXiv:1602.08918](#)].
- [24] X.-Q. Hu, S.-P. Jin, and Z.-J. Xiao, *Semileptonic decays $B_c \rightarrow (\eta_c, J/\psi) l \bar{\nu}_l$ in the "PQCD + Lattice" approach*, *Chin. Phys. C* **44** (2020) 023104, [[arXiv:1904.07530](#)].
- [25] D. Leljak, B. Melic, and M. Patra, *On lepton flavour universality in semileptonic $B_c \rightarrow \eta_c, J/\psi$ decays*, *JHEP* **05** (2019) 094, [[arXiv:1901.08368](#)].
- [26] C.-T. Tran, M. A. Ivanov, J. G. Körner, and P. Santorelli, *Implications of new physics in the decays $B_c \rightarrow (J/\psi, \eta_c) \tau \nu$* , *Phys. Rev. D* **97** (2018) 054014, [[arXiv:1801.06927](#)].
- [27] S. Tang, S. Jia, P. Maris, and J. P. Vary, *Semileptonic decay of B_c to η_c and J/ψ on the light front*, *Phys. Rev. D* **104** (2021) 016002, [[arXiv:2011.05454](#)].
- [28] E. E. Jenkins, M. E. Luke, A. V. Manohar, and M. J. Savage, *Semileptonic B_c decay and heavy quark spin symmetry*, *Nucl. Phys. B* **390** (1993) 463, [[hep-ph/9204238](#)].
- [29] P. Colangelo and F. De Fazio, *Using heavy quark spin symmetry in semileptonic B_c decays*, *Phys. Rev. D* **61** (2000) 034012, [[hep-ph/9909423](#)].
- [30] P. Colangelo, F. De Fazio, and F. Loparco, *Role of $B_c^+ \rightarrow B_{s,d}^{(*)} \bar{\ell} \nu_\ell$ in the Standard Model and in the search for BSM signals*, *Phys. Rev. D* **103** (2021) 075019, [[arXiv:2102.05365](#)].
- [31] P. Colangelo, F. De Fazio, and F. Loparco, *$c \rightarrow uv\bar{\nu}$ transitions of B_c mesons: 331 model facing Standard Model null tests*, *Phys. Rev. D* **104** (2021) 115024, [[arXiv:2107.07291](#)].
- [32] G. P. Lepage, L. Magnea, C. Nakhleh, U. Magnea, and K. Hornbostel, *Improved nonrelativistic QCD for heavy quark physics*, *Phys. Rev. D* **46** (1992) 4052, [[hep-lat/9205007](#)].
- [33] G. T. Bodwin, E. Braaten, and G. P. Lepage, *Rigorous QCD analysis of inclusive annihilation and production of heavy quarkonium*, *Phys. Rev. D* **51** (1995) 1125, [[hep-ph/9407339](#)]. [Erratum: *Phys. Rev. D* **55**, 5853 (1997)].
- [34] A. Gunawardana and G. Paz, *On HQET and NRQCD Operators of Dimension 8 and Above*, *JHEP* **07** (2017) 137, [[arXiv:1702.08904](#)].
- [35] N. Brambilla, A. Pineda, J. Soto, and A. Vairo, *Effective Field Theories for Heavy Quarkonium*, *Rev. Mod. Phys.* **77** (2005) 1423, [[hep-ph/0410047](#)].
- [36] N. Brambilla, Y. Sumino, and A. Vairo, *Quarkonium spectroscopy and perturbative QCD: A New perspective*, *Phys. Lett. B* **513** (2001) 381, [[hep-ph/0101305](#)].

- [37] N. Brambilla, Y. Sumino, and A. Vairo, *Quarkonium spectroscopy and perturbative QCD: Massive quark loop effects*, *Phys. Rev. D* **65** (2002) 034001, [[hep-ph/0108084](#)].
- [38] J. Aebischer and B. Grinstein, *Standard Model prediction of the B_c lifetime*, *JHEP* **07** (2021) 130, [[arXiv:2105.02988](#)].
- [39] J. Aebischer and B. Grinstein, *A novel determination of the B_c lifetime*, [arXiv:2108.10285](#).
- [40] W. Buchmuller and D. Wyler, *Effective Lagrangian Analysis of New Interactions and Flavor Conservation*, *Nucl. Phys. B* **268** (1986) 621.
- [41] V. Cirigliano, J. Jenkins, and M. Gonzalez-Alonso, *Semileptonic decays of light quarks beyond the Standard Model*, *Nucl. Phys. B* **830** (2010) 95, [[arXiv:0908.1754](#)].
- [42] J. Aebischer and J. Kumar, *Flavour violating effects of Yukawa running in SMEFT*, *JHEP* **09** (2020) 187, [[arXiv:2005.12283](#)].
- [43] P. Colangelo, F. De Fazio, and F. Loparco, *Probing New Physics with $\bar{B} \rightarrow \rho(770) \ell^- \bar{\nu}_\ell$ and $\bar{B} \rightarrow a_1(1260) \ell^- \bar{\nu}_\ell$* , *Phys. Rev. D* **100** (2019) 075037, [[arXiv:1906.07068](#)].
- [44] A. F. Falk, H. Georgi, B. Grinstein, and M. B. Wise, *Heavy Meson Form-factors From QCD*, *Nucl. Phys. B* **343** (1990) 1.
- [45] A. F. Falk and M. Neubert, *Second order power corrections in the heavy quark effective theory. 1. Formalism and meson form-factors*, *Phys. Rev. D* **47** (1993) 2965, [[hep-ph/9209268](#)].
- [46] A. K. Leibovich, Z. Ligeti, I. W. Stewart, and M. B. Wise, *Semileptonic B decays to excited charmed mesons*, *Phys. Rev. D* **57** (1998) 308, [[hep-ph/9705467](#)].
- [47] A. F. Falk and B. Grinstein, *Power corrections to leading logs and their application to heavy quark decays*, *Phys. Lett. B* **247** (1990) 406–411.
- [48] A. F. Falk and B. Grinstein, *Heavy meson pair production in $e^+ e^-$ annihilation from the static quark effective theory*, *Phys. Lett. B* **249** (1990) 314–320.
- [49] M. Neubert, *Short distance expansion of heavy quark currents*, *Phys. Rev. D* **46** (1992) 2212–2227.
- [50] M. Neubert, *Reparametrization invariance and the expansion of currents in the heavy quark effective theory*, *Phys. Lett. B* **306** (1993) 357, [[hep-ph/9302269](#)].
- [51] A. F. Falk, *Hadrons of arbitrary spin in the heavy quark effective theory*, *Nucl. Phys. B* **378** (1992) 79.
- [52] R. Casalbuoni, A. Deandrea, N. Di Bartolomeo, R. Gatto, F. Feruglio, and G. Nardulli, *Hadronic transitions among quarkonium states in a soft exchange approximation. Chiral breaking and spin symmetry breaking processes*, *Phys. Lett. B* **309** (1993) 163, [[hep-ph/9304280](#)].

- [53] R. Casalbuoni, A. Deandrea, N. Di Bartolomeo, R. Gatto, F. Feruglio, and G. Nardulli, *Effective Lagrangian for quarkonia and light mesons in a soft-exchange-approximation*, *Phys. Lett. B* **302** (1993) 95.
- [54] F. De Fazio, *Radiative transitions of heavy quarkonium states*, *Phys. Rev. D* **79** (2009) 054015, [[arXiv:0812.0716](#)]. [Erratum: *Phys.Rev.D* 83, 099901 (2011)].
- [55] T. Mannel and G. A. Schuler, *Heavy quarkonium effective theory*, *Z. Phys. C* **67** (1995) 159, [[hep-ph/9410333](#)].
- [56] T. Mannel and G. A. Schuler, *HQanti-QET: An Effective theory approach to heavy quarkonia decays*, *Phys. Lett. B* **349** (1995) 181, [[hep-ph/9412337](#)].
- [57] C. Bourrely, B. Machet, and E. de Rafael, *Semileptonic Decays of Pseudoscalar Particles ($M \rightarrow M' \ell \nu_\ell$) and Short Distance Behavior of Quantum Chromodynamics*, *Nucl. Phys. B* **189** (1981) 157–181.
- [58] L. Lellouch, *Lattice constrained unitarity bounds for $\bar{B}^0 \rightarrow \pi^+ \ell^- \bar{\nu}_\ell$ decays*, *Nucl. Phys. B* **479** (1996) 353–391, [[hep-ph/9509358](#)].
- [59] C. G. Boyd, B. Grinstein, and R. F. Lebed, *Model independent determinations of $\bar{B} \rightarrow D \ell$, $D^* \ell \bar{\nu}$ form-factors*, *Nucl. Phys. B* **461** (1996) 493–511, [[hep-ph/9508211](#)].
- [60] I. Caprini and M. Neubert, *Improved bounds for the slope and curvature of $\bar{B} \rightarrow D(*) \ell \bar{\nu}$ form-factors*, *Phys. Lett. B* **380** (1996) 376–384, [[hep-ph/9603414](#)].
- [61] C. G. Boyd, B. Grinstein, and R. F. Lebed, *Precision corrections to dispersive bounds on form-factors*, *Phys. Rev. D* **56** (1997) 6895–6911, [[hep-ph/9705252](#)].
- [62] I. Caprini, L. Lellouch, and M. Neubert, *Dispersive bounds on the shape of $\bar{B} \rightarrow D(*) \ell \bar{\nu}$ form-factors*, *Nucl. Phys. B* **530** (1998) 153–181, [[hep-ph/9712417](#)].
- [63] M. Di Carlo, G. Martinelli, M. Naviglio, F. Sanfilippo, S. Simula, and L. Vittorio, *Unitarity bounds for semileptonic decays in lattice QCD*, *Phys. Rev. D* **104** (2021) 054502, [[arXiv:2105.02497](#)].
- [64] G. Martinelli, S. Simula, and L. Vittorio, *Constraints for the semileptonic $B \rightarrow D(*)$ form factors from lattice QCD simulations of two-point correlation functions*, *Phys. Rev. D* **104** (2021) 094512, [[arXiv:2105.07851](#)].
- [65] **HPQCD** Collaboration, L. J. Cooper, C. T. Davies, J. Harrison, J. Komijani, and M. Wingate, *$B_c \rightarrow B_{s(d)}$ form factors from lattice QCD*, *Phys. Rev. D* **102** (2020) 014513, [[arXiv:2003.00914](#)].