

A FAMILY OF HOMOGENEOUS OPERATORS IN THE COWEN-DOUGLAS CLASS OVER THE POLY-DISC

PRAHLLAD DEB AND SOMNATH HAZRA

ABSTRACT. We construct a large family of positive-definite kernels $K : \mathbb{D}^n \times \mathbb{D}^n \rightarrow M(r, \mathbb{C})$, holomorphic in the first variable and anti-holomorphic in the second, that are quasi-invariant with respect to the subgroup $\text{Möb} \times \cdots \times \text{Möb}$ (n times) of the bi-holomorphic automorphism group of \mathbb{D}^n . The adjoint of the n -tuples of multiplication operators by the co-ordinate functions on the Hilbert spaces \mathcal{H}_K determined by K is then homogeneous with respect to this subgroup. We show that these n -tuples are irreducible, are in the Cowen-Douglas class $B_r(\mathbb{D}^n)$ and that they are mutually pairwise unitarily inequivalent.

1. INTRODUCTION

Let Möb denote the bi-holomorphic automorphism group of the unit disc $\mathbb{D} := \{z \in \mathbb{C} : |z| < 1\}$. A bounded linear operator on a complex separable Hilbert space \mathcal{H} is said to be homogeneous if the spectrum $\sigma(T)$ of T is contained in $\overline{\mathbb{D}}$ and, $g(T)$ is unitarily equivalent to T for every g in the Möbius group Möb . Indeed, since $g \in \text{Möb}$ is a rational function with pole outside the closed disc $\overline{\mathbb{D}}$, it follows that $g(T)$ is well defined whenever the spectrum $\sigma(T)$ of T is contained in $\overline{\mathbb{D}}$. In a number of articles [1, 2, 6, 7, 9, 11], the class of homogeneous operators has been studied extensively.

The notion of a homogeneous operator has a natural generalization to commuting tuples of operators. Let Ω be a bounded symmetric domain in \mathbb{C}^n and G be a subgroup of the bi-holomorphic automorphism group $\text{Aut}(\Omega)$ of Ω . A commuting n -tuple of operators (T_1, T_2, \dots, T_n) is said to be homogeneous with respect to G if the Taylor joint spectrum of (T_1, T_2, \dots, T_n) lies in $\overline{\Omega}$ and $g(T_1, T_2, \dots, T_n)$, defined by the holomorphic functional calculus, is unitarily equivalent with (T_1, T_2, \dots, T_n) for all $g \in G$. In [8], Korányi and Misra described all irreducible homogeneous tuples of operators in the Cowen-Douglas class over the open unit ball $\mathbb{B}^n := \{z = (z_1, \dots, z_n) \in \mathbb{C}^n : \sum |z_i|^2 < 1\}$, see also [10]. In this generality, the focus has been on the study of commuting tuples of homogeneous operators in the Cowen-Douglas class over an irreducible bounded symmetric domain. In a recent paper, we initiated the study of n -tuples of operators homogeneous under the group $\text{Möb}^n := \text{Möb} \times \cdots \times \text{Möb}$. The group Möb^n is evidently a subgroup of the bi-holomorphic automorphism group of \mathbb{D}^n and acting transitively on \mathbb{D}^n .

In the recent paper [5], all irreducible n -tuples in $B_r(\mathbb{D}^n)$, $r = 1, 2, 3$, homogeneous with respect to the group Möb^n (respectively, $\text{Aut}(\mathbb{D}^n)$) have been listed modulo unitary equivalence. It has been also noted, in the same article, that the representations associated to the n -tuples of operators in $B_r(\mathbb{D}^n)$ homogeneous with respect to Möb^n , $r = 1, 2, 3$, must be multiplicity-free.

2020 *Mathematics Subject Classification.* Primary: 47B13, 47B32, Secondary: 20C25, 53C07.

Key words and phrases. Cowen-Douglas class, Homogeneous operators, Hermitian holomorphic homogeneous vector bundles, Curvature.

The research of the first named author was supported by J C Bose National Fellowship of G. Misra (DSTO 1984) followed by a postdoctoral fellowship at Ben-Gurion University of the Negev, Israel (850053497). The research of the second named author was supported by a post-doctoral Fellowship of the NBHM followed by GA CR grant no. 21-27941S.

In general, the classification modulo unitary equivalence of all homogeneous tuples of operators in $B_r(\mathbb{D}^n)$ for $r > 3$ is rather complicated. Recall that if T_i , $i = 1, \dots, n-1$ are homogeneous operators in $B_1(\mathbb{D})$ and S is an irreducible homogeneous operator in $B_r(\mathbb{D})$, $r = 1$ or 2 , then setting $\hat{T}_i = I \otimes \dots \otimes I \otimes T_i \otimes I \otimes \dots \otimes I$ and $\hat{S} := I \otimes \dots \otimes I \otimes S$, we infer that operators of the form $(\hat{T}_1, \dots, \hat{T}_{n-1}, \hat{S})$ are the only irreducible n -tuples of operators (up to unitary equivalence) in $B_r(\mathbb{D}^n)$, $r = 1, 2$, homogeneous with respect to Möb^n (cf. [5, Theorem 3.1, Theorem 6.7]).

Continuing in the same manner, for $r \geq 3$, examples of irreducible n -tuples $(\hat{T}_{n_1}, \dots, \hat{T}_{n_p})$, in $B_r(\mathbb{D}^n)$ homogeneous under Möb^n can be produced by taking tensor product, namely,

$$(1.1) \quad (\hat{T}_{n_1}, \dots, \hat{T}_{n_p}), \text{ with } \hat{T}_{n_i} = I \otimes \dots \otimes I \otimes T_{n_i} \otimes I \otimes \dots \otimes I, \quad T_{n_i} \in B_{r_i}(\mathbb{D}^{n_i})$$

such that $n_1 + \dots + n_p = n$, $r_1 \dots r_p = r$. Unlike the case when $r = 1$ or 2 , it was shown in [5, Theorem 7.8] that there are n -tuples in $B_3(\mathbb{D}^n)$ homogeneous under Möb^n are not necessarily of the form in (1.1) if we insist that $n_i = 1$, $1 \leq i \leq n$. The specific example for $r = 3$ given in that paper, however, is of the form (1.1) if one of the n_i is allowed to be > 1 . Never the less, we don't know if every irreducible n -tuple in $B_r(\mathbb{D}^n)$ homogeneous under Möb^n is of the form (1.1).

It is useful to also give an alternative description of some of these operators, we study in this paper, as n -tuples of multiplication of operators on a reproducing kernel Hilbert space. This is the class obtained by setting $n_i = 1$, $i = 1, \dots, n$ and $r_1 = \dots = r_{n-1} = 1$ and $r_n = r$. For $\lambda_1, \dots, \lambda_n > 0$, let $\mathcal{H}^{(\lambda_i)}(\mathbb{D})$ be the reproducing kernel Hilbert space with the reproducing kernel $K^{(\lambda_i)}(z, w) = (1 - z\bar{w})^{-\lambda_i}$, $z, w \in \mathbb{D}$ and $A^{(\lambda_n, \mu)}$ be the reproducing kernel Hilbert space determined by the positive definite kernel $B^{(\lambda_n, \mu)}$ as in [6, Equation (4.3)]. Then the adjoint of the n -tuple of multiplication operators by the co-ordinate functions on the reproducing kernel Hilbert space $\mathcal{H}^{(\lambda_1)} \otimes \mathcal{H}^{(\lambda_2)} \otimes \dots \otimes \mathcal{H}^{(\lambda_{n-1})} \otimes A^{(\lambda_n, \mu)}$ is in $B_r(\mathbb{D}^n)$, it is irreducible and homogeneous with respect to Möb^n and are mutually unitarily inequivalent.

Here we construct a family of irreducible n -tuples of operators in $B_r(\mathbb{D}^n)$ homogeneous with respect to Möb^n that is not of the form (1.1) if all the n_i 's are required to be 1. This, we hope, would be a first step in any attempt to classify homogeneous n -tuples in $B_r(\mathbb{D}^n)$, $r > 3$. The description of the Hilbert spaces in [5, Section 7] prompts a natural generalisation of the Γ map leading to construction of n -tuples of operators in $B_r(\mathbb{D}^n)$, $r \geq 3$, in a manner very similar to the one in [6].

In Section 3, for a fixed but arbitrary r -tuples of positive real numbers $\mu := (\mu_1, \dots, \mu_r)$, we introduce "the Γ map" depending on μ , from $\oplus_{0 \leq \beta \leq \alpha} (\otimes_{i=1}^n \mathcal{H}^{(\lambda_i + 2\beta_i)}(\mathbb{D}))$ taking values in $\text{Hol}(\mathbb{D}^n, \mathbb{C}^r)$. Here $\lambda = (\lambda_1, \dots, \lambda_n)$ is a fixed but arbitrary n -tuple of positive real numbers, $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_n) \in (\mathbb{N} \cup \{0\})^n$ such that the cardinality of the set $\Lambda = \{\beta \in (\mathbb{N} \cup \{0\})^n : \beta \leq \alpha\}$ is r , where " \leq " denotes the graded co-lexicographic ordering on $(\mathbb{N} \cup \{0\})^n$. Choosing the inner product in the image of Γ that makes Γ unitary, we see that the image of Γ is a reproducing kernel Hilbert space $\mathcal{H}^{(\lambda, \mu)}$. The reproducing kernel of the Hilbert space $\mathcal{H}^{(\lambda, \mu)}$ is computed explicitly. We then prove that the n -tuple of multiplication operators $\mathbf{M}^{(\lambda, \mu)} := (M_{z_1}, M_{z_2}, \dots, M_{z_n})$ by the co-ordinate functions on $\mathcal{H}^{(\lambda, \mu)}$ is Möb^n -homogeneous by obtaining the associated multiplier representation of Möb^n explicitly. We also establish that the adjoint of the n -tuple of multiplication operators $\mathbf{M}^{(\lambda, \mu)}$ is in $B_r(\mathbb{D}^n)$. In Section 4, we show that the tuple of multiplication operators $\mathbf{M}^{(\lambda, \mu)}$ on $\mathcal{H}^{(\lambda, \mu)}$ is irreducible. In the final section, we prove that $\mathbf{M}^{(\lambda, \mu)}$ and $\mathbf{M}^{(\lambda', \mu')}$ are unitarily equivalent if and only if $\lambda = \lambda'$ and $\mu = \mu'$.

2. PRELIMINARIES

Let \mathbb{D} be the open unit disc in \mathbb{C} and Möb be the set of all bi-holomorphic automorphisms $z \mapsto e^{i\theta} \frac{z-a}{1-\bar{a}z}$, $a \in \mathbb{D}$, $\theta \in [0, 2\pi)$ of \mathbb{D} . Note that $\text{SU}(1, 1) := \left\{ \begin{pmatrix} a & b \\ \bar{b} & \bar{a} \end{pmatrix} : |a|^2 - |b|^2 = 1 \right\}$ is a two fold

cover of Möb. Consequently, the universal covering space $\widetilde{\text{Möb}}$ of Möb is same as that of $\text{SU}(1, 1)$. Let $\mathbb{D}^n := \{(z_1, \dots, z_n) : |z_1|, \dots, |z_n| < 1\}$ and $\text{Möb}^n := \text{Möb} \times \dots \times \text{Möb}$. It is known that bi-holomorphic automorphisms $\text{Aut}(\mathbb{D}^n)$ of \mathbb{D}^n is the semi-direct product $\text{Möb}^n \rtimes \mathfrak{S}_n$, where \mathfrak{S}_n is the permutation group on n symbols. As we have said before, the group Möb^n acts transitively on \mathbb{D}^n . In this paper, we study n - tuples of multiplication operators homogeneous under the group Möb^n .

Let \mathcal{H}_K be a Hilbert space of holomorphic functions on \mathbb{D}^n taking values in \mathbb{C}^r for some $r \in \mathbb{N}$ possessing a reproducing kernel K satisfying the reproducing property: for each $z \in \mathbb{D}^n$, $\xi \in \mathbb{C}^r$ and $f \in \mathcal{H}_K$,

$$(2.1) \quad \langle f, K(\cdot, z)\xi \rangle_{\mathcal{H}_K} = \langle f(z), \xi \rangle_{\mathbb{C}^r}.$$

Note that $K(\cdot, z)\xi \in \mathcal{H}_K$ for each $z \in \mathbb{D}^n$ and $\xi \in \mathbb{C}^r$ and as is well-known, the linear span of the set $\{K(\cdot, z)\xi : z \in \mathbb{D}^n, \xi \in \mathbb{C}^r\}$ is dense in \mathcal{H}_K . Also, by differentiating both sides of the Equation (2.1), we have that

$$(2.2) \quad \langle f, \bar{\partial}_i^l K(\cdot, z)\xi \rangle_{\mathcal{H}_K} = \langle \partial_i^l f(z), \xi \rangle_{\mathbb{C}^r}, \quad f \in \mathcal{H}_K, \quad 1 \leq i \leq n, \quad l \in \mathbb{N}, \quad \text{and } \xi \in \mathbb{C}^r.$$

Recall from [3, 4] that the adjoint \mathbf{M}^* of the n - tuple of multiplication operators \mathbf{M} by the co-ordinate functions on \mathcal{H}_K is in the Cowen-Douglas class $B_r(\mathbb{D}^n)$ over \mathbb{D}^n if the operator $D_{\mathbf{M}^*} : \mathcal{H}_K \rightarrow \mathcal{H}_K \oplus \dots \oplus \mathcal{H}_K$ defined by $D_{\mathbf{M}^*} f = (M_{z_1}^* f, \dots, M_{z_n}^* f)$, $f \in \mathcal{H}_K$ satisfies the following properties:

- $\dim \ker D_{\mathbf{M}^* - zI} = r$, $z \in \mathbb{D}^n$;
- $\text{ran} D_{\mathbf{M}^* - zI}$ is closed in $\mathcal{H}_K \oplus \dots \oplus \mathcal{H}_K$;
- $\bigvee_{z \in \mathbb{D}^n} \ker D_{\mathbf{M}^* - zI}$ is dense in \mathcal{H}_K .

In this case, there is a hermitian holomorphic vector bundle $E_{\mathbf{M}^*} := \{(z, v) : v \in \ker D_{\mathbf{M}^* - zI}\} \subset \mathbb{D}^n \times \mathcal{H}_K$ defined over \mathbb{D}^n such that the unitary equivalence class of the n - tuple \mathbf{M}^* and the (local) equivalence class of the hermitian holomorphic vector bundle $E_{\mathbf{M}^*}$ determine each other. Given two reproducing kernels $K_1, K_2 : \mathbb{D}^n \rightarrow M(r, \mathbb{C})$ with the property that M_i^* on \mathcal{H}_{K_i} , $i = 1, 2$ are in $B_r(\mathbb{D}^n)$, it follows that M_1^* and M_2^* are unitarily equivalent if and only if there exists a holomorphic function $\Phi : \mathbb{D}^n \rightarrow \text{GL}(r, \mathbb{C})$ such that $K_1(z, w) = \Phi(z)K_2(z, w)\Phi(w)^*$ which is same as the fact that the hermitian holomorphic vector bundles $E_{M_1^*}$ and $E_{M_2^*}$ are unitarily equivalent over \mathbb{D}^n .

If in addition, the n - tuple of multiplication operators $\mathbf{M} = (M_{z_1}, \dots, M_{z_n})$ by the co-ordinate functions on \mathcal{H}_K is homogeneous with respect to Möb^n - that is, for each $\phi = (\phi_1, \dots, \phi_n) \in \text{Möb}^n$, $\phi(\mathbf{M}) = (M_{\phi_1}, \dots, M_{\phi_n})$, where $\phi(\mathbf{M})$ is defined by the usual holomorphic functional calculus, is unitarily equivalent to \mathbf{M} - then the hermitian holomorphic vector bundle $E_{\mathbf{M}^*}$ is homogeneous with respect to Möb^n as well. A hermitian holomorphic vector bundle $\pi : E_{\mathbf{M}^*} \rightarrow \mathbb{D}^n$ is said to be *homogeneous* with respect to a subgroup G of the group of bi-holomorphic automorphisms of \mathbb{D}^n if for every $\phi \in G$, there is an isometric bundle automorphism $\hat{\phi} : E_{\mathbf{M}^*} \rightarrow E_{\mathbf{M}^*}$ such that $\phi \circ \pi = \pi \circ \hat{\phi}$. In this paper, the group G is fixed once for all to be the subgroup Möb^n of the group of bi-holomorphic automorphisms of \mathbb{D}^n . Moreover, it follows from Theorem 4.1 in [5] that the universal covering group $\widetilde{\text{Möb}}^n$ acts on $E_{\mathbf{M}^*}$ uniquely by isometric bundle automorphisms.

Also, we recall from [5, Section 4] that the homogeneity of $\mathbf{M}^* \in B_r(\mathbb{D}^n)$ makes the reproducing kernel K *quasi-invariant*, that is,

$$(2.3) \quad K(z, w) = J(\tilde{\phi}, z)K(\phi(z), \phi(w))J(\tilde{\phi}, w)^*, \quad z, w \in \mathbb{D}^n, \quad \tilde{\phi} \in \widetilde{\text{Möb}}^n,$$

where $\phi = p(\tilde{\phi})$, $p : \widetilde{\text{Möb}}^n \rightarrow \text{Möb}^n$ is the universal covering map and for each $\tilde{\phi} \in \widetilde{\text{Möb}}^n$, $J(\tilde{\phi}, \cdot) : \mathbb{D}^n \rightarrow \text{GL}(r, \mathbb{C})$ is a holomorphic mapping. Further, if the function J satisfies the

“co-cycle” identity

$$J(\tilde{\phi}\tilde{\psi}, z) = J(\tilde{\psi}, z)J(\tilde{\phi}, \psi(z)), \quad \tilde{\phi}, \tilde{\psi} \in \widetilde{\text{Möb}}^n, \quad \text{and } z \in \mathbb{D}^n,$$

we say that J is a *co-cycle*. If K is quasi-invariant and J is a co-cycle, then it is easy to verify that the map $U : \widetilde{\text{Möb}}^n \times \mathcal{H}_K \rightarrow \mathcal{H}_K$ defined by

$$(2.4) \quad (U(\tilde{\phi}^{-1})f)(z) := J(\tilde{\phi}, z)f(\phi(z))$$

is a unitary representation of $\widetilde{\text{Möb}}^n$ onto \mathcal{H}_K .

3. CONSTRUCTION OF HILBERT SPACES

In this section, we construct a family of reproducing kernel Hilbert spaces consisting of holomorphic functions on \mathbb{D}^n taking values in \mathbb{C}^r for a given $r \in \mathbb{N}$. The reproducing kernels associated to these Hilbert spaces are computed explicitly. We then prove that the n -tuples of multiplication operators by the co-ordinate functions on these Hilbert spaces are bounded. Finally, the adjoint of these n -tuples of multiplication operators are shown to be in $B_r(\mathbb{D}^n)$.

For a given $r \in \mathbb{N}$, fix an element $\alpha = (\alpha_1, \dots, \alpha_n) \in (\mathbb{N} \cup \{0\})^n$ so that the cardinality $|\Lambda|$ of the set $\Lambda = \{\beta \in (\mathbb{N} \cup \{0\})^n : \beta \leq \alpha\}$ is r . Here we are using the graded co-lexicographic ordering on $(\mathbb{N} \cup \{0\})^n$.

For each $\beta = (\beta_1, \dots, \beta_n) \in \Lambda$ and $\lambda = (\lambda_1, \dots, \lambda_n)$ with each $\lambda_i > 0$, consider the reproducing kernel Hilbert space $\mathcal{H}^{(\lambda+2\beta)} := \bigotimes_{i=1}^n \mathcal{H}^{(\lambda_i+2\beta_i)}(\mathbb{D})$ of holomorphic functions on \mathbb{D}^n with the reproducing kernel

$$(3.1) \quad K^{(\lambda+2\beta)}(z, w) = \prod_{i=1}^n (1 - z_i \bar{w}_i)^{-(\lambda_i+2\beta_i)}, \quad z, w \in \mathbb{D}^n.$$

Define the mapping $\Gamma_\beta : \mathcal{H}^{(\lambda+2\beta)} \rightarrow \text{Hol}(\mathbb{D}^n, \mathbb{C}^r)$ for $\beta \in \Lambda$ as follows

$$\Gamma_\beta(f) = \left(\left(\binom{\gamma}{\beta} \frac{\partial^{\gamma-\beta} f}{(\lambda+2\beta)_{\gamma-\beta}} \right)_{0 \leq \gamma \leq \alpha} \right).$$

Note that if $\gamma_i < \beta_i$ for some i , then $\binom{\gamma}{\beta} = 0$. It follows from the definition that each Γ_β is injective. We define the mapping Γ on $\bigoplus_{0 \leq \beta \leq \alpha} \mathcal{H}^{(\lambda+2\beta)}$ taking values in $\text{Hol}(\mathbb{D}^n, \mathbb{C}^r)$ by

$$(f_\beta)_{0 \leq \beta \leq \alpha} \mapsto \sum_{0 \leq \beta \leq \alpha} \mu_\beta \Gamma_\beta(f_\beta)$$

where μ_β with $0 \leq \beta \leq \alpha$ are all positive real numbers and $\mu_0 = 1$. Note from the definition that Γ is an injective linear transformation. Let $\mathcal{H}^{(\lambda, \mu)}$ denote the image of Γ . Being injective, Γ defines an inner product on $\mathcal{H}^{(\lambda, \mu)}$ as follows

$$\langle \Gamma((f_\beta)_{0 \leq \beta \leq \alpha}), \Gamma((g_\beta)_{0 \leq \beta \leq \alpha}) \rangle := \langle (f_\beta)_{0 \leq \beta \leq \alpha}, (g_\beta)_{0 \leq \beta \leq \alpha} \rangle$$

making Γ a unitary transformation onto $\mathcal{H}^{(\lambda, \mu)}$. We now prove that $\mathcal{H}^{(\lambda, \mu)}$ is a reproducing kernel Hilbert space by first showing that the image $\text{Im}(\Gamma_\beta)$ of Γ_β is so for each $\beta \in \Lambda$.

Proposition 3.1. *Let $\beta \in \Lambda$. Then $\text{Im}(\Gamma_\beta)$ is a reproducing kernel Hilbert space with the reproducing kernel*

$$K_\beta(z, w) = \left(\left(\frac{\binom{s}{\beta} \binom{t}{\beta}}{(\lambda+2\beta)_{s-\beta} (\lambda+2\beta)_{t-\beta}} \partial^{s-\beta} \bar{\partial}^{t-\beta} K^{(\lambda+2\beta)}(z, w) \right)_{0 \leq s, t \leq \alpha} \right), \quad z, w \in \mathbb{D}^n$$

where $K^{(\lambda+2\beta)}(z, w)$ is as in (3.1).

Proof. Let $\{e_\theta \in \mathbb{C}^r : \theta \in \Lambda\}$ be the standard ordered basis for \mathbb{C}^r . Note that for $f_\beta \in \mathcal{H}^{(\lambda+2\beta)}$,

$$\begin{aligned} \langle \Gamma_\beta f_\beta, K_\beta(\cdot, w)e_\theta \rangle &= \left\langle \Gamma_\beta f_\beta, \Gamma_\beta \frac{\binom{\theta}{\beta}}{(\lambda+2\beta)_{\theta-\beta}} \bar{\partial}^{\theta-\beta} K^{(\lambda+2\beta)}(\cdot, w) \right\rangle \\ &= \frac{\binom{\theta}{\beta}}{(\lambda+2\beta)_{\theta-\beta}} \partial^{\theta-\beta} f_\beta(w) \\ &= \langle \Gamma_\beta f_\beta(w), e_\theta \rangle \end{aligned}$$

completing the proof. \square

Corollary 3.2. $\mathcal{H}^{(\lambda, \mu)}$ is a reproducing kernel Hilbert space with the reproducing kernel

$$(3.2) \quad K^{(\lambda, \mu)}(z, w) = \sum_{0 \leq \beta \leq \alpha} \mu_\beta^2 K_\beta(z, w), \quad z, w \in \mathbb{D}^n.$$

Next, we prove that the reproducing kernel $K^{(\lambda, \mu)}$ is quasi-invariant with respect to $\widetilde{\text{Möb}}^n$, that is, $K^{(\lambda, \mu)}$ satisfies the equation (2.3) for some co-cycle $J : \widetilde{\text{Möb}}^n \times \mathbb{D}^n \rightarrow \text{GL}(r, \mathbb{C})$. This can be achieved by producing an unitary multiplier representation of $\widetilde{\text{Möb}}^n$ onto $\mathcal{H}^{(\lambda, \mu)}$ defined by the equation (2.4) with J as the multiplier.

In the case of reproducing kernel Hilbert spaces of scalar valued functions on \mathbb{D} (that is, for $n = r = 1$), the unitary multiplier representations of Möb are well known. Since these are the building blocks of the Hilbert spaces of our interest, we first describe them. These are the elements of holomorphic discrete series representation D_λ^+ of Möb acting on the weighted Bergman space $\mathcal{H}^{(\lambda)}(\mathbb{D})$ possessing a reproducing kernel $B^{(\lambda)}(z, w) = (1 - z\bar{w})^{-\lambda}$ for $z, w \in \mathbb{D}$ and $\lambda > 0$.

For $g \in \text{Möb}$ and $t \in \mathbb{R}_{>0}$, taking the principal branch of power function we can uniquely define $g'(z)^t$ as a holomorphic function on \mathbb{D} . Now for $\tilde{g} \in \widetilde{\text{Möb}}$ with $p(\tilde{g}) = g \in \text{Möb}$ (p is the universal covering map), the multiplier $j_t(\tilde{g}, z) = g'(z)^t$ defines on $\mathcal{H}^{(t)}(\mathbb{D})$ the unitary representation $D_+^{(t)}$ as follows:

$$D_+^{(t)}(\tilde{g}^{-1})(f) := (g')^{\frac{t}{2}}(f \circ g), \quad f \in \mathcal{H}^{(t)}(\mathbb{D}), \quad \tilde{g} \in \widetilde{\text{Möb}}.$$

As a consequence, for $f_\beta \in \mathcal{H}^{(\lambda+2\beta)}$ and $\tilde{\phi} \in \widetilde{\text{Möb}}^n$,

$$D_+^{(\lambda+2\beta)}(\tilde{\phi}^{-1})(f_\beta) = (\phi')^{\frac{\lambda+2\beta}{2}}(f_\beta \circ \phi) = \prod_{i=1}^n (\phi'_i)^{\frac{\lambda_i+2\beta_i}{2}}(f_{\beta_i} \circ \phi_i)$$

defines a unitary multiplier representation of $\widetilde{\text{Möb}}^n$ onto $\mathcal{H}^{(\lambda+2\beta)}$ for each $\beta \in \Lambda$. Therefore, the direct sum of these representations can be transferred to $\mathcal{H}^{(\lambda, \mu)}$ by the map Γ . We show that this is a multiplier representation.

We need a relation between $g''(z)$ and $g'(z)$, for $g \in \text{Möb}$, in what follows. Note that for $g \in \text{Möb}$,

$$(3.3) \quad g''(z) = -2c_g g'(z)^{\frac{3}{2}}$$

where c_g is independent of z and the meaning of $g'(z)^{\frac{3}{2}}$ is as defined above.

Proposition 3.3. For $\lambda = (\lambda_1, \dots, \lambda_n)$ with each $\lambda_i > 0$, the image of $\bigoplus_{0 \leq \beta \leq \alpha} D_+^{(\lambda+2\beta)}$ under the mapping Γ is a multiplier representation of $\widetilde{\text{Möb}}^n$ onto $\mathcal{H}^{(\lambda, \mu)}$ with the multiplier

$$(3.4) \quad J(\tilde{\phi}, z)_{\theta_\eta} = \begin{cases} \binom{\theta}{\eta} (-c_\phi)^{\theta-\eta} (\phi')^{\frac{\lambda+\theta+\eta}{2}}(z) & \text{if } 0 \leq \eta_i \leq \theta_i, \quad i = 1, \dots, n \\ 0 & \text{otherwise.} \end{cases}$$

Proof. We begin with the observation that it is enough to show, for any elementary tensors $f_\beta = f_{\beta_1} \otimes \cdots \otimes f_{\beta_n} \in \mathcal{H}^{(\lambda+2\beta)}$ with $f_{\beta_i} \in \mathcal{H}^{(\lambda_i+2\beta_i)}$, $i = 1, \dots, n$, the following identity:

$$\Gamma_\beta(D_+^{(\lambda+2\beta)}(\tilde{\phi}^{-1})f_\beta) = J(\tilde{\phi}, \cdot)\Gamma_\beta(f_\beta), \quad \beta \in \Lambda, \quad \text{and } \tilde{\phi} \in \widetilde{\text{Möb}}^n.$$

Now using the definition of Γ_β note that

$$(3.5) \quad \Gamma_\beta(D_+^{(\lambda+2\beta)}(\tilde{\phi}^{-1})f_\beta) = \left(\binom{\gamma}{\beta} \frac{\partial^{\gamma-\beta}((\phi')^{\frac{\lambda+2\beta}{2}} f_\beta \circ \phi)}{(\lambda+2\beta)_{\gamma-\beta}} \right)_{0 \leq \gamma \leq \alpha}$$

where

$$\partial^{\gamma-\beta}((\phi')^{\frac{\lambda+2\beta}{2}} f_\beta \circ \phi) = \prod_{i=1}^n \partial_i^{\gamma_i-\beta_i} \left((\phi'_i)^{\frac{\lambda_i+2\beta_i}{2}} f_{\beta_i} \circ \phi_i \right).$$

For $\gamma \in \Lambda$ with $\gamma_i < \beta_i$ for some $1 \leq i \leq n$, since $\binom{\gamma_i}{\beta_i} = 0$ it is seen that $(\Gamma_\beta(f))_\gamma = 0$. Now using the formula (cf. [6, Lemma 3.1])

$$\partial^k((g')^l f \circ g) = \sum_{i=0}^k \binom{k}{i} (2l+i)_{k-i} (-c_g)^{k-i} (g')^{l+\frac{k+i}{2}} (f^{(i)} \circ g)$$

where $g'' = -2c_g(g')^{\frac{3}{2}}$ for $g \in \text{Möb}$, we have that

$$\begin{aligned} & \prod_{i=1}^n \partial_i^{\gamma_i-\beta_i} \left((\phi'_i)^{\frac{\lambda_i+2\beta_i}{2}} f_{\beta_i} \circ \phi_i \right) \\ &= \prod_{i=1}^n \sum_{t_i=0}^{\gamma_i-\beta_i} \binom{\gamma_i-\beta_i}{t_i} (\lambda_i+2\beta_i+t_i)_{\gamma_i-\beta_i-t_i} (-c_{\phi_i})^{\gamma_i-\beta_i-t_i} \\ & \times \left((\partial_i^{t_i} f_{\beta_i}) \circ \phi_i \right) (\phi'_i)^{\frac{\lambda_i+\gamma_i+\beta_i+t_i}{2}} \\ &= \prod_{i=1}^n \sum_{t_i+\beta_i=\beta_i}^{\gamma_i} \binom{\gamma_i-\beta_i}{t_i+\beta_i-\beta_i} (\lambda_i+\beta_i+t_i+\beta_i)_{\gamma_i-(t_i+\beta_i)} (-c_{\phi_i})^{\gamma_i-(t_i+\beta_i)} \\ & \times \left((\partial_i^{t_i+\beta_i-\beta_i} f_{\beta_i}) \circ \phi_i \right) (\phi'_i)^{\frac{\lambda_i+\gamma_i+(t_i+\beta_i)}{2}} \\ &= \prod_{i=1}^n \sum_{s_i=\beta_i}^{\gamma_i} \binom{\gamma_i-\beta_i}{s_i-\beta_i} (\lambda_i+\beta_i+s_i)_{\gamma_i-s_i} (-c_{\phi_i})^{\gamma_i-s_i} \left((\partial_i^{s_i-\beta_i} f_{\beta_i}) \circ \phi_i \right) (\phi'_i)^{\frac{\lambda_i+\gamma_i+s_i}{2}} \\ &= \sum_{s_i=\beta_i}^{\gamma_i} \prod_{i=1}^n \binom{\gamma_i-\beta_i}{s_i-\beta_i} (\lambda_i+\beta_i+s_i)_{\gamma_i-s_i} (-c_{\phi_i})^{\gamma_i-s_i} \left((\partial_i^{s_i-\beta_i} f_{\beta_i}) \circ \phi_i \right) (\phi'_i)^{\frac{\lambda_i+\gamma_i+s_i}{2}} \\ &= \sum_{s=\beta}^{\gamma} \binom{\gamma-\beta}{s-\beta} (\lambda+\beta+s)_{\gamma-s} (-c_\phi)^{\gamma-s} (\phi')^{\frac{\lambda+\gamma+s}{2}} \left((\partial^{s-\beta} f_\beta) \circ \phi \right). \end{aligned}$$

Here s denotes the tuple (s_1, \dots, s_n) . Thus this computation together with (3.5) yield that for $0 \leq \beta \leq \alpha$,

$$\begin{aligned} & (\Gamma_\beta(D_+^{(\lambda+2\beta)}(\tilde{\phi}^{-1})f_\beta))_\gamma \\ &= \frac{\binom{\gamma}{\beta}}{(\lambda+2\beta)_{\gamma-\beta}} \sum_{s=\beta}^{\gamma} \binom{\gamma-\beta}{s-\beta} (\lambda+\beta+s)_{\gamma-s} (-c_\phi)^{\gamma-s} (\phi')^{\frac{\lambda+\gamma+s}{2}} (\partial^{s-\beta} f_\beta) \circ \phi \\ &= \frac{1}{(\lambda+2\beta)_{\gamma-\beta}} \sum_{s=\beta}^{\gamma} \binom{\gamma}{s} \binom{s}{\beta} (\lambda+\beta+s)_{\gamma-s} (-c_\phi)^{\gamma-s} (\phi')^{\frac{\lambda+\gamma+s}{2}} (\partial^{s-\beta} f_\beta) \circ \phi \end{aligned}$$

where the last equality holds since $\binom{\gamma}{\beta} \binom{\gamma-\beta}{s-\beta} = \binom{\gamma}{s} \binom{s}{\beta}$. Consequently, we have that

$$(3.6) \quad \Gamma_\beta(D_+^{(\lambda+2\beta)}(\tilde{\phi}^{-1})f_\beta)(z) = J(\tilde{\phi}, z)\Gamma_\beta(f_\beta)$$

where $J(\tilde{\phi}, z)_{\theta\eta}$ is as given in (3.4). □

Theorem 3.4. *Let $K^{(\lambda, \mu)}$ be the reproducing kernel of the Hilbert space $\mathcal{H}^{(\lambda, \mu)}$. Then*

$$(3.7) \quad K^{(\lambda, \mu)}(z, w) = J(\tilde{\phi}, z)K^{(\lambda, \mu)}(\phi(z), \phi(w))J(\tilde{\phi}, w)^*, \quad \tilde{\phi} \in \widetilde{\text{Möb}}^n, \quad z, w \in \mathbb{D}^n,$$

where J is the co-cycle defined by the equation (3.4).

Proof. Let $\mathcal{U}(\mathcal{H}^{(\lambda, \mu)})$ be the set of unitary operators on $\mathcal{H}^{(\lambda, \mu)}$. We then define the mapping $U : \widetilde{\text{Möb}}^n \rightarrow \mathcal{U}(\mathcal{H}^{(\lambda, \mu)})$ by the formula

$$U(\tilde{\phi}^{-1})f := J(\tilde{\phi}, \cdot)f \circ \phi, \quad \tilde{\phi} \in \widetilde{\text{Möb}}^n, \quad f \in \mathcal{H}^{(\lambda, \mu)}.$$

Observe from Proposition 3.3 that U is a unitary representation and consequently, the equation (3.7) follows. □

A canonical decomposition of the reproducing kernel $K^{(\lambda, \mu)}$ can be obtained using the identity (3.7) and the expression of the co-cycle J given in (3.4). In order to obtain this decomposition, we need some notations. For $1 \leq i \leq n$, denote \tilde{S}_i to be the i -th shift defined as follows:

$$(3.8) \quad \tilde{S}_i e_\theta = (\theta_i + 1)e_{\theta+\varepsilon_i}, \quad 0 \leq \theta \leq \alpha,$$

and $D(z\bar{w})$ is the diagonal operator $D(z\bar{w})e_\theta = (1 - z\bar{w})^{-\theta}e_\theta$, $0 \leq \theta \leq \alpha$ where $\{\varepsilon_i : 1 \leq i \leq n\}$ and $\{e_\theta : \theta \in \Lambda\}$ are the standard ordered bases of \mathbb{C}^n and \mathbb{C}^r , respectively. For $z \in \mathbb{D}^n$, let $\tilde{\phi}_z \in \widetilde{\text{Möb}}^n$ with $p(\tilde{\phi}_z) = \phi_z \in \text{Möb}^n$ be the automorphism $\phi_z = (\phi_{z_1}, \dots, \phi_{z_n})$ where p is the universal covering map and $\phi_{z_i}(w_i) = \frac{w_i - z_i}{1 - \bar{z}_i w_i}$, $w_i \in \mathbb{D}$, $1 \leq i \leq n$. Consequently, $c_{\phi_{z_i}} = -\frac{\bar{z}_i}{\sqrt{1 - |z_i|^2}}$. The following lemma provides a canonical decomposition of the co-cycle J introduced in (3.4) in terms of the shift operators \tilde{S}_i , $1 \leq i \leq n$ and the diagonal matrix $D(z\bar{w})$.

Lemma 3.5. *The co-cycle J defined in (3.4) turns out to be*

$$(3.9) \quad J(\phi_z, z) = (1 - |z|^2)^{-\frac{\lambda}{2}} D(|z|^2) \exp \left(\sum_{i=1}^n \bar{z}_i \tilde{S}_i \right).$$

Proof. We begin by recalling from the equation (3.4) that

$$J(\tilde{\phi}, z)_{\theta\eta} = \binom{\theta}{\eta} (-c_\phi)^{\theta-\eta} (\phi')^{\frac{\lambda+\theta+\eta}{2}}(z)$$

whenever $0 \leq \eta_i \leq \theta_i$. Substituting $\phi = \phi_z$ in this equation we have, for $0 \leq \eta_i \leq \theta_i$ and $1 \leq i \leq n$, that

$$J(\tilde{\phi}_z, z)_{\theta\eta} = \binom{\theta}{\eta} \left(\frac{\bar{z}}{\sqrt{1-|z|^2}} \right)^{\theta-\eta} \left(\frac{1-|z|^2}{(1-|z|^2)^2} \right)^{\frac{\lambda+\theta+\eta}{2}} = (1-|z|^2)^{-\frac{\lambda}{2}} \binom{\theta}{\eta} \bar{z}^{\theta-\eta} (1-|z|^2)^{-\theta}.$$

On the other hand, note that

$$\begin{aligned} D(|z|^2) \exp \left(\sum_{i=1}^n \bar{z}_i \tilde{S}_i \right) e_\eta &= D(|z|^2) \exp(\bar{z}_n \tilde{S}_n) \cdots \exp(\bar{z}_1 \tilde{S}_1) e_\eta \\ &= D(|z|^2) \sum_{k=0}^{\alpha} \bar{z}^k \binom{\eta+k}{k} e_{\eta+k} \\ &= D(|z|^2) \sum_{\theta=\eta}^{\alpha} \binom{\theta}{\eta} \bar{z}^{\theta-\eta} e_\theta \\ &= \sum_{\theta=\eta}^{\alpha} \binom{\theta}{\eta} \bar{z}^{\theta-\eta} (1-|z|^2)^{-\theta} e_\theta \end{aligned}$$

where the second last equality is obtained by replacing $\eta+k$ to θ and using the identity $\binom{\theta}{\theta-\eta} = \binom{\theta}{\eta}$. Thus it proves the desired identity. \square

Proposition 3.6. *The reproducing kernel $K^{(\lambda, \mu)}$ of the Hilbert space $\mathcal{H}^{(\lambda, \mu)}$ is of the form*

$$K^{(\lambda, \mu)}(z, w) = (1-z\bar{w})^{-\lambda} D(z\bar{w}) \exp \left(\sum_{i=1}^n \bar{w}_i \tilde{S}_i \right) K^{(\lambda, \mu)}(0, 0) \exp \left(\sum_{i=1}^n z_i \tilde{S}_i^* \right) D(z\bar{w})$$

for all $z, w \in \mathbb{D}^n$ where $K^{(\lambda, \mu)}(0, 0)$ is a positive diagonal matrix.

Proof. The proof follows from Lemma 3.5 and Equation (3.7). \square

We now prove the boundedness of the multiplication operators on $\mathcal{H}^{(\lambda, \mu)}$ using the following well-known lemma. Recall that the multiplication operator $M^{(\epsilon)}$ on $\mathcal{H}^{(\epsilon)}(\mathbb{D})$ defined by $(M^{(\epsilon)}f)(z) = zf(z)$ is bounded for any $\epsilon > 0$. Consequently, the reproducing kernel $K^{(\epsilon)}$ of the Hilbert space $\mathcal{H}^{(\epsilon)}(\mathbb{D})$ satisfies the inequality (3.10) in Lemma 3.7 for any $\epsilon > 0$. We use this technique to show that the n -tuple of the multiplication operators by the co-ordinate functions on $\mathcal{H}^{(\lambda, \mu)}$ are bounded.

Lemma 3.7. *Let \mathcal{H}_K be a reproducing kernel Hilbert space with the reproducing kernel K on some domain in \mathbb{C}^n . Then for $1 \leq j \leq n$, the multiplication operator M_{z_j} is bounded if and only if there exists a positive constant c_j such that*

$$(3.10) \quad (c_j - z_j \bar{w}_j) K(\mathbf{z}, \mathbf{w}) \geq 0.$$

Theorem 3.8. *The multiplication operators M_{z_1}, \dots, M_{z_n} by the co-ordinate functions on $\mathcal{H}^{(\lambda, \mu)}$ are bounded.*

Proof. We begin by recalling that the multiplication operators M_{z_1}, \dots, M_{z_n} by the co-ordinate functions on the reproducing kernel Hilbert space $\mathcal{H}^{(\epsilon)}(\mathbb{D}^n)$ with the reproducing kernel $K^{(\epsilon)}(z, w) = \prod_{i=1}^n (1-z_i \bar{w}_i)^{-\epsilon_i}$ for $z, w \in \mathbb{D}^n$ are bounded for all $\epsilon = (\epsilon_1, \dots, \epsilon_n) \in \mathbb{R}_{>0}^n$. Consequently, it follows from Lemma 3.7 that for each $i = 1, \dots, n$, there exists a positive constant $c_i > 0$ such that the inequality in (3.10) holds with $K = K^{(\epsilon)}$. In view of this observation together with Lemma 3.7, it is enough to write down the kernel $K^{(\lambda, \mu)}$ as a product of $K^{(\epsilon)}$ and a positive definite kernel K_1 on \mathbb{D}^n for some $\epsilon_1, \dots, \epsilon_n > 0$. Note that the expression of $K^{(\lambda, \mu)}$ obtained in Theorem 3.6

suggests $K^{(\lambda-\epsilon, \mu')}$ to be a possible choice for K_1 provided $K^{(\lambda-\epsilon, \mu')}(0, 0)$ is same as $K^{(\lambda, \mu)}(0, 0)$ for some positive real numbers μ'_β for $0 \leq \beta \leq \alpha$ with $\mu'_0 = 1$ and $\epsilon > 0$. In rest of the proof, we show the existence of such positive real numbers μ'_β for $0 \leq \beta \leq \alpha$ and $\epsilon > 0$.

For $0 \leq \theta \leq \alpha$, recall from the equation (3.2) that the θ -th entry of the diagonal matrix $K^{(\lambda, \mu)}(0, 0)$ is

$$K^{(\lambda, \mu)}(0, 0)_{\theta\theta} = \sum_{0 \leq \beta \leq \theta} \mu_0^2 K_\beta(0, 0)_{\theta\theta}$$

with $\mu_0 = 1$ and

$$K_\beta(z, w) = \left(\left(\frac{\binom{s}{\beta} \binom{t}{\beta}}{(\lambda + 2\beta)_{s-\beta} (\lambda + 2\beta)_{t-\beta}} \partial^{s-\beta} \bar{\partial}^{t-\beta} K^{(\lambda+2\beta)}(z, w) \right) \right)_{0 \leq s, t \leq \alpha}.$$

Thus it follows that for each $0 \leq \theta \leq \alpha$,

$$K_\beta(0, 0)_{\theta\theta} = \frac{\binom{\theta}{\beta}^2 (\theta - \beta)!}{(\lambda + 2\beta)_{\theta-\beta}}$$

and consequently,

$$(3.11) \quad K^{(\lambda, \mu)}(0, 0)_{\theta\theta} = \sum_{0 \leq \beta \leq \theta} \mu_\beta^2 \frac{\binom{\theta}{\beta}^2 (\theta - \beta)!}{(\lambda + 2\beta)_{\theta-\beta}}.$$

Let $[K^{(\lambda, \mu)}(0, 0)]$ be the column vector whose θ -th entry is $K^{(\lambda, \mu)}(0, 0)_{\theta\theta}$ for $0 \leq \theta \leq \alpha$. Then observe from the equation (3.11) that

$$(3.12) \quad [K^{(\lambda, \mu)}(0, 0)] = L(\lambda)(\mu^2)$$

where μ^2 is the column vector with μ_β^2 in its β -th position for $0 \leq \beta \leq \alpha$ and $L(\lambda)$ is the lower triangular matrix whose $\theta\beta$ -th entry is $\frac{\binom{\theta}{\beta}^2 (\theta - \beta)!}{(\lambda + 2\beta)_{\theta-\beta}}$ for $0 \leq \beta \leq \theta$. Also, note that $L(\lambda)$ is invertible and continuous in λ verifying that $L(\lambda - \epsilon)^{-1} L(\lambda)(\mu^2)$ is a vector with all of its entries being positive, for sufficiently small $\epsilon = (\epsilon_1, \dots, \epsilon_n)$ with each $\epsilon_i > 0$. Thus it completes the proof with the choice $(\mu')^2 = L(\lambda - \epsilon)^{-1} L(\lambda)(\mu^2)$. \square

We conclude this section by showing that the adjoint of the n - tuple of multiplication operators $(M_{z_1}, \dots, M_{z_n})$ by the co-ordinate functions on $\mathcal{H}^{(\lambda, \mu)}$ is in the Cowen-Douglas class $B_r(\mathbb{D}^n)$. From now on, this n - tuple $(M_{z_1}, \dots, M_{z_n})$ will be denoted by $\mathbf{M}^{(\lambda, \mu)}$.

Theorem 3.9. *The adjoint of the n -tuple of multiplication operators $\mathbf{M}^{(\lambda, \mu)}$ is in $B_r(\mathbb{D}^n)$ where r is the cardinality of the set $\Lambda = \{\beta \in (\mathbb{N} \cup \{0\})^n : \beta \leq \alpha\}$.*

Proof. Since by the construction of Hilbert space $\mathcal{H}^{(\lambda, \mu)}$, $\mathbf{M}^{(\lambda, \mu)*}$ is similar to the adjoint of multiplication operators on $\bigoplus_{0 \leq \beta \leq \alpha} \mathcal{H}^{(\lambda+2\beta)}$, it follows from the fact that the Cowen-Douglas class $B_r(\mathbb{D}^n)$ is invariant under the similarity. \square

4. IRREDUCIBILITY

In this section, we prove that the n - tuples of multiplication operators $\mathbf{M}^{(\lambda, \mu)}$ are irreducible. The lemma below, modelled after Lemma 5.1 in [6], can be proved exactly in the same way as in the original proof, so it is omitted.

Lemma 4.1. *Suppose that the n - tuple of multiplication operators $\mathbf{M} = (M_{z_1}, \dots, M_{z_n})$ by the co-ordinate functions on a reproducing kernel Hilbert space \mathcal{H} with the reproducing kernel K is in $B_r(\mathbb{D}^n)$. If there exists an orthogonal projection X commuting with \mathbf{M} then*

$$\Phi_X(z)K(z, w) = K(z, w)\overline{\Phi_X(w)}^{tr}$$

for some holomorphic function $\Phi_X : \mathbb{D}^n \rightarrow \mathbb{C}^{r \times r}$ with $\Phi_X^2 = \Phi_X$.

Thus $\mathbf{M}^{(\lambda, \mu)}$ on $\mathcal{H}^{(\lambda, \mu)}$ is irreducible if and only if there is no non-trivial projection X_0 on \mathbb{C}^r satisfying

$$X_0 K_0^{(\lambda, \mu)}(z, 0)^{-1} K_0^{(\lambda, \mu)}(z, w) K_0^{(\lambda, \mu)}(0, w)^{-1} = K_0^{(\lambda, \mu)}(z, 0)^{-1} K_0^{(\lambda, \mu)}(z, w) K_0^{(\lambda, \mu)}(0, w)^{-1} X_0$$

where $K_0^{(\lambda, \mu)}(z, w) = K^{(\lambda, \mu)}(0, 0)^{-\frac{1}{2}} K^{(\lambda, \mu)}(z, w) K^{(\lambda, \mu)}(0, 0)^{-\frac{1}{2}}$. Let

$$(4.1) \quad \hat{K}^{(\lambda, \mu)}(z, w) = K_0^{(\lambda, \mu)}(z, 0)^{-1} K_0^{(\lambda, \mu)}(z, w) K_0^{(\lambda, \mu)}(0, w)^{-1}$$

which is called the normalized kernel associated to K at origin.

Theorem 4.2. *The tuple of multiplication operators $\mathbf{M}^{(\lambda, \mu)}$ on $\mathcal{H}^{(\lambda, \mu)}$ are irreducible.*

Proof. In view of the discussion above, we show that there is no non-trivial projection commuting with the normalized kernel $\hat{K}^{(\lambda, \mu)}(z, w)$ for $z, w \in \mathbb{D}^n$ associated to the reproducing kernel $K^{(\lambda, \mu)}(z, w)$. Recall that $\Lambda = \{\theta \in (\mathbb{N} \cup \{0\})^n : \theta \leq \alpha\}$ with $|\Lambda| = r$ and $\{e_\theta : \theta \in \Lambda\}$ is the standard ordered basis of \mathbb{C}^r . Denote the set $\{e_\theta : \theta \in \Lambda\}$ by \mathcal{E} . For $1 \leq i \leq n$, let $\Lambda_0^i = \{\theta \in \Lambda : \theta_i = 0\} = \{\theta^{(0, i)}, \theta^{(1, i)}, \dots, \theta^{(k, i)}\}$ where $\theta^{(p, i)} \leq \theta^{(q, i)}$ if $p \leq q$. Then observe that the set $\mathcal{S}_{(\theta, i)} = \{\tilde{S}_i^j e_\theta : j \geq 0\}$, $\theta \in \Lambda_0^i$ is invariant under \tilde{S}_i . Since $1 \leq i \leq n$ is arbitrary but fixed for the rest of the proof, from now on we write \mathcal{S}_θ in place of $\mathcal{S}_{(\theta, i)}$. Note that

$$\Lambda = \bigcup_{\theta \in \Lambda_0^i} \mathcal{S}_\theta.$$

Let P be a permutation on \mathcal{E} such that P maps \mathcal{E} to the ordered basis

$$P\mathcal{E} = \left\{ S_i^j e_{\theta^{(0, i)}} : j \geq 0 \right\} \cup \dots \cup \left\{ S_i^j e_{\theta^{(k, i)}} : j \geq 0 \right\}$$

with $S_i e_\beta = e_{\beta + \epsilon_i}$ where we consider the ordering $S_i^l e_{\theta^{(p, i)}} < S_i^j e_{\theta^{(q, i)}}$ if $p < q$, or if $l < j$ whenever $p = q$.

For $z = (0, \dots, z_i, \dots, 0)$ and $w = (0, \dots, w_i, \dots, 0)$ in \mathbb{D}^n , let

$$A(z_i, w_i) = \hat{K}^{(\lambda, \mu)}(z, w)$$

and observe that

$$PA(z_i, w_i)P^{-1} = \bigoplus_{\theta \in \Lambda_0^i} (1 - z_i \bar{w}_i)^{-m(\theta, i)} \exp(-z_i \tilde{S}_{(\theta, i)}^*) B_{(\theta, i)}^{-1} \tilde{D}_\theta(z_i \bar{w}_i) \exp(\bar{w}_i \tilde{S}_{(\theta, i)}) \times \\ B_{(\theta, i)} \exp(z_i \tilde{S}_{(\theta, i)}^*) \tilde{D}_\theta(z_i \bar{w}_i) B_{(\theta, i)}^{-1} \exp(-\bar{w}_i \tilde{S}_{(\theta, i)}),$$

where $B_{(\theta, i)} = K^{(\lambda, \mu)}(0, 0)|_{\mathcal{S}_\theta}$, $m(\theta, i) = |\mathcal{S}_\theta| - 1$, $\tilde{S}_{(\theta, i)} = \tilde{S}_i|_{\mathcal{S}_\theta}$ and $\tilde{D}_\theta(z_i \bar{w}_i)$ is the diagonal operator defined as $\tilde{D}_\theta(z_i \bar{w}_i) S_i^j e_\theta = (1 - z_i \bar{w}_i)^{m(\theta, i) - j} e_{\theta + j\epsilon_i}$. We denote the expression

$$\exp(-z_i \tilde{S}_{(\theta, i)}^*) B_{(\theta, i)}^{-1} \tilde{D}_\theta(z_i \bar{w}_i) \exp(\bar{w}_i \tilde{S}_{(\theta, i)}) B_{(\theta, i)} \exp(z_i \tilde{S}_{(\theta, i)}^*) \tilde{D}_\theta(z_i \bar{w}_i) B_{(\theta, i)}^{-1} \exp(-\bar{w}_i \tilde{S}_{(\theta, i)})$$

by $A_\theta(z_i, w_i)$ and note that

$$\begin{aligned} (1 - z_i \bar{w}_i)^{-m(\theta, i)} A_\theta(z_i, w_i) &= \left(\sum_{n \geq 0} a_{(\theta, i)}(n) (z_i \bar{w}_i)^n \right) \left(\sum_{p \geq 0} b_{(\theta, i)}(p) z_i^{p+1} \bar{w}_i^p + \dots \right) \\ &= \sum_{q \geq 0} c_{(\theta, i)}(q) z_i^{q+1} \bar{w}_i^q + \dots \end{aligned}$$

where $c_{(\theta, i)}(q) = \sum_{l=0}^q a_{(\theta, i)}(l) b_{(\theta, i)}(q-l)$. It follows from [6, Lemma 5.4] that each $b_{(\theta, i)}(p)$, $p \geq 0$, is a shift on $\mathbb{C}^{|\mathcal{S}_\theta|}$ which in turn implies that each $c_{(\theta, i)}(q)$, $q \geq 0$, is indeed a shift since each $a_{(\theta, i)} \in \mathbb{C}$.

For $0 \leq l \leq m(\theta, i)$, let $\mathcal{P}_l = \{p \geq 0 : b_{(\theta, i)}(p)_{l, l+1} \neq 0\}$ which can be seen from [6, Lemma 5.4] to be a non-empty set. Let $n_l = \min \mathcal{P}_l$. We then have that

$$(c_{(\theta, i)}(n_l))_{l, l+1} = \left(\sum_{n=0}^{n_l} a_{(\theta, i)}(n) b_{(\theta, i)}(n_l - n) \right)_{l, l+1} \neq 0.$$

Thus for each $1 \leq i \leq n$, $\theta \in \Lambda_0^i$ and $0 \leq l \leq m(\theta, i)$, there exists r co-efficient matrices $\oplus_{\theta \in \Lambda_0^i} c_{(\theta, i)}(n_l)$ of $PA(z_i, w_i)P^{-1}$ whose $(l, l+1)$ -th entry in the θ 'th block is non-zero.

Let X be a projection commuting with $\hat{K}^{(\lambda, \mu)}$. Then note that $Y := PXP^{-1}$ is also a projection commuting with $P\hat{K}^{(\lambda, \mu)}P^{-1}$ since P is a permutation matrix on the orthonormal basis \mathcal{E} of \mathbb{C}^r . In particular, Y commutes with $\oplus_{\theta \in \Lambda_0^i} c_{(\theta, i)}(n_l)$ for all $1 \leq i \leq n$, $\theta \in \Lambda_0^i$ and $0 \leq l \leq m(\theta, i)$. Thus Y must be the identity operator verifying that so is X . \square

5. INEQUIVALENCE

The objective of this section is to show that the operators in the family $\{\mathbf{M}^{(\lambda, \mu)} : \lambda \in \mathbb{R}_{>0}^n, \mu \in \mathbb{R}_{>0}^r, \mu_0 = 1\}$ are mutually inequivalent. Recall from Theorem 3.9 that for every $\lambda \in \mathbb{R}_{>0}^n$ and $\mu \in \mathbb{R}_{>0}^r$, the adjoint of the n -tuple $\mathbf{M}^{(\lambda, \mu)}$ is in $B_r(\mathbb{D}^n)$. Consequently, $\mathbf{M}^{(\lambda, \mu)}$ gives rise to a hermitian holomorphic vector bundle $E^{(\lambda, \mu)}$ over \mathbb{D}^n with the hermitian structure induced from the reproducing kernel $K^{(\lambda, \mu)}$ as is well known (cf. [3]). Then for each $1 \leq i, j \leq n$, the ij -th component of the curvature of the bundle $E^{(\lambda, \mu)}$ is defined by the formula

$$\mathcal{K}^{(\lambda, \mu)}(z)_{ij} = \bar{\partial}_j(K^{(\lambda, \mu)}(z, z))^{-1} \partial_i K^{(\lambda, \mu)}(z, z), \quad z \in \mathbb{D}^n.$$

Lemma 5.1. *If $\mathbf{M}^{(\lambda, \mu)}$ and $\mathbf{M}^{(\lambda', \mu')}$ are unitarily equivalent then $\lambda = \lambda'$.*

Proof. Since $\mathbf{M}^{(\lambda, \mu)}$ and $\mathbf{M}^{(\lambda', \mu')}$ are unitarily equivalent, it follows (cf. [3]) that $E^{(\lambda, \mu)}$ and $E^{(\lambda', \mu')}$ are isomorphic as hermitian holomorphic vector bundles. Consequently, the curvature components of $E^{(\lambda, \mu)}$ and $E^{(\lambda', \mu')}$ are similar. Note that the metric of the bundle $E^{(\lambda, \mu)}$ (resp. $E^{(\lambda', \mu')}$) is $K^{(\lambda, \mu)}(z, z)$ (resp. $K^{(\lambda', \mu')}(z, z)$), $z \in \mathbb{D}^n$.

A direct computation shows that

$$(5.1) \quad \mathcal{K}_{11}^{(\lambda, \mu)}(0) = \bar{\partial}_1 \left(K^{(\lambda, \mu)}(z, z) \partial_1 K^{(\lambda, \mu)}(z, z) \right) \Big|_{z=0} = \lambda_1 I + 2D_1 - \tilde{S}_1^* B^{-1} \tilde{S}_1 B + B^{-1} \tilde{S}_1 B \tilde{S}_1^*,$$

where $B = K^{(\lambda, \mu)}(0, 0)$ and $D_1 := \bar{\partial}_1 \partial_1 D(|z|^2) \Big|_{z=0}$. Evaluating $\mathcal{K}_{11}^{(\lambda, \mu)}(0)$ on e_θ , $0 \leq \theta \leq \alpha$, we obtain

$$\mathcal{K}_{11}^{(\lambda, \mu)}(0) e_\theta = \left(\lambda_1 + 2\theta_1 - \frac{(\theta_1 + 1)^2 b_\theta}{b_{\theta + \epsilon_1}} + \frac{\theta_1^2 b_{\theta - \epsilon_1}}{b_\theta} \right) e_\theta,$$

where b_θ is the θ 'th diagonal entry of B . Thus we have

$$\mathrm{Tr} \mathcal{K}_{11}^{(\lambda, \mu)}(0) = r\lambda_1 + 2 \sum_{\theta \in \Lambda} \theta_1.$$

Similarly, we have

$$\mathrm{Tr} \mathcal{K}_{11}^{(\lambda', \mu')}(0) = r\lambda'_1 + 2 \sum_{\theta \in \Lambda} \theta_1.$$

Since $\mathcal{K}_{11}^{(\lambda, \mu)}(0)$ and $\mathcal{K}_{11}^{(\lambda', \mu')}(0)$ are similar, equating $\mathrm{Tr} \mathcal{K}_{11}^{(\lambda, \mu)}(0)$ and $\mathrm{Tr} \mathcal{K}_{11}^{(\lambda', \mu')}(0)$, we obtain $\lambda_1 = \lambda'_1$. A similar computation gives us $\lambda_i = \lambda'_i$ for all $2 \leq i \leq n$. \square

Theorem 5.2. *If $M^{(\lambda, \mu)}$ and $M^{(\lambda', \mu')}$ are unitarily equivalent then $\lambda = \lambda'$ and $\mu = \mu'$.*

Proof. In view of Lemma 5.1, it is enough to show that $\mu = \mu'$ whenever $M^{(\lambda, \mu)}$ and $M^{(\lambda, \mu')}$ are unitarily equivalent. Since by Theorem 3.9 both $M^{(\lambda, \mu)}$ and $M^{(\lambda, \mu')}$ are in $B_r(\mathbb{D}^n)$ and they are unitarily equivalent, it follows that the hermitian holomorphic vector bundles $E^{(\lambda, \mu)}$ and $E^{(\lambda, \mu')}$ are isomorphic. Consequently, there exists a holomorphic function Φ on \mathbb{D}^n taking values in $\mathrm{GL}(r, \mathbb{C})$ such that

$$(5.2) \quad K^{(\lambda, \mu')}(z, w) = \Phi(z)K^{(\lambda, \mu)}(z, w)\Phi(w)^*, \text{ for all } z, w \in \mathbb{D}^n.$$

Further, since both $M^{(\lambda, \mu)}$ and $M^{(\lambda, \mu')}$ are homogeneous with respect to $\mathrm{Möb}^n$, and $\widetilde{\mathrm{Möb}}^n$ action on $E^{(\lambda, \mu)}$ is unique (cf. [7, Theorem 2.1]), it follows that the cocycles J and J' associated to $E^{(\lambda, \mu)}$ and $E^{(\lambda, \mu')}$, respectively, satisfy

$$(5.3) \quad J(\tilde{g}, z) = \Phi(g(z))^{-1}J'(\tilde{g}, z)\Phi(z), \quad \tilde{g} \in \widetilde{\mathrm{Möb}}^n, z \in \mathbb{D}^n.$$

Indeed, note that the expression $\Phi(g(\cdot))^{-1}J'(\tilde{g}, \cdot)\Phi(\cdot)$ gives rise to an action of $\widetilde{\mathrm{Möb}}^n$ on $E^{(\lambda, \mu)}$. In particular, for $g = (g_{z_1}, \dots, g_{z_n}) \in \mathrm{Möb}^n$ with $g_{z_i}(w_i) = \frac{w_i - z_i}{1 - \bar{z}_i w_i}$, $i = 1, \dots, n$, we have from the Equation (3.9) that

$$(1 - |z|^2)^{-\frac{\lambda}{2}} D(|z|^2) \exp\left(\sum_{i=1}^n \bar{z}_i \tilde{S}_i\right) = (1 - |z|^2)^{-\frac{\lambda}{2}} \Phi(0)^{-1} D(|z|^2) \exp\left(\sum_{i=1}^n \bar{z}_i \tilde{S}_i\right) \Phi(z),$$

or equivalently,

$$\Phi(z) = \exp\left(-\sum_{i=1}^n \bar{z}_i \tilde{S}_i\right) D(|z|^2)^{-1} \Phi(0) D(|z|^2) \exp\left(\sum_{i=1}^n \bar{z}_i \tilde{S}_i\right).$$

Note from this expression of $\Phi(z)$ that $\partial_1^{p_1} \dots \partial_n^{p_n} \Phi(z)|_{z=0} = 0$ for all $(p_1, \dots, p_n) \in (\mathbb{N} \cup \{0\})^n$ with at least one $p_j \neq 0$. Consequently, it follows that

$$\Phi(z) = \Phi(0), \quad z \in \mathbb{D}^n$$

implying that

$$(5.4) \quad D(|z|^2) \exp\left(\sum_{i=1}^n \bar{z}_i \tilde{S}_i\right) \Phi(0) = \Phi(0) D(|z|^2) \exp\left(\sum_{i=1}^n \bar{z}_i \tilde{S}_i\right).$$

Now equating the coefficients of $|z_i|^2$ for each $i = 1, \dots, n$, in the Taylor expansion of both sides of this equation we have that

$$\frac{\partial^2}{\partial z_i \partial \bar{z}_i} D(|z|^2)|_{z=0} \Phi(0) = \Phi(0) \frac{\partial^2}{\partial z_i \partial \bar{z}_i} D(|z|^2)|_{z=0}, \quad i = 1, \dots, n.$$

Since $D(|z|^2)$ is the diagonal operator $D(|z|^2)e_\theta = (1 - |z|^2)^{-\theta}e_\theta$, $0 \leq \theta \leq \alpha$, the equation above implies that $\Phi(0)$ commutes with the diagonal matrices D_1, \dots, D_n where for $i = 1, \dots, n$, D_i is

the diagonal matrix whose β -th diagonal entry is β_i for any $\beta = (\beta_1, \dots, \beta_n) \in \Lambda$. Consequently, $\Phi(0)$ also commutes with all the linear combinations of the matrices D_1, \dots, D_n . Note that there exist $c_1, \dots, c_n \in \mathbb{C}$ such that the diagonal matrix $c_1 D_1 + \dots + c_n D_n$ has distinct diagonal entries. Thus $\Phi(0)$ is a diagonal matrix. Moreover, by comparing the \bar{z}_i -th coefficients \tilde{S}_i for each $i = 1, \dots, n$, in the Taylor expansion of both sides of the equation (5.4), it is seen that $\Phi(0)$ commutes with \tilde{S}_i , $i = 1, \dots, n$. Therefore, it follows from the definition of \tilde{S}_i , $i = 1, \dots, n$, there exists a non-zero $c \in \mathbb{C}$ such that $\Phi(0) = cI$ where I is the $r \times r$ identity matrix.

Finally, since $\Phi(0)$ satisfies the equation (5.2), we have that $K^{\lambda, \mu}(0, 0) = K^{(\lambda', \mu')}(0, 0)$. Therefore, the equality $\mu = \mu'$ holds from the equation (3.11) completing the proof. \square

We recall the construction of a natural class of irreducible n - tuple of operators in $B_r(\mathbb{D}^n)$ that are homogeneous with respect to Möbⁿ from the Introduction, namely,

$$(5.5) \quad (\hat{T}_{n_1}, \dots, \hat{T}_{n_p}), \text{ with } \hat{T}_{n_i} = I \otimes \dots \otimes I \otimes T_{n_i} \otimes I \otimes \dots \otimes I, \quad T_{n_i} \in B_{r_i}(\mathbb{D}^{n_i})$$

such that $n_1 + \dots + n_p = n$, $r_1 \dots r_p = r$. However, as promised in the Introduction, we now show that the n - tuples $M^{(\lambda, \mu)}$ are not of the form (5.5) if we insist that $n_i = 1$, $1 \leq i \leq n$. To establish this, we first prove a lemma.

Lemma 5.3. *Suppose $\alpha \neq \epsilon_1$. Then there exists $1 \leq i < j \leq n$ and an element θ in Λ such that $\theta - \epsilon_i + \epsilon_j$ is in Λ , but $\theta + \epsilon_j$ is not in Λ .*

Proof. We divide the proof into two cases. In the first case, we prove the existence of such θ, i, j by assuming $\alpha = (|\alpha|, 0, \dots, 0)$. In the second case, we assume that $\alpha \neq (|\alpha|, 0, \dots, 0)$ and prove that such θ, i, j exist.

Case 1: Suppose $\alpha = (|\alpha|, 0, \dots, 0)$. Since $\alpha \neq \epsilon_1$, it follows that $|\alpha| > 1$. Consider $\theta = \alpha - \epsilon_1$, $i = 1$ and any $j > 1$. Then we have $\theta - \epsilon_1 + \epsilon_j \in \Lambda$, but $\theta + \epsilon_j \notin \Lambda$.

Case 2: Assume that $\alpha \neq (|\alpha|, 0, \dots, 0)$. There exists $k > 1$ such that $\alpha_k \geq 1$. Consider $\theta = (|\alpha|, 0, \dots, 0)$, $i = 1$ and $j = k$. Since $|\theta + \epsilon_j| = |\alpha| + 1$, it follows that $\theta + \epsilon_j \notin \Lambda$.

We claim that $\theta - \epsilon_1 + \epsilon_k \in \Lambda$. On the contrary, assume $\theta - \epsilon_1 + \epsilon_k > \alpha$. Note that $|\theta - \epsilon_1 + \epsilon_k| = |\alpha|$. Since $\alpha_k \geq 1$, we have $0 \leq (\theta - \epsilon_1 + \epsilon_k)_l \leq \alpha_l$ for every $l \geq 2$ and therefore, it follows from our assumption that $(\theta - \epsilon_1 + \epsilon_k)_l = \alpha_l$ for all $l \geq 2$. Also, the inequality $\theta - \epsilon_1 + \epsilon_k > \alpha$ implies that $\alpha_1 < (\theta - \epsilon_1 + \epsilon_k)_1 = |\alpha| - 1$. Thus

$$|\alpha| = \sum_{l=1}^n \alpha_l = \alpha_1 + \alpha_k < |\alpha| - 1 + 1 = |\alpha|,$$

which is a contradiction. This proves the claim.

The verification of these two cases completes the proof of the lemma. \square

Theorem 5.4. *For each $1 \leq i \leq n$, assume that adjoint of the multiplication operator on a reproducing kernel Hilbert space \mathcal{H}_i with the reproducing kernel K_i is a homogeneous operator in $B_{r_i}(\mathbb{D})$ and denote M to be the n - tuple of multiplication operators on $\otimes_{i=1}^n \mathcal{H}_i$. If the adjoint of $M^{(\lambda, \mu)}$ is in $B_r(\mathbb{D}^n)$ with $r > 2$ then M and $M^{(\lambda, \mu)}$ are unitarily inequivalent.*

Proof. Since M^* is in $B_{\prod_i r_i}(\mathbb{D}^n)$, it follows that if $\prod_i r_i \neq r$, then M and $M^{(\lambda, \mu)}$ are not unitarily equivalent. Therefore we assume that $\prod_i r_i = r$. Note that the adjoint of both the tuples $M^{(\lambda, \mu)}$ and M are in $B_r(\mathbb{D}^n)$. Thus if $M^{(\lambda, \mu)}$ and M are unitarily equivalent, then the corresponding curvature components of $\hat{K}^{(\lambda, \mu)}$ and \hat{K} are similar, where $\hat{K}^{(\lambda, \mu)}$ and \hat{K} are normalized kernels of $K^{(\lambda, \mu)}$ and K at origin, respectively (cf. (4.1)).

A direct computation shows that

$$(5.6) \quad \hat{\mathcal{K}}_{ij}^{(\lambda, \mu)}(0) = \bar{\partial}_j \partial_i \hat{K}^{(\lambda, \mu)}(z, z)|_{z=0} = B^{-\frac{1}{2}} \tilde{S}_j B \tilde{S}_i^* B^{-\frac{1}{2}} - B^{\frac{1}{2}} \tilde{S}_i^* B^{-1} \tilde{S}_j B^{\frac{1}{2}},$$

for $1 \leq i \neq j \leq n$, where $B = K^{(\lambda, \mu)}(0, 0)$ and \tilde{S}_i is as in the equation (3.8). By Lemma 5.3, there exists a θ in Λ and $1 \leq p \neq q \leq n$ such that $\theta - \epsilon_p + \epsilon_q \in \Lambda$, but $\theta + \epsilon_q \notin \Lambda$. This implies that $B^{-\frac{1}{2}}\tilde{S}_q B \tilde{S}_p^* B^{-\frac{1}{2}} e_\theta \neq 0$, but $B^{\frac{1}{2}}\tilde{S}_p^* B^{-1}\tilde{S}_q B^{\frac{1}{2}} e_\theta = 0$. Therefore, we have $\hat{\mathcal{K}}_{pq}^{(\lambda, \mu)}(0) \neq 0$.

On the other hand, since $\hat{K} = \otimes_{i=1}^n \hat{K}_i$, it follows that $\hat{\mathcal{K}}_{ij}(0) = \bar{\partial}_j \partial_i \hat{K}(z, z)|_{z=0} = 0$, for any $1 \leq i \neq j \leq n$. This implies that $\hat{\mathcal{K}}_{pq}(0)$ and $\hat{\mathcal{K}}_{pq}^{(\lambda, \mu)}(0)$ are not similar. Therefore, M and $\mathbf{M}^{(\lambda, \mu)}$ are not unitarily equivalent. \square

Acknowledgement. The authors are grateful to Professor Gadadhar Misra for his invaluable comments and suggestions in preparation of this article.

REFERENCES

- [1] B. Bagchi and G. Misra, *Homogeneous operators and projective representations of the Möbius group: a survey*, Proc. Indian Acad. Sci. Math. Sci. **111** (2001), no. 4, 415–437. [1](#)
- [2] ———, *The homogeneous shifts*, J. Funct. Anal. **204** (2003), no. 2, 293–319. [1](#)
- [3] M. J. Cowen and R. G. Douglas, *Operators possessing an open set of eigenvalues*, Colloq. Math. Soc. János Bolyai **I,II** (1980), no. 35, 323–341. [3](#), [11](#)
- [4] R. E. Curto and N. Salinas, *Generalized Bergman kernels and the Cowen-Douglas theory*, Amer. J. Math. **106** (1984), no. 2, 447–488. [3](#)
- [5] P. Deb and S. Hazra, *Homogeneous hermitian holomorphic vector bundles and operators in the cowen-douglas class over the poly-disc*, J. Math. Anal. Appl., <https://doi.org/10.1016/j.jmaa.2022.126031>. [1](#), [2](#), [3](#)
- [6] A. Korányi and G. Misra, *Homogeneous operators on Hilbert spaces of holomorphic functions*, J. Funct. Anal. **254** (2008), no. 9, 2419–2436. [1](#), [2](#), [6](#), [9](#), [11](#)
- [7] ———, *A classification of homogeneous operators in the Cowen-Douglas class*, Adv. Math. **226** (2011), no. 6, 5338–5360. [1](#), [12](#)
- [8] A. Korányi and G. Misra, *Homogeneous Hermitian holomorphic vector bundles and the Cowen-Douglas class over bounded symmetric domains*, C. R. Math. Acad. Sci. Paris **354** (2016), no. 3, 291–295. [1](#)
- [9] G. Misra and S. S. Roy, *On the irreducibility of a class of homogeneous operators*, System theory, the Schur algorithm and multidimensional analysis, Oper. Theory Adv. Appl., vol. 176, pp. 165–198. [1](#)
- [10] G. Misra and H. Upmeyer, *Homogeneous vector bundles and intertwining operators for symmetric domains*, Adv. Math. **303** (2016), 1077–1121. [1](#)
- [11] S. S. Roy, *Homogeneous operators, jet construction and similarity*, Complex Anal. Oper. Theory **5** (2011), no. 1, 261–281. [1](#)
- [12] D. R. Wilkins, *Homogeneous vector bundles and Cowen-Douglas operators*, Internat. J. Math. **4** (1993), no. 3, 503–520.

(P. Deb) DEPARTMENT OF MATHEMATICS, BEN-GURION UNIVERSITY OF THE NEGEV, BEER-SHEVA, 84105, ISRAEL

Email address, P.Deb: prahllad.deb@gmail.com

(S. Hazra) MATHEMATICAL INSTITUTE IN OPAVA, SILESIAN UNIVERSITY IN OPAVA, NA RYBNICKU 626/1, 74601, OPAVA, CZECHIA

Email address, S. Hazra: somnath.hazra.2008@gmail.com