

The Multisecretary problem with many types

Omar Besbes

Columbia Business School, New York, NY, 10027
ob2105@columbia.edu

Yash Kanoria

Columbia Business School, New York, NY, 10027
ykanoria@gmail.com

Akshit Kumar

Columbia Business School, New York, NY, 10027
ak4599@columbia.edu

We study the multisecretary problem with capacity to hire up to B out of T candidates, and values drawn i.i.d. from a distribution F on $[0, 1]$. We investigate achievable regret performance, where the latter is defined as the difference between the performance of an oracle with perfect information of future types (values) and an online policy. While the case of distributions over a few discrete types is well understood, very little is known when there are many types, with the exception of the special case of a uniform distribution of types. In this work we consider a larger class of distributions which includes the few discrete types as a special case. We first establish the insufficiency of the common certainty equivalent heuristic for distributions with many types and “gaps” (intervals) of absent types; even for simple deviations from the uniform distribution, it leads to regret $\Theta(\sqrt{T})$, as large as that of a non-adaptive algorithm. We introduce a new algorithmic principle which we call “conservativeness with respect to gaps” (CwG), and use it to design an algorithm that applies to any distribution. We establish that the proposed algorithm yields optimal regret scaling of $\tilde{\Theta}(T^{1/2-1/(2(\beta+1))})$ for a broad class of distributions with gaps, where β quantifies the mass accumulation of types around gaps. We recover constant regret scaling for the special case of a bounded number of types ($\beta = 0$ in this case). In most practical network revenue management problems, the number of types is large and the current certainty equivalent heuristics scale poorly with the number of types. The new algorithmic principle called *Conservatism w.r.t Gaps* (CwG) that we developed, can pave the way for progress on handling many types for the broader class of network revenue management problems like order fulfillment and online matching.

Key words: multi-secretary problem, revenue management, regret analysis, online algorithms

1. Introduction

The classical secretary problem was introduced by Cayley (1875) and Moser (1956) and is one in which a decision maker is presented with a sequence of T independent values representing the candidate abilities and the decision maker must hire exactly one candidate for the secretary position. The decision maker interviews the candidates sequentially and after interviewing each candidate, the ability of the candidate is revealed; at which point the decision maker must make an irrevocable hire or reject decision, before the next candidate in the sequence can be interviewed. If a candidate is rejected, the decision maker moves on to interviewing the next candidate and if a candidate is hired, then the process terminates and the remaining candidates are automatically rejected. The decision maker is aware of the number of candidates T and the distribution of the

candidate abilities. The decision maker faces an optimal stopping problem where the goal of the decision maker is to maximize the expected ability of the hired candidate or equivalently minimize regret which is the difference between the expected ability of the most capable candidate and the candidate chosen by the decision maker. For a survey on the extensions of the classical secretary problem, refer to Freeman (1983), Ferguson (1989).

The multi-secretary variant of the above problem is one where instead of hiring just one secretary, the decision maker has a budget to hire B secretaries. The multi-secretary variant of the above problem was initially studied by Kleywegt and Papastavrou (1998) and Kleinberg (2005). Recently, Arlotto and Gurvich (2019) study achievable regret where the latter is defined as the expected difference between the sum of the ability of the B most capable candidates and the sum of the ability of the candidates hired by the decision maker. Arlotto and Gurvich (2019) showed that, when the distribution of types is discrete, regret is bounded uniformly for all values of number of candidates T and the hiring budget B , where the constant may scale with the reciprocal of the minimum probability mass on any type. In order to prove this result, they devise an adaptive policy called the Budget-Ratio (BR) policy where they compare the ratio of the remaining budget to the remaining number of candidates to interview and make the hire/reject decision by comparing the budget ratio to some fixed thresholds. This performance guarantee, in conjunction with a lower bound on regret from Kleinberg (2005) yields a tight understanding of the class of distributions over a few discrete types.

When there are many types, e.g., a continuum of types, very little is known. The only known result on regret with many types is that of Bray (2019), who considers the multi-secretary problem when the candidates abilities are distributed as independent uniform random variables over the continuous and bounded support $[0, 1]$. He showed that in this case, instead of the regret being uniformly bounded, the best possible scaling for this distribution of types is $\Theta(\log T)$.

This motivates us to ask the following questions in the present paper: *How does regret scale as a function of the underlying distribution in the presence of many types? What algorithmic approach allows to achieve optimal regret scaling with many types?*

The workhorse algorithmic approach used in the literature, including to prove the constant regret bound in the case of finite types (Arlotto and Gurvich 2019) and the logarithmic bound in the case of a uniform distribution over types (Bray 2019), is to use variants of a re-solving heuristic called the *certainty equivalent (CE) heuristic* where the idea is to address a stochastic optimization problem by replacing all the random variables about the future by their expectations and to use the solution of the optimization problem to make allocation decisions at each time step. Applied to the multisecretary problem, the CE heuristic computes a budget ratio, which is the ratio of the remaining hiring budget to the number of remaining candidates, and hires the candidate if the

complementary CDF of the candidate ability is less than the budget ratio, else the candidate is rejected; accordingly we term the inverse complementary CDF of the budget ratio the *CE threshold*. Arlotto and Gurvich (2019) additionally employ a smart tie-breaking rule for determining whether to admit a (discrete) type which is at the current CE threshold.

In the present work, we generalize the settings of Arlotto and Gurvich (2019) and Bray (2019) and study the multi-secretary problem under general distributions on $[0, 1]$ which allow for many types, ranging from a few discrete types to a continuum of types, together with gaps (intervals of absent types), and identify how the nature of the distribution affects the (best possible) scaling of regret. We refer to intervals where the probability mass is zero as *gaps*, and refer to intervals where the probability mass is positive as *mass clusters*. This generalization has both important practical and theoretical motivations. From a practical perspective, many settings which resemble the multi-secretary problem such as hiring, and closely related problems such as fulfillment problems, the types may indeed belong to a continuum and not all types (quality levels or demand locations) are present. The algorithmic insights we develop here may port over to broader classes of problems. From a theoretical perspective, the family of distributions we consider generalizes the distributions considered in prior work, and encompasses a broad range of distributions.

We summarize our main contributions as follows:

- We first establish that for any non-atomic type distribution with a gap, the certainty equivalent (CE) policy regret scales as $\Omega(\sqrt{T})$ (cf. Theorem 2). This is in contrast to the prior finding of (Bray 2019) that the regret of CE is $\Theta(\log T)$ for the uniform distribution. One or more gaps in the distribution cause the regret to increase to $\Omega(\sqrt{T})$. We note that this result applies to non-atomic distributions and hence the deterioration we prove is not due to a poor tie breaking rule (as is commonly documented in the case of a few discrete types), but to the presence of a gap in the type space. As such, this result highlights a new practical feature of distributions that could significantly affect how one should operate.
- In the presence of gaps, we establish a universal lower bound (for any policy) on the growth rate of the regret as a function of how mass accumulates around gaps. In particular, we establish that any policy must incur $\Omega(T^{1/2-1/(2(\beta+1))})$ where $\beta \geq 0$ quantifies the mass accumulation of types around gaps (cf. Theorem 3).
- To operate with distributions which have gaps, we introduce a new algorithmic principle which we call *Conservativeness with respect to gaps* (CwG). We devise a new CwG policy which makes a crucial modification to the CE policy. The idea is that if at any time the CE threshold is close to the boundary of a gap, CwG instead uses the gap as the acceptance threshold to avoid incurring large regret in the future (large regret would otherwise occur in the event that the CE threshold eventually falls on the opposite side of that gap).

- We establish that for a broad class of distributions with gaps, which we refer to as (β, ε_0) -clustered distributions (cf. Definition 1), the algorithm we devised is near-optimal, in that its worst case regret scales as $\tilde{O}(T^{1/2-1/(2(\beta+1))})$, matching the scaling of the lower bound in T up to polylogarithmic terms (cf. Theorem 4).

We also highlight here that for the case of a few discrete types, our algorithm recovers bounded regret, as in Arlotto and Gurvich (2019) (cf. Corollary 1). For the uniform distribution on $[0, 1]$, CwG is identical to CE since there are no gaps, and we recover the $O(\log T)$ result of Bray (2019).

In terms of techniques used, for the upper bound, we build upon the compensated coupling technique framework in Vera and Banerjee (2021). The novel aspect of our analysis include the *conservativeness with respect to gaps* principle and an analysis which can encompass a large class of distributions as opposed to just discrete distributions in Arlotto and Gurvich (2019), Vera and Banerjee (2021), Bumpensanti and Wang (2020) or some specific non-atomic distributions (eg. uniform) Bray (2019). For the lower bound in Theorem 3, we build on the ideas in Arlotto and Gurvich (2019) and generalize these. We consider two events of $\Omega(1)$ probability, and show that an optimal online algorithm can do well only on one event or the other but not on both the events. In doing so, we are able to characterize the “hardness” of problem instances through the parameter β which is a measure of local mass accumulation around the gaps.

The multi-secretary problem is a special case of a broader class of network revenue management (NRM) problems, or more broadly dynamic resource constrained reward collection problems; see Balseiro et al. (2021) for a recent survey and unified modeling framework for this class of problems. There is a wide variety of applications in auction theory Kleinberg (2005), online resource allocation Kleywegt and Papastavrou (1998), Talluri et al. (2004), order fulfillment Jasin and Sinha (2015), among others. Note that this literature typically assumes a few discrete demand and supply types. In the context of auctions, we can think of the multi-secretary problem as having B copies of an identical good with the arriving candidates bidding on the good, and the decision maker aims to sell the goods in the most profitable manner.

Vera and Banerjee (2021), Vera et al. (2021) generalized the arguments in Arlotto and Gurvich (2019) to a broader class of online packing and online matching problems and proved a uniform regret guarantee across all values of capacity B and time horizon T . They developed a technique called *compensated coupling* and used it to prove a constant regret guarantee without requiring any non-degeneracy assumptions. Bumpensanti and Wang (2020) also proved constant regret guarantees for a class of NRM problems however their algorithm and proof techniques differ significantly from that of Vera and Banerjee (2021), Vera et al. (2021). While all these papers impressively establish constant regret bounds, all of them assume a few discrete types, and their regret bounds scale polynomially in the number of types. However in most practical systems, the number of types

is in fact large. For example, in the multi-secretary problem, the set of possible candidates abilities could be very large. In the context of fulfillment problems, there are more than 40,000 zip codes in the United States and most of these are demand locations (i.e., demand types) for e-commerce websites like Amazon, Walmart, Best Buy, etc. While the regret guarantees obtained in previous work do not grow with demand volume, they scale poorly (polynomially) in the number of demand types. Our work on the one-dimensional case, i.e., the multisecretary problem, may pave the way for progress on handling many types in broader problem classes.

2. Model

Consider the multi-secretary hiring problem with a time horizon of T , i.e., at each time step $t \in [T]$, a decision maker is presented with a candidate with ability θ_t . Given a hiring budget of B , the decision maker can select at most B candidates so as to maximize the total expected ability of those selected up to and including the time T . At each time $t \in [T]$, once the candidate is interviewed and the ability θ_t becomes known, the decision maker must either hire the candidate or reject the candidate. A hired candidate cannot be rejected later and a rejected candidate cannot be hired later, making all the decisions irrevocable. In the case $B > T$, the hiring problem is trivial and hence for a fixed time horizon $T \in \mathbb{N}$, we will only be interested in hiring budgets $B \in [T]$. The candidate abilities $\theta_1, \theta_2, \dots, \theta_T$ are independent and identically distributed with a cumulative distribution function F with support in $[0, 1]$; and F is known to the decision maker. All the results will apply to any distribution with bounded support through appropriate scaling.

A policy is said to be *feasible* if the number of selected candidates is at most the hiring budget B and it is said to be an *online* (non-anticipating) policy if the decision on the t -th candidate is based only on the ability θ_t of the candidate at time t , the ability of the past candidates, $\{\theta_j\}_{j=1}^{t-1}$ and the history of the hire/reject decisions up to the time t .

Let U_1, U_2, \dots, U_T be a sequence of random variables that are independent and uniformly distributed over $[0, 1]$ and independent of the abilities $\theta_1, \theta_2, \dots, \theta_T$. (The U s will allow us to accommodate randomized policies.) Define the filtration $\mathcal{F}_t = \sigma(\theta_1, U_1, \theta_2, U_2, \dots, \theta_t, U_t)$ for all $t \in [T]$. A feasible online policy π is a sequence of $\{\mathcal{F}_t : t \in [T]\}$ -measurable binary random variables $\{\mu_1^\pi, \mu_2^\pi, \dots, \mu_T^\pi\}$ such that $\sum_{t=1}^T \mu_t^\pi \leq B$ almost surely. Here $\mu_t^\pi = 1$ means that the candidate interviewed at time t is hired. We define the set of feasible online policies as $\Pi(B, T)$.

For any feasible and online policy $\pi \in \Pi(B, T)$, define $A_t^\pi = \sum_{k=1}^t \theta_k \mu_k^\pi, \forall t \in [T]$ to be the accumulated ability of candidates hired up to the time t . The total expected ability under a policy $\pi \in \Pi(B, T)$ is given by $V_{\text{on}}^\pi(B, T) = \mathbb{E}[A_T^\pi] = \mathbb{E}\left[\sum_{t=1}^T \theta_t \mu_t^\pi\right]$. Fix $T \in \mathbb{N}$ and $B \in [T]$, the multi-secretary problem is to maximize the total expected ability given by $V_{\text{on}}^*(B, T) = \sup_{\pi \in \Pi(B, T)} V_{\text{on}}^\pi$.

Next we consider the offline, full-information version of the multi-secretary problem in which the candidate abilities $\theta_1, \theta_2, \dots, \theta_T$ are known before the hiring season begins. In the offline setting,

the problem essentially reduces to solving an integer program given as $V(B, T; \theta_1, \theta_2, \dots, \theta_T) = \max_{\mu} \{ \sum_{t=1}^T \theta_t \mu_t : (\mu_1, \mu_2, \dots, \mu_T) \in \{0, 1\}^n \text{ and } \sum_{t=1}^T \mu_t \leq B \}$ and the total expected ability by the offline problem is given as $V_{\text{off}}^*(B, T) = \mathbb{E} [V(B, T; \theta_1, \theta_2, \dots, \theta_T)]$. Since on each sample path of $\theta_1, \theta_2, \dots, \theta_T$, the offline problem chooses the top B candidates, we have that $V_{\text{off}}^*(B, T) \geq V_{\text{on}}^\pi(B, T), \forall \pi \in \Pi(B, T)$ and $\forall B \in [T]$. To measure the performance of a feasible online policy $\pi \in \Pi(B, T)$, we consider the offline problem as a benchmark and define the (expected) regret of the policy π as the difference between the expected value of the offline problem and the expected value attained by the policy π i.e., $\text{Regret}(B, T; \pi) \triangleq V_{\text{off}}^*(B, T) - V_{\text{on}}^\pi(B, T)$. We also define the (minimum achievable, expected) regret as the difference between the expected value of the offline problem and the expected value under the optimal online policy $\pi^* \in \Pi(B, T)$.

$$\text{Regret}(B, T) = \inf_{\pi \in \Pi(B, T)} \text{Regret}(B, T; \pi) = V_{\text{off}}^*(B, T) - V_{\text{on}}^*(B, T).$$

In what follows, we will focus on characterizing the growth rate of $\text{Regret}(B, T)$ as a function of T and the characteristics of the distribution of types.

3. Failure of the certainty equivalent policy under many types with gaps

One of the most popular algorithmic approaches in the network revenue management literature is the certainty-equivalent (CE) heuristic where the idea is to repeatedly solve for a deterministic approximation of a stochastic optimization problem by replacing random quantities by their expectations. When the number of types is small, versions of CE have been used in the literature Arlotto and Gurvich (2019), Bumpensanti and Wang (2020), Vera et al. (2021), Vera and Banerjee (2021), Vera et al. (2020) to prove constant regret bounds. The success of these algorithms is predicated on breaking ties appropriately. With a continuum of types, ties are a zero probability event. As such, it might seem “sufficient” to simply apply CE. Indeed, in the special case of a uniform distribution, Bray (2019) used CE to achieve $\mathcal{O}(\log T)$ regret, and showed that this is the best scaling achievable. Let `ce` denote the Certainty Equivalent (CE) policy (which is formally defined in Algorithm 1 below). We formally state the result of Bray (2019) in Theorem 1.

THEOREM 1 (Bray (2019)). *Suppose the candidate-ability distribution F is Uniform($[0, 1]$). For all $T \in \mathbb{N}$ and $B \in [T]$, there exists a policy $\text{CE} \in \Pi(B, T)$ and a universal constant $C_1 < \infty$ such that $\text{Regret}(B, T; \text{CE}) \leq C_1 \log T$. Furthermore, for any online policy $\pi \in \Pi(B, T)$, there exists a time horizon $T_0 < \infty$ and a constant $C_2 > 0$ such that $\text{Regret}(B, T) \geq C_2 \log T$ for all $T \geq T_0$.*

Next, we explore the performance of CE for more general non-atomic type distributions. Note that we will only consider non-atomic distributions F in this section, hence the tie-breaking rule at

the threshold will not matter. Let $\mu_1^{\text{ce}}, \mu_2^{\text{ce}}, \dots, \mu_T^{\text{ce}}$ denote the sequence of hiring decisions under the CE policy and define $B_t = B - \sum_{k=1}^t \mu_k^{\text{ce}}$ as the remaining budget after the time t . The idea of the CE policy is as follows: before the arrival of the candidate at time $t + 1$, given the remaining budget B_t and the number of remaining candidates being $T - t$, one selects a threshold Θ_{t+1}^* such that the expected number of candidates with ability greater than the threshold Θ_{t+1}^* in the remaining $T - t$ time periods matches the remaining budget of B_t . Given the distribution of the candidate ability F , the CE policy essentially solves for Θ_{t+1}^* such that $(T - t)(1 - F(\Theta_{t+1}^*)) = B_t$. After observing the candidate ability θ_{t+1} at time $t + 1$, the CE policy hires the candidates if $\theta_{t+1} \geq \Theta_{t+1}^*$, else the candidate is rejected. The CE policy is formalized in Algorithm 1.

Algorithm 1: Certainty Equivalent (CE) policy

Input: Time Horizon T , Hiring Budget B , distribution F

Initialize: $B_0 = B$

for $t = 1$ **to** T **do**

$$p_{t-1}^{\text{ce}} = 1 - \frac{B_{t-1}}{T-t+1}$$

Observe a candidate of ability θ_t

if $F(\theta_t) \geq p_{t-1}^{\text{ce}}$ **then**

Hire the candidate with ability θ_t and $B_t \leftarrow B_{t-1} - 1$

else

Reject the candidate with ability θ_t and $B_t \leftarrow B_{t-1}$

end

end

THEOREM 2 (Failure of CE Policy). Fix any $\eta \in (0, 1)$ and $\varepsilon \in (0, 1/2]$. Suppose the candidate-ability distribution F is any non-atomic distribution that has a gap of length at least η , i.e., $\exists c \in (0, 1 - \eta)$ such that $F(c) = F((c + \eta)^-)$, and such that there is mass at least ε on each side of the gap $\min\{F(c), 1 - F(c + \eta)\} \geq \varepsilon$. Then for the CE policy defined in Algorithm 1, there exists $T_0 \equiv T_0(\varepsilon) < \infty$, a constant $C \equiv C(\eta, \varepsilon) > 0$ and $B \in [T]$ such that $\text{Regret}(B, T; \text{CE}) \geq C\sqrt{T}$ for all $T \geq T_0$.

From Theorem 1 it follows that the regret of CE is logarithmic for uniform type distribution, however Theorem 2 shows that the regret of the CE policy increases dramatically if there is a gap in the types, even when one maintains the uniform distribution of types (or any other distribution) outside of the gap. As a matter of fact, the regret scaling is as large as that of a non-adaptive policy.

Argument for a specific distribution. Next we present the proof of Theorem 2 for a specific distribution. In Appendix EC.1, we describe how this proof extends to all distributions covered by the theorem. Our argument will illuminate the Achilles heel of the CE policy and point towards how to improve upon the CE policy. Let us consider the simplest departure from the uniform distribution considered in Theorem 1. Consider the distribution $F_0 \sim \text{Uniform}([0, 1/4] \cup [3/4, 1])$ where we have a gap in the interval $(1/4, 3/4)$ and the density is uniform in the intervals $[0, 1/4]$ and $[3/4, 1]$. (Note that Theorem 2 allows arbitrary distributions containing a non-trivial gap; only for ease of exposition we discuss the F_0 distribution here.) The probability mass in the intervals $[0, 1/4]$ and $[3/4, 1]$ is $1/2$ each. We assume that the budget $B = \lceil T/2 \rceil$. The number of candidates that arrive with ability in the interval $[3/4, 1]$ is distributed as $\text{Binomial}(T, 1/2)$ and hence the number of candidates with ability in $[3/4, 1]$ on any sample path is typically $\lceil T/2 \rceil \pm \Theta(\sqrt{T})$. The offline (i.e., hindsight) optimal essentially chooses the B -th largest value in the sample path of the candidate abilities as the threshold and accepts candidates with ability greater than or equal to the threshold. Due to the discrepancy mentioned above, the B -th largest value (the offline threshold) on any sample path could either be in the interval $[0, 1/4]$ or in the interval $[3/4, 1]$ and this information is not known to the CE policy a priori. We observe that the CE threshold quantile p_t^{ce} is in fact a martingale with respect to the filtration \mathcal{F}_t (see Lemma EC.1 in the appendix) and the martingale difference at time t has a magnitude proportional to $1/(T-t)$, corresponding to the change in the threshold from time step $t-1$ to t . In fact, to leading order, p_t^{ce} is an unbiased random walk with steps which are independent $(\text{Bernoulli}(1/2) - 1/2)/(T-t)$. Therefore, the expected distance of the threshold p_t^{ce} from $p_0^{\text{ce}} = 1/2$ is of the order of $1/\sqrt{T-t}$. As is usual with an unbiased random walk/Brownian motion, there exists a universal constant $\nu > 0$ such that for each $t \in (T/3, 2T/3)$, with probability at least ν , the following event $E_t = E_{t,1} \cap E_{t,2}$ occurs: the CE threshold quantile p_t^{ce} is in the interval $[0, 1/2 - 1/\sqrt{T}]$ (call this $E_{t,1}$) whereas the offline threshold quantile p_t^{og} at time t is larger than $1/2$ (call this $E_{t,2}$). Under $E_{t,1}$, independent of the past with probability $1/\sqrt{T}$ the ability of the next arrival $\theta_{t+1} \in [1/4 - 1/(2\sqrt{T}), 1/4]$ (call this $E_{t,3}$; we observed $\mathbb{P}(E_{t,3}|E_{t,1}) = 1/\sqrt{T}$) so that θ_{t+1} is accepted by CE but would have been rejected by offline, and since $E_{t,3}$ leads to acceptance it only increases (slightly) the likelihood that $E_{t,2}$ will occur

$$\begin{aligned} \mathbb{P}(E_{t,2}|E_{t,1}) &\leq \mathbb{P}(E_{t,2}|E_{t,1}, E_{t,3}) \\ \Rightarrow \mathbb{P}(E_t \cap E_{t,3}) &= \mathbb{P}(E_{t,1})\mathbb{P}(E_{t,3}|E_{t,1})\mathbb{P}(E_{t,2}|E_{t,1}, E_{t,3}) \\ &\geq (1/\sqrt{T})\mathbb{P}(E_{t,1})\mathbb{P}(E_{t,2}|E_{t,1}) = \mathbb{P}(E)/\sqrt{T} \geq \nu/\sqrt{T}. \end{aligned}$$

And $E_t \cap E_{t,3}$ leads to an additional regret under CE of at least the size of the gap $3/4 - 1/4 = 1/2$ due to the decision at time $t+1$ (since some future arrival with value $\geq 3/4$ will need to be rejected

because of the acceptance of $\theta_{t+1} \in [1/4 - 1/(2\sqrt{T}), 1/4]$ at $t+1$). Hence the total expected regret under CE is at least $\sum_{t=T/3}^{2T/3} \nu/(2\sqrt{T}) = \nu\sqrt{T}/6$, i.e., $\Omega(\sqrt{T})$.

We wish to emphasize here that the fundamental driver of the $\Omega(\sqrt{T})$ regret scaling as for the distribution F_0 is the presence of a gap, as is clear from the informal argument above. To clarify this point further, let us contrast the distribution F_0 which has a gap, with the uniform distribution considered in Bray (2019) which does not have a gap. The argument regarding the threshold p_t^{ce} being a martingale and in expectation order $1/\sqrt{T-t}$ away from $p_0^{\text{ce}} = 1/2$ is still valid in the case of the uniform distribution. The only difference now is that in the case of the uniform distribution, the cost of a mistake is also of the order of $1/\sqrt{T-t}$ and hence the expected compensation at time t is of the order $1/\sqrt{T-t} \times 1/\sqrt{T-t} = 1/(T-t)$. Summing this over $t = 1, 2, \dots, T$ results in a regret of $\mathcal{O}(\log T)$.

Remark As a prelude to the achievability results to come, we note that a loss of $\Omega(\sqrt{T})$ can, in fact, be avoided when facing a distribution such as $F_0 \sim \text{Uniform}([0, 1/4] \cup [3/4, 1])$ with gaps. In particular, our main achievability result will imply that for the distribution F_0 , if one uses the Conservative wrt Gaps (CwG) policy that we devise, the regret scales as $\mathcal{O}((\log T)^2)$ (see Corollary 2) which is a drastic improvement upon the performance of the CE policy, and is near optimal. The main idea is that when p_t^{ce} is close to a gap quantile, CE tends to make costly mistakes, and we avoid such mistakes by instead using the nearby gap quantile itself as the threshold quantile.

4. Main Results

While the CE policy saw its performance deteriorate in the presence of gaps in the type space, it is not clear what performance is achievable as a function of the type distribution under many (e.g., a continuum of) types, and what algorithmic approach can achieve near optimal performance. To answer these questions, we will anchor our analysis around a general family of distributions which allow for gaps in the type space and can capture as special cases discrete distributions as well as the uniform distribution.

We next define (β, ε_0) -clustered distributions. For any $q \in [0, 1]$, we define $F^{-1}(q) \triangleq \inf\{v : F(v) \geq q\}$.

DEFINITION 1 ((β, ε_0) -CLUSTERED DISTRIBUTIONS). Fix $\beta \in [0, \infty)$ and $\varepsilon_0 \in (0, 1]$. A distribution F is said to be (β, ε_0) -clustered if there exists $n \in \mathbb{N} \cup \{0\}$ and *gap quantiles* $q_0^* = 0 < q_1^* < \dots < q_n^* < q_{n+1}^* = 1$ such that we have

- (a) (Cluster “density” requirement) $\forall i \in [n+1], \forall q, \tilde{q} \in (q_{i-1}^*, q_i^*]$, we have that $|F^{-1}(q) - F^{-1}(\tilde{q})| \leq |q - \tilde{q}|^{\frac{1}{\beta+1}}$.
- (b) (Cluster size requirement) $q_i^* - q_{i-1}^* \geq \varepsilon_0, \forall i \in [n+1]$.

Let $\mathcal{F}_{\beta, \varepsilon_0}$ denote the class of (β, ε_0) -clustered distributions. Note that for any $\beta \in [0, \infty)$, the requirement (a) implies that for values $u < v$ such that $F^{-1}((q_{i-1}^*)^+) < u < v < F^{-1}(q_i^*)$, we have $F(u) < F(v)$, i.e., there is strictly positive probability mass in the interval of values $(u, v]$. One can show this property by contradiction. Suppose $F(u) = F(v) = q$. If $q = q_i^*$, then $F^{-1}(q_i^*) \leq v$ by definition of $F^{-1}(\cdot)$ so we have a contradiction. Hence $q < q_i^*$. Let $\tilde{q} \triangleq q + \min(q_i^* - q, (v - u)^{\beta+1})$. Then $\tilde{q} \leq q_i^*$ by definition of \tilde{q} , and $F^{-1}(\tilde{q}) > v = u + (v - u) \geq F^{-1}(q) + (v - u)$ by definition of $F^{-1}(\cdot)$ and hence $F^{-1}(\tilde{q}) - F^{-1}(q) > v - u \geq (\tilde{q} - q)^{1/(\beta+1)}$, i.e., requirement (a) is violated, which is a contradiction. This holds for every (positive length) subinterval of $H_i \triangleq [F^{-1}((q_{i-1}^*)^+), F^{-1}(q_i^*)]$, for all $i \in [n+1]$, where $F^{-1}(q^+) \triangleq \lim_{\epsilon \rightarrow 0^+} F^{-1}(q + \epsilon)$. Motivated by this “positive mass everywhere” property, we call the (H_i) ’s *mass clusters* or just *clusters*. We will call *gaps* the complementary intervals $G_i \triangleq (F^{-1}(q_i^*), F^{-1}((q_i^*)^+))$ for $i \in [n]$, and the intervals at the extremes $G_0 = [0, F^{-1}(0^+))$, $G_{n+1} = (F^{-1}(1), 1]$, since they contain no probability mass.

The requirement (a) can be thought of as a within-cluster “density” requirement, which becomes weaker as β increases; we can think of β as quantifying the within-cluster mass density (with a decreasing relationship). Requirement (b) is a cluster size requirement, ε_0 being the minimum cluster size; this requirement becomes more stringent as ε_0 increases. The policy we will propose will take the gap quantiles as an input, and our performance guarantee will be stronger for smaller β and larger ε_0 . Accordingly, we will find it advantageous to only include gap quantiles q_i which correspond to positive-length intervals (gaps) in value space $F^{-1}((q_{i-1}^*)^+) < F^{-1}(q_i^*)$ since including a gap quantile q for which $F^{-1}((q_{i-1}^*)^+) = F^{-1}(q_i^*)$ will hurt our ability to satisfy the cluster size requirement (b), without helping to satisfy the cluster density requirement (a). Hence, G_1 through G_n will be non-trivial (positive length) gaps, whereas G_0 and G_{n+1} may be empty or may be intervals of non-zero length at the extremes.

Next we present some examples of (β, ε_0) -clustered distributions including discrete distributions, as well the uniform distribution, along with the appropriate choices of gap quantiles.

EXAMPLE 1 (UNIFORM DISTRIBUTION). The uniform distribution over $[0, 1]$, considered in Bray (2019) in the context of the multi-secretary problem, is a $(0, 1)$ -clustered distribution with $n = 0$ gaps. The gap quantiles are only the trivial ones $q_0^* = 0$ and $q_1^* = 1$. There is a single mass cluster $H_1 = [0, 1]$ with mass 1, which clearly satisfies the cluster density requirement with $\beta = 0$.

EXAMPLE 2 (DISCRETE DISTRIBUTIONS). Consider a discrete distribution (as studied in Arlotto and Gurvich 2019). Let the support $\{a_1, a_2, \dots, a_m\}$ with probability masses $\{f_1, f_2, \dots, f_m\}$. Assume that $0 \leq a_1 < a_2 < \dots < a_m \leq 1$. We make use of the natural choice of gap quantiles $q_i^* = \sum_{j=1}^i f_j$ for all $i \in [m-1]$, leading to gaps $G_0 = [0, a_1)$, $G_i = (a_i, a_{i+1}) \forall i \in [m-1]$, $G_m = (a_m, 1]$ and clusters $H_i = \{a_i\} \forall i \in [m]$. Now for $q, \tilde{q} \in (q_{i-1}^*, q_i^*)$, we have that $|F^{-1}(q) - F^{-1}(\tilde{q})| = 0 \leq |q - \tilde{q}|$, i.e., the cluster density requirement is satisfied for $\beta = 0$. Defining

$\varepsilon_0 \triangleq \min\{f_1, f_2, \dots, f_m\}$ the cluster size requirement is satisfied. Therefore the discrete distribution belongs to the class of $(0, \varepsilon_0)$ -clustered distributions where $\varepsilon_0 = \min_{i \in [m]} f_i$ is the minimum probability mass in the support.

EXAMPLE 3 (A CLASS OF BIMODAL DISTRIBUTIONS). An example of a (β, ε_0) -clustered distribution with $n = 1$ gap (with gap quantile $q_1^* = 1/2$), for general $\beta \geq 0$ and $\varepsilon_0 = 1/2$, which we will make use of to prove our lower bound results is presented below:

$$F_\beta(x) = \begin{cases} -2 \cdot 4^\beta \cdot (1/4 - x)^{\beta+1} + 1/2 & 0 \leq x \leq 1/4 \\ 1/2 & 1/4 \leq x \leq 3/4 \\ 2 \cdot 4^\beta \cdot (x - 3/4)^{\beta+1} + 1/2 & 3/4 \leq x \leq 1 \end{cases} \quad (1)$$

$$F_\beta^{-1}(q) = \begin{cases} 1/4 - (1 - 2q)^{\frac{1}{\beta+1}}/4, & 0 \leq q \leq 1/2 \\ (2q - 1)^{\frac{1}{\beta+1}}/4 + 3/4, & 1/2 < q \leq 1 \end{cases} \quad (2)$$

Using the F_β^{-1} in (2), we can easily verify that F_β is indeed is a $(\beta, 1/2)$ -clustered distribution, with one gap $G_1 = (1/4, 3/4)$ and clusters $H_1 = [0, 1/4]$ and $H_2 = [3/4, 1]$. Refer to Figure 1 for a plot of the density f_β and the CDF F_β of the $(\beta, 1/2)$ -clustered distribution defined in (1).




Figure 1 (Left) PDF f_β for $\beta = 0, 1, 2$. Notice the gap from $1/4$ to $3/4$, (Right) CDF F_β for $\beta = 0, 1, 2$.

4.1. Lower bound on Performance

THEOREM 3 (universal lower bound). Consider any $\beta \in [0, \infty)$ and $\varepsilon_0 \in (0, 1/2]$. Then there exists a candidate-ability distribution $F \in \mathcal{F}_{\beta, \varepsilon_0}$, a time horizon $T_0 \equiv T_0(\beta, \varepsilon_0) < \infty$, a positive constant $C \equiv C(\beta, \varepsilon_0)$ such that for all $T \geq T_0$ and for any online policy $\pi \in \Pi(B, T)$, we have that,

$$\sup_{B \in [T]} \text{Regret}(B, T; \pi) \geq CT^{\frac{1}{2} - \frac{1}{2(\beta+1)}} \mathbb{1}\{\beta > 0\} + C \log T \mathbb{1}\{\beta = 0\}.$$

Discussion of Theorem 3 This result establishes an impossibility result: it says that for any fixed $\beta \in [0, \infty)$ and $\varepsilon_0 \in (0, 1/2]$, there exists a distribution $F \in \mathcal{F}_{\beta, \varepsilon_0}$ for which no online policy can achieve a better regret scaling than the one pointed out in Theorem 3. This lower bound also highlights that the fundamental limits of the regret scaling are governed by the parameter β which characterizes the curvature of the distribution around the gap boundaries. We observe that as $\beta \uparrow \infty$, the scaling of regret approaches \sqrt{T} ; therefore, no matter the online policy, it will suffer the same regret as that of a non adaptive policy. Hence β can be seen as characterizing the “hardness” of an instance. The parameter β has a physical interpretation as well. It captures how mass accumulates in the type space. For some intuition, consider the F_β distribution described in (1) and consider the gap boundary at $3/4$. As we move a distance δ from the boundary point $3/4$, the probability mass accrued grows as $C\delta^{\beta+1}$ for some universal constant $C > 0$. Alternately, to accrue a probability mass of ε , we need to move a distance $C\varepsilon^{1/(\beta+1)}$ away from the boundary $3/4$. Therefore as β increases, the distance one needs to travel to collect a probability mass of ε also increases and this property is what makes the instances harder as β increases. We elaborate on this property in the formal proof in the Appendix EC.2.

Proof Sketch We defer the formal proof of Theorem 3 to Appendix EC.2 and provide here a proof sketch. For $\beta = 0$, our lower bound follows from Theorem 1. For the case of $\beta > 0$, we consider the distributions F_β defined in (1). We consider two events of $\Omega(1)$ probability - one is the perturbation of the other and show that on both events, the online algorithm will make $\Omega(\sqrt{T})$ mistakes and minimum compensation to be paid for these mistakes is $\Omega(T^{-1/2(\beta+1)})$. To determine the minimum compensation amount $\Omega(T^{-1/2(\beta+1)})$, we leverage how the mass accumulates over space (characterized by Definition 1). Combining the two gives us that the regret scales as $\Omega(T^{(1/2)-1/(2(\beta+1))})$.

4.2. Near-optimal algorithm: Conservativeness with respect to Gaps

4.2.1. Conservative with respect to Gaps (CwG) policy We observe that CE policy breaks down for distributions with many types and “gaps” (intervals) of absent types; it leads to regret scaling as $\Theta(\sqrt{T})$, as large as that of a non-adaptive algorithm. We identified that the main driver for the \sqrt{T} regret of the CE policy is the presence of gaps. To solve this issue, we introduce a new algorithmic principle which we call “conservativeness with respect to gaps” (CwG), and use it to provably achieve near optimal regret scalings for the (β, ε_0) -clustered distributions which allow for gaps. The idea of CwG is that if there is a risk that the acceptance threshold based on CE will move across a given gap in the future, then CwG uses that gap as the acceptance threshold instead of using the CE-based threshold. Based on the CwG principle, we devise a new policy called the Conservative with respect to Gaps (CwG) policy which we present in Algorithm 2.

Algorithm 2: Conservativeness with respect to Gaps (CwG)

Input: Time Horizon T , Hiring Budget B , (β, ε_0) -clustered dist. F with gaps $G_i = (a_i, b_i)$

Initialize: $B_0 = B, q_i^* = F(a_i) = F(b_i^-), \forall i \in [n], \tilde{T} = \max\{0, T - \lfloor 16 \log(1/\varepsilon_0)/\varepsilon_0^2 \rfloor\}$

for $t = 1$ **to** \tilde{T} **do**

$$p_{t-1}^{\text{ce}} = 1 - \frac{B_{t-1}}{T-t+1}$$

$$\mathcal{S}_{t-1} = \left\{ i : p_{t-1}^{\text{ce}} \in \mathcal{B} \left(q_i^*, \sqrt{2 \frac{\log(T-t+1)}{T-t+1}} \right) \right\}$$

if $\mathcal{S}_{t-1} = \emptyset$ **then**

$$p_{t-1}^{\text{cwg}} = p_{t-1}^{\text{ce}}$$

else

$$j_{t-1}^* = \arg \min_{i \in \mathcal{S}_{t-1}} |p_{t-1}^{\text{ce}} - q_i^*|$$

$$p_{t-1}^{\text{cwg}} = q_{j_{t-1}^*}^*$$

end

Observe a candidate of ability θ_t and form the set $\mathcal{I}_t = \{q \in [0, 1] : F^{-1}(q) = \theta_t\}$

Let X_t be a uniform sample from the set \mathcal{I}_t

if $X_t \geq p_{t-1}^{\text{cwg}}$ **and** $B_{t-1} > 0$ **then**

Hire the candidate and $B_t \leftarrow B_{t-1} - 1$

else

Reject the candidate and $B_t \leftarrow B_{t-1}$

end

end

$$\text{Define } p_{\tilde{T}}^{\text{ce}} = 1 - \frac{B_{\tilde{T}}}{T-\tilde{T}}$$

for $t = \tilde{T} + 1$ **to** T **do**

Observe a candidate of ability θ_t and form the set $\mathcal{I}_t = \{q \in [0, 1] : F^{-1}(q) = \theta_t\}$

Let X_t be a uniform sample from the set \mathcal{I}_t

if $X_t \geq p_{\tilde{T}}^{\text{ce}}$ **and** $B_{t-1} > 0$ **then**

Hire the candidate and $B_t \leftarrow B_{t-1} - 1$

else

Reject the candidate and $B_t \leftarrow B_{t-1}$

end

end

The algorithm essentially operates in two phases. For simplicity, assume that $T \geq \lfloor 16 \log(1/\varepsilon_0)/\varepsilon_0^2 \rfloor$, then for the first $\tilde{T} \triangleq T - \lfloor 16 \log(1/\varepsilon_0)/\varepsilon_0^2 \rfloor$ steps we make use of CwG principle where we shift the re-solving threshold p_t^{ce} to one of the quantiles corresponding to the boundary of a gap if the re-solving threshold p_t^{ce} is close to any one of them and use this conservative threshold p_t^{cwg} as our acceptance threshold. Note that this policy works for arbitrary distributions. For simplicity, let us just discuss the case of non-atomic and atomic distributions separately and the insights would translate to any general distribution as well. For a non-atomic distribution, after

observing the type θ_t , we form the set \mathcal{I}_t and since for a non-atomic distribution, F^{-1} is increasing, we have that \mathcal{I}_t is a singleton and hence $X_t = F(\theta_t)$. For an atomic distribution, after observing a type θ_t , we form a set \mathcal{I}_t which is one of the interval Q_i discussed in Example 2. If $p_t^{\text{cwg}} \notin Q_i$ then the hire/reject decision is unambiguous. The only case of ambiguity is when $p_t^{\text{cwg}} \in Q_i$, in which case we make use of randomization to break ties where we consider a random sample X_t drawn uniformly from the interval Q_i and hire the candidate only if the random sample X_t is greater than or equal to p_t^{cwg} . In the final $\lceil 16 \log(1/\varepsilon_0)/\varepsilon_0^2 \rceil$ time steps, the radius $\sqrt{2 \log(T-t)/(T-t)}$ by which we measure the closeness of CE threshold p_t^{ce} and the quantiles $\{q_i^*\}_{i=1}^n$ becomes too large and we can no longer accurately set the CwG quantile threshold p_t^{cwg} . In that case, we simply make use of the static allocation policy where we solve for a threshold $p_{\tilde{T}}^{\text{ce}}$ at time \tilde{T} and use that for the rest of the remaining $\lceil 16 \log(1/\varepsilon_0)/\varepsilon_0^2 \rceil$ time steps.

4.2.2. Performance Analysis

THEOREM 4. *There is a universal constant $C < \infty$ such that the following occurs. For any $\beta \in [0, \infty)$ and $\varepsilon_0 \in (0, 1]$, suppose the candidate-ability distribution F with associated gaps is (β, ε_0) -clustered. Then for all $T \in \mathbb{N}$ and for all $B \in [T]$, the regret of our CwG policy (Algorithm 2) is upper bounded as*

$$\text{Regret}(B, T; \text{CwG}) \leq C \left(1 + \frac{1}{\beta}\right) (\log T)^{\frac{1}{2} + \frac{1}{2(\beta+1)}} T^{\frac{1}{2} - \frac{1}{2(1+\beta)}} \cdot \mathbb{1}\{\beta > 0\} + C(\log T)^2 + C \frac{\sqrt{\log(1/\varepsilon_0)}}{\varepsilon_0}$$

Theorem 4 follows from Theorem 5 by setting $\delta = 0$ and the proof of Theorem 5 is deferred to Section 5.5.

Discussion of Theorem 4 By comparing the upper bound in Theorem 4 to the lower bound in Theorem 3, we note that the scaling of the upper bound matches the scaling of the lower bound in T up to a polylogarithmic factor and hence the proposed CwG policy is near-optimal. In the case of the CE policy, we had identified that the main driver of the regret scaling as $\Theta(\sqrt{T})$ is the presence of gaps in the distribution of candidate abilities. We now have established in Theorem 4 that one can overcome the difficulty introduced by gaps in the distribution by using the CwG principle that we devised.

REMARK 1. The discrete distribution considered in Example 2 belongs the class of $(0, \varepsilon_0)$ -clustered distributions and hence from Theorem 4, it follows that $\text{Regret}(B, T) = \mathcal{O}((\log T)^2)$. However recall from Example 2 that for discrete distributions we have that $\forall i \in [n+1], \forall q, \tilde{q} \in Q_i, |F^{-1}(q) - F^{-1}(\tilde{q})| = 0$ and this distinguishes the discrete distribution from the general $(0, \varepsilon_0)$ -clustered distributions. This distinction allows us to obtain stronger regret guarantees than the one implied by Theorem 4 and recover the result of Arlotto and Gurvich (2019).

COROLLARY 1. *Suppose the candidate-ability distribution is F where F is a discrete distribution with support $\{a_1, a_2, \dots, a_m\}$ and probability masses $\{f_1, f_2, \dots, f_m\}$. Define $\varepsilon_0 = \min\{f_1, f_2, \dots, f_m\}$. Then, for all $T \in \mathbb{N}$ and for all $B \in [T]$, we have $\text{Regret}(B, T) \leq \text{Regret}(B, T; \text{CwG}) \leq C\sqrt{\log(1/\varepsilon_0)}/\varepsilon_0$ for the universal constant $C < \infty$ in Theorem 4.*

The proof of Corollary 1 follows by modifying the analysis presented for Theorem 4 and we defer the details of the modifications to Section 5.6.

COROLLARY 2. *Suppose the candidate-ability distribution is F_0 where F_0 is defined in (1) for $\beta = 0$. Then we have that for all $T \in \mathbb{N}$ and for all $B \in [T]$ the regret of our CwG policy is upper bounded as $\text{Regret}(B, T; \text{CwG}) \leq C(\log T)^2$ for the universal constant $C < \infty$ in Theorem 4.*

This corollary follows immediately from Theorem 4 by setting $\beta = 0$.

Discussion of Corollary 2 The distribution $F_0 \sim \text{Uniform}([0, 1/4] \cup [3/4, 1])$ is possibly the simplest variant of the uniform distribution with a gap. Corollary 2 shows that regret scales as $\mathcal{O}((\log T)^2)$ for the distribution F_0 which is a “hard” instance for CE policy. This is a significant improvement from the $\Omega(\sqrt{T})$ regret scaling of the CE policy for the same distribution F_0 and the regret of the CwG policy for the F_0 distribution is only a $\log T$ factor off compared to the regret for the uniform distribution (cf. Theorem 1). The key takeaway message from Corollary 2 in conjunction with Theorem 2 is that the presence of gaps is not a fundamental driver for the deterioration in regret and one can overcome the difficulty posed by gaps by using the CwG principle that we discussed above.

4.2.3. Extension to many small discrete types The definition of the (β, ε_0) -clustered distributions (Definition 1) precludes the clusters H_i from containing intervals of non-zero length such that the probability mass in such an interval is zero. However, we can, in fact, extend our definition of (β, ε_0) -clustered distribution to include what we refer to as *many small discrete types* (with intervals containing no probability mass between them). The guarantees of Theorem 4 extend to the setting with many small discrete types. To get some intuition for the setting with many small discrete types, let us consider an example. Consider the distribution F_0 defined in (1) for $\beta = 0$. We have a non-atomic distribution over the intervals $[0, 1/4]$ and $[3/4, 1]$. We can discretize the interval $[0, 1/4]$ into $1/(4\delta)$ grid points located at $\{0, \delta, 2\delta, \dots, 1/4\}$ such that $\mathbb{P}(i\delta) = 2\delta$. This would constitute a setting with many small discrete types since there are a large number of atomic types (separated by small empty intervals) and the probability mass of each type is small, i.e., it is proportional to δ . As $\delta \rightarrow 0$, we recover the F_0 distribution in the limit. A similar discretization can be done for $[3/4, 1]$ as well. Such a discretization of continuum of types into many small discrete types can be done for general (β, ε_0) -clustered distributions. The discretization may not even be

uniformly spaced. We require that maximum distance between the discretized grid points be at most δ . In such discretizations, we have some flexibility in choosing which empty intervals to classify as “gaps”. In the case of many small types, if the size of the empty intervals (due to discretization) is at most δ then we may consider the entire clump of these many small types as belonging to one cluster say H_i and hence one quantile interval say Q_i . We present a generalization of Definition 1 which we call $(\beta, \varepsilon_0, \delta)$ -clustered distribution.

DEFINITION 2 ($(\beta, \varepsilon_0, \delta)$ -CLUSTERED DISTRIBUTIONS). Fix $\beta \in [0, \infty)$, $\varepsilon_0 \in (0, 1]$ and $\delta \in [0, 1]$. A distribution F is said to be $(\beta, \varepsilon_0, \delta)$ -clustered if there exists $n \in \mathbb{N} \cup \{0\}$ and *gap quantiles* $q_0^* = 0 < q_1^* < \dots < q_n^* < q_{n+1}^* = 1$ such that we have

- (a) (Generalized cluster “density” requirement) $\forall i \in [n + 1], \forall q, \tilde{q} \in (q_{i-1}^*, q_i^*)$, we have that

$$|F^{-1}(q) - F^{-1}(\tilde{q})| \leq |q - \tilde{q}|^{\frac{1}{\beta+1}} + \delta.$$
- (b) (Cluster size requirement) $q_i^* - q_{i-1}^* \geq \varepsilon_0, \forall i \in [n + 1]$.

THEOREM 5. *There is a universal constant $C < \infty$ such that the following occurs. For any $\beta \in [0, \infty)$ and $\varepsilon_0 \in (0, 1]$, suppose the candidate-ability distribution F with associated gaps is $(\beta, \varepsilon_0, \delta)$ -clustered with many small types separated by distance at most δ . Then for all $T \in \mathbb{N}$ and for all $B \in [T]$, the regret of the CwG policy is upper bounded as*

$$\begin{aligned} \text{Regret}(B, T; \text{CwG}) &\leq C(1 + 1/\beta)(\log T)^{\frac{1}{2} + \frac{1}{2(\beta+1)}} T^{\frac{1}{2} - \frac{1}{2(1+\beta)}} \cdot \mathbb{1}\{\beta > 0\} + C(\log T)^2 \\ &\quad + C\delta\sqrt{T \log T} + C\sqrt{\log(1/\varepsilon_0)}/\varepsilon_0 \end{aligned}$$

The proof of Theorem 5 is provided in Section 5.5. Note that as $\delta \rightarrow 0$, we obtain a continuum of types as the limit of the many small types and we recover our regret bound in Theorem 4. Consider scaling δ in the parameter T : if $\delta = o(1/\sqrt{T})$, we again recover a regret bound with the same scaling as that in Theorem 4.

4.3. Numerical Simulations

We simulate the CE and CwG policy for different distributions with the total number of candidates T ranging from 100 to 100,000 and consider that the hiring budget scales as $B = T/2$. In Figure 2 on the left, we compare the performance of the CE and CwG policy for the candidate ability distribution $F_0 \sim \text{Uniform}([0, 1/4] \cup [3/4, 1])$.

We observe a drastic difference in the performance of the two policies. For the CE policy, the regret scales as $\Theta(\sqrt{T})$ which aligns with our theoretical result in Theorem 2. For the CwG policy, regret scales as $\mathcal{O}((\log T)^2)$ which is in line with Corollary 2.

In Figure 2 on the right, we observe the regret scaling of the CwG policy for different distributions and plot them on a linear-log scale. The distributions we consider are discrete distribution parameterized by ε , the uniform distribution and $F_0 \sim \text{Uniform}([0, 1/4] \cup [3/4, 1])$. We consider a




Figure 2 (Left) The plot compares the performance of the CE and CwG policies on $F_0 \sim \text{Uniform}([0, 1/4] \cup [3/4, 1])$. (Right) The plot contains the regret scaling of the CwG policy under different distributions.

three point discrete distribution with support $\{0.25, 0.5, 0.75\}$ and the probability masses given as $\{0.5 - \epsilon_0/2, \epsilon_0, 0.5 - \epsilon_0/2\}$. We consider $\epsilon_0 = \{1/3, 1/12\}$ and observe that the regret is uniformly bounded in terms of the scaling parameters (B, T) and is inversely proportional to the minimum probability mass ϵ_0 , which is in line with Corollary 1. For the $\text{Uniform}([0, 1])$ distribution, we observe that the regret for the CwG policy scales as $\Theta(\log T)$ which matches Theorem 1. For the distribution $F_0 \sim \text{Uniform}([0, 1/4] \cup [3/4, 1])$, we verify that the regret scales as $\mathcal{O}((\log T)^2)$.

5. Analysis of our CwG Policy

This section provides a proof of our general achievability result, Theorem 5. (Theorem 4 is an immediate corollary of Theorem 5, obtained by setting $\delta = 0$.)

5.1. Outline

Recall $\tilde{T} = T - 16 \log(1/\epsilon_0)/\epsilon_0^2$ in Algorithm 2 (the CwG policy). Our algorithm operates in two phases, the first phase include time steps t such that $1 \leq t \leq \tilde{T}$, while the second phase consists of the remaining time steps t such that $\tilde{T} + 1 \leq t \leq T$. Since the second phase is easier to analyse, we will start with bounding the regret incurred in the second phase. Recall that the CwG policy (Algorithm 2) in the last $16 \log(1/\epsilon_0)/\epsilon_0^2$ time steps, makes use of the static allocation policy where we solve for the CE quantile threshold p_T^{ce} and thereafter use the time invariant quantile threshold p_T^{ce} . Using a well known fact in the network revenue management literature Cooper (2002), Gallego and Van Ryzin (1994), Reiman and Wang (2008), Bumpensanti and Wang (2020), we know that the regret accrued under a static allocation policy is upper bounded by $C\sqrt{\text{horizon length}}$, where $C < \infty$ is some universal constant. Since the CwG policy (Algorithm 2)

employs the static allocation policy for the last $16 \log(1/\varepsilon_0^2)/\varepsilon_0$ periods, we get the $C \sqrt{\log(1/\varepsilon_0)}/\varepsilon_0$ summand in the regret bound in Theorem 5.

The analysis of first phase makes use of the regret decomposition using the compensated coupling framework introduced by Vera and Banerjee (2021). Vera and Banerjee (2021) showed that the (expected) regret can be decomposed as the sum of (expected) compensations that need to be provided to the offline-to-go (OTG) policy so that it takes that same action as the online policy. To do this for our CwG policy, we analyze two thresholds: the CwG quantile threshold p_t^{cwg} and the OTG quantile threshold p_t^{otg} . We bound the expected compensation at time t for $1 \leq t \leq \tilde{T}$ and we do so by dividing the analysis into two cases: (a) $1 - B_t/\tau > 4\sqrt{\log \tau/\tau}$ and (b) $1 - B_t/\tau \leq 4\sqrt{\log \tau/\tau}$. At any time either one of the cases arises. We bound the expected compensation for each of the two cases. The analysis for both cases uses almost the same recipe. We show that with high probability the CwG quantile threshold p_t^{cwg} and OTG quantile threshold p_t^{otg} belong to the same cluster (Lemma 4) and with high probability the difference between p_t^{cwg} and p_t^{otg} is bounded above by a constant times $\sqrt{\log \tau/\tau}$ (Lemma 3). Compensation is needed at time t only if there is an arrival θ_t such that its quantile $F(\theta_t)$ lies between the two thresholds p_t^{cwg} and p_t^{otg} and the amount of compensation is bounded by $|F^{-1}(p_t^{\text{cwg}}) - F^{-1}(p_t^{\text{otg}})|$ (refer to Lemma 5). Using Lemma 3, 4 and 5, we show that the expected compensation in both the cases (a) and (b) is upper bounded by $\tilde{O}((\log \tau/\tau)^{\frac{1}{2} + \frac{1}{2(1+\beta)}})$ where $\tau = T - t$. Summing this expected compensation from $\tau = T$ to $\tau_0 = 16 \log(1/\varepsilon_0)/\varepsilon_0^2$, gives us that regret is bounded as $\tilde{O}(T^{\frac{1}{2} - \frac{1}{2(\beta+1)}})$, leading to Theorem 5.

5.2. Compensated Coupling and Regret Decomposition

To analyze the performance of the CwG policy that we devised, we will make use of the general regret decomposition provided in Lemma 1 of Vera and Banerjee (2021), which applies to any online policy. Denote a sample path of candidate ability quantiles as $\vec{\omega} = \{\omega_1, \omega_2, \dots, \omega_T\}$. The idea of compensated coupling framework introduced in Vera and Banerjee (2021) is to compare the OFFLINE and ONLINE policy acting on the same sample path, by compensating the OFFLINE policy whenever the action under the ONLINE policy differs from the optimal action under the OFFLINE policy just enough that the OFFLINE policy follows (takes the same action as) the ONLINE policy at each time step (this is the ‘‘coupling’’), and earns the same reward as it would have without the compensation. Let B_t denote the remaining budget under the CwG policy after time t and let $\partial R_t(B_{t-1}, \mu_t^{\text{cwg}})[\vec{\omega}]$ denote the minimum compensation that needs to be provided to the offline policy so that it takes the same action as the one taken by the online CwG policy at time t . Let $a_t^{\text{off}}[\vec{\omega}]$ be the set of actions which are optimal under the offline policy at time t . Define $Q_t(B_{t-1}) = \{\vec{\omega} : \mu_t^{\text{cwg}} \notin a_t^{\text{off}}[\vec{\omega}]\}$ to be the set of sample paths where the action of the OFFLINE and the ONLINE policy differ given that the remaining hiring budget at the end of time $t - 1$ is B_{t-1} .

Compensate the OFFLINE policy at time t only if $\mu_t^{\text{cwg}} \notin a_t^{\text{off}}[\vec{\omega}]$. We can decompose the regret under the CwG policy as follows.

LEMMA 1 (**Vera and Banerjee (2021)**). *For all $T \in \mathbb{N}$ and for all $B \in [T]$, consider the CwG policy and let B_t be the resultant hiring budget process under the CwG policy with $B_0 = B$, then*

$$\text{Regret}(B, T; \text{CwG}) = \sum_{t=1}^T \mathbb{E} [\partial R_t(B_{t-1}, \mu_t^{\text{cwg}})[\vec{\omega}] \cdot \mathbb{1}_{Q_t(B_{t-1})}[\vec{\omega}]] \quad (3)$$

We refer the readers to Vera and Banerjee (2021) for a proof of Lemma 1. See Appendix EC.3.1 for more details. We will use the notation $\mathcal{R}_t = \partial R_t(B_{t-1}, \mu_t^{\text{cwg}})[\vec{\omega}] \cdot \mathbb{1}_{Q_t(B_{t-1})}[\vec{\omega}]$ and drop the dependence on $\vec{\omega}$ for notational convenience.

5.3. OFFLINE-TO-GO Policy and OTG Threshold

Define $\vec{\omega}^{t+1} = \{\omega_{t+1}, \omega_{t+2}, \dots, \omega_T\}$. Let $\vec{\omega}^{t+1}(x)$ denote the x -th largest value quantile in $\vec{\omega}^{t+1}$ for integer x . Define $q_t^l \triangleq \vec{\omega}^{t+1}(B_t + 1)$, $q_t^u \triangleq \vec{\omega}^{t+1}(B_t)$. Denote the corresponding values by $l_t = F^{-1}(q_t^l)$, $u_t = F^{-1}(q_t^u)$. Since the principle of compensated coupling is to persuade the offline policy to take the same action as the online policy using (just) sufficient compensations, the offline policy at time t may look different from the offline policy initially. To distinguish between the two, at any time t , we will instead refer to the offline policy as the *Offline To Go* (OTG) policy, which due to coupling follows the same actions as the online policy up till time t , and then from time $t + 1$ onwards takes the optimal offline decision with arrivals in $\vec{\omega}^{t+1}$ given a remaining budget B_t where the budget B_t is determined using the online policy since up till t . Given the CwG quantile threshold p_t^{cwg} , we define $p_t^{\text{otg}} \triangleq \arg \max_{x \in [q_t^l, q_t^u]} |p_t^{\text{cwg}} - x|$ when $B_t > 0$ otherwise define $p_t^{\text{otg}} = 1$. The reason to adopt this particular p_t^{cwg} -dependent definition of p_t^{otg} is that the compensation needed at time step t will now be bounded above by the separation between the CwG threshold and the OTG threshold in value space (Lemma 5 below).

5.4. Preliminaries and Helper Lemmas

We begin by introducing some helper lemmas which we will use to prove the regret bound. We defer the proof of these lemmas to the electronic companion EC.3. Let t denote the number of time steps elapsed and $\tau \triangleq T - t$ denote the remaining number of times steps. Assume that $T \geq 16 \log(1/\varepsilon_0)/\varepsilon_0^2$ and define $\tilde{T} = T - \lfloor 16 \log(1/\varepsilon_0)/\varepsilon_0^2 \rfloor$. Define the following events for $t \leq \tilde{T}$:

$$A_{1,t} = \{|p_t^{\text{ce}} - p_t^{\text{otg}}| \leq \sqrt{2 \log \tau / \tau}\}, \quad (4)$$

$$A_{2,t} = \{|p_t^{\text{cwg}} - p_t^{\text{otg}}| \leq 3\sqrt{\log \tau / \tau}\}, \quad (5)$$

$$A_{3,t} = \cup_{i=1}^{n+1} \{p_t^{\text{cwg}} \in \bar{Q}_i, p_t^{\text{otg}} \in \bar{Q}_i\}, \quad (6)$$

where $\bar{Q}_i = [q_{i-1}^*, q_i^*]$ and n denotes the number of gaps. The interpretation of $A_{3,t}$ is that the CwG policy threshold and the OTG policy threshold belong (weakly) to the same mass cluster. The following lemmas show that these three events are very likely to occur:

LEMMA 2. For any $t \leq \tilde{T}$, consider the event $A_{1,t}$ defined in (4). We have that $\mathbb{P}(A_{1,t}^c) \leq 2/\tau^4$.

LEMMA 3. For any $t \leq \tilde{T}$, consider the event $A_{2,t}$ defined in (5). We have that $\mathbb{P}(A_{2,t}^c) \leq 2/\tau^4$.

LEMMA 4. For any $t \leq \tilde{T}$, consider the event $A_{3,t}$ defined in (6). We have that $\mathbb{P}(A_{3,t}^c) \leq 2n(n+1)/\tau^4$, where n is the number of gaps.

Let $q_t^\theta = F(\theta_t)$ be the quantile of the candidate ability θ_t at time t . If $p_{t-1}^{\text{cwg}} < q_{t-1}^l$ then we have that $p_{t-1}^{\text{otg}} = q_{t-1}^u$ and compensation is needed only if $q_t^\theta \in [p_{t-1}^{\text{cwg}}, p_{t-1}^{\text{otg}}]$. If $p_{t-1}^{\text{cwg}} > q_{t-1}^u$ then we have that $p_{t-1}^{\text{otg}} = q_{t-1}^l$ and compensation is needed only if $q_t^\theta \in [p_{t-1}^{\text{otg}}, p_{t-1}^{\text{cwg}}]$. If $p_{t-1}^{\text{cwg}} \in (q_{t-1}^l, q_{t-1}^u)$, then no compensation is required.

LEMMA 5. Let $q_t^\theta = F(\theta_t)$ denote the quantile corresponding to θ_t . Compensation needs to be provided only if $q_t^\theta \in (\min\{p_{t-1}^{\text{cwg}}, p_{t-1}^{\text{otg}}\}, \max\{p_{t-1}^{\text{cwg}}, p_{t-1}^{\text{otg}}\})$; let $\partial R_t(B_{t-1}, \mu_t^{\text{cwg}})$ denote the compensation. Then we have that $\partial R_t(B_{t-1}, \mu_t^{\text{cwg}}) \leq \max\{F^{-1}(p_{t-1}^{\text{cwg}}) - F^{-1}(p_{t-1}^{\text{otg}}), F^{-1}(p_{t-1}^{\text{otg}}) - F^{-1}((p_{t-1}^{\text{cwg}})^+)\}$.

5.5. Proof of Theorem 5

Define $\tilde{T} = T - 16\log(1/\varepsilon_0)/\varepsilon_0^2$ and define $\tau_0 = 16\log(1/\varepsilon_0)/\varepsilon_0^2$. Consider some time $t \leq \tilde{T}$ and let $\tau = T - t$ denote the time to go. Recall that $p_t^{\text{cwg}} \in \mathcal{F}_t$. Now p_t^{otg} depends on the candidate abilities $\{\omega_{t+1}, \dots, \omega_T\}$, but only via the B_t -th largest quantile q_t^u and $(B_t + 1)$ -th largest quantile is q_t^l . To facilitate our analysis, we employ the so-called principle of deferred decisions, and initially only reveal q_t^u and q_t^l (in addition to the history up to time t), which is enough to uniquely determine p_t^{otg} . Given this information, we know that the periods to go include a random subset of $B_t - 1$ quantiles located above q_t^u (these quantiles are i.i.d. uniform in $[q_t^u, 1]$), and the remaining $\tau - B_t - 1$ quantiles to go are below q_t^l (these quantiles are i.i.d. uniform in $[0, q_t^l]$). If $p_t^{\text{cwg}} \in (q_t^l, q_t^u)$, then no compensation is required. Let us bound the contribution to the regret (compensation at time $t + 1$) of sample paths with $p_t^{\text{cwg}} \leq q_t^l$; we will use the notation $\tilde{\mathcal{R}}_{t+1}$ for this regret. (A similar bound can be shown analogously for the contribution to the regret of sample paths satisfying the symmetric counterpart condition $p_t^{\text{cwg}} \geq q_t^u$; we omit the details for the sake of brevity.) Let q_{t+1}^θ be the candidate ability quantile at time $t + 1$ (the candidate ability is $\theta_{t+1} = F^{-1}(q_{t+1}^\theta)$). Compensation is needed only if $q_{t+1}^\theta \in [p_t^{\text{cwg}}, q_t^u]$. This event occurs if the realized quantile q_{t+1}^θ is $q_{t+1}^\theta = q_t^l$ or $q_{t+1}^\theta \in [p_t^{\text{cwg}}, q_t^l]$ is one of the $\tau - B_t - 1$ lower quantiles (call this event E_t). The probability of the event E_t is given as $\mathbb{P}(E_t) = \mathbb{I}_{\{p_t^{\text{cwg}} \leq q_t^l\}}/\tau + (\tau - B_t - 1)(q_t^l - p_t^{\text{cwg}})_+/(q_t^l\tau)$. From Lemma 5, it follows that the compensation under the event E_t is bounded above by $F^{-1}(q_t^u) - F^{-1}((p_t^{\text{cwg}})^+)$. Hence the expected compensation at $t + 1$ conditioned on $\mathcal{F}_t, q_t^l, q_t^u$ is bounded above as

$$\mathbb{E} \left[\tilde{\mathcal{R}}_{t+1} | \mathcal{F}_t, q_t^l, q_t^u \right] \leq \frac{|F^{-1}(q_t^u) - F^{-1}((p_t^{\text{cwg}})^+)|}{\tau} + \frac{(q_t^l - p_t^{\text{cwg}})|F^{-1}(q_t^u) - F^{-1}((p_t^{\text{cwg}})^+)|(\tau - B_t - 1)}{q_t^l\tau}. \quad (7)$$

Next we need to bound the ratio $(\tau - B_t - 1)/(q_t^l \tau)$. Now at any time $t \leq T - \lfloor \tau_0 \rfloor$, exactly one of the following two complementary cases occurs: (a) $1 - B_t/\tau > 4\sqrt{\log \tau/\tau}$ or (b) $1 - B_t/\tau \leq 4\sqrt{\log \tau/\tau}$, where recall $\tau = T - t$. Recall the event $A_{3,t}$ defined in (6), which is identical to

$$A_{3,t} = \{q_t^l, q_t^u, p_t^{\text{cwg}} \text{ are quantiles belonging (weakly) to the same cluster}\}.$$

Next, we will establish an upper bound on (7) for each of the cases (a) and (b).

Case (a): $1 - B_t/\tau > 4\sqrt{\log \tau/\tau}$ Define the following events:

$$A_{4,t} \triangleq \{q_t^l \geq (1/2)(1 - B_t/\tau)\},$$

$$A_{5,t} \triangleq A_{3,t} \cap A_{4,t}.$$

Under the event $A_{3,t}$, from Definition 2 (a) it follows that $|F^{-1}(q_t^u) - F^{-1}((p_t^{\text{cwg}})^+)| \leq |q_t^u - p_t^{\text{cwg}}|^{\frac{1}{1+\beta}} + \delta$. Now, on the event $A_{4,t}$, we have that $(\tau - B_t - 1)/(\tau q_t^l) \leq 2$. We have that

$$\begin{aligned} \mathbb{P}(A_{4,t}^c | B_t, \text{Case (a)}) &= \mathbb{P}(q_t^l < (1/2)(1 - B_t/\tau)) \leq \mathbb{P}(\text{Binomial}(\tau, (1/2)(1 - B_t/\tau)^-) \geq \tau - B_t - 1) \\ &\leq \exp(-\Omega(\tau - B_t)) \leq C/(\tau - B_t)^4 \leq C/\tau^2. \end{aligned}$$

where the last inequality follows from the case assumption that $\tau - B_t \geq 4\sqrt{\tau \log \tau}$ and the inequality is true for some appropriately defined constant $C < \infty$. It follows that

$$\mathbb{P}(A_{4,t}^c | \text{Case (a)}) \leq C/\tau^2. \quad (8)$$

Using (7), and the definitions of the events $A_{3,t}$ and $A_{4,t}$, we have that

$$\mathbb{E}[\tilde{\mathcal{R}}_{t+1} | \mathcal{F}_t, q_t^l, q_t^u] \leq \mathbb{I}_{A_{3,t}} \cdot [|q_t^u - p_t^{\text{cwg}}|^{\frac{1}{1+\beta}} + \delta] / \tau + 2\mathbb{I}_{A_{3,t}} \mathbb{I}_{A_{4,t}} [|q_t^u - p_t^{\text{cwg}}|^{1+\frac{1}{1+\beta}} + |q_t^u - p_t^{\text{cwg}}| \delta] \quad (9)$$

$$\begin{aligned} &+ \mathbb{I}_{A_{3,t}^c} + \mathbb{I}_{A_{4,t}^c}, \\ &\leq |q_t^u - p_t^{\text{cwg}}|^{\frac{1}{1+\beta}} / \tau + \delta / \tau + 2|q_t^u - p_t^{\text{cwg}}|^{1+\frac{1}{1+\beta}} + 2|q_t^u - p_t^{\text{cwg}}| \delta + \mathbb{I}_{A_{4,t}^c} + \mathbb{I}_{A_{3,t}^c}, \quad (10) \end{aligned}$$

where the first inequality follows from $q_t^l - p_t^{\text{cwg}} \leq q_t^u - p_t^{\text{cwg}}$, and the second inequality follows from the fact that $\mathbb{I}_{A_{3,t}}, \mathbb{I}_{A_{4,t}} \leq 1$. Using the definition of the event $A_{2,t}$ in (5) and Lemma 3, we have that for all $\alpha \in [1, 2]$, we have

$$\mathbb{E}[|q_t^u - p_t^{\text{cwg}}|^\alpha] \leq \mathbb{E}[\mathbb{I}_{A_{2,t}} |q_t^u - p_t^{\text{cwg}}|^\alpha + \mathbb{I}_{A_{2,t}^c}] \leq 3^\alpha (\log \tau / \tau)^{\alpha/2} + 2/\tau^4 \leq 2 \cdot 3^\alpha (\log \tau / \tau)^{\alpha/2}. \quad (11)$$

Taking expectations on both sides of (10), we obtain that

$$\begin{aligned} \mathbb{E}[\tilde{\mathcal{R}}_{t+1} | \text{case(a)}] &\stackrel{(i)}{\leq} \mathbb{E} \left[|q_t^u - p_t^{\text{cwg}}|^{\frac{1}{\beta+1}} \right] / \tau + \delta / \tau + 2\mathbb{E} \left[|q_t^u - p_t^{\text{cwg}}|^{1+\frac{1}{\beta+1}} \right] + 2\mathbb{E} [|q_t^u - p_t^{\text{cwg}}|] \delta \\ &+ \mathbb{P}(A_{3,t}^c | \text{case(a)}) + \mathbb{P}(A_{4,t}^c | \text{case(a)}), \\ &\stackrel{(ii)}{\leq} 6 (\log \tau / \tau)^{\frac{1}{2(\beta+1)}} / \tau + \delta / \tau + 36 (\log \tau / \tau)^{\frac{1}{2} + \frac{1}{2(\beta+1)}} + 6\delta \sqrt{\log \tau / \tau} + \mathbb{P}(A_{3,t}^c | \text{case(a)}) + C/\tau^2, \quad (12) \end{aligned}$$

where inequality (i) follows from the taking expectation on both sides, and inequality (ii) follows from using (11) for the first, third and fourth summands, and the sixth summand follows from (8).

Case (b): $1 - B_t/\tau \leq 4\sqrt{\log \tau/\tau}$ The case assumption $1 - B_t/\tau \leq 4\sqrt{\log \tau/\tau}$ implies that $(\tau - B_t - 1)/\tau \leq 4\sqrt{\log \tau/\tau}$, and obviously we have $(q_t^l - p_t^{\text{cwg}})/q_t^l \leq 1$. Therefore the second term in the RHS of (7) is bounded above as

$$|q_t^l - p_t^{\text{cwg}}| |F^{-1}(q_t^u) - F^{-1}((p_t^{\text{cwg}})^+)| (\tau - B_t - 1) / (q_t^l \tau) \leq 4\sqrt{\log \tau/\tau} |F^{-1}(q_t^u) - F^{-1}((p_t^{\text{cwg}})^+)|$$

Therefore we can upper bound $\mathbb{E}[\tilde{\mathcal{R}}_{t+1} | \mathcal{F}_t, q_t^l, q_t^u, \text{case (b)}]$ as

$$\begin{aligned} & \mathbb{E}[\tilde{\mathcal{R}}_{t+1} | \mathcal{F}_t, q_t^l, q_t^u, \text{case (b)}] \\ & \leq \mathbb{I}_{A_{3,t}} \left[|q_t^u - p_t^{\text{cwg}}|^{\frac{1}{1+\beta}} / \tau + \delta / \tau \right] + \mathbb{I}_{A_{3,t}} \left[4\sqrt{\log \tau/\tau} |q_t^u - p_t^{\text{cwg}}|^{\frac{1}{\beta+1}} + 4\sqrt{\log \tau/\tau} \delta \right] + \mathbb{I}_{A_{3,t}^c}, \\ & \leq |q_t^u - p_t^{\text{cwg}}|^{\frac{1}{1+\beta}} / \tau + \delta / \tau + 4\sqrt{\log \tau/\tau} |q_t^u - p_t^{\text{cwg}}|^{\frac{1}{\beta+1}} + 4\delta\sqrt{\log \tau/\tau} + \mathbb{I}_{A_{3,t}^c}, \end{aligned}$$

Taking expectations on both sides we get that

$$\begin{aligned} & \mathbb{E}[\tilde{\mathcal{R}}_{t+1} | \text{Case (b)}] \\ & \leq \mathbb{E} \left[|q_t^u - p_t^{\text{cwg}}|^{\frac{1}{1+\beta}} / \tau + \delta / \tau + 4\sqrt{\log \tau/\tau} \mathbb{E} \left[|q_t^u - p_t^{\text{cwg}}|^{\frac{1}{1+\beta}} \right] + 4\delta\sqrt{\log \tau/\tau} + \mathbb{P}(A_{3,t}^c | \text{Case (b)}) \right], \\ & \leq 6(\log \tau/\tau)^{\frac{1}{2(1+\beta)}} / \tau + \delta / \tau + 24(\log \tau/\tau)^{\frac{1}{2} + \frac{1}{2(1+\beta)}} + 4\delta\sqrt{\log \tau/\tau} + \mathbb{P}(A_{3,t}^c | \text{Case (b)}), \end{aligned} \quad (13)$$

where the second inequality follows from the fact that the first and the second term are bounded by (11). This completes our analysis of case (b).

From Lemma 4, it follows that $\mathbb{P}(A_{3,t}^c) \leq 2n(n+1)/\tau^4$. We have that

$$\sum_{t=1}^{T-\lfloor \tau_0 \rfloor} \mathbb{P}(A_{3,t}^c) \leq \sum_{\tau=\tau_0}^T 2n(n+1)/\tau^4 \leq n(n+1)/\tau_0^3 \leq n(n+1)\varepsilon_0^6 \stackrel{(\star)}{\leq} 1 \quad (14)$$

where (\star) follows since if there are n gaps, there are $n+1$ clusters, and hence $\varepsilon_0 \leq 1/(n+1)$.

Fix $t \leq T - \lfloor \tau_0 \rfloor$. We now take the expectation over the case which arises (case (a) or case (b)). Combining (12) and (13), for a constant $C < \infty$ we have that

$$\mathbb{E}[\tilde{\mathcal{R}}_{t+1}] \leq C(\log \tau/\tau)^{\frac{1}{2} + \frac{1}{2(\beta+1)}} + C\delta\sqrt{\log \tau/\tau} + \mathbb{P}(A_{3,t}^c).$$

An identical bound holds for the regret contribution from sample paths where $p_t^{\text{cwg}} \geq q_t^u$, by a symmetric argument. As a result, we can bound the expected total regret at time $t+1$ as per $\mathbb{E}[\mathcal{R}_{t+1}] \leq 2\mathbb{E}[\tilde{\mathcal{R}}_{t+1}]$. Now summing this bound from $\tau = T$ to $\tau = \tau_0$, we have, using (14) that

$$\sum_{t=0}^{T-\lfloor \tau_0 \rfloor} \mathbb{E}[\mathcal{R}_{t+1}] \leq C' \left[(1 + 1/\beta)(\log T)^{\frac{1}{2} + \frac{1}{2(\beta+1)}} T^{\frac{1}{2} - \frac{1}{2(1+\beta)}} \cdot \mathbb{1}\{\beta > 0\} + (\log T)^2 + \delta\sqrt{T \log T} \right].$$

Finally, consider time steps t such that $\tilde{T} + 1 \leq t \leq T$. In the last $16 \log(1/\varepsilon_0)/\varepsilon_0^2$ time steps, we make use of the static allocation policy and as noted before the regret accrued during the static allocation policy is upper bounded by $C\sqrt{\tau_0} = C\sqrt{\log(1/\varepsilon_0)}/\varepsilon_0$ for some universal constant $C < \infty$. Combining the two parts completes the proof. \square

5.6. Proof of Corollary 1

The discrete distribution considered is a $(\beta = 0, \varepsilon_0, \delta = 0)$ -clustered distribution for $\varepsilon_0 = \min_{1 \leq i \leq m} \{f_i\}$. As done for the general case above, our analysis for the case of discrete distributions as considered in the Example 2 also follows in two parts. The regret accrued during the second part due to the static allocation policy is upper bounded by $C\sqrt{\log(1/\varepsilon_0)}/\varepsilon_0$ for some universal constant $C < \infty$. Next we will consider the first part. The argument for the first part will mirror the analysis presented in the proof of Theorem 5 except for one important improvement we make for this special case. Consider the regret contribution of sample paths satisfying $p_t^{\text{cwg}} \leq q_t^l$ as we did previously. (Again, there is an analogous analysis for the symmetric condition $p_t^{\text{cwg}} > q_t^u$, which we omit to avoid repetition.) The only but important distinction in the case of discrete distributions is that on the event $A_{3,t}$, which is that q_t^l, q_t^u and p_t^{cwg} are quantiles belonging to the same cluster, the compensation is given as $F^{-1}(q_t^u) - F^{-1}((p_t^{\text{cwg}})^+)$, however for discrete distributions, we have that $F^{-1}(q_t^u) - F^{-1}((p_t^{\text{cwg}})^+) = 0$. Previously, in the general case, we had upper bounded $F^{-1}(q_t^u) - F^{-1}((p_t^{\text{cwg}})^+)$ by $|q_t^u - p_t^{\text{cwg}}|^{1/(1+\beta)} + \delta$ using Definition 2. Because of $F^{-1}(q_t^u) - F^{-1}((p_t^{\text{cwg}})^+) = 0$, we have that the expected compensation $\mathbb{E}[\mathcal{R}_t]$ is now bounded above by $\mathbb{P}(A_{3,t}^c) + C/\tau^2$. Summing this upper bound from $\tau = \tau_0$ to $\tau = T$, we get that the summation is upper bounded by a universal constant $C < \infty$ using (14). Combining the regret accrued in the two parts, we get the required result. \square

6. Conclusion

In this work, we consider the multi-secretary problem with many types. In particular we generalize the setting studied in Arlotto and Gurvich (2019), Bray (2019) to a large class of distributions namely (β, ε_0) -clustered distributions with gaps. The gaps are well motivated in practice, not only in multi-secretary problems but also in more general network revenue management (NRM) problems like online matching or order fulfillment problems. Arlotto and Gurvich (2019) in their concluding remarks had posed the question of whether bounded regret is achievable also with continuous distributions. We not only answer this question in the negative but also characterize how the regret scales as a function of the spatial accumulation of mass around gaps quantified by β and the minimum probability mass ε_0 .

While we study the multi-secretary problem, which is a special one-dimensional case of the general NRM problem, we believe that this is an important step and our idea of *Conservativeness with respect to gaps* can be extended to a more general class of NRM problems with many types, with some effort. One such interesting application area to explore is the order fulfillment problem with finite number of fulfillment centers and demand (types) spread over some multi-dimensional space with regions of gaps in space.

Finally, another possible direction to explore is the setting where the candidate ability distribution is unknown to the decision maker. For the multi-secretary problem with finite types, Arlotto and Gurvich (2019) conjecture that the regret should scale as $\log T$ and this is achieved using an algorithm which uses the first $\mathcal{O}(\log T)$ steps to learn the distribution and then computes the threshold and use the Budget-Ratio policy with the learnt distribution. A similar idea can be applied to continuous distributions with no gaps and the density being positive everywhere, since the problem there can be to learn the value of the order statistics and use them as thresholds to make hire/reject decisions. However in the case where the distribution has gaps in the type space, such a result does not apply and hence an open question is how the regret scales for unknown distributions with gaps.

References

- Arlotto A, Gurvich I (2019) Uniformly bounded regret in the multiselection problem. *Stochastic Systems* 9(3):231–260.
- Balseiro S, Besbes O, Pizarro D (2021) Survey of dynamic resource constrained reward collection problems: Unified model and analysis. *Available at SSRN 3963265* .
- Bray R (2019) Does the multiselection problem always have bounded regret? *Available at SSRN 3497056* .
- Bumpensanti P, Wang H (2020) A re-solving heuristic with uniformly bounded loss for network revenue management. *Management Science* 66(7):2993–3009.
- Cayley A (1875) Mathematical questions with their solutions. *The Educational Times* 23:18–19.
- Cooper WL (2002) Asymptotic behavior of an allocation policy for revenue management. *Operations Research* 50(4):720–727.
- Ferguson TS (1989) Who solved the secretary problem? *Statistical science* 4(3):282–289.
- Freeman P (1983) The secretary problem and its extensions: A review. *International Statistical Review/Revue Internationale de Statistique* 189–206.
- Gallego G, Van Ryzin G (1994) Optimal dynamic pricing of inventories with stochastic demand over finite horizons. *Management science* 40(8):999–1020.
- Jasin S, Sinha A (2015) An lp-based correlated rounding scheme for multi-item ecommerce order fulfillment. *Operations Research* 63(6):1336–1351.
- Kleinberg R (2005) A multiple-choice secretary algorithm with applications to online auctions. *Proceedings of the sixteenth annual ACM-SIAM symposium on Discrete algorithms*, 630–631 (Citeseer).
- Kleywegt AJ, Papastavrou JD (1998) The dynamic and stochastic knapsack problem. *Operations research* 46(1):17–35.
- Moser L (1956) On a problem of Cayley. *Scripta Math* 22:289–292.

- Reiman MI, Wang Q (2008) An asymptotically optimal policy for a quantity-based network revenue management problem. *Mathematics of Operations Research* 33(2):257–282.
- Talluri KT, Van Ryzin G, Van Ryzin G (2004) *The theory and practice of revenue management*, volume 1 (Springer).
- Vera A, Arlotto A, Gurvich I, Levin E (2020) Dynamic resource allocation: The geometry and robustness of constant regret. Technical report, Working paper.
- Vera A, Banerjee S (2021) The bayesian prophet: A low-regret framework for online decision making. *Management Science* 67(3):1368–1391.
- Vera A, Banerjee S, Gurvich I (2021) Online allocation and pricing: Constant regret via bellman inequalities. *Operations Research* .

Electronic Companion

EC.1. Proof of Theorem 2

The following lemma justifies a claim we made in the argument in the main paper.

LEMMA EC.1. *The threshold quantile under the CE policy $(p_t^{\text{ce}})_{t=0}^{T-1}$ is a martingale with respect to the filtration $(\mathcal{F}_t)_{t=0}^{T-1}$.*

Proof of Lemma EC.1 Without loss of generality, let us assume that $\mathbb{P}([0, c]) = \varepsilon$. Recall that $p_t^{\text{ce}} = 1 - B_t/(T-t)$ is the certainty equivalent quantile threshold defined in Algorithm 1 and define $\Delta_t = p_t^{\text{ce}} - p_{t-1}^{\text{ce}}, \forall t \geq 1$. We have that $\Delta_t = X_t - X_{t-1} = p_t^{\text{ce}} - p_{t-1}^{\text{ce}}$. At time t , the candidate is hired with probability $1 - p_{t-1}^{\text{ce}}$ and rejected with probability p_{t-1}^{ce} and hence with probability $1 - p_{t-1}^{\text{ce}}$, the budget $B_t = B_{t-1} - 1$ and with probability p_{t-1}^{ce} , the budget $B_t = B_{t-1}$. (We use here the fact that F is assumed to be non-atomic, and so tie-breaking is not needed with probability 1.) Therefore we write

$$\Delta_t = \begin{cases} \frac{B_{t-1}}{T-(t-1)} - \frac{B_{t-1}-1}{T-t} = -\frac{B_{t-1}}{(T-t)(T-t+1)} + \frac{1}{T-t}, & \text{w.p. } 1 - p_{t-1}^{\text{ce}}, \\ \frac{B_{t-1}}{T-(t-1)} - \frac{B_{t-1}}{T-t} = -\frac{B_{t-1}}{(T-t)(T-t+1)}, & \text{w.p. } p_{t-1}^{\text{ce}}. \end{cases}$$

Computing the conditional expectation of Δ_t , we get that

$$\begin{aligned} \mathbb{E}[\Delta_t | \mathcal{F}_{t-1}] &= \left(\frac{B_{t-1}}{T-(t-1)} - \frac{B_{t-1}-1}{T-t} \right) (1 - p_{t-1}^{\text{ce}}) + \left(1 - \frac{B_{t-1}}{T-t} - p_{t-1}^{\text{ce}} \right) p_{t-1}^{\text{ce}} \\ &= \frac{1 - p_{t-1}^{\text{ce}}}{T-t} + \frac{B_{t-1}}{T-t+1} - \frac{B_{t-1}}{T-t} \\ &= \frac{B_{t-1}}{(T-t+1)(T-t)} - \frac{B_{t-1}}{(T-t+1)(T-t)} = 0 \end{aligned}$$

Therefore we have that $(p_t^{\text{ce}})_{t=0}^{T-1}$ is a martingale with respect to the filtration $(\mathcal{F}_t)_{t=0}^{T-1}$. \square

Proof of Theorem 2 The argument provided in the main paper for the specific distribution F_0 generalizes to arbitrary non-atomic F with a gap of size $\eta > 0$ and probability mass at least $\varepsilon > 0$ on each side of the gap as follows:

- Suppose the gap is located at quantile $q \in [\varepsilon, 1 - \varepsilon]$.
- We consider the case $B = qT$.
- We employ the same argument with the generalized definitions of the events as follows

$$\begin{aligned} E_{t,1} &\triangleq \{p_t^{\text{ce}} < q - 1/\sqrt{T}\} \\ E_{t,2} &\triangleq \{p_t^{\text{otg}} > q\} \\ E_t &\triangleq E_{t,1} \cap E_{t,2} \\ E_{t,3} &\triangleq \{\theta_t \in [F^{-1}(q - 1/\sqrt{T}), F^{-1}(q)]\}. \end{aligned}$$

Here the offline (hindsight) threshold quantile p_t^{otg} is taken to be *any* threshold quantile that implements the offline optimal solution, given the history under CE up to period t .

- Our argument again establishes that for each $t \in [T/3, 2T/3]$ we have $\mathbb{P}(E_t \cap E_{t,3}) \geq \nu/\sqrt{T}$ for some $\nu = \nu(\varepsilon) > 0$. Lemma EC.1 formalizes the martingale property. Hence, to make the argument formal, we have only to establish that the martingale steps are of size $\Theta(1/T)$ for $t \leq 2T/3$, which we do now. Note from the the proof of Lemma EC.1 that

$$|\Delta_t| \leq 1/(T-t) \leq 3/T \quad \text{for } t \leq 2T/3, \quad (\text{EC.1})$$

i.e., we have shown the required upper bound on the martingale step size. We now establish the lower bound on the step size. Using the Azuma-Hoeffding inequality, since the step size is bounded above by $3/T$, w.p. $1 - \exp(-\Omega(T))$, $|p_t^{\text{ce}} - p_0^{\text{ce}}| = |p_t^{\text{ce}} - (1-q)| \leq \varepsilon/2$. It follows that, whp, for $t \leq 2T/3$, we have $|p_t^{\text{ce}} - (1-q)| \leq \varepsilon/2 \Rightarrow p_t^{\text{ce}} \in [\varepsilon/2, 1 - \varepsilon/2] \Leftrightarrow B_t/(T-t) \in [\varepsilon/2, 1 - \varepsilon/2]$, and hence the martingale step size is bounded below as $|\Delta_t| \geq \min(-\frac{B_{t-1}}{(T-t)(T-t+1)} + \frac{1}{T-t}, \frac{B_{t-1}}{(T-t)(T-t+1)}) \geq \varepsilon/(T-t) \geq \varepsilon/T$ for all $t \leq 2T/3$.

- Each mistake $E_t \cap E_{t,3}$ adds regret at least as large as the gap size $\eta > 0$.
- Summing over $t \in [T/3, 2T/3]$, we deduce that CE suffers from expected regret at least $\eta\nu\sqrt{T}/3$.

This completes the proof. \square

EC.2. Proof of Theorem 3

Proof of Theorem 3. For a given T large enough, it suffices to choose a value of B and a distribution $F \in \mathcal{F}_{\beta, \varepsilon_0}$ such that the regret of the optimal online policy is lower bounded by $CT^{\frac{1}{2} - \frac{1}{2(\beta+1)}} \mathbb{1}\{\beta > 0\} + C \log T \mathbb{1}\{\beta = 0\}$ for appropriately chosen constant $C > 0$.

We will first consider the case of $\beta = 0$. We consider the distribution F to be the uniform distribution over $[0, 1]$. We can verify that for the uniform distribution, $\beta = 0$ and we have that $\text{Uniform}([0, 1]) \in \mathcal{F}_{\beta, \varepsilon_0}$ for all $0 < \varepsilon_0 \leq 1/2$. Using Bray (2019), and in particular Theorem 1, we know that $\text{Regret}(B, T) \geq C \log T$ for all $T \geq T_0$ and $C > 0$ for $B = \lceil T/2 \rceil$ and the uniform distribution ($\beta = 0$).

Next we will focus on the case when $\beta > 0$. For the lower bound, we will consider the (β, ε_0) -clustered distributions F_β parameterized by β and $\varepsilon_0 \leq 1/2$ given in (1). Using F_β^{-1} defined in (2), we can easily verify that F_β is a (β, ε_0) -clustered distribution for $0 < \varepsilon_0 \leq 1/2$. Again we assume that $B = \lceil T/2 \rceil$. Next we will setup some notation. The support of the distribution F_β is $[0, 1/4] \cup [3/4, 1]$. Define $\Delta_\beta \triangleq T^{-\frac{1}{2(\beta+1)}}/4$. We will divide this support into the following intervals: $I_1 = [0, 1/4 - R_1\Delta_\beta]$, $I_{2a} = [1/4 - R_1\Delta_\beta, 1/4 - \Delta_\beta]$, $I_{3a} = [1/4 - \Delta_\beta, 1/4]$, $I_{3b} = [3/4, 3/4 + \Delta_\beta]$, $I_{2b} = [3/4 + \Delta_\beta, 3/4 + R_2\Delta_\beta]$, $I_4 = [3/4 + R_2\Delta_\beta, 1]$. Due to the choice of Δ_β , we have that measure μ_β on the interval I_{3a}, I_{3b} is $\mu_\beta(I_{3a}) = \mu_\beta(I_{3b}) = T^{-1/2}/2$. The values of R_1 and R_2 are chosen in such a way that measure μ_β of the interval I_{2a} and I_{2b} is $\mu_\beta(I_{2a}) = T^{-1/2}/128$

and $\mu_\beta(I_{2b}) = T^{-1/2}/128$. Note that R_1, R_2 are only functions of β and do not depend on T and $R_1, R_2 > 1$. Define $I_2 = I_{2,a} \cup I_{2,b}, I_3 = I_{3,a} \cup I_{3,b}$.

Let $Z_i^{t_1:t_2}$ denote the number of candidates arrivals with their abilities in the interval I_i in the time periods $\{t_1, t_1 + 1, \dots, t_2\}$. Consider the following events:

$$\begin{aligned} L_T &= \left\{ T/4 - \sqrt{T}/2 \leq Z_4^{1:\lfloor T/2 \rfloor} \leq T/4 \right\} \cap \left\{ T/4 - 4\sqrt{T} \leq Z_4^{\lfloor T/2 \rfloor + 1:T} \leq T/4 - 3\sqrt{T} \right\}, \\ H_T &= \left\{ T/4 - \sqrt{T}/2 \leq Z_4^{1:\lfloor T/2 \rfloor} \leq T/4 \right\} \cap \left\{ T/4 + \sqrt{T}/2 \leq Z_4^{\lfloor T/2 \rfloor + 1:T} \right\}, \\ A_T &= \left\{ \sqrt{T}/2 \leq Z_3^T \leq \min\{3Z_3^{T/2}, 2\sqrt{T}\} \right\} = \left\{ (1/2)\mathbb{E}[Z_3^T] \leq Z_3^T \leq \min\{3Z_3^{T/2}, 2\mathbb{E}[Z_3^T]\} \right\}, \\ M_T &= \left\{ \sqrt{T}/128 \leq Z_2^T \leq \sqrt{T}/32 \right\} = \left\{ (1/2)\mathbb{E}[Z_2^T] \leq Z_2^T \leq 2\mathbb{E}[Z_2^T] \right\}. \end{aligned}$$

On the event $H_T \cap A_T \cap M_T$, we have that $Z_4^{1:T} \geq T/2 \geq B$ and hence the offline policy would hire all the candidates with ability in $[3/4 + \Delta, 1]$ and reject all the candidates with ability in $I_1 \cup I_2 \cup I_3 \cup [\frac{3}{4} + R_2\Delta_\beta, \frac{3}{4} + \Delta]$, for some $\Delta \geq R_2\Delta_\beta$. On the flip side, on the event $L_T \cap A_T \cap M_T$, we have that $Z_4^{1:T} \leq T/2 - 3\sqrt{T}$ and hence the offline policy hires all the candidates with abilities in the interval I_4, I_3, I_2 and a subset of the interval I_1 given by $[1/4 - \Delta, 1/4 - R_1\Delta_\beta]$ to exhaust the budget B , where $\Delta \geq R_1\Delta_\beta$.

Next we want to show that the events $H_T \cap A_T \cap M_T$ and $L_T \cap A_T \cap M_T$ are non-trivial events i.e, there exists positive constants α and T_0 such that $\mathbb{P}(H_T \cap A_T \cap M_T), \mathbb{P}(L_T \cap A_T \cap M_T) \geq \alpha > 0$ for all $T \geq T_0$. To do this, we will make use of some lemmas stated below. The proofs of the following lemma essentially follow from an application of the Berry-Esseen theorem and hence are omitted.

LEMMA EC.2. $\exists 0 < \alpha_l, T_l < \infty$ such that for all $T \geq T_l$, we have that $\mathbb{P}(L_T) \geq \alpha_l$.

LEMMA EC.3. $\exists 0 < \alpha_h, T_h < \infty$ such that for all $T \geq T_h$, we have that $\mathbb{P}(H_T) \geq \alpha_h$.

LEMMA EC.4. Let $\alpha \triangleq \min\{\alpha_h, \alpha_l\}$. $\exists T_a < \infty$ such that for all $T \geq T_a$, we have $\mathbb{P}(A_T^c) \leq \alpha/4$.

LEMMA EC.5. Let $\alpha \triangleq \min\{\alpha_h, \alpha_l\}$. $\exists T_m < \infty$ such that for all $T \geq T_m$, we have $\mathbb{P}(M_T^c) \leq \alpha/4$.

Define $T_0 \triangleq \max\{T_l, T_h, T_a, T_m\}$ and we have that for all $T \geq T_0$,

$$\begin{aligned} \mathbb{P}(L_T \cap A_T \cap M_T) &= \mathbb{P}(A_T \cap M_T) \mathbb{P}(L_T | A_T \cap M_T) \\ &= \mathbb{P}(L_T) - \mathbb{P}(L_T | (A_T \cap M_T)^c) \mathbb{P}(A_T^c \cup M_T^c) \\ &\geq \mathbb{P}(L_T) - (\mathbb{P}(A_T^c) + \mathbb{P}(M_T^c)) \geq \alpha/2, \end{aligned}$$

where the last inequality follows from Lemmas EC.2, EC.4 and EC.5. Similarly, we have that for all $T \geq T_0$, $\mathbb{P}(H_T \cap A_T \cap M_T) \geq \alpha/2$.

Consider the random variable $\Lambda(B, T) = \sum_{t=1}^T \theta_t \mu_t^{\text{off}} - \sum_{t=1}^T \theta_t \mu_t^{\pi^*}$ that tracks the difference between the performance of the offline policy and the performance of any online policy $\pi^* \in \Pi(B, T)$.

We have that $\text{Regret}(B, T) = \mathbb{E}[\Lambda(B, T)]$. Let $S_3^{\star, \lfloor T/2 \rfloor}$ denote the number of candidates with ability in the interval I_3 up until time $\lfloor T/2 \rfloor$ that were hired under the online policy. Define the event

$$C_T = \left\{ S_3^{\star, \lfloor T/2 \rfloor} \geq Z_3^{1:T}/4 \right\}.$$

On the event $C_T^c \cap L_T \cap A_T \cap M_T$, we have that the online policy incorrectly (with respect the offline policy) rejects at least $\sqrt{T}/4$ candidates with abilities in I_3 . This is because we have that $C_T^c = \{S_3^{\star, \lfloor T/2 \rfloor} < Z_3^{1:T}/4\}$ and under the offline policy, we should hire all the candidates with ability in I_3 under the event $L_T \cap A_T$. Since $Z_3^{1:T} \leq 3Z_3^{\lfloor T/2 \rfloor}$ on the event A_T , under the event C_T^c , we have that $S_3^{\star, \lfloor T/2 \rfloor} < 3Z_3^{\lfloor T/2 \rfloor}/4$. Therefore the number of incorrect decisions is at least $Z_3^{\lfloor T/2 \rfloor}/4 \geq Z_3^{1:T}/12 \geq \sqrt{T}/24$. Due to these mistakes in the first half, the online algorithm will have to hire at least $\sqrt{T}/24$ candidates with ability in $I_1 \cup I_{2a}$. Essentially the online optimal policy due to the mistakes in the first half, will have to substitute the candidates that it rejected in the interval I_3 by hiring more candidates in the interval $I_1 \cup I_{2a}$ than what are hired by the offline algorithm. In particular, the online optimal policy will have to hire at least $\sqrt{T}/24$ more candidates in the interval $I_1 \cup I_{2a}$. Now since we want a lower bound on the regret, let us assume that all the candidates that were rejected were in the interval I_{3a} so that we can get a lower bound. If we considered that candidates with ability in the interval I_{3b} were also rejected, then compared to their substitutes in $I_1 \cup I_{2a}$ that will be hired, the compensation will be at least $1/2$ due to the gap $[1/4, 3/4]$, hence to get a lower bound, we just assume that the hiring mistakes only happened for the candidate with ability in I_{3a} . Now the number of arrivals in the interval I_{2a} is at most $\sqrt{T}/32$ under the event M_T and hence at least $\sqrt{T}/96$ additional candidates will need to be hired from the interval I_1 . Because of this the compensation that needs to be provided for each of the $\sqrt{T}/96$ mistakes is at least $(R_1 - 1)\Delta_\beta$. Hence we have that

$$\mathbb{E}[\Lambda(B, T) \mathbb{1}(C_T^c \cap L_T \cap A_T \cap H_T)] \geq (\sqrt{T}/96)(R_1 - 1)\Delta_\beta \mathbb{P}(C_T^c \cap L_T \cap A_T \cap H_T)$$

For each sequence in $\{\theta_1, \dots, \theta_T\} \in C_T \cap L_T \cap A_T \cap M_T$, we can construct a sequence in $C_T \cap H_T \cap A_T \cap M_T$. To do so, we keep the first $\lfloor T/2 \rfloor$ arrivals the same and do not make any changes to the values in the intervals I_2 and I_3 . Under the event L_T , we have that $Z_4^{\lfloor T/2 \rfloor + 1:T} \in \left[T/4 - 4\sqrt{T}, T/4 - 3\sqrt{T} \right]$ whereas under H_T , we have that $Z_4^{\lfloor T/2 \rfloor + 1:T} \geq T/4 + \sqrt{T}/2$. Therefore in the time steps $\{\lfloor T/2 \rfloor + 1, \dots, T\}$, we need to replace at most $9\sqrt{T}/2$ values in the interval I_1 with values in the interval I_4 which can be done by taking a mirror image across the line $x = 1/2$. Now this can be done as long as the number of arrivals in I_1 under the event $C_T \cap L_T \cap A_T \cap M_T$ (denoted by $Z_1^{\lfloor T/2 \rfloor + 1:T}$) is atleast $9\sqrt{T}/2$. Under the event $C_T \cap L_T \cap A_T \cap M_T$, we have that

$$\begin{aligned} Z_1^{\lfloor T/2 \rfloor + 1:T} &= (T - \lfloor T/2 \rfloor) - \left(Z_2^{\lfloor T/2 \rfloor + 1:T} + Z_3^{\lfloor T/2 \rfloor + 1:T} + Z_4^{\lfloor T/2 \rfloor + 1:T} \right) \\ &\geq (T - \lfloor T/2 \rfloor) - \left(Z_2^{1:T} + Z_3^{1:T} + Z_4^{\lfloor T/2 \rfloor + 1:T} \right), \\ &\geq T/4 + \sqrt{T}/2, \end{aligned}$$

where the last inequality follows from the fact that $Z_2^{1:T} \leq \sqrt{T}/32$, $Z_3^{1:T} \leq 2\sqrt{T}$ and $Z_4^{\lfloor T/2 \rfloor + 1:T} \leq T/4 - 3\sqrt{T}$. Therefore, for T large enough, we can replace values from I_1 to I_4 to construct a sequence in $C_T \cap H_T \cap A_T \cap M_T$, we call this resultant set D_T and since the probability of arrival in the interval I_1 and I_4 are bounded away from zero, we have that there exists $\hat{\alpha} > 0$ such that $\mathbb{P}(D_T) \geq \hat{\alpha} \mathbb{P}(C_T \cap L_T \cap A_T \cap M_T)$ and $\hat{\alpha}$ does not depend on T . Since the two sequences share the same first $\lfloor T/2 \rfloor$ elements, we have that $S_3^{*, \lfloor T/2 \rfloor} \geq Z_3^{1:T}/4$ also on the set D_T . Now under $H_T \cap A_T \cap M_T$, the offline algorithm would not hire any candidate with ability in the interval I_3 however the optimal online algorithm conditional on the event $C_T \cap A_T$, the online policy makes at least $Z_3^{1:T}/4 \geq \sqrt{T}/8$ mistakes. Again to get a lower bound on the regret, let us assume that all the mistakes happen for candidates with ability in $I_{3,b}$. Now this means that for the last $\lfloor T/2 \rfloor$ periods the optimal online algorithm will have to reject at least $\sqrt{T}/8$ candidates in the interval I_4 . Using an analogous argument for the previous case, we have that

$$\mathbb{E}[\Lambda(B, T) \mathbb{1}(D_T)] \geq (\sqrt{T}/8)(R_2 - 1) \Delta_\beta \mathbb{P}(D_T) \geq (\sqrt{T}/8)(R_2 - 1) \hat{\alpha} \mathbb{P}(C_T \cap L_T \cap A_T \cap M_T),$$

Since $\mathbb{P}(L_T \cap A_T \cap M_T) = \mathbb{P}(C_T^c \cap L_T \cap A_T \cap M_T) + \mathbb{P}(C_T \cap L_T \cap A_T \cap M_T)$, we have that

$$\begin{aligned} & \max\{\mathbb{P}(C_T^c \cap L_T \cap A_T \cap M_T), \hat{\alpha} \mathbb{P}(C_T \cap L_T \cap A_T \cap M_T)\} \\ & \geq \min\{\hat{\alpha}, 1\} (\mathbb{P}(C_T^c \cap L_T \cap A_T \cap M_T) + \mathbb{P}(C_T \cap L_T \cap A_T \cap M_T)) / 2 \\ & = \min\{\hat{\alpha}, 1\} \mathbb{P}(L_T \cap A_T \cap M_T) / 2. \end{aligned}$$

Let $\mathcal{C} = \min\{R_1 - 1, R_2 - 1\}$, we have that for all $T \geq T_0$ and $B = \lfloor T/2 \rfloor$

$$\begin{aligned} \text{Regret}(B, T) & \geq \max\left\{ \mathbb{E}[\Lambda(B, T) \mathbb{1}(C_T^c \cap L_T \cap A_T \cap M_T)], \mathbb{E}[\Lambda(B, T) \mathbb{1}(D_T)] \right\} \\ & \geq \mathcal{C} \Delta_\beta (\sqrt{T}/96) \max\{\mathbb{P}(C_T^c \cap L_T \cap A_T \cap M_T), \hat{\alpha} \mathbb{P}(C_T \cap L_T \cap A_T \cap M_T)\} \\ & \geq \mathcal{C} \Delta_\beta (\sqrt{T}/96) \min\{\hat{\alpha}, 1\} \mathbb{P}(L_T \cap A_T \cap M_T) / 2 \\ & \geq \tilde{\mathcal{C}} T^{\frac{1}{2} - \frac{1}{2(\beta+1)}} \end{aligned}$$

where $\tilde{\mathcal{C}} \equiv \mathcal{C}(\beta, \alpha, \hat{\alpha})$. This completes the proof. \square

EC.2.1. Proofs of Helper Lemmas for Theorem 3

EC.2.1.1. Proof of Lemma EC.2

Proof of Lemma EC.2 We have that the event $L_T = \left\{ T/4 - \sqrt{T}/2 \leq Z_4^{1:\lfloor T/2 \rfloor} \leq T/4 \right\} \cap \left\{ T/4 - 4\sqrt{T} \leq Z_4^{\lfloor T/2 \rfloor + 1:T/2} \leq T/4 - 3\sqrt{T} \right\} := L_T^1 \cap L_T^2$. Since the arrivals are i.i.d, we have that the events L_T^1 and L_T^2 are independent and hence we have that $\mathbb{P}(L_T) = \mathbb{P}(L_T^1) \mathbb{P}(L_T^2)$. We have that

$\mathbb{E} \left[Z_4^{1:\lfloor T/2 \rfloor} \right] \approx T/4 - \sqrt{T}/4 - \sqrt{T}/256$ and $\mathbb{E} \left[Z_4^{\lfloor T/2 \rfloor + 1:T} \right] \approx T/4 - \sqrt{T}/4 - \sqrt{T}/256$. Define $p := 1/2 - T^{-1/2}/2 - T^{-1/2}/128$. Then we have that

$$\begin{aligned} \mathbb{P}(L_T^1) &= \mathbb{P} \left(\mathbb{E} \left[Z_4^{1:\lfloor T/2 \rfloor} \right] - 63\sqrt{T}/256 \leq Z_4^{1:\lfloor T/2 \rfloor} \leq \mathbb{E} \left[Z_4^{1:\lfloor T/2 \rfloor} \right] + 65\sqrt{T}/256 \right), \\ &= \mathbb{P} \left(-63\sqrt{T}/256 \leq Z_4^{1:\lfloor T/2 \rfloor} - \mathbb{E} \left[Z_4^{1:\lfloor T/2 \rfloor} \right] \leq 65\sqrt{T}/256 \right), \\ &= \mathbb{P} \left(-\frac{63\sqrt{2}}{256\sqrt{p(1-p)}} \leq \frac{Z_4^{1:\lfloor T/2 \rfloor} - \mathbb{E} \left[Z_4^{1:\lfloor T/2 \rfloor} \right]}{\sqrt{\lfloor T/2 \rfloor p(1-p)}} \leq \frac{65\sqrt{2}}{256\sqrt{p(1-p)}} \right) \end{aligned}$$

Note that there exists a $T_{l'}$ such that for all $T \geq T_{l'}$, we have that $1/16 \leq p(1-p) \leq 1/4$ and since $Z_4^{1:\lfloor T/2 \rfloor} \sim \text{Bin}(\lfloor T/2 \rfloor, p)$, we can write $\bar{Z} := \left(Z_4^{1:\lfloor T/2 \rfloor} - \mathbb{E} \left[Z_4^{1:\lfloor T/2 \rfloor} \right] \right) / \sqrt{\lfloor T/2 \rfloor p(1-p)} = \sum_{t=1}^{\lfloor T/2 \rfloor} X_i$ where X_i is i.i.d with $\mathbb{E}[X_i] = 0, \mathbb{E}[X_i^2] = \sigma^2 = p(1-p) > 0$ and $\mathbb{E}[|X_i|^3] \leq \rho < \infty$ for some constant ρ . Define $\gamma_1 = -63\sqrt{2}/128, \gamma_2 = 65\sqrt{2}/128$. Then for all $T \geq T_{l'}$, we have that

$$\mathbb{P}(L_T^1) \stackrel{(a)}{\geq} \mathbb{P}(\gamma_1 \leq \bar{Z} \leq \gamma_2) \stackrel{(b)}{=} \mathbb{P}(\bar{Z} \leq \gamma_2) - \mathbb{P}(\bar{Z} \leq \gamma_1) \stackrel{(c)}{\geq} \Phi(\gamma_2) - \frac{C_1}{\sqrt{T}} - \Phi(\gamma_1) - \frac{C_2}{\sqrt{T}},$$

where (a) follows from the fact that for $T \geq T_{l'}$, $-\frac{63\sqrt{2}}{256\sqrt{p(1-p)}} \leq \gamma_1 < \gamma_2 \leq \frac{65\sqrt{2}}{256\sqrt{p(1-p)}}$, (b) follows trivially, (c) follows from the Berry-Esseen theorem. Since $\gamma_1 < \gamma_2$, we have that $\Phi(\gamma_2) - \Phi(\gamma_1) > 2\epsilon$ for some constant $\epsilon > 0$ independent of T and hence we can choose a $T_{l''}$ such that for all $T \geq \max\{T_{l'}, T_{l''}\}$, we have that $\Phi(\gamma_2) - \frac{C_1}{\sqrt{T}} - \Phi(\gamma_1) - \frac{C_2}{\sqrt{T}} \geq \epsilon$.

We can repeat an analogous argument and show that for T large enough we have that $\mathbb{P}(L_T^2) \geq \epsilon'$. Since $\mathbb{P}(L_T) = \mathbb{P}(L_T^1) \mathbb{P}(L_T^2)$, we have that there exists $T_l < \infty$ such that for all $T \geq T_l$, we have that $\mathbb{P}(L_T) \geq \epsilon\epsilon' := \alpha_l$. \square

EC.2.1.2. Proof of Lemma EC.3

Proof of Lemma EC.3 The proof of Lemma EC.3 follows analogously to the proof of Lemma EC.2. \square

EC.2.1.3. Proof of Lemma EC.4

Proof of Lemma EC.4 For simplicity of notation, let $Z_3^T := Z_3^{1:T}$ and $Z_3^{\lfloor T/2 \rfloor} := Z_3^{1:\lfloor T/2 \rfloor}$. We have that $A_T^c = \{Z_3^T < \sqrt{T}/2\} \cup \{Z_3^T > \min\{3Z_3^{\lfloor T/2 \rfloor}, 2\mathbb{E}[Z_3^T]\}\}$. Denote by $X_3^{\lfloor T/2 \rfloor}$ as the number of arrivals that happen in the interval I_3 in the last $\lfloor T/2 \rfloor$ time periods. We can write $Z_3^T = Z_3^{\lfloor T/2 \rfloor} + X_3^{\lfloor T/2 \rfloor}$ where $Z_3^{\lfloor T/2 \rfloor}$ is independent of $X_3^{\lfloor T/2 \rfloor}$ due to the i.i.d arrivals. We have that $\mathbb{E}[Z_3^T] = \sqrt{T}$ and $\mathbb{E}[X_3^{\lfloor T/2 \rfloor}] = T^{-1/2}\lfloor T/2 \rfloor \approx \sqrt{T}/2$ and $\mathbb{E}[Z_3^{\lfloor T/2 \rfloor}] = T^{-1/2}\lfloor T/2 \rfloor \approx \sqrt{T}/2$. Now we have that

$$\begin{aligned} \mathbb{P} \left(Z_3^T > \min\{3Z_3^{\lfloor T/2 \rfloor}, 2\mathbb{E}[Z_3^T]\} \right) &= \mathbb{P} \left(Z_3^{\lfloor T/2 \rfloor} + X_3^{\lfloor T/2 \rfloor} > \min\{3Z_3^{\lfloor T/2 \rfloor}, 2\mathbb{E}[Z_3^T]\} \right) \\ &= \mathbb{P} \left(X_3^{\lfloor T/2 \rfloor} > \min\{2Z_3^{\lfloor T/2 \rfloor}, 2\mathbb{E}[Z_3^T] - Z_3^{\lfloor T/2 \rfloor}\} \right) \end{aligned}$$

To simplify notation, we will denote $Z_3^{\lfloor T/2 \rfloor} := Z$ and $X_3^{\lfloor T/2 \rfloor} := X$. Furthermore, we will assume that $\mathbb{E}[Z] = \mathbb{E}[X] = \mu$ and that $\mathbb{E}[Z_3^T] = 2\mu$. Also define $p_A = T^{-1/2}$. Then we have that

$$\mathbb{P}(X > \min\{2Z, 4\mu - Z\}) = \mathbb{P}(X - \mu > \min\{2Z - \mu, 3\mu - Z\})$$

Define the event $F = \{3\mu/4 \leq Z \leq 4\mu/3\}$ and we have that $F^c = \{Z < 3\mu/4\} \cup \{Z > 4\mu/3\}$. The event F implies that $\min\{2Z - \mu, 3\mu - Z\} = 2Z - \mu \geq \mu/2$. Hence we have that

$$\begin{aligned} \mathbb{P}(X - \mu > \min\{2Z - \mu, 3\mu - Z\}) &= \mathbb{P}(X - \mu > \min\{2Z - \mu, 3\mu - Z\} | F) \mathbb{P}(F) \\ &\quad + \mathbb{P}(X - \mu > \min\{2Z - \mu, 3\mu - Z\} | F^c) \mathbb{P}(F^c) \\ &\leq \mathbb{P}(X - \mu > \mu/2) + \mathbb{P}(Z < 3\mu/4) + \mathbb{P}(Z > 4\mu/3) \end{aligned}$$

Now we have that

$$\begin{aligned} \mathbb{P}(X - \mu > \mu/2) &= \mathbb{P}\left(\frac{X - \mu}{\sqrt{(T/2)p_A(1-p_A)}} > \frac{\mu}{2\sqrt{(T/2)p_A(1-p_A)}}\right) \leq \mathbb{P}\left(\frac{X - \mu}{\sqrt{(T/2)p_A(1-p_A)}} > \frac{\sqrt{2}\sqrt[4]{T}}{4}\right) \\ \mathbb{P}(Z - \mu < -\mu/4) &= \mathbb{P}\left(\frac{Z - \mu}{\sqrt{(T/2)p_A(1-p_A)}} < \frac{-\mu}{4\sqrt{(T/2)p_A(1-p_A)}}\right) \leq \mathbb{P}\left(\frac{Z - \mu}{\sqrt{(T/2)p_A(1-p_A)}} < \frac{-\sqrt{2}\sqrt[4]{T}}{4}\right) \\ \mathbb{P}(Z - \mu > \mu/3) &= \mathbb{P}\left(\frac{Z - \mu}{\sqrt{(T/2)p_A(1-p_A)}} > \frac{\mu}{3\sqrt{(T/2)p_A(1-p_A)}}\right) \leq \mathbb{P}\left(\frac{Z - \mu}{\sqrt{(T/2)p_A(1-p_A)}} > \frac{\sqrt{2}\sqrt[4]{T}}{3}\right) \end{aligned}$$

Using the Berry-Esseen theorem, for a universal constant $C < \infty$, we have that

$$\begin{aligned} \left| \mathbb{P}\left(\frac{X - \mu}{\sqrt{(T/2)p_A(1-p_A)}} > \frac{\sqrt{2}\sqrt[4]{T}}{4}\right) - \bar{\Phi}\left(\frac{\sqrt{2}\sqrt[4]{T}}{4}\right) \right| &\leq \frac{C}{\sqrt[4]{T}}, \\ \left| \mathbb{P}\left(\frac{Z - \mu}{\sqrt{(T/2)p_A(1-p_A)}} < \frac{-\sqrt{2}\sqrt[4]{T}}{4}\right) - \bar{\Phi}\left(\frac{-\sqrt{2}\sqrt[4]{T}}{4}\right) \right| &\leq \frac{C}{\sqrt[4]{T}}, \\ \left| \mathbb{P}\left(\frac{Z - \mu}{\sqrt{(T/2)p_A(1-p_A)}} > \frac{\sqrt{2}\sqrt[4]{T}}{3}\right) - \bar{\Phi}\left(\frac{\sqrt{2}\sqrt[4]{T}}{3}\right) \right| &\leq \frac{C}{\sqrt[4]{T}} \end{aligned}$$

Using the fact that $\Phi(-x) = \bar{\Phi}(x)$ and the fact that $\bar{\Phi}(x) \leq C \exp(-cx^2)/x$, for some constants c, C , we have that

$$\begin{aligned} \mathbb{P}(X - \mu > \mu/2) &\leq \frac{C}{\sqrt[4]{T}} + \frac{C \exp(-c\sqrt{T})}{\sqrt[4]{T}} \\ \mathbb{P}(Z - \mu < -\mu/4) &\leq \frac{C}{\sqrt[4]{T}} + \frac{C \exp(-c\sqrt{T})}{\sqrt[4]{T}} \\ \mathbb{P}(Z - \mu > \mu/3) &\leq \frac{C}{\sqrt[4]{T}} + \frac{C \exp(-c\sqrt{T})}{\sqrt[4]{T}} \end{aligned}$$

Combining the above three inequalities, for some constants c, C , we have that,

$$\mathbb{P}\left(Z_3^T > \min\{3Z_3^{\lfloor T/2 \rfloor}, 2\mathbb{E}[Z_3^T]\}\right) \leq \frac{C}{\sqrt[4]{T}} + \frac{C \exp(-c\sqrt{T})}{\sqrt[4]{T}}$$

We can similarly show that $\mathbb{P}\left(Z_3^T < \sqrt{T}/2\right) \leq C/\sqrt[4]{T} + C \exp(-c\sqrt{T})/\sqrt[4]{T}$. Therefore for some other constants c, C , we have that

$$\mathbb{P}(A_T^c) \leq \mathbb{P}\left(Z_3^T < \sqrt{T}/2\right) + \mathbb{P}\left(Z_3^T > \min\{3Z_3^{\lfloor T/2 \rfloor}, 2\mathbb{E}[Z_3^T]\}\right) \leq C/\sqrt[4]{T} + C \exp(-c\sqrt{T})/\sqrt[4]{T}$$

Hence we have that for all $T \geq T_a$, we have that $\mathbb{P}(A_T^c) \leq \alpha/4$ and this completes the proof. \square

EC.2.1.4. Proof of Lemma EC.5

Proof of Lemma EC.5 The proof of Lemma EC.5 follows analogously to the proof of Lemma EC.4. \square

EC.3. Some Details and Proofs Related to Section 5

EC.3.1. Details of Compensated Coupling

Let B_t denote the remaining budget after time t . For any $\vec{\omega}$, let $V_t^{\text{off}}(B_t)[\vec{\omega}]$ denote the maximum of the total ability of candidates selected by the offline algorithm from time t onwards, given that the remaining budget is B_t . The deterministic dynamic program faced by the offline algorithm can be written as

$$\begin{aligned} V_t^{\text{off}}(B_t)[\vec{\omega}] &= \max_{a \in \mathcal{A}(B_t)} \{r(B_t, a, \omega_{t+1}) + V_{t+1}^{\text{off}}(\mathcal{T}(B_t, a, \omega_{t+1}))[\vec{\omega}]\}, \quad \forall t \in \{0, 1, \dots, T-2\} \\ V_{T-1}^{\text{off}}(B_{T-1})[\vec{\omega}] &= \max_{a \in \mathcal{A}(B_{T-1})} \{r(B_{T-1}, a, \omega_T)\} \end{aligned}$$

where $\mathcal{A}(B) = \{\text{hire, reject}\}$, $\forall B > 0$ and $\mathcal{A}(0) = \{\text{reject}\}$ and $r(B_t, \text{hire}, \omega_t) = \omega_t$, $r(B_t, \text{reject}, \omega_t) = 0$ and $\mathcal{T}(B_t, \text{hire}, \omega_t) = B_t - 1$, $\mathcal{T}(B_t, \text{reject}, \omega_t) = B_t$. Hence we have that $V_{\text{off}}^*(B, T) = \mathbb{E}[V_0^{\text{off}}(B)[\vec{\omega}]]$. Similarly, we can define a dynamic program for the CwG policy. For any $\vec{\omega}$, let $V_t^{\pi^{\text{cwg}}}(B_t)[\vec{\omega}]$ denote the total ability of the hired candidates from time t onwards, given that the remaining budget is B_t at time t . Then the DP corresponding to the policy $\pi^{\text{cwg}} = \{\mu_1^{\text{cwg}}, \mu_2^{\text{cwg}}, \dots, \mu_T^{\text{cwg}}\}$ is given as

$$\begin{aligned} V_t^{\pi^{\text{cwg}}}(B_t)[\vec{\omega}] &= r(B_t, \mu_{t+1}^{\text{cwg}}, \omega_{t+1}) + V_{t+1}^{\pi^{\text{cwg}}}(\mathcal{T}(B_t, \mu_{t+1}^{\text{cwg}}, \omega_{t+1})), \quad \forall t \in \{0, 1, \dots, T-2\} \\ V_{T-1}^{\pi^{\text{cwg}}}(B_{T-1})[\vec{\omega}] &= r(B_{T-1}, \mu_T^{\text{cwg}}, \omega_T) \end{aligned}$$

Note that the action μ_t^{cwg} is a function of the remaining budget B_t but for notational convenience, we do not make that dependence explicit. We have that $V_{\text{on}}^{\pi^{\text{cwg}}}(B, T) = \mathbb{E}[V_0^{\pi^{\text{cwg}}}(B)[\vec{\omega}]]$. Next we discuss the intuition behind the compensated coupling framework developed by Vera and Banerjee (2021). For a given sample path $\vec{\omega}$, suppose that both the offline algorithm and the CwG policy are at time t with the same remaining budget B_t . Let $a_t^{\text{off}}[\vec{\omega}]$ be the set of actions which are optimal

under the offline algorithm at time t . If $\mu_t^{\text{cwg}} \notin a_t^{\text{off}}[\vec{\omega}]$, then it could be the case that choosing μ_t^{cwg} at time t leads to a lower future total reward for the offline algorithm than choosing an action in $a_t^{\text{off}}[\vec{\omega}]$. In order to persuade the offline algorithm to take the same action as that of the CwG policy, the idea is to *compensate* the offline algorithm such that choosing μ_t^{cwg} will make the future reward no less than that by choosing an action in $a_t^{\text{off}}[\vec{\omega}]$. Let $\partial R_t(B_{t-1}, \mu_t^{\text{cwg}})[\vec{\omega}]$ denote the compensation that needs to be provided to the offline algorithm so that it takes the same action as the one determined by the online CwG policy at time t . From Vera and Banerjee (2021), we know that

$$\partial R_{t+1}(B_t, \mu_{t+1}^{\text{cwg}})[\vec{\omega}] = V_t^{\text{off}}(B_t)[\vec{\omega}] - (r(B_t, \mu_{t+1}^{\text{cwg}}, \omega_{t+1}) + V_{t+1}^{\text{off}}(\mathcal{T}(B_t, \mu_{t+1}^{\text{cwg}}, \omega_{t+1}))[\vec{\omega}])$$

Define $Q_t(B_{t-1}) = \{\vec{\omega} : \mu_t^{\text{cwg}} \notin a_t^{\text{off}}[\vec{\omega}]\}$ to be the set of sample paths where the action of the offline algorithm and the online algorithm differ given that the remaining hiring budget at time t is B_t . We only compensate the offline algorithm at time t if $\mu_t^{\text{cwg}} \notin a_t^{\text{off}}[\vec{\omega}]$. Based on this discussion, we can decompose the regret under the CwG policy as

LEMMA EC.6 (Vera and Banerjee (2021)). *For all $T \in \mathbb{N}$ and for all $B \in [T]$, consider the CwG policy as described in Algorithm 2 and let B_t be the resultant hiring budget process under the CwG policy with $B_0 = B$, then we have that*

$$\text{Regret}(B, T; \text{CwG}) = \sum_{t=0}^{T-1} \mathbb{E} [\partial R_{t+1}(B_t, \mu_{t+1}^{\text{cwg}})[\vec{\omega}] \cdot \mathbb{1}_{Q_{t+1}(B_t)}[\vec{\omega}]]$$

We refer the readers to Vera and Banerjee (2021) for a proof of the above lemma.

EC.3.2. Proof of Lemma 2

Proof of Lemma 2. Let us assume that $p_t^{\text{ce}} \geq p_t^{\text{otg}} + \sqrt{2 \log \tau / \tau}$. Now conditional on B_t and given the knowledge of p_t^{otg} , we know that there are B_t arrivals with quantile larger than p_t^{otg} and $\tau - B_t$ arrivals with quantiles less than p_t^{otg} . Let $X_t \triangleq \text{Ber}(\tau, (p_t^{\text{ce}} - \sqrt{2 \log \tau / \tau})^+)$ with $\mathbb{E}[X_t | B_t] = (\tau - B_t - \sqrt{2 \log \tau / \tau})^+$. Then we have that

$$\mathbb{P}\left(p_t^{\text{ce}} \geq p_t^{\text{otg}} + \sqrt{2 \log \tau / \tau} \mid B_t\right) \leq \mathbb{P}\left(X_t \geq \tau - B_t \mid B_t\right) \leq \mathbb{P}\left(X_t - \mathbb{E}[X_t | B_t] \geq \sqrt{2 \log \tau / \tau} \mid B_t\right) \leq 1/\tau^4$$

where the last inequality follows from the Hoeffding inequality. It follows that $\mathbb{P}(p_t^{\text{ce}} \geq p_t^{\text{otg}} + \sqrt{2 \log \tau / \tau}) \leq 1/\tau^4$. Analogously, we can show the same for the case of $p_t^{\text{otg}} \geq p_t^{\text{ce}} + \sqrt{2 \log \tau / \tau}$. \square

EC.3.3. Proof of Lemma 3

Proof of Lemma 3. We have that $|p_t^{\text{cwg}} - p_t^{\text{otg}}| = |p_t^{\text{cwg}} - p_t^{\text{ce}} + p_t^{\text{ce}} - p_t^{\text{otg}}| \leq |p_t^{\text{cwg}} - p_t^{\text{ce}}| + |p_t^{\text{ce}} - p_t^{\text{otg}}|$. By the definition of the algorithm we have that $|p_t^{\text{cwg}} - p_t^{\text{ce}}| \leq \sqrt{2 \log \tau / \tau}$. Now conditional on B_t , we have that event $A_{1,t}$ implies the event $A_{2,t}$ and hence we have that $A_{2,t}^c$ implies $A_{1,t}^c$ which implies that $\mathbb{P}(A_{2,t}^c | B_t) \leq \mathbb{P}(A_{1,t}^c | B_t)$. Using the proof of Lemma 2, we have that $\mathbb{P}(A_{2,t}^c | B_t) \leq 2/\tau^4$ and the claim of the lemma follows. \square

EC.3.4. Proof of Lemma 4

Proof of Lemma 4. We have that $A_{3,t}^c = \cup_{i,j:Q_i^\circ \cap Q_j^\circ = \emptyset} \{p_t^{\text{cwg}} \in Q_i, p_t^{\text{otg}} \in Q_j\}$ where A° denotes the interior of the set A . Consider the event $\{p_t^{\text{cwg}} \in Q_i, p_t^{\text{otg}} \in Q_j\}$ such that $Q_i^\circ \cap Q_j^\circ = \emptyset$. From the definition of p_t^{cwg} in Algorithm 2 and the fact that $Q_i^\circ \cap Q_j^\circ = \emptyset$ implies that $|p_t^{\text{ce}} - p_t^{\text{otg}}| \geq \sqrt{2 \log \tau / \tau}$. Using Lemma 2 and the union bound completes the proof. \square

EC.3.5. Proof of Lemma 5

Proof of Lemma 5. Assume that $p_t^{\text{cwg}} \leq q_t^l$, then according to the definition of p_t^{otg} defined in Section 5.3, we have that $p_t^{\text{otg}} = q_t^u$. Compensation is provided only if $q_t^\theta \in [p_t^{\text{cwg}}, p_t^{\text{otg}}]$. Suppose that is the case, then we have that $F^{-1}((p_t^{\text{cwg}})^+) \leq \theta_t \leq F^{-1}(p_t^{\text{otg}}) = F^{-1}(q_t^u) = u_t$. The CwG policy would accept the candidate with ability θ_t since $\theta_t \geq F^{-1}((p_t^{\text{cwg}})^+)$ where as the OTG would want to reject the candidate with ability θ_t , because in the future it knows that it can select a candidate with ability at least $u_t \geq \theta_t$. Hence to persuade the OTG, we need to compensate it $u_t - \theta_t = F^{-1}(p_t^{\text{otg}}) - \theta_t$ and maximum compensation can hence be $F^{-1}(p_t^{\text{otg}}) - F^{-1}((p_t^{\text{cwg}})^+)$. An analogous analysis can be done for the case when $p_t^{\text{cwg}} \geq q_t^u$ which follows similarly. \square