

MINIMAL UNITARY DILATIONS FOR COMMUTING CONTRACTIONS

SOURAV PAL AND PRAJAKTA SAHASRABUDDHE

ABSTRACT. For commuting contractions T_1, \dots, T_n acting on a Hilbert space \mathcal{H} with $T = \prod_{i=1}^n T_i$, we show that (T_1, \dots, T_n) dilates to commuting isometries (V_1, \dots, V_n) on the minimal isometric dilation space of T with $V = \prod_{i=1}^n V_i$ being the minimal isometric dilation of T if and only if (T_1^*, \dots, T_n^*) dilates to commuting isometries (Y_1, \dots, Y_n) on the minimal isometric dilation space of T^* with $Y = \prod_{i=1}^n Y_i$ being the minimal isometric dilation of T^* . Then, we prove an analogue of this result for unitary dilations of (T_1, \dots, T_n) and its adjoint. We find a necessary and sufficient condition such that (T_1, \dots, T_n) possesses a unitary dilation (W_1, \dots, W_n) on the minimal unitary dilation space of T with $W = \prod_{i=1}^n W_i$ being the minimal unitary dilation of T . We show an explicit construction of such a unitary dilation on both Schäffer and Sz. Nagy-Foias minimal unitary dilation spaces of T . Also, we show that a relatively weaker hypothesis is necessary and sufficient for the existence of such a unitary dilation when T is a C_0 contraction, i.e. when $T^{*n} \rightarrow 0$ strongly as $n \rightarrow \infty$. We construct a different unitary dilation for (T_1, \dots, T_n) when T is a C_0 contraction.

CONTENTS

1. Introduction	1
2. A few preparatory results	3
3. Schäffer-type minimal unitary dilation	5
4. Minimal unitary dilation when the product is a C_0 contraction	18
5. Sz. Nagy-Foias type minimal unitary dilation	20
6. Appendix	23
7. Data availability statement	30
References	30

1. INTRODUCTION

Throughout the paper all operators are bounded linear operators acting on complex Hilbert spaces. A contraction is an operator with norm not greater than 1. We begin with the definitions of isometric and unitary dilations of a tuple of commuting contractions.

Definition 1.1. Let (T_1, \dots, T_n) be a tuple of commuting contractions acting on a Hilbert space \mathcal{H} . A commuting tuple of unitaries (W_1, \dots, W_n) acting on a Hilbert space \mathcal{H}' is said to be a *unitary dilation* of (T_1, \dots, T_n) if \mathcal{H} can be realized as a closed linear subspace of \mathcal{H}' and for any

2010 *Mathematics Subject Classification.* 47A15, 47A20, 47B35, 47B38.

Key words and phrases. Unitary dilation, Minimal dilation space.

The first named author is supported by the Seed Grant of IIT Bombay, the CPDA and the MATRICS Award (Award No. MTR/2019/001010) of Science and Engineering Research Board (SERB), India. The second named author has been supported by the Ph.D Fellowship of Council of Scientific and Industrial Research (CSIR), India.

non-negative integers k_1, \dots, k_n we have

$$T_1^{k_1} \dots T_n^{k_n} = P_{\mathcal{H}}(W_1^{k_1} \dots W_n^{k_n})|_{\mathcal{H}},$$

where $P_{\mathcal{H}} : \mathcal{H}' \rightarrow \mathcal{H}$ is the orthogonal projection. Moreover, such a unitary dilation is called *minimal* if

$$\mathcal{H}' = \overline{\text{Span}} \{W_1^{t_1} \dots W_n^{t_n} h : h \in \mathcal{H}, t_1, \dots, t_n \in \mathbb{Z}\}.$$

Similarly, a commuting tuple of isometries (V_1, \dots, V_n) acting on a Hilbert space \mathcal{H} is said to be an *isometric dilation* of (T_1, \dots, T_n) if \mathcal{H} can be realized as a closed linear subspace of \mathcal{H} and for any non-negative integers k_1, \dots, k_n we have

$$T_1^{k_1} \dots T_n^{k_n} = P_{\mathcal{H}}(V_1^{k_1} \dots V_n^{k_n})|_{\mathcal{H}}.$$

Moreover, such an isometric dilation is called *minimal* if

$$\mathcal{H} = \overline{\text{Span}} \{V_1^{k_1} \dots V_n^{k_n} h : h \in \mathcal{H}, k_1, \dots, k_n \in \mathbb{N} \cup \{0\}\}.$$

A path-breaking work due to Sz. Nagy, [36] established that a contraction always dilates to a unitary. The result of Sz. Nagy was generalized by Ando, [3] to a pair of commuting contractions. A pair of commuting contractions possesses an unconditional unitary dilation. Later, Parrott showed by a counter example that a triple of commuting contractions may not dilate to a triple of commuting unitaries, see [42]. In a more general operator theoretic language rational dilation succeeds on the closed unit disk $\overline{\mathbb{D}}$ and on the closed bidisc $\overline{\mathbb{D}^2}$ and fails on the closed polydisc $\overline{\mathbb{D}^n}$ when $n \geq 3$. Thus, for $n \geq 3$ history witnessed only conditional unitary dilations for commuting contractions. Some notable works in this direction are due to Agler [1], Brehmer [18], Curto, Vasilescu [21, 22], Ball, Li, Timotin, Trent [9], Ball, Trent, Vinnikov [11], Eschmeier [27], Müller, Vasilescu [35], Burdak [19], Barik, Das, Sarkar [12] and many others, see the reference list and the references therein.

In this article we study isometric and unitary dilations of a tuple of commuting contractions (T_1, \dots, T_n) on the minimal isometric and minimal unitary dilation spaces (which are always unique upto unitaries) of $T = \prod_{i=1}^n T_i$. Our first main result is the following.

Theorem 1.2. *Let $T_1, \dots, T_n \in \mathcal{B}(\mathcal{H})$ be commuting contractions and let \mathcal{H} be the minimal isometric dilation space for their product $T = \prod_{i=1}^n T_i$. Then (T_1, \dots, T_n) possesses an isometric dilation (V_1, \dots, V_n) on \mathcal{H} with $V = \prod_{i=1}^n V_i$ being the minimal isometric dilation of T if and only if (T_1^*, \dots, T_n^*) possesses an isometric dilation (Y_1, \dots, Y_n) on \mathcal{H}_* with $Y = \prod_{i=1}^n Y_i$ being the minimal isometric dilation of T , where \mathcal{H}_* is the minimal isometric dilation space for T^* .*

In our second main result, which is stated below, we find a necessary and sufficient condition such that (T_1, \dots, T_n) dilates to a commuting unitary tuple (W_1, \dots, W_n) on the minimal unitary dilation space \mathcal{H}' of T with $\prod_{i=1}^n W_i = W$ being the minimal unitary dilation of T .

Theorem 1.3. *Let $T_1, \dots, T_n \in \mathcal{B}(\mathcal{H})$ be commuting contractions, $T = \prod_{j=1}^n T_j$ and $T_i' = \prod_{j \neq i} T_j$ for $1 \leq i \leq n$.*

- (a) *If $\widetilde{\mathcal{H}_0}$ is a minimal unitary dilation space for T , then (T_1, \dots, T_n) possesses a unitary dilation (W_1, \dots, W_n) on $\widetilde{\mathcal{H}_0}$ with $W = \prod_{i=1}^n W_i$ being the minimal unitary dilation of T if and only if there exist unique projections P_1, \dots, P_n and unique commuting unitaries U_1, \dots, U_n in $\mathcal{B}(\mathcal{D}_T)$ with $\prod_{i=1}^n U_i = I$ such that the following hold for $i = 1, \dots, n$:*

$$(1) D_T T_i = P_i^\perp U_i^* D_T + P_i U_i^* D_T T, \quad$$

- (2) $P_i^\perp U_i^* P_j^\perp U_j^* = P_j^\perp U_j^* P_i^\perp U_i^*$,
 - (3) $U_i P_i U_j P_j = U_j P_j U_i P_i$,
 - (4) $D_T U_i P_i U_i^* D_T = D_{T_i}^2$,
 - (5) $P_1 + U_1^* P_2 U_1 + U_1^* U_2^* P_3 U_2 U_1 + \dots + U_1^* U_2^* \dots U_{n-1}^* P_n U_{n-1} \dots U_2 U_1 = I_{\mathcal{D}_T}$.
- (b) *Such a unitary dilation is minimal and unique in the sense that if (X_1, \dots, X_n) on $\widetilde{\mathcal{K}}_1$ is another unitary dilation of (T_1, \dots, T_n) such that the product $X = \prod_{i=1}^n X_i$ is a minimal unitary dilation of T , then there is a unitary $\rho : \widetilde{\mathcal{K}}_0 \rightarrow \widetilde{\mathcal{K}}_1$ such that $(X_1, \dots, X_n) = (\rho * W_1 \rho, \dots, \rho * W_n \rho)$.*

This is Theorem 3.8 in this paper. We show an explicit construction of such a unitary dilation on the Schäffer's minimal unitary dilation space $\widetilde{\mathcal{K}}_0 = l^2(\mathcal{D}_T) \oplus \mathcal{H} \oplus l^2(\mathcal{D}_{T^*})$ of T , where $\mathcal{D}_T = \overline{\text{Ran}}(I - T^* T)^{\frac{1}{2}}$ and $\mathcal{D}_{T^*} = \overline{\text{Ran}}(I - T T^*)^{\frac{1}{2}}$. Note that upto a unitary $\widetilde{\mathcal{K}}_0$ is the smallest Hilbert space on which (T_1, \dots, T_n) can have such a unitary dilation and the reason is that it is the minimal unitary dilation space of the product $\prod_{i=1}^n T_i = T$.

In [41], the authors of this article show that (T_1, \dots, T_n) dilates to a commuting tuple of isometries (V_1, \dots, V_n) on the minimal isometric dilation space \mathcal{H} of T with $V = \prod_{i=1}^n V_i$ being the minimal isometric dilation of T if and only if the conditions (1) – (5) of Theorem 1.3 hold. Naturally, any unitary extension of the subnormal tuple (V_1, \dots, V_n) becomes a unitary dilation for (T_1, \dots, T_n) . The most interesting fact about the unitary dilation in Theorem 1.3 is that without any additional hypothesis (V_1, \dots, V_n) on \mathcal{H} admits a minimal unitary extension (W_1, \dots, W_n) on \mathcal{H}' , the minimal unitary dilation space for T .

We also show in Theorem 5.3 that such a unitary dilation can be constructed with the conditions (1) – (4) only, though we do not have an exact converse part then. We construct a special unitary dilation for (T_1, \dots, T_n) when the product T is a C_0 contraction, i.e. $T^{*n} \rightarrow 0$ strongly as $n \rightarrow \infty$. Indeed, in Theorem 4.3 with an explicit construction we show that an analogue of Theorem 1.3 can be obtained with a weaker hypothesis when T is a C_0 contraction. This is another main result of this paper. The other achievement of this paper is Theorem 5.2 in which we construct an explicit unitary dilation for (T_1, \dots, T_n) on the Sz. Nagy-Foias minimal unitary dilation space of T when T is a completely non-unitary (c.n.u.) contraction, i.e. when T is missing a unitary summand in its canonical decomposition. We accumulate a few preparatory results in Section 2.

2. A FEW PREPARATORY RESULTS

We begin with a famous result due to Douglas, Muhly and Pearcy. This result will play a major role in the proof of the main results of this paper.

Proposition 2.1 ([26], Proposition 2.2). *For $i = 1, 2$, let T_i be a contraction on a Hilbert space \mathcal{H}_i and let X be an operator from \mathcal{H}_2 to \mathcal{H}_1 . A necessary and sufficient condition that the operator on $\mathcal{H}_1 \oplus \mathcal{H}_2$ defined by the matrix $\begin{bmatrix} T_1 & X \\ 0 & T_2 \end{bmatrix}$ be a contraction is that there exists a contraction C mapping \mathcal{H}_2 into \mathcal{H}_1 such that $X = (I - T_1 T_1^*)^{\frac{1}{2}} C (I - T_2^* T_2)^{\frac{1}{2}}$.*

The following theorem is another general and useful result in operator theory.

Theorem 2.2 ([17], Lemma 13). *Let (R_1, \dots, R_{n-1}, U) on \mathcal{H} be a dilation of (S_1, \dots, S_{n-1}, P) on \mathcal{H} , where P is a contraction on \mathcal{H} and U on \mathcal{H} is the Schäffer minimal unitary dilation of P . Then*

for all $j = 1, 2, \dots, n-1$, R_j admits a matrix representation of the form

$$\begin{bmatrix} * & * & * \\ 0 & S_j & * \\ 0 & 0 & * \end{bmatrix}$$

with respect to the decomposition $\mathcal{K} = l^2(\mathcal{D}_T) \oplus \mathcal{H} \oplus l^2(\mathcal{D}_T^*)$.

In [15], Berger, Coburn and Lebow found the most popular factorization of a pure isometry. This is stated as Theorem 3.5 in this paper. Later Bercovici, Douglas and Foias proved a finer version of that result which we are going to state below.

Lemma 2.3 (Bercovici, Douglas and Foias, [14]). *Let U_1, \dots, U_n be unitaries on Hilbert space \mathcal{H} and P_1, \dots, P_n be orthogonal projections on \mathcal{H} . For $1 \leq i \leq n$, let $V_i = M_{U_i P_i^\perp + z U_i P_i}$. Then (V_1, \dots, V_n) defines a commuting n -tuple of isometries with $\prod_{i=1}^n V_i = M_z$ if and only if the following conditions are satisfied.*

- (1) $U_i U_j = U_j U_i$ for all $1 \leq i < j \leq n$,
- (2) $U_1 \dots U_n = I_{\mathcal{H}}$,
- (3) $P_j + U_j^* P_i U_j = P_i + U_i^* P_j U_i \leq I_{\mathcal{H}}$, for all $i \neq j$ and
- (4) $P_1 + U_1^* P_2 U_1 + U_1^* U_2^* P_3 U_2 U_1 + \dots + U_1^* U_2^* \dots U_{n-1}^* P_n U_{n-1} \dots U_2 U_1 = I_{\mathcal{H}}$.

The following result also appeared in [14].

Lemma 2.4 ([14], Lemma 2.2). *Consider unitary operators U, U_1, U_2 and orthogonal projections P, P_1 and P_2 on Hilbert space \mathcal{H} . If $V_{U,P}, V_{U_1,P_1}$ and V_{U_2,P_2} on $H^2(\mathcal{H})$ are defined as $V_{U,P} = M_{P^\perp U^* + z P U^*}$, $V_{U_1,P_1} = M_{P_1^\perp U_1^* + z P_1 U_1^*}$ and $V_{U_2,P_2} = M_{P_2^\perp U_2^* + z P_2 U_2^*}$, then the following are equivalent.*

- (i) $V_{U,P} = V_{U_1,P_1} V_{U_2,P_2}$,
- (ii) $U = U_1 U_2$ and $P = P_1 + U_1^* P_2 U_1$.

Let us state a pair of lemma that are analogous to Lemmas 2.3 & 2.4 respectively. In fact they provide factorization of similar kind for co-analytic symbols. The results can be proved using similar techniques as in the proof of Lemmas 2.3 & 2.4 in [14] and thus we skip the proofs. These results will be used in the proof of the main unitary dilation theorem.

Lemma 2.5. *Let U_1, \dots, U_n be unitaries and P_1, \dots, P_n be orthogonal projections on \mathcal{H} . For each $1 \leq i \leq n$, let $V_i = M_{U_i Q_i^\perp + \bar{z} U_i P_i}$. Then (V_1, \dots, V_n) defines a commuting n -tuple of co-isometries with $\prod_{i=1}^n V_i = M_{\bar{z}}$ if and only if the following conditions are satisfied.*

- (1) $U_i U_j = U_j U_i$ for all $1 \leq i < j \leq n$,
- (2) $U_1 \dots U_n = I_{\mathcal{H}}$,
- (3) $P_j + U_j^* P_i U_j = P_i + U_i^* P_j U_i \leq I_{\mathcal{H}}$, for all $i \neq j$ and
- (4) $P_1 + U_1^* P_2 U_1 + U_1^* U_2^* P_3 U_2 U_1 + \dots + U_1^* U_2^* \dots U_{n-1}^* P_n U_{n-1} \dots U_2 U_1 = I_{\mathcal{H}}$.

Lemma 2.6. *Consider unitary operators U, U_1, U_2 and orthogonal projections P, P_1 and P_2 on Hilbert space \mathcal{H} . If $W_{U,P}, W_{U_1,P_1}$ and W_{U_2,P_2} are Toeplitz operators on $H^2(\mathcal{H})$ defined by $W_{U,P} = M_{U P^\perp + \bar{z} U P}$, $W_{U_1,P_1} = M_{U_1 P_1^\perp + \bar{z} U_1 P_1}$ and $W_{U_2,P_2} = M_{U_2 P_2^\perp + \bar{z} U_2 P_2}$, then the following are equivalent.*

- (i) $W_{U,P} = W_{U_1,P_1} W_{U_2,P_2}$,
- (ii) $U = U_1 U_2$ and $P = P_2 + U_2^* P_1 U_2$.

3. SCHÄFFER-TYPE MINIMAL UNITARY DILATION

In this Section, we find a necessary and sufficient condition such that a tuple of commuting contraction (T_1, \dots, T_n) dilates to a tuple of commuting unitaries acting on the minimal unitary dilation space of $T = \prod_{i=1}^n T_i$. The dilation is minimal and we also show an explicit construction of such a minimal unitary dilation on the Schäffer's minimal space. In [41], we have constructed an isometric dilation in such scenario. We recall that result here. This will be frequently used throughout the paper.

Theorem 3.1 ([41], Theorem 3.4). *Let $T_1, \dots, T_n \in \mathcal{B}(\mathcal{H})$ be commuting contractions, $T = \prod_{i=1}^n T_i$ and $T'_i = \prod_{j \neq i} T_j$ for $1 \leq i \leq n$.*

(a) *If \mathcal{K} is the minimal isometric dilation space of T , then (T_1, \dots, T_n) possesses an isometric dilation (V_1, \dots, V_n) on \mathcal{K} with $V = \prod_{i=1}^n V_i$ being the minimal isometric dilation of T if and only if there are unique orthogonal projections P_1, \dots, P_n and unique commuting unitaries U_1, \dots, U_n in $\mathcal{B}(\mathcal{D}_T)$ with $\prod_{i=1}^n U_i = I_{\mathcal{D}_T}$ such that the following conditions are satisfied for each $i = 1, \dots, n$:*

- (1) $D_T T_i = P_i^\perp U_i^* D_T + P_i U_i^* D_T T$,
- (2) $P_i^\perp U_i^* P_j^\perp U_j^* = P_j^\perp U_j^* P_i^\perp U_i^*$,
- (3) $U_i P_i U_j P_j = U_j P_j U_i P_i$,
- (4) $D_T U_i P_i U_i^* D_T = D_{T_i}^2$,
- (5) $P_1 + U_1^* P_2 U_1 + U_1^* U_2^* P_3 U_2 U_1 + \dots + U_1^* U_2^* \dots U_{n-1}^* P_n U_{n-1} \dots U_2 U_1 = I_{\mathcal{D}_T}$.

(b) *Such an isometric dilation is minimal and unique in the sense that if (W_1, \dots, W_n) on \mathcal{K}_1 and (Y_1, \dots, Y_n) on \mathcal{K}_2 are two isometric dilations of (T_1, \dots, T_n) such that $W = \prod_{i=1}^n W_i$ and $Y = \prod_{i=1}^n Y_i$ are minimal isometric dilations of T on \mathcal{K}_1 and \mathcal{K}_2 respectively, then there is a unitary $\tilde{U} : \mathcal{K}_1 \rightarrow \mathcal{K}_2$ such that $(W_1, \dots, W_n) = (\tilde{U}^* Y_1 \tilde{U}, \dots, \tilde{U}^* Y_n \tilde{U})$.*

We have from [41] an existence of an isometric dilation on the minimal dilation space of T with an assumption of four out of five conditions of Theorem 3.1. This is a weaker version of Theorem 3.1 where we miss an appropriate converse part.

Theorem 3.2 ([41], Theorem 3.5). *Let $T_1, \dots, T_n \in \mathcal{B}(\mathcal{H})$ be commuting contractions, $T'_i = \prod_{j \neq i} T_j$ for all $1 \leq i \leq n$ and $T = \prod_{i=1}^n T_i$. Then (T_1, \dots, T_n) possesses an isometric dilation on the minimal isometric dilation space of T , if there are projections P_1, \dots, P_n and commuting unitaries U_1, \dots, U_n in $\mathcal{B}(\mathcal{D}_T)$ such that the following conditions hold for $i = 1, \dots, n$:*

- (1) $D_T T_i = P_i^\perp U_i^* D_T + P_i U_i^* D_T T$,
- (2) $P_i^\perp U_i^* P_j^\perp U_j^* = P_j^\perp U_j^* P_i^\perp U_i^*$,
- (3) $U_i P_i U_j P_j = U_j P_j U_i P_i$,
- (4) $D_T U_i P_i U_i^* D_T = D_{T_i}^2$.

Conversely, if (T_1, \dots, T_n) possesses an isometric dilation $(\hat{V}_1, \dots, \hat{V}_n)$ with $V = \prod_{i=1}^n V_i$ being the minimal isometric dilation of T , then there are unique projections P_1, \dots, P_n and unique commuting unitaries U_1, \dots, U_n in $\mathcal{B}(\mathcal{D}_T)$ satisfying the conditions (1) – (4) above.

The next few lemmas are useful for proving the main results of this Section. Note that the next lemma is proved in [16] in a more general setting.

Lemma 3.3. *Let \mathcal{H} be a Hilbert space and let $A, B \in \mathcal{B}(\mathcal{H})$ be commuting contractions. Then there is a unique contraction $C \in \mathcal{B}(\mathcal{D}_{AB})$ such that $D_B^2 A = D_{AB} C D_{AB}$.*

Proof. Let us consider the operator Y on $\mathcal{H} \oplus \mathcal{H}$ defined by the matrix $Y = \begin{bmatrix} A^*B^* & D_B^2A \\ 0 & AB \end{bmatrix}$. Since $\begin{bmatrix} B^* & D_B^2 \\ 0 & B \end{bmatrix}$ is a contraction by Proposition 2.1 and since A is a contraction, it follows that

$$Y = \begin{bmatrix} A^*B^* & D_B^2A \\ 0 & AB \end{bmatrix} = \begin{bmatrix} B^* & D_B^2 \\ 0 & B \end{bmatrix} \begin{bmatrix} A^* & 0 \\ 0 & A \end{bmatrix}$$

is a contraction. So, again by Proposition 2.1, there is a contraction $F \in \mathcal{B}(\mathcal{H})$ such that

$$D_B^2A = D_{AB}FD_{AB}.$$

Suppose the matrix of F with respect to the decomposition $\mathcal{H} = \mathcal{D}_{AB} \oplus \ker(D_{AB})$ is $\begin{bmatrix} F_{11} & F_{12} \\ F_{21} & F_{22} \end{bmatrix}$. Since $D_B^2A = D_{AB}FD_{AB}$, it follows that D_B^2A vanishes on $\ker(D_{AB})$. Therefore, the matrix of D_B^2A with respect to decomposition $\mathcal{H} = \mathcal{D}_{AB} \oplus \ker(D_{AB})$ is $\begin{bmatrix} D_B^2A & 0 \\ 0 & 0 \end{bmatrix}$. So, we have $D_B^2A = D_{AB}F_{11}D_{AB}$, where $F_{11} \in \mathcal{B}(\mathcal{D}_{AB})$ is a contraction.

For the uniqueness part, let there be two contractions $C, C_1 \in \mathcal{B}(\mathcal{D}_{AB})$ satisfying $D_B^2A = D_{AB}CD_{AB}$. Then $D_{AB}(C - C_1)D_{AB} = 0$ and consequently $\langle (C - C_1)D_{AB}h, D_{AB}h \rangle = 0$ for any $h \in \mathcal{H}$. This shows that $C = C_1$ and the proof is complete. ■

The following result can be found in the literature (e.g. [10]), though here we include a shorter proof for the sake of completeness.

Lemma 3.4. *Let $T_1, \dots, T_n \in \mathcal{B}(\mathcal{H})$ be commuting contractions. Let $T = \prod_{j=1}^n T_j$, and $T'_i = \prod_{j \neq i} T_j$ for $1 \leq i \leq n$. Then $D_T T_i = F_i D_T + F_i^* D_T T$ and $D_T T'_i = F'_i D_T + F_i^* D_T T$, where $F_i, F'_i \in \mathcal{B}(\mathcal{D}_T)$ are unique solutions of $D_{T'_i}^2 T_i = D_T X_i D_T$ and $D_{T'_i}^2 T'_i = D_T X'_i D_T$ respectively for each i .*

Proof. Let $G = F_i^* D_T T + F_i D_T - D_T T_i$. Then G is defined from $\mathcal{H} \rightarrow \mathcal{D}_T$. Since F_i, F'_i are unique solutions of $D_{T'_i}^2 T_i = D_T X_i D_T$ and $D_{T'_i}^2 T'_i = D_T X'_i D_T$ respectively for each i , we have that

$$D_T G = D_T F_i^* D_T T + D_T F_i D_T - D_T^2 T_i = (T_i'^* - T^* T_i) T + (T_i - T_i'^* T) - T_i + T^* T T_i = 0.$$

Now for any $h, h' \in \mathcal{H}$, $\langle Gh, D_T h' \rangle = \langle D_T Gh, h' \rangle = 0$. Hence $G = 0$ and consequently $D_T T_i = F_i D_T + F_i^* D_T T$. We can have a similar proof for $D_T T'_i = F'_i D_T + F_i^* D_T T$. ■

A refined Berger-Coburn-Lebow Model Theorem. In [15], Berger, Coburn and Lebow found the following remarkable factorization of a pure isometry. In Theorem 3.10 in [40], the first named author of this paper make a slight refinement of that result in the following manner.

Theorem 3.5 (Berger-Coburn-Lebow, [15]). *Let V_1, \dots, V_n be commuting isometries on \mathcal{H} such that $V = \prod_{i=1}^n V_i$ is a pure isometry. Let $V'_i = \prod_{j \neq i} V_j$ for $1 \leq i \leq n$. Then, there exist projections P_1, \dots, P_n and unitaries U_1, \dots, U_n in $\mathcal{B}(\mathcal{D}_{V^*})$ such that*

$$(V_1, \dots, V_n, V) \equiv (T_{P_1^\perp U_1 + z P_1 U_1}, \dots, T_{P_n^\perp U_n + z P_n U_n}, T_z) \text{ on } H^2(\mathcal{D}_{V^*}).$$

Moreover, $U_i^* P_i^\perp, P_i U_i$ are unique operators such that $V_i^* - V_i' V^* = D_{V^*} U_i^* P_i^\perp D_{V^*}$ and $V_i'^* - V_i V^* = D_{V^*} P_i U_i D_{V^*}$ respectively for each $i = 1, \dots, n$.

We state a lemma below and its proof will take cues from the proof of Theorem 3.1 from [41]. This will be useful.

Lemma 3.6. *Let $T_1, \dots, T_n \in \mathcal{B}(\mathcal{H})$ be commuting contractions, $T = \prod_{i=1}^n T_i$ and $T'_i = \prod_{j \neq i} T_j$ for $1 \leq i \leq n$. Let \mathcal{K} be the minimal isometric dilation space of T . If (T_1, \dots, T_n) possesses an isometric dilation (V_1, \dots, V_n) on \mathcal{K} with $V = \prod_{i=1}^n V_i$ being the minimal isometric dilation of T , then the following hold:*

- (1) $D_T T'_i = U_i P_i D_T + U_i P_i^\perp D_T T$,
- (2) $D_T P_i^\perp D_T = D_{T'_i}^2$.

Proof. Since (T_1, \dots, T_n) possesses an isometric dilation (V_1, \dots, V_n) on \mathcal{K} with $V = \prod_{i=1}^n V_i$ being the minimal isometric dilation of T , by part-(b) of Theorem 3.1, we can assume without loss of generality that

$$V_i = \begin{bmatrix} T_i & 0 & 0 & 0 & \dots \\ P_i U_i^* D_T & P_i^\perp U_i^* & 0 & 0 & \dots \\ 0 & P_i U_i^* & P_i^\perp U_i^* & 0 & \dots \\ 0 & 0 & P_i U_i^* & P_i^\perp U_i^* & \dots \\ \dots & \dots & \dots & \dots & \dots \end{bmatrix} \text{ and } V = \begin{bmatrix} T & 0 & 0 & 0 & \dots \\ D_T & 0 & 0 & 0 & \dots \\ 0 & I & 0 & 0 & \dots \\ 0 & 0 & I & 0 & \dots \\ \dots & \dots & \dots & \dots & \dots \end{bmatrix}, \text{ for } 1 \leq i \leq n,$$

following the construction of the isometric dilation as in Theorem 3.1 from [41]. Here P_1, \dots, P_n are unique projections and U_1, \dots, U_n are unique commuting unitaries on \mathcal{D}_T such that $\prod_{i=1}^n U_i = I$ and the conditions (1) – (5) of the Theorem 3.1 are satisfied. Let $V'_i = \prod_{j \neq i} V_j$ for any $i = 1, \dots, n$. Since V_1, \dots, V_n and $V = \prod_{i=1}^n V_i$ are isometries, we have $V'_i = V_i^* V$. Hence for any $1 \leq i \leq n$, we obtain

$$\begin{aligned} V'_i = V_i^* V &= \begin{bmatrix} T_i^* & D_T U_i P_i & 0 & 0 & \dots \\ 0 & U_i P_i^\perp & U_i P_i & 0 & \dots \\ 0 & 0 & U_i P_i^\perp & U_i P_i & \dots \\ 0 & 0 & 0 & U_i P_i^\perp & \dots \\ \dots & \dots & \dots & \dots & \dots \end{bmatrix} \begin{bmatrix} T & 0 & 0 & 0 & \dots \\ D_T & 0 & 0 & 0 & \dots \\ 0 & I & 0 & 0 & \dots \\ 0 & 0 & I & 0 & \dots \\ \dots & \dots & \dots & \dots & \dots \end{bmatrix} \\ &= \begin{bmatrix} T_i^* T_i + D_T U_i P_i D_T & 0 & 0 & 0 & \dots \\ U_i P_i^\perp D_T & U_i P_i & 0 & 0 & \dots \\ 0 & U_i P_i^\perp & U_i P_i & 0 & \dots \\ 0 & 0 & U_i P_i^\perp & U_i P_i & \dots \\ \dots & \dots & \dots & \dots & \dots \end{bmatrix} \\ &= \begin{bmatrix} T'_i & 0 & 0 & 0 & \dots \\ U_i P_i^\perp D_T & U_i P_i & 0 & 0 & \dots \\ 0 & U_i P_i^\perp & U_i P_i & 0 & \dots \\ 0 & 0 & U_i P_i^\perp & U_i P_i & \dots \\ \dots & \dots & \dots & \dots & \dots \end{bmatrix}, \end{aligned}$$

where the equation $T_i^* T_i + D_T U_i P_i D_T = T'_i$ follows from (a') in the (\Rightarrow) part of the proof of Theorem 3.1 from [41]. Thus, (2, 1) entry of $V'_i V = V V'_i$ gives us that $D_T T'_i = U_i P_i D_T + U_i P_i^\perp D_T T$ for any $i = 1, \dots, n$. Since V'_i is an isometry, the (1, 1) entry of $V'_i^* V'_i = I$ gives $D_T P_i^\perp D_T = I - T_i^* T_i = D_{T'_i}^2$. ■

We now show that the existence of an isometric dilation for a tuple of commuting contractions (T_1, \dots, T_n) on the minimal isometric dilation space of their product $T = \prod_{i=1}^n T_i$ is equivalent to the existence of a minimal isometric dilation for (T_1^*, \dots, T_n^*) on the minimal isometric dilation space of T^* . This is one of the main results of this article.

Theorem 3.7. *Let $T_1, \dots, T_n \in \mathcal{B}(\mathcal{H})$ be commuting contractions and let \mathcal{K} be the minimal isometric dilation space for their product $T = \prod_{i=1}^n T_i$. Then (T_1, \dots, T_n) possesses an isometric dilation (V_1, \dots, V_n) on \mathcal{K} with $V = \prod_{i=1}^n V_i$ being the minimal isometric dilation of T if and only if (T_1^*, \dots, T_n^*) possesses an isometric dilation (Y_1, \dots, Y_n) on \mathcal{K}_* with $Y = \prod_{i=1}^n Y_i$ being the minimal isometric dilation of T^* , where \mathcal{K}_* is the minimal isometric dilation space for T^* .*

Proof. It suffices to prove one side of the theorem, because, the other side follows by an analogous argument. So, we assume that (T_1, \dots, T_n) possesses an isometric dilation (V_1, \dots, V_n) on \mathcal{K} with $V = \prod_{i=1}^n V_i$ being the minimal isometric dilation of T . Then by Theorem 3.1, there are unique projections P_1, \dots, P_n and unique commuting unitaries U_1, \dots, U_n in $\mathcal{B}(\mathcal{D}_T)$ with $\prod_{i=1}^n U_i = I$ such that the following conditions are satisfied for $1 \leq i \leq n$:

- (1) $D_T T_i = P_i^\perp U_i^* D_T + P_i U_i^* D_T T$,
- (2) $P_i^\perp U_i^* P_j^\perp U_j^* = P_j^\perp U_j^* P_i^\perp U_i^*$,
- (3) $U_i P_i U_j P_j = U_j P_j U_i P_i$,
- (4) $D_T U_i P_i U_i^* D_T = D_{T_i}^2$,
- (5) $P_1 + U_1^* P_2 U_1 + U_1^* U_2^* P_3 U_2 U_1 + \dots + U_1^* U_2^* \dots U_{n-1}^* P_n U_{n-1} \dots U_2 U_1 = I_{\mathcal{D}_T}$.

To prove that (T_1^*, \dots, T_n^*) possesses an isometric dilation (Y_1, \dots, Y_n) on \mathcal{K}_* with $Y = \prod_{i=1}^n Y_i$ being the minimal isometric dilation of T^* . By Theorem 3.1 it suffices if we show the existence of unique projections Q_1, \dots, Q_n and unique commuting unitaries $\tilde{U}_1, \dots, \tilde{U}_n$ in $\mathcal{B}(\mathcal{D}_{T^*})$ with $\prod_{i=1}^n \tilde{U}_i = I$ satisfying

- (1') $D_{T^*} T_i^* = Q_i^\perp \tilde{U}_i^* D_{T^*} + Q_i \tilde{U}_i^* D_{T^*} T^*$,
- (2') $Q_i^\perp \tilde{U}_i^* Q_j^\perp \tilde{U}_j^* = Q_j^\perp \tilde{U}_j^* Q_i^\perp \tilde{U}_i^*$,
- (3') $\tilde{U}_i Q_i \tilde{U}_j Q_j = \tilde{U}_j Q_j \tilde{U}_i Q_i$,
- (4') $D_{T^*} \tilde{U}_i Q_i \tilde{U}_i^* D_{T^*} = D_{T_i^*}^2$,
- (5') $Q_1 + \tilde{U}_1^* Q_2 \tilde{U}_1 + \dots + \tilde{U}_1^* \dots \tilde{U}_{n-1}^* Q_n \tilde{U}_{n-1} \dots \tilde{U}_1 = I_{\mathcal{D}_{T^*}}$,

for $1 \leq i \leq n$. Let $T'_i = \prod_{j \neq i} T_j$ for $i = 1, \dots, n$. As in the proof of Theorem 3.1, if we denote $F_i = P_i^\perp U_i^*$ and $F'_i = U_i P_i$, then we have $F_i^* F_i + F'_i F_i'^* = I_{\mathcal{D}_T} = F_i F_i^* + F_i'^* F'_i$ and $F_i F_i' = F_i' F_i = 0$. Also, condition-(4) leads to $D_T F_i F_i'^* D_T = D_{T_i}^2$ for $1 \leq i \leq n$ and condition-(1) gives

$$D_T T_i = F_i D_T + F_i'^* D_T T. \quad (3.1)$$

Lemma 3.3 guarantees the existence of operators $G_1, \dots, G_n, G'_1, \dots, G'_n \in \mathcal{B}(\mathcal{D}_{T^*})$ satisfying

$$D_{T^*} G_i D_{T^*} = D_{T_i^*}^2 T_i^*, \text{ \& \ } D_{T^*} G'_i D_{T^*} = D_{T_i^*}^2 T_i'^* \quad (3.2)$$

and Lemma 3.4 further shows that they satisfy

$$D_{T^*} T_i^* = G_i D_{T^*} + G_i'^* D_{T^*} T^*. \quad (3.3)$$

Following the converse part of the proof of Theorem 3.1 we have from (a) and (a') the identities $D_{T_i'}^2 T_i = T_i - T_i'^* T = D_T F_i D_T$ and $D_{T_i'}^2 T_i' = T_i' - T_i'^* T = D_T F_i' D_T$ respectively. Again, Lemma 3.3 guarantees the uniqueness of such F_i and F_i' . Thus, we have that $F_i = P_i^\perp U_i^*$ and $F_i' = U_i P_i$ satisfy

$$D_T F_i D_T = D_{T_i'}^2 T_i \quad \& \quad D_T F_i' D_T = D_{T_i'}^2 T_i'. \quad (3.4)$$

From Lemma 3.4 we have that F_i, F'_i satisfy

$$D_T T'_i = F'_i D_T + F_i^* D_T T. \quad (3.5)$$

Also, from Lemma 3.6 we have that

$$D_T F_i F_i^* D_T = D_{T'_i}^2. \quad (3.6)$$

Set $\tilde{U}_i = G_i^* + G'_i$, $\tilde{U}'_i = G_i^* - G'_i$ and $Q_i = \frac{1}{2}(I - \tilde{U}_i^* \tilde{U}_i)$. We prove that \tilde{U}_i is a unitary and Q_i is a projection for each $1 \leq i \leq n$. Then we have

$$G_i G_i^* + G_i'^* G'_i = I_{\mathcal{D}_{T^*}} = G_i^* G_i + G'_i G_i'^* \quad \& \quad G_i G'_i = G'_i G_i = 0. \quad (3.7)$$

See the Appendix for a proof of (3.7). Now

$$\tilde{U}_i^* \tilde{U}_i = (G_i + G_i'^*)(G_i^* + G_i'^*) = G_i G_i^* + G_i G_i'^* + G_i'^* G_i^* + G_i'^* G_i' = G_i G_i^* + G_i'^* G_i' = I$$

and

$$\tilde{U}_i \tilde{U}_i^* = (G_i^* + G_i')(G_i + G_i'^*) = G_i^* G_i + G_i^* G_i'^* + G_i' G_i + G_i' G_i'^* = G_i^* G_i + G_i' G_i'^* = I.$$

Hence \tilde{U}_i is a unitary for each i . Observe that $G_i = (\tilde{U}_i^* + \tilde{U}_i'^*)/2$ and $G'_i = (\tilde{U}_i - \tilde{U}_i')/2$. Hence $G'_i G_i = 0$ implies that $\tilde{U}_i \tilde{U}_i'^* = \tilde{U}_i' \tilde{U}_i^*$ and $G_i G'_i = 0$ implies that $\tilde{U}_i^* \tilde{U}_i' = \tilde{U}_i'^* \tilde{U}_i$. Thus $Q_i = \frac{1}{2}(I - \tilde{U}_i^* \tilde{U}_i')$. Also, we have that

$$Q_i^2 = \frac{1}{2}(I - \tilde{U}_i^* \tilde{U}_i') \frac{1}{2}(I - \tilde{U}_i^* \tilde{U}_i') = \frac{1}{4}(I - 2\tilde{U}_i^* \tilde{U}_i' + I) = Q_i \quad \& \quad Q_i^* = \frac{1}{2}(I - \tilde{U}_i^* \tilde{U}_i') = Q_i.$$

Therefore, each Q_i is a projection. It follows from here that $G'_i = \tilde{U}_i Q_i$, $G_i = Q_i^\perp \tilde{U}_i^*$ for each i . Substituting the values of G_i, G'_i in (3.3) and (6.1) (see the Appendix) we obtain our desired identities (1') and (4') respectively. In order to have (2'), (3') it suffices if we prove $[G_i, G_j] = [G'_i, G'_j] = 0$ for all $1 \leq i, j \leq n$. We have from Theorem 3.1 that (V_1, \dots, V_n) is an isometric dilation of (T_1, \dots, T_n) . Note that

$$D_{T^*}(G_i^* T - T F_i) D_T = D_{T^*} G_i^* D_{T^*} T - T D_T F_i D_T = (T_i - T T_i'^*) T - T (T_i - T_i'^* T) = 0.$$

Since $T D_T = D_{T^*} T$ i.e. T maps \mathcal{D}_T into \mathcal{D}_{T^*} , we have that

$$G_i^* T|_{\mathcal{D}_T} = T F_i|_{\mathcal{D}_T}. \quad (3.8)$$

Using (3.4), (3.2) and the fact that $T^* D_{T^*} = D_T T^*$ we have that

$$D_T F_i' T^* D_{T^*} + D_{T^*} G_i D_{T^*} = D_{T'_i}^2 T_i'^* T^* + D_{T_i'^*}^2 T_i^* = T_i'^* T^* - T_i'^* T_i T_i'^* T^* + T_i^* - T_i'^* T_i'^* T_i^* = T_i^* D_{T^*}^2.$$

Therefore, for all $i = 1, \dots, n$ we have

$$D_{T^*} G_i = T_i^* D_{T^*} - D_T F_i' T^*. \quad (3.9)$$

Again, by the commutativity of $V_1, \dots, V_n, V'_1, \dots, V'_n$, the operators F_i, F_j, F'_i, F'_j satisfy

$$[F_i, F_j] = [F'_i, F'_j] = 0 \text{ and } [F_i^*, F'_j] = [F_j^*, F'_i]. \quad (3.10)$$

Now we prove that $[G_i, G_j] = [G'_i, G'_j] = 0$ and $[G_i, G_j^*] = [G_j, G_i^*]$ for all $1 \leq i < j \leq n$. First note that F'_i 's satisfy these commutator relations. The space \mathcal{H} can be decomposed into $\mathcal{H}_1 \oplus \mathcal{H}_2$ such that $V|_{\mathcal{H}_1}$ is a pure isometry and $V|_{\mathcal{H}_2}$ is a unitary. We show that $\mathcal{H}_1, \mathcal{H}_2$ are reducing subspaces for each V_i . If $V_i = \begin{bmatrix} A_i & B_i \\ C_i & D_i \end{bmatrix}$ and $V = \begin{bmatrix} V_{K1} & 0 \\ 0 & V_{K2} \end{bmatrix}$ with respect to the decomposition $\mathcal{H} = \mathcal{H}_1 \oplus \mathcal{H}_2$, then $V_i V = V V_i$ implies that $B_i V_{K2} = V_{K1} B_i$ and $C_i V_{K1} = V_{K2} C_i$. Therefore, for all $k \in \mathbb{N}$, $V_{K2}^* B_i^* =$

$B_i^* V_{K1}^{*k}$ and $V_{K1}^{*n} C_i^* = C_i^* V_{K2}^{*k}$. Now V_{K1} is a pure isometry and thus for each $h \in \mathcal{H}$, $\|B_i^* V_{K1}^{*k} h\| \rightarrow 0$. Again V_{K2} is a unitary and so we have $\|V_{K2}^{*k} B_i^* h\| = \|B_i^* h\|$. Therefore, $V_{K2}^{*k} B_i^* = B_i^* V_{K1}^{*k}$ would imply that $B_i = 0$. Similarly $C_i = 0$ for each $i = 1, \dots, n$. Hence let $V_i = V_{i1} \oplus V_{i2}$ for all $1 \leq i \leq n$. Now $V_{K1} = \prod_{i=1}^n V_{i1}$. Thus, by Theorem 3.5 there exist commuting unitaries $\tilde{U}_{11}, \dots, \tilde{U}_{n1}$ and projections Q_{11}, \dots, Q_{n1} in $\mathcal{B}(\mathcal{D}_{V_{K1}^*})$ such that $(V_{11}, \dots, V_{n1}) \equiv (M_{\tilde{U}_{11} Q_{11}^\perp + z \tilde{U}_{11} Q_{11}}, \dots, M_{\tilde{U}_{n1} Q_{n1}^\perp + z \tilde{U}_{n1} Q_{n1}})$ and that

$$D_{V_{K1}^*} Q_{i1}^\perp \tilde{U}_{i1}^* D_{V_{K1}^*} = D_{V_{i1}^*}^2 V_i^*, \quad (3.11)$$

$$D_{V_{K1}^*} \tilde{U}_{i1} Q_{i1} D_{V_{K1}^*} = D_{V_{i1}^*}^2 V_i'^* \quad (3.12)$$

for each $i = 1, \dots, n$. The fact that $(M_{\tilde{U}_{11} Q_{11}^\perp + z \tilde{U}_{11} Q_{11}}, \dots, M_{\tilde{U}_{n1} Q_{n1}^\perp + z \tilde{U}_{n1} Q_{n1}})$ is a commuting tuple gives

$$[\tilde{U}_{i1} Q_{i1}^\perp, \tilde{U}_{j1} Q_{j1}^\perp] = 0 = [\tilde{U}_{i1} Q_{i1}, \tilde{U}_{j1} Q_{j1}], [\tilde{U}_{i1} Q_{i1}^\perp, \tilde{U}_{j1} Q_{j1}] = [\tilde{U}_{j1} Q_{j1}^\perp, \tilde{U}_{i1} Q_{i1}], \quad (3.13)$$

for $1 \leq i \leq n$. Again, since $\prod_{i=1}^n M_{\tilde{U}_{i1} Q_{i1}^\perp + z \tilde{U}_{i1} Q_{i1}} = M_z$, we have that

$$Q_{11} + \tilde{U}_{11}^* Q_{21} \tilde{U}_{11} + \tilde{U}_{11}^* \tilde{U}_{21}^* Q_{31} \tilde{U}_{21} \tilde{U}_{11} + \dots + \tilde{U}_{11}^* \tilde{U}_{21}^* \dots \tilde{U}_{(n-1)1}^* Q_{n1} \tilde{U}_{(n-1)1} \dots \tilde{U}_{21} \tilde{U}_{11} = I_{\mathcal{D}_{V_{K1}^*}} \quad (3.14)$$

and that $\prod_{i=1}^n \tilde{U}_{i1} = I_{\mathcal{D}_{V_{K1}^*}}$. Also, V_{i2}, \dots, V_{n2} are unitary on \mathcal{H}_2 . It follows that

$$D_{V_i'^*}^2 = I_{\mathcal{H}} - \begin{bmatrix} V_{i1}' & 0 \\ 0 & V_{i2}' \end{bmatrix} \begin{bmatrix} V_{i1}'^* & 0 \\ 0 & V_{i2}'^* \end{bmatrix} = \begin{bmatrix} I_{\mathcal{H}_1} - V_{i1}' V_{i1}'^* & 0 \\ 0 & I_{\mathcal{H}_2} - V_{i2}' V_{i2}'^* \end{bmatrix} = \begin{bmatrix} I_{\mathcal{H}_1} - V_{i1}' V_{i1}'^* & 0 \\ 0 & 0 \end{bmatrix}.$$

Therefore, $D_{V_i'^*} = D_{V_{i1}'^*} \oplus 0$. Similarly we can prove that $D_{V_i^*} = D_{V_{i1}^*} \oplus 0$ for all $i = 1, \dots, n$, with respect to the above decomposition of \mathcal{H} . So $D_{V^*} = D_{V_{K1}^*} \oplus 0$. Substituting this we have from (3.11) and (3.12)

$$D_{V_i'^*}^2 V_i^* = (D_{V_{i1}'^*}^2 \oplus 0)(V_{i1}^* \oplus V_{i2}^*) = D_{V_{i1}'^*}^2 V_{i1}^* \oplus 0 = D_{V_{K1}^*} Q_{i1}^\perp \tilde{U}_{i1}^* D_{V_{K1}^*} \oplus 0 = D_{V^*} (Q_{i1}^\perp \tilde{U}_{i1}^* \oplus 0) D_{V^*}.$$

Hence, for $1 \leq i \leq n$ we have

$$D_{V_i'^*}^2 V_i^* = D_{V^*} (Q_{i1}^\perp \tilde{U}_{i1}^* \oplus 0) D_{V^*} \quad (3.15)$$

and similarly we have

$$D_{V_i^*}^2 V_i'^* = D_{V^*} (\tilde{U}_{i1} Q_{i1} \oplus 0) D_{V^*}. \quad (3.16)$$

Note that if (V, \mathcal{H}) is the minimal isometric dilation of a contraction (T, \mathcal{H}) then the dimensions of \mathcal{D}_{V^*} and \mathcal{D}_{T^*} are equal. Indeed, if $X : \mathcal{D}_{T^*} \rightarrow \mathcal{D}_{V^*}$ is defined as $XD_{T^*}h = D_{V^*}h$ for all $h \in \mathcal{H}$ and is extended continuously to the closure, then X is a unitary (see [13]). We briefly recall the proof here. Since V is the minimal isometric dilation of T , we have

$$\mathcal{H} = \overline{\text{span}}\{V^k h : k \geq 0, h \in \mathcal{H}\}.$$

Now, for $n \in \mathbb{N}$ and $h \in \mathcal{H}$ we have $D_{V^*}^2 V^n h = (I - VV^*)V^n h = 0$. Therefore, we have $D_{V^*} V^n h = 0$ for any $n \in \mathbb{N}$ and $h \in \mathcal{H}$. Thus, $\mathcal{D}_{V^*} = \overline{D_{V^*} \mathcal{H}} = \overline{D_{V^*} \mathcal{H}}$. In fact

$$\|D_{V^*} h\|^2 = \{(I - VV^*)h, h\} = \|h\|^2 - \|V^* h\|^2 = \|h\|^2 - \|T^* h\|^2 = \|D_{T^*} h\|^2.$$

Therefore, X as defined above is a unitary. We have that

$$D_{V^*} X G_i X^* D_{V^*} = D_{V_{i1}'^*}^2 V_i^* \text{ and } D_{V^*} X G_i' X^* D_{V^*} = D_{V_{i1}^*}^2 V_i'^*. \quad (3.17)$$

See the Appendix for a proof of (3.17). Thus, by the uniqueness argument as in Lemma 3.3, we have from (3.15), (3.16) and (3.17) that $X G_i X^* = Q_{i1}^\perp \tilde{U}_{i1}^* \oplus 0$ and $X G_i' X^* = \tilde{U}_{i1} Q_{i1} \oplus 0$. This is same

as saying that $XG_iX^* = Q_{iK}^\perp \tilde{U}_{iK}^*$ and $XG'_iX^* = \tilde{U}_{iK}Q_{iK}$, where $\tilde{U}_{iK} = \begin{bmatrix} \tilde{U}_{i1} & 0 \\ 0 & I_{\mathcal{H}_2} \end{bmatrix}$ and $Q_{iK} = Q_{i1} \oplus 0$ with respect to the decomposition $\mathcal{D}_{V_{K1}^*} \oplus 0$ of \mathcal{D}_{V^*} . Also, from (3.13) it is clear that

$$[\tilde{U}_{iK}Q_{iK}^\perp, \tilde{U}_{jK}Q_{jK}^\perp] = 0 = [\tilde{U}_{iK}Q_{iK}, \tilde{U}_{jK}Q_{jK}], [\tilde{U}_{iK}Q_{iK}^\perp, \tilde{U}_{jK}Q_{jK}] = [\tilde{U}_{jK}Q_{jK}^\perp, \tilde{U}_{iK}Q_{iK}].$$

Thus, we have $[G_i^*, G'_j] = [G_j^*, G'_i]$, $[G_i, G_i] = 0 = [G'_i, G'_j]$. So, we have $\tilde{U}_i\tilde{U}_j = \tilde{U}_j\tilde{U}_i$. From (3.14) and the fact that $\prod_{i=1}^n \tilde{U}_{i1} = I_{\mathcal{D}_{V_{K1}^*}}$, it follow that $\prod_{i=1}^n \tilde{U}_i = I_{\mathcal{D}_{T^*}}$. Since $G'_i = \tilde{U}_iQ_i$, $G_i = Q_i^\perp \tilde{U}_i^*$, (3.14) guarantees that condition-(5) holds. The uniqueness of Q_i and \tilde{U}_i for each i follows from the uniqueness of G_i and G'_i . The proof is now complete. \blacksquare

Now we are in a position to present the main unitary dilation theorem which is another main result of this paper.

Theorem 3.8. *Let $T_1, \dots, T_n \in \mathcal{B}(\mathcal{H})$ be commuting contractions, $T = \prod_{j=1}^n T_j$ and $T'_i = \prod_{j \neq i} T_j$ for $1 \leq i \leq n$.*

- (a) *If $\widetilde{\mathcal{K}_0}$ is a minimal unitary dilation space for T , then (T_1, \dots, T_n) possesses a unitary dilation (W_1, \dots, W_n) on $\widetilde{\mathcal{K}_0}$ with $W = \prod_{i=1}^n W_i$ being the minimal unitary dilation of T if and only if there exist unique projections P_1, \dots, P_n and unique commuting unitaries U_1, \dots, U_n in $\mathcal{B}(\mathcal{D}_T)$ with $\prod_{i=1}^n U_i = I$ such that the following hold for $i = 1, \dots, n$:*
 - (1) $D_T T_i = P_i^\perp U_i^* D_T + P_i U_i^* D_T T$,
 - (2) $P_i^\perp U_i^* P_j^\perp U_j^* = P_j^\perp U_j^* P_i^\perp U_i^*$,
 - (3) $U_i P_i U_j P_j = U_j P_j U_i P_i$,
 - (4) $D_T U_i P_i U_i^* D_T = D_{T_i}^2$,
 - (5) $P_1 + U_1^* P_2 U_1 + U_1^* U_2^* P_3 U_2 U_1 + \dots + U_1^* U_2^* \dots U_{n-1}^* P_n U_{n-1} \dots U_2 U_1 = I_{\mathcal{D}_T}$.
- (b) *Such a unitary dilation is minimal and unique in the sense that if (X_1, \dots, X_n) on $\widetilde{\mathcal{K}_1}$ is another unitary dilation of (T_1, \dots, T_n) such that the product $X = \prod_{i=1}^n X_i$ is a minimal unitary dilation of T , then there is a unitary $\rho : \widetilde{\mathcal{K}_0} \rightarrow \widetilde{\mathcal{K}_1}$ such that $(X_1, \dots, X_n) = (\rho * W_1 \rho, \dots, \rho * W_n \rho)$.*

Proof. Part-(a). (The \Leftarrow part). Suppose there are unique projections P_1, \dots, P_n and unique commuting unitaries U_1, \dots, U_n in $\mathcal{B}(\mathcal{D}_T)$ with $\prod_{i=1}^n U_i = I$ satisfying the operator identities (1) – (5) for all $1 \leq i \leq n$. We explicitly construct a unitary dilation (W_1, \dots, W_n) on $\widetilde{\mathcal{K}_0}$ of (T_1, \dots, T_n) such that the product $W = \prod_{i=1}^n W_i$ becomes a minimal unitary dilation of the product of the contractions $T = \prod_{i=1}^n T_i$. Evidently, by Theorem 3.1, (T_1, \dots, T_n) possesses an isometric dilation on the minimal isometric dilation space \mathcal{K}_0 of T , especially when $\mathcal{K}_0 = \mathcal{H} \oplus l^2(\mathcal{D}_T)$, the Schäffer's minimal isometric dilation space of T , (T_1, \dots, T_n) dilates to a tuple of commuting isometries (V_1, \dots, V_n) on \mathcal{K}_0 , where

$$V_i = \begin{bmatrix} T_i & 0 & 0 & 0 & \dots \\ P_i U_i^* D_T & P_i^\perp U_i^* & 0 & 0 & \dots \\ 0 & P_i U_i^* & P_i^\perp U_i^* & 0 & \dots \\ 0 & 0 & P_i U_i^* & P_i^\perp U_i^* & \dots \\ \dots & \dots & \dots & \dots & \dots \end{bmatrix}, \quad 1 \leq i \leq n,$$

and $\prod_{i=1}^n V_i = V$ is the Schäffer's minimal isometric dilation of T . Again, since (T_1, \dots, T_n) possesses an isometric dilation (V_1, \dots, V_n) on \mathcal{H}_0 with $\prod_{i=1}^n V_i = V$ being the minimal isometric dilation of T , it follows from Theorem 3.7 that (T_1^*, \dots, T_n^*) possesses an isometric dilation (Y_1, \dots, Y_n) on the minimal isometric dilation space \mathcal{H}_* of T^* with $\prod_{i=1}^n Y_i = Y$ being the minimal isometric dilation of T^* . Therefore, Theorem 3.1 guarantees the existence of a set of unique orthogonal projections Q_1, \dots, Q_n and unique commuting unitaries $\tilde{U}_1, \dots, \tilde{U}_n$ in $\mathcal{B}(\mathcal{D}_{T^*})$ with $\prod_{i=1}^n \tilde{U}_i = I$ such that the following identities hold for all $i = 1, \dots, n$:

$$\begin{aligned} (1') \quad & D_{T^*} T_i^* = Q_i^\perp \tilde{U}_i^* D_{T^*} + Q_i \tilde{U}_i^* D_{T^*} T^*, \\ (2') \quad & Q_i^\perp \tilde{U}_i^* Q_j^\perp \tilde{U}_j^* = Q_j^\perp \tilde{U}_j^* Q_i^\perp \tilde{U}_i^*, \\ (3') \quad & \tilde{U}_i Q_i \tilde{U}_j Q_j = \tilde{U}_j Q_j \tilde{U}_i Q_i, \\ (4') \quad & D_{T^*} \tilde{U}_i Q_i \tilde{U}_i^* D_{T^*} = D_{T_i^*}^2, \\ (5') \quad & Q_1 + \tilde{U}_1^* Q_2 \tilde{U}_1 + \dots + \tilde{U}_1^* \dots \tilde{U}_{n-1}^* Q_n \tilde{U}_{n-1} \dots \tilde{U}_1 = I_{\mathcal{D}_{T^*}}. \end{aligned}$$

Since (V_1, \dots, V_n) is an isometric dilation of (T_1, \dots, T_n) on the minimal isometric dilation space of T with $\prod_{i=1}^n V_i = V$ being the minimal isometric dilation of T , the (\Rightarrow) part of the Theorem 3.1 is applicable. From conditions (a) and (a') in the proof of (\Rightarrow) part of the Theorem 3.1, we have the identities $D_{T_i'}^2 T_i = T_i - T_i'^* T = D_T F_i D_T$ and $D_{T_i'}^2 T_i' = T_i' - T_i'^* T = D_T F_i' D_T$ respectively, where $F_i = P_i^\perp U_i^*$ and $F_i' = U_i P_i$. Thus, for $1 \leq i \leq n$ we have

$$D_T U_i P_i D_T = T_i' - T_i'^* T = D_{T_i'}^2 T_i' \quad \& \quad D_T P_i^\perp U_i^* D_T = T_i - T_i'^* T = D_{T_i'}^2 T_i. \quad (3.18)$$

An analogous argument holds for (T_1^*, \dots, T_n^*) and thus we have

$$D_{T^*} \tilde{U}_i Q_i D_{T^*} = T_i'^* - T_i T^* = D_{T_i'^*}^2 T_i'^* \quad \& \quad D_{T^*} Q_i^\perp \tilde{U}_i D_{T^*} = T_i^* - T_i'^* T^* = D_{T_i'^*}^2 T_i^*. \quad (3.19)$$

It is well-known from Sz.-Nagy-Foias theory (see [13]) that any two minimal unitary dilations of a contraction are unitarily equivalent. Thus, without loss of generality we consider the Schäffer's minimal unitary dilation space \mathcal{H}_0 of T , where

$$\mathcal{H}_0 = l^2(\mathcal{D}_T) \oplus \mathcal{H} \oplus l^2(\mathcal{D}_{T^*}) = \dots \oplus \mathcal{D}_T \oplus \mathcal{D}_T \oplus \mathcal{D}_T \oplus \mathcal{H} \oplus \mathcal{D}_{T^*} \oplus \mathcal{D}_{T^*} \oplus \mathcal{D}_{T^*} \oplus \dots$$

and construct a unitary dilation on \mathcal{H}_0 for (T_1, \dots, T_n) . Let us define W_1, \dots, W_n on $\mathcal{H}_0 = l^2(\mathcal{D}_T) \oplus \mathcal{H} \oplus l^2(\mathcal{D}_{T^*})$ by

$$W_i = \left[\begin{array}{cccc|c|cccc} \ddots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \dots & P_i^\perp U_i^* & P_i U_i^* & 0 & 0 & 0 & 0 & 0 \dots \\ \dots & 0 & P_i^\perp U_i^* & P_i U_i^* & 0 & 0 & 0 & 0 \dots \\ \dots & 0 & 0 & P_i^\perp U_i^* & P_i U_i^* D_T & -P_i U_i^* T^* & 0 & 0 \dots \\ \hline \dots & 0 & 0 & 0 & T_i & D_{T^*} \tilde{U}_i Q_i & 0 & 0 \dots \\ \hline & & & 0 & 0 & \tilde{U}_i Q_i^\perp & \tilde{U}_i Q_i & 0 \dots \\ & & & & & 0 & \tilde{U}_i Q_i^\perp & \tilde{U}_i Q_i \dots \\ & & & & & 0 & 0 & \tilde{U}_i Q_i^\perp \dots \\ & & & & & \vdots & \vdots & \vdots \ddots \end{array} \right], \quad 1 \leq i \leq n.$$

We prove that (W_1, \dots, W_n) is a unitary dilation of (T_1, \dots, T_n) . Note that the spaces $\mathcal{H}_0' = l^2(\mathcal{D}_T) \oplus \mathcal{H}$ and $\mathcal{H}_0 = \mathcal{H} \oplus l^2(\mathcal{D}_T)$ are isomorphic by the canonical unitary that maps $\xi \oplus h$ to $h \oplus \xi$, where $\xi \in l^2(\mathcal{D}_T)$ and $h \in \mathcal{H}$. Hence, this canonically gives a unitary

$$\phi : l^2(\mathcal{D}_T) \oplus \mathcal{H} \oplus l^2(\mathcal{D}_{T^*}) \rightarrow \mathcal{H}_0 \oplus l^2(\mathcal{D}_{T^*}). \quad (3.20)$$

Now if $\tilde{W}_i = \phi W_i \phi^*$ on $\mathcal{H}_0 \oplus l^2(\mathcal{D}_{T^*})$ is the replica of W_i for each i , then it suffices to show that $(\tilde{W}_1, \dots, \tilde{W}_n)$ is a unitary dilation of (T_1, \dots, T_n) . It is evident that with respect to the decomposition $\mathcal{H}_0 \oplus l^2(\mathcal{D}_{T^*})$,

$$\tilde{W}_i = \phi W_i \phi^* = \begin{bmatrix} V_i & D_i \\ 0 & E_i \end{bmatrix}, \quad (1 \leq i \leq n) \quad (3.21)$$

where $D_i : l^2(\mathcal{D}_{T^*}) \rightarrow \mathcal{H}_0$ and $E_i : l^2(\mathcal{D}_{T^*}) \rightarrow l^2(\mathcal{D}_{T^*})$ are the following operators:

$$D_i = \begin{bmatrix} D_{T^*} \tilde{U}_i Q_i & 0 & 0 & \cdots \\ -P_i U_i^* T^* & 0 & 0 & \cdots \\ 0 & 0 & 0 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix} \quad \& \quad E_i = \begin{bmatrix} \tilde{U}_i Q_i^\perp & \tilde{U}_i Q_i & 0 & \cdots \\ 0 & \tilde{U}_i Q_i^\perp & \tilde{U}_i Q_i & \cdots \\ 0 & 0 & \tilde{U}_i Q_i^\perp & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix}.$$

Evidently $\tilde{W}_i|_{\mathcal{H}_0} = V_i$ for each i and thus it suffices to show that $(\tilde{W}_1, \dots, \tilde{W}_n)$ is a commuting tuple of unitaries, because, then $(\tilde{W}_1, \dots, \tilde{W}_n)$ becomes a unitary extension of (V_1, \dots, V_n) and hence a unitary dilation of (T_1, \dots, T_n) .

Step 1. First we prove that $(\tilde{W}_1, \dots, \tilde{W}_n)$ is a commuting tuple. For each i, j we have,

$$\tilde{W}_i \tilde{W}_j = \begin{bmatrix} V_i V_j & V_i D_j + D_i E_j \\ 0 & E_i E_j \end{bmatrix} \quad \text{and} \quad \tilde{W}_j \tilde{W}_i = \begin{bmatrix} V_j V_i & V_j D_i + D_j E_i \\ 0 & E_j E_i \end{bmatrix}.$$

Thus, $\tilde{W}_i \tilde{W}_j = \tilde{W}_j \tilde{W}_i$ if and only if the following conditions hold:

- (i) $V_i V_j = V_j V_i$,
- (ii) $V_i D_j + D_i E_j = V_j D_i + D_j E_i$,
- (iii) $E_i E_j = E_j E_i$.

The condition-(i) follows from Theorem 3.1. We prove condition-(iii). First we simplify condition-(2') in the statement of this theorem using condition-(3') and the fact that $U_i U_j = U_j U_i$. We have $\tilde{U}_i^* Q_j \tilde{U}_j^* + Q_i \tilde{U}_i^* \tilde{U}_j^* = \tilde{U}_j^* Q_i \tilde{U}_i^* + Q_j \tilde{U}_j^* \tilde{U}_i^*$. Since \tilde{U}_i, \tilde{U}_j are commuting unitaries, this further gives

$$\tilde{U}_i^* Q_j \tilde{U}_i + Q_i = \tilde{U}_j^* Q_i \tilde{U}_j + Q_j. \quad (3.22)$$

The condition $\tilde{U}_i^* Q_j \tilde{U}_i + Q_i = \tilde{U}_j^* Q_i \tilde{U}_j + Q_j \leq I$ then follows from condition-(5'). Note that E_i on $l^2(\mathcal{D}_{T^*})$ is equivalent to $T_{\tilde{U}_i Q_i^\perp + \tilde{U}_i Q_i}$ on $H^2(\mathcal{D}_{T^*})$. Hence by Lemma 2.5, (iii) holds. The proof of (ii), i.e.

$$V_i D_j + D_i E_j = V_j D_i + D_j E_i \quad (3.23)$$

is technical and is given in the Appendix. Therefore, $(\tilde{W}_1, \dots, \tilde{W}_n)$ is a commuting tuple.

Step 2. Now we prove that \tilde{W}_i is a unitary for each $i = 1, 2, \dots, n$. First we note that $\tilde{W}_i^* \tilde{W}_i = I$ if and only if

$$\begin{bmatrix} V_i^* V_i & V_i^* D_i \\ D_i^* V_i & D_i^* D_i + E_i^* E_i \end{bmatrix} = \begin{bmatrix} I & 0 \\ 0 & I \end{bmatrix}$$

which holds if and only if

- (a) $V_i^* V_i = I_{\mathcal{H}_0}$,
- (b) $V_i^* D_i = 0$ on $l^2(\mathcal{D}_{T^*})$,
- (c) $D_i^* D_i + E_i^* E_i = I_{l^2(\mathcal{D}_{T^*})}$.

Clearly, (a) follows as each V_i is an isometry. For showing (b) note that

$$V_i^* D_i = \begin{bmatrix} T_i^* D_{T^*} \tilde{U}_i Q_i - D_T U_i P_i U_i^* T^* & 0 & 0 & \dots \\ -U_i P_i^\perp P_i U_i^* T^* & 0 & 0 & \dots \\ 0 & 0 & 0 & \dots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix}.$$

Clearly, $U_i P_i^\perp P_i U_i^* T^* = U_i (I - P_i) P_i U_i^* T^* = 0$ as P_i is a projection. Now

$$(T_i^* D_{T^*} \tilde{U}_i Q_i - D_T U_i P_i U_i^* T^*) D_{T^*} = T_i^* (T_i'^* - T_i T^*) - D_T U_i P_i U_i^* D_T T^* = T_i^* (T_i'^* - T_i T^*) - D_{T_i}^2 T^* = 0,$$

where the first and the second equality follow from (3.19) and condition-(4) of the theorem respectively. This proves (b). For proving (c) we observe that $Q_i Q_i^\perp = 0$, $\tilde{U}_i^* \tilde{U}_i = I$ and $Q_i + Q_i^\perp = I$. Further the $(1, 1)$ entry of $D_i^* D_i + E_i^* E_i$ with respect to the decomposition $\mathcal{D}_{T^*} \oplus \mathcal{D}_{T^*} \oplus \dots$ is $Q_i \tilde{U}_i^* D_{T^*}^2 \tilde{U}_i Q_i + T U_i P_i U_i^* T^* + Q_i^\perp \tilde{U}_i^* \tilde{U}_i Q_i^\perp$. So, we have

$$\begin{aligned} Q_i \tilde{U}_i^* D_{T^*}^2 \tilde{U}_i Q_i + T U_i P_i U_i^* T^* + Q_i^\perp \tilde{U}_i^* \tilde{U}_i Q_i^\perp &= Q_i \tilde{U}_i^* \tilde{U}_i Q_i - Q_i \tilde{U}_i^* T T^* \tilde{U}_i Q_i + T U_i P_i U_i^* T^* + Q_i^\perp \\ &= Q_i + Q_i^\perp \quad [\text{by (6.2)}] \\ &= I. \end{aligned}$$

Thus, (c) holds. We now prove $\tilde{W}_i \tilde{W}_i^* = I$. Note that $\tilde{W}_i \tilde{W}_i^* = I$ if and only if

$$\begin{bmatrix} V_i V_i^* + D_i D_i^* & D_i E_i^* \\ E_i D_i^* & E_i E_i^* \end{bmatrix} = \begin{bmatrix} I & 0 \\ 0 & I \end{bmatrix} \quad (3.24)$$

and a proof to this is similar to that of $\tilde{W}_i^* \tilde{W}_i = I$ and is given in the Appendix. Thus, \tilde{W}_i is a unitary for each $i = 1, \dots, n$. Hence, $(\tilde{W}_1, \dots, \tilde{W}_n)$ is a unitary dilation of (T_1, \dots, T_n) .

Step 3. Now we prove that $\prod_{i=1}^n W_i = W$, where W is the Schäffer's minimal unitary dilation of T , i.e.

$$W = \left[\begin{array}{cccc|c|cccc} \ddots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \dots & 0 & I & 0 & 0 & 0 & 0 & 0 & \dots \\ \dots & 0 & 0 & I & 0 & 0 & 0 & 0 & \dots \\ \dots & 0 & 0 & 0 & D_T & -T^* & 0 & 0 & \dots \\ \hline \dots & 0 & 0 & 0 & T & D_{T^*} & 0 & 0 & \dots \\ \hline \dots & 0 & 0 & 0 & 0 & 0 & I & 0 & \dots \\ \dots & 0 & 0 & 0 & 0 & 0 & 0 & I & \dots \\ \dots & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{array} \right]. \quad (3.25)$$

Let $\tilde{W} = \phi W \phi^*$ on $\mathcal{K}_0 \oplus l^2(\mathcal{D}_{T^*})$ where, ϕ is as in (3.20). Then, evidently

$$\tilde{W} = \begin{bmatrix} V & D \\ 0 & E \end{bmatrix}, \quad (1 \leq k \leq n) \quad (3.26)$$

where, V is the Schäffer's minimal isometric dilation of T , $D : l^2(\mathcal{D}_{T^*}) \rightarrow \mathcal{H}_0$ and $E : l^2(\mathcal{D}_{T^*}) \rightarrow l^2(\mathcal{D}_{T^*})$ are the following operators:

$$D = \begin{bmatrix} D_{T^*} & 0 & 0 & \cdots \\ -T^* & 0 & 0 & \cdots \\ 0 & 0 & 0 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix} \quad \& \quad E = \begin{bmatrix} 0 & I & 0 & \cdots \\ 0 & 0 & I & \cdots \\ 0 & 0 & 0 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix}. \quad (3.27)$$

It follows from (3.21) that,

$$\prod_{i=1}^n \tilde{W}_i = \prod_{i=1}^n \phi W_i \phi^* = \phi \left(\prod_{i=1}^n W_i \right) \phi^*.$$

Therefore, $W = \prod_{i=1}^n W_i$ if and only if $\tilde{W} = \phi W \phi^* = \prod_{i=1}^n \tilde{W}_i$. Hence it suffices to prove that $\tilde{W} = \prod_{i=1}^n \tilde{W}_i$. It was proved by Bercovici, Douglas and Foias in [14], all terms that are involved in condition-(5) (and in condition-(5')) are mutually orthogonal projections. This is because, the sum of projections is a projection if and only if they are mutually orthogonal. Suppose, for any $1 \leq k \leq n$,

$$\underline{T}_k = T_1 \dots T_k, \underline{U}_k = U_1 \dots U_k, \underline{P}_k = P_1 + \underline{U}_1^* P_2 \underline{U}_1 + \dots + \underline{U}_{k-1}^* P_k \underline{U}_{k-1},$$

$$\tilde{\underline{U}}_k = \tilde{U}_1 \dots \tilde{U}_k, \underline{Q}_k = Q_1 + \tilde{\underline{U}}_1^* Q_2 \tilde{\underline{U}}_1 + \dots + \tilde{\underline{U}}_{k-1}^* Q_k \tilde{\underline{U}}_{k-1},$$

where $U_k, P_k \in \mathcal{B}(\mathcal{D}_T)$ are as in the hypothesis of this theorem and $\tilde{U}_k, Q_k \in \mathcal{B}(\mathcal{D}_{T^*})$ are as obtained in the beginning of the proof satisfying conditions (1') – (5'). Evidently each \underline{U}_k is a unitary and each \underline{P}_k is a projection. Further, it follows from the hypothesis of this theorem and from the condition (5') that,

$$\underline{T}_n = T, \underline{U}_n = I_{\mathcal{D}_T}, \underline{P}_n = I_{\mathcal{D}_T}, \tilde{\underline{U}}_n = I_{\mathcal{D}_{T^*}}, \quad \text{and} \quad \underline{Q}_n = I_{\mathcal{D}_{T^*}}. \quad (3.28)$$

Define

$$\underline{V}_k = \begin{bmatrix} \underline{T}_k & 0 & 0 & 0 & \cdots \\ \underline{P}_k \underline{U}_k^* D_T & \underline{P}_k^\perp \underline{U}_k^* & 0 & 0 & \cdots \\ 0 & \underline{P}_k \underline{U}_k^* & \underline{P}_k^\perp \underline{U}_k^* & 0 & \cdots \\ 0 & 0 & \underline{P}_k \underline{U}_k^* & \underline{P}_k^\perp \underline{U}_k^* & \cdots \\ \cdots & \cdots & \cdots & \cdots & \cdots \end{bmatrix}, \quad 1 \leq k \leq n.$$

Therefore, we have from (3.28) that $\underline{V}_n = V$, where V is the Schäffer's minimal isometric dilation of T . Note that for all $k = 1, \dots, n$, the operators V_k and \underline{V}_k have the following block- matrix form with respect to the decomposition $\mathcal{H}_0 = \mathcal{H} \oplus l^2(\mathcal{D}_T)$:

$$V_k = \begin{bmatrix} T_k & 0 \\ C_k & S_k \end{bmatrix}, \quad \underline{V}_k = \begin{bmatrix} \underline{T}_k & 0 \\ \underline{C}_k & \underline{S}_k \end{bmatrix},$$

where,

$$C_k = \begin{bmatrix} P_k U_k^* D_T \\ 0 \\ 0 \\ \vdots \end{bmatrix} : \mathcal{H} \rightarrow l^2(\mathcal{D}_T), \quad S_k = \begin{bmatrix} P_k^\perp U_k^* & 0 & 0 & \cdots \\ P_k U_k^* & P_k^\perp U_k^* & 0 & \cdots \\ 0 & P_k U_k^* & P_k^\perp U_k^* & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix} : l^2(\mathcal{D}_T) \rightarrow l^2(\mathcal{D}_T)$$

$$\underline{C}_k = \begin{bmatrix} \underline{P}_k \underline{U}_k^* D_T \\ 0 \\ 0 \\ \vdots \end{bmatrix} : \mathcal{H} \rightarrow l^2(\mathcal{D}_T), \quad \underline{S}_k = \begin{bmatrix} \underline{P}_k^\perp \underline{U}_k^* & 0 & 0 & \cdots \\ \underline{P}_k \underline{U}_k^* & \underline{P}_k^\perp \underline{U}_k^* & 0 & \cdots \\ 0 & \underline{P}_k \underline{U}_k^* & \underline{P}_k^\perp \underline{U}_k^* & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix} : l^2(\mathcal{D}_T) \rightarrow l^2(\mathcal{D}_T).$$

Let

$$\tilde{\underline{W}}_k = \begin{bmatrix} \underline{V}_k & \underline{D}_k \\ 0 & \underline{E}_k \end{bmatrix}, \quad (1 \leq k \leq n)$$

where $\underline{D}_k : l^2(\mathcal{D}_{T^*}) \rightarrow \mathcal{K}_0$ and $\underline{E}_k : l^2(\mathcal{D}_{T^*}) \rightarrow l^2(\mathcal{D}_{T^*})$ are the following operators:

$$\underline{D}_k = \begin{bmatrix} D_{T^*} \tilde{\underline{U}}_k \underline{Q}_k & 0 & 0 & \cdots \\ -\underline{P}_k \underline{U}_k^* T^* & 0 & 0 & \cdots \\ 0 & 0 & 0 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix} \quad \& \quad \underline{E}_k = \begin{bmatrix} \tilde{\underline{U}}_k \underline{Q}_k^\perp & \tilde{\underline{U}}_k \underline{Q}_k & 0 & \cdots \\ 0 & \tilde{\underline{U}}_k \underline{Q}_k^\perp & \tilde{\underline{U}}_k \underline{Q}_k & \cdots \\ 0 & 0 & \tilde{\underline{U}}_k \underline{Q}_k^\perp & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix}.$$

Again by (3.26), (3.27) and (3.28), we have

$$\tilde{\underline{W}}_n = \tilde{\underline{W}}. \quad (3.29)$$

We now show that $\tilde{\underline{W}}_k \tilde{\underline{W}}_{k+1} = \tilde{\underline{W}}_{k+1}$. Considering the block matrices of these operators with respect to the decomposition $\mathcal{K}_0 \oplus l^2(\mathcal{D}_{T^*})$ it suffices if we prove that

$$\begin{bmatrix} \underline{V}_k \underline{V}_{k+1} & \underline{V}_k \underline{D}_{k+1} + \underline{D}_k \underline{E}_{k+1} \\ 0 & \underline{E}_k \underline{E}_{k+1} \end{bmatrix} = \begin{bmatrix} \underline{V}_{k+1} & \underline{D}_{k+1} \\ 0 & \underline{E}_{k+1} \end{bmatrix}, \quad 1 \leq k \leq n-1.$$

From Step 3 in the proof of Theorem 3.1, we have that $\underline{V}_k \underline{V}_{k+1} = \underline{V}_{k+1}$ for $1 \leq k \leq n-1$. Again, considering the block matrices of \underline{V}_k , \underline{V}_{k+1} and \underline{V}_{k+1} with respect to decomposition $\mathcal{K}_0 = \mathcal{H} \oplus l^2(\mathcal{D}_T)$ we have $\underline{S}_{k+1} = \underline{S}_k \underline{S}_{k+1}$. This identity and a simple calculation lead to

$$\underline{P}_k \underline{U}_k^* \underline{P}_{k+1} \underline{U}_{k+1}^* = 0 \quad \& \quad \underline{P}_k^\perp \underline{U}_k^* \underline{P}_{k+1} \underline{U}_{k+1}^* + \underline{P}_k \underline{U}_k^* \underline{P}_{k+1}^\perp \underline{U}_{k+1}^* = \underline{P}_{k+1} \underline{U}_{k+1}^*. \quad (3.30)$$

Again, it is clear from the definition that $\tilde{\underline{U}}_{k+1} = \tilde{\underline{U}}_k \tilde{\underline{U}}_{k+1}$ and $\underline{Q}_{k+1} = \underline{Q}_k + \tilde{\underline{U}}_k^* \underline{Q}_{k+1} \tilde{\underline{U}}_k$. Further it can inductively be proved that

$$\underline{Q}_k + \tilde{\underline{U}}_k^* \underline{Q}_{k+1} \tilde{\underline{U}}_k = \underline{Q}_{k+1} + \tilde{\underline{U}}_{k+1}^* \underline{Q}_k \tilde{\underline{U}}_{k+1}. \quad (3.31)$$

Hence, it follows from Lemma 2.6 that $\underline{E}_{k+1} = \underline{E}_k \underline{E}_{k+1}$. Thus,

$$\tilde{\underline{U}}_k \underline{Q}_k \tilde{\underline{U}}_{k+1} \underline{Q}_{k+1} = 0 \quad (3.32)$$

and

$$\tilde{\underline{U}}_k \underline{Q}_k^\perp \tilde{\underline{U}}_{k+1} \underline{Q}_{k+1} + \tilde{\underline{U}}_k \underline{Q}_k \tilde{\underline{U}}_{k+1} \underline{Q}_{k+1}^\perp = \tilde{\underline{U}}_{k+1} \underline{Q}_{k+1}. \quad (3.33)$$

It remains to show that

$$\underline{V}_k \underline{D}_{k+1} + \underline{D}_k \underline{E}_{k+1} = \underline{D}_{k+1}. \quad (3.34)$$

A proof to this is technical and is given in the Appendix. Hence, for each k we have that $\tilde{\underline{W}}_k \tilde{\underline{W}}_{k+1} = \tilde{\underline{W}}_{k+1}$. Thus, recursively we have $\tilde{\underline{W}}_n = \prod_{i=1}^n \tilde{\underline{W}}_i$. It then follows from (3.29), that $\tilde{\underline{W}} = \tilde{\underline{W}}_n = \prod_{i=1}^n \tilde{\underline{W}}_i$, as required.

(The \Rightarrow part). Suppose $(\widehat{W}_1, \dots, \widehat{W}_n)$ is a unitary dilation of (T_1, \dots, T_n) on a minimal unitary dilation space \mathcal{K}' of T with $\prod_{i=1}^n \widehat{W}_i = \widehat{W}$ being the minimal unitary dilation of T . Then

$$\mathcal{K}' = \overline{\text{span}}\{\widehat{W}^n h : h \in \mathcal{H}, n \in \mathbb{Z}\}.$$

Let W as in (3.25) be the Schäffer's minimal unitary dilation of T on the Schäffer's minimal space $\widetilde{\mathcal{K}}_0$. So,

$$\widetilde{\mathcal{K}}_0 = \overline{\text{span}}\{W^n h : h \in \mathcal{H}, n \in \mathbb{Z}\}.$$

Therefore, the map $\tau : \widetilde{\mathcal{K}}_0 \rightarrow \mathcal{K}'$ defined by $\tau(W^n h) = \widehat{W}^n(h)$ is a unitary which is identity on \mathcal{H} . Thus \mathcal{H} is a reducing subspace for τ and consequently $\tau = \begin{bmatrix} I_{\mathcal{H}} & 0 \\ 0 & \tau_2 \end{bmatrix}$, for some unitary τ_2 , with respect to the decomposition $\mathcal{H} \oplus (\widetilde{\mathcal{K}}_0 \ominus \mathcal{H})$ of $\widetilde{\mathcal{K}}_0$ and $\mathcal{H} \oplus (\mathcal{K}' \ominus \mathcal{H})$ of \mathcal{K}' . Evidently, $(\tau^* \widehat{W}_1 \tau, \dots, \tau^* \widehat{W}_n \tau)$ is a commuting tuple of unitaries dilating (T_1, \dots, T_n) on \mathcal{K}' with $\prod_{i=1}^n \tau^* \widehat{W}_i \tau = W$ being the Schäffer's minimal unitary dilation of T as in (3.25). It is clear from (3.25) that W has the following block-matrix form with respect to the decomposition $\widetilde{\mathcal{K}}_0 = l^2(\mathcal{D}_T) \oplus \mathcal{H} \oplus l^2(\mathcal{D}_T^*)$:

$$\begin{bmatrix} * & * & * \\ 0 & T & * \\ 0 & 0 & * \end{bmatrix}.$$

Suppose $W_i = \tau^* \widehat{W}_i \tau$ for $i = 1, \dots, n$. Then Lemma 2.2 tells us that with respect to the decomposition $l^2(\mathcal{D}_T) \oplus \mathcal{H} \oplus l^2(\mathcal{D}_T^*)$ of $\widetilde{\mathcal{K}}_0$, each W_j has a matrix representation of the form

$$\begin{bmatrix} * & * & * \\ 0 & T_j & * \\ 0 & 0 & * \end{bmatrix}.$$

It is obvious from the block-matrix form that $\mathcal{H} \oplus l^2(\mathcal{D}_T)$ is an invariant subspace for each W_j and for W . Set $V_j := W_j|_{\mathcal{H} \oplus l^2(\mathcal{D}_T)} = \begin{bmatrix} T_j & 0 \\ * & * \end{bmatrix}$ and $V = W|_{\mathcal{H} \oplus l^2(\mathcal{D}_T)}$. Then, V is the Schäffer's minimal isometric dilation of T on the Schäffer's minimal isometric dilation space $\mathcal{H} \oplus l^2(\mathcal{D}_T)$ and (V_1, \dots, V_n) is a commuting isometric tuple that dilates (T_1, \dots, T_n) . Since $\prod_{i=1}^n W_i = W$, a simple computation shows that $\prod_{i=1}^n V_i = V$. Thus, (V_1, \dots, V_n) is an isometric dilation of (T_1, \dots, T_n) with $\prod_{i=1}^n V_i = V$ being the minimal isometric dilation of T . Therefore, it follows from Theorem 3.1 that there are unique orthogonal projections P_1, \dots, P_n and unique commuting unitaries U_1, \dots, U_n in $\mathcal{B}(\mathcal{D}_P)$ with $\prod_{i=1}^n U_i = I_{\mathcal{D}_P}$ satisfying the conditions (1) – (5) of this theorem.

Part-(b). Since $W = \prod_{i=1}^n W_i$ on $\widetilde{\mathcal{K}}_0$ is a minimal unitary dilation of $T = \prod_{i=1}^n T_i$, it follows from the definition of minimality that the dilation (W_1, \dots, W_n) of (T_1, \dots, T_n) is minimal. Thus,

$$\widetilde{\mathcal{K}}_0 = \overline{\text{Span}}\{W^n h : h \in \mathcal{H} \text{ \& } n \in \mathbb{Z}\} = \overline{\text{Span}}\{W_1^{d_1} \dots W_n^{d_n} h : h \in \mathcal{H} \text{ \& } d_1, \dots, d_n \in \mathbb{Z}\}.$$

If (X_1, \dots, X_n) on $\widetilde{\mathcal{K}}_1$ is a unitary dilation of (T_1, \dots, T_n) with the product $X = \prod_{i=1}^n X_i$ being a minimal unitary dilation of T , then this is a minimal unitary dilation of (T_1, \dots, T_n) and consequently we have

$$\widetilde{\mathcal{K}}_1 = \overline{\text{Span}}\{X^n h : h \in \mathcal{H} \text{ \& } n \in \mathbb{Z}\} = \overline{\text{Span}}\{X_1^{d_1} \dots X_n^{d_n} h : h \in \mathcal{H} \text{ \& } d_1, \dots, d_n \in \mathbb{Z}\}.$$

Evidently, the unitary $\rho : \widetilde{\mathcal{H}}_0 \rightarrow \widetilde{\mathcal{H}}_1$ that maps $W^n h$ to $X^n h$ also maps $W_1^{d_1} \dots W_n^{d_n} h$ to $X_1^{d_1} \dots X_n^{d_n} h$ and this gives $X_i = \rho^* W_i \rho$ for $1 \leq i \leq n$ so that the product $X = \prod_{i=1}^n X_i = \rho^* W \rho$. The proof is now complete. ■

Theorem 3.1 shows that a tuple of commuting contraction (T_1, \dots, T_n) admits an isometric dilation (V_1, \dots, V_n) on the minimal isometric dilation space of $T = \prod_{i=1}^n T_i$ if we assume the exact five conditions as in Theorem 3.8. Now being a subnormal tuple, (V_1, \dots, V_n) always extends to a commuting unitary tuple which must be a unitary dilation of (T_1, \dots, T_n) . The main achievement here is that we found such a unitary extension on the minimal unitary dilation space of T without assuming any additional conditions. Needless to mention that Theorem 3.7 plays the central role in determining this.

Remark 3.9. The minimal unitary dilation of Theorem 3.8 is not unconditional even for $n = 2$, though Ando's theorem tells us that every pair of commuting contractions (T_1, T_2) dilates to a pair of commuting unitaries without any conditions on (T_1, T_2) . As we have seen in Example 3.6 in [41] that if we choose

$$T_1 = \begin{pmatrix} 0 & 0 & 0 \\ 1/3 & 0 & 0 \\ 0 & 1/3\sqrt{3} & 0 \end{pmatrix} \quad \& \quad T_2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ -1/\sqrt{3} & 0 & 0 \end{pmatrix},$$

then condition-(4) of Theorem 3.8 is not satisfied and consequently (T_1, T_2) does not dilate to a pair of commuting unitaries (W_1, W_2) on the minimal unitary dilation space of $T_1 T_2$ with $W_1 W_2$ being the minimal unitary dilation of $T_1 T_2$.

We conclude this Section with the following analogue of Theorem 3.7 for unitary dilation.

Theorem 3.10. *Let $T_1, \dots, T_n \in \mathcal{B}(\mathcal{H})$ be commuting contractions and let $\widetilde{\mathcal{H}}$ be the minimal unitary dilation space for $T = \prod_{i=1}^n T_i$. Then (T_1, \dots, T_n) possesses a unitary dilation (W_1, \dots, W_n) on $\widetilde{\mathcal{H}}$ with $W = \prod_{i=1}^n W_i$ being the minimal unitary dilation of T if and only if (T_1^*, \dots, T_n^*) possesses a unitary dilation (Z_1, \dots, Z_n) on $\widetilde{\mathcal{H}}_*$ with $Z = \prod_{i=1}^n Z_i$ being the minimal unitary dilation of T^* , where $\widetilde{\mathcal{H}}_*$ is the minimal unitary dilation space for T^* .*

Proof. Suppose (T_1, \dots, T_n) possesses a unitary dilation (W_1, \dots, W_n) on $\widetilde{\mathcal{H}}$ with W being the minimal unitary dilation of T . Then (T_1, \dots, T_n) satisfies conditions (1) – (5) of Theorem 3.8 and hence by Theorem 3.1, it admits an isometric dilation (V_1, \dots, V_n) on the minimal isometric dilation space \mathcal{H} of T with $V = \prod_{i=1}^n V_i$ being the minimal isometric dilation of T . So, by Theorem 3.7, (T_1^*, \dots, T_n^*) possesses an isometric dilation (Y_1, \dots, Y_n) on the minimal isometric dilation space \mathcal{H}_* of T^* with $Y = \prod_{i=1}^n Y_i$ being the minimal isometric dilation of T^* . Again applying Theorem 3.1 on (T_1^*, \dots, T_n^*) we have that there are unique orthogonal projections Q_1, \dots, Q_n and unique commuting unitaries $\widetilde{U}_1, \dots, \widetilde{U}_n$ in $\mathcal{B}(\mathcal{D}_{T^*})$ with $\prod_{i=1}^n \widetilde{U}_i = I_{\mathcal{D}_{T^*}}$ satisfying the hypotheses of Theorem 3.1. Thus, with the same hypotheses we apply Theorem 3.8 to obtain a unitary dilation (Z_1, \dots, Z_n) of (T_1^*, \dots, T_n^*) on the minimal unitary dilation space $\widetilde{\mathcal{H}}_*$ of T^* . ■

4. MINIMAL UNITARY DILATION WHEN THE PRODUCT IS A C_0 CONTRACTION

In this Section, we show that only four out of five conditions of Theorem 3.8 are necessary and sufficient for the existence of a unitary dilation of a tuple of commuting contraction (T_1, \dots, T_n)

when the product $T = \prod_{i=1}^n T_i$ is a C_0 contraction. First we recall a pair of isometric dilation theorems in this setting from [41].

Theorem 4.1. *Let T_1, \dots, T_n be commuting contractions on a Hilbert space \mathcal{H} such that their product $T = \prod_{i=1}^n T_i$ is a C_0 contraction. Then (T_1, \dots, T_n) possesses an isometric dilation (V_1, \dots, V_n) on \mathcal{H} with $V = \prod_{i=1}^n V_i$ being a minimal isometric dilation of T if and only if there are unique orthogonal projections P_1, \dots, P_n and unique commuting unitaries U_1, \dots, U_n in $\mathcal{B}(\mathcal{D}_{T^*})$ with $\prod_{i=1}^n U_i = I_{\mathcal{D}_{T^*}}$ such that the following hold for $i = 1, \dots, n$:*

- (1) $D_{T^*} T_i^* = P_i^\perp U_i^* D_{T^*} + P_i U_i^* D_{T^*} T^*$,
- (2) $P_i^\perp U_i^* P_j^\perp U_j^* = P_j^\perp U_j^* P_i^\perp U_i^*$,
- (3) $U_i P_i U_j P_j = U_j P_j U_i P_i$,
- (4) $P_1 + U_1^* P_2 U_1 + U_1^* U_2^* P_3 U_2 U_1 + \dots + U_1^* U_2^* \dots U_{n-1}^* P_n U_{n-1} \dots U_2 U_1 = I_{\mathcal{D}_{T^*}}$.

We have also seen in [41] that an isometric dilation can be constructed with conditions (1) – (3) only of Theorem 4.1. We present the result below.

Theorem 4.2. *Let $T_1, \dots, T_n \in \mathcal{B}(\mathcal{H})$ be commuting contractions such that $T = \prod_{i=1}^n T_i$ is a C_0 contraction. Let $T_i' = \prod_{j \neq i} T_j$ for $1 \leq i \leq n$. Then (T_1, \dots, T_n) possesses an isometric dilation on the minimal isometric dilation space of T , if there are projections P_1, \dots, P_n and commuting unitaries U_1, \dots, U_n in $\mathcal{B}(\mathcal{D}_{T^*})$ such that the following hold for $i = 1, \dots, n$:*

- (1) $D_T T_i = P_i^\perp U_i^* D_T + P_i U_i^* D_T T$,
- (2) $P_i^\perp U_i^* P_j^\perp U_j^* = P_j^\perp U_j^* P_i^\perp U_i^*$,
- (3) $U_i P_i U_j P_j = U_j P_j U_i P_i$.

Conversely, if a commuting tuple of contractions (T_1, \dots, T_n) , with the product $T = \prod_{i=1}^n T_i$ being pure, possesses an isometric dilation $(\hat{V}_1, \dots, \hat{V}_n)$, where $\hat{V} = \prod_{i=1}^n \hat{V}_i$ is the minimal isometric dilation of T , then there are unique projections P_1, \dots, P_n and unique commuting unitaries U_1, \dots, U_n in $\mathcal{B}(\mathcal{D}_{T^})$ satisfying the conditions (1) – (3) above.*

We now present the unitary dilation theorem and this is the main result of this Section.

Theorem 4.3. *Let T_1, \dots, T_n be commuting contractions on a Hilbert space \mathcal{H} such that their product $T = \prod_{i=1}^n T_i$ is a C_0 contraction. Then (T_1, \dots, T_n) possesses a unitary dilation (W_1, \dots, W_n) on \mathcal{H} with $W = \prod_{i=1}^n W_i$ being the minimal unitary dilation of T on \mathcal{H} if and only if there are unique projections Q_1, \dots, Q_n and unique commuting unitaries $\tilde{U}_1, \dots, \tilde{U}_n$ in $\mathcal{B}(\mathcal{D}_{T^*})$ such that $\prod_{i=1}^n \tilde{U}_i = I_{\mathcal{D}_{T^*}}$ and the following conditions hold for $i = 1, \dots, n$:*

- (1) $D_{T^*} T_i^* = Q_i^\perp \tilde{U}_i^* D_{T^*} + Q_i \tilde{U}_i^* D_{T^*} T^*$,
- (2) $Q_i^\perp \tilde{U}_i^* Q_j^\perp \tilde{U}_j^* = Q_j^\perp \tilde{U}_j^* Q_i^\perp \tilde{U}_i^*$,
- (3) $\tilde{U}_i Q_i \tilde{U}_j Q_j = \tilde{U}_j Q_j \tilde{U}_i Q_i$,
- (4) $Q_1 + \tilde{U}_1^* Q_2 \tilde{U}_1 + \tilde{U}_1^* \tilde{U}_2^* Q_3 \tilde{U}_2 \tilde{U}_1 + \dots + \tilde{U}_1^* \tilde{U}_2^* \dots \tilde{U}_{n-1}^* Q_n \tilde{U}_{n-1} \dots \tilde{U}_2 \tilde{U}_1 = I_{\mathcal{D}_{T^*}}$.

Proof. First suppose that there are projections Q_1, \dots, Q_n and commuting unitaries $\tilde{U}_1, \dots, \tilde{U}_n$ in $\mathcal{B}(\mathcal{D}_{T^*})$ satisfying the given conditions. Then following the proof of Theorem 4.1 from [41] we have that the Toeplitz operator tuple $(T_{\tilde{U}_1 Q_1^\perp + z \tilde{U}_1 Q_1}, \dots, T_{\tilde{U}_n Q_n^\perp + z \tilde{U}_n Q_n})$ on $H^2(\mathcal{D}_{T^*})$ is an isometric dilation of (T_1, \dots, T_n) with their product $\prod_{i=1}^n T_{\tilde{U}_i Q_i^\perp + z \tilde{U}_i Q_i} = T_z$ being the minimal isometric dilation of the C_0 contraction T . It is obvious that $(W_1, \dots, W_n) = (M_{\tilde{U}_1 Q_1^\perp + z \tilde{U}_1 Q_1}, \dots, M_{\tilde{U}_n Q_n^\perp + z \tilde{U}_n Q_n})$ on $L^2(\mathcal{D}_{T^*})$ is a unitary extension of $(M_{\tilde{U}_1 Q_1^\perp + z \tilde{U}_1 Q_1}, \dots, M_{\tilde{U}_n Q_n^\perp + z \tilde{U}_n Q_n})$ on $H^2(\mathcal{D}_{T^*})$. So, (W_1, \dots, W_n)

on $L^2(\mathcal{D}_{T^*})$ is a unitary dilation of (T_1, \dots, T_n) . Needless to mention that $\prod_{i=1}^n W_i = M_z$ is the minimal unitary dilation of T .

Conversely, suppose (W_1, \dots, W_n) is a unitary dilation of (T_1, \dots, T_n) on the minimal unitary dilation space $\widetilde{\mathcal{H}}$ of T with $\prod_{i=1}^n W_i = W$ being the minimal unitary dilation of T . Without loss of generality let $\widetilde{\mathcal{H}} = l^2(\mathcal{D}_T) \oplus \mathcal{H} \oplus l^2(\mathcal{D}_{T^*})$. Then by converse part of Theorem 3.8, there are unique projections P_1, \dots, P_n and unique commuting unitaries U_1, \dots, U_n in $\mathcal{B}(\mathcal{D}_T)$ such that $\prod_{i=1}^n U_i = I_{\mathcal{D}_T}$ and conditions (1) – (5) in the statement of Theorem 3.8 are satisfied. Thus (\Leftarrow) part of Theorem 3.1 tells us that (T_1, \dots, T_n) possesses an isometric dilation (V_1, \dots, V_n) on the minimal isometric dilation space of T with $\prod_{i=1}^n V_i$ being the minimal isometric dilation of T . Then by Theorem 3.7, (T_1^*, \dots, T_n^*) possesses an isometric dilation (Y_1, \dots, Y_n) on the minimal isometric dilation space of T^* with $\prod_{i=1}^n Y_i$ being the minimal isometric dilation of T^* . Hence, by an application of Theorem 3.1 again we have that there are unique projections Q_1, \dots, Q_n and unique commuting unitaries $\tilde{U}_1, \dots, \tilde{U}_n$ in $\mathcal{B}(\mathcal{D}_{T^*})$ such that $\prod_{i=1}^n \tilde{U}_i = I_{\mathcal{D}_{T^*}}$ such that the conditions (1) – (4) are satisfied. Hence the proof is complete. ■

We now present an analogue of Theorem 4.2 for a unitary dilation when the product T is a C_0 contraction.

Theorem 4.4. *Let $T_1, \dots, T_n \in \mathcal{B}(\mathcal{H})$ be commuting contractions such that $T = \prod_{i=1}^n T_i$ is a C_0 contraction. Let $T'_i = \prod_{j \neq i} T_j$ for $1 \leq i \leq n$. Then (T_1, \dots, T_n) possesses a unitary dilation on the minimal unitary dilation space of T , if there are projections P_1, \dots, P_n and commuting unitaries U_1, \dots, U_n in $\mathcal{B}(\mathcal{D}_{T^*})$ such that the following hold for $i = 1, \dots, n$:*

- (1) $D_T T_i = P_i^\perp U_i^* D_T + P_i U_i^* D_T T$,
- (2) $P_i^\perp U_i^* P_j^\perp U_j^* = P_j^\perp U_j^* P_i^\perp U_i^*$,
- (3) $U_i P_i U_j P_j = U_j P_j U_i P_i$.

Conversely, if a commuting tuple of contractions (T_1, \dots, T_n) , with the product $T = \prod_{i=1}^n T_i$ being a C_0 contraction, possesses a unitary dilation (W_1, \dots, W_n) , where $W = \prod_{i=1}^n W_i$ is the minimal unitary dilation of T , then there are unique projections P_1, \dots, P_n and unique commuting unitaries U_1, \dots, U_n in $\mathcal{B}(\mathcal{D}_{T^})$ satisfying the conditions (1) – (3) above.*

Proof. The existence of such a set of projections and unitaries guarantees the existence of an isometric dilation (V_1, \dots, V_n) on $H^2(\mathcal{D}_{T^*})$ of (T_1, \dots, T_n) by Theorem 4.2. Following the proof of Theorem 4.1 from in [41], we see that $V_i = T_{U_i P_i^\perp + z U_i P_i}$ for $1 \leq i \leq n$. Thus, if we set $W_i := M_{U_i P_i^\perp + z U_i P_i}$ on $L^2(\mathcal{D}_{T^*})$, then (W_1, \dots, W_n) is a unitary extension of (V_1, \dots, V_n) and hence is a unitary dilation of (T_1, \dots, T_n) . The converse part follows from Theorem 4.3. ■

5. SZ. NAGY-FOIAS TYPE MINIMAL UNITARY DILATION

Suppose $T \in \mathcal{B}(\mathcal{H})$ is a c.n.u. contraction and V on \mathcal{H}_0 is the minimal isometric dilation of T . By Wold decomposition there are reducing subspaces $\mathcal{H}_{01}, \mathcal{H}_{02}$ of V such that $\mathcal{H}_0 = \mathcal{H}_{01} \oplus \mathcal{H}_{02}$, $V|_{\mathcal{H}_{01}}$ is the unilateral shift and $V|_{\mathcal{H}_{02}}$ is a unitary. Then \mathcal{H}_{01} can be identified with $H^2(\mathcal{D}_{T^*})$ and \mathcal{H}_{02} can be identified with $\overline{\Delta_T(L^2(\mathcal{D}_T))}$, where $\Delta_T(t) = [I_{\mathcal{D}_T} - \Theta_T(e^{it})^* \Theta_T(e^{it})]^{1/2}$, where Θ_T is the characteristic function of the contraction T . For further details see Chapter-VI of [13]. Thus,

$\mathcal{K}_0 = \mathcal{K}_{01} \oplus \mathcal{K}_{02}$ can be identified with $\mathbb{K}_+ = H^2(\mathcal{D}_{T^*}) \oplus \overline{\Delta_T(L^2(\mathcal{D}_T))}$. Also, V on \mathcal{K}_0 can be realized as $M_z \oplus M_{e^{it}}|_{\overline{\Delta_T(L^2(\mathcal{D}_T))}}$. Thus, there is a unitary

$$\tau = \tau_1 \oplus \tau_2 : \mathcal{K}_{01} \oplus \mathcal{K}_{02} \rightarrow (H^2 \otimes \mathcal{D}_{T^*}) \oplus \overline{\Delta_T(L^2(\mathcal{D}_T))} := \widetilde{\mathbb{K}}_+ \quad (5.1)$$

such that V on \mathcal{K}_0 can be realized as $(M_z \otimes I_{\mathcal{D}_{T^*}}) \oplus M_{e^{it}}|_{\overline{\Delta_T(L^2(\mathcal{D}_T))}}$ on $\widetilde{\mathbb{K}}_+$.

If (T_1, \dots, T_n) is a commuting tuple of contractions on \mathcal{H} satisfying the hypotheses of Theorem 3.2, then it dilates to a commuting tuple of isometries (V_1, \dots, V_n) on minimal isometric dilation space \mathcal{K}_0 of T . Now by Wold decomposition of commuting isometries, we have that \mathcal{K}_{01} and \mathcal{K}_{02} are reducing subspaces for each V_i and that

$$V_{i2} = V_i|_{\mathcal{K}_{02}} \quad (5.2)$$

is a unitary for $1 \leq i \leq n$. We have from [41] the following useful analogue of Theorem 3.2 where we consider the Sz. Nagy-Foias minimal isometric dilation space of T .

Theorem 5.1 ([41], Theorem 6.1). *Let (T_1, \dots, T_n) be a tuple of commuting contractions acting on \mathcal{H} such that $T = \prod_{i=1}^n T_i$ is a c.n.u. contraction. Suppose there are projections P_1, \dots, P_n and commuting unitaries U_1, \dots, U_n in $\mathcal{B}(\mathcal{D}_T)$ satisfying*

- (1) $D_T T_i = P_i^\perp U_i^* D_T + P_i U_i^* D_T T$
- (2) $P_i^\perp U_i^* P_j^\perp U_j^* = P_j^\perp U_j^* P_i^\perp U_i^*$,
- (3) $U_i P_i U_j P_j = U_j P_j U_i P_i$,
- (4) $D_T U_i P_i U_i^* D_T = D_{T_i}^2$,

for $1 \leq i < j \leq n$. Then there are projections Q_1, \dots, Q_n and commuting unitaries $\tilde{U}_1, \dots, \tilde{U}_n$ in $\mathcal{B}(\mathcal{D}_{T^*})$ such that (T_1, \dots, T_n) dilates to the tuple of commuting isometries $(\tilde{V}_{11} \oplus \tilde{V}_{12}, \dots, \tilde{V}_{n1} \oplus \tilde{V}_{n2})$ on $\widetilde{\mathbb{K}}_+ = H^2 \otimes \mathcal{D}_{T^*} \oplus \overline{\Delta_T(L^2(\mathcal{D}_T))}$, where

$$\begin{aligned} \tilde{V}_{i1} &= I \otimes \tilde{U}_i Q_i^\perp + M_z \otimes \tilde{U}_i Q_i, \\ \tilde{V}_{i2} &= \tau_2 V_{i2} \tau_2^*, \end{aligned}$$

for unitaries τ_2 and V_{i2} as in (5.1) and (5.2) respectively for $1 \leq i \leq n$.

Now we present an analogue of this isometric dilation theorem in the unitary dilation setting. This is a main result of this Section.

Theorem 5.2. *Let (T_1, \dots, T_n) be a tuple of commuting contractions acting on \mathcal{H} such that $T = \prod_{i=1}^n T_i$ is a c.n.u. contraction. Suppose there are projections P_1, \dots, P_n and commuting unitaries U_1, \dots, U_n in $\mathcal{B}(\mathcal{D}_T)$ satisfying*

- (1) $D_T T_i = P_i^\perp U_i^* D_T + P_i U_i^* D_T T$
- (2) $P_i^\perp U_i^* P_j^\perp U_j^* = P_j^\perp U_j^* P_i^\perp U_i^*$,
- (3) $U_i P_i U_j P_j = U_j P_j U_i P_i$,
- (4) $D_T U_i P_i U_i^* D_T = D_{T_i}^2$,

for $1 \leq i < j \leq n$. Then there are projections Q_1, \dots, Q_n and commuting unitaries $\tilde{U}_1, \dots, \tilde{U}_n$ in $\mathcal{B}(\mathcal{D}_{T^*})$ such that (T_1, \dots, T_n) dilates to a tuple of commuting unitaries $(\tilde{W}_{11} \oplus \tilde{W}_{12}, \dots, \tilde{W}_{n1} \oplus \tilde{W}_{n2})$ on $\widetilde{\mathbb{K}} = L^2 \otimes \mathcal{D}_{T^*} \oplus \overline{\Delta_T(L^2(\mathcal{D}_T))}$, where

$$\begin{aligned} \tilde{W}_{i1} &= I \otimes \tilde{U}_i Q_i^\perp + M_z \otimes \tilde{U}_i Q_i, \\ \tilde{W}_{i2} &= \tau_2 V_{i2} \tau_2^*, \end{aligned}$$

for unitaries τ_2 and V_{i2} as in (5.1) and (5.2) respectively for $1 \leq i \leq n$.

Proof. Since there are projections $P_1, \dots, P_n \in \mathcal{B}(\mathcal{D}_T)$ and commuting unitaries $U_1, \dots, U_n \in \mathcal{B}(\mathcal{D}_T)$ satisfying conditions (1) – (4), we have by Theorem 5.1 that (T_1, \dots, T_n) has an isometric dilation $(\tilde{V}_{11} \oplus \tilde{V}_{12}, \dots, \tilde{V}_{n1} \oplus \tilde{V}_{n2})$ on $\tilde{\mathbb{K}}_+ = H^2 \otimes \mathcal{D}_T^* \oplus \overline{\Delta_T(L^2(\mathcal{D}_T))}$, where

$$\begin{aligned}\tilde{V}_{i1} &= I \otimes \tilde{U}_i Q_i^\perp + M_z \otimes \tilde{U}_i Q_i, \\ \tilde{V}_{i2} &= \tau_2 V_{i2} \tau_2^*, \quad (1 \leq i \leq n)\end{aligned}$$

for a unitary $\tau_2 : \mathcal{H}_{02} \rightarrow \overline{\Delta_T(L^2(\mathcal{D}_T))}$ and a unitary V_{i2} on \mathcal{H}_{02} as in (5.1) and (5.2) respectively. Let us consider for $i = 1, \dots, n$ the following operators on $\tilde{\mathbb{K}} = L^2 \otimes \mathcal{D}_T^* \oplus \overline{\Delta_T(L^2(\mathcal{D}_T))}$:

$$\begin{aligned}\tilde{W}_{i1} &= I \otimes \tilde{U}_i Q_i^\perp + M_z \otimes \tilde{U}_i Q_i, \\ \tilde{W}_{i2} &= \tau_2 V_{i2} \tau_2^*.\end{aligned}$$

Evidently, $(\tilde{W}_{11} \oplus \tilde{W}_{12}, \dots, \tilde{W}_{n1} \oplus \tilde{W}_{n2})$ is a unitary extension of $(\tilde{V}_{11} \oplus \tilde{V}_{12}, \dots, \tilde{V}_{n1} \oplus \tilde{V}_{n2})$ and consequently $(\tilde{W}_{11} \oplus \tilde{W}_{12}, \dots, \tilde{W}_{n1} \oplus \tilde{W}_{n2})$ is a unitary dilation of (T_1, \dots, T_n) . ■

We conclude this paper with a weaker version of Theorem 3.8 that assumes conditions (1) – (4) of Theorem 3.8 for having a unitary dilation of (T_1, \dots, T_n) . We have already seen in Theorem 5.2 that a c.n.u. tuple of commuting contractions (T_1, \dots, T_n) dilates to commuting unitaries on the minimal unitary dilation space of $T = \prod_{i=1}^n T_i$ with these four conditions. However, because of such weaker hypotheses we will not have a proper converse part.

Theorem 5.3. *Let $T_1, \dots, T_n \in \mathcal{B}(\mathcal{H})$ be commuting contractions, $T_i' = \prod_{j \neq i} T_j$ for all $1 \leq i \leq n$ and $T = \prod_{i=1}^n T_i$. Then (T_1, \dots, T_n) possesses a unitary dilation on the minimal unitary dilation space of T , if there are projections P_1, \dots, P_n and commuting unitaries U_1, \dots, U_n in $\mathcal{B}(\mathcal{D}_T)$ such that the following hold for $i = 1, \dots, n$:*

- (1) $D_T T_i = P_i^\perp U_i^* D_T + P_i U_i^* D_T T$,
- (2) $P_i^\perp U_i^* P_j^\perp U_j^* = P_j^\perp U_j^* P_i^\perp U_i^*$,
- (3) $U_i P_i U_j P_j = U_j P_j U_i P_i$,
- (4) $D_T U_i P_i U_i^* D_T = D_{T_i}^2$.

Conversely, if (T_1, \dots, T_n) possesses a unitary dilation $(\hat{W}_1, \dots, \hat{W}_n)$ with $W = \prod_{i=1}^n W_i$ being the minimal unitary dilation of T , then there are unique projections P_1, \dots, P_n and unique commuting unitaries U_1, \dots, U_n in $\mathcal{B}(\mathcal{D}_T)$ satisfying the conditions (1) – (4) above.

Proof. Let (T_1, \dots, T_n) satisfy the conditions (1) – (4). Suppose $T = U \oplus \tilde{T}$ is the canonical decomposition of T with respect to $\mathcal{H} = \mathcal{H}_1 \oplus \mathcal{H}_2$, where \mathcal{H}_1 is the maximal reducing subspace of T such that $U = T|_{\mathcal{H}_1}$ is a unitary and $\tilde{T} = T|_{\mathcal{H}_2}$ is a c.n.u. contraction. Then, it is well-known (e.g. see Lemma 2.2 in [27] or Theorem 3.7 in [39]) that $\mathcal{H}_1, \mathcal{H}_2$ are common reducing subspaces for T_1, \dots, T_n and $(T_1|_{\mathcal{H}_1}, \dots, T_n|_{\mathcal{H}_1})$ is a tuple of commuting unitaries, whereas $(T_1|_{\mathcal{H}_2}, \dots, T_n|_{\mathcal{H}_2})$ is a c.n.u. tuple, i.e. a tuple of commuting contractions whose product is a c.n.u. contraction. Since $D_T \equiv D_{\tilde{T}}$ and $\mathcal{D}_{\tilde{T}} \subseteq \mathcal{H}_2$, we have that the projections P_1, \dots, P_n and commuting unitaries U_1, \dots, U_n satisfy analogues of conditions (1) & (4) respectively with D_T and T_i being replaced by $D_{\tilde{T}}$ and $T_i|_{\mathcal{H}_2}$ respectively. Now, Theorem 5.2 tells us that the c.n.u. tuple $(T_1|_{\mathcal{H}_2}, \dots, T_n|_{\mathcal{H}_2})$ can be dilated to commuting unitaries say (W_1, \dots, W_n) on the Sz. Nagy-Foias minimal unitary dilation space for \tilde{T} . It is merely mentioned that there is a

unitary τ from the Sz. Nagy-Foias minimal unitary space to the Schäffer's minimal unitary dilation space $l^2(\mathcal{D}_{\tilde{T}}) \oplus \mathcal{H}_2 \oplus l^2(\mathcal{D}_{\tilde{T}^*})$ of \tilde{T} . Clearly $l^2(\mathcal{D}_{\tilde{T}}) \oplus \mathcal{H}_2 \oplus l^2(\mathcal{D}_{\tilde{T}^*})$ can be identified with $l^2(\mathcal{D}_T) \oplus \mathcal{H}_2 \oplus l^2(\mathcal{D}_{T^*})$. Thus, (W_1, \dots, W_n) can be identified with a tuple of commuting unitaries say $(\tilde{W}_1, \dots, \tilde{W}_n)$ on $l^2(\mathcal{D}_T) \oplus l^2(\mathcal{D}_{T^*}) \oplus \mathcal{H}_2$ dilating $(T_1|_{\mathcal{H}_2}, \dots, T_n|_{\mathcal{H}_2})$. It is evident that $(\tilde{W}_1 \oplus T_1|_{\mathcal{H}_1}, \dots, \tilde{W}_n \oplus T_n|_{\mathcal{H}_1})$ is a unitary dilation of (T_1, \dots, T_n) on the minimal unitary dilation space $l^2(\mathcal{D}_T) \oplus l^2(\mathcal{D}_{T^*}) \oplus \mathcal{H}_2 \oplus \mathcal{H}_1 \equiv l^2(\mathcal{D}_T) \oplus \mathcal{H} \oplus l^2(\mathcal{D}_{T^*})$ of T . The converse part follows from Theorem 3.8. ■

Note that the class of commuting contractions that dilate to commuting unitaries by satisfying four conditions of Theorem 5.3 is strictly larger than the class satisfying the five conditions of Theorem 3.8, though they are being dilated to the same space. In this context we would like to recall Example 5.4 from [41]. Indeed, if we consider

$$T_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, T_2 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \text{ and } T_3 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \text{ on } \mathbb{C}^3,$$

we see that the commuting triple (T_1, T_2, T_3) satisfies conditions (1) – (4) but fails to meet condition (5) of Theorem 3.8. Thus, this triple possesses a unitary dilation (U_1, U_2, U_3) on the minimal unitary dilation space of $T_1 T_2 T_3$ but the product $U_1 U_2 U_3$ is not the minimal unitary dilation of $T_1 T_2 T_3$.

6. APPENDIX

Proof of (3.7). We have that

$$D_T F_i D_T + D_T^* G_i' D_T^* T = D_{T_i'}^2 T_i + D_{T_i'}^2 T_i'^* T = T_i - T_i'^* T_i' T_i + T_i'^* T - T_i T_i'^* T_i'^* T = T_i D_T^2.$$

Applying the relation $T D_T = D_T^* T$ we have $D_T F_i D_T + D_T^* G_i' T D_T = T_i D_T^2$. Thus, we have $D_T F_i = T_i D_T - D_T^* G_i' T|_{\mathcal{D}_T}$. Similarly, we can have $D_T F_i' D_T + D_T^* G_i T D_T = T_i' D_T^2$, which implies $D_T F_i' = T_i' D_T - D_T^* G_i T|_{\mathcal{D}_T}$. Now $D_T F_i' F_i'^* D_T = D_{T_i'}^2$ leads to the following:

$$\begin{aligned} & (T_i' D_T - D_T^* G_i T)(D_T T_i'^* - T^* G_i^* D_T^*) - D_{T_i'}^2 = 0 \\ \implies & T_i' D_T^2 T_i'^* - T_i' T^* D_T^* G_i^* D_T^* - D_T^* G_i D_T^* T T_i'^* + D_T^* G_i T T^* G_i^* D_T^* - D_{T_i'}^2 = 0 \\ & \quad [\text{by } T^* D_T^* = D_T T^*, T D_T = D_T^* T] \\ \implies & T_i' T_i'^* - T_i' T^* T T_i'^* - T_i' T^* D_T^* G_i^* D_T^* - T_i^* T T_i'^* + T_i' T^* T T_i'^* \\ & \quad + D_T G_i T T^* G_i^* D_T^* - I + T_i^* T_i = 0 \quad [\text{since } D_T^* G_i D_T^* = D_{T_i'}^2 T_i^* = T_i^* - T_i' T^*] \\ \implies & D_T G_i T T^* G_i^* D_T^* + T_i^* T_i - T_i^* T T_i'^* - T_i' T^* D_T^* G_i^* D_T^* = I - T_i' T_i'^* \\ \implies & D_T^* G_i T T^* G_i^* D_T^* + T_i^* (T_i - T T_i'^*) - T_i' T^* D_T^* G_i^* D_T^* = D_{T_i'}^2 \\ \implies & D_T^* G_i T T^* G_i^* D_T^* + T_i^* D_T^* G_i^* D_T^* - T_i' T^* D_T^* G_i^* D_T^* = D_{T_i'}^2 \\ \implies & D_T^* G_i T T^* G_i^* D_T^* + (T_i^* - T_i' T^*) D_T^* G_i^* D_T^* = D_{T_i'}^2 \\ \implies & D_T^* G_i T T^* G_i^* D_T^* + D_T^* G_i D_T^* D_T^* G_i^* D_T^* = D_{T_i'}^2 \\ \implies & D_T^* G_i G_i^* D_T^* = D_{T_i'}^2. \end{aligned}$$

Similarly, from $D_T F_i F_i^* D_T = D_{T_i'}^2$ and $D_T F_i = T_i D_T - D_{T^*} G_i' T|_{\mathcal{D}_T}$ we obtain

$$D_{T^*} G_i' G_i^* D_{T^*} = D_{T_i'}^2. \quad (6.1)$$

As a consequence we have the following.

$$\begin{aligned} D_{T^*} G_i' G_i^* D_{T^*} &= (T_i^* D_{T^*} - D_T F_i' T^*) G_i' D_{T^*} = T_i^* D_{T^*} G_i' D_{T^*} - D_T F_i' F_i^* D_T T^* \\ &= T_i^* (T_i'^* - T_i T^*) - D_T F_i' F_i^* D_T T^* \\ &= T^* - T_i^* T_i T^* - D_T F_i' F_i^* D_T T^* \\ &= D_{T_i'}^2 T^* - D_T F_i' F_i^* D_T T^* \\ &= 0. \end{aligned}$$

Therefore, $G_i G_i' = 0$. Similarly we can prove that $G_i' G_i = 0$. Again note that

$$\begin{aligned} D_{T^*} (G_i G_i^* + G_i'^* G_i') D_{T^*} &= D_{T^*} G_i G_i^* D_{T^*} + D_{T^*} G_i'^* T T^* G_i' D_{T^*} + D_{T^*} G_i'^* D_{T^*} D_{T^*} G_i' D_{T^*} \\ &= D_{T_i'}^2 + D_{T^*} T F_i' F_i^* T^* D_{T^*} + (T_i' - T T_i^*) (T_i'^* - T_i T^*) \\ &= I - T_i' T_i'^* + T D_{T_i'}^2 T^* + T_i' T_i'^* - T T^* - T T^* + T T_i^* T_i T^* \\ &= I + T T^* - T T_i^* T_i T^* - 2 T T^* + T T_i^* T_i T^* \\ &= D_{T^*}^2. \end{aligned}$$

Similarly, we can prove that $D_{T^*} (G_i^* G_i + G_i' G_i'^*) D_{T^*} = D_{T^*}^2$. So, we obtain

$$G_i^* G_i + G_i' G_i'^* = I = G_i G_i^* + G_i'^* G_i'.$$

Proof of (3.17). Note that we already have $D_{V^*} V^k h = 0$ for all $h \in \mathcal{H}$, $k \in \mathbb{N}$. Therefore, for all $h \in \mathcal{H}$ and $k \in \mathbb{N}$ we have

$$D_{V_i'^*}^2 V_i^* V^k h = (I - V_i' V_i'^*) V_i' V^{k-1} h = (V_i' - V_i' V_i'^* V_i') V^{k-1} h = 0.$$

Thus, (3.17) holds for all vectors in $\overline{\text{span}}\{V^k h : k \in \mathbb{N}, h \in \mathcal{H}\}$. As V is an isometry, $D_{V^*}^2 = D_{V^*}$ and thus (3.15) tells us that $D_{V_i'^*}^2 V_i^* = 0$ on $\mathcal{N}(\mathcal{D}_T)$ and thus $D_{V_i'^*}^2 V_i^* = D_{V^*} D_{V_i'^*}^2 V_i^* D_{V^*}$. For $h, h' \in \mathcal{H}$, we have that

$$\begin{aligned} \langle D_{V^*} X G_i X^* D_{V^*} D_{V^*} h, D_{V^*} h' \rangle &= \langle D_{V^*}^2 X G_i X^* D_{V^*}^2 h, h' \rangle = \langle D_{V^*} X G_i X^* D_{V^*} h, h' \rangle \\ &= \langle G_i X^* D_{V^*} h, X^* D_{V^*} h' \rangle \\ &= \langle G_i D_{T^*} h, D_{T^*} h' \rangle \\ &\quad [\text{since } X D_{T^*} = D_{V^*} \text{ \& } X^* X = I] \\ &= \langle D_{T^*} G_i D_{T^*} h, h' \rangle \\ &= \langle (T_i^* - T_i' T^*) h, h' \rangle \\ &= \langle V_i^* h, h' \rangle - \langle V^* h, V_i'^* h' \rangle \\ &= \langle (V_i^* - V_i' V^*) h, h' \rangle \\ &= \langle D_{V_i'^*}^2 V_i^* h, h' \rangle \\ &= \langle D_{V^*} D_{V_i'^*}^2 V_i^* D_{V^*} h, h' \rangle \\ &= \langle D_{V_i'^*}^2 V_i^* D_{V^*} h, D_{V^*} h' \rangle. \end{aligned}$$

Hence, the first identity in (3.17) holds. Similarly, one can prove the other identity.

Proof of (3.23). For proving (ii) let us first observe the following:

$$\begin{aligned}
 V_i D_j + D_i E_j &= \begin{bmatrix} T_i & 0 & 0 & 0 & \cdots \\ P_i U_i^* D_T & P_i^\perp U_i^* & 0 & 0 & \cdots \\ 0 & P_i U_i^* & P_i^\perp U_i^* & 0 & \cdots \\ 0 & 0 & P_i U_i^* & P_i^\perp U_i^* & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix} \begin{bmatrix} D_{T^*} \tilde{U}_j Q_j & 0 & 0 & \cdots \\ -P_j U_j^* T^* & 0 & 0 & \cdots \\ 0 & 0 & 0 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix} \\
 &+ \begin{bmatrix} D_{T^*} \tilde{U}_i Q_i & 0 & 0 & \cdots \\ -P_i U_i^* T^* & 0 & 0 & \cdots \\ 0 & 0 & 0 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix} \begin{bmatrix} \tilde{U}_j Q_j^\perp & \tilde{U}_j Q_j & 0 & \cdots \\ 0 & \tilde{U}_j Q_j^\perp & \tilde{U}_j Q_j & \cdots \\ 0 & 0 & \tilde{U}_j Q_j^\perp & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix} \\
 &= \begin{bmatrix} T_i D_{T^*} \tilde{U}_j Q_j + D_{T^*} \tilde{U}_i Q_i \tilde{U}_j Q_j^\perp & D_{T^*} \tilde{U}_i Q_i \tilde{U}_j Q_j & 0 & \cdots \\ P_i U_i^* D_T D_{T^*} \tilde{U}_j Q_j - P_i^\perp U_i^* P_j U_j^* T^* - P_i U_i^* T^* \tilde{U}_j Q_j^\perp & P_i U_i^* T^* \tilde{U}_j Q_j & 0 & \cdots \\ -P_i U_i^* P_j U_j^* T^* & 0 & 0 & \cdots \\ 0 & 0 & 0 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix}
 \end{aligned}$$

and

$$\begin{aligned}
 V_j D_i + D_j E_i &= \begin{bmatrix} T_j & 0 & 0 & 0 & \cdots \\ P_j U_j^* D_T & P_j^\perp U_j^* & 0 & 0 & \cdots \\ 0 & P_j U_j^* & P_j^\perp U_j^* & 0 & \cdots \\ 0 & 0 & P_j U_j^* & P_j^\perp U_j^* & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix} \begin{bmatrix} D_{T^*} \tilde{U}_i Q_i & 0 & 0 & \cdots \\ -P_i U_i^* T^* & 0 & 0 & \cdots \\ 0 & 0 & 0 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix} \\
 &+ \begin{bmatrix} D_{T^*} \tilde{U}_j Q_j & 0 & 0 & \cdots \\ -P_j U_j^* T^* & 0 & 0 & \cdots \\ 0 & 0 & 0 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix} \begin{bmatrix} \tilde{U}_i Q_i^\perp & \tilde{U}_i Q_i & 0 & \cdots \\ 0 & \tilde{U}_i Q_i^\perp & \tilde{U}_i Q_i & \cdots \\ 0 & 0 & \tilde{U}_i Q_i^\perp & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix} \\
 &= \begin{bmatrix} T_j D_{T^*} \tilde{U}_i Q_i + D_{T^*} \tilde{U}_j Q_j \tilde{U}_i Q_i^\perp & D_{T^*} \tilde{U}_j Q_j \tilde{U}_i Q_i & 0 & \cdots \\ P_j U_j^* D_T D_{T^*} \tilde{U}_i Q_i - P_j^\perp U_j^* P_i U_i^* T^* - P_j U_j^* T^* \tilde{U}_i Q_i^\perp & P_j U_j^* T^* \tilde{U}_i Q_i & 0 & \cdots \\ -P_j U_j^* P_i U_i^* T^* & 0 & 0 & \cdots \\ 0 & 0 & 0 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix}.
 \end{aligned}$$

Clearly $P_i U_i^* P_j U_j^* T^* = P_j U_j^* P_i U_i^* T^*$ and $D_{T^*} \tilde{U}_i Q_i \tilde{U}_j Q_j = D_{T^*} \tilde{U}_j Q_j \tilde{U}_i Q_i$ follow from (3) and (3)' respectively. Next observe that

$$D_{T^*} T U_i P_i D_T = T D_T U_i P_i D_T = T(T_i' - T_i^* T) = (T_i' - T T_i^*) T = D_{T^*} Q_i \tilde{U}_i^* D_{T^*} T = D_{T^*} Q_i \tilde{U}_i^* T D_T.$$

Since both $T U_i P_i$ and $Q_i \tilde{U}_i^* T$ map \mathcal{D}_T into \mathcal{D}_{T^*} , we have that

$$T U_i P_i = Q_i \tilde{U}_i^* T|_{\mathcal{D}_T}. \quad (6.2)$$

Therefore, we have

$$P_j U_j^* T^* \tilde{U}_i Q_i = T^* \tilde{U}_j Q_j \tilde{U}_i Q_i = T^* \tilde{U}_i Q_i \tilde{U}_j Q_j = P_i U_i^* T^* \tilde{U}_j Q_j$$

So, for proving $V_i D_j + D_i E_j = V_j D_i + D_j E_i$, it suffices to show

- (a) $T_i D_{T^*} \tilde{U}_j Q_j + D_{T^*} \tilde{U}_i Q_i \tilde{U}_j Q_j^\perp = T_j D_{T^*} \tilde{U}_i Q_i + D_{T^*} \tilde{U}_j Q_j \tilde{U}_i Q_i^\perp$,
- (b) $P_i U_i^* D_T D_{T^*} \tilde{U}_j Q_j - P_i^\perp U_i^* P_j U_j^* T^* - P_i U_i^* T^* \tilde{U}_j Q_j^\perp = P_j U_j^* D_T D_{T^*} \tilde{U}_i Q_i - P_j^\perp U_j^* P_i U_i^* T^* - P_j U_j^* T^* \tilde{U}_i Q_i^\perp$.

For proving (a) we first show that

$$\tilde{U}_i Q_i^\perp \tilde{U}_j Q_j + \tilde{U}_i Q_i \tilde{U}_j Q_j^\perp = \tilde{U}_j Q_j^\perp \tilde{U}_i Q_i + \tilde{U}_j Q_j \tilde{U}_i Q_i^\perp. \quad (6.3)$$

Note that

$$\begin{aligned} \tilde{U}_i Q_i^\perp \tilde{U}_j Q_j + \tilde{U}_i Q_i \tilde{U}_j Q_j^\perp &= \tilde{U}_i \tilde{U}_j Q_j - \tilde{U}_i Q_i \tilde{U}_j Q_j + \tilde{U}_i Q_i \tilde{U}_j - \tilde{U}_i Q_i \tilde{U}_j Q_j \\ &= \tilde{U}_i \tilde{U}_j (Q_j + \tilde{U}_j^* Q_i \tilde{U}_j) - 2 \tilde{U}_i Q_i \tilde{U}_j Q_j \\ &= \tilde{U}_j \tilde{U}_i (Q_i + \tilde{U}_i^* Q_j \tilde{U}_i) - 2 \tilde{U}_i Q_i \tilde{U}_j Q_j \quad [\text{From (3.22).}] \\ &= \tilde{U}_j \tilde{U}_i Q_i + \tilde{U}_j Q_j \tilde{U}_i - \tilde{U}_j Q_j \tilde{U}_i Q_i - \tilde{U}_j Q_j \tilde{U}_i Q_i \\ &= \tilde{U}_j Q_j^\perp \tilde{U}_i Q_i + \tilde{U}_j Q_j \tilde{U}_i Q_i^\perp. \end{aligned}$$

Now we prove (a) using conditions-(1'), (3') and (6.3) in the following way.

$$\begin{aligned} &T_i D_{T^*} \tilde{U}_j Q_j + D_{T^*} \tilde{U}_i Q_i \tilde{U}_j Q_j^\perp \\ &= D_{T^*} \tilde{U}_i Q_i^\perp \tilde{U}_j Q_j + T D_{T^*} \tilde{U}_i Q_i \tilde{U}_j Q_j + D_{T^*} \tilde{U}_i Q_i \tilde{U}_j Q_j^\perp \quad [\text{by condition } - (1')] \\ &= D_{T^*} (\tilde{U}_i Q_i^\perp \tilde{U}_j Q_j + \tilde{U}_i Q_i \tilde{U}_j Q_j^\perp) + T D_{T^*} \tilde{U}_i Q_i \tilde{U}_j Q_j \\ &= D_{T^*} (\tilde{U}_j Q_j^\perp \tilde{U}_i Q_i + \tilde{U}_j Q_j \tilde{U}_i Q_i^\perp) + T D_{T^*} \tilde{U}_j Q_j \tilde{U}_i Q_i \quad [\text{by (6.3) and condition } - (3')] \\ &= D_{T^*} \tilde{U}_j Q_j^\perp \tilde{U}_i Q_i + T D_{T^*} \tilde{U}_j Q_j \tilde{U}_i Q_i + D_{T^*} \tilde{U}_j Q_j \tilde{U}_i Q_i^\perp \\ &= (D_{T^*} \tilde{U}_j Q_j^\perp + T D_{T^*} \tilde{U}_j Q_j) \tilde{U}_i Q_i + D_{T^*} \tilde{U}_j Q_j \tilde{U}_i Q_i^\perp \\ &= T_j D_{T^*} \tilde{U}_i Q_i + D_{T^*} \tilde{U}_j Q_j \tilde{U}_i Q_i^\perp. \quad [\text{by condition } - (1')] \end{aligned}$$

This proves (a). Before proving (b) note that by an argument similar to that in (6.3), we can have

$$P_i U_i^* P_j^\perp U_j^* + P_i^\perp U_i^* P_j U_j^* = P_j U_j^* P_i^\perp U_i^* + P_j^\perp U_j^* P_i U_i^*. \quad (6.4)$$

Using (3.18), (3.19) we have that

$$\begin{aligned} D_T U_i P_i D_T + D_{T^*} Q_i^\perp \tilde{U}_i^* T D_T &= D_{T_i}^2 T_i' + D_{T_i'^*} T_i^* T \\ &= T_i' - T_i^* T + T_i^* T - T_i' T^* T \\ &= T_i' D_T^2. \end{aligned}$$

Hence,

$$D_T U_i P_i = T_i' D_T - D_{T^*} Q_i^\perp \tilde{U}_i^* T|_{\mathcal{D}_T}. \quad (6.5)$$

Further we observe that

$$\begin{aligned}
& (P_i U_i^* D_T D_{T^*} - P_i^\perp U_i^* T^* - D_T D_{T^*} \tilde{U}_i Q_i + T^* \tilde{U}_i Q_i^\perp) D_{T^*} \\
&= P_i U_i^* D_T D_{T^*}^2 - P_i^\perp U_i^* T^* D_{T^*} - D_T D_{T^*} \tilde{U}_i Q_i D_{T^*} + T^* \tilde{U}_i Q_i^\perp D_{T^*} \\
&= P_i U_i^* D_T - P_i U_i^* D_T T T^* - P_i^\perp U_i^* D_T T^* - D_T D_{T^*} \tilde{U}_i Q_i D_{T^*} + T^* \tilde{U}_i Q_i^\perp D_{T^*} \quad [\text{From } T^* D_{T^*} = D_T T^*] \\
&= P_i U_i^* D_T - (P_i U_i^* D_T T + P_i^\perp U_i^* D_T) T^* - D_T D_{T^*} \tilde{U}_i Q_i D_{T^*} + T^* \tilde{U}_i Q_i^\perp D_{T^*} \\
&= P_i U_i^* D_T - D_T T_i T^* - D_T D_{T_i^*}^2 T_i'^* + T^* \tilde{U}_i Q_i^\perp D_{T^*} \quad [\text{From condition-(1) and (3.19)}] \\
&= P_i U_i^* D_T - D_T (T_i T^* + T_i'^* - T_i T_i^* T_i'^*) + T^* \tilde{U}_i Q_i^\perp D_{T^*} \\
&= P_i U_i^* D_T - D_T T_i'^* + T^* \tilde{U}_i Q_i^\perp D_{T^*} \\
&= 0. \quad [\text{From (6.5)}]
\end{aligned}$$

Therefore, we have

$$P_i U_i^* D_T D_{T^*} - P_i^\perp U_i^* T^*|_{\mathcal{D}_{T^*}} = D_T D_{T^*} \tilde{U}_i Q_i - T^* \tilde{U}_i Q_i^\perp. \quad (6.6)$$

Now we prove (b). We have

$$\begin{aligned}
& P_i U_i^* D_T D_{T^*} \tilde{U}_j Q_j - P_i^\perp U_i^* P_j U_j^* T^* - P_i U_i^* T^* \tilde{U}_j Q_j^\perp \\
&= P_i U_i^* (D_T D_{T^*} \tilde{U}_j Q_j - T^* \tilde{U}_j Q_j^\perp) - P_i^\perp U_i^* P_j U_j^* T^* \quad [\text{by (6.6)}] \\
&= P_i U_i^* (P_j U_j^* D_T D_{T^*} - P_j^\perp U_j^* T^*) - P_i^\perp U_i^* P_j U_j^* T^* \quad [\text{by (6.2)}] \\
&= P_i U_i^* P_j U_j^* D_T D_{T^*} - P_i U_i^* P_j^\perp U_j^* T^* - P_i^\perp U_i^* P_j U_j^* T^* \\
&= P_j U_j^* P_i U_i^* D_T D_{T^*} - (P_i U_i^* P_j^\perp U_j^* + P_i^\perp U_i^* P_j U_j^*) T^* \\
&= P_j U_j^* P_i U_i^* D_T D_{T^*} - (P_j U_j^* P_i^\perp U_i^* + P_j^\perp U_j^* P_i U_i^*) T^* \quad [\text{by (6.4)}] \\
&= P_j U_j^* (P_i U_i^* D_T D_{T^*} - P_i^\perp U_i^* T^*) - P_j^\perp U_j^* P_i U_i^* T^* \\
&= P_j U_j^* (D_T D_{T^*} \tilde{U}_i Q_i - T^* \tilde{U}_i Q_i^\perp) - P_j^\perp U_j^* P_i U_i^* T^* \quad [\text{by (6.6)}] \\
&= P_j U_j^* D_T D_{T^*} \tilde{U}_i Q_i - P_j^\perp U_j^* P_i U_i^* T^* - P_j U_j^* T^* \tilde{U}_i Q_i^\perp.
\end{aligned}$$

Proof of (3.24). Note that (3.24) holds if and only if

- (a') $V_i V_i^* + D_i D_i^* = I_{\mathcal{H}_0}$,
- (b') $D_i E_i^* = 0$,
- (c') $E_i E_i^* = I_{l^2(\mathcal{D}_{T^*})}$.

First we observe that the matrix of $E_i E_i^*$ with respect to the decomposition $\mathcal{D}_{T^*} \oplus \mathcal{D}_{T^*} \oplus \dots$ is

$$\begin{bmatrix}
\tilde{U}_i Q_i^\perp \tilde{U}_i^* + \tilde{U}_i Q_i \tilde{U}_i^* & \tilde{U}_i Q_i Q_i^\perp \tilde{U}_i^* & 0 & 0 & \dots \\
\tilde{U}_i Q_i^\perp Q_i \tilde{U}_i^* & \tilde{U}_i Q_i^\perp \tilde{U}_i^* + \tilde{U}_i Q_i \tilde{U}_i^* & \tilde{U}_i Q_i Q_i^\perp \tilde{U}_i^* & 0 & \dots \\
0 & \tilde{U}_i Q_i^\perp Q_i \tilde{U}_i^* & \tilde{U}_i Q_i^\perp \tilde{U}_i^* + \tilde{U}_i Q_i \tilde{U}_i^* & \tilde{U}_i Q_i Q_i^\perp \tilde{U}_i^* & \dots \\
0 & 0 & \tilde{U}_i Q_i^\perp Q_i \tilde{U}_i^* & \tilde{U}_i Q_i^\perp \tilde{U}_i^* + \tilde{U}_i Q_i \tilde{U}_i^* & \dots \\
\vdots & \vdots & \vdots & \vdots & \ddots
\end{bmatrix}.$$

Since $\tilde{U}_i \tilde{U}_i^* = I$, $Q_i + Q_i^\perp = I$ and $Q_i^\perp Q_i = 0$, we have (c'). The $(1, 1)$ entry of $D_i E_i^*$ is $D_T^* \tilde{U}_i Q_i Q_i^\perp \tilde{U}_i^* = 0$ and the $(2, 1)$ entry is $-P U_i^* T^* Q_i^\perp \tilde{U}_i^*$. So, we have from (6.2)

$$-P U_i^* T^* Q_i^\perp \tilde{U}_i^* = -T^* \tilde{U}_i Q_i Q_i^\perp \tilde{U}_i^* = 0.$$

All other entries of $D_i E_i^*$ are equal to 0. This proves (b'). For proving (a) we first observe that,

$$V_i V_i^* = \begin{bmatrix} T_i T_i^* & T_i D_T U_i P_i & 0 & 0 & \dots \\ P_i U_i^* D_T T_i^* & P_i U_i^* D_T^2 U_i P_i + P_i^\perp U_i^* U_i P_i^\perp & P_i^\perp U_i^* U_i P_i & 0 & \dots \\ 0 & P_i U_i^* U_i P_i^\perp & P_i U_i^* U_i P_i + P_i^\perp U_i^* U_i P_i^\perp & P_i^\perp U_i^* U_i P_i & \dots \\ 0 & 0 & P_i U_i^* U_i P_i^\perp & P_i U_i^* U_i P_i + P_i^\perp U_i^* U_i P_i^\perp & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}.$$

Using the fact that $U_i^* U_i = I$, $P_i^\perp P_i = 0$ and $P_i + P_i^\perp = I$ we obtain

$$V_i V_i^* = \begin{bmatrix} T_i T_i^* & T_i D_T U_i P_i & 0 & 0 & \dots \\ P_i U_i^* D_T T_i^* & I - P_i U_i^* T^* T U_i P_i & 0 & 0 & \dots \\ 0 & 0 & I & 0 & \dots \\ 0 & 0 & 0 & I & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}.$$

Hence matrix of $V_i V_i^* + D_i D_i^*$ with respect to the decomposition $\mathcal{H} \oplus \mathcal{D}_T \oplus \mathcal{D}_T \oplus \dots$ is

$$\begin{bmatrix} T_i T_i^* + D_T^* \tilde{U}_i Q_i \tilde{U}_i^* D_T^* & T_i D_T U_i P_i - D_T^* \tilde{U}_i Q_i T U_i P_i & 0 & 0 & \dots \\ P_i U_i^* D_T T_i^* - P_i U_i^* T^* D_T^* \tilde{U}_i Q_i & I & 0 & 0 & \dots \\ 0 & 0 & I & 0 & \dots \\ 0 & 0 & 0 & I & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}.$$

Here the $(1, 1)$ entry is equal to the identity by (4'). Thus, it remains to prove that

$$T_i D_T U_i P_i - D_T^* \tilde{U}_i Q_i T U_i P_i|_{\mathcal{D}_T} = 0.$$

Note that

$$\begin{aligned} & T_i D_T U_i P_i D_T - D_T^* \tilde{U}_i Q_i T U_i P_i D_T \\ &= T_i (T_i' - T_i^* T) - D_T^* \tilde{U}_i Q_i \tilde{U}_i^* T D_T \quad [\text{by (6.2)}] \\ &= T_i T_i' - T_i T_i^* T - D_T^* \tilde{U}_i Q_i \tilde{U}_i^* D_T^* T \\ &= T - T_i T_i^* T - (I - T_i T_i^*) T \quad [\text{by condition-(4')}] \\ &= 0. \end{aligned}$$

Proof of (3.34). Note that

$$\underline{V}_k D_{k+1} + \underline{D}_k E_{k+1} = \begin{bmatrix} \underline{T}_k D_T^* \tilde{U}_{k+1} Q_{k+1} + D_T^* \underline{\tilde{U}}_k \underline{Q}_k \tilde{U}_{k+1} Q_{k+1}^\perp & D_T^* \underline{\tilde{U}}_k \underline{Q}_k \tilde{U}_{k+1} Q_{k+1} & 0 & \dots \\ \underline{P}_k U_k^* D_T D_T^* \tilde{U}_{k+1} Q_{k+1} - \underline{P}_k^\perp U_k^* P_{k+1} U_{k+1}^* T^* - \underline{P}_k U_k^* T^* \tilde{U}_{k+1} Q_{k+1}^\perp & -\underline{P}_k U_k^* T^* \tilde{U}_{k+1} Q_{k+1} & 0 & \dots \\ -\underline{P}_k U_k^* P_{k+1} U_{k+1}^* T^* & 0 & 0 & \dots \\ 0 & 0 & 0 & \dots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix}.$$

In the above block-matrix the $(3, 1)$ entry is equal to 0. This is because, from (3.30) we have $\underline{P}_k \underline{U}_k^* \underline{P}_{k+1} \underline{U}_{k+1}^* = 0$. Again, the $(2, 2)$ entry is equal to 0 as we have from (6.2) that

$$\underline{P}_k \underline{U}_k^* T^* \tilde{U}_{k+1} \underline{Q}_{k+1} = \underline{P}_k \underline{U}_k^* \underline{P}_{k+1} \underline{U}_{k+1}^* T^*.$$

It follows from (3.32) that the $(1, 2)$ entry

$$D_{T^*} \tilde{U}_k \underline{Q}_k \tilde{U}_{k+1} \underline{Q}_{k+1} = 0.$$

Hence $\underline{V}_k D_{k+1} + \underline{D}_k E_{k+1} = \underline{D}_{k+1}$ if and only if

- (a) $\underline{T}_k D_{T^*} \tilde{U}_{k+1} \underline{Q}_{k+1} + D_{T^*} \tilde{U}_k \underline{Q}_k \tilde{U}_{k+1} \underline{Q}_{k+1}^\perp = D_{T^*} \tilde{U}_{k+1} \underline{Q}_{k+1}$ (as maps from \mathcal{D}_{T^*} to \mathcal{H}) and
- (b) $\underline{P}_k \underline{U}_k^* D_{T^*} \tilde{U}_{k+1} \underline{Q}_{k+1} - \underline{P}_k^\perp \underline{U}_k^* \underline{P}_{k+1} \underline{U}_{k+1}^* T^* - \underline{P}_k \underline{U}_k^* T^* \tilde{U}_{k+1} \underline{Q}_{k+1}^\perp = -\underline{P}_{k+1} \underline{U}_{k+1}^* T^*$ (as maps from \mathcal{D}_{T^*} to \mathcal{D}_T).

First we show that for $m = 1, \dots, n$,

$$\underline{T}_m D_{T^*} = D_{T^*} \tilde{U}_m \underline{Q}_m^\perp + T D_{T^*} \tilde{U}_m \underline{Q}_m. \quad (6.7)$$

We prove this inductively as follows. From condition-(1') we have that (6.7) holds for $m = 1$. Suppose it holds for $m = k$ for some $k \in \mathbb{N}$. We will prove that it holds for $m = k + 1$. Note that

$$\begin{aligned} \underline{T}_{k+1} D_{T^*} &= \underline{T}_{k+1} (D_{T^*} \tilde{U}_k \underline{Q}_k^\perp + T D_{T^*} \tilde{U}_k \underline{Q}_k) \\ &= \underline{T}_{k+1} D_{T^*} \tilde{U}_k \underline{Q}_k^\perp + \underline{T}_{k+1} T D_{T^*} \tilde{U}_k \underline{Q}_k \\ &= (D_{T^*} \tilde{U}_{k+1} \underline{Q}_{k+1}^\perp + T D_{T^*} \tilde{U}_{k+1} \underline{Q}_{k+1}) \tilde{U}_k \underline{Q}_k^\perp + T (D_{T^*} \tilde{U}_{k+1} \underline{Q}_{k+1}^\perp + T D_{T^*} \tilde{U}_{k+1} \underline{Q}_{k+1}) \tilde{U}_k \underline{Q}_k \\ &\quad [\text{by condition} - (1')] \\ &= D_{T^*} \tilde{U}_{k+1} \underline{Q}_{k+1}^\perp \tilde{U}_k \underline{Q}_k^\perp + T D_{T^*} (\tilde{U}_{k+1} \underline{Q}_{k+1} \tilde{U}_k \underline{Q}_k^\perp + \tilde{U}_{k+1} \underline{Q}_{k+1}^\perp \tilde{U}_k \underline{Q}_k) \\ &\quad + T^2 D_{T^*} \tilde{U}_{k+1} \underline{Q}_{k+1} \tilde{U}_k \underline{Q}_k \quad [\text{by (3.32), (3.33)}] \\ &= D_{T^*} \tilde{U}_{k+1} \underline{Q}_{k+1}^\perp + T D_{T^*} \tilde{U}_{k+1} \underline{Q}_{k+1}. \end{aligned}$$

Hence by induction (6.7) holds for all $k = 1, \dots, n$. For proving (a) we first observe that

$$\begin{aligned} &\underline{T}_k D_{T^*} \tilde{U}_{k+1} \underline{Q}_{k+1} + D_{T^*} \tilde{U}_k \underline{Q}_k \tilde{U}_{k+1} \underline{Q}_{k+1}^\perp \\ &= \underline{T}_k D_{T^*} \tilde{U}_{k+1} \underline{Q}_{k+1} - D_{T^*} \tilde{U}_k \underline{Q}_k^\perp \tilde{U}_{k+1} \underline{Q}_{k+1} + D_{T^*} \tilde{U}_{k+1} \underline{Q}_{k+1} \\ &= (\underline{T}_k D_{T^*} - D_{T^*} \tilde{U}_k \underline{Q}_k^\perp) \tilde{U}_{k+1} \underline{Q}_{k+1} + D_{T^*} \tilde{U}_{k+1} \underline{Q}_{k+1} \\ &= T D_{T^*} \tilde{U}_k \underline{Q}_k \tilde{U}_{k+1} \underline{Q}_{k+1} + D_{T^*} \tilde{U}_{k+1} \underline{Q}_{k+1} \quad [\text{by (6.7)}] \\ &= D_{T^*} \tilde{U}_{k+1} \underline{Q}_{k+1}. \quad [\text{by (3.32)}] \end{aligned}$$

This proves (a). Now for proving (b) we have

$$\begin{aligned} &\underline{P}_k \underline{U}_k^* D_{T^*} \tilde{U}_{k+1} \underline{Q}_{k+1} - \underline{P}_k^\perp \underline{U}_k^* \underline{P}_{k+1} \underline{U}_{k+1}^* T^* - \underline{P}_k \underline{U}_k^* T^* \tilde{U}_{k+1} \underline{Q}_{k+1}^\perp \\ &= \underline{P}_k \underline{U}_k^* (D_{T^*} \tilde{U}_{k+1} \underline{Q}_{k+1} - T^* \tilde{U}_{k+1} \underline{Q}_{k+1}^\perp) - \underline{P}_k^\perp \underline{U}_k^* \underline{P}_{k+1} \underline{U}_{k+1}^* T^* \\ &= \underline{P}_k \underline{U}_k^* \underline{P}_{k+1} \underline{U}_{k+1}^* D_{T^*} \tilde{U}_{k+1} \underline{Q}_{k+1} - \underline{P}_k \underline{U}_k^* \underline{P}_{k+1} \underline{U}_{k+1}^* T^* - \underline{P}_k^\perp \underline{U}_k^* \underline{P}_{k+1} \underline{U}_{k+1}^* T^* \quad [\text{by (6.6)}] \\ &= -\underline{P}_{k+1} \underline{U}_{k+1}^* T^*. \quad [\text{by (3.30)}] \end{aligned}$$

This proves (b).

7. DATA AVAILABILITY STATEMENT

- (1) Data sharing is not applicable to this article as no datasets were generated or analysed during the current study.
- (2) In case any datasets are generated during and/or analysed during the current study, they must be available from the corresponding author on reasonable request.

REFERENCES

- [1] J. Agler, The Arveson extension theorem and coanalytic models, *Integral Equations Operator Theory*, 5 (1982), 608 – 631.
- [2] C. Ambrozie and V. Müller, Commutative dilation theory, *Operator Theory*, ed. by D. Alpay (Springer, Berlin, 2014).
- [3] T. Ando, On a pair of commutative contractions, *Acta Sci. Math. (Szeged)*, 24 (1963), 88 – 90.
- [4] T. Ando, Unitary dilation for a triple of commuting contractions, *Bull. Acad. Polon. Sci. Sr. Sci. Math. Astronom. Phys.*, 24 (1976), 851 – 853.
- [5] A. Arias and G. Popescu, Noncommutative interpolation and Poisson transforms, *Israel J. Math.*, 115 (2000), 205 – 234.
- [6] A. Arias and G. Popescu, Noncommutative interpolation and Poisson transforms II, *Houston J. Math.*, 25 (1999), 79 – 98.
- [7] W. Arveson, Subalgebras of C^* -algebras. II, *Acta Math.*, 128 (1972), 271 – 308.
- [8] B. Bagchi, T. Bhattacharyya and G. Misra, Some thoughts on Andos theorem and Parrotts example, *Linear Algebra Appl.*, 341 (2002), 357 – 367.
- [9] J. A. Ball, W. S. Li, D. Timotin and T. T. Trent, A commutant lifting theorem on the polydisc, *Indiana Univ. Math. J.*, 48 (1999), 653 – 675.
- [10] J. A. Ball and H. Sau, Functional models for commuting Hilbert-space contractions, *Operator theory, operator algebras and their interactions with geometry and topology, Ronald G. Douglas memorial volume*, 11 – 54, *Oper. Theory Adv. Appl.*, 278, Birkhuser/Springer, Cham, [2020].
- [11] J. Ball, T. Trent and V. Vinnikov, Interpolation and commutant lifting for multipliers on reproducing kernel Hilbert spaces, *Operator theory and analysis, Oper. Theory Adv. Appl.*, 122, Birkhäuser, Basel, (2001), 89 – 138.
- [12] S. Barik, B. K. Das, K. J. Haria and J. Sarkar, Isometric dilations and von Neumann inequality for a class of tuples in the polydisc, *Trans. Amer. Math. Soc.*, 372 (2019), 1429 – 1450.
- [13] H. Bercovici, C. Foias, L. Kerchy and B. Sz.-Nagy, Harmonic analysis of operators on Hilbert space, Universitext, Springer, New York, (2010).
- [14] H. Bercovici, R.G. Douglas and C. Foias, On the classification of multi-isometries, *Acta Sci.Math. (Szeged)*, 72 (2006), 639 – 661.
- [15] C. A. Berger, L. A. Coburn and A. Lebow, Representation and index theory for C^* -algebras generated by commuting isometries, *J. Functional Analysis*, 27 (1978), 51 – 99.
- [16] T. Bhattacharyya, The tetrablock as a spectral set, *Indiana Univ. Math. J.*, 63 (2014), no. 6, 1601 – 1629.

- [17] T. Bhattacharyya and H. Sau, Explicit and unique construction of tetrablock unitary dilation in a certain case, *Complex Anal. Oper. Theory*, 10 (2016), 749 – 768.
- [18] S. Brehmer, Über vetauschbare Kontraktionen des Hilbertschen Raumes, *Acta Sci. Math. (Szeged)*, 22 (1961), 106 – 111.
- [19] Z. Burdak, On the model and invariant subspaces for pairs of commuting isometries, *Integral Equations Operator Theory*, 91 (2019), 1 – 23.
- [20] Z. Ceauescu and I. Suciu, Isometric dilations of commuting contractions. I, *J. Operator Theory*, 12 (1984), 65 – 88.
- [21] R. Curto and F. H. Vasilescu, Standard operator models in the polydisc, *Indiana Univ. Math. J.*, 42 (1993), 791 – 810.
- [22] R. Curto and F. H. Vasilescu, Standard operator models in the polydisc II., *Indiana Univ. Math. J.*, 44 (1995), 727 – 746.
- [23] B. K. Das and J. Sarkar, Ando dilations, von Neumann inequality, and distinguished varieties, *J. Funct. Anal.*, 272 (2017), 2114 – 2131.
- [24] B. K. Das, S. Sarkar and J. Sarkar, Factorizations of contractions, *Adv. Math.*, 322 (2017), 186 – 200.
- [25] S. Dey, Standard commuting dilations and liftings, *Colloq. Math.*, 126 (2012), 87 – 94.
- [26] R. G. Douglas, P. S. Muhly and C. Pearcy, Lifting commuting operators, *Michigan Math. J.*, 15 (1968), 385 – 395.
- [27] J. Eschmeier, Invariant subspaces for commuting contractions, *J. Operator Theory*, 45 (2001), 413 – 443.
- [28] C. Foias and B. Sz.-Nagy, Analyse harmonique des opérateurs de l’espace de Hilbert, *Akademiai Kiado, Budapest*, (1967).
- [29] A. Grinshpan, D.S. Kaliuzhnyi-Verbovetskyi, V. Vinnikov and H.J. Woerdeman, Classes of tuples of commuting contractions satisfying the multivariable von Neumann inequality, *J. Funct. Anal.*, 256 (2009), 3035 – 3054.
- [30] P. R. Halmos, Normal dilations and extensions of operators, *Summa Brasil. Math.*, 2 (1950), 125 – 134.
- [31] J. A. Holbrook, Schur norms and the multivariate von Neumann inequality. Recent advances in operator theory and related topics (Szeged, 1999), 375 – 386, *Oper. Theory Adv. Appl.*, 127, Birkhuser, Basel, (2001).
- [32] G. Knese, The von Neumann inequality for 3×3 matrices, *Bull. London Math. Soc.*, 48 (2016), 53 – 57.
- [33] E. Levy and Orr Moshe Shalit, Dilation theory in finite dimensions: the possible, the impossible and the unknown, *Rocky Mountain J. Math.*, 44 (2014), 203 – 221.
- [34] J. McCarthy and Orr Shalit, Unitary n-dilations for tuples of commuting matrices, *Proc. Amer. Math. Soc.*, 141 (2013), 563 – 571.
- [35] V. Müller and F. H. Vasilescu, Standard models for some commuting multishifts, *Proc. Amer. Math. Soc.*, 117 (1993), 979 – 989.
- [36] B. Sz.-Nagy, Sur les contractions de l’espace de Hilbert, *Acta Sci. Math. (Szeged)*, 15 (1953), 87 – 92.
- [37] B. Sz.-Nagy, On Schäffers construction of unitary dilations, *Ann. Univ. Sci. Budapest. Eötvös Sect. Math.*, 3(4) (1960/1961), 343 – 346.
- [38] D. Opřla, A generalization of Andô’s theorem and Parrotts example, *Proc. Amer. Math. Soc.*, 134 (2006), 2703 – 2710.

- [39] S. Pal, Common reducing subspaces and decomposition of contractions, *Forum Math.*, To appear.
- [40] S. Pal, Triangular tetrablock-contractions, factorization of contractions, dilation and subvarieties, <https://arxiv.org/abs/2204.11387>.
- [41] S. Pal and P. Sahasrabuddhe, Minimal isometric dilations and operator models for the polydisc, <https://arxiv.org/abs/2204.11391>.
- [42] S. Parrott, Unitary dilations for commuting contractions, *Pacific J. Math.*, 34 (1970), 481 – 490.
- [43] M. Ptak, Unitary dilations of multiparameter semigroups of operators, *Ann. Polon. Math.*, 45 (1985), 237 – 243.
- [44] J. J. Schäffer, On unitary dilations of contractions, *Proc. Amer. Math. Soc.*, 6 (1955), 322.
- [45] Orr M. Shalit, Dilation theory: a guided tour. Operator theory, functional analysis and applications, 551 – 623, *Oper. Theory Adv. Appl.*, 282, Birkhuser/Springer, Cham, (2021).
- [46] J. Stochel and F.H. Szafraniec, Unitary dilation of several contractions, *Oper. Theory Adv. Appl.*, 127 (2001), 585 – 598.
- [47] D. Timotin, Regular dilations and models for multicontractions, *Indiana Univ. Math. J.*, 47 (1998), 671 – 684.
- [48] N. Varopoulos, On an inequality of von Neumann and an application of the metric theory of tensor products to operator theory, *J. Funct. Anal.*, 16 (1974), 83 – 100.
- [49] F. H. Vasilescu, An operator-valued Poisson kernel, *J. funct. analysis*, 110 (1992), 47 – 72.
- [50] F. H. Vasilescu, Positivity conditions and standard models for commuting multioperators. Multivariable operator theory (Seattle, WA, 1993), 347 – 365, *Contemp. Math.*, 185, Amer. Math. Soc., Providence, RI, (1995).
- [51] J. von Neumann, Eine Spektraltheorie für allgemeine Operatoren eines unitären Raumes, *Math. Nachr.*, 4 (1951), 258 – 281.

(Sourav Pal) MATHEMATICS DEPARTMENT, INDIAN INSTITUTE OF TECHNOLOGY BOMBAY, POWAI, MUMBAI - 400076, INDIA.

Email address: sourav@math.iitb.ac.in

(Prajakta Sahasrabuddhe) MATHEMATICS DEPARTMENT, INDIAN INSTITUTE OF TECHNOLOGY BOMBAY, POWAI, MUMBAI - 400076, INDIA.

Email address: prajakta@math.iitb.ac.in