

Gauge bosons masses in the context of the Supersymmetric $SU(3)_C \otimes SU(3)_L \otimes U(1)_N$ Model

M. C. Rodriguez

Grupo de Física Teórica e Matemática Física

Departamento de Física

Universidade Federal Rural do Rio de Janeiro - UFRRJ

BR 465 Km 7, 23890-000, Seropédica - RJ

Brasil

Abstract

We analyze the boson masses and their mixing in the Minimal Supersymmetric $SU(3)_C \otimes SU(3)_L \otimes U(1)_N$ Model, and we will show all the numerical results are in agreement with actual current experimental limits.

PACS number(s): 12.60. Jv

Keywords: Supersymmetric models

1 Introduction

Recently the CDF Collaboration at Fermilab presented its new High-precision measurement, with an accuracy of $\sim 10^{-4}$, of the W -boson mass [1]

$$(M_W)_{CDF} = (80.4335 \pm 0.0094) \text{ GeV}, \quad (1)$$

this measurement represent an excess bigger than 6σ in relation to the more precise value (*conservative* scenario) of the Standard Model (SM) [2, 3] which is given by [4]

$$(M_W)_{SM} = (80.3505 \pm 0.0077) \text{ GeV}. \quad (2)$$

Clearly if this result, presented by the CDF, is confirmed by other experimental collaborations, this mean a new indication of physics beyond the SM.

From the theoretical point of view, the SM cannot be a fundamental theory since it has so many questions like that of the number of families do

not have an answer in its context. One of these possibilities to solve this problem is that, at energies of a few TeVs, the gauge symmetry may be

$$SU(3)_C \otimes SU(3)_L \otimes U(1)_N, \quad (3)$$

(or it is more known as M331 for shortness) instead of that of the SM [5, 6]. They are interesting possibilities for the physics at the TeV scale and we can also accomodate the CDF data [7]. When σ_1 get non zero value the neutrinos get the following Majorana mass term [8]

$$\frac{G_{ab}}{2\sqrt{2}} < \sigma_1^0 > . \quad (4)$$

The supersymmetric version of the 331 minimal model, or it is more known as MSUSY331 for shortness, was considered [9, 10, 11, 12]. In this model beyond the anti-sextet S , we need to introduce a Sextet S' Higgs boson, to cancel chiral anomalies generated by the superpartners of S in similar way as we have to add a new doublet scalar at Minimal Supersymmetric Standard Model (MSSM).

When all the neutral fields in sextet and anti-sextet get vev, the neutrinos obtain mass via the see-saw mechanism, due to the mix between higgsinos with the usual leptons, which is given by:

$$\mu_{0i} \hat{L}_i \hat{\eta}' + \lambda_{2ij} (\epsilon \hat{L}_i \hat{L}_j \hat{\eta}) + \lambda_{3ij} (\hat{L}_i \hat{S} \hat{L}_j) + \lambda_{4i} (\epsilon \hat{L}_i \hat{\chi} \hat{\rho}), \quad (5)$$

in a similar way to what we have done [13]. It must be explore it in the future.

The vaccum expectation value of this new scalar field, S' , can explain the shift on the W mass as we will shown below. We can decompose our scalars fields in $SU(2) \otimes U(1)$ representations [14, 15, 16]

$$\mathbf{6}_0 \rightarrow \mathbf{3}_2 \oplus \mathbf{2}_{-1} \oplus \mathbf{1}_{-4} \quad (6)$$

where

$$\begin{aligned} S &= \begin{pmatrix} T & \frac{\Phi_S}{\sqrt{2}} \\ \frac{\Phi_S^T}{\sqrt{2}} & H_2^{--} \end{pmatrix} \sim (\mathbf{1}, \mathbf{6}^*, 0), \\ T &= \begin{pmatrix} \sigma_1^0 & \frac{h_1^+}{\sqrt{2}} \\ \frac{h_1^+}{\sqrt{2}} & H_1^{--} \end{pmatrix} \sim (\mathbf{1}, \mathbf{3}, +2), \\ \Phi_S &= \begin{pmatrix} h_2^+ \\ \sigma_2^0 \end{pmatrix} \sim (\mathbf{1}, \mathbf{2}, -1), \quad H_2^{--} \sim (\mathbf{1}, \mathbf{1}, -4). \end{aligned} \quad (7)$$

Φ_S is the Higgs doublet of SM while T was considered at [17]. One of main goal at Large Hadron Collider (LHC) is to find new physics beyond the SM. Then it is useful to predict the masses of new particles arising in some interesting models. We want to present a detailed analyses of the gauge bosons masses in similar way as we have done in the scalar sector [18, 19]. We have also presented a previous mass spectrum of this model, without the Sextet [12, 20], in this article we have also studied the processes

$$g, d \rightarrow U^{--} + J, \quad g, u \rightarrow U^{--} + j_\alpha. \quad (8)$$

These exotic quarks have charge $(5/3)e$ and $-(4/3)e$, respectively. These processes was suggested by Alexandre Belyaev and its signature is llX and it can be detected at LHC if they really exist in nature.

We will present in next Section, Sec.(2), the Minimal Supersymmetric 331 Model, (MSUSY331), and our numerical results are present at Sec.(3). Our conclusions are presented at the last Section of this article. We present in appendices the analytical analysis of the mixtures of the gauge bosons of this model and some numerical analyses about the mixing in the neutral boson sector..

2 Minimal Supersymmetric 331 Model (MSUSY331)

The scalars in this model are given by [11, 12]:

$$\begin{aligned} \eta &= \begin{pmatrix} \eta^0 \\ \eta_1^- \\ \eta_2^+ \end{pmatrix} \sim (\mathbf{1}, \mathbf{3}, 0), \quad \rho = \begin{pmatrix} \rho^+ \\ \rho^0 \\ \rho^{++} \end{pmatrix} \sim (\mathbf{1}, \mathbf{3}, +1), \\ \chi &= \begin{pmatrix} \chi^- \\ \chi^{--} \\ \chi^0 \end{pmatrix} \sim (\mathbf{1}, \mathbf{3}, -1), \quad \tilde{\chi}' = \begin{pmatrix} \tilde{\chi}'^{++} \\ \tilde{\chi}'^{+-} \\ \tilde{\chi}'^0 \end{pmatrix} \sim (\mathbf{1}, \mathbf{3}^*, +1), \\ \tilde{\eta}' &= \begin{pmatrix} \tilde{\eta}'^0 \\ \tilde{\eta}'^+ \\ \tilde{\eta}'^- \end{pmatrix} \sim (\mathbf{1}, \mathbf{3}^*, 0), \quad \tilde{\rho}' = \begin{pmatrix} \tilde{\rho}'^- \\ \tilde{\rho}'^0 \\ \tilde{\rho}'^{--} \end{pmatrix} \sim (\mathbf{1}, \mathbf{3}^*, -1), \\ S &= \begin{pmatrix} \sigma_1^0 & \frac{h_2^+}{\sqrt{2}} & \frac{h_1^-}{\sqrt{2}} \\ \frac{h_2^+}{\sqrt{2}} & H_1^{++} & \frac{\sigma_2^0}{\sqrt{2}} \\ \frac{h_1^-}{\sqrt{2}} & \frac{\sigma_2^0}{\sqrt{2}} & H_2^{--} \end{pmatrix} \sim (\mathbf{1}, \mathbf{6}^*, 0), \end{aligned}$$

$$\tilde{S}' = \begin{pmatrix} \tilde{\sigma}_1'^0 & \frac{\tilde{h}_2'^-}{\sqrt{2}} & \frac{\tilde{h}_1'^+}{\sqrt{2}} \\ \frac{\tilde{h}_2'^-}{\sqrt{2}} & \tilde{H}_1'^{--} & \frac{\tilde{\sigma}_2'^0}{\sqrt{2}} \\ \frac{\tilde{h}_1'^+}{\sqrt{2}} & \frac{\tilde{\sigma}_2'^0}{\sqrt{2}} & \tilde{H}_2'^{++} \end{pmatrix} \sim (\mathbf{1}, \mathbf{6}, 0). \quad (9)$$

The charged gauge bosons of this model are:

$$\begin{aligned} W_m^\pm(x) &= -\frac{1}{\sqrt{2}}(V_m^1(x) \mp iV_m^2(x)), \\ V_m^\pm(x) &= -\frac{1}{\sqrt{2}}(V_m^4(x) \pm iV_m^5(x)), \\ U_m^{\pm\pm}(x) &= -\frac{1}{\sqrt{2}}(V_m^6(x) \pm iV_m^7(x)). \end{aligned} \quad (10)$$

while the neutral gauge bosons are the photon

$$A_m = \sin \theta_W (V_m^3 - \sqrt{3}V_m^8) + \sqrt{1 - 4 \sin^2 \theta_W} V_m, \quad (11)$$

where θ_W is the Weinberg angel and we have also two massives neutral bosons defined as

$$\begin{aligned} Z_m &= \cos \theta_W V_m^3 + \sqrt{3} \tan \theta_W \sin \theta_W V_m^8 - \tan \theta_W \sqrt{1 - 4 \sin^2 \theta_W} V_m, \\ Z'_m &= \frac{1}{\cos \theta_W} \left[\sqrt{1 - 4 \sin^2 \theta_W} V_m^8 + \sqrt{3} \sin \theta_W V_m \right]. \end{aligned} \quad (12)$$

In order to get the gauge bosons masses we have to calculate

$$\begin{aligned} \mathcal{L}_{Higgs}^{Escalar} &= (\mathcal{D}_m \eta)^\dagger (\mathcal{D}^m \eta) + (\mathcal{D}_m \rho)^\dagger (\mathcal{D}^m \rho) + (\mathcal{D}_m \chi)^\dagger (\mathcal{D}^m \chi) + (\overline{\mathcal{D}}_m \eta')^\dagger (\overline{\mathcal{D}}^m \eta') \\ &+ (\overline{\mathcal{D}}_m \rho')^\dagger (\overline{\mathcal{D}}^m \rho') + (\overline{\mathcal{D}}_m \chi')^\dagger (\overline{\mathcal{D}}^m \chi') + Tr[(\mathcal{D}_m S)^\dagger (\mathcal{D}^m S)] \\ &+ Tr[(\overline{\mathcal{D}}_m S')^\dagger (\overline{\mathcal{D}}^m S')]. \end{aligned} \quad (13)$$

The mass of charged gauge boson are given by [12, 20]

$$\begin{aligned} M_W^2 &= \frac{g^2}{4} \left[v_\eta^2 + v_\rho^2 + v_{\sigma_2}^2 + v_{\eta'}^2 + v_{\rho'}^2 + 2(v_{\sigma_1}^2 + v_{\sigma'_1}^2) + v_{\sigma'_2}^2 \right], \\ M_V^2 &= \frac{g^2}{4} \left[v_\eta^2 + v_\chi^2 + v_{\sigma_2}^2 + v_{\eta'}^2 + v_{\chi'}^2 + 2(v_{\sigma_1}^2 + v_{\sigma'_1}^2) + v_{\sigma'_2}^2 \right], \\ \delta_{WV} &= \frac{g^2}{\sqrt{2}} (v_{\sigma_1} v_{\sigma_2} + v_{\sigma'_1} v_{\sigma'_2}), \\ M_U^2 &= \frac{g^2}{4} (v_\rho^2 + v_\chi^2 + 4v_{\sigma_2}^2 + v_{\rho'}^2 + v_{\chi'}^2 + 4v_{\sigma'_2}^2). \end{aligned} \quad (14)$$

This new charged vector boson, V^\pm , is best known in the literature as being W' and in recent analyzes its masses have been considered in the following range [21]

$$0.25 \leq M_{W'} \leq 2 \text{ TeV}. \quad (15)$$

The charged gauge bosons can mixing. We present the analytical analysis of this mixture in Sec.(A). We also have, as in SM, the usual Z -boson and an extra neutral gauge boson known as Z' -boson and in the approximation that $v_\chi \simeq v_{\chi'}$ and they are bigger than the others vev of this model we get the following values [12]

$$\begin{aligned} M_Z^2 &\approx \frac{1}{2} \left(\frac{g^2 + 4g'^2}{g^2 + 3g'^2} \right) (v_\eta^2 + v_\rho^2 + v_{\sigma_2}^2 + v_{\eta'}^2 + v_{\rho'}^2 + v_{\sigma'_2}^2), \\ M_{Z'}^2 &\approx \frac{2}{3} (g^2 + 3g'^2) (v_\chi^2 + v_{\chi'}^2), \end{aligned} \quad (16)$$

using the first expression from Eq.(41), together with M_Z^2 , we can write

$$\frac{M_Z^2}{M_{W'}^2} = \frac{1 + 4t^2}{1 + 3t^2}, \quad (17)$$

where. we have defined the new parameter t in the following way:

$$t \equiv \frac{g'}{g}, \quad (18)$$

but from the SM we know the following result

$$\frac{M_Z^2}{M_W^2} = \frac{1}{1 - \sin^2 \theta_W}, \quad (19)$$

where θ_W is the Weak mixing angle. When we impose Eq.(19) is the same results are presented in Eq.(19), we get the famous relation

$$t^2 = \frac{\sin^2 \theta_W}{1 - 4 \sin^2 \theta_W}, \quad (20)$$

It imply

$$\sin^2 \theta_W < \frac{1}{4}, \quad (21)$$

with a Landau pole ¹ in

$$\sin^2 \theta_W(\mu) = \frac{1}{4}, \quad (22)$$

this result is in agreement with the actual experimental bound, given by

$$\sin^2 \theta_W = 1 - \frac{M_W^2}{M_Z^2} = 0.223562, \quad (23)$$

our neutral gauge bosons can mixing, the analytical results without this approximation is presented at Sec.(B). The limits on the Z' mass is given by [21]

$$1.0 \leq M_{Z'} \leq 4.5 \text{ TeV}. \quad (24)$$

3 Numerical Analyses for CDF Experimental Results.

The masses of the W -boson and Z -boson receive a new tree-level contribution given by:

$$\delta M_W^2 = \frac{g^2}{4} [2(v_{\sigma_1}^2 + v_{\sigma'_1}^2) + v_{\sigma'_2}^2], \quad \delta M_Z^2 = \frac{g^2}{4 \cos^2 \theta_W} [4(v_{\sigma_1}^2 + v_{\sigma'_1}^2) + v_{\sigma'_2}^2], \quad (25)$$

therefore, as first prevision of this model is

$$\delta M_W^2 \neq \delta M_Z^2, \quad (26)$$

our analytical results are in agreement with the results obtained for the M331 as presented by the references [8, 23, 24]. In order, to explain the W -mass anomaly we have to impose:

$$\left| \sqrt{2(v_{\sigma_1}^2 + v_{\sigma'_1}^2) + v_{\sigma'_2}^2} \right| = \frac{2}{g} \sqrt{|(M_W)_{SM}^2 - (M_W)_{CDF}^2|} = 11.19 \text{ GeV}. \quad (27)$$

This new contribution does not have any restrictions coming from the fermion mass explanation.

¹Exists an energy scale μ where the model loses its perturbative regime, see [22].

Recently it was proposed this shift can be easily explained by a real triplet Higgs boson [25, 26] and the triplet vev, v_T , is around 10 GeV. We can also use an extra complex triplet Higgs boson with $v_T \approx 3.2$ GeV [27] to explain this anomaly. There is also, an interesting proposal to explain this anomaly using the Minimal R -Symmetric extension of MSSM, known as MRSSM [28, 29, 30] and in this case $|v_T| \leq 4$ GeV [29]. The mechanism is based on the fact that in this model a new scalar field is introduced in the Triplet representation and it is not necessary to generate mass for the fermions [29, 30], the vev of this new scalar can be around 3 GeV [31].

By another hand, for the ρ -parameter, in our model, it is hold the following expression:

$$\begin{aligned}\rho &= \left(\frac{M_W^2}{M_Z^2 \cos^2 \theta_W} \right) = \frac{1+R}{1+R'}, \\ R &= 2 \left(\frac{v_{\sigma_1}^2 + v_{\sigma'_1}^2}{v_{MP}^2} \right) + \frac{v_{\sigma'_2}^2}{v_{MP}^2}, \quad R' = 4 \left(\frac{v_{\sigma_1}^2 + v_{\sigma'_1}^2}{v_{MP}^2} \right) + \frac{v_{\sigma'_2}^2}{v_{MP}^2}, \\ \rho &= 1 + R - R' = 1 - 2 \left(\frac{v_{\sigma_1}^2 + v_{\sigma'_1}^2}{v_{MP}^2} \right).\end{aligned}\tag{28}$$

remember the experimental measurement of this parameter is given by: $\rho = 0.9998 \pm 0.0008$, then we must satisfy the following inequality:

$$\sqrt{v_{\sigma_1}^2 + v_{\sigma'_1}^2} < \sqrt{\frac{0.0008}{2}} v_{MP} < 5 \text{ GeV}.\tag{29}$$

So we can explain the new CDF data for the W -mass, see Eq.(27), and the ρ -parameter, Eq.(29), if the new vev of this model are of the order of a few GeV. First result can be taken all the vev from the triplet equal to zero, it means

$$v_{\sigma'_1}^2 = v_{\sigma'_2}^2 = 0 \text{ GeV},\tag{30}$$

it imply $\rho = 1$ and also

$$\begin{aligned}\delta M_Z^2 &= \frac{\delta M_W^2}{\cos^2 \theta_W} \Rightarrow \delta M_Z^2 = 17.187592 \text{ GeV}^2, \\ v_{\sigma'_2} &= 11.1942 \text{ GeV}.\end{aligned}\tag{31}$$

When we take the

$$v_{\sigma'_1}^2 = v_{\sigma'_2}^2 = 0 \text{ GeV},\tag{32}$$

our Eq.(29) give the upper limit $v_{\sigma_1}^2 < 5 \text{ GeV}$ the same result given in [23], but in order to also satisfy Eq.(27), we must have

$$v_{\sigma_1} = 7.91552 \text{ GeV}, \quad (33)$$

therefore, this solution is not possible inside m331 model, but it is possible in MSUSY331.

We must have all new vev different of zero. We can use Eqs.(27,29) for fix $v_{\sigma'_2}$:

$$v_{\sigma'_2} = \sqrt{\frac{4\delta M_W^2}{g} - \text{erro} * v_{MP}^2} = 8.7691841 \text{ GeV}. \quad (34)$$

Using Eq.(29) we can define v_{σ_1} :

$$v_{\sigma_1} = \sqrt{\frac{2\delta M_W^2}{g} - \frac{v_{\sigma'_2}^2}{2} - v_{\sigma'_1}^2} \quad (35)$$

our results is shown in Fig.(1). Now we get the following numerical results:

$$\begin{aligned} \rho &= 0.9992, \\ \delta M_Z^2 &= 28.8278 \text{ GeV}^2, \Rightarrow (M_Z)_{CDF} = 91.3181 \text{ GeV}. \end{aligned} \quad (36)$$




Figure 1: Possible values for v_{σ_1} as function of $v_{\sigma'_1}$, in the minimal supersymmetric $SU(3)_C \otimes SU(3)_L \otimes U(1)_N$ Model, see Eq.(35).

We will consider, for the charged gauge bosons, the following two case:

$v_{\sigma'_2}$ GeV	M_U GeV	M_V GeV	$M_{Z'}$ GeV	$\delta_{ZZ'}$ GeV ²
12.0	468.506	461.584	2023.910	-21956.1
12.5	468.512	461.586	2023.910	-21956.0
13.0	468.518	461.587	2023.910	-21955.8
13.5	468.524	461.589	2023.910	-21955.6
14.0	468.530	461.590	2023.910	-21955.5
14.5	468.536	461.592	2023.910	-21955.3
15.0	468.543	461.593	2023.910	-21955.1
15.5	468.550	461.595	2023.910	-21954.9
16.0	468.557	461.597	2023.910	-21954.7

Table 1: Masses of new gauge bosons for some values of $v_{\sigma'_2}$ but $v_{\sigma_1} = v_{\sigma'_1} = 0$ GeV; $v_\chi = v_{\chi'} = 1000$ GeV and the values of M_V , see Eq.(15), and the values of $M_{Z'}$, Eq.(24) and they satisfy the bounds presented by Eqs.(15,24).

- a-) Only $v_{\sigma'_2} \neq 0$, in this case the gauge bosons W and V are the physical ones; Our numerical results are presented in Tabs.(1,2). Under this hypotheese, the mixing parameter is zero, see Eq.(41), then the bosons W and V are the physical ones. In this scenario we can easily conclude the following hierarque in the masses are

$$M_{Z'} > M_U > M_V, \quad (37)$$

this result is in agreement with the previous analysis presented in [20].

- b-) Consider all new vev are non zero, where W_1 and W_2 are the physical gauge bosons. Our results can be found in Tabs.(3,4). In this case we get the following results

- $M_{W_1} \cong M_W$ and $M_{W_2} \cong M_V$;
- $M_U = 468.476$ GeV and $M_{Z'} = 2023.910$ GeV.

The Eq.(37) is still hold in this case.

4 Conclusions

We show that in the context of Minimal Supersymmetric $SU(3)_C \otimes SU(3)_L \otimes U(1)_N$ Model, MSUSY331, we can explain both the new measurement on

$v_{\chi'}$ GeV	M_U GeV	M_V GeV	$M_{Z'}$ GeV
1000	468.498	461.582	2023.91
1500	593.809	588.368	2579.43
2000	734.150	729.757	3199.01
2500	882.379	878.726	3851.85
3000	1035.110	1032.000	4523.57
3500	1190.620	1187.910	5206.88
4000	1347.930	1345.550	5897.75
4500	1506.500	1504.360	6593.79
5000	1665.950	1664.020	7293.54

Table 2: Masses for all new gauge bosons of this model for some values of $v_{\chi'}$ with $v_{\sigma'_2} = 11.1942$ GeV; $v_{\chi} = 1000$ GeV; $v_{\sigma_1} = v_{\sigma'_1} = 0$ GeV and they are in agreement with the bounds at Eqs.(15,24). The values for $\delta_{ZZ'} = -21956.4$ GeV².

v_{σ_1} GeV	M_W GeV	M_V GeV	δ_{WV} GeV ²
0.0	80.4332	461.582	12.9285
0.5	80.4335	461.582	14.4345
1.0	80.4345	461.582	15.9406
1.5	80.4361	461.582	17.4467
2.0	80.4385	461.583	18.9528
2.5	80.4414	461.583	20.4588
3.0	80.4451	461.584	21.9649
3.5	80.4494	461.585	23.4710
4.0	80.4544	461.586	24.9771
4.5	80.4600	461.587	26.4831
5.0	80.4663	461.588	27.9892
5.5	80.4732	461.589	29.4953

Table 3: Masses of single charged gauge bosons and δ_{WV} for some values of v_{σ_1} for $v_{\chi} = v_{\chi'} = 1000$ GeV; $v_{\sigma'_1} = 4.89453$ GeV; $v_{\sigma'_2} = 8.7691841$ GeV.

$v_{\chi'}$ GeV	M_W GeV	M_V GeV	M_{W_1} GeV	M_{W_2} GeV
1000	80.4335	461.582	80.4335	461.582
1500	80.4335	588.368	80.4335	588.368
2000	80.4335	729.757	80.4335	729.757
2500	80.4335	878.726	80.4335	878.726
3000	80.4335	1032.000	80.4335	1032.000
3500	80.4335	1187.91	80.4335	1187.91
4000	80.4335	1345.55	80.4335	1345.55
4500	80.4335	1504.36	80.4335	1504.36
5000	80.4335	1664.02	80.4335	1664.02

Table 4: Masses of M_W , M_V , M_{W_1} and M_{W_2} , see Eq.(46), for some values of $v_{\chi'}$ for $v_\chi = 1000$ GeV; $v_{\sigma_1} 0.5$ GeV; $v_{\sigma'_1} = 4.89453$ GeV; $v_{\sigma'_2} = 8.7691841$ GeV.

the W -boson mass as well as the ρ -parameter if Eqs.(27,29) are satisfied simultaneously as we have showed in Eqs.(31,36) and also in our Fig.(1). We also calculated the masses of all the gauge bosons of this model and $M_{Z'} > M_U > M_V$ and their values are in agreement with the actual experimental data given at Eqs.(15,24).

Acknowledgments

We would like thanks V. Pleitez for useful discussions above 331 models and about this interesting research topic. We also to thanks IFT for the nice hospitality during my several visit to perform my studies about the severals 331 Models and also for done this article.

A Mixing between W and V .

We notice, from Eq.(41), if $v_{\sigma_1} \neq 0$ the gauge bosons W and V are no longer the physical ones, in agreement with the results presented at [14, 15, 7]. They can mixing and the physical bosons are $W_{1,2}^\pm$ and their masses are

$$\begin{aligned}
M_{W_1}^2 &= \frac{1}{2} \left(M_W^2 + M_V^2 - \sqrt{(M_W^2 - M_V^2)^2 + 4\delta_{WV}^2} \right), \\
M_{W_2}^2 &= \frac{1}{2} \left(M_W^2 + M_V^2 + \sqrt{(M_W^2 - M_V^2)^2 + 4\delta_{WV}^2} \right). \quad (38)
\end{aligned}$$

The physical eigenstates are defined as

$$\begin{pmatrix} W_{1m}^\pm \\ W_{2m}^\pm \end{pmatrix} = \begin{pmatrix} \cos \theta^\pm & -\sin \theta^\pm \\ \sin \theta^\pm & \cos \theta^\pm \end{pmatrix} \begin{pmatrix} W_m^\pm \\ V_m^\pm \end{pmatrix}, \quad (39)$$

the mixing angle is given by

$$\tan^2 \theta^\pm = \frac{M_{W_1}^2 - M_W^2}{M_{W_2}^2 - M_V^2}. \quad (40)$$

B Mixing between Z and Z' .

The neutral gauge bosons Z and Z' can mix and it is given by

$$\mathcal{L}^{\text{neutra}} = \begin{pmatrix} Z_m & Z'_m \end{pmatrix} \begin{pmatrix} M_Z^2 & \delta_{ZZ'} \\ \delta_{ZZ'} & M_{Z'}^2 \end{pmatrix} \begin{pmatrix} Z_m \\ Z'_m \end{pmatrix}, \quad (41)$$

where

$$\begin{aligned} M_Z^2 &= \frac{g^2 v_{MP}^2}{4 \cos^2 \theta_W}, \\ M_{Z'}^2 &= \frac{g^2}{4} \left[\frac{4v_{MP}^2}{3s} + \frac{2t^2 U^2}{s} + \frac{4s}{3} (w^2 + w'^2) \right], \\ \delta_{ZZ'} &= \frac{g^2}{4\sqrt{3}h_w s} [(V^2 - U^2) - 6t^2 U^2], \end{aligned} \quad (42)$$

compare those values with our result presented on Eqs.(16). We have defined

$$\begin{aligned} V^2 &= v^2 + 2y^2 + z^2 + v'^2 + 2y'^2 + z'^2, \\ U^2 &= u^2 + u'^2, \\ t^2 &= \frac{\sin^2 \theta_W}{1 - 4 \sin^2 \theta_W}, \\ s &= 1 + 3t^2, \\ v_{MP}^2 &= V^2 + U^2, \end{aligned} \quad (43)$$

to understand how to get the third expression above see Eqs.(19,20). We can show the following relations

$$\cos \theta_W = \frac{\sqrt{s}}{\sqrt{s + t^2}},$$

$$\begin{aligned}
\sin \theta_W &= \frac{t}{\sqrt{s+t^2}}, \\
\sqrt{s+t^2} &= \left(\sqrt{h_W} \right)^{-1}, \\
h_W &= 1 - 4 \sin^2 \theta_W.
\end{aligned} \tag{44}$$

The physical mass of physical gauge bosons are

$$\begin{pmatrix} Z_{1m}^0 \\ Z_{2m}^0 \end{pmatrix} = \begin{pmatrix} \cos \theta^0 & -\sin \theta^0 \\ \sin \theta^0 & \cos \theta^0 \end{pmatrix} \begin{pmatrix} Z_m^0 \\ (Z')_m^0 \end{pmatrix}, \tag{45}$$

where the mixing angle in this sector is

$$\tan^2 \theta^0 = \frac{M_{Z_1}^2 - M_Z^2}{M_{Z_2}^2 - M_{Z'}^2}, \tag{46}$$

where

$$\begin{aligned}
M_{Z_1}^2 &= \frac{1}{2} \left(M_Z^2 + M_{Z'}^2 + \sqrt{(M_Z^2 - M_{Z'}^2)^2 + 4\delta_{ZZ'}^2} \right), \\
M_{Z_2}^2 &= \frac{1}{2} \left(M_Z^2 + M_{Z'}^2 - \sqrt{(M_Z^2 - M_{Z'}^2)^2 + 4\delta_{ZZ'}^2} \right).
\end{aligned} \tag{47}$$

C Preliminar Numerical Analysis for Z and Z'

For our numerical analyses, at tree-level, we will apply the constraint²

$$V_\eta^2 + V_\rho^2 + 2V_2^2 = (246 \text{ GeV})^2, \tag{48}$$

coming from M_W , where, we have defined

$$\begin{aligned}
V_\eta^2 &= v_\eta^2 + v_{\eta'}^2, \quad V_\rho^2 = v_\rho^2 + v_{\rho'}^2, \\
V_2^2 &= v_{\sigma_2}^2 + v_{\sigma_2'}^2.
\end{aligned} \tag{49}$$

Assuming that

$$v_\eta = 20, \quad v_{\sigma_2} = 10, \quad v_{\eta'} = v_{\rho'} = 1 \text{ GeV}, \tag{50}$$

²As we have choose at [11].

The value of v_ρ is fixed by Eq.(2), we get

$$v_\rho = 245.20 \text{ GeV}. \quad (51)$$

As an example as the mixing of Z and Z' we use the vev defined in Eq.(50) together the following vev

$$\begin{aligned} v_{\sigma_1} &= v_{\sigma'_1} = v_{\sigma'_2} = 0 \text{ GeV}, \\ v_\chi &= v_{\chi'} = 1000 \text{ GeV}, \end{aligned} \quad (52)$$

in Eq.(42) to get the following results³

$$\begin{aligned} M_Z &= 91.1875 \text{ GeV}, \quad M_{Z'} = 2023.91 \text{ GeV}, \\ \delta_{ZZ'} &= -21958 \text{ GeV}^2, \end{aligned} \quad (53)$$

the value for Z' mass is in agreement with Eq.(24). Repare $\delta_{ZZ'} \neq 0$, therefore Z and Z' are not physical gauge boson. The masses of physical gauge bosons $Z_{1,2}$, using Eq.(47), are⁴

$$M_{Z_1} = 91.1875 \text{ GeV}, \quad M_{Z_2} = 2023.91 \text{ GeV}. \quad (54)$$

Therefore the bosons Z and Z' , as a first approximation, can be considered as being the physical states, and their masses given at Eq.(42).

References

- [1] T. Aaltonen *et al* (CDF Collaboration), *High-precision measurement of the W boson mass with the CDF II detector*, *Science* **376**, 170, (2022).
- [2] S. J. L. Rosner, *Resource letter: The Standard model and beyond*, *Am. J. Phys.* **71**, 302, (2003).
- [3] A. S. Kronfeld and C. Quigg, *Resource Letter: Quantum Chromodynamics*, *Am. J. Phys.* **78**, 1081, (2010).

³The parameter $\delta_{ZZ'}$ has no dependence in v_χ and $v_{\chi'}$, see Eq.(42), do it we will shown only this specific case, see also our Tabs.(1,2).

⁴Compare $\delta_{ZZ'}$ with M_Z^2 and $M_{Z'}^2$, and it is smaller therefore the mixing parameter is almost negligible.

- [4] J. de Blas, M. Pierini, L. Reina and L. Silvestrini, *Impact of the recent measurements of the top-quark and W-boson masses on electroweak precision fits*, [arXiv:2204.04204].
- [5] F. Pisano and V. Pleitez, *An $SU(3) \otimes U(1)$ model for electroweak interactions*, *Phys. Rev.***D46**, 410, (1992).
- [6] R. Foot, O. F. Hernandez, F. Pisano and V. Pleitez, *Lepton masses in an $SU(3)_L \otimes U(1)_N$ gauge model*, *Phys. Rev. D* **47**, 4158, (1993).
- [7] D. Van Loi and P. Van Dong, *Novel effects of the W-boson mass shift in the 3-3-1 model*, *Eur. Phys. J.***C83**, 56, (2023); [arXiv:2206.10100 [hep-ph]].
- [8] F. Pisano and S. S. Sharma, *Majoron emitting neutrinoless double beta decay in the electroweak chiral gauge extensions*, *Phys. Rev.***D57**, 5670, (1998).
- [9] T. V. Duong and E. Ma, *Supersymmetric $SU(3) \otimes U(1)$ Gauge Model: Higgs Structure at the Electroweak Energy Scale*, *Phys. Lett.***B316**, 307 (1993).
- [10] H. N. Long and P. B. Pal, *Nucleon instability in a supersymmetric $SU(3)_C \otimes SU(3)_L \otimes U(1)$ model*, *Mod. Phys. Lett.***A13**, 2355, (1998).
- [11] J. C. Montero, V. Pleitez and M. C. Rodriguez, *A Supersymmetric 3-3-1 model*, *Phys. Rev. D***65**, 035006, (2002).
- [12] M. Capdequi-Peyranère and M.C. Rodriguez, *Charginos and neutralinos production at 3-3-1 supersymmetric model in $e^- e^-$ scattering*, *Phys. Rev. D* **65**, 035001 (2002).
- [13] J. C. Montero, V. Pleitez and M. C. Rodriguez, *Lepton masses in a supersymmetric 3-3-1 model*, *Phys. Rev.***D65**, 095008, (2002).
- [14] J. T. Liu and D. Ng, *Lepton flavor changing processes and CP violation in the 331 model*, *Phys. Rev.***D50**, 548, (1994), [arXiv:hep-ph/9401228 [hep-ph]].
- [15] J. C. Montero, C. A. de Sousa Pires and V. Pleitez, *Spontaneous breaking of a global symmetry in a 331 model*, *Phys. Rev.***D60**, 115003, (1999), [arXiv:hep-ph/9903251 [hep-ph]].

- [16] G. De Conto, A. C. B. Machado and V. Pleitez, *Minimal 3-3-1 model with a spectator sextet*, *Phys. Rev.***D92**, 075031, (2015), [arXiv:1505.01343 [hep-ph]].
- [17] G. B. Gelmini and M. Roncadelli, *Left-Handed Neutrino Mass Scale and Spontaneously Broken Lepton Number*, *Phys. Lett.***B99**, 411 (1981).
- [18] M. C. Rodriguez, *The Scalar Potential of Supersymmetric $SU(3)_C \otimes SU(3)_L \otimes U(1)_N$ Model*, [arXiv:2111.15393 [hep-ph]].
- [19] M. C. Rodriguez, *Scalar sector in the minimal supersymmetric 3-3-1 model*, *Int. J. Mod. Phys.***A21**, 4303, (2006).
- [20] M. C. Rodriguez, *Mass Spectrum in the Minimal Supersymmetric 3-3-1 model*, *J. Mod. Phys.***2**, 1193, (2011).
- [21] P. Osland, A. A. Pankov and I. A. Serenkova, *Bounds on the mass and mixing of Z' and W' bosons decaying into different pairings of W, Z , or Higgs bosons using CMS data at the LHC*, [arXiv:2206.01438] [hep-ph].
- [22] A. G. Dias, R. Martinez and V. Pleitez, *Concerning the Landau pole in 3-3-1 models*, *Eur. Phys. J.***C39**, 101, (2005); [arXiv:hep-ph/0407141 [hep-ph]].
- [23] J. C. Montero, C. A. de Sousa Pires and V. Pleitez, *Comment on ‘Majoron emitting neutrinoless double beta decay in the electroweak chiral gauge extensions’*, *Phys. Rev.***60**, 098701, (1999).
- [24] S. S. Sharma and F. Pisano, *Reply to ‘comment on: ‘Majoron emitting neutrinoless double beta decay in the electroweak chiral gauge extensions’*, *Phys. Rev.***60**, 098702, (1999).
- [25] T. Blank and W. Hollik, *Precision observables in $SU(2) \times U(1)$ models with an additional Higgs triplet*, *Nucl. Phys.***B514**, 113-134, (1998); [arXiv:hep-ph/9703392 [hep-ph]].
- [26] M. C. Chen, S. Dawson and T. Krupovnickas, *Higgs triplets and limits from precision measurements*, *Phys. Rev.***D74**, 035001, (2006); [arXiv:hep-ph/0604102 [hep-ph]].
- [27] J. L. Evans, T. T. Yanagida and N. Yokozaki, *W boson mass anomaly and grand unification*, [arXiv:2205.03887] [hep-ph].

- [28] G. D. Kribs, E. Poppitz, and N. Weiner, *Flavor in supersymmetry with an extended R-symmetry*, *Phys. Rev.* **D78**, 055010, (2008).
- [29] P. Diessner, J. Kalinowski, W. Kotlarski, and D. Stöckinger, *Higgs boson mass and electroweak observables in the MRSSM*, *JHEP* **12**, 124, (2014).
- [30] P. Diessner and G. Weiglein, *Precise prediction for the W boson mass in the MRSSM*, *JHEP* **07**, 011, (2019).
- [31] P. Athron, M. Bach, D. H. J. Jacon, W. Kotlarski, D. Stöckinger and A. Voigt, *Precise calculation of the W boson pole mass beyond the Standard Model with FlexibleSUSY*, [arXiv:2204.05285] [hep-ph].