

SPHERICAL CONTACT TORIC MANIFOLDS

HUI LI

ABSTRACT. Let (M, α) be a $2n + 1$ -dimensional connected compact contact toric manifold of Reeb type. Suppose the contact form α is regular, we find conditions under which M is homeomorphic to S^{2n+1} .

Let (M, α) be a contact manifold of dimension $2n + 1$. If an $n + 1$ -dimensional torus T^{n+1} acts on M effectively, preserving the contact form α , then M is called a **contact toric manifold**. If the Reeb vector field of the contact form α is generated by a one parameter subgroup of T^{n+1} , then the T^{n+1} -manifold M is called a **contact toric manifold of Reeb type**. Compact contact toric manifolds are classified by Lerman [11]. Most of the compact contact toric manifolds are of Reeb type.

Suppose (M, α) is a connected compact contact toric manifold of dimension $2n + 1$. Let \mathfrak{t} be the Lie algebra of the torus T^{n+1} , and \mathfrak{t}^* the dual Lie algebra. The **contact moment map** $\phi: M \rightarrow \mathfrak{t}^*$ is defined to be

$$\langle \phi(x), X \rangle = \alpha_x(X_M(x)), \quad \forall x \in M, \text{ and } \forall X \in \mathfrak{t},$$

where X_M is the vector field on M generated by the X -action. The **moment cone** of ϕ is defined to be

$$C(\phi) = \{t\phi(x) \mid t \geq 0, x \in M\}.$$

It is known ([3, 11] etc.) that, if M is of Reeb type, then the moment map image is an n -dimensional convex polytope lying in a hyperplane away from the origin, and $C(\phi)$ is a cone over the polytope, it is a *strictly convex rational polyhedral cone* of dimension $n + 1$. *Strictly convex* means that $C(\phi)$ contains no linear subspaces of \mathfrak{t}^* of positive dimension, *polyhedral* means that $C(\phi)$ is a cone over a polytope, and *rational* means that the normal vectors of the facets of the cone lie in the integral lattice of \mathfrak{t} .

Let (M, α) be a contact manifold. The contact form α is called **regular** if the Reeb flow of α generates a *free* circle action. The sphere S^{2n+1} with its standard contact form α_0 (the contact form induced from the standard symplectic form on \mathbb{C}^{n+1}) is a contact manifold with a regular contact form (see [3, Example 4.9] for example). The sphere (S^{2n+1}, α_0) is a contact toric manifold of Reeb type, the T^{n+1} action is the one induced from the standard T^{n+1} action on \mathbb{C}^{n+1} . The moment map image of (S^{2n+1}, α_0) is the standard simplex in \mathbb{R}^{n+1} , and the moment cone is the cone over the standard simplex, which is the nonnegative coordinate zone. For a contact toric manifold (M, α) of Reeb type, we can always find a suitable contact form α' , which defines the same contact structure as α , such that its Reeb flow generates a locally free circle action (see [3]).

Key words and phrases. contact toric manifold, Reeb type, K-contact manifold, symplectic toric manifold, cohomology ring, first Chern class.

2010 classification. Primary: 53D10, 53D20; Secondary: 55N10, 57R20.

In this paper, we study compact contact toric manifolds of Reeb type with regular contact forms, we give some sufficient conditions under which the manifolds are homeomorphic to S^{2n+1} .

Theorem. *Let (M, α) be a $2n + 1$ -dimensional connected compact contact toric manifold of Reeb type. Suppose the contact form is regular so that the Reeb flow generates a free S^1 action. If the orbit space $N = M/S^1$ of the Reeb flow satisfies any one of the following two conditions, then M is homeomorphic to S^{2n+1} .*

- (1) $H^*(N; \mathbb{Z}) = \mathbb{Z}[x]/x^{n+1}$, where $x = [\Omega]$, with $\pi: M \rightarrow N$ and $\pi^*\Omega = d\alpha$.
- (2) $c_1(N) = (n + 1)[\Omega]$.

Proof of Theorem. Since the Reeb flow corresponds to a free S^1 -action, $N = M/S^1$ is a smooth manifold; it is a compact symplectic toric manifold of dimension $2n$. Since the fixed points of the T^n -action on N are isolated, we may choose a subcircle action of the T^n -action on N such that its moment map has an isolated point as its minimum, by [13], N is simply connected, i.e.,

$$\pi_1(N) = 1.$$

$\pi: M \rightarrow N$ is a principal S^1 -bundle. By the theorems of Boothby and Wang [2, Theorem 1], and Kobayashi [9], α defines a connection form on M , $\pi^*\Omega$ is the curvature form of this connection, $[\Omega]$ is an *integral* class, and it is the first Chern class of this circle bundle.

First, suppose N satisfies condition (1), i.e., the integral cohomology ring of N is

$$H^*(N; \mathbb{Z}) = \mathbb{Z}[x]/x^{n+1}, \text{ where } x = [\Omega].$$

This first implies that $[\Omega] \in H^2(N; \mathbb{Z})$ is a generator. We look at the homotopy exact sequence of the S^1 -bundle $\pi: M \rightarrow N$

$$\cdots \rightarrow \pi_2(N) \rightarrow \pi_1(S^1) \rightarrow \pi_1(M) \rightarrow \pi_1(N) \rightarrow \cdots.$$

By considering the universal S^1 -bundle $ES^1 \rightarrow BS^1$, a classifying map $N \rightarrow BS^1$, and a diagram chasing, we see that the map

$$\pi_2(N) = H_2(N; \mathbb{Z}) = H^2(N; \mathbb{Z}) = \mathbb{Z} \rightarrow \pi_1(S^1) = H_1(S^1; \mathbb{Z}) = H^1(S^1; \mathbb{Z}) = \mathbb{Z}$$

is multiplication by the first Chern class $[\Omega] \in H^2(N; \mathbb{Z}) = \mathbb{Z}$ of the S^1 -bundle $\pi: M \rightarrow N$ (we need $\pi_1(N) = 1$ for the claim $\pi_2(N) = H_2(N; \mathbb{Z})$). Since $[\Omega]$ is primitive, (up to a sign) the above map is multiplication by 1, so the map

$$\pi_2(N) \rightarrow \pi_1(S^1)$$

is surjective; since we also have $\pi_1(N) = 1$, by the above exact sequence, we get

$$\pi_1(M) = 1.$$

We now look at the Gysin exact sequence for the circle bundle $\pi: M \rightarrow N$,

$$\cdots \rightarrow H^*(M; \mathbb{Z}) \rightarrow H^{*-1}(N; \mathbb{Z}) \rightarrow H^{*+1}(N; \mathbb{Z}) \rightarrow H^{*+1}(M; \mathbb{Z}) \rightarrow \cdots,$$

by the above integral cohomology ring structure of N , we know that the odd cohomology groups of N are 0 and the map $H^{*-1}(N; \mathbb{Z}) \rightarrow H^{*+1}(N; \mathbb{Z})$ is an isomorphism for each $1 \leq * \leq 2n - 1$, so we get

$$H^*(M; \mathbb{Z}) = \mathbb{Z} \text{ when } * = 0, 2n + 1, \text{ and } H^*(M; \mathbb{Z}) = 0 \text{ when } * \neq 0, 2n + 1.$$

We proved that M has the integral cohomology ring of S^{2n+1} .

Since M is simply connected, and has the integral cohomology ring of S^{2n+1} , by the generalized Poincaré conjecture [8, 20], M is homeomorphic to S^{2n+1} . (For dimension 3, by the classification in [11] for compact connected contact toric manifolds of Reeb type, the manifolds are diffeomorphic to the lens spaces. Since M is simply connected, M is diffeomorphic to S^3 . This is an alternative argument for dimension 3.)

Next suppose N satisfies condition (2), i.e., $c_1(N) = (n + 1)[\Omega]$. The manifold (N, Ω) , as a symplectic toric manifold, by Delzant's theorem ([4]), is equivariantly symplectomorphic to the symplectic reduced space of the Kähler manifold \mathbb{C}^m (for some $m > 0$) by a torus action which preserves the Kähler structure of \mathbb{C}^m . By [6, Theorem 3.5], this symplectic reduced space is a Kähler manifold. Hence (N, Ω) possesses a Kähler structure compatible with Ω . Since N is Kähler, and $c_1(N) = (n + 1)[\Omega]$ with $[\Omega]$ being an integral Kähler class, by a theorem of Kobayashi and Ochiai ([10]), N is biholomorphic to $\mathbb{C}\mathbb{P}^n$. The class $[\Omega]$ must be primitive, since if it was not, then $c_1(N) \neq c_1(\mathbb{C}\mathbb{P}^n)$, contradicting to N being biholomorphic to $\mathbb{C}\mathbb{P}^n$. Then N has the integral cohomology ring of $\mathbb{C}\mathbb{P}^n$, with $[\Omega]$ being the degree 2 generator; the rest of the proof then may go as in the last case. Or alternatively, once we know that N is biholomorphic to $\mathbb{C}\mathbb{P}^n$, and the first Chern class of the S^1 -bundle $M \rightarrow N$ is $[\Omega]$, with $[\Omega]$ being primitive integral, then as in the last case, we get that $\pi_1(M) = 1$; by the classification of Kobayashi ([9, Theorem 8]), when the total space is simply connected, an S^1 -bundle over $\mathbb{C}\mathbb{P}^n$ is classified by its first Chern class, hence M is homeomorphic to S^{2n+1} . \square

Corollary. *Let (M, α) be a $2n + 1$ -dimensional connected compact contact toric manifold of Reeb type. Suppose the contact form α is regular so that the Reeb flow generates a free S^1 action. If the orbit space $N = M/S^1$ satisfies any one of the two conditions in the theorem, then the moment cone of M has exactly $n + 1$ number of facets, and the primitive integral normal vectors of the facets span a \mathbb{Z} basis of \mathbb{Z}^{n+1} .*

Proof. When the orbit space N satisfies any one of the two conditions of our theorem, by the proof, N has the same Betti numbers as $\mathbb{C}\mathbb{P}^n$, i.e., $b_{2i}(N) = 1$ and $b_{2i+1}(N) = 0$ for any $0 \leq 2i \leq 2n$.

The symplectic manifold (N, Ω) has an effective $T^n = T^{n+1}/S^1$ action. Let $\bar{\phi}$ be the moment map of the T^n -action on N . The fixed points of the T^n action on N are isolated, and the n -dimensional moment polytope $\bar{\phi}(N)$ is the convex hull of the images of the T^n fixed points ([1, 7]). Since $d\alpha = \pi^*\Omega$, the polytopes $\bar{\phi}(N)$ and $\phi(M)$ differ by a constant in \mathfrak{t}^* .

We choose a generic element $\eta \in \mathfrak{t}' = \text{Lie}(T^n)$. Then the moment map component $\bar{\phi}^\eta$ is a perfect Morse function on N . Together with the fact on the Betti numbers of N , we know that $\bar{\phi}^\eta$ has $n + 1$ number of critical points, which are precisely the fixed points of the one parameter subgroup $\exp(t\eta)$ -action, which are precisely the same as the fixed points of the T^n -action on N . So the moment polytope $\bar{\phi}(N)$ has $n + 1$ number of vertices, and then it has $n + 1$ number of facets. Then $\phi(M)$ has $n + 1$ number of facets, and the moment cone $C(\phi)$ has $n + 1$ number of facets.

By our theorem, the assumptions imply that $\pi_1(M) = 1$. Let v_1, \dots, v_{n+1} be the primitive integral normal vectors of the facets of the moment cone, by [14], up to a sign,

$$\pi_1(M) = \det[v_1, \dots, v_{n+1}].$$

Hence $\{v_1, \dots, v_{n+1}\}$ forms a \mathbb{Z} -basis of \mathbb{Z}^{n+1} . \square

A Riemannian metric g on a contact manifold (M, α) is said to be *adapted* to α if g is preserved by the Reeb flow of α , and if there exists an almost complex structure J on $\ker(\alpha)$ such that g , J and $d\alpha$ are compatible on the symplectic subbundle $\ker(\alpha)$. In such a case, we call (α, g) a K-contact structure on M , or we call (M, α, g) a **K-contact manifold**. A compact K-contact manifold (M, α, g) admits a torus action preserving α , where the torus is the closure of the Reeb flow of α in the isometry group of M (which is compact by [15]). If the compact K-contact manifold (M, α, g) is of dimension $2n + 1$, then the dimension of the torus is bounded from above by $n + 1$ ([16]). If (M, α) is a contact toric manifold of Reeb type, then we can choose an invariant Riemannian metric g adapted to α such that M is given a K-contact structure (α, g) , hence is a K-contact manifold (see for example [12]). So compact contact toric manifolds of Reeb type belong to the category of K-contact manifolds.

The sphere (S^{2n+1}, α_0) with the standard contact form α_0 admits a (standard) K-contact structure. A perturbation of α_0 can give us a different K-contact structure on S^{2n+1} such that the Reeb flow has exactly $n + 1$ number of closed orbits (which are circles) [18], this is the minimal number of closed Reeb orbits that a $2n + 1$ -dimensional compact K-contact manifold can have.

Let (M, α) be a $2n + 1$ -dimensional compact connected contact toric manifold of Reeb type with a regular contact form. Under the condition of our theorem, by the corollary, the moment cone of (M, α) has exactly $n + 1$ number of facets, or the moment polytope $\phi(M)$ has $n + 1$ number of facets. We identify the moment polytope of the $T^n = T^{n+1}/S^1$ -action on $N = M/S^1$ with $\phi(M)$, and identify the Lie algebra of T^n with a subalgebra of \mathfrak{t} . We choose a generic element $\eta \in \mathfrak{t}' = \text{Lie}(T^n)$, then $X = (\eta, \xi)$ is a generic element in \mathfrak{t} , where ξ is the generator of the Lie algebra of the subgroup S^1 . Let X_M be the generating vector field of $X = (\eta, \xi)$ in M . Then the flow of X_M has exactly $n + 1$ number of closed orbits. By the choice of X and by [21, Theorem 7.2], (may also see the proof of [18, Prop. 1]), we can find a contact form α' on M whose Reeb vector field is X_M . Then we have a (different) K-contact structure (α', g') on M with exactly $n + 1$ number of closed Reeb orbits. Hence the manifold we study in our theorem is a compact connected manifold of dimension $2n + 1$ with a K-contact structure which has exactly $n + 1$ number of closed Reeb orbits (minimal number of closed Reeb orbits). Compact connected K-contact manifolds of dimension $2n + 1$ with exactly $n + 1$ number of closed Reeb orbits are studied in [17, 18, 19], where the theorems claim that such manifolds are finitely covered by S^{2n+1} , are homeomorphic to S^{2n+1} when they are simply connected. However, a counter example of this claim is given in [5], the 7-dimensional Stiefel manifold $V_2(\mathbb{R}^5)$, which is simply connected and is not homeomorphic to S^7 .

REFERENCES

- [1] M. Atiyah, *Convexity and commuting Hamiltonians*, Bull. Lond. Math. Soc. **14** (1982), 1-15.
- [2] W. M. Boothby and H. C. Wang, *On contact manifolds*, Annals of Math., **68**, No. 3, 1958, 721-734.
- [3] C. P. Boyer and K. Galicki, *A note on toric contact geometry*, J. Geom. Phys. **35** (2000), 288-298.
- [4] T. Delzant, *Hamiltoniens périodiques et images convexes de l'application moment*, Bull. Soc. Math. France **116** (1988), no. 3, 315-339.

- [5] O. Goertsches, H. Nozawa, and D. Töben, *Equivariant cohomology of K-contact manifolds*, Math. Ann. (2012) **354**, 1555-1582.
- [6] V. Guillemin and S. Sternberg, *Geometric quantization and multiplicities of group representations*, Invent. Math. **67** (1982), 515-538.
- [7] V. Guillemin and S. Sternberg, *Convexity properties of the moment mapping*, Invent. Math. **67** (1982), no. 3, 491-513.
- [8] B. Kleiner and J. Lott, *Notes on Perelman's papers*, Geometry and Topology **12** (2008), 2587-2858.
- [9] S. Kobayashi, *Principal fibre bundles with the 1-dimensional toroidal group*, Tôhoku Math. J. **8** (1956), 29-45.
- [10] S. Kobayashi and T. Ochiai, *Charaterizations of complex projective spaces and hyperquadrics*, J. Math. Kyoto Univ. **13-1** (1973), 31-47.
- [11] E. Lerman, *Contact toric manifolds*, Journal of Symp. Geom., **1**, no. 4, 785-828, 2002.
- [12] E. Lerman, *Homotopy groups of K-contact toric manifolds*, Trans. Ameri. Math. Soc., **356**, no. 10, 4075-4083, 2004.
- [13] H. Li, π_1 of Hamiltonian S^1 -manifolds, Proceedings of Ameri. Math. Soc., vol. **131**, no. 11, 3579-3582, 2003.
- [14] H. Li, *The fundamental groups of contact toric manifolds*, J. of symp. geom., Vol **18**, No. 3, 815-818, 2020.
- [15] S. B. Myers and N. E. Steenrod, *The group of isometries of a Riemannian manifold*, Ann. Math. **40**(2), 400-416 (1939).
- [16] P. Rukimbira, *The dimension of leaf closures of K-contact flows*, Ann. Global Anal. Geom. **12**, 103-108 (1994).
- [17] P. Rukimbira, *On K-contact manifolds with minimal number of closed characteristics*, Proceedings of the Ameri. Math. Soc., vol. **127**, no. 11, 3345-3351, 1999.
- [18] P. Rukimbira, *Spherical rigidity via contact dynamics*, Bull. Belg. Math. Soc. **7** (2000), 563-569.
- [19] P. Rukimbira, *Correction to Spherical rigidity via contact dynamics*, Bull. Belg. Math. Soc. **8** (2001), 147-153.
- [20] S. Smale, *Generalized Poincaré's conjecture in dimensions greater than four*, Ann. of Math. (2) **74** (1961), 391-406.
- [21] K. Yano and M. Kon, *Structures on Manifolds*, Series in Pure Mathematics, World Scientific **3** (1984).

SCHOOL OF MATHEMATICAL SCIENCES, SOOCHOW UNIVERSITY, SUZHOU, 215006, CHINA.

Email address: hui.li@suda.edu.cn