# Thermodynamics of hairy accelerating black holes in gauged supergravity and beyond

Yi Wang and Jie Ren

School of Physics, Sun Yat-sen University, Guangzhou, 510275, China E-mail: wangy973@mail2.sysu.edu.cn, renjie7@mail.sysu.edu.cn

ABSTRACT: We study the thermodynamics of accelerating asymptotically AdS black holes with scalar hair in four-dimensional Einstein-Maxwell-dilaton theories (U(1)-charged), which can be embedded in gauged supergravities for specific dilaton coupling constants. These accelerating black holes are described by charged dilaton C-metrics, which have unique features but are almost unexplored before. Thermodynamics of slowly accelerating black holes is generalized to include scalar hair. We find that properly accounting for mixed boundary conditions for the scalar field leads to standard consistent first law and other thermodynamic relations. We compute the dual stress-energy tensor and the mass through holographic renormalization, which is associated with the boundary conditions of the scalar field. We also find that the AMD mass is the same as the holographic mass. Along the same line, we also obtain the thermodynamics of  $U(1)^2$ -charged accelerating black holes.

# Contents

1	Introduction	1
<b>2</b>	Accelerating black holes in Einstein-Maxwell-dilaton systems	3
3	Holographic Renormalization	5
	3.1 Holographic dictionary for mixed boundary conditions	5
	3.2 Holographic mass of accelerating charged dilaton black holes	7
4	Thermodynamics of slowly accelerating charged dilaton black holes	9
5	Thermodynamics of $U(1)^2$ -charged accelerating black holes	14
6	Summary and Discussion	17

#### 1 Introduction

The C-metric is a class of exact solutions of Einstein field equations, and the original one was found by Hermann Weyl in 1917 [1]. This name came from the C slot of the classification by Ehlers and Kundt in 1963 [2]. A breakthrough work by Kinnersley and Walker in 1970 [3] reveals that this solution describes a pair of accelerating black holes apart from each other. The generalization of this solution was done by Plebański-Demiański in 1976 [4], which includes rotation and cosmological constant. For more interpretations of this solution see [5, 6] for references. Based on the AdS/CFT correspondence, this solution also has fruitful applications. For instance, the construction of exact black holes on the branes [7, 8], the plasma ball [9], the black funnels and droplets [10, 11], and the quantum BTZ black hole [12].

Black holes are perhaps the most intriguing objects in our universe, connecting the bridge between classical and quantum gravity. Especially this link is highlighted by the early discoveries that black hole as a thermal object obeys the laws of thermodynamics [13–15], and the black hole chemistry [16] gives more diversity to these thermal objects. Although the study of the C-metric together with the black hole thermodynamics has a long history, the thermodynamics of the accelerating black holes has been formulated for small acceleration in AdS spacetime in recent years [17–20]. Later, the authors of [21, 22] formulated extended thermodynamics for slowly accelerating black holes in AdS spacetime, in which it is even allowed to vary the tension along each axis independently. A key ingredient of their works is a rescaling parameter  $\alpha$  for the time coordinate as pointed out before in [23] for Kerr AdS black holes. Last year, the authors of [24] generalized previous results to include magnetic charge and allow the variation of Newton's constant in the

first law of thermodynamics. From the holographic perspective, varying Newton's constant together with the AdS radius can keep the corresponding CFT unchanging [25, 26].

More generally, asymptotically anti-de Sitter (AdS) black holes with scalar hair in gauged supergravity are valuable in the AdS/CFT correspondence. The charged dilaton C-metrics found in [27, 28] are significant generalizations of the original C-metric. They describe asymptotically AdS accelerating black holes with scalar hair in Einstein-Maxwelldilaton systems. The scalar potential of these systems can be expressed by a superpotential, and these theories can be embedded in gauged supergravities for specific dilaton coupling constants. Even without acceleration, the planar black hole solutions of these systems have rich properties of distinctive IR geometries [28, 29]. Thus, generalizing these hairy black holes with an acceleration is expected to have more fruitful applications. Like the original C-metric, the charged dilaton C-metrics also have a conical singularity along one or both polar axes. This conical singularity can be replaced by a magnetic flux tube [30] or a cosmic string (topological defect) [31], giving a precise explanation of the mechanism of the acceleration. Furthermore, these solutions can also be generalized to multiply-charged accelerating black holes [32].

Here we consider the thermodynamics of slowly accelerating black holes in asymptotically AdS spacetime [33] with scalar hair and electric charge. Since there are no accelerating horizons, the thermodynamics of these systems are clear. If a scalar field exists, however, the thermodynamics will become more fascinating. On the one hand, it was proposed that the first law of hairy black hole thermodynamics will be modified with a "scalar charge" [34–36]. On the other hand, these papers [37, 38] show that properly accounting for a new contribution from the asymptotic value of the scalar field for total energy (more general boundary conditions for scalar field) leads to unmodified thermodynamics. It will be very interesting to analyze our solutions along this line. See section 3 for details. Without introducing a scalar charge, we drive the first law of thermodynamics [20, 24] and the Smarr relation [39, 40] for hairy slowly accelerating black holes as follows:

$$\delta M = T\delta S + V\delta P + \sum_{I=1}^{2} \Phi_{I}\delta Q_{I} - \lambda_{+}\delta\mu_{+} - \lambda_{-}\delta\mu_{-} - \eta \frac{\delta G_{4}}{G_{4}}, \qquad (1.1)$$

$$M = 2(TS - PV) + \sum_{I=1}^{2} \Phi_{I}Q_{I}, \qquad (1.2)$$

where

$$\eta = M - TS - \sum_{I=1}^{2} \Phi_{I} Q_{I} - PV + \lambda_{+} \mu_{+} + \lambda_{-} \mu_{-}.$$
(1.3)

If we employ the duality relation as in [26]:

$$C = k \frac{\ell^2}{16\pi G_4},$$
 (1.4)

after some simple algebra, we can rewrite the first law (1.1) as follows

$$\delta M = T\delta S + \sum_{I=1}^{2} \Phi_I \delta Q_I + V_c \delta P + \mu_c \delta C - \lambda_+ \delta \mu_+ - \lambda_- \delta \mu_-, \qquad (1.5)$$

where

$$V_c = V + \frac{\eta}{2P}, \qquad \mu_c = \frac{\eta}{2C}.$$
(1.6)

This generalizes the result of [26] with conical defects in the bulk spacetime.

This paper is organized as follows. In section 2, we introduce the solutions of accelerating charged dilaton black holes. In section 3, we take a brief review on boundary counterterms, boundary conditions for the scalar field, and the holographic dictionary, then obtain the holographic stress-energy tensor of these solutions to compute the holographic mass. In section 4, we derive the standard thermodynamic laws for slowly accelerating electrically charged dilaton black holes. In section 5, we give the thermodynamics of  $U(1)^2$ -charged accelerating black holes. In section 6, we summarize our results and discuss generalizations to black funnels and droplets [10, 11].

#### 2 Accelerating black holes in Einstein-Maxwell-dilaton systems

Before discussing the black hole thermodynamics, we introduce the solutions of accelerating charged dilaton black holes, which can be found in [27, 28]. The bulk action is

$$S_{\text{bulk}} = \frac{1}{16\pi G_4} \int_{\mathcal{M}} d^{3+1}x \sqrt{-G} \left( R - \frac{1}{2} (\partial \phi)^2 - V(\phi) - \frac{1}{4} Z(\phi) \mathcal{F}^2 \right), \quad (2.1)$$

where  $\mathcal{F} = dA$ ,  $Z(\phi) = e^{-\zeta \phi}$ , and the scalar potential is

$$V(\phi) = -\frac{2e^{-\frac{\phi}{\zeta}} \left(\zeta^2 (3\zeta^2 - 1) + 8\zeta^2 e^{\frac{1+\zeta^2}{2\zeta}\phi} + (3-\zeta^2) e^{\zeta\phi + \frac{\phi}{\zeta}}\right)}{(1+\zeta^2)^2 \ell^2},$$
(2.2)

which can also be expressed with a superpotential

$$V = \left(\frac{dW}{d\phi}\right)^2 - \frac{3}{4}W^2, \qquad W = \frac{\sqrt{8}}{(1+\zeta^2)\ell} \left(e^{\frac{1}{2}\zeta\phi} + \zeta^2 e^{-\frac{1}{2\zeta}\phi}\right).$$
(2.3)

Here  $\zeta$  is the dilaton coupling constant, and  $\ell$  is the AdS radius. When  $\zeta = \sqrt{3}$ , 1,  $1/\sqrt{3}$  and 0, these solutions can be embedded in gauged supergravities. In particular, the  $\zeta = 0$  case corresponds to RN-AdS accelerating black hole, which has been explored before, and the goal of this paper is to analyze the other solutions. The equations of motion are invariant under an electric-magnetic duality transformation and  $\phi \to -\phi$  [30], and the discussion of the magnetic-charged dilation black holes is similar to electric-charged ones. Therefore, we will only focus on electrically charged solutions.

The line elements describing accelerating black holes with scalar hair and electric charge can be written as follows:

$$ds^{2} = \frac{1}{a^{2}(x-y)^{2}} \left[ f_{x}^{\frac{2\zeta^{2}}{1+\zeta^{2}}} \left( -\frac{F_{y}}{\alpha^{2}} dt'^{2} + \frac{dy^{2}}{F_{y}} \right) + h_{y}^{\frac{2\zeta^{2}}{1+\zeta^{2}}} \left( \frac{dx^{2}}{G_{x}} + \frac{G_{x}}{K^{2}} d\varphi^{2} \right) \right], \quad (2.4)$$

with

$$F_{y} = -(1 - y^{2})(1 + 2acy)h_{y}^{\frac{1-\zeta^{2}}{1+\zeta^{2}}} + \frac{1}{a^{2}\ell^{2}}h_{y}^{\frac{2\zeta^{2}}{1+\zeta^{2}}},$$

$$G_{x} = (1 - x^{2})(1 + 2acx)f_{x}^{\frac{1-\zeta^{2}}{1+\zeta^{2}}},$$

$$h_{y} = 1 + aby, \qquad f_{x} = 1 + abx,$$

$$A = \frac{2}{\alpha}\sqrt{\frac{2bc}{1+\zeta^{2}}}(y - y_{h})dt, \qquad e^{\zeta\phi} = \left(\frac{h_{y}}{f_{x}}\right)^{\frac{2\zeta^{2}}{1+\zeta^{2}}}.$$
(2.5)

The AdS boundary of these solutions are at x = y, and the boundary metrics are

$$ds_{\rm bdy}^2 = f_x^{\frac{2\zeta^2}{1+\zeta^2}} \left( -\frac{F_x}{\alpha^2} dt'^2 + \frac{f_x^{\frac{2\zeta^2}{1+\zeta^2}}}{a^2\ell^2 F_x G_x} dx^2 + \frac{G_x}{K^2} d\varphi^2 \right).$$
(2.6)

2

The coordinate transformation  $y = -\frac{1}{ar}$ ,  $x = \cos(\theta)$ , t' = at takes the metrics to another form

$$ds^{2} = \frac{1}{H(r,\theta)^{2}} \left[ f(\theta)^{\frac{2\zeta^{2}}{1+\zeta^{2}}} \left( -\frac{F(r)}{\alpha^{2}} dt^{2} + \frac{dr^{2}}{F(r)} \right) + r^{2}h(r)^{\frac{2\zeta^{2}}{1+\zeta^{2}}} \left( \frac{d\theta^{2}}{G(\theta)} + \frac{\sin^{2}\theta G(\theta)d\varphi^{2}}{K^{2}} \right) \right],$$
(2.7)

with

$$H = 1 + ar \cos \theta,$$

$$F = (1 - a^{2}r^{2})(1 - \frac{2c}{r})h^{\frac{1-\zeta^{2}}{1+\zeta^{2}}} + \frac{r^{2}}{\ell^{2}}h^{\frac{2\zeta^{2}}{1+\zeta^{2}}},$$

$$G = (1 + 2ac \cos \theta)f^{\frac{1-\zeta^{2}}{1+\zeta^{2}}},$$

$$h = 1 - \frac{b}{r}, \qquad f = 1 + ab \cos \theta,$$

$$A = \frac{2}{\alpha}\sqrt{\frac{2bc}{1+\zeta^{2}}} \left(\frac{1}{r_{h}} - \frac{1}{r}\right)dt, \qquad e^{\zeta\phi} = \left(\frac{h}{f}\right)^{\frac{2\zeta^{2}}{1+\zeta^{2}}}.$$
(2.8)

The parameters appearing in the metrics c and a are usual parameters related to the mass and acceleration of the black holes. While the scalar field is associated with the parameter b, the parameters b and c determine the gauge potential together. When b vanishes, the metrics reduce to usual accelerating black hole in AdS [4, 5] without any charges. In the limit  $\ell \to \infty$ , up to a scalar field rescaling, these solutions recover accelerating dilaton black holes in string theory. In particular, when  $\zeta < 1$  these solutions correspond to pair

creation of dilaton black holes [30]. And another limit  $a \to 0$  recovers usual dilaton black holes in asymptotically AdS spacetime [41].

Analogous to previous works [19–22, 24], to obtain correct thermodynamic quantities for slowly accelerating charged dilaton black holes in asymptotically AdS spacetime, it is vital to rescale the time coordinate with  $\alpha$ . The parameter K carries the information of conical deficit of these spacetime at two axes, and then it gives the tension defined by [20, 24] (where  $\theta_{+} = 0, \theta_{-} = \pi$ )

$$\mu_{\pm} = \frac{\delta_{\pm}}{8\pi G_4} = \frac{1}{4G_4} \left[ 1 - \frac{G(\theta_{\pm})}{K} \right] = \frac{1}{4G_4} \left[ 1 - \frac{(1\pm 2ac)(1\pm ab)^{\frac{1-\zeta^2}{1+\zeta^2}}}{K} \right].$$
(2.9)

Finally, to maintain the Lorentz signature for the metrics describing slowly accelerating black holes with scalar hair and electric charge, we need to constrain the range of parameters as follows:

- It is obvious that the conformal boundary of these spacetime are located at  $r_{bdy} = -1/a \cos \theta$ . To obtain slowly accelerating solutions (no acceleration horizon), we must have  $a \cos \theta \le 1/r \le 1/r_h$  and  $F(r_{bdy}) > 0$  [33].
- For the  $\theta$ -coordinate, we require  $G(\theta) > 0$  and  $f(\theta) > 0$  on  $\theta \in [0, \pi]$ , which lead to  $ac < \frac{1}{2}$  and ab < 1.
- The spacetime is singular when  $r \to 0$  or  $r \to b$ , and the later requires  $r_h > b$  coincidence with the restriction on h(r) > 0. To have a black hole without naked singularity, we require F(r) have at least one root in the range  $r \in (b, 1/a)$ .

For these ranges of the parameters, these metrics correspond to slowly accelerating black holes with electric charge and scalar hair in asymptotically AdS spacetime. Consequently, we can discuss the thermodynamics of these black holes in equilibrium without ambiguity.

#### 3 Holographic Renormalization

As our systems include a scalar field, there are more subtleties. In particular, properly renormalizing the stress-energy tensor with mixed boundary conditions for the scalar field is important for correctly giving the thermodynamic relations of these solutions. In this section, we will take a brief review of boundary counterterms, boundary conditions for the scalar field, and the holographic dictionary, then calculate the holographic stress-energy tensor and mass for these solutions. For more details, see [37].

#### 3.1 Holographic dictionary for mixed boundary conditions

To obtain the counterterms, the Fefferman-Graham (FG) [42] expansion is needed near the AdS boundary. In the FG gauge, the metric takes the following form:

$$ds^{2} = \frac{\ell^{2}}{z^{2}} \left( dz^{2} + g_{ij}(x, z) dx^{i} dx^{j} \right), \qquad (3.1)$$

with

$$g_{ij}(x,z) = g(x)_{(0)ij} + zg(x)_{(1)ij} + z^2 g(x)_{(2)ij} + \cdots,$$
  

$$A_i(x,z) = A_i(x)_{(0)} + zA_i(x)_{(1)} + z^2 A_i(x)_{(2)} + \cdots,$$
  

$$\phi(x,z) = z^{\Delta_-} \varphi(x,z) = z^{\Delta_-} \left(\varphi(x)_{(0)} + z\varphi(x)_{(1)} + z^2 \varphi(x)_{(2)} + \cdots\right).$$
(3.2)

Defining the following constants [37]

$$V_k \equiv \lim_{r \to r_{\rm bdy}} V^{(k)}(\phi), \qquad Z_k \equiv \lim_{r \to r_{\rm bdy}} Z^{(k)}(\phi), \tag{3.3}$$

where  $V^{(k)}(\phi)$  and  $Z^{(k)}(\phi)$  are the k-th derivative of  $V(\phi)$  and  $Z(\phi)$  respectively. In our systems, we have  $\Delta_{-} = 1$  and the scalar field vanishes at the boundary, so the scalar potential has the asymptotic expansion near the boundary

$$V(\phi) = -\frac{6}{\ell^2} - \frac{\phi^2}{\ell^2} + \frac{1 - 4\zeta^2 + \zeta^4}{24\zeta^2\ell^2}\phi^4 + \cdots$$
 (3.4)

Then the AdS mass of the scalar field is

$$m_{\phi}^2 = V_2 = -\frac{2}{\ell^2},\tag{3.5}$$

which lies in the range

$$-\frac{3^2}{4} + 1 \ge \ell^2 m_\phi^2 \ge -\frac{3^2}{4}.$$
(3.6)

Therefore, we can consider more general (Neumann or mixed) boundary conditions for the scalar field, which can lead to the first law of thermodynamics without any scalar charge [37].

In the FG expansion, the equations of motion determine the relation of the coefficients.<sup>1</sup> As a consequence, these relations lead to covariant boundary counterterms at the cutoff  $z = \epsilon$  as follows:

$$S_{\rm ct} = -\frac{1}{16\pi G_4} \int_{z=\epsilon} d^3x \sqrt{-\gamma} \left(\frac{4}{\ell} + \ell R[\gamma] + \frac{1}{2\ell}\phi^2\right),\tag{3.8}$$

where  $\gamma_{ij}$  is the induced metric on the boundary. With this counterterm, we can holographically renormalize the action for an asymptotically AdS spacetime with the Gibbons-Hawking term

$$S_{\rm GH} = \frac{1}{16\pi G_4} \int_{z=\epsilon} d^3 x \sqrt{-\gamma} \ 2\mathcal{K},\tag{3.9}$$

<sup>1</sup>The relation of the coefficients of the FG expansion is

$$g(x)_{(1)ij} = 0,$$

$$g(x)_{(2)ij} + R_{(0)ij} - \frac{1}{4}R_{(0)}g(x)_{(0)ij} + \frac{1}{8}\varphi(x)^{2}_{(0)}g(x)_{(0)ij} = 0,$$

$$g(x)^{jk}_{(0)} \left(D_{(0)j}g(x)_{(2)ki} - D_{(0)i}g(x)_{(2)jk}\right) - \frac{1}{2}\varphi(x)_{(0)}\partial_{i}\varphi(x)_{(0)} = 0,$$

$$\varphi(x)_{(2)} + \frac{1}{2}\Box_{(0)}\varphi(x)_{(0)} + \frac{1}{2}\mathrm{Tr}\left(g^{-1}_{(0)}g_{(2)}\right)\varphi(x)_{(0)} - \frac{1}{12}\ell^{2}V_{4}\varphi(x)^{3}_{(0)} = 0,$$
(3.7)

where  $D_{(0)i}$ ,  $\Box_{(0)}$  and  $R^{(0)}$  are covariant derivative, Laplacian, and Ricci scalar with respect to  $g(x)_{(0)ij}$ .

where  $\mathcal{K}$  is the trace of extrinsic curvature with respect to the induced metric. Then the renormalized action independent of the cutoff  $\epsilon$  is

$$S_{\rm ren} = \lim_{\epsilon \to 0} \left( S_{\rm bulk} + S_{\rm GH} + S_{\rm ct} \right), \tag{3.10}$$

where  $S_{\text{bulk}}$  is the action (2.1) with r being integrated to  $z = \epsilon$ .

Since the scalar field admits more general boundary conditions, the choice [37]

$$J_{\mathcal{F}_{\varphi}} = -\ell^2 \varphi_{(1)} - \mathcal{F}_{\varphi}'(\varphi_{(0)}), \qquad (3.11)$$

where  $\mathcal{F}_{\varphi}(\varphi_{(0)}) = \frac{1}{3}x_1\varphi_{(0)}^3$ , corresponds to a triple-trace deformation in the dual CFT [43–45]. Specifying the value of  $x_1$  is important for correctly giving the laws of thermodynamics for these solutions, and this choice also requires an additional term for the on-shell action as follows:

$$S_{\mathcal{F}_{\varphi}} = \frac{1}{16\pi G_4} \int_{z=\epsilon} d^3x \sqrt{-g_{(0)}} \Big( J_{\mathcal{F}_{\varphi}}\varphi_{(0)} + \mathcal{F}_{\varphi}(\varphi_{(0)}) \Big).$$
(3.12)

Then the modified one-point functions are

$$\langle \mathcal{J}_{i} \rangle_{\mathrm{M}} = \frac{1}{16\pi G_{4}} Z_{0} A_{i(1)},$$

$$\langle \mathcal{O}_{\Delta_{-}} \rangle_{\mathrm{M}} = \frac{1}{16\pi G_{4}} \varphi_{(0)},$$

$$\langle \mathcal{T}_{ij} \rangle_{\mathrm{M}} = \frac{1}{16\pi G_{4}} \Big[ 3\ell^{2} g_{(3)ij} + \Big( \mathcal{F}_{\varphi}(\varphi_{(0)}) - \varphi_{(0)} \mathcal{F}_{\varphi}'(\varphi_{(0)}) \Big) g_{(0)ij} \Big],$$

$$(3.13)$$

and the local Ward identities are

$$D_{(0)i} \langle \mathcal{J}^{i} \rangle_{\mathrm{M}} = 0,$$

$$D_{(0)}^{j} \langle \mathcal{T}_{ij} \rangle_{\mathrm{M}} = \langle \mathcal{O}_{\Delta_{-}} \rangle_{\mathrm{M}} \partial_{i} J_{\mathcal{F}_{\varphi}} + \langle \mathcal{J}^{j} \rangle_{\mathrm{M}} \mathcal{F}_{ij}^{(0)},$$

$$\langle \mathcal{T}^{i}_{i} \rangle_{\mathrm{M}} = \frac{1}{16\pi G_{4}} \Big[ 2\varphi_{(0)} J_{\mathcal{F}_{\varphi}} + 3\mathcal{F}_{\varphi}(\varphi_{(0)}) - \varphi_{(0)} \mathcal{F}_{\varphi}'(\varphi_{(0)}) \Big].$$
(3.14)

We are going to see that our solutions fit perfectly in this framework in the next subsection.

# 3.2 Holographic mass of accelerating charged dilaton black holes

With these techniques, we calculate the holographic mass from the boundary stress-energy tensor. First, it is necessary to use the FG coordinates, and we take the same coordinate transformation as in [22]:

$$\frac{1}{r} = -a\rho - \sum_{n=1}^{4} F_n(\rho) z^n, \quad \cos \theta = \rho + \sum_{n=1}^{4} G_n(\rho) z^n.$$
(3.15)

The functions  $F_n(\rho)$  and  $G_n(\rho)$  can be solved by requiring  $g_{zz} \equiv \ell^2/z^2$  and  $g_{z\rho} \equiv 0$  at each order of z in the FG gauge (3.1), and the leading order gives

$$G_1(\rho) = -\frac{a\ell^2 F_1(\rho) X(\rho)}{\gamma_b(\rho)^2},$$
(3.16)

where

$$X(\rho) = (1 - \rho^2)(1 + 2ac\rho),$$
  

$$\gamma_b(\rho) = \sqrt{(1 + ab\rho)^{\frac{3\zeta^2 - 1}{1 + \zeta^2}} - a^2\ell^2 X(\rho)}.$$
(3.17)

Note that the square root requires  $\gamma_b^2(\rho) > 0$ , which is the same as  $F(r_{\text{bdy}}) > 0$ . To simplify calculations, we choose  $F_1(\rho) = -\frac{\gamma_b^2(\rho)}{\alpha}$ , and then the boundary metric can be written as

$$ds_{(0)}^{2} = -\frac{(1+ab\rho)^{\frac{3-5\zeta^{2}}{1+\zeta^{2}}}\gamma_{b}^{2}(\rho)}{\ell^{4}}dt^{2} + \frac{\alpha^{2}}{\ell^{2}X(\rho)\gamma_{b}(\rho)^{2}}d\rho^{2} + \frac{(1+ab\rho)^{\frac{3-5\zeta^{2}}{1+\zeta^{2}}}\alpha^{2}X(\rho)}{K^{2}\ell^{2}}d\varphi^{2}.$$
 (3.18)

Since the scalar field admits more general boundary conditions in our systems, we take the mixed boundary conditions for these solutions. The value of  $x_1$  is

$$x_1 = \frac{1 - \zeta^2}{4\zeta} \ell^2, \tag{3.19}$$

which is specified by requiring that the holographic mass is the same as the AMD mass (4.19) below, and this result can also lead to consistent thermodynamic laws. In particular, this condition is also the same as non-accelerating black holes of these system [28]. As a result, the expectation value of the stress-energy tensor (3.13) is

$$\langle \mathcal{T}_{i}^{j} \rangle_{\mathrm{M}} = \frac{\ell^{4} (1+ab\rho)^{\frac{3\zeta^{2}-5}{1+\zeta^{2}}}}{16\pi G_{4}\alpha^{3}(1+\zeta^{2})^{2}} \mathrm{diag}\left[-\omega_{E}, (1+\zeta^{2})\omega_{\rho}, \omega_{\varphi}\right], \qquad (3.20)$$

with

$$\begin{split} \omega_{1} &= 1 - \zeta^{2} + 12ac\rho, \qquad X_{b}(\rho) = X(\rho)(1 + ab\rho)^{\frac{1-3\zeta^{2}}{1+\zeta^{2}}}, \\ \omega_{2} &= (3 - \zeta^{2})\rho + ac\left(-3\zeta^{4} + \zeta^{2}(1 + 15\rho^{2}) + 27\rho^{2}\right), \\ \omega_{3} &= 3\zeta^{4} - \zeta^{2}\left(1 + \rho^{2}(1 - 8ac\rho)\right) + 3\rho^{2}(1 + 8ac\rho). \\ \omega_{E} &= (1 + \zeta^{2})\left(2c(1 + \zeta^{2}) + b\omega_{1}\right)\left(2 - 3a^{2}\ell^{2}X_{b}(\rho)\right) \\ &+ 2ab^{2}\left[2ac\zeta^{2}(1 - \zeta^{2}) + 2(1 - \zeta^{2})\rho + 6ac(3 + \zeta^{2})\rho^{2} - a^{2}\ell^{2}X_{b}(\rho)\omega_{2}\right] \\ &+ a^{2}b^{3}\left[2\left(\zeta^{4} - \zeta^{2}(1 + \rho^{2}) + \rho^{2} + 8ac\rho^{3}\right) - a^{2}\ell^{2}X_{b}(\rho)\omega_{3}\right], \qquad (3.21) \\ \omega_{\rho} &= 2c(1 + \zeta^{2}) + b\omega_{1} + 2ab^{2}\left(\rho - ac(\zeta^{2} - 9\rho^{2})\right) + a^{2}b^{3}(\zeta^{2} + \rho^{2} + 8ac\rho^{3}), \\ \omega_{\varphi} &= (1 + \zeta^{2})\left(2c(1 + \zeta^{2}) + b\omega_{1}\right)\left(1 - 3a^{2}\ell^{2}X_{b}(\rho)\omega_{2}\right] \\ &+ a^{2}b^{3}\left[(1 + \zeta^{2})\left(\rho - ac(\zeta^{2} - 9\rho^{2})\right) - a^{2}\ell^{2}X_{b}(\rho)\omega_{3}\right]. \end{split}$$

Then the following integration gives the holographic mass

$$M = \int -\langle \mathcal{T}_t^t \rangle_{\mathrm{M}} \sqrt{-g_{(0)}} \, d\rho d\varphi$$
$$= \frac{c + \frac{1-\zeta^2}{1+\zeta^2} \frac{b}{2} + \frac{1}{8} a \ell^2 \left( (1-2ac)^2 (1-ab)^{2\frac{1-\zeta^2}{1+\zeta^2}} - (1+2ac)^2 (1+ab)^{2\frac{1-\zeta^2}{1+\zeta^2}} \right)}{\alpha K G_4}, \qquad (3.22)$$

this integration requires ab < 1, which is satisfied with our parameter constraint. When b equals zero, we can find the boundary metric (3.18) and the stress-energy tensor (3.20) are all same as in [22] respectively.

The asymptotic expansions of the gauge potential and the scalar field are

$$A_t(\rho, z) = A_t(\rho)_{(0)} + zA_t(\rho)_{(1)} + z^2 A_t(\rho)_{(2)} + \cdots,$$
  

$$\phi(\rho, z) = z \left(\varphi(\rho)_{(0)} + z\varphi(\rho)_{(1)} + z^2\varphi(\rho)_{(2)} + \cdots\right),$$
(3.23)

with

$$\begin{aligned} A_{t}(\rho)_{(0)} &= \frac{2}{\alpha r_{h}} \sqrt{\frac{2bc}{1+\zeta^{2}}} (1+ar_{h}\rho), \\ A_{t}(\rho)_{(1)} &= -\frac{2}{\alpha^{2}} \sqrt{\frac{2bc}{1+\zeta^{2}}} \gamma_{b}(\rho)^{2}, \end{aligned}$$
(3.24)  
$$\varphi(\rho)_{(0)} &= -\frac{2b\zeta(1+ab\rho)^{2\frac{\zeta^{2}-1}{1+\zeta^{2}}}}{\alpha(1+\zeta^{2})}, \\ \varphi(\rho)_{(1)} &= \frac{b\zeta(1+ab\rho)^{\frac{\zeta^{2}-3}{1+\zeta^{2}}}}{\alpha^{2}(1+\zeta^{2})} \Big[ a\ell^{2} \left(2\rho + 4a^{2}bc\rho^{3} + a(b-2c+b\rho^{2}+6c\rho^{2})\right) \\ &+ b\frac{\zeta^{2}-1}{1+\zeta^{2}} (1+ab\rho)^{\frac{3\zeta^{2}-1}{1+\zeta^{2}}} \Big]. \end{aligned}$$

Then we can easily get the dual theory one-point functions (3.13) and check the local Ward identities (3.14) with the stress-energy tensor (3.20). With  $\varphi(\rho)_{(2)}$ ,  $g(\rho)_{(1)ij}$  and  $g(\rho)_{(2)ij}$ , we can also check (3.7), but it is not illuminating to write everything down. We leave the discussion of these holographic results in the last section.

#### 4 Thermodynamics of slowly accelerating charged dilaton black holes

We begin this section by writing down thermodynamic quantities first:

$$M = \frac{c + a_1 \frac{b}{2} + \frac{1}{8} a \ell^2 \left( (1 - 2ac)^2 (1 - ab)^{2a_1} - (1 + 2ac)^2 (1 + ab)^{2a_1} \right)}{\alpha K G_4},$$

$$\alpha = \sqrt{a \alpha K G_4 M}, \qquad S = \frac{\pi r_h^2 h(r_h)^{1-a_1}}{K G_4 \left( 1 - a^2 r_h^2 \right)},$$

$$T = \frac{h(r_h)^{a_1}}{4\pi \alpha r_h^2} \left[ 2c(1 - a^2 r_h^2) + \frac{r_h^2 \left( 2r_h - b \left( 1 + 2a_1 + (1 - 2a_1)a^2 r_h^2 \right) \right)}{\ell^2 h(r_h)^{2a_1} (1 - a^2 r_h^2)} \right],$$

$$Q = \frac{\sqrt{(1 + a_1)bc}}{2K G_4}, \qquad \Phi = \frac{2\sqrt{(1 + a_1)bc}}{\alpha r_h}, \qquad P = \frac{3}{8\pi \ell^2 G_4}, \qquad (4.1)$$

$$V = \frac{4\pi \ell^2}{3\alpha K} \left[ c(1 + \frac{a_1b}{r_h}) - \alpha K M G_4 + \frac{r_h^2 \left( 2r_h - b \left( 1 + 2a_1 + (1 - 2a_1)a^2 r_h^2 \right) \right)}{2\ell^2 h(r_h)^{2a_1 - 1} (1 - a^2 r_h^2)^2} \right],$$

$$\lambda_{\pm} = \frac{1}{\alpha L_1} \left[ L_2 \left( L_3 L_{3\pm} - L_4 L_{4\pm} + L_5 L_{5\pm} \right) - \left( L_6 L_{3\pm} + L_7 L_{5\pm} + L_8 L_{8\pm} \right) L_9 \right],$$

with

$$\begin{split} a_{1} &= \frac{1-\zeta^{2}}{1+\zeta^{2}}, \quad h(r_{h}) = 1 - \frac{b}{r_{h}}, \quad L_{1\pm} = a_{1}^{2}b^{2}(1\mp 2ac), \quad L_{2\pm} = a_{1}b(1\mp ab)(2ac\mp 1), \\ L_{1} &= 4a\ell^{2}r_{h}(1-a^{2}r_{h}^{2})^{3}(1-a^{2}b^{2})^{a_{1}} \left[b^{2}\left(a_{1}^{2}-4a^{2}(a_{1}^{2}-1)c^{2}\right)-4c^{2}\right], \\ L_{2} &= (1-a^{2}r_{h}^{2})h(r_{h})^{-2a_{1}}\left(a^{2}br_{h}^{2}(a_{1}-1)-(a_{1}+1)b+2r_{h}\right), \\ L_{3} &= 2a^{2}\ell^{2}r_{h}^{2}(2c-r_{h})h(r_{h})^{2a_{1}}, \quad L_{4} = 2\ell^{2}c(1-a^{2}r_{h}^{2})h(r_{h})^{2a_{1}}, \\ L_{5} &= (a_{1}-1)br_{h}^{2}-a_{1}b\ell^{2}(1-a^{2}r_{h}^{2})(1-\frac{2c}{r_{h}})h(r_{h})^{2a_{1}-1}, \quad L_{6} = 2a^{2}r_{h}^{3}h(r_{h}), \\ L_{7} &= (a_{1}-1)b(1-a^{2}r_{h}^{2}), \quad L_{8} = 2ar_{h}(1-a^{2}r_{h}^{2})h(r_{h}), \\ L_{6\pm} &= (1\pm ab)(1\mp ab)^{a_{1}}, \quad L_{7\pm} = a^{2}\ell^{2}(2ac\pm 1)(1\pm ab)^{a_{1}}(1-a^{2}b^{2})^{a_{1}}, \\ L_{9} &= 2\ell^{2}c(1-a^{2}r_{h}^{2})^{2}+r_{h}^{2}\left[2r_{h}-b\left(a^{2}(1-2a_{1})r_{h}^{2}+2a_{1}+1\right)\right]h(r_{h})^{-2a_{1}}, \\ L_{3\pm} &= L_{6\pm}\left[2aL_{1\pm}-L_{2\pm}+2c(1\mp ab)(2ac\pm 1)\left(a^{2}\ell^{2}(1\pm ab)^{2a_{1}}-1\right)\right] \\ &+a_{1}bL_{7\pm}\left(4a^{2}c(b-c)-1\right), \\ L_{4\pm} &= L_{6\pm}\left[L_{2\pm}-2aL_{1\pm}+2c(ab\mp 1)\left(1\mp 6ac+a^{2}\ell^{2}(1\pm 2ac)(1\pm ab)^{2a_{1}}\right)\right] \\ &-a_{1}bL_{7\pm}\left(4a^{2}c(b-3c)+1\right), \\ L_{5\pm} &= -L_{6\pm}\left(2aL_{1\pm}+L_{2\pm}+2c(ab\mp 1)(1\pm 2ac)\right) \\ &+L_{7\pm}\left[2c+b\left(4a^{2}a_{1}c(c-b)-2a^{2}bc+a_{1}\right)\right], \\ L_{8\pm} &= L_{6\pm}\left(L_{1\pm}-4c^{2}(1\mp ab)\right)+2abca_{1}(b-2c)L_{7\pm}. \end{split}$$

Consequently, together with the tensions  $\mu_{\pm}$  defined by (2.9) we can quickly check both the first law (1.1) or (1.5) and Smarr relation (1.2) are satisfied.

The value of  $\alpha$  for arbitrary  $\zeta$  can not be directly calculated. However, when  $\zeta = 1, 1/\sqrt{3}, \sqrt{3}$ , which cases can be embedded in gauged supergravities, we find that the corresponding  $\alpha$  can be obtained by the holographic method below (4.9) with different stress-energy tensors which are related to different boundary conditions for the scalar field. Thus we write the  $\alpha$  and its corresponding  $\lambda_{\pm}$  of these three cases down as follows and discuss the issue of good/bad limit related to them:

• 
$$\zeta = 1$$

The  $\alpha$  in (4.1) and its corresponding  $\lambda_{\pm}$  are

$$\alpha = \sqrt{ac(1 - a^2\ell^2)},$$
  

$$\lambda_{\pm} = \frac{1}{2\alpha} \left( \frac{\mp 1 - ab(1 \mp 2ar_h)}{a(1 - a^2r_h^2)} + \frac{2r_h \mp a\left(\ell^2(1 \pm ar_h)^2 - r_h^2\right)}{1 \pm ar_h} \right).$$
(4.3)

This  $\alpha$  has a bad limit:  $\alpha \to 0$  as  $a \to 0$ . A better  $\alpha$  that exists a good limit  $(a \to 0, \alpha \to 1)$  and its corresponding  $\lambda_{\pm}$  are

$$\alpha = \sqrt{1 - a^2 \ell^2}, \qquad \lambda_{\pm} = \frac{1}{\alpha} \left( \frac{r_h}{1 \pm a r_h} - c \mp a \ell^2 - \frac{b}{2} \frac{1 \mp a r_h}{1 \pm a r_h} \right). \tag{4.4}$$

Although the  $\alpha$  in (4.1) does not have a good limit (see more details later), it is highly nontrivial to formulate the first law. The nonuniqueness of  $\alpha$  is related to the nonuniqueness of the solution of PDEs below we solve. Different  $\alpha$  leads to different  $\lambda_{\pm}$ . In particular, we can obtain the better  $\alpha$  only for the  $\zeta = 1$  solution.

• 
$$\zeta = 1/\sqrt{3}$$

$$\alpha = \sqrt{a\left(c + \frac{b}{4}\right)\left(1 - a^{2}\ell^{2}\left(1 + \frac{a^{2}bc^{2}}{c + \frac{b}{4}}\right)\right)},$$
  

$$\lambda_{\pm} = \frac{\sqrt{1 \pm ab}}{\alpha} \left(\frac{2ar_{h} \mp 1}{2a(1 - a^{2}r_{h}^{2})} - a^{2}\ell^{2}c(1 \pm \frac{1}{ar_{h}})\right).$$
(4.5)

•  $\zeta = \sqrt{3}$  $\alpha = \sqrt{\frac{a\left(c - \frac{b}{4}\right)\left(1 - a^{2}\ell^{2}\left(1 + \frac{b^{2}}{\ell^{2}} - \frac{a^{2}bc^{2}}{c - \frac{b}{4}}\right)\right)}{1 - a^{2}b^{2}}},$   $\lambda_{\pm} = \frac{1}{\alpha\sqrt{1 \pm ab}}\left(\frac{(r_{h} - b)(1 - a^{2}br_{h})}{1 - a^{2}r_{h}^{2}} - a^{2}\ell^{2}c(1 \pm \frac{1}{ar_{h}}) \mp \frac{1 + a^{2}br_{h}(a^{2}br_{h} - 2)}{2a(1 - a^{2}r_{h}^{2})}\right).$ (4.6)

Now, let us turn towards how to obtain all of these thermodynamic quantities.

• α:

When  $\zeta = 1, 1/\sqrt{3}$ , and  $\sqrt{3}$ , we can use the holographic method [21, 22] to specify  $\alpha$ . Write

$$\delta g_{(0)}^{ij} = \frac{\delta g_{(0)}^{ij}}{\delta K} \delta K + \frac{\delta g_{(0)}^{ij}}{\delta a} \delta a + \frac{\delta g_{(0)}^{ij}}{\delta b} \delta b + \frac{\delta g_{(0)}^{ij}}{\delta c} \delta c, \qquad (4.7)$$

$$\delta\mu_{\pm} = \frac{\delta\mu_{\pm}}{\delta K}\delta K + \frac{\delta\mu_{\pm}}{\delta a}\delta a + \frac{\delta\mu_{\pm}}{\delta b}\delta b + \frac{\delta\mu_{\pm}}{\delta c}\delta c, \qquad (4.8)$$

keep  $\delta \ell = 0$ ,  $\delta \mu_{\pm} = 0$ , and calculate

$$\delta S = \int_{\partial \mathcal{M}} d^3x \sqrt{-g_{(0)}} \langle \mathcal{T}_{ij} \rangle \delta g_{(0)}^{ij}.$$
(4.9)

Then we can require  $\delta S = 0$  to solve PDEs to obtain  $\alpha$  as a function of a, b, c, and  $\ell$ , and the results are (4.3), (4.4), (4.5), (4.6). We can see that these three cases are sufficient for us to guess an  $\alpha$  for arbitrary  $\zeta$ .

In particular, we can obtain  $\alpha$  only for the  $\zeta = 1/\sqrt{3}$  case with the corresponding stress-energy tensor (3.20). When  $\zeta = 1$  or  $\sqrt{3}$ , this stress-energy tensor can not give  $\alpha$  consistently, but we find that the  $\alpha$  for these two cases can be obtained by other stress-energy tensors. These stress-energy tensors are related to Dirichlet boundary conditions for  $\zeta = 1$  and unknown boundary conditions for  $\zeta = \sqrt{3}$ :

$$\zeta = \frac{1}{\sqrt{3}} \quad \Rightarrow \quad \langle \mathcal{T}_{ij} \rangle_{\mathrm{M}} = \frac{\ell^2}{16\pi G_4} \Big( 3g_{(3)ij} - \frac{2}{3} \frac{x_1}{\ell^2} \varphi^3_{(0)} g_{(0)ij} \Big), \tag{4.10}$$

$$\zeta = 1 \quad \Rightarrow \quad \langle \mathcal{T}_{ij} \rangle_{\mathrm{D}} = \frac{\ell^2}{16\pi G_4} \Big( 3g_{(3)ij} + \varphi_{(0)}\varphi_{(1)}g_{(0)ij} \Big), \tag{4.11}$$

$$\zeta = \sqrt{3} \quad \Rightarrow \quad \langle \mathcal{T}_{ij} \rangle_{\rm un} = \frac{\ell^2}{16\pi G_4} \Big( 3g_{(3)ij} + \frac{2}{3}\varphi_{(0)}\varphi_{(1)}g_{(0)ij} \Big). \tag{4.12}$$

Since the  $\zeta = 1$  case admits Dirichlet boundary conditions too, it is not surprised that the  $\langle \mathcal{T}_{ij} \rangle_{\rm D}$  can yield the same correct mass for this case. Although the boundary conditions for  $\langle \mathcal{T}_{ij} \rangle_{\rm un}$  are unclear, we can exactly obtain the same mass as the mixed conditions from  $\langle \mathcal{T}_{ij} \rangle_{\rm un}$  for arbitrary  $\zeta$ , and the differences of the local Ward identities for unknown boundary conditions are

$$\langle \mathcal{T}^i{}_i \rangle_{\rm un} = 0, \tag{4.13}$$

$$D_{(0)}^{j} \langle \mathcal{T}_{ij} \rangle_{\mathrm{un}} = \frac{\ell^2}{16\pi G_4} \left( \frac{2}{3} \varphi_{(1)} \partial_i \varphi_{(0)} - \frac{1}{3} \varphi_{(0)} \partial_i \varphi_{(1)} \right) + \langle \mathcal{J}^j \rangle_{\mathrm{M}} \mathcal{F}_{ij}^{(0)}.$$
(4.14)

Compared with mixed boundary conditions, this boundary condition preserves the conformal symmetry ( $\langle \mathcal{T}^i_i \rangle_{\mathrm{M}} \neq 0$ ). Further understanding on  $\langle \mathcal{T}_{ij} \rangle_{\mathrm{un}}$  is necessary. When c = 0 and b = 0, these solutions reduce to pure AdS. In order to get pure AdS in Rindler spacetime, it is necessary to take the following coordinate transformation [33]:

$$1 + \frac{R^2}{\ell^2} = \frac{1 + (1 - a^2 \ell^2) r^2 / \ell^2}{(1 - a^2 \ell^2) H^2}, \quad R \sin \vartheta = \frac{r \sin \theta}{H}.$$
 (4.15)

This takes the bulk spacetime to AdS in global coordinates

$$ds_{\rm AdS}^2 = -\left(1 + \frac{R^2}{\ell^2}\right)dt^2 + \frac{dR^2}{1 + \frac{R^2}{\ell^2}} + R^2\left(d\vartheta^2 + \sin^2\vartheta\frac{d\phi^2}{K^2}\right),\tag{4.16}$$

where we have chosen  $\alpha = \sqrt{1 - a^2 \ell^2}$ . However, when b and c equal zero, our  $\alpha$  for arbitrary  $\zeta$  in (4.1) can not recover this corresponding value. The better  $\alpha$  in (4.4) can recover this corresponding value and the differences are only different  $\lambda_{\pm}$ .

In contrast to [22], when  $\zeta = 1$ , the corresponding better  $\alpha = \sqrt{1 - a^2 \ell^2}$  is independent of the electric charge. The limit  $b \to 0$  takes these solutions to Schwarzchild accelerating black hole in AdS. However, except for the better  $\alpha$  in (4.4), this limit can not take our  $\alpha$  to the corresponding results in [21, 22], but the consequence of this difference is only a shift for the thermodynamic lengths  $(\lambda_{\pm})$ . We think that there exists better  $\alpha$  for arbitrary  $\zeta$ , which can be reduced to the corresponding results when we take all limits mentioned before. For example, when  $\zeta = 0$ , these solutions reduce to RN-AdS accelerating black hole. Compared with the  $\alpha$  in (4.1):

$$\alpha = \sqrt{a\left(c + \frac{b}{2}\right)\left(1 - a^2\ell^2(1 + 2a^2bc)\right)},$$
(4.17)

the corresponding better  $\alpha$  is

$$\alpha = \sqrt{\left(1 + 2a^2bc\right)\left(1 - a^2\ell^2(1 + 2a^2bc)\right)},\tag{4.18}$$

which exists a good limit. However, for arbitrary  $\zeta$ , it is not guaranteed to have such limits, which is similar to the limit  $\ell \to \infty$  can not take the thermodynamics of accelerating black hole in AdS to flat spacetime, which is mentioned in [22]. • *M*:

Following the main idea of [21, 22], we check the holographic mass with the conformal mass by Ashtekar-Magnon-Das (AMD) [46, 47]. First we perform a conformal transformation on the bulk metric,  $\bar{g}_{\mu\nu} = \bar{H}^2 g_{\mu\nu}$ , which leads to no divergence near the boundary. Then we can obtain the conserved charge with the integration of a conserved current related to a Killing vector  $\xi$ :

$$Q(\xi) = \frac{\ell}{8\pi G_4} \lim_{\bar{H}\to 0} \oint \frac{\ell^2}{\bar{H}} N^{\alpha} N^{\beta} \bar{C}^{\nu}{}_{\alpha\mu\beta} \xi_{\nu} d\bar{S}^{\mu}, \qquad (4.19)$$

where  $\bar{H} = \ell H r^{-1}$ ,  $N_{\mu} = \partial_{\mu} \bar{H}$  is the normal vector of the boundary,  $\bar{C}^{\mu}{}_{\alpha\nu\beta}$  is the Weyl tensor of  $\bar{g}_{\mu\nu}$ , and the tangent surface element of  $\bar{H} = 0$  is

$$d\bar{S}_{\mu} = \delta^{t}_{\mu} \frac{\ell^{2} f^{\frac{4\zeta^{2}}{1+\zeta^{2}}} d(\cos\theta) d\phi}{\alpha K}.$$
(4.20)

The conformal mass is given by  $M = Q(\partial_t)$ , which yields the same result as the holographic method.

• T and S:

To obtain the Hawking temperature of slowly accelerating black holes with charge and scalar hair, we use the Euclidean method as normal:

$$ds_{\tau,r}^2 \propto \left[\frac{F(r)}{\alpha^2} d\tau^2 + \frac{dr^2}{F(r)}\right] \quad \Rightarrow \quad T = \frac{F'(r_h)}{4\pi\alpha}.$$
(4.21)

The entropy is obtained simply by the horizon area formula [48] as:

$$S = \frac{\text{Area}}{4G_4} = \frac{1}{4KG_4} \int \frac{r^2 h^{\frac{2\zeta^2}{1+\zeta^2}} \sin\theta}{H^2} d\theta d\phi \Big|_{r=r_h} = \frac{\pi r_h^2 (1-\frac{b}{r_h})^{\frac{2\zeta^2}{1+\zeta^2}}}{KG_4 \left(1-a^2 r_h^2\right)}.$$
 (4.22)

• Q and  $\Phi$ :

The electric charge is obtained by Gauss's law integral and the chemical potential is obtained by Hawking-Ross prescription [49]

$$Q = \frac{1}{16\pi G_4} \int_{H=0} *\mathcal{F} = \frac{1}{2KG_4} \sqrt{\frac{2bc}{1+\zeta^2}},$$
(4.23)

$$\Phi = \frac{1}{16\pi G_4 Q\beta} \int_{\partial \mathcal{M}} \sqrt{\gamma} n_i \mathcal{F}^{ij} A_j = \frac{2}{\alpha r_h} \sqrt{\frac{2bc}{1+\zeta^2}}, \qquad (4.24)$$

where  $\beta$  is the inverse temperature (periodicity of Euclidean time),  $\gamma_{ij}$  is the induced metric, and  $n^i$  is the outward pointing unit normal vector of the boundary ( $\partial \mathcal{M} \equiv \{H = 0\}$ ).

•  $V, \eta$  and  $\lambda_{\pm}$ :

Finally, the condition  $F(r_h) = 0$  is used to derive the thermodynamic variables V,  $\eta$  and  $\lambda_{\pm}$ . Because the physical quantities  $P = 3/8\pi \ell^2 G_4$  (related to the cosmological constant and Newton's constant) and  $\mu_{\pm}$  (tensions of deficit) are fixed, we can utilize the following relation to construct the first law and obtain the remaining quantities:

$$\frac{\partial F_{r_h}}{\partial r_h}\delta r_h + \frac{\partial F_{r_h}}{\partial c}\delta c + \frac{\partial F_{r_h}}{\partial b}\delta b + \frac{\partial F_{r_h}}{\partial \ell}\delta \ell + \frac{\partial F_{r_h}}{\partial a}\delta a + \frac{\partial F_{r_h}}{\partial K}\delta K = 0.$$
(4.25)

After replacing  $\delta r_h$  with  $\delta S$  and  $F'(r_h)$  with  $4\pi\alpha T$ , fixing the term  $T\delta S$ , and replacing others with  $\delta M$ ,  $\delta Q$ ,  $\delta \mu_{\pm}$ ,  $\delta G_4$  and  $\delta P$ , we construct the first law (1.1). The coefficients related with  $\delta P$ ,  $\delta Q$  and  $\delta \mu_{\pm}$  are what we need in (4.1), (4.2) and (4.4). The coefficient of  $\delta G_4$  is correctly  $\eta/G_4$  at the same time.

# 5 Thermodynamics of $U(1)^2$ -charged accelerating black holes

Following the analysis of accelerating charged dilaton black holes, we can easily obtain the thermodynamics of  $U(1)^2$ -charged accelerating black holes. The bulk action is

$$S_{\text{bulk}} = \frac{1}{16\pi G_4} \int_{\mathcal{M}} d^{3+1}x \sqrt{-G} \left( R - \frac{1}{2} (\partial \phi)^2 - V(\phi) - \frac{1}{4} \sum_{I=1}^2 Z_I(\phi) \mathcal{F}_I^2 \right), \tag{5.1}$$

where  $\mathcal{F}_I = dA_I$ ,  $Z_1(\phi) = e^{\phi/\zeta}$ ,  $Z_2(\phi) = e^{-\zeta\phi}$ , and the scalar potential is the same as (2.2). When  $\zeta = 1/\sqrt{3}$ , 1,  $\sqrt{3}$ , these theories can be embedded in gauged STU supergravities.

These theories admit the  $U(1)^2$ -charged accelerating black hole solutions as follows [32]:

$$ds^{2} = \frac{1}{(1+arx)^{2}} \left[ f(x)^{j} \left( -\frac{F(r)}{\alpha^{2}} dt^{2} + \frac{dr^{2}}{F(r)} \right) + r^{2} h(r)^{j} \left( \frac{dx^{2}}{G(x)} + \frac{G(x)d\varphi^{2}}{K^{2}} \right) \right], \quad (5.2)$$

with

$$j = \frac{2\zeta^2}{1+\zeta^2}, \quad h(r) = 1 - \frac{q}{r}, \quad f(x) = 1 + aqx,$$
  

$$F(r) = (1 - a^2r^2)(1 - \frac{2m}{r} + \frac{e^2}{r^2})h(r)^{-j} + \frac{r^2}{\ell^2}h(r)^j,$$
  

$$G(x) = (1 - x^2)(1 + 2amx + a^2e^2x^2)f(x)^{-j},$$
  

$$A_1 = \frac{\sqrt{2j(1 - a^2q^2)(e^2 - q(2m - q))}}{\alpha} \left(\frac{1}{r_h h(r_h)} - \frac{1}{rh(r)}\right)dt,$$
  

$$A_2 = \frac{e\sqrt{2(2 - j)}}{\alpha} \left(\frac{1}{r_h} - \frac{1}{r}\right)dt, \quad \phi = \sqrt{j(2 - j)}\log\left(\frac{h(r)}{f(x)}\right).$$
  
(5.3)

When  $A_1 = 0$ , these solutions reduce to accelerating charged dilaton black holes. The limit  $q \to 0$  takes these solutions to RN-AdS accelerating black hole.

The FG expansion (3.15) of these solutions leads to the boundary metric as follows:

$$ds_{(0)}^2 = -\frac{\gamma_q^2(\rho)}{\ell^4 (1+aq\rho)^{4j}} dt^2 + \frac{\alpha^2}{\ell^2 X_q(\rho)\gamma_q(\rho)^2} d\rho^2 + \frac{\alpha^2 X_q(\rho)}{K^2 \ell^2 (1+aq\rho)^{4j}} d\varphi^2, \qquad (5.4)$$

where

$$X_q(\rho) = (1 - \rho^2)(1 + 2am\rho + a^2 e^2 \rho^2),$$
  

$$\gamma_q(\rho) = \sqrt{(1 + aq\rho)^{2j} - a^2 \ell^2 X_q(\rho)}.$$
(5.5)

The stress-energy tensor (4.12) is

$$\langle \mathcal{T}_{i}^{j} \rangle_{\rm un} = \frac{\ell^{4} (1 + aq\rho)^{2j - 3} \omega_{e}}{48\pi G_{4} \alpha^{3}} \operatorname{diag} \left[ -\left(2\gamma_{q}^{2} - a^{2}\ell^{2} X_{q}\right), (1 + aq\rho)^{2j}, \left(\gamma_{q}^{2} - 2a^{2}\ell^{2} X_{q}\right) \right], \quad (5.6)$$

and the stress-energy tensor (3.13) or (4.10) is

$$\langle \mathcal{T}_{i}^{j} \rangle_{\mathrm{M}} = \langle \mathcal{T}_{i}^{j} \rangle_{\mathrm{un}} + \frac{\ell^{4} (1 + aq\rho)^{4j-3} aj(2-j)\omega_{q}q^{2}}{24\pi G_{4}\alpha^{3}} \mathrm{diag}\left[1, 1, 1\right],$$
(5.7)

where

$$\begin{aligned} \omega_4 &= (1 - a^2 e^2)j + 6a(j-1)m\rho + 6a^2 e^2(j-2)\rho^2, \\ \omega_5 &= j(2j-1)\rho + am \left(3(j-2)(2j-3)\rho^2 - j(2j+5)\right) \\ &+ a^2 e^2 \rho \left(j-2j^2 + 2(j-2)(2j-9)\rho^2\right), \end{aligned} \tag{5.8} \\ \omega_6 &= 2j(1+j) + am\rho \left(j(2j-1) + (j-2)(2j-3)\rho^2\right) + 2a^2 e^2(j-3)(j-2)\rho^4, \\ \omega_e &= 6m + 12ae^2\rho - 3q\omega_4 + aq^2\omega_5 + a^2q^3\omega_6, \\ \omega_q &= \rho + a \left[q - ae^2\rho + ae^2\rho^3(2 + aq\rho) + m \left(3\rho^2 - 1 + aq\rho(1+\rho^2)\right)\right]. \end{aligned}$$

We find that these two stress-energy tensors yield the same correct mass as the conformal method (4.19), which means that the mass integration (3.22) related to the  $\omega_q$  term is zero. With these results, we can quickly check that the trace of the  $\langle \mathcal{T}_i^j \rangle_{\text{un}}$  is zero but the trace of the  $\langle \mathcal{T}_i^j \rangle_{\text{un}}$  is not. We can also check that the limit  $q \to 0$  takes these two results to reduce to the corresponding result in [22]. The differences of the dictionary (3.13) and the local Ward identities (3.14) are only one more vector field involved, but we are not going to write down the asymptotic expansions of the gauge and scalar field this time.

The thermodynamic quantities of these solutions are:

$$\begin{split} \Xi &= 1 + a^2 e^2, \qquad \Theta = \frac{e^2 - q(2m - q)}{1 - a^2 q^2}, \qquad \mu_{\pm} = \frac{1}{4G_4} \Big[ 1 - \frac{\Xi \pm 2am}{K(1 \pm aq)^j} \Big], \\ M &= \frac{2m \Big( 1 + (j - 1)a^2 q^2 \Big) - jq\Xi}{2\alpha KG_4 (1 - a^2 q^2)} - \frac{a\ell^2 \Big( \frac{(\Xi + 2am)^2}{(1 + aq)^{2j}} - \frac{(\Xi - 2am)^2}{(1 - aq)^{2j}} \Big)}{8\alpha KG_4}, \qquad \alpha = \sqrt{a\alpha KG_4 M}, \\ T &= \frac{h(r_h)^{-j}}{2\pi \alpha r_h^2} \Big[ (1 - a^2 r_h^2) (m - \frac{e^2}{r_h}) + \frac{r_h^2 \Big( r_h + qj(1 - a^2 r_h^2) - q \Big)}{\ell^2 h(r_h)^{1 - 2j} (1 - a^2 r_h^2)} \Big], \qquad \alpha = \sqrt{a\alpha KG_4 M}, \\ S &= \frac{\pi r_h^2 h(r_h)^j}{KG_4 \left( 1 - a^2 r_h^2 \right)}, \qquad Q_1 = \frac{\sqrt{2j\Theta}}{4KG_4}, \qquad \Phi_1 = \frac{\sqrt{2j\Theta}(1 - a^2 qr_h)}{\alpha r_h h(r_h)}, \qquad (5.9) \\ Q_2 &= \frac{e\sqrt{2(2 - j)}}{4KG_4}, \qquad \Phi_2 = \frac{e\sqrt{2(2 - j)}}{\alpha r_h}, \qquad P = \frac{3}{8\pi \ell^2 G_4}, \\ V &= \frac{4\pi \ell^2}{3\alpha K} \Big[ m - \alpha KMG_4 - \frac{je^2}{2r_h} + \frac{(1 - a^2 qr_h)j\Theta}{2r_h h(r_h)} + \frac{r_h^2 \Big( 2r_h - q \left( 2 - j - (1 - 2a^2 r_h^2) j \right) \Big)}{2\ell^2 h(r_h)^{1 - 2j} (1 - a^2 r_h^2)^2} \Big]. \end{split}$$

With these results, we can check that the Smarr relation (1.2) is satisfied. Since we can always construct the first law (1.1) from  $\delta F(r_h) = 0$  (4.25) with these thermodynamic quantities, and there are no simple expressions for the thermodynamic lengths ( $\lambda_{\pm}$ ) for arbitrary coupling constant. So we only write down  $\lambda_{\pm}$  for these cases, which can be embedded in gauged STU supergravities.

The thermodynamic lengths for  $\zeta=1,\,1/\sqrt{3},\,\sqrt{3}$  are:

•  $\zeta = 1$ 

The better  $\alpha$  which exists a good limit, and its corresponding thermodynamic lengths are

$$\alpha = \frac{\sqrt{\left(\Xi - 2a^2mq\right)\left(1 - a^2\ell^2(\Xi + \frac{q^2}{\ell^2} - 2a^2mq)\right)}}{1 - a^2q^2},$$

$$\lambda_{\pm} = \frac{Y_{0\pm} \pm (Y_1 + Y_{1\pm})q + (Y_{2\pm} \mp Y_2)q^2 + Y_{3\pm}q^3 + Y_{4\pm}q^4 + Y_{5\pm}q^5 - Y_6q^6}{\alpha Y(1 - a^2q^2)},$$
(5.10)

where

$$\begin{split} Y &= 2r_h^2 (1 - a^2 r_h^2) \Big[ \ell^2 (1 - a^2 r_h^2) \Big( 1 + a^2 (e^2 h(r_h) - qr_h) \Big) - a^2 r_h^3 q h(r_h)^2 \Big], \\ Y_{\pm} &= \ell^2 r_h \Big( ar_h (ar_h \mp 2\Xi) + 3\Xi - 2 \Big) \mp 2a \ell^4 (1 - a^2 r_h^2) \Xi^2, \\ Y_{0\pm} &= -r_h (1 - a^2 r_h^2) \Big( e^2 \ell^2 + r_h (r_h^3 - Y_{\pm}) \Big), \quad Y_1 = 4a^3 \ell^4 \Xi r_h (e^2 + r_h^2) (1 - a^2 r_h^2)^2, \\ Y_{1\pm} &= 2a \ell^2 r_h^2 (1 - a^2 r_h^2) \Big( r_h + ar_h^2 (3ar_h \mp 2) + ae^2 (2ar_h \mp) (1 + a^2 r_h^2) \Big) \\ &\pm 2r_h^4 (1 \mp ar_h) \Big( 1 \pm ar_h (1 \mp ar_h) \Big), \quad Y_2 = 2a^5 \ell^4 (e^2 + r_h^2)^2 (1 - a^2 r_h^2)^2, \quad (5.11) \\ Y_{2\pm} &= \mp 2a^2 \ell^2 r_h (1 - a^2 r_h^2) \Big\{ 2ar_h^3 (2 + a^2 r_h^2) + e^2 \Big[ ar_h \Big( 1 + ar_h (5ar_h \pm 1) \Big) \mp 1 \Big] \Big\} \\ &+ r_h^3 \Big[ a^2 r_h^2 \Big( 7 \mp 2ar_h (3 + a^2 r_h^2) \Big) - 1 \Big], \\ Y_{3\pm} &= 2a^2 r_h^2 \Big\{ a\ell^2 (1 - a^2 r_h^2) \Big[ ar_h^2 (2 \pm 3ar_h) \pm r_h + ae^2 \Big( 1 + ar_h (ar_h \pm 4) \Big) \Big] \\ &\pm r_h^2 \Big[ ar_h \Big( 3 + ar_h (3ar_h \pm 1) \Big) \mp 3 \Big] \Big\}, \\ Y_{4\pm} &= a^2 r_h \Big\{ a^2 \ell^2 (ar_h \mp 1) (ar_h \pm 1)^3 (e^2 + r_h^2) + r_h^2 \Big[ 2 \mp ar_h \Big( 2 + ar_h (1 \pm ar_h) (ar_h \pm 5) \Big) \Big] \Big\}, \\ Y_{5\pm} &= 2a^4 r_h^4 \Big( 2 + ar_h (ar_h \pm 1) \Big), \qquad Y_6 &= a^4 r_h^3 (1 + a^2 r_h^2). \end{split}$$

The  $\alpha$  in (5.9) and its corresponding  $\lambda_{\pm}$  are as follows:

$$\alpha = \sqrt{\frac{a\left(2m - q\Xi\right)\left(1 - a^{2}\ell^{2}(\Xi + \frac{q^{2}}{\ell^{2}} - 2a^{2}mq)\right)}{2(1 - a^{2}q^{2})^{2}}},$$

$$\lambda_{\pm} = \frac{1}{2\alpha(1 \pm aq)} \left[\frac{ar_{h}(1 \mp ar_{h})(2 \pm ar_{h}) \mp 1}{a(1 - a^{2}r_{h}^{2})} \mp a\ell^{2}\left(1 \pm ar_{h}\right)\left(1 \pm \frac{ae^{2}}{r_{h}}\right) - q\frac{2 \pm a\left(q - 4r_{h} + aqr_{h}(ar_{h} \mp 1)\right)}{1 - a^{2}r_{h}^{2}}\right].$$
(5.12)

• 
$$\zeta = 1/\sqrt{3}$$
  
 $\alpha = \sqrt{\frac{a\left[4m\left(1 - a^{2}\ell^{2}(\Xi + \frac{q^{2}}{2\ell^{2}} - a^{2}mq)\right) - \Xi(1 - a^{2}\ell^{2}\Xi)q\right]}{4(1 - a^{2}q^{2})}},$  (5.13)  
 $\lambda_{\pm} = \frac{1}{2\alpha\sqrt{1 \pm ab}}\left(r_{h} - a^{2}\ell^{2}(r_{h} + \frac{e^{2}}{r_{h}}) \pm \frac{1 - a^{2}\ell^{2}\Xi}{a} + \frac{ar_{h} \mp 2}{a(1 - a^{2}r_{h}^{2})} - \frac{q(1 \mp 2ar_{h})}{1 - a^{2}r_{h}^{2}}\right).$   
•  $\zeta = \sqrt{3}$   
 $\alpha = \sqrt{\frac{a\left(2m(2 + a^{2}q^{2}) - 3q\Xi\right)}{4(1 - a^{2}q^{2})} - \frac{1}{8}a^{2}\ell^{2}\left(\frac{(\Xi + 2am)^{2}}{(1 + aq)^{3}} - \frac{(\Xi - 2am)^{2}}{(1 - aq)^{3}}\right)}{(1 - aq)^{3}},$   
 $\lambda_{\pm} = \frac{1}{2\alpha\sqrt{(1 \pm ab)^{3}}}\left[r_{h} - a^{2}\ell^{2}(r_{h} + \frac{e^{2}}{r_{h}}) \pm \frac{1 - a^{2}\ell^{2}\Xi}{a} + \frac{ar_{h} \mp 2}{a(1 - a^{2}r_{h}^{2})}\right] + q\frac{ar_{h}\left(aq(3 \pm 2aq) \pm 6\right) - 3(1 \pm aq) - qa^{3}r_{h}^{2}(aq \pm 3)}{1 - a^{2}r_{h}^{2}}\right].$ 

There is no doubt that one can analyze the  $U(1)^4$ -charged accelerating black holes [32] along this line.

#### 6 Summary and Discussion

To summarize, we have derived the thermodynamic laws for slowly accelerating charged dilaton black holes in four-dimensional Einstein-Maxwell-dilaton theories for arbitrary dilaton coupling constant. These systems can be embedded in gauged supergravities when  $\zeta = 0, 1/\sqrt{3}, 1, \text{ and } \sqrt{3}$ . The thermodynamics of slowly accelerating black holes is generalized to include scalar hair, which does not introduce a scalar charge. We also find that the holographic mass associated with appropriate boundary conditions for the scalar agrees with the AMD mass. These results can be straightforwardly generalized to  $U(1)^2$ -charged accelerating black holes. We have also computed the holographic stress-energy tensor, which can be explained as an energy-momentum tensor of perfect fluid plus a non-hydrodynamic correction [21, 22, 50], and other dual CFT one-point functions.

Our results open a way to analyze the phase structure of slowly accelerating black holes with gauge and scalar filed in asymptotically AdS spacetime. Even for the non-accelerating black hole in the  $\zeta = 1$  case, rich phase structures (e.g. zeroth order phase transition) are obtained [37]. Thus, the general accelerating hairy black hole is expected to have richer phases. Since there is thermodynamic instability for non-accelerating charged dilaton black holes when  $\zeta > \zeta_{max}$  [51], with our results one can also study the thermodynamic stability or instability for hairy accelerating black holes.

Two key points of our work are properly considering mixed boundary conditions for the scalar field and rescaling the time coordinate with parameter  $\alpha$ . However, we only find the better  $\alpha$  for the  $\zeta = 1$  solution. The better  $\alpha$  for arbitrary  $\zeta$  is still an unsolved problem,

or maybe it is similar to the fact that the limit  $\ell \to \infty$  can not take the thermodynamics of accelerating black hole in AdS to flat spacetime. Although the hairy accelerating black holes emit gravitational and electromagnetic radiation at infinity as usual accelerating black holes [52, 53], the standard thermodynamic analysis is still valid; the thermodynamic quantities we obtained are related to the comoving observer, who can not receive any radiation.

Since there are gauge and scalar field in the bulk, apart from the gravitational field described by the boundary metric, there are also a source  $J_{\mathcal{F}_{\varphi}}$  of dual CFT operator, one or two non-dynamical electromagnetic field  $\mathcal{F}_{ij}^{(0)}$ , and one or two current  $\mathcal{J}^i$  generated by the quantum field. Since the Lorentz force, the source of dual theory, and the gravitational field counterbalance each other in the background geometry, there is a stationary configuration of plasma or fluid. In particular, the constraints  $\gamma_b^2(\rho) > 0$  and  $\gamma_q^2(\rho) > 0$  shift the other horizons beyond our parameter space. If these conditions are allowed to violate, the black funnels and droplets [10, 11] with scalar hair will appear, we expect that properly taking into account the parameter restrictions also can take these systems to thermal equilibrium. Constructing black funnels and droplets with scalar hair, the thermodynamics of pair hairy accelerating black holes associated with boost mass [54], and the thermodynamics of pair creation of dilaton black holes [30] (hairy accelerating black holes in asymptotically flat spacetime) are also interesting projects for the future.

# Acknowledgments

This work is supported in part by the NSF of China under Grant No. 11905298.

#### References

- [1] H. Weyl, On the theory of gravitation, Ann. Phys. 54 (1917) 117.
- [2] J. Ehlers, W. Kundt, Exact solutions of the gravitational field equations, in Gravitation: an introduction to current research, edited by L. Witten (Wiley, New York, London, 1962).
- W. Kinnersley and M. Walker, Uniformly accelerating charged mass in general relativity, Phys. Rev. D 2 (1970), 1359-1370
- [4] J. F. Plebanski and M. Demianski, Rotating, charged, and uniformly accelerating mass in general relativity, Annals Phys. 98 (1976), 98-127
- [5] O. J. C. Dias and J. P. S. Lemos, Pair of accelerated black holes in anti-de Sitter background: AdS C metric, Phys. Rev. D 67 (2003), 064001 [arXiv:hep-th/0210065].
- [6] J. B. Griffiths, P. Krtous and J. Podolsky, Interpreting the C-metric, Class. Quant. Grav. 23 (2006), 6745-6766 [arXiv:gr-qc/0609056].
- [7] R. Emparan, G. T. Horowitz and R. C. Myers, *Exact description of black holes on branes*, JHEP 01 (2000), 007 [arXiv:hep-th/9911043].
- [8] R. Emparan, G. T. Horowitz and R. C. Myers, Exact description of black holes on branes. 2. Comparison with BTZ black holes and black strings, JHEP 01 (2000), 021 [arXiv:hep-th/9912135].
- [9] R. Emparan and G. Milanesi, Exact Gravitational Dual of a Plasma Ball, JHEP 08 (2009), 012 [arXiv:0905.4590].

- [10] V. E. Hubeny, D. Marolf and M. Rangamani, Black funnels and droplets from the AdS C-metrics, Class. Quant. Grav. 27 (2010), 025001 [arXiv:0909.0005].
- [11] M. M. Caldarelli, O. J. C. Dias, R. Monteiro and J. E. Santos, Black funnels and droplets in thermal equilibrium, JHEP 05 (2011), 116 [arXiv:1102.4337].
- [12] R. Emparan, A. M. Frassino and B. Way, *Quantum BTZ black hole*, JHEP **11** (2020), 137 [arXiv:2007.15999].
- [13] J. D. Bekenstein, Black holes and entropy, Phys. Rev. D 7, 2333 (1973).
- [14] J. D. Bekenstein, Generalized second law of thermodynamics in black hole physics, Phys. Rev. D 9, 3292 (1974).
- [15] S. W. Hawking, *Particle Creation by Black Holes*, Commun. Math. Phys. 43, 199 (1975)
   Erratum: [Commun. Math. Phys. 46, 206 (1976)].
- [16] D. Kubiznak, R. B. Mann and M. Teo, Black hole chemistry: thermodynamics with Lambda, Class. Quant. Grav. 34 (2017) no.6, 063001 [arXiv:1608.06147].
- [17] M. Astorino, Thermodynamics of Regular Accelerating Black Holes, Phys. Rev. D 95 (2017) no.6, 064007 [arXiv:1612.04387].
- [18] M. Astorino, CFT Duals for Accelerating Black Holes, Phys. Lett. B 760 (2016), 393-405 [arXiv:1605.06131].
- [19] M. Appels, R. Gregory and D. Kubiznak, *Thermodynamics of Accelerating Black Holes*, Phys. Rev. Lett. **117** (2016) no.13, 131303 [arXiv:1604.08812].
- [20] M. Appels, R. Gregory and D. Kubiznak, Black Hole Thermodynamics with Conical Defects, JHEP 05 (2017), 116 [arXiv:1702.00490].
- [21] A. Anabalón, M. Appels, R. Gregory, D. Kubizňák, R. B. Mann and A. Ovgün, *Holographic Thermodynamics of Accelerating Black Holes*, Phys. Rev. D 98 (2018) no.10, 104038 [arXiv:1805.02687].
- [22] A. Anabalón, F. Gray, R. Gregory, D. Kubizňák and R. B. Mann, Thermodynamics of Charged, Rotating, and Accelerating Black Holes, JHEP 04 (2019), 096 [arXiv:1811.04936].
- [23] G. W. Gibbons, M. J. Perry and C. N. Pope, The First law of thermodynamics for Kerr-anti-de Sitter black holes, Class. Quant. Grav. 22 (2005), 1503-1526 [arXiv:hep-th/0408217].
- [24] D. Cassani, J. P. Gauntlett, D. Martelli and J. Sparks, Thermodynamics of accelerating and supersymmetric AdS4 black holes, Phys. Rev. D 104 (2021) no.8, 086005 [arXiv:2106.05571].
- [25] M. R. Visser, Holographic Thermodynamics Requires a Chemical Potential for Color, [arXiv:2101.04145].
- [26] W. Cong, D. Kubiznak and R. B. Mann, Thermodynamics of AdS Black Holes: Critical Behavior of the Central Charge, Phys. Rev. Lett. 127 (2021) no.9, 091301 [arXiv:2105.02223].
- [27] H. Lü and J. F. Vazquez-Poritz, Dynamic C-metrics in gauged supergravities, Phys. Rev. D 91 (2015) 064004 [arXiv:1408.3124].
- [28] J. Ren, Phase transitions of hyperbolic black holes in anti-de Sitter space, arXiv:1910.06344.
- [29] J. Ren and W. Zheng, Analytic AC conductivities from holography, Phys. Rev. D 105 (2022) no.6, 066013 [arXiv:2109.07481].

- [30] F. Dowker, J. P. Gauntlett, D. A. Kastor and J. H. Traschen, Pair creation of dilaton black holes, Phys. Rev. D 49 (1994) 2909 [hep-th/9309075].
- [31] R. Gregory and M. Hindmarsh, Smooth metrics for snapping strings, Phys. Rev. D 52 (1995), 5598-5605 [arXiv:gr-qc/9506054].
- [32] H. Lü and J. F. Vázquez-Poritz, C-metrics in Gauged STU Supergravity and Beyond, JHEP 12 (2014), 057 [arXiv:1408.6531].
- [33] J. Podolsky, Accelerating black holes in anti-de Sitter universe, Czech. J. Phys. 52 (2002), 1-10 [arXiv:gr-qc/0202033].
- [34] G. W. Gibbons, R. Kallosh and B. Kol, Moduli, scalar charges, and the first law of black hole thermodynamics, Phys. Rev. Lett. 77 (1996), 4992-4995 [arXiv:hep-th/9607108].
- [35] H. Lü, Y. Pang and C. N. Pope, AdS Dyonic Black Hole and its Thermodynamics, JHEP 11 (2013), 033 [arXiv:1307.6243].
- [36] H. Lu, C. N. Pope and Q. Wen, Thermodynamics of AdS Black Holes in Einstein-Scalar Gravity, JHEP 03 (2015), 165 [arXiv:1408.1514].
- [37] M. M. Caldarelli, A. Christodoulou, I. Papadimitriou and K. Skenderis, *Phases of planar AdS black holes with axionic charge*, JHEP 04 (2017), 001 [arXiv:1612.07214].
- [38] D. Astefanesei, R. Ballesteros, D. Choque and R. Rojas, Scalar charges and the first law of black hole thermodynamics, Phys. Lett. B 782 (2018), 47-54 [arXiv:1803.11317].
- [39] L. Smarr, Mass formula for Kerr black holes, [erratum: Phys. Rev. Lett. 30 (1973), 521-521]
   Phys. Rev. Lett. 30 (1973), 71-73
- [40] D. Kastor, S. Ray and J. Traschen, Enthalpy and the Mechanics of AdS Black Holes, Class. Quant. Grav. 26 (2009), 195011 [arXiv:0904.2765].
- [41] C. J. Gao and S. N. Zhang, Dilaton black holes in de Sitter or Anti-de Sitter universe, Phys. Rev. D 70 (2004), 124019 [arXiv:hep-th/0411104].
- [42] C. Fefferman and C. Robin Graham, Conformal Invariants in Elie Cartan et les Mathématiques d'aujourd 'hui (Astérisque, 1985) Numero Hors Serie, 95-116, see also: [arXiv:0710.0919].
- [43] E. Witten, Multitrace operators, boundary conditions, and AdS/CFT correspondence, [arXiv:hep-th/0112258].
- [44] W. Mueck, An Improved correspondence formula for AdS/CFT with multitrace operators, Phys. Lett. B 531 (2002), 301-304 [arXiv:hep-th/0201100].
- [45] I. Papadimitriou, Multi-Trace Deformations in AdS/CFT: Exploring the Vacuum Structure of the Deformed CFT, JHEP 05 (2007), 075 [arXiv:hep-th/0703152].
- [46] A. Ashtekar and A. Magnon, Asymptotically anti-de Sitter space-times, Class. Quant. Grav. 1 (1984), L39-L44
- [47] A. Ashtekar and S. Das, Asymptotically Anti-de Sitter space-times: Conserved quantities, Class. Quant. Grav. 17 (2000), L17-L30 [arXiv:hep-th/9911230].
- [48] C. Herdeiro, B. Kleihaus, J. Kunz and E. Radu, On the Bekenstein-Hawking area law for black objects with conical singularities, Phys. Rev. D 81 (2010), 064013 [arXiv:0912.3386].
- [49] S. W. Hawking and S. F. Ross, Duality between electric and magnetic black holes, Phys. Rev. D 52 (1995), 5865-5876 [arXiv:hep-th/9504019].

- [50] G. Bernardi de Freitas and H. S. Reall, Algebraically special solutions in AdS/CFT, JHEP 06 (2014), 148 [arXiv:1403.3537].
- [51] A. Sheykhi, M. H. Dehghani and S. H. Hendi, *Thermodynamic instability of charged dilaton black holes in AdS spaces*, Phys. Rev. D 81 (2010), 084040 [arXiv:0912.4199].
- [52] J. Podolsky, M. Ortaggio and P. Krtous, Radiation from accelerated black holes in an anti-de Sitter universe, Phys. Rev. D 68 (2003), 124004 [arXiv:gr-qc/0307108].
- [53] J. Podolsky and H. Kadlecova, Radiation generated by accelerating and rotating charged black holes in (anti-)de Sitter space, Class. Quant. Grav. 26 (2009), 105007 [arXiv:0903.3577].
- [54] K. Dutta, S. Ray and J. Traschen, Boost mass and the mechanics of accelerated black holes, Class. Quant. Grav. 23 (2006), 335-352 [arXiv:hep-th/0508041].