The normalized Laplacian spectrum of *n*-polygon graphs and its applications

Tengjie Chen^{a,b} Zhenhua Yuan^{a,b} Junhao Peng^{a,b}

^aSchool of Math and Information Science, Guangzhou University, Guangzhou 510006, China.

^bGuangdong Provincial Key Laboratory co-sponsored by province and city of Information Security Technology, Guangzhou University, Guangzhou 510006, China.

Abstract

Given an arbitrary connected G, the *n*-polygon graph $\tau_n(G)$ is obtained by adding a path with length $n \ (n \ge 2)$ to each edge of graph G, and the iterated *n*-polygon graphs $\tau_n^g(G) \ (g \ge 0)$, is obtained from the iteration $\tau_n^g(G) = \tau_n(\tau_n^{g-1}(G))$, with initial condition $\tau_n^0(G) = G$. In this paper, a method for calculating the eigenvalues of normalized Laplacian matrix for graph $\sigma_n(G)$ is presented if the eigenvalues of normalized Laplacian matrix for graph G is given firstly. Then, the normalized Laplacian spectrums for the graph $\tau_n(G)$ and the graphs $\tau_n^g(G) \ (g \ge 0)$ can also be derived. Finally, as applications, we calculate the multiplicative degree-Kirchhoff index, Kemeny's constant and the number of spanning trees for the graph $\tau_n(G)$ and the graphs $\tau_n^g(G)$ by exploring their connections with the normalized Laplacian spectrum, exact results for these quantities are obtained.

Key words: Normalized Laplacian spectrum, Multiplicative degree-Kirchhoff index, Kemeny's constant, The number of spanning trees. *PACS:* 05.40.Fb, 05.60.Cd

1 Introduction

The spectra of the adjacency matrices, Laplacian and the normalized Laplacian for a graph is of particular important since many important structural and dynamical properties of the graph can be obtained from the eigenvalues

Email address: pengjh@gzhu.edu.cn (Junhao Peng).

and eigenvectors of these matrices [1,2]. For example, the spectra of these matrices provides information on degree distribution, the multiplicative degree-Kirchhoff index, community structure, local clustering, total number of links, the number of spanning trees and etc. [2-4]. Also, first-passage properties of the networks [5,6], such as mean first-passage time, the mean hitting time, the mixing time and the Kemeny's const, can be expressed in terms of the eigenvalues of the Laplacian and normalized Laplacian [2,7,8]. In the past several years, there is a particular interest in the study of spectra for the normalized Laplacian matrices of different graphs [9-17].

Recently, a class of iterated graphs obtained by replacing each edge of a 'base graph' with a given structure, have attracted lots attention [18, 19], Since these graphs not only exhibit rich structural properties, such as self-similarity, fractal [19–21], scale-free and small-world properties [18, 22], but also show distinctive dynamic properties [20–24].

For an arbitrary connected 'base graph' G, if the normalized Laplacian spectrum of graph G is known, how can we calculate the normalized Laplacian spectrum and related quantities of graph $\tau(G)$, which is obtained from Gby replacing each edge of G with a given structure. It is an interesting topic with wide applications and lots of results were obtained. Examples include the graph $\tau(G)$ obtained by replacing every edges of G with a geometry, such as triangles [25–27], a quadrilateral [28–30], a pentagon [31]. However, for more general graphs $\tau_n(G)$ obtained by replacing every edges of G by a (n + 1)polygon $(n \geq 2)$, related topic is more difficult and general result for the normalized Laplacian spectrum $\tau_n(G)$ is still unknown.

To fill this gap, we consider the case while every edges of G are replaced by a (n + 1)-polygon. We analyze the relation between the normalized Laplacian spectrum of the graph $\tau_n(G)$ and that of graph G and present detailed spectrum of the graph $\tau_n(G)$ for any $n \ge 2$. Using our results recursively, we also obtain the normalized Laplacian spectrum of the iterated graph $\tau_n^g(G)$ $(g \ge 0)$, where $\tau_n^0(G) = G$, and $\tau_n^g(G) = \tau_n(\tau_n^{g-1}(G))$. As applications, we also analyze the multiplicative degree-Kirchhoff index, the Kemeny's const and the number of spanning trees of the graph $\tau_n^g(G)$. Exactly results for these quantities are also presented.

This paper is organized as follows. First, in Sec. 2, we present related definitions and notations which will appear in the manuscript. In Sec. 3, we present some classical results we quote and some preliminary results which will be used in the manuscript. Then, in Sec. 4, we present the main results on the normalized Laplacian spectrum of the graph $\tau_n(G)$, and in Sec. 5, we calculate the multiplicative degree-Kirchhoff index, the Kemeny's const and the number of spanning trees of the graph $\tau_n^t(G)$ and $\tau_n^g(G)$. The detailed proof of our results are presented in Sec.6-10 and Appendix A-C.

2 Definitions and notations

Let G = (V(G), E(G)) be a undirected graph with vertex set V(G) and edge set E(G), $N_0 = |V(G)|$ represent the total number of nodes of G and $E_0 = |E(G)|$ be the total number of edges of graph G. For any two nodes $i, j \in V(G)$, if there is an edge between node i and j in E(G), we say i is a neighbor of j or i and j are adjacent (i.e., $i \sim j$). If e is an edge with end-vertices i and j, we say edge e is incident to nodes i and j, or nodes i and j are incident with edge e. The degree of vertex i, referred to as d_i , is the number of edges incident to node i.

Definition 2.1 [2] For any undirected graph G, the Laplacian matrix of G is defined as

$$L(G) = D(G) - A(G),$$

where $D(G) = diag(d_1, d_2, \dots, d_{N_0})$ is the degree diagonal matrix of G and $A(G) = (A_{ij})_{N_0 \times E_0}$ is the adjacency matrix of G, with

$$A_{ij} = \begin{cases} 1 \ i \sim j \\ 0 \ others \end{cases}$$
(1)

Definition 2.2 [2,32] For any undirected graph G, the normalized Laplacian of G is defined to be

$$\mathcal{L}_G = D(G)^{-\frac{1}{2}} L(G) D(G)^{-\frac{1}{2}} = I - D(G)^{-\frac{1}{2}} A(G) D(G)^{-\frac{1}{2}}.$$
 (2)

Let δ_{ij} be the Kronecker delta function. Then, the (i, j)-th entry of the matrix \mathcal{L}_G can be written as

$$\mathcal{L}_G(i,j) = \delta_{ij} - \frac{A_{ij}}{\sqrt{d_i d_j}}.$$
(3)

Definition 2.3 [2] For any graph undirected G with N_0 vertexes, the normalized Laplacian spectrum of graph G is defined to be

$$\sigma(G) = \{\lambda_1, \lambda_2, \cdots, \lambda_{N_0}\}, \qquad (4)$$

where λ_i $(i = 1, 2, \dots, N_0)$ are the eigenvalues of \mathcal{L}_G .

Definition 2.4 [3] Let G be a undirected graph, with vertex set $V(G) = \{1, 2, \dots, N_0\}$ and edge set $E(G) = \{e_1, e_2, \dots, e_{E_0}\}$. Then the incidence matrix of graph G is defined by

$$B = (b_{ij})_{N_0 \times E_0} \,,$$



Fig. 1. The construction method of the *n*-polygon graph. The *n*-polygon graph of G, denoted by $\tau_n(G)$, is obtained from G by replacing each edge ij of G with the (n + 1)-polygon on the right-hand side of the arrow.

where

$$b_{ij} = \begin{cases} 1, \text{ if vertex } i \text{ and edge } e_j \text{ are incident} \\ 0, \text{ otherwise} \end{cases}.$$

Definition 2.5 [3, 7] Let G' be a directed graph, $V(G') = \{1, 2, \dots, N_0\}$ and directed edge set $E(G') = \{e'_1, e'_2, \dots, e'_{E_0}\}$. Then the incidence matrix of directed graph G' is defined by

$$B' = \left(b'_{ij}\right)_{N_0 \times E_0},$$

where

$$b_{ij}^{'} = \begin{cases} -1, \text{ if vertex } i \text{ is the source vertex of edge } e_{j}^{'} \\ 0, \text{ if vertex } i \text{ is not incident on edge } e_{j}^{'} \\ 1, \text{ if vertex } i \text{ is the target vertex of edge } e_{j}^{'} \end{cases}$$

Definition 2.6 [3, 7] Let G' be a directed graph, G' is called as weakly connected if its underlying undirected graph (i.e. a graph obtained by replacing each directed edge of G' with an undirected edge) is connected.

Definition 2.7 The n-polygon $(n \ge 2)$ graph of G, denoted by $\tau_n(G)$, is the graph obtained by adding a path with length n to each edge of G.

In other word, τ_n is a operator which turns each edge of a graph into a (n+1)-polygon. Thus, $\tau_n(G)$ is a graph obtained by replacing each edge ij of G with a (n+1)-polygon, as shown in Fig. 1.

Definition 2.8 For any g > 0, the iterated n-polygon graph with generation $g \ (g \ge 1)$, referred to as $\tau_n^g(G)$, is defined as the graph obtained through the iteration $\tau_n^g(G) = \tau_n(\tau_n^{g-1}(G))$, with initial condition $\tau^0(G) = G$.

Let N_g and E_g be the total number of vertices and edges of graph $\tau_n^g(G)$ respectively.

For any g > 0, we have

$$N_g = N_{g-1} + (n-1)E_{g-1}, \ E_g = (n+1)E_{g-1}.$$

Therefore

$$N_g = N_0 + (n-1)\frac{(n+1)^g - 1}{n}E_0,$$
(5)

and

$$E_g = (n+1)^g E_0. (6)$$

3 Preliminaries

In this section, we present some classical results we quote and some preliminary results which will be used to derive our main results in the manuscript.

Lemma 3.1 [3] For any connected undirected graph G, let $\lambda_1 \leq \lambda_2 \leq \cdots \leq \lambda_{N_0}$ be the eigenvalues of \mathcal{L}_G , Then

- (i) $\lambda_1 = 0$, and for any $2 \le i \le N_0$, $0 < \lambda_i \le 2$;
- (ii) $\lambda_{N_0} = 2$ if and only if G is bipartite;
- (iii) G is bipartite if and only if both λ_i and $2 \lambda_i$ are eigenvalues of \mathcal{L}_G and $m_{\mathcal{L}_G}(\lambda_i) = m_{\mathcal{L}_G}(2 \lambda_i)$, where $m_{\mathcal{L}_G}(\lambda_i)$ is the multiplicity of λ_i which is the eigenvalue of \mathcal{L}_G .

Therefore, all the eigenvalues of \mathcal{L}_G are non-negative. Given the normalized Laplacian spectrum of graph G, the multiplicative degree-Kirchhoff index, the Kemeny's constant and the number of spanning trees of graph G can be expressed as follows.

Lemma 3.2 [2, 3, 7] Let G be a connected undirected graph with N_0 vertices and E_0 edges, $0 = \lambda_1 < \lambda_2 \leq \cdots \leq \lambda_{N_0}$ are the eigenvalues of the normalized Laplacian \mathcal{L}_G . Then,

(i) the multiplicative degree-Kirchhoff index of G can be written as

$$Kf'(G) = 2E_0 \sum_{i=2}^{N_0} \frac{1}{\lambda_i};$$

(ii) the Kemeny's constant of G is

$$K(G) = \sum_{i=2}^{N_0} \frac{1}{\lambda_i};$$

(iii) the number $N_{st}(G)$ of spanning trees of G is

$$N_{st}(G) = \frac{1}{2E_0} \left(\prod_{i=1}^{N_0} d_i\right) \prod_{i=2}^{N_0} \lambda_i.$$

Lemma 3.3 [33]Let $f(x) = b_n x^n + b_{n-1} x^{n-1} + \cdots + b_1 x + b_0$ ($b_n \neq 0$) be a polynomial of degree n and r_i ($i = 1, 2, \cdots, n$) be the roots of f(x) = 0. Then Vieta's formulas states that

$$\prod_{i=1}^{n} r_i = (-1)^n \frac{b_0}{b_n},$$
$$\sum_{i=1}^{n} \frac{1}{r_i} = -\frac{b_1}{b_0}.$$

and

Lemma 3.4 [2] Let B be the incidence matrix of a connected graph G with
$$N_0$$
 vertices. Then

$$rank(B) = \begin{cases} N_0 - 1 \text{ if } G \text{ is bipartite,} \\ N_0 & \text{if } G \text{ is non-bipartite.} \end{cases}$$

Lemma 3.5 [3, 7] Let B' be the incidence matrix of weakly connected directed graph G' with N_0 vertices. Then rank $(B') = N_0 - 1$.

Lemma 3.6 Let u be an arbitrary real number, and $\{a_n(\mu)\}_{n\geq -1}$ is a series which defined by the recursive equation $a_n(\mu) = 2(1-\mu)a_{n-1}(\mu) - a_{n-2}(\mu)$, with initial conditions $a_{-1}(\mu) = 0$ and $a_0(\mu) = 1$. Then

(i) for any $n \ge 1$, $a_n(2-\mu) = (-1)^n a_n(\mu)$; (ii) if $\mu = 0$ $a_n(0) = n+1$, (7)

and if $\mu = 2$,

$$a_n(2) = (-1)^n (n+1);$$
 (8)

(iii) for any $n \ge 0$, $a_n(\mu)$ is an nth order polynomial in μ , which can be written as

$$a_n(\mu) = \sum_{i=0}^n a_n^{(i)} \mu^i$$

where $a_n^{(i)}$ is the coefficient of μ^i ; further more,

$$a_n^{(0)} = n + 1, (9)$$

$$a_n^{(1)} = -\frac{n^3 + 3n^2 + 2n}{3},\tag{10}$$

$$a_n^{(n)} = (-1)^n \, 2^n; \tag{11}$$

(iv) for any $n \ge 2$, $a_n(\mu)$ and $1 + a_n(\mu)$ can be expanded as

$$a_{n}(\mu) = \begin{cases} \left(a_{\frac{n-1}{2}+1}(\mu) - a_{\frac{n-1}{2}-1}(\mu)\right) a_{\frac{n-1}{2}}(\mu) & n \text{ is odd} \\ \left(a_{\frac{n}{2}}(\mu) - a_{\frac{n}{2}-1}(\mu)\right) \left(a_{\frac{n}{2}}(\mu) + a_{\frac{n}{2}-1}(\mu)\right) n \text{ is even} \end{cases}, \quad (12)$$

and

$$1+a_{n}(\mu) = \begin{cases} \left(a_{\frac{n+1}{2}-1}(\mu) - a_{\frac{n+1}{2}-2}(\mu)\right) \left(a_{\frac{n+1}{2}}(\mu) + a_{\frac{n+1}{2}-1}(\mu)\right) & n \text{ is odd} \\ \left(a_{\frac{n}{2}}(\mu) - a_{\frac{n}{2}-2}(\mu)\right) a_{\frac{n}{2}}(\mu) & n \text{ is even} \end{cases}$$
(13)

The proof of Lemma 3.6 is presented in Appendix A.

Lemma 3.7 If $n \ (n \ge 3)$ is an odd number, and $\{a_k(\mu)\}_{k\ge 0}$ is a series defined in Lemma 3.6, we have the following results.

Let μ be an arbitrary root of equation $a_{\frac{n-1}{2}}(\mu) - a_{\frac{n-1}{2}-1}(\mu) = 0$. Then

$$1 - \frac{a_{\frac{n-1}{2}+1}(\mu) - a_{\frac{n-1}{2}-1}(\mu)}{a_{\frac{n-1}{2}}(\mu) - a_{\frac{n-1}{2}-2}(\mu)} = 2,$$
(14)

$$a_{n-2}(\mu) = 1. (15)$$

((i) Let μ be an arbitrary root of equation $a_{\frac{n-1}{2}}(\mu) + a_{\frac{n-1}{2}-1}(\mu) = 0$. Then

$$1 - \frac{a_{\frac{n-1}{2}+1}(\mu) - a_{\frac{n-1}{2}-1}(\mu)}{a_{\frac{n-1}{2}}(\mu) - a_{\frac{n-1}{2}-2}(\mu)} = 0,$$
(16)

$$a_{n-2}(\mu) = -1. \tag{17}$$

(iii) Let μ be an arbitrary root of equation $a_{\frac{n-1}{2}}(\mu) = 0$. Then

$$1 - \frac{a_{\frac{n-1}{2}+1}(\mu) - a_{\frac{n-1}{2}-1}(\mu)}{a_{\frac{n-1}{2}}(\mu) - a_{\frac{n-1}{2}-2}(\mu)} \in (-\infty, 0) \cup (2, +\infty), \qquad (18)$$

$$a_{n-1}(\mu) = -1$$
 and $a_{n-2}(\mu) = -2(1-\mu)$. (19)

The proof of the Lemma is presented in Appendix B.

Lemma 3.8 If $n \ (n \ge 2)$ is an even number, and $\{a_k(\mu)\}_{k\ge 0}$ is a series defined in Lemma 3.6, we have the following results.

Let μ be an arbitrary root of equation $a_{\frac{n}{2}}(\mu) - a_{\frac{n-2}{2}}(\mu) = 0$. Then

$$1 - \frac{a_{\frac{n}{2}}(\mu) - a_{\frac{n}{2}-1}(\mu)}{a_{\frac{n}{2}-1}(\mu) - a_{\frac{n}{2}-2}(\mu)} = 2,$$
(20)

$$a_{n-2}(\mu) = 1.$$
 (21)

((i) Let μ be an arbitrary root of equation $a_{\frac{n}{2}-1}(\mu) = 0$. Then

$$1 - \frac{a_{\frac{n}{2}}(\mu) - a_{\frac{n}{2}-1}(\mu)}{a_{\frac{n}{2}-1}(\mu) - a_{\frac{n}{2}-2}(\mu)} = 0, \qquad (22)$$

$$a_{n-2}(\mu) = -1. (23)$$

(iii) Let μ be an arbitrary root of equation of equation $a_{\frac{n}{2}}(\mu) + a_{\frac{n}{2}-1}(\mu) = 0$. Then

$$1 - \frac{a_{\frac{n}{2}}(\mu) - a_{\frac{n}{2}-1}(\mu)}{a_{\frac{n}{2}-1}(\mu) - a_{\frac{n}{2}-2}(\mu)} \in (-\infty, 0) \cup (2, +\infty), \qquad (24)$$

$$a_{n-1}(\mu) = -1 \quad and \quad a_{n-2}(\mu) = -2(1-\mu).$$
 (25)

The proof of the Lemma is presented in Appendix C.

4 Main Results

For any connected undirected graph G with N_0 vertices, if the normalized Laplacian spectrum $\sigma_G = \{\lambda_1, \lambda_2, \dots, \lambda_{N_0}\}$ is known, we will present the normalized Laplacian spectrum of n-polygon graph $\tau_n(G)$, defined by Definition 2.7 in this section. Firstly, in Lemma 4.1, we present the general relation between the eigenvalues of \mathcal{L}_G and the eigenvalues of $\mathcal{L}_{\tau_n(G)}$, where \mathcal{L}_G and $\mathcal{L}_{\tau_n(G)}$ are the normalized Laplacian matrices for graph G and graph $\tau_n(G)$ respectively. Two Remarks follow Lemma 4.1 to clarify the detailed formulas of the relations in the case n is odd and in the case n is even. Then, in Theorem 4.2 and Theorem 4.3, we present the way to derive the eigenvalues of $\mathcal{L}_{\tau_n(G)}$. By using Theorem 4.2 and Theorem 4.3 recursively, we can also derive the eigenvalues of graph $\tau_n^g(G)$ $(n \ge 2, g \ge 1)$, defined in Definition 2.8, which is the iterated n-polygon graph of G with generation g.

Lemma 4.1 For any integer $n \ (n \geq 2)$, let μ be a real number such that $a_{n-1}(\mu) \neq 0$ and $a_{n-1}(\mu) + 1 \neq 0$, where $\{a_k(\mu)\}$ is a series defined in Lemma 3.6. Then, μ is an eigenvalue of $\mathcal{L}_{\tau_n(G)}$ with multiplicity $k \ (k > 0)$ if and only if $\lambda \equiv 1 - \frac{a_n(\mu)}{1+a_{n-1}(\mu)}$ is an eigenvalue of \mathcal{L}_G with multiplicity k and $\lambda \neq 0$, $\lambda \neq 2$.

The proof of this Lemma is presented in Sec. 6.

Remark 1 If n is an odd number, replacing $a_n(\mu)$ and $1 + a_{n-1}(\mu)$ from Eqs.(12) and (13) respectively, we get

$$\lambda \equiv 1 - \frac{a_n(\mu)}{1 + a_{n-1}(\mu)} = 1 - \frac{\left(a_{\frac{n-1}{2}+1}(\mu) - a_{\frac{n-1}{2}-1}(\mu)\right)a_{\frac{n-1}{2}}(\mu)}{\left(a_{\frac{n-1}{2}}(\mu) - a_{\frac{n-1}{2}-2}(\mu)\right)a_{\frac{n-1}{2}}(\mu)} = 1 - \frac{a_{\frac{n-1}{2}+1}(\mu) - a_{\frac{n-1}{2}-1}(\mu)}{a_{\frac{n-1}{2}}(\mu) - a_{\frac{n-1}{2}-2}(\mu)},$$
(26)

and $1 + a_{n-1}(\mu) \neq 0$ is equivalent to $a_{\frac{n-1}{2}} - a_{\frac{n-1}{2}-2} \neq 0$ and $a_{\frac{n-1}{2}} \neq 0$.

Remark 2 If n is an even number, replacing $a_n(\mu)$ and $1 + a_{n-1}(\mu)$ from Eqs.(12) and (13) respectively, we get

$$\lambda \equiv 1 - \frac{a_n(\mu)}{1 + a_{n-1}(\mu)} = 1 - \frac{\left(a_{\frac{n}{2}}(\mu) - a_{\frac{n}{2}-1}(\mu)\right) \left(a_{\frac{n}{2}}(\mu) + a_{\frac{n}{2}-1}(\mu)\right)}{\left(a_{\frac{n}{2}-1}(\mu) - a_{\frac{n}{2}-2}(\mu)\right) \left(a_{\frac{n}{2}}(\mu) + a_{\frac{n}{2}-1}(\mu)\right)} = 1 - \frac{a_{\frac{n}{2}}(\mu) - a_{\frac{n}{2}-1}(\mu)}{a_{\frac{n}{2}-1}(\mu) - a_{\frac{n}{2}-2}(\mu)},$$
(27)

and $1 + a_{n-1}(\mu) \neq 0$ is equivalent to $a_{\frac{n}{2}-1}(\mu) - a_{\frac{n}{2}-2}(\mu) \neq 0$ and $a_{\frac{n}{2}}(\mu) + a_{\frac{n}{2}-1}(\mu) \neq 0$.

Theorem 4.2 Let G be a connected graph with N_0 vertices and E_0 edges, and $\tau_n(G)$ be the n-polygon graph of G, where $n \geq 3$ and n is odd. The eigenvalues for the normalized Laplacian $\mathcal{L}_{\tau_n(G)}$ can be obtained in the following way.

(i) 0 is the eigenvalue of $\mathcal{L}_{\tau_n(G)}$ with the multiplicity 1. If G is bipartite, 2 is the eigenvalue of $\mathcal{L}_{G(n)}$ with the multiplicity 1.

(ii)Let μ be an arbitrary root of equation $a_{\frac{n-1}{2}}(\mu) = 0$, then μ is a eigenvalue of $\mathcal{L}_{\tau_n(G)}$ with multiplicity N_0 .

(iii)Let μ be an arbitrary root of equation $a_{\frac{n-1}{2}}(\mu) + a_{\frac{n-1}{2}-1}(\mu) = 0$, then μ is a eigenvalue of $\mathcal{L}_{\tau_n(G)}$ with multiplicity $E_0 - N_0 + 1$.

(iv)In the case G is non-bipartite, let μ be an arbitrary root of equation $a_{\frac{n-1}{2}}(\mu) - a_{\frac{n-1}{2}-1}(\mu) = 0$, then μ is a eigenvalue of $\mathcal{L}_{\tau_n(G)}$ with multiplicity $E_0 - N_0$.

(v) In the case G is bipartite, let μ be an arbitrary root of equation $a_{\frac{n-1}{2}}(\mu) - a_{\frac{n-1}{2}-1}(\mu) = 0$, then μ is a eigenvalue of $\mathcal{L}_{\tau_n(G)}$ with multiplicity $E_0 - N_0 + 1$. (vi)Let λ be an arbitrary eigenvalue of \mathcal{L}_G such that $\lambda \neq 0$ and $\lambda \neq 2$, $\mu_i(\lambda)$, $(i = 1, 2, \dots, \frac{n+1}{2})$ be the roots of equation

$$1 - \frac{a_{\frac{n-1}{2}+1}(x) - a_{\frac{n-1}{2}-1}(x)}{a_{\frac{n-1}{2}}(x) - a_{\frac{n-1}{2}-2}(x)} = \lambda.$$
(28)

Then, $\mu_i(\lambda)$, $(i = 1, 2, \cdots, \frac{n+1}{2})$ are eigenvalues of $\mathcal{L}_{\tau_n(G)}$ with $m_{\mathcal{L}_G}(\lambda) = m_{\mathcal{L}_{\tau_n(G)}}(\mu_i(\lambda)), i = 1, 2, \cdots, \frac{n+1}{2}$.

The proof of Theorem 4.2 is presented in Sec. 7.

Theorem 4.3 Let G be a connected graph with N_0 vertices and E_0 edges, and $\tau_n(G)$ be the n-polygon graph of G, where $n \ge 2$ and n is even. The eigenvalues for the normalized Laplacian $\mathcal{L}_{\tau_n(G)}$ can be obtained in the following way.

(i) 0 is the eigenvalue of $\mathcal{L}_{\tau_n(G)}$ with the multiplicity 1.

(ii)Let μ be an arbitrary root of equation $a_{\frac{n}{2}}(\mu) + a_{\frac{n}{2}-1}(\mu) = 0$, then μ is eigenvalue of $\mathcal{L}_{\tau_n(G)}$ with the multiplicity N_0 .

(iii)Let μ be an arbitrary root of equation $a_{\frac{n}{2}-1}(\mu) = 0$, then μ is a eigenvalue

of $\mathcal{L}_{\tau_n(G)}$ with multiplicity $E_0 - N_0 + 1$. (iv)In the case G is non-bipartite, let μ be an arbitrary root of equation $a_{\frac{n}{2}}(\mu) - a_{\frac{n}{2}-1}(\mu) = 0$, then μ is a eigenvalue of $\mathcal{L}_{\tau_n(G)}$ with multiplicity $E_0 - N_0$. (v)In the case G is bipartite, let μ be an arbitrary root of equation $a_{\frac{n}{2}}(\mu) - a_{\frac{n}{2}-1}(\mu) = 0$, then μ is a eigenvalue of $\mathcal{L}_{\tau_n(G)}$ with multiplicity $E_0 - N_0 + 1$. (vi)Let λ be an arbitrary eigenvalue of \mathcal{L}_G such that $\lambda \neq 0$ and $\lambda \neq 2$, $\mu_i(\lambda)$, ($i = 1, 2, \cdots, \frac{n}{2}$) be the roots of equation

$$1 - \frac{a_{\frac{n}{2}}(x) - a_{\frac{n}{2}-1}(x)}{a_{\frac{n}{2}-1}(x) - a_{\frac{n}{2}-2}(x)} = \lambda.$$
(29)

Then, $\mu_i(\lambda)$, $(i = 1, 2, \cdots, \frac{n}{2})$ are eigenvalues of $\mathcal{L}_{\tau_n(G)}$ with $m_{\mathcal{L}_G}(\lambda) = m_{\mathcal{L}_{\tau_n(G)}}(\mu_i(\lambda)), i = 1, 2, \cdots, \frac{n}{2}$.

The proof of the Theorem is presented in Sec. 8.

Remark 3 By using Theorem 4.2 and Theorem 4.3, we can obtain the complete normalized Laplacian spectrum $\sigma(\tau_n(G))$ of n-polygon graph $\tau_n(G)$. We can also derive the complete normalized Laplacian spectrum of graph $\tau_n^g(G)$ $(n \ge 2, g \ge 1)$, defined in Definition 2.8, which is the iterated n-polygon graph of G with generation g, by using Theorem 4.2 and Theorem 4.3 recursively. In fact, let n = 2 and n = 4 in Theorem 4.3 respectively, we can recover the results obtained in Ref. [25] and Ref. [31] respectively; let n = 3 in Theorem 4.2, we can recover the results obtained in Ref. [30].

5 Applications

Recalling that the multiplicative degree-Kirchhoff index, the Kemeny's const and the number of spanning trees can be expressed as functions of the normalized Laplacian spectrum, as shown in Lemma 3.2, we obtain the following Theorems.

Theorem 5.1 Let G be a connected undirected graph with N_0 vertices and E_0 edges, $\tau_n(G)$ $(n \ge 2)$ be the n-polygon graph of G and $\tau_n^g(G)$ $(n \ge 2, g \ge 1)$, defined in Definition 2.8, be the iterated n-polygon graph of G with generation g, and Kf'(G), Kf'($\tau_n(G)$), Kf'($\tau_n^g(G)$) represent the multiplicative degree-Kirchhoff indexes of graph G, $\tau_n(G)$, $\tau_n^g(G)$ respectively. Then, for any $n \ge 2$,

$$Kf'(\tau_n(G)) = (n^2 + n) Kf'(G) + \frac{2}{3}(n+1)(n^2 - 1) E_0^2 - \frac{2}{3}(n^2 - 1) E_0 N_0 - \frac{1}{6}(n^2 - 1)(n-2) E_0,$$
(30)

and for $n \geq 2$ and $g \geq 1$,

$$Kf'(\tau_n^g(G)) = \left(n^2 + n\right)^g Kf'(G) - \frac{1}{3}(n-2)(n+1)^g(n^g-1)E_0 + \frac{2(n-1)}{3n}(n+1)^g\left[(n+1)^g(n^2+1) - n^{g+2} - 1\right]E_0^2 - \frac{2}{3}(n+1)^g(n^g-1)E_0N_0.$$
(31)

The proof of Theorem 5.1 is presented in Sec. 9.

Theorem 5.2 Let G be a connected undirected graph with N_0 vertices and E_0 edges, and K(G), $K(\tau_n(G))$, $K(\tau_n^g(G))$ represent the Kemeny's constant of graph G, $\tau_n(G)$, $\tau_n^g(G)$ respectively. Then, for any $n \ge 2$,

$$K(\tau_n(G)) = nK(G) + \frac{1}{3}(n^2 - 1)E_0 - \frac{1}{3}(n - 1)N_0 - \frac{1}{12}(n - 1)(n - 2),$$

and for $n \geq 2$ and $g \geq 1$,

$$K\left(\tau_{n}^{g}(G)\right) = n^{g}K\left(G\right) - \frac{1}{3}\left(n^{g} - 1\right)N_{0} - \frac{1}{12}\left(n - 2\right)\left(n^{g} - 1\right) + \left[\frac{(n-1)\left(n^{2} + 1\right)}{3n}\left(n + 1\right)^{g} + \frac{1}{3}\left(-n^{3} + n^{2} + 1\right)n^{g-1} - \frac{1}{3}\right]E_{0}.$$

This Theorem can be obtained directly from Theorem 5.1 and Lemma 3.2.

Theorem 5.3 Let G be a connected undirected graph with N_0 vertices and E_0 edges and $N_{st}(G)$, $N_{st}(\tau_n(G))$, $N_{st}(\tau_n^g(G))$ represent the number of spanning trees of graph G, $\tau_n(G)$, $\tau_n^g(G)$ respectively. Then, for $n \geq 2$,

$$N_{st}(\tau_n(G)) = (n+1)^{N_0-1} n^{E_0-N_0+1} \cdot N_{st}(G), \qquad (32)$$

and for $n \geq 2$ and $g \geq 1$,

$$N_{st}\left(\tau_{n}^{g}(G)\right) = (n+1)^{\frac{(n-1)\left[(n+1)^{g}-ng-1\right]}{n^{2}}E_{0}+gN_{0}-g} \cdot n^{\frac{(n+1)^{g}+(n-1)ng-1}{n^{2}}E_{0}-gN_{0}+g} \cdot N_{st}\left(G\right).$$
(33)

The proof of Theorem 5.3 is presented in Sec. 10.

6 The proof of Lemma 4.1

First, we prove the "only if" part of this Lemma. For an arbitrary real number μ , if $a_{n-1}(\mu) \neq 0$ and $a_{n-1}(\mu) + 1 \neq 0$ and μ is an eigenvalue of $\mathcal{L}_{\tau_n(G)}$ with multiplicity k (k > 0), we will show that $\lambda \equiv 1 - \frac{a_n(\mu)}{1 + a_{n-1}(\mu)}$ is an eigenvalue of \mathcal{L}_G with multiplicity k.

Let $\vec{v} = (v_1, v_2, \cdots, v_{N_1})^T$ be an eigenvector with respect to the eigenvalue μ of $\tau_n(G)$. Thus

$$\mathcal{L}_{\tau_n(G)}\vec{v} = \mu\vec{v}.\tag{34}$$

Replacing every entries of $\mathcal{L}_{\tau_n(G)}$ from Eq. (3), we find, for any vertex q of graph $\tau_n(G)$, the corresponding entry v_q in \vec{v} satisfies

$$(1-\mu) v_q = \sum_{t=1}^{N_1} \frac{A_{\tau_n(G)}(q,t)}{\sqrt{d'_q d'_t}} v_t,$$
(35)

where $A_{\tau_n(G)}(q,t)$ is the (q,t) entry for the adjacency matrix of $\tau_n(G)$, d'_q and d'_t are the degree for vertexes q and t in graph $\tau_n(G)$ respectively.

Let V(G) and $V(\tau_n(G))$ be the vertexes set of graph G and $\tau_n(G)$ respectively and $V_N = \{v : v \in V(\tau_n(G)), v \notin V(G)\}$. We have $V(G) \subset V(\tau_n(G))$ and $V(\tau_n(G)) = V_N \cup V(G)$. We will show the relation between the corresponding entries of \vec{v} for nodes in V(G) and those for nodes in V_N .

For an arbitrary edge of graph G, let i and j be the two ends of the edge. Recalling the construction of the graph $\tau_n(G)$, as shown in Fig. 1, the edge is replacing with a (n + 1)-polygon, whose nodes are labeled as i, i'_1 , i'_2 , \cdots , i'_{n-2} , i'_{n-1} , j.

Let $N_0^i \subseteq V(G)$ denote the set for neighbors of vertex *i* in graph *G* and $N_N^i \subseteq V_N$ denote the set of the new neighbors of vertex *i* in graph $\tau_n(G)$. Note that $d'_i = 2d_i$ and $d'_j = 2d_j$. We can rewritte Eq. (35) as

$$(1-\mu) v_{i} = \sum_{i_{1}' \in N_{N}^{i}} \frac{1}{\sqrt{d_{i}'d_{i_{1}'}'}} v_{i_{1}'} + \sum_{j \in N_{O}^{i}} \frac{1}{\sqrt{d_{i}'d_{j}'}} v_{j}$$
$$= \sum_{i_{1}' \in N_{N}^{i}} \frac{1}{2\sqrt{d_{i}}} v_{i_{1}'} + \sum_{j \in N_{O}^{i}} \frac{1}{2\sqrt{d_{i}d_{j}}} v_{j}.$$
(36)

Similarly, for any vertex $i'_1 \in N_N^i$,

$$(1-\mu) v_{i_1'} = \frac{1}{\sqrt{d_{i_1'}' d_{i_2'}'}} v_{i_2'} + \frac{1}{\sqrt{d_{i_1'}' d_i'}} v_i$$
$$= \frac{1}{2} v_{i_2'} + \frac{1}{2\sqrt{d_i}} v_i,$$
(37)

for the vertex $i'_{n-1} \in N_N^j$,

$$(1-\mu)v_{i'_{n-1}} = \frac{1}{2}v_{i'_{n-2}} + \frac{1}{2\sqrt{d_j}}v_j, \tag{38}$$

and for the vertex $i_k^{'}$ $(2 \le k \le n-2)$,

$$(1-\mu) v_{i'_{k}} = \frac{1}{2} \left(v_{i'_{k-1}} + v_{i'_{k+1}} \right).$$
(39)

Eliminating the variables $v_{i'_2}, v_{i'_3}, \cdots, v_{i'_{n-2}}$ in Eqs. (37) and (39), we have

$$a_{n-2}(\mu)v_{i_1'} = v_{i_{n-1}'} + a_{n-3}(\mu)\frac{v_i}{\sqrt{d_i}},\tag{40}$$

where $\{a_n(\mu)\}\$ is just the series defined in Lemma 3.6.

Similarly, eliminating the variables $v_{i'_2}, v_{i'_3}, \cdots, v_{i'_{n-1}}$ in Eqs. (37), (38) and (39), we get

$$a_{n-1}(\mu)v_{i_1'} = a_{n-2}(\mu)\frac{1}{\sqrt{d_i}}v_i + \frac{1}{\sqrt{d_j}}v_j.$$
(41)

Therefore, in the case $a_{n-1}(\mu) \neq 0$, multiplying both sides of the Eq. (36) by a factor $a_{n-1}(\mu)$, and replacing $a_{n-1}(\mu)v_{i'_1}$ from Eq.(41), we have

$$2(1-\mu)a_{n-1}(\mu)v_i = a_{n-2}(\mu)v_i + \sum_{j \in N_O^i} \frac{1+a_{n-1}(\mu)}{\sqrt{d_i d_j}}v_j, \qquad (42)$$

which is equivalent to

$$a_n(\mu)v_i = \sum_{j \in N_O^i} \frac{1 + a_{n-1}(\mu)}{\sqrt{d_i d_j}} v_j.$$
(43)

In the case $a_{n-1}(\mu) \neq 0$ and $1 + a_{n-1}(\mu) \neq 0$, Eq. (43) is also equal to

$$\frac{a_n(\mu)}{1+a_{n-1}(\mu)}v_i = \sum_{j \in N_O^i} \frac{1}{\sqrt{d_i d_j}}v_j = \sum_{j \in V(G)} \frac{A_{ij}}{\sqrt{d_i d_j}}v_j,$$
(44)

which implies that $\frac{a_n(\mu)}{1+a_{n-1}(\mu)}$ is just an eigenvalue of the matrix $D(G)^{-\frac{1}{2}}A(G)D(G)^{-\frac{1}{2}}$. Therefore, in the case $a_{n-1}(\mu) \neq 0$ and $1 + a_{n-1}(\mu) \neq 0$, $\lambda \equiv 1 - \frac{a_n(\mu)}{1+a_{n-1}(\mu)}$ is an eigenvalue of the matrix $\mathcal{L}_G = I - D(G)^{-\frac{1}{2}}A(G)D(G)^{-\frac{1}{2}}$. Further more, Eqs. (37), (38) and (39) also show that the entries $v_{i'_k}$ $(k = 1, 2, \dots, n-1)$ of \vec{v} are completely decided by entries v_i and v_j . This is to say, the dimension of solution space for linear equation $\mathcal{L}_{\tau_n(G)}\vec{v} = \mu\vec{v}$ is the same as that for linear equation $\mathcal{L}_G\vec{v'} = \lambda\vec{v'}$. Therefore, $\lambda \equiv 1 - \frac{a_n(\mu)}{1+a_{n-1}(\mu)}$ is an eigenvalue of the matrix \mathcal{L}_G with the same multiplicity as the eigenvalue μ of $\mathcal{L}_{\tau_n(G)}$.

As shown in Lemma 3.7 and Lemma 3.8, if $a_{n-1}(\mu) \neq 0$ and $1 + a_{n-1}(\mu) \neq 0$, $\lambda \equiv 1 - \frac{a_n(\mu)}{1 + a_{n-1}(\mu)} \neq 0$ and $\lambda \equiv 1 - \frac{a_n(\mu)}{1 + a_{n-1}(\mu)} \neq 2$.

Therefore, we obtain the "only if" part of this Lemma. Then we prove the "if" part of this Lemma.

Let λ be an arbitrary eigenvalue of \mathcal{L}_G with multiplicity k (k > 0), and $\vec{v'} = (v_1, v_2, \cdots, v_{N_0})^T$ be an eigenvector with respect to the eigenvalue λ of \mathcal{L}_G , i.e.,

$$\mathcal{L}_G \vec{v'} = \lambda \vec{v'},\tag{45}$$

which can be rewritten as

$$(1 - \lambda) v_i = \sum_{j \in V(G)} \frac{A_{ij}}{\sqrt{d_i d_j}} v_j = \sum_{j \in N_O^i} \frac{1}{\sqrt{d_i d_j}} v_j,$$
(46)

for any node $i \in V(G)$. Therefore for any μ which satisfies $\lambda = 1 - \frac{a_n(\mu)}{1 + a_{n-1}(\mu)}$, Eq. (44) holds and $1 + a_{n-1}(\mu) \neq 0$.

Note that $\lambda \neq 0$ and $\lambda \neq 2$. As proved in Lemma 3.7 and Lemma 3.8, we get $a_{n-1}(\mu) \neq 0$, and for any node $i \in V(G)$, Eq. (42) holds.

For any edge with ends i and j, let v_i and v_j be the corresponding two entries in $\vec{v'}$ and let

$$v_{i_1'} = \frac{a_{n-2}(\mu)}{a_{n-1}(\mu)} \frac{1}{\sqrt{d_i}} v_i + \frac{1}{a_{n-1}(\mu)\sqrt{d_j}} v_j, \tag{47}$$

$$v_{i_2'} = 2\left(1-\mu\right)v_{i_1'} - \frac{1}{\sqrt{d_i}}v_i,\tag{48}$$

and for any $k \ (2 \le k \le n-2)$,

$$v_{i'_{k+1}} = 2\left(1-\mu\right)v_{i'_{k}} - v_{i'_{k-1}}.$$
(49)

In this way, we can obtain a N_1 -dimensional vector \vec{v} , whose entries satisfy Eqs. (36)-(39). Then, for any node $q \in V(\tau_n(G))$, the entry v_q in \vec{v} satisfies Eq. (35).

Therefore, the vector \vec{v} satisfies Eq. (34) and μ is an eigenvalue of $\mathcal{L}_{\tau_n(G)}$. Note that the corresponding eigenvectors \vec{v} of μ is in one-to-one correspondence with

eigenvectors $\vec{v'}$ of λ . Thus the eigenvalue μ of $\mathcal{L}_{\tau_n(G)}$ has the same multiplicity as eigenvalue λ of \mathcal{L}_G .

This ends the proof.

7 Proof of Theorem 4.2

(i)It is obvious from Lemma 3.1.

(ii) For any root μ of equation $a_{\frac{n-1}{2}}(\mu) = 0$, we can obtain from Lemma 3.7 (see Eqs.(19)) that

$$a_{n-1}(\mu) = -1$$
, and $a_{n-2}(\mu) = -2(1-\mu)$.

Inserting the two equations into Eq.(41), we have

$$v_{i_1'} = \frac{2(1-\mu)}{\sqrt{d_i}} v_i - \frac{1}{\sqrt{d_j}} v_j.$$
(50)

Therefore, for any node $i \in V(G)$,

$$\sum_{i_{1}' \in N_{N}^{i}} \frac{1}{\sqrt{d_{i}'d_{i_{1}'}'}} v_{i_{1}'} + \sum_{j \in V(G)} \frac{1}{\sqrt{d_{i}'d_{j}'}} v_{j}$$

$$= \sum_{i_{1}' \in N_{N}^{i}} \frac{1}{2\sqrt{d_{i}}} \left[\frac{2(1-\mu)}{\sqrt{d_{i}}} v_{i} - \frac{1}{\sqrt{d_{j}}} v_{j} \right] + \sum_{j \in N_{O}^{i}} \frac{1}{2\sqrt{d_{i}d_{j}}} v_{j}$$

$$= \sum_{j \in N_{O}^{i}} \left[-\frac{1}{2\sqrt{d_{i}d_{j}}} v_{j} + \frac{2(1-\mu)}{2d_{i}} v_{i} \right] + \sum_{j \in N_{O}^{i}} \frac{1}{2\sqrt{d_{i}d_{j}}} v_{j}$$

$$= \sum_{j \in N_{O}^{i}} \frac{(1-\mu)}{d_{i}} v_{i}$$

$$= (1-\mu)v_{i}, \qquad (51)$$

where N_N^i and N_O^i , which represent the neighbors of node *i* in V_N and V(G) respectively, are defined in Lemma 4.1. Thus, Eq.(36) holds no matter what v_i is.

Let $\vec{v'} = (v_1, v_2, \dots, v_{N_0})^T$ be an arbitrary vector. Eq. (51) inform us that Eq.(36) holds if μ is a root of equation $a_{\frac{n-1}{2}}(\mu) = 0$. For an arbitrary edge of graph G, let nodes i and j be the two ends of the edge. Considering the (n + 1)-polygon as shown in the right-hand side of Fig. 1, and calculating

 $v_{i'_1}, v_{i'_2}, \dots, v_{i'_{n-1}}$ by using Eqs.(47), (48), (49). In this way, we obtain a N_1 dimensional vector \vec{v} , whose entries satisfy Eqs. (36)-(39). Then, for any node $q \in V(\tau_n(G))$, the entry v_q in \vec{v} satisfies Eq. (35).

Therefore, the vector \vec{v} satisfies Eq. (34) and μ is an eigenvalue of $\mathcal{L}_{\tau_n(G)}$. Further more, there are strict one-to-one correspondences be the eigenvectors \vec{v} of μ and $\vec{v'}$, which is an arbitrary vector in N_0 -dimensional space. Thus the multiplicity of the eigenvalue μ is N_0 .

(iii) Let μ be an arbitrary root of equation $a_{\frac{n-1}{2}}(\mu) + a_{\frac{n-1}{2}-1}(\mu) = 0$. Then

$$a_{n-1}(\mu) = \left[a_{\frac{n-1}{2}}(\mu) - a_{\frac{n-1}{2}-1}(\mu)\right] \left[a_{\frac{n-1}{2}}(\mu) + a_{\frac{n-1}{2}-1}(\mu)\right] = 0,$$

and we can obtain from Lemma 3.7 (see Eq. (17)) that

$$a_{n-2}\left(\mu\right) = -1.$$

Therefore,

$$a_{n-3}(\mu) = 2(1-\mu)a_{n-2}(\mu) - a_{n-1}(\mu) = -2(1-\mu)$$

Replacing a_{n-1} and a_{n-2} with 0 and -1 in Eq.(41) respectively, for any two nodes i and j of graph G, if $i \sim j$,

$$\frac{v_i}{\sqrt{d_i}} = \frac{v_j}{\sqrt{d_j}}.$$
(52)

Since G is a connected graph, Eq. (52) shows that $\frac{v_i}{\sqrt{d_i}}$ is a const for any $i \in V(G)$. Let $\theta \equiv \frac{v_i}{\sqrt{d_i}}$, for any $i \in V(G)$. Eqs. (40) and (36) can be rewritten as

$$v_{i'_1} + v_{i'_{n-1}} = 2 (1 - \mu) \theta, \quad i'_1 \in N_N^i, i'_{n-1} \in N_N^j.$$
(53)

$$\sum_{i'_{1} \in N^{i}_{M}} v_{i'_{1}} = [2(1-\mu) - 1] \theta d_{i}, \quad i \in V(G).$$
(54)

Calculating $\sum_{i \in V(G)} \sum_{i'_1 \in N_N^i} v_{i'_1}$ by using Eq.(53), we have

$$\sum_{i \in V(G)} \sum_{i_1' \in N_N^i} v_{i_1'} = \frac{1}{2} \sum_{i \sim j, i_1' \in N_N^i, i_{n-1}' \in N_N^j} \left(v_{i_1'} + v_{i_{n-1}'} \right) = 2 \left(1 - \mu \right) \theta E_0.$$
(55)

On the other hand, Calculating $\sum_{i \in V(G)} \sum_{i'_1 \in N_N^i} v_{i'_1}$ by using Eq.(54), we get

$$\sum_{i \in V(G)} \sum_{i_1' \in N_N^i} v_{i_1'} = [2(1-\mu) - 1] \theta \sum_{i \in V(G)} d_i = [4(1-\mu) - 2] \theta E_0.$$
(56)

Therefore, $[4(1-\mu)-2]\theta = 2(1-\mu)\theta$, which leads to $\mu = 0$ or $\theta = 0$. If $\mu = 0$, replacing $a_{\frac{n-1}{2}}(\mu)$ and $a_{\frac{n-1}{2}-1}(\mu)$ from Eq.(7), we have

$$a_{\frac{n-1}{2}}(\mu) + a_{\frac{n-1}{2}-1}(\mu) = \frac{n-1}{2} + 1 + \frac{n-1}{2} - 1 + 1 = n \neq 0.$$

Thus, $\mu \neq 0$, and $\theta = 0$, which leads to

$$v_i = 0, \tag{57}$$

for any node i of graph G.

Substituting v_i with 0, Eq.(36) can be rewritten as

$$\sum_{i_1' \in N_N^i} v_{i_1'} = 0, \tag{58}$$

for any node $i \in V(G)$.

Similarly, replacing v_i and $a_{n-2}(\mu)$ with 0 and -1 respectively in Eq. (40), we get

$$v_{i_1'} + v_{i_{n-1}'} = 0. (59)$$

Further more, if n is odd and $v_i = 0$, in any (n+1)-polygon of $\tau_n(G)$ as shown in Fig. 1, we find, Eqs. (37), (38) and (39) are equivalent to

$$v_{i_{t+1}} + v_{i_{n-t-1}} = 0, \quad t = 0, \ 1, \ 2, \ \cdots, \ \frac{n-1}{2} - 1,$$
 (60)

$$a_t(\mu_i)v_{i'_1} = v_{i'_{t+1}}, \quad t = 1, 2, \cdots, \frac{n-1}{2} - 1.$$
 (61)

Therefore, Eq. (35) holds if and only if Eqs. (57), (58), (60) and (82) hold for any (n + 1)-polygon of graph $\tau_n(G)$.

Note that the total number of (n + 1)-polygons in graph $\tau_n(G)$ is E_0 . For an arbitrary (n + 1)-polygon, P_r , of graph $\tau_n(G)$, let $v_{i'_1} = -v_{i'_{n-1}} = x_r$, and let $X = (x_1, x_2, \dots, x_{E_0})^T$. We find Eqs.(58) and (59) are equivalent to B'X = 0, where B' is the incident matrix of weakly connected directed graph G' (see Definition 2.5 and 2.6).

Similarly, for an arbitrary (n + 1)-polygon, P_r , of graph $\tau_n(G)$, let

$$v_{i'_{t+1}} = -v_{i'_{n-t-1}} = y_r^{(t)}, \ t = 1, \ 2, \ \dots, \ \frac{n-1}{2} - 1,$$

and let $Y^{(t)} = (y_1^{(t)}, y_2^{(t)}, \dots, y_{E_0}^{(t)})^T$, we find, Eqs. (60) and (82) can be rewritten as

$$Y^{(t)} - a_t(\mu)X = 0, \quad t = 1, 2, \cdots, \frac{n-1}{2} - 1.$$

Therefore, there are strict one-to-one correspondences between the roots of Eq. (34) and roots of the equation

$$\begin{pmatrix} I \ 0 \ \cdots \ 0 \ -a_1(\mu) \ I \\ 0 \ I \ \cdots \ 0 \ -a_2(\mu) \ I \\ \vdots \ \vdots \ \ddots \ \vdots \ \vdots \\ 0 \ 0 \ \cdots \ I \ -a_{\frac{n-1}{2}-1}(\mu) \ I \\ 0 \ 0 \ \cdots \ 0 \ B' \end{pmatrix} \begin{pmatrix} Y^{(1)} \\ Y^{(2)} \\ \vdots \\ Y^{\binom{n-1}{2}-1} \\ X \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix},$$
(62)

where I is an identity matrix of order E_0 and B' is the incidence matrix of weakly connected directed graph G' whose underlying undirected graph (all edges replaced by undirected edges) is G.

Because $rank(B') = N_0 - 1$ (see Lemma 3.5). Therefore the rank for the coefficient matrix of linear equation (62) is $\frac{n-3}{2}E_0 + N_0$ and the dimension for the vector space spanned by the roots of linear equation (62) is $E_0 - N_0 + 1$. Thus the dimension for the eigenspace of μ is also $E_0 - N_0 + 1$ and the multiplicity for eigenvalue μ is $E_0 - N_0 + 1$.

(iv) Let μ be an arbitrary root of equation $a_{\frac{n-1}{2}}(\mu) - a_{\frac{n-1}{2}-1}(\mu) = 0$. Then

$$a_{n-1}(\mu) = \left[a_{\frac{n-1}{2}}(\mu) - a_{\frac{n-1}{2}-1}(\mu)\right] \left[a_{\frac{n-1}{2}}(\mu) + a_{\frac{n-1}{2}-1}(\mu)\right] = 0,$$

and we can obtain from Lemma 3.7 (see Eq. (B.5)) that

$$a_{n-2}\left(\mu\right) = 1.$$

Replacing a_{n-1} and a_{n-1} with 0 and 1 in Eq.(41), for any two nodes *i* and *j* of graph *G*, if $i \sim j$,

$$\frac{v_i}{\sqrt{d_i}} = -\frac{v_j}{\sqrt{d_j}}.$$
(63)

Note that G is non-bipartite, there is at least an odd cycle C in G [8,34]. Let i_1, i_2, \dots, i_k (k is odd) be the nodes series of cycle C. Eq. (63) informs us that

$$\frac{v_{i_1}}{\sqrt{d_{i_1}}} = -\frac{v_{i_2}}{\sqrt{d_{i_2}}} = \frac{v_{i_3}}{\sqrt{d_{i_3}}} = \dots = \frac{v_{i_k}}{\sqrt{d_{i_k}}} = -\frac{v_{i_1}}{\sqrt{d_{i_1}}}.$$

Therefore $v_{i_1} = v_{i_2} = \cdots = v_{i_k} = 0$. Since G is connected, Eq. (63) informs us,

$$v_i = 0. (64)$$

for any node i of graph G.

Substituting v_i with 0, Eq.(36) can be rewritten as

$$\sum_{i_1' \in N_N^i} v_{i_1'} = 0, \tag{65}$$

for any node i of graph G.

Similarly, replacing v_i and $a_{n-2}(\mu)$ with 0 and 1 respectively in Eq. (40), we get

$$v_{i_1'} - v_{i_{n-1}'} = 0. ag{66}$$

Further more, if n is odd and $v_i = 0$ for any node $i \in V(G)$, we find, Eqs. (37), (38) and (39) are equivalent to

$$v_{i'_{t+1}} - v_{i'_{n-t-1}} = 0, \quad t = 0, \ 1, \ 2, \ \cdots, \ \frac{n-1}{2} - 1,$$
 (67)

$$a_t(\mu_i)v_{i_1'} = v_{i_{t+1}'}, \quad t = 1, 2, \cdots, \frac{n-1}{2} - 1.$$
 (68)

Therefore, Eq. (35) holds if and only if Eqs. (64), (65), (67) and (89) hold for any (n + 1)-polygon of graph $\tau_n(G)$.

Note that the total number of (n + 1)-polygons in graph $\tau_n(G)$ is E_0 . For an arbitrary (n + 1)-polygon, P_r , of graph $\tau_n(G)$, let $v_{i'_1} = v_{i'_{n-1}} = x_r$, and let $X = (x_1, x_2, \dots, x_{E_0})^T$. We find Eqs.(65) and (66) are equivalent to BX = 0, where B is the incident matrix of graph G.

Similarly, for an arbitrary (n + 1)-polygon, P_r , of graph $\tau_n(G)$, let

$$v_{i'_{t+1}} = v_{i'_{n-t-1}} = y_r^{(t)}, \ t = 1, \ 2, \ \dots, \ \frac{n-1}{2} - 1,$$

and let $Y^{(t)} = (y_1^{(t)}, y_2^{(t)}, \dots, y_{E_0}^{(t)})^T$, we find, Eqs. (67) and (89) can be rewritten as

$$Y^{(t)} - a_t(\mu)X = 0, \quad t = 1, \ 2, \ \cdots, \ \frac{n-1}{2} - 1.$$

Therefore, there are strict one-to-one correspondences between the roots of

Eq. (34) and roots of the equation

$$\begin{pmatrix} I \ 0 \ \cdots \ 0 \ -a_1(\mu) \ I \\ 0 \ I \ \cdots \ 0 \ -a_2(\mu) \ I \\ \vdots \ \vdots \ \ddots \ \vdots \ \vdots \\ 0 \ 0 \ \cdots \ I \ -a_{\frac{n-1}{2}-1}(\mu) \ I \\ 0 \ 0 \ \cdots \ 0 \ B \end{pmatrix} \begin{pmatrix} Y^{(1)} \\ Y^{(2)} \\ \vdots \\ Y^{\binom{n-1}{2}-1} \\ X \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix},$$
(69)

where B is the incident matrix of G and I is identity matrix of order E_0 .

Because G is non-bipartite, $rank(B) = N_0$ (see Lemma 3.4). Therefore the rank for the coefficient matrix of linear equation (69) is $\frac{n-3}{2}E_0 + N_0$ and the dimension for the vector space spanned by the roots of linear equation (69) is $E_0 - N_0$. Thus the dimension for the eigenspace of μ is also $E_0 - N_0$ and the multiplicity for eigenvalue μ is $E_0 - N_0$.

(v) Let μ be an arbitrary root of equation $a_{\frac{n-1}{2}}(\mu) - a_{\frac{n-1}{2}-1}(\mu) = 0$. That is to say

$$a_{\frac{n-1}{2}}(\mu) = a_{\frac{n-1}{2}-1}(\mu).$$

As shown in Lemma 3.6, for any $n \ge 1$, $a_n(2-\mu) = (-1)^n a_n(\mu)$. Therefore,

$$a_{\frac{n-1}{2}}(2-\mu) = (-1)^{\frac{n-1}{2}} a_{\frac{n-1}{2}}(\mu)$$

= $(-1)^{\frac{n-1}{2}} a_{\frac{n-1}{2}-1}(\mu)$
= $(-1)^{\frac{n-1}{2}} \times (-1)^{\frac{n-1}{2}-1} a_{\frac{n-1}{2}-1}(2-\mu)$
= $(-1)^{n-2} a_{\frac{n-1}{2}-1}(2-\mu)$
= $(-1)a_{\frac{n-1}{2}-1}(2-\mu).$ (70)

Therefore, $2 - \mu$ is a root of equation $a_{\underline{n-1}}(\mu) + a_{\underline{n-1}}(\mu) = 0$.

Therefore $2 - \mu$ is a eigenvalue $\tau_n(G)$ with multiplicity $E_0 - N_0 + 1$ (see (iii) of this Theorem).

Note that G is bipartite, then $\tau_n(G)$ is also bipartite. We can obtain from Lemma 3.6 that μ is also a eigenvalue $\tau_n(G)$ with multiplicity $E_0 - N_0 + 1$.

(vi) It is just the result of Lemma 4.1.

The proof is completed.

8 Proof of Theorem 4.3

This theorem can be proved in the same way as Theorem 4.2.

(i) It is obvious from Lemma 3.1.

(ii) Similar to the proof of Theorem4.2(ii). For any root μ of equation $a_{\frac{n}{2}}(\mu) + a_{\frac{n}{2}-1}(\mu) = 0$, we can obtain from Lemma 3.8 (see Eq.(25)) that

$$a_{n-1}(\mu) - 1$$
, and $a_{n-2}(\mu) = -2(1-\mu)$.

Inserting the two equations into Eq.(41), we have

$$v_{i_1'} = \frac{2(1-\mu)}{\sqrt{d_i}} v_i - \frac{1}{\sqrt{d_j}} v_j.$$
(71)

Therefore, for any node $i \in V(G)$,

$$\sum_{i_{1}' \in N_{N}^{i}} \frac{1}{\sqrt{d_{i}'d_{i_{1}'}'}} v_{i_{1}'} + \sum_{j \in V(G)} \frac{1}{\sqrt{d_{i}'d_{j}'}} v_{j}$$

$$= \sum_{i_{1}' \in N_{N}^{i}} \frac{1}{2\sqrt{d_{i}}} \left[\frac{2(1-\mu)}{\sqrt{d_{i}}} v_{i} - \frac{1}{\sqrt{d_{j}}} v_{j} \right] + \sum_{j \in N_{O}^{i}} \frac{1}{2\sqrt{d_{i}d_{j}}} v_{j}$$

$$= \sum_{j \in N_{O}^{i}} \left[-\frac{1}{2\sqrt{d_{i}d_{j}}} v_{j} + \frac{2(1-\mu)}{2d_{i}} v_{i} \right] + \sum_{j \in N_{O}^{i}} \frac{1}{2\sqrt{d_{i}d_{j}}} v_{j}$$

$$= \sum_{j \in N_{O}^{i}} \frac{(1-\mu)}{d_{i}} v_{i}$$

$$= (1-\mu)v_{i}, \qquad (72)$$

where N_N^i and N_O^i , which represent the neighbors of node *i* in V_N and V(G) respectively, are defined in Lemma 4.1. Thus, Eq.(36) holds no matter what v_i is.

Let $\vec{v'} = (v_1, v_2, \dots, v_{N_0})^T$ be an arbitrary vector. Eq. (72) inform us that Eq.(36) holds if μ is a root of equation $a_{\frac{n-1}{2}}(\mu) = 0$. For an arbitrary edge of graph G, let nodes i and j be the two ends of the edge. Considering the (n + 1)-polygon as shown in the right-hand side of Fig. 1, and calculating $v_{i'_1}, v_{i'_2}, \dots, v_{i'_{n-1}}$ by using Eqs.(47), (48), (49). In this way, we obtain a N_1 dimensional vector \vec{v} , whose entries satisfy Eqs. (36)-(39). Then, for any node $q \in V(\tau_n(G))$, the entry v_q in \vec{v} satisfies Eq. (35).

Therefore, the vector \vec{v} satisfies Eq. (34) and μ is an eigenvalue of $\mathcal{L}_{\tau_n(G)}$. Further more, there are strict one-to-one correspondences be the eigenvectors \vec{v} of μ and $\vec{v'}$, which is an arbitrary vector in N_0 -dimensional space. Thus the multiplicity of the eigenvalue μ is N_0 .

(iii) Let μ be an arbitrary root of equation $a_{\frac{n}{2}-1}(\mu) = 0$. Then

$$a_{n-1}(\mu) = [a_{\frac{n}{2}}(\mu) - a_{\frac{n}{2}-2}(\mu)]a_{\frac{n}{2}-1}(\mu) = 0,$$

and we can obtain from Lemma 3.8 (see Eq. (23)) that

$$a_{n-2}\left(\mu\right) = -1.$$

Therefore,

$$a_{n-3}(\mu) = 2(1-\mu)a_{n-2}(\mu) - a_{n-1}(\mu) = -2(1-\mu)$$

Replacing a_{n-1} and a_{n-2} with 0 and -1 in Eq.(41) respectively, for any two nodes i and j of graph G, if $i \sim j$,

$$\frac{v_i}{\sqrt{d_i}} = \frac{v_j}{\sqrt{d_j}}.$$
(73)

Since G is a connected graph, Eq. (73) shows that $\frac{v_i}{\sqrt{d_i}}$ is a const for any $i \in V(G)$. Let $\theta \equiv \frac{v_i}{\sqrt{d_i}}$, for any $i \in V(G)$. Eqs. (40) and (36) can be rewritten as

$$v_{i'_1} + v_{i'_{n-1}} = 2 (1 - \mu) \theta, \quad i'_1 \in N^i_N, i'_{n-1} \in N^j_N.$$
(74)

$$\sum_{i_1' \in N_M^i} v_{i_1'} = [2(1-\mu) - 1] \,\theta d_i, \quad i \in V(G).$$
(75)

Calculating $\sum_{i \in V(G)} \sum_{i'_1 \in N_N^i} v_{i'_1}$ by using Eq.(74), we have

$$\sum_{i \in V(G)} \sum_{i_1' \in N_N^i} v_{i_1'} = \frac{1}{2} \sum_{i \sim j, i_1' \in N_N^i, i_{n-1}' \in N_N^j} \left(v_{i_1'} + v_{i_{n-1}'} \right) = 2 \left(1 - \mu \right) \theta E_0.$$
(76)

On the other hand, Calculating $\sum_{i \in V(G)} \sum_{i'_1 \in N_N^i} v_{i'_1}$ by using Eq.(75), we get

$$\sum_{i \in V(G)} \sum_{i_1' \in N_N^i} v_{i_1'} = \left[2\left(1-\mu\right)-1\right] \theta \sum_{i \in V(G)} d_i = \left[4\left(1-\mu\right)-2\right] \theta E_0.$$
(77)

Therefore, $[4(1-\mu)-2]\theta = 2(1-\mu)\theta$, which leads to $\mu = 0$ or $\theta = 0$.

If $\mu = 0$, replacing $a_{\frac{n}{2}-1}(\mu)$ from Eq.(7), we have

$$a_{\frac{n}{2}-1}(\mu) = \frac{n}{2} - 1 + 1 = \frac{n}{2} \neq 0.$$

Thus, $\mu \neq 0$, and $\theta = 0$, which leads to

$$v_i = 0, \tag{78}$$

for any node i of graph G.

Substituting v_i with 0, Eq.(36) can be rewritten as

$$\sum_{i_1' \in N_N^i} v_{i_1'} = 0, \tag{79}$$

for any node $i \in V(G)$.

Similarly, replacing v_i and $a_{n-2}(\mu)$ with 0 and -1 respectively in Eq. (40), we get

$$v_{i_1'} + v_{i_{n-1}'} = 0. ag{80}$$

Further more, if n is odd and $v_i = 0$, in any (n+1)-polygon of $\tau_n(G)$ as shown in Fig. 1, we find, Eqs. (37), (38) and (39) are equivalent to

$$v_{i'_{t+1}} + v_{i'_{n-t-1}} = 0, \quad t = 0, \ 1, \ 2, \ \cdots, \ \frac{n}{2} - 1,$$
 (81)

$$a_t(\mu_i)v_{i'_1} = v_{i'_{t+1}}, \quad t = 1, 2, \cdots, \frac{n}{2} - 1.$$
 (82)

Therefore, Eq. (35) holds if and only if Eqs. (78), (79), (81) and (82) hold for any (n + 1)-polygon of graph $\tau_n(G)$.

Note that the total number of (n + 1)-polygons in graph $\tau_n(G)$ is E_0 . For an arbitrary (n + 1)-polygon, P_r , of graph $\tau_n(G)$, let $v_{i'_1} = -v_{i'_{n-1}} = x_r$, and let $X = (x_1, x_2, \dots, x_{E_0})^T$. We find Eqs.(79) and (80) are equivalent to B'X = 0, where B' is the incident matrix of weakly connected directed graph G' (see Definition 2.5 and 2.6).

Similarly, for an arbitrary (n + 1)-polygon, P_r , of graph $\tau_n(G)$, let

$$v_{i'_{t+1}} = -v_{i'_{n-t-1}} = y_r^{(t)}, \ t = 1, \ 2, \ \dots, \ \frac{n}{2} - 1,$$

and let $Y^{(t)} = (y_1^{(t)}, y_2^{(t)}, \dots, y_{E_0}^{(t)})^T$, we find, Eqs. (81) and (82) can be rewritten as

$$Y^{(t)} - a_t(\mu)X = 0, \quad t = 1, 2, \cdots, \frac{n}{2} - 1$$

Therefore, there are strict one-to-one correspondences between the roots of Eq. (34) and roots of the equation

$$\begin{pmatrix} I \ 0 \ \cdots \ 0 \ -a_1 \ (\mu) \ I \\ 0 \ I \ \cdots \ 0 \ -a_2 \ (\mu) \ I \\ \vdots \ \vdots \ \ddots \ \vdots \ \vdots \\ 0 \ 0 \ \cdots \ I \ -a_{\frac{n}{2}-2} \ (\mu) \ I \\ 0 \ 0 \ \cdots \ 0 \ B' \end{pmatrix} \begin{pmatrix} Y^{(1)} \\ Y^{(2)} \\ \vdots \\ Y^{(\frac{n}{2}-2)} \\ X \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix},$$
(83)

where I is an identity matrix of order E_0 and B' is the incidence matrix of weakly connected directed graph G' whose underlying undirected graph (all edges replaced by undirected edges) is G.

Because $rank(B') = N_0 - 1$ (see Lemma 3.5). Therefore the rank for the coefficient matrix of linear equation (83) is $\frac{n-4}{2}E_0 + N_0$ and the dimension for the vector space spanned by the roots of linear equation (83) is $E_0 - N_0 + 1$. Thus the dimension for the eigenspace of μ is also $E_0 - N_0 + 1$ and the multiplicity for eigenvalue μ is $E_0 - N_0 + 1$.

(iv) Let μ be an arbitrary root of equation $a_{\frac{n}{2}}(\mu) - a_{\frac{n}{2}-2}(\mu) = 0$. Then

$$a_{n-1}(\mu) = [a_{\frac{n}{2}}(\mu) - a_{\frac{n}{2}-2}(\mu)]a_{\frac{n}{2}-1}(\mu) = 0,$$

and we can obtain from Lemma 3.8 (see Eq. (21)) that

$$a_{n-2}\left(\mu\right) = 1.$$

Replacing a_{n-1} and a_{n-1} with 0 and 1 in Eq.(41), for any two nodes *i* and *j* of graph *G*, if $i \sim j$,

$$\frac{v_i}{\sqrt{d_i}} = -\frac{v_j}{\sqrt{d_j}}.$$
(84)

Note that G is non-bipartite, there is at least an odd cycle C in G [8]. Let i_1 , i_2, \dots, i_k (k is odd) be the nodes series of cycle C. Eq. (84) informs us that

$$\frac{v_{i_1}}{\sqrt{d_{i_1}}} = -\frac{v_{i_2}}{\sqrt{d_{i_2}}} = \frac{v_{i_3}}{\sqrt{d_{i_3}}} = \dots = \frac{v_{i_k}}{\sqrt{d_{i_k}}} = -\frac{v_{i_1}}{\sqrt{d_{i_1}}}.$$

Therefore $v_{i_1} = v_{i_2} = \cdots = v_{i_k} = 0$. Since G is connected, Eq. (84) informs us,

$$v_i = 0. \tag{85}$$

for any node i of graph G.

Substituting v_i with 0, Eq.(36) can be rewritten as

$$\sum_{i_1' \in N_N^i} v_{i_1'} = 0, \tag{86}$$

for any node i of graph G.

Similarly, replacing v_i and $a_{n-2}(\mu)$ with 0 and 1 respectively in Eq. (40), we get

$$v_{i_1'} - v_{i_{n-1}'} = 0. (87)$$

Further more, if n is odd and $v_i = 0$ for any node $i \in V(G)$, we find, Eqs. (37), (38) and (39) are equivalent to

$$v_{i_{t+1}'} - v_{i_{n-t-1}'} = 0, \quad t = 0, \ 1, \ 2, \ \cdots, \ \frac{n}{2} - 1,$$
 (88)

$$a_t(\mu_i)v_{i'_1} = v_{i'_{t+1}}, \quad t = 1, \ 2, \ \cdots, \ \frac{n}{2} - 1.$$
 (89)

Therefore, Eq. (35) holds if and only if Eqs. (85), (86), (88) and (89) hold for any (n + 1)-polygon of graph $\tau_n(G)$.

Note that the total number of (n + 1)-polygons in graph $\tau_n(G)$ is E_0 . For an arbitrary (n + 1)-polygon, P_r , of graph $\tau_n(G)$, let $v_{i'_1} = v_{i'_{n-1}} = x_r$, and let $X = (x_1, x_2, \dots, x_{E_0})^T$. We find Eqs.(86) and (87) are equivalent to BX = 0, where B is the incident matrix of graph G.

Similarly, for an arbitrary (n + 1)-polygon, P_r , of graph $\tau_n(G)$, let

$$v_{i'_{t+1}} = v_{i'_{n-t-1}} = y_r^{(t)}, \ t = 1, \ 2, \ \dots, \ \frac{n}{2} - 1,$$

and let $Y^{(t)} = (y_1^{(t)}, y_2^{(t)}, \dots, y_{E_0}^{(t)})^T$, we find, Eqs. (88) and (89) can be rewritten as

$$Y^{(t)} - a_t(\mu)X = 0, \quad t = 1, 2, \cdots, \frac{n-1}{2} - 1$$

Therefore, there are strict one-to-one correspondences between the roots of Eq. (34) and roots of the equation

$$\begin{pmatrix} I \ 0 \ \cdots \ 0 \ -a_1 \ (\breve{\mu}_i) \ I \\ 0 \ I \ \cdots \ 0 \ -a_2 \ (\breve{\mu}_i) \ I \\ \vdots \ \vdots \ \ddots \ \vdots \ \vdots \\ 0 \ 0 \ \cdots \ I \ -a_{\frac{n}{2}-1} \ (\breve{\mu}_i) \ I \\ 0 \ 0 \ \cdots \ 0 \ B \end{pmatrix} \begin{pmatrix} Y^{(1)} \\ Y^{(2)} \\ \vdots \\ Y^{(\frac{n}{2}-1)} \\ X \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix},$$
(90)

where B is the incident matrix of G and I is identity matrix of order E_0 .

Because G is non-bipartite, $rank(B) = N_0$ (see Lemma 3.4). Therefore the rank for the coefficient matrix of linear equation (90) is $\frac{n-2}{2}E_0 + N_0$ and the dimension for the vector space spanned by the roots of linear equation (90) is $E_0 - N_0$. Thus the dimension for the eigenspace of μ is also $E_0 - N_0$ and the multiplicity for eigenvalue μ is $E_0 - N_0$.

(v) Let μ be an arbitrary root of equation $a_{\frac{n}{2}}(\mu) - a_{\frac{n}{2}-2}(\mu) = 0$. Then

$$a_{n-1}(\mu) = [a_{\frac{n}{2}}(\mu) - a_{\frac{n}{2}-2}(\mu)]a_{\frac{n}{2}-1}(\mu) = 0,$$

and we can obtain from Lemma 3.8 (see Eq. (21)) that

$$a_{n-2}\left(\mu\right) = 1.$$

Replacing a_{n-1} and a_{n-2} with 0 and 1 in Eq.(41), for any two nodes *i* and *j* of graph *G*, if $i \sim j$,

$$\frac{v_i}{\sqrt{d_i}} = -\frac{v_j}{\sqrt{d_j}}.$$
(91)

Note that G is bipartite whose vertex set can be partitioned into two disjoint sets X and Y, i.e., $V(G) = X \cup Y$. By Eq.(91),we have

$$\frac{v_i}{\sqrt{d_i}} = \begin{cases} \theta, i \in X\\ -\theta, i \in Y \end{cases}$$
(92)

If $i \in X$ and $i \in Y$, Eq.(36) can be rewritten as, respectively,

$$\sum_{i_1' \in N_N^i} v_{i_1'} = [2(1-\mu)+1] \,\theta d_i, \quad i \in X,$$
(93)

$$\sum_{i_1' \in N_N^i} v_{i_1'} = -\left[2\left(1-\mu\right)+1\right] \theta d_i, \quad i \in Y.$$
(94)

If $i \in X$, Eq.(40) can be rewritten as

$$v_{i'_{1}} - v_{i'_{n-1}} = 2 (1 - \mu) \theta, \quad i'_{1} \in N_{N}^{i}, i'_{n-1} \in N_{N}^{j}.$$
(95)

By calculating the difference of new neighbor nodes of nodes in bipartite graph G, we have

$$\sum_{i \in X} \sum_{i_1' \in N_N^i} v_{i_1'} - \sum_{i \in Y} \sum_{i_1' \in N_N^i} v_{i_1'} = \sum_{i \in X} \left(v_{i_1'} - v_{i_{n-1}'} \right).$$
(96)

Therefore,

$$[4(1-\mu)+2]\theta E_0 = 2(1-\mu)\theta E_0.$$
(97)

Thus, $\theta = 0$, which leads to

$$v_i = 0, \tag{98}$$

for any node i of graph G.

Similarity to the proof of (iv), we find, in the case G is bipartite, there are strict one-to-one correspondences between the roots of Eq. (34) and roots of the equation Eq. (90). However, different with non-bipartite graph, here $rank(B) = N_0 - 1$ (see Lemma 3.4). Then, the dimension for the eigenspace of μ is $E_0 - N_0 + 1$ and the multiplicity for eigenvalue μ is $E_0 - N_0 + 1$.

(vi)It is just the result of Lemma 4.1.

The proof is completed.

9 Proof of Theorem 5.1

Firstly, we prove Eq. (30) holds in the case n is an odd number.

Let λ ($\lambda \neq 0$ and $\lambda \neq 2$) be an arbitrary eigenvalue of \mathcal{L}_G and $\mu_i(\lambda)$ ($i = 1, 2, \dots, \frac{n+1}{2}$) be the roots of equation

$$1 - \frac{a_{\frac{n-1}{2}+1}(x) - a_{\frac{n-1}{2}-1}(x)}{a_{\frac{n-1}{2}}(x) - a_{\frac{n-1}{2}-2}(x)} = \lambda.$$
(99)

Since $a_{\frac{n-1}{2}}(x) - a_{\frac{n-1}{2}-2}(x) \neq 0$, Eq.(99) can be rewritten as

$$f(x) = a_{\frac{n+1}{2}}(x) - (1-\lambda)a_{\frac{n-1}{2}}(x) - a_{\frac{n-3}{2}}(x) + (1-\lambda)a_{\frac{n-5}{2}}(x) = 0, \quad (100)$$

which is a the polynomial equation of degree $\frac{n+1}{2}$. Let b_i be the coefficient of x^i $(i = 1, 2, \dots, \frac{n+1}{2})$ for polynomial f(x). According to Lemma 3.6(iv), we get

$$b_{0} = a_{\frac{n+1}{2}}^{(0)} (x) - (1-\lambda) a_{\frac{n-1}{2}}^{(0)} (x) - a_{\frac{n-3}{2}}^{(0)} (x) + (1-\lambda) a_{\frac{n-5}{2}}^{(0)} (x)$$

= $\frac{n+1}{2} + 1 - (1-\lambda) \left(\frac{n-1}{2} + 1\right) - \left(\frac{n-3}{2} + 1\right) + (1-\lambda) \left(\frac{n-5}{2} + 1\right)$
= $2\lambda,$ (101)

$$b_{1} = a_{\frac{n+1}{2}}^{(1)} (x) - (1-\lambda) a_{\frac{n-1}{2}}^{(1)} (x) - a_{\frac{n-3}{2}}^{(1)} (x) + (1-\lambda) a_{\frac{n-5}{2}}^{(1)} (x)$$

$$= -\frac{\left(\frac{n+1}{2}\right)^{3} + 3\left(\frac{n+1}{2}\right)^{2} + 2\left(\frac{n+1}{2}\right)}{3} + (1-\lambda) \frac{\left(\frac{n-1}{2}\right)^{3} + 3\left(\frac{n-1}{2}\right)^{2} + 2\left(\frac{n-1}{2}\right)}{3}$$

$$+ \frac{\left(\frac{n-3}{2}\right)^{3} + 3\left(\frac{n-3}{2}\right)^{2} + 2\left(\frac{n-3}{2}\right)}{3} - (1-\lambda) \frac{\left(\frac{n-5}{2}\right)^{3} + 3\left(\frac{n-5}{2}\right)^{2} + 2\left(\frac{n-5}{2}\right)}{3}$$

$$= -\frac{\left(n-1\right)^{2}}{2}\lambda - 2n, \qquad (102)$$

and

$$b_{\frac{n+1}{2}} = a_{\frac{n+1}{2}}^{\left(\frac{n+1}{2}\right)}(x) = (-1)^{\frac{n+1}{2}} 2^{\frac{n+1}{2}}.$$
(103)

By using Lemma 3.3, we have

$$\sum_{i=1}^{\frac{n+1}{2}} \frac{1}{\mu_i(\lambda)} = -\frac{b_1}{b_0} = \frac{n}{\lambda} + \left(\frac{n-1}{2}\right)^2.$$
 (104)

Similarly, let $\check{\mu}_i(i=1,2,\cdots,\frac{n-1}{2})$ be the roots of equation $a_{\frac{n-1}{2}} - a_{\frac{n-1}{2}-1} = 0$, $\check{\mu}_i(i=\frac{n-1}{2}+1,\frac{n-1}{2}+2,\cdots,n-1)$ be the roots of equation $a_{\frac{n-1}{2}} + a_{\frac{n-1}{2}-1} = 0$, and $\hat{\mu}_i(i=1,2,\cdots,\frac{n-1}{2})$ be the roots of equation $a_{\frac{n-1}{2}} = 0$. Rewriting these equations as polynomial equation and using Lemma 3.3, we have

$$\sum_{i=1}^{\frac{n-1}{2}} \frac{1}{\check{\mu}_i} = \frac{n^2 - 1}{4},\tag{105}$$

$$\sum_{i=\frac{n-1}{2}+1}^{n-1} \frac{1}{\check{\mu}_i} = \frac{n^2 - 1}{12},\tag{106}$$

and

$$\sum_{i=1}^{\frac{n-1}{2}} \frac{1}{\hat{\mu}_i} = \frac{n^2 + 2n - 3}{12}.$$
 (107)

Let $0 = \lambda_1 < \lambda_2 \leq \lambda_3 \leq \cdots \leq \lambda_{N_0}$ be the eigenvalues of the normalized Laplacian \mathcal{L}_G . If G is non-bipartite, Theorem 4.2 informs us that the normalized Laplacian spectrum of graph $\tau_n(G)$ can be written as

$$\sigma\left(\tau_{n}(G)\right) = \bigcup_{j=2}^{N_{0}} \left[\bigcup_{i=1}^{\frac{n+1}{2}} \{\mu_{i}\left(\lambda_{j}\right)\}\right] \bigcup_{i=1}^{\frac{n-1}{2}} \underbrace{\{\hat{\mu}_{i}, \hat{\mu}_{i}, \cdots, \hat{\mu}_{i}\}}_{N_{0}} \bigcup_{i=1}^{\frac{n-1}{2}} \underbrace{\{\check{\mu}_{i}, \check{\mu}_{i}, \cdots, \check{\mu}_{i}\}}_{E_{0}-N_{0}} \left\{\underbrace{\check{\mu}_{i}, \check{\mu}_{i}, \cdots, \check{\mu}_{i}}_{E_{0}-N_{0}+1} \cup \{0\}\right\}.$$
(108)

Therefore, in the case n is an odd number and G is non-bipartite, by using Lemma 3.2(i) and Eqs (104)-(107), we have

$$Kf'(\tau_n(G)) = 2E_1 \sum_{u \in \sigma(\tau_n(G)), \ \mu \neq 0} \frac{1}{\mu}$$

=2E_1 $\left[\sum_{i=2}^{N_0} \sum_{j=1}^{\frac{n+1}{2}} \frac{1}{\mu_j(\lambda_i)} + \sum_{i=1}^{\frac{n-1}{2}} \frac{1}{\hat{\mu}_i} N_0 + \sum_{i=1}^{\frac{n-1}{2}} \frac{1}{\hat{\mu}_i} (E_0 - N_0) + \sum_{i=\frac{n-1}{2}+1}^{n-1} \frac{1}{\hat{\mu}_i} (E_0 - N_0 + 1) \right]$
=2E_1 $\left[\sum_{i=2}^{N_0} \left(\frac{n}{\lambda_i} + \left(\frac{n-1}{2} \right)^2 \right) + \frac{n^2 + 2n - 3}{12} N_0 + \frac{n^2 - 1}{4} (E_0 - N_0) + \frac{n^2 - 1}{12} (E_0 - N_0 + 1) \right]$
=2(n+1) E_0 $\left[n \sum_{i=2}^{N_0} \frac{1}{\lambda_i} + \frac{n^2 - 1}{3} E_0 - \frac{n-1}{3} N_0 - \frac{(n-1)(n-2)}{6} \right]$
= (n^2 + n) Kf'(G) + $\frac{2}{3} (n+1) (n^2 - 1) E_0^2 - \frac{2}{3} (n^2 - 1) E_0 N_0 - \frac{1}{6} (n^2 - 1) (n-2) E_0.$ (109)

Thus Eq. (30) holds in the case n is an odd number and G is non-bipartite.

If G is bipartite, Theorem 4.2 informs us that the normalized Laplacian spectrum of graph $\tau_n(G)$ can be written as

$$\sigma\left(\tau_{n}(G)\right) = \bigcup_{j=2}^{N_{0}-1} \left[\bigcup_{i=1}^{\frac{n+1}{2}} \{\mu_{i}\left(\lambda_{j}\right)\}\right] \bigcup_{i=1}^{\frac{n-1}{2}} \underbrace{\{\hat{\mu}_{i}, \hat{\mu}_{i}, \cdots, \hat{\mu}_{i}\}}_{N_{0}} \bigcup_{i=1}^{\frac{n-1}{2}} \underbrace{\{\check{\mu}_{i}, \check{\mu}_{i}, \cdots, \check{\mu}_{i}\}}_{E_{0}-N_{0}+1} \left[\bigcup_{i=\frac{n-1}{2}+1}^{n-1} \underbrace{\{\check{\mu}_{i}, \check{\mu}_{i}, \cdots, \check{\mu}_{i}\}}_{E_{0}-N_{0}+1} \cup \{0\} \cup \{2\}.$$
(110)

Similarly, we can also verify Eq. (30) holds in the case n is an odd number and G is bipartite. Therefore, Eq. (30) holds in the case n is an odd number.

Then, we prove Eq. (30) holds in the case n is an even number.

Let λ ($\lambda \neq 0$ and $\lambda \neq 2$) be an arbitrary eigenvalue of \mathcal{L}_G and $\mu_i(\lambda)$ ($i = 1, 2, \dots, \frac{n}{2}$) be the roots of equation

$$1 - \frac{a_{\frac{n}{2}}(x) - a_{\frac{n}{2}-1}(x)}{a_{\frac{n}{2}-1}(x) - a_{\frac{n}{2}-2}(x)} = \lambda.$$
 (111)

Similar to the proof of Eq. (104), we have

$$\sum_{i=1}^{\frac{n}{2}} \frac{1}{\mu_i(\lambda)} = \frac{n}{\lambda} + \frac{n^2 - 2n}{4}.$$
 (112)

Similarly, let $\check{\mu}_i(i=1,2,\cdots,\frac{n}{2})$ be the roots of equation $a_{\frac{n}{2}} - a_{\frac{n}{2}-2} = 0$, $\check{\mu}_i(i=\frac{n}{2}+1,\frac{n}{2}+2,\cdots,n-1)$ be the roots of equation $a_{\frac{n}{2}-1} = 0$, and $\hat{\mu}_i(i=1,2,\cdots,\frac{n}{2})$ be the roots of equation $a_{\frac{n}{2}} + a_{\frac{n}{2}-1} = 0$. We have

$$\sum_{i=1}^{\frac{n}{2}} \frac{1}{\breve{\mu}_i} = \frac{n^2}{4},\tag{113}$$

$$\sum_{i=\frac{n}{2}+1}^{n-1} \frac{1}{\breve{\mu}_i} = \frac{n^2 - 4}{12},\tag{114}$$

and

$$\sum_{i=1}^{\frac{n}{2}} \frac{1}{\hat{\mu}_i} = \frac{n^2 + 2n}{12}.$$
(115)

Let $0 = \lambda_1 < \lambda_2 \leq \lambda_3 \leq \cdots \leq \lambda_{N_0}$ be the eigenvalues of the normalized Laplacian \mathcal{L}_G . Similar to the proof of Eq. (109), in the case *n* is an even number and *G* is non-bipartite, we have

$$Kf'(\tau_{n}(G)) = 2E_{1}\left[\sum_{i=2}^{N_{0}}\sum_{j=1}^{\frac{n}{2}}\frac{1}{\mu_{j}(\lambda_{i})} + \sum_{i=1}^{\frac{n}{2}}\frac{1}{\tilde{\mu}_{i}}N_{0} + \sum_{i=1}^{\frac{n}{2}}\frac{1}{\tilde{\mu}_{i}}(E_{0} - N_{0}) + \sum_{i=\frac{n}{2}+1}^{n-1}\frac{1}{\tilde{\mu}_{i}}(E_{0} - N_{0} + 1)\right] = 2E_{1}\left[\sum_{i=2}^{N_{0}}\left(\frac{n}{\lambda_{i}} + \frac{n^{2} - 2n}{4}\right) + \frac{n^{2} + 2n}{12}N_{0} + \frac{n^{2}}{4}(E_{0} - N_{0}) + \frac{n^{2} - 4}{12}(E_{0} - N_{0} + 1)\right] = 2(n+1)E_{0}\left[n\sum_{i=2}^{N_{0}}\frac{1}{\lambda_{i}} + \frac{n^{2} - 1}{3}E_{0} - \frac{n-1}{3}N_{0} - \frac{(n-1)(n-2)}{6}\right] = (n^{2} + n)Kf'(G) + \frac{2}{3}(n+1)(n^{2} - 1)E_{0}^{2} - \frac{2}{3}(n^{2} - 1)E_{0}N_{0} - \frac{1}{6}(n^{2} - 1)(n-2)E_{0}.$$
(116)

Thus Eq. (30) holds in the case n is an even number and G is non-bipartite. Similarly, we can also verify Eq. (30) holds in the case n is an even number and G is bipartite. Therefore, Eq. (30) holds for any n ($n \ge 2$).

Next, we will prove Eq. (31) holds for any $n \ge 2$ and $g \ge 1$.

Recalling the definition of the iterated *n*-polygon graph with generation g (see Definition 2.8), for any $g \ge 1$, $\tau_n^g(G) = \tau_n(\tau_n^{g-1}(G))$. Therefore, we can obtain from from Eq.(30) that, for any $g \ge 1$,

$$Kf'(\tau_n^g(G)) = (n^2 + n) Kf'(\tau_n^{g-1}(G)) + \frac{2}{3}(n+1)(n^2 - 1) E_{g-1}^2 - \frac{2}{3}(n^2 - 1) E_{g-1}N_{g-1} - \frac{1}{6}(n^2 - 1)(n-2) E_{g-1}.$$
 (117)

Using Eq.(117) recursively and replacing N_g from Eq.(5), we have

$$\begin{split} &Kf'\left(\tau_n^g(G)\right) \\ = \left(n^2 + n\right)^g Kf'\left(G\right) + \frac{2}{3}\left(n+1\right)\left(n^2 - 1\right)\sum_{i=0}^{g-1} \left(n^2 + n\right)^i E_{g-1-i}^2 - \frac{2}{3}\left(n^2 - 1\right)\cdot \\ &\sum_{i=0}^{g-1} \left(n^2 + n\right)^i E_{g-1-i} N_{g-1-i} - \frac{1}{3}\left(n^2 - 1\right)\left(n-2\right)\sum_{i=0}^{g-1} \left(n^2 + n\right)^i E_{g-1-i} \\ = \left(n^2 + n\right)^g Kf'\left(G\right) + \frac{2}{3}\left(n+1\right)\left(n^2 - 1\right)\left[\left(n+1\right)^{2g-1} - n^g\left(n+1\right)^{g-1}\right] E_0^2 - \frac{2}{3}\left(n^2 - 1\right)\left\{\frac{\left(n+1\right)^{g-1}\left(n^g - 1\right)}{n-1} E_0 N_0 + \frac{\left(n+1\right)^{g-1}\left[\left(n-1\right)\left(n+1\right)^g - n^{g+1} + 1\right]}{n} E_0^2\right\} \\ &- \frac{1}{3}\left(n^2 - 1\right)\left(n-2\right)\frac{\left(n+1\right)^{g-1}\left(n^g - 1\right)}{n-1} E_0 \\ = \left(n^2 + n\right)^g Kf'\left(G\right) + \frac{2\left(n-1\right)}{3n}\left(n+1\right)^g \left[\left(n+1\right)^g \left(n^2 + 1\right) - n^{g+2} - 1\right] E_0^2 \\ &- \frac{1}{3}\left(n-2\right)\left(n+1\right)^g \left(n^g - 1\right) E_0 - \frac{2}{3}\left(n+1\right)^g \left(n^g - 1\right) E_0 N_0. \end{split}$$

Then we obtain Eq. (31).

10 Proof of Theorem 5.3

Firstly, we prove Eq. (32) holds in the case n is an odd number.

Let λ ($\lambda \neq 0$ and $\lambda \neq 2$) be an arbitrary eigenvalue of \mathcal{L}_G and $\mu_i(\lambda)$ ($i = 1, 2, \dots, \frac{n}{2}$) be the roots of equation

$$1 - \frac{a_{\frac{n-1}{2}+1}(x) - a_{\frac{n-1}{2}-1}(x)}{a_{\frac{n-1}{2}}(x) - a_{\frac{n-1}{2}-2}(x)} = \lambda.$$
 (118)

Note that Eq.(118) can be rewritten as Eq. (100), which is a polynomial equation, with some coefficients shown as Eqs.(101) and (103). By using Lemma 3.3, we have

$$\prod_{i=1}^{\frac{n+1}{2}} \mu_i(\lambda) = (-1)^n \frac{b_0}{b_{\frac{n+1}{2}}} = \frac{\lambda}{2^{\frac{n-1}{2}}}.$$
(119)

Similarly, let $\check{\mu}_i(i=1,2,\cdots,\frac{n-1}{2})$ be the roots of equation $a_{\frac{n-1}{2}} - a_{\frac{n-1}{2}-1} = 0$, $\check{\mu}_i(i=\frac{n-1}{2}+1,\frac{n-1}{2}+2,\cdots,n-1)$ be the roots of equation $a_{\frac{n-1}{2}} + a_{\frac{n-1}{2}-1} = 0$, and $\hat{\mu}_i(i=1,2,\cdots,\frac{n-1}{2})$ be the roots of equation $a_{\frac{n-1}{2}} = 0$. Rewriting these equations as polynomial equation and using Lemma 3.3, we have

$$\prod_{i=1}^{\frac{n-1}{2}} \check{\mu}_i = \frac{1}{2^{\frac{n-1}{2}}},\tag{120}$$

$$\prod_{i=\frac{n-1}{2}+1}^{n-1}\check{\mu}_i = \frac{n}{2^{\frac{n-1}{2}}},\tag{121}$$

and

$$\prod_{i=1}^{\frac{n-1}{2}} \hat{\mu}_i = \frac{n+1}{2^{\frac{n+1}{2}}}.$$
(122)

Let $0 = \lambda_1 < \lambda_2 \leq \lambda_3 \leq \cdots \leq \lambda_{N_0}$ be the eigenvalues of the normalized Laplacian \mathcal{L}_G . If G is non-bipartite, Note that the normalized Laplacian spectrum $\sigma(\tau_n(G))$ of graph $\tau_n(G)$ are shown in Eq. (108), we have

$$\prod_{\mu \in \sigma(\tau_n(G)), \mu \neq 0} \mu \\
= \left(\prod_{i=1}^{\frac{n-1}{2}} \hat{\mu}_i\right)^{N_0} \times \left(\prod_{i=1}^{\frac{n-1}{2}} \check{\mu}_i\right)^{E_0 - N_0} \times \left(\prod_{i=\frac{n-1}{2}+1}^{n-1} \check{\mu}_i\right)^{E_0 - N_0 + 1} \times \prod_{i=2}^{N_0} \prod_{j=1}^{\frac{n-1}{2}} \mu_j \left(\lambda_i\right) \\
= \left(\frac{n+1}{2^{\frac{n+1}{2}}}\right)^{N_0} \times \left(\frac{1}{2^{\frac{n-1}{2}}}\right)^{E_0 - N_0} \times \left(\frac{n}{2^{\frac{n-1}{2}}}\right)^{E_0 - N_0 + 1} \times \prod_{i=2}^{N_0} \frac{\lambda_i}{2^{\frac{n-1}{2}}} \\
= \frac{n^{E_0 - N_0 + 1}}{(2^{n-1})^{E_0}} \left(\frac{n+1}{2}\right)^{N_0} \times \prod_{i=2}^{N_0} \lambda_i.$$
(123)

Therefore,

$$\frac{N_{st}(G(n))}{N_{st}(G)} = \frac{2^{N_1} \prod_{\mu \in \sigma(\tau_n(G)), \mu \neq 0} \mu}{(n+1) \prod_{i=2}^{N_0} \lambda_i} = n^{E_0 - N_0 + 1} (n+1)^{N_0 - 1}.$$
 (124)

Thus, Eq. (32) holds in the case n is an odd number and G is non-bipartite.

In the case n is an odd number and G is bipartite, recalling the normalized Laplacian spectrum $\sigma(\tau_n(G))$ of graph $\tau_n(G)$ are shown in Eq. (110), we can also verify that Eq. (32) holds. Therefore Eq. (32) holds in the case n is an odd number.

Then, we prove Eq. (32) holds in the case n is an even number.

Let λ ($\lambda \neq 0$ and $\lambda \neq 2$) be an arbitrary eigenvalue of \mathcal{L}_G and $\mu_i(\lambda)$ (i =

1, 2, \cdots , $\frac{n}{2}$) be the roots of equation

$$1 - \frac{a_{\frac{n}{2}}(x) - a_{\frac{n}{2}-1}(x)}{a_{\frac{n}{2}-1}(x) - a_{\frac{n}{2}-2}(x)} = \lambda.$$
 (125)

Similar to the proof of Eq. (119), we have

$$\prod_{i=1}^{\frac{n}{2}} \mu_i\left(\lambda\right) = \frac{\lambda}{2^{\frac{n}{2}}}.$$
(126)

Similarly, let $\check{\mu}_i(i=1,2,\cdots,\frac{n}{2})$ be the roots of equation $a_{\frac{n}{2}} - a_{\frac{n}{2}-2} = 0$, $\check{\mu}_i(i=\frac{n}{2}+1,\frac{n}{2}+2,\cdots,n-1)$ be the roots of equation $a_{\frac{n}{2}-1} = 0$, and $\hat{\mu}_i(i=1,2,\cdots,\frac{n}{2})$ be the roots of equation $a_{\frac{n}{2}} + a_{\frac{n}{2}-1} = 0$. We have

$$\prod_{i=1}^{\frac{n}{2}} \check{\mu}_i = \frac{1}{2^{\frac{n}{2}-1}},\tag{127}$$

$$\prod_{i=\frac{n}{2}+1}^{n-1} \breve{\mu}_i = \frac{n}{2^{\frac{n}{2}}},\tag{128}$$

and

$$\prod_{i=1}^{\frac{n}{2}} \hat{\mu}_i = \frac{n+1}{2^{\frac{n}{2}}}.$$
(129)

Calculating the normalized Laplacian spectrum $\sigma(\tau_n(G))$ of graph $\tau_n(G)$ by using Theorem 4.3 in the case *n* is even number, we can also get Eq. (124). Thus, Eq. (32) holds in the case *n* is an even number.

Recalling the definition of the iterated *n*-polygon graph with generation g (see Definition 2.8), for any $g \ge 1$, $\tau_n^g(G) = \tau_n(\tau_n^{g-1}(G))$. Therefore, we can obtain from from Eq.(32) that, for any $g \ge 1$,

$$N_{st}\left(\tau_n^g(G)\right) = (n+1)^{N_{g-1}-1} n^{E_{g-1}-N_{g-1}+1} \cdot N_{st}\left(\tau_n^{g-1}(G)\right).$$
(130)

By using Eq.(130) recursively, we obtain Eq.(33).

11 Conclusion

Given the normalized Laplacian spectrum of an arbitrary connected graph G, we have obtained the normalized Laplacian spectrum of the graph $\tau_n(G)$, which is obtained by replacing each edge of G with a (n+1)-polygon $(n \ge 2)$. Then, the normalized Laplacian spectrum of the graphs $\tau_n^g(G)$ $(g \ge 0)$ can

also be obtained. As applications, we also calculated the multiplicative degree-Kirchhoff index, the Kemeny's const and the number of spanning trees for graphs $\tau_n^g(G)$ ($g \ge 0$). Note that 'edge replacing' is a common graph operator which were widely used to construct different meaningful networks. The results obtained here would be helpful for network optimization.

Acknowledgement(s)

The authors would like to express their sincere gratitude to all former researchers who present original results they quote in this manuscript.

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

The work is supported by the National Natural Science Foundation of China (Grant No. 61873069 and 61772147).

ORCID

Junhao Peng, https://orcid.org/0000-0001-8839-7070

References

- L. Lovász, Random walks on graphs: A survey, in: D. Miklós, V. T. Sós, T. Szönyi (Eds.), Combinatorics, Paul Erdös is Eighty, Vol. 2, János Bolyai Mathematical Society, Budapest, 1996, pp. 353–398.
- [2] D. M. Cvetković, P. Rowlinson, S. Simić, An introduction to the theory of graph spectra, Vol. 75, Cambridge University Press, London, 2010.
- [3] F. R. Chung, F. C. Graham, Spectral graph theory, no. 92, American Mathematical Soc., 1997.

- [4] R. Mehatari, A. Banerjee, Effect on normalized graph laplacian spectrum by motif attachment and duplication, Appl. Math. Comput. 261 (2015) 382–387.
- [5] S. Redner, A Guide to First-Passage Processes, Cambridge University Press, UK, 2007.
- [6] K. W. Wen, L. F. Huang, Q. Wang, J. S. Yu, Modulation of first-passage time for gene expression via asymmetric cell division, Int. J. Biomath. 12 (4) (2019) 97–114.
- [7] S. H. Wang, Graph Theory, The Chinese science publishing company, 2004.
- [8] A. E. Brouwer, A. E. Brouwer, Spectra of Graphs, Springer New York, 2012.
- [9] M. Cavers, S. Fallat, S. Kirkland, On the normalized Laplacian energy and general Randić index R_{-1} of graphs, Linear Multilinear Algebra 433 (1) (2010) 172–190.
- [10] Y. Lin, Z. Z. Zhang, Random walks in weighted networks with a perfect trap: An application of Laplacian spectra, Phys. Rev. E 87 (6) (2013) 062140. doi:10.1103/PhysRevE.87.062140.
- [11] Z. Z. Zhang, Y. Lin, X. Y. Guo, Eigenvalues for the transition matrix of a smallworld scale-free network: Explicit expressions and applications, Phys. Rev. E 91 (2015) 062808.
- [12] J. Huang, S. C. Li, On the normalized laplacians spectrum, degree-kirchhoff index and the spanning trees of graphs, Bull. Aust. Math. Soc 91 (3) (2015) 353–367.
- [13] Z. Z. Zhang, Y. Lin, X. Y. Guo, Eigenvalues for the transition matrix of a smallworld scale-free network: Explicit expressions and applications, Phys. Rev. E 91 (2015) 062808.
- [14] R. Sharma, B. Adhikari, A. Mishra, Structural and spectral properties of corona graphs, Discrete Applied Mathematics 228 (2017) 14–31.
- [15] Y. Qi, Z. Z. Zhang, Spectral Properties of Extended Sierpinski Graphs and Their Applications, IEEE Trans. Netw. Sci. Eng. 6 (3) (2019) 512–522.
- [16] C. L. He, S. C. Li, W. J. Luo, L. Q. Sun, Calculating the normalized laplacian spectrum and the number of spanning trees of linear pentagonal chains, Journal of Computational and Applied Mathematics 344 (2018) 381–393.
- [17] G.-S. Cheon, J. S. Kim, S. A. Mojallal, Spectral properties of pascal graphs, Linear Multilinear Algebra 66 (7) (2018) 1403–1417.
- [18] S. N. Dorogovtsev, A. V. Goltsev, J. F. F. Mendes, Pseudofractal scale-free web, Phys. Rev. E 65 (2002) 066122. doi:10.1103/PhysRevE.65.066122.
- [19] H. D. Rozenfeld, S. Havlin, D. Ben-Avraham, Fractal and transfractal recursive scale-free nets, New J. Phys. 9 (175).

- [20] L. Gao, J. H. Peng, C. M. Tang, A. P. Riascos, Trapping efficiency of random walks on weighted scale-free trees, J. Stat. Mech. 2021 (6) (2021) 063405.
- [21] J. H. Peng, E. Agliari, Scaling laws for diffusion on (trans)fractal scale-free networks, Chaos 27 (8) (2017) 083108.
- [22] J. H. Peng, E. Agliari, Z. Z. Zhang, Exact calculations of first-passage properties on the pseudofractal scale-free web, Chaos 25 (7) (2015) 073118.
- [23] Z. Z. Zhang, Y. Lin, Y. J. Ma, Effect of trap position on the efficiency of trapping in treelike scale-free networks, J. Phys. A 44 (7) (2011) 075102.
- [24] E. Agliari, Exact mean first-passage time on the T-graph, Phys. Rev. E 77 (1) (2008) 011128. doi:10.1103/PhysRevE.77.011128.
- [25] P. C. Xie, Z. Z. Zhang, F. Comellas, On the spectrum of the normalized laplacian of iterated triangulations of graphs, Appl. Math. Comput. 273 (2016) 1123– 1129.
- [26] H. Y. Chen, Hitting times for random walks on subdivision and triangulation graphs, Linear Multilinear Algebra 66 (1) (2018) 117–130.
- [27] C. Wang, Z. Guo, S. Li, Expected hitting times for random walks on the ktriangle graph and their applications, Appl. Math. Comput. 338 (2018) 698– 710.
- [28] Z. Guo, S. Li, X. Liu, X. Mei, Expected hitting times for random walks on the diamond hierarchical graphs involving some classical parameters, Linear Multilinear Algebra 69 (10) (2021) 1841–1857.
- [29] D. Q. Li, Y. P. Hou, The normalized laplacian spectrum of quadrilateral graphs and its applications, Appl. Math. Comput. 297 (2017) 180–188.
- [30] J. Huang, S. H. Li, Expected hitting times for random walks on quadrilateral graphs and their applications, Linear Multilinear Algebra 66 (12) (2018) 2389– 2408.
- [31] X. J. Xu, P. W. Wang, Z. P. Wang, The normalized laplacian spectrum of pentagonal graphs and its applications, Journal of Mathematical Research with Applications 039 (004) (2019) 343–352.
- [32] X. D. Zhang, A Matrix Algebra Approach to Artificial Intelligence, Springer Nature Singapore Pte Ltd, 2020.
- [33] P. Borwein, T. Erdelyi, Polynomials and polynomial inequalities, Springer, 1997.
- [34] A. Jeribi, Spectral theory and applications of linear operators and block operator matrices, Springer International Publishing, 2015.

A The proof of the Lemma 3.6

Here, we use a_k to represent $a_k(\mu)$ to lighten the notations. Firstly, we solve the recursive relation which series $\{a_n\}$ satisfies and present the general formula of $\{a_n\}$. Let $\beta = 2(1-\mu)$. Then the recursive relation of series $\{a_n(\mu)\}$ can be rewritten as

$$a_n = \beta a_{n-1} - a_{n-2}, \tag{A.1}$$

with initial conditions $a_{-1} = 0$, $a_0 = 1$. The characteristic equation of equation (A.1) is $x^2 - \beta x + 1 = 0$, with roots $x = \frac{\beta \pm i \sqrt{4-\beta^2}}{2}$. Then $\{a_n(\mu)\}$ can be rewritten as

$$a_n = c_1 \left(\frac{\beta + i\sqrt{4 - \beta^2}}{2}\right)^n + c_2 \left(\frac{\beta - i\sqrt{4 - \beta^2}}{2}\right)^n.$$
 (A.2)

By using the initial conditions $a_{-1} = 0$, $a_0 = 1$, we have

$$\begin{cases} a_{-1} = c_1 \left(\frac{\beta + i\sqrt{4-\beta^2}}{2}\right)^{-1} + c_2 \left(\frac{\beta - i\sqrt{4-\beta^2}}{2}\right)^{-1} = 0 \\ a_0 = c_1 + c_2 = 1 \end{cases}, \quad (A.3)$$

which yields

$$\begin{cases} c_1 = \frac{1}{2} - \frac{i\beta}{2\sqrt{4-\beta^2}} \\ c_2 = \frac{1}{2} + \frac{i\beta}{2\sqrt{4-\beta^2}} \end{cases}.$$
 (A.4)

Therefore,

$$a_{n}(\mu) = \left(\frac{1}{2} - \frac{i\beta}{2\sqrt{4-\beta^{2}}}\right) \left(\frac{\beta + i\sqrt{4-\beta^{2}}}{2}\right)^{n} + \left(\frac{1}{2} + \frac{i\beta}{2\sqrt{4-\beta^{2}}}\right) \left(\frac{\beta - i\sqrt{4-\beta^{2}}}{2}\right)^{n} \equiv a_{n}(\beta),$$
(A.5)

where $\beta = 2(1 - \mu)$. Then, we prove the results of Lemma 3.6 item-by-item.

(i) Note that $\beta = 2(1 - \mu)$. Then $2[1 - (2 - \mu)] = 2(\mu - 1) = -\beta$. Therefore,

$$a_{n}(2-\mu) = a_{n}(-\beta)$$

$$= \left(\frac{1}{2} - \frac{-i\beta}{2\sqrt{4-\beta^{2}}}\right) \left(\frac{-\beta + i\sqrt{4-\beta^{2}}}{2}\right)^{n} + \left(\frac{1}{2} + \frac{-i\beta}{2\sqrt{4-\beta^{2}}}\right) \left(\frac{-\beta - i\sqrt{4-\beta^{2}}}{2}\right)^{n}$$

$$= \left(-1\right)^{n} \left[\left(\frac{1}{2} + \frac{i\beta}{2\sqrt{4-\beta^{2}}}\right) \left(\frac{\beta - i\sqrt{4-\beta^{2}}}{2}\right)^{n} + \left(\frac{1}{2} - \frac{i\beta}{2\sqrt{4-\beta^{2}}}\right) \left(\frac{\beta + i\sqrt{4-\beta^{2}}}{2}\right)^{n}\right]$$

$$= \left(-1\right)^{n} a_{n}(\beta) = \left(-1\right)^{n} a_{n}(\mu).$$

(ii) Note that $\beta = 2(1 - \mu)$. Then

$$\beta = \begin{cases} 2 & \text{if } \mu = 0 \\ -2 & \text{if } \mu = 2 \end{cases}$$

Replacing β with 2 and -2 respectively in Eq. (A.5), we get Eqs. (7) and (8).

(iii) Firstly, we prove $a_n(\mu)$ is a polynomial of degree *n* while $(n \ge 0)$ by mathematical induction.

Step 1: for n = 0 and n = 1,

$$a_0(\mu) = 1, \tag{A.6}$$

$$a_1(\mu) = -2\mu + 2. \tag{A.7}$$

It is obvious that $a_0(\mu)$ and $a_1(\mu)$ are polynomials of degree 0 and 1 respectively.

Step 2: for any $n \ge 2$, we assume $a_{n-1}(\mu)$ and $a_{n-2}(\mu)$ are polynomials of degree n-1 and n-2 respectively.

$$a_2(\mu) = 4\mu^2 - 8\mu + 3. \tag{A.8}$$

Note that $a_n(\mu) = 2(1-\mu)a_{n-1}(\mu) - a_{n-2}(\mu)$. Then, the degree of polynomial $a_n(\mu)$ is just 1 plus the degree of polynomial $a_{n-1}(\mu)$. Therefore $a_n(\mu)$ is a polynomial of degree n, which can be expressed as

$$a_n(\mu) = \sum_{i=0}^n a_n^{(i)} \mu^i,$$
 (A.9)

where $a_n^{(i)}$ is the coefficient of μ^i .

Replacing $a_n(\mu)$, $a_{n-1}(\mu)$ and $a_{n-2}(\mu)$ from Eq. (A.9) in recursive equation $a_n(\mu) = 2(1-\mu)a_{n-1}(\mu) - a_{n-2}(\mu)$, we find $\{a_n^{(0)}\}$ satisfies the following recurrence relation

$$a_n^{(0)} = 2a_{n-1}^{(0)} - a_{n-2}^{(0)}, \tag{A.10}$$

with initial conditions $a_0^{(0)} = 1$ and $a_1^{(0)} = 2$.

By solving the recursive equation as shown in Eq. (A.10), we obtain Eq. (9).

Similarly, we find $\{a_n^{(1)}\}$ satisfies the following recurrence relation

$$a_n^{(1)} = -2a_{n-1}^{(0)} + 2a_{n-1}^{(1)} - a_{n-2}^{(1)} = -2n + 2a_{n-1}^{(1)} - a_{n-2}^{(1)},$$
(A.11)

with initial conditions $a_1^{(1)} = -2$, $a_2^{(1)} = -8$; and $\{a_n^{(n)}\}$ satisfies the following recurrence relation

$$a_n^{(n)} = -2a_{n-1}^{(n-1)}, \tag{A.12}$$

with initial condition $a_0^{(0)} = 1$. By solving the recurrence relation as shown in Eqs. (A.11) and (A.12), we obtain Eqs. (10) and (11).

(iv) Here, we prove Eq. (12) by mathematical induction and Eq. (13) can also be obtained similarly.

Step 1: Eq.(12) is true for n = 2 and n = 3.

$$a_2 = \beta a_1 - a_0 = a_1^2 - a_0^2 = (a_1 + a_0)(a_1 - a_0),$$

and

$$a_3 = \beta a_2 - a_1 = (a_2 - a_0) a_1.$$

Step 2: assume Eq.(12) is true for n = 2t and n = 2t + 1 (t is an arbitrary nonnegative integer), i.e.,

$$a_{2t} = (a_t - a_{t-1}) (a_t + a_{t-1}),$$
$$a_{2t+1} = (a_{t+1} - a_{t-1}) a_t.$$

Then,

$$a_{2t+2} = \beta a_{2t+1} - a_{2t}$$

= $\beta (a_{t+1} - a_{t-1}) a_t - (a_t^2 - a_{t-1}^2)$
= $\beta a_{t+1} a_t - \beta a_{t-1} a_t - a_t^2 + a_{t-1}^2$
= $\beta a_{t+1} a_t - a_t^2 - a_{t-1} (\beta a_t - a_{t-1})$
= $\beta a_{t+1} a_t - a_t^2 - a_{t-1} a_{t+1}$
= $a_{t+1} (\beta a_t - a_{t-1}) - a_t^2$
= $a_{t+1}^2 - a_t^2$
= $(a_{t+1} + a_t)(a_{t+1} - a_t)$,

and

$$a_{2t+3} = \beta a_{2t+2} - a_{2t+1}$$

= $\beta \left(a_{t+1}^2 - a_t^2 \right) - (a_{t+1} - a_{t-1}) a_t$
= $\beta a_{t+1}^2 - \beta a_t^2 - a_{t+1} a_t + a_{t-1} a_t$
= $a_{t+1} \left(\beta a_{t+1} - a_t \right) - a_t \left(\beta a_t - a_{t-1} \right)$
= $a_{t+1} a_{t+2} - a_t a_{t+1}$
= $a_{t+1} \left(a_{t+2} - a_t \right)$.

Therefore Eq.(12) is true for n = 2t + 2 and n = 2t + 3. Thus Eq.(12) is true for any $n \ge 2$.

Similarly, we can also prove Eq.(13) by mathematical induction.

B Proof of Lemma 3.7

(i) Let $n \ (n \ge 3)$ be an arbitrary odd number and μ be an arbitrary root of equation $a_{\frac{n-1}{2}}(\mu) - a_{\frac{n-1}{2}-1}(\mu) = 0$. Therefore,

$$a_{\frac{n-1}{2}-1}(\mu) = a_{\frac{n-1}{2}}(\mu) = 2(1-\mu)a_{\frac{n-1}{2}-1}(\mu) - a_{\frac{n-1}{2}-2}(\mu).$$
(B.1)

Then

$$[2(1-\mu)-1]a_{\frac{n-1}{2}-1}(\mu) = a_{\frac{n-1}{2}-2}(\mu).$$
(B.2)

Replacing $a_{\frac{n-1}{2}+1}(\mu)$ and $a_{\frac{n-1}{2}}(\mu)$ with $2(1-\mu)a_{\frac{n-1}{2}}(\mu) - a_{\frac{n-1}{2}-1}(\mu)$ and $2(1-\mu)a_{\frac{n-1}{2}-1}(\mu) - a_{\frac{n-1}{2}-2}(\mu)$ in $1 - \frac{a_{\frac{n-1}{2}+1}-a_{\frac{n-1}{2}-1}}{a_{\frac{n-1}{2}}-a_{\frac{n-1}{2}-2}}$ respectively, we obtain

$$1 - \frac{a_{\frac{n-1}{2}+1}(\mu) - a_{\frac{n-1}{2}-1}(\mu)}{a_{\frac{n-1}{2}}(\mu) - a_{\frac{n-1}{2}-2}(\mu)} = 1 - \frac{2(1-\mu)a_{\frac{n-1}{2}}(\mu) - a_{\frac{n-1}{2}-1}(\mu) - a_{\frac{n-1}{2}-1}(\mu)}{2(1-\mu)a_{\frac{n-1}{2}-1}(\mu) - a_{\frac{n-1}{2}-2}(\mu) - a_{\frac{n-1}{2}-2}(\mu)}$$
$$= 1 - \frac{[2(1-\mu)-2]a_{\frac{n-1}{2}-1}(\mu)}{\{2(1-\mu)-2[2(1-\mu)-1]\}a_{\frac{n-1}{2}-1}(\mu)}$$
$$= 2.$$
(B.3)

One can easy obtain from Eq.(12) that

$$a_{n-1} = \left(a_{\frac{n-1}{2}} - a_{\frac{n-1}{2}-1}\right) \left(a_{\frac{n-1}{2}} + a_{\frac{n-1}{2}-1}\right) = 0.$$
(B.4)

By using Eqs.(12) and (13), we have

$$a_{n-2}(\mu) = a_{\frac{n-2-1}{2}}(\mu) \left(a_{\frac{n-2-1}{2}+1}(\mu) - a_{\frac{n-2-1}{2}-1}(\mu) \right)$$

= $a_{\frac{n-1}{2}-1}(\mu) \left(a_{\frac{n-1}{2}}(\mu) - a_{\frac{n-1}{2}-2}(\mu) \right)$
= $a_{\frac{n-1}{2}}(\mu) \left(a_{\frac{n-1}{2}}(\mu) - a_{\frac{n-1}{2}-2}(\mu) \right)$
= $1 + a_{n-1}(\mu)$
= $1.$ (B.5)

(ii) Let μ be an arbitrary root of equation $a_{\frac{n-1}{2}}(\mu) + a_{\frac{n-1}{2}-1}(\mu) = 0$. Then

$$-a_{\frac{n-1}{2}-1}(\mu) = a_{\frac{n-1}{2}}(\mu) = 2(1-\mu)a_{\frac{n-1}{2}-1}(\mu) - a_{\frac{n-1}{2}-2}(\mu)$$
(B.6)

Thus

$$[2(1-\mu)+1]a_{\frac{n-1}{2}-1}(\mu) = a_{\frac{n-1}{2}-2}(\mu).$$
(B.7)

Replacing
$$a_{\frac{n-1}{2}+1}(\mu)$$
 and $a_{\frac{n-1}{2}}(\mu)$ with $2(1-\mu)a_{\frac{n-1}{2}}(\mu) - a_{\frac{n-1}{2}-1}(\mu)$ and
 $2(1-\mu)a_{\frac{n-1}{2}-1}(\mu) - a_{\frac{n-1}{2}-2}(\mu)$ in $1 - \frac{a_{\frac{n-1}{2}+1}-a_{\frac{n-1}{2}-1}}{a_{\frac{n-1}{2}}-a_{\frac{n-1}{2}-2}}$, we obtain
 $1 - \frac{a_{\frac{n-1}{2}+1}(\mu) - a_{\frac{n-1}{2}-1}(\mu)}{a_{\frac{n-1}{2}}(\mu) - a_{\frac{n-1}{2}-2}(\mu)} = 1 - \frac{2(1-\mu)a_{\frac{n-1}{2}}(\mu) - a_{\frac{n-1}{2}-1}(\mu) - a_{\frac{n-1}{2}-1}(\mu)}{2(1-\mu)a_{\frac{n-1}{2}-1}(\mu) - a_{\frac{n-1}{2}-2}(\mu) - a_{\frac{n-1}{2}-2}(\mu)}$
 $= 1 - \frac{-[2(1-\mu)+2]a_{\frac{n-1}{2}-1}(\mu)}{\{2(1-\mu)-2[2(1-\mu)+1]\}a_{\frac{n-1}{2}-1}(\mu)}$
 $= 0$ (B.8)

Note that $a_{n-1} = \left(a_{\frac{n-1}{2}} - a_{\frac{n-1}{2}-1}\right) \left(a_{\frac{n-1}{2}} + a_{\frac{n-1}{2}-1}\right) = 0$. By using Eqs.(12) and (13), we have

$$a_{n-2}(\mu) = a_{\frac{n-2-1}{2}}(\mu) \left(a_{\frac{n-2-1}{2}+1}(\mu) - a_{\frac{n-2-1}{2}-1}(\mu) \right)$$

= $a_{\frac{n-1}{2}-1}(\mu) \left(a_{\frac{n-1}{2}}(\mu) - a_{\frac{n-1}{2}-2}(\mu) \right)$
= $-a_{\frac{n-1}{2}}(\mu) \left(a_{\frac{n-1}{2}}(\mu) - a_{\frac{n-1}{2}-2}(\mu) \right)$
= $-(1 + a_{n-1}(\mu))$
= -1 (B.9)

(iii) Let μ be an arbitrary root of equation $a_{\frac{n-1}{2}}(\mu) = 0$. Then

$$0 = a_{\frac{n-1}{2}}(\mu) = 2(1-\mu)a_{\frac{n-1}{2}-1}(\mu) - a_{\frac{n-1}{2}-2}(\mu), \qquad (B.10)$$

which yields

$$2(1-\mu) a_{\frac{n-1}{2}-1}(\mu) = a_{\frac{n-1}{2}-2}(\mu).$$

Replacing $a_{\frac{n-1}{2}+1}(\mu)$ and $a_{\frac{n-1}{2}-2}(\mu)$ with $2(1-\mu)a_{\frac{n-1}{2}}(\mu) - a_{\frac{n-1}{2}-1}(\mu)$ and $2(1-\mu)a_{\frac{n-1}{2}-1}(\mu)$ in $1 - \frac{a_{\frac{n-1}{2}+1}-a_{\frac{n-1}{2}-1}}{a_{\frac{n-1}{2}-2}}$ respectively, we obtain

$$1 - \frac{a_{\frac{n-1}{2}+1}(\mu) - a_{\frac{n-1}{2}-1}(\mu)}{a_{\frac{n-1}{2}}(\mu) - a_{\frac{n-1}{2}-2}(\mu)} = 1 - \frac{2(1-\mu)a_{\frac{n-1}{2}}(\mu) - a_{\frac{n-1}{2}-1}(\mu) - a_{\frac{n-1}{2}-1}(\mu)}{-2(1-\mu)a_{\frac{n-1}{2}-1}(\mu)} = 1 - \frac{1}{1-\mu}.$$
(B.11)

Since $\mu \in (0, 2)$, we have $1 - \frac{1}{1-\mu} \in (-\infty, 0) \cup (2, +\infty)$.

It is easy to verify from Eq.(12) that

$$a_n(\mu) = \left(a_{\frac{n-1}{2}+1}(\mu) - a_{\frac{n-1}{2}-1}(\mu)\right)a_{\frac{n-1}{2}}(\mu) = 0.$$

Thus

$$1 = 1 + a_{n}(\mu)$$

= $\left(a_{\frac{n+1}{2}-1}(\mu) - a_{\frac{n+1}{2}-2}(\mu)\right) \left(a_{\frac{n+1}{2}}(\mu) + a_{\frac{n+1}{2}-1}(\mu)\right)$
= $\left(a_{\frac{n-1}{2}}(\mu) - a_{\frac{n-1}{2}-1}(\mu)\right) \left[2(1-\mu)a_{\frac{n-1}{2}}(\mu) - a_{\frac{n-1}{2}-1}(\mu) + a_{\frac{n-1}{2}}(\mu)\right]$
= $\left(a_{\frac{n-1}{2}-1}(\mu)\right)^{2}$. (B.12)

Therefore

$$a_{n-1}(\mu) = \left(a_{\frac{n-1}{2}}(\mu) - a_{\frac{n-1}{2}-1}(\mu)\right) \left(a_{\frac{n-1}{2}}(\mu) + a_{\frac{n-1}{2}-1}(\mu)\right)$$
$$= -\left(a_{\frac{n-1}{2}-1}(\mu)\right)^{2}$$
$$= -1.$$
(B.13)

Replacing $a_n(\mu)$ and $a_{n-1}(\mu)$ with 0 and -1 in $a_n(\mu) = 2(1-\mu)a_{n-1}(\mu) - a_{n-2}(\mu)$, we have

$$a_{n-2}(\mu) = -2(1-\mu).$$
 (B.14)

C Proof of Lemma 3.8

(i) Let $n \ (n \ge 2)$ be an arbitrary even number and μ be an arbitrary root of equation $a_{\frac{n}{2}}(\mu) - a_{\frac{n-2}{2}}(\mu) = 0$. Then $a_{\frac{n}{2}}(\mu) = a_{\frac{n-2}{2}}(\mu) = 0$ and

$$a_{n-1}(\mu) = \left(a_{\frac{n}{2}}(\mu) - a_{\frac{n}{2}-2}(\mu)\right)a_{\frac{n}{2}-1}(\mu) = 0.$$
(C.1)

Replacing $a_{\frac{n}{2}-2}(\mu)$ with $a_{\frac{n}{2}}(\mu)$ in $1 - \frac{a_{\frac{n}{2}+1}-a_{\frac{n}{2}-1}}{a_{\frac{n}{2}-1}-a_{\frac{n}{2}-2}}$, we obtain

$$1 - \frac{a_{\frac{n}{2}}(\mu) - a_{\frac{n}{2}-1}(\mu)}{a_{\frac{n}{2}-1}(\mu) - a_{\frac{n}{2}-2}(\mu)} = 1 - \frac{a_{\frac{n}{2}}(\mu) - a_{\frac{n}{2}-1}(\mu)}{a_{\frac{n}{2}-1}(\mu) - a_{\frac{n}{2}}(\mu)} = 2.$$
 (C.2)

By using Eqs.(12) and (13), we have

$$a_{n-2}(\mu) = \left(a_{\frac{n}{2}-1}(\mu) - a_{\frac{n}{2}-2}(\mu)\right) \left(a_{\frac{n}{2}-1}(\mu) + a_{\frac{n}{2}-2}(\mu)\right)$$

= $\left(a_{\frac{n}{2}-1}(\mu) - a_{\frac{n}{2}-2}(\mu)\right) \left(a_{\frac{n}{2}-1}(\mu) + a_{\frac{n}{2}}(\mu)\right)$
= $1 + a_{n-1}(\mu)$
= $1.$ (C.3)

(ii) Let μ be an arbitrary root of equation $a_{\frac{n}{2}-1}(\mu) = 0$. Replacing $a_{\frac{n}{2}}(\mu)$ with $2(1-\mu)a_{\frac{n}{2}-1}(\mu) - a_{\frac{n}{2}-2}(\mu)$ in $1 - \frac{a_{\frac{n}{2}-a_{\frac{n}{2}-1}}}{a_{\frac{n}{2}-1}-a_{\frac{n}{2}-2}}$, we obtain

$$1 - \frac{a_{\frac{n}{2}}(\mu) - a_{\frac{n}{2}-1}(\mu)}{a_{\frac{n}{2}-1}(\mu) - a_{\frac{n}{2}-2}(\mu)}$$

=1 - $\frac{2(1-\mu)a_{\frac{n}{2}-1}(\mu) - a_{\frac{n}{2}-2}(\mu) - a_{\frac{n}{2}-1}(\mu)}{a_{\frac{n}{2}-1}(\mu) - a_{\frac{n}{2}-2}(\mu)}$
=1 - $\frac{-a_{\frac{n}{2}-2}(\mu)}{-a_{\frac{n}{2}-2}(\mu)}$
=0. (C.4)

By using Eqs.(12) and (13) and noticing that $a_{n-1}(\mu) = 0$, we have

$$a_{n-2}(\mu) = \left(a_{\frac{n}{2}-1}(\mu) - a_{\frac{n}{2}-2}(\mu)\right) \left(a_{\frac{n}{2}-1}(\mu) + a_{\frac{n}{2}-2}(\mu)\right)$$
$$= -\left(a_{\frac{n}{2}-1}(\mu) - a_{\frac{n}{2}-2}(\mu)\right) \left(a_{\frac{n}{2}-1}(\mu) + a_{\frac{n}{2}}(\mu)\right)$$
$$= -\left(1 + a_{n-1}(\mu)\right)$$
$$= -1.$$
(C.5)

(iii) Let μ be an arbitrary root of equation of equation $a_{\frac{n}{2}}(\mu) + a_{\frac{n}{2}-1}(\mu) = 0$. We have

$$-a_{\frac{n}{2}-1}(\mu) = a_{\frac{n}{2}}(\mu) = 2(1-\mu)a_{\frac{n}{2}-1}(\mu) - a_{\frac{n}{2}-2}(\mu), \qquad (C.6)$$

which yields

$$a_{\frac{n}{2}-2}(\mu) = [2(1-\mu)+1]a_{\frac{n}{2}-1}(\mu).$$

Replacing $a_{\frac{n}{2}}(\mu)$ and $a_{\frac{n}{2}-2}(\mu)$ with $a_{\frac{n}{2}-1}(\mu)$ and $[2(1-\mu)+1]a_{\frac{n}{2}-1}(\mu)$ in $1 - \frac{a_{\frac{n}{2}} - a_{\frac{n}{2}-1}}{a_{\frac{n}{2}-1} - a_{\frac{n}{2}-2}}$ respectively, we have

$$1 - \frac{a_{\frac{n}{2}}(\mu) - a_{\frac{n}{2}-1}(\mu)}{a_{\frac{n}{2}-1}(\mu) - a_{\frac{n}{2}-2}(\mu)} = 1 - \frac{-2a_{\frac{n}{2}-1}(\mu)}{a_{\frac{n}{2}-1}(\mu) - [2(1-\mu)+1]a_{\frac{n}{2}-1}(\mu)}$$
$$= 1 - \frac{1}{1-\mu}.$$
 (C.7)

Since $\mu \in (0, 2)$, we have $1 - \frac{1}{1-\mu} \in (-\infty, 0) \cup (2, +\infty)$.

It is easy to verify

$$a_n(\mu) = \left[a_{\frac{n}{2}}(\mu) - a_{\frac{n}{2}-1}(\mu)\right] \left[a_{\frac{n}{2}}(\mu) + a_{\frac{n}{2}-1}(\mu)\right] = 0.$$

Then

$$a_{n-1}(\mu) = \left(a_{\frac{n}{2}}(\mu) - a_{\frac{n}{2}-2}(\mu)\right) a_{\frac{n}{2}-1}(\mu)$$

= $-\left(a_{\frac{n}{2}}(\mu) - a_{\frac{n}{2}-2}(\mu)\right) a_{\frac{n}{2}}(\mu)$
= $-(1 + a_n(\mu))$
= $-1.$ (C.8)

Replacing $a_n(\mu)$ and $a_{n-1}(\mu)$ with 0 and -1 in $a_n(\mu) = 2(1-\mu)a_{n-1}(\mu) - a_{n-2}(\mu)$ respectively, we have

$$a_{n-2}(\mu) = -2(1-\mu).$$
 (C.9)

The proof is completed.