On integrability of a third-order complex nonlinear wave equation

SERGEI SAKOVICH

Institute of Physics, National Academy of Sciences of Belarus sergsako@gmail.com

Abstract

We show that the new third-order complex nonlinear wave equation, introduced recently by Müller-Hoissen [arXiv:2202.04512], does not pass the Painlevé test for integrability. We find two reductions of this equation, one integrable and one non-integrable, whose solutions jointly cover all solutions of the original equation.

1 Introduction

The following new third-order complex nonlinear wave equation was introduced recently by Müller-Hoissen [1]:

$$\left(\frac{f_{xt}}{f}\right)_t + 2\left(ff^*\right)_x = 0,\tag{1}$$

where f(x,t) is a complex function of two real variables, subscripts denote respective derivatives, and the asterisk stands for the complex conjugate. This nonlinear wave equation (1) was called a completely integrable partial differential equation in [1], and its multi-soliton solutions were obtained there.

Let us note, however, that this new nonlinear equation was studied in [1] not in its original form (1) but in the form of the following system of two equations:

$$a_t = (ff^*)_x, \qquad f_{xt} + 2af = 0,$$
(2)

where a(x,t) is a real function. Being the first "negative flow" of the nonlinear Schrödinger equation's hierarchy, this nonlinear system (2) is integrable and possesses multi-soliton solutions. The integrable system (2) is obviously a reduction of the new nonlinear wave equation (1), in the sense that all solutions of (2) (except for f = 0, of course) are solutions of (1) as well, because (1) follows from (2) through elimination of the dependent variable a. In other words, the new equation (1) possesses multi-soliton solutions because it possesses an integrable reduction. But is the new equation (1) integrable itself? No Lax pair of (1) is known.

In the present paper, we study the integrability of the new equation (1) by means of the singularity analysis (a.k.a. the Painlevé analysis) in its version for partial differential equations [2, 3]. In Section 2, we show that the nonlinear equation (1) does not pass the Painlevé test for integrability. In Section 3, we find two reductions of the nonlinear equation (1), one integrable and one non-integrable, whose solutions jointly cover all solutions of the original equation. Section 4 contains concluding remarks.

2 Singularity analysis

In our experience, the Painlevé analysis is a reliable and convenient method to test the integrability of nonlinear wave equations, including high-order, nonevolutionary, multi-component and high-dimensional ones [4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19]. The reliability of the Painlevé test for integrability has been empirically verified by the analysis of wide classes of nonlinear equations, such as fifth-order Korteweg–de Vries type equations [20], bilinear equations [21], coupled Korteweg–de Vries equations [22, 23, 24, 25], coupled higher-order nonlinear Schrödinger equations [26], generalized Ito equations [27], sixth-order nonlinear wave equations [28], seventh-order Korteweg–de Vries type equations [29], etc.

To start the singularity analysis of the nonlinear equation (1), we rewrite this equation and its complex conjugate

$$\left(\frac{f_{xt}^*}{f^*}\right)_t + 2\left(ff^*\right)_x = 0\tag{3}$$

as the following system of two polynomial equations:

$$ff_{xtt} - f_t f_{xt} + 2f^2 (fg)_x = 0,$$

$$gg_{xtt} - g_t g_{xt} + 2g^2 (fg)_x = 0,$$
(4)

where g stands for f^* . From now on, we consider f(x,t) and g(x,t) as two mutually independent complex functions of two complex variables x and t.

A singularity manifold $\phi(x,t) = 0$ is non-characteristic for this system (4) if $\phi_x \phi_t \neq 0$, and we take $\phi_t = 1$ without loss of generality,

$$\phi = t + \psi(x), \qquad \psi_x \neq 0, \tag{5}$$

where $\psi(x)$ is arbitrary. Substitution of the expansions

$$f = f_0(x)\phi^p + \dots + f_r(x)\phi^{p+r} + \dots,$$

$$g = g_0(x)\phi^q + \dots + g_r(x)\phi^{q+r} + \dots$$
(6)

to the nonlinear system (4) determines the leading exponents p and q (i.e., the dominant behavior of solutions f and g near the singularity manifold $\phi = 0$) and the resonances r (i.e., the positions, where arbitrary functions can enter the expansions). In this way, we find the following one branch to study further:

$$p = q = -1, \qquad r = -1, 0, 2, 2, 3, 4,$$
(7)

where r = -1 corresponds to the arbitrariness of $\psi(x)$ in (5).

For this branch (7), we try to represent the general solution of (4) by the Laurent type expansions

$$f = \sum_{i=0}^{\infty} f_i(x)\phi^{i-1}, \qquad g = \sum_{i=0}^{\infty} g_i(x)\phi^{i-1}, \tag{8}$$

with ϕ given by (5). We substitute (8) to (4), collect terms with ϕ^{n-5} , for $n = 0, 1, 2, 3, \ldots$ separately, and obtain in this way the following.

For n = 0, where we have a single resonance due to (7), we get the expression for g_0 ,

$$g_0 = -\frac{1}{f_0},$$
(9)

while the function $f_0(x)$ remains arbitrary.

For n = 1, there is no resonance, and we get the expressions

$$f_1 = -\frac{f_0'}{2\psi'}, \qquad g_1 = -\frac{f_0'}{2f_0^2\psi'},$$
(10)

where the prime denotes the derivative with respect to x.

For n = 2, we have a double resonance, the functions $f_2(x)$ and $g_2(x)$ remain arbitrary, the compatibility conditions are identically satisfied by (9) and (10), and no restrictions for $\psi(x)$ and $f_0(x)$ appear.

For n = 3, where we have a single resonance, we get the expression for g_3 ,

$$g_{3} = \frac{f_{3}}{f_{0}^{2}} - \frac{g_{2}f_{0}'}{2f_{0}\psi'} + \frac{(f_{0}')^{3}}{2f_{0}^{4}(\psi')^{3}} + \frac{f_{2}'}{2f_{0}^{2}\psi'} - \frac{g_{2}'}{\psi'} + \frac{(f_{0}')^{2}\psi''}{2f_{0}^{3}(\psi')^{4}} - \frac{f_{0}'f_{0}''}{2f_{0}^{3}(\psi')^{3}},$$
(11)

the function $f_3(x)$ remains arbitrary, but the following nontrivial compatibility condition appears:

$$\left(\frac{f_2}{f_0} + f_0 g_2\right)' = 0. \tag{12}$$

The fact that we have got a nontrivial compatibility condition at a resonance means that we have to modify our expansions of solutions by additional logarithmic terms. Consequently, the nonlinear equation (1) does not pass the Painlevé test for integrability.

3 Two reductions

Let us return to the nonlinear equation (1) and consider it together with its complex conjugate (3). From these two equations, we get the relation

$$\left(\frac{f_{xt}}{f} - \frac{f_{xt}^*}{f^*}\right)_t = 0 \tag{13}$$

satisfied by any solution of (1). We introduce two real functions of two real variables, u(x, t) and v(x, t), such that

$$\frac{f_{xt}}{f} = u(x,t) + \mathrm{i}v(x,t),\tag{14}$$

where $i^2 = -1$. Then, it follows from (13) and (14) that $v_t = 0$, and (14) takes the form

$$\frac{f_{xt}}{f} = u(x,t) + \mathrm{i}w(x),\tag{15}$$

where w(x) is a real function of one real variable.

It was pointed out in [1] that the nonlinear equation (1) is invariant under the transformation

$$x \mapsto h(x),\tag{16}$$

where h(x) is an arbitrary function. In other words, if a function f(x,t) is a solution of the nonlinear equation (1), then f(h(x),t) is also a solution of (1), for any function h(x).

Under the transformation (16), the function w(x) in the relation (15) changes in the following way:

$$w(x) \mapsto h'(x) w(h(x)), \tag{17}$$

where the prime denotes the derivative. Therefore, for any solution f(x, t) which corresponds to any nonzero function w(x) in (15), the integral

$$x = \int w(h) \, dh \tag{18}$$

determines (at least, locally) the function h(x) of the transformation (16), such that the transformed solution f(h(x), t) corresponds to w = 1. Consequently, we can use the relation (15) in the form

$$\frac{f_{xt}}{f} = u(x,t) + ik, \qquad k = 0, 1,$$
(19)

provided that, in the case of k = 1, every single solution f(x, t) represents the whole its equivalence class f(h(x), t) with any function h(x).

The original nonlinear equation (1) together with the relation (19) give us the following system of two equations:

$$f_{xt} - (u + ik)f = 0,$$

$$u_t + 2(ff^*)_x = 0, \qquad k = 0, 1,$$
(20)

where f(x,t) is a complex function of two real variables, but u(x,t) is a real function of two real variables. The two cases of the system (20), with k = 0 and with k = 1, are two distinct reductions of the original equation (1). Solutions of these two reductions jointly cover all solutions of (1), provided that every solution f(x,t) of the system (20) with k = 1 is generalized as f(h(x),t), with any function h(x). (Note that the system (20) with k = 1 is not invariant under the transformation (16), whereas the system (20) with k = 0 is invariant.) The system (20) with k = 0 is just the system (2) studied in [1], the correspondence being u = -2a. The system (20) with k = 1 is something new.

Let us apply the Painlevé test for integrability to the system (20). We combine (20) with its complex conjugate, denote f^* as g, treat f and g as mutually independent, complexify all variables, and consider the following system of three equations:

$$f_{xt} - (u + ik)f = 0,$$

$$g_{xt} - (u - ik)g = 0,$$

$$u_t + 2(fg)_x = 0, \qquad k = 0, 1,$$
(21)

where f(x,t), g(x,t) and u(x,t) are complex function of two complex variables.

We find from (21) that, near any non-characteristic singularity manifold $\phi(x,t) = 0$ with ϕ given by (5) and arbitrary $\psi(x)$, the leading exponents of f, g and u are -1, -1 and -2, respectively, and the resonances are

$$r = -1, 0, 2, 3, 4, \tag{22}$$

where r = -1 corresponds to the arbitrariness of $\psi(x)$ in (5). Then we substitute the expansions

$$f = \sum_{i=0}^{\infty} f_i(x)\phi^{i-1}, \qquad g = \sum_{i=0}^{\infty} g_i(x)\phi^{i-1}, u = \sum_{i=0}^{\infty} u_i(x)\phi^{i-2}$$
(23)

to the system (21), collect terms with ϕ^{n-3} , for $n = 0, 1, 2, 3, \ldots$ separately, and obtain in this way the following.

For n = 0, we get the expressions

$$g_0 = -\frac{1}{f_0}, \qquad u_0 = 2\psi',$$
(24)

the function $f_0(x)$ remains arbitrary, and no nontrivial compatibility condition appears at this resonance.

For n = 1, where we have no resonance, we get the expressions

$$f_1 = -\frac{f_0'}{2\psi'}, \qquad g_1 = -\frac{f_0'}{2f_0^2\psi'}, \qquad u_1 = 0.$$
 (25)

For n = 2, we get the expressions

$$f_2 = -\frac{f_0(u_2 + ik)}{2\psi'}, \qquad g_2 = \frac{u_2 - ik}{2f_0\psi'},$$
(26)

the function $u_2(x)$ remains arbitrary, and no nontrivial compatibility condition appears at this resonance.

For n = 3, we get the expressions

$$u_{3} = -\frac{u_{2}'}{2\psi'} + \frac{(u_{2} + ik)\psi''}{2(\psi')^{2}},$$

$$g_{3} = \frac{f_{3}}{f_{0}^{2}} - \frac{ikf_{0}'}{2f_{0}^{2}(\psi')^{2}} + \frac{(f_{0}')^{3}}{2f_{0}^{4}(\psi')^{3}} - \frac{3u_{2}'}{4f_{0}(\psi')^{2}}$$

$$-\frac{ik\psi''}{4f_{0}(\psi')^{3}} + \frac{3u_{2}\psi''}{4f_{0}(\psi')^{3}} + \frac{(f_{0}')^{2}\psi''}{2f_{0}^{3}(\psi')^{4}} - \frac{f_{0}'f_{0}''}{2f_{0}^{3}(\psi')^{3}},$$
(27)

and the function $f_3(x)$ remains arbitrary. However, the following compatibility condition appears at this resonance:

$$k\psi'' = 0, (28)$$

which is not satisfied identically if k = 1. Consequently, the system (20) with k = 1 does not pass the Painlevé test for integrability.

If k = 0, the compatibility condition (28) is satisfied identically, and we continue computations. For n = 4, we obtain explicit expressions for u_4 and g_4 (cumbersome ones, therefore omitted here), the function $f_4(x)$ remains arbitrary, and no nontrivial compatibility condition appears at this resonance. Consequently, the system (20) with k = 0 has passed the Painlevé test for integrability (and this is an expected result, of course).

4 Conclusion

In this paper, we studied the integrability of the third-order complex nonlinear wave equation (1), introduced recently in [1]. We used the Painlevé test for integrability and obtained the following results.

The nonlinear equation (1) fails to pass the Painlevé test for integrability. Therefore we believe that one cannot find any good Lax pair for this equation.

The nonlinear equation (1) possesses two reductions, one integrable and one non-integrable, whose solutions jointly cover all solutions of this equation.

The integrable reduction is the system (20) with k = 0, studied in [1] in the form (2). This reduction corresponds to the condition $f_{xt}/f = f_{xt}^*/f^*$ imposed on solutions of (1). This integrable reduction is just what provides the original equation (1) with multi-soliton solutions.

The non-integrable reduction is the system (20) with k = 1. It corresponds to the condition $f_{xt}/f \neq f_{xt}^*/f^*$ imposed on solutions of (1). Solutions of this non-integrable reduction must be generalized by the arbitrary coordinate transformation (16), in order to cover all those solutions of (1) which are not solutions of the integrable reduction.

It seems to be an interesting future problem to find any explicit solutions of the new (non-integrable) system (20) with k = 1. We believe that there are no multi-soliton solutions, and that the travelling waves (if there are any) interact inelastically.

References

- [1] F. Müller-Hoissen, A vectorial binary Darboux transformation for the first member of the negative part of the AKNS hierarchy, arXiv:2202.04512v3.
- [2] J. Weiss, M. Tabor, G. Carnevale, The Painlevé property for partial differential equations, J. Math. Phys. 24 (1983) 522–526.
- [3] M. Tabor, Chaos and Integrability in Nonlinear Dynamics: An Introduction, Wiley, New York, 1989.
- [4] S.Yu. Sakovich, Painlevé analysis and Bäcklund transformations of Doktorov– Vlasov equations, J. Phys. A: Math. Gen. 27 (1994) L33–L38.

- [5] S.Yu. Sakovich, Painlevé analysis of new soliton equations by Hu, J. Phys. A: Math. Gen. 27 (1994) L503–L505.
- [6] S.Yu. Sakovich, On zero-curvature representations of evolution equations, J. Phys. A: Math. Gen. 28 (1995) 2861–2869.
- [7] S.Yu. Sakovich, Painlevé analysis of a higher-order nonlinear Schrödinger equation, J. Phys. Soc. Jpn. 66 (1997) 2527–2529.
- [8] S.Yu. Sakovich, On integrability of a (2+1)-dimensional perturbed KdV equation, J. Nonlinear Math. Phys. 5 (1998) 230–233; arXiv:solv-int/9805012.
- [9] A. Karasu-Kalkanlı, S.Yu. Sakovich, Bäcklund transformation and special solutions for the Drinfeld–Sokolov–Satsuma–Hirota system of coupled equations, J. Phys. A: Math. Gen. 34 (2001) 7355–7358; arXiv:nlin/0102001.
- [10] A. Karasu-Kalkanlı, S.Yu. Sakovich, İ. Yurduşen, Integrability of Kersten– Krasil'shchik coupled KdV–mKdV equations: singularity analysis and Lax pair, J. Math. Phys. 44 (2003) 1703–1708; arXiv:nlin/0206046.
- [11] A. Karasu-Kalkanh, S. Sakovich, Singularity analysis of a spherical Kadomtsev– Petviashvili equation, J. Phys. Soc. Jpn. 74 (2005) 505–507; arXiv:nlin/0404037.
- [12] S. Sakovich, Enlarged spectral problems and nonintegrability, Phys. Lett. A 345 (2005) 63-68; arXiv:nlin/0504037.
- [13] S. Sakovich, Integrability of the vector short pulse equation, J. Phys. Soc. Jpn. 77 (2008) 123001; arXiv:0801.3179.
- [14] S. Sakovich, Singularity analysis and integrability of a Burgers-type system of Foursov, Symmetry Integr. Geom. Methods Appl. 7 (2011) 002; arXiv:1010.5709.
- S. Sakovich, On two aspects of the Painlevé analysis, Int. J. Analysis 2013 (2013) 172813; arXiv:solv-int/9909027.
- [16] S. Sakovich, Integrability study of a four-dimensional eighth-order nonlinear wave equation, Nonlinear Phenom. Complex Syst. 20 (2017) 267–271; arXiv:1607.08408.
- [17] S. Sakovich, On a new avatar of the sine-Gordon equation, Nonlinear Phenom. Complex Syst. 21 (2018) 62–68; arXiv:1703.04678.
- [18] S. Sakovich, A new Painlevé-integrable equation possessing KdV-type solitons, Nonlinear Phenom. Complex Syst. 22 (2019) 299–304; arXiv:1907.01324.
- [19] S. Sakovich, Integrability of one bilinear equation: singularity analysis and dimension, Nonlinear Phenom. Complex Syst. 24 (2021) 311–316; arXiv:2109.02073.
- [20] H. Harada, S. Oishi, A new approach to completely integrable partial differential equations by means of the singularity analysis, J. Phys. Soc. Jpn. 54 (1985) 51–56.
- [21] B. Grammaticos, A. Ramani, J. Hietarinta, A search for integrable bilinear equations: The Painlevé approach, J. Math. Phys. 31 (1990) 2572–2578.
- [22] A. Karasu-Kalkanlı, Painlevé classification of coupled Korteweg–de Vries systems, J. Math. Phys. 38 (1997) 3616–3622.
- [23] S.Yu. Sakovich, Coupled KdV equations of Hirota–Satsuma type, J. Nonlinear Math. Phys. 6 (1999) 255–262; arXiv:solv-int/9901005.

- [24] S.Yu. Sakovich, Addendum to: Coupled KdV equations of Hirota–Satsuma type, J. Nonlinear Math. Phys. 8 (2001) 311–312; arXiv:nlin/0104072.
- [25] S. Sakovich, A note on the Painlevé property of coupled KdV equations, Int. J. Part. Diff. Eqns. 2014 (2014) 125821; arXiv:nlin/0402004.
- [26] S.Yu. Sakovich, T. Tsuchida, Symmetrically coupled higher-order nonlinear Schrödinger equations: singularity analysis and integrability, J. Phys. A: Math. Gen. 33 (2000) 7217–7226; arXiv:nlin/0006004.
- [27] A. Karasu-Kalkanlı, A. Karasu, S.Yu. Sakovich, Integrability of a generalized Ito system: the Painlevé test, J. Phys. Soc. Jpn. 70 (2001) 1165–1166; arXiv:nlin/0102030.
- [28] A. Karasu-Kalkanh, A. Karasu, A. Sakovich, S. Sakovich, R. Turhan, A new integrable generalization of the Korteweg–de Vries equation, J. Math. Phys. 49 (2008) 073516; arXiv:0708.3247.
- [29] G.Q. Xu, The integrability for a generalized seventh-order KdV equation: Painlevé property, soliton solutions, Lax pairs and conservation laws, Phys. Scr. 89 (2014) 125201.