Snowmass White Paper: Generalized Symmetries in Quantum Field Theory and Beyond

Clay Córdova^{1*}, Thomas T. Dumitrescu^{2†}, Kenneth Intriligator^{3‡}, Shu-Heng Shao^{4§}

¹Enrico Fermi Institute and Kadanoff Center for Theoretical Physics, University of Chicago

²Mani L. Bhaumik Institute for Theoretical Physics, UCLA

³Department of Physics, University of California, San Diego

⁴C.N. Yang Institute for Theoretical Physics, Stony Brook University

Abstract

Symmetry plays a central role in quantum field theory. Recent developments include symmetries that act on defects and other subsystems, and symmetries that are categorical rather than group-like. These generalized notions of symmetry allow for new kinds of anomalies that constrain dynamics. We review some transformative instances of these novel aspects of symmetry in quantum field theory, and give a broad-brush overview of recent applications.

May 2022

*e-mail: clayc@uchicago.edu

†e-mail: tdumitrecu@physics.ucla.edu

‡e-mail: keni@ucsd.edu

§e-mail: shu-heng.shao@stonybrook.edu

Contents

1	Introduction From Ordinary to Generalized Symmetries					
2						
	2.1	Ordinary Symmetries				
	2.2	Higher Form Symmetries				
	2.3	Higher Group Symmetries				
	2.4	Non-Invertible Symmetries				
	2.5	Subsystem Symmetries and Fractons				
3	Anomalies and Dynamical Consequences of Symmetry					
	3.1	Anomalies and Anomaly Matching				
	3.2	Applications to QFT Dynamics				

1 Introduction

Symmetry has long been a transformative, development-guiding notion in physics. Examples include Noether's theorems [1, 2] connecting symmetries and conservation laws, and Landau's symmetry-based theory of phase transitions [3]. Transitions between quantum states are constrained by the Wigner-Eckart theorem, which applies to quantum systems whose Hamiltonian is invariant under a symmetry group. Here we will discuss symmetry in the context of quantum field theory (QFT), where additional structures such as locality or Lorentz invariance lead to a rich set of refinements and generalizations of the notion of symmetry. The resulting generalized symmetries can act on lower-dimensional defects and subsystems within the QFT, and their properties are in general captured by higher categories rather than by groups.

Given that symmetry is such a vast subject, this overview will necessarily be an incomplete sketch of a few recent, selected lines of development, with many omissions. As advertised we primarily focus on symmetries in QFT, with emphasis on renormalization group (RG) flows and anomalies. The later are intimately related to the possibility of projectively realizing symmetries in quantum theory. For a different recent survey from a condensed matter perspective, see for instance [4].

The fundamental forces are described via local gauge interactions. In that context, gauge invariance reflects a convenient but unphysical redundancy introduced to satisfy the requirements of locality and Lorentz invariance in relativistic QFT; as such it is exact, but has no observable consequences, e.g. it can famously not be spontaneously broken [5].

By contrast, global symmetries imply relations between observable physical quantities; as such they may be spontaneously broken, and they are useful even if only approximate. Examples include the approximate unbroken $SU(3)_{flavor}$ and $SU(2)_{isospin}$ symmetries of QCD, as well as the approximate discrete symmetries C and P. QCD spontaneously breaks its approximate chiral quark flavor symmetries; the pions are the associated (pseudo-) Nambu-Goldstone bosons (NGBs). Such considerations led to the discovery of the Adler-Bell-Jackiw (ABJ) anomaly $[6,7]^2$ and the closely related explicit violation of ABJ-anomalous symmetries by instantons [10]. Global symmetries can powerfully constrain QFT dynamics, even in strongly coupled theories. 't Hooft noted in [12] that anomalies involving only global currents (unlike the ABJ case) are obstructions to gauging and they must be constant on renormalization group (RG) flows; such anomalies are often now often referred to as 't Hooft anomalies and will be reviewed here.

¹ This paper is one of Landau's famous contributions during his tenure at the Ukrainian Physico-Technical Institute in Kharkiv.

² It was recently pointed out that the ABJ anomaly actually leads to non-invertible global symmetries [8,9].

³ See [11] for an influential account of symmetry in the development of QFT.

Global symmetries are termed accidental (or emergent) when they arise or enhance in the deep IR and violated by higher dimension operators. Examples are $U(1)_B$ baryon and $U(1)_L$ lepton number, which are preserved by the dimension 4 operators of the Standard Model (SM) but are violated by higher dimension operators (as well as by electroweak instantons).⁴ All global symmetries are expected to be accidental in quantum gravity, see for instance the recent review [13] and references therein.

There have been many attempts to generalize global symmetries in relativistic QFT, e.g. by combining internal and Lorentz symmetries, or by contemplating conserved currents of higher spin $s \ge 3.5$ The latter famously arise in the context of integrable QFTs in d=2 spacetime dimensions, as in [14]. For d>2, the Coleman-Mandula (CM) theorem [15] implies that higher spin conserved currents essentially render the theory free.

There are two interesting, broad classes of exceptions to the CM theorem: conformal field theories, where the Poincaré group is extended to the conformal group SO(d,2),⁶ and supersymmetric field theories [17]; these two classes intersect in the superconformal theories [18]. We will not specifically focus on these classes of theories here; they are covered in other Snowmass white papers (see e.g. [19,20]). Instead, we will survey other large classes of generalized symmetries that are not ruled out by the CM-theorem and that require neither conformal symmetry nor supersymmetry (though they do occur in conformal and supersymmetric theories).

The remainder of this white paper is organized as follows:

In section 2 we review basic aspects of ordinary symmetries (section 2.1) before surveying four kinds of generalized global symmetries: higher form symmetries (section 2.2), higher group symmetries (section 2.3), non-invertible symmetries (section 2.4) and subsystem symmetries (section 2.5). The first three generalizations can arise in relativistic systems, while the fourth one is only known to occur in non-relativistic contexts, notably in condensed matter models of fractons.

Section 3 is concerned with 't Hooft anomaly matching and applications to QFT dynamics. In section 3.1 we review the notion of an 't Hooft anomaly for ordinary global symmetries, as well as the closely associated concepts of anomaly matching and anomaly inflow. We then outline how these anomalies extend to the generalized symmetries introduced in section 2. In section 3.2 we survey some recent applications of anomaly matching for ordinary and global symmetries.

2 From Ordinary to Generalized Symmetries

2.1 Ordinary Symmetries

Here we summarize a few standard facts about ordinary global symmetries that will set the stage for subsequent generalizations. Throughout this paper, d stands for the spacetime dimension.

In a QFT with an ordinary, or 0-form, global symmetry $G^{(0)}$, there is a topological defect $U^{(0)}(g,\Sigma_{d-1})$ for every group element $g\in G^{(0)}$ that is supported on a closed codimension-1 submanifold Σ_{d-1} of spacetime.⁷ The fusion of the $U^{(0)}(g,\Sigma_{d-1})$ is governed by the group law for $G^{(0)}$.⁸ An elementary example is $G^{(0)}=U(1)$, in which case Noether's theorem implies a conserved 1-form current $J^{(1)}$,

$$d * j^{(1)} = 0. (1)$$

⁴ The combination $U(1)_{B-L}$ is exactly preserved in the SM, but is expected to be violated by physics beyond the SM (BSM).

⁵ These are symmetric tensors of the Lorentz group with $s \ge 3$ indices. By contrast, the stress tensor $T_{\mu\nu}$ is a two-index Lorentz tensor of spin s = 2.

⁶ A CFT analogue of the CM theorem was proved in [16].

⁷ We sometimes call $U^{(0)}(g,\Sigma_{d-1})$ a symmetry defect. For an exposition of this perspective, see for instance [21] and references therein. Throughout we use $X^{(p)}$ to indicate that X is a p-form.

⁸ If $G^{(0)}$ has an 't Hooft anomaly, it shows up in more complicated configurations of the symmetry defects (see for instance [21]).

We can then construct a conserved charge $Q^{(0)}(\Sigma_{d-1})$, whose exponential gives the $U(g,\Sigma_{d-1})$,

$$Q^{(0)}(\Sigma_{d-1}) = \int_{\Sigma_{d-1}} *j^{(1)} , \qquad U^{(0)}(g = e^{i\alpha}, \Sigma_{d-1}) = \exp\left(i\alpha Q^{(0)}(\Sigma_{d-1})\right) . \tag{2}$$

Current conservation and Stokes' theorem imply that the dependence on Σ_{d-1} is topological, as long as the deformation of Σ_{d-1} does not collide with other insertions. The group multiplication law follows from the additivity of the charge. In the most familiar setup, Σ_{d-1} is a spatial slice in Lorentzian signature; in Euclidean signature Σ_{d-1} can be any suitable closed codimension-1 cycle. Some comments are in order:

• If $G^{(0)}$ is non-Abelian and connected, the group law for the $U^{(0)}(g, \Sigma_{d-1})$ can be deduced from the fact that the OPE of the currents $j_a^{(1)}$ encodes the structure constants f_{abc} of the Lie algebra of $G^{(0)}$,

$$j_a^{(1)}(x)j_b^{(1)}(0) \sim C(x)f_{abc}j_c^{(1)}(0) + \cdots,$$
 (3)

with C(x) a known, singular function depending on the Lorentz indices of the currents (which we suppress).

- If $G^{(0)}$ is not connected (e.g. it could be a discrete group) then some of the symmetry defects $U^{(0)}(g, \Sigma_{d-1})$ are not associated with conserved currents; nevertheless they typically exist and have the properties outlined above. This is a generalization of Noether's theorem.
- $G^{(0)}$ naturally acts on local operators by linking them with $U^{(0)}(g,\Sigma_{d-1})$. These operators in turn create states (e.g. point particles) charged under $G^{(0)}$. More generally, $G^{(0)}$ acts on the Hilbert space \mathscr{H} in Lorentzian signature.
- If the vacuum is preserved by a subgroup $H^{(0)} \subset G^{(0)}$, then the $G^{(0)}$ is spontaneously broken to $H^{(0)}$. This leads to a vacuum manifold $\mathscr{M} = G^{(0)}/H^{(0)}$, and hence to Nambu-Goldstone (NG) bosons (if \mathscr{M} has positive dimension) or domain walls between the disconnected components of \mathscr{M} . The Hilbert space \mathscr{H} in each vacuum furnishes a representation of $H^{(0)}$.
- A QFT_d with $G^{(0)}$ symmetry can be coupled to a non-dynamical background $G^{(0)}$ connection $A^{(1)}$ in a way that preserves background gauge invariance. If $G^{(0)}$ has an 't Hooft anomaly this requires placing the theory at the boundary of an invertible anomaly theory $\mathcal{A}_{d+1}[A^{(1)}]$ that depends on the $G^{(0)}$ connection. The symmetry can then be gauged by path-integrating over $A^{(1)}$.

2.2 Higher Form Symmetries

CM-like theorems famously limit possibly generalizations of $G^{(0)}$ symmetry in relativistic QFT (see introduction; see below for a discussion of non-relativistic systems). One such generalization, suggested by experience from string theory and SUSY QFT, 11 was spelled out in [24, 21]:12 replace the 1-form current $j^{(1)}$ by a (p+1)-form current $J^{(p)}$ satisfying

$$d * J^{(p+1)} = 0. (4)$$

⁹ See for instance [22] for a recent discussion of possible failure modes, with references.

 $^{^{10}}$ If one insists on a strictly *d*-dimensional description, the anomaly spoils background gauge-invariance in a mild and familiar way, e.g. the partition function is multiplied by complex phases under $G^{(0)}$ gauge transformations.

¹¹ In string theory, *p*-form symmetries (like all symmetries) are naturally gauged. In SUSY QFT's they can be global symmetries, and the associated charges can extend the SUSY algebra (see [23] and references therein).

¹² See [25–32] for a highly incomplete sampling of later work focusing on higher-form symmetries.

As before we can construct a charge $Q^{(p)} = \int_{\Sigma_{d-p-1}} *J^{(p+1)}$ by integrating over a codimension-(p+1) cycle, and a symmetry defect $U^{(p)}(g=e^{i\alpha},\Sigma_{d-p-1})=\exp(i\alpha Q^{(p)}(\Sigma_{d-p-1}))$, both of which depend topologically on Σ_{d-p-1} . The symmetry defects satisfy a U(1) group law, and we say that they generate a U(1) p-form symmetry, denoted $U(1)^{(p)}$. An important example is free Maxwell theory in any spacetime dimension d, based on a dynamical U(1) connection $a^{(1)}$ with field strength $f^{(2)}=da^{(1)}$. The Maxwell equations,

$$d * f^{(2)} = 0, \ df^{(2)} = 0,$$
 (5)

can be read as conservation equations for an electric 2-form current $f^{(2)}$ and a magnetic (d-2)-form current $f^{(2)}$. Thus the theory has $U(1)_e^{(1)} \times U(1)_m^{(d-3)}$ symmetry.¹³ Some comments are in order:

- In analogy with 0-form symmetries, we can also consider discrete $(p \ge 1)$ -form symmetries $G^{(p)}$, which are necessarily abelian. For instance, adding fields of electric charge N to Maxwell theory breaks $U(1)_e^{(1)} \to \mathbb{Z}_N^{(1)}$. Another important example is the $\mathbb{Z}_N^{(1)}$ center symmetry of pure SU(N) Yang-Mills theory (and more generally, the center symmetry of any pure gauge theory). Due to their importance in gauge theory, we will mostly focus on 1-form symmetries, but the discussion holds for all p-form symmetries with $p \ge 1$.
- All 1-form symmetries $G^{(1)}$ naturally act on line defects by linking them with the codimension-2 topological symmetry defects $U^{(1)}(g,\Sigma_{d-2})$. For instance, the line operators charged under the $\mathbb{Z}_N^{(1)}$ center symmetry of SU(N) gauge theory are the Wilson lines. However, the 1-form symmetry defects cannot link local operators, and hence such symmetries commute with local operators, as well as with the point particle states they create or with the S-matrix of these particles. This is how $(p \ge 1)$ -form symmetries evade CM-like theorems.
- 1-form symmetries can spontaneously break. For instance, in dimension $d \ge 4$ the photon of theories with an abelian Coulomb phase, $d \ge 4$ the photon of theories with an abelian Coulomb phase, $d \ge 4$ the photon of theories with an abelian Coulomb phase, $d \ge 4$ the photon of theories with an abelian Coulomb phase, $d \ge 4$ the photon of theories with an abelian Coulomb phase, $d \ge 4$ the photon of theories with an abelian Coulomb phase, $d \ge 4$ the photon of theories with an abelian Coulomb phase, $d \ge 4$ the photon of theories with an abelian Coulomb phase, $d \ge 4$ the photon of theories with an abelian Coulomb phase, $d \ge 4$ the photon of theories with an abelian Coulomb phase, $d \ge 4$ the photon of theories with an abelian Coulomb phase, $d \ge 4$ the photon of theories with an abelian Coulomb phase, $d \ge 4$ the photon of the photon of $d \ge 4$ the photon of the photon of $d \ge 4$ the photon of the photon $d \ge 4$ the photon of the photon $d \ge 4$ the photon of $d \ge 4$ the p

¹³ The statement that these are compact U(1) symmetries is related to the fact that $a^{(1)}$ is a U(1) connection, with quantized electric charges and magnetic fluxes. The $U(1)_e^{(1)}$ and $U(1)_m^{(d-3)}$ symmetries have a mixed 't Hooft anomaly [21] that plays an important role in many applications.

 $^{^{14}}$ This is because the associated codimension-(p+1) symmetry defects can always be exchanged without crossing, leading to abelian fusion rules

¹⁵ In these cases the reduction of the center symmetry to $\mathbb{Z}_N^{(1)}$ can be understood in terms of screening of the Wilson lines by electrically charged matter.

¹⁶ Similar, long familiar results apply in d=3, involving ordinary rather than higher-form symmetry: the 3d photon can be dualized to a compact scalar, which is the NG boson of spontaneously broken $U(1)_m^{(0)}$ [33]. In d=3, confinement arises either if $U(1)_m^{(0)}$ is restored by the vacuum, or if it is explicitly broken, e.g. by 3d instantons (4d monopoles). Explicit breaking gives a mass to the pseudo-NG boson scalar photon [33, 34]. For spontaneously broken $U(1)_m^{(p)}$ with p>0, the spin-p NG boson does not get a mass from explicit breaking of the symmetry, see [4] and references cited therein for this *robustness* of higher-form symmetry. For d=4, the photon can be viewed as the NG boson of spontaneous breaking of either $U(1)_e^{(1)}$ or $U(1)_m^{(1)}$.

¹⁷ The low-energy theory has the IR-free photon of Maxwell theory. The photon can have IR irrelevant, higher-derivative interactions if $U(1)_{\ell}^{(1)}$ is accidental; e.g. this happens for supersymmetric theories on their Coulomb branch.

¹⁸ This extends Polyakov's description [38], making the thermal circle there inessential [21].

¹⁹ See [4] for a recent discussion of generalized symmetries through the lens of condensed matter physics.

- A suitable background field for a $G^{(1)}$ symmetry is a 2-form gauge field $B^{(2)}$, 20 with its standard 1-form gauge transformations $B^{(2)} \to B^{(2)} + d\Lambda^{(1)}$. As for ordinary symmetries, the theory is invariant under such transformations up to possible 't Hooft anomalies, which can be cancelled by a suitable invertible anomaly theory $\mathcal{A}_{d+1}[B^{(2)}]$. In the absence of 't Hooft anomalies, the $G^{(1)}$ symmetry can be gauged by path-integrating over $B^{(2)}$.
- Typically, gauging a p-form symmetry $G^{(p)}$ in d spacetime dimensions gives rise to a new theory with a $\tilde{G}^{(q)}$ global symmetry, whose degree q depends on whether G is discrete or continuous. For instance, gauging a $\mathbb{Z}_n^{(p)}$ symmetry in d dimensions typically gives rise to a new theory with a $\mathbb{Z}_n^{(q=d-p-2)}$ symmetry. This generalizes the familiar twisted sector (or quantum) symmetry of two-dimensional orbifolds, where d=2 and p=0.
- More generally, one can gauge a discrete *p*-form symmetry only along a codimension-*q* submanifold in spacetime. This is known as higher gauging [41] of a *p*-form global symmetry. Higher gauging does not change the bulk of the QFT, but generates a codimension-*q* topological defect, known as a condensation defect [42–46,41,47]. Such condensation defects are generally not invertible, i.e. they generate a non-invertible global symmetry (see section 2.4 below).

2.3 Higher Group Symmetries

Higher group symmetries arise when p-form symmetries $G^{(p)}$ of different form degrees are present and can affect each other in various ways. Roughly speaking, if p < q then $G^{(p)}$ can act on $G^{(q)}$, while if p > q the background field $B^{(p+1)}$ for $G^{(p)}$ can shift under background gauge transformations of the background field $B^{(q+1)}$ for $G^{(q)}$. The precise form of the shift is determined by certain group cohomology (or Postnikov) classes.²³

Higher-group gauge symmetry has a long history in string theory, most famously in the context of the Green-Schwarz (GS) mechanism and its generalizations. The fact that higher groups can also furnish global symmetries of QFTs is a more recent observation that is actively being explored and applied in many contexts, see [51–54, 25, 55, 56, 40, 57–80] for an incomplete sampling of the recent literature.²⁴

Rather than discuss higher-group symmetries in general, we will focus on a concrete and elementary example: a 2-group symmetry involving $G^{(0)} = U(1)_A^{(0)}$ and $G^{(1)} = U(1)_B^{(1)}$. The relevant Postnikov class resides in $H^3(G^{(0)}, G^{(1)})$, and we assume that it is non-vanishing. In this case the fusion of the $U(1)_A^{(0)}$ current $J_A^{(1)}$ with itself contains the $U(1)_B^{(1)}$ current $J_B^{(2)}$. Schematically,

$$j^{(1)}(x)_A j_A^{(1)}(0) \sim K(x) J_R^{(2)}(0) + \cdots,$$
 (6)

where K(x) is a known, singular function depending on the Lorentz indices of the currents. Its overall normalization is determined by the Postnikov class. This fusion rule is reminiscent of the non-Abelian fusion rule for $G^{(0)}$ currents above, except that it involves currents of different form degree. The structure of this OPE forces us to deform the algebra of background gauge transformations. In particular, we find that

²⁰ If $G^{(1)}$ is discrete, then $B^{(2)}$ is valued in $H^2(M_d, G^{(1)})$, where M_d is the spacetime manifold.

²¹ Here $\Lambda^{(1)}$ is the 1-form gauge parameter. Again a refinement is required if $G^{(1)}$ is discrete.

²² More care is required if $G^{(p)}$ has 't Hooft anomalies or if p = 0 and G is non-Abelian. See for instance [39, 40].

²³ See for instance [48–50] for a mathematical exposition of higher groups and related subjects.

²⁴ Note that higher-form global symmetries are a special case of higher-group global symmetries.

²⁵ Here we assume that d > 2. For comments on the d = 2 case see for instance [25].

²⁶ Such examples can be engineered in 4d Abelian gauge theories with fermionic mater, where the Postnikov data is determined by certain mixed triangle anomaly coefficients involving two $j_A^{(1)}$ currents and one abelian gauge field. In this case $U(1)_B^{(1)}$ is the magnetic 1-form symmetry of that gauge field (assuming no magnetic matter).

the background field $B^{(2)}$ that couples to $J_B^{(2)}$ receives a background GS shift depending on the background field $A^{(1)}$ that couples to $J_A^{(1)}$, as well as on the Postnikov class. Additional details can be found in [58,64].

2.4 Non-Invertible Symmetries

As reviewed in Section 2.1, in relativistic systems, the modern characterization of an ordinary global symmetry is in terms of its topological symmetry operators and defects of codimension one in spacetime. A natural question then arises: While every ordinary global symmetry element is associated with a topological defect, is the converse true? This turns out *not* to be the case: there are topological defects not associated with any ordinary global symmetry. These defects are *non-invertible* because there is no inverse in the fusion algebra and their fusion can involve more than one operator.

In 1+1d, non-invertible topological lines are ubiquitous, especially in rational CFTs and integrable field theories.²⁷ In rational CFT [81, 84, 85], there is a general construction of topological lines that commute with the extended chiral algebra [86, 87]. The simplest example is the Kramers-Wannier duality line in the Ising CFT [88–90]. The fusion of two non-invertible duality lines \mathcal{N} gives the sum of the identity line and the \mathbb{Z}_2 symmetry line η :

$$\mathcal{N} \times \mathcal{N} = 1 + \eta. \tag{7}$$

As we bring the (local) order operator past this line, it becomes a (non-local) disorder operator on the other side.

In recent years, it has been advocated that these non-invertible topological defects should be viewed as a generalization of ordinary global symmetries [39, 91]. The fusion and crossing properties of these topological lines are mathematically captured by a fusion category. The generalized gauging of these lines in 1+1d has been developed in [90, 92, 93, 39, 94–96]. When there is an obstruction to gauging, the corresponding anomalies lead to non-trivial dynamical constraints on the renormalization group flows [91, 97, 94, 98]. The relation to fermionic anomalies has been discussed in [99–103].

In higher spacetime dimensions, there are also examples of non-invertible defects of various codimensions. In 2+1d, the non-abelian anyons in a low-energy TQFT, described by the theory of modular tensor categories, can be viewed as the non-invertible generalization of one-form global symmetries (see [104–107, 41] for recent discussions). Recently, the non-invertible Kramer-Wannier duality defects have been generalized to 3+1d gauge theories [108, 109, 102, 110, 47, 111, 112] via the gauging of higher-form global symmetries. More generally, in any theory with a higher-form symmetry, the most fundamental non-invertible symmetries are condensation defects [42–46] arising from *higher gauging* [41]. Condensation defects in 2+1d were systematically analyzed in [41], generalizing the results of [85, 113]. The non-invertible fusion "coefficients" of these *n*-dimensional topological defects are generally not numbers, but *n*-dimensional TQFTs [41,47]. More recently, infinitely many non-invertible global symmetries were identified in the Standard Model and axion models from ABJ anomalies [8,9].

Non-invertible defects have also found applications in quantum gravity in the context of the completeness hypothesis [114–118]. They have also been realized on the lattice [119–125,108,126–128], sometimes under the name of "algebraic higher symmetry" [129,45].

2.5 Subsystem Symmetries and Fractons

A novel kind of global symmetry, known as *subsystem global symmetry*, has featured prominently in many condensed matter systems, including the gapless model of [130] and many gapped fracton models

 $^{^{27}}$ The search for hidden symmetry has a long history in these theories. For instance, in rational CFTs any primary operator g with a fusion rule that only contains the identity operator on the right-hand side leads to an invertible abelian symmetry of the theory that can be used to extend the chiral algebra [81–83].

[131–133]. (See [134–136] for reviews on fractons, and [137] for a Snowmass white paper on field theories for fractons.)

Subsystem symmetry is similar to the higher-form symmetry, but they differ in important ways (see [138, 139] for related discussions). For both kinds of global symmetries, the conserved charges are supported on closed, higher-codimensional manifolds $\mathscr S$ in space. Unlike the higher-form symmetry charge, the charge of a subsystem symmetry depends not only on the topology of $\mathscr S$, but possibly also on the shape and the location of $\mathscr S$. Furthermore, the charge might only be allowed to be on certain $\mathscr S$ but not all manifolds of a given codimension. On a lattice, the number of independent conserved charges grows with the number of sites, leading to a huge global symmetry in the continuum limit.

Just like ordinary global symmetries, subsystem global symmetries can be spontaneously broken [140–146]. They can also have 't Hooft anomalies [144, 147, 148], which are obstructions to gauging them. These anomalies are captured by Subsystem Symmetry Protected Topological (SSPT) phases in one higher dimension [149–154,148,155]. The symmetry fractionalization of subsystem symmetries has been recently discussed in [156].

Fractors are a large class of lattice models with subsystem global and gauge symmetries. In 3+1 dimensions, fractor models are robust topological phases with many peculiar properties, including infinite ground state degeneracy that is sub-extensive in the system size, and excitations with restricted mobility. Because of these unusual characteristics, gapped fractor models cannot be described by conventional TQFT. Some fractor models can be realized by gauging a subsystem symmetry [133, 157–160, 144, 161]. In the low-energy limit, they are typically described by certain exotic higher-rank tensor gauge theories [162–170, 158, 171–178, 138, 179–181, 142–144, 182–191].

The most significant consequence of subsystem symmetry is that the low-energy observables are sensitive to details of the short-distance physics. For instance, in fracton models the large ground state degeneracy is enforced by the subsystem global symmetries and their 't Hooft anomalies [144, 148]. Similarly, the restricted mobility of the fracton excitations is traced to the selection rules arising from a subsystem global symmetry, the *time-like global symmetry*, that acts on defects, rather than operators [190]. This surprising UV/IR mixing [142, 145, 192–194], which is reminiscent of that in quantum field theory on a non-commutative space [195], is the main reason why these exotic systems defy a conventional continuum field theory description [158, 196, 197, 142–144, 198, 199, 182, 161, 200, 147, 186, 145, 201, 202].

A summary of the various kinds of symmetry described above is given in Table 1.

Properties of	Ordinary	Higher-form	Non-invertible	Subsystem
symmetry operator	symmetry	symmetry	symmetry	symmetry
Codimension	1	> 1	≥ 1	> 1
in spacetime				
Topological	yes	yes	yes	not completely
				but conserved in time
Fusion rule	group	group	category	group

Table 1: Properties of symmetry defects for various kinds of global symmetry [203].

3 Anomalies and Dynamical Consequences of Symmetry

3.1 Anomalies and Anomaly Matching

The strong force, confinement, dualities, and emergent degrees of freedom illustrate how renormalization group (RG) flows can lead to surprising interconnections between classically very different UV descriptions of IR physics, or between the UV and IR limits of RG flows. A key feature of any concept of symmetry is how it survives continuous, symmetry preserving deformations of the theory. This includes in particular renormalization group flows: if at short distances a model supports a given symmetry, then the same symmetry must be present in any long distance effective field theory.²⁸

This idea of global symmetry matching between the ultraviolet and the infrared becomes remarkably powerful when extended to include 't Hooft anomalies of global symmetries. 't Hooft's pioneering idea [12] was to view such anomalies as an obstruction to gauging a global symmetry, and to argue that they must be invariant under the renormalization group by coupling to a suitable set of spectator (i.e. weakly interacting) fermions. Subsequently, this idea of anomaly matching was understood as constraints on various structure functions of conserved current correlation functions [204, 205].

The modern point of view of anomalies is that of anomaly inflow [206] (see also [207]) where an anomaly of a d-dimensional quantum field theory with spacetime X_d is presented via a theory \mathcal{A}_{d+1} in d+1spacetime dimensions, with bulk spacetime Y_{d+1} such that the original spacetime is the boundary of the bulk spacetime, $X_d = \partial Y_{d+1}$. In some contexts, e.g. [206, 208, 209], the d+1-dimensional anomaly theory \mathcal{A}_{d+1} has a clear physical interpretation, with the lower dimensional anomalous theory arising as a defect in the ambient physical system.²⁹ This point of view is conceptually similar to that of symmetry protected topological phases in condensed matter physics, where a trivially gapped system in bulk gives rise to anomalous edge modes on its boundary. From a more abstract perspective, the anomaly theory generalizes the concept of projective representations, which describe simple anomalies in quantum mechanics (see e.g. [26]), to the quantum field theory setting. The anomaly theory can be taken as the definition of the anomaly, and this notation can be applied to generalized symmetries. The anomaly theory \mathcal{A}_{d+1} has been significantly developed and applied in recent years in the high energy, condensed matter, and mathematics communities. This includes results for familiar continuous global symmetries [214–220], discrete global symmetries [221–225,56,226–230], higher-form and higher-group symmetries [52,54,58,59,231,74], and subsystem symmetries [144, 147, 148], families of quantum field theories [232–241], as well as structural developments [242–244, 40, 99, 245–249, 55, 250, 251].

For continuous global symmetries in even d, the anomaly theory \mathcal{A}_{d+1} is often a Chern-Simons-type TQFT, as in the descent formalism of [252] and references therein. More generally, global or non-perturbative anomalies can be described via the η -invariant [253] which can also be described via inflow [254, 255]. In the context of $AdS_{d+1} \leftrightarrow CFT_d$ duality, the anomaly theory \mathcal{A}_{d+1} is a topological subsector of the AdS_{d+1} bulk gravity theory [256]. The anomaly theory \mathcal{A}_{d+1} for a traditional QFT merely encodes an overall phase, so it is associated with a one-dimensional Hilbert space. The anomaly theory in such cases is called *invertible* [257, 258] (see also [55, 259, 260]).³⁰

²⁸ Of course symmetries can be enhanced in the IR, by additional, accidental components. Moreover, symmetries can decouple in the IR if they only act non-trivially on degrees of freedom that have been decoupled from the low-energy theory. Most precisely, there is a homomorphism from the symmetry group of the UV to that of the IR.

 $^{^{29}}$ If the anomalous symmetries are gauged on the defect, the bulk theory is physical and cannot be decoupled; there is no "genuine" d dimensional theory in this case. For 't Hooft anomalies, the dynamics of the bulk d+1 dimensional theory can decoupled, leaving only the anomaly. A classic example are the d+1 dimensional Wess-Zumino-Witten terms for the pions of 4d QCD in the chiral symmetry broken phase [210,211] and 2d non-linear sigma models [212]. Another example are the chiral, domain wall fermions of [213], where the defect arises via spatially dependent backgrounds (the Dirac mass) in the bulk.

There are also more exotic *d*-dimensional QFTs, sometimes called relative [261] or meta or non-genuine theories, where the \mathcal{A}_{d+1} theory plays a more essential role than just a projective phase and is instead associated with a multi-dimensional Hilbert

3.2 Applications to QFT Dynamics

There are a wide variety of applications of anomalies and symmetry to the dynamics of quantum field theory. Due to their robust nature, symmetries and their associated anomalies can often be reliably computed in weakly coupled limits of a theory, and then applied to constrain the dynamics of strongly coupled phases. For ordinary continuous global symmetries, non-zero 't Hooft anomalies preclude a mass gap: the IR theory must have massless chiral fields that saturate the anomalies in the unbroken case, or massless NGBs (with a WZW-type term) in the spontaneously broken case. For discrete global symmetries, non-zero 't Hooft anomalies can also in some cases be matched by a gapped TQFT (if an obstruction [265,266] vanishes). 't Hooft anomaly matching played a key role in the discovery of electromagnetic dualities in four-dimensional gauge theories [267–269] and has since become an essential field theory technique. Modern applications are inspired by this example but expand the logic to the more general concepts of symmetry described above, and hence to the more general class of invertible field theories encoding their anomalies. Below we highlight several classes of examples.

Higher form symmetry has been crucial in understanding the infrared behavior of Chern-Simons matter theories in three dimensions. In this context one also finds a close link to fundamental concepts in three-dimensional topological field theory where one-form symmetry generators can be interpreted as abelian anyons [24, 21, 270–273, 28, 274, 275]. The language of higher form symmetry then gives a way to understand the properties of these line defects in a broader context of non-topological field theories. In particular this has given rise to an explosion of new level-rank and boson-fermion dualities for such interacting phases [276, 273, 277–284], as well as new results for the phase diagrams of strongly interacting gauge theories [285–288, 279, 281, 289, 290, 283, 291].

In four spacetime dimensions there have also been extensive recent applications of symmetry to constrain the dynamics of gauge theories [53,271,292–298,60,299–304,31,63,305–310,69,311,72,312,310,313–316]. In these gauge theory contexts, one-form symmetry finds a natural utilization as the symmetry governing the transition between confinement and deconfinement [21]. A highlight is the identification of an anomaly at $\theta = \pi$ in certain pure Yang-Mills theories implying non-trivial long-distance dynamics of these models [26]. There is also a natural link to the three-dimensional Chern-Simons matter results which often manifest on domain walls in four dimensions [317–322]. Further there is a sharp interplay between anomalies, spontaneous symmetry breaking, and the mass gap in such theories [323–327, 265, 266, 328].

The extensive appearance of higher-form symmetry in quantum field theory means that frequently one finds higher group global symetries, which have their own anomalies and dynamical implications. Higher-group global symmetry has also been extensively explored in five and six-dimensional quantum field theory and little string theories [329, 32, 61, 30, 330, 71, 331, 75, 76, 79, 332], where in the latter continuous higher symmetry is particularly natural due to the presence of a conserved string number [333, 64]. In particular, these ideas have been applied to better understand certain supersymmetric dualities [334–336,65,337–339], and to derive universal constraints on renormalization group flows [340–342].

Similar to the anomaly constraints from an ordinary symmetry, the non-invertible symmetries also have dramatic consequences on renormalization group flows. In two-dimensional QFT, invertible and non-invertible topological lines have been explored extensively in [39,91,343,97,246,101,94,344–351,229,98, 352,96,353] in recent years with various dynamical applications. In particular, using a modular invariance argument, it was shown in [91] (see also [97] for generalizations) that the existence of certain non-invertible topological lines is incompatible with a trivially gapped phase. There are also recent applications to 2d adjoint QCD, clarifying the infrared behavior [94]. In four-dimensional theories the dynamical implications of non-invertible symmetry are just beginning to emerge with recent results on anomalies of such symmetries [109,47] and applications to Yang-Mills theory [102].

space. This was seen in the context of 2d chiral RCFTs, where the bulk is a 3d Chern-Simons theory [81, 262]. Other examples include certain 6d $\mathcal{N} = (2,0)$ theories, which arise at the boundary of 7d Chern-Simons type theories, see e.g. [263, 21, 264, 32].

Acknowledgements

We are grateful to our collaborators and colleagues who have helped shape this exciting subject, and to the organizers of the formal theory Snowmass process, L. Rastelli, and D. Poland, for inviting us to contribute this white paper. CC, TD, and KI acknowledge support from the Simons Collaboration on Global Categorical Symmetries. In addition CC is supported by the US Department of Energy DE-SC0021432; TD is supported by a DOE Early Career Award under DE-SC0020421 and the Mani L. Bhaumik Presidential Chair in Theoretical Physics at UCLA; KI is supported by DOE award DE-SC0009919 and Simons Foundation award 568420.

References

- [1] E. Noether, *Invariant Variation Problems*, *Gott. Nachr.* **1918** (1918) 235–257, [physics/0503066].
- [2] N. Byers, E. Noether's discovery of the deep connection between symmetries and conservation laws, in Symposium on the Heritage of Emmy Noether, 7, 1998. physics/9807044.
- [3] L. D. Landau, On the theory of phase transitions, Zh. Eksp. Teor. Fiz. 7 (1937) 19–32.
- [4] J. McGreevy, Generalized Symmetries in Condensed Matter, arXiv: 2204.03045.
- [5] S. Elitzur, *Impossibility of Spontaneously Breaking Local Symmetries*, *Phys. Rev. D* **12** (1975) 3978–3982.
- [6] S. L. Adler, Axial vector vertex in spinor electrodynamics, Phys. Rev. 177 (1969) 2426–2438.
- [7] J. S. Bell and R. Jackiw, A PCAC puzzle: $\pi^0 \rightarrow \gamma \gamma$ in the σ model, Nuovo Cim. A **60** (1969) 47–61.
- [8] Y. Choi, H. T. Lam, and S.-H. Shao, *Non-invertible Global Symmetries in the Standard Model*, arXiv:2205.05086.
- [9] C. Córdova and K. Ohmori, *Non-Invertible Chiral Symmetry and Exponential Hierarchies*, arXiv:2205.06243.
- [10] G. 't Hooft, How Instantons Solve the U(1) Problem, Phys. Rept. **142** (1986) 357–387.
- [11] S. Coleman, *Aspects of Symmetry: Selected Erice Lectures*. Cambridge University Press, Cambridge, U.K., 1985.
- [12] G. 't Hooft, Naturalness, chiral symmetry, and spontaneous chiral symmetry breaking, NATO Sci. Ser. B **59** (1980) 135–157.
- [13] D. Harlow, B. Heidenreich, M. Reece, and T. Rudelius, *The Weak Gravity Conjecture: A Review*, arXiv:2201.08380.
- [14] A. B. Zamolodchikov and A. B. Zamolodchikov, Factorized s Matrices in Two-Dimensions as the Exact Solutions of Certain Relativistic Quantum Field Models, Annals Phys. 120 (1979) 253–291.
- [15] S. R. Coleman and J. Mandula, *All Possible Symmetries of the S Matrix*, *Phys. Rev.* **159** (1967) 1251–1256.

- [16] J. Maldacena and A. Zhiboedov, *Constraining Conformal Field Theories with A Higher Spin Symmetry*, J. Phys. A **46** (2013) 214011, [arXiv:1112.1016].
- [17] R. Haag, J. T. Lopuszanski, and M. Sohnius, *All Possible Generators of Supersymmetries of the s Matrix*, *Nucl. Phys. B* **88** (1975) 257.
- [18] W. Nahm, Supersymmetries and their Representations, Nucl. Phys. B 135 (1978) 149.
- [19] P. C. Argyres, J. J. Heckman, K. Intriligator, and M. Martone, *Snowmass White Paper on SCFTs*, arXiv: 2202.07683.
- [20] D. Poland and D. Simmons-Duffin, *Snowmass White Paper: The Numerical Conformal Bootstrap*, in 2022 *Snowmass Summer Study*, 3, 2022. arXiv:2203.08117.
- [21] D. Gaiotto, A. Kapustin, N. Seiberg, and B. Willett, *Generalized Global Symmetries*, *JHEP* **02** (2015) 172, [arXiv:1412.5148].
- [22] D. Harlow and H. Ooguri, Symmetries in quantum field theory and quantum gravity, Commun. Math. Phys. 383 (2021), no. 3 1669–1804, [arXiv:1810.05338].
- [23] T. T. Dumitrescu and N. Seiberg, *Supercurrents and Brane Currents in Diverse Dimensions*, *JHEP* **07** (2011) 095, [arXiv:1106.0031].
- [24] A. Kapustin and N. Seiberg, *Coupling a QFT to a TQFT and Duality*, *JHEP* **04** (2014) 001, [arXiv:1401.0740].
- [25] E. Sharpe, Notes on generalized global symmetries in QFT, Fortsch. Phys. 63 (2015) 659–682, [arXiv:1508.04770].
- [26] D. Gaiotto, A. Kapustin, Z. Komargodski, and N. Seiberg, *Theta, Time Reversal, and Temperature*, *JHEP* **05** (2017) 091, [arXiv:1703.00501].
- [27] D. M. Hofman and N. Iqbal, Generalized global symmetries and holography, SciPost Phys. 4 (2018), no. 1 005, [arXiv:1707.08577].
- [28] P.-S. Hsin, H. T. Lam, and N. Seiberg, Comments on One-Form Global Symmetries and Their Gauging in 3d and 4d, SciPost Phys. 6 (2019), no. 3 039, [arXiv:1812.04716].
- [29] P.-S. Hsin and S.-H. Shao, Lorentz Symmetry Fractionalization and Dualities in (2+1)d, SciPost Phys. 8 (2020) 018, [arXiv:1909.07383].
- [30] D. R. Morrison, S. Schafer-Nameki, and B. Willett, *Higher-Form Symmetries in 5d*, *JHEP* **09** (2020) 024, [arXiv:2005.12296].
- [31] M. Del Zotto, I. n. García Etxebarria, and S. S. Hosseini, *Higher form symmetries of Argyres-Douglas theories*, *JHEP* **10** (2020) 056, [arXiv:2007.15603].
- [32] S. Gukov, P.-S. Hsin, and D. Pei, *Generalized global symmetries of T[M] theories. Part I, JHEP* **04** (2021) 232, [arXiv:2010.15890].
- [33] A. M. Polyakov, Quark Confinement and Topology of Gauge Groups, Nucl. Phys. B 120 (1977) 429–458.

- [34] I. Affleck, J. A. Harvey, and E. Witten, *Instantons and (Super)Symmetry Breaking in (2+1)-Dimensions, Nucl. Phys. B* **206** (1982) 413–439.
- [35] A. Kovner and B. Rosenstein, New look at QED in four-dimensions: The Photon as a Goldstone boson and the topological interpretation of electric charge, Phys. Rev. D 49 (1994) 5571–5581, [hep-th/9210154].
- [36] D. M. Hofman and N. Iqbal, Goldstone modes and photonization for higher form symmetries, SciPost Phys. 6 (2019), no. 1 006, [arXiv:1802.09512].
- [37] E. Lake, Higher-form symmetries and spontaneous symmetry breaking, arXiv:1802.07747.
- [38] A. M. Polyakov, Thermal Properties of Gauge Fields and Quark Liberation, Phys. Lett. B 72 (1978) 477–480.
- [39] L. Bhardwaj and Y. Tachikawa, *On finite symmetries and their gauging in two dimensions*, *JHEP* **03** (2018) 189, [arXiv:1704.02330].
- [40] Y. Tachikawa, On gauging finite subgroups, SciPost Phys. 8 (2020), no. 1 015, [arXiv:1712.09542].
- [41] K. Roumpedakis, S. Seifnashri, and S.-H. Shao, *Higher Gauging and Non-invertible Condensation Defects*, arXiv:2204.02407.
- [42] L. Kong and X.-G. Wen, *Braided fusion categories, gravitational anomalies, and the mathematical framework for topological orders in any dimensions*, arXiv:1405.5858.
- [43] D. V. Else and C. Nayak, Cheshire charge in (3+1)-dimensional topological phases, Phys. Rev. B **96** (2017), no. 4 045136, [arXiv:1702.02148].
- [44] D. Gaiotto and T. Johnson-Freyd, Condensations in higher categories, arXiv:1905.09566.
- [45] L. Kong, T. Lan, X.-G. Wen, Z.-H. Zhang, and H. Zheng, *Algebraic higher symmetry and categorical symmetry a holographic and entanglement view of symmetry*, *Phys. Rev. Res.* **2** (2020), no. 4 043086, [arXiv:2005.14178].
- [46] T. Johnson-Freyd, (3+1)D topological orders with only a \mathbb{Z}_2 -charged particle, arXiv:2011.11165.
- [47] Y. Choi, C. Córdova, P.-S. Hsin, H. T. Lam, and S.-H. Shao, *Non-invertible Condensation, Duality, and Triality Defects in 3+1 Dimensions*, arXiv:2204.09025.
- [48] J. C. Baez and A. D. Lauda, Higher-dimensional algebra v: 2-groups, Version 3 (2004) 423–491.
- [49] J. Baez and U. Schreiber, Higher gauge theory: 2-connections on 2-bundles, hep-th/0412325.
- [50] U. Schreiber and K. Waldorf, Connections on non-Abelian Gerbes and their Holonomy, Theory Appl. Categ. 28 (2013) 476–540, [arXiv:0808.1923].
- [51] A. Kapustin and R. Thorngren, *Topological Field Theory on a Lattice, Discrete Theta-Angles and Confinement, Adv. Theor. Math. Phys.* **18** (2014), no. 5 1233–1247, [arXiv:1308.2926].
- [52] A. Kapustin and R. Thorngren, *Higher symmetry and gapped phases of gauge theories*, arXiv:1309.4721.

- [53] S. Gukov and A. Kapustin, *Topological Quantum Field Theory, Nonlocal Operators, and Gapped Phases of Gauge Theories*, arXiv:1307.4793.
- [54] R. Thorngren and C. von Keyserlingk, *Higher SPT's and a generalization of anomaly in-flow*, arXiv:1511.02929.
- [55] D. Gaiotto and T. Johnson-Freyd, *Symmetry Protected Topological phases and Generalized Cohomology*, *JHEP* **05** (2019) 007, [arXiv:1712.07950].
- [56] L. Bhardwaj, D. Gaiotto, and A. Kapustin, *State sum constructions of spin-TFTs and string net constructions of fermionic phases of matter*, *JHEP* **04** (2017) 096, [arXiv:1605.01640].
- [57] C. Delcamp and A. Tiwari, From gauge to higher gauge models of topological phases, JHEP 10 (2018) 049, [arXiv:1802.10104].
- [58] C. Córdova, T. T. Dumitrescu, and K. Intriligator, *Exploring 2-Group Global Symmetries*, *JHEP* **02** (2019) 184, [arXiv:1802.04790].
- [59] F. Benini, C. Córdova, and P.-S. Hsin, On 2-Group Global Symmetries and their Anomalies, JHEP 03 (2019) 118, [arXiv:1803.09336].
- [60] Y. Tanizaki and M. Ünsal, *Modified instanton sum in QCD and higher-groups*, *JHEP* **03** (2020) 123, [arXiv:1912.01033].
- [61] I. Bah, F. Bonetti, and R. Minasian, *Discrete and higher-form symmetries in SCFTs from wrapped M5-branes*, *JHEP* **03** (2021) 196, [arXiv:2007.15003].
- [62] L. Bhardwaj and S. Schäfer-Nameki, *Higher-form symmetries of 6d and 5d theories*, *JHEP* **02** (2021) 159, [arXiv:2008.09600].
- [63] P.-S. Hsin and H. T. Lam, *Discrete theta angles, symmetries and anomalies, SciPost Phys.* **10** (2021), no. 2 032, [arXiv:2007.05915].
- [64] C. Córdova, T. T. Dumitrescu, and K. Intriligator, 2-Group Global Symmetries and Anomalies in Six-Dimensional Quantum Field Theories, JHEP **04** (2021) 252, [arXiv:2009.00138].
- [65] M. Del Zotto and K. Ohmori, 2-Group Symmetries of 6D Little String Theories and T-Duality, Annales Henri Poincare 22 (2021), no. 7 2451–2474, [arXiv:2009.03489].
- [66] N. Iqbal and N. Poovuttikul, 2-group global symmetries, hydrodynamics and holography, arXiv:2010.00320.
- [67] Y. Hidaka, M. Nitta, and R. Yokokura, *Global 3-group symmetry and 't Hooft anomalies in axion electrodynamics*, *JHEP* **01** (2021) 173, [arXiv:2009.14368].
- [68] T. D. Brennan and C. Córdova, Axions, higher-groups, and emergent symmetry, JHEP **02** (2022) 145, [arXiv:2011.09600].
- [69] Y. Lee, K. Ohmori, and Y. Tachikawa, *Matching higher symmetries across Intriligator-Seiberg duality*, *JHEP* **10** (2021) 114, [arXiv:2108.05369].
- [70] F. Apruzzi, L. Bhardwaj, J. Oh, and S. Schafer-Nameki, *The Global Form of Flavor Symmetries and 2-Group Symmetries in 5d SCFTs*, arXiv:2105.08724.

- [71] F. Apruzzi, F. Bonetti, I. n. G. Etxebarria, S. S. Hosseini, and S. Schafer-Nameki, *Symmetry TFTs from String Theory*, arXiv:2112.02092.
- [72] L. Bhardwaj, 2-Group Symmetries in Class S, arXiv:2107.06816.
- [73] F. Apruzzi, L. Bhardwaj, D. S. W. Gould, and S. Schafer-Nameki, 2-Group Symmetries and their Classification in 6d, arXiv:2110.14647.
- [74] P.-S. Hsin, W. Ji, and C.-M. Jian, Exotic Invertible Phases with Higher-Group Symmetries, arXiv:2105.09454.
- [75] M. Del Zotto, I. n. G. Etxebarria, and S. Schafer-Nameki, 2-Group Symmetries and M-Theory, arXiv: 2203.10097.
- [76] M. Cvetič, J. J. Heckman, M. Hübner, and E. Torres, *0-Form, 1-Form and 2-Group Symmetries via Cutting and Gluing of Orbifolds*, arXiv:2203.10102.
- [77] E. Sharpe, An introduction to decomposition, arXiv:2204.09117.
- [78] T. Pantev, D. Robbins, E. Sharpe, and T. Vandermeulen, *Orbifolds by 2-groups and decomposition*, arXiv:2204.13708.
- [79] M. Del Zotto, J. J. Heckman, S. N. Meynet, R. Moscrop, and H. Y. Zhang, *Higher Symmetries of 5d Orbifold SCFTs*, arXiv:2201.08372.
- [80] M. Del Zotto and I. n. G. Etxebarria, Global Structures from the Infrared, arXiv: 2204.06495.
- [81] G. W. Moore and N. Seiberg, *Classical and Quantum Conformal Field Theory*, *Commun. Math. Phys.* **123** (1989) 177.
- [82] K. A. Intriligator, Bonus Symmetry in Conformal Field Theory, Nucl. Phys. B 332 (1990) 541–565.
- [83] A. N. Schellekens and S. Yankielowicz, *Extended Chiral Algebras and Modular Invariant Partition Functions*, *Nucl. Phys. B* **327** (1989) 673–703.
- [84] G. W. Moore and N. Seiberg, Taming the Conformal Zoo, Phys. Lett. B 220 (1989) 422–430.
- [85] J. Fuchs, I. Runkel, and C. Schweigert, *TFT construction of RCFT correlators 1. Partition functions*, *Nucl. Phys. B* **646** (2002) 353–497, [hep-th/0204148].
- [86] E. P. Verlinde, Fusion Rules and Modular Transformations in 2D Conformal Field Theory, Nucl. Phys. B **300** (1988) 360–376.
- [87] V. B. Petkova and J. B. Zuber, *Generalized twisted partition functions*, *Phys. Lett. B* **504** (2001) 157–164, [hep-th/0011021].
- [88] J. Frohlich, J. Fuchs, I. Runkel, and C. Schweigert, *Kramers-Wannier duality from conformal defects*, *Phys. Rev. Lett.* **93** (2004) 070601, [cond-mat/0404051].
- [89] J. Frohlich, J. Fuchs, I. Runkel, and C. Schweigert, *Duality and defects in rational conformal field theory*, *Nucl. Phys.* **B763** (2007) 354–430, [hep-th/0607247].
- [90] J. Frohlich, J. Fuchs, I. Runkel, and C. Schweigert, *Defect lines, dualities, and generalised orbifolds*, in *Proceedings, 16th International Congress on Mathematical Physics (ICMP09): Prague, Czech Republic, August 3-8, 2009*, 2009. arXiv:0909.5013.

- [91] C.-M. Chang, Y.-H. Lin, S.-H. Shao, Y. Wang, and X. Yin, *Topological Defect Lines and Renormalization Group Flows in Two Dimensions*, *JHEP* **01** (2019) 026, [arXiv:1802.04445].
- [92] N. Carqueville and I. Runkel, *Orbifold completion of defect bicategories*, *Quantum Topol.* **7** (2016) 203, [arXiv:1210.6363].
- [93] I. Brunner, N. Carqueville, and D. Plencner, A quick guide to defect orbifolds, Proc. Symp. Pure Math. 88 (2014) 231–242, [arXiv:1310.0062].
- [94] Z. Komargodski, K. Ohmori, K. Roumpedakis, and S. Seifnashri, *Symmetries and strings of adjoint QCD*₂, *JHEP* **03** (2021) 103, [arXiv:2008.07567].
- [95] D. Gaiotto and J. Kulp, Orbifold groupoids, JHEP 02 (2021) 132, [arXiv:2008.05960].
- [96] T.-C. Huang, Y.-H. Lin, and S. Seifnashri, Construction of two-dimensional topological field theories with non-invertible symmetries, arXiv:2110.02958.
- [97] R. Thorngren and Y. Wang, Fusion Category Symmetry I: Anomaly In-Flow and Gapped Phases, arXiv:1912.02817.
- [98] R. Thorngren and Y. Wang, Fusion Category Symmetry II: Categoriosities at c = 1 and Beyond, arXiv: 2106.12577.
- [99] R. Thorngren, *Anomalies and Bosonization, Commun. Math. Phys.* **378** (2020), no. 3 1775–1816, [arXiv:1810.04414].
- [100] W. Ji, S.-H. Shao, and X.-G. Wen, *Topological Transition on the Conformal Manifold, Phys. Rev. Res.* 2 (2020), no. 3 033317, [arXiv:1909.01425].
- [101] Y.-H. Lin and S.-H. Shao, *Duality Defect of the Monster CFT*, *J. Phys. A* **54** (2021), no. 6 065201, [arXiv:1911.00042].
- [102] J. Kaidi, K. Ohmori, and Y. Zheng, *Kramers-Wannier-like Duality Defects in (3+1)D Gauge Theories*, *Phys. Rev. Lett.* **128** (2022), no. 11 111601, [arXiv:2111.01141].
- [103] I. M. Burbano, J. Kulp, and J. Neuser, Duality Defects in E₈, arXiv:2112.14323.
- [104] J. Kaidi, Z. Komargodski, K. Ohmori, S. Seifnashri, and S.-H. Shao, *Higher central charges and topological boundaries in 2+1-dimensional TQFTs*, arXiv:2107.13091.
- [105] M. Buican and R. Radhakrishnan, Galois orbits of TQFTs: symmetries and unitarity, JHEP **01** (2022) 004, [arXiv:2109.02766].
- [106] M. Yu, Gauging Categorical Symmetries in 3d Topological Orders and Bulk Reconstruction, arXiv:2111.13697.
- [107] F. Benini, C. Copetti, and L. Di Pietro, *Factorization and Global Symmetries in Holography*, arXiv:2203.09537.
- [108] M. Koide, Y. Nagoya, and S. Yamaguchi, Non-invertible topological defects in 4-dimensional \mathbb{Z}_2 pure lattice gauge theory, arXiv:2109.05992.
- [109] Y. Choi, C. Córdova, P.-S. Hsin, H. T. Lam, and S.-H. Shao, *Non-Invertible Duality Defects in 3+1 Dimensions*, arXiv:2111.01139.

- [110] Y. Hayashi and Y. Tanizaki, Non-invertible self-duality defects of Cardy-Rabinovici model and mixed gravitational anomaly, arXiv:2204.07440.
- [111] L. Bhardwaj, L. Bottini, S. Schafer-Nameki, and A. Tiwari, *Non-Invertible Higher-Categorical Symmetries*, arXiv:2204.06564.
- [112] J. Kaidi, G. Zafrir, and Y. Zheng, *Non-Invertible Symmetries of* $\mathcal{N}=4$ *SYM and Twisted Compactification*, arXiv:2205.01104.
- [113] A. Kapustin and N. Saulina, Surface operators in 3d Topological Field Theory and 2d Rational Conformal Field Theory, arXiv:1012.0911.
- [114] T. Rudelius and S.-H. Shao, *Topological Operators and Completeness of Spectrum in Discrete Gauge Theories*, *JHEP* **12** (2020) 172, [arXiv:2006.10052].
- [115] B. Heidenreich, J. McNamara, M. Montero, M. Reece, T. Rudelius, and I. Valenzuela, Non-Invertible Global Symmetries and Completeness of the Spectrum, JHEP **09** (2021) 203, [arXiv:2104.07036].
- [116] J. McNamara, Gravitational Solitons and Completeness, arXiv:2108.02228.
- [117] C. Córdova, K. Ohmori, and T. Rudelius, Generalized Symmetry Breaking Scales and Weak Gravity Conjectures, arXiv:2202.05866.
- [118] G. Arias-Tamargo and D. Rodriguez-Gomez, *Non-Invertible Symmetries from Discrete Gauging and Completeness of the Spectrum*, arXiv:2204.07523.
- [119] U. Grimm and G. M. Schutz, *The Spin 1/2 XXZ Heisenberg chain, the quantum algebra* U(q)[sl(2)], and duality transformations for minimal models, J. Statist. Phys. **71** (1993) 921–964, [hep-th/0111083].
- [120] A. Feiguin, S. Trebst, A. W. W. Ludwig, M. Troyer, A. Kitaev, Z. Wang, and M. H. Freedman, Interacting anyons in topological quantum liquids: The golden chain, Phys. Rev. Lett. 98 (2007), no. 16 160409, [cond-mat/0612341].
- [121] M. Hauru, G. Evenbly, W. W. Ho, D. Gaiotto, and G. Vidal, *Topological conformal defects with tensor networks*, *Phys. Rev. B* **94** (2016), no. 11 115125, [arXiv:1512.03846].
- [122] D. Aasen, R. S. K. Mong, and P. Fendley, *Topological Defects on the Lattice I: The Ising model, J. Phys. A* **49** (2016), no. 35 354001, [arXiv:1601.07185].
- [123] M. Buican and A. Gromov, Anyonic Chains, Topological Defects, and Conformal Field Theory, Commun. Math. Phys. **356** (2017), no. 3 1017–1056, [arXiv:1701.02800].
- [124] D. Aasen, P. Fendley, and R. S. K. Mong, *Topological Defects on the Lattice: Dualities and Degeneracies*, arXiv:2008.08598.
- [125] K. Inamura, On lattice models of gapped phases with fusion category symmetries, arXiv:2110.12882.
- [126] T.-C. Huang, Y.-H. Lin, K. Ohmori, Y. Tachikawa, and M. Tezuka, *Numerical evidence for a Haagerup conformal field theory*, arXiv:2110.03008.

- [127] R. Vanhove, L. Lootens, M. Van Damme, R. Wolf, T. Osborne, J. Haegeman, and F. Verstraete, *A critical lattice model for a Haagerup conformal field theory*, arXiv:2110.03532.
- [128] Y. Liu, Y. Zou, and S. Ryu, Operator fusion from wavefunction overlaps: Universal finite-size corrections and application to Haagerup model, arXiv:2203.14992.
- [129] W. Ji and X.-G. Wen, Categorical symmetry and noninvertible anomaly in symmetry-breaking and topological phase transitions, Phys. Rev. Res. 2 (2020), no. 3 033417, [arXiv:1912.13492].
- [130] A. Paramekanti, L. Balents, and M. P. A. Fisher, *Ring exchange, the exciton bose liquid, and bosonization in two dimensions, Phys. Rev. B* **66** (Aug, 2002) 054526.
- [131] C. Chamon, Quantum glassiness in strongly correlated clean systems: An example of topological overprotection, Phys. Rev. Lett. **94** (Jan, 2005) 040402.
- [132] J. Haah, Local stabilizer codes in three dimensions without string logical operators, Phys. Rev. A 83 (Apr, 2011) 042330.
- [133] S. Vijay, J. Haah, and L. Fu, Fracton Topological Order, Generalized Lattice Gauge Theory and Duality, Phys. Rev. B 94 (2016), no. 23 235157, [arXiv:1603.04442].
- [134] R. M. Nandkishore and M. Hermele, *Fractons, Ann. Rev. Condensed Matter Phys.* **10** (2019) 295–313, [arXiv:1803.11196].
- [135] M. Pretko, X. Chen, and Y. You, Fracton Phases of Matter, Int. J. Mod. Phys. A **35** (2020), no. 06 2030003, [arXiv:2001.01722].
- [136] K. T. Grosvenor, C. Hoyos, F. Peña Benitez, and P. Surówka, *Space-dependent symmetries and fractons*, arXiv:2112.00531.
- [137] T. Brauner, S. A. Hartnoll, P. Kovtun, H. Liu, M. Mezei, A. Nicolis, R. Penco, S.-H. Shao, and D. T. Son, *Snowmass White Paper: Effective Field Theories for Condensed Matter Systems*, in 2022 Snowmass Summer Study, 3, 2022. arXiv:2203.10110.
- [138] N. Seiberg, Field Theories With a Vector Global Symmetry, SciPost Phys. 8 (2020), no. 4 050, [arXiv:1909.10544].
- [139] M. Qi, L. Radzihovsky, and M. Hermele, *Fracton phases via exotic higher-form symmetry-breaking*, *Annals Phys.* **424** (2021) 168360, [arXiv:2010.02254].
- [140] C. D. Batista and Z. Nussinov, Generalized Elitzur's theorem and dimensional reduction, Phys. Rev. B 72 (2005) 045137, [cond-mat/0410599].
- [141] D. A. Johnston, M. Mueller, and W. Janke, *Plaquette Ising models, degeneracy and scaling, European Physical Journal Special Topics* **226** (Mar., 2017) [arXiv:1612.00060].
- [142] N. Seiberg and S.-H. Shao, Exotic Symmetries, Duality, and Fractons in 2+1-Dimensional Quantum Field Theory, SciPost Phys. **10** (2021), no. 2 027, [arXiv:2003.10466].
- [143] N. Seiberg and S.-H. Shao, Exotic U(1) Symmetries, Duality, and Fractons in 3+1-Dimensional Quantum Field Theory, SciPost Phys. **9** (2020), no. 4 046, [arXiv:2004.00015].
- [144] N. Seiberg and S.-H. Shao, Exotic \mathbb{Z}_N symmetries, duality, and fractons in 3+1-dimensional quantum field theory, SciPost Phys. 10 (2021), no. 1 003, [arXiv:2004.06115].

- [145] P. Gorantla, H. T. Lam, N. Seiberg, and S.-H. Shao, Low-energy limit of some exotic lattice theories and UV/IR mixing, Phys. Rev. B 104 (2021), no. 23 235116, [arXiv:2108.00020].
- [146] J. Distler, A. Karch, and A. Raz, Spontaneously Broken Subsystem Symmetries, arXiv:2110.12611.
- [147] P. Gorantla, H. T. Lam, N. Seiberg, and S.-H. Shao, A modified Villain formulation of fractons and other exotic theories, J. Math. Phys. **62** (2021), no. 10 102301, [arXiv:2103.01257].
- [148] F. J. Burnell, T. Devakul, P. Gorantla, H. T. Lam, and S.-H. Shao, *Anomaly Inflow for Subsystem Symmetries*, arXiv:2110.09529.
- [149] R. Raussendorf, C. Okay, D.-S. Wang, D. T. Stephen, and H. P. Nautrup, *Computationally universal phase of quantum matter*, *Phys. Rev. Lett.* **122** (Mar, 2019) 090501.
- [150] Y. You, T. Devakul, F. J. Burnell, and S. L. Sondhi, *Subsystem symmetry protected topological order*, *Phys. Rev. B* **98** (Jul, 2018) 035112.
- [151] Y. You, T. Devakul, F. J. Burnell, and S. L. Sondhi, *Symmetric Fracton Matter: Twisted and Enriched, Annals Phys.* **416** (2020) 168140, [arXiv:1805.09800].
- [152] T. Devakul, D. J. Williamson, and Y. You, *Classification of subsystem symmetry-protected topological phases*, *Phys. Rev. B* **98** (2018), no. 23 235121, [arXiv:1808.05300].
- [153] D. T. Stephen, H. P. Nautrup, J. Bermejo-Vega, J. Eisert, and R. Raussendorf, *Subsystem symmetries, quantum cellular automata, and computational phases of quantum matter, Quantum* **3** (May, 2019) 142.
- [154] T. Devakul, W. Shirley, and J. Wang, Strong planar subsystem symmetry-protected topological phases and their dual fracton orders, Phys. Rev. Res. 2 (2020), no. 1 012059, [arXiv:1910.01630].
- [155] S. Yamaguchi, Gapless edge modes in (4+1)-dimensional topologically massive tensor gauge theory and anomaly inflow for subsystem symmetry, arXiv:2110.12861.
- [156] D. T. Stephen, A. Dua, J. Garre-Rubio, D. J. Williamson, and M. Hermele, *Fractionalization of subsystem symmetries in two dimensions*, arXiv:2203.13244.
- [157] D. J. Williamson, Fractal symmetries: Ungauging the cubic code, Phys. Rev. B 94 (2016), no. 15 155128, [arXiv:1603.05182].
- [158] K. Slagle and Y. B. Kim, Quantum Field Theory of X-Cube Fracton Topological Order and Robust Degeneracy from Geometry, Phys. Rev. B 96 (2017), no. 19 195139, [arXiv:1708.04619].
- [159] W. Shirley, K. Slagle, and X. Chen, *Foliated fracton order from gauging subsystem symmetries*, *SciPost Phys.* **6** (2019), no. 4 041, [arXiv:1806.08679].
- [160] A. Gromov, Towards classification of Fracton phases: the multipole algebra, Phys. Rev. X 9 (2019), no. 3 031035, [arXiv:1812.05104].
- [161] P. Gorantla, H. T. Lam, N. Seiberg, and S.-H. Shao, fcc lattice, checkerboards, fractons, and quantum field theory, Phys. Rev. B 103 (2021), no. 20 205116, [arXiv:2010.16414].

- [162] Z.-C. Gu and X.-G. Wen, A Lattice bosonic model as a quantum theory of gravity, gr-qc/0606100.
- [163] C. Xu, Gapless bosonic excitation without symmetry breaking: An algebraic spin liquid with soft gravitons, Physical Review B 74 (Dec, 2006) [cond-mat/0609595].
- [164] S. Pankov, R. Moessner, and S. L. Sondhi, *Resonating singlet valence plaquettes*, *Physical Review B* **76** (Sep, 2007) [arXiv:0705.0846].
- [165] C. Xu and C. Wu, Resonating plaquette phases in su(4) heisenberg antiferromagnet, Physical Review B 77 (Apr, 2008) [arXiv:0801.0744].
- [166] Z.-C. Gu and X.-G. Wen, *Emergence of helicity +- 2 modes (gravitons) from qbit models*, *Nucl. Phys. B* **863** (2012) 90–129, [arXiv:0907.1203].
- [167] A. Rasmussen, Y.-Z. You, and C. Xu, Stable Gapless Bose Liquid Phases without any Symmetry, arXiv e-prints (Jan., 2016) arXiv:1601.08235, [arXiv:1601.08235].
- [168] M. Pretko, Subdimensional Particle Structure of Higher Rank U(1) Spin Liquids, Phys. Rev. B 95 (2017), no. 11 115139, [arXiv:1604.05329].
- [169] M. Pretko, Generalized Electromagnetism of Subdimensional Particles: A Spin Liquid Story, Phys. Rev. B 96 (2017), no. 3 035119, [arXiv:1606.08857].
- [170] A. Prem, M. Pretko, and R. Nandkishore, *Emergent Phases of Fractonic Matter*, *Phys. Rev. B* **97** (2018), no. 8 085116, [arXiv:1709.09673].
- [171] M. Pretko and L. Radzihovsky, *Symmetry-Enriched Fracton Phases from Supersolid Duality*, *Phys.Rev.Lett.* **121** (Dec., 2018) 235301, [arXiv:1808.05616].
- [172] H. Ma and M. Pretko, *Higher-rank deconfined quantum criticality at the Lifshitz transition and the exciton Bose condensate*, *Phys.Rev.B* **98** (Sept., 2018) 125105, [arXiv:1803.04980].
- [173] D. Bulmash and M. Barkeshli, *The Higgs Mechanism in Higher-Rank Symmetric U*(1) *Gauge Theories*, *Phys. Rev. B* **97** (2018), no. 23 235112, [arXiv:1802.10099].
- [174] H. Ma, M. Hermele, and X. Chen, Fracton topological order from the Higgs and partial-confinement mechanisms of rank-two gauge theory, Phys.Rev.B 98 (July, 2018) 035111, [arXiv:1802.10108].
- [175] K. Slagle, A. Prem, and M. Pretko, Symmetric Tensor Gauge Theories on Curved Spaces, Annals Phys. 410 (2019) 167910, [arXiv:1807.00827].
- [176] M. Pretko, The Fracton Gauge Principle, Phys. Rev. B 98 (2018), no. 11 115134, [arXiv:1807.11479].
- [177] D. J. Williamson, Z. Bi, and M. Cheng, Fractonic Matter in Symmetry-Enriched U(1) Gauge Theory, Phys. Rev. B 100 (2019), no. 12 125150, [arXiv:1809.10275].
- [178] Y. You, T. Devakul, S. L. Sondhi, and F. J. Burnell, *Fractonic Chern-Simons and BF theories*, *Physical Review Research* **2** (May, 2020) 023249, [arXiv:1904.11530].

- [179] Y. You, F. J. Burnell, and T. L. Hughes, *Multipolar Topological Field Theories: Bridging Higher Order Topological Insulators and Fractons*, *Phys. Rev. B* **103** (2021), no. 24 245128, [arXiv:1909.05868].
- [180] A. Gromov, A Duality Between U(1) Haah Code and 3D Smectic A Phase, arXiv:2002.11817.
- [181] A. Gromov, A. Lucas, and R. M. Nandkishore, *Fracton hydrodynamics*, *Phys. Rev. Res.* **2** (2020), no. 3 033124, [arXiv:2003.09429].
- [182] K. Slagle, Foliated Quantum Field Theory of Fracton Order, Phys. Rev. Lett. 126 (2021), no. 10 101603, [arXiv:2008.03852].
- [183] A. Karch and A. Raz, Reduced Conformal Symmetry, JHEP 04 (2021) 182, [arXiv:2009.12308].
- [184] S. Yamaguchi, Supersymmetric quantum field theory with exotic symmetry in 3+1 dimensions and fermionic fracton phases, PTEP **2021** (2021), no. 6 063B04, [arXiv:2102.04768].
- [185] H. Geng, S. Kachru, A. Karch, R. Nally, and B. C. Rayhaun, Fractons and Exotic Symmetries from Branes, Fortsch. Phys. 69 (2021), no. 11-12 2100133, [arXiv:2108.08322].
- [186] P.-S. Hsin and K. Slagle, Comments on foliated gauge theories and dualities in 3+1d, SciPost Phys. 11 (2021), no. 2 032, [arXiv:2105.09363].
- [187] Y.-H. Du, U. Mehta, D. X. Nguyen, and D. T. Son, *Volume-preserving diffeomorphism as nonabelian higher-rank gauge symmetry*, arXiv:2103.09826.
- [188] L. Bidussi, J. Hartong, E. Have, J. Musaeus, and S. Prohazka, *Fractons, dipole symmetries and curved spacetime*, arXiv:2111.03668.
- [189] A. Jain and K. Jensen, Fractons in curved space, arXiv:2111.03973.
- [190] P. Gorantla, H. T. Lam, N. Seiberg, and S.-H. Shao, *Global Dipole Symmetry, Compact Lifshitz Theory, Tensor Gauge Theory, and Fractons*, arXiv:2201.10589.
- [191] A. Pérez and S. Prohazka, *Asymptotic symmetries and soft charges of fractons*, arXiv:2203.02817.
- [192] Y. You, J. Bibo, T. L. Hughes, and F. Pollmann, Fractonic critical point proximate to a higher-order topological insulator: How does UV blend with IR?, arXiv:2101.01724.
- [193] Z. Zhou, X.-F. Zhang, F. Pollmann, and Y. You, Fractal Quantum Phase Transitions: Critical Phenomena Beyond Renormalization, arXiv:2105.05851.
- [194] Y. You and R. Moessner, Fractonic plaquette-dimer liquid beyond renormalization, arXiv:2106.07664.
- [195] S. Minwalla, M. Van Raamsdonk, and N. Seiberg, *Noncommutative perturbative dynamics*, *JHEP* **02** (2000) 020, [hep-th/9912072].
- [196] D. Bulmash and M. Barkeshli, Generalized U(1) Gauge Field Theories and Fractal Dynamics, arXiv:1806.01855.
- [197] K. Slagle, D. Aasen, and D. Williamson, Foliated Field Theory and String-Membrane-Net Condensation Picture of Fracton Order, SciPost Phys. 6 (2019), no. 4 043, [arXiv:1812.01613].

- [198] P. Gorantla, H. T. Lam, N. Seiberg, and S.-H. Shao, *More Exotic Field Theories in 3+1 Dimensions*, *SciPost Phys.* **9** (2020) 073, [arXiv:2007.04904].
- [199] D. Aasen, D. Bulmash, A. Prem, K. Slagle, and D. J. Williamson, *Topological Defect Networks for Fractors of all Types, Phys. Rev. Res.* **2** (2020) 043165, [arXiv:2002.05166].
- [200] T. Rudelius, N. Seiberg, and S.-H. Shao, Fractons with Twisted Boundary Conditions and Their Symmetries, Phys. Rev. B 103 (2021), no. 19 195113, [arXiv:2012.11592].
- [201] J. Distler, M. Jafry, A. Karch, and A. Raz, *Interacting Fractons in 2+1-Dimensional Quantum Field Theory*, arXiv:2112.05726.
- [202] E. Lake, RG and Stability in the Exciton Bose Liquid, arXiv:2110.02986.
- [203] S.-H. Shao, Symmetries and their generalizations in topological phases of matter, Strings 2021 Review Talk.
- [204] Y. Frishman, A. Schwimmer, T. Banks, and S. Yankielowicz, *The Axial Anomaly and the Bound State Spectrum in Confining Theories*, *Nucl. Phys. B* **177** (1981) 157–171.
- [205] S. R. Coleman and B. Grossman, 't Hooft's Consistency Condition as a Consequence of Analyticity and Unitarity, Nucl. Phys. B 203 (1982) 205–220.
- [206] C. G. Callan, Jr. and J. A. Harvey, *Anomalies and Fermion Zero Modes on Strings and Domain Walls*, *Nucl. Phys. B* **250** (1985) 427–436.
- [207] L. D. Faddeev and S. L. Shatashvili, *Algebraic and Hamiltonian Methods in the Theory of Nonabelian Anomalies, Teor. Mat. Fiz.* **60** (1984) 206–217.
- [208] D. Freed, J. A. Harvey, R. Minasian, and G. W. Moore, *Gravitational anomaly cancellation for M theory five-branes*, *Adv. Theor. Math. Phys.* **2** (1998) 601–618, [hep-th/9803205].
- [209] J. A. Harvey, TASI 2003 lectures on anomalies, 9, 2005. hep-th/0509097.
- [210] E. Witten, Global Aspects of Current Algebra, Nucl. Phys. B 223 (1983) 422–432.
- [211] E. Witten, Current Algebra, Baryons, and Quark Confinement, Nucl. Phys. B 223 (1983) 433-444.
- [212] E. Witten, Nonabelian Bosonization in Two-Dimensions, Commun. Math. Phys. **92** (1984) 455–472.
- [213] D. B. Kaplan, A Method for simulating chiral fermions on the lattice, Phys. Lett. B 288 (1992) 342–347, [hep-lat/9206013].
- [214] R. Thorngren, Framed Wilson Operators, Fermionic Strings, and Gravitational Anomaly in 4d, JHEP 02 (2015) 152, [arXiv:1404.4385].
- [215] K. Ohmori, H. Shimizu, Y. Tachikawa, and K. Yonekura, *Anomaly polynomial of general 6d SCFTs*, *PTEP* **2014** (2014), no. 10 103B07, [arXiv:1408.5572].
- [216] J. Wang, X.-G. Wen, and E. Witten, *A New SU(2) Anomaly*, *J. Math. Phys.* **60** (2019), no. 5 052301, [arXiv:1810.00844].
- [217] Y. Lee, K. Ohmori, and Y. Tachikawa, *Revisiting Wess-Zumino-Witten terms*, *SciPost Phys.* **10** (2021), no. 3 061, [arXiv:2009.00033].

- [218] Y. Lee and Y. Tachikawa, Some comments on 6D global gauge anomalies, PTEP 2021 (2021), no. 8 08B103, [arXiv:2012.11622].
- [219] M. M. Anber and E. Poppitz, Generalized 't Hooft anomalies on non-spin manifolds, JHEP **04** (2020) 097, [arXiv:2002.02037].
- [220] Y. Lee and K. Yonekura, Global anomalies in 8d supergravity, arXiv: 2203.12631.
- [221] A. Kapustin and R. Thorngren, *Anomalies of discrete symmetries in three dimensions and group cohomology*, *Phys. Rev. Lett.* **112** (2014), no. 23 231602, [arXiv:1403.0617].
- [222] A. Kapustin and R. Thorngren, *Anomalies of discrete symmetries in various dimensions and group cohomology*, arXiv:1404.3230.
- [223] A. Kapustin, R. Thorngren, A. Turzillo, and Z. Wang, Fermionic Symmetry Protected Topological Phases and Cobordisms, JHEP 12 (2015) 052, [arXiv:1406.7329].
- [224] J. C. Wang, Z.-C. Gu, and X.-G. Wen, Field theory representation of gauge-gravity symmetry-protected topological invariants, group cohomology and beyond, Phys. Rev. Lett. 114 (2015), no. 3 031601, [arXiv:1405.7689].
- [225] D. Gaiotto and A. Kapustin, *Spin TQFTs and fermionic phases of matter*, *Int. J. Mod. Phys. A* **31** (2016), no. 28n29 1645044, [arXiv:1505.05856].
- [226] M. Guo, P. Putrov, and J. Wang, *Time reversal, SU(N) Yang–Mills and cobordisms: Interacting topological superconductors/insulators and quantum spin liquids in 3+1D, Annals Phys.* **394** (2018) 244–293, [arXiv:1711.11587].
- [227] M. Guo, K. Ohmori, P. Putrov, Z. Wan, and J. Wang, Fermionic Finite-Group Gauge Theories and Interacting Symmetric/Crystalline Orders via Cobordisms, Commun. Math. Phys. 376 (2020), no. 2 1073–1154, [arXiv:1812.11959].
- [228] D. Tong and C. Turner, Notes on 8 Majorana Fermions, SciPost Phys. Lect. Notes 14 (2020) 1, [arXiv:1906.07199].
- [229] A. Grigoletto and P. Putrov, *Spin-cobordisms*, surgeries and fermionic modular bootstrap, arXiv:2106.16247.
- [230] P. B. Smith, A. Karasik, N. Lohitsiri, and D. Tong, *On discrete anomalies in chiral gauge theories*, *JHEP* **01** (2022) 112, [arXiv:2106.06402].
- [231] C.-T. Hsieh, Y. Tachikawa, and K. Yonekura, *Anomaly Inflow and p-Form Gauge Theories*, *Commun. Math. Phys.* **391** (2022), no. 2 495–608, [arXiv:2003.11550].
- [232] Y. Kikuchi and Y. Tanizaki, *Global inconsistency, 't Hooft anomaly, and level crossing in quantum mechanics*, *PTEP* **2017** (2017), no. 11 113B05, [arXiv:1708.01962].
- [233] R. Thorngren, Topological Terms and Phases of Sigma Models, arXiv:1710.02545.
- [234] Y. Tachikawa and K. Yonekura, *Anomalies involving the space of couplings and the Zamolodchikov metric*, *JHEP* **12** (2017) 140, [arXiv:1710.03934].

- [235] Y. Tanizaki and T. Sulejmanpasic, Anomaly and global inconsistency matching: θ -angles, $SU(3)/U(1)^2$ nonlinear sigma model, SU(3) chains and its generalizations, Phys. Rev. B **98** (2018), no. 11 115126, [arXiv:1805.11423].
- [236] N. Seiberg, Y. Tachikawa, and K. Yonekura, *Anomalies of Duality Groups and Extended Conformal Manifolds*, *PTEP* **2018** (2018), no. 7 073B04, [arXiv:1803.07366].
- [237] C.-T. Hsieh, Y. Tachikawa, and K. Yonekura, *Anomaly of the Electromagnetic Duality of Maxwell Theory*, *Phys. Rev. Lett.* **123** (2019), no. 16 161601, [arXiv:1905.08943].
- [238] C. Córdova, D. S. Freed, H. T. Lam, and N. Seiberg, *Anomalies in the Space of Coupling Constants and Their Dynamical Applications I, SciPost Phys.* **8** (2020), no. 1 001, [arXiv:1905.09315].
- [239] C. Córdova, D. S. Freed, H. T. Lam, and N. Seiberg, *Anomalies in the Space of Coupling Constants and Their Dynamical Applications II*, SciPost Phys. 8 (2020), no. 1 002, [arXiv:1905.13361].
- [240] P.-S. Hsin, A. Kapustin, and R. Thorngren, *Berry Phase in Quantum Field Theory: Diabolical Points and Boundary Phenomena*, *Phys. Rev. B* **102** (2020) 245113, [arXiv:2004.10758].
- [241] Y. Choi and K. Ohmori, *Higher Berry Phase of Fermions and Index Theorem*, arXiv:2205.02188.
- [242] A. Kapustin, Symmetry Protected Topological Phases, Anomalies, and Cobordisms: Beyond Group Cohomology, arXiv:1403.1467.
- [243] A. Kapustin, Bosonic Topological Insulators and Paramagnets: a view from cobordisms, arXiv:1404.6659.
- [244] N. Seiberg and E. Witten, *Gapped Boundary Phases of Topological Insulators via Weak Coupling*, *PTEP* **2016** (2016), no. 12 12C101, [arXiv:1602.04251].
- [245] Z. Wan and J. Wang, Higher anomalies, higher symmetries, and cobordisms I: classification of higher-symmetry-protected topological states and their boundary fermionic/bosonic anomalies via a generalized cobordism theory, Ann. Math. Sci. Appl. 4 (2019), no. 2 107–311, [arXiv:1812.11967].
- [246] C. Córdova, K. Ohmori, S.-H. Shao, and F. Yan, *Decorated* Z₂ symmetry defects and their time-reversal anomalies, *Phys. Rev. D* **102** (2020), no. 4 045019, [arXiv:1910.14046].
- [247] K. Yonekura, General anomaly matching by Goldstone bosons, JHEP **03** (2021) 057, [arXiv:2009.04692].
- [248] I. Hason, Z. Komargodski, and R. Thorngren, Anomaly Matching in the Symmetry Broken Phase: Domain Walls, CPT, and the Smith Isomorphism, SciPost Phys. 8 (2020), no. 4 062, [arXiv:1910.14039].
- [249] R. Thorngren and Y. Wang, Anomalous symmetries end at the boundary, JHEP **09** (2021) 017, [arXiv:2012.15861].
- [250] D. Delmastro, D. Gaiotto, and J. Gomis, *Global anomalies on the Hilbert space*, *JHEP* 11 (2021) 142, [arXiv:2101.02218].
- [251] A. Cherman, T. Jacobson, and M. Neuzil, *Universal Deformations*, *SciPost Phys.* **12** (2022) 116, [arXiv:2111.00078].

- [252] L. Alvarez-Gaume and P. H. Ginsparg, *The Structure of Gauge and Gravitational Anomalies*, *Annals Phys.* **161** (1985) 423. [Erratum: Annals Phys. 171, 233 (1986)].
- [253] X.-z. Dai and D. S. Freed, *eta invariants and determinant lines*, *J. Math. Phys.* **35** (1994) 5155–5194, [hep-th/9405012]. [Erratum: J.Math.Phys. 42, 2343–2344 (2001)].
- [254] K. Yonekura, Dai-Freed theorem and topological phases of matter, JHEP **09** (2016) 022, [arXiv:1607.01873].
- [255] E. Witten and K. Yonekura, *Anomaly Inflow and the η-Invariant*, in *The Shoucheng Zhang Memorial Workshop*, 9, 2019. arXiv:1909.08775.
- [256] E. Witten, Anti-de Sitter space and holography, Adv. Theor. Math. Phys. 2 (1998) 253–291, [hep-th/9802150].
- [257] D. S. Freed, Anomalies and Invertible Field Theories, Proc. Symp. Pure Math. 88 (2014) 25–46, [arXiv:1404.7224].
- [258] D. S. Freed, Short-range entanglement and invertible field theories, arXiv:1406.7278.
- [259] K. Yonekura, On the cobordism classification of symmetry protected topological phases, Commun. Math. Phys. 368 (2019), no. 3 1121–1173, [arXiv:1803.10796].
- [260] M. Yamashita and K. Yonekura, *Differential models for the Anderson dual to bordism theories and invertible QFT's, I,* arXiv:2106.09270.
- [261] D. S. Freed and C. Teleman, *Relative quantum field theory, Commun. Math. Phys.* **326** (2014) 459–476, [arXiv:1212.1692].
- [262] S. Elitzur, G. W. Moore, A. Schwimmer, and N. Seiberg, *Remarks on the Canonical Quantization of the Chern-Simons-Witten Theory*, *Nucl. Phys. B* **326** (1989) 108–134.
- [263] E. Witten, *AdS / CFT correspondence and topological field theory*, *JHEP* **12** (1998) 012, [hep-th/9812012].
- [264] S. Monnier, The anomaly field theories of six-dimensional (2,0) superconformal theories, Adv. Theor. Math. Phys. 22 (2018) 2035–2089, [arXiv:1706.01903].
- [265] C. Córdova and K. Ohmori, *Anomaly Obstructions to Symmetry Preserving Gapped Phases*, arXiv:1910.04962.
- [266] C. Córdova and K. Ohmori, *Anomaly Constraints on Gapped Phases with Discrete Chiral Symmetry*, *Phys. Rev. D* **102** (2020), no. 2 025011, [arXiv:1912.13069].
- [267] N. Seiberg, *Electric magnetic duality in supersymmetric nonAbelian gauge theories*, *Nucl. Phys. B* **435** (1995) 129–146, [hep-th/9411149].
- [268] K. A. Intriligator and N. Seiberg, *Duality, monopoles, dyons, confinement and oblique confinement in supersymmetric SO(N(c)) gauge theories, Nucl. Phys. B* **444** (1995) 125–160, [hep-th/9503179].
- [269] K. A. Intriligator and N. Seiberg, *Lectures on supersymmetric gauge theories and electric-magnetic duality*, *Nucl. Phys. B Proc. Suppl.* **45BC** (1996) 1–28, [hep-th/9509066].

- [270] Y. Tachikawa and K. Yonekura, On time-reversal anomaly of 2+1d topological phases, PTEP 2017 (2017), no. 3 033B04, [arXiv:1610.07010].
- [271] Y. Tachikawa and K. Yonekura, *More on time-reversal anomaly of 2+1d topological phases*, *Phys. Rev. Lett.* **119** (2017), no. 11 111603, [arXiv:1611.01601].
- [272] F. J. Burnell, Anyon condensation and its applications, Ann. Rev. Condensed Matter Phys. 9 (2018) 307–327, [arXiv:1706.04940].
- [273] P.-S. Hsin and N. Seiberg, *Level/rank Duality and Chern-Simons-Matter Theories*, *JHEP* **09** (2016) 095, [arXiv:1607.07457].
- [274] Y. Lee and Y. Tachikawa, A study of time reversal symmetry of abelian anyons, JHEP **07** (2018) 090, [arXiv:1805.02738].
- [275] D. Delmastro and J. Gomis, Symmetries of Abelian Chern-Simons Theories and Arithmetic, JHEP **03** (2021) 006, [arXiv:1904.12884].
- [276] O. Aharony, Baryons, monopoles and dualities in Chern-Simons-matter theories, JHEP **02** (2016) 093, [arXiv:1512.00161].
- [277] O. Aharony, F. Benini, P.-S. Hsin, and N. Seiberg, *Chern-Simons-matter dualities with SO and USp gauge groups*, *JHEP* **02** (2017) 072, [arXiv:1611.07874].
- [278] F. Benini, P.-S. Hsin, and N. Seiberg, Comments on global symmetries, anomalies, and duality in (2 + 1)d, JHEP **04** (2017) 135, [arXiv:1702.07035].
- [279] C. Córdova, P.-S. Hsin, and N. Seiberg, Global Symmetries, Counterterms, and Duality in Chern-Simons Matter Theories with Orthogonal Gauge Groups, SciPost Phys. 4 (2018), no. 4 021, [arXiv:1711.10008].
- [280] F. Benini, *Three-dimensional dualities with bosons and fermions*, *JHEP* **02** (2018) 068, [arXiv:1712.00020].
- [281] C. Córdova, P.-S. Hsin, and N. Seiberg, *Time-Reversal Symmetry, Anomalies, and Dualities in* (2+1)d, *SciPost Phys.* **5** (2018), no. 1 006, [arXiv:1712.08639].
- [282] F. Benini and S. Benvenuti, $\mathcal{N} = 1$ QED in 2 + 1 dimensions: dualities and enhanced symmetries, JHEP **05** (2021) 176, [arXiv:1804.05707].
- [283] C. Córdova, P.-S. Hsin, and K. Ohmori, *Exceptional Chern-Simons-Matter Dualities*, *SciPost Phys.* **7** (2019), no. 4 056, [arXiv:1812.11705].
- [284] V. Bashmakov, J. Gomis, Z. Komargodski, and A. Sharon, *Phases of* $\mathcal{N} = 1$ theories in 2 + 1 dimensions, *JHEP* **07** (2018) 123, [arXiv:1802.10130].
- [285] D. S. Freed, Z. Komargodski, and N. Seiberg, *The Sum Over Topological Sectors and* θ *in the* 2+1-Dimensional \mathbb{CP}^1 σ -Model, Commun. Math. Phys. **362** (2018), no. 1 167–183, [arXiv:1707.05448].
- [286] Z. Komargodski and N. Seiberg, A symmetry breaking scenario for QCD₃, JHEP **01** (2018) 109, [arXiv:1706.08755].

- [287] Z. Komargodski, A. Sharon, R. Thorngren, and X. Zhou, Comments on Abelian Higgs Models and Persistent Order, SciPost Phys. 6 (2019), no. 1 003, [arXiv:1705.04786].
- [288] J. Gomis, Z. Komargodski, and N. Seiberg, *Phases Of Adjoint QCD*₃ *And Dualities, SciPost Phys.* **5** (2018), no. 1 007, [arXiv:1710.03258].
- [289] Z. Komargodski, T. Sulejmanpasic, and M. Ünsal, Walls, anomalies, and deconfinement in quantum antiferromagnets, Phys. Rev. B 97 (2018), no. 5 054418, [arXiv:1706.05731].
- [290] C. Choi, D. Delmastro, J. Gomis, and Z. Komargodski, *Dynamics of QCD*₃ with Rank-Two Quarks And Duality, JHEP **03** (2020) 078, [arXiv:1810.07720].
- [291] A. Armoni, T. T. Dumitrescu, G. Festuccia, and Z. Komargodski, *Metastable vacua in large-N QCD*₃, *JHEP* **01** (2020) 004, [arXiv:1905.01797].
- [292] P. Putrov, J. Wang, and S.-T. Yau, *Braiding Statistics and Link Invariants of Bosonic/Fermionic Topological Quantum Matter in 2+1 and 3+1 dimensions*, *Annals Phys.* **384** (2017) 254–287, [arXiv:1612.09298].
- [293] H. Shimizu and K. Yonekura, *Anomaly constraints on deconfinement and chiral phase transition*, *Phys. Rev. D* **97** (2018), no. 10 105011, [arXiv:1706.06104].
- [294] A. Cherman, S. Sen, M. Unsal, M. L. Wagman, and L. G. Yaffe, *Order parameters and color-flavor center symmetry in QCD*, *Phys. Rev. Lett.* **119** (2017), no. 22 222001, [arXiv:1706.05385].
- [295] I. n. García-Etxebarria and M. Montero, *Dai-Freed anomalies in particle physics*, *JHEP* **08** (2019) 003, [arXiv:1808.00009].
- [296] K. Aitken, A. Cherman, and M. Ünsal, *Dihedral symmetry in SU(N) Yang-Mills theory*, *Phys. Rev.* D **100** (2019), no. 8 085004, [arXiv:1804.05845].
- [297] C. Córdova and T. T. Dumitrescu, Candidate Phases for SU(2) Adjoint QCD₄ with Two Flavors from $\mathcal{N} = 2$ Supersymmetric Yang-Mills Theory, arXiv:1806.09592.
- [298] Z. Wan, J. Wang, and Y. Zheng, New higher anomalies, SU(N) Yang–Mills gauge theory and \mathbb{CP}^{N-1} sigma model, Annals Phys. **414** (2020) 168074, [arXiv:1812.11968].
- [299] K. Yonekura, Anomaly matching in QCD thermal phase transition, JHEP **05** (2019) 062, [arXiv:1901.08188].
- [300] J. Wang, Y.-Z. You, and Y. Zheng, Gauge enhanced quantum criticality and time reversal deconfined domain wall: SU(2) Yang-Mills dynamics with topological terms, Phys. Rev. Res. 2 (2020), no. 1 013189, [arXiv:1910.14664].
- [301] Z. Wan, J. Wang, and Y. Zheng, Higher anomalies, higher symmetries, and cobordisms II: Lorentz symmetry extension and enriched bosonic / fermionic quantum gauge theory, Ann. Math. Sci. Appl. **05** (2020), no. 2 171–257, [arXiv:1912.13504].
- [302] Z. Wan and J. Wang, Higher anomalies, higher symmetries, and cobordisms III: QCD matter phases anew, Nucl. Phys. B 957 (2020) 115016, [arXiv:1912.13514].
- [303] Z. Wan and J. Wang, Beyond Standard Models and Grand Unifications: Anomalies, Topological Terms, and Dynamical Constraints via Cobordisms, JHEP 07 (2020) 062, [arXiv:1910.14668].

- [304] A. Cherman, T. Jacobson, S. Sen, and L. G. Yaffe, *Higgs-confinement phase transitions with fundamental representation matter*, *Phys. Rev. D* **102** (2020), no. 10 105021, [arXiv:2007.08539].
- [305] E. Poppitz and F. D. Wandler, *Topological terms and anomaly matching in effective field theories on* $\mathbb{R}^3 \times \mathbb{S}^1$:. *Part I. Abelian symmetries and intermediate scales, JHEP* **01** (2021) 091, [arXiv:2009.14667].
- [306] M. M. Anber and S. Baker, *Natural inflation, strong dynamics, and the role of generalized anomalies*, *Phys. Rev. D* **102** (2020), no. 10 103515, [arXiv:2008.05491].
- [307] S. S. Razamat and D. Tong, *Gapped Chiral Fermions*, *Phys. Rev. X* **11** (2021), no. 1 011063, [arXiv:2009.05037].
- [308] M. M. Anber, Condensates and anomaly cascade in vector-like theories, JHEP **03** (2021) 191, [arXiv:2101.04132].
- [309] M. M. Anber, S. Hong, and M. Son, New anomalies, TQFTs, and confinement in bosonic chiral gauge theories, JHEP **02** (2022) 062, [arXiv:2109.03245].
- [310] M. M. Anber and E. Poppitz, Nonperturbative effects in the Standard Model with gauged 1-form symmetry, JHEP 12 (2021) 055, [arXiv:2110.02981].
- [311] M. Ünsal, TQFT at work for IR-renormalons, resurgence and Lefschetz decomposition, arXiv:2106.14971.
- [312] A. A. Cox, E. Poppitz, and F. D. Wandler, *The mixed 0-form/1-form anomaly in Hilbert space:* pouring the new wine into old bottles, *JHEP* **10** (2021) 069, [arXiv:2106.11442].
- [313] L. Bhardwaj, S. Giacomelli, M. Hübner, and S. Schäfer-Nameki, *Relative Defects in Relative Theories: Trapped Higher-Form Symmetries and Irregular Punctures in Class S*, arXiv:2201.00018.
- [314] J. Wang, Z. Wan, and Y.-Z. You, Cobordism and Deformation Class of the Standard Model, arXiv:2112.14765.
- [315] Y. Tanizaki and M. Ünsal, Center vortex and confinement in Yang-Mills theory and QCD with anomaly-preserving compactifications, PTEP 2022 (2022) 04, [arXiv:2201.06166].
- [316] J. Wang and Y.-Z. You, Symmetric Mass Generation, arXiv: 2204.14271.
- [317] D. Gaiotto, Z. Komargodski, and N. Seiberg, *Time-reversal breaking in QCD*₄, walls, and dualities in 2 + 1 dimensions, *JHEP* **01** (2018) 110, [arXiv:1708.06806].
- [318] V. Bashmakov, F. Benini, S. Benvenuti, and M. Bertolini, *Living on the walls of super-QCD*, *SciPost Phys.* **6** (2019), no. 4 044, [arXiv:1812.04645].
- [319] M. M. Anber and E. Poppitz, *Domain walls in high-T SU(N) super Yang-Mills theory and QCD(adj)*, *JHEP* **05** (2019) 151, [arXiv:1811.10642].
- [320] A. A. Cox, E. Poppitz, and S. S. Y. Wong, *Domain walls and deconfinement: a semiclassical picture of discrete anomaly inflow, JHEP* **12** (2019) 011, [arXiv:1909.10979].

- [321] M. M. Anber and E. Poppitz, *Deconfinement on axion domain walls*, *JHEP* **03** (2020) 124, [arXiv:2001.03631].
- [322] D. Delmastro and J. Gomis, *Domain walls in 4d\mathcal{N} = 1 SYM*, *JHEP* **03** (2021) 259, [arXiv:2004.11395].
- [323] I. n. García-Etxebarria, H. Hayashi, K. Ohmori, Y. Tachikawa, and K. Yonekura, 8d gauge anomalies and the topological Green-Schwarz mechanism, JHEP 11 (2017) 177, [arXiv:1710.04218].
- [324] J. Wang, X.-G. Wen, and E. Witten, Symmetric Gapped Interfaces of SPT and SET States: Systematic Constructions, Phys. Rev. X 8 (2018), no. 3 031048, [arXiv:1705.06728].
- [325] Z. Wan and J. Wang, Adjoint QCD₄, Deconfined Critical Phenomena, Symmetry-Enriched Topological Quantum Field Theory, and Higher Symmetry-Extension, Phys. Rev. D **99** (2019), no. 6 065013, [arXiv:1812.11955].
- [326] J. Wang, K. Ohmori, P. Putrov, Y. Zheng, Z. Wan, M. Guo, H. Lin, P. Gao, and S.-T. Yau, Tunneling Topological Vacua via Extended Operators: (Spin-)TQFT Spectra and Boundary Deconfinement in Various Dimensions, PTEP 2018 (2018), no. 5 053A01, [arXiv:1801.05416].
- [327] R. Kobayashi, K. Ohmori, and Y. Tachikawa, *On gapped boundaries for SPT phases beyond group cohomology*, *JHEP* **11** (2019) 131, [arXiv:1905.05391].
- [328] R. Thorngren, *Topological quantum field theory, symmetry breaking, and finite gauge theory in* 3+1D, Phys. Rev. B **101** (2020), no. 24 245160, [arXiv:2001.11938].
- [329] F. Albertini, M. Del Zotto, I. n. García Etxebarria, and S. S. Hosseini, *Higher Form Symmetries and M-theory*, *JHEP* **12** (2020) 203, [arXiv:2005.12831].
- [330] F. Apruzzi, M. van Beest, D. S. W. Gould, and S. Schäfer-Nameki, *Holography, 1-form symmetries, and confinement, Phys. Rev. D* **104** (2021), no. 6 066005, [arXiv:2104.12764].
- [331] M. Hubner, D. R. Morrison, S. Schafer-Nameki, and Y.-N. Wang, *Generalized Symmetries in F-theory and the Topology of Elliptic Fibrations*, arXiv:2203.10022.
- [332] J. J. Heckman, C. Lawrie, L. Lin, H. Y. Zhang, and G. Zoccarato, 6d SCFTs, Center-Flavor Symmetries, and Stiefel-Whitney Compactifications, arXiv: 2205.03411.
- [333] C. Córdova, T. T. Dumitrescu, and K. Intriligator, *Multiplets of Superconformal Symmetry in Diverse Dimensions*, *JHEP* **03** (2019) 163, [arXiv:1612.00809].
- [334] I. Bah and E. Nardoni, Structure of Anomalies of 4d SCFTs from M5-branes, and Anomaly Inflow, JHEP 03 (2019) 024, [arXiv:1803.00136].
- [335] I. Bah, F. Bonetti, R. Minasian, and E. Nardoni, *Class & Anomalies from M-theory Inflow*, *Phys. Rev. D* **99** (2019), no. 8 086020, [arXiv:1812.04016].
- [336] I. Bah, F. Bonetti, R. Minasian, and P. Weck, *Anomaly Inflow Methods for SCFT Constructions in Type IIB*, *JHEP* **02** (2021) 116, [arXiv:2002.10466].
- [337] L. Bhardwaj, M. Hubner, and S. Schafer-Nameki, *1-form Symmetries of 4d N=2 Class S Theories*, *SciPost Phys.* **11** (2021) 096, [arXiv:2102.01693].

- [338] L. Bhardwaj, M. Hubner, and S. Schafer-Nameki, *Liberating Confinement from Lagrangians:* 1-form Symmetries and Lines in 4d N=1 from 6d N=(2,0), SciPost Phys. **12** (2022) 040, [arXiv:2106.10265].
- [339] L. Bhardwaj, Global Form of Flavor Symmetry Groups in 4d N=2 Theories of Class S, arXiv:2105.08730.
- [340] C. Córdova, T. T. Dumitrescu, and K. Intriligator, *Anomalies, renormalization group flows, and the a-theorem in six-dimensional* (1, 0) theories, JHEP **10** (2016) 080, [arXiv:1506.03807].
- [341] J. J. Heckman and T. Rudelius, *Evidence for C-theorems in 6D SCFTs*, *JHEP* **09** (2015) 218, [arXiv:1506.06753].
- [342] C. Córdova, T. T. Dumitrescu, and K. Intriligator, $\mathcal{N} = (1, 0)$ anomaly multiplet relations in six dimensions, JHEP **07** (2020) 065, [arXiv:1912.13475].
- [343] Y.-H. Lin and S.-H. Shao, *Anomalies and Bounds on Charged Operators*, *Phys. Rev. D* **100** (2019), no. 2 025013, [arXiv:1904.04833].
- [344] M. Yu, Symmetries and anomalies of (1+1)d theories: 2-groups and symmetry fractionalization, *JHEP* **08** (2021) 061, [arXiv:2010.01136].
- [345] C.-M. Chang and Y.-H. Lin, Lorentzian dynamics and factorization beyond rationality, JHEP 10 (2021) 125, [arXiv:2012.01429].
- [346] S. Pal and Z. Sun, *High Energy Modular Bootstrap, Global Symmetries and Defects, JHEP* **08** (2020) 064, [arXiv:2004.12557].
- [347] M. Nguyen, Y. Tanizaki, and M. Ünsal, Noninvertible 1-form symmetry and Casimir scaling in 2D Yang-Mills theory, Phys. Rev. D 104 (2021), no. 6 065003, [arXiv:2104.01824].
- [348] E. Sharpe, *Topological operators, noninvertible symmetries and decomposition*, arXiv:2108.13423.
- [349] S. Hegde and D. P. Jatkar, *Defect Partition Function from TDLs in Commutant Pairs*, arXiv:2101.12189.
- [350] Y.-H. Lin and S.-H. Shao, \mathbb{Z}_N symmetries, anomalies, and the modular bootstrap, Phys. Rev. D 103 (2021), no. 12 125001, [arXiv:2101.08343].
- [351] K. Inamura, Topological field theories and symmetry protected topological phases with fusion category symmetries, JHEP **05** (2021) 204, [arXiv:2103.15588].
- [352] T.-C. Huang and Y.-H. Lin, *Topological Field Theory with Haagerup Symmetry*, arXiv:2102.05664.
- [353] S. Gukov, D. Pei, C. Reid, and A. Shehper, Symmetries of 2d TQFTs and Equivariant Verlinde Formulae for General Groups, arXiv:2111.08032.