

# Comparison on the criticality parameters for two supercritical branching processes in random environments

Xiequan Fan<sup>\*,1</sup>, Haijuan Hu<sup>b</sup>, Hao Wu<sup>1</sup>, Yinna Ye<sup>c</sup>

<sup>a</sup>Center for Applied Mathematics, Tianjin University, Tianjin 300072, China

<sup>b</sup>School of Mathematics and Statistics, Northeastern University at Qinhuangdao, Qinhuangdao, China

<sup>c</sup>Department of Applied Mathematics, School of Science, Xi'an Jiaotong-Liverpool University, 215123 Suzhou, China

## Abstract

Let  $\{Z_{1,n}, n \geq 0\}$  and  $\{Z_{2,n}, n \geq 0\}$  be two supercritical branching processes in different random environments, with criticality parameters  $\mu_1$  and  $\mu_2$  respectively. It is known that  $\frac{1}{n} \ln Z_{1,n} \rightarrow \mu_1$  and  $\frac{1}{m} \ln Z_{2,m} \rightarrow \mu_2$  in probability as  $m, n \rightarrow \infty$ . In this paper, we are interested in the comparison on the two criticality parameters. To this end, we prove a non-uniform Berry-Esseen's bound and Cramér's moderate deviations for  $\frac{1}{n} \ln Z_{1,n} - \frac{1}{m} \ln Z_{2,m}$  as  $m, n \rightarrow \infty$ . An application is also given for constructing confidence interval for  $\mu_1 - \mu_2$ .

*Keywords:* Branching processes; Random environment; Comparison on the criticality parameters; Berry-Esseen's bound; Cramér's moderate deviations

*2000 MSC:* primary 60J80; 60K37; secondary 60F10

## 1. Introduction

The branching process in a random environment (BPRE) was first introduced by Smith and Wilkinson [21] to model the growth of a population in an independent and identically distributed (iid) random environment. Various limit theorems for BPRE have been obtained: basic results for extinction probabilities and limit theorems for BPRE can be found in Athreya and Karlin [2, 3]. For subcritical BPRE, researches focus on the study of the survival probability and conditional limit theorems: see, for instance, Vatutin [23], Afanasyev *et al.* [1], Vatutin and Zheng [24] and Bansaye and Vatutin [5]. While, for supercritical BPRE, a number of researches have studied central limit theorem, moderate and large deviations; see, for instance, Böinghoff and Kersting [8], Bansaye and Böinghoff [4], Huang and Liu [17], Nakashima [20], Böinghoff [7], and Grama *et al.* [15], Fan *et al.* [13] and Gao [14]. See also Wang and Liu [25] and Huang *et al.* [18] for BPRE with immigrations. Despite the fact that the limit theorems for BPRE are well studied, there are no results for comparison on the criticality parameters. The objective of the paper is to fit up this gap.

Let  $(\xi_1, \xi_2)^T = ((\xi_{1,n}, \xi_{2,n})^T)_{n \geq 0}$  be a sequence of iid two dimensional random vectors, where  $(\xi_{1,n}, \xi_{2,n})^T \in \mathbb{R}^2$  stands for the random environment at time  $n$ . Notice that for give  $n$ ,  $\xi_{1,n}$  and  $\xi_{2,n}$

\*Corresponding author.

*E-mail:* fanxiequan@hotmail.com (X. Fan)

may not be independent. For any  $n \in \mathbb{N}$ , each realization of  $\xi_{1,n}$  corresponds to a probability law  $p(\xi_{1,n}) = \{p_i(\xi_{1,n}) : i \in \mathbb{N}\}$ , that is  $p_i(\xi_{1,n}) \geq 0$  and  $\sum_{i=0}^{\infty} p_i(\xi_{1,n}) = 1$ . Similarly, each realization of  $\xi_{2,n}$  corresponds to a probability law  $p(\xi_{2,n})$ . Let  $\{Z_{1,n}, n \geq 0\}$  and  $\{Z_{2,n}, n \geq 0\}$  be two branching processes in the random environment  $\xi_1$  and  $\xi_2$ , respectively. Then  $\{Z_{1,n}, n \geq 0\}$  and  $\{Z_{2,n}, n \geq 0\}$  can be described as follows: for  $n \geq 0$ ,

$$Z_{1,0} = 1, \quad Z_{1,n+1} = \sum_{i=1}^{Z_{1,n}} X_{1,n,i}, \quad Z_{2,0} = 1, \quad Z_{2,n+1} = \sum_{i=1}^{Z_{2,n}} X_{2,n,i},$$

where  $X_{1,n,i}$  and  $X_{2,n,i}$  are the number of offspring of the  $i$ -th individual in generation  $n$  with environments  $\xi_{1,n}$  and  $\xi_{2,n}$ , respectively. Denote  $\mathbb{P}_{\xi_1, \xi_2}$  the conditional probability when the environment  $(\xi_1, \xi_2)^T$  is given, and  $\tau$  the joint law of the environment  $(\xi_1, \xi_2)^T$ . Then

$$\mathbb{P}(dx_1, dx_2, dy_1, dy_2) = \mathbb{P}_{\xi_1, \xi_2}(dx_1, dx_2)\tau(dy_1, dy_2)$$

is the joint law of the two branching processes in random environment. Usually, the conditional probabilities  $\mathbb{P}_{\xi_1}$  and  $\mathbb{P}_{\xi_2}$  are called the quenched laws, while the total probability  $\mathbb{P}$  is called the annealed law. In particular, if  $\xi_1$  and  $\xi_2$  are independent, then we have  $\mathbb{P}_{\xi_1, \xi_2}(dx_1, dx_2) = \mathbb{P}_{\xi_1}(dx_1)\mathbb{P}_{\xi_2}(dx_2)$  and  $\tau(dy_1, dy_2) = \tau(dy_1)\tau(dy_2)$ , where  $\mathbb{P}_{\xi_1}(dx_1)\tau(dy_1)$  and  $\mathbb{P}_{\xi_2}(dx_2)\tau(dy_2)$  are respectively the annealed laws with respect to the branching processes  $\{Z_{1,n}, n \geq 0\}$  and  $\{Z_{2,n}, n \geq 0\}$ . In the sequel,  $\mathbb{E}_{\xi_1, \xi_2}$  and  $\mathbb{E}$  denote the expectations with respect to  $\mathbb{P}_{\xi_1, \xi_2}$  and  $\mathbb{P}$ , respectively. For any  $n \geq 1$ , set

$$m_{1,n}^{(p)} = \sum_{k=0}^{\infty} k^p p_k(\xi_{1,n}), \quad m_{2,n}^{(p)} = \sum_{k=0}^{\infty} k^p p_k(\xi_{2,n}), \quad \Pi_{1,n} = \prod_{i=0}^{n-1} m_{1,i}, \quad \Pi_{2,n} = \prod_{i=0}^{n-1} m_{2,i},$$

with the convention that  $\Pi_{1,0} = \Pi_{2,0} = 1$ . Clearly,  $(m_{1,n}^{(p)})_{n \geq 0}$  and  $(m_{2,n}^{(p)})_{n \geq 0}$  are two sequences of iid random variables. For simplicity of notations, write

$$m_{1,n} = m_{1,n}^{(1)} \quad \text{and} \quad m_{2,n} = m_{2,n}^{(1)},$$

and denote

$$\begin{aligned} X_{1,n} &= \ln m_{1,n}, & X_{2,n} &= \ln m_{2,n}, & \mu_1 &= \mathbb{E}X_{1,0}, & \mu_2 &= \mathbb{E}X_{2,0}, \\ \sigma_1^2 &= \text{Var}(X_{1,n}), & \sigma_2^2 &= \text{Var}(X_{2,n}), & \rho &= \frac{\text{Cov}(X_{1,n}, X_{2,n})}{\sigma_1 \sigma_2}, \end{aligned}$$

where  $\mu_1$  and  $\mu_2$  are known as the criticality parameters for BPREs  $\{Z_{1,n}, n \geq 0\}$  and  $\{Z_{2,n}, n \geq 0\}$ , respectively. In particular, if  $\xi_1$  and  $\xi_2$  are independent, we have  $\rho = 0$ . To avoid the environments  $\xi_1$  and  $\xi_2$  are degenerated, assume that  $0 < \sigma_1, \sigma_2 < \infty$ . Write  $\ln^+ x = \max\{\ln x, 0\}$ . Throughout the paper, we introduce the following conditions of Grama *et al.* [15]:

$$\mathbb{E} \left[ \frac{Z_{1,1}}{m_{1,0}} \ln^+ Z_{1,1} + \frac{Z_{2,1}}{m_{2,0}} \ln^+ Z_{2,1} \right] < \infty \tag{1.1}$$

and

$$p_0(\xi_{1,0}) = p_0(\xi_{2,0}) = 0, \quad a.s. \tag{1.2}$$

The assumption (1.2) implies that each individual has at least one offspring. The assumptions (1.1) and (1.2) together implies that the processes  $\{Z_{1,n}, n \geq 0\}$  and  $\{Z_{2,n}, n \geq 0\}$  are both supercritical, and satisfy that  $\mu_1, \mu_2 > 0$  and  $\mathbb{P}(Z_{1,n} \rightarrow \infty) = \mathbb{P}(Z_{2,n} \rightarrow \infty) = 1$ . See Athreya and Karlin [3] and Tanny [22].

In this paper, we are interested in the comparison between criticality parameters  $\mu_1$  and  $\mu_2$ . Consider the following common hypothesis testing:

$$H_0 : \mu_1 = \mu_2 \quad \longleftrightarrow \quad H_1 : \mu_1 \neq \mu_2.$$

It is known that  $\frac{1}{n} \ln Z_{1,n} \rightarrow \mu_1$  and  $\frac{1}{m} \ln Z_{2,m} \rightarrow \mu_2$  in probability as  $m \wedge n \rightarrow \infty$ . Thus, to answer the hypothesis testing, it is critical to know the asymptotic distribution of  $\frac{1}{n} \ln Z_{1,n} - \frac{1}{m} \ln Z_{2,m}$ , which is the main purpose of this paper. Throughout the paper, we assume either

$$\rho \in [0, 1) \quad \text{or} \quad \frac{\sigma_1}{n} \neq \frac{\sigma_2}{m}.$$

The conditions  $\rho = 1$  and  $\frac{\sigma_1}{n} = \frac{\sigma_2}{m}$  together implies that  $\frac{X_{1,1-\mu_1}}{n} = \frac{X_{2,1-\mu_2}}{m}$  a.s., which should be avoided. The main results are presented in Section 2. Let us introduce them briefly. Define

$$R_{m,n} = \frac{\frac{1}{n} \ln Z_{1,n} - \mu_1 - \frac{1}{m} \ln Z_{2,m} + \mu_2}{\sqrt{\frac{1}{n}\sigma_1^2 + \frac{1}{m}\sigma_2^2 - 2\rho\sigma_1\sigma_2\frac{m \wedge n}{mn}}}, \quad n, m \in \mathbb{N}.$$

Firstly, Theorem 2.1 presents the central limit theorem (CLT) for  $R_{m,n}$ : for all  $x \in \mathbb{R}$ , it holds

$$\lim_{m \wedge n \rightarrow \infty} \mathbb{P}(R_{m,n} \leq x) = \Phi(x), \quad (1.3)$$

where  $\Phi(x)$  is the standard normal distribution function. Secondly, under the moment condition  $\mathbb{E}[X_{1,0}^{2+\delta} + X_{2,0}^{2+\delta}] < \infty$  with  $\rho \in (0, 1]$  and the condition  $\mathbb{E}\left[\frac{Z_{1,1}^p}{m_{1,0}^p} + \frac{Z_{2,1}^p}{m_{2,0}^p}\right] < \infty$  with  $p > 1$ , Theorem 2.2 gives a non-uniform Berry-Esseen bound for  $R_{m,n}$ : for any  $\delta' \in (0, \delta)$  and all  $x \in \mathbb{R}$ ,

$$\left| \mathbb{P}(R_{m,n} \leq x) - \Phi(x) \right| \leq \frac{C}{(m \wedge n)^{\delta/2}} \frac{1}{1 + |x|^{1+\delta'}}. \quad (1.4)$$

By Lemma 4.3 and (4.10) in the paper, it seems that  $R_{m,n}$  only has a finite moment of order  $1 + \delta'$ ,  $\delta' \in (0, \delta)$ , under the stated conditions, which explains why the non-uniform Berry-Esseen bound is of order  $|x|^{-1-\delta'}$  instead of order  $|x|^{-2-\delta}$  as  $x \rightarrow \infty$ . The inequality (1.4) implies the following Berry-Esseen bound:

$$\sup_{x \in \mathbb{R}} \left| \mathbb{P}(R_{m,n} \leq x) - \Phi(x) \right| \leq \frac{C}{(m \wedge n)^{\delta/2}}. \quad (1.5)$$

In particular, when  $m \rightarrow \infty$ , we have  $\frac{1}{m} \ln Z_{2,m} \rightarrow \mu_2$  in probability, which leads to  $R_{m,n} \rightarrow \frac{\ln Z_{1,n} - n\mu_1}{\sigma_1 \sqrt{n}}$  in probability, and then inequality (1.5) recovers the Berry-Esseen bound of Grama *et al.* [15], that is,

$$\sup_{x \in \mathbb{R}} \left| \mathbb{P}\left(\frac{\ln Z_{1,n} - n\mu_1}{\sigma_1 \sqrt{n}} \leq x\right) - \Phi(x) \right| \leq \frac{C}{n^{\delta/2}}.$$

Finally, we establish Cramér's moderate deviations. Assuming Cramér's condition  $\mathbb{E}[e^{\lambda_0 X_{1,0}} + e^{\lambda_0 X_{2,0}}] < \infty$  for a constant  $\lambda_0 > 0$  and  $\mathbb{E}\left[\frac{Z_{1,1}^p}{m_{1,0}} + \frac{Z_{2,1}^p}{m_{2,0}}\right] < \infty$  for a constant  $p > 1$ , Theorem 2.3 shows that for all  $0 \leq x \leq C^{-1}\sqrt{m \wedge n}$ ,

$$\left| \ln \frac{\mathbb{P}(R_{m,n} \geq x)}{1 - \Phi(x)} \right| \leq C \frac{1 + x^3}{\sqrt{m \wedge n}}. \quad (1.6)$$

From (1.6), we obtain the following equivalence to the normal tail: it holds

$$\frac{\mathbb{P}(R_{m,n} \geq x)}{1 - \Phi(x)} = 1 + o(1) \quad (1.7)$$

uniformly for  $0 \leq x = o((m \wedge n)^{1/6})$  as  $m \wedge n \rightarrow \infty$ . When  $m \rightarrow \infty$ , it is easy to see that (1.7) holds with  $R_{m,n}$  replaced by  $\frac{\ln Z_{1,n} - n\mu_1}{\sigma_1 \sqrt{n}}$ . Such type Cramér's moderate deviations for  $\frac{\ln Z_{1,n} - n\mu_1}{\sigma_1 \sqrt{n}}$  have been established by Grama *et al.* [15].

The paper is organized as follows. Our main results are stated and discussed in Section 2. In Section 3, an application of our results to construction of confidence intervals for  $\mu_1 - \mu_2$  is demonstrated. The proofs of the main results are given in Section 4.

Throughout the paper,  $c$  and  $C$ , probably supplied with some indices, denote respectively a small positive constant and a large positive constant. Their values may vary from line to line. For two sequences of positive numbers  $(a_n)_{n \geq 1}$  and  $(b_n)_{n \geq 1}$ , we write  $a_n \asymp b_n$  if there exists a positive constant  $C$  such that for all  $n$ , it holds  $C^{-1}b_n \leq a_n \leq Cb_n$ .

## 2. Main results

We will need the following conditions.

**A1.** There exists a constant  $\delta \in (0, 1]$  such that

$$\mathbb{E}[X_{1,0}^{2+\delta} + X_{2,0}^{2+\delta}] < \infty.$$

**A2.** There exist a constant  $p > 1$  such that

$$\mathbb{E}\left[\frac{Z_{1,1}^p}{m_{1,0}^p} + \frac{Z_{2,1}^p}{m_{2,0}^p}\right] < \infty.$$

Denote  $\Phi(x)$  the standard normal distribution function. Let

$$V_{m,n,\rho} = \sqrt{\frac{1}{n}\sigma_1^2 + \frac{1}{m}\sigma_2^2 - 2\rho\sigma_1\sigma_2\frac{m \wedge n}{m n}}, \quad (2.1)$$

and define

$$R_{m,n} := \frac{\frac{1}{n} \ln Z_{1,n} - \mu_1 - \frac{1}{m} \ln Z_{2,m} + \mu_2}{V_{m,n,\rho}}, \quad n, m \in \mathbb{N}. \quad (2.2)$$

We have the following CLT for  $R_{m,n}$ .

**Theorem 2.1.** For all  $x \in \mathbb{R}$ , it holds

$$\lim_{m \wedge n \rightarrow \infty} \mathbb{P}(R_{m,n} \leq x) = \Phi(x). \quad (2.3)$$

The following theorem gives a non-uniform Berry-Esseen bound for  $R_{m,n}$ .

**Theorem 2.2.** Assume that conditions **A1** and **A2** are satisfied. Let  $\delta'$  be a constant such that  $\delta' \in (0, \delta)$ . Then for all  $x \in \mathbb{R}$ ,

$$\left| \mathbb{P}(R_{m,n} \leq x) - \Phi(x) \right| \leq \frac{C}{(m \wedge n)^{\delta/2}} \frac{1}{1 + |x|^{1+\delta'}}. \quad (2.4)$$

**Remark 2.1.** By (4.10) and Lemma 4.3, under conditions **A1** and **A2**,  $R_{m,n}$  has a finite moment of order  $1 + \delta'$ ,  $\delta' \in (0, \delta)$ , which explains why the non-uniform Berry-Esseen bound (2.4) is of order  $|x|^{-1-\delta'}$  instead of order  $|x|^{-2-\delta}$  as  $x \rightarrow \infty$ .

Denote

$$d_w(X) = \int_{-\infty}^{+\infty} |\mathbb{P}(X \leq x) - \Phi(x)| dx$$

the Wasserstein-1 distance between the random variable  $X$  and the standard normal random variable. The following corollary is a direct consequence of Theorem 2.2, which gives a convergence rate of  $R_{m,n}$  to the standard normal random variable in the Wasserstein-1 distance.

**Corollary 2.1.** Assume that conditions **A1** and **A2** are satisfied. Then

$$d_w(R_{m,n}) \leq \frac{C}{(m \wedge n)^{\delta/2}}.$$

By Theorem 2.2, we also have the following Berry-Esseen bounds for  $R_{m,n}$ .

**Corollary 2.2.** Assume that conditions **A1** and **A2** are satisfied. Then

$$\sup_{x \in \mathbb{R}} \left| \mathbb{P}(R_{m,n} \leq x) - \Phi(x) \right| \leq \frac{C}{(m \wedge n)^{\delta/2}}. \quad (2.5)$$

Notice that  $\frac{1}{m} \ln Z_{2,m} \rightarrow \mu_2$  in probability, and thus

$$R_{\infty,n} := \lim_{m \rightarrow \infty} R_{m,n} = \frac{\ln Z_{1,n} - n\mu_1}{\sigma_1 \sqrt{n}}$$

in probability. Then when  $m \rightarrow \infty$ , Corollary 2.2 recovers the Berry-Esseen bound established by Grama *et al.* [15], that is,

$$\sup_{x \in \mathbb{R}} \left| \mathbb{P}\left(\frac{\ln Z_{1,n} - n\mu_1}{\sigma_1 \sqrt{n}} \leq x\right) - \Phi(x) \right| \leq \frac{C}{n^{\delta/2}}.$$

It is known that the convergence rate of the last Berry-Esseen bound coincides the best possible one for iid random variables with finite moments of order  $2 + \delta$ .

Next, we are going to establish Cramér's moderate deviations for  $R_{m,n}$ . To this end, we need the following conditions.

**A3.** The random variables  $X_{1,0}$  and  $X_{2,0}$  have exponential moments, i.e. there exists a constant  $\lambda_0 > 0$  such that

$$\mathbb{E}[e^{\lambda_0 X_{1,0}} + e^{\lambda_0 X_{2,0}}] < \infty.$$

**A4.** There exists a constant  $p > 1$  such that

$$\mathbb{E}\left[\frac{Z_{1,1}^p}{m_{1,0}} + \frac{Z_{2,1}^p}{m_{2,0}}\right] < \infty.$$

We have the following Cramér's moderate deviations for  $R_{m,n}$ .

**Theorem 2.3.** *Assume that conditions **A3** and **A4** are satisfied. Then for all  $0 \leq x \leq c\sqrt{m \wedge n}$ ,*

$$\left| \ln \frac{\mathbb{P}(R_{m,n} \geq x)}{1 - \Phi(x)} \right| \leq C \frac{1 + x^3}{\sqrt{m \wedge n}}. \quad (2.6)$$

Thanks to the symmetry between  $m$  and  $n$ , Theorems 2.1, 2.2 and 2.3 remain valid when  $R_{m,n}$  is replaced by  $-R_{m,n}$ . Notice that  $-R_{m,n} = R_{n,m}$ .

By an argument similar to the proof of Corollary 2.2 in [12], it is easy to see that Theorem 2.3 implies the following moderate deviation principle (MDP) result for  $R_{m,n}$ .

**Corollary 2.3.** *Assume that conditions **A3** and **A4** are satisfied. Let  $a_n$  be any sequence of real numbers satisfying  $a_n \rightarrow \infty$  and  $a_n/\sqrt{m \wedge n} \rightarrow 0$  as  $m \wedge n \rightarrow \infty$ . Then, for each Borel set  $B$ , the following inequalities hold*

$$\begin{aligned} - \inf_{x \in B^\circ} \frac{x^2}{2} &\leq \liminf_{n \rightarrow \infty} \frac{1}{a_n^2} \mathbb{P}\left(\frac{R_{m,n}}{a_n} \in B\right) \\ &\leq \limsup_{n \rightarrow \infty} \frac{1}{a_n^2} \ln \mathbb{P}\left(\frac{R_{m,n}}{a_n} \in B\right) \leq - \inf_{x \in \bar{B}} \frac{x^2}{2}, \end{aligned} \quad (2.7)$$

where  $B^\circ$  and  $\bar{B}$  denote the interior and the closure of  $B$ , respectively.

From Theorem 2.3, using the inequality  $|e^y - 1| \leq e^C|y|$  valid for  $|y| \leq C$ , we obtain the following result about the uniform equivalence to the normal tail.

**Corollary 2.4.** *Assume that conditions **A3** and **A4** are satisfied. Then it holds*

$$\frac{\mathbb{P}(R_{m,n} \geq x)}{1 - \Phi(x)} = 1 + o(1) \quad (2.8)$$

uniformly for  $x \in [0, o((m \wedge n)^{1/6})]$  as  $m \wedge n \rightarrow \infty$ . The result remains valid when  $\frac{\mathbb{P}(R_{m,n} \geq x)}{1 - \Phi(x)}$  is replaced by  $\frac{\mathbb{P}(R_{m,n} \leq -x)}{\Phi(-x)}$ .

Notice that when  $\{Z_{2,n}, n \geq 0\}$  is an independent copy of  $\{Z_{1,n}, n \geq 0\}$ , we have  $\mu_1 = \mu_2, \sigma_1 = \sigma_2$  and  $\rho = 0$ . Then, for  $m = n$ , it holds  $R_{n,n} = \frac{\ln Z_{1,n} - \ln Z_{2,n}}{\sqrt{2n}\sigma_1}$ . Consequently, by Theorem 2.2, under conditions **A1** and **A2** it holds for all  $x \in \mathbb{R}$ ,

$$\left| \mathbb{P}\left(\frac{\ln Z_{1,n} - \ln Z_{2,n}}{\sqrt{2n}\sigma_1} \leq x\right) - \Phi(x) \right| \leq \frac{C}{n^{\delta/2}} \frac{1}{1 + |x|^{1+\delta'}}.$$

By Corollary 2.4, under conditions **A3** and **A4** we have

$$\frac{\mathbb{P}\left(\frac{\ln Z_{1,n} - \ln Z_{2,n}}{\sqrt{2n\sigma_1}} \geq x\right)}{1 - \Phi(x)} = 1 + o(1)$$

uniformly for  $x \in [0, o(n^{1/6})]$  as  $n \rightarrow \infty$ .

### 3. Applications to construction of confidence intervals

When parameters  $\sigma_1, \sigma_2$ , and  $\rho$  are known, we can apply Theorems 2.2 and 2.3 to construct confidence intervals for  $\mu_1 - \mu_2$ .

**Proposition 3.1.** *Let  $\kappa_{m,n} \in (0, 1)$ . Consider the following two groups of conditions:*

**B1.** *The conditions of Theorem 2.2 hold and*

$$|\ln \kappa_{m,n}| = o(\ln(m \wedge n)), \quad \text{as } m \wedge n \rightarrow \infty. \quad (3.1)$$

**B2.** *The conditions of Theorem 2.3 hold and*

$$|\ln \kappa_{m,n}| = o((m \wedge n)^{1/3}), \quad \text{as } m \wedge n \rightarrow \infty. \quad (3.2)$$

*If either **B1** or **B2** holds, then for  $n$  large enough,  $[A_{m,n}, B_{m,n}]$  is the confidence interval of  $\mu_1 - \mu_2$  with confidence level  $1 - \kappa_{m,n}$ , where*

$$A_{m,n} = \frac{1}{n} \ln Z_{1,n} - \frac{1}{m} \ln Z_{2,m} - V_{m,n,\rho} \Phi^{-1}\left(1 - \frac{\kappa_{m,n}}{2}\right)$$

and

$$B_{m,n} = \frac{1}{n} \ln Z_{1,n} - \frac{1}{m} \ln Z_{2,m} + V_{m,n,\rho} \Phi^{-1}\left(1 - \frac{\kappa_{m,n}}{2}\right).$$

*Proof.* Assume **B1**. By Theorem 2.2, as  $m \wedge n \rightarrow \infty$ , we have

$$\frac{\mathbb{P}(R_{m,n} > x)}{1 - \Phi(x)} = 1 + o(1) \quad \text{and} \quad \frac{\mathbb{P}(R_{m,n} < -x)}{\Phi(-x)} = 1 + o(1) \quad (3.3)$$

uniformly for  $0 \leq x = o(\sqrt{\ln(m \wedge n)})$ . For  $p \searrow 0$ , the quantile function of the standard normal distribution has the following asymptotic expansion

$$\Phi^{-1}(p) = -\sqrt{\ln \frac{1}{p^2} - \ln \ln \frac{1}{p^2} - \ln(2\pi)} + o(1).$$

In particular, when  $\kappa_{m,n}$  satisfies the condition (3.1), the upper  $(1 - \frac{\kappa_{m,n}}{2})$ -th quantile of standard normal distribution satisfies  $\Phi^{-1}(1 - \frac{\kappa_{m,n}}{2}) = -\Phi^{-1}(\frac{\kappa_{m,n}}{2}) = O(\sqrt{|\ln \kappa_{m,n}|})$ , is of order  $o(\sqrt{\ln(m \wedge n)})$ . Then, applying the last equality to (3.3), we have as  $m \wedge n \rightarrow \infty$ ,

$$\mathbb{P}\left(R_{m,n} > \Phi^{-1}\left(1 - \frac{\kappa_{m,n}}{2}\right)\right) \sim \frac{\kappa_{m,n}}{2} \quad (3.4)$$

and

$$\mathbb{P}\left(R_{m,n} < -\Phi^{-1}\left(1 - \frac{\kappa_{m,n}}{2}\right)\right) \sim \frac{\kappa_{m,n}}{2}. \quad (3.5)$$

Therefore, as  $m \wedge n \rightarrow \infty$ ,

$$\mathbb{P}\left(-\Phi^{-1}\left(1 - \frac{\kappa_{m,n}}{2}\right) \leq R_{m,n} \leq \Phi^{-1}\left(1 - \frac{\kappa_{m,n}}{2}\right)\right) \sim 1 - \kappa_{m,n}, \quad (3.6)$$

which implies  $\mu_1 - \mu_2 \in [A_{m,n}, B_{m,n}]$  with probability  $1 - \kappa_{m,n}$  for  $m \wedge n$  large enough.

Now, assume **B2**. By Theorem 2.3, as  $m \wedge n \rightarrow \infty$ , we have

$$\frac{\mathbb{P}(R_{m,n} > x)}{1 - \Phi(x)} = 1 + o(1) \quad \text{and} \quad \frac{\mathbb{P}(R_{m,n} < -x)}{\Phi(-x)} = 1 + o(1) \quad (3.7)$$

uniformly for  $0 \leq x = o((m \wedge n)^{1/6})$ . When  $\kappa_{m,n}$  satisfies the condition (3.2), the upper  $(1 - \frac{\kappa_{m,n}}{2})$ -th quantile of the standard normal distribution satisfies  $\Phi^{-1}(1 - \frac{\kappa_{m,n}}{2}) = -\Phi^{-1}(\frac{\kappa_{m,n}}{2}) = O(\sqrt{|\ln \kappa_{m,n}|})$ , which is of order  $o((m \wedge n)^{1/6})$ . Then by (3.7), we get, as  $m \wedge n \rightarrow \infty$ ,

$$\mathbb{P}\left(-\Phi^{-1}\left(1 - \frac{\kappa_{m,n}}{2}\right) \leq R_{m,n} \leq \Phi^{-1}\left(1 - \frac{\kappa_{m,n}}{2}\right)\right) \sim 1 - \kappa_{m,n}.$$

From the equality above, the result still holds.  $\square$

When  $\{Z_{2,n}, n \geq 0\}$  is an independent copy of  $\{Z_{1,n}, n \geq 0\}$ , we can apply Theorems 2.2 and 2.3 to construct confidence intervals for  $\sigma_1$ .

**Proposition 3.2.** *Let  $\kappa_{n,n} \in (0, 1)$ . If either **B1** or **B2** holds, then for  $n$  large enough,  $[A_n, B_n]$  is the confidence interval of  $\sigma_1^2$  with confidence level  $1 - \kappa_{n,n}$ , where*

$$A_n = \frac{(\ln Z_{1,n} - \ln Z_{2,n})^2}{2n\chi_{1-\frac{1}{2}\kappa_{n,n}}^2(1)} \quad \text{and} \quad B_n = \frac{(\ln Z_{1,n} - \ln Z_{2,n})^2}{2n\chi_{\frac{1}{2}\kappa_{n,n}}^2(1)}$$

with  $\chi_q^2(1)$  the  $q$ -quantiles for chi-squared distribution with 1 degrees of freedom.

*Proof.* Assume **B1**. By Theorem 2.2, as  $n \rightarrow \infty$ , we have

$$\frac{\mathbb{P}\left(\frac{(\ln Z_{1,n} - \ln Z_{2,n})^2}{2n\sigma_1^2} > x\right)}{\mathbb{P}(\chi^2(1) \geq x)} = 1 + o(1) \quad (3.8)$$

uniformly for  $0 \leq x = o(\sqrt{\ln n})$ . Then, applying the last equality to (3.8), we have, as  $n \rightarrow \infty$ ,

$$\mathbb{P}\left(\chi_{\frac{1}{2}\kappa_{n,n}}^2(1) \leq \frac{(\ln Z_{1,n} - \ln Z_{2,n})^2}{2n\sigma_1^2} \leq \chi_{1-\frac{1}{2}\kappa_{n,n}}^2(1)\right) \sim 1 - \kappa_{n,n}, \quad (3.9)$$

which implies  $\sigma_1^2 \in [A_n, B_n]$  with probability  $1 - \kappa_{n,n}$  for  $n$  large enough.

Assume **B2**. Similar argument holds. This completes the proof of Proposition 3.2.  $\square$



#### 4. Proofs of Theorems

For  $l = 1, 2$ , denote the normalized population size

$$W_{l,n} = \frac{Z_{l,n}}{\prod_{l,n}}, \quad n \geq 0.$$

Then  $(W_{1,n})_{n \geq 0}$  and  $(W_{2,n})_{n \geq 0}$  are both nonnegative martingales under the annealed law  $\mathbb{P}$ , with respect to the natural filtration  $\mathcal{F}_0 = \sigma\{\xi_1, \xi_2\}$ ,  $\mathcal{F}_n = \sigma\{\xi_1, \xi_2, X_{1,k,i}, X_{2,k,i}, 0 \leq k \leq n-1, i \geq 1\}$ ,  $n \geq 1$ . Then, by Doob's convergence theorem, the limit

$$W_{l,\infty} = \lim_{n \rightarrow \infty} W_{l,n}$$

exists  $\mathbb{P}$ -a.s. and, by Fatou's lemma, satisfies  $\mathbb{E}W_{l,\infty} \leq 1$ . The conditions (1.1) and (1.2) together implies that  $\mathbb{P}(W_{l,n} > 0) = \mathbb{P}(Z_{l,n} \rightarrow \infty) = \lim_{n \rightarrow \infty} \mathbb{P}(Z_{l,n} > 0) = 1$ , and that the martingale  $W_{l,n}$  converges to  $W_{l,\infty}$  in  $\mathbb{L}^1(\mathbb{P})$  (see Athreya and Karlin [3] and also Tanny [22]).

For simplicity of notations, without loss of generality, we assume that  $m \leq n$ . In the sequel, denote

$$\eta_{m,n,i} = \frac{X_{1,i-1} - \mu_1}{n V_{m,n,\rho}}, \quad i = 1, \dots, n, \quad \text{and} \quad \eta_{m,n,n+j} = -\frac{X_{2,j-1} - \mu_2}{m V_{m,n,\rho}}, \quad j = 1, \dots, m.$$

Then  $R_{m,n}$  can be rewritten in the following form

$$R_{m,n} = \sum_{i=1}^{n+m} \eta_{m,n,i} + \frac{\ln W_{1,n}}{n V_{m,n,\rho}} - \frac{\ln W_{2,m}}{m V_{m,n,\rho}}. \quad (4.10)$$

Set

$$Y_i = \eta_{m,n,i} + \eta_{m,n,n+i}, \quad i = 1, \dots, m, \quad \text{and} \quad Y_i = \eta_{m,n,i}, \quad i = m+1, \dots, n.$$

Then  $(Y_i)_{1 \leq i \leq n}$  is a finite sequence of centered and independent random variables, and satisfies

$$\sum_{i=1}^n Y_i = \sum_{i=1}^{n+m} \eta_{m,n,i} \quad \text{and} \quad \sum_{i=1}^n \mathbb{E}Y_i^2 = 1.$$

Moreover, we have

$$\text{Var}(Y_i) \asymp \frac{1}{m}, \quad i = 1, \dots, m, \quad \text{and} \quad \text{Var}(Y_i) \asymp \frac{m}{n^2}, \quad i = m+1, \dots, n,$$

as  $m \rightarrow \infty$ .

##### 4.1. Proof of Theorem 2.1

Without loss of generality, we assume that  $m \leq n$ . Recall that  $0 < \sigma_1, \sigma_2 < \infty$ , and that  $(Y_i)_{1 \leq i \leq n}$  satisfies

$$\sum_{i=1}^n Y_i = \sum_{i=1}^{n+m} \eta_{m,n,i} \quad \text{and} \quad \sum_{i=1}^n \mathbb{E}Y_i^2 = 1.$$

Notice that

$$V_{m,n,\rho} \asymp \frac{1}{\sqrt{m}} \quad \text{and} \quad \max_{1 \leq i \leq n} \text{Var}(Y_i) \rightarrow 0, \quad m \rightarrow \infty.$$

Thus by CLT for independent random variables, we have  $\sum_{i=1}^{n+m} \eta_{m,n,i}$  converges in distribution to the standard normal distribution as  $m \rightarrow \infty$ . Recall that for  $l = 1, 2$ ,  $W_{l,n}$  converges to  $W_{l,\infty}$  in  $\mathbb{L}^1(\mathbb{P})$  as  $n \rightarrow \infty$ , thus  $\frac{\ln W_{1,n}}{n V_{m,n,\rho}}$  and  $\frac{\ln W_{2,m}}{m V_{m,n,\rho}}$  both convergence in probability to 0 as  $m \rightarrow \infty$ . Hence, by (4.10), we have  $R_{m,n}$  converges in distribution to the normal distribution. This completes the proof of Theorem 2.1.

#### 4.2. Preliminary Lemmas for Theorem 2.2

In the proof of Theorem 2.2, we need the following non-uniform Berry-Esseen bound of Bikelis [6]. See also Chen and Shao [9] for more general results.

**Lemma 4.1.** *Let  $(Y_i)_{1 \leq i \leq n}$  be independent random variables satisfying  $\mathbb{E}Y_i = 0$  and  $\mathbb{E}|Y_i|^{2+\delta} < \infty$  for some positive constant  $\delta \in (0, 1]$  and all  $1 \leq i \leq n$ . Assume that  $\sum_{i=1}^n \mathbb{E}Y_i^2 = 1$ . Then for all  $x \in \mathbb{R}$ ,*

$$\left| \mathbb{P}\left(\sum_{i=1}^n Y_i \leq x\right) - \Phi(x) \right| \leq \frac{C}{1 + |x|^{2+\delta}} \sum_{i=1}^n \mathbb{E}|Y_i|^{2+\delta}.$$

Consider the (conditional) Laplace transforms of  $W_{1,\infty}$  and  $W_{2,\infty}$  as follows: for all  $t \geq 0$ ,

$$\phi_{i,\xi}(t) = \mathbb{E}_\xi e^{-tW_{i,\infty}} \quad \text{and} \quad \phi_i(t) = \mathbb{E}\phi_{i,\xi}(t) = \mathbb{E}e^{-tW_{i,\infty}}, \quad i = 1, 2.$$

Clearly, as  $W_{i,\infty} \geq 0$   $\mathbb{P}$ -a.s. we have  $\phi_i(t) \in (0, 1], i = 1, 2$ . Moreover, we have the following bounds for  $\phi_i(t), i = 1, 2$ , as  $t \rightarrow \infty$ .

**Lemma 4.2.** *Assume that conditions A1 and A2 are satisfied. Then for  $i = 1, 2$ , it holds*

$$\phi_i(t) \leq \frac{C}{1 + (\ln^+ t)^{1+\delta}}, \quad t \rightarrow \infty.$$

*Proof.* Set  $i = 1, 2$ . Define  $T^n$  the shift operator by  $T^n(\xi_{i,0}, \xi_{i,1}, \dots) = (\xi_{i,n}, \xi_{i,n+1}, \dots)$ , for  $n \geq 1$ . Then we get for a fixed  $k \geq 0$ ,

$$\Pi_{i,n}(T^k \xi_i) = m_{i,k} m_{i,k+1} \cdots m_{i,k+n-1}.$$

In particular, we have  $\Pi_{i,n} = \Pi_{i,n}(T^0 \xi_i)$ . Next, we use the method of Grama *et al.* [16] to complete the proof. From (3.15) in [16], it is proven that for any  $t > 0$  and  $n \geq 1$ ,

$$\begin{aligned} \phi_i(t) &\leq \mathbb{E}\phi_i\left(\frac{t}{\Pi_{i,n}}\right) \prod_{j=0}^{n-1} \left(p_1(\xi_{i,j}) + (1 - p_1(\xi_{i,j}))\beta_K\right) \\ &\quad + \frac{1}{K} \mathbb{E}\left[\phi_{i,T^n \xi_i}\left(\frac{t}{\Pi_{i,n}}\right) \left(\mathbb{E}_{T^n \xi_i} W_{i,\infty}^p\right)\right] + \mathbb{P}\left(\frac{t}{\Pi_{i,n}} < t_K\right), \end{aligned} \quad (4.11)$$

where  $t_K := (CK)^{-1/(p-1)}$ ,  $C$  and  $K$  are positive constants, such that

$$\beta_K := 1 - (1 - 1/p)t_K \in (0, 1).$$

From (3.18) in [16], it is proven the following inequality

$$\mathbb{E} \left[ \phi_{i, T^n \xi_i} \left( \frac{t}{\Pi_{i,n}} \right) \mathbb{E}_{T^n \xi_i} W_{i,\infty}^p \right] \leq \mathbb{E} \phi_i \left( \frac{t}{\Pi_{i,n}} \right) + \sum_{k=0}^{\infty} \mathbb{E} \frac{\phi_i \left( \frac{t}{\Pi_{i,k+1}(T^n \xi_i) \Pi_{i,n}(\xi_i)} \right) m_k^{(p)}(T^n \xi_i)}{\Pi_{i,k}^{p-1}(T^n \xi_i) m_k^p(T^n \xi_i)}. \quad (4.12)$$

Let  $p \in (1, 2]$ . Given  $n, K$ , let  $Y_{i,n,K}$  be a positive random variable whose distribution is defined by

$$\begin{aligned} \mathbb{E} g(Y_{i,n,K}) &= \frac{1}{q_{i,n,K}} \left[ \mathbb{E} g \left( \frac{1}{\Pi_{i,n}} \right) \prod_{j=0}^{n-1} (p_1(\xi_{i,j}) + (1 - p_1(\xi_{i,j}))\beta_K) \right. \\ &\quad \left. + \frac{1}{K} \mathbb{E} g \left( \frac{1}{\Pi_{i,n}} \right) + \frac{1}{K} \sum_{k=0}^{\infty} \mathbb{E} \frac{g \left( \frac{1}{\Pi_{i,k+1}(T^n \xi_i) \Pi_{i,n}(\xi_i)} \right) m_{i,k}^{(p)}(T^n \xi_i)}{\Pi_{i,k}^{p-1}(T^n \xi_i) m_{i,k}^p(T^n \xi_i)} \right], \end{aligned} \quad (4.13)$$

for all bounded and measurable function  $g$ , where  $q_{i,n,K}$  is the normalizing constant (to make  $\mathbb{E} g(Y_{i,n,K}) = 1$ , when  $g = 1$ ) defined by

$$\begin{aligned} q_{i,n,K} &= \mathbb{E} \left[ \prod_{j=0}^{n-1} (p_1(\xi_{i,j}) + (1 - p_1(\xi_{i,j}))\beta_K) \right] + \frac{1}{K} \left[ 1 + \sum_{k=0}^{\infty} \mathbb{E} \frac{1}{\Pi_{i,k}^{p-1}(T^n \xi_i)} \frac{m_{i,k}^{(p)}(T^n \xi_i)}{m_{i,k}^p(T^n \xi_i)} \right] \\ &= \left[ \mathbb{E} (p_1(\xi_{i,0}) + (1 - p_1(\xi_{i,0}))\beta_K) \right]^n + \frac{1}{K} \left[ 1 + \frac{1}{1 - \mathbb{E} m_{i,0}^{-(p-1)}} \mathbb{E} \left( \frac{m_{i,0}^{(p)}}{m_{i,0}^p} \right) \right]. \end{aligned} \quad (4.14)$$

The last line follows by (3.21) in [16]. Combining (4.11)-(4.13) together, we have

$$\phi_i(t) \leq q_{i,n,K} \mathbb{E} \phi_i(Y_{i,n,K} t) + \mathbb{P} \left( \frac{t}{\Pi_{i,n}} < t_K \right). \quad (4.15)$$

Choose  $A_i$  such that  $\ln A_i > \mu_i$ . Clearly, we have  $A_i > 1$ . When  $A_i^{n+1} \leq t < A_i^{n+2}$ , let  $K = K_{i,n} = ((n+1) \ln A_i)^{1+\delta}$ , then we have  $\lim_{n \rightarrow \infty} \sqrt[n]{t_{K_{i,n}}} = 1$ . By Nagaev's inequality (see Corollary 2.5 in [11]), for  $n$  large enough,

$$\mathbb{P} \left( \frac{t}{\Pi_{i,n}} < t_{K_{i,n}} \right) = \mathbb{P} \left( \Pi_{i,n} > \frac{t}{t_{K_{i,n}}} \right) \leq \mathbb{P} \left( S_{i,n} - n\mu_i > n \left( \ln \frac{A_i}{\sqrt[n]{t_{K_{i,n}}}} - \mu_i \right) \right) \leq \frac{C}{n^{1+\delta}}, \quad (4.16)$$

where  $S_{i,n} = \sum_{j=1}^n X_{i,j-1}$ . When  $t \geq A_i^2$ , let  $n$  be a positive integer such that  $A_i^{n+1} \leq t < A_i^{n+2}$ , so we get

$$n > \frac{\ln t}{\ln A_i} - 2.$$

Thus, by (4.16), for any  $A_i^{n+1} \leq t < A_i^{n+2}$ ,

$$\mathbb{P} \left( \frac{t}{\Pi_{i,n}} < t_{K_{i,n}} \right) \leq \frac{C}{n^{1+\delta}} < C \left( \frac{\ln t}{\ln A_i} - 2 \right)^{-1-\delta} \leq \frac{C}{(\ln t)^{1+\delta}}. \quad (4.17)$$

Notice that  $0 \leq \phi_i(t) \leq 1$  ( $t > 0$ ). By (4.15) and (4.17), we have

$$\phi_i(t) \leq \sum_{n=1}^{\infty} q_{i,n,K_{i,n}} \mathbf{1}_{\{A_i^{n+1} \leq t < A_i^{n+2}\}} + \frac{C}{1 + (\ln^+ t)^{1+\delta}}. \quad (4.18)$$

From the definition (4.14) of  $q_{i,n,K}$ , when  $A_i^{n+1} \leq t < A_i^{n+2}$ , we have

$$q_{i,n,K_{i,n}} \leq \frac{C}{1 + (\ln^+ t)^{1+\delta}}. \quad (4.19)$$

Finally, by (4.18) and (4.19), we obtain for any  $t > 0$ ,

$$\phi_i(t) \leq \frac{C}{1 + (\ln^+ t)^{1+\delta}}.$$

This completes the proof of Lemma 4.2.  $\square$

We have the following result for the  $\mathbb{L}^p(\mathbb{P})$  moments of  $\ln W_{i,\infty}$  and  $\ln W_{i,n}$ .

**Lemma 4.3.** *Assume conditions A1 and A2 are satisfied. Then for  $i = 1, 2$  and  $q \in (1, 1 + \delta)$ , the following two inequalities hold*

$$\mathbb{E} |\ln W_{i,\infty}|^q < \infty \quad (4.20)$$

and

$$\sup_{n \in \mathbb{N}} \mathbb{E} |\ln W_{i,n}|^q < \infty. \quad (4.21)$$

*Proof.* Set  $i = 1, 2$ . Consider the following truncation on  $\mathbb{E} |\ln W_{i,\infty}|^q$ , that is

$$\mathbb{E} |\ln W_{i,\infty}|^q = \mathbb{E} |\ln W_{i,\infty}|^q \mathbf{1}_{\{W_{i,\infty} > 1\}} + \mathbb{E} |\ln W_{i,\infty}|^q \mathbf{1}_{\{W_{i,\infty} \leq 1\}}. \quad (4.22)$$

For the first term in the right-hand side of the equality above, we have

$$\mathbb{E} |\ln W_{i,\infty}|^q \mathbf{1}_{\{W_{i,\infty} > 1\}} \leq C \mathbb{E} W_{i,\infty} < \infty. \quad (4.23)$$

For the second term, we have

$$\begin{aligned} \mathbb{E} |\ln W_{i,\infty}|^q \mathbf{1}_{\{W_{i,\infty} \leq 1\}} &= q \int_1^{\infty} \frac{1}{t} (\ln t)^{q-1} \mathbb{P}(W_{i,\infty} \leq t^{-1}) dt \\ &\leq q e \int_1^{\infty} \frac{\phi_i(t)}{t} (\ln t)^{q-1} dt \\ &= q e \left( \int_1^e \frac{\phi_i(t)}{t} (\ln t)^{q-1} dt + \int_e^{\infty} \frac{\phi_i(t)}{t} (\ln t)^{q-1} dt \right). \end{aligned} \quad (4.24)$$

The last inequality above can be obtained by Markov's inequality, i.e.,

$$\mathbb{P}(W_{i,\infty} \leq t^{-1}) \leq e \mathbb{E} e^{-tW_{i,\infty}} = e \phi_i(t).$$

Clearly, it holds

$$\int_1^e \frac{\phi_i(t)}{t} (\ln t)^{q-1} dt < \infty. \quad (4.25)$$

From Lemma 4.2 and  $q < 1 + \delta$ , we have

$$\int_e^\infty \frac{\phi_i(t)}{t} (\ln t)^{q-1} dt \leq C \int_e^\infty \frac{1}{t(\ln t)^{2+\delta-q}} dt < \infty. \quad (4.26)$$

Substituting (4.25) and (4.26) into (4.24), we get

$$\mathbb{E} |\ln W_{i,\infty}|^q \mathbf{1}_{\{W_{i,\infty} \leq 1\}} < \infty. \quad (4.27)$$

Therefore, by (4.22), (4.23) and (4.27), we get (4.20).

Next, we give a proof for (4.21). Since  $x \mapsto |\ln^q(x) \mathbf{1}_{\{x \leq 1\}}|$ ,  $q > 1$ , is a non-negative and convex function, by Lemma 2.1 in [17], we have

$$\sup_{n \in \mathbb{N}} \mathbb{E} |\ln W_{i,n}|^q \mathbf{1}_{\{W_{i,n} \leq 1\}} = \mathbb{E} |\ln W_{i,\infty}|^q \mathbf{1}_{\{W_{i,\infty} \leq 1\}}.$$

With the similar truncation as  $\mathbb{E} |\ln W_{i,\infty}|^q$ , by (4.27), we get

$$\begin{aligned} \sup_{n \in \mathbb{N}} \mathbb{E} |\ln W_{i,n}|^q &= \sup_{n \in \mathbb{N}} \left( \mathbb{E} |\ln W_{i,n}|^q \mathbf{1}_{\{W_{i,n} > 1\}} + \mathbb{E} |\ln W_{i,n}|^q \mathbf{1}_{\{W_{i,n} \leq 1\}} \right) \\ &\leq \sup_{n \in \mathbb{N}} \mathbb{E} |\ln W_{i,n}|^q \mathbf{1}_{\{W_{i,n} > 1\}} + \sup_{n \in \mathbb{N}} \mathbb{E} |\ln W_{i,n}|^q \mathbf{1}_{\{W_{i,n} \leq 1\}} \\ &\leq C \mathbb{E} W_{i,\infty} + \mathbb{E} |\ln W_{i,\infty}|^q \mathbf{1}_{\{W_{i,\infty} \leq 1\}} < \infty. \end{aligned}$$

This completes the proof of Lemma 4.3.  $\square$

The following lemma is a simple consequence of Lemma 2.4 of Grama *et al.* [15].

**Lemma 4.4.** *Assume that conditions A1 and A2 are satisfied. Then there exists constant  $\gamma \in (0, 1)$ , such that*

$$\mathbb{E} |\ln W_{1,n} - \ln W_{1,\infty}| + \mathbb{E} |\ln W_{2,n} - \ln W_{2,\infty}| \leq C \gamma^n.$$

In the proof of Theorem 2.2, the following lemma plays an important role.

**Lemma 4.5.** *Assume that conditions A1 and A2 are satisfied. Let  $\delta'$  be a constant such that  $\delta' \in (0, \delta)$ . Then for all  $x \in \mathbb{R}$ ,*

$$\mathbb{P} \left( R_{m,n} \leq x, \sum_{i=1}^{n+m} \eta_{m,n,i} \geq x \right) \leq \frac{C}{(m \wedge n)^{\delta/2}} \frac{1}{1 + |x|^{1+\delta'}} \quad (4.28)$$

and

$$\mathbb{P} \left( R_{m,n} \geq x, \sum_{i=1}^{n+m} \eta_{m,n,i} \leq x \right) \leq \frac{C}{(m \wedge n)^{\delta/2}} \frac{1}{1 + |x|^{1+\delta'}}. \quad (4.29)$$

*Proof.* We only give a proof for inequality (4.28), as inequality (4.29) can be proved in the same way. Without loss of generality, we may assume that  $m \leq n$ .

First, we show that inequality (4.28) holds for  $x \leq -Cm^{1/2}$  with some positive constant  $C$ . Recall

$$R_{m,n} = \frac{\ln Z_{1,n} - n\mu_1}{nV_{m,n,\rho}} - \frac{\ln Z_{2,m} - m\mu_2}{mV_{m,n,\rho}}, \quad m, n \in \mathbb{N}. \quad (4.30)$$

Then we have for all  $x \in \mathbb{R}$ ,

$$\mathbb{P}\left(R_{m,n} \leq x, \sum_{i=1}^{n+m} \eta_{m,n,i} \geq x\right) \leq \mathbb{P}\left(R_{m,n} \leq x\right) \leq P_1 + P_2, \quad (4.31)$$

where

$$P_1 = \mathbb{P}\left(\frac{\ln Z_{1,n} - n\mu_1}{nV_{m,n,\rho}} \leq \frac{x}{2}\right) \quad \text{and} \quad P_2 = \mathbb{P}\left(-\frac{\ln Z_{2,m} - m\mu_2}{mV_{m,n,\rho}} \leq \frac{x}{2}\right).$$

Since  $Z_{1,n} \geq 1$   $\mathbb{P}$ -a.s. and  $V_{m,n,\rho} \asymp m^{-1/2}$  as  $m \rightarrow \infty$ , there exists a positive constant  $C$  such that

$$\frac{\ln Z_{1,n} - n\mu_1}{nV_{m,n,\rho}} > -\frac{\mu_1}{V_{m,n,\rho}} > -\frac{1}{2}Cm^{1/2} \quad \mathbb{P}\text{-a.s.},$$

and thus  $P_1 = 0$  for all  $x \leq -Cm^{1/2}$ . For  $P_2$ , by Lemma 4.1, Markov's inequality and the fact  $\mathbb{E}W_{2,m} = 1$ , we have for all  $x \leq -Cm^{1/2}$ ,

$$\begin{aligned} P_2 &\leq \mathbb{P}\left(\sum_{j=1}^m \eta_{m,n,n+j} \leq -\frac{|x|}{4}\right) + \mathbb{P}\left(\frac{\ln W_{2,m}}{mV_{m,n,\rho}} \geq \frac{|x|}{4}\right) \\ &\leq \mathbb{P}\left(\sum_{j=1}^m \eta_{m,n,n+j} \leq -\frac{|x|}{4}\right) + \exp\left\{-\frac{|x|}{4}mV_{m,n,\rho}\right\} \mathbb{E}W_{2,m} \\ &\leq \frac{C}{m^{\delta/2}} \frac{1}{1 + |x|^{2+\delta}}. \end{aligned} \quad (4.32)$$

Hence, inequality (4.28) holds for all  $x \leq -Cm^{1/2}$ .

Next, we show that inequality (4.28) holds for all  $x \geq Cm^{1/2}$ . For all  $x \geq 0$ , we have

$$\mathbb{P}\left(R_{m,n} \leq x, \sum_{i=1}^{n+m} \eta_{m,n,i} \geq x\right) \leq \mathbb{P}\left(\sum_{i=1}^{n+m} \eta_{m,n,i} \geq x\right).$$

Applying Lemma 4.1 to the right-hand-side of the last inequality, we have for all  $x \geq 0$ ,

$$\begin{aligned} \mathbb{P}\left(R_{m,n} \leq x, \sum_{i=1}^{n+m} \eta_{m,n,i} \geq x\right) &\leq 1 - \Phi(x) \\ &\quad + \frac{C_1}{1 + |x|^{2+\delta}} \left( \sum_{i=1}^m \mathbb{E}|\eta_{m,n,i} + \eta_{m,n,n+i}|^{2+\delta} + \sum_{i=m+1}^n \mathbb{E}|\eta_{m,n,i}|^{2+\delta} \right). \end{aligned}$$

Using the inequality

$$(a + b)^{2+\delta} \leq 2^{1+\delta}(|a|^{2+\delta} + |b|^{2+\delta}), \quad a, b \in \mathbb{R},$$

we deduce that for all  $x \geq 0$ ,

$$\begin{aligned} \mathbb{P}\left(R_{m,n} \leq x, \sum_{i=1}^{n+m} \eta_{m,n,i} \geq x\right) \\ \leq 1 - \Phi(x) + \frac{C_1}{1 + |x|^{2+\delta}} \left( \sum_{i=1}^n \mathbb{E}|\eta_{m,n,i}|^{2+\delta} + \sum_{i=1}^m \mathbb{E}|\eta_{m,n,n+i}|^{2+\delta} \right). \end{aligned}$$

Notice that  $V_{m,n,\rho} \asymp m^{-1/2}$  as  $m \rightarrow \infty$ . Then, by the inequalities

$$\frac{1}{\sqrt{2\pi}(1+x)} e^{-x^2/2} \leq 1 - \Phi(x) \leq \frac{1}{\sqrt{\pi}(1+x)} e^{-x^2/2}, \quad x \geq 0, \quad (4.33)$$

we have for all  $x \geq Cm^{1/2}$ ,

$$\begin{aligned} \mathbb{P}\left(R_{m,n} \leq x, \sum_{i=1}^{n+m} \eta_{m,n,i} \geq x\right) &\leq \frac{C_2}{m^{\delta/2}} \frac{1}{1+|x|^{2+\delta}} + \frac{C_3}{1+|x|^{2+\delta}} \left( \frac{n}{n^{2+\delta} m^{-1-\delta/2}} + \frac{m}{m^{2+\delta} m^{-1-\delta/2}} \right) \\ &\leq \frac{C_4}{m^{\delta/2}} \frac{1}{1+|x|^{2+\delta}}. \end{aligned}$$

Thus, inequality (4.28) holds for all  $x \geq Cm^{1/2}$ .

To end the proof of lemma, in the sequel we show that (4.28) holds for all  $|x| < Cm^{1/2}$ . Consider the following notations for all  $0 \leq k \leq m-1$

$$\begin{aligned} Y_{m,n,k} &= \sum_{i=k+1}^n \eta_{m,n,i} + \sum_{j=k+1}^m \eta_{m,n,n+j}, \quad \tilde{Y}_{m,n,k} = Y_{m,n,0} - Y_{m,n,k}, \\ H_{m,n,k} &= \frac{\ln W_{1,k}}{n V_{m,n,\rho}} - \frac{\ln W_{2,k}}{m V_{m,n,\rho}} \quad \text{and} \quad D_{m,n,k} = \frac{\ln W_{1,n}}{n V_{m,n,\rho}} - \frac{\ln W_{2,m}}{m V_{m,n,\rho}} - H_{m,n,k}. \end{aligned}$$

Set  $\alpha_m = m^{-\delta/2}$  and  $k = [m^{1-\delta/2}]$ , where  $[t]$  stands for the largest integer less than  $t$ . From (4.10), we deduce that for all  $x \in \mathbb{R}$ ,

$$\begin{aligned} \mathbb{P}\left(R_{m,n} \leq x, \sum_{i=1}^{n+m} \eta_{m,n,i} \geq x\right) &= \mathbb{P}\left(Y_{m,n,0} + H_{m,n,k} + D_{m,n,k} \leq x, Y_{m,n,0} \geq x\right) \\ &\leq \mathbb{P}\left(Y_{m,n,0} + H_{m,n,k} \leq x + \alpha_m, Y_{m,n,0} \geq x\right) \\ &\quad + \mathbb{P}\left(|D_{m,n,k}| \geq \alpha_m\right). \end{aligned} \quad (4.34)$$

For the tail probability  $\mathbb{P}(|D_{m,n,k}| \geq \alpha_m)$ , by Markov's inequality and Lemma 4.4, there exists a

constant  $\gamma \in (0, 1)$  such that for all  $-m < x < m$ ,

$$\begin{aligned}
\mathbb{P}(|D_{m,n,k}| > \alpha_m) &\leq \frac{\mathbb{E}|D_{m,n,k}|}{\alpha_m} \\
&= \frac{m^{\delta/2}}{V_{m,n,\rho}} \mathbb{E} \left| \left[ \left( \frac{\ln W_{1,n}}{n} - \frac{\ln W_{1,\infty}}{n} \right) - \left( \frac{\ln W_{2,n}}{m} - \frac{\ln W_{2,\infty}}{m} \right) \right] \right. \\
&\quad \left. - \left[ \left( \frac{\ln W_{1,k}}{n} - \frac{\ln W_{1,\infty}}{n} \right) - \left( \frac{\ln W_{2,k}}{m} - \frac{\ln W_{2,\infty}}{m} \right) \right] \right| \\
&\leq \frac{m^{\delta/2}}{V_{m,n,\rho}} \left( \mathbb{E} \left| \frac{\ln W_{1,n}}{n} - \frac{\ln W_{1,\infty}}{n} \right| + \mathbb{E} \left| \frac{\ln W_{2,n}}{m} - \frac{\ln W_{2,\infty}}{m} \right| \right. \\
&\quad \left. + \mathbb{E} \left| \frac{\ln W_{1,k}}{n} - \frac{\ln W_{1,\infty}}{n} \right| + \mathbb{E} \left| \frac{\ln W_{2,k}}{m} - \frac{\ln W_{2,\infty}}{m} \right| \right) \\
&\leq C_1 m^{(1+\delta)/2} \left( \frac{1}{n} \gamma^n + \frac{1}{m} \gamma^n + \frac{1}{n} \gamma^k + \frac{1}{m} \gamma^k \right) \\
&\leq \frac{C_2}{m^{\delta/2}} \frac{1}{1 + |x|^{2+\delta}}. \tag{4.35}
\end{aligned}$$

Next, we give an estimation for the first term at the right-hand side of (4.34). Let

$$G_{m,n,k}(x) = \mathbb{P}(Y_{m,n,k} \leq x) \quad \text{and} \quad v_k(ds, dt) = \mathbb{P}(\tilde{Y}_{m,n,k} \in ds, H_{m,n,k} \in dt).$$

Since  $Y_{m,n,k}$  and  $(\tilde{Y}_{m,n,k}, H_{m,n,k})$  are independent, we have

$$\begin{aligned}
&\mathbb{P}(Y_{m,n,0} + H_{m,n,k} \leq x + \alpha_m, Y_{m,n,0} \geq x) \\
&= \mathbb{P}(Y_{m,n,k} + \tilde{Y}_{m,n,k} + H_{m,n,k} \leq x + \alpha_m, Y_{m,n,k} + \tilde{Y}_{m,n,k} \geq x) \\
&= \iint \mathbb{P}(Y_{m,n,k} + s + t \leq x + \alpha_m, Y_{m,n,k} + s \geq x) v_k(ds, dt) \\
&= \iint \mathbf{1}_{\{t \leq \alpha_m\}} \left( G_{m,n,k}(x - s - t + \alpha_m) - G_{m,n,k}(x - s) \right) v_k(ds, dt). \tag{4.36}
\end{aligned}$$

Denote  $C_{m,n,k}^2 = \text{Var}(Y_{m,n,k})$ , then it holds  $C_{m,n,k} = 1 + O(k/n) \rightarrow 1$  as  $m \rightarrow \infty$ . By Lemma 4.1, we



obtain for all  $x \in \mathbb{R}$ ,

$$\begin{aligned}
& \left| \mathbb{P} \left( \frac{Y_{m,n,k}}{C_{m,n,k}} \leq \frac{x}{C_{m,n,k}} \right) - \Phi \left( \frac{x}{C_{m,n,k}} \right) \right| \\
& \leq \frac{C_1}{1 + |x/C_{m,n,k}|^{2+\delta}} \left( \sum_{j=k+1}^m \mathbb{E} \left| \frac{\eta_{m,n,i}}{C_{m,n,k}} + \frac{\eta_{m,n,n+i}}{C_{m,n,k}} \right|^{2+\delta} + \sum_{i=m+1}^n \mathbb{E} \left| \frac{\eta_{m,n,i}}{C_{m,n,k}} \right|^{2+\delta} \right) \\
& \leq \frac{C_2}{1 + |x|^{2+\delta}} \left( \sum_{j=k+1}^m \mathbb{E} \left| \frac{X_{2,j-1} - \mu_2}{mV_{m,n,\rho}} \right|^{2+\delta} + \sum_{i=k+1}^n \mathbb{E} \left| \frac{X_{1,i-1} - \mu_1}{nV_{m,n,\rho}} \right|^{2+\delta} \right) \\
& \leq \frac{C_3}{1 + |x|^{2+\delta}} \left( \frac{m}{m^{2+\delta} m^{-1-\delta/2}} + \frac{n}{n^{2+\delta} m^{-1-\delta/2}} \right) \\
& \leq \frac{C_4}{m^{\delta/2}} \frac{1}{1 + |x|^{2+\delta}}.
\end{aligned}$$

By the last inequality, we deduce that for all  $x \in \mathbb{R}$ ,

$$\begin{aligned}
|G_{m,n,k}(x) - \Phi(x)| & \leq \left| \mathbb{P} \left( \frac{Y_{m,n,k}}{C_{m,n,k}} \leq \frac{x}{C_{m,n,k}} \right) - \Phi \left( \frac{x}{C_{m,n,k}} \right) \right| + \left| \Phi \left( \frac{x}{C_{m,n,k}} \right) - \Phi(x) \right| \\
& \leq \frac{C}{m^{\delta/2}} \frac{1}{1 + |x|^{2+\delta}} + \exp \left\{ -\frac{x^2}{2} \right\} \left( \frac{x}{C_{m,n,k}} - x \right) \\
& \leq \frac{C}{m^{\delta/2}} \frac{1}{1 + |x|^{2+\delta}} + C \frac{k}{n} \frac{1}{1 + |x|^{2+\delta}} \\
& \leq \frac{C}{m^{\delta/2}} \frac{1}{1 + |x|^{2+\delta}}.
\end{aligned}$$

Therefore, we have for all  $x \in \mathbb{R}$ ,

$$\mathbb{P} \left( Y_{m,n,0} + H_{m,n,k} \leq x + \alpha_m, Y_{m,n,0} \geq x \right) \leq J_1 + J_2 + J_3, \tag{4.37}$$

where

$$J_1 = \iint \mathbf{1}_{\{t \leq \alpha_m\}} |\Phi(x - s - t + \alpha_m) - \Phi(x - s)| v_k(ds, dt),$$

$$J_2 = \frac{C}{m^{\delta/2}} \iint \mathbf{1}_{\{t \leq \alpha_m\}} \frac{1}{1 + |x - s|^{2+\delta}} v_k(ds, dt)$$

and

$$J_3 = \frac{C}{m^{\delta/2}} \iint \mathbf{1}_{\{t \leq \alpha_m\}} \frac{1}{1 + |x - s - t|^{2+\delta}} v_k(ds, dt).$$

For  $J_1$ , by the mean value theorem, we have for all  $x \in \mathbb{R}$ ,

$$\begin{aligned}
& \mathbf{1}_{\{t \leq \alpha_m\}} |\Phi(x - s - t + \alpha_m) - \Phi(x - s)| \\
& \leq C |\alpha_m - t| \exp \left\{ -\frac{x^2}{8} \right\} + |\alpha_m - t| \mathbf{1}_{\{|s| \geq 1 + \frac{1}{4}|x|\}} + |\alpha_m - t| \mathbf{1}_{\{|t| \geq 1 + \frac{1}{4}|x|\}}
\end{aligned}$$

and

$$J_1 \leq J_{11} + J_{12} + J_{13}, \quad (4.38)$$

where

$$\begin{aligned} J_{11} &= C \iint |\alpha_m - t| \exp\left\{-\frac{x^2}{8}\right\} v_k(ds, dt), \\ J_{12} &= \iint |\alpha_m - t| \mathbf{1}_{\{|s| \geq 1 + \frac{1}{4}|x|\}} v_k(ds, dt) \end{aligned}$$

and

$$J_{13} = \iint |\alpha_m - t| \mathbf{1}_{\{|t| \geq 1 + \frac{1}{4}|x|\}} v_k(ds, dt).$$

By Lemma 4.3, it is obvious that for all  $x \in \mathbb{R}$ ,

$$\begin{aligned} J_{11} &\leq C_1 \exp\left\{-\frac{x^2}{8}\right\} \left(\alpha_m + \mathbb{E}|H_{m,n,k}|\right) \\ &\leq \frac{C_2}{m^{\delta/2}} \frac{1}{1 + |x|^{2+\delta}}. \end{aligned} \quad (4.39)$$

For  $J_{12}$ , we have the following estimation for all  $x \in \mathbb{R}$ ,

$$J_{12} \leq \alpha_m \mathbb{P}\left(|\tilde{Y}_{m,n,k}| \geq 1 + \frac{1}{4}|x|\right) + \mathbb{E}|H_{m,n,k}| \mathbf{1}_{\{|\tilde{Y}_{m,n,k}| \geq 1 + \frac{1}{4}|x|\}}.$$

Denote  $\tilde{C}_{m,n,k}^2 = \text{Var}(\tilde{Y}_{m,n,k})$ , then we have  $\tilde{C}_{m,n,k}^2 \asymp \frac{1}{m^{\delta/2}}$ . Let  $\delta' \in (0, \delta)$ . By Lemma 4.1, we deduce that for all  $x \in \mathbb{R}$ ,

$$\begin{aligned} \mathbb{P}\left(|\tilde{Y}_{m,n,k}| \geq 1 + \frac{1}{4}|x|\right) &= \mathbb{P}\left(\frac{\tilde{Y}_{m,n,k}}{\tilde{C}_{m,n,k}} \geq \frac{1 + |x|/4}{\tilde{C}_{m,n,k}}\right) + \mathbb{P}\left(\frac{\tilde{Y}_{m,n,k}}{\tilde{C}_{m,n,k}} \leq -\frac{1 + |x|/4}{\tilde{C}_{m,n,k}}\right) \\ &\leq 1 - \Phi\left(\frac{1 + |x|/4}{\tilde{C}_{m,n,k}}\right) + \Phi\left(-\frac{1 + |x|/4}{\tilde{C}_{m,n,k}}\right) \\ &\quad + \frac{C}{1 + \left|\frac{1 + |x|/4}{\tilde{C}_{m,n,k}}\right|^{2+\delta}} \sum_{i=1}^k \mathbb{E}\left|\frac{\eta_{m,n,i} + \eta_{m,n,n+i}}{\tilde{C}_{m,n,k}}\right|^{2+\delta} \\ &\leq \frac{C_1}{1 + |x|^{2+\delta}} \frac{1}{m^{\delta(2+\delta)/4}} \frac{1}{k^{\delta/2}} \\ &\leq \frac{C_2}{1 + |x|^{2+\delta}} \frac{1}{m^{\delta}} \end{aligned} \quad (4.40)$$

and, by Hölder's inequality with  $\tau = 1 + \frac{\delta + \delta'}{2 + 2\delta - \delta'}$  and  $\iota$  satisfying  $\frac{1}{\tau} + \frac{1}{\iota} = 1$ , it holds for all  $|x| \leq Cm^{1/2}$ ,

$$\begin{aligned} \mathbb{E}|H_{m,n,k}| \mathbf{1}_{\{|\tilde{Y}_{m,n,k}| \geq 1 + \frac{1}{4}|x|\}} &\leq \left( \mathbb{E}|H_{m,n,k}|^\tau \right)^{1/\tau} \mathbb{P} \left( |\tilde{Y}_{m,n,k}| \geq 1 + \frac{1}{4}|x| \right)^{1/\iota} \\ &\leq C \frac{1}{m^{1/2}} \left( \frac{C_1}{1 + |x|^{2+\delta}} \frac{1}{m^\delta} \right)^{1/\iota} \\ &\leq \frac{C}{m^{\delta/2}} \frac{1}{1 + |x|^{1+\delta'}}. \end{aligned}$$

Hence, we have for all  $|x| \leq Cm^{1/2}$ ,

$$J_{12} \leq \frac{C_3}{m^{\delta/2}} \frac{1}{1 + |x|^{1+\delta'}}.$$

For  $J_{13}$ , we have for all  $x \in \mathbb{R}$ ,

$$J_{13} \leq \alpha_m \mathbb{P} \left( |H_{m,n,k}| \geq 1 + \frac{1}{4}|x| \right) + \mathbb{E}|H_{m,n,k}| \mathbf{1}_{\{|H_{m,n,k}| \geq 1 + \frac{1}{4}|x|\}}.$$

By Lemma 4.3 with  $p' = 1 + \delta/2$  and Markov's inequality, we deduce that for all  $|x| \leq Cm^{1/2}$ ,

$$\begin{aligned} \mathbb{P} \left( |H_{m,n,k}| \geq 1 + \frac{1}{4}|x| \right) &\leq \frac{4^{p'}}{1 + |x|^{p'}} \mathbb{E}|H_{m,n,k}|^{p'} \leq \frac{C}{1 + |x|^{p'}} \frac{1}{m^{p'/2}} \\ &\leq \frac{C}{1 + |x|^{2+\delta}}, \end{aligned} \tag{4.41}$$

and, by Lemma 4.3 with  $p'' = \frac{1}{2}(\delta + \delta')$ ,

$$\begin{aligned} \mathbb{E}|H_{m,n,k}| \mathbf{1}_{\{|H_{m,n,k}| \geq 1 + \frac{1}{4}|x|\}} &\leq \frac{C_1}{1 + |x|^{p''}} \mathbb{E}|H_{m,n,k}|^{1+p''} \leq \frac{C_2}{1 + |x|^{p''}} \frac{1}{m^{(1+p'')/2}} \\ &\leq \frac{C_3}{m^{\delta/2}} \frac{1}{1 + |x|^{1+\delta'}}. \end{aligned}$$

Hence, we have for all  $|x| \leq Cm^{1/2}$ ,

$$J_{13} \leq \frac{C}{m^{\delta/2}} \frac{1}{1 + |x|^{1+\delta'}}.$$

Returning to (4.38), we get for all  $|x| \leq Cm^{1/2}$ ,

$$J_1 \leq \frac{C}{m^{\delta/2}} \frac{1}{1 + |x|^{1+\delta'}}. \tag{4.42}$$

Next, we consider  $J_2$ . By an argument similar to the proof of (4.40), we have for all  $x \in \mathbb{R}$ ,

$$\begin{aligned}
J_2 &= \frac{C_1}{m^{\delta/2}} \int \int \mathbf{1}_{\{t \leq \alpha_m\}} \frac{1}{1 + |x - s|^{2+\delta}} v_k(ds, dt) \\
&\leq \frac{C_1}{m^{\delta/2}} \left( \int_{|s| < 1 + |x|/2} \frac{1}{1 + |x - s|^{2+\delta}} v_k(ds) + \int_{|s| \geq 1 + |x|/2} \frac{1}{1 + |x - s|^{2+\delta}} v_k(ds) \right) \\
&\leq \frac{C_2}{m^{\delta/2}} \left[ \frac{1}{1 + |x/2|^{2+\delta}} + \mathbb{P} \left( \left| \frac{\tilde{Y}_{m,n,k}}{\tilde{C}_{m,n,k}} \right| > \frac{1 + |x|/2}{\tilde{C}_{m,n,k}} \right) \right] \\
&\leq \frac{C_3}{m^{\delta/2}} \frac{1}{1 + |x|^{2+\delta}}.
\end{aligned} \tag{4.43}$$

For  $J_3$ , by some arguments similar to that of (4.40) and (4.41), we have for all  $|x| \leq Cm^{1/2}$ ,

$$\begin{aligned}
J_3 &= \frac{C_1}{m^{\delta/2}} \int \int \mathbf{1}_{\{t \leq \alpha_m\}} \frac{1}{1 + |x - s - t|^{2+\delta}} v_k(ds, dt) \\
&\leq \frac{C_1}{m^{\delta/2}} \left( \int \int_{|s+t| \leq 2 + |x|/2} \frac{1}{1 + |x/2|^{2+\delta}} v_k(ds, dt) + \int \int_{|s| > 1 + |x|/4} v_k(ds, dt) + \int \int_{|t| > 1 + |x|/4} v_k(ds, dt) \right) \\
&\leq \frac{C_2}{m^{\delta/2}} \left[ \frac{1}{1 + |x/2|^{2+\delta}} + \mathbb{P} \left( \left| \frac{\tilde{Y}_{m,n,k}}{\tilde{C}_{m,n,k}} \right| > \frac{1 + |x|/4}{\tilde{C}_{m,n,k}} \right) + \mathbb{P} \left( |H_{m,n,k}| > 1 + \frac{|x|}{4} \right) \right] \\
&\leq \frac{C_3}{m^{\delta/2}} \frac{1}{1 + |x|^{2+\delta}}.
\end{aligned} \tag{4.44}$$

Applying the inequalities (4.42)-(4.44) to (4.37), we get for all  $|x| \leq Cm^{1/2}$ ,

$$\mathbb{P} \left( Y_{m,n,0} + H_{m,n,k} \leq x + \alpha_m, Y_{m,n,0} \geq x \right) \leq \frac{C}{m^{\delta/2}} \frac{1}{1 + |x|^{1+\delta'}}. \tag{4.45}$$

Combining (4.34), (4.35) and (4.45) together, we get (4.28) for all  $|x| \leq Cm^{1/2}$ . This completes the proof of Lemma 4.5.  $\square$

### 4.3. Proof of Theorem 2.2

We are now in a position to end the proof of Theorem 2.2. By Lemma 4.1 and the fact  $V_{m,n,\rho} \asymp \sqrt{m^{-1} + n^{-1}}$ , we have for all  $x \in \mathbb{R}$ ,

$$\begin{aligned}
\left| \mathbb{P} \left( \sum_{i=1}^{n+m} \eta_{m,n,i} \leq x \right) - \Phi(x) \right| &\leq \frac{C_1}{1 + |x|^{2+\delta}} \left( \sum_{i=1}^m \mathbb{E} |\eta_{m,n,i} + \eta_{m,n,n+i}|^{2+\delta} + \sum_{i=m+1}^n \mathbb{E} |\eta_{m,n,i}|^{2+\delta} \right) \\
&\leq \frac{C_2}{1 + |x|^{2+\delta}} \sum_{i=1}^{m+n} \mathbb{E} |\eta_{m,n,i}|^{2+\delta} \\
&\leq \frac{C_3}{1 + |x|^{2+\delta}} \left( \frac{n}{n^{2+\delta} \left( \frac{1}{n} + \frac{1}{m} \right)^{(2+\delta)/2}} + \frac{m}{m^{2+\delta} \left( \frac{1}{n} + \frac{1}{m} \right)^{(2+\delta)/2}} \right) \\
&\leq \frac{C}{(m \wedge n)^{\delta/2}} \frac{1}{1 + |x|^{2+\delta}}.
\end{aligned} \tag{4.46}$$

Notice that

$$\begin{aligned}
\mathbb{P}(R_{m,n} \leq x) &= \mathbb{P}\left(R_{m,n} \leq x, \sum_{i=1}^{n+m} \eta_{m,n,i} \leq x\right) + \mathbb{P}\left(R_{m,n} \leq x, \sum_{i=1}^{n+m} \eta_{m,n,i} > x\right) \\
&= \mathbb{P}\left(\sum_{i=1}^{n+m} \eta_{m,n,i} \leq x\right) - \mathbb{P}\left(R_{m,n} > x, \sum_{i=1}^{n+m} \eta_{m,n,i} \leq x\right) \\
&\quad + \mathbb{P}\left(R_{m,n} \leq x, \sum_{i=1}^{n+m} \eta_{m,n,i} > x\right).
\end{aligned}$$

Applying (4.46) to the last equality, we deduce that for all  $x \in \mathbb{R}$ ,

$$\begin{aligned}
\left|\mathbb{P}(R_{m,n} \leq x) - \Phi(x)\right| &\leq \frac{C}{(m \wedge n)^{\delta/2}} \frac{1}{1 + |x|^{2+\delta}} + \mathbb{P}\left(R_{m,n} > x, \sum_{i=1}^{n+m} \eta_{m,n,i} \leq x\right) \\
&\quad + \mathbb{P}\left(R_{m,n} \leq x, \sum_{i=1}^{n+m} \eta_{m,n,i} > x\right).
\end{aligned} \tag{4.47}$$

By Lemma 4.5, it follows that for all  $x \in \mathbb{R}$ ,

$$\left|\mathbb{P}(R_{m,n} \leq x) - \Phi(x)\right| \leq \frac{C}{(m \wedge n)^{\delta/2}} \frac{1}{1 + |x|^{1+\delta'}}. \tag{4.48}$$

This completes the proof of Theorem 2.2.  $\square$

#### 4.4. Preliminary Lemmas for Theorem 2.3

To prove Theorem 2.3, we shall make use of the following lemma (see Theorem 3.1 of Grama *et al.* [15]). The lemma shows that conditions **A3** and **A4** imply the existence of a harmonic moment of positive order  $\alpha > 0$ .

**Lemma 4.6.** *Assume that conditions **A3** and **A4** are satisfied. There exists a constants  $a_0 > 0$  such that for all  $\alpha \in (0, a_0)$ , the following inequalities hold*

$$\mathbb{E}W_{1,\infty}^{-\alpha} + \mathbb{E}W_{2,\infty}^{-\alpha} < \infty \tag{4.49}$$

and

$$\sup_{n \in \mathbb{N}} (\mathbb{E}W_{1,n}^{-\alpha} + \mathbb{E}W_{2,n}^{-\alpha}) < \infty. \tag{4.50}$$

*Proof.* We give an alternative proof for Theorem 3.1 of Grama *et al.* [15]. Let  $i = 1, 2$ . By the fact that

$$W_{i,\infty}^{-\alpha} = \frac{1}{\Gamma(\alpha)} \int_0^\infty e^{-tW_{i,\infty}} t^{\alpha-1} dt,$$

we have

$$\begin{aligned}
\mathbb{E}W_{i,\infty}^{-\alpha} &= \frac{1}{\Gamma(\alpha)} \int_0^\infty \phi_i(t) t^{\alpha-1} dt \\
&= \frac{1}{\Gamma(\alpha)} \left( \int_0^1 \phi_i(t) t^{\alpha-1} dt + \int_1^\infty \phi_i(t) t^{\alpha-1} dt \right),
\end{aligned} \tag{4.51}$$

where  $\Gamma$  is the gamma function. For the first term in the above bracket, since  $0 \leq \phi_i(t) \leq 1$  ( $t \geq 0$ ), then we have for all  $a_0 > 0$ ,

$$\int_0^1 \phi_i(t) t^{a_0-1} dt \leq C \int_0^1 t^{a_0-1} dt < \infty. \quad (4.52)$$

For the second term, it is necessary to prove that there exists a positive constant  $a_0$  such that for all  $t > 0$ ,

$$\phi_i(t) \leq \frac{C}{1+t^{a_0}}. \quad (4.53)$$

Next, we use the method of Grama *et al.* [16] to complete the proof of (4.53). Let  $Y_{i,n,K}$  be defined in (4.13). By (4.15), we have for all  $t > 0$ ,

$$\phi_i(t) \leq q_{i,n,K} \mathbb{E} \phi_i(Y_{i,n,K} t) + \mathbb{P} \left( \frac{t}{\Pi_{i,n}} < t_K \right). \quad (4.54)$$

Let  $p \in (1, 2]$ . By (4.13), we have

$$\begin{aligned} q_{i,n,K} \mathbb{E} \tilde{Y}_{i,n,K}^{-a} &= \mathbb{E} \left[ \Pi_{i,n}^a \prod_{j=0}^{n-1} (p_1(\xi_{i,j}) + (1-p_1(\xi_{i,j}))\beta_K) \right] + \frac{1}{K} \mathbb{E} \Pi_{i,n}^a \\ &\quad + \frac{1}{K} \mathbb{E} \left[ \Pi_{i,n}^a(\xi_i) \sum_{k=0}^{\infty} \frac{\Pi_{i,k+1}^a(T^n \xi_i)}{\Pi_{i,k}^{p-1}(T^n \xi_i)} \left( \frac{m_{i,k}^{(p)}(T^n \xi_i)}{m_{i,k}^p(T^n \xi_i)} \right) \right]. \end{aligned}$$

Since  $\Pi_{i,n}(\xi_i)$  is independent of  $\Pi_{i,k+1}(T^n \xi_i)$  and  $\Pi_{i,k}(T^n \xi_i)$  under  $\mathbb{P}$  for any  $k \geq 0$ ,

$$\begin{aligned} q_{i,n,K} \mathbb{E} \tilde{Y}_{i,n,K}^{-a} &= \mathbb{E} \left[ \Pi_{i,n}^a \prod_{j=0}^{n-1} (p_1(\xi_{i,j}) + (1-p_1(\xi_{i,j}))\beta_K) \right] \\ &\quad + \frac{1}{K} \mathbb{E} \Pi_{i,n}^a \mathbb{E} \left[ 1 + \sum_{k=0}^{\infty} \frac{\Pi_{i,k+1}^a(T^n \xi_i)}{\Pi_{i,k}^{p-1}(T^n \xi_i)} \left( \frac{m_{i,k}^{(p)}(T^n \xi_i)}{m_{i,k}^p(T^n \xi_i)} \right) \right] \\ &= \mathbb{E} \prod_{j=0}^{n-1} [m_{i,j}^a (p_1(\xi_{i,j}) + (1-p_1(\xi_{i,j}))\beta_K)] \\ &\quad + \frac{1}{K} \mathbb{E} \Pi_{i,n}^a \mathbb{E} \left[ 1 + \sum_{k=0}^{\infty} \frac{\Pi_{i,k+1}^a(T^n \xi_i)}{\Pi_{i,k}^{p-1}(T^n \xi_i)} \left( \frac{m_{i,k}^{(p)}(T^n \xi_i)}{m_{i,k}^p(T^n \xi_i)} \right) \right] \\ &= \left\{ \mathbb{E} [m_{i,0}^a (p_1(\xi_{i,0}) + (1-p_1(\xi_{i,0}))\beta_K)] \right\}^n \\ &\quad + \frac{(\mathbb{E} m_{i,0}^a)^n}{K} \left[ 1 + \frac{1}{1 - \mathbb{E} m_{i,0}^{a-(p-1)}} \mathbb{E} \left( \frac{m_{i,0}^a m_{i,0}^{(p)}}{m_{i,0}^p} \right) \right]. \end{aligned} \quad (4.55)$$

Thus, it follows from the dominated convergence theorem and (4.55) that

$$q_{i,n,K} \mathbb{E} \tilde{Y}_{i,n,K}^{-a} \xrightarrow{a \downarrow 0} q_{i,n,K} \xrightarrow{n \rightarrow \infty} \frac{1}{K} (1+C) \xrightarrow{K \rightarrow \infty} 0.$$

Then, we take  $n_0, K_0$  and  $a_0 \in (0, \lambda_0)$  small enough such that  $q_{i, n_0, K_0} \mathbb{E} \tilde{Y}_{i, n_0, K_0}^{-a_0} < 1$ . By (4.54) and Markov's inequality, we can get

$$\begin{aligned} \phi_i(t) &\leq q_{i, n_0, K_0} \mathbb{E} \phi(\tilde{Y}_{i, n_0, K_0} t) + \mathbb{P}\left(\frac{t}{\Pi_{i, n_0}} < t_{K_0}\right) \\ &\leq q_{i, n_0, K_0} \mathbb{E} \phi(\tilde{Y}_{i, n_0, K_0} t) + \frac{C_{n_0, K_0}}{t^{a_0}}. \end{aligned} \quad (4.56)$$

Notice that  $0 \leq \phi_i(t) \leq 1$  ( $t > 0$ ). Finally, by (4.56) and Lemma 4.1 in [19], we have for all  $t > 0$ ,

$$\phi_i(t) \leq \frac{C}{1 + t^{a_0}}.$$

Let  $0 < \alpha < a_0$ , we have

$$\int_1^\infty \phi_i(t) t^{\alpha-1} dt \leq C \int_1^\infty t^{\alpha-a_0-1} dt < \infty. \quad (4.57)$$

By (4.51), (4.52) and (4.57), inequality (4.49) holds.

It remains to prove (4.50) now. Since  $x \mapsto x^{-\alpha}$  ( $\alpha > 0$ ,  $x > 0$ ) is a non-negative convex function. Then by Lemma 2.1 in [17], we have

$$\sup_{n \in \mathbb{N}} \mathbb{E} W_{i, n}^{-\alpha} = \mathbb{E} W_{i, \infty}^{-\alpha} < \infty.$$

This completes the proof of Lemma 4.6. □

Under conditions **A3** and **A4**, we have the following counterpart of Lemma 4.5.

**Lemma 4.7.** *Assume that conditions **A3** and **A4** are satisfied. Then for all  $|x| \leq \sqrt{\ln(m \wedge n)}$ ,*

$$\mathbb{P}\left(R_{m, n} \leq x, \sum_{i=1}^{n+m} \eta_{m, n, i} \geq x\right) \leq C \frac{1+x^2}{\sqrt{m \wedge n}} \exp\left\{-\frac{1}{2}x^2\right\} \quad (4.58)$$

and

$$\mathbb{P}\left(R_{m, n} \geq x, \sum_{i=1}^{n+m} \eta_{m, n, i} \leq x\right) \leq C \frac{1+x^2}{\sqrt{m \wedge n}} \exp\left\{-\frac{1}{2}x^2\right\}. \quad (4.59)$$

*Proof.* As conditions **A3** and **A4** together implies conditions **A1** and **A2**, when  $|x| \leq 1$ , the inequalities (4.58) and (4.59) are simple consequences of Lemma 4.5. Thus we only need to prove the inequalities (4.58) and (4.59) for all  $1 \leq |x| \leq \sqrt{\ln(m \wedge n)}$ . In the sequel, we only give a proof for inequality (4.58) with  $1 \leq |x| \leq \sqrt{\ln(m \wedge n)}$ , as inequality (4.59) with  $1 \leq |x| \leq \sqrt{\ln(m \wedge n)}$  can be proved in the same way.

Without loss of generality, we may assume that  $m \leq n$ . Recall the notations  $Y_{m, n, k}$ ,  $\tilde{Y}_{m, n, k}$ ,  $H_{m, n, k}$  and  $D_{m, n, k}$  defined in the proof of Lemma 4.5. Set  $\alpha_m = m^{-1/2}$  and  $k = \lceil m^{1/2} \rceil$ . From (4.34), it holds for all  $x \in \mathbb{R}$ ,

$$\begin{aligned} \mathbb{P}\left(R_{m, n} \leq x, \sum_{i=1}^{n+m} \eta_{m, n, i} \geq x\right) &\leq \mathbb{P}\left(Y_{m, n, 0} + H_{m, n, k} \leq x + \alpha_m, Y_{m, n, 0} \geq x\right) \\ &\quad + \mathbb{P}\left(|D_{m, n, k}| \geq \alpha_m\right). \end{aligned} \quad (4.60)$$

For  $\mathbb{P}(|D_{m,n,k}| \geq \alpha_m)$ , by an argument similar to (4.35), we have for all  $1 \leq |x| \leq \sqrt{\ln m}$ ,

$$\mathbb{P}(|D_{m,n,k}| > \alpha_m) \leq C \frac{x^2}{\sqrt{m}} \exp \left\{ -\frac{1}{2}x^2 \right\}. \quad (4.61)$$

Next, we give an estimation for the first term of bound (4.60). Recall the notations  $G_{m,n,k}(x)$ ,  $v_k(ds, dt)$  and  $C_{m,n,k}^2$  defined in the proof of Lemma 4.5, then we have

$$\begin{aligned} & \mathbb{P}(Y_{m,n,0} + H_{m,n,k} \leq x + \alpha_m, Y_{m,n,0} \geq x) \\ &= \iint \mathbf{1}_{\{t \leq \alpha_m\}} \left( G_{m,n,k}(x - s - t + \alpha_m) - G_{m,n,k}(x - s) \right) v_k(ds, dt) \end{aligned}$$

and  $C_{m,n,k} = 1 + O(1/\sqrt{m}) \rightarrow 1$  as  $m \rightarrow \infty$ . Using Cramér's moderate deviations (for  $|x| \leq m^{1/6}$ ) and Bernstein's inequality (for  $|x| > m^{1/6}$ ) for independent random variables, we have the following non-uniform Berry-Esseen's bound: for all  $x \in \mathbb{R}$ ,

$$\begin{aligned} & \left| \mathbb{P} \left( \frac{Y_{m,n,k}}{C_{m,n,k}} \leq \frac{x}{C_{m,n,k}} \right) - \Phi \left( \frac{x}{C_{m,n,k}} \right) \right| \\ & \leq C_1 \exp \left\{ -\frac{x^2}{2(1 + \frac{C}{\sqrt{m}}|x|)} \right\} \left( 1 + \left( \frac{|x|}{C_{m,n,k}} \right)^2 \right) \left( \sum_{j=k+1}^m \mathbb{E} \left| \frac{\eta_{m,n,i}}{C_{m,n,k}} + \frac{\eta_{m,n,n+i}}{C_{m,n,k}} \right|^3 + \sum_{i=m+1}^n \mathbb{E} \left| \frac{\eta_{m,n,i}}{C_{m,n,k}} \right|^3 \right) \\ & \leq C_2 \frac{1 + |x|^2}{\sqrt{m}} \exp \left\{ -\frac{x^2}{2(1 + \frac{C}{\sqrt{m}}|x|)} \right\}. \end{aligned}$$

By the last inequality, we deduce that for all  $x \in \mathbb{R}$ ,

$$\begin{aligned} |G_{m,n,k}(x) - \Phi(x)| & \leq \left| \mathbb{P} \left( \frac{Y_{m,n,k}}{C_{m,n,k}} \leq \frac{x}{C_{m,n,k}} \right) - \Phi \left( \frac{x}{C_{m,n,k}} \right) \right| + \left| \Phi \left( \frac{x}{C_{m,n,k}} \right) - \Phi(x) \right| \\ & \leq C_2 \frac{1 + x^2}{\sqrt{m}} \exp \left\{ -\frac{x^2}{2(1 + \frac{C}{\sqrt{m}}|x|)} \right\} + \exp \left\{ -\frac{x^2}{2C_{m,n,k}^2} \right\} \left| \frac{x}{C_{m,n,k}} - x \right| \\ & \leq C_2 \frac{1 + x^2}{\sqrt{m}} \exp \left\{ -\frac{x^2}{2(1 + \frac{C}{\sqrt{m}}|x|)} \right\} + C_3 \frac{|x|}{\sqrt{m}} \exp \left\{ -\frac{x^2}{2C_{m,n,k}^2} \right\} \\ & \leq C_4 \frac{1 + x^2}{\sqrt{m}} \exp \left\{ -\frac{x^2}{2(1 + \frac{C}{\sqrt{m}}|x|)} \right\}. \end{aligned}$$

Therefore, we have for all  $x \in \mathbb{R}$ ,

$$\mathbb{P}(Y_{m,n,0} + H_{m,n,k} \leq x + \alpha_m, Y_{m,n,0} \geq x) \leq J_1 + J_2 + J_3, \quad (4.62)$$

where

$$\begin{aligned} J_1 &= \iint \mathbf{1}_{\{t \leq \alpha_m\}} \left| \Phi(x - s - t + \alpha_m) - \Phi(x - s) \right| v_k(ds, dt), \\ J_2 &= C \iint \mathbf{1}_{\{t \leq \alpha_m\}} \frac{1 + |x - s|^2}{\sqrt{m}} \exp \left\{ -\frac{(x - s)^2}{2(1 + \frac{C}{\sqrt{m}}|x - s|)} \right\} v_k(ds, dt) \end{aligned}$$



and

$$J_3 = C \iint \mathbf{1}_{\{t \leq \alpha_m\}} \frac{1 + |x - s - t|^2}{\sqrt{m}} \exp \left\{ - \frac{(x - s - t)^2}{2(1 + \frac{C}{\sqrt{m}}|x - s - t|)} \right\} v_k(ds, dt).$$

Denote  $\tilde{C}_{m,n,k}^2 = \text{Var}(\tilde{Y}_{m,n,k})$ , then it holds  $\tilde{C}_{m,n,k}^2 = O(1/\sqrt{m})$  as  $m \rightarrow \infty$ . For the upper bound of  $J_1$ , by the mean value theorem, we have for all  $1 \leq |x| \leq \sqrt{\ln m}$ ,

$$\begin{aligned} \mathbf{1}_{\{t \leq \alpha_m\}} |\Phi(x - s - t + \alpha_m) - \Phi(x - s)| &\leq C|\alpha_m - t| \exp \left\{ - \frac{x^2}{2(1 + \frac{C}{\sqrt{m}}|x|)} \right\} \\ &\quad + |\alpha_m - t| \mathbf{1}_{\{|s| \geq 2|x|\tilde{C}_{m,n,k}\}} + |\alpha_m - t| \mathbf{1}_{\{|t| \geq C_0|x|\tilde{C}_{m,n,k}\}}, \end{aligned}$$

which leads to

$$J_1 \leq J_{11} + J_{12} + J_{13}, \quad (4.63)$$

where

$$\begin{aligned} J_{11} &= C \iint |\alpha_m - t| \exp \left\{ - \frac{x^2}{2(1 + \frac{C}{\sqrt{m}}|x|)} \right\} v_k(ds, dt), \\ J_{12} &= \iint |\alpha_m - t| \mathbf{1}_{\{|s| \geq 2|x|\tilde{C}_{m,n,k}\}} v_k(ds, dt) \end{aligned}$$

and

$$J_{13} = \iint |\alpha_m - t| \mathbf{1}_{\{|t| \geq C_0|x|\tilde{C}_{m,n,k}\}} v_k(ds, dt).$$

By Lemma 4.3, it is obvious that for all  $1 \leq |x| \leq \sqrt{\ln m}$ ,

$$\begin{aligned} J_{11} &\leq C_1 \left( \alpha_m + \mathbb{E}|H_{m,n,k}| \right) \exp \left\{ - \frac{x^2}{2(1 + \frac{C}{\sqrt{m}}|x|)} \right\} \\ &\leq \frac{C_2}{\sqrt{m}} \exp \left\{ - \frac{x^2}{2(1 + \frac{C}{\sqrt{m}}|x|)} \right\}. \end{aligned}$$

For  $J_{12}$ , we have the following estimation for all  $1 \leq |x| \leq \sqrt{\ln m}$ ,

$$J_{12} \leq \alpha_m \mathbb{P} \left( |\tilde{Y}_{m,n,k}| \geq 2|x|\tilde{C}_{m,n,k} \right) + \mathbb{E}|H_{m,n,k}| \mathbf{1}_{\{|\tilde{Y}_{m,n,k}| \geq 2|x|\tilde{C}_{m,n,k}\}}.$$

By Bernstein's inequality, we deduce that for all  $x \in \mathbb{R}$ ,

$$\begin{aligned} \mathbb{P} \left( |\tilde{Y}_{m,n,k}| \geq 2|x|\tilde{C}_{m,n,k} \right) &= \mathbb{P} \left( \frac{\tilde{Y}_{m,n,k}}{\tilde{C}_{m,n,k}} \geq 2|x| \right) + \mathbb{P} \left( \frac{\tilde{Y}_{m,n,k}}{\tilde{C}_{m,n,k}} \leq -2|x| \right) \\ &\leq 2 \exp \left\{ - \frac{(2x)^2}{2(1 + \frac{C}{\sqrt{m}}|x|)} \right\} \end{aligned}$$

and, by Cauchy-Schwarz's inequality,

$$\begin{aligned}
\mathbb{E}|H_{m,n,k}| \mathbf{1}_{\{|\tilde{Y}_{m,n,k}| \geq 2|x|\tilde{C}_{m,n,k}\}} &\leq \left( \mathbb{E}|H_{m,n,k}|^2 \right)^{1/2} \mathbb{P}\left( |\tilde{Y}_{m,n,k}| \geq 2|x|\tilde{C}_{m,n,k} \right)^{1/2} \\
&\leq \frac{C}{\sqrt{m}} \left( 2 \exp \left\{ -\frac{(2x)^2}{2(1 + \frac{C}{\sqrt{m}}|x|)} \right\} \right)^{1/2} \\
&\leq \frac{C_1}{\sqrt{m}} \exp \left\{ -\frac{x^2}{2(1 + \frac{C}{\sqrt{m}}|x|)} \right\}.
\end{aligned}$$

Hence, we have for all  $1 \leq |x| \leq \sqrt{\ln m}$ ,

$$J_{12} \leq \frac{C}{\sqrt{m}} \exp \left\{ -\frac{x^2}{2(1 + \frac{C}{\sqrt{m}}|x|)} \right\}.$$

For  $J_{13}$ , we have for all  $1 \leq |x| \leq \sqrt{\ln m}$ ,

$$J_{13} \leq \alpha_m \mathbb{P}\left( |H_{m,n,k}| \geq C_0|x|\tilde{C}_{m,n,k} \right) + \mathbb{E}\left[ |H_{m,n,k}| \mathbf{1}_{\{|H_{m,n,k}| \geq C_0|x|\tilde{C}_{m,n,k}\}} \right].$$

Notice that  $V_{m,n,\rho} \asymp \frac{1}{\sqrt{m}}$  and  $\tilde{C}_{m,n,k} \asymp \frac{1}{m^{1/4}}$ . It is easy to see that for all  $1 \leq |x| \leq \sqrt{\ln m}$ ,

$$\begin{aligned}
\mathbb{P}\left( |H_{m,n,k}| \geq C_0|x|\tilde{C}_{m,n,k} \right) &\leq \mathbb{P}\left( \left| \frac{\ln W_{1,k}}{n V_{m,n,\rho}} \right| \geq \frac{1}{2}C_0|x|\tilde{C}_{m,n,k} \right) + \mathbb{P}\left( \left| \frac{\ln W_{2,k}}{m V_{m,n,\rho}} \right| \geq \frac{1}{2}C_0|x|\tilde{C}_{m,n,k} \right) \\
&\leq T_1 + T_2.
\end{aligned}$$

By Lemma 4.6 and Markov's inequality, we have for all  $1 \leq |x| \leq \sqrt{\ln m}$ ,

$$\begin{aligned}
T_1 &\leq \mathbb{P}\left( \frac{\ln W_{1,k}}{n V_{m,n,\rho}} \geq \frac{1}{2}C_0|x|\tilde{C}_{m,n,k} \right) + \mathbb{P}\left( \frac{\ln W_{1,k}}{n V_{m,n,\rho}} \leq -\frac{1}{2}C_0|x|\tilde{C}_{m,n,k} \right) \\
&\leq \mathbb{P}\left( W_{1,k} \geq \exp\left\{ \frac{1}{2}C_0|x|n V_{m,n,\rho}\tilde{C}_{m,n,k} \right\} \right) + \mathbb{P}\left( W_{1,k}^{-1} \geq \exp\left\{ \frac{1}{2}C_0|x|n V_{m,n,\rho}\tilde{C}_{m,n,k} \right\} \right) \\
&\leq \mathbb{E}[W_{1,k}] \exp \left\{ -\frac{1}{2}C_0|x|n V_{m,n,\rho}\tilde{C}_{m,n,k} \right\} + \mathbb{E}[W_{1,k}^{-\alpha}] \exp \left\{ -\frac{1}{2}\alpha C_0|x|n V_{m,n,\rho}\tilde{C}_{m,n,k} \right\} \\
&\leq C \exp \left\{ -\frac{1}{2}x^2 \right\},
\end{aligned}$$

with  $C_0$  larger enough. Similarly, we have for all  $1 \leq |x| \leq \sqrt{\ln m}$ ,

$$T_2 \leq C \exp \left\{ -\frac{1}{2}x^2 \right\}.$$

Hence, we get for all  $1 \leq |x| \leq \sqrt{\ln m}$ ,

$$\mathbb{P}\left( |H_{m,n,k}| \geq C_0|x|\tilde{C}_{m,n,k} \right) \leq C \exp \left\{ -\frac{1}{2}x^2 \right\}.$$

Clearly, by Lemma 4.6 and the inequality  $|\ln x|^2 \leq C_\alpha(x + x^{-\alpha})$  for all  $\alpha, x > 0$ , it holds

$$\mathbb{E}|H_{m,n,k}|^2 \leq \frac{C_1}{m} \left( \mathbb{E}W_{1,k} + \mathbb{E}W_{1,k}^{-\alpha} + \mathbb{E}W_{2,k} + \mathbb{E}W_{2,k}^{-\alpha} \right) \leq \frac{C_2}{m}.$$

By Markov's inequality and Cauchy-Schwarz's inequality, we deduce that for all  $1 \leq |x| \leq \sqrt{\ln m}$ ,

$$\begin{aligned} \mathbb{E} \left[ |H_{m,n,k}| \mathbf{1}_{\{|H_{m,n,k}| \geq C_0|x|\tilde{C}_{m,n,k}\}} \right] &\leq \mathbb{E} \left[ |H_{m,n,k}| \mathbf{1}_{\{W_{1,k} \geq \exp\{\frac{1}{2}C_0|x|n V_{m,n,\rho}\tilde{C}_{m,n,k}\}\}} \right] \\ &\quad + \mathbb{E} \left[ |H_{m,n,k}| \mathbf{1}_{\{W_{2,k} \geq \exp\{\frac{1}{2}C_0|x|m V_{m,n,\rho}\tilde{C}_{m,n,k}\}\}} \right] \\ &\quad + \mathbb{E} \left[ |H_{m,n,k}| \mathbf{1}_{\{W_{1,k}^{-1} \geq \exp\{\frac{1}{2}C_0|x|n V_{m,n,\rho}\tilde{C}_{m,n,k}\}\}} \right] \\ &\quad + \mathbb{E} \left[ |H_{m,n,k}| \mathbf{1}_{\{W_{2,k}^{-1} \geq \exp\{\frac{1}{2}C_0|x|m V_{m,n,\rho}\tilde{C}_{m,n,k}\}\}} \right] \\ &\leq \exp \left\{ -\frac{1}{4}C_0|x|n V_{m,n,\rho}\tilde{C}_{m,n,k} \right\} \mathbb{E}[W_{1,k}^{1/2} |H_{m,n,k}|] \\ &\quad + \exp \left\{ -\frac{1}{4}C_0|x|m V_{m,n,\rho}\tilde{C}_{m,n,k} \right\} \mathbb{E}[W_{2,k}^{1/2} |H_{m,n,k}|] \\ &\quad + \exp \left\{ -\frac{1}{4}\alpha C_0|x|n V_{m,n,\rho}\tilde{C}_{m,n,k} \right\} \mathbb{E}[W_{1,k}^{-\alpha/2} |H_{m,n,k}|] \\ &\quad + \exp \left\{ -\frac{1}{4}\alpha C_0|x|m V_{m,n,\rho}\tilde{C}_{m,n,k} \right\} \mathbb{E}[W_{2,k}^{-\alpha/2} |H_{m,n,k}|] \\ &\leq \exp \left\{ -\frac{1}{4}C_0|x|n V_{m,n,\rho}\tilde{C}_{m,n,k} \right\} (\mathbb{E}W_{1,k})^{1/2} (\mathbb{E}|H_{m,n,k}|^2)^{1/2} \\ &\quad + \exp \left\{ -\frac{1}{4}C_0|x|m V_{m,n,\rho}\tilde{C}_{m,n,k} \right\} (\mathbb{E}W_{2,k})^{1/2} (\mathbb{E}|H_{m,n,k}|^2)^{1/2} \\ &\quad + \exp \left\{ -\frac{1}{4}\alpha C_0|x|n V_{m,n,\rho}\tilde{C}_{m,n,k} \right\} (\mathbb{E}W_{1,k}^{-\alpha})^{1/2} (\mathbb{E}|H_{m,n,k}|^2)^{1/2} \\ &\quad + \exp \left\{ -\frac{1}{4}\alpha C_0|x|m V_{m,n,\rho}\tilde{C}_{m,n,k} \right\} (\mathbb{E}W_{2,k}^{-\alpha})^{1/2} (\mathbb{E}|H_{m,n,k}|^2)^{1/2} \\ &\leq \frac{C}{\sqrt{m}} \exp \left\{ -\frac{1}{2}x^2 \right\}, \end{aligned}$$

with  $C_0$  larger enough. Hence, we have for all  $1 \leq |x| \leq \sqrt{\ln m}$ ,

$$J_{13} \leq \frac{C}{\sqrt{m}} \exp \left\{ -\frac{1}{2}x^2 \right\}.$$

Returning to (4.63), we get for all  $1 \leq |x| \leq \sqrt{\ln m}$ ,

$$J_1 \leq \frac{C}{\sqrt{m}} \exp \left\{ -\frac{x^2}{2(1 + \frac{C}{\sqrt{m}}|x|)} \right\}. \quad (4.64)$$

For  $J_2$ , by an argument similar to the proof of (4.64), we have for all  $1 \leq |x| \leq \sqrt{\ln m}$ ,

$$\begin{aligned}
J_2 &= C_1 \iint \mathbf{1}_{\{t \leq \alpha_m\}} \frac{1 + |x - s|^2}{\sqrt{m}} \exp \left\{ -\frac{(x - s)^2}{2(1 + \frac{C_2}{\sqrt{m}}|x - s|)} \right\} v_k(ds, dt) \\
&\leq \frac{C_3}{\sqrt{m}} \left( \int_{|s| \leq |x| \tilde{C}_{m,n,k}} (1 + x^2) \exp \left\{ -\frac{x^2}{2(1 + \frac{C_4}{\sqrt{m}}|x|)} \right\} v_k(ds) + \int_{|s| > |x| \tilde{C}_{m,n,k}} (1 + x^2) v_k(ds) \right) \\
&\leq \frac{C_3}{\sqrt{m}} \left[ (1 + x^2) \exp \left\{ -\frac{x^2}{2(1 + \frac{C_4}{\sqrt{m}}|x|)} \right\} + \mathbb{P} \left( \left| \frac{\tilde{Y}_{m,n,k}}{\tilde{C}_{m,n,k}} \right| > |x| \right) \right] \\
&\leq \frac{C}{\sqrt{m}} (1 + x^2) \exp \left\{ -\frac{x^2}{2(1 + \frac{C}{\sqrt{m}}|x|)} \right\}. \tag{4.65}
\end{aligned}$$

Similarly, for  $J_3$ , we have for all  $1 \leq |x| \leq \sqrt{\ln m}$ ,

$$\begin{aligned}
J_3 &= C_1 \iint \mathbf{1}_{\{t \leq \alpha_m\}} \frac{1 + |x - s - t|^2}{\sqrt{m}} \exp \left\{ -\frac{(x - s - t)^2}{2(1 + \frac{C}{\sqrt{m}}|x - s - t|)} \right\} v_k(ds, dt) \\
&\leq \frac{C_2}{\sqrt{m}} \left( \iint (1 + x^2) \exp \left\{ -\frac{x^2}{2(1 + \frac{C}{\sqrt{m}}|x|)} \right\} v_k(ds, dt) + \iint_{|s| > |x| \tilde{C}_{m,n,k}} v_k(ds, dt) \right. \\
&\quad \left. + \iint_{|t| > C_0|x| \tilde{C}_{m,n,k}} v_k(ds, dt) \right) \\
&\leq \frac{C_2}{\sqrt{m}} \left( (1 + x^2) \exp \left\{ -\frac{x^2}{2(1 + \frac{C}{\sqrt{m}}|x|)} \right\} + \mathbb{P} \left( \left| \frac{\tilde{Y}_{m,n,k}}{\tilde{C}_{m,n,k}} \right| > |x| \right) + \mathbb{P} \left( |H_{m,n,k}| > C_0|x| \tilde{C}_{m,n,k} \right) \right) \\
&\leq \frac{C_4}{\sqrt{m}} (1 + x^2) \exp \left\{ -\frac{x^2}{2(1 + \frac{C}{\sqrt{m}}|x|)} \right\}. \tag{4.66}
\end{aligned}$$

Applying the inequalities (4.64)-(4.66) to (4.62), we get for all  $1 \leq |x| \leq \sqrt{\ln m}$ ,

$$\begin{aligned}
\mathbb{P} \left( Y_{m,n,0} + H_{m,n,k} \leq x + \alpha_m, Y_{m,n,0} \geq x \right) &\leq \frac{C_1}{\sqrt{m}} (1 + x^2) \exp \left\{ -\frac{x^2}{2(1 + \frac{C}{\sqrt{m}}|x|)} \right\} \\
&\leq \frac{C_1}{\sqrt{m}} (1 + x^2) \left( 1 + \frac{C_2}{\sqrt{m}} |x|^3 \right) \exp \left\{ -\frac{x^2}{2} \right\} \\
&\leq \frac{C_3}{\sqrt{m}} (1 + x^2) \exp \left\{ -\frac{x^2}{2} \right\}. \tag{4.67}
\end{aligned}$$

Combining (4.60), (4.61) and (4.67) together, we get (4.58) for all  $1 \leq |x| \leq \sqrt{\ln m}$ . This completes the proof of Lemma 4.7.  $\square$

#### 4.5. Proof of Theorem 2.3

We give a proof of Theorem 2.3 for the case of  $\frac{\mathbb{P}(R_{m,n} \geq x)}{1 - \Phi(x)}$ ,  $x \geq 0$ . Thanks to the symmetry between  $m$  and  $n$ , the case of  $\frac{\mathbb{P}(-R_{m,n} \geq x)}{\Phi(-x)}$  can be proved in the similar way. To prove Theorem 2.3, we start

with the proofs of Lemmas 4.8 and 4.9, and conclude Theorem 2.3 by combining Lemmas 4.8 and 4.9 together. To avoid trivial case, we assume that  $m \wedge n \geq 2$ .

The following lemma gives the upper bound of Theorem 2.3.

**Lemma 4.8.** *Assume that conditions **A3** and **A4** are satisfied. Then it holds for all  $0 \leq x \leq c\sqrt{m \wedge n}$ ,*

$$\ln \frac{\mathbb{P}(R_{m,n} \geq x)}{1 - \Phi(x)} \leq C \frac{1 + x^3}{\sqrt{m \wedge n}}. \quad (4.68)$$

*Proof.* First, we consider the the case  $0 \leq x \leq \sqrt{\ln(m \wedge n)}$ . Notice that

$$\begin{aligned} \mathbb{P}(R_{m,n} \geq x) &= \mathbb{P}\left(R_{m,n} \geq x, \sum_{i=1}^{n+m} \eta_{m,n,i} \geq x\right) + \mathbb{P}\left(R_{m,n} \geq x, \sum_{i=1}^{n+m} \eta_{m,n,i} < x\right) \\ &\leq \mathbb{P}\left(\sum_{i=1}^{n+m} \eta_{m,n,i} \geq x\right) + \mathbb{P}\left(R_{m,n} \geq x, \sum_{i=1}^{n+m} \eta_{m,n,i} < x\right). \end{aligned}$$

Applying Cramér's moderate deviations for independent random variables (cf. inequality (1) in [10]) to the last equality, we deduce that for all  $0 \leq x \leq c\sqrt{m \wedge n}$ ,

$$\mathbb{P}(R_{m,n} \geq x) \leq (1 - \Phi(x)) \left(1 + C \frac{1 + x^3}{\sqrt{m \wedge n}}\right) + \mathbb{P}\left(R_{m,n} \geq x, \sum_{i=1}^{n+m} \eta_{m,n,i} < x\right).$$

By Lemma 4.7 and (4.33), it follows that for all  $0 \leq x \leq \sqrt{\ln(m \wedge n)}$ ,

$$\mathbb{P}(R_{m,n} \geq x) \leq (1 - \Phi(x)) \left(1 + C \frac{1 + x^3}{\sqrt{m \wedge n}}\right).$$

By the last inequality and the inequality  $1 + x \leq e^x$ , we get (4.68) for all  $0 \leq x \leq \sqrt{\ln(m \wedge n)}$ .

Next, we consider the case  $\sqrt{\ln(m \wedge n)} \leq x \leq c\sqrt{m \wedge n}$ . Clearly, it holds for all  $x \in \mathbb{R}$ ,

$$\begin{aligned} \mathbb{P}(R_{m,n} \geq x) &= \mathbb{P}\left(\sum_{i=1}^{n+m} \eta_{m,n,i} + \frac{\ln W_{1,n}}{n V_{m,n,\rho}} - \frac{\ln W_{2,m}}{m V_{m,n,\rho}} \geq x\right) \\ &\leq I_1 + I_2 + I_3, \end{aligned} \quad (4.69)$$

where

$$\begin{aligned} I_1 &= \mathbb{P}\left(\sum_{i=1}^{n+m} \eta_{m,n,i} \geq x \left(1 - \frac{(\frac{1}{n} + \frac{1}{m\alpha})x}{V_{m,n,\rho}}\right)\right), \\ I_2 &= \mathbb{P}\left(\frac{\ln W_{1,n}}{n V_{m,n,\rho}} \geq \frac{x^2}{n V_{m,n,\rho}}\right) \quad \text{and} \quad I_3 = \mathbb{P}\left(-\frac{\ln W_{2,m}}{m V_{m,n,\rho}} \geq \frac{x^2}{m\alpha V_{m,n,\rho}}\right) \end{aligned}$$

with  $\alpha$  given by Lemma 4.6. Next, we give some estimations for  $I_1, I_2$  and  $I_3$ . By condition **A3**,  $\sum_{i=1}^{n+m} \eta_{m,n,i}$  is a sum of independent random variables with finite moment generating functions. By

Cramér's moderate deviations for independent random variables (cf. [10]), we obtain for all  $1 \leq x \leq c\sqrt{m \wedge n}$ ,

$$\begin{aligned} I_1 &\leq \left(1 - \Phi\left(x\left(1 - \frac{(\frac{1}{n} + \frac{1}{m\alpha})x}{V_{m,n,\rho}}\right)\right)\right) \exp\left\{\frac{C}{\sqrt{m+n}}\left(x\left(1 - \frac{(\frac{1}{n} + \frac{1}{m\alpha})x}{V_{m,n,\rho}}\right)\right)^3\right\} \\ &\leq \left(1 - \Phi\left(x\left(1 - \frac{(\frac{1}{n} + \frac{1}{m\alpha})x}{V_{m,n,\rho}}\right)\right)\right) \exp\left\{C\frac{x^3}{\sqrt{m \wedge n}}\right\}. \end{aligned}$$

Using the inequalities (4.33), we deduce that for all  $x \geq 1$  and  $\varepsilon_n \in (0, \frac{1}{2}]$ ,

$$\begin{aligned} \frac{1 - \Phi(x(1 - \varepsilon_n))}{1 - \Phi(x)} &= 1 + \frac{\int_{x(1-\varepsilon_n)}^x \frac{1}{\sqrt{2\pi}} e^{-t^2/2} dt}{1 - \Phi(x)} \leq 1 + \frac{\frac{1}{\sqrt{2\pi}} e^{-x^2(1-\varepsilon_n)^2/2} x \varepsilon_n}{\frac{1}{\sqrt{2\pi(1+x)}} e^{-x^2/2}} \\ &\leq 1 + Cx^2 \varepsilon_n \exp\{Cx^2 \varepsilon_n\} \\ &\leq \exp\{2Cx^2 \varepsilon_n\}. \end{aligned} \tag{4.70}$$

Hence, by the fact  $V_{m,n,\rho} \asymp \frac{1}{\sqrt{m}}$ , it holds for all  $1 \leq x \leq c\sqrt{m \wedge n}$ ,

$$I_1 \leq \left(1 - \Phi(x)\right) \exp\left\{C\frac{x^3}{\sqrt{m \wedge n}}\right\}. \tag{4.71}$$

By Markov's inequality, it is easy to see that for all  $x \geq \sqrt{\ln(m \wedge n)}$ ,

$$\begin{aligned} I_2 &= \mathbb{P}\left(W_{1,n} \geq \exp\{x^2\}\right) \\ &\leq \exp\{-x^2\} \mathbb{E}W_{1,n} = \exp\{-x^2\} \\ &\leq C\frac{1+x}{\sqrt{m \wedge n}} \left(1 - \Phi(x)\right) \end{aligned} \tag{4.72}$$

and

$$\begin{aligned} I_3 &= \mathbb{P}\left(W_{2,m} \leq \exp\{-\alpha^{-1}x^2\}\right) \\ &\leq \exp\{-x^2\} \mathbb{E}W_{2,m}^{-\alpha} \leq C \exp\{-x^2\} \\ &\leq C\frac{1+x}{\sqrt{m \wedge n}} \left(1 - \Phi(x)\right). \end{aligned} \tag{4.73}$$

Combining (4.71)-(4.73) together, we obtain for all  $\sqrt{\ln(m \wedge n)} \leq x \leq c\sqrt{m \wedge n}$ ,

$$\begin{aligned} \mathbb{P}\left(R_{m,n} \geq x\right) &\leq \left(1 - \Phi(x)\right) \exp\left\{C_1\frac{x^3}{\sqrt{m \wedge n}}\right\} + C_2\frac{(1+x)}{\sqrt{m \wedge n}} \left(1 - \Phi(x)\right) \\ &\leq \left(1 - \Phi(x)\right) \exp\left\{C_3\frac{x^3}{\sqrt{m \wedge n}}\right\}, \end{aligned}$$

which implies the desired inequality for all  $\sqrt{\ln(m \wedge n)} \leq x \leq c\sqrt{m \wedge n}$ .  $\square$

The following lemma gives the lower bound of Theorem 2.3.

**Lemma 4.9.** *Assume that conditions **A3** and **A4** are satisfied. Then it holds for all  $0 \leq x \leq c\sqrt{m \wedge n}$ ,*

$$\ln \frac{\mathbb{P}(R_{m,n} \geq x)}{1 - \Phi(x)} \geq -C \frac{1 + x^3}{\sqrt{m \wedge n}}. \quad (4.74)$$

*Proof.* The proof of lower bound is similar to the proof of upper bound. For instance, to prove (4.74) for all  $\sqrt{\ln(m \wedge n)} \leq x \leq c\sqrt{m \wedge n}$ , we only need to notice that

$$\begin{aligned} \mathbb{P}\left(R_{m,n} \geq x\right) &= \mathbb{P}\left(\sum_{i=1}^{n+m} \eta_{m,n,i} + \frac{\ln W_{1,n}}{n V_{m,n,\rho}} - \frac{\ln W_{2,m}}{m V_{m,n,\rho}} \geq x\right) \\ &\geq I_1 - I_2 - I_3, \end{aligned}$$

where

$$\begin{aligned} I_1 &= \mathbb{P}\left(\sum_{i=1}^{n+m} \eta_{m,n,i} \geq x \left(1 + \frac{(\frac{1}{n\alpha} + \frac{1}{m})x}{V_{m,n,\rho}}\right)\right), \\ I_2 &= \mathbb{P}\left(-\frac{\ln W_{1,n}}{n V_{m,n,\rho}} \geq \frac{x^2}{n\alpha V_{m,n,\rho}}\right) \quad \text{and} \quad I_3 = \mathbb{P}\left(\frac{\ln W_{2,m}}{m V_{m,n,\rho}} \geq \frac{x^2}{m V_{m,n,\rho}}\right) \end{aligned}$$

with  $\alpha$  given by Lemma 4.6. The remanning of the proof is similar to the argument of Lemma 4.8.  $\square$

## Acknowledgements

Fan would like to thank professor Quansheng Liu for his helpful discussion on the harmonic moments for branching processes in a random environment. This work has been partially supported by the National Natural Science Foundation of China (Grant no. 11971063).

## References

- [1] Afanasyev, V.I., Böinghoff, C., Kersting, G. and Vatutin, V.A. (2014). Conditional limit theorems for intermediately subcritical branching processes in random environment. *Ann. Inst. Henri Poincaré Probab. Stat.* **50**(2): 602–627.
- [2] Athreya, K.B. and Karlin, S. (1971). On branching processes with random environments: I: Extinction probabilities. *Ann. Math. Stat.* **42**(5): 1499–1520.
- [3] Athreya, K.B. and Karlin, S. (1971). Branching processes with random environments: II: Limit theorems. *Ann. Math. Stat.* **42**(6), 1843–1858.
- [4] Bansaye, V. and Böinghoff, C. (2011). Upper large deviations for branching processes in random environment with heavy tails. *Electron. J. Probab.* **16**(69): 1900–1933.
- [5] Bansaye, V. and Vatutin, V. (2017). On the survival probability for a class of subcritical branching processes in random environment. *Bernoulli* **23**(1): 58–88.

- [6] Bikelis, A. (1966). On estimates of the remainder term in the central limit theorem. *Lith. Math. J.* **6**(3): 323–346.
- [7] Böinghoff, C. (2014). Limit theorems for strongly and intermediately supercritical branching processes in random environment with linear fractional offspring distributions. *Stochastic Process. Appl.* **124**(11): 3553–3577.
- [8] Böinghoff, C and Kersting, C. (2010). Upper large deviations of branching processes in a random environment for offspring distributions with geometrically bounded tails. *Stochastic Process. Appl.* **120**(10): 2064–2077.
- [9] Chen, L.H.Y. and Shao, Q.M. (2001). A non-uniform Berry-Esseen bound via Stein’s method. *Probab. Theory Relat. Fields* **120**(2): 236–254.
- [10] Fan, X., Grama, I. and Liu, Q. (2013). Cramér large deviation expansions for martingales under Bernstein’s condition. *Stochastic Process. Appl.* **123**: 3919–3942.
- [11] Fan, X., Grama, I. and Liu, Q. (2017). Deviation inequalities for martingales with applications *J. Math. Anal. Appl.* **448**(1): 538–566.
- [12] Fan, X., Grama, I., Liu, Q., Shao, Q.M. (2019). Self-normalized Cramér type moderate deviations for martingales. *Bernoulli* **25**(4A): 2793–2823.
- [13] Fan, X., Hu, H. and Liu, Q. (2020). Uniform Cramér moderate deviations and Berry-Esseen bounds for a supercritical branching process in a random environment. *Front. Math. China* **15**(5): 891–914.
- [14] Gao, Z.Q. (2021). Exact convergence rate in the central limit theorem for a branching process in a random environment. *Stat. Probab. Letters* **178**: 109194.
- [15] Grama, I., Liu, Q. and Miqueu, E. (2017). Berry-Esseen bound and Cramér large deviation expansion for a supercritical branching process in a random environment. *Stochastic Process. Appl.* **127**(4): 1255–1281.
- [16] Grama, I., Liu, Q. and Miqueu, E. (2021). Asymptotic of the distribution and harmonic moments for a supercritical branching process in a random environment. (hal-03416307).
- [17] Huang, C. and Liu, Q. (2012). Moments, moderate and large deviations for a branching process in a random environment. *Stochastic Process. Appl.* **122**(2): 522–545.
- [18] Huang, C., Wang, C. and Wang, X. (2022). Moments and large deviations for supercritical branching processes with immigration in random environments. *Acta Math. Sci.* **42**(1): 49–72.
- [19] Liu, Q. (1999). Asymptotic properties of supercritical age-dependent branching processes and homogeneous branching random walks. *Stochastic Process. Appl.* **82**(1): 61–87.
- [20] Nakashima, M. (2013). Lower deviations of branching processes in random environment with geometrical offspring distributions. *Stochastic Process Appl* **123**(9): 3560–3587.



- [21] Smith, W.L. and Wilkinson, W.E. (1969). On branching processes in random environments. *Ann. Math. Stat.* **40**(3): 814–827.
- [22] Tanny, D. (1988). A necessary and sufficient condition for a branching process in a random environment to grow like the product of its means. *Stochastic Process. Appl.* **28**(1): 123–139.
- [23] Vatutin, V. A. A refinement of limit theorems for the critical branching processes in random environment, in: Workshop on Branching Processes and their Applications. Lect Notes Stat Proc 197 (pp. 3-19), Springer, Berlin, 2010.
- [24] Vatutin, V. and Zheng, X. (2012). Subcritical branching processes in random environment without Cramer condition. *Stochastic Process. Appl.* **122**: 2594–2609.
- [25] Wang, Y. and Liu, Q. (2017). Limit theorems for a supercritical branching process with immigration in a random environment. *Sci. China Math.* **60**(12): 2481–2502.