KHINCHIN FAMILIES, SET CONSTRUCTIONS, PARTITIONS AND EXPONENTIALS

ALICIA CANTÓN, JOSÉ L. FERNÁNDEZ, PABLO FERNÁNDEZ, AND VÍCTOR J. MACIÁ

ABSTRACT. In this paper, we give a simple criterion to verify that functions of the form e^g are in the Hayman class when g is a power series with nonnegative coefficients. Thus, using the Hayman and Báez-Duarte formulas, we obtain asymptotics for the coefficients of generating functions that arise in many examples of set construction in analytic combinatorics. This new criterion greatly simplifies that obtained previously by the authors.

1. INTRODUCTION

The present paper is a follow-up to [3]. There, following the lead of Hayman [12], Rosenbloom [15] and Báez-Duarte [2], a basic theory of the so called Khinchin families is laid out. The notions of Gaussian and strongly Gaussian power series (which include power series in the Hayman class) are developed, and it is shown how the asymptotic formulas of Hayman and Báez-Duarte (see Theorem 4.9) provide a convenient way of handling the asymptotic of coefficients of strongly Gaussian power series.

In this context, a particularly interesting question consists of establishing the asymptotic behavior of the coefficients of a power series f that is written as $f = e^g$, where g is a power series with nonnegative coefficients.

For instance, in Combinatorics, the set construction, both labeled and unlabeled, is concerned with the combinatorial class of sets formed with objects drawn from a given combinatorial class. The generating function of the class of sets is of the form $f = e^g$, where typically g is a power series with nonnegative coefficients. The generating function of the Bell numbers and the partition function are examples of functions arising from the set construction. See Section 3 of the present paper, or consult the comprehensive treatise [7] of Flajolet and Sedgewick.

Hayman, in [12], deals with this question in the following particular cases: when g (not necessarily with nonnegative coefficients) already belongs to the Hayman class, and when g is a polynomial with nonnegative coefficients that obey certain divisibility properties. See the beginning of Section 5 for details.

⁽Alicia Cantón) DEPARTAMENTO DE MATEMÁTICA E INFORMÁTICA APLICADAS A LAS INGENIERÍAS CIVIL Y NAVAL, ETSIN, UNIVERSIDAD POLITÉCNICA DE MADRID, AVENIDA DE LA MEMORIA 4, CIUDAD UNIVERSITARIA, 28040 MADRID, SPAIN

⁽José L. Fernández, Pablo Fernández and Víctor J. Maciá) DEPARTAMENTO DE MATEMÁTICAS, FACULTAD DE CIENCIAS, UNIVERSIDAD AUTÓNOMA DE MADRID, CIUDAD UNIVERSITARIA DE CAN-TOBLANCO S/N, 28049 MADRID, SPAIN.

E-mail addresses: alicia.cantonCupm.es, joseluis.fernandezCuam.es,

pablo.fernandez@uam.es, victor.macia@uam.es.

²⁰²⁰ Mathematics Subject Classification. Primary: 30B10. Secondary: 05A16, 11P82, 60F99.

Key words and phrases. Khinchin families, Hayman admissible functions, exponentials, set constructions, partitions, analytic combinatorics, asymptotic formulae, local central limit theorem.

In Theorem 4.1 of [3], a basic criterion is presented that can be used to establish conditions on the power series g with nonnegative coefficients which imply that $f = e^g$ is in the Hayman class and, therefore, is strongly Gaussian. This criterion, combined with the Hayman and Báez-Duarte asymptotic formulas, gives asymptotic results for the coefficients of a large and varied collection of power series.

In this paper, this criterion is greatly simplified and its applicability expanded. Theorems 5.5 and 5.7, the main results of the present paper, exhibit conditions on the (nonnegative) coefficients of a power series g which guarantee that e^g is in the Hayman class. The criteria mentioned above are written directly in terms of the growth of the coefficients of g. Compare with Theorem 4.1 in [3], that is reproduced as Theorem G in this paper.

By appealing to Theorems 5.5 and 5.7, the verification that the generating functions of many combinatorial set constructions is in the Hayman class becomes immediate; asymptotic formulas for their coefficients follow simply from the Hayman and Báez-Duarte asymptotic formulas.

Although this paper is a follow-up to [3], we intend this presentation to be selfcontained. The most salient and relevant features of the theory of Khinchin families are described within the present paper; nonetheless, we refer to [3] for a detailed treatment.

1.1. Notation and some preliminaries. The symbol $a_n \sim b_n$ as $n \to \infty$ means that $a_n/b_n \to 1$ as $n \to \infty$, while $a_n \simeq b_n$ as $n \to \infty$ means that $1/C \leq a_n/b_n < C$ for some C > 0.

We use \mathbf{E} , \mathbf{V} and \mathbf{P} to denote expectation, variance and probability generically.

For random variables X and Y, we write $X \stackrel{d}{=} Y$ to signify that X and Y have the same distribution.

If $(Z_n)_{n\geq 1}$ is a sequence of random variables and Z is another random variable, the notation $Z_n \xrightarrow{d} Z$ as $n \to \infty$ means convergence in distribution, which by Levy's theorem is equivalent to pointwise convergence of characteristic functions, i.e.,

$$\lim_{n \to \infty} \mathbf{E}(e^{i\theta Z_n}) = \mathbf{E}(e^{i\theta Z}), \quad \text{for every } \theta \in \mathbb{R}.$$

For sums of power of divisors of integers, we use the following notations. For any integer $c \ge 0$, we denote with $\sigma_c(m)$ the sum of the *c*th powers of the divisors of *m*:

$$\sigma_c(m) = \sum_{j|m} j^c$$
, for $m \ge 1$.

Moreover, $\sigma_c^{\text{odd}}(m)$ denotes the analogous sum but restricted to the odd divisors of m:

$$\sigma_c^{\text{odd}}(m) = \sum_{j \mid m, j \text{ odd}} j^c, \quad \text{for } m \ge 1.$$

1.2. **Plan of the paper.** Section 2 covers the basic background material on Khinchin families, the framework of the present paper, while Section 4 focuses on Gaussian and strongly Gaussian Khinchin families and the Hayman class. The reference [3] covers Khinchin families at length.

Section 3 describes the fundamental set constructions from the point of view of Khinchin families and the exponential function, which furnishes the basic context of application of the results of this paper.

The basic criteria for verifying that the exponential of a power series of nonnegative coefficients is in the Hayman class are the main results of Section 5.

Finally, in Section 6 we briefly recall the procedure to obtain asymptotic formulas of coefficients of strongly Gaussian power series.

2. Khinchin families

We denote by \mathcal{K} the class of *nonconstant* power series $f(z) = \sum_{n=0}^{\infty} a_n z^n$ with positive radius of convergence, which have *nonnegative* Taylor coefficients, $a_n \ge 0$, for each $n \ge 0$, and such that $a_0 > 0$.

The Khinchin family of such a power series $f \in \mathcal{K}$ with radius of convergence R > 0 is the family of random variables $(X_t)_{t \in [0,R)}$ with values in $\{0, 1, \ldots\}$ and with mass functions given by

$$\mathbf{P}(X_t = n) = \frac{a_n t^n}{f(t)}, \text{ for each } n \ge 0 \text{ and } t \in (0, R).$$

The variable X_0 of the family is defined as $X_0 \equiv 0$. Notice that f(t) > 0 for each $t \in [0, R)$.

Any Khinchin family is continuous in distribution in [0, R), in the sense that if a sequence $(s_n)_{n\geq 1} \subset [0, R)$ converges to $s_0 \in [0, R)$, then $X_{s_n} \stackrel{d}{\longrightarrow} X_{s_0}$, as $n \to \infty$ (see, for instance, [6]). Observe that no hypothesis upon joint distribution of the variables X_t is considered; $(X_t)_{t\in[0,R)}$ is a family, not a process.

2.1. Basic properties. For the basic theory of Khinchin families (results, proofs, examples and applications), we refer the reader to [3]. Here we describe the specific aspects of the theory to be used in the present paper.

2.1.1. Mean and variance functions. For the mean and variance of X_t we reserve the notation $m(t) = \mathbf{E}(X_t)$ and $\sigma^2(t) = \mathbf{V}(X_t)$, for $t \in [0, R)$. In terms of f, the mean and the variance of X_t may be written as

$$m(t) = \frac{tf'(t)}{f(t)}, \qquad \sigma^2(t) = tm'(t), \quad \text{for } t \in [0, R).$$

For each $t \in (0, R)$, the variable X_t is not a constant, and so $\sigma^2(t) > 0$. Consequently, m(t) is strictly increasing in [0, R), though, in general, $\sigma(t)$ is not increasing. We denote

$$M_f = \lim_{t \uparrow R} m(t) \,.$$

Whenever $M_f = +\infty$, for each integer $n \ge 0$, we use t_n to denote the unique $t_n \in [0, R)$, such that $m(t_n) = n$.

2.1.2. Normalization and characteristic functions. For each $t \in (0, R)$, the normalization of X_t is

$$\breve{X}_t \triangleq \frac{X_t - m(t)}{\sigma(t)} \cdot$$

The characteristic function of the variable X_t may be written in terms of the power series f itself as

$$\mathbf{E}(e^{i\theta X_t}) = \frac{f(te^{i\theta})}{f(t)}, \quad \text{for } t \in (0, R) \text{ and } \theta \in \mathbb{R},$$

while for its normalized version X_t we have that

$$\mathbf{E}(e^{i\theta X_t}) = \mathbf{E}(e^{i\theta X_t/\sigma(t)}) e^{-i\theta m(t)/\sigma(t)}, \quad \text{for } t \in (0, R) \text{ and } \theta \in \mathbb{R},$$

and so,

$$|\mathbf{E}(e^{i\theta X_t})| = |\mathbf{E}(e^{i\theta X_t/\sigma(t)})|, \text{ for } t \in (0, R) \text{ and } \theta \in \mathbb{R}.$$

2.1.3. Fulcrum F of f. The holomorphic function f does not vanish on the real interval [0, R), and so, it does not vanish in a simply connected region containing that interval. We may consider $\ln f$, a holomorphic branch of the logarithm of f which is real on [0, R), and the function F, which we shall call the fulcrum of f, defined and holomorphic in a region containing $(-\infty, \ln R)$, and which is given by

$$F(z) = \ln f(e^z).$$

If f does not vanish anywhere in the disk $\mathbb{D}(0, R)$, then the fulcrum F of f is defined in the whole half plane $\{z \in \mathbb{C} : \Re z < \ln R\}$. In this paper, this situation of f nonvanishing in $\mathbb{D}(0, R)$ is the most interesting. In this case $f(z) = e^{g(z)}$, where g is a function holomorphic in $\mathbb{D}(0, R)$ and $g(0) \in \mathbb{R}$, and the fulcrum F of f may be written as

$$F(z) = g(e^z)$$
, for z such that $\Re z < \ln R$.

The mean and variance function of f may be expressed in terms of its fulcrum F as

$$m(t) = F'(s)$$
 and $\sigma^2(t) = F''(s)$, for $s < \ln R$ and $t = e^s$.

2.2. Hayman's identity. For a power series $f(z) = \sum_{n=0}^{\infty} a_n z^n$ in \mathcal{K} , Cauchy's formula for the coefficient a_n may be written in terms of the characteristic function of its Khinchin family $(X_t)_{t \in [0,R)}$ as

$$a_n = \frac{f(t)}{2\pi t^n} \int_{|\theta| < \pi} \mathbf{E}(e^{i\theta X_t}) e^{-i\theta n} d\theta, \quad \text{for each } t \in (0, R) \text{ and } n \ge 1.$$

In terms of the characteristic function of the normalized variable \check{X}_t , Cauchy's formula becomes

$$a_n = \frac{f(t)}{2\pi t^n \sigma(t)} \int_{|\theta| < \pi\sigma(t)} \mathbf{E}(e^{i\theta \breve{X}_t}) e^{-i\theta(n-m(t))/\sigma(t)} d\theta \quad \text{for } t \in (0, R) \text{ and } n \ge 1.$$

If $M_f = \infty$, we may take for each $n \ge 1$ the (unique) radius $t_n \in (0, R)$ so that $m(t_n) = n$, to write

(2.1)
$$a_n = \frac{f(t_n)}{2\pi t_n^n \sigma(t_n)} \int_{|\theta| < \pi \sigma(t_n)} \mathbf{E}(e^{i\theta \check{X}_{t_n}}) \, d\theta \,, \quad \text{for each } n \ge 1 \,,$$

which we call Hayman's identity.

This identity (2.1), which is just Cauchy's formula with an appropriate choice of radius t_n , neatly encapsulates, in fact, the saddle point method.

2.3. Basic Khinchin families. The most basic collections of probability distributions in $\{0, 1, ...\}$, i.e., Bernoulli and binomial, geometric and negative binomial and Poisson, are (the most basic) Khinchin families. A quick review follows; see more details, for example, in Section 2.1.6 of [3]. (a) The Khinchin family of the function f(z) = 1 + z consists of the Bernoulli variables.

In this case, $R = \infty$, and the mean and variance functions are m(t) = t/(1+t)and $\sigma^2(t) = t/(1+t)^2$. For each t > 0, the random variable X_t is a *Bernoulli* variable with parameter p = t/(1+t).

- (b) Let $f(z) = (1+z)^N$, with integer $N \ge 1$. This is the binomial case. In this case, $R = \infty$, and the mean and variance functions are m(t) = Nt/(1+t)and $\sigma^2(t) = Nt/(1+t)^2$. For each t > 0, the random variable X_t is a binomial variable with parameters N and p = t/(1+t).
- (c) The function f(z) = 1/(1-z) corresponds to the geometric case. In this case R = 1, and the mean and variance functions are m(t) = t/(1-t)and $\sigma^2(t) = t/(1-t)^2$. For each $t \in (0,1)$, the random variable X_t is a geometric variable (number of failures until first success) of parameter 1-t, that is, $\mathbf{P}(X_t = k) = t^k(1-t)$ for $k \ge 0$.
- (d) Let $f(z) = 1/(1-z)^N$, with integer $N \ge 1$; the negative binomial case. In this case R = 1, and the mean and variance functions are m(t) = Nt/(1-t)and $\sigma^2(t) = Nt/(1-t)^2$. For each $t \in (0,1)$, the random variable X_t is a negative binomial variable of parameters $N \ge 1$ and p = 1 - t.
- (e) The Khinchin family of the exponential function $f(z) = e^z$ consists of the Poisson variables: the *Poisson case*.

In this case $R = \infty$, and the mean and variance functions are m(t) = t and $\sigma^2(t) = t$. For each t > 0, the random variable X_t in its Khinchin family is a *Poisson* variable with parameter t.

2.4. A couple of power series comparisons. In the proofs of Theorems 5.5 and 5.7, the main results of this paper, we will resort to the following asymptotics for a couple of series.

Proposition 2.1. For $\beta > 0$,

$$\sum_{n=1}^{\infty} n^{\beta-1} t^n \sim \Gamma(\beta) \frac{1}{(1-t)^{\beta}}, \quad as \ t \uparrow 1.$$

Proof. The binomial expansion gives that

$$\frac{1}{(1-z)^{\beta}} = \sum_{n=0}^{\infty} \frac{\Gamma(n+\beta)}{\Gamma(\beta) n!} z^n, \quad \text{for } |z| < 1.$$

And Stirling's formula gives, for $\beta > 0$, that

(2.2)
$$\frac{\Gamma(n+\beta)}{\Gamma(\beta) n!} \sim \frac{n^{\beta-1}}{\Gamma(\beta)}, \quad \text{as } n \to \infty.$$

Thus,

$$\sum_{n=1}^{\infty} n^{\beta-1} t^n \sim \Gamma(\beta) \frac{1}{(1-t)^{\beta}}, \quad \text{as } t \uparrow 1.$$

Remark 2.2 (Moments of geometric random variables). The power series comparison of Proposition 2.1 translates into a comparison of moments of the Khinchin family $(X_t)_{t \in [0,1)}$ of 1/(1-z), i.e., of geometric variables. Namely, the following: for $\beta > 0$,

$$\mathbf{E}(X_t^{\beta}) \sim \Gamma(\beta+1) \mathbf{E}(X_t)^{\beta}$$
, as $t \uparrow 1$.

For $\beta \in (-1,0)$, the same result holds for the variables \tilde{X}_t , the X_t conditioned at being positive, given by $\mathbf{P}(\tilde{X}_t = k) = t^{k-1}(1-t)$ for $k \ge 1$ and $t \in (0,1)$.

Proposition 2.3. For $\beta \geq 0$, we have that

$$\sum_{n=0}^{\infty} n^{\beta} \frac{t^n}{n!} \sim t^{\beta} e^t \quad and \quad \sum_{n=1}^{\infty} \frac{1}{n^{\beta}} \frac{t^n}{n!} \sim \frac{e^t}{t^{\beta}} \quad as \ t \to \infty.$$

For a proof of Proposition 2.3, we refer the reader to [4].

Remark 2.4 (Moments of Poisson random variables). Proposition 2.3 is actually a statement about positive moments of the Poisson variables X_t , and also about negative moments of the conditioned Poisson variables \tilde{X}_t , as the mean t tends to ∞ . The conditioned Poisson variables \tilde{X}_t are given by

$$\mathbf{P}(\tilde{X}_t = k) = \frac{e^{-t}}{1 - e^{-t}} \frac{t^k}{k!}, \text{ for } k \ge 1 \text{ and } t > 0.$$

This moments estimation is the following: if $(X_t)_{t \in [0,1)}$ is the Khinchin family of e^z , then for any $\beta \ge 0$,

$$\mathbf{E}(X_t^\beta) \sim t^\beta$$
, as $t \to \infty$

and for any $\beta < 0$,

$$\mathbf{E}(\tilde{X}_t^\beta) \sim t^\beta$$
, as $t \to \infty$.

See also [13] for precise asymptotic *expansions* of the negative moments of Poisson variables.

Remark 2.5 (Khinchin families and clans). We refer the reader to [4] for more information on the asymptotic comparison of $\mathbf{E}(X_t^{\beta})$ with $\mathbf{E}(X_t)^{\beta}$, as $t \uparrow R$, for a given Khinchin family (X_t) , which is behind the notion of a *Khinchin clan*.

3. Set construction and Khinchin families

The set construction of combinatorial classes fits nicely within the framework of Khinchin families. As a general reference for combinatorial classes and operations with them, we strongly suggest Chapter II in Flajolet–Sedgewick [7]. See also [1], where the constructions below are *presented* as *decomposable structures*.

We shall abbreviate 'exponential generating function' by egf, and 'ordinary generating function' by ogf.

Most of the (ordinary or exponential) generating functions of sets of combinatorial classes are exponentials of power series with nonnegative coefficients, *the object of interest* of this paper.

We will verify by means of Theorems 5.5 and 5.7 that most of the set constructions give rise to generating functions which are in the Hayman class, see Section 4.3, so that there are asymptotic formulas for their coefficients given by the Hayman and Báez-Duarte asymptotic formulas of Section 4.2.1.

3.1. Labeled combinatorial classes and sets. If $g(z) = \sum_{n=1}^{\infty} (b_n/n!) z^n$ is the exponential generating function (egf) of a labeled combinatorial class \mathcal{G} (no object of size 0), then $f(z) = e^{g(z)} = \sum_{n=0}^{\infty} (a_n/n!) z^n$ is the egf of the labeled class of sets formed with the objects of \mathcal{G} , termed assemblies in [1].

Next, we consider first sets of the basic classes: sets, lists and cycles, and then sets of rooted trees and of functions.

3.1.1. Sets of sets. Let \mathcal{G} be the labeled class of nonempty sets: $b_n = 1$, for each $n \geq 1$, and $b_0 = 0$. Its eqf is $g(z) = e^z - 1$. Then

$$f(z) = \exp(e^z - 1) = \sum_{n=0}^{\infty} \frac{B_n}{n!} z^n$$

is the eqf of sets of sets, or, equivalently, of partitions of sets, see [7], p. 107. Here, B_n is the *n*th Bell number, which counts the number of partitions of the set $\{1, \ldots, n\}$. In this case $R = \infty$, and the mean and variance function are $m(t) = te^t$ and $\sigma^2(t) = t(t+1)e^t$.

The characteristic function of the Khinchin family of f is given by

$$\mathbf{E}(e^{i heta X_t}) = \exp(e^{te^{i heta}} - e^t), \quad ext{for } heta \in \mathbb{R} ext{ and } t > 0,$$

and thus, for $\theta \in \mathbb{R}$ and t > 0,

(3.1)
$$\mathbf{E}(e^{i\theta \breve{X}_t}) = \exp\left(e^{te^{i\theta e^{-t/2}/\sqrt{t(t+1)}}} - e^t - i\theta\sqrt{t/(t+1)}e^{t/2}\right).$$

• The class \mathcal{G} of pointed sets (i.e., sets with one marked element) has egf $g(z) = ze^z$. The class of sets of pointed sets is isomorphic to the class of idempotent maps; its egf f is given by $f(z) = \exp(ze^z)$. See [7], p. 131.

3.1.2. Sets of lists. For the labeled class \mathcal{G} of (nonempty) lists, the function g is just g(z) = z/(1-z). And the function $f(z) = \exp(z/(1-z))$ is the eqf of the sets of lists, the so called fragmented permutations. See [7], p. 125.

3.1.3. Sets of cycles. The labeled class \mathcal{G} of (nonempty) cycles has $g(z) = \ln(1/(1-z))$ as egf. The function $f(z) = \exp(g(z)) = 1/(1-z)$ is the egf of the sets of cycles, or, in other terms, of the permutations.

• The length of the cycles could be restricted. Thus, for integer $k \ge 1$,

$$\exp\left(\sum_{n\leq k} z^n/n\right)$$
 or $\exp\left(\sum_{n\geq k} z^n/n\right)$

are the egfs of permutations such that all the cycles in their cycle decomposition have length at most k, or at least k, respectively. We may also consider, for $k \ge 1$,

$$\exp\Big(\sum_{d\ge 1,\,d|k} z^d/d\Big),\,$$

which is the eqf of the permutations σ such that σ^k is the identity.

3.1.4. Sets of trees and sets of functions.

• Sets of trees (forests). The class \mathcal{G} of rooted (labeled) trees has egf $g(z) = \sum_{n=1}^{\infty} (n^{n-1}/n!) z^n$. See [7], Section II.5.1; this is Cayley's theorem. The class of forests (sets) of rooted (labeled) trees has then egf $f = e^g$.

Cayley's theorem also shows that the eqf of the class of unrooted (labeled) trees is $g(z) = \sum_{n=1}^{\infty} (n^{n-2}/n!) z^n$.

• Sets of functions. The class \mathcal{G} of functions has egf $g(z) = \sum_{n=1}^{\infty} (n^n/n!) z^n$. See [7], Section II.5.2. The class of sets of functions has then egf $f = e^g$.

3.2. Unlabeled combinatorial classes and sets. We split the discussion into multisets and (proper) sets. See Section I.2.2 in [7] and [1] as general references for the set constructions of unlabeled combinatorial classes.

3.2.1. Multisets of unlabeled combinatorial classes. If $C(z) = \sum_{n=1}^{\infty} c_n z^n$ is the ordinary generating function (ogf) of a combinatorial (unlabeled) class \mathcal{G} (no object of size 0), then

$$f(z) = \prod_{j=1}^{\infty} \frac{1}{(1-z^j)^{c_j}} = \exp\left(\sum_{n=1}^{\infty} C(z^n)/n\right)$$

is the ogf of the class of sets formed with the objects of \mathcal{G} , termed multisets in [1].

We may write $f(z) = e^{g(z)}$, where g is the power series

$$g(z) = \sum_{j,k \ge 1} c_j \frac{z^{kj}}{k} = \sum_{m=1}^{\infty} \left(\sum_{j|m} jc_j \right) \frac{z^m}{m} \triangleq \sum_{m=1}^{\infty} b_m z^m.$$

Observe that the power series g has nonnegative coefficients:

$$b_m = \frac{1}{m} \sum_{j|m} jc_j$$
, for $m \ge 1$

• The ogf of partitions, the partition function, given by

$$P(z) = \prod_{j=1}^{\infty} \frac{1}{1-z^j} = \sum_{n=0}^{\infty} p(n) z^n, \quad \text{for } z \in \mathbb{D},$$

is in \mathcal{K} . Here p(n) is the number of partitions of the integer $n \geq 1$.

The mean and variance functions of its Khinchin family are not so direct. In this instance, C is C(z) = z/(1-z), as $c_n = 1$ for $n \ge 1$ (one object of weight n, for each $n \ge 1$). And thus g has coefficients $b_m = \sigma_1(m)/m$, where $\sigma_1(m)$ is the sum of the divisors of the integer m.

For general $C(z) = \sum_{n=1}^{\infty} c_n z^n$ as above, the corresponding f is the ogf of the colored partitions; c_j different colors for part j.

3.2.2. Sets of unlabeled combinatorial classes. If again we write $C(z) = \sum_{n=1}^{\infty} c_n z^n$ for the ordinary generating function (ogf) of a combinatorial (unlabeled) class \mathcal{G} (no object of size 0), then

$$f(z) = \prod_{j=1}^{\infty} (1+z^j)^{c_j} = \exp\Big(\sum_{n=1}^{\infty} (-1)^{n+1} C(z^n)/n\Big),$$

is the ogf of the class of (proper) sets formed with the objects of \mathcal{G} , termed *selections* in [1].

We may write $f(z) = e^{g(z)}$, where g is the power series

$$g(z) = \sum_{j,k\geq 1} c_j \frac{z^{kj} (-1)^{k+1}}{k} = \sum_{m=1}^{\infty} \left(\sum_{jk=m} jc_j (-1)^{k+1} \right) \frac{z^m}{m} \cdot$$

In general, the power series g could have negative coefficients; this is the case, for instance, for $f(z) = (1+z)^5(1+z^2)$.

But for the sequence of coefficients of C given by $c_j = j^{c-1}, j \ge 1$, where c is an integer $c \ge 1$, the coefficients of g are nonnegative; in fact, the mth coefficient of g is

$$\frac{1}{m}\,\sigma_c^{\rm odd}(m)\,\omega(m)\,,$$

where $\omega(m)$ is given by

$$\omega(m) = \frac{2^c - 2}{2^c - 1} 2^{\chi(m)c} + \frac{1}{2^c - 1}, \quad \text{for } m \ge 1,$$

 $\chi(m)$ is the highest integer exponent so that $2^{\chi(m)} \mid m$, and $\sigma_c^{\text{odd}}(m)$ is the sum of the *c*th powers of the odd divisors of *m*. This is so because of the identity

(3.2)
$$\sum_{jk=m} j^c (-1)^{k+1} = \left(\sum_{j|m, j \text{ odd}} j^c\right) \omega(m), \quad \text{for } m \ge 1.$$

To verify (3.2), observe first that as functions of m, both summations in (3.2) are multiplicative. Write m as $m = 2^{\chi(m)}s$, with s odd, and observe that for $m = 2^r$, with $r \ge 1$, the summation on the left is $\omega(m)$ and the summation on the right is 1, while for m = s odd, the two sums coincide and $\omega(s) = 1$.

We may also write

j

$$\sum_{k=m} j^c (-1)^{k+1} = \sigma_c(m) - 2\sigma_c(m/2), \quad \text{for } m \ge 1,$$

with the understanding that $\sigma_c(m/2) = 0$ if m is odd. For the identity (3.2) and a variety of relations among a number of diverse sums on divisors, we refer to [9].

More generally, if the coefficient c_j of C is given by $c_j = R(j)$, for $j \ge 1$, where R(z) is a polynomial with nonnegative integer coefficients, then the coefficients of g are nonnegative.

• The ogf of *partitions with distinct parts*, given by

$$Q(z) = \prod_{j=1}^{\infty} (1+z^j) = \sum_{n=0}^{\infty} q(n) z^n, \quad \text{for } z \in \mathbb{D},$$

is in \mathcal{K} . Here q(n) is the number of partitions into distinct parts of the integer $n \geq 1$.

In this particular instance, the power series C is C(z) = z/(1-z), as $c_n = 1$, for $n \ge 1$, and the power series g is

$$g(z) = \sum_{m=1}^{\infty} \frac{\sigma_1(m) - 2\sigma_1(m/2)}{m} z^m$$

4. Gaussian Khinchin families and Hayman class

Definition 4.1. A power series $f \in \mathcal{K}$ and its Khinchin family $(X_t)_{t \in [0,R)}$ are termed *Gaussian* if the normalized sequence (\check{X}_t) converges in distribution, as $t \uparrow R$, to the standard normal or, equivalently, if

$$\lim_{t\uparrow R} \mathbf{E}(e^{\imath\theta\check{X}_t}) = e^{-\theta^2/2}, \quad \text{for each } \theta \in \mathbb{R}.$$

Among the basic Khinchin families considered in Section 2.3, only the family associated to the exponential e^z is Gaussian.

For the functions f(z) = 1 + z (Bernoulli case) or $f(z) = (1+z)^N$ (binomial case), the corresponding (\check{X}_t) converges in distribution, as $t \to \infty$, to the constant 0. In fact, for any polynomial in \mathcal{K} , the corresponding (\check{X}_t) converges in distribution, as $t \to \infty$, towards the constant 0.

For the function f(z) = 1/(1-z) (geometric case), (X_t) converges in distribution, as $t \uparrow 1$, towards a variable Z, where Z + 1 is an exponential variable of parameter 1. For $f(z) = 1/(1-z)^N$, with integer $N \ge 1$ (negative binomial case), it can be verified that (X_t) converges in distribution, as $t \uparrow 1$, towards a variable Z_N , where $Z_N + \sqrt{N}$ follows a Gamma distribution with shape parameter N and rate parameter \sqrt{N} (or scale parameter $1/\sqrt{N}$).

Thus, in all the cases considered above, (\check{X}_t) converges in distribution as $t \uparrow R$, but only for the exponential e^z the limit is the (standard) normal distribution. See details in Section 3.1 of [3].

4.1. Gaussianity of exponentials. The following Theorem A is the basic criterion for gaussianity in terms of the fulcrum F of f (of Section 2.1.3); it originates in Hayman's [12], but we refer to Theorem 3.2 in [3] for a proof.

Theorem A. If $f \in \mathcal{K}$ has radius of convergence R > 0 and vanishes nowhere in $\mathbb{D}(0, R)$, and if for the fulcrum F of f one has

(4.1)
$$\lim_{s\uparrow \ln R} \frac{\sup_{\phi\in\mathbb{R}} \left| F^{\prime\prime\prime}(s+i\phi) \right|}{F^{\prime\prime}(s)^{3/2}} = 0,$$

then f is Gaussian.

Notice that Theorem A applies whenever $f \in \mathcal{K}$ is of the form $f = e^g$ for some power series g, that may have negative Taylor coefficients. As registered in the following Theorem B, if $f = e^g$ and g has nonnegative coefficients, a simpler condition on g implies the gaussianity of $f = e^g$.

Theorem B. Let $f \in \mathcal{K}$ be such that $f = e^g$, where g has radius of convergence R > 0and nonnegative coefficients. If g is a polynomial of degree 1 or g satisfies

(4.2)
$$\lim_{t\uparrow R} \frac{g''(t)}{g''(t)^{3/2}} = 0,$$

then f is Gaussian.

For a proof, see Theorem 3.3 in [3].

• For the exponential function $f(z) = e^z$, we have that its fulcrum is $F(z) = e^z$, and its gaussianity follows readily from Theorem A.

• More generally, if $B(z) = \sum_{j=0}^{N} b_j z^j$ is a polynomial of degree N such that $e^B \in \mathcal{K}$, then e^B is Gaussian. For, in this case, $F(z) = B(e^z)$, and for $z = s + i\phi$ we have that

$$|F'''(z)| = \left|\sum_{j=0}^{N} b_j j^3 e^{jz}\right| \le \sum_{j=0}^{N} |b_j| j^3 e^{js} = O(e^{Ns}),$$

while

(4.3)
$$F''(s) = \sum_{j=0}^{N} b_j \, j^2 e^{js} \sim b_N \, N^2 e^{Ns} \quad \text{as } s \uparrow \infty \,,$$

and gaussianity follows from Theorem A. Observe that (4.3) implies that b_N is real and $b_N > 0$; besides, since $e^B \in \mathcal{K}$, the coefficients of the polynomial B must be real numbers.

• For $f(z) = \exp(e^z - 1)$, the eqf of the class of sets of sets, and for $f(z) = \exp(z/(1-z))$, the eqf of the class of sets of lists, Theorem A (or Theorem B) gives readily that both are Gaussian. Similarly, the function $f(z) = \exp(ze^z)$, the eqf of the class of sets of pointed sets, is Gaussian.

• Again directly from Theorem A or Theorem B, we see that the functions $f(z) = \exp(1/(1-z)^{\gamma})$, with $\gamma > 0$, are all Gaussian. The functions

$$f(z) = \exp\left(\sum_{n=1}^{\infty} n^{\alpha} z^n\right), \text{ for } |z| < 1,$$

with $\alpha > -1$ are also Gaussian. Indeed, the fulcrum F of such an f is

$$F(z) = \sum_{n=1}^{\infty} n^{\alpha} e^{nz}, \quad \text{for } \Re z < 0.$$

Then, for s < 0 and $r = e^s$, we have that

$$\sup_{\phi \in \mathbb{R}} |F'''(s + \imath \phi)| = \sum_{n=1}^{\infty} n^{\alpha+3} r^n, \quad \text{while} \quad |F''(s)| = \sum_{n=1}^{\infty} n^{\alpha+2} r^n.$$

By appealing to Proposition 2.1, we deduce that condition (4.1) is satisfied. Observe that R = 1, in this case.

• In the same vein, the eqf of sets of functions $f(z) = \exp\left(\sum_{n=1}^{\infty} (n^n/n!)z^n\right)$, with R = 1/e, is also Gaussian. Similarly, and more generally,

$$f(z) = \exp\left(\sum_{n=1}^{\infty} \frac{n^{n-\delta}}{n!} z^n\right)$$

is Gaussian, for $0 \le \delta < 1/2$. To see this, observe (Stirling's formula) that

$$\frac{n^{n-\delta}}{n!} \asymp \frac{e^n}{n^{\delta+1/2}}, \quad \text{as } n \to \infty.$$

and argue as above with $\alpha = -\delta - 1/2$.

• The partition function P and the ogf Q of partitions into distinct parts are seen to be Gaussian as a consequence of Theorem A; but see Section 5 for the stronger statement that P and Q are both in the Hayman class.

Now we turn our attention to a couple of non-examples.

• The egf $f(z) = \exp(\ln(1/(1-z)))$ of the class of sets of cycles, which is simply f(z) = 1/(1-z), the egf of permutations, is not Gaussian, as we have mentioned at the beginning of this section. The condition of Theorem A is, of course, not satisfied; in fact, for the corresponding fulcrum $F(z) = -\ln(1-e^z)$, for $\Re z < 0$, it holds that

$$\lim_{s \uparrow 0} \frac{\sup_{\phi \in \mathbb{R}} |F'''(s+i\phi)|}{F''(s)^{3/2}} = 2.$$

Analogously, the class of sets of cycles of length at least k is not Gaussian, for any $k \ge 1$.

• The egf $f(z) = \exp\left(\sum_{n=1}^{\infty} (n^{n-1}/n!)z^n\right)$ of forests of rooted trees is not Gaussian. In fact, we are going to check that the characteristic functions of its normalized Khinchin family converge in distribution to the constant 1.

The power series f has radius of convergence R = 1/e, and

$$\lim_{t\uparrow 1/e} f(t) = \exp\left(\sum_{n=1}^{\infty} \frac{n^{n-1}}{n! e^n}\right) < +\infty,$$

since

$$\frac{n^{n-1}}{n!\,e^n} \asymp \frac{1}{n^{3/2}} \quad \text{as } n \to \infty.$$

In particular, the function f extends to be continuous in $cl(\mathbb{D}(0, 1/e))$. Now

$$m(t) = \sum_{n=1}^{\infty} \frac{n^n}{n!} t^n$$
, for $t \in (0, 1/e)$,

and Proposition 2.1 gives that

$$m(t/e) = \sum_{n=1}^{\infty} \frac{n^n}{n! e^n} t^n \asymp \frac{1}{(1-t)^{1/2}}, \text{ as } t \uparrow 1.$$

Also

$$\sigma^{2}(t) = \sum_{n=1}^{\infty} \frac{n^{n+1}}{n!} t^{n}, \quad \text{for } t \in (0, 1/e) ,$$

and Proposition 2.1 gives that

$$\sigma^2(t/e) = \sum_{n=1}^{\infty} \frac{n^{n+1}}{n! \, e^n} \, t^n \asymp \frac{1}{(1-t)^{3/2}} \,, \quad \text{as } t \uparrow 1 \,.$$

Thus

$$\frac{\sigma(t/e)}{m(t/e)} \asymp \frac{1}{(1-t)^{1/4}}\,, \quad \text{as } t\uparrow 1\,,$$

and so, in particular,

(4.4)
$$\lim_{t\uparrow 1/e}\frac{m(t)}{\sigma(t)}=0\,.$$

Now for $\theta \in \mathbb{R}$ we may write

$$\mathbf{E}(e^{i\theta\breve{X}_t}) = \frac{f(te^{i\theta/\sigma(t)})}{f(t)} e^{-i\theta m(t)/\sigma(t)} .$$

The first factor of this expression of the characteristic function of \check{X}_t tends towards 1 because $\lim_{t\uparrow 1} \sigma(t) = +\infty$ and the continuity of f on $\operatorname{cl}(\mathbb{D}(0, 1/e))$, while the second factor tends towards 1, as a consequence of (4.4). And thus, $\lim_{t\uparrow 1/e} \mathbf{E}(e^{i\theta\check{X}_t}) = 1$, for any $\theta \in \mathbb{R}$.

This means that X_t tends in distribution towards the constant 0, and not to the Gaussian distribution.

Similarly, and more generally, for

$$f(z) = \exp\left(\sum_{n=1}^{\infty} \frac{n^{n-\delta}}{n!} z^n\right)$$

the corresponding (\check{X}_t) tends in distribution, as $t \uparrow 1$, towards the constant 0, for $1/2 < \delta \leq 5/2$, and thus, the efg of the sets of functions is Gaussian, but the egfs of sets of trees and of sets of rooted trees are not Gaussian.

Question 1. Is

$$f(z) = \exp\left(\sum_{n=1}^{\infty} \frac{n^{n-1/2}}{n!} z^n\right)$$

Gaussian? It is not strongly Gaussian (see the definition below) since for its Khinchin family we have that

$$\lim_{t \uparrow 1/e} \frac{m(t)}{\sigma(t)} = (2\pi)^{-1/4} \, ,$$

while for strongly Gaussian power series it is always the case that that limit is $+\infty$. See formula (4.6) below.

4.2. Strongly Gaussian Khinchin families. The notion of strongly Gaussian power series f in \mathcal{K} was introduced by Báez-Duarte in [2].

Definition 4.2. A power series $f \in \mathcal{K}$ and its Khinchin family $(X_t)_{t \in [0,R)}$ are termed strongly Gaussian if

a)
$$\lim_{t\uparrow R} \sigma(t) = +\infty$$
, and b) $\lim_{t\uparrow R} \int_{|\theta| < \pi\sigma(t)} \left| \mathbf{E}(e^{i\theta \breve{X}_t}) - e^{-\theta^2/2} \right| d\theta = 0.$

The exponential $f(z) = e^z$ is strongly Gaussian. In this case $\sigma(t) = \sqrt{t}$. This strong gaussianity follows from the gaussianity of e^z and dominated convergence using the bound

$$\left|\mathbf{E}(e^{i\theta \check{X}_t})\right| = e^{t(\cos(\theta/\sqrt{t})-1)} \le e^{-2\theta^2/\pi^2}, \quad \text{for } \theta \in \mathbb{R}, \, t > 0 \text{ such that } |\theta| < \pi\sqrt{t}.$$

As we will see in Theorem C, strongly Gaussian power series are Gaussian. Thus the power series $(1+z)^N$ and $1/(1-z)^N$, for $N \ge 1$, are all not strongly Gaussian.

For strongly Gaussian power series, we have the following key Theorem C. It appears in [12] for power series in the Hayman class. See Theorem A in [3] for a proof under the more general asymption of strongly Gaussian power series.

Theorem C (Hayman's central limit theorem). If the power series $f(z) = \sum_{n=0}^{\infty} a_n z^n$ in \mathcal{K} is strongly Gaussian, then

(4.5)
$$\lim_{t \uparrow R} \sup_{n \in \mathbb{Z}} \left| \frac{a_n t^n}{f(t)} \sqrt{2\pi} \sigma(t) - e^{-(n-m(t))^2/(2\sigma^2(t))} \right| = 0.$$

Moreover,

$$\lim_{t\uparrow R} \mathbf{P}(\breve{X}_t \le b) = \Phi(b), \quad \text{for every } b \in \mathbb{R},$$

and so, (\breve{X}_t) converges in distribution towards the standard normal and f is Gaussian.

In this statement, $a_n = 0$, for n < 0.

By considering n = -1 in (4.5), it follows that

(4.6)
$$\lim_{t\uparrow R} \frac{m(t)}{\sigma(t)} = +\infty$$

and, in particular, that $M_f = \infty$ for every $f \in \mathcal{K}$ that is strongly Gaussian.

The function $f(z) = e^{z^2}$ is Gaussian, because of Theorem A. Its variance function $\sigma^2(t) = 4t^2$ tends towards ∞ as $t \to \infty$, but f is not strongly Gaussian, since its Taylor coefficients of odd order are null and do not satisfy the asymptotic formula (4.7) below.

4.2.1. *Coefficients of strongly Gaussian power series.* For strongly Gaussian power series, we have the following asymptotic formula for its coefficients.

Theorem D (Hayman's asymptotic formula). If $f(z) = \sum_{n=0}^{\infty} a_n z^n$ in \mathcal{K} is strongly Gaussian, then

(4.7)
$$a_n \sim \frac{1}{\sqrt{2\pi}} \frac{f(t_n)}{t_n^n \sigma(t_n)}, \quad as \ n \to \infty.$$

In the asymptotic formula above, t_n is given by $m(t_n) = n$, for each $n \ge 1$. This asymptotic formula follows readily from Theorem C, or alternatively, from Hayman's formula (2.1) and strong gaussianity.

For the exponential function $f(z) = e^z$, one has m(t) = t and $\sigma(t) = \sqrt{t}$, for $t \ge 0$, and $t_n = n$ for $n \ge 1$. The asymptotic formula above gives

$$\frac{1}{n!} \sim \frac{1}{\sqrt{2\pi}} \frac{e^n}{n^n \sqrt{n}}, \quad \text{as } n \to \infty,$$

that is Stirling's formula.

Actually, if ω_n is a good approximation of t_n , in the sense that

$$\lim_{n \to \infty} \frac{m(\omega_n) - n}{\sigma(\omega_n)} = 0$$

then from Hayman's formula (2.1) and strong gaussianity we have that

(4.8)
$$a_n \sim \frac{1}{\sqrt{2\pi}} \frac{f(\omega_n)}{\sigma(\omega_n)\,\omega_n^n}, \quad \text{as } n \to \infty.$$

In general, precise expressions for the t_n are rare, since inverting m(t) is usually complicated. But, fortunately, in practice, one can do with a certain asymptotic approximation due to Báez-Duarte, [2], which we now describe.

Suppose that $f \in \mathcal{K}$ is strongly Gaussian. Assume that $\widetilde{m}(t)$ is continuous and monotonically increasing to $+\infty$ in [0, R) and that $\widetilde{m}(t)$ is a good approximation of m(t) in the sense that

(4.9)
$$\lim_{t\uparrow R} \frac{m(t) - \widetilde{m}(t)}{\sigma(t)} = 0.$$

Let τ_n be defined by $\widetilde{m}(\tau_n) = n$, for each $n \ge 1$.

Theorem E (Báez-Duarte asymptotic formula). With the notations above, if $f(z) = \sum_{n=0}^{\infty} a_n z^n$ in \mathcal{K} is strongly Gaussian and (4.9) is satisfied, then

$$a_n \sim \frac{1}{\sqrt{2\pi}} \frac{f(\tau_n)}{\tau_n^n \, \sigma(\tau_n)}, \quad as \ n \to \infty.$$

This follows readily from (4.8).

Besides, if $\tilde{\sigma}(t)$ is such that $\sigma(t) \sim \tilde{\sigma}(t)$ as $t \uparrow R$, we may further write

(4.10)
$$a_n \sim \frac{1}{\sqrt{2\pi}} \frac{f(\tau_n)}{\tau_n^n \,\widetilde{\sigma}(\tau_n)}, \quad \text{as } n \to \infty.$$

In practice, using (4.10) requires approximating m by \tilde{m} to obtain τ_n , and then obtaining good enough approximations of σ and f on (0, R) to produce asymptotic formulas of $\sigma(\tau_n)$ and $f(\tau_n)$.

4.3. Hayman class. The class of Hayman consists of power series f in \mathcal{K} which satisfy some concrete and verifiable conditions which imply that f is strongly Gaussian, see Theorem F below.

Definition 4.3. A power series $f \in \mathcal{K}$ is in the Hayman class (or is Haymanadmissible or just H-admissible) if for a certain function $h: [0, R) \to (0, \pi]$, the following conditions are satisfied:

(4.11) (major arc):
$$\lim_{t\uparrow R} \sup_{|\theta| \le h(t)\,\sigma(t)} \left| \mathbf{E}(e^{i\theta X_t}) \, e^{\theta^2/2} - 1 \right| = 0 \,,$$

(4.12) (minor arc):
$$\lim_{t\uparrow R} \sigma(t) \sup_{h(t)\sigma(t) \le |\theta| \le \pi\sigma(t)} |\mathbf{E}(e^{i\theta \check{X}_t})| = 0,$$

and

(4.13) (variance condition):
$$\lim_{t\uparrow R} \sigma(t) = \infty.$$

We refer to the function h in the definition above as a *cut* between a *major* arc and a *minor* arc.

Some authors include in the Hayman class power series with a finite number of negative coefficients. This is not the case in this paper.

For f in the Hayman class, the characteristic function of \check{X}_t is uniformly approximated by $e^{-\theta^2/2}$ in the major arc, while it is uniformly $o(1/\sigma(t))$ in the minor arc.

Observe that condition (4.12) may be written more simply and in terms of f itself as the requirement that

$$\lim_{t\uparrow R} \sigma(t) \sup_{h(t) \le |\theta| \le \pi} |\mathbf{E}(e^{i\theta X_t})| = \lim_{t\uparrow R} \sigma(t) \sup_{h(t) \le |\theta| \le \pi} \frac{|f(te^{i\theta})|}{f(t)} = 0.$$

Theorem F. Power series in the Hayman class are strongly Gaussian.

In a certain sense, the above theorem places the conditions for Hayman class as a criterion for being strongly Gaussian. For a proof, see Theorem 3.7 in [3]. Thus we have that being in the Hayman class implies strong gaussianity, which in turn implies gaussianity. In applications, we shall always check that the power series belongs to the Hayman class.

We refer to [3], and, of course, to [12], for examples and further properties of the Hayman class.

5. Exponentials and the Hayman class

Along this section, we consider nonconstant power series $g(z) = \sum_{n=0}^{\infty} b_n z^n$ with *nonnegative coefficients* and radius of convergence R > 0; it is not required that g(0) > 0. We are interested in conditions on g which guarantee that $f = e^g$, which is in \mathcal{K} , is in the Hayman class.

There are two results of Hayman in [12] along this line.

- (a) If g is a power series in the Hayman class, then $f = e^g$ is in the Hayman class; see Theorem VI in [12].
- (b) If B is a nonconstant polynomial with nonnegative coefficients and such that $Q_B = \gcd\{n \ge 1 : b_n > 0\} = 1$, then e^B is in the Hayman class.

Remark 5.1. In [14], it is shown that if g is in the Hayman class, then $f = e^g$ satisfies the stronger (than Hayman) conditions of Harris and Schoenfeld, [11], which allow to obtain full asymptotic expansions of coefficientes and not just asymptotic formulas.

Remark 5.2. In Theorem X of [12], Hayman proves the following stronger version of (b): Let $B(z) = \sum_{n=0}^{N} b_n z^n$ be a nonconstant polynomial with real coefficients such that for each d > 1 there exists m, not a multiple of d, such that $b_m \neq 0$, and such that if m(d) is the largest such m, then $b_{m(d)} > 0$. If e^B is in \mathcal{K} , then e^B is in the Hayman class.

For a direct proof of E2), we refer to Proposition 5.1 in [3].

Theorem G, stated below, which is Theorem 4.1 in [3], exhibits conditions on the function g which ensure that f is in the Hayman class.

Further, we describe in Theorem 5.3 a 'practical' approach to verify that a power series satisfies the conditions of Theorem G.

Finally, Theorems 5.5 and 5.7 give easily verifiable conditions on the coefficients b_n of g that imply that $f = e^g$ is in the Hayman class. These theorems may be directly applied to the generating functions of sets constructions (and also to other exponentials) to check that they belong to the Hayman class.

Let $(X_t)_{t \in [0,R)}$ be the Khinchin family of $f = e^g$. The mean and variance functions of f, written in terms of g, are

(5.1)
$$m(t) = \frac{tf'(t)}{f(t)} = tg'(t) = \sum_{n=1}^{\infty} nb_n t^n, \text{ for } t \in (0, R),$$

and

(5.2)
$$\sigma^{2}(t) = tm'(t) = tg'(t) + t^{2}g''(t) = \sum_{n=1}^{\infty} n^{2}b_{n} t^{n}, \text{ for } t \in (0, R).$$

Since g has nonnegative coefficients, the variance function $\sigma^2(t)$ of f is increasing in [0, R). Observe also that $m(t) \leq \sigma^2(t)$, for $t \in [0, R)$.

The variance condition (4.13) required for f to be in the Hayman class translates readily in the following condition in terms of g:

$$\lim_{t\uparrow R} \left(tg'(t) + t^2 g''(t) \right) = +\infty.$$

To properly handle the minor and major arc conditions in terms of g, we introduce

(5.3)
$$\omega_g(t) \triangleq \frac{1}{6} (b_1 t + 8b_2 t^2 + \frac{9}{2} t^3 g'''(t)), \quad \text{for } t \in (0, R)$$

The following theorem is Theorem 4.1 in [3].

Theorem G. Let g be a nonconstant power series with radius of convergence R and nonnegative coefficients.

If the variance condition

(5.4)
$$\lim_{t\uparrow R} \left(tg'(t) + t^2 g''(t) \right) = +\infty$$

is satisfied and there is a cut function $h: [0, R) \to (0, \pi)$ so that

(5.5)
$$\lim_{t \uparrow R} \omega_g(t) h(t)^3 = 0.$$

and

(5.6)
$$\lim_{t\uparrow R} \sigma(t) \exp\left(\sup_{h(t)\leq |\theta|\leq \pi} \Re g(te^{i\theta}) - g(t)\right) = 0$$

hold, then $f = e^g$ is in the Hayman class.

Condition (5.5) of Theorem G gives that the cut function h fulfills condition (4.11) on the major arc. Also, condition (5.6) of Theorem G, which involves h and g, implies that condition (4.12) on the minor arc is satisfied.

Condition (5.6) on the minor arc is the most delicate to check. In practice, it depends on properly bounding $\sup_{\omega \le |\theta| \le \pi} \Re g(te^{i\theta}) - g(t)$ for general ω , as it is exhibited in the following variant, actually a corollary, of Theorem G.

Theorem 5.3. Let g be a nonconstant power series with radius of convergence R and nonnegative coefficients, and let $h: [0, R) \to (0, \pi)$ be a cut function.

Assume that the variance condition (5.4) and the condition on the major arc (5.5) of Theorem G are satisfied.

Assume further that there are positive functions U, V defined in (t_0, R) , for some $t_0 \in (0, R)$, where U takes values in $(0, \pi]$ and V in $(0, \infty)$, and such that

(5.7)
$$\sup_{|\theta| \ge \omega} \left(\Re g(te^{i\theta}) - g(t) \right) \le -V(t) \,\omega^2 \,, \quad \text{for } \omega \le U(t) \text{ and } t \in (t_0, R) \,,$$

and

(5.8)
$$h(t) \le U(t), \text{ for } t \in (t_0, R) \text{ and } \lim_{t \uparrow R} \sigma(t) e^{-V(t)h(t)^2} = 0.$$

Then $f = e^g$ is in the Hayman class.

Proof. The pair of conditions (5.7) and (5.8) together imply condition (5.6). This is so since the first half of (5.8) allows us to take $\omega = h(t)$ in (5.7) to obtain

$$\exp\left(\sup_{|\theta| \ge h(t)} \Re g(te^{i\theta}) - g(t)\right) \le \exp\left(-V(t)h(t)^2\right), \quad \text{for } t \in (t_0, R),$$

and to then apply the second half of (5.8).

Notice further that if for a function V defined in (t_0, R) , for some $t_0 \in (0, R)$, and taking values in $(0, +\infty)$, we have that

(5.9)
$$\Re g(te^{i\theta}) - g(t) \le -V(t) \theta^2, \quad \text{for } |\theta| \le \pi \text{ and } t \in (t_0, R),$$

then we may set $U \equiv \pi$, and the cut h is just required to satisfy

(5.10)
$$\lim_{t \uparrow R} \sigma(t) e^{-V(t)h(t)^2} = 0.$$

Thus these two conditions (5.9) and (5.10) imply the conditions (5.7) and (5.8) of Theorem 5.3.

Next we will exhibit easily verifiable requirements on the coefficients of g to obtain functions V and U so that the conditions (5.7) and (5.8) hold, or, simply, a function V so that the conditions (5.9) and (5.10) are satisfied.

The announced requirements differ if the power series g is entire or has finite radius of convergence; we split the discussion accordingly.

5.1. Case g entire, $R = \infty$. Here, g is an entire power series with nonnegative coefficients.

Lemma 5.4. If an entire power series g with nonnegative coefficients satisfies, for some $\beta > 0$ and B > 0, that

(5.11)
$$\Re g(te^{i\theta}) - g(t) \le B(e^{\beta t \cos \theta} - e^{\beta t}), \quad \text{for } t > 0 \text{ and } |\theta| \le \pi,$$

then condition (5.9) is satisfied with $V(t) = Ce^{\beta t}$ for some constant C > 0 depending only on B and β , and condition (5.10) requires that $\lim_{t\to\infty} \sigma(t) \exp(-Ce^{\beta t}h(t)^2) = 0$. *Proof.* For t > 0 and $\beta > 0$, the convexity of the exponential function $x \mapsto e^{\beta tx}$ in the interval [-1, 1] gives

$$e^{\beta t} - e^{\beta t \cos \theta} \ge (1 - \cos \theta) \frac{e^{\beta t} - e^{-\beta t}}{2}, \quad \text{for } \theta \in [-\pi, \pi].$$

Thus, we have, for $t \ge 1$, that

$$e^{\beta t \cos \theta} - e^{\beta t} \le (\cos \theta - 1) \frac{e^{\beta t} - e^{-\beta t}}{2} \le e^{\beta t} \frac{1 - e^{-2\beta}}{2} (\cos \theta - 1) \le -\frac{1 - e^{-2\beta}}{\pi^2} e^{\beta t} \theta^2.$$

In the second inequality we have used that $t \ge 1$; the inequality $1 - \cos x \ge 2x^2/\pi^2$, valid for $x \in [-\pi, \pi]$, yields the last step.

Theorem 5.5. Let
$$g(z) = \sum_{n=0}^{\infty} b_n z^n$$
 satisfy
(5.12) $B \frac{\beta^n}{n!} \le b_n \le L \frac{\lambda^n}{n!}$ for $n \ge 1$

for some constants B, L > 0, and $\beta, \lambda \ge 0$ such that $2\lambda < 3\beta$.

Then g satisfies the requirements of Theorem 5.3 and therefore $f = e^g$ is in the Hayman class.

Proof. Condition (5.12) gives that the power series g has radius of convergence $R = +\infty$. Besides, condition (5.12) and Proposition 2.3 show that the variance $\sigma^2(t)$ satisfies that $t^2 e^{\beta t} = O(\sigma^2(t))$ and that $\sigma^2(t) = O(t^2 e^{\lambda t})$, and that ω_g satisfies that $\omega_g(t) = O(t^3 e^{\lambda t})$, as $t \to \infty$.

The variance condition (5.4) is thus obviously satisfied. The major arc condition (5.5) holds with the cut function $h(t) = e^{-\alpha t}$, where $\alpha \in (\lambda/3, \beta/2)$, since $3\alpha > \lambda$.

Also, for t > 0 and $|\theta| \le \pi$, we have that

$$\Re g(te^{i\theta}) - g(t) = \sum_{n=1}^{\infty} b_n t^n (\cos n\theta - 1) \le B \sum_{n=1}^{\infty} \frac{(\beta t)^n}{n!} (\cos n\theta - 1)$$
$$= B \left(\Re e^{\beta te^{i\theta}} - e^{\beta t} \right) \le B (|e^{\beta te^{i\theta}}| - e^{\beta t}) = B (e^{\beta t \cos \theta} - e^{\beta t}).$$

and, consequently, condition (5.11) of Lemma 5.4 is satisfied.

Condition (5.9) is satisfied with the choice $V(t) = Ce^{\beta t}$, t > 1, and (5.10) holds because $2\alpha < \beta$. As mentioned above, these two conditions imply the conditions (5.7) and (5.8) of Theorem 5.3, and thus we conclude that f is in the Hayman class.

5.2. Case g not entire, $R < \infty$. Here, we will repeatedly appeal to Proposition 2.1.

Lemma 5.6. If a power series g with nonnegative coefficients and radius of convergence R = 1 satisfies, for some $\beta > 0$ and B > 0, that

(5.13)
$$\Re g(te^{i\theta}) - g(t) \le B\left(\frac{1}{|1 - te^{i\theta}|^{\beta}} - \frac{1}{(1 - t)^{\beta}}\right), \text{ for } t \in (0, 1) \text{ and } |\theta| \le \pi,$$

then

 $\sup_{|\theta| \ge \omega} \Re g(te^{i\theta}) - g(t) \le -C \, \frac{1}{(1-t)^{2+\beta}} \, \omega^2 \,, \quad for \ t \in (1/2, 1) \ and \ 0 \le \omega \le D(1-t) \,,$

where C > 0 and D > 0 depend only on β and B.

And, in particular, if we set $V(t) = C/(1-t)^{2+\beta}$, and U(t) = D(1-t), for $t \in (1/2, 1)$, then condition (5.7) is satisfied and condition (5.8) requires that

$$\lim_{t \to \infty} \sigma(t) \exp\left(-Ch(t)^2/(1-t)^{2+\beta}\right) = 0.$$

Proof. Let $\beta > 0$. For $t \in (0, 1)$ and $|\theta| \leq \pi$, we have that

$$\Re g(te^{i\theta}) - g(t) \le \frac{B}{(1-t)^{\beta}} \left(\left(\left| \frac{1-t}{1-te^{i\theta}} \right|^2 \right)^{\beta/2} - 1 \right).$$

Thus, for $t \in (1/2, 1)$ and $0 \le \omega < (1 - t)$ we have that

$$\sup_{|\theta| \ge \omega} \Re g(te^{i\theta}) - g(t) \le \frac{B}{(1-t)^{\beta}} \Big(\Big(1 - C\frac{1}{(1-t)^2} \,\omega^2 \Big)^{\beta/2} - 1 \Big) \,,$$

and, for $\omega < D_{\beta}(1-t)$, for $D_{\beta} > 0$ appropriately small, and some constant $C_{\beta} > 0$, we have that

$$\sup_{|\theta| \ge \omega} \Re g(te^{i\theta}) - g(t) \le -\frac{B}{(1-t)^{\beta}} C_{\beta} \frac{1}{(1-t)^2} \omega^2.$$

Theorem 5.7. Let $g(z) = \sum_{n=0}^{\infty} b_n z^n$ satisfy

(5.14)
$$B \frac{n^{\beta}}{R^{n}} \le b_{n} \le L \frac{n^{\lambda}}{R^{n}}, \quad for \ n \ge 1,$$

for some constants B, L > 0, finite radius R > 0 and $\beta, \lambda > -1$ such that $2\lambda < 3\beta + 1$.

Then g satisfies the requirements of Theorem 5.3 and therefore $f = e^g$ is in the Hayman class.

Proof. Because of (5.14), the power series g has radius of convergence R. By considering g(Rz), we may assume that R = 1. Appealing to Proposition 2.1, we see that the variance function of $f = e^g$ satisfies that $1/(1-t)^{3+\beta} = O(\sigma^2(t))$ and that $\sigma^2(t) = O(1/(1-t)^{3+\lambda})$, and that ω_g satisfies that $\omega_g(t) = O(1/(1-t)^{4+\lambda})$ as $t \uparrow 1$.

The variance condition (5.4) (or directly (4.13)) is obviously satisfied.

We propose a cut $h(t) = (1 - t)^{\alpha}$, where $\alpha \in (\lambda/3 + 4/3, \beta/2 + 3/2)$. The major arc condition (5.5) holds since $3\alpha > \lambda + 4$.

For $t \in (0, 1)$ and $|\theta| \leq \pi$, we have that

$$\begin{aligned} \Re g(te^{i\theta}) - g(t) &= \sum_{n=1}^{\infty} b_n t^n (\cos n\theta - 1) \le B \sum_{n=1}^{\infty} n^{\beta} t^n (\cos n\theta - 1) \\ &\le B_{\beta} \sum_{n=1}^{\infty} \frac{\Gamma(n+\beta+1)}{\Gamma(\beta+1)n!} t^n (\cos n\theta - 1) = B_{\beta} \Big(\Re \frac{1}{(1-te^{i\theta})^{\beta+1}} - \frac{1}{(1-t)^{\beta+1}} \Big) \\ &\le B_{\beta} \Big(\Big| \frac{1}{1-te^{i\theta}} \Big|^{\beta+1} - \Big(\frac{1}{1-t} \Big)^{\beta+1} \Big) ,\end{aligned}$$

where we have appealed to the comparison (2.2) within Proposition 2.1. Consequently, condition (5.13) of Lemma 5.6 does hold.

If we set $V(t) = C/(1-t)^{\beta+3}$, for $t \in (1/2, 1)$, (5.7) is satisfied and (5.8) requires that $\lim_{t\to\infty} \sigma(t) \exp(-Ch(t)^2/(1-t)^{\beta+3}) = 0$, which does hold since $2\alpha < \beta + 3$. Thus the conclusion follows from Theorem 5.3

Thus the conclusion follows from Theorem 5.3.

5.3. Applications of Theorems 5.5 and 5.7.

5.3.1. Sets of labeled classes.

• The egf e^{e^z-1} of the Bell numbers, enumerating sets of sets, is $f = e^g$ with $b_n = 1/n!$, for $n \ge 1$.

The egf e^{ze^z} of sets of pointed sets, is $f = e^g$ with $b_n = n/n!$, for $n \ge 1$.

Both satisfy the hypothesis of Theorem 5.5 and, thus, in particular they are in the Hayman class.

• The power series $f(z) = e^{z/(1-z)}$, egf of sets of lists, has g(z) = z/(1-z) with $b_n = 1$, for $n \ge 1$. The coefficients of the function g satisfy the hypothesis of Theorem 5.7 and, thus, in particular, $f = e^g$ is in the Hayman class.

In general, $f(z) = e^{z/(1-z)^{\gamma}}$ is in the Hayman class for $\gamma > 0$.

But, $f(z) = e^{\ln 1/(1-z)} = 1/(1-z)$, the egf of sets of cycles is not even Gaussian, as shown in Section 4.1.

• The eqf f of sets of functions is $f = e^g$ with $g(z) = \sum_{n=1}^{\infty} (n^n/n!) z^n$. In this case, the coefficients b_n of g satisfy (5.14) with R = 1/e and $\beta = \lambda = -1/2$, and therefore f is in the Hayman class.

In general, the exponential of $\sum_{n=1}^{\infty} (n^{n-\alpha}/n!) z^n$ is in the Hayman class if $0 \le \alpha < 1/2$. As we have seen in Section 4.1, for $\alpha = 1/2$ it is not strongly Gaussian, and for $\alpha > 1/2$ is not even Gaussian.

5.3.2. Sets of unlabeled classes.

• For the ogf P of partitions, we have that $P = e^g$, with

$$g(z) = \sum_{n=1}^{\infty} \frac{\sigma_1(n)}{n} \, z^n \,,$$

where $\sigma_1(n)$ denotes the sum of divisors of the positive integer n. The coefficients $b_n = \sigma_1(n)/n$ satisfy, in this case,

$$1 \leq b_n \leq D_{\varepsilon} n^{\varepsilon}$$
,

for each $\varepsilon > 0$ and some constant $D_{\varepsilon} > 0$. See Theorem 322 in [10]. (Actually, we may bound from above with $\ln \ln n$, see Theorem 323 in [10]). Therefore the function gsatisfies the hypothesis of Theorem 5.7 and this means, in particular, that $P = e^g$ is in the Hayman class.

The same argument gives that the infinite product $\prod_{j=1}^{\infty} 1/(1-z^j)^{c_j}$, where the c_j are integers satisfying $1 \le c_j \le c$, for some constant c > 1, is in the Hayman class.

• For the ogf Q of partitions into distinct parts, given by

$$Q(z) \stackrel{(1)}{=} \prod_{j=1}^{\infty} (1+z^j), \quad \text{or alternatively, by} \quad Q(z) \stackrel{(2)}{=} \prod_{j=0}^{\infty} \frac{1}{1-z^{2j+1}},$$

we have that $Q = e^g$, where

$$g(z) \stackrel{\text{by (1)}}{=} \sum_{k,j \ge 1} \frac{(-1)^{k+1}}{k} z^{kj} \stackrel{\text{by (2)}}{=} \sum_{k \ge 1; j \ge 0} \frac{1}{k} z^{k(2j+1)} = \sum_{n=1}^{\infty} \frac{\sigma_1^{\text{odd}}(n)}{n} z^n \,,$$

where $\sigma_1^{\text{odd}}(n)$ registers the sum of the odd divisors of n.

The coefficients $b_n = \sigma_1^{\text{odd}}(n)/n$ satisfy

(5.15)
$$\frac{1}{n} \le b_n \le D_{\varepsilon} n^{\varepsilon} \,,$$

for each $\varepsilon > 0$ and some constant $D_{\varepsilon} > 0$. The inequality on the left holds simply because 1 | n, while the inequality on the right holds because $\sigma_1^{\text{odd}}(n) \leq \sigma_1(n)$ and Theorem 322 in [10].

These bounds, though, are not within reach of Theorem 5.7, and for the function Q we will verify directly the hypothesis of Theorem 5.3.

Using (5.15), Proposition 2.1 and formula (5.2), we have for the variance function $\sigma^2(t)$ of $f = e^g$ that

$$\frac{1}{(1-t)^2} = O(\sigma^2(t)) \quad \text{and} \quad \sigma^2(t) = O\left(\frac{1}{(1-t)^{3+\varepsilon}}\right), \quad \text{ as } t \uparrow 1,$$

and for the function $\omega_g(t)$ that $\omega_g(t) = O(1/(1-t)^{4+\varepsilon})$, as $t \uparrow 1$.

The variance condition (5.4) (or directly (4.13)) is obviously satisfied. We propose a cut $h(t) = (1-t)^{\alpha}$, where $\alpha \in (4/3, 3/2)$.

The major arc condition (5.5) holds since $3\alpha > 4 + \varepsilon$, for appropriate small ε . For $t \in (0, 1)$ and $|\theta| \leq \pi$, and $z = te^{i\theta}$, we have that

$$\left|\frac{1+te^{i\theta}}{1+t}\right|^2 = 1 + \frac{2t(\cos\theta - 1)}{(1+t)^2} \le 1 + \frac{t}{2}(\cos\theta - 1) \le e^{(t/2(\cos\theta - 1))} = e^{(\Re z - |z|)/2},$$

and, so,

$$\frac{|Q(z)|}{Q(|z|)} \le \exp\left(\frac{1}{4}\left(\Re\frac{z}{1-z} - \frac{|z|}{1-|z|}\right)\right) = \exp\left(\frac{1}{4}\left(\Re\frac{1}{1-z} - \frac{1}{1-|z|}\right)\right),$$

and, consequently,

$$\Re g(z) - g(|z|) = \ln \frac{|Q(z)|}{Q(|z|)} \le \frac{1}{4} \left(\Re \frac{1}{1-z} - \frac{1}{1-|z|} \right) \le \left| \frac{1}{1-z} \right| - \frac{1}{1-|z|}.$$

Lemma 5.6 gives us $V(t) = C/(1-t)^3$ and U(t) = D(1-t). Now, condition (5.8) is satisfied since $\alpha < 3/2$.

Thus, the function g satisfies the conditions of Theorem 5.3, and, in particular, $Q = e^g$ is in the Hayman class.

• For the ogf M of plane partitions (see [7], p. 580),

$$M(z) = \prod_{j=1}^{\infty} \frac{1}{(1-z^j)^j} \,,$$

we have that $M = e^g$, with g given by

$$g(z) = \sum_{n=1}^{\infty} \frac{\sigma_2(n)}{n} z^n \,,$$

where $\sigma_2(n)$ denotes the sum of the squares of the divisors of the integer $n \ge 1$. For each $\varepsilon > 0$, there is a constant $C_{\varepsilon} > 0$ such that

$$n \le \frac{\sigma_2(n)}{n} \le C_{\varepsilon} n^{1+\varepsilon}$$
, for each $n \ge 1$.

This follows since $\sigma_2(n) \leq n\sigma_1(n)$ and $\sigma_1(n) \leq C_{\varepsilon}n^{1+\varepsilon}$. Thus we see that g satisfies the conditions of Theorem 5.7, and, in particular, we obtain that $M = e^g$ is in the Hayman class.

Likewise, and more generally, we see that for integer $c \ge 0$, the ogf of colored partitions,

$$\prod_{j=1}^{\infty} \frac{1}{(1-z^j)^{j^c}},$$

where each part j appears in j^c different colors, is in the Hayman class. Observe that $n^c \leq \sigma_{c+1}(n)/n \leq C_{\varepsilon} n^{c+\varepsilon}$, for $n \geq 1$.

Question 2. For the number $p_s(n)$ of partitions into squares (i.e., partitions whose parts are whole squares) we have that

$$S(z) \triangleq \prod_{j=1}^{\infty} \frac{1}{1-z^{j^2}} = \sum_{n=0}^{\infty} p_s(n) \, z^n \, .$$

Is S Gaussian or strongly Gaussian? See [8] and [16]. Theorem 5.7 is not applicable: the corresponding g is

$$g(z) = \sum_{m=1}^{\infty} \frac{1}{m} \left(\sum_{j^2 \mid m} j^2 \right) z^m ,$$

and $\sum_{j^2|m} j^2 = 1$ for any *m* which is a product of distinct primes.

Same question for

$$\prod_{j=1}^{\infty} \frac{1}{1-z^{2^j}},$$

whose coefficients count the number of partitions whose parts are powers of 2. See [5]. Theorem 5.7 is not applicable, either.

Question 3. The infinite product $\prod_{j=1}^{\infty} (1+z^j)^{j^c}$ has nonnegative coefficients. Is it Hayman?

6. On asymptotic formulas of coefficients

Once you know that a power series f is in the Hayman class, and thus that f is strongly Gaussian, you may use the asymptotic formula of Hayman of Theorem D, or even better, the asymptotic formula of Báez-Duarte of Theorem E, to yield an asymptotic formula for the coefficients of f.

As a token of the general approach, consider the partition function P. Hayman's formula gives us that

(6.1)
$$p(n) \sim \frac{1}{\sqrt{2\pi}} \frac{P(t_n)}{t_n^n \sigma(t_n)}, \quad \text{as } n \to \infty,$$

where m(t) and $\sigma^2(t)$ are the mean and variance functions of P and t_n is such that $m(t_n) = n$.

As such, this formula (6.1) is too implicit, of course, on three counts: t_n is not explicit, and hardly ever is, and, besides, the formula involves $\sigma(t_n)$ and $P(t_n)$.

We proceed by appealing to the Báez-Duarte formula (4.10), and by obtaining, on the one hand, asymptotic formulas for τ_n and $\tilde{\sigma}(\tau_n)$, and for $P(\tau_n)$ on the other hand, as follows (see the details in Section 6.3.1 of [3]).

Concerning τ_n and $\sigma(\tau_n)$. By means of Euler's summation, we may approximate the mean function m(t) by

$$m(e^{-s}) \sim \frac{\zeta(2)}{s^2} \triangleq \widetilde{m}(e^{-s}), \text{ as } s \downarrow 0,$$

and $\sigma^2(t)$ by

$$\sigma^2(e^{-s}) \sim \frac{2\zeta(2)}{s^3} \triangleq \widetilde{\sigma}^2(e^{-s}), \quad \text{as } s \downarrow 0,$$

while checking that condition (4.9) is satisfied.

Now if τ_n is given by $\tau_n = e^{-s_n}$ and $s_n = \sqrt{\zeta(2)/n}$ so that $\widetilde{m}(\tau_n)$, we may appeal to the formula of Báez-Duarte (4.10) and write

(6.2)
$$p(n) \sim \frac{1}{\sqrt{2\pi}} \frac{P(\tau_n)}{\tau_n^n \,\widetilde{\sigma}(\tau_n)}, \quad \text{as } n \to \infty.$$

Concerning $P(\tau_n)$. Finally, Euler's summation again gives that

$$P(e^{-s}) \sim \frac{1}{\sqrt{2\pi}} \sqrt{s} e^{\zeta(2)/s}, \quad \text{as } s \downarrow 0,$$

and thus

$$P(e^{-s_n}) \sim \frac{1}{\sqrt{2\pi}} \sqrt{s_n} e^{\zeta(2)/s_n}, \quad \text{as } n \to \infty.$$

Upon substitution in (6.2), the Hardy–Ramanujan asymptotic formula for partitions follows:

$$p(n) \sim \frac{1}{4\sqrt{3}} \frac{1}{n} e^{2\sqrt{\zeta(2)}\sqrt{n}}$$
, as $n \to \infty$.

In short, this is the basic procedure for obtaining asymptotic formulas for coefficients of power series in the Hayman class. For a number of detailed examples, including the partition function above, general set constructions, and a few others, we refer the reader again to [3].

Funding information. Research of J. L. Fernández and P. Fernández supported by Fundación Akusmatika. Research of V. Maciá was partially funded by grant MTM2017-85934-C3-2-P2 of Ministerio de Economía y Competitividad of Spain and European Research Council Advanced Grant 834728.

References

- [1] ARRIATA, R., BARBOUR, A. D. AND TAVARÉ, S.: Logarithmic combinatorial structures: a probabilistic approach. European Mathematical Society, 2003.
- [2] BÁEZ-DUARTE, L.: Hardy-Ramanujan's asymptotic formula for partitions and the central limit theorem. Adv. Math. 125 (1997), no. 1, 114–120.
- [3] CANTÓN, A., FERNÁNDEZ, J. L., FERNÁNDEZ, P. AND MACIÁ, V.: Khinchin families and Hayman class. Comput. Methods Funct. Theory 21 (2021), no. 4, 851–904.
- [4] CANTÓN, A., FERNÁNDEZ, J. L., FERNÁNDEZ, P. AND MACIÁ, V.: A note on Khinchin clans. In preparation, 2022.
- [5] DE BRUIJN, N. G.: On Mahler's partition problem. Indag. Math. 10 (1948), 210–220.
- [6] FERNÁNDEZ, J.L. AND MACIÁ, V.: Khinchin families and large powers. Preprint 2022, arXiv: 2201.11746.
- [7] FLAJOLET, P. AND SEDGEWICK, R.: Analytic combinatorics. Cambridge University Press, 2009.
- [8] GAFNI, A.: Power partitions. J. Number Theory 163 (2016), 19–42.
- [9] GLAISHER, J. W. L.: On certain sums of products of quantities depending upon the divisors of a number. *Mess. Math.*, 15 (1885), 1–20.
- [10] HARDY, G. H. AND WRIGHT, E. M.: An introduction to the theory of numbers. Fifth edition. Oxford University Press, 1978.
- [11] HARRIS, B. AND SCHOENFELD, L.: Asymptotic expansions for the coefficients of analytic functions. *Illinois J. Math.* 12 (1968), 264–277.
- [12] HAYMAN, W. K.: A generalisation of Stirling's formula. J. Reine Angew. Math. 196 (1956), 67–95.
- [13] JONES, C. M. AND ZHIGLJAVSKY, A. A.: Approximating the negative moments of the Poisson distribution. *Statist. Probab. Lett.* 66 (2004), no. 2, 171–181.
- [14] ODLYZKO, A. M. AND RICHMOND, L. B.: Asymptotic expansions for the coefficients of analytic generating functions. *Aequationes Math.* 28 (1985), no. 1-2, 50–63.

- [15] ROSENBLOOM, P. C.: Probability and entire functions. In Studies in mathematical analysis and related topics, 325–332. Stanford Univ. Press, Stanford, Calif., 1962.
- [16] VAUGHAN, R. C.: Squares: Additive questions and partitions. Int. J. Number Theory 11 (2015), no. 5, 1367–1409.