

Hybrid k_T -factorization and impact factors at NLO

Andreas van Hameren^{a1}, Leszek Motyka^{b2},
and Grzegorz Ziarko^{a3}

^a *Institute of Nuclear Physics Polish Academy of Sciences,
PL-31342 Kraków, Poland*

^b *Institute of Theoretical Physics, Jagiellonian University,
S. Łojasiewicza 11, 30-348 Kraków, Poland*

June 6, 2022

Abstract

In the hybrid k_T -factorization formula, one initial-state parton momentum is space-like and carries non-vanishing transverse components, while the other is on-shell. We promote this factorization formula to next-to-leading order. Studying the partonic cross section, we identify all soft and collinear divergencies in the real and virtual contribution, and recognize that all non-cancelling ones can be attributed to PDF evolution, evolution kernel, and target impact factors. In result, we construct a framework that may be used to compute NLO impact factors in general. In particular, we recover known expressions for inclusive NLO quark-and gluon impact factor corrections.

¹hameren@ifj.edu.pl

²leszek.motyka@uj.edu.pl

³grzegorz.ziarko@ifj.edu.pl

Contents

1	Introduction	2
2	Main result	4
3	Unfamiliar real contribution	14
4	Unfamiliar virtual contribution	21
5	Familiar virtual contribution	28
6	Familiar real contribution	30
7	Summary	34
A	Spinors and transverse momenta	43
B	Color representation	44
C	Collinear splitting functions	46
D	The triple auxiliary parton limit	49
E	Some integrals	53

1 Introduction

The high energy factorization approach to scattering amplitudes has played a seminal role in developments of theory and phenomenology of strong interactions since quantum chromodynamics was discovered [1–6], [7–9], [10–14, 14–25]. The cornerstone of this factorization, closely related to an even earlier concept of Regge factorization, may be traced back to the Lorentz dilation of time between the projectile and target frames in ultrarelativistic particle collisions. Due to this effect, the dynamics in the hadronic projectile and in the hadronic target is governed by vastly different time scales and the dynamics approximately factorizes. From theory side, the high energy factorization approach is based on isolating the leading powers of the total hadron–hadron collision energy squared, S , in the scattering amplitudes. Then, in the perturbative approach to the high energy scattering amplitudes one finds perturbative corrections that are enhanced by powers of logarithms of the energy, L_S^n , where $L_S = \ln(S/S_0)$, with S_0 being a fixed and moderate scale, and the total collision energy squared $S \gg S_0$ can be arbitrarily large. The formal all order resummations of these enhanced perturbative corrections are realized within the BFKL scheme [1–6], [10–14, 14–25]. In general, the Regge factorization principle allows to factorize

the dynamics in the projectile A phase space region, in the target B region and in the intermediate (exchange) region. The scattering amplitude \mathcal{A} may be written symbolically as

$$\mathcal{A} = iS\Phi_A \otimes \mathcal{G}(S) \otimes \Phi_B,$$

where Φ_A and Φ_B are process dependent impact factors, $\mathcal{G}(S)$ is an universal exchange part, the convolutions denoted by \otimes are performed over transverse momenta, and the other details were omitted as we focus only on the structural features here.

In QCD the BFKL resummation framework was constructed for the gluonic exchange $\mathcal{G}(S)$ at the leading logarithmic (LL) and the next-to-leading logarithmic accuracy (NLL), where terms $(\alpha_s L_S)^n$ and $\alpha_s^{n+1} L_S^n$ are resummed correspondingly. In recent years within $\mathcal{N} = 4$ supersymmetric Young-Mills theories this approach was further developed to reach the next-to-next-to-leading logarithmic accuracy [26–28]. In order to fully use the potential of the BFKL approach in phenomenological applications at the presently highest available NLL accuracy, it is necessary to include the impact factors $\Phi_{A,B}$ at the NLO accuracy. Such computations are highly non-trivial in the high energy limit. To this date they have been performed for about a dozen of processes yielding interesting predictions mostly for high energy hadron colliders, like the Large Hadron Collider, RHIC and Tevatron, and for electron–proton colliders: HERA and the Electron–Ion Collider.

The calculations of the impact factors at the NLO started from inclusive quark and gluon impact factors [29–31], then the forward jet impact factors were calculated [32–37]. Based on these results applications of the NLL BFKL to collider data was developed, to start with numerous detailed analyses of forward jet cross-sections in hadron colliders [38–44]. Following those pioneering studies the formalism was applied to other high energy scattering and particle production processes. In particular: for exclusive vector meson photoproduction [45–48], inclusive and diffractive dijets [49–51] and multi-particle final states [52, 53]. The inclusive γ^* impact factor was calculated as well [54–57], that allowed to analyze the DIS processes [58, 59] and total $\gamma^*\gamma^*$ cross-sections [60]. Other examples are the forward hadroproduction of heavy quarks [61–63], dihadrons [64, 65], the Higgs boson [66–68] and heavy baryons [69]. So far the calculations of the NLO impact factors were performed process–by–process in an explicit form. In the present letter we propose a computational framework that allows to obtain in a straightforward way the NLO impact factors from NLO collinear scattering amplitudes.

Complementary to the quest for higher logarithmic accuracy in BFKL resummation and higher perturbative precision for impact factors is the application high energy factorization to define and calculate scattering processes involving space-like initial-state gluons and several final-state jet and/or particles. The idea is to have a factorized formula with parton density functions (PDFs) and a parton level cross section, both however depending on transverse momentum components (the k_T) of the initial-state gluons besides the components parallel to the initial-state hadron momenta. Basically, the formula for heavy quark production established in [8, 70] is generalized by replacing the partonic cross section by the one for other processes, involving jets, massive vector bosons, etc. The calculations are at tree-level, for which the main hurdle to take is the gauge invariant definition of tree-level amplitudes with space-like initial state partons, which

can be achieved with the help of effective vertices from Lipatov’s effective action [71, 72], or with the auxiliary parton method [73]. Originally, the PDFs were imagined to undergo BFKL evolution, but the factorized formula has also been applied with k_T -dependent PDFs obtained from “unintegration procedures” [74, 75], and from non-linear evolution [76, 77].

Recently, calculations within this kind of factorization have been performed at next-to-leading order (NLO) for processes with one particle in the final state [67, 78, 79]. Moving to NLO inevitably leads to infrared divergences in real and virtual contributions which need to cancel or treated within a renormalization procedure. In collinear factorization this is rigorously established partly through renormalization of the PDFs. In this paper we show how the factorization formula with k_T dependence can be promoted to NLO within the auxiliary parton method.

More specifically, we consider the so-called hybrid factorization formula, for which one initial-state parton is light-like, and its associated PDF only depends only on the longitudinal momentum fraction, while the other is space-like and its associated PDF also depends on k_T . We will identify the divergences in real and virtual contributions that do not manifestly cancel, and we will show that they are process independent. With the process we mean here the one with the initial-state space-like gluon. The divergences are *not* independent of the auxiliary partons, which may a scattering gluon or quark. We will, however, show that the difference between the choice of auxiliary partons is process independent.

In a heuristic picture, the auxiliary parton plays a role of a target, with the dynamics that is Regge factorized from the dynamics of the transition in the projectile region. Such a picture is more or less explicitly present in most of the LO calculations of the impact factors, and at the LO it is straightforward and simple. A generalization of this method to the NLO amplitudes is more demanding. We will see, however, that our result fits the interpretation in which the NLO auxiliary parton scattering amplitude and the LL exchange (related to the BFKL kernel) amplitude taken at $\mathcal{O}(\alpha_s)$ are factorized from the suitable NLO on-shell amplitude, to result with the explicit expression for the NLO impact factor. Again, it is important to stress that both the NLO auxiliary parton impact factors and the $\mathcal{O}(\alpha_s)$ LL exchange amplitude are universal. Hence, the explicit expressions for these contributions may be used to obtain the NLO impact factor for any process in the projectile region. This method, including the explicit form of the quark and gluon NLO impact factors, is the main result of this paper.

2 Main result

The context of our work is the hybrid k_T -type factorized cross section formula for hadron scattering, which at lowest order schematically looks like

$$d\sigma^{(0)} = \int dx_{\text{in}} d^2k_{\perp} d\bar{x}_{\text{in}} F(x_{\text{in}}, |k_{\perp}|) f(\bar{x}_{\text{in}}) dB^*(x_{\text{in}}, k_{\perp}, \bar{x}_{\text{in}}) . \quad (1)$$

The functions $F(x_{\text{in}}, |k_{\perp}|)$ and $f(\bar{x}_{\text{in}})$ are parton density functions (PDFs), both depending on a longitudinal momentum fraction, and only one of them depending on transverse momentum. Of course both typically depend also on an additional scale and on the parton type. Our attention

however goes to the, here Born-level, differential partonic cross section dB^* , which is special compared to collinear factorization in the sense that one of its initial-state partons is space-like, and has momentum

$$\mathbf{k}_{\text{in}}^\mu = x_{\text{in}} P_A^\mu + \mathbf{k}_\perp^\mu, \quad (2)$$

where P_A is the light-like hadron momentum and \mathbf{k}_\perp is transverse, *i.e.* $\mathbf{k}_\perp \cdot P_A = 0$. The ‘ \star ’ in dB^* highlights the \mathbf{k}_\perp -dependence. At the lowest order it is straightforward to define and calculate dB^* , and essentially only involves tree-level amplitudes, and final-state momenta that are well-separated by a number of phase space cuts.

The NLO contribution looks, again rather schematically, like

$$\begin{aligned} d\sigma^{(1)} = \int dx_{\text{in}} d^2k_\perp d\bar{x}_{\text{in}} \left\{ F(x_{\text{in}}, |\mathbf{k}_\perp|) f(\bar{x}_{\text{in}}) \left[dV^*(x_{\text{in}}, \mathbf{k}_\perp, \bar{x}_{\text{in}}) + dR^*(x_{\text{in}}, \mathbf{k}_\perp, \bar{x}_{\text{in}}) \right] \right. \\ \left. + \left[F^{(1)}(x_{\text{in}}, |\mathbf{k}_\perp|) f(\bar{x}_{\text{in}}) + F(x_{\text{in}}, |\mathbf{k}_\perp|) f^{(1)}(\bar{x}_{\text{in}}) \right] \text{dB}^*(x_{\text{in}}, \mathbf{k}_\perp, \bar{x}_{\text{in}}) \right\}. \quad (3) \end{aligned}$$

Here, dV^* represents the virtual contribution involving one-loop amplitudes, and dR^* the real contribution involving extra phase space integrals. Both carry one more power of the strong coupling constant than dB^* , but are combined with the same PDFs as at the lowest order. In the second line of Eq. (3) the functions $f^{(1)}$, $F^{(1)}$ are higher order corrections, and carry an extra power of the coupling constant.

The loop integrals in dV^* and the phase space integrals in dR^* cause soft and collinear divergences. Within collinear factorization, these divergences cancel up to certain collinear divergences, which we denote Δ_{coll} and $\Delta_{\bar{\text{coll}}}$, that are associated with the variables x_{in} and \bar{x}_{in} . The well-known formula for $\Delta_{\bar{\text{coll}}}$ is given in Eq. (45). It is absorbed into $f^{(1)}(\bar{x}_{\text{in}})$, or at least the factorization prescription can be interpreted as such. The same happens with Δ_{coll} and the correction to the other collinear PDF that appears instead of $F(x_{\text{in}}, |\mathbf{k}_\perp|)$ in the collinear factorization formula. For the hybrid formula above, we again find $\Delta_{\bar{\text{coll}}}$, which can be treated the same way, and we find a different but similar Δ_{coll}^* (Eq. (47)), which now must be absorbed into $F^{(1)}(x_{\text{in}}, |\mathbf{k}_\perp|)$.

In the following, we will identify extra non-cancelling divergencies in dV^* and dR^* , which we call *unfamiliar* contributions Δ_{unf} (Eq. (38)). They break a notion of universality that exists at Born level. They are proportional to the LO cross section, and must be interpreted as corrections to the impact factor associated with the space-like gluon. So the structure we eventually find is

$$\begin{aligned} d\sigma^{(1)} = \int dx_{\text{in}} d^2k_\perp d\bar{x}_{\text{in}} \left\{ F(x_{\text{in}}, |\mathbf{k}_\perp|) f(\bar{x}_{\text{in}}) \left[dV^*(x_{\text{in}}, \mathbf{k}_\perp, \bar{x}_{\text{in}}) + dR^*(x_{\text{in}}, \mathbf{k}_\perp, \bar{x}_{\text{in}}) \right]_{\text{cancelling}} \right. \\ \left. + \left[F^{(1)}(x_{\text{in}}, |\mathbf{k}_\perp|) + F(x_{\text{in}}, |\mathbf{k}_\perp|) \Delta_{\text{unf}} + \Delta_{\text{coll}}^* \right] f(\bar{x}_{\text{in}}) \text{dB}^*(x_{\text{in}}, \mathbf{k}_\perp, \bar{x}_{\text{in}}) \right. \\ \left. + F(x_{\text{in}}, |\mathbf{k}_\perp|) \left[f^{(1)}(\bar{x}_{\text{in}}) + \Delta_{\bar{\text{coll}}} \right] \text{dB}^*(x_{\text{in}}, \mathbf{k}_\perp, \bar{x}_{\text{in}}) \right\}. \quad (4) \end{aligned}$$

The term $dR^*(x_{\text{in}}, \mathbf{k}_\perp, \bar{x}_{\text{in}})$ inside the cancelling contribution only involves tree-level amplitudes. It requires numerical phase space integration, and the IR divergencies can be dealt with using methods known in literature, *e.g.* the subtraction method of [80]. Only a special subtraction

term related to the space-like gluon is necessary, which will be addressed in this paper. The term $dV^*(x_{\text{in}}, k_{\perp}, \bar{x}_{\text{in}})$ requires one-loop amplitudes. For processes that are not too complicated, they can rather straightforwardly be obtained from analytic expressions for on-shell one-loop amplitudes with one more external leg, and in this paper we describe the terms appearing in this procedure that that must go to $\Delta_{\text{unf}} dB^*(x_{\text{in}}, k_{\perp}, \bar{x}_{\text{in}})$.

The procedure mentioned above is the *auxiliary parton method* which will be described in this section. For tree-level amplitudes, the necessity for analytic expressions of on-shell amplitudes can be avoided and the amplitudes with a space-like gluon can be constructed directly using eikonal Feynman rules [73]. In the case of one-loop amplitudes, an approach to avoid the necessity for analytic expressions of on-shell one-loop amplitudes is described in [81].

2.1 Notation

In order to proceed, we need to establish some notation. We introduce the decomposition of a general four vector K^{μ} as

$$K^{\mu} = \xi_K P^{\mu} + \bar{\xi}_K \bar{P}^{\mu} + K_{\perp}^{\mu} \quad (5)$$

where the momenta P and \bar{P} have positive energy and satisfy

$$P^2 = \bar{P}^2 = 0 \quad , \quad 2P \cdot \bar{P} = v^2 > 0 \quad , \quad P \cdot K_{\perp} = \bar{P} \cdot K_{\perp} = 0 \quad (6)$$

so

$$\xi_K = \frac{\bar{P} \cdot K}{P \cdot \bar{P}} \quad , \quad \bar{\xi}_K = \frac{P \cdot K}{P \cdot \bar{P}} \quad (7)$$

The momenta P and \bar{P} are the directions of the scattering hadron momenta, and we prefer to let them keep a scale rather than to normalize them. The coordinates ξ and $\bar{\xi}$ are dimensionless and strictly positive for positive-energy momenta. We need to distinguish between variables ξ and direction P versus variables x and momentum P_A satisfying

$$\xi_K P^{\mu} = x_K P_A^{\mu} \quad , \quad (8)$$

as will become clear later. For the $\bar{\xi}$ variables and \bar{P} there is no such issue, and \bar{P} can in fact be considered the hadron momentum itself and we use the variable

$$\bar{x}_K = \bar{\xi}_K \quad (9)$$

from now on.

The infinitesimal volume of a light-like momentum K in terms of the decomposition becomes

$$d^4K \delta(K^2) = d\xi_K d\bar{x}_K d^2K_{\perp} \frac{1}{2\xi_K} \delta\left(\bar{x}_K - \frac{|K_{\perp}|^2}{v^2 \xi_K}\right) \quad (10)$$

We will use the same notation for K_{\perp} as a two-dimensional Euclidean vector, and for its embedding in Minkowski space. We will use the absolute value notation to make clear that a square is

positive. We introduce a notation for the n -particle parton-level differential cross section including a tree-level on-shell squared matrix element and the flux factor

$$d\Sigma_n^*(k_{\text{in}}, k_{\bar{\text{in}}}; \{p_i\}_{i=1}^n) = \left(\prod_{i=1}^n d^4 p_i \delta(p_i^2 - m_i^2) \right) \delta^4 \left(k_{\text{in}} + k_{\bar{\text{in}}} - \sum_{i=1}^n p_i \right) \frac{|\overline{\mathcal{M}}^*|^2(k_{\text{in}}, k_{\bar{\text{in}}}; \{p_i\}_{i=1}^n)}{4\xi_{\text{in}}\bar{\chi}_{\bar{\text{in}}}\mathcal{P}\cdot\bar{\mathcal{P}}}. \quad (11)$$

Let us state immediately that in this write-up,

$$\text{the symbol } \mathcal{M} \text{ always refers to a } \textit{tree-level} \text{ amplitude.} \quad (12)$$

The arguments in Σ_n^* and $|\overline{\mathcal{M}}^*|^2$ are separated into initial-state and final-state ones by the semi-colon. The star highlights that there is a space-like initial state, so

$$k_{\text{in}}^\mu = \xi_{\text{in}}\mathcal{P}^\mu + k_\perp^\mu, \quad k_{\bar{\text{in}}}^\mu = \bar{\chi}_{\bar{\text{in}}}\bar{\mathcal{P}}^\mu. \quad (13)$$

If $|k_\perp| = 0$ and both initial states are light-like, then the star is omitted. The squared matrix element $|\overline{\mathcal{M}}^*|^2$ is assumed to be summed over final-state colors and spins, and averaged over initial-state colors and spins. One might say that the subscript n in Σ_n^* is superfluous because its arguments are listed explicitly, just like in $|\overline{\mathcal{M}}^*|^2$, but we keep it for later convenience.

In order to define a differential Born-level cross section, we also need a ‘‘jet function’’ that guarantees that final-state partons cannot become arbitrarily soft or arbitrarily collinear to each other or to the initial-state momenta if this causes a singularity. Realize that this includes the case when a final-state momentum becomes collinear to \mathcal{P} . Also squared matrix elements with a space-like initial-state can have such a collinear singularity. We write

$$dB^*(k_{\text{in}}, k_{\bar{\text{in}}}; \{p_i\}_{i=1}^n) = \frac{1}{|k_\perp|^2} d\Sigma_n^*(k_{\text{in}}, k_{\bar{\text{in}}}; \{p_i\}_{i=1}^n) J_B(\{p_i\}_{i=1}^n). \quad (14)$$

Whereas the ‘‘space-like gluon propagator’’ was taken out in the definition of the squared matrix element to have a smooth on-shell limit, it appears to be most convenient to include the factor $1/|k_\perp|^2$ in the definition of the partonic cross section. We include possible non-parton momenta in the argument list of the jet function $J_B(\{p_i\}_{i=1}^n)$ and simply assumes that it selects only parton momenta.

In order to obtain the real contribution at NLO, the partonic scattering process is augmented with one more final-state parton, and the jet function is advanced in that it incorporates an at most one-step recombination of parton momenta, so it produces both the resolved situation with one more jet than at Born level, and the unresolved situation with the same number of jets. We denote it

$$J_R(\{p_i\}_{i=1}^{n+1}). \quad (15)$$

We deliberately equip the jet functions with the labels B for ‘‘Born’’ and R for ‘‘real’’ rather than n and $n+1$, to highlight that J_R is not merely a generalization of J_B regarding phase space cuts. We mention explicitly that also in the space-like initial-state kinematics, the jet function considers a

phase space configuration unresolved if a final-state momentum becomes collinear to P . Given this jet function, we define what we call the *familiar real contribution* as

$$dR^{\text{fam}}(k_{\text{in}}, k_{\overline{\text{in}}}; \{p_i\}_{i=1}^{n+1}) = \frac{\alpha_\epsilon \mu^{2\epsilon}}{\pi_\epsilon} \frac{1}{|k_\perp|^2} d\Sigma_{n+1}^*(k_{\text{in}}, k_{\overline{\text{in}}}; \{p_i\}_{i=1}^{n+1}) J_R(\{p_i\}_{i=1}^{n+1}). \quad (16)$$

This is what one naively would write down as the real radiation contribution given the Born formula. Of course, actual integration of this differential cross section leads to soft and collinear divergencies, but these can be dealt with using existing methods that can be found in literature of NLO calculations. Only the collinear singularity associated with the space-like initial-state is different than usual, and we will return to this later.

The overall constant in Eq. (16) gives the correction factor compared to Born level considering that we are using dimensional regularization, we assume $g_s = 1$ inside the squared matrix element, and considering the fact that we omitted factors of 2π in our definition of the phase space in Eq. (11) compared to usual definitions found in literature. The symbols in Eq. (16) represent

$$\pi_\epsilon = \frac{\pi^{1-\epsilon}}{\Gamma(1-\epsilon)} \quad , \quad \alpha_\epsilon = \frac{\alpha_s}{2\pi} \frac{(4\pi)^\epsilon}{\Gamma(1-\epsilon)} \quad , \quad \epsilon = \frac{4 - \text{dim}}{2}. \quad (17)$$

In other words, both the Born and the real, and later the virtual, contribution are missing an overall factor $(2\pi)^{4-3n} g_s^{2n} = (2\pi)^{4-3n} (4\pi\alpha_s)^n$.

2.2 The auxiliary parton method

As mentioned in the introduction, the tree-level squared matrix element with the space-like initial-state gluon can be defined with the auxiliary parton method proposed in [73] and most recently applied in [82]. Let the desired parton-level process be

$$g^*(k_{\text{in}}) \omega_{\overline{\text{in}}}(k_{\overline{\text{in}}}) \rightarrow \omega_1(p_1) \omega_2(p_2) \cdots \omega_n(p_n) \quad (18)$$

where the ω_i are the type of partons or particles involved besides the space-like gluon. In the auxiliary parton method, this process is obtained from the process

$$q(k_1(\Lambda)) \omega_{\overline{\text{in}}}(k_{\overline{\text{in}}}) \rightarrow q(k_2(\Lambda)) \omega_1(p_1) \omega_2(p_2) \cdots \omega_n(p_n) \quad (19)$$

where the momenta $k_1(\Lambda)$ and $k_2(\Lambda)$ are parametrized with Λ in such a way that $k_1^2 = k_2^2 = 0$ for any value of Λ , and $k_1 - k_2 = k_{\text{in}} + \mathcal{O}(\Lambda^{-1})$. In fact, in publications so far the parametrization was such that even the $\mathcal{O}(\Lambda^{-1})$ is absent, but here we will allow this irrelevant term, which allows us to use the parametrization

$$k_1^\mu = \Lambda P^\mu \quad , \quad k_2^\mu = p_\Lambda^\mu = (\Lambda - \xi_{\text{in}}) P^\mu - k_\perp^\mu + \frac{|k_\perp|^2}{(\Lambda - \xi_{\text{in}}) \nu^2} \bar{P}^\mu. \quad (20)$$

The process of Eq. (19) with the auxiliary quarks is on-shell, and its squared matrix element is well-defined and it is known how to calculate it. The squared matrix element of the desired process with the space-like gluon is obtained by taking

$$\Lambda \rightarrow \infty. \quad (21)$$

In [81] and [82] it was noted that instead of an auxiliary scattering quark, also an auxiliary scattering gluon can be used. At the level of squared matrix elements summed over color, one just needs to include a different overall factor to correct for the difference in color representation:

$$\frac{1}{g_s^2 C_{\text{aux}}} \frac{\xi_{\text{in}}^2 |\mathbf{k}_\perp|^2}{\Lambda^2} |\overline{\mathcal{M}}^{\text{aux}}|^2(\Lambda P, \mathbf{k}_{\overline{\text{in}}}; \mathbf{p}_\Lambda, \{\mathbf{p}_i\}_{i=1}^n) \xrightarrow{\Lambda \rightarrow \infty} |\overline{\mathcal{M}}^*|^2(\mathbf{k}_{\text{in}}, \mathbf{k}_{\overline{\text{in}}}; \{\mathbf{p}_i\}_{i=1}^n) \quad (22)$$

with

$$C_{\text{aux-q}} = \frac{N_c^2 - 1}{N_c}, \quad C_{\text{aux-g}} = 2N_c. \quad (23)$$

We call this *auxiliary parton universality* and we will see that it does not hold anymore at NLO. Another property that will appear not to hold anymore is the *smooth on-shell limit*

$$\lim_{z \rightarrow 0} \int \frac{d^2 \mathbf{k}_\perp}{\pi} \delta(|\mathbf{k}_\perp|^2 - z) |\overline{\mathcal{M}}^*|^2(\xi_{\text{in}} P + \mathbf{k}_\perp, \mathbf{k}_{\overline{\text{in}}}; \{\mathbf{p}_i\}_{i=1}^n) = |\overline{\mathcal{M}}|^2(\xi_{\text{in}} P, \mathbf{k}_{\text{in}}; \{\mathbf{p}_i\}_{i=1}^n), \quad (24)$$

where the right-hand-side is the squared matrix element for the process with an on-shell initial-state gluon instead of the space-like gluon.

2.3 Scaling behavior

As explicated in [83], rather than a function of ξ_{in} and \mathbf{k}_\perp , the space-like tree level amplitude must be seen as a function of \mathbf{k}_{in} and the light-like *direction* P with the single property that $\mathbf{k}_{\text{in}} \cdot P = 0$. Interpreted like that, the amplitude satisfies the scaling relation

$$\mathcal{A}_{\text{tree}}^*(\lambda P, \mathbf{k}_{\text{in}}, \mathbf{k}_{\overline{\text{in}}}; \{\mathbf{p}_i\}_{i=1}^n) = \lambda \mathcal{A}_{\text{tree}}^*(P, \mathbf{k}_{\text{in}}, \mathbf{k}_{\overline{\text{in}}}; \{\mathbf{p}_i\}_{i=1}^n). \quad (25)$$

The factor ξ_{in}^2 in Eq. (22) is included exactly in order to match the direction $P \rightarrow \xi_{\text{in}} P$ with the on-shell limit momentum. This could also have been achieved by defining the auxiliary momenta of Eq. (20) as $\Lambda \xi_{\text{in}} P$ and $(\Lambda - 1) \xi_{\text{in}} P - \dots$ etc., but that would be inconvenient for the following.

We see that the amplitude, and thus the squared matrix element, but in fact also dB^* do not depend on ξ_{in} or P separately, but only on the combination $\xi_{\text{in}} P$. We can scale

$$\xi_{\text{in}} \rightarrow x_{\text{in}} = \lambda^{-1} \xi_{\text{in}} \quad \text{and} \quad P \rightarrow P_\Lambda = \lambda P \quad (26)$$

without changing dB^* . This “scale” λ is set by introducing an integration over initial-state variables as

$$\int d^4 \mathbf{k}_{\text{in}} \mathcal{F}(\mathbf{k}_{\text{in}}) dB^*(\mathbf{k}_{\text{in}}, \mathbf{k}_{\overline{\text{in}}}; \{\mathbf{p}_i\}_{i=1}^n) \quad (27)$$

with

$$\mathcal{F}(\mathbf{k}_{\text{in}}) = \int \frac{d^2 \mathbf{k}_\perp}{\pi} \frac{dx_{\text{in}}}{x_{\text{in}}} F(x_{\text{in}}, |\mathbf{k}_\perp|) \delta^4(\mathbf{k}_{\text{in}} - x_{\text{in}} P_\Lambda - \mathbf{k}_\perp) \quad (28)$$

where $F(x_{\text{in}}, |\mathbf{k}_\perp|)$ is the \mathbf{k}_\perp -dependent PDF, or TMD. This introduces the physical hadron momentum and its fraction.

2.4 Unfamiliar contributions at NLO

Considering that we already introduced a real contribution we designated *familiar*, the reader may correctly expect that there is also an *unfamiliar* contribution. It occurs in the auxiliary parton method because there is a region in phase space where the radiative gluon should be allowed to participate in the ‘‘consumption’’ of the large P-component Λ when $\Lambda \rightarrow \infty$. We calculate the contribution in Section 3, and it turns out to violate the three tree-level properties of auxiliary parton universality, the smooth on-shell limit, and the smooth large Λ limit. The *unfamiliar real contribution* is proportional to the Born level result and given by

$$dR^{*\text{unf}}(k_{\text{in}}, k_{\bar{\text{in}}}; \{p_i\}_{i=1}^n) = \mathcal{R}_{\text{aux}} dB^*(k_{\text{in}}, k_{\bar{\text{in}}}; \{p_i\}_{i=1}^n), \quad (29)$$

with

$$\mathcal{R}_{\text{aux}} = \alpha_\epsilon N_c \left(\frac{\mu^2}{|k_\perp|^2} \right)^\epsilon \left[\frac{4}{\epsilon} \ln \eta + \bar{\mathcal{R}}_{\text{aux}} \right] + \mathcal{O}(\epsilon) + \mathcal{O}(\Lambda^{-1}), \quad (30)$$

and

$$\bar{\mathcal{R}}_{\text{aux-q}} = \frac{3}{\epsilon} - \frac{2\pi^2}{3} + \frac{7}{2} - \frac{1}{N_c^2} \left[\frac{1}{\epsilon^2} + \frac{3}{2\epsilon} + 4 \right], \quad (31)$$

$$\bar{\mathcal{R}}_{\text{aux-g}} = \frac{1}{\epsilon^2} + \frac{11}{\epsilon} - \frac{2\pi^2}{3} + \frac{67}{9} - \frac{n_f}{N_c} \left[\frac{2}{3\epsilon} + \frac{10}{9} - \frac{1}{N_c^2} \left(\frac{1}{3\epsilon} - \frac{1}{6} \right) \right]. \quad (32)$$

The auxiliary-gluon case includes the contribution for the real process with an initial-state auxiliary gluon and a quark-antiquark pair in the final state, and n_f is the number of quark flavors included. The correct separation of this contribution from the familiar contribution necessitates the parameter η to be specified later.

The virtual contribution at NLO lives in the same phase space as the Born contribution, but involves the interference between the tree-level amplitude and the one-loop amplitude instead of the tree-level squared matrix element. The paper [81] addresses the calculation of one-loop amplitudes with a space-like gluon using the auxiliary parton method, and it is observed that they contain $\ln \Lambda$. They also violate auxiliary parton universality and the smooth on-shell limit, and in Section 4, we will argue that the complete virtual contribution causing these violations is again proportional to the Born contribution, and given by

$$dV^{*\text{unf}}(\xi_{\text{in}}, k_\perp, \bar{x}_{\bar{\text{in}}}; \{p_i\}_{i=1}^n) = \alpha_\epsilon N_c \text{Re}(\mathcal{V}_{\text{aux}}) dB^*(\xi_{\text{in}}, k_\perp, \bar{x}_{\bar{\text{in}}}; \{p_i\}_{i=1}^n), \quad (33)$$

with

$$\mathcal{V}_{\text{aux}} = \left(\frac{\mu^2}{|k_\perp|^2} \right)^\epsilon \left[\frac{2}{\epsilon} \ln \frac{\Lambda}{\xi_{\text{in}}} - i\pi + \bar{\mathcal{V}}_{\text{aux}} \right] + \mathcal{O}(\epsilon) + \mathcal{O}(\Lambda^{-1}), \quad (34)$$

and

$$\bar{\mathcal{V}}_{\text{aux-q}} = \frac{1}{\epsilon} \frac{13}{6} + \frac{\pi^2}{3} + \frac{80}{18} + \frac{1}{N_c^2} \left[\frac{1}{\epsilon^2} + \frac{3}{2\epsilon} + 4 \right] - \frac{n_f}{N_c} \left[\frac{2}{3\epsilon} + \frac{10}{9} \right], \quad (35)$$

$$\bar{\mathcal{V}}_{\text{aux-g}} = -\frac{1}{\epsilon^2} + \frac{\pi^2}{3}. \quad (36)$$

The parameter n_f is the number of flavors running in quark loops. We call this the *unfamiliar virtual contribution*. The rest of the virtual contribution we call *familiar*, and is independent of the auxiliary partons used, has a smooth on-shell limit, and behaves regularly with Λ . While the last claim will not exceed the status of conjecture in this publication, the former two will get more solid ground.

2.5 Interpretation of the unfamiliar contributions

Combining the unfamiliar contributions and organizing them in a suggestive manner, we can write

$$dR^{*\text{unf}} + dV^{*\text{unf}} = \Delta_{\text{unf}} dB^* , \quad (37)$$

where

$$\Delta_{\text{unf}} = \frac{\alpha_\epsilon N_c}{\epsilon} \left(\frac{\mu^2}{|k_\perp|^2} \right)^\epsilon [2J_{\text{univ}} + J_{\text{aux}} + 2J_{\text{remnant}}] , \quad (38)$$

with

$$J_{\text{univ}} = \frac{11}{6} - \frac{n_f}{3N_c} - \frac{\mathcal{K}}{N_c}(-\epsilon) \quad \text{writing} \quad \mathcal{K} = N_c \left(\frac{67}{18} - \frac{\pi^2}{6} \right) - \frac{5n_f}{9} , \quad (39)$$

and

$$J_{\text{aux-q}} = \frac{3}{2} - \frac{1}{2}(-\epsilon) \quad , \quad J_{\text{aux-g}} = \frac{11}{6} + \frac{n_f}{3N_c^3} + \frac{n_f}{6N_c^3}(-\epsilon) , \quad (40)$$

while

$$J_{\text{remnant}} = 2 \ln \eta + \ln \frac{\Lambda}{\xi_{\text{in}}} . \quad (41)$$

We can now compare with the results in [29] for the corrections to impact factors. Realize the difference in the definition of parameter for dimensional regularization $\epsilon = -\epsilon$. We see that $J_{\text{aux-q}} + J_{\text{univ}}$ is identical to their equation (4.9), and $J_{\text{aux-g}} + J_{\text{univ}}$ is identical to their equation (5.11). We introduced the symbol \mathcal{K} for the same quantity defined in their equation (4.10). The factor of 2 for J_{univ} in Eq. (38) appears because our virtual result is not UV-subtracted. In other words, $\epsilon \tilde{V}_{\text{aux}} - J_{\text{univ}}$ gives (two times) the virtual expressions between the square brackets in (4.6) and (5.9) in [29].

Having identified the impact factor contributions, we are left with J_{remnant} . We will see that the consistent choice of the parameter η necessary to correctly separate the familiar and unfamiliar real contributions demands that the scaling relation of Eq. (26) is set by $\lambda = \Lambda$, and is then given by $\eta = 1 - x_{\text{in}}$. This setting of the scaling also implies that $\Lambda/\xi_{\text{in}} = 1/x_{\text{in}}$, and we find

$$J_{\text{remnant}} = 2 \ln(1 - x_{\text{in}}) + \ln \frac{1}{x_{\text{in}}} . \quad (42)$$

The $\ln(1 - x_{\text{in}})$ term is subleading in the power expansion in x_{in} , and it drops out in the high energy limit. The remaining contribution from Eq. (42) to Δ_{unf} proportional to $\ln(1/x_{\text{in}})/\epsilon$ does

not depend on the auxiliary parton (is universal) and cancels against a corresponding singular term in the real familiar contribution that is isolated in Eq. (48). Hence we interpret this remnant term not to be a part of the auxiliary parton impact factor, but move it to Δ_{coll}^* , discussed in the next subsection, so it belongs to the evolution of the gluon transverse momentum dependent distribution (TMD).

2.6 Non-cancelling familiar contributions

The unfamiliar contributions are those with the non-smooth on-shell limit, therefore familiar ones have a smooth on-shell limit. We must specify here that the smoothness really only refers to the $1/\epsilon$ divergencies. The familiar real contribution may still develop finite logarithms of $|k_\perp|$, but they are eventually integrable underneath the k_\perp -integral.

Since the familiar contributions have a smooth on-shell limit, the (non)-cancellation of their divergencies are expected happen just as in the on-shell limit. In Section 5 and Section 6 we show that this is indeed the case and that the only non-cancelling ones are of the collinear kind related to the initial-state partons. At the on-shell \bar{P} -side they are of the well-known form, at the space-like P -side they are slightly different.

Let us repeat the usual treatment of the non-cancelling collinear divergencies. The tree-level squared matrix elements in the familiar real contribution have singularities when the radiative gluon becomes collinear to \bar{P} . Part of the divergent contribution can only be written in a form such that they are proportional to the Born result if the convolution with the PDF is included. This contribution can then be written as the Born cross section, but with the collinear PDF $f(\bar{x}_{\text{in}})$ replaced by

$$-\frac{\alpha_\epsilon}{\epsilon} \int_{\bar{x}_{\text{in}}}^1 dz \mathcal{P}_{\text{in}}^{\text{reg}}(z) \frac{1}{z} f\left(\frac{\bar{x}_{\text{in}}}{z}\right) \quad (43)$$

where $\mathcal{P}_{\text{in}}^{\text{reg}}(z)$ is the regulated splitting function coming from the collinear limit of the tree-level squared matrix element. We must mention here that it includes the color factor that we later in this work will often take out. So at this point, it just has its singularity at $z = 1$ regulated via the plus-prescription, but has no terms proportional to $\delta(1 - z)$. The plus-prescription takes out the soft-collinear contribution which *does* cancel against the virtual contribution. This term can then be interpreted within the subtractive renormalization of the PDF, at which point the term $\gamma_{\text{in}}\delta(1 - z)$, coming from the non-cancelling collinear virtual contribution, is added to the splitting function. The well-known coefficients are

$$\gamma_q = \gamma_{\bar{q}} = \frac{3}{2} \frac{N_c^2 - 1}{2N_c} \quad , \quad \gamma_g = \frac{11N_c - 2n_f}{6} \quad (44)$$

This has to happen in conjunction with ultraviolet (UV) renormalization, because within dimensional regularization, collinear and UV divergencies typically cancel to a large degree among themselves in one-loop amplitudes, and the collinear divergencies have to be “put back” in the form of UV counterterms. So in short, in the $\overline{\text{MS}}$ scheme

$$\Delta_{\text{coll}} = -\frac{\alpha_\epsilon}{\epsilon} \int_{\bar{x}_{\text{in}}}^1 dz [\mathcal{P}_{\text{in}}^{\text{reg}}(z) + \gamma_{\text{in}}\delta(1 - z)] \frac{1}{z} f\left(\frac{\bar{x}_{\text{in}}}{z}\right) \quad (45)$$

is imagined to be absorbed into the NLO PDF.

The tree-level squared matrix element also has singularities when the radiative gluon becomes collinear to P , and in Section 6 we find that the associated non-cancelling contribution equivalent to the above can again be written as the Born cross section, but the k_\perp -dependent PDF $F(x_{\text{in}}, |k_\perp|)$ replaced by

$$-\frac{\alpha_\epsilon}{\epsilon} \int_{x_{\text{in}}}^1 dz \frac{2N_c}{[1-z]_+} \frac{1}{z} F\left(\frac{x_{\text{in}}}{z}, |k_\perp|\right). \quad (46)$$

Regarding the non-cancelling familiar virtual contributions, we will show in Section 5 that they are identical to the ones in the on-shell limit, so we see that

$$\Delta_{\text{coll}}^* = -\frac{\alpha_\epsilon}{\epsilon} \int_{x_{\text{in}}}^1 dz \left[\frac{2N_c}{[1-z]_+} + \gamma_g \delta(1-z) \right] \frac{1}{z} F\left(\frac{x_{\text{in}}}{z}, |k_\perp|\right) \quad (47)$$

must be absorbed into the k_\perp -dependent NLO PDF.

There is a $\ln(1/x_{\text{in}})/\epsilon$ hidden in Eq. (46) which we would like to combine with the one from Eq. (42). The reason why it appears in Eq. (46) is that we insisted on letting the plus-distribution act on the integrand including $1/z$, just like in the collinear case for which the *regularized* splitting function appears. It brings the familiar contribution to a form for which we know how the other $1/\epsilon$ poles cancel. As an illustration, not letting the plus-distribution act on $1/z$ would remove x_{in} from $x_{\text{in}}S = 2x_{\text{in}}P_A \cdot \bar{P}$ appearing on the right-hand side of Eq. (149), generating unwanted, unprotected, occurrences of $\ln S$. We can undo this, and write

$$\begin{aligned} \int_{x_{\text{in}}}^1 dz \frac{1}{[1-z]_+} \frac{1}{z} F\left(\frac{x_{\text{in}}}{z}, |k_\perp|\right) \\ = \int_{x_{\text{in}}}^1 \frac{dz}{z(1-z)} \left[F\left(\frac{x_{\text{in}}}{z}, |k_\perp|\right) - F(x_{\text{in}}, |k_\perp|) \right] + \ln\left(\frac{1}{x_{\text{in}}}\right) F(x_{\text{in}}, |k_\perp|). \end{aligned} \quad (48)$$

Now, the second term on the right-hand side cancels against Eq. (42). The integral on the right-hand side does not generate $\ln x_{\text{in}}$ anymore, as can easily be seen with the variable substitution $z = 1/(1+y/x_{\text{in}})$:

$$\int_{x_{\text{in}}}^1 \frac{dz}{z(1-z)} \left[F\left(\frac{x_{\text{in}}}{z}, |k_\perp|\right) - F(x_{\text{in}}, |k_\perp|) \right] = \int_0^{1-x_{\text{in}}} \frac{dy}{y} \left[F(x_{\text{in}}+y, |k_\perp|) - F(x_{\text{in}}, |k_\perp|) \right]. \quad (49)$$

Since the contribution looks very similar to the collinear case, it makes sense to consider a scenario like employed in [79], and write the function F using a convolution involving a collinear PDF as

$$F(x_{\text{in}}, |k_\perp|) = \int_{x_{\text{in}}}^1 \frac{dz}{z} \mathcal{C}(z, |k_\perp|) f\left(\frac{x_{\text{in}}}{z}\right). \quad (50)$$

We will not digress on the meaning of the function $\mathcal{C}(z, |k_\perp|)$ as it is beyond the scope of this work, but we do observe that

$$\begin{aligned} \int_{x_{\text{in}}}^1 dz \frac{1}{[1-z]_+} \frac{1}{z} F\left(\frac{x_{\text{in}}}{z}, |k_\perp|\right) \\ = \int_{x_{\text{in}}}^1 \frac{dz}{z} \mathcal{C}(z, |k_\perp|) \left[\int_{x_{\text{in}}/z}^1 dy \frac{1}{[1-y]_+} \frac{1}{y} f\left(\frac{x_{\text{in}}}{yz}\right) + \ln\left(\frac{1-x_{\text{in}}}{1-x_{\text{in}}/z}\right) f\left(\frac{x_{\text{in}}}{z}\right) \right], \end{aligned} \quad (51)$$




Figure 1: Kinematics of the different contributions for the example in which the auxiliary partons are quarks, the collinear initial state parton is a gluon, and the real-radiation parton is also a gluon.

and see that this non-cancelling contribution can be interpreted in terms of evolution of the collinear PDF.

3 Unfamiliar real contribution

Originally, the auxiliary parton method was introduced just as a trick to obtain helicity amplitudes with space-like partons. In order to go to NLO, we need to lift the method to cross section level including phase space. First, we need to introduce some more notation. We will frequently take a few arguments explicitly out of the list $\{p_i\}_{i=1}^n$ in a differential phase space, and give them different symbols so we can write their differential volumes explicitly. So, for example we can write

$$d\Sigma_{n+2}^*(k_{\text{in}}, k_{\overline{\text{in}}}; \{p_i\}_{i=1}^{n+2}) = d^4q \delta(q^2) d^4r \delta(r^2) \frac{d\Sigma_{n+2}^*}{dq dr}(k_{\text{in}}, k_{\overline{\text{in}}}; q, r, \{p_i\}_{i=1}^n), \quad (52)$$

implying $q = p_{n+1}$ and $r = p_{n+2}$. Prepending an expression like on the right-hand side with an integral symbol means we are only integrating over these explicit variables. The essential observation is now that the squared matrix element relation of Eq. (22) can be formulated in terms of differential cross sections as follows:

$$\frac{2\xi_{\text{in}}}{C_{\text{aux}}} \int d^4q \delta(q^2) \delta(\xi_{\text{in}} + \xi_q - \Lambda) \delta^2(k_{\perp} + q_{\perp}) \frac{d\Sigma_{n+1}^{\text{aux}}}{dq}(\Lambda P, k_{\overline{\text{in}}}; q, \{p_i\}_{i=1}^n) J_B(\{p_i\}_{i=1}^n) \xrightarrow{\Lambda \rightarrow \infty} dB^*(k_{\text{in}}, k_{\overline{\text{in}}}; \{p_i\}_{i=1}^n). \quad (53)$$

On the left-hand side there is the on-shell differential cross section for the process with auxiliary partons, with auxiliary initial-state momentum ΛP and auxiliary final-state momentum q that does *not* enter the jet function. Consequently, it is the only final-state momentum for which the P component can become arbitrarily large, “consuming” the large initial-state component Λ . This is expressed by the delta-function with argument $\xi_{\text{in}} + \xi_q - \Lambda$. Realize that the flux factor inside $d\Sigma_{n+1}^{\text{aux}}$ on the left-hand side has a factor Λ , playing the role of a denominator factor in Eq. (22). The other one comes from the $1/\xi_q$ in the differential volume of q . The factor $1/|k_\perp|^2$ in the definition of Eq. (14) was of course chosen to accommodate the factor in Eq. (22), and the explicit factor ξ_{in} together with the flux factor balance the ξ_{in}^2 in Eq. (22). The factor 2 on the left-hand side above corrects the factor 1/2 in the differential volume of q .

In the case of auxiliary gluons, one could argue that any possible final-state gluon could take the auxiliary role. This could be imagined to be facilitated by the jet function, choosing any gluon, but only one at a time, to be allowed to have large ξ variable. So if the Born process (on the right-hand side of Eq. (53)) involves n_g final-state gluons, then on the left-hand side there would be $n_g + 1$ contributions. This would effectively increase the symmetry factor $1/(n_g + 1)!$ inside $d\Sigma_{n+1}^{\text{aux}}$ to $1/n_g!$, so we can just as well assume that one specific gluon on the left-hand side is the auxiliary one, and assume that $d\Sigma_{n+1}^{\text{aux}}$ has a factor $1/n_g!$ from the start.

In order to introduce the unfamiliar real contribution, let us however start with the simple situation of an auxiliary quark-antiquark pair, and a Born level final state that does not involve any gluons. As mentioned earlier, the unfamiliar contribution appears because we should allow the radiative gluon to take part in the “consumption” of the large P component Λ . For clarity, we use a separate symbol r instead of p_{n+1} for the momentum of the radiative gluon. In order to let r contribute to the consumption of Λ and the provision of k_\perp , we add ξ_r and r_\perp as arguments to the auxiliary delta functions compared to Eq. (53), and define the *unfamiliar real contribution* as:

$$\begin{aligned}
& dR^{\text{unf}}(k_{\text{in}}, k_{\overline{\text{in}}}; \{p_i\}_{i=1}^n) \\
& \stackrel{\Lambda \rightarrow \infty}{=} \frac{\alpha_\epsilon \mu^{2\epsilon}}{\pi_\epsilon} \frac{2\xi_{\text{in}}}{C_{\text{aux}}} \int d^{4-2\epsilon} q \delta(q^2) d^{4-2\epsilon} r \delta(r^2) \delta(\xi_{\text{in}} + \xi_q + \xi_r - \Lambda) \delta^{2-2\epsilon}(k_\perp + q_\perp + r_\perp) \\
& \quad \times \frac{d\Sigma_{n+2}^{\text{aux}}}{dq dr}(\Lambda P, k_{\overline{\text{in}}}; r, q, \{p_i\}_{i=1}^n) J_B(\{p_i\}_{i=1}^n). \quad (54)
\end{aligned}$$

Now, with a similar reasoning as in the Born case, we can argue that the formula can be used also in case of a Born process with final-state gluons, by considering the auxiliary and/or radiative one as special and including a factor $1/n_g!$ in $d\Sigma_{n+2}^{\text{aux}}$ from the start. Additionally, when both are gluons, the auxiliary and the radiative one are distinguished by $\xi_r < \xi_q$, as we will see later.

While ξ_r is allowed to become large in Eq. (54), it is also allowed to become arbitrarily small, a phase space region we want to reserve for the familiar contribution. Thus we need to subtract something from the unfamiliar contribution as defined in Eq. (54), which we will address Section 3.1.

In order to proceed, we write the momenta q and r explicitly in the Sudakov decomposition

and denote

$$\Lambda' = \Lambda - \xi_{\text{in}} \quad (55)$$

so we can scale the variables $\xi_q = \Lambda' x_q$ and $\xi_r = \Lambda' x_r$ to get

$$\begin{aligned} & dR^{\text{unf}}(k_{\text{in}}, k_{\text{in}}; \{p_i\}_{i=1}^n) \\ & \stackrel{\Lambda \rightarrow \infty}{=} \frac{\alpha_\epsilon \mu^{2\epsilon}}{\pi_\epsilon} \frac{2\xi_{\text{in}}}{C_{\text{aux}}} \int_0^1 \frac{dx_q}{2x_q} \int_0^{\bar{x}_{\text{in}}} d\bar{x}_q \int d^{2-2\epsilon} q_\perp \delta\left(\bar{x}_q - \frac{|q_\perp|^2}{v^2 x_q \Lambda'}\right) \\ & \quad \times \int_0^1 \frac{dx_r}{2x_r} \int_0^{\bar{x}_{\text{in}}} d\bar{x}_r \int d^{2-2\epsilon} r_\perp \delta\left(\bar{x}_r - \frac{|r_\perp|^2}{v^2 x_r \Lambda'}\right) \\ & \quad \times \Lambda'^{-1} \delta(x_q + x_r - 1) \delta^{2-2\epsilon}(k_\perp + q_\perp + r_\perp) \\ & \quad \times \frac{d\Sigma_{n+2}^{\text{aux}}}{dq dr} ((\Lambda' + \xi_{\text{in}})P, \bar{x}_{\text{in}}; x_r \Lambda' P + r_\perp + \bar{x}_r \bar{P}, x_q \Lambda' P + q_\perp + \bar{x}_q \bar{P}, \{p_i\}_{i=1}^n) J_B(\{p_i\}_{i=1}^n). \end{aligned} \quad (56)$$

Inside $d\Sigma_{n+2}^{\text{aux}}$ there is now a tree-level squared matrix element with three partons with large coefficients for P . Realize that \bar{x}_q and \bar{x}_r are suppressed with Λ due to the on-shell conditions. We calculate this triple- Λ limit in Appendix D and find to be

$$\begin{aligned} & \frac{1}{C_{\text{aux}}} |\overline{\mathcal{M}}^{\text{aux}}|^2 ((\Lambda' + \xi_{\text{in}})P, k_{\text{in}}; x_r \Lambda' P + r_\perp + \bar{x}_r \bar{P}, x_q \Lambda' P + q_\perp + \bar{x}_q \bar{P}, \{p_i\}_{i=1}^n) \\ & \stackrel{\Lambda' \rightarrow \infty}{\longrightarrow} \mathcal{Q}_{\text{aux}}(x_q, q_\perp, x_r, r_\perp) \frac{\Lambda'^2 |\overline{\mathcal{M}}^*|^2(\xi_{\text{in}}P - q_\perp - r_\perp, k_{\text{in}}; \{p_i\}_{i=1}^n)}{\xi_{\text{in}}^2 |q_\perp + r_\perp|^2}, \end{aligned} \quad (57)$$

with

$$\begin{aligned} \mathcal{Q}_{\text{aux}}(x_q, q_\perp, x_r, r_\perp) &= x_q x_r \mathcal{P}_{\text{aux}}(x_q, x_r) |q_\perp + r_\perp|^2 \\ & \times \left[\frac{c_{\bar{q}}}{|q_\perp|^2 |r_\perp|^2} + \frac{1}{x_r |q_\perp|^2 + x_q |r_\perp|^2 - x_q x_r |q_\perp + r_\perp|^2} \left(\frac{c_r x_r^2}{|r_\perp|^2} + \frac{c_q x_q^2}{|q_\perp|^2} \right) \right]. \end{aligned} \quad (58)$$

The auxiliary parton dependent quantities are

$$\text{auxiliary quarks: } \mathcal{P}_{\text{aux}}(x_q, x_r) = \frac{1 + x_q^2 - \epsilon x_r^2}{x_r}, \quad c_q = c_{\bar{q}} = N_c, \quad c_r = \frac{-1}{N_c}, \quad (59)$$

$$\text{auxiliary gluons: } \mathcal{P}_{\text{aux}}(x_q, x_r) = \frac{1 + x_q^4 + x_r^4}{x_q x_r}, \quad c_q = c_{\bar{q}} = c_r = N_c, \quad (60)$$

$$\text{gluon to quarks: } \mathcal{P}_{\text{aux}}(x_q, x_r) = 1 - \frac{2x_q x_r}{1 - \epsilon}, \quad c_q = c_r = \frac{1}{2}, \quad c_{\bar{q}} = \frac{-1}{2N_c^2}. \quad (61)$$

The ‘‘gluon to quarks’’ refers to the real process with an initial-state auxiliary gluon and a quark-antiquark pair in the final state. We keep the labels q and r for the final-state partons in this case.

In the following, we prefer to work with a parameter for dimensional regularization that is imagined to be positive, and we introduce

$$\bar{\epsilon} = -2\epsilon. \quad (62)$$

Then, at this stage, we find (remember the flux factor)

$$\begin{aligned}
& dR^{\text{unf}}(k_{\text{in}}, k_{\overline{\text{in}}}; \{p_i\}_{i=1}^n) \\
& \xrightarrow{\Lambda \rightarrow \infty} \frac{2a_\epsilon}{\pi_\epsilon \mu^{\bar{\epsilon}}} \int_0^1 \frac{dx_q}{2x_q} \int_0^{\bar{x}_{\overline{\text{in}}}} d\bar{x}_q \int d^{2+\bar{\epsilon}} q_\perp \delta\left(\bar{x}_q - \frac{|q_\perp|^2}{v^2 x_q \Lambda'}\right) \\
& \quad \times \int_0^1 \frac{dx_r}{2x_r} \int_0^{\bar{x}_{\overline{\text{in}}}} d\bar{x}_r \int d^{2+\bar{\epsilon}} r_\perp \delta\left(\bar{x}_r - \frac{|r_\perp|^2}{v^2 x_r \Lambda'}\right) \Lambda'^{-1} \delta(x_q + x_r - 1) \delta^{2+\bar{\epsilon}}(k_\perp + q_\perp + r_\perp) \\
& \quad \times \mathcal{Q}_{\text{aux}}(x_q, q_\perp, x_r, r_\perp) \frac{\xi_{\text{in}}^2}{\Lambda'} \frac{\Lambda'^2 d\Sigma_n^*(\xi_{\text{in}} P + k_\perp, \bar{x}_{\overline{\text{in}}}; \{p_i\}_{i=1}^n)}{\xi_{\text{in}}^2 |k_\perp|^2} J_B(\{p_i\}_{i=1}^n).
\end{aligned} \tag{63}$$

After performing the \bar{x}_q, \bar{x}_r integrals, we find Eq. (29) with

$$\begin{aligned}
\mathcal{R}_{\text{aux}}(\xi_{\text{in}}, k_\perp, \mu, \Lambda, \bar{x}_{\overline{\text{in}}}) &= \frac{2a_\epsilon}{\pi_\epsilon \mu^{\bar{\epsilon}}} \int_0^1 \frac{dx_q}{2x_q} \int d^{2+\bar{\epsilon}} q_\perp \theta(0 < |q_\perp|^2 < v^2 \bar{x}_{\overline{\text{in}}} x_q \Lambda') \\
& \quad \times \int_0^1 \frac{dx_r}{2x_r} \int d^{2+\bar{\epsilon}} r_\perp \theta(0 < |r_\perp|^2 < v^2 \bar{x}_{\overline{\text{in}}} x_r \Lambda') \\
& \quad \times \delta^{2+\bar{\epsilon}}(k_\perp + q_\perp + r_\perp) \delta(x_q + x_r - 1) \mathcal{Q}_{\text{aux}}(x_q, q_\perp, x_r, r_\perp).
\end{aligned} \tag{64}$$

Understanding that $x_q + x_r = 1$ and $k_\perp + q_\perp + r_\perp = 0$, we can write

$$\begin{aligned}
\mathcal{Q}_{\text{aux}}(x_q, q_\perp, x_r, r_\perp) &= x_q x_r \mathcal{P}_{\text{aux}}(x_q, x_r) |k_\perp|^2 \left(\frac{c_{\bar{q}}}{|r_\perp|^2 |r_\perp + k_\perp|^2} + \frac{c_q (1 - x_r)^2}{|r_\perp + k_\perp|^2 |r_\perp + x_r k_\perp|^2} \right. \\
& \quad \left. + \frac{c_r x_r^2}{|r_\perp|^2 |r_\perp + x_r k_\perp|^2} \right), \tag{65}
\end{aligned}$$

and we see that the explicit $1/x_q$ and $1/x_r$ in the integral cancel, and that the only possible singularities with respect to x_q, x_r are in $\mathcal{P}_{\text{aux}}(x_q, x_r)$. In case of auxiliary quarks, there is only a singularity with respect to x_r , and in case of auxiliary gluons, we demand that $x_r < x_q$. Remember that we distinguish between the gluons and must not include a factor 2. Because of this, and the fact that the q_\perp, r_\perp integrals are UV finite, we can safely take $\Lambda \rightarrow \infty$ inside the $\theta(0 < |q_\perp|^2 < v^2 \bar{x}_{\overline{\text{in}}} x_q \Lambda')$, and eliminate the q_\perp integral and the x_q integral. Thus we find

$$\begin{aligned}
\mathcal{R}_{\text{aux}} &= \frac{a_\epsilon}{2\pi_\epsilon \mu^{\bar{\epsilon}}} \int_0^u dx \mathcal{P}_{\text{aux}}(1 - x, x) \int d^{2+\bar{\epsilon}} r_\perp \theta(0 < |r_\perp|^2 < v^2 x \bar{x}_{\overline{\text{in}}} \Lambda') \\
& \quad \times |k_\perp|^2 \left(\frac{c_{\bar{q}}}{|r_\perp|^2 |r_\perp + k_\perp|^2} + \frac{c_q (1 - x)^2}{|r_\perp + k_\perp|^2 |r_\perp + x k_\perp|^2} + \frac{c_r x^2}{|r_\perp|^2 |r_\perp + x k_\perp|^2} \right). \tag{66}
\end{aligned}$$

In order for the formulas to accommodate for the phase space partition for the auxiliary gluon case, we write an upper integration limit “u” for the x_r -integration, with

$$u = 1 \text{ for the auxiliary quark case, and } u = 1/2 \text{ for the auxiliary-gluon case.} \tag{67}$$

Now let us take out the pole in \mathcal{P}_{aux} explicitly, and write

$$\mathcal{P}_{\text{aux}}(1-x, x) = \frac{\mathcal{P}_{\text{aux}}^{\text{pole}}}{x} + \mathcal{P}_{\text{aux}}^{\text{rest}}(x). \quad (68)$$

For the terms with $\mathcal{P}_{\text{aux}}^{\text{rest}}$, we can safely take $\Lambda \rightarrow \infty$ also in $\theta(0 < |r_{\perp}|^2 < v^2 x \bar{x}_{\text{in}} \Lambda')$. Then we can use Eq. (254), and find that the part with $\mathcal{P}_{\text{aux}}^{\text{rest}}$ is given by

$$\mathcal{R}_{\text{aux}}^{\text{rest}} = \frac{2\alpha_{\epsilon}}{\bar{\epsilon}} \left(\frac{|k_{\perp}|}{\mu} \right)^{\bar{\epsilon}} \int_0^u dx \mathcal{P}_{\text{aux}}^{\text{rest}}(x) (c_{\bar{q}} + c_q (1-x)^{\bar{\epsilon}} + c_r x^{\bar{\epsilon}}), \quad (69)$$

which we will calculate later. Regarding the terms with $\mathcal{P}_{\text{aux}}^{\text{pole}}/x$ in Eq. (66), we observe that for the one with c_r , we can scale r_{\perp} with x to get

$$\mathcal{R}_{\text{aux}}^{\text{pole},r} = \frac{\alpha_{\epsilon}}{2\pi_{\epsilon} \mu^{\bar{\epsilon}}} c_r \int_0^u dx \frac{\mathcal{P}_{\text{aux}}^{\text{pole}}}{x} x^{\bar{\epsilon}} \int d^{2+\bar{\epsilon}} r_{\perp} \theta(0 < x|r_{\perp}|^2 < v^2 x \bar{x}_{\text{in}} \Lambda') \frac{|k_{\perp}|^2}{|r_{\perp}|^2 |r_{\perp} + k_{\perp}|^2}. \quad (70)$$

We see that the singularity in x is protected by $\bar{\epsilon}$, and that $x \rightarrow 0$ does not conflict with $\Lambda \rightarrow \infty$ in the integration boundary, so also here we can take the latter limit, and find

$$\mathcal{R}_{\text{aux}}^{\text{pole},r} = \frac{2\alpha_{\epsilon}}{\bar{\epsilon}} \left(\frac{|k_{\perp}|}{\mu} \right)^{\bar{\epsilon}} c_r \frac{\mathcal{P}_{\text{aux}}^{\text{pole}}}{\bar{\epsilon}} u^{\bar{\epsilon}}. \quad (71)$$

For the other terms in Eq. (66) with $\mathcal{P}_{\text{aux}}^{\text{pole}}$, we write

$$\mathcal{R}_{\text{aux}}^{\text{pole},q} + \mathcal{R}_{\text{aux}}^{\text{pole},\bar{q}} = \mathcal{R}_{\text{aux}}^{\text{pole},0} + \mathcal{R}_{\text{aux}}^{\text{pole},1}, \quad (72)$$

where

$$\mathcal{R}_{\text{aux}}^{\text{pole},0} = \frac{\alpha_{\epsilon}}{2\pi_{\epsilon} \mu^{\bar{\epsilon}}} |k_{\perp}|^2 [c_{\bar{q}} + c_q] \int_0^u dx \frac{\mathcal{P}_{\text{aux}}^{\text{pole}}}{x} \int d^{2+\bar{\epsilon}} r_{\perp} \frac{\theta(0 < |r_{\perp}|^2 < v^2 x \bar{x}_{\text{in}} \Lambda')}{|r_{\perp}|^2 |r_{\perp} + k_{\perp}|^2}, \quad (73)$$

and

$$\begin{aligned} \mathcal{R}_{\text{aux}}^{\text{pole},1} &= \frac{\alpha_{\epsilon}}{2\pi_{\epsilon} \mu^{\bar{\epsilon}}} |k_{\perp}|^2 c_q \int_0^u dx \frac{\mathcal{P}_{\text{aux}}^{\text{pole}}}{x} \int d^{2+\bar{\epsilon}} r_{\perp} \theta(0 < |r_{\perp}|^2 < v^2 x \bar{x}_{\text{in}} \Lambda') \\ &\quad \times \left(\frac{(1-x)^2}{|r_{\perp} + k_{\perp}|^2 |r_{\perp} + x k_{\perp}|^2} - \frac{1}{|r_{\perp}|^2 |r_{\perp} + k_{\perp}|^2} \right). \end{aligned} \quad (74)$$

We see that the integrand of $\mathcal{R}_{\text{aux}}^{\text{pole},1}$ does not have a singularity in x , and we can take $\Lambda \rightarrow \infty$ in the integration boundary. Thus we find

$$\begin{aligned} \mathcal{R}_{\text{aux}}^{\text{pole},1} &= \frac{2\alpha_{\epsilon}}{\bar{\epsilon}} \left(\frac{|k_{\perp}|}{\mu} \right)^{\bar{\epsilon}} c_q \int_0^u dx \frac{\mathcal{P}_{\text{aux}}^{\text{pole}}}{x} [(1-x)^{\bar{\epsilon}} - 1] \\ &= 2\alpha_{\epsilon} \left(\frac{|k_{\perp}|}{\mu} \right)^{\bar{\epsilon}} c_q \mathcal{P}_{\text{aux}}^{\text{pole}} [-\text{Li}_2(u) + \mathcal{O}(\bar{\epsilon})]. \end{aligned} \quad (75)$$

For $\mathcal{R}_{\text{aux}}^{\text{pole},0}$ finally, we use Eq. (256) to find

$$\mathcal{R}_{\text{aux}}^{\text{pole},0} = 2\alpha_\epsilon \left(\frac{|k_\perp|}{\mu} \right)^{\bar{\epsilon}} (c_{\bar{q}} + c_q) \mathcal{P}_{\text{aux}}^{\text{pole}} \left[\frac{1}{\bar{\epsilon}^2} - \frac{1}{\bar{\epsilon}} \ln \frac{|k_\perp|^2}{u\nu^2 \bar{\chi}_{\text{in}} \Lambda'} + \frac{1}{4} \text{Li}_2 \left(\frac{|k_\perp|^2}{u\nu^2 \bar{\chi}_{\text{in}} \Lambda'} \right) \right]. \quad (76)$$

So combined, including the assumption

$$c_{\bar{q}} = c_q \quad (77)$$

which holds both for the auxiliary quarks and for auxiliary gluons, we find

$$\begin{aligned} \mathcal{R}_{\text{aux}}^{\text{pole}} &= 2\alpha_\epsilon \left(\frac{|k_\perp|}{\mu} \right)^{\bar{\epsilon}} \mathcal{P}_{\text{aux}}^{\text{pole}} \\ &\times \left\{ c_q \left[\frac{2}{\bar{\epsilon}^2} - \frac{2}{\bar{\epsilon}} \ln \frac{|k_\perp|^2}{u\nu^2 \bar{\chi}_{\text{in}} \Lambda'} + \frac{1}{2} \text{Li}_2 \left(\frac{|k_\perp|^2}{u\nu^2 \bar{\chi}_{\text{in}} \Lambda'} \right) - \text{Li}_2(u) \right] + c_r \frac{u^{\bar{\epsilon}}}{\bar{\epsilon}^2} \right\}. \end{aligned} \quad (78)$$

Realize that the $\text{Li}_2(|k_\perp|^2/(u\nu^2 \bar{\chi}_{\text{in}} \Lambda')) \rightarrow 0$ for $\Lambda \rightarrow 0$.

3.1 Subtraction of double-counted contributions

The unfamiliar real contribution calculated above includes the phase space region for which ξ_r is allowed to become small, which is already included in the familiar contribution, and we need to subtract something to correct this. In Section 6, we will see that it is the contribution of Eq. (73), but up to a certain integration limit $u = \eta$ for χ . That is, we need to subtract Eq. (76) at $u = \eta$. Thus we have, taking $\Lambda \rightarrow \infty$ inside the Li_2 functions,

$$\mathcal{R}_{\text{aux}}^{\text{pole}} - \mathcal{R}_{\text{aux}}^{\text{subt}} = 2\alpha_\epsilon \left(\frac{|k_\perp|}{\mu} \right)^{\bar{\epsilon}} \mathcal{P}_{\text{aux}}^{\text{pole}} \left\{ c_q \left[\frac{2}{\bar{\epsilon}} \ln \frac{u}{\eta} - \text{Li}_2(u) \right] + c_r \frac{u^{\bar{\epsilon}}}{\bar{\epsilon}^2} \right\}, \quad (79)$$

with now again $u = 1$ for auxiliary quarks, and $u = 1/2$ for auxiliary gluons. It must be stressed that it is essential that the subtraction term establishes the disappearance of both Λ and $1/\bar{\epsilon}^2$ in Eq. (76). The remaining $1/\bar{\epsilon}^2$ -term in Eq. (79) cancels against virtual contributions. As explained in Section 6, we want to mention already here that in order for this to work out, the scale setting the relation between ξ and P versus χ and P_Λ in Section 2.3 must be $\lambda = \Lambda$.

3.2 Auxiliary quarks

The collinear splitting function is given by

$$\mathcal{P}_{qg}(1-x, x) = \frac{1 + (1-x)^2 + \frac{1}{2} \delta \bar{\epsilon} x^2}{x} = \frac{2}{x} + x - 2 + \frac{1}{2} \delta \bar{\epsilon} x, \quad (80)$$

with $\delta = 1$ (remember that $\bar{\epsilon} = -2\epsilon$). So we find, putting $u = 1$ in Eq. (69),

$$\mathcal{R}_{\text{aux-q}}^{\text{rest}} = \frac{2\alpha_\epsilon}{\bar{\epsilon}} \left(\frac{|k_\perp|}{\mu} \right)^{\bar{\epsilon}} \left\{ [c_{\bar{q}} + c_q + c_r] \left(-\frac{3}{2} + \frac{\delta}{4} \bar{\epsilon} \right) + c_q \frac{5}{4} \bar{\epsilon} + c_r \frac{7}{4} \bar{\epsilon} + \mathcal{O}(\bar{\epsilon}^2) \right\}. \quad (81)$$

Putting the coefficients

$$c_{\bar{q}} = c_q = N_c \quad , \quad c_r = -1/N_c \quad , \quad (82)$$

and forgetting about $\mathcal{O}(\bar{\epsilon}^2)$, we get

$$\mathcal{R}_{\text{aux-q}}^{\text{rest}} = 2\alpha_\epsilon N_c \left(\frac{|k_\perp|}{\mu} \right)^{\bar{\epsilon}} \left\{ -\frac{3}{\bar{\epsilon}} + \frac{5+2\delta}{4} - \frac{1}{N_c^2} \left[-\frac{3}{2\bar{\epsilon}} + \frac{7+\delta}{4} \right] \right\}. \quad (83)$$

Adding Eq. (79) with $\mathcal{P}_{\text{aux}}^{\text{pole}} = 2$ and $u = 1$, we get

$$\mathcal{R}_{\text{aux-q}} = 2\alpha_\epsilon N_c \left(\frac{|k_\perp|}{\mu} \right)^{\bar{\epsilon}} \left\{ -\frac{1}{\bar{\epsilon}} \left[3 + 4 \ln \eta \right] - \frac{\pi^2}{3} + \frac{5+2\delta}{4} - \frac{1}{N_c^2} \left[\frac{2}{\bar{\epsilon}^2} - \frac{3}{2\bar{\epsilon}} + \frac{7+\delta}{4} \right] \right\}, \quad (84)$$

and putting $\bar{\epsilon} = -2\epsilon$ and $\delta = 1$, we find Eq. (31).

3.3 Auxiliary gluons

Now the collinear splitting function is given by

$$\mathcal{P}_{gq}(1-x, x) = \frac{1+x^4 + (1-x)^4}{x(1-x)} = \frac{2}{x} + 2 \left(\frac{1}{1-x} - 2 + x - x^2 \right). \quad (85)$$

Putting $u = 1/2$ in Eq. (69), we find

$$\begin{aligned} \mathcal{R}_{\text{aux-g}}^{\text{rest}} &= \frac{2\alpha_\epsilon}{\bar{\epsilon}} \left(\frac{|k_\perp|}{\mu} \right)^{\bar{\epsilon}} \left\{ [c_{\bar{q}} + c_q + c_r] \left(2 \ln 2 - \frac{11}{6} \right) \right. \\ &\quad \left. + c_q \left(\frac{131}{72} - \frac{132}{72} \ln 2 - \ln^2 2 \right) \bar{\epsilon} + c_r \left(\frac{137}{72} + \frac{132}{72} \ln 2 - \ln^2 2 - \frac{\pi^2}{6} \right) \bar{\epsilon} + \mathcal{O}(\bar{\epsilon}^2) \right\}. \end{aligned} \quad (86)$$

Setting $c_{\bar{q}} = c_q = c_r = N_c$ and forgetting about the $\mathcal{O}(\bar{\epsilon}^2)$, we get

$$\mathcal{R}_{\text{aux-g}}^{\text{rest}} = 2\alpha_\epsilon N_c \left(\frac{|k_\perp|}{\mu} \right)^{\bar{\epsilon}} \left\{ \frac{3}{\bar{\epsilon}} \left(2 \ln 2 - \frac{11}{6} \right) + \frac{67}{18} - \frac{\pi^2}{6} - 2 \ln^2 2 \right\}. \quad (87)$$

Adding Eq. (79) with $\mathcal{P}_{\text{aux}}^{\text{pole}} = 2$ and $u = 1/2$, we get up to $\mathcal{O}(\bar{\epsilon}^0)$

$$\mathcal{R}_{\text{aux-g}} = 2\alpha_\epsilon N_c \left(\frac{|k_\perp|}{\mu} \right)^{\bar{\epsilon}} \left\{ \frac{2}{\bar{\epsilon}^2} - \frac{1}{\bar{\epsilon}} \left[\frac{11}{2} + 4 \ln \eta \right] - \frac{\pi^2}{3} + \frac{67}{18} \right\}, \quad (88)$$

and putting $\bar{\epsilon} = -2\epsilon$, we get

$$\mathcal{R}_{\text{aux-g}} = \alpha_\epsilon N_c \left(\frac{\mu^2}{|k_\perp|^2} \right)^\epsilon \left\{ \frac{1}{\epsilon^2} + \frac{1}{\epsilon} \left[\frac{11}{2} + 4 \ln \eta \right] - \frac{2\pi^2}{3} + \frac{67}{9} \right\}. \quad (89)$$

3.4 Initial-state gluon and final-state quark-antiquark pair

Now the collinear splitting function is given by

$$\mathcal{P}_{gg}(1-x, x) = 1 - \frac{2x(1-x)}{1 + \frac{1}{2}\delta\bar{\epsilon}} = 1 - 2x + 2x^2 + (x - x^2)\delta\bar{\epsilon} + \mathcal{O}(\bar{\epsilon}^2). \quad (90)$$

Putting $u = 1$ in Eq. (69), we find

$$\mathcal{R}^{\text{rest}} = \frac{2\alpha_\epsilon}{\bar{\epsilon}} \left(\frac{|k_\perp|}{\mu} \right)^{\bar{\epsilon}} \left\{ [c_{\bar{q}} + c_q + c_r] \left(\frac{2}{3} + \frac{\delta}{6}\bar{\epsilon} \right) + [c_q + c_r] \frac{-13}{18}\bar{\epsilon} + \mathcal{O}(\bar{\epsilon}^2) \right\}. \quad (91)$$

We keep the labels r, q for final-state variables, now the quark-antiquark pair, so we can still use Eq. (58) instead of Eq. (251). Thus, we need to set $c_{\bar{q}} = -1/(2N_c^2)$ and $c_q = c_r = 1/2$. Forgetting about $\mathcal{O}(\bar{\epsilon}^2)$, we then find

$$\mathcal{R}_{\text{aux-g}}^{\text{q}\bar{\text{q}}} = \mathcal{R}^{\text{rest}} = 2\alpha_\epsilon \left(\frac{|k_\perp|}{\mu} \right)^{\bar{\epsilon}} \left\{ \frac{2}{3\bar{\epsilon}} - \frac{13}{18} + \frac{\delta}{6} - \frac{1}{N_c^2} \left(\frac{1}{3\bar{\epsilon}} + \frac{\delta}{12} \right) \right\}. \quad (92)$$

Putting $\bar{\epsilon} = -2\epsilon$, we get

$$\mathcal{R}_{\text{aux-g}}^{\text{q}\bar{\text{q}}} = \alpha_\epsilon \left(\frac{\mu^2}{|k_\perp|^2} \right)^\epsilon \left\{ -\frac{2}{3\epsilon} - \frac{13}{9} + \frac{\delta}{3} + \frac{1}{N_c^2} \left(\frac{1}{3\epsilon} - \frac{\delta}{6} \right) \right\}. \quad (93)$$

4 Unfamiliar virtual contribution

The unfamiliar real contributions have the general form

$$dR^{\text{unf}} = \alpha_\epsilon N_c \left(\frac{\mu^2}{|k_\perp|^2} \right)^\epsilon \left\{ \frac{A}{\epsilon^2} + \frac{B}{\epsilon} + C \right\} dB^*, \quad (94)$$

where the coefficients A, B, C depend on the auxiliary parton type. So besides violating parton universality, they also clearly violate the smooth on-shell limit. We may expect that there are also virtual contributions that violate these tree-level properties and in fact we find that they also violate the smooth limit $\Lambda \rightarrow \infty$, while having the general form of Eq. (94). In the following we derive the relevant virtual contributions, which we again call *unfamiliar*. *We conjecture that there are no other Λ -dependent virtual contributions than the one we identify here.*

Regarding phase space, the virtual contribution is the same as the Born contribution. Instead of the tree-level squared matrix element, it however contains the interference between the tree-level amplitude and the one-loop amplitude. The non-smooth on-shell limit suggests to have a closer look at it in one-loop amplitudes, in particular the limit of $|k_\perp| \rightarrow 0$ before $\Lambda \rightarrow \infty$. For, maybe large, but fixed value of Λ , this limit corresponds to the collinear limit for the auxiliary parton momenta: with the momenta of Eq. (20), we have

$$2k_1 \cdot k_2 = 2(-\Lambda P) \cdot p_\Lambda = \frac{-\Lambda}{\Lambda - \xi_{\text{in}}} |k_\perp|^2 \rightarrow 0 \quad \text{for} \quad |k_\perp| \rightarrow 0. \quad (95)$$

We let the symbol k_1 refer to a negative energy momentum here to be consistent with the literature in the following. The collinear limit of one-loop amplitudes has been studied in [84, 85]. There, color decompositions of amplitudes are employed, and it is noted that collinear limits can be studied on so-called *primitive amplitudes*. These do not explicitly depend on color, and correspond to a fixed ordering of the external legs.

Let us however start with tree-level. In particular, we will show now that the two limits $\Lambda \rightarrow \infty$ and $|k_\perp| \rightarrow 0$ commute. Following the formalism of [85], we have at tree-level the collinear factorization formula

$$\mathcal{A}_{\text{tree}}(1^{h_1}, 2^{h_2}, 3, \dots, n) \xrightarrow{1||2} \sum_{h=\pm} \text{Split}_{-h}^{\text{tree}}(1^{h_1}, 2^{h_2}) \mathcal{A}_{\text{tree}}(K^h, 3, \dots, n) \quad (96)$$

where $\mathcal{A}_{\text{tree}}$ is a primitive amplitude (which at tree-level is identical to a so called *partial amplitude* and therefore directly associated with a color structure), and the adjacent (incoming) momenta k_1, k_2 become collinear with $k_1 + k_2 \rightarrow K$. The superscripts h indicate helicities. The splittings are typically represented as a function of the splitting variable z in

$$k_1 \rightarrow zK \quad , \quad k_2 \rightarrow (1-z)K . \quad (97)$$

For our situation, we have $K = -\xi_{\text{in}}P$ and $z = \Lambda/\xi_{\text{in}}$, so our splitting variable is not bounded by 1. Therefore, it is more convenient to use the representation of the splitting functions in Appendix C in terms of more general momentum fractions x and y such that

$$k_1 \rightarrow xK \quad , \quad k_2 \rightarrow yK . \quad (98)$$

Inserting into those formulas

$$x \leftarrow -\Lambda \quad \text{and} \quad y \leftarrow (\Lambda - \xi_{\text{in}}) , \quad (99)$$

one can easily see that only the ones for which the auxiliary helicities are opposite survive for large Λ , and that they become $\Lambda/(\xi_{\text{in}}\kappa)$ and $\Lambda/(\xi_{\text{in}}\kappa^*)$. The quantities κ, κ^* are proportional to $|k_\perp|$, see Appendix A. Thus, we find

$$\begin{aligned} \mathcal{A}_{\text{tree}}((-\Lambda P)^{h_1}, p_\Lambda^{h_2}, 3, \dots, n) &\xrightarrow{|k_\perp| \rightarrow 0} \frac{\Lambda}{\xi_{\text{in}}\kappa} \mathcal{A}_{\text{tree}}((-\xi_{\text{in}}P)^+, 3, \dots, n) \\ &+ \frac{\Lambda}{\xi_{\text{in}}\kappa^*} \mathcal{A}_{\text{tree}}((-\xi_{\text{in}}P)^-, 3, \dots, n) + \mathcal{O}(\Lambda^0) , \end{aligned} \quad (100)$$

which indeed is $\Lambda/(\xi_{\text{in}}|k_\perp|)$ times the on-shell limit $|k_\perp| \rightarrow 0$ of the space-like amplitude $\mathcal{A}_{\text{tree}}^*(-\xi_{\text{in}}P - k_\perp, 3, \dots, n)$, see for example Section 5 in [82].

The one-loop factorization formula is a bit more complicated, and following [85] it is given by

$$\begin{aligned} \mathcal{A}_{\text{loop}}(1^{h_1}, 2^{h_2}, 3, \dots, n) &\xrightarrow{1||2} \sum_{h=\pm} \left\{ \text{Split}_{-h}^{\text{tree}}(1^{h_1}, 2^{h_2}) \mathcal{A}_{\text{loop}}(K^h, 3, \dots, n) \right. \\ &\left. + \text{Split}_{-h}^{\text{loop}}(1^{h_1}, 2^{h_2}) \mathcal{A}_{\text{tree}}(K^h, 3, \dots, n) \right\} . \end{aligned} \quad (101)$$

Remember that we are looking for contributions of the type of Eq. (94), which are clearly absent in the first term within curly brackets in Eq. (101): they are not in the tree-level splitting function, and the loop amplitude is on-shell and independent of k_\perp . Thus our interest goes to the second contributions. The one-loop splitting functions are given by

$$\text{Split}_{-h}^{\text{loop}}(1^{h_1}, 2^{h_2}) = g_s^2 c_\Gamma \times \text{Split}_{-h}^{\text{tree}}(1^{h_1}, 2^{h_2}) \times \mathcal{V}(-h, 1^{h_1}, 2^{h_2}), \quad (102)$$

with

$$g_s^2 c_\Gamma = \frac{g_s^2}{(4\pi)^{2-\epsilon}} \frac{\Gamma(1+\epsilon)\Gamma^2(1-\epsilon)}{\Gamma(1-2\epsilon)} = \frac{\alpha_\epsilon}{2} + \mathcal{O}(\epsilon^3). \quad (103)$$

The opposite-helicity \mathcal{V} functions can be found in [85] to be

$$\begin{aligned} \mathcal{V}_{g \rightarrow \bar{q}_1 q_2}(h, 1^\pm, 2^\mp) = & \left(\frac{\mu^2}{-s_{12}} \right)^\epsilon \left\{ \frac{1}{\epsilon} \left(\ln[z(1-z)] + \frac{13}{6} \right) - \frac{1}{2} \ln^2 \frac{z}{1-z} - \frac{\pi^2}{6} + \frac{83}{18} - \frac{\delta_R}{6} \right. \\ & \left. + \frac{1}{N_c^2} \left[\frac{1}{\epsilon^2} + \frac{3}{2\epsilon} + \frac{7+\delta_R}{2} \right] - \frac{n_f}{N_c} \left[\frac{2}{3\epsilon} + \frac{10}{9} \right] \right\} + \mathcal{O}(\epsilon) \quad (104) \end{aligned}$$

with $\delta_R = 1$ in the 't Hooft-Veltman scheme of dimensional regularization and $\delta_R = 0$ for dimensional reduction, and

$$\mathcal{V}_{g \rightarrow g_1 g_2}(h, 1^\pm, 2^\mp) = \left(\frac{\mu^2}{-s_{12}} \right)^\epsilon \left\{ -\frac{1}{\epsilon^2} + \frac{1}{\epsilon} \ln[z(1-z)] - \frac{1}{2} \ln^2 \frac{z}{1-z} - \frac{\pi^2}{6} \right\} + \mathcal{O}(\epsilon). \quad (105)$$

The invariant $s_{12} = (k_1 + k_2)^2$ is to be taken before the collinear limit of course. Since these functions do not actually depend on the values of the helicities, and we saw that at tree-level the collinear limit commutes with the large Λ limit, we are tempted to identify the unfamiliar part of the one-loop amplitude as

$$\mathcal{A}_{\text{loop}}^{\star \text{unf}}(-\xi_{\text{in}} \mathbf{P} - \mathbf{k}_\perp, 3, \dots, n) = \frac{\alpha_\epsilon}{2} \mathcal{V}_{\text{aux}} \mathcal{A}_{\text{tree}}^{\star}(-\xi_{\text{in}} \mathbf{P} - \mathbf{k}_\perp, 3, \dots, n) \quad (106)$$

where we obtain the expressions for \mathcal{V}_{aux} in Eq. (35) and Eq. (36) by just inserting $z \leftarrow \Lambda/\xi_{\text{in}}$ into Eq. (104) and Eq. (105). Realize that $\Lambda/\xi_{\text{in}} > 1$ *does* generate imaginary parts in the logarithms.

Simply replacing the on-shell tree-level amplitude with the space-like one to arrive at Eq. (106) seems a bit sketchy and we will give it more solid ground in Section 4.1. First however, we want to address color. In the beginning of Section 4.5 in [85] it is pointed out that *it is only the leading partial amplitude, $A_{5;1}$ which has the same color structure as the tree-level amplitude, that receives splitting contributions from $\text{Split}^{\text{loop}}$* . Therefore, color can be treated the same as at tree-level, and the formula for the unfamiliar contribution can directly be lifted to virtual contribution at cross section level. This includes a factor N_c coming with the leading partial amplitudes (see Equation (2.5-6) and (2.10) in [85]), and the factor of 2 for the virtual contribution being the interference between tree-level and one-loop. Thus we find

$$dV^{\star \text{unf}} = \alpha_\epsilon N_c \text{Re}(\mathcal{V}_{\text{aux}}) dB^{\star}. \quad (107)$$

Figure 2: Example of the decomposition of a planar one-loop amplitude with 5 external legs into scalar functions times their coefficients. One-point scalar functions are not included, and the decomposition only holds to and including $\mathcal{O}(\epsilon^0)$. The rational terms are remainders that do not contain logarithms of the invariants of the external momenta, but cannot be captured within the scalar function decomposition. Essentially, to each possible cyclicly ordered partition of $(1, 2, 3, 4, 5)$ into 4, 3 or 2 groups belongs a term.

4.1 One-loop amplitudes in the auxiliary parton approach

In this section, we justify using the space-like tree-level amplitude itself instead of its on-shell limit in Eq. (106). This requires a repetition of some technical aspects regarding the calculation of one-loop amplitudes.

The application of the auxiliary parton approach in one-loop amplitudes has been studied in [81], in particular in the context of integrand-based methods to calculate these [86, 87]. One-loop amplitudes can be decomposed in terms of scalar integrals, and for NLO calculations it suffices to include integrals with at most 4 propagator denominators, so-called box integrals of the type

$$\begin{aligned}
 & \begin{array}{c} K_2 \left(\begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array} \right) \left(\begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array} \right) K_3 \\ K_1 \left(\begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array} \right) \left(\begin{array}{c} \vdots \\ \vdots \\ \vdots \end{array} \right) K_4 \end{array} = \frac{(2\pi\mu)^{2\epsilon}}{i(2\pi)^4 c_\Gamma} \int \frac{d^{4-2\epsilon} \ell}{\ell^2 (\ell + K_1)^2 (\ell + K_1 + K_2)^2 (\ell + K_1 + K_2 + K_3)^2} \\
 & \tag{108}
 \end{aligned}$$

where $K_{1,2,3}$ are sums of external momenta, and momentum conservation implies $K_4 = -K_1 - K_2 - K_3$. Realize that these integrals are invariant under shifts $\ell \rightarrow \ell + K$ of the integration momentum. Fewer denominator factors lead to triangle, bubble, and tadpole integrals. These integrals evaluate, within dimensional regularization, to (poly)logarithms of ratios of invariants from the external momenta, while the coefficients for these integrals are “tree-level objects”, that




Figure 3: Scalar one-loop functions contributing to the limit when 1 and 2 become collinear.

is ratios of contractions of momenta and polarization vectors. The list of possible scalar integrals can simply be deduced from the possible kinematical channels implied by the amplitude, and the integrand-based methods extract the coefficients from the one-loop integrand. The latter is the function of the loop momentum ℓ that needs to be integrated over in order to arrive at the one-loop amplitude:

$$\begin{array}{c} 1 \\ \diagup \\ \bigcirc \\ \diagdown \\ n \end{array} \begin{array}{c} 2 \\ \diagdown \\ \bigcirc \\ \diagup \\ n-1 \end{array} \begin{array}{c} 3 \\ \vdots \\ \vdots \end{array} = \int d^{4-2\epsilon} \ell \frac{\mathcal{N}(\ell)}{\ell^2(\ell + p_1)^2(\ell + p_1 + p_2)^2 \cdots (\ell + p_1 + p_2 + \cdots + p_{n-1})}. \quad (109)$$

We wrote the integrand here as a numerator polynomial $\mathcal{N}(\ell)$ divided by all possible ℓ -dependent denominator factors for a planar one-loop amplitude. The integrand-based methods use carefully chosen explicit values of ℓ to extract numerical values of the coefficients for the scalar integrals from the integrand. These methods silently assume that the integrand is calculated in the Feynman gauge in order to avoid other than quadratic denominator factors.

In [84, 85], $\text{Split}^{\text{loop}}$ was calculated by realizing that only contributions from scalar integrals of the type in Fig. 3 need to be taken into account, that is besides the bubble only scalar integrals with the two external lines that are becoming collinear attached to the loop on their own. The box integrals must have at most one off-shell leg. Contributions from box integrals with two off-shell legs that satisfy this criterion do not survive the collinear limit. The fact that the box integrals contribute hints at the main difficulty the authors in the cited work resolved, namely that not only graphs with the factorized form of a one-loop part times a tree-level blob contribute. For these factorized graphs, the prescription to simply replace the limiting on-shell amplitude by the off-shell amplitude in Eq. (106) is obviously correct. In the following we argue that it is also correct for the collection of other graphs.

At first sight, it seems natural to combine the integrand-based methods with the auxiliary parton method by carefully keeping the dependence on Λ when evaluating the scalar functions and possibly encounter logarithms of Λ in those, while taking the limit $\Lambda \rightarrow \infty$ on the integrand, or rather taking the leading power, and extracting the coefficients after this limit. The integrand will, like the tree-level amplitude, be proportional to Λ , and this factor is removed by the (square root of the) prescription of Eq. (22). Realize that this integrand, just like tree-level amplitudes, does not depend on the type of auxiliary partons. In [81] however, it was observed

that in particular the triangle integral

$$\begin{array}{c} \nearrow p_\Lambda \\ \square \\ \nwarrow -\Lambda P \end{array} \left(\begin{array}{c} \vdots \\ \leftarrow k \end{array} \right) = \frac{(2\pi\mu)^{2\epsilon}}{i(2\pi)^4 c_\Gamma} \int d^{4-2\epsilon} \ell \frac{\Lambda}{\ell^2(\ell - \Lambda P)^2(\ell - k)^2} = \frac{-\Lambda}{\epsilon^2 |k_\perp|^2} \left(\frac{\mu^2}{|k_\perp|^2} \right)^\epsilon \quad (110)$$

obstructs this naive approach, because it violates the natural scaling with Λ : while its integrand behaves as $\mathcal{O}(\Lambda^0)$, the integral is proportional to Λ . We apply the convention for scalar functions to include a factor Λ in the numerator of the integrand for every denominator factor that behaves as $\mathcal{O}(\Lambda)$, so Λ effectively acts as a regulator of linear denominator factors. As a result of the ‘‘anomalous’’ behavior of this scalar integral, its correct coefficient cannot be found by taking the leading power in Λ on the one-loop integrand, and the subleading terms need to be included. This then leads to a contribution that depends on the type of auxiliary partons.

Notice that the triangle above is exactly the one appearing in the collinear, or here on-shell, limit. The box integrals are, up to $\mathcal{O}(\Lambda^{-1})$ and $\mathcal{O}(\epsilon)$ given by

$$\begin{array}{c} \nearrow p_\Lambda \\ \square \\ \nwarrow -\Lambda P \end{array} \begin{array}{c} \nearrow p_3 \\ \searrow p_4 \end{array} = \frac{1}{2P \cdot p_4 |k_\perp|^2} \left\{ \left(\frac{\mu^2}{|k_\perp|^2} \right)^\epsilon \left[\frac{4}{\epsilon^2} - \frac{2}{\epsilon} \ln \frac{2\Lambda P \cdot p_4}{|k_\perp|^2} \right] - \pi^2 \right\}, \quad (111)$$

$$\begin{array}{c} \nearrow p_\Lambda \\ \square \\ \nwarrow -\Lambda P \end{array} \begin{array}{c} \nearrow \vdots \\ \searrow p_4 \end{array} \left(\begin{array}{c} \vdots \\ \leftarrow K_3 \end{array} \right) = \frac{1}{2P \cdot p_4 |k_\perp|^2} \left\{ \left(\frac{\mu^2}{|k_\perp|^2} \right)^\epsilon \left[\frac{2}{\epsilon^2} - \frac{2}{\epsilon} \ln \frac{2\Lambda P \cdot p_4}{-K_3^2} \right] \right. \\ \left. + 2\text{Li}_2 \left(1 + \frac{|k_\perp|^2}{K_3^2} \right) - \frac{2\pi^2}{3} \right\}. \quad (112)$$

We use the convention that momenta indicated by the letter p are light-like. Also, $P \cdot p_4$ must be imagined to have a small negative imaginary part, while K_3^2 has a small positive imaginary part. The formulas have the necessary factor $(\mu^2/|k_\perp|^2)^\epsilon$. In fact, the only other scalar function that does, besides the bubble, is

$$\begin{array}{c} \nearrow p_\Lambda \\ \square \\ \nwarrow -\Lambda P \end{array} \begin{array}{c} \nearrow \vdots \\ \searrow \vdots \end{array} \left(\begin{array}{c} \vdots \\ \leftarrow K_3 \end{array} \right) \left(\begin{array}{c} \vdots \\ \leftarrow K_4 \end{array} \right) = \frac{1}{2P \cdot K_4 |k_\perp|^2} \left\{ \left(\frac{\mu^2}{|k_\perp|^2} \right)^\epsilon \left[\frac{1}{\epsilon^2} - \frac{2}{\epsilon} \ln \frac{\Lambda P \cdot K_4}{\mu^2} - \frac{1}{\epsilon} \ln \frac{\mu^4}{K_3^2 K_4^2} \right] \right. \\ \left. - \frac{1}{2} \ln^2 \frac{K_3^2}{K_4^2} - \frac{2\pi^2}{3} \right\}. \quad (113)$$

As mentioned before, the contribution to the one-loop amplitude of this function times its coefficient is irrelevant in the limit $|k_\perp| \rightarrow 0$, but we will need it in the following.

As can be seen, the box functions do scale correctly with Λ and their coefficients can be extracted from the one-loop integrand after the limit $\Lambda \rightarrow \infty$. Consequently, the result will

not depend on the type of auxiliary partons, but we still need to include part of it in what we call unfamiliar contribution, both because they do contribute to the non-smooth on-shell limit, and because they carry $\ln \Lambda$. Now it is important to note that for any of these functions (and realize that K_3, K_4 are sums of external momenta and there is a function for each combination) the relevant part regarding $\ln \Lambda$ is almost the same, namely

$$\frac{-1}{2P \cdot K_4 |k_\perp|^2} \left(\frac{\mu^2}{|k_\perp|^2} \right)^\epsilon \frac{2 \ln \Lambda}{\epsilon}. \quad (114)$$

Only the factor $1/P \cdot K_4$ (or $1/P \cdot p_4$) varies.

The calculation of the coefficients of these box functions has been addressed in Section 5.1 of [81]. Thanks to momentum conservation they can be uniquely labeled by the momentum K_4 . The pair of specific choices of the loop momentum to extract the coefficients are ℓ_1, ℓ_2 , satisfying the conditions that

$$\ell_{1,2}^2 = P \cdot \ell_{1,2} = (\ell_{1,2} - k)^2 = (\ell_{1,2} - K_4)^2 = 0. \quad (115)$$

Here, the momentum $k = \xi_{\text{in}} P + k_\perp$ has positive energy, so it differs by a minus sign with [81]. The coefficients are given by

$$C(K_4) = \frac{1}{2} [\text{Res}(K_4; \ell_1) + \text{Res}(K_4; \ell_2)], \quad (116)$$

with

$$\text{Res}(K_4; \ell_{1,2}) = -\Lambda |k_\perp|^2 \sum_{h=-,+} \text{Diagram} \quad (117)$$

The blobs are on-shell tree-level amplitudes, and the sum is over the helicities of the “internal” gluon. The expression in [81] has Λ^2 but here we follow the convention to include one factor Λ in the box function. The expression above then scales with Λ as expected for a tree-level object.

The reader may have noticed that the above looks like a term in the Britto-Cachazo-Feng (BCF) decomposition of the tree-level amplitude [88, 89]. Decomposing the transverse momentum k_\perp into polarization vectors

$$k^\mu = \xi_{\text{in}} P^\mu + k_\perp^\mu, \quad k_\perp^\mu = -\frac{\kappa}{\sqrt{2}} e_-^\mu - \frac{\kappa^*}{\sqrt{2}} e_+^\mu \quad (118)$$

the specific loop momenta are given by

$$\ell_1^\mu = z_1 P^\mu - \frac{\kappa}{\sqrt{2}} e_-^\mu, \quad \ell_2^\mu = z_2 P^\mu - \frac{\kappa^*}{\sqrt{2}} e_+^\mu \quad (119)$$

where z_1, z_2 are such that $(\ell_{1,2} - K_4)^2 = 0$. So we see that P plays the role of the “shift vector” to accommodate the on-shellness of the internal line. In other words, we have

$$\sum_{K_4} \frac{\text{Res}(K_4; \ell_1)}{\left(\frac{\kappa}{\sqrt{2}} e_- + K_4 \right)^2} = -\Lambda |k_\perp|^2 \mathcal{A}_{\text{tree}} \left(\left(\frac{\kappa}{\sqrt{2}} e_- \right)^-, \left(\frac{\kappa^*}{\sqrt{2}} e_+ - \xi_{\text{in}} P \right)^+, 3, \dots, n \right) \quad (120)$$

where the sum is over all possibilities of

$$K_4 = p_n \quad , \quad K_4 = p_n + p_{n-1} \quad , \dots \quad (121)$$

With ℓ_2 the amplitude with e_+ is obtained instead. Realize that these are the momenta, and that P plays the role of “polarization vector”.

We are however not interested in this result. Rather, we are interested in the sum on the left-hand side of Eq. (120), but instead with that denominator, we want the terms to be multiplied with (114). And indeed, it turns out that

$$\sum_{K_4} \frac{\text{Res}(K_4; \ell_1)}{2P \cdot K_4 |k_\perp|^2} = -\Lambda \mathcal{A}_{\text{tree}}^* (-\xi_{\text{in}} P - k_\perp, 3, \dots, n) . \quad (122)$$

So far, we were not able to prove this remarkable result, but we were able to confirm it numerically for amplitudes with up to a several number of external gluons. Incidentally, the result offers an alternative way to compute tree-level amplitudes with a space-like gluon. Realize that this formula is not to be interpreted recursively. The blobs in Eq. (117) are “normal” on-shell tree-level amplitudes.

Both ℓ_1 and ℓ_2 give the same result, so within the integrand-based framework, the so-called “tilde coefficients” of the box function vanish, and the triangle coefficient can be calculated directly from the “triple cut”. These then relatively straightforwardly lead to the space-like tree-level amplitude again. So eventually, these findings support the prescription to replace the on-shell limit with the space-like tree-level amplitude leading to Eq. (106).

5 Familiar virtual contribution

The divergent structure of on-shell one-loop QCD amplitudes is universal and well known [90]. As is common when presenting this structure, we consider the UV-subtracted contribution. It allows us to write down the complete divergent contribution, that is the sum of familiar and unfamiliar. We will, however, be able to distinguish the two.

If we stick to the terms relevant to auxiliary partons, we have for the divergent virtual contribution in case of auxiliary quarks, explicitly denoted q and \bar{q} , the following expression times α_ϵ :

$$\begin{aligned} & -2 \frac{C_F}{e^2} |\overline{\mathcal{M}}_{\text{aux-}q}^*|^2 \\ & + \frac{2}{\epsilon} \left\{ \ln \left(\frac{\mu^2}{|k_\perp|^2} \right) (\overline{\mathcal{M}}_{\text{aux-}q}^*)_{q\bar{q}}^2 + \sum_{i \neq q, \bar{q}} \left[\ln \left(\frac{\mu^2}{2\Lambda P \cdot p_i} \right) (\overline{\mathcal{M}}_{\text{aux-}q}^*)_{iq}^2 + \ln \left(\frac{\mu^2}{-2\Lambda P \cdot p_i} \right) (\overline{\mathcal{M}}_{\text{aux-}q}^*)_{i\bar{q}}^2 \right] \right\} \\ & - 2 \frac{3C_F}{2e} |\overline{\mathcal{M}}_{\text{aux-}q}^*|^2 . \end{aligned} \quad (123)$$

The squared matrix elements involve a space-like gluon, but carry the auxiliary label because they are still in the auxiliary quark color representation. The large- Λ limit and the division be

C_{aux} are assumed to be already performed. Eventually, the real part must be taken for the virtual contribution, but we want to keep the imaginary part explicit here for later comparison. The first line contains the two soft-collinear terms associated with the auxiliary quarks. The second line contains the other soft terms associated with those. The sums are over external partons. Realize that in literature these sums are typically written over double indices, say i and j , with $i \neq j$ rather than $i > j$, explaining the factor 2 above. Allowing for a slight abuse of notation incorrectly suggesting positivity, $(\overline{\mathcal{M}}_{\text{aux-q}}^*)_{ij}^2$ is the color-correlated squared matrix element involving external partons i and j . The last line represents the two collinear terms associated with the auxiliary quarks. In Appendix B we calculate

$$|\overline{\mathcal{M}}_{\text{aux-q}}^*|^2 = |\overline{\mathcal{M}}^*|^2 \quad , \quad (\overline{\mathcal{M}}_{\text{aux-q}}^*)_{q\bar{q}}^2 = \frac{1}{2N_c} |\overline{\mathcal{M}}^*|^2 \quad , \quad (124)$$

and

$$(\overline{\mathcal{M}}_{\text{aux-q}}^*)_{i\bar{q}}^2 + (\overline{\mathcal{M}}_{\text{aux-q}}^*)_{i\bar{q}}^2 = (\overline{\mathcal{M}}^*)_{i\star}^2 \quad . \quad (125)$$

where the latter is the correlator involving the space-like gluon, independent of the auxiliary color representation. Using color conservation, we have

$$\sum_{i \neq q, \bar{q}} (\overline{\mathcal{M}}_{\text{aux-q}}^*)_{i\bar{q}}^2 = -(\overline{\mathcal{M}}_{\text{aux-q}}^*)_{q\bar{q}}^2 - (\overline{\mathcal{M}}_{\text{aux-q}}^*)_{\bar{q}q}^2 = -\left(\frac{1}{2N_c} + C_F\right) |\overline{\mathcal{M}}^*|^2 = -\frac{N_c}{2} |\overline{\mathcal{M}}^*|^2 \quad (126)$$

and the same for $q \leftrightarrow \bar{q}$. So we have the divergent virtual contribution

$$\begin{aligned} & -2 \frac{C_F}{\epsilon^2} |\overline{\mathcal{M}}^*|^2 \\ & + \frac{1}{N_c \epsilon} \ln\left(\frac{\mu^2}{|k_\perp|^2}\right) |\overline{\mathcal{M}}^*|^2 + \frac{N_c}{\epsilon} \left[2 \ln\left(\frac{\Lambda}{\xi_{\text{in}}}\right) - i\pi\right] |\overline{\mathcal{M}}^*|^2 + \frac{2}{\epsilon} \sum_{i \neq \star} \ln\left(\frac{\mu^2}{2\xi_{\text{in}} \mathbf{P} \cdot \mathbf{p}_i}\right) (\overline{\mathcal{M}}^*)_{i\star}^2 \\ & - 3 \frac{C_F}{\epsilon} |\overline{\mathcal{M}}^*|^2 \quad . \end{aligned} \quad (127)$$

We introduced ξ_{in} here to make sure that the combination $\xi_{\text{in}} \mathbf{P}$ appears rather than \mathbf{P} on its own. After some reorganization, we can write

$$\begin{aligned} & -\frac{C_A}{\epsilon^2} |\overline{\mathcal{M}}^*|^2 + \frac{2}{\epsilon} \sum_{i \neq \star} \ln\left(\frac{\mu^2}{2\xi_{\text{in}} \mathbf{P} \cdot \mathbf{p}_i}\right) (\overline{\mathcal{M}}^*)_{i\star}^2 - \frac{11N_c - 2n_f}{6\epsilon} |\overline{\mathcal{M}}^*|^2 \\ & + \frac{N_c}{\epsilon} \left\{ \frac{1}{N_c^2} \left[\frac{1}{\epsilon} + \ln\left(\frac{\mu^2}{|k_\perp|^2}\right) \right] + 2 \ln\left(\frac{\Lambda}{\xi_{\text{in}}}\right) - i\pi + \frac{1}{3} + \frac{3}{2N_c^2} - \frac{n_f}{3N_c} \right\} |\overline{\mathcal{M}}^*|^2 \quad . \end{aligned} \quad (128)$$

The first line is what we call the divergent familiar virtual contribution related to the space-like gluon. It looks exactly as if the space-like gluon were at the on-shell limit, but with the space-like tree-level squared matrix element. The second line is the unfamiliar contribution, and corresponds to the divergent terms in Eq. (34) and Eq. (35). Realize that those results are UV un-subtracted, while the results above are, so we need to add $(11N_c - 2n_f)/(6\epsilon) |\overline{\mathcal{M}}^*|^2$ back to the result here in order to match.

In case of auxiliary gluons, labelled A and B, we have α_ϵ times

$$\begin{aligned}
& -2 \frac{C_A}{\epsilon^2} |\overline{\mathcal{M}}_{\text{aux-g}}^*|^2 \\
& + \frac{2}{\epsilon} \left\{ \ln \left(\frac{\mu^2}{|k_\perp|^2} \right) (\overline{\mathcal{M}}_{\text{aux-g}}^*)_{AB}^2 + \sum_{i \neq A, B} \left[\ln \left(\frac{\mu^2}{2\Lambda P \cdot p_i} \right) (\overline{\mathcal{M}}_{\text{aux-g}}^*)_{iA}^2 + \ln \left(\frac{\mu^2}{-2\Lambda P \cdot p_i} \right) (\overline{\mathcal{M}}_{\text{aux-g}}^*)_{iB}^2 \right] \right\} \\
& - 2 \frac{11N_c - 2n_f}{6\epsilon} |\overline{\mathcal{M}}_{\text{aux-g}}^*|^2. \tag{129}
\end{aligned}$$

In Appendix B we calculate

$$|\overline{\mathcal{M}}_{\text{aux-g}}^*|^2 = |\overline{\mathcal{M}}^*|^2, \quad (\overline{\mathcal{M}}_{\text{aux-g}}^*)_{AB}^2 = -\frac{N_c}{2} |\overline{\mathcal{M}}^*|^2, \tag{130}$$

and

$$(\overline{\mathcal{M}}_{\text{aux-g}}^*)_{iA}^2 + (\overline{\mathcal{M}}_{\text{aux-g}}^*)_{iB}^2 = (\overline{\mathcal{M}}^*)_{i\star}^2. \tag{131}$$

Realize again that we assume the squared matrix elements to be divided by C_{aux} already. Using color conservation, we now get

$$\sum_{i \neq A, B} (\overline{\mathcal{M}}_{\text{aux-g}}^*)_{iB}^2 = -\left(-\frac{N_c}{2} + C_A\right) |\overline{\mathcal{M}}^*|^2 = -\frac{N_c}{2} |\overline{\mathcal{M}}^*|^2, \tag{132}$$

and the divergent contribution becomes

$$\begin{aligned}
& -2 \frac{C_A}{\epsilon^2} |\overline{\mathcal{M}}^*|^2 \\
& - \frac{N_c}{\epsilon} \ln \left(\frac{\mu^2}{|k_\perp|^2} \right) |\overline{\mathcal{M}}^*|^2 + \frac{N_c}{\epsilon} \left[2 \ln \left(\frac{\Lambda}{\xi_{\text{in}}} \right) - i\pi \right] |\overline{\mathcal{M}}^*|^2 + \frac{2}{\epsilon} \sum_{i \neq \star} \ln \left(\frac{\mu^2}{2\xi_{\text{in}} P \cdot p_i} \right) (\overline{\mathcal{M}}^*)_{i\star}^2 \\
& - \frac{11N_c - 2n_f}{3\epsilon} |\overline{\mathcal{M}}^*|^2, \tag{133}
\end{aligned}$$

and after reorganization

$$\begin{aligned}
& -\frac{C_A}{\epsilon^2} |\overline{\mathcal{M}}^*|^2 + \frac{2}{\epsilon} \sum_{i \neq \star} \ln \left(\frac{\mu^2}{2\xi_{\text{in}} P \cdot p_i} \right) (\overline{\mathcal{M}}^*)_{i\star}^2 - \frac{11N_c - 2n_f}{6\epsilon} |\overline{\mathcal{M}}^*|^2 \\
& - \frac{N_c}{\epsilon} \left\{ \frac{1}{\epsilon} + \ln \left(\frac{\mu^2}{|k_\perp|^2} \right) - 2 \ln \left(\frac{\Lambda}{\xi_{\text{in}}} \right) + i\pi \right\} |\overline{\mathcal{M}}^*|^2 - \frac{11N_c - 2n_f}{6\epsilon} |\overline{\mathcal{M}}^*|^2. \tag{134}
\end{aligned}$$

We see the same structure as with the auxiliary quarks, with the second line now matching the divergent part of Eq. (34) and Eq. (36) after adding $(11N_c - 2n_f)/(6\epsilon) |\overline{\mathcal{M}}^*|^2$.

6 Familiar real contribution

The familiar real contribution becomes divergent when a radiative gluon becomes soft or collinear to another parton. The soft limit of a gluon in QCD squared matrix elements is universal and

well-known [91]. Furthermore, this limit commutes with the large- Λ limit. Terms related to the auxiliary partons, let us take quarks, when a final-state gluon with momentum r becomes soft are given by

$$\frac{(\mathbf{P} \cdot \mathbf{p}_i)}{(\mathbf{P} \cdot \mathbf{r})(\mathbf{r} \cdot \mathbf{p}_i)} (\overline{\mathcal{M}}_{\text{aux-q}}^*)_{iq}^2 + \frac{(\mathbf{P} \cdot \mathbf{p}_i)}{(\mathbf{P} \cdot \mathbf{r})(\mathbf{r} \cdot \mathbf{p}_i)} (\overline{\mathcal{M}}_{\text{aux-q}}^*)_{i\bar{q}}^2 \quad (135)$$

which clearly follows from both the large- Λ limit on the usual on-shell soft expression, and from taking the soft limit on the amplitude after the large- Λ limit constructed with eikonal Feynman rules. Using the relations for the color-correlated squared matrix elements from the previous section again, we find

$$\frac{(\xi_{\text{in}} \mathbf{P} \cdot \mathbf{p}_i)}{(\xi_{\text{in}} \mathbf{P} \cdot \mathbf{r})(\mathbf{r} \cdot \mathbf{p}_i)} (\overline{\mathcal{M}}^*)_{i\star}^2, \quad (136)$$

that is again the pattern that the space-like gluon should be treated as if it were at the on-shell limit, but with the space-like LO squared matrix element.

Besides the other usual soft and collinear singularities, there is also a singularity when a radiative gluon becomes collinear to the momentum P . Appendix C.1 we derive the limit

$$|\overline{\mathcal{M}}^*|^2(\xi_{\text{in}} \mathbf{P} + \mathbf{k}_\perp, \mathbf{k}_{\overline{\text{in}}}; \mathbf{r}, \{\mathbf{p}_i\}_{i=1}^n) \xrightarrow{r \rightarrow \xi_r P} \frac{2N_c}{\mathbf{P} \cdot \mathbf{r}} \frac{\xi_{\text{in}}^2}{\xi_r (\xi_{\text{in}} - \xi_r)^2} |\overline{\mathcal{M}}^*|^2((\xi_{\text{in}} - \xi_r) \mathbf{P} + \mathbf{k}_\perp, \mathbf{k}_{\overline{\text{in}}}; \{\mathbf{p}_i\}_{i=1}^n). \quad (137)$$

For comparison, the usual collinear limit, for example with the parton on the on-shell side, looks like

$$|\overline{\mathcal{M}}^*|^2(\mathbf{k}_{\text{in}}, \bar{\mathbf{x}}_{\overline{\text{in}}} \bar{\mathbf{P}}; \mathbf{r}, \{\mathbf{p}_i\}_{i=1}^n) \xrightarrow{r \rightarrow \bar{x}_r \bar{\mathbf{P}}} \frac{C_{\overline{\text{in}}}}{\bar{\mathbf{P}} \cdot \mathbf{r}} \frac{\mathcal{P}_{\overline{\text{in}}}(1 - \bar{x}_r / \bar{x}_{\overline{\text{in}}})}{\bar{x}_{\overline{\text{in}}} - \bar{x}_r} |\overline{\mathcal{M}}^*|^2(\mathbf{k}_{\text{in}}, (\bar{\mathbf{x}}_{\overline{\text{in}}} - \bar{x}_r) \bar{\mathbf{P}}; \{\mathbf{p}_i\}_{i=1}^n), \quad (138)$$

where $\mathcal{P}_{\overline{\text{in}}}(z)$ is the non-regularized splitting function associated with the on-shell initial-state parton, with the color factor $C_{\overline{\text{in}}}$ taken out explicitly. The singular behavior leads to the non-cancelling collinear divergence absorbed into the NLO correction to the collinear PDF following the well-know factorization prescription. Here we investigate how the divergence looks like in the space-like case.

It turns out that we can be a bit more ambitious in the definition of the collinear limit, and instead of Eq. (137), we can write down a formula that does not compromise phase space, very much like subtraction terms constructed to render real-radiation phase space integrals finite in NLO calculations. This means, that we need to accommodate for the momentum recoil when the momentum r is not exactly at the collinear limit. Fortunately, we are privileged compared to collinear factorization in that we have initial-state transverse momentum that can be used for the recoil, and we can simply write

$$|\overline{\mathcal{M}}^*|^2(\xi_{\text{in}} \mathbf{P} + \mathbf{k}_\perp, \mathbf{k}_{\overline{\text{in}}}; \mathbf{r}, \{\mathbf{p}_i\}_{i=1}^n) \longrightarrow \frac{2N_c}{\mathbf{P} \cdot \mathbf{r}} \frac{\xi_{\text{in}}^2}{\xi_r (\xi_{\text{in}} - \xi_r)^2} |\overline{\mathcal{M}}^*|^2((\xi_{\text{in}} - \xi_r) \mathbf{P} + \mathbf{k}_\perp - \mathbf{r}_\perp, (\bar{\mathbf{x}}_{\overline{\text{in}}} - \bar{x}_r) \bar{\mathbf{P}}; \{\mathbf{p}_i\}_{i=1}^n). \quad (139)$$

Now, $\{p_i\}_{i=1}^n$ can be seen as identical on the left-hand side and the right-hand side for any momentum r , not only at the limit $r = \xi_r P$. We will make one adjustments to the formula, that will still give the correct limit, namely we will neglect \bar{x}_r on the right-hand side of Eq. (139).

In the following, we will employ integration over initial-state variables, and prefer to write formulas in terms of x and P_A already instead of ξ and P (Section 2.3). We will use the symbol

$$S = 2P_A \cdot \bar{P} \quad (140)$$

but it will appear only multiplied with an x variable. Notice that the factor in front of the squared matrix element on the right-hand side of Eq. (139) is invariant under rescaling of the ξ variables and P . We insert the formula, now in terms of x variables and P_A , into the cross section formula, to get (remember the flux factor)

$$\begin{aligned} & dR_{\text{coll}}^{\text{fam}}(k_{\text{in}}, k_{\bar{\text{in}}}; \{p_i\}_{i=1}^n) \\ &= \frac{a_e}{\pi_e \mu^{\bar{e}}} \frac{1}{|k_{\perp}|^2} \int d^{4+\bar{e}}r \delta(r^2) \theta(x_r < x_{\text{in}}) 2N_c \frac{1}{P_A \cdot r} \frac{x_{\text{in}}^2}{x_r(x_{\text{in}} - x_r)^2} \\ & \quad \times \frac{x_{\text{in}} - x_r}{x_{\text{in}}} d\Sigma_n^*(x_{\text{in}} - x_r, k_{\perp} - r_{\perp}, \bar{x}_{\bar{\text{in}}}; \{p_i\}_{i=1}^n) J_B(\{p_i\}_{i=1}^n) \\ &= \frac{2a_e N_c}{\pi_e \mu^{\bar{e}}} \int_0^{x_{\text{in}}} \frac{dx_r}{x_r} \int \frac{d^{2+\bar{e}}r_{\perp}}{|r_{\perp}|^2} \frac{|k_{\perp} - r_{\perp}|^2}{|k_{\perp}|^2} \theta(|r_{\perp}|^2 < S\bar{x}_{\bar{\text{in}}}x_r) \\ & \quad \times \frac{x_{\text{in}}}{x_{\text{in}} - x_r} dB^*(x_{\text{in}} - x_r, k_{\perp} - r_{\perp}, \bar{x}_{\bar{\text{in}}}; \{p_i\}_{i=1}^n). \quad (141) \end{aligned}$$

Now we put this underneath an integral over x_{in} and k_{\perp} with an extra function $F(x_{\text{in}}, k_{\perp})$

$$\begin{aligned} & \int_0^1 \frac{dx_{\text{in}}}{x_{\text{in}}} \int d^2k_{\perp} F(x_{\text{in}}, k_{\perp}) dR_{\text{coll}}^{\text{fam}}(x_{\text{in}}P_A + k_{\perp}, k_{\bar{\text{in}}}; \{p_i\}_{i=1}^n) \\ &= \frac{2a_e N_c}{\pi_e \mu^{\bar{e}}} \int_0^1 \frac{dy}{y} \int d^2k_{\perp} \int_y^1 \frac{dz}{z(1-z)} \int \frac{d^{2+\bar{e}}r_{\perp}}{|r_{\perp}|^2} \frac{|k_{\perp}|^2}{|k_{\perp} + r_{\perp}|^2} \theta\left(|r_{\perp}|^2 < \frac{S\bar{x}_{\bar{\text{in}}}y(1-z)}{z}\right) \\ & \quad \times F\left(\frac{y}{z}, k_{\perp} + r_{\perp}\right) dB^*(y, k_{\perp}, \bar{x}_{\bar{\text{in}}}; \{p_i\}_{i=1}^n) \\ &= \int_0^1 \frac{dx_{\text{in}}}{x_{\text{in}}} \int d^2k_{\perp} \tilde{F}(x_{\text{in}}, k_{\perp}) dB^*(x_{\text{in}}, k_{\perp}, \bar{x}_{\bar{\text{in}}}; \{p_i\}_{i=1}^n) \quad (142) \end{aligned}$$

with

$$\begin{aligned} \tilde{F}(x_{\text{in}}, k_{\perp}) &= \frac{2a_e N_c}{\pi_e \mu^{\bar{e}}} \int_{x_{\text{in}}}^1 \frac{dz}{z(1-z)} \int \frac{d^{2+\bar{e}}r_{\perp}}{|r_{\perp}|^2} \frac{|k_{\perp}|^2}{|k_{\perp} + r_{\perp}|^2} \theta\left(|r_{\perp}|^2 < \frac{S\bar{x}_{\bar{\text{in}}}x_{\text{in}}(1-z)}{z}\right) \\ & \quad \times F\left(\frac{x_{\text{in}}}{z}, k_{\perp} + r_{\perp}\right). \quad (143) \end{aligned}$$

Now we can compare this formula with Equation (2.21) in [79], which was interpreted in the context of multi-Regge kinematics evolution. The factor $|k_{\perp}|^2/|k_{\perp} + r_{\perp}|^2$ is missing in [79], but we could also get rid of it by re-defining $F(x_{\text{in}}, k_{\perp}) \rightarrow |k_{\perp}|^2 F(x_{\text{in}}, k_{\perp})$. Notice that neglecting \bar{x}_r

on the right-hand side of Eq. (143) is equivalent to crossing out q_1^+ in Equation (2.17) in [79]. In the following, we prefer to stick to the equation above, while realizing that

$$F(x, 0) = 0, \quad (144)$$

so there is no singularity at $r_\perp = -k_\perp$.

First of all, we observe that Eq. (143) does not conflict with the demand that x_{in} is small. This demand can be stated as the condition that $F(x, k_\perp) = 0$ for $x > x_{\text{max}}$ and some $x_{\text{max}} \ll 1$. This means that the integral only contributes for $x_{\text{in}}/z < x_{\text{max}}$, that is for $z > x_{\text{in}}/x_{\text{max}}$. Since $z < 1$, this then implies that x_{in} must be smaller than x_{max} for the integral, and thus $\tilde{F}(x_{\text{in}}, k_\perp)$, not to vanish.

Let us change integration variable $z = 1/(1 + y/x_{\text{in}})$ in Eq. (143) to write

$$\tilde{F}(x_{\text{in}}, k_\perp) = \frac{2a_\epsilon N_c}{\pi_\epsilon \mu^{\bar{\epsilon}}} \int_0^{1-x_{\text{in}}} \frac{dy}{y} \int_0^{S\bar{x}_{\text{in}}y} d^{2+\bar{\epsilon}} r_\perp \frac{|k_\perp|^2}{|r_\perp|^2 |r_\perp + k_\perp|^2} F(x_{\text{in}} + y, k_\perp + r_\perp), \quad (145)$$

where we introduced the notation

$$\int_0^U d^{2+\bar{\epsilon}} r_\perp \equiv \int d^{2+\bar{\epsilon}} r_\perp \theta(|r_\perp|^2 < U). \quad (146)$$

Realizing that $y \rightarrow 0$ implies $|r_\perp| \rightarrow 0$, as can be seen by substituting $r_\perp \rightarrow \sqrt{y} r_\perp$ so y disappears from the integration limit for r_\perp , one could imagine that the $1/\bar{\epsilon}$ divergencies can be isolated via subtraction as

$$\begin{aligned} \tilde{F}(x_{\text{in}}, k_\perp) &= \frac{2a_\epsilon N_c}{\pi_\epsilon \mu^{\bar{\epsilon}}} \int_0^{1-x_{\text{in}}} \frac{dy}{y} \int_0^{S\bar{x}_{\text{in}}y} d^{2+\bar{\epsilon}} r_\perp \frac{|k_\perp|^2}{|r_\perp|^2 |r_\perp + k_\perp|^2} [F(x_{\text{in}} + y, k_\perp + r_\perp) - F(x_{\text{in}}, k_\perp)] \\ &+ F(x_{\text{in}}, k_\perp) \frac{2a_\epsilon N_c}{\pi_\epsilon \mu^{\bar{\epsilon}}} \int_0^{1-x_{\text{in}}} \frac{dy}{y} \int_0^{S\bar{x}_{\text{in}}y} d^{2+\bar{\epsilon}} r_\perp \frac{|k_\perp|^2}{|r_\perp|^2 |r_\perp + k_\perp|^2}. \end{aligned} \quad (147)$$

This, however, does not work because the integral on the first line above still contains a $1/\bar{\epsilon}$ divergence due to the now unprotected singularity at $r_\perp = -k_\perp$. The second integral *does* contain the complete $1/\bar{\epsilon}^2$ contribution, and this exercise was still useful because the reader may have noticed that this integral looks exactly like Eq. (73). Realize that there, despite having dedicated symbols, in fact $\mathcal{P}_{\text{aux}}^{\text{pole}} = 2$ and $c_{\bar{q}} = c_q = N_c$. In order to equate the contributions, we must only assume that $S = \Lambda v^2$, implying that the relation between ξ and P versus x and P_A in Eq. (26) is set by $\lambda = \Lambda$. Then we find that the correct value of η in Section 3.1 is the upper limit for the integral here, so $\eta = 1 - x_{\text{in}}$.

In order to isolate all $1/\bar{\epsilon}$ divergencies properly, we add and subtract identical terms in

Eq. (143) instead, and write

$$\begin{aligned}
\tilde{F}(x_{\text{in}}, k_{\perp}) &= \frac{2a_{\epsilon} N_c}{\pi_{\epsilon} \mu^{\bar{\epsilon}}} \int_{x_{\text{in}}}^1 \frac{dz}{z(1-z)} \int_0^{S\bar{x}_{\text{in}}x_{\text{in}}(1-z)/z} d^{2+\bar{\epsilon}} r_{\perp} \frac{|k_{\perp}|^2}{|r_{\perp}|^2 |r_{\perp} + k_{\perp}|^2} \\
&\quad \times \left[F\left(\frac{x_{\text{in}}}{z}, k_{\perp} + r_{\perp}\right) - \frac{|r_{\perp} + k_{\perp}|^2}{|k_{\perp}|^2} F\left(\frac{x_{\text{in}}}{z}, k_{\perp}\right) \right] \\
&+ \frac{2a_{\epsilon} N_c}{\pi_{\epsilon} \mu^{\bar{\epsilon}}} \int_{x_{\text{in}}}^1 \frac{dz}{z(1-z)} \int_0^{S\bar{x}_{\text{in}}x_{\text{in}}(1-z)/z} d^{2+\bar{\epsilon}} r_{\perp} \frac{1}{|r_{\perp}|^2} \left[F\left(\frac{x_{\text{in}}}{z}, k_{\perp}\right) - zF(x_{\text{in}}, k_{\perp}) \right] \\
&+ \frac{2a_{\epsilon} N_c}{\pi_{\epsilon} \mu^{\bar{\epsilon}}} \int_{x_{\text{in}}}^1 \frac{dz}{1-z} \int_0^{S\bar{x}_{\text{in}}x_{\text{in}}(1-z)/z} d^{2+\bar{\epsilon}} r_{\perp} \frac{1}{|r_{\perp}|^2} F(x_{\text{in}}, k_{\perp}) . \tag{148}
\end{aligned}$$

The first integral is finite for $\bar{\epsilon} \rightarrow 0$. The last integral is proportional the Born result, and given by $F(x_{\text{in}}, k_{\perp})$ times

$$\begin{aligned}
\frac{2a_{\epsilon} N_c}{\pi_{\epsilon} \mu^{\bar{\epsilon}}} \int_{x_{\text{in}}}^1 \frac{dz}{1-z} \int_0^{S\bar{x}_{\text{in}}x_{\text{in}}(1-z)/z} d^{2+\bar{\epsilon}} r_{\perp} \frac{1}{|r_{\perp}|^2} &= 2a_{\epsilon} N_c \left(\frac{S\bar{x}_{\text{in}}x_{\text{in}}(1-x_{\text{in}})}{\mu^2} \right)^{\bar{\epsilon}/2} \frac{4}{\bar{\epsilon}^2} \\
&- 2a_{\epsilon} N_c \text{Li}_2(1-x_{\text{in}}) + \mathcal{O}(\bar{\epsilon}) . \tag{149}
\end{aligned}$$

The divergent contribution is of the soft kind and cancels against the corresponding virtual contribution.

The factor z in front of $F(x_{\text{in}}, k_{\perp})$ in the second integral in Eq. (148) seems arbitrary, but is included so the divergent part of this integral can be written in terms of a plus-distribution, and the integral becomes

$$\begin{aligned}
\frac{4a_{\epsilon} N_c}{\bar{\epsilon}} \left(\frac{S\bar{x}_{\text{in}}x_{\text{in}}(1-x_{\text{in}})}{\mu^2} \right)^{\bar{\epsilon}/2} \int_{x_{\text{in}}}^1 \frac{dz}{[1-z]_+} \frac{1}{z} F\left(\frac{x_{\text{in}}}{z}, k_{\perp}\right) \\
+ 2a_{\epsilon} N_c \int_{x_{\text{in}}}^1 \frac{dz}{1-z} \ln \frac{1-z}{z} \left[\frac{1}{z} F\left(\frac{x_{\text{in}}}{z}, k_{\perp}\right) - F(x_{\text{in}}, k_{\perp}) \right] + \mathcal{O}(\bar{\epsilon}) . \tag{150}
\end{aligned}$$

The divergent part

$$- \frac{2a_{\epsilon} N_c}{\epsilon} \int_{x_{\text{in}}}^1 \frac{dz}{[1-z]_+} \frac{1}{z} F\left(\frac{x_{\text{in}}}{z}, k_{\perp}\right) , \tag{151}$$

now written in terms of $\epsilon = -\bar{\epsilon}/2$ corresponds to Eq. (46).

7 Summary

In the present paper we extended the auxiliary parton method for obtaining the impact factors in high energy factorization to the next-to-leading order. The main results are the promotion of the hybrid factorization formula to NLO, and a calculational scheme that allows to obtain the NLO impact factors from suitable helicity amplitudes with on-shell partons. We also rederived NLO impact factors for the quark and the gluon obtained at first in Ref. [29].

In the derivations presented here a bottom-up approach was taken, based on exact NLO expressions of general collinear partonic amplitudes in QCD in the high energy limit, *i.e.* by taking the leading term in the $1/\Lambda$ power expansion, where the large parameter Λ proportional to the large total energy squared S available in a particle collision. We found that the auxiliary parton method in the large Λ limit and inclusion of NLO corrections do not commute. It means, that the auxiliary parton line cannot be factored out in the high energy limit before it is treated at NLO accuracy. In order to make this point explicit we introduced nomenclature motivated by the structure of the tree level expressions for scattering amplitudes in the high energy limit: contributions with the auxiliary parton treated at the LO approximation were named *familiar*, while NLO contributions to the auxiliary parton impact factors were called *unfamiliar*.

The main guiding principle of the analysis described in the paper was following how the infrared divergencies emerging in the on-shell expressions are organized in the hybrid factorization scheme. To be more specific, we verified that the NLO expressions including the infrared singularities, are consistent with the Regge factorization: the singularities and the NLO corrections may be separated into an impact factor of the target (in our case — the auxiliary parton) and to the remaining contribution coming from the projectile and from the in-between region, corresponding to an evolution kernel absorbing terms enhanced by a large $\ln \Lambda$.

Using this factorization one may proceed further and split off the evolution part from the familiar contribution, and in this way obtain the projectile impact factor from the on-shell amplitudes. In particular, we showed that the familiar contribution obtained in this procedure is universal w.r.t. the auxiliary parton flavor — that is the same for an auxiliary quark and gluon. Hence, the projectile impact factor coming from the proposed scheme is universal as well. This universality holds for virtual and real NLO corrections independently.

We analyzed cancellation of ϵ -poles in the dimensional regularization, coming both from infrared and ultraviolet divergencies. The infrared singularities were found to cancel in the NLO target impact factor and the remaining part of NLO corrections separately when combined with the infrared singularities coming from the initial state parton lines, and the ultraviolet poles cancelled out after performing the renormalization. In the *familiar* part of the amplitudes we found terms corresponding to one step of the evolution of transverse momentum dependent parton distribution function. The real emission part of this evolution equation is given by Eq. (143). This integral kernel closely resembles the LO BFKL kernel with a difference in the longitudinal momentum part: the $1/z$ factor present in the BFKL equation is replaced by $1/(z(1-z))$ in Eq. (143), and the transverse momentum integration is limited in Eq. (143). This difference occurs because we did not isolate the leading $\ln S$ approximation, but kept all the NLO contribution at the leading power of $1/S$. The $1/(1-z)$ factor in the evolution kernel (141) is a manifestation of the soft gluon corrections. Essentially the same equation was found in [79], where it was related to the last emission in a cascade of rapidity-ordered gluons. Of course the present results are still fixed order results and some more work is necessary to obtain the evolution equation that would resum both BFKL and soft gluon logarithms in a rigorous manner.

The results obtained here open a way to systematically extract the NLO impact factors for hybrid factorization from on-shell helicity amplitudes. In fact, the extraction of the virtual NLO

contributions to the impact factors is straightforward and boils down to analytic subtractions of universal virtual terms described above from corresponding NLO virtual corrections to on-shell amplitudes. The computation of the real emission corrections to impact factors still remains to be done for each process separately, using the same scheme that was applied to compute the real radiation corrections to the auxiliary quark and gluon.

Acknowledgments

GZ is supported by grant no. 2019/35/B/ST2/03531 of the Polish National Science Centre. AvH and LM gratefully acknowledge support of the Polish National Science Center (NCN) grant No. 2017/27/B/ST2/02755.

References

- [1] L. N. Lipatov, *Reggeization of the Vector Meson and the Vacuum Singularity in Nonabelian Gauge Theories*, *Sov. J. Nucl. Phys.* **23** (1976) 338–345.
- [2] E. A. Kuraev, L. N. Lipatov, and V. S. Fadin, *Multi - Reggeon Processes in the Yang-Mills Theory*, *Sov. Phys. JETP* **44** (1976) 443–450.
- [3] E. A. Kuraev, L. N. Lipatov, and V. S. Fadin, *The Pomeron Singularity in Nonabelian Gauge Theories*, *Sov. Phys. JETP* **45** (1977) 199–204.
- [4] I. I. Balitsky and L. N. Lipatov, *The Pomeron Singularity in Quantum Chromodynamics*, *Sov. J. Nucl. Phys.* **28** (1978) 822–829.
- [5] V. S. Fadin and L. N. Lipatov, *High-Energy Production of Gluons in a QuasimultiRegge Kinematics*, *JETP Lett.* **49** (1989) 352.
- [6] L. N. Lipatov, *Small x physics in perturbative QCD*, *Phys. Rept.* **286** (1997) 131–198, [hep-ph/9610276].
- [7] S. Catani, M. Ciafaloni, and F. Hautmann, *GLUON CONTRIBUTIONS TO SMALL x HEAVY FLAVOR PRODUCTION*, *Phys. Lett. B* **242** (1990) 97–102.
- [8] S. Catani, M. Ciafaloni, and F. Hautmann, *High-energy factorization and small x heavy flavor production*, *Nucl. Phys. B* **366** (1991) 135–188.
- [9] S. Catani and F. Hautmann, *High-energy factorization and small x deep inelastic scattering beyond leading order*, *Nucl. Phys. B* **427** (1994) 475–524, [hep-ph/9405388].

- [10] V. S. Fadin, M. I. Kotsky, and R. Fiore, *Gluon Reggeization in QCD in the next-to-leading order*, *Phys. Lett. B* **359** (1995) 181–188.
- [11] V. S. Fadin, R. Fiore, and M. I. Kotsky, *Gluon Regge trajectory in the two loop approximation*, *Phys. Lett. B* **387** (1996) 593–602, [hep-ph/9605357].
- [12] V. S. Fadin and L. N. Lipatov, *Radiative corrections to QCD scattering amplitudes in a multi - Regge kinematics*, *Nucl. Phys. B* **406** (1993) 259–292.
- [13] V. S. Fadin, R. Fiore, and A. Quartarolo, *Quark contribution to the reggeon - reggeon - gluon vertex in QCD*, *Phys. Rev. D* **50** (1994) 5893–5901, [hep-th/9405127].
- [14] V. S. Fadin and R. Fiore, *Non-forward NLO BFKL kernel*, *Phys. Rev. D* **72** (2005) 014018, [hep-ph/0502045].
- [15] V. S. Fadin, R. Fiore, and M. I. Kotsky, *Gribov's theorem on soft emission and the reggeon-reggeon - gluon vertex at small transverse momentum*, *Phys. Lett. B* **389** (1996) 737–741, [hep-ph/9608229].
- [16] V. S. Fadin, R. Fiore, A. Flachi, and M. I. Kotsky, *Quark - anti-quark contribution to the BFKL kernel*, *Phys. Lett. B* **422** (1998) 287–293, [hep-ph/9711427].
- [17] V. Del Duca, *Real next-to-leading corrections to the multi - gluon amplitudes in the helicity formalism*, *Phys. Rev. D* **54** (1996) 989–1009, [hep-ph/9601211].
- [18] V. Del Duca and C. R. Schmidt, *Virtual next-to-leading corrections to the impact factors in the high-energy limit*, *Phys. Rev. D* **57** (1998) 4069–4079, [hep-ph/9711309].
- [19] V. Del Duca and C. R. Schmidt, *Virtual next-to-leading corrections to the Lipatov vertex*, *Phys. Rev. D* **59** (1999) 074004, [hep-ph/9810215].
- [20] V. S. Fadin and L. N. Lipatov, *BFKL pomeron in the next-to-leading approximation*, *Phys. Lett. B* **429** (1998) 127–134, [hep-ph/9802290].
- [21] G. Camici and M. Ciafaloni, *Irreducible part of the next-to-leading BFKL kernel*, *Phys. Lett. B* **412** (1997) 396–406, [hep-ph/9707390]. [Erratum: *Phys.Lett.B* 417, 390 (1998)].
- [22] M. Ciafaloni and G. Camici, *Energy scale(s) and next-to-leading BFKL equation*, *Phys. Lett. B* **430** (1998) 349–354, [hep-ph/9803389].
- [23] M. Ciafaloni and D. Colferai, *The BFKL equation at next-to-leading level and beyond*, *Phys. Lett. B* **452** (1999) 372–378, [hep-ph/9812366].
- [24] I. Balitsky and G. A. Chirilli, *Next-to-leading order evolution of color dipoles*, *Phys. Rev. D* **77** (2008) 014019, [0710.4330].

- [25] G. P. Salam, *An Introduction to leading and next-to-leading BFKL*, *Acta Phys. Polon. B* **30** (1999) 3679–3705, [hep-ph/9910492].
- [26] A. V. Kotikov and L. N. Lipatov, *NLO corrections to the BFKL equation in QCD and in supersymmetric gauge theories*, *Nucl. Phys. B* **582** (2000) 19–43, [hep-ph/0004008].
- [27] N. Gromov, F. Levkovich-Maslyuk, and G. Sizov, *Pomeron Eigenvalue at Three Loops in $\mathcal{N} = 4$ Supersymmetric Yang-Mills Theory*, *Phys. Rev. Lett.* **115** (2015), no. 25 251601, [1507.04010].
- [28] S. Caron-Huot and M. Herranen, *High-energy evolution to three loops*, *JHEP* **02** (2018) 058, [1604.07417].
- [29] M. Ciafaloni and D. Colferai, *K factorization and impact factors at next-to-leading level*, *Nucl. Phys. B* **538** (1999) 187–214, [hep-ph/9806350].
- [30] V. S. Fadin, R. Fiore, M. I. Kotsky, and A. Papa, *The Gluon impact factors*, *Phys. Rev. D* **61** (2000) 094005, [hep-ph/9908264].
- [31] V. S. Fadin, R. Fiore, M. I. Kotsky, and A. Papa, *The Quark impact factors*, *Phys. Rev. D* **61** (2000) 094006, [hep-ph/9908265].
- [32] J. Bartels, D. Colferai, and G. P. Vacca, *The NLO jet vertex for Mueller-Navelet and forward jets: The Quark part*, *Eur. Phys. J. C* **24** (2002) 83–99, [hep-ph/0112283].
- [33] J. Bartels, D. Colferai, and G. P. Vacca, *The NLO jet vertex for Mueller-Navelet and forward jets: The Gluon part*, *Eur. Phys. J. C* **29** (2003) 235–249, [hep-ph/0206290].
- [34] M. Hentschinski and A. Sabio Vera, *NLO jet vertex from Lipatov’s QCD effective action*, *Phys. Rev. D* **85** (2012) 056006, [1110.6741].
- [35] F. Caporale, D. Y. Ivanov, B. Murdaca, and A. Papa, *Mueller-Navelet small-cone jets at LHC in next-to-leading BFKL*, *Nucl. Phys. B* **877** (2013) 73–94, [1211.7225].
- [36] G. Chachamis, M. Hentschinski, J. D. Madrigal Martínez, and A. Sabio Vera, *Next-to-leading order corrections to the gluon-induced forward jet vertex from the high energy effective action*, *Phys. Rev. D* **87** (2013), no. 7 076009, [1212.4992].
- [37] M. Hentschinski, J. D. M. Martínez, B. Murdaca, and A. Sabio Vera, *The gluon-induced Mueller–Tang jet impact factor at next-to-leading order*, *Nucl. Phys. B* **889** (2014) 549–579, [1409.6704].
- [38] A. Sabio Vera, *The Effect of NLO conformal spins in azimuthal angle decorrelation of jet pairs*, *Nucl. Phys. B* **746** (2006) 1–14, [hep-ph/0602250].
- [39] D. Colferai, F. Schwennsen, L. Szymanowski, and S. Wallon, *Mueller Navelet jets at LHC - complete NLL BFKL calculation*, *JHEP* **12** (2010) 026, [1002.1365].

- [40] F. Caporale, D. Y. Ivanov, B. Murdaca, A. Papa, and A. Perri, *The next-to-leading order jet vertex for Mueller-Navelet and forward jets revisited*, *JHEP* **02** (2012) 101, [1112.3752].
- [41] D. Y. Ivanov and A. Papa, *The next-to-leading order forward jet vertex in the small-cone approximation*, *JHEP* **05** (2012) 086, [1202.1082].
- [42] B. Ducloue, L. Szymanowski, and S. Wallon, *Confronting Mueller-Navelet jets in NLL BFKL with LHC experiments at 7 TeV*, *JHEP* **05** (2013) 096, [1302.7012].
- [43] B. Ducloué, L. Szymanowski, and S. Wallon, *Evidence for high-energy resummation effects in Mueller-Navelet jets at the LHC*, *Phys. Rev. Lett.* **112** (2014) 082003, [1309.3229].
- [44] F. Caporale, D. Y. Ivanov, B. Murdaca, and A. Papa, *Mueller–Navelet jets in next-to-leading order BFKL: theory versus experiment*, *Eur. Phys. J. C* **74** (2014), no. 10 3084, [1407.8431]. [Erratum: *Eur.Phys.J.C* 75, 535 (2015)].
- [45] D. Y. Ivanov, M. I. Kotsky, and A. Papa, *The Impact factor for the virtual photon to light vector meson transition*, *Eur. Phys. J. C* **38** (2004) 195–213, [hep-ph/0405297].
- [46] D. Y. Ivanov and A. Papa, *Electroproduction of two light vector mesons in the next-to-leading approximation*, *Nucl. Phys. B* **732** (2006) 183–199, [hep-ph/0508162].
- [47] D. Y. Ivanov and A. Papa, *Electroproduction of two light vector mesons in next-to-leading BFKL: Study of systematic effects*, *Eur. Phys. J. C* **49** (2007) 947–955, [hep-ph/0610042].
- [48] R. Boussarie, A. V. Grabovsky, D. Y. Ivanov, L. Szymanowski, and S. Wallon, *Next-to-Leading Order Computation of Exclusive Diffractive Light Vector Meson Production in a Saturation Framework*, *Phys. Rev. Lett.* **119** (2017), no. 7 072002, [1612.08026].
- [49] P. Taels, T. Altinoluk, G. Beuf, and C. Marquet, *Dijet photoproduction at low x at next-to-leading order and its back-to-back limit*, 2204.11650.
- [50] P. Caucal, F. Salazar, and R. Venugopalan, *Dijet impact factor in DIS at next-to-leading order in the Color Glass Condensate*, *JHEP* **11** (2021) 222, [2108.06347].
- [51] R. Boussarie, A. V. Grabovsky, L. Szymanowski, and S. Wallon, *Towards a complete next-to-logarithmic description of forward exclusive diffractive dijet electroproduction at HERA: real corrections*, *Phys. Rev. D* **100** (2019), no. 7 074020, [1905.07371].
- [52] R. Boussarie, A. V. Grabovsky, L. Szymanowski, and S. Wallon, *On the one loop $\gamma^{(*)} \rightarrow q\bar{q}$ impact factor and the exclusive diffractive cross sections for the production of two or three jets*, *JHEP* **11** (2016) 149, [1606.00419].

- [53] K. Roy and R. Venugopalan, *NLO impact factor for inclusive photon+dijet production in $e + A$ DIS at small x* , *Phys. Rev. D* **101** (2020), no. 3 034028, [1911.04530].
- [54] J. Bartels, D. Colferai, S. Gieseke, and A. Kyrieleis, *NLO corrections to the photon impact factor: Combining real and virtual corrections*, *Phys. Rev. D* **66** (2002) 094017, [hep-ph/0208130].
- [55] V. S. Fadin, D. Y. Ivanov, and M. I. Kotsky, *On the calculation of the NLO virtual photon impact factor*, *Nucl. Phys. B* **658** (2003) 156–174, [hep-ph/0210406].
- [56] I. Balitsky and G. A. Chirilli, *Photon impact factor in the next-to-leading order*, *Phys. Rev. D* **83** (2011) 031502, [1009.4729].
- [57] I. Balitsky and G. A. Chirilli, *Photon impact factor and k_T -factorization for DIS in the next-to-leading order*, *Phys. Rev. D* **87** (2013), no. 1 014013, [1207.3844].
- [58] G. Beuf, *Dipole factorization for DIS at NLO: Combining the $q\bar{q}$ and $q\bar{q}g$ contributions*, *Phys. Rev. D* **96** (2017), no. 7 074033, [1708.06557].
- [59] H. Hänninen, T. Lappi, and R. Paatelainen, *One-loop corrections to light cone wave functions: the dipole picture DIS cross section*, *Annals Phys.* **393** (2018) 358–412, [1711.08207].
- [60] D. Y. Ivanov, B. Murdaca, and A. Papa, *The $\gamma^*\gamma^*$ total cross section in next-to-leading order BFKL and LEP2 data*, *JHEP* **10** (2014) 058, [1407.8447].
- [61] M. Ciafaloni and G. Rodrigo, *Heavy quark impact factor at next-to-leading level*, *JHEP* **05** (2000) 042, [hep-ph/0004033].
- [62] F. G. Celiberto, D. Y. Ivanov, B. Murdaca, and A. Papa, *High-energy resummation in heavy-quark pair photoproduction*, *Phys. Lett. B* **777** (2018) 141–150, [1709.10032].
- [63] A. D. Bolognino, F. G. Celiberto, M. Fucilla, D. Y. Ivanov, and A. Papa, *High-energy resummation in heavy-quark pair hadroproduction*, *Eur. Phys. J. C* **79** (2019), no. 11 939, [1909.03068].
- [64] F. G. Celiberto, D. Y. Ivanov, B. Murdaca, and A. Papa, *High energy resummation in dihadron production at the LHC*, *Phys. Rev. D* **94** (2016), no. 3 034013, [1604.08013].
- [65] F. G. Celiberto, D. Y. Ivanov, B. Murdaca, and A. Papa, *Dihadron production at the LHC: full next-to-leading BFKL calculation*, *Eur. Phys. J. C* **77** (2017), no. 6 382, [1701.05077].
- [66] F. G. Celiberto, D. Y. Ivanov, M. M. A. Mohammed, and A. Papa, *High-energy resummed distributions for the inclusive Higgs-plus-jet production at the LHC*, *Eur. Phys. J. C* **81** (2021), no. 4 293, [2008.00501].

- [67] M. Hentschinski, K. Kutak, and A. van Hameren, *Forward Higgs production within high energy factorization in the heavy quark limit at next-to-leading order accuracy*, *Eur. Phys. J. C* **81** (2021), no. 2 112, [2011.03193]. [Erratum: *Eur.Phys.J.C* 81, 262 (2021)].
- [68] F. G. Celiberto, M. Fucilla, D. Y. Ivanov, M. M. A. Mohammed, and A. Papa, *The next-to-leading order Higgs impact factor in the infinite top-mass limit*, 2205.02681.
- [69] F. G. Celiberto, M. Fucilla, D. Y. Ivanov, and A. Papa, *High-energy resummation in Λ_c baryon production*, *Eur. Phys. J. C* **81** (2021), no. 8 780, [2105.06432].
- [70] J. C. Collins and R. K. Ellis, *Heavy quark production in very high-energy hadron collisions*, *Nucl. Phys. B* **360** (1991) 3–30.
- [71] L. N. Lipatov, *Gauge invariant effective action for high-energy processes in QCD*, *Nucl. Phys. B* **452** (1995) 369–400, [hep-ph/9502308].
- [72] E. N. Antonov, L. N. Lipatov, E. A. Kuraev, and I. O. Cherednikov, *Feynman rules for effective Regge action*, *Nucl. Phys. B* **721** (2005) 111–135, [hep-ph/0411185].
- [73] A. van Hameren, P. Kotko, and K. Kutak, *Helicity amplitudes for high-energy scattering*, *JHEP* **01** (2013) 078, [1211.0961].
- [74] M. A. Kimber, A. D. Martin, and M. G. Ryskin, *Unintegrated parton distributions*, *Phys. Rev. D* **63** (2001) 114027, [hep-ph/0101348].
- [75] F. Hautmann, H. Jung, A. Lelek, V. Radescu, and R. Zlebcik, *Collinear and TMD Quark and Gluon Densities from Parton Branching Solution of QCD Evolution Equations*, *JHEP* **01** (2018) 070, [1708.03279].
- [76] I. Balitsky, *Operator expansion for high-energy scattering*, *Nucl. Phys. B* **463** (1996) 99–160, [hep-ph/9509348].
- [77] Y. V. Kovchegov, *Unitarization of the BFKL pomeron on a nucleus*, *Phys. Rev. D* **61** (2000) 074018, [hep-ph/9905214].
- [78] M. A. Nefedov, *Computing one-loop corrections to effective vertices with two scales in the EFT for Multi-Regge processes in QCD*, *Nucl. Phys. B* **946** (2019) 114715, [1902.11030].
- [79] M. A. Nefedov, *Towards stability of NLO corrections in High-Energy Factorization via Modified Multi-Regge Kinematics approximation*, *JHEP* **08** (2020) 055, [2003.02194].
- [80] S. Catani and M. H. Seymour, *A General algorithm for calculating jet cross-sections in NLO QCD*, *Nucl. Phys. B* **485** (1997) 291–419, [hep-ph/9605323]. [Erratum: *Nucl.Phys.B* 510, 503–504 (1998)].

- [81] A. van Hameren, *Calculating off-shell one-loop amplitudes for k_T -dependent factorization: a proof of concept*, 1710.07609.
- [82] E. Blanco, A. van Hameren, P. Kotko, and K. Kutak, *All-plus helicity off-shell gauge invariant multigluon amplitudes at one loop*, *JHEP* **12** (2020) 158, [2008.07916].
- [83] A. van Hameren, *BCFW recursion for off-shell gluons*, *JHEP* **07** (2014) 138, [1404.7818].
- [84] Z. Bern and G. Chalmers, *Factorization in one loop gauge theory*, *Nucl. Phys. B* **447** (1995) 465–518, [hep-ph/9503236].
- [85] Z. Bern, V. Del Duca, W. B. Kilgore, and C. R. Schmidt, *The infrared behavior of one loop QCD amplitudes at next-to-next-to leading order*, *Phys. Rev. D* **60** (1999) 116001, [hep-ph/9903516].
- [86] G. Ossola, C. G. Papadopoulos, and R. Pittau, *Reducing full one-loop amplitudes to scalar integrals at the integrand level*, *Nucl. Phys. B* **763** (2007) 147–169, [hep-ph/0609007].
- [87] W. T. Giele, Z. Kunszt, and K. Melnikov, *Full one-loop amplitudes from tree amplitudes*, *JHEP* **04** (2008) 049, [0801.2237].
- [88] R. Britto, F. Cachazo, and B. Feng, *New recursion relations for tree amplitudes of gluons*, *Nucl. Phys. B* **715** (2005) 499–522, [hep-th/0412308].
- [89] R. Britto, F. Cachazo, B. Feng, and E. Witten, *Direct proof of tree-level recursion relation in Yang-Mills theory*, *Phys. Rev. Lett.* **94** (2005) 181602, [hep-th/0501052].
- [90] Z. Kunszt, A. Signer, and Z. Trocsanyi, *Singular terms of helicity amplitudes at one loop in QCD and the soft limit of the cross-sections of multiparton processes*, *Nucl. Phys. B* **420** (1994) 550–564, [hep-ph/9401294].
- [91] A. Bassetto, M. Ciafaloni, and G. Marchesini, *Jet Structure and Infrared Sensitive Quantities in Perturbative QCD*, *Phys. Rept.* **100** (1983) 201–272.
- [92] M. L. Mangano and S. J. Parke, *Multiparton amplitudes in gauge theories*, *Phys. Rept.* **200** (1991) 301–367, [hep-th/0509223].
- [93] R. K. Ellis and J. C. Sexton, *QCD Radiative Corrections to Parton Parton Scattering*, *Nucl. Phys. B* **269** (1986) 445–484.

A Spinors and transverse momenta

Given a light-like momentum \mathbf{p} , we define the spinors as four-component vectors with two vanishing components:

$$|p] = \begin{pmatrix} L(\mathbf{p}) \\ \mathbf{0} \end{pmatrix}, \quad L(\mathbf{p}) = \frac{1}{\sqrt{|\mathbf{p}_0 + \mathbf{p}_3|}} \begin{pmatrix} -\mathbf{p}_1 + i\mathbf{p}_2 \\ \mathbf{p}_0 + \mathbf{p}_3 \end{pmatrix} \quad (152)$$

$$|p\rangle = \begin{pmatrix} \mathbf{0} \\ R(\mathbf{p}) \end{pmatrix}, \quad R(\mathbf{p}) = \frac{\sqrt{|\mathbf{p}_0 + \mathbf{p}_3|}}{\mathbf{p}_0 + \mathbf{p}_3} \begin{pmatrix} \mathbf{p}_0 + \mathbf{p}_3 \\ \mathbf{p}_1 + i\mathbf{p}_2 \end{pmatrix}. \quad (153)$$

The ‘‘dual’’ spinors are defined as

$$[p| = ((\mathcal{E}L(\mathbf{p}))^T, \mathbf{0}) \quad , \quad \langle p| = (\mathbf{0}, (\mathcal{E}^T R(\mathbf{p}))^T) \quad , \quad \text{where} \quad \mathcal{E} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}. \quad (154)$$

Defined as such, their dyadic products satisfy the relation

$$|p\rangle[p| + |p]\langle p| = \not{p} = \gamma_\mu p^\mu \quad (155)$$

where the γ -matrices are in the Weyl representation with

$$\gamma^5 \equiv i\gamma^0\gamma^1\gamma^2\gamma^3 = \begin{pmatrix} -\mathbf{1} & \mathbf{0} \\ \mathbf{0} & \mathbf{1} \end{pmatrix}. \quad (156)$$

For the spinor products, we also use the notation

$$\langle p||q\rangle = \langle pq\rangle = (pq)_- \quad , \quad [p||q] = [pq] = (pq)_+. \quad (157)$$

The most important relations for the spinor products are antisymmetry

$$\langle pq\rangle = -\langle qp\rangle \quad , \quad [pq] = -[qp] \quad (158)$$

and

$$\langle pq\rangle[qp] = (p + q)^2. \quad (159)$$

For a momentum r_\perp transverse to two light-like momenta P and \bar{P} that are not collinear, we define

$$\kappa_r^* = \frac{\langle P|\not{r}_\perp|\bar{P}\rangle}{[P\bar{P}]} \quad , \quad \kappa_r = \frac{\langle \bar{P}|\not{r}_\perp|P\rangle}{\langle \bar{P}P\rangle}. \quad (160)$$

Realize that κ_r, κ_r^* are invariant under replacement of \bar{P} with any other light-like momentum that is not collinear to P [83]. Using polarization vectors

$$e_+^\mu = \frac{\langle \bar{P}|\gamma^\mu|P\rangle}{\sqrt{2}\langle \bar{P}P\rangle} \quad , \quad e_-^\mu = \frac{\langle P|\gamma^\mu|\bar{P}\rangle}{\sqrt{2}[P\bar{P}]}, \quad (161)$$

satisfying $e_- \cdot e_- = e_+ \cdot e_+ = 0$, $e_- \cdot e_+ = -1$, we can then write

$$r_\perp^\mu = -\frac{\kappa_r^*}{\sqrt{2}} e_+^\mu - \frac{\kappa_r}{\sqrt{2}} e_-^\mu. \quad (162)$$

The inner product of two transverse moment is given by

$$r_\perp \cdot q_\perp = \frac{e_- \cdot e_+}{2} (\kappa_r \kappa_q^* + \kappa_r^* \kappa_q) = -\frac{\kappa_r \kappa_q^* + \kappa_r^* \kappa_q}{2}. \quad (163)$$

B Color representation

The relation of Eq. (22) holds as such after summation over color degrees of freedom of the external partons. Before squaring and summation, the color-dependent amplitude has a color representation that comes natural with the auxiliary partons used. In case of auxiliary quarks, the (anti-)fundamental color indices of these constitute the color of the space-like gluon in a color-flow-type of representation. Let us write the color indices explicitly with an i for the final-state auxiliary quark, and j for the initial-state auxiliary quark. We consider the situation with n more gluons. In [73] it was shown that indeed (employing Einstein summation)

$$\mathcal{M}_{\text{aux-q}}^{*a_1 a_2 \dots a_n ji} \delta_{ij} = 0, \quad (164)$$

as required for the color pair to represent a single gluon. The auxiliary label only refers to the color content and regarding the kinematics the Λ limit is already performed. The amplitude can be decomposed in terms of partial amplitudes \mathcal{A}^* as

$$\mathcal{M}_{\text{aux-q}}^{*a_1 a_2 \dots a_n ji} = 2^{n/2} \sum_{\sigma \in S_n} (\mathrm{T}^{a_{\sigma(1)}} \mathrm{T}^{a_{\sigma(2)}} \dots \mathrm{T}^{a_{\sigma(n)}})_{ji} \mathcal{A}^*(g^*, \sigma(1), \sigma(2), \dots, \sigma(n)). \quad (165)$$

In [81] it was shown that the partial amplitudes obtained with auxiliary quarks are the same as the ones obtained with auxiliary gluons, that is why we do not give them the auxiliary label. We immediately see that

$$\begin{aligned} \mathcal{M}_{\text{aux-q}}^{*a_1 a_2 \dots a_n ji} \sqrt{2} \mathrm{T}_{ij}^b &= 2^{(n+1)/2} \sum_{\sigma \in S_n} \mathrm{Tr}(\mathrm{T}^b \mathrm{T}^{a_{\sigma(1)}} \mathrm{T}^{a_{\sigma(2)}} \dots \mathrm{T}^{a_{\sigma(n)}}) \mathcal{A}^*(g^*, \sigma(1), \sigma(2), \dots, \sigma(n)) \\ &= \mathcal{M}^{*b a_1 a_2 \dots a_n} \end{aligned} \quad (166)$$

gives the color decomposition of $(n+1)$ -gluon amplitude, one of which happens to be space-like with color index b . The relation can be inverted as, suppressing the irrelevant a_i indices,

$$\mathcal{M}^{*b} \sqrt{2} \mathrm{T}_{kl}^b = \mathcal{M}_{\text{aux-q}}^{*ji} 2 \mathrm{T}_{ij}^b \mathrm{T}_{kl}^b = \mathcal{M}_{\text{aux-q}}^{*ji} \left(\delta_{il} \delta_{kj} - \frac{1}{N_c} \delta_{ij} \delta_{kl} \right) = \mathcal{M}_{\text{aux-q}}^{*kl}. \quad (167)$$

It must be mentioned that the partial amplitudes for auxiliary gluons are only the same as for auxiliary quarks in case the auxiliary gluons are adjacent. It is however not difficult to understand that partial amplitudes for which the auxiliary gluons are not adjacent vanish in the large Λ limit because they have an extra power of Λ in the denominator. Let the auxiliary gluons be number $n+1$ and $n+2$. The full amplitude can be decomposed as

$$\begin{aligned} \mathcal{M}_{\text{aux-g}}^{*a_1 a_2 \dots a_{n+2}} &= \\ &2^{n/2+1} \sum_{\sigma \in S_n} \mathrm{Tr}(\mathrm{T}^{a_{n+2}} \mathrm{T}^{a_{\sigma(1)}} \dots \mathrm{T}^{a_{\sigma(n)}} \mathrm{T}^{a_{n+1}}) \mathcal{A}_{\text{aux-g}}^*(n+2, \sigma(1), \dots, \sigma(n), n+1) \\ &+ 2^{n/2+1} \sum_{\sigma \in S_n} \mathrm{Tr}(\mathrm{T}^{a_{n+2}} \mathrm{T}^{a_{n+1}} \mathrm{T}^{a_{\sigma(1)}} \dots \mathrm{T}^{a_{\sigma(n)}}) \mathcal{A}_{\text{aux-g}}^*(n+2, n+1, \sigma(1), \dots, \sigma(n)). \end{aligned} \quad (168)$$

Usually, an $(n + 2)$ -gluon amplitude has $(n + 1)!$ terms, but we excluded the permutations for which the auxiliary gluons are not adjacent, so $2(n!)$ remain. Also, the partial amplitudes carry the auxiliary label. In terms of the eikonal Feynman rules for the tree-level space-like amplitude, it is however not difficult to see that exchanging the role of the auxiliary partons effectively means replacing $P \leftrightarrow -P$, and leads to an overall minus sign. Thus we can write

$$\begin{aligned} \mathcal{M}_{\text{aux-g}}^{*a_1 a_2 \dots a_{n+2}} &= 2^{n/2+1} \sum_{\sigma \in S_n} \text{Tr}([T^{a_{n+1}}, T^{a_{n+2}}] T^{a_{\sigma(1)}} \dots T^{a_{\sigma(n)}}) \mathcal{A}^*(g^*, \sigma(1), \dots, \sigma(n)) \\ &= 2^{n/2+1} i f^{a_{n+1} a_{n+2} b} \sum_{\sigma \in S_n} \text{Tr}(T^b T^{a_{\sigma(1)}} \dots T^{a_{\sigma(n)}}) \mathcal{A}^*(g^*, \sigma(1), \dots, \sigma(n)) . \end{aligned} \quad (169)$$

We conclude that, suppressing irrelevant indices now,

$$\mathcal{M}_{\text{aux-q}}^{*ji} = \sqrt{2} T_{ij}^b \mathcal{M}^{*b} \quad , \quad \mathcal{M}_{\text{aux-g}}^{*a_{n+1} a_{n+2}} = \sqrt{2} i f^{a_{n+1} a_{n+2} b} \mathcal{M}^{*b} . \quad (170)$$

Squaring and summing over color degrees of freedom, we get

$$\sum_{\text{color}} |\mathcal{M}_{\text{aux-q}}^{*ji}|^2 = 2 T_{ij}^b T_{ij}^{c\dagger} \sum_{\text{color}} \mathcal{M}^{*b} \mathcal{M}^{*c\dagger} = \sum_{\text{color}} |\mathcal{M}^{*b}|^2 , \quad (171)$$

where we used the \dagger to indicate complex conjugation. Averaging over initial-state color, we get

$$\frac{1}{N_c} \sum_{\text{color}} |\mathcal{M}_{\text{aux-q}}^{*ji}|^2 = \frac{N_c^2 - 1}{N_c} \frac{1}{N_c^2 - 1} \sum_{\text{color}} |\mathcal{M}^{*b}|^2 \quad (172)$$

where we put the natural color average factor for the space-like gluon. For the gluon case, we get

$$\sum_{\text{color}} |\mathcal{M}_{\text{aux-g}}^{*ab}|^2 = 2 f^{abe} f^{abd} \sum_{\text{color}} \mathcal{M}^{*e} \mathcal{M}^{*d\dagger} = 2 N_c \sum_{\text{color}} |\mathcal{M}^{*e}|^2 , \quad (173)$$

so the same times $2N_c$. Averaging over initial-state color, we get

$$\frac{1}{N_c^2 - 1} \sum_{\text{color}} |\mathcal{M}_{\text{aux-g}}^{*ab}|^2 = 2 N_c \frac{1}{N_c^2 - 1} \sum_{\text{color}} |\mathcal{M}^{*c}|^2 . \quad (174)$$

We also need color-correlated squared matrix elements. Highlighting only the color indices for the relevant parton, they are [80]

$$\text{quark: } \mathcal{M}^{i\dagger} T_{i\bar{i}}^a \mathcal{M}^{\bar{i}} \quad , \quad \text{antiquark: } \mathcal{M}^{j\dagger} (-T_{j\bar{j}}^a) \mathcal{M}^{\bar{j}} \quad , \quad \text{gluon: } \mathcal{M}^{b\dagger} (i f^{bac}) \mathcal{M}^c . \quad (175)$$

For the sum of squared matrix elements with color correlators for the auxiliary lines with equal coefficients (put to 1 here), we get

$$\begin{aligned} \mathcal{M}_{\text{aux-q}}^{*ji\dagger} T_{i\bar{i}}^a \mathcal{M}_{\text{aux-q}}^{*j\bar{i}} + \mathcal{M}_{\text{aux-q}}^{*ji\dagger} (-T_{j\bar{j}}^a) \mathcal{M}_{\text{aux-q}}^{*j\bar{i}} &= 2 \mathcal{M}^{*b\dagger} T_{ij}^{b\dagger} T_{i\bar{i}}^a T_{j\bar{j}}^c \mathcal{M}^{*c} - 2 \mathcal{M}^{*b\dagger} T_{ij}^{b\dagger} T_{j\bar{j}}^a T_{i\bar{i}}^c \mathcal{M}^{*c} \\ &= \mathcal{M}^{*b\dagger} (i f^{bac}) \mathcal{M}^{*c} , \end{aligned} \quad (176)$$

as desired. The one term with both auxiliary quark correlators is given by

$$\mathcal{M}_{\text{aux-q}}^{\star j i \dagger} T_{i\bar{i}}^a (-T_{j\bar{j}}^a) \mathcal{M}_{\text{aux-q}}^{\star \bar{i}} = -2\mathcal{M}^{\star b \dagger} T_{ij}^{b \dagger} \frac{1}{2} \left(\delta_{ij} \delta_{\bar{i}\bar{j}} - \frac{1}{N_c} \delta_{i\bar{i}} \delta_{j\bar{j}} \right) T_{\bar{i}\bar{j}}^c \mathcal{M}^{\star c} = \frac{1}{2N_c} \mathcal{M}^{\star b \dagger} \mathcal{M}^{\star b}. \quad (177)$$

With auxiliary gluons, we get, using the Jacobi identity,

$$\begin{aligned} \mathcal{M}_{\text{aux-g}}^{\star b_1 b_2 \dagger} (if^{b_1 a c_1}) \mathcal{M}_{\text{aux-g}}^{\star c_1 b_2} + \mathcal{M}_{\text{aux-g}}^{\star b_1 b_2 \dagger} (if^{b_2 a c_2}) \mathcal{M}_{\text{aux-g}}^{\star b_1 c_2} \\ = 2i\mathcal{M}^{\star d \dagger} f^{b_1 b_2 d} f^{b_1 a c_1} f^{c_1 b_2 e} \mathcal{M}^{\star e} + 2i\mathcal{M}^{\star d \dagger} f^{b_1 b_2 d} f^{b_2 a c_2} f^{b_1 c_2 e} \mathcal{M}^{\star e} \\ = 2N_c \mathcal{M}^{\star d \dagger} (if^{d a e}) \mathcal{M}^{\star e}, \end{aligned} \quad (178)$$

so again $2N_c$ times the desired result, like for the non-correlated square. The term with both correlators is

$$\begin{aligned} \mathcal{M}_{\text{aux-g}}^{\star b_1 b_2 \dagger} (if^{b_1 a c_1}) (if^{b_2 a c_2}) \mathcal{M}_{\text{aux-g}}^{\star c_1 c_2} = -2\mathcal{M}^{\star d \dagger} f^{b_1 b_2 d} f^{b_1 a c_1} f^{b_2 a c_2} f^{c_1 c_2 e} \mathcal{M}^{\star e} \\ = 2N_c \frac{-N_c}{2} \mathcal{M}^{\star d \dagger} \mathcal{M}^{\star d}. \end{aligned} \quad (179)$$

C Collinear splitting functions

The collinear limit of a momentum pair p_i^μ, p_j^μ in an amplitude can be derived by factoring out the obvious singular part, and then replacing

$$p_i^\mu \rightarrow (1-z)p_{ij}^\mu, \quad p_j^\mu \rightarrow zp_{ij}^\mu. \quad (180)$$

Consider for example the MHV amplitude (we omit the overall imaginary unit)

$$\mathcal{A}_n(1^-, 2^+, 3^-, 4^+, \dots, n^+) = \frac{\langle p_1 p_3 \rangle^4}{\langle p_1 p_2 \rangle \langle p_2 p_3 \rangle \langle p_3 p_4 \rangle \cdots \langle p_{n-1} p_n \rangle \langle p_n p_1 \rangle} \quad (181)$$

which is singular when p_1^μ, p_2^μ become collinear. Performing the procedure mentioned above, we get

$$\begin{aligned} \mathcal{A}_n(1^-, 2^+, 3^-, 4^+, \dots, n^+) \rightarrow \frac{1}{\langle p_1 p_2 \rangle} \frac{(1-z)^2 \langle p_{12} p_3 \rangle^4}{\sqrt{z} \langle p_{12} p_3 \rangle \langle p_3 p_4 \rangle \cdots \langle p_{n-1} p_n \rangle \sqrt{1-z} \langle p_n p_1 \rangle} \\ = \frac{1}{\langle p_1 p_2 \rangle} \frac{(1-z)^2}{\sqrt{z(1-z)}} \mathcal{A}_{n-1}((12)^-, 3^-, 4^+, \dots, n^+) \end{aligned} \quad (182)$$

which corresponds to Eq.(6.15) in [92] (because the all-plus amplitude vanishes).

Let us consider a slightly more general procedure, in which we take the limits $p_i^\mu \rightarrow xp^\mu$, $p_j^\mu \rightarrow yp^\mu$, and then correct the factorized amplitude so it contains momentum $(x+y)p^\mu$. For the example above we get

$$\begin{aligned} \mathcal{A}_n(1^-, 2^+, 3^-, 4^+, \dots, n^+) \rightarrow \frac{1}{\langle p_1 p_2 \rangle} \frac{x^2}{\sqrt{xy}} \frac{\langle pp_3 \rangle^4}{\langle pp_3 \rangle \langle p_3 p_4 \rangle \cdots \langle p_{n-1} p_n \rangle \langle p_n p \rangle} \\ = \frac{1}{\langle p_1 p_2 \rangle} \frac{x^2}{\sqrt{xy}(x+y)} \frac{\langle (x+y)pp_3 \rangle^4}{\langle (x+y)pp_3 \rangle \langle p_3 p_4 \rangle \cdots \langle p_{n-1} p_n \rangle \langle p_n (x+y)p \rangle} \\ = \frac{1}{\langle p_1 p_2 \rangle} \frac{x^2}{\sqrt{xy}(x+y)} \mathcal{A}_{n-1}((x+y)p^-, 3^-, 4^+, \dots, n^+). \end{aligned} \quad (183)$$

We get the previous formula by simply setting $x = 1 - z$ and $y = z$. In our definition of the spinors, they scale with not-necessarily positive fractions as

$$|xp\rangle = \sqrt{|x|} |p\rangle \quad , \quad |xp] = \frac{x}{\sqrt{|x|}} |p] = \text{sgn}(x) \sqrt{|x|} |p] \quad , \quad (184)$$

so for the previous splitting function, the fractions x , y , and $(x + y)$ just need to be replaced by their absolute values. Let us also consider the explicit example with an anti-MHV amplitude. We get

$$\begin{aligned} \mathcal{A}_n(1^-, 2^+, 3^+, 4^-, \dots, n^-) &= \frac{[p_3 p_2]^4}{[p_1 p_n][p_n p_{n-1}] \cdots [p_4 p_3][p_3 p_2][p_2 p_1]} \\ &\rightarrow \frac{1}{[p_2 p_1]} \frac{\text{sgn}(xy) y^2}{\sqrt{|xy|}} \frac{[p_3 p]^4}{[p p_n][p_n p_{n-1}] \cdots [p_4 p_3][p_3 p]} \\ &= \frac{-1}{[p_1 p_2]} \frac{\text{sgn}(xy) y^2}{\sqrt{|xy|} |x + y|} \mathcal{A}_{n-1}((x + y)^+, 3^+, 4^-, \dots, n^-) \quad . \quad (185) \end{aligned}$$

We can summarize

$$\text{Split}^{g \rightarrow g_1 g_2}((x + y)^- \rightarrow 1_x^-, 2_y^+) = \frac{1}{(12)_-} \frac{x^2}{\sqrt{|xy|} |x + y|} \quad (186)$$

$$\text{Split}^{g \rightarrow g_1 g_2}((x + y)^+ \rightarrow 1_x^-, 2_y^+) = \frac{-1}{(12)_+} \frac{\text{sgn}(xy) y^2}{\sqrt{|xy|} |x + y|} \quad , \quad (187)$$

where the $(1_x^+, 2_y^-)$ functions are obtained by exchanging $x^2 \leftrightarrow y^2$ in the numerators. Realize that here the splitting functions indicate the amplitude helicities, while in [85] the collinear limit is indicated with the opposite helicity. The other non-vanishing ones are

$$\text{Split}^{g \rightarrow g_1 g_2}((x + y)^+ \rightarrow 1_x^+, 2_y^+) = \frac{1}{(12)_-} \frac{1}{\sqrt{|xy|} |x + y|} \quad (188)$$

$$\text{Split}^{g \rightarrow g_1 g_2}((x + y)^- \rightarrow 1_x^-, 2_y^-) = \frac{-1}{(12)_+} \frac{\text{sgn}(xy)}{\sqrt{|xy|} |x + y|} \quad . \quad (189)$$

For squared amplitudes summed over helicities, the gluon splitting function becomes

$$\mathcal{Q}_{gg}(x, y) = \frac{x^4 + y^4 + (x + y)^4}{|xy|(x + y)^2} = \frac{1 + (y/x)^4 + (1 + y/x)^4}{|y/x|(1 + y/x)^2} = \frac{\mathcal{P}_{gg}(|1 + y/x|)}{|1 + y/x|} \quad . \quad (190)$$

For the splitting of a quark or anti-quark into a quark or anti-quark and a gluon we with $p_{q/\bar{q}} \rightarrow$

$xp_{q/\bar{q}}$ and $p_g \rightarrow yp_{q/\bar{q}}$ are given by

$$\text{Split}(q_{x+y}^\mp \rightarrow q_x^\pm g_y^\pm) = \sigma_\mp(y) \frac{\pm\sqrt{|x+y|}}{\sqrt{|y|} (p_q p_g)_\mp} \quad (191)$$

$$\text{Split}(q_{x+y}^\mp \rightarrow q_x^\pm g_y^\mp) = \sigma_\pm(y) \frac{\mp|x|}{\sqrt{|y||x+y|} (p_q p_g)_\pm} \quad (192)$$

$$\text{Split}(\bar{q}_{x+y}^\pm \rightarrow \bar{q}_x^\mp g_y^\mp) = \sigma_\pm(y) \frac{\mp\sqrt{|x+y|}}{\sqrt{|y|} (p_g p_{\bar{q}})_\pm} \quad (193)$$

$$\text{Split}(\bar{q}_{x+y}^\pm \rightarrow \bar{q}_x^\mp g_y^\pm) = \sigma_\mp(y) \frac{\pm|x|}{\sqrt{|y||x+y|} (p_g p_{\bar{q}})_\mp} \quad (194)$$

where

$$\sigma_+(y) = 1 \quad , \quad \sigma_-(y) = \text{sgn}(y) \quad , \quad (195)$$

leading to the splitting function

$$Q_{qq}(x, y) = \frac{1}{|y|} \left(|x+y| + \frac{x^2}{|x+y|} \right) = \frac{1 + (1 + y/x)^2}{|y/x| |1 + y/x|} = \frac{\mathcal{P}_{qq}(|1 + y/x|)}{|1 + y/x|} \quad (196)$$

The remaining splitting functions are given by

$$\text{Split}(g_{x+y}^\mp \rightarrow \bar{q}_x^\pm q_y^\mp) = \frac{1}{(p_{\bar{q}} p_q)_\mp} \frac{|y|}{|x+y|} \quad (197)$$

$$\text{Split}(g_{x+y}^\mp \rightarrow \bar{q}_x^\mp q_y^\pm) = \frac{1}{(p_{\bar{q}} p_q)_\mp} \frac{|x|}{|x+y|} \quad (198)$$

C.1 Collinear limit with P for space-like amplitudes

We extract the splitting functions by considering the example of MHV amplitudes. For a space-like gluon with momentum $k = p_1 + k_\perp$ we have [83],

$$\mathcal{A}_n^*(p_1 + k_\perp, 2^+, 3^-, 4^+, \dots, n^+) = \frac{1}{\kappa^*} \frac{\langle p_1 p_3 \rangle^4}{\langle p_1 p_2 \rangle \langle p_2 p_3 \rangle \langle p_3 p_4 \rangle \cdots \langle p_{n-1} p_n \rangle \langle p_n p_1 \rangle} \quad (199)$$

As such, the amplitude is singular for $k_\perp \rightarrow 0$ and still requires a factor $|k_\perp|$. We, however, prefer to define amplitudes like this and not to break their analytic structure. The necessary factor is included when constructing the squared matrix element.

We can easily extract the limit $p_1 \rightarrow xP$ and $p_2 \rightarrow yP$

$$\begin{aligned} & \mathcal{A}_n^*(p_1 + k_\perp, 2^+, 3^-, 4^+, \dots, n^+) \\ & \xrightarrow{p_1 \rightarrow xP, p_2 \rightarrow yP} \frac{1}{\langle p_1 p_2 \rangle} \frac{1}{\kappa^*} \frac{x^2 \langle P p_3 \rangle^4}{\sqrt{|y|} \langle P p_3 \rangle \langle p_3 p_4 \rangle \cdots \langle p_{n-1} p_n \rangle \sqrt{|x|} \langle p_n P \rangle} \\ & = \frac{1}{\langle p_1 p_2 \rangle} \frac{1}{\sqrt{|xy||x+y|}} \frac{1}{\kappa^*} \frac{\langle (x+y) P | p_3 \rangle^4}{\langle (x+y) P | p_3 \rangle \langle p_3 p_4 \rangle \cdots \langle p_{n-1} p_n \rangle \langle p_n | (x+y) P \rangle} \\ & = \frac{1}{\langle P | p_2 \rangle} \frac{|x|}{\sqrt{|y||x+y|}} \mathcal{A}_{n-1}^*((x+y)P + k_\perp, 3^-, 4^+, \dots, n^+) \quad (200) \end{aligned}$$

For the opposite helicity there is $\sqrt{|y|}/y$ instead of $1/\sqrt{|y|}$, and also $1/[p_2P] = -1/[Pp_2]$. Thus we find

$$\text{Split}(g^*((x+y)P + k_\perp) \rightarrow g^*(xP + k_\perp), g^+(yP)) = \frac{1}{(\text{Pr})_-} \frac{|x|}{\sqrt{|y|}|x+y|} \quad (201)$$

$$\text{Split}(g^*((x+y)P + k_\perp) \rightarrow g^*(xP + k_\perp), g^-(yP)) = \frac{-1}{(\text{Pr})_+} \frac{\text{sgn}(y)|x|}{\sqrt{|y|}|x+y|}. \quad (202)$$

For squared matrix elements with a space-like initial-state gluon, this leads to the collinear limit for a final-state gluon with momentum r (and taking $g_s = 1$)

$$|\overline{\mathcal{M}}^*|^2(x_k, k_\perp, \bar{x}; r, \{p_{i=1}^n\}) \xrightarrow{r \rightarrow x_r P} 2N_c \frac{1}{P \cdot r} \frac{x_k^2}{x_r(x_k - x_r)^2} |\overline{\mathcal{M}}^*|^2((x_k - x_r), k_\perp, \bar{x}; \{p_{i=1}^n\}). \quad (203)$$

D The triple auxiliary parton limit

Let us start with a simple example of the auxiliary parton method applied to amplitudes. Consider the partial amplitude for a process involving a quark-antiquark pair and 3 gluons.

$$\mathcal{A}(q^-, g_1^-, g_2^+, g_3^+, \bar{q}^+) = \frac{\langle q1 \rangle^3 \langle \bar{q}1 \rangle}{\langle \bar{q}q \rangle \langle q1 \rangle \langle 12 \rangle \langle 23 \rangle \langle 3\bar{q} \rangle}. \quad (204)$$

The quark-antiquark pair take the role of auxiliary partons, which means they get the spinors

$$|\bar{q}\rangle = \sqrt{\Lambda + x} |P\rangle, \quad |q\rangle = \sqrt{\Lambda} |P\rangle + \frac{\kappa^*}{\sqrt{\Lambda} \langle P\bar{P} \rangle} |\bar{P}\rangle. \quad (205)$$

We shifted Λ by an amount x for convenience. Contrary to the construction in [82], these spinors will give the sum of the quark and antiquark momenta a non-zero component for \bar{P}^μ , which however still vanishes for $\Lambda \rightarrow \infty$. Important is that

$$\langle \bar{q}q \rangle = \kappa^* + \mathcal{O}(\Lambda^{-1}) \quad (206)$$

does not vanish or become large for $\Lambda \rightarrow \infty$, and the amplitude scales as Λ due to the occurrence of $|\bar{q}\rangle$ and $|q\rangle$ in other spinor products. Inserting the spinors, we find

$$\mathcal{A}(q^-, g_1^-, g_2^+, g_3^+, \bar{q}^+) \rightarrow \frac{\Lambda \langle P1 \rangle^4}{\kappa^* \langle P1 \rangle \langle 12 \rangle \langle 23 \rangle \langle 3P \rangle} = \Lambda \mathcal{A}^*(g^*, g_1^-, g_2^+, g_3^+). \quad (207)$$

For the triple-parton limit, we distribute Λ over the quark and a gluon with spinors

$$|q\rangle = \sqrt{z\Lambda} |P\rangle + \frac{\kappa_q^*}{\sqrt{z\Lambda} \langle P\bar{P} \rangle} |\bar{P}\rangle, \quad |r\rangle = \sqrt{(1-z)\Lambda} |P\rangle + \frac{\kappa_r^*}{\sqrt{(1-z)\Lambda} \langle P\bar{P} \rangle} |\bar{P}\rangle. \quad (208)$$

We see that if we choose r to refer to 2 in Eq. (204), then it will produce a factor Λ in the denominator and give a subleading contribution. If we choose 1, then the spinor products in the

numerator will not produce powers of Λ . In order to produce a leading contribution, it must refer to 3, and inserting the spinors, we get

$$\begin{aligned}
\mathcal{A}(q^-, g_1^-, g_2^+, g_r^+, \bar{q}^+) &= \frac{\langle q1 \rangle^3 \langle \bar{q}1 \rangle}{\langle \bar{q}q \rangle \langle q1 \rangle \langle 12 \rangle \langle 2r \rangle \langle r\bar{q} \rangle} \\
&\rightarrow \frac{(\sqrt{z\Lambda} \langle P1 \rangle)^3 (\sqrt{\Lambda} \langle P1 \rangle)}{\frac{\kappa_q^*}{\sqrt{z}} (\sqrt{z\Lambda} \langle P1 \rangle) \langle 12 \rangle (\sqrt{(1-z)\Lambda} \langle 2P \rangle) \frac{-\kappa_r^*}{\sqrt{1-z}}} \\
&= \frac{-z^{3/2} \Lambda \langle P1 \rangle^4}{\kappa_q^* \kappa_r^* \langle P1 \rangle \langle 12 \rangle \langle 2P \rangle} = -z^{3/2} \frac{\kappa_q^* + \kappa_r^*}{\kappa_q^* \kappa_r^*} \Lambda \mathcal{A}(g^*, g_1^-, g_2^+), \quad (209)
\end{aligned}$$

where the factor $\kappa_q^* + \kappa_r^*$ occurs, because the space-like gluon in $\Lambda(g^*, g_1^-, g_2^+)$ has transverse momentum $q_\perp^\mu + r_\perp^\mu$, and thus the amplitude a factor $\kappa_{q+r}^* = \kappa_q^* + \kappa_r^*$ in the denominator.

Below we give all possible ‘‘splitting functions’’. The complete construction of momenta and spinors is

$$K_{\bar{q}}^\mu = -(\chi + \Lambda)P^\mu, \quad K_q^\mu = z_q \Lambda P^\mu + q_\perp^\mu - \frac{q_\perp^2}{sz_q \Lambda} \bar{p}^\mu, \quad K_r^\mu = z_r \Lambda P^\mu + r_\perp^\mu - \frac{r_\perp^2}{sz_r \Lambda} \bar{p}^\mu, \quad (210)$$

with

$$z_r + z_q = 1. \quad (211)$$

So we have, to $\mathcal{O}(\Lambda^{-1})$,

$$K_{\bar{q}}^\mu + K_q^\mu + K_r^\mu = -\chi P^\mu + q_\perp^\mu + r_\perp^\mu, \quad (K_{\bar{q}} + K_q + K_r)^2 = (q_\perp + r_\perp)^2 \quad (212)$$

and

$$s_{q\bar{q}} = (K_{\bar{q}} + K_q)^2 = \frac{q_\perp^2}{z_q}, \quad s_{r\bar{q}} = (K_{\bar{q}} + K_r)^2 = \frac{r_\perp^2}{z_r} \quad (213)$$

$$s_{rq} = (K_q + K_r)^2 = (q_\perp + r_\perp)^2 - \frac{q_\perp^2}{z_q} - \frac{r_\perp^2}{z_r}. \quad (214)$$

It is important to observe that these three invariants do not depend on Λ and do not produce a factor of Λ in propagator denominators. The spinors of the momenta are give by

$$|\bar{q}\rangle = \sqrt{\Lambda + \chi} |P\rangle, \quad |\bar{q}] = -\sqrt{\Lambda + \chi} |P] \quad (215)$$

$$|q\rangle = \sqrt{z_q \Lambda} |P\rangle + \frac{\kappa_q^*}{\sqrt{z_q \Lambda} \langle P\bar{P} \rangle} |\bar{P}\rangle, \quad |q] = \sqrt{z_q \Lambda} |P] + \frac{\kappa_q}{\sqrt{z_q \Lambda} [\bar{P}P]} |\bar{P}] \quad (216)$$

$$|r\rangle = \sqrt{z_r \Lambda} |P\rangle + \frac{\kappa_r^*}{\sqrt{z_r \Lambda} \langle P\bar{P} \rangle} |\bar{P}\rangle, \quad |r] = \sqrt{z_r \Lambda} |P] + \frac{\kappa_r}{\sqrt{z_r \Lambda} [\bar{P}P]} |\bar{P}] \quad (217)$$

with products

$$\begin{aligned}
\langle q\bar{q} \rangle &= -\frac{\kappa_q^*}{\sqrt{z_q}}, \quad [\bar{q}q] = \frac{\kappa_q}{\sqrt{z_q}} \\
\langle r\bar{q} \rangle &= -\frac{\kappa_r^*}{\sqrt{z_r}}, \quad [\bar{q}r] = \frac{\kappa_r}{\sqrt{z_r}} \\
\langle r q \rangle &= \sqrt{\frac{z_r}{z_q}} \kappa_q^* - \sqrt{\frac{z_q}{z_r}} \kappa_r^*, \quad [qr] = \sqrt{\frac{z_r}{z_q}} \kappa_q - \sqrt{\frac{z_q}{z_r}} \kappa_r
\end{aligned} \quad (218)$$

It is convenient to introducing the functions

$$\text{Sqr}(\bar{q}, q, r) = \frac{\kappa_q + \kappa_r}{[\bar{q}q][qr][r\bar{q}]} \quad , \quad \text{Ang}(\bar{q}, q, r) = \frac{\kappa_q^* + \kappa_r^*}{\langle \bar{q}q \rangle \langle qr \rangle \langle r\bar{q} \rangle} . \quad (219)$$

D.1 Auxiliary quark-antiquark pair plus a gluon

We have

$$\mathcal{A}(q, g_r, \dots, \bar{q}) \rightarrow \Lambda \mathcal{S}(\bar{q}, q, g_r) \mathcal{A}^*(g^*, \dots) \quad (220)$$

$$\mathcal{A}(q, \dots, g_r, \bar{q}) \rightarrow \Lambda \mathcal{S}(g_r, \bar{q}, q) \mathcal{A}^*(g^*, \dots) \quad (221)$$

The logic in the arguments of \mathcal{S} is that g is next to \bar{q} or next to q while keeping the cyclic order. The function \mathcal{S} for non-vanishing helicities are

$$\mathcal{S}(\bar{q}^+, q^-, g_r^-) = \text{Sqr}(\bar{q}, q, r) \frac{z_q^{1/2}[r\bar{q}]}{\sqrt{z_r}} \quad , \quad \mathcal{S}(\bar{q}^-, q^+, g_r^+) = \text{Ang}(\bar{q}, q, r) \frac{z_q^{1/2}\langle r\bar{q} \rangle}{\sqrt{z_r}} \quad , \quad (222)$$

$$\mathcal{S}(\bar{q}^+, q^-, g_r^+) = \text{Ang}(\bar{q}, q, r) \frac{z_q^{3/2}\langle r\bar{q} \rangle}{\sqrt{z_r}} \quad , \quad \mathcal{S}(\bar{q}^-, q^+, g_r^-) = \text{Sqr}(\bar{q}, q, r) \frac{z_q^{3/2}[r\bar{q}]}{\sqrt{z_r}} \quad , \quad (223)$$

$$\mathcal{S}(g_r^-, \bar{q}^+, q^-) = \text{Sqr}(\bar{q}, q, r) \frac{-[qr]}{\sqrt{z_r}} \quad , \quad \mathcal{S}(g_r^+, \bar{q}^-, q^+) = \text{Ang}(\bar{q}, q, r) \frac{\langle qr \rangle}{\sqrt{z_r}} \quad , \quad (224)$$

$$\mathcal{S}(g_r^+, \bar{q}^+, q^-) = \text{Ang}(\bar{q}, q, r) \frac{z_q \langle qr \rangle}{\sqrt{z_r}} \quad , \quad \mathcal{S}(g_r^-, \bar{q}^-, q^+) = \text{Sqr}(\bar{q}, q, r) \frac{-z_q [qr]}{\sqrt{z_r}} \quad . \quad (225)$$

In general an amplitude for $(n + 1)$ gluons and a quark-antiquark pair can for any given helicity configuration be decomposed as

$$\mathcal{M}_{\text{tree}}^{a_1 a_2 \dots a_{n+1} j i} = 2^{\frac{n+1}{2}} \sum_{\sigma \in S_{n+1}} (T^{a_{\sigma(1)}} T^{a_{\sigma(2)}} \dots T^{a_{\sigma(n+1)}})_{ji} \mathcal{A}(q, g_{\sigma(1)}, g_{\sigma(2)}, \dots, g_{\sigma(n+1)}, \bar{q}) \quad (226)$$

where \mathcal{A} without color indices is the color-ordered partial amplitude. Now, gluon number $n + 1$ is the radiative one, and only when it is the first or the last in the argument list, then the partial amplitude contributes in the joint large- Λ limit for the radiative gluon and the auxiliary pair:

$$\mathcal{M}_{\text{tree}}^{a_1 a_2 \dots a_n a_r j i} \rightarrow \Lambda \mathcal{S}(\bar{q}, q, g_r) \sqrt{2} (T^{a_r} \Omega^*)_{ji} + \Lambda \mathcal{S}(g_r, \bar{q}, q) \sqrt{2} (\Omega^* T^{a_r})_{ji} \quad , \quad (227)$$

where

$$\Omega_{lk}^* = 2^{\frac{n}{2}} \sum_{\sigma \in S_n} (T^{a_{\sigma(1)}} T^{a_{\sigma(2)}} \dots T^{a_{\sigma(n)}})_{lk} \mathcal{A}(g^*, g_{\sigma(1)}, g_{\sigma(2)}, \dots, g_{\sigma(n)}) \quad (228)$$

is the full colored amplitude with 1 space-like and n on-shell gluons. The right-hand side of Eq. (227) must be squared and summed over color and helicities to obtain the formula for squared matrix elements. We find

$$|\overline{\mathcal{M}}|^2(\Lambda + x, \bar{x}; K_r, K_q, \{p_i\}) \xrightarrow{\Lambda \rightarrow \infty} C_{\text{aux}} \mathcal{Q}_{\text{aux}}(z_q, q_{\perp}, z_r, r_{\perp}) \frac{\Lambda^2 |\overline{\mathcal{M}}^*|^2(x, -q_{\perp} - r_{\perp}, \bar{x}; \{p_i\})}{x^2 |q_{\perp} + r_{\perp}|^2} \quad , \quad (229)$$

with

$$\mathcal{Q}_{\text{aux}}(z_q, \mathbf{q}_\perp, z_r, \mathbf{r}_\perp) = \mathcal{P}_{\text{aux}}(z_q, z_r) |\mathbf{q}_\perp + \mathbf{r}_\perp|^2 \left(\frac{c_{\bar{q}}}{|s_{r\bar{q}}| |s_{q\bar{q}}|} + \frac{c_q z_q}{|s_{rq}| |s_{q\bar{q}}|} + \frac{c_r z_r}{|s_{rq}| |s_{r\bar{q}}|} \right), \quad (230)$$

and

$$\mathcal{C}_{\text{aux}} = \frac{N_c^2 - 1}{N_c}, \quad c_{\bar{q}} = c_q = N_c, \quad c_r = -\frac{1}{N_c}, \quad (231)$$

and collinear splitting function

$$\mathcal{P}_{\text{aux}}(z_q, z_r) = \mathcal{P}_{qq}(z_q, z_r) = \frac{1 + z_q^2 - \epsilon z_r^2}{z_r}. \quad (232)$$

The term proportional to ϵ from dimensional regularization was inferred from a cross-check using the explicit expressions for relevant squared matrix elements in [93].

D.2 Auxiliary gluon pair plus a gluon

We keep the labels \bar{q} and q for the auxiliary pair, and have

$$\mathcal{A}(g_q, g_r, \dots, g_{\bar{q}}) \rightarrow \Lambda \mathcal{S}(g_{\bar{q}}, g_q, g_r) \mathcal{A}^*(g^*, \dots) \quad (233)$$

$$\mathcal{A}(g_q, \dots, g_r, g_{\bar{q}}) \rightarrow \Lambda \mathcal{S}(g_r, g_{\bar{q}}, g_q) \mathcal{A}^*(g^*, \dots) \quad (234)$$

with

$$\mathcal{S}(g_{\bar{q}}^+, g_q^-, g_r^-) = \text{Sqr}(\bar{q}, q, r) \frac{-z_q^{1/2} [r\bar{q}]}{\sqrt{z_q z_r}} \quad (235)$$

$$\mathcal{S}(g_{\bar{q}}^+, g_q^-, g_r^+) = \text{Ang}(\bar{q}, q, r) \frac{z_q^{5/2} \langle r\bar{q} \rangle}{\sqrt{z_q z_r}} \quad (236)$$

$$\mathcal{S}(g_{\bar{q}}^+, g_q^+, g_r^-) = \text{Ang}(\bar{q}, q, r) \frac{z_q^{1/2} z_r^2 \langle r\bar{q} \rangle}{\sqrt{z_q z_r}} \quad (237)$$

$$\mathcal{S}(g_r^-, g_{\bar{q}}^+, g_q^-) = \text{Sqr}(\bar{q}, q, r) \frac{[qr]}{\sqrt{z_q z_r}} \quad (238)$$

$$\mathcal{S}(g_r^+, g_{\bar{q}}^+, g_q^-) = \text{Ang}(\bar{q}, q, r) \frac{z_q^2 \langle qr \rangle}{\sqrt{z_q z_r}} \quad (239)$$

$$\mathcal{S}(g_r^-, g_{\bar{q}}^+, g_q^+) = \text{Ang}(\bar{q}, q, r) \frac{z_r^2 \langle qr \rangle}{\sqrt{z_q z_r}} \quad (240)$$

For squared matrix elements, we find Eq. (229) and Eq. (230) again, but with

$$\mathcal{C}_{\text{aux}} = 2N_c, \quad c_{\bar{q}} = c_q = c_r = N_c, \quad (241)$$

and collinear splitting function

$$\mathcal{P}_{gg}(z_q, z_r) = \frac{1 + z_r^4 + z_q^4}{z_r z_q}. \quad (242)$$

D.3 Initial-state gluon with final-state quark-antiquark pair

For the triple limit, we also need to include this case, which has no double limit equivalent. We will still use the label r for the gluon, but realize that now for $|r\rangle$ there is a minus sign $|r\rangle \rightarrow -\Lambda|P\rangle$, and a momentum fraction $z_{\bar{q}}$ will be associated with the final-state anti-quark.

$$\mathcal{S}_g(\bar{q}^+, q^-, g_r^-) = \text{Sqr}(\bar{q}, q, r) \left(-z_q^{1/2} z_{\bar{q}} [r\bar{q}] \right) \quad (243)$$

$$\mathcal{S}_g(\bar{q}^+, q^-, g_r^+) = \text{Ang}(\bar{q}, q, r) \left(z_q^{3/2} \langle r\bar{q} \rangle \right) \quad (244)$$

$$\mathcal{S}_g(g_r^-, \bar{q}^+, q^-) = \text{Sqr}(\bar{q}, q, r) \left(-z_q^{3/2} [qr] \right) \quad (245)$$

$$\mathcal{S}_g(g_r^+, \bar{q}^+, q^-) = \text{Ang}(\bar{q}, q, r) \left(z_q z_{\bar{q}}^{1/2} \langle qr \rangle \right) \quad (246)$$

$$\mathcal{S}_g(\bar{q}^-, q^+, g_r^+) = \text{Ang}(\bar{q}, q, r) \left(z_q^{1/2} z_{\bar{q}} \langle r\bar{q} \rangle \right) \quad (247)$$

$$\mathcal{S}_g(\bar{q}^-, q^+, g_r^-) = \text{Sqr}(\bar{q}, q, r) \left(-z_q^{3/2} [r\bar{q}] \right) \quad (248)$$

$$\mathcal{S}_g(g_r^+, \bar{q}^-, q^+) = \text{Ang}(\bar{q}, q, r) \left(z_{\bar{q}}^{3/2} \langle qr \rangle \right) \quad (249)$$

$$\mathcal{S}_g(g_r^-, \bar{q}^-, q^+) = \text{Sqr}(\bar{q}, q, r) \left(-z_q z_{\bar{q}}^{1/2} [qr] \right) \quad (250)$$

The functions Ang and Sqr now contain $\kappa_{\bar{q}}, \kappa_{\bar{q}}^*$ instead of κ_r, κ_r^* . For squared matrix elements, we find Eq. (229) with

$$\mathcal{P}_{\text{aux}}(z_q, q_{\perp}, z_{\bar{q}}, \bar{q}_{\perp}) = \mathcal{P}_{\text{ab}}(z_q, z_r) |q_{\perp} + \bar{q}_{\perp}|^2 \left(\frac{c_r}{|s_{\bar{q}r}| |s_{qr}|} + \frac{c_q z_q}{|s_{\bar{q}q}| |s_{qr}|} + \frac{c_{\bar{q}} z_{\bar{q}}}{|s_{\bar{q}q}| |s_{\bar{q}r}|} \right), \quad (251)$$

and

$$C_{\text{aux}} = \frac{N_c^2 - 1}{N_c}, \quad c_{\bar{q}} = c_q = \frac{1}{2}, \quad c_r = -\frac{1}{2N_c}, \quad (252)$$

and collinear splitting function

$$\mathcal{P}_{qg}(z_q, z_r) = 1 - \frac{z_q z_{\bar{q}}}{1 - \epsilon}. \quad (253)$$

E Some integrals

The following integrals appear in this paper

$$I_A = \int d^{2+\bar{\epsilon}} r_{\perp} \frac{\theta(0 < r_{\perp}^2 < \infty)}{r_{\perp}^2 |r_{\perp} + k_{\perp}|^2} = \frac{\pi_{\epsilon} |k_{\perp}|^{\bar{\epsilon}}}{|k_{\perp}|^2} \frac{4}{\bar{\epsilon}} + \mathcal{O}(\bar{\epsilon}^2), \quad (254)$$

and

$$\begin{aligned} I_B &= \int d^{2+\bar{\epsilon}} r_{\perp} \frac{\theta(0 < r_{\perp}^2 < v^2)}{r_{\perp}^2 |r_{\perp} + k_{\perp}|^2}, \quad \frac{|k_{\perp}|^2}{v^2} < 1 \\ &= \frac{\pi_{\epsilon} |k_{\perp}|^{\bar{\epsilon}}}{|k_{\perp}|^2} \left\{ \frac{4}{\bar{\epsilon}} + \ln \left(1 - \frac{|k_{\perp}|^2}{v^2} \right) \right\} + \mathcal{O}(\bar{\epsilon}) \\ &= \frac{\pi_{\epsilon} v^{\bar{\epsilon}}}{|k_{\perp}|^2} \left\{ \frac{4}{\bar{\epsilon}} + 2 \ln \frac{|k_{\perp}|^2}{v^2} + \ln \left(1 - \frac{|k_{\perp}|^2}{v^2} \right) \right\} + \mathcal{O}(\bar{\epsilon}), \end{aligned} \quad (255)$$

and

$$\begin{aligned}
I_C &= \int_0^u \frac{dx}{x} \int d^{2+\bar{\epsilon}} r_\perp \frac{\theta(0 < r_\perp^2 < vx)}{r_\perp^2 |r_\perp + \mathbf{k}_\perp|^2} \quad , \quad \frac{|\mathbf{k}_\perp|^2}{uv} < 1 \\
&= \frac{\pi_\epsilon |\mathbf{k}_\perp|^{\bar{\epsilon}}}{|\mathbf{k}_\perp|^2} \left\{ \frac{4}{\bar{\epsilon}^2} - \frac{4}{\bar{\epsilon}} \ln \frac{|\mathbf{k}_\perp|^2}{uv} + \text{Li}_2 \left(\frac{|\mathbf{k}_\perp|^2}{uv} \right) \right\} + \mathcal{O}(\bar{\epsilon}) .
\end{aligned} \tag{256}$$