

MULTILEVEL PICARD APPROXIMATION ALGORITHM FOR SEMILINEAR PARTIAL INTEGRO-DIFFERENTIAL EQUATIONS AND ITS COMPLEXITY ANALYSIS

ARIEL NEUFELD AND SIZHOU WU

ABSTRACT. In this paper we introduce a multilevel Picard approximation algorithm for semilinear parabolic partial integro-differential equations (PIDEs). We prove that the numerical approximation scheme converges to the unique viscosity solution of the PIDE under consideration. To that end, we derive a Feynman-Kac representation for the unique viscosity solution of the semilinear PIDE, extending the classical Feynman-Kac representation for linear PIDEs. Furthermore, we show that the algorithm does not suffer from the curse of dimensionality, i.e. the computational complexity of the algorithm is bounded polynomially in the dimension d and the prescribed reciprocal of the accuracy ε .

1. INTRODUCTION

Nonlinear partial integro-differential equations (PIDEs) have many important applications in finance, physics, biology, economics, and engineering; we refer to [17, 18, 23, 51] and the references therein, as well as [13] for an excellent survey on applications of nonlocal PDEs in various fields.

In the last years, several papers proposed modern numerical methods to solve nonlinear PDEs involving either neural networks [7, 27, 28, 31, 36, 37, 49, 53, 54], or multilevel Monte Carlo methods [8, 10, 11, 26, 32, 38, 39, 40], which are able to approximately solve high-dimensional nonlinear PDEs in up to 10^7 dimensions. Moreover, it has been proven in [8, 11, 26, 38, 39, 40], under some (typically Lipschitz-) continuity conditions on the corresponding coefficients of the PDE under consideration, that multilevel Picard approximations do overcome the curse of dimensionality in the sense that the computational complexity of the algorithms is bounded polynomially in the dimension d and the prescribed (reciprocal of the) accuracy ε .

However, the development of numerical methods and their complexity analysis for PIDEs is only at its infancy and is still emerging. [33, 34] proposed both machine learning based and Monte Carlo based methods for solving *linear* PIDEs and showed that their algorithms do not suffer from the curse of dimensionality. In [4] the deep Galerkin algorithm of [54] has been extended to PIDEs. In [58] physics informed neural networks for solving nonlinear PIDEs have been constructed. Deep neural network based algorithms for solving nonlinear PIDEs have been presented in [15, 30]. However, none of these papers provide a complexity analysis of their algorithms. Moreover, very recently, [13] introduced for non-local nonlinear PDEs with Neumann boundary conditions both a deep splitting algorithm as well as a multilevel Picard approximation algorithm to numerically solve the PDEs under consideration. However, no convergence analysis nor a complexity analysis of their algorithms has been provided. Moreover, the nonlocal term is expressed through a probability measure, which corresponds to a normalized Lévy measure of *finite activity*.

In this paper, we develop a multilevel Picard approximation (MLP) algorithm for semilinear PIDEs, and provide both a convergence analysis and a complexity analysis of the methodology. We show that the algorithm does not suffer from the curse of dimensionality, i.e. the computational complexity of the algorithm is bounded polynomially in the dimension d and the prescribed (reciprocal of the) accuracy ε . Similar to the ideas in [8, 11, 26, 38, 39, 40], the approach of developing such an approximation method for PIDEs is the following. Consider the following nonlinear PIDE on $[0, T] \times \mathbb{R}^d$ satisfying

Key words and phrases. Multilevel Picard approximation, Nonlinear PIDE, overcoming the curse of dimensionality, Monte Carlo methods, Feynman-Kac representation.

Financial support by the Nanyang Assistant Professorship Grant (NAP Grant) *Machine Learning based Algorithms in Finance and Insurance* is gratefully acknowledged.

$u^d(T, x) = g^d(x)$ and

$$\begin{aligned} \frac{\partial}{\partial t} u^d(t, x) + \langle \nabla u^d(t, x), \mu^d(t, x) \rangle + \frac{1}{2} \text{Trace}(\sigma^d(t, x) [\sigma^d(t, x)]^T \text{Hess}_x(u^d)(t, x)) + f^d(t, x, u^d(t, x)) \\ + \int_{\mathbb{R}^d} \left(u^d(t, x + \eta_t^d(z, x)) - u^d(t, x) - \langle \nabla u^d(t, x), \eta_t^d(z, x) \rangle \right) \nu^d(dz) = 0. \end{aligned} \quad (1.1)$$

In this paper, we restrict ourselves to semilinear parabolic PIDEs which are only nonlinear in their solutions $u(t, x)$ but not in their derivatives. To be able to obtain a converge result of our numerical method, we first prove that under suitable conditions on the coefficients (see Section 2.1), the PIDE has a unique viscosity solution, and its unique solution admits a Feynman-Kac representation

$$u^d(t, x) = \mathbb{E} \left[g^d(X_T^{d,t,x}) \right] + \int_t^T \mathbb{E} \left[f^d(s, X_s^{d,t,x}, u^d(s, X_s^{d,t,x})) \right] ds, \quad (1.2)$$

where

$$dX_s^{d,t,x} = \mu^d(s, X_{s-}^{d,t,x}) ds + \sigma^d(s, X_{s-}^{d,t,x}) dW_s^d + \int_{\mathbb{R}^d} \eta_s^d(X_{s-}^{d,t,x}, z) \tilde{\pi}^d(dz, ds), \quad (1.3)$$

with $\tilde{\pi}^d(dz, dt) := \pi^d(dz, dt) - \nu^d(dz) \otimes dt$ being the compensated Poisson random measure of $\pi^d(dz, dt)$. The comparison result is obtained by an application of the result in [44]. The existence of a viscosity solution and its Feynman-Kac representation is established by first obtaining the existence of smooth solutions and their Feynman-Kac representations for PIDEs with smooth coefficients [21] and then applying a limit argument involving mollifiers as well as a stochastic fixed point argument based on [12, 39]; see Section 5.

Motivated by the Feynman-Kac representation (1.2), one can then define the following map (from a suitable space of functions)

$$(\Phi^d \circ v^d)(t, x) := \mathbb{E} \left[g^d(X_T^{d,t,x}) \right] + \int_t^T \mathbb{E} \left[f^d(s, X_s^{d,t,x}, v^d(s, X_s^{d,t,x})) \right] ds$$

for which the solution u^d of the PIDE is a fixpoint. This motivates us to define a sequence of Picard iterations

$$u_k^d(t, x) = (\Phi^d \circ u_{k-1}^d)(t, x).$$

The above sequence of Picard iterations is well studied, see e.g. [39], for the case without jumps, where it was shown that under appropriate assumptions on the coefficients of the PDE under consideration, Φ^d forms a contraction mapping on some Banach space and $\lim_{k \rightarrow \infty} u_k^d(t, x) = u^d(t, x)$ for all $(t, x) \in [0, T] \times \mathbb{R}^d$ (see, e.g., Proposition 2.2 in [39] for details). Furthermore, notice that

$$\begin{aligned} u_k^d(t, x) &= u_1^d(t, x) + \sum_{l=1}^{k-1} \left[u_{l+1}^d(t, x) - u_l^d(t, x) \right] \\ &= (\Phi^d \circ u_0^d)(t, x) + \sum_{l=1}^{k-1} \left[(\Phi^d \circ u_l^d)(t, x) - (\Phi^d \circ u_{l-1}^d)(t, x) \right] \\ &= \mathbb{E} \left[g^d(X_T^{d,t,x}) \right] + \sum_{l=0}^{k-1} \int_t^T \mathbb{E} \left[f^d(s, X_s^{d,t,x}, u_l^d(s, X_s^{d,t,x})) \right. \\ &\quad \left. - \mathbf{1}_{\mathbb{N}}(l) f^d(s, X_s^{d,t,x}, u_{l-1}^d(s, X_s^{d,t,x})) \right] ds \end{aligned} \quad (1.4)$$

with $u_0^d \equiv 0$. Then, as suggested by Hutzenthaler et al. (see Theorem 1.1 in [39] for PDEs without non-local operators), one can introduce a multilevel Picard approximation scheme of $u^d(t, x)$ by replacing $(X_s^{d,t,x})_{s \in [t, T]}$ in (1.4) with its (time-) discretized approximation, and by approximating the expectations and time integrals in (1.4) by Monte Carlo methods. To be able to simulate the jumps of the SDE in (1.3), we propose to use the Monte Carlo approximation method of the compensator integral as in [34, Section 4.4]. Since Euler-discretizations and Monte Carlo approximations do not suffer from the curse of dimensionality, we are then able to show that the corresponding complexity of the algorithm only grows polynomially in the dimension d and the prescribed (reciprocal of the) accuracy ε , extending the results in [8, 11, 26, 38, 39, 40] to PIDEs.

The remainder of this paper is organised as follows. In Sections 2.1 and 2.2, we introduce the settings of the MLP approximations, and formulate the main results of this paper (see Theorems 2.9 and 2.10). Some pseudo-codes for MLP approximations are collected in Section 2.3. In Section 3, we show some lemmas for important estimates and properties of SDEs with jumps and their Euler approximations, which will be used in Sections 5 and 7. In Section 4, some useful results for stochastic fixed point equations are presented. In Section 5, we investigate the existence and uniqueness of viscosity solutions of semilinear PIDE (1.1), and establish a Feynman-Kac formula for semilinear PIDEs (see Proposition 5.16). Section 6 presents an error bound for MLP approximations not involving Euler scheme which is taken from [39]. The proof of Theorems 2.9 and 2.10 are given in Section 7.

Notation. In conclusion, we introduce some notations used throughout this paper. We denote by \mathbb{N} and \mathbb{N}_0 the set of all positive integers and the set of all natural numbers, respectively, and denote by \mathbb{Z} the set of all integers. For each $d \in \mathbb{N}$, the $d \times d$ identity matrix is denoted by \mathbf{I}_d . For each $d \in \mathbb{N}$, we use \mathbb{S}^d to denote the space of $d \times d$ symmetric matrices, and for matrices $A, B \in \mathbb{S}^d$ the notation $A \geq B$ means $A - B$ is positive semi-definite. For each $d \in \mathbb{N}$ and any vectors $a, b \in \mathbb{R}^d$, we denote by $\langle a, b \rangle$ the scalar product of a and b , and denote by $\|a\|$ the euclidean norm of a . For each $d \in \mathbb{N}$ and every matrix $A \in \mathbb{R}^{d \times d}$, we denote by $\|A\|_F$ the Frobenius norm of A , and we use A^{ij} to denote the element on the i -th row and j -th column of A for $i, j = 1, \dots, d$. For each $d \in \mathbb{N}$ and every matrix $A \in \mathbb{R}^{d \times d}$, we denote by A^T the transpose of A . For any metric spaces (E, d_E) and (F, d_F) , we denote by $C(E, F)$ the set of continuous functions from E to F . For every topological space E , denote by $\mathcal{B}(E)$ the Borel σ -algebra of E . For all measurable spaces (X_1, Σ_1) and (X_2, Σ_2) , we denote by $\mathcal{M}(\Sigma_1, \Sigma_2)$ the set of Σ_1/Σ_2 -measurable functions from X_1 to X_2 . For all $a, b \in \mathbb{R}$, we use the notations $a \wedge b := \min\{a, b\}$ and $a \vee b := \max\{a, b\}$. For any set B , we use $\mathbf{1}_B$ to denote the indicator function of B .

Let \mathbb{T} denote either $(0, T)$ or $[0, T]$, and denote by $LSC(\mathbb{T} \times \mathbb{R}^d)$ ($USC(\mathbb{T} \times \mathbb{R}^d)$) the space of lower (upper) semicontinuous functions $u : \mathbb{T} \times \mathbb{R}^d \rightarrow \mathbb{R}$. We denote by $LSC_1(\mathbb{T} \times \mathbb{R}^d)$ ($USC_1(\mathbb{T} \times \mathbb{R}^d)$) the space of functions $u \in LSC(\mathbb{T} \times \mathbb{R}^d)$ ($USC(\mathbb{T} \times \mathbb{R}^d)$) satisfying the linear growth condition

$$\sup_{(t,x) \in \mathbb{T} \times \mathbb{R}^d} \frac{|u(t, x)|}{1 + \|x\|} < \infty, \quad (1.5)$$

and we denote by $C_1(\mathbb{T} \times \mathbb{R}^d)$ the space of continuous functions $u : \mathbb{T} \times \mathbb{R}^d \rightarrow \mathbb{R}$ satisfying (1.5). Moreover, $SC(\mathbb{T} \times \mathbb{R}^d) := LSC(\mathbb{T} \times \mathbb{R}^d) \cup USC(\mathbb{T} \times \mathbb{R}^d)$, and $SC_1(\mathbb{T} \times \mathbb{R}^d) := LSC_1(\mathbb{T} \times \mathbb{R}^d) \cup USC_1(\mathbb{T} \times \mathbb{R}^d)$. We use the notation $C^{1,2}(\mathbb{T} \times \mathbb{R}^d)$ to denote the space of once in time $t \in \mathbb{T}$ and twice in space $x \in \mathbb{R}^d$ continuously differentiable functions $u : \mathbb{T} \times \mathbb{R}^d \rightarrow \mathbb{R}$, and we also use the notation $C_1^{1,2}(\mathbb{T} \times \mathbb{R}^d) := C^{1,2}(\mathbb{T} \times \mathbb{R}^d) \cap C_1(\mathbb{T} \times \mathbb{R}^d)$. Moreover, we denote by $C_1^2(\mathbb{R}^d)$ the space of twice continuously differentiable functions $u : \mathbb{R}^d \rightarrow \mathbb{R}$ satisfying (1.5) (without dependence on t). The notation $C_c^\infty(\mathbb{R}^d)$ means the space of infinitely differentiable real-valued functions on \mathbb{R}^d with compact support.

2. MULTILEVEL PICARD APPROXIMATION SCHEME: SETTING AND MAIN RESULTS

2.1. Setting. Let $T > 0$ be a fixed constant, and let (Ω, \mathcal{F}, P) be a complete probability space equipped with a filtration $\mathbb{F} := (\mathcal{F}_t)_{t \in [0, T]}$ satisfying the usual conditions. For each $d \in \mathbb{N}$ we are given an \mathbb{R}^d -valued standard \mathbb{F} -Brownian motion denoted by $(W_t^d)_{t \in [0, T]}$. For each $d \in \mathbb{N}$ let $\nu^d(dz)$ be a Lévy measure on \mathbb{R}^d , and let $\pi^d(dz, dt)$ be a \mathbb{F} -Poisson random measure on $([0, T] \times \mathbb{R}^d, \mathcal{B}([0, T]) \otimes \mathcal{B}(\mathbb{R}^d))$ with intensity measure $\nu^d(dz) \otimes dt$, and we denote by $\tilde{\pi}^d(dz, dt) := \pi^d(dz, dt) - \nu^d(dz) \otimes dt$ the compensated Poisson random measure of $\pi^d(dz, dt)$. Moreover, for each $d \in \mathbb{N}$ let $\mu^d = (\mu^{d,1}, \dots, \mu^{d,d}) \in C([0, T] \times \mathbb{R}^d, \mathbb{R}^d)$, $\sigma^d = (\sigma^{d,ij})_{i,j \in \{1,2,\dots,d\}} \in C([0, T] \times \mathbb{R}^d, \mathbb{R}^{d \times d})$, and $\eta^d = (\eta^{d,1}, \dots, \eta^{d,d}) \in C([0, T] \times \mathbb{R}^d \times \mathbb{R}^d, \mathbb{R}^d)$. Then for each $d \in \mathbb{N}$ and $(t, x) \in [0, T] \times \mathbb{R}^d$ consider the following stochastic differential equation (SDE) on $[t, T]$

$$dX_s^{d,t,x} = \mu^d(s, X_{s-}^{d,t,x}) ds + \sigma^d(s, X_{s-}^{d,t,x}) dW_s^d + \int_{\mathbb{R}^d} \eta_s^d(X_{s-}^{d,t,x}, z) \tilde{\pi}^d(dz, ds) \quad (2.1)$$

with initial condition $X_t^{d,t,x} = x$. Let $g^d \in C(\mathbb{R}^d, \mathbb{R})$ and $f^d \in C([0, T] \times \mathbb{R}^d \times \mathbb{R}, \mathbb{R})$. Then we make the following assumptions for the coefficient functions and Lévy measures.

Regularity and growth conditions.

Assumption 2.1 (Lipschitz and linear growth conditions). *There exists constants $L, p \in (0, \infty)$ satisfying for all $d \in \mathbb{N}$, $x, y \in \mathbb{R}^d$, $v, w \in \mathbb{R}$, and $t \in [0, T]$ that*

$$T|f^d(t, x, v) - f^d(t, y, w)| + |g^d(x) - g^d(y)| \leq L^{1/2}(T|v - w| + T^{-1/2}\|x - y\|), \quad (2.2)$$

$$\|\mu^d(t, x) - \mu^d(t, y)\|^2 + \|\sigma^d(t, x) - \sigma^d(t, y)\|_F^2 + \int_{\mathbb{R}^d} \|\eta_t^d(x, z) - \eta_t^d(y, z)\|^2 \nu^d(dz) \leq L\|x - y\|^2, \quad (2.3)$$

$$\|\mu^d(t, x)\|^2 + \|\sigma^d(t, x)\|_F^2 + \int_{\mathbb{R}^d} \|\eta_t^d(x, z)\|^2 \nu^d(dz) \leq Ld^p(1 + \|x\|^2), \quad (2.4)$$

and

$$|T \cdot f^d(t, x, 0)|^2 + |g^d(x)|^2 \leq L(d^p + \|x\|^2). \quad (2.5)$$

Assumption 2.2 (Pointwise Lipschitz and integrability conditions). *For each $d \in \mathbb{N}$ there exists a constant $C_d \in (0, \infty)$ such that for all $x, x' \in \mathbb{R}^d$, $t \in [0, T]$, and $z \in \mathbb{R}^d$*

$$\|\eta_t^d(x, z)\|^2 \leq C_d(1 \wedge \|z\|^2), \quad \|\eta_t^d(x, z) - \eta_t^d(x', z)\|^2 \leq C_d\|x - x'\|^2(1 \wedge \|z\|^2). \quad (2.6)$$

Assumption 2.3. *For all $(t, x) \in [0, T] \times \mathbb{R}^d$ and $z \in \mathbb{R}^d$ we assume that $D_x \eta_t(x, z)$ exists, where $D_x \eta_t(x, z)$ denotes the Jacobian matrix of $\eta_t(x, z)$ with respect to x . Moreover, assume that there exists a constant $\lambda > 0$ such that for all $(t, x) \in [0, T] \times \mathbb{R}^d$, $z \in \mathbb{R}^d$, and $\gamma \in [0, 1]$*

$$\lambda \leq |\det(\mathbf{I}_d + \gamma D_x \eta_t(x, z))|. \quad (2.7)$$

Remark 2.4. *Note that Assumption 2.3 is needed to show the existence and uniqueness of linear PIDEs whose coefficient functions are infinitely differentiable and compactly supported as functions of $x \in \mathbb{R}^d$ (see Proposition 5.15). Moreover, note that Assumption 2.3 is satisfied in the Lévy case where $\eta_t(x, z) = z$ or $\eta_t(x, z) = z \mathbf{1}_{\{\|z\| \leq 1\}}$ for all $(t, x) \in [0, T] \times \mathbb{R}^d$ and $z \in \mathbb{R}^d$.*

Assumption 2.5. *There exist constants $K, q \in (0, \infty)$ satisfying for all $d \in \mathbb{N}$, $\delta \in (0, 1)$, and $(t, x) \in [0, T] \times \mathbb{R}^d$ that*

$$\int_{\mathbb{R}^d} (1 \wedge \|z\|^2) \nu^d(dz) \leq Kd^p \quad \text{and} \quad \int_{\|z\| \leq \delta} \|\eta_t^d(x, z)\|^2 \nu^d(dz) \leq Kd^p \delta^q (1 + \|x\|^2). \quad (2.8)$$

Note that Assumptions 2.1 and 2.2 guarantees that for each $d \in \mathbb{N}$, the SDE in (2.1) has a unique solution satisfying that

$$\mathbb{E} \left[\sup_{s \in [t, T]} \|X_s^{d, t, x}\|^2 \right] < C(1 + \|x\|^2), \quad (2.9)$$

where C is a constant only depending on L, d, p , and T (see, e.g., Theorem 3.1 and Theorem 3.2 in [47], or Lemma 114 and Theorem 117 in [55]). The next assumption is needed to obtain the rate of convergence of the Euler approximation of SDE (2.1), in the case that the coefficients μ^d, σ^d and η^d depend on time t (see Lemma 3.6 in Section 3).

Assumption 2.6 (Temporal 1/2-Hölder condition). *There exist constants $L_1, L_2 > 0$ satisfying for all $d \in \mathbb{N}$, $x \in \mathbb{R}^d$, $z \in \mathbb{R}^d$, and $(s, s') \in [0, T]^2$ that*

$$\|\mu^d(s, x) - \mu^d(s', x)\|^2 + \|\sigma^d(s, x) - \sigma^d(s', x)\|_F^2 \leq L_1|s - s'|, \quad (2.10)$$

and

$$\int_{\mathbb{R}^d} \|\eta_s^d(x, z) - \eta_{s'}^d(x, z)\|^2 \nu^d(dz) \leq L_2|s - s'|. \quad (2.11)$$

For each $d \in \mathbb{N}$ let

$$F^d : \mathcal{M}(\mathcal{B}([0, T] \times \mathbb{R}^d), \mathcal{B}(\mathbb{R})) \rightarrow \mathcal{M}(\mathcal{B}([0, T] \times \mathbb{R}^d), \mathcal{B}(\mathbb{R}))$$

be the operator such that

$$[0, T] \times \mathbb{R}^d \ni (t, x) \mapsto (F^d(v))(t, x) := f^d(t, x, v(t, x)), \quad v \in \mathcal{M}(\mathcal{B}([0, T] \times \mathbb{R}^d), \mathcal{B}(\mathbb{R})).$$

Time Discretization Approximations. Let $\Theta = \cup_{n \in \mathbb{N}} \mathbb{Z}^n$ be an index set which we will use for the families of independent random variables needed for the Monte Carlo approximations. For each $m \in \mathbb{N}$, let $W^{m,\theta} = (W^{m,\theta,1}, \dots, W^{m,\theta,d}) : [0, T] \times \Omega \rightarrow \mathbb{R}^m, \theta \in \Theta$, be independent \mathbb{R}^m -valued standard \mathbb{F} -Brownian motions. For each $d \in \mathbb{N}$, let $\pi^{d,\theta}(dz, ds), \theta \in \Theta$, be independent \mathbb{F} -Poisson random measures on $\mathbb{R}^d \times [0, T]$ with intensity measures $\nu^d(dz) \otimes dt$, respectively, and denote by $\tilde{\pi}^{d,\theta}(dz, ds) := \pi^{d,\theta}(dz, ds) - \nu^d(dz) \otimes dt$ the compensated Poisson random measures of $\pi^{d,\theta}(dz, ds)$. For each $d \in \mathbb{N}$ and $(t, x) \in [0, T] \times \mathbb{R}^d$, let $(X_s^{d,\theta,t,x})_{s \in [t, T]} : [t, T] \times \Omega \rightarrow \mathbb{R}^d, \theta \in \Theta$, be $\mathcal{B}([0, T]) \otimes \mathcal{F}/\mathcal{B}(\mathbb{R}^d)$ -measurable functions satisfying for all $\theta \in \Theta$ and $s \in [t, T]$ almost surely that $X_t^{d,\theta,t,x} = x$ and

$$dX_s^{d,\theta,t,x} = \mu^d(s, X_{s-}^{d,\theta,t,x}) ds + \sigma^d(s, X_{s-}^{d,\theta,t,x}) dW_s^{d,\theta} + \int_{\mathbb{R}^d} \eta_s^d(X_{s-}^{d,\theta,t,x}, z) \tilde{\pi}^{d,\theta}(dz, ds). \quad (2.12)$$

For each $d \in \mathbb{N}, N \in \mathbb{N}$, and $(t, x) \in [0, T] \times \mathbb{R}^d$, let $(Y_s^{d,\theta,t,x,N})_{s \in [t, T]} : [t, T] \times \Omega \rightarrow \mathbb{R}^d, \theta \in \Theta$, be measurable functions satisfying for all $n \in \mathbb{N}_0, s \in [t + \frac{n(T-t)}{N}, t + \frac{(n+1)(T-t)}{N}] \cap [t, T]$ that $Y_t^{d,\theta,t,x,N} = x$ and

$$\begin{aligned} Y_s^{d,\theta,t,x,N} &= Y_{t+\frac{n(T-t)}{N}}^{d,\theta,t,x,N} + \mu^d\left(t + \frac{n(T-t)}{N}, Y_{t+\frac{n(T-t)}{N}}^{d,\theta,t,x,N}\right) \left[s - \left(t + \frac{n(T-t)}{N}\right)\right] \\ &\quad + \sigma^d\left(t + \frac{n(T-t)}{N}, Y_{t+\frac{n(T-t)}{N}}^{d,\theta,t,x,N}\right) \left(W_s^{d,\theta} - W_{t+\frac{n(T-t)}{N}}^{d,\theta}\right) \\ &\quad + \int_{t+\frac{n(T-t)}{N}}^s \int_{\mathbb{R}^d} \eta_{t+\frac{n(T-t)}{N}}^d\left(Y_{t+\frac{n(T-t)}{N}}^{d,\theta,t,x,N}, z\right) \tilde{\pi}^{d,\theta}(dz, dr). \end{aligned} \quad (2.13)$$

To ease notations we define

$$\kappa_N(s) := t + \frac{\lfloor N(s-t)/(T-t) \rfloor \cdot (T-t)}{N}, \quad s \in [t, T],$$

where $\lfloor y \rfloor := \max\{n \in \mathbb{N}_0 : n \leq y\}$ for $y \in [0, \infty)$. Then for all $d, N \in \mathbb{N}, \theta \in \Theta$, and $(t, x) \in [0, T] \times \mathbb{R}^d$, (2.13) can be written as

$$\begin{aligned} dY_s^{d,\theta,t,x,N} &= \mu^d\left(\kappa_N(s-), Y_{\kappa_N(s-)}^{d,\theta,t,x,N}\right) ds + \sigma^d\left(\kappa_N(s-), Y_{\kappa_N(s-)}^{d,\theta,t,x,N}\right) dW_s^{d,\theta} \\ &\quad + \int_{\mathbb{R}^d} \eta_{\kappa_N(s-)}^d\left(Y_{\kappa_N(s-)}^{d,\theta,t,x,N}, z\right) \tilde{\pi}^{d,\theta}(dz, ds), \quad s \in [t, T]. \end{aligned} \quad (2.14)$$

Although (2.14) provides a first approximation to (2.12), one is not able to directly simulate the items of the integrals with respect to the compensated Poisson random measures. To find an appropriate approximation of (2.12) that allows us to simulate it explicitly, we follow the idea of L.Gonon and C. Schwab (see, Sections 4.2 and 4.4 in [34]), and introduce the following approximation procedures. For each $d \in \mathbb{N}$ and $\delta \in (0, 1)$ we define the set A_δ^d by

$$A_\delta^d := \{z \in \mathbb{R}^d : \|z\| \geq \delta\},$$

and also define a probability measure ν_δ^d on $(\mathbb{R}^d, \mathcal{B}(\mathbb{R}^d))$ by

$$\nu_\delta^d(B) := \frac{\nu^d(B \cap A_\delta^d)}{\nu^d(A_\delta^d)}, \quad B \in \mathcal{B}(\mathbb{R}^d).$$

Then, for each $d \in \mathbb{N}, (t, x) \in [0, T] \times \mathbb{R}^d$, and $\delta \in (0, 1)$, let $(Y_s^{d,\theta,t,x,N,\delta})_{s \in [t, T]} : [t, T] \times \Omega \rightarrow \mathbb{R}^d, \theta \in \Theta$, be measurable functions satisfying that $Y_t^{d,\theta,t,x,N,\delta} = x$ and almost surely for all $s \in [t, T]$

$$\begin{aligned} dY_s^{d,\theta,t,x,N,\delta} &= \mu^d\left(\kappa_N(s-), Y_{\kappa_N(s-)}^{d,\theta,t,x,N,\delta}\right) ds + \sigma^d\left(\kappa_N(s-), Y_{\kappa_N(s-)}^{d,\theta,t,x,N,\delta}\right) dW_s^{d,\theta} \\ &\quad + \int_{A_\delta^d} \eta_{\kappa_N(s-)}^d\left(Y_{\kappa_N(s-)}^{d,\theta,t,x,N,\delta}, z\right) \pi^{d,\theta}(dz, ds) \\ &\quad - \int_{A_\delta^d} \eta_{\kappa_N(s-)}^d\left(Y_{\kappa_N(s-)}^{d,\theta,t,x,N,\delta}, z\right) \nu^d(dz) ds. \end{aligned} \quad (2.15)$$

Furthermore, for each $d, N, \mathcal{M} \in \mathbb{N}$ and $\delta \in (0, 1)$ let $V_{i,j}^{d,\theta,N,\delta,\mathcal{M}}, \theta \in \Theta, i = 0, 1, 2, \dots, N, j = 1, 2, \dots, \mathcal{M}$, be independent \mathbb{R}^d -valued random variables with identical distribution ν_δ^d , independent

of $(W^{d,\theta}, \pi^{d,\theta})_{(\theta,d) \in \Theta \times \mathbb{N}}$. For each $d \in \mathbb{N}$, $(t, x) \in [0, T] \times \mathbb{R}^d$, $N, \mathcal{M} \in \mathbb{N}$, and $\delta \in (0, 1)$, let $(Y_s^{d,\theta,t,x,N,\delta,\mathcal{M}})_{s \in [t,T]} : [t, T] \times \Omega \rightarrow \mathbb{R}^d$, $\theta \in \Theta$, be measurable functions satisfying that $Y_t^{d,\theta,t,x,N,\delta,\mathcal{M}} = x$ and almost surely for all $s \in [t, T]$

$$\begin{aligned} dY_s^{d,\theta,t,x,N,\delta,\mathcal{M}} &= \mu^d \left(\kappa_N(s-), Y_{\kappa_N(s-)}^{d,\theta,t,x,N,\delta,\mathcal{M}} \right) ds + \sigma^d \left(\kappa_N(s-), Y_{\kappa_N(s-)}^{d,\theta,t,x,N,\delta,\mathcal{M}} \right) dW_s^{d,\theta} \\ &\quad + \int_{A_\delta^d} \eta_{\kappa_N(s-)}^d \left(Y_{\kappa_N(s-)}^{d,\theta,t,x,N,\delta,\mathcal{M}}, z \right) \pi^{d,\theta}(dz, ds) \\ &\quad - \left(\frac{\nu^d(A_\delta^d)}{\mathcal{M}} \sum_{j=1}^{\mathcal{M}} \eta_{\kappa_N(s-)}^d \left(Y_{\kappa_N(s-)}^{d,\theta,t,x,N,\delta,\mathcal{M}}, V_{(\kappa_N(s-)-t)N,j}^{d,\theta,N,\delta,\mathcal{M}} \right) \right) ds. \end{aligned}$$

Multilevel Picard Approximation Scheme. Next, we introduce the multi-level Picard approximations. Let $\xi^\theta : \Omega \rightarrow [0, 1]$, $\theta \in \Theta$, be i.i.d. uniformly distributed random variables, and assume that $(\xi^\theta)_{\theta \in \Theta}$ is independent of $(W^{d,\theta}, \pi^{d,\theta})_{(\theta,d) \in \Theta \times \mathbb{N}}$. Let $\mathcal{R}^\theta : [0, T] \times \Omega \rightarrow [0, T]$, $\theta \in \Theta$, satisfy that $\mathcal{R}_t^\theta = t + (T-t)\xi^\theta$ for all $t \in [0, T]$ and $\theta \in \Theta$. For each $d \in \mathbb{N}$, $M, N \in \mathbb{N}$, $\theta \in \Theta$, $(t, x) \in [0, T] \times \mathbb{R}^d$, and $s \in [t, T]$, let $(Y_s^{(d,\theta,t,x,N,\delta,\mathcal{M},l,i)})_{(l,i) \in \mathbb{N} \times (\mathbb{Z}/\{0\})}$ and $(\mathcal{R}_t^{(\theta,l,i)})_{(l,i) \in \mathbb{N} \times \mathbb{N}_0}$ be independent copies of $Y_s^{d,\theta,t,x,N,\delta,\mathcal{M}}$ and \mathcal{R}_t^θ , respectively.

For each $d \in \mathbb{N}$, $n \in \mathbb{N}_0$, $M, N, \mathcal{M} \in \mathbb{N}$, $\theta \in \Theta$, $\delta \in (0, 1)$, let $U_{n,M,N}^{d,\theta,\delta,\mathcal{M}} : [0, T] \times \mathbb{R}^d \times \Omega \rightarrow \mathbb{R}$ be measurable functions satisfying for all $(t, x) \in \mathbb{R}^d$ that

$$\begin{aligned} U_{n,M,N}^{d,\theta,\delta,\mathcal{M}}(t, x) &= \frac{\mathbf{1}_{\mathbb{N}}(n)}{M^n} \sum_{i=1}^{M^n} g^d \left(Y_T^{(d,\theta,t,x,N,\delta,\mathcal{M},0,-i)} \right) \\ &\quad + \sum_{l=0}^{n-1} \frac{(T-t)}{M^{n-l}} \left[\sum_{i=1}^{M^{n-l}} \left(F \left(U_{l,M,N}^{(d,\theta,\delta,\mathcal{M},l,i)} \right) - \mathbf{1}_{\mathbb{N}}(l) F \left(U_{l-1,M,N}^{(d,\theta,\delta,\mathcal{M},-l,i)} \right) \right) \left(\mathcal{R}_t^{(\theta,l,i)}, Y_{\mathcal{R}_t^{(\theta,l,i)}}^{(d,\theta,t,x,N,\delta,\mathcal{M},l,i)} \right) \right], \end{aligned} \quad (2.16)$$

where $(U_{n,M,N}^{d,\theta,\delta,\mathcal{M},l,i})_{(l,i) \in \mathbb{Z} \times \mathbb{N}_0}$ are independent copies of $U_{n,M,N}^{d,\theta,\delta,\mathcal{M}}$.

Viscosity solutions of PIDEs. For every $d \in \mathbb{N}$, let $G^d : (0, T) \times \mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d \times \mathbb{S}^d \times C^2(\mathbb{R}^d) \rightarrow \mathbb{R}$ be a functional defined for all $(t, x, r, y, A, \phi) \in (0, T) \times \mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d \times \mathbb{S}^d \times C^2(\mathbb{R}^d)$ by

$$\begin{aligned} G^d(t, x, r, y, A, \phi) &:= - \langle y, \mu^d(t, x) \rangle - \frac{1}{2} \text{Trace}(\sigma^d(t, x) [\sigma^d(t, x)]^T A) - f^d(t, x, r) \\ &\quad - \int_{\mathbb{R}^d} \left(\phi(x + \eta_t^d(x, z)) - \phi(x) - \langle \nabla_x \phi(x), \eta_t^d(x, z) \rangle \right) \nu^d(dz). \end{aligned} \quad (2.17)$$

Then for every $d \in \mathbb{N}$ we consider the semilinear PIDE of parabolic type

$$- \frac{\partial}{\partial t} u^d(t, x) + G^d(t, x, u^d(t, x), \nabla_x u^d(t, x), \text{Hess}_x u^d(t, x), u^d(t, \cdot)) = 0 \quad \text{on } (0, T) \times \mathbb{R}^d, \quad (2.18)$$

$$u^d(T, x) = g^d(x) \quad \text{on } \mathbb{R}^d. \quad (2.19)$$

We use the following definition of viscosity solutions of PIDE (2.18) (cf. Definition 2.1 in [44]).

Definition 2.7. Let $d \in \mathbb{N}$. A function $u^d \in USC_1((0, T) \times \mathbb{R}^d)$ ($u^d \in LSC_1((0, T) \times \mathbb{R}^d)$) is called a viscosity subsolution (supersolution) of PIDE (2.18) if for every $(t, x) \in (0, T) \times \mathbb{R}^d$ and $\varphi \in C^{1,2}((0, T) \times \mathbb{R}^d)$ such that $\varphi(t, x) = u^d(t, x)$, and $u^d \leq \varphi$ ($u^d \geq \varphi$), we have that

$$- \frac{\partial}{\partial t} \varphi(t, x) + G^d(t, x, \varphi(t, x), \nabla_x \varphi(t, x), \text{Hess}_x \varphi(t, x), \varphi(t, \cdot)) \leq 0 \quad (\geq 0).$$

A function $u^d : (0, T) \times \mathbb{R}^d \rightarrow \mathbb{R}$ is said to be a viscosity solution of PIDE (2.18) if u is both a viscosity subsolution and a viscosity supersolution of (2.18).

Remark 2.8. Let $d \in \mathbb{N}$. If $u^d \in C_1([0, T] \times \mathbb{R}^d)$ is a viscosity solution of (2.18) with $u^d(T, x) = g^d(x)$ for $x \in \mathbb{R}^d$, then the function $v^d(t, x) := u^d(T-t, x)$, $(t, x) \in [0, T] \times \mathbb{R}^d$, satisfies the follows.

(i) $v^d(0, x) = g^d(x)$ for $x \in \mathbb{R}^d$.

- (ii) For every $(t, x) \in (0, T) \times \mathbb{R}^d$ and $\varphi \in C^{1,2}((0, T) \times \mathbb{R}^d)$ such that $\varphi(t, x) = v^d(t, x)$ and $v^d \leq \varphi$ ($v^d \geq \varphi$), we have

$$\frac{\partial}{\partial t} \varphi(t, x) + G^d(t, x, \varphi(t, x), \nabla_x \varphi(t, x), \text{Hess}_x \varphi(t, x), \varphi(t, \cdot)) \leq 0 \quad (\geq 0).$$

The converse holds as well.

2.2. Main results. The following theorem shows that there exists a unique viscosity solution of PIDE (2.18), and that this solution satisfies a Feynman-Kac representation. This allows to prove that our MLP approximation scheme converges to the unique viscosity solution of PIDE (2.18).

Theorem 2.9. Assume Setting 2.1. Then the following holds.

- (i) For each $d \in \mathbb{N}$, $(t, x) \in [0, T] \times \mathbb{R}^d$, and $\theta \in \Theta$ there exists a unique \mathbb{F} -adapted \mathbb{R}^d -valued càdlàg process $(X_s^{d,\theta,t,x})_{s \in [t, T]}$ such that (2.12) holds.
- (ii) For each $d \in \mathbb{N}$ there exists a unique Borel function $u^d : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$ satisfying for all $(t, x) \in [0, T] \times \mathbb{R}^d$ that

$$\mathbb{E} \left[|g^d(X_T^{d,0,t,x})| \right] + \int_t^T \mathbb{E} \left[|f^d(s, X_s^{d,0,t,x}, u^d(s, X_s^{d,0,t,x}))| \right] ds + \sup_{y \in \mathbb{R}^d} \sup_{s \in [0, T]} \left(\frac{|u^d(s, y)|}{\sqrt{d^p + \|y\|^2}} \right) < \infty,$$

and

$$u^d(t, x) = \mathbb{E} \left[g^d(X_T^{d,0,t,x}) \right] + \int_t^T \mathbb{E} \left[f^d(s, X_s^{d,0,t,x}, u^d(s, X_s^{d,0,t,x})) \right] ds.$$

- (iii) For each $d \in \mathbb{N}$ there exists a unique viscosity solution $v^d \in C_1([0, T] \times \mathbb{R}^d)$ of PIDE (2.18) with $v^d(T, x) = g^d(x)$ for all $x \in \mathbb{R}^d$.
- (iv) It holds for all $d \in \mathbb{N}$ and $(t, x) \in [0, T] \times \mathbb{R}^d$ that $v^d(t, x) = u^d(t, x)$.
- (v) There exists a positive constant $c = c(T, L, L_1, L_2, K)$ satisfying for all $d \in \mathbb{N}$, $(t, x) \in [0, T] \times \mathbb{R}^d$, $\delta \in (0, 1)$, $n \in \mathbb{N}_0$, and $M, N, \mathcal{M} \in \mathbb{N}$ with $\mathcal{M} \geq \delta^{-2} K d^p$ that

$$\begin{aligned} & \left(\mathbb{E} \left[|U_{n,M,N}^{d,0,\delta,\mathcal{M}}(t, x) - u^d(t, x)|^2 \right] \right)^{1/2} \\ & \leq c(d^p + \|x\|^2) \left[e^{M/2} M^{-n/2} e^{2nTL^{1/2}} + (N^{-1}T + \delta^q d^p + \delta^{-2} K d^p \mathcal{M}^{-1})^{-1/2} \right]. \end{aligned} \quad (2.20)$$

The next theorem shows that the MLP approximation scheme defined by (2.16) can overcome the curse of dimensionality. To describe the computational complexity, for each $d \in \mathbb{N}$, $n \in \mathbb{N}_0$, and $M \in \mathbb{N}$ we introduce a natural number $\mathfrak{C}_{n,M}^{(d)}$ to denote the sum of the number of function evaluations of g^d , of the number of function evaluations of μ^d , of the number of function evaluations of σ^d , of the number of function evaluations of η^d , and of the number of realizations of scalar random variables used to obtain one realization of the MLP approximation algorithm in (2.16). Moreover, for each $d \in \mathbb{N}$ we use $\mathfrak{g}^{(d)}$ to denote the number of function evaluations of g^d , $\mathfrak{f}^{(d)}$ to denote the number of function evaluations of f^d , $\mathfrak{e}^{(d)}$ to denote the sum of the number of realizations of scalar random variables generated, of the number of function evaluations of μ^d , of the number of function evaluations of σ^d , and of the number of function evaluations of η^d .

Theorem 2.10. Assume the settings in Section 2.1. Moreover, for each $d \in \mathbb{N}$, let $\mathfrak{e}^{(d)}$, $\mathfrak{g}^{(d)}$, $\mathfrak{f}^{(d)}$, $n, M \in \mathbb{N}$, $\mathfrak{C}_{n,M}^{(d)} \in \mathbb{N}$, satisfy for all $n, M \in \mathbb{N}$ that

$$\mathfrak{C}_{n,M}^{(d)} \leq M^n (M^M \mathfrak{e}^{(d)} + \mathfrak{g}^{(d)}) \mathbf{1}_{\mathbb{N}}(n) + \sum_{l=0}^{n-1} [M^{n-l} (M^M \mathfrak{e}^{(d)} + \mathfrak{f}^{(d)} + \mathfrak{C}_{l,M}^{(d)} + \mathfrak{C}_{l-1,M}^{(d)})]. \quad (2.21)$$

Then, for all $d \in \mathbb{N}$ and $n \in \mathbb{N}$ we have

$$\sum_{k=1}^{n+1} \mathfrak{C}_{k,k}^{(d)} \leq 12(3\mathfrak{e}^{(d)} + \mathfrak{g}^{(d)} + 2\mathfrak{f}^{(d)}) 36^n n^{2n}. \quad (2.22)$$

Moreover, for each $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$, and $x \in \mathbb{R}^d$ there exists a positive integer $\mathbf{n}^d(x, \varepsilon) \geq 2$ such that for all $\gamma \in (0, 1]$ and $t \in [0, T]$,

$$\sup_{n \in [\mathbf{n}^d(x, \varepsilon), \infty) \cap \mathbb{N}} \left(\mathbb{E} \left[\left| U_{n, n, n^n}^{d, 0, n^{-n/q}, n^{n+2n/q}} K d^p(t, x) - u^d(t, x) \right|^2 \right] \right)^{1/2} < \varepsilon, \quad (2.23)$$

and

$$\begin{aligned} \left(\sum_{n=1}^{\mathbf{n}^d(x, \varepsilon)} \mathfrak{C}_{n, n}^{(d)} \right) \varepsilon^{\gamma+4} &\leq 12(3\mathfrak{e}^{(d)} + \mathfrak{g}^{(d)} + 2\mathfrak{f}^{(d)}) [c(d^p + \|x\|^2)]^{\gamma+4} \\ &\cdot \sup_{n \in \mathbb{N}} \left(36^n n^{-\gamma n/2} (e^{n/2} e^{2nTL^{1/2}} + (1+T+d^p)^{1/2})^{\gamma+4} \right) < \infty, \end{aligned} \quad (2.24)$$

where $c = c(T, L, L_1, L_2, K)$ is the constant introduced in (v) in Theorem 2.9.

2.3. Pseudocode. In this subsection we provide a pseudocode to show how the multilevel Picard approximations (2.16) can be implemented.

Algorithm 1 Multilevel Picard Approximation

```

1: function MLP( $t, x, n, M, N, \delta, \mathcal{M}$ )
2:    $t_1(j) \leftarrow (T-t)/N$  for all  $j \in \{0, \dots, N-1\}$ ; ▷ the length of time intervals in the time discretization
3:   for  $i \leftarrow 1$  to  $M^n$  do
4:      $X(i) \leftarrow x$ ;
5:     Generate  $N$  realizations  $W(j) \in \mathbb{R}^d, j \in \{0, \dots, N-1\}$ , of i.i.d. standard normal random vectors;
6:     Generate  $N$  realizations  $P(j) \in \mathbb{N}_0, j \in \{0, \dots, N-1\}$ , of i.i.d. Poisson random variables with parameter  $\nu^d(A_\delta^d)(T-t)N^{-1}$ ;
       (see, e.g., (1.8.3) in [52])
7:     for  $k \leftarrow 0$  to  $N-1$  do
8:       Generate  $P(k)$  realizations  $Z(j) \in \mathbb{R}^d, j \in \{1, \dots, P(k)\}$ , of i.i.d. random vectors with distribution  $\nu_\delta^d(dz)$ ;
9:       Generate  $\mathcal{M}$  realizations  $V(j) \in \mathbb{R}^d, j \in \{1, 2, \dots, \mathcal{M}\}$ , of i.i.d. random vectors with distribution  $\nu_\delta^d(dz)$ ;
10:       $X(i) \leftarrow X(i) + \mu^d(t+k(T-t)/N, X(i)) + \sigma^d(t+k(T-t)/N, X(i))\sqrt{t_1(k)} \cdot W(k)$ 
         $+ \sum_{j=1}^{P(k)} \eta_{t+k(T-t)/N}^d(X(i), Z(j)) - \frac{t_1(k)\nu^d(A_\delta^d)}{\mathcal{M}} \sum_{j=1}^{\mathcal{M}} \eta_{t+(T-t)k/N}^d(X(i), V(j))$ ;
11:    end for
12:  end for
13:  if  $n > 0$  then
14:     $u = \frac{1}{M^n} \sum_{i=1}^{M^n} g(X(i))$ ;
15:  else
16:     $u = 0$ ;
17:  end if
18:  for  $l \leftarrow 0$  to  $n-1$  do
19:     $Y(i) \leftarrow x$  for all  $i \in \{1, \dots, M^{n-l}\}$ ;
20:    Generate  $M^{n-l}$  realizations  $\mathcal{R}(i) \in [t, T], i \in 1, \dots, M^{n-l}$ , of i.i.d. random variables uniformly distributed on  $[t, T]$ ;
21:    for  $i \leftarrow 1$  to  $M^{n-l}$  do
22:       $S(i) \leftarrow \lfloor (\mathcal{R}(i) - t)N/(T-t) \rfloor + 1$ ; ▷ total time points of the time discretization of SDE
23:       $t_2(j) \leftarrow (T-t)/N$  for all  $j \in \{0, \dots, S(i)-2\}$ ; ▷ the length of the first  $S(i)-1$  time intervals
24:       $t_2(S(i)-1) \leftarrow \mathcal{R}(i) - \lfloor t + N^{-1}(T-t)(S(i)-1) \rfloor$ ; ▷ the length of the last time interval
25:      Generate  $S(i)$  realizations  $W(j) \in \mathbb{R}^d, j \in \{0, \dots, S(i)-1\}$ , of independent standard normal random vectors;
26:      Generate  $S(i)-1$  realizations  $P(j) \in \mathbb{N}_0, j \in \{0, \dots, S(i)-2\}$ , of i.i.d. Poisson random variables with parameter
         $\nu^d(A_\delta^d)(T-t)N^{-1}$ ;
27:      Generate one realization  $P(S_2(i)-1) \in \mathbb{N}_0$  of Poisson random variable with parameter  $\nu^d(A_\delta^d)t_2(S(i)-1)$ ;
28:      for  $k \leftarrow 0$  to  $S(i)-1$  do
29:        Generate  $P(k)$  realizations  $Z(j) \in \mathbb{R}^d, j \in \{1, \dots, P(k)\}$ , of i.i.d. random vectors with distribution  $\nu_\delta^d(dz)$ ;
30:        Generate  $\mathcal{M}$  realizations  $V(j) \in \mathbb{R}^d, j \in \{1, 2, \dots, \mathcal{M}\}$ , of i.i.d. random vectors with distribution  $\nu_\delta^d(dz)$ ;
31:         $Y(i) \leftarrow Y(i) + \mu^d(t+k(T-t)/N, Y(i)) + \sigma^d(t+k(T-t)/N, Y(i))\sqrt{t_2(k)} \cdot W(k)$ 
           $+ \sum_{j=1}^{P(k)} \eta_{t+k(T-t)/N}^d(Y(i), Z(j)) - \frac{t_2(k)\nu^d(A_\delta^d)}{\mathcal{M}} \sum_{j=1}^{\mathcal{M}} \eta_{t+k(T-t)/N}^d(Y(i), V(j))$ ;
32:      end for
33:    end for
34:     $u \leftarrow u + \frac{T-t}{M^{n-l}} \sum_{i=1}^{M^{n-l}} f(\mathcal{R}(i), Y(i), \text{MLP}(\mathcal{R}(i), Y(i), l, M, N, \delta, \mathcal{M}))$ ;
35:    if  $l > 0$  then
36:       $u \leftarrow u - \frac{T-t}{M^{n-l}} \sum_{i=1}^{M^{n-l}} f(\mathcal{R}(i), Y(i), \text{MLP}(\mathcal{R}(i), Y(i), l-1, M, N, \delta, \mathcal{M}))$ ;
37:    end if
38:  end for
39:  return  $u$ ;
40: end function

```

3. PRELIMINARIES: DIMENSION-DEPENDING BOUNDS FOR SDEs WITH JUMPS

Throughout this chapter, we assume the settings in Section 2.1, and fix d and θ . To ease notations, we will omit the superscripts d and θ for the notations introduced in Section 2.1 (e.g., for each $(t, x) \in [0, T] \times \mathbb{R}^d$, and $s \in [t, T]$, $X_s^{d,\theta,t,x}$ will be denoted by $X_s^{t,x}$). We follow the idea of [34] which provides dimension-dependent bounds for (time discretization of) SDEs with jumps but without time dependence for the coefficients.

3.1. Bounds for non-discretized SDEs.

Lemma 3.1. *For all $(t, x) \in \mathbb{R}^d$ and $s \in [t, T]$ it holds that*

$$\mathbb{E}[\|X_s^{t,x}\|^2] \leq (6Ld^p(s-t) + \|x\|^2)e^{(1+6L)(s-t)}. \quad (3.1)$$

Proof. We fix $(t, x) \in [0, T] \times \mathbb{R}^d$ throughout this proof. For $X^{t,x} = (X^{t,x,1}, \dots, X^{t,x,d})$, applying Itô's formula (see. e.g. Theorem 3.1 in [35]) to $\|X^{t,x}\|^2$ yields that almost surely for all $s \in [t, T]$

$$\begin{aligned} \|X_s^{t,x}\|^2 &= \|x\|^2 + \int_t^s \left(2\langle X_{r-}^{t,x}, \mu(r, X_{r-}^{t,x}) \rangle + \|\sigma(r, X_{r-}^{t,x})\|_F^2 \right) dr \\ &\quad + 2 \int_t^s \sum_{i=1}^d \sum_{j=1}^d X_{r-}^{t,x,i} \sigma^{ij}(r, X_{r-}^{t,x}) dW_r^j + 2 \int_t^s \int_{\mathbb{R}^d} \langle X_{r-}^{t,x}, \eta_r(X_{r-}^{t,x}, z) \rangle \tilde{\pi}(dz, dr) \\ &\quad + \int_t^s \int_{\mathbb{R}^d} \|\eta_r(X_{r-}^{t,x}, z)\|^2 \pi(dz, dr). \end{aligned} \quad (3.2)$$

Next, for each $n \in \mathbb{N}$ define the following stopping time

$$\begin{aligned} \tau_n^{t,x} &:= \inf \left\{ s \geq t : \int_t^s \sum_{i=1}^m \sum_{j=1}^d |2X_r^{t,x,i} \sigma^{ij}(r, X_r^{t,x})|^2 dr \right. \\ &\quad \left. + \int_t^s \int_{\mathbb{R}^d} \sum_{i=1}^d |2\eta_r^i(X_r^{t,x}, z) X_r^{t,x,i}|^2 \nu(dz) dr \geq n \right\} \wedge T. \end{aligned} \quad (3.3)$$

Then, with (3.3) we observe that the stochastic integrals with respect to the Brownian motion and the compensated Poisson random measure, respectively, on the right hand side of (3.2) are true martingales on $[0, \tau_n^{t,x}]$ for each $n \in \mathbb{N}$. Thus, by (3.2) we obtain for all $s \in [t, T]$ and $n \in \mathbb{N}$ that

$$\begin{aligned} \mathbb{E} \left[\left\| X_{s \wedge \tau_n^{t,x}}^{t,x} \right\|^2 \right] &= \|x\|^2 + 2\mathbb{E} \left[\int_t^{s \wedge \tau_n^{t,x}} \langle X_r^{t,x}, \mu(r, X_r^{t,x}) \rangle dr \right] + \mathbb{E} \left[\int_t^{s \wedge \tau_n^{t,x}} \|\sigma(r, X_r^{t,x})\|_F^2 dr \right] \\ &\quad + \mathbb{E} \left[\int_t^{s \wedge \tau_n^{t,x}} \int_{\mathbb{R}^d} \|\eta_r(X_r^{t,x}, z)\|^2 \nu(dz) dr \right]. \end{aligned} \quad (3.4)$$

Considering (2.3), (2.4), and (3.4), by Young's inequality and the fact that $(a+b)^2 \leq 2a^2 + 2b^2$ for all $a, b \in \mathbb{R}$, we have for all $s \in [t, T]$ and $n \in \mathbb{N}$ that

$$\begin{aligned} \mathbb{E} \left[\left\| X_{s \wedge \tau_n^{t,x}}^{t,x} \right\|^2 \right] &\leq \|x\|^2 + \int_t^s \mathbb{E} \left[\left\| X_{r \wedge \tau_n^{t,x}}^{t,x} \right\|^2 \right] dr \\ &\quad + 2\mathbb{E} \left[\int_t^{s \wedge \tau_n^{t,x}} \|\mu(r, X_r^{t,x}) - \mu(r, 0)\|^2 dr \right] + 2 \int_t^s \|\mu(r, 0)\|^2 dr \\ &\quad + 2\mathbb{E} \left[\int_t^{s \wedge \tau_n^{t,x}} \|\sigma(r, X_r^{t,x}) - \sigma(r, 0)\|_F^2 dr \right] + 2 \int_t^s \|\sigma(r, 0)\|_F^2 dr \\ &\quad + 2\mathbb{E} \left[\int_t^{s \wedge \tau_n^{t,x}} \int_{\mathbb{R}^d} \|\eta_r(X_r^{t,x}, z) - \eta_r(0, z)\|^2 \nu(dz) dr \right] \\ &\quad + 2 \int_t^s \int_{\mathbb{R}^d} \|\eta_r(0, z)\|^2 \nu(dz) dr \\ &\leq \|x\|^2 + 6Ld^p(s-t) + (1+6L) \int_t^s \mathbb{E} \left[\left\| X_{r \wedge \tau_n^{t,x}}^{t,x} \right\|^2 \right] dr. \end{aligned}$$

Therefore, by (2.9) and Grönwall's lemma we have for all $s \in [t, T]$ and $n \in \mathbb{N}$ that

$$\mathbb{E} \left[\left\| X_{s \wedge \tau_n^{t,x}}^{t,x} \right\|^2 \right] \leq (6Ld^p(s-t) + \|x\|^2) e^{(1+6L)(s-t)}.$$

Finally, the application of Fatou's lemma yields for all $s \in [t, T]$ that

$$\mathbb{E} \left[\|X_s^{t,x}\|^2 \right] \leq \liminf_{n \rightarrow \infty} \mathbb{E} \left[\left\| X_{s \wedge \tau_n^{t,x}}^{t,x} \right\|^2 \right] \leq (6Ld^p(s-t) + \|x\|^2) e^{(1+6L)(s-t)},$$

which completes the proof of this lemma. \square

Lemma 3.2. *There exist a constant ρ only depending on L and T , satisfying for all $t \in [0, T]$, $s \in [t, T]$, and $x, y \in \mathbb{R}^d$ that*

$$\mathbb{E} \left[\|X_s^{t,x} - X_s^{t,y}\|^2 \right] \leq 4\|x - y\|^2 e^{\rho(s-t)}.$$

Proof. We first fix $t \in [0, T]$ and $x, y \in \mathbb{R}^d$. Notice for every $s \in [t, T]$ that

$$\begin{aligned} X_s^{t,x} - X_s^{t,y} &= x - y + \int_t^s \left(\mu(r, X_{r-}^{t,x}) - \mu(r, X_{r-}^{t,y}) \right) dr + \int_t^s \left(\sigma(r, X_{r-}^{t,x}) - \sigma(r, X_{r-}^{t,y}) \right) dW_r \\ &\quad + \int_t^s \int_{\mathbb{R}^d} \left(\eta_r(X_{r-}^{t,x}, z) - \eta_r(X_{r-}^{t,y}, z) \right) \tilde{\pi}(dz, dr). \end{aligned}$$

Then by Jensen's inequality, Itô's isometry, and the fact that $(\sum_{i=1}^4 a_i)^2 \leq 4 \sum_{i=1}^4 a_i^2$ for $a_i \in \mathbb{R}$, $i = 1, 2, 3, 4$, it holds for all $s \in [t, T]$ that

$$\begin{aligned} \mathbb{E} \left[\|X_s^{t,x} - X_s^{t,y}\|^2 \right] &\leq 4\|x - y\|^2 + 4(s-t) \int_t^s \mathbb{E} \left[\|\mu(r, X_r^{t,x}) - \mu(r, X_r^{t,y})\|^2 \right] dr \\ &\quad + 4 \int_t^s \mathbb{E} \left[\|\sigma(r, X_r^{t,x}) - \sigma(r, X_r^{t,y})\|_F^2 \right] dr \\ &\quad + 4 \int_t^s \int_{\mathbb{R}^d} \mathbb{E} \left[\|\eta_r(X_r^{t,x}, z) - \eta_r(X_r^{t,y}, z)\|^2 \right] \nu(dz) dr. \end{aligned}$$

Thus, by (2.3) we obtain for all $s \in [t, T]$ that

$$\begin{aligned} \mathbb{E} \left[\|X_s^{t,x} - X_s^{t,y}\|^2 \right] &\leq 4\|x - y\|^2 + 4(s-t)L \int_t^s \mathbb{E} \left[\|X_r^{t,x} - X_r^{t,y}\|^2 \right] dr \\ &\quad + 8L \int_t^s \mathbb{E} \left[\|X_r^{t,x} - X_r^{t,y}\|^2 \right] dr. \end{aligned}$$

Therefore, by (2.9) and Grönwall's lemma, we have for all $s \in [t, T]$ that

$$\mathbb{E} \left[\|X_s^{t,x} - X_s^{t,y}\|^2 \right] \leq 4\|x - y\|^2 e^{4L(T+2)(s-t)},$$

which completes the proof of this lemma. \square

Lemma 3.3. *There exists a positive constant c only depending on T and L satisfying for all $(t, x) \in [0, T] \times \mathbb{R}^d$, $s \in [t, T]$, and $s' \in [s, T]$ that*

$$\mathbb{E} \left[\|X_{s'}^{t,x} - X_s^{t,x}\|^2 \right] \leq c [(s' - s) + (s' - s)^2] (d^p + \|x\|^2). \quad (3.5)$$

Proof. We notice that almost surely for all $(t, x) \in [0, T] \times \mathbb{R}^d$, $s \in [t, T]$ and $s' \in [s, T]$ that

$$X_{s'}^{t,x} = X_s^{t,x} + \int_s^{s'} \mu(r, X_{r-}^{t,x}) dr + \int_s^{s'} \sigma(r, X_{r-}^{t,x}) dW_r + \int_s^{s'} \int_{\mathbb{R}^d} \eta_r(X_{r-}^{t,x}, z) \tilde{\pi}(dz, dr).$$

Hence, by Itô's isometry, Hölder's inequality, (2.3), (2.4) and Lemma 3.1 it holds for all $(t, x) \in [0, T] \times \mathbb{R}^d$, $s \in [t, T]$ and $s' \in [s, T]$ that

$$\begin{aligned} \mathbb{E} \left[\|X_{s'}^{t,x} - X_s^{t,x}\|^2 \right] &\leq 3(s' - s) \int_s^{s'} \mathbb{E} \left[\|\mu(r, X_r^{t,x})\|^2 \right] dr + 3 \int_s^{s'} \mathbb{E} \left[\|\sigma(r, X_r^{t,x})\|_F^2 \right] dr \\ &\quad + 3 \int_s^{s'} \mathbb{E} \left[\int_{\mathbb{R}^d} \|\eta_r(X_r^{t,x}, z)\|^2 \nu(dz) \right] dr \end{aligned}$$

$$\begin{aligned}
&\leq 3(s' - s) \int_s^{s'} \mathbb{E} \left[2 \|\mu(r, X_r^{t,x}) - \mu(r, 0)\|^2 + 2 \|\mu(r, 0)\|^2 \right] dr \\
&\quad + 3 \int_s^{s'} \mathbb{E} \left[2 \|\sigma(r, X_r^{t,x}) - \sigma(r, 0)\|_F^2 + 2 \|\sigma(r, 0)\|^2 \right] dr \\
&\quad + 3 \int_s^{s'} \mathbb{E} \left[\int_{\mathbb{R}^d} \left(2 \|\eta_r(X_r^{t,x}, z) - \eta_r(0, z)\|^2 + 2 \|\eta_r(0, z)\|^2 \right) \nu(dz) \right] dr \\
&\leq 3(s' - s) \int_s^{s'} (2L\mathbb{E} [\|X_r^{t,x}\|^2] + 2Ld^p) dr \\
&\quad + 6 \int_s^{s'} (2L\mathbb{E} [\|X_r^{t,x}\|^2] + 2Ld^p) dr \\
&\leq c(s' - s)^2(d^p + \|x\|^2) + c(s' - s)(d^p + \|x\|^2)
\end{aligned} \tag{3.6}$$

where $c = 12L(1 + 6LT)e^{(1+6L)T}$. \square

Lemma 3.4. For all $t \in [0, T]$, $t' \in [t, T]$, $s \in [t', T]$, and $x, x' \in \mathbb{R}^d$ it holds that

$$\begin{aligned}
\mathbb{E} \left[\left\| X_s^{t,x} - X_s^{t',x'} \right\|^2 \right] &\leq 9(\|x - x'\| + |t - t'| + |t - t'|^{1/2})^2 \cdot \exp\{6LT(T + 4)\} \\
&\quad \cdot \left[1 + 3L^{1/2}d^{p/2}(1 + \|x'\|^2)^{1/2} \right]^2.
\end{aligned} \tag{3.7}$$

Proof. We follow the idea of the proof of Lemma 3.6 in [9], and fix $t \in [0, T]$, $t' \in [t, T]$, $s \in [t', T]$, and $x, x' \in \mathbb{R}^d$ throughout this proof. One first notice that almost surely

$$\begin{aligned}
X_s^{t,x} - X_s^{t',x'} &= X_{t'}^{t,x} - x' + \int_{t'}^s \left(\mu(r, X_{r-}^{t,x}) - \mu(r, X_{r-}^{t',x'}) \right) dr \\
&\quad + \int_{t'}^s \left(\sigma(r, X_{r-}^{t,x}) - \sigma(r, X_{r-}^{t',x'}) \right) dW_r + \int_{t'}^s \int_{\mathbb{R}^d} \left(\eta_r(X_{r-}^{t,x}, z) - \eta_r(X_{r-}^{t',x'}, z) \right) \tilde{\pi}(dz, dr).
\end{aligned}$$

Then by Minkowski's inequality and Hölder's inequality we have

$$\begin{aligned}
\left(\mathbb{E} \left[\left\| X_s^{t,x} - X_s^{t',x'} \right\|^2 \right] \right)^{1/2} &\leq \left(\mathbb{E} \left[\left\| X_{t'}^{t,x} - x' \right\|^2 \right] \right)^{1/2} \\
&\quad + \left(\int_{t'}^s (s - t') \mathbb{E} \left[\left\| \mu(r, X_{r-}^{t,x}) - \mu(r, X_{r-}^{t',x'}) \right\|^2 \right] dr \right)^{1/2} \\
&\quad + \left(\mathbb{E} \left[\left\| \int_{t'}^s \left(\sigma(r, X_{r-}^{t,x}) - \sigma(r, X_{r-}^{t',x'}) \right) dW_r \right\|^2 \right] \right)^{1/2} \\
&\quad + \left(\mathbb{E} \left[\left\| \int_{t'}^s \int_{\mathbb{R}^d} \left(\eta_r(X_{r-}^{t,x}, z) - \eta_r(X_{r-}^{t',x'}, z) \right) \tilde{\pi}(dz, dr) \right\|^2 \right] \right)^{1/2}
\end{aligned}$$

Therefore, by Itô's isometry and (2.3) we obtain

$$\begin{aligned}
\left(\mathbb{E} \left[\left\| X_s^{t,x} - X_s^{t',x'} \right\|^2 \right] \right)^{1/2} &\leq \left(\mathbb{E} \left[\left\| X_{t'}^{t,x} - x' \right\|^2 \right] \right)^{1/2} \\
&\quad + L^{1/2} \left((s - t') \int_{t'}^s \mathbb{E} \left[\left\| X_{r-}^{t,x} - X_{r-}^{t',x'} \right\|^2 \right] dr \right)^{1/2} \\
&\quad + \left(\int_{t'}^s \mathbb{E} \left[\left\| \sigma(r, X_{r-}^{t,x}) - \sigma(r, X_{r-}^{t',x'}) \right\|_F^2 \right] dr \right)^{1/2} \\
&\quad + \left(\int_{t'}^s \mathbb{E} \left[\int_{\mathbb{R}^d} \left\| \eta_r(X_{r-}^{t,x}, z) - \eta_r(X_{r-}^{t',x'}, z) \right\|^2 \nu(dz) \right] dr \right)^{1/2} \\
&\leq \left(\mathbb{E} \left[\left\| X_{t'}^{t,x} - x' \right\|^2 \right] \right)^{1/2} + L^{1/2} \left((s - t') \int_{t'}^s \mathbb{E} \left[\left\| X_{r-}^{t,x} - X_{r-}^{t',x'} \right\|^2 \right] dr \right)^{1/2}
\end{aligned}$$

$$+ 2L^{1/2} \left(\int_{t'}^s \mathbb{E} \left[\left\| X_r^{t,x} - X_r^{t',x'} \right\|^2 \right] dr \right)^{1/2}.$$

Then by the fact that for all $a, b, c \in \mathbb{R}$ it holds that $(a + b + c)^2 \leq 3(a^2 + b^2 + c^2)$ and Hölder's inequality we have

$$\begin{aligned} \mathbb{E} \left[\left\| X_s^{t,x} - X_s^{t',x'} \right\|^2 \right] &\leq 3\mathbb{E} \left[\left\| X_{t'}^{t,x} - x' \right\|^2 \right] + 3LT \int_{t'}^s \mathbb{E} \left[\left\| X_r^{t,x} - X_r^{t',x'} \right\|^2 \right] dr \\ &\quad + 12L \int_{t'}^s \mathbb{E} \left[\left\| X_r^{t,x} - X_r^{t',x'} \right\|^2 \right] dr \\ &= 3\mathbb{E} \left[\left\| X_{t'}^{t,x} - x' \right\|^2 \right] + 3L(T+4) \int_{t'}^s \mathbb{E} \left[\left\| X_r^{t,x} - X_r^{t',x'} \right\|^2 \right] dr \end{aligned} \quad (3.8)$$

Moreover, by (3.8), (2.9), and Grönwall's lemma we get

$$\mathbb{E} \left[\left\| X_s^{t,x} - X_s^{t',x'} \right\|^2 \right] \leq 3\mathbb{E} \left[\left\| X_{t'}^{t,x} - x' \right\|^2 \right] \cdot \exp\{3LT(T+4)\}. \quad (3.9)$$

Furthermore, we notice that almost surely

$$X_{t'}^{t,x} - x' = x - x' + \int_t^{t'} \mu(r, X_{r-}^{t,x}) dr + \int_t^{t'} \sigma(r, X_{r-}^{t,x}) dW_r + \int_t^{t'} \int_{\mathbb{R}^d} \eta_r(X_{r-}^{t,x}, z) \tilde{\pi}(dz, dr).$$

Hence, by Minkowski's inequality and Hölder's inequality we get

$$\begin{aligned} &\left(\mathbb{E} \left[\left\| X_{t'}^{t,x} - x' \right\|^2 \right] \right)^{1/2} \leq \|x - x'\| + \left((t' - t) \int_t^{t'} \mathbb{E} \left[\left\| \mu(r, X_r^{t,x}) \right\|^2 \right] dr \right)^{1/2} \\ &+ \left(\mathbb{E} \left[\left\| \int_t^{t'} \sigma(r, X_{r-}^{t,x}) dW_r \right\|^2 \right] \right)^{1/2} + \left(\mathbb{E} \left[\left\| \int_t^{t'} \int_{\mathbb{R}^d} \eta_r(X_{r-}^{t,x}, z) \tilde{\pi}(dz, dr) \right\|^2 \right] \right)^{1/2}. \end{aligned} \quad (3.10)$$

Moreover, by Itô's isometry, (3.10) implies that

$$\begin{aligned} &\left(\mathbb{E} \left[\left\| X_{t'}^{t,x} - x' \right\|^2 \right] \right)^{1/2} \leq \|x - x'\| + \left((t' - t) \int_t^{t'} \mathbb{E} \left[\left\| \mu(r, X_r^{t,x}) \right\|^2 \right] dr \right)^{1/2} \\ &+ \left(\int_t^{t'} \mathbb{E} \left[\left\| \sigma(r, X_r^{t,x}) \right\|_F^2 \right] dr \right)^{1/2} + \left(\int_t^{t'} \mathbb{E} \left[\int_{\mathbb{R}^d} \left\| \eta_r(X_r^{t,x}, z) \right\|^2 \nu(dz) \right] dr \right)^{1/2}. \end{aligned}$$

Thus, applying Minkowski's inequality, (2.3), and (2.4) yields

$$\begin{aligned} &\left(\mathbb{E} \left[\left\| X_{t'}^{t,x} - x' \right\|^2 \right] \right)^{1/2} \\ &\leq \|x - x'\| + \left((t' - t) \int_t^{t'} \left\| \mu(r, x') \right\|^2 dr \right)^{1/2} + \left((t' - t) \int_t^{t'} \mathbb{E} \left[\left\| \mu(r, X_r^{t,x}) - \mu(r, x') \right\|^2 \right] dr \right)^{1/2} \\ &+ \left(\int_t^{t'} \left\| \sigma(r, x') \right\|_F^2 dr \right)^{1/2} + \left(\int_t^{t'} \mathbb{E} \left[\left\| \sigma(r, X_r^{t,x}) - \sigma(r, x') \right\|_F^2 \right] dr \right)^{1/2} \\ &+ \left(\int_t^{t'} \mathbb{E} \left[\int_{\mathbb{R}^d} \left\| \eta_r(X_r^{t,x}, z) - \eta_r(x', z) \right\|^2 \nu(dz) \right] dr \right)^{1/2} + \left(\int_t^{t'} \int_{\mathbb{R}^d} \left\| \eta_r(x', z) \right\|^2 \nu(dz) dr \right)^{1/2} \\ &\leq \|x - x'\| + \left((t' - t) \int_t^{t'} \left\| \mu(r, x') \right\|^2 dr \right)^{1/2} + L^{1/2} \left((t' - t) \int_t^{t'} \mathbb{E} \left[\left\| X_r^{t,x} - x' \right\|^2 \right] dr \right)^{1/2} \\ &+ \left(\int_t^{t'} \left\| \sigma(r, x') \right\|_F^2 dr \right)^{1/2} + 2L^{1/2} \left(\int_t^{t'} \mathbb{E} \left[\left\| X_r^{t,x} - x' \right\|^2 \right] dr \right)^{1/2} \end{aligned}$$

$$+ \left(\int_t^{t'} \int_{\mathbb{R}^d} \|\eta_r(x', z)\|^2 \nu(dz) dr \right)^{1/2}.$$

Then using Hölder's inequality, and the fact that for all $a, b, c \in \mathbb{R}$ it holds that $(a + b + c)^2 \leq 3a^2 + 3b^2 + 3c^2$, we have

$$\begin{aligned} \mathbb{E} \left[\left\| X_{t'}^{t,x} - x' \right\|^2 \right] &\leq 3L(T+4) \int_t^{t'} \mathbb{E} \left[\left\| X_r^{t,x} - x' \right\|^2 \right] dr \\ &\quad + 3 \left[\|x - x'\| + \left(|t - t'| \cdot \int_t^{t'} \|\mu(r, x')\|^2 dr \right)^{1/2} + \left(\int_t^{t'} \|\sigma(r, x')\|_F^2 dr \right)^{1/2} \right. \\ &\quad \left. + \left(\int_t^{t'} \int_{\mathbb{R}^d} \|\eta_r(x', z)\|^2 \nu(dz) dr \right)^{1/2} \right]^2. \end{aligned} \quad (3.11)$$

This together with (2.4) imply that

$$\begin{aligned} \mathbb{E} \left[\left\| X_{t'}^{t,x} - x' \right\|^2 \right] &\leq 3L(T+4) \int_t^{t'} \mathbb{E} \left[\left\| X_r^{t,x} - x' \right\|^2 \right] dr \\ &\quad + 3 \left[\|x - x'\| + |t - t'| \cdot L^{1/2} d^{p/2} (1 + \|x'\|^2)^{1/2} \right. \\ &\quad \left. + |t - t'|^{1/2} \cdot L^{1/2} d^{p/2} (1 + \|x'\|^2)^{1/2} + |t - t'|^{1/2} L^{1/2} d^{p/2} (1 + \|x'\|^2)^{1/2} \right]^2. \end{aligned}$$

Hence, we have

$$\begin{aligned} \mathbb{E} \left[\left\| X_{t'}^{t,x} - x' \right\|^2 \right] &\leq 3L(T+4) \int_t^{t'} \mathbb{E} \left[\left\| X_r^{t,x} - x' \right\|^2 \right] dr + 3(\|x - x'\| + |t - t'| + |t - t'|^{1/2})^2 \\ &\quad \cdot \left[1 + 3L^{1/2} d^{p/2} (1 + \|x'\|^2)^{1/2} \right]^2 \end{aligned}$$

Therefore, by (2.9) and Grönwall's lemma we have

$$\begin{aligned} \mathbb{E} \left[\left\| X_{t'}^{t,x} - x' \right\|^2 \right] &\leq 3(\|x - x'\| + |t - t'| + |t - t'|^{1/2})^2 \cdot \exp\{3LT(T+4)\} \\ &\quad \cdot \left[1 + 3L^{1/2} d^{p/2} (1 + \|x'\|^2)^{1/2} \right]^2. \end{aligned} \quad (3.12)$$

Hence, combining (3.12) with (3.9) yields (3.7), which completes the proof of this lemma. \square

From now on, for ease of notation, we define $\Lambda = \{(t, s) \in [0, T]^2 : t \leq s\}$.

Corollary 3.5. *For each $(t, x) \in [0, T] \times \mathbb{R}^d$ and $s \in [t, T]$, let $(t_k, s_k, x_k)_{k=1}^\infty$ be a sequence such that $(t_k, s_k, x_k) \in \Lambda \times \mathbb{R}^d$ for all $k \in \mathbb{N}$ and $\lim_{k \rightarrow \infty} (t_k, s_k, x_k) = (t, s, x)$. Then for any $\varepsilon > 0$ it holds that*

$$\lim_{k \rightarrow \infty} \mathbb{P} \left(\left\| X_s^{t,x} - X_{s_k}^{t_k, x_k} \right\| \geq \varepsilon \right) = 0. \quad (3.13)$$

Proof. Fix $\varepsilon \in (0, \infty)$, $(t, x) \in [0, T] \times \mathbb{R}^d$, $s \in [t, T]$ and a sequence $(t_k, s_k, x_k)_{k=1}^\infty$ satisfying that $(t_k, s_k, x_k) \in \Lambda \times \mathbb{R}^d$ for all $k \in \mathbb{N}$, and $\lim_{k \rightarrow \infty} (t_k, s_k, x_k) = (t, s, x)$. By Chebyshev inequality and the fact that $(a + b)^2 \leq 2a^2 + 2b^2$ for all $a, b \in \mathbb{R}$, we have

$$\begin{aligned} \mathbb{P} \left(\left\| X_s^{t,x} - X_{s_k}^{t_k, x_k} \right\| \geq \varepsilon \right) &\leq \varepsilon^{-2} \mathbb{E} \left[\left\| X_s^{t,x} - X_{s_k}^{t_k, x_k} \right\|^2 \right] \\ &\leq 2\varepsilon^{-2} \mathbb{E} \left[\left\| X_s^{t,x} - X_{s_k}^{t,x} \right\|^2 \right] + 2\varepsilon^{-2} \mathbb{E} \left[\left\| X_{s_k}^{t,x} - X_{s_k}^{t_k, x_k} \right\|^2 \right]. \end{aligned}$$

Therefore, the application of Lemma 3.3 and Lemma 3.4 yields

$$\begin{aligned} \mathbb{P} \left(\left\| X_s^{t,x} - X_{s_k}^{t_k, x_k} \right\| \geq \varepsilon \right) &\leq 2\varepsilon^{-2} \cdot c[(s_k - s) + (s_k - s)^2](d^p + \|x\|^2) \\ &\quad + 2\varepsilon^{-2} \cdot 9 \left(\|x_k - x\| + |t_k - t| + |t_k - t|^{1/2} \right)^2 \exp\{6LT(T+4)\} \\ &\quad \cdot \left[1 + 3L^{1/2} d^{p/2} (1 + \|x_k\|^2 + \|x\|^2)^{1/2} \right]^2, \end{aligned} \quad (3.14)$$

where c is the positive constant introduced in (3.5) in Lemma 3.3. Finally, passing limit as $k \rightarrow \infty$ in (3.14) gives (3.13). \square

3.2. Bounds for Euler approximations.

Lemma 3.6 (Euler Approximation). *There exist positive constants $C_1 = C_1(T, L)$, $C_2 = C_2(T, L, L_1, L_2)$, $\rho_1 = \rho_1(L)$, and $\rho_2 = \rho_2(L)$ satisfying for all $N \in \mathbb{N}$ and $(t, x) \in [0, T] \times \mathbb{R}^d$ that*

$$\mathbb{E} \left[\sup_{s \in [t, T]} \|Y_s^{t, x, N}\|^2 \right] \leq C_1(d^p + \|x\|^2)e^{\rho_1(T-t)} \quad (3.15)$$

and

$$\mathbb{E} \left[\sup_{s \in [t, T]} \|X_s^{t, x} - Y_s^{t, x, N}\|^2 \right] \leq C_2(d^p + \|x\|^2)N^{-1}e^{\rho_2(T-t)}. \quad (3.16)$$

Proof. We fix $(t, x) \in [0, T] \times \mathbb{R}^d$ and $N \in \mathbb{N}$ throughout this proof. For $Y^{t, x, N} = (Y^{t, x, N, 1}, \dots, Y^{t, x, N, d})$, applying Itô's formula to $\|Y_s^{t, x, N}\|^2$ shows almost surely for all $s \in [t, T]$ that

$$\begin{aligned} \|Y_s^{t, x, N}\|^2 &= \|x\|^2 + 2 \int_t^s \left(\langle Y_r^{t, x, N}, \mu(\kappa_N(r), Y_{\kappa_N(r)}^{t, x, N}) \rangle + \frac{1}{2} \left\| \sigma(\kappa_N(r), Y_{\kappa_N(r)}^{t, x, N}) \right\|_F^2 \right) dr \\ &\quad + 2 \sum_{i=1}^d \sum_{j=1}^d \int_t^s Y_{r-}^{t, x, N, i} \sigma^{ij}(\kappa_N(r-), Y_{\kappa_N(r-)}^{t, x, N}) dW_r^j \\ &\quad + 2 \int_t^s \int_{\mathbb{R}^d} \langle Y_{r-}^{t, x, N}, \eta_{\kappa_N(r-)}(Y_{\kappa_N(r-)}^{t, x, N}, z) \rangle \tilde{\pi}(dz, dr) \\ &\quad + \int_t^s \int_{\mathbb{R}^d} \left\| \eta_{\kappa_N(r-)}(Y_{\kappa_N(r-)}^{t, x, N}, z) \right\|^2 \pi(dz, dr). \end{aligned}$$

Then, by Davis' inequality (see, e.g., Theorem VII.90 and VII (90.1) in [22]) we obtain for every $s \in [t, T]$ that

$$a_N(s) := \mathbb{E} \left[\sup_{r \in [t, s]} \|Y_r^{t, x, N}\|^2 \right] \leq \|x\|^2 + \sum_{i=1}^4 A_i(s), \quad (3.17)$$

where

$$\begin{aligned} A_1(s) &:= 2\mathbb{E} \left[\int_t^s \left(\left| \langle Y_r^{t, x, N}, \mu(\kappa_N(r), Y_{\kappa_N(r)}^{t, x, N}) \rangle \right| + \frac{1}{2} \left\| \sigma(\kappa_N(r), Y_{\kappa_N(r)}^{t, x, N}) \right\|_F^2 \right) dr \right], \\ A_2(s) &:= 8\mathbb{E} \left[\left(\int_t^s \left\| \left[\sigma(\kappa_N(r), Y_{\kappa_N(r)}^{t, x, N}) \right]^T Y_{\kappa_N(r)}^{t, x, N} \right\|^2 dr \right)^{1/2} \right], \\ A_3(s) &:= 8\mathbb{E} \left[\left(\int_t^s \int_{\mathbb{R}^d} \left| \langle Y_{r-}^{t, x, N}, \eta_{\kappa_N(r-)}(Y_{\kappa_N(r-)}^{t, x, N}, z) \rangle \right|^2 \pi(dz, dr) \right)^{1/2} \right], \\ A_4(s) &:= \mathbb{E} \left[\int_t^s \int_{\mathbb{R}^d} \left\| \eta_{\kappa_N(r)}(Y_{\kappa_N(r)}^{t, x, N}, z) \right\|^2 \nu(dz) dr \right]. \end{aligned}$$

Furthermore, by Cauchy-Schwarz inequality, Young's inequality, and (2.4) we have for all $s \in [t, T]$ that

$$\begin{aligned} A_1(s) &\leq \mathbb{E} \left[\int_t^s \left(\|Y_r^{t, x, N}\|^2 + \left\| \mu(\kappa_N(r), Y_{\kappa_N(r)}^{t, x, N}) \right\|^2 + \left\| \sigma(\kappa_N(r), Y_{\kappa_N(r)}^{t, x, N}) \right\|_F^2 \right) dr \right] \\ &\leq \mathbb{E} \left[\int_t^s \left(\|Y_r^{t, x, N}\|^2 + 2 \left\| \mu(\kappa_N(r), Y_{\kappa_N(r)}^{t, x, N}) - \mu(\kappa_N(r), 0) \right\|^2 + 2 \left\| \mu(\kappa_N(r), 0) \right\|^2 \right. \right. \\ &\quad \left. \left. + 2 \left\| \sigma(\kappa_N(r), Y_{\kappa_N(r)}^{t, x, N}) - \sigma(\kappa_N(r), 0) \right\|_F^2 + 2 \left\| \sigma(\kappa_N(r), 0) \right\|_F^2 \right) dr \right] \\ &\leq \mathbb{E} \left[\int_t^s \left(\|Y_r^{t, x, N}\|^2 + 4L \left\| Y_{\kappa_N(r)}^{t, x, N} \right\|^2 + 4Ld^p \right) dr \right] \\ &\leq 4TLd^p + (1 + 4L) \int_t^s a_N(r) dr. \end{aligned} \quad (3.18)$$

Similarly, by Cauchy-Schwarz inequality, Young's inequality, (2.3), and (2.4) we get for all $s \in [t, T]$ and $\varepsilon > 0$ that

$$\begin{aligned}
A_2(s) &\leq 8\mathbb{E} \left[\left(\int_t^s \left\| Y_{\kappa_N(r)}^{t,x,N} \right\|^2 \left\| \sigma(\kappa_N(r), Y_{\kappa_N(r)}^{t,x,N}) \right\|_F^2 dr \right)^{1/2} \right] \\
&\leq 8\mathbb{E} \left[\sup_{r \in [t,s]} (\|Y_r^{t,x,N}\|) \left(\int_t^s \left(2 \left\| \sigma(\kappa_N(r), Y_{\kappa_N(r)}^{t,x,N}) - \sigma(\kappa_N(r), 0) \right\|_F^2 + 2 \left\| \sigma(\kappa_N(r), 0) \right\|_F^2 \right) dr \right)^{1/2} \right] \\
&\leq 4\varepsilon a_N(s) + \frac{4}{\varepsilon} \mathbb{E} \left[\int_t^s \left(2L \left\| Y_{\kappa_N(r)}^{t,x,N} \right\|^2 + 2Ld^p \right) dr \right] \\
&\leq 4\varepsilon a_N(s) + \frac{8TLd^p}{\varepsilon} + \frac{8L}{\varepsilon} \int_t^s a_N(r) dr, \tag{3.19}
\end{aligned}$$

and

$$\begin{aligned}
A_3(s) &\leq 8\mathbb{E} \left[\sup_{r \in [t,s]} (\|Y_r^{t,x,N}\|) \left(\int_t^s \int_{\mathbb{R}^d} \left\| \eta_{\kappa_N(r-)}(Y_{\kappa_N(r-)}^{t,x,N}, z) \right\|^2 \pi(dz, dr) \right)^{1/2} \right] \\
&\leq 4\varepsilon a_N(s) + \frac{4}{\varepsilon} A_4(s). \tag{3.20}
\end{aligned}$$

Moreover, by (2.3) and (2.4) we have for all $s \in [t, T]$ that

$$\begin{aligned}
A_4(s) &\leq \mathbb{E} \left[\int_t^s \int_{\mathbb{R}^d} \left(2 \left\| \eta_{\kappa_N(r)}(Y_{\kappa_N(r)}^{t,x,N}, z) - \eta_{\kappa_N(r)}(0, z) \right\|^2 + 2 \left\| \eta_{\kappa_N(r)}(0, z) \right\|^2 \right) \nu(dz) dr \right] \\
&\leq \mathbb{E} \left[\int_t^s \left(2L \left\| Y_{\kappa_N(r)}^{t,x,N} \right\|^2 + 2Ld^p \right) dr \right] \\
&\leq 2TLd^p + 2L \int_t^s a_N(r) dr. \tag{3.21}
\end{aligned}$$

Then, combining (3.17) and (3.18)-(3.21) yields for all $s \in [t, T]$ that

$$a_N(s) \leq \|x\|^2 + 8\varepsilon a_N(s) + 2TLd^p(3 + 8/\varepsilon) + (1 + 6L + 16L/\varepsilon) \int_t^s a_N(r) dr. \tag{3.22}$$

Define a sequence of stopping times $\{\tau_k^N\}_{k=1}^\infty$ by

$$\tau_k^N := \inf\{s \geq t : \|Y_s^{t,x,N}\| \geq k\} \wedge T, \quad k \in \mathbb{N}$$

Note that for all $k \in \mathbb{N}$, $Y^{t,x,N}$ has no fixed time of discontinuity. Thus, we have that

$$\mathbb{E} \left[\sup_{s \in [t, T]} \left\| Y_{s \wedge \tau_k^N}^{t,x,N} \right\|^2 \right] \leq k^2 + \mathbb{E} \left[\left\| Y_{\tau_k^N}^{t,x,N} - Y_{\tau_k^N-}^{t,x,N} \right\|^2 \right] = k^2$$

Hence, by taking $\varepsilon = 1/9$ and replacing s by $s \wedge \tau_k^N$ in (3.22), as well as using Grönwall's lemma we obtain

$$a_N(s \wedge \tau_k^N) \leq C_1(d^p + \|x\|^2)e^{\rho_1(s-t)} \quad \text{for all } s \in [t, T] \text{ and } k \in \mathbb{N},$$

where $C_1 := 9 \cdot \max\{150LT, 1\}$ and $\rho_1 := 9(1 + 150L)$. Then the application of Fatou's lemma gives

$$a_N(T) \leq \liminf_{k \rightarrow \infty} a_N(T \wedge \tau_k^N) \leq C_1(d^p + \|x\|^2)e^{\rho_1(T-t)},$$

which proves (3.15).

Next, we show the proof of (3.16), which is analogous to the proof of (3.15). For every $s \in [t, T]$, and $i, j \in \{1, \dots, d\}$, we define

$$\tilde{\mu}_s^N := \mu(s, X_{s-}^{t,x}) - \mu(\kappa_N(s-), Y_{\kappa_N(s-)}^{t,x,N}), \quad \tilde{\eta}_s^N(z) := \eta_s(X_{s-}^{t,x}, z) - \eta_{\kappa_N(s-)}(Y_{\kappa_N(s-)}^{t,x,N}, z),$$

and

$$\tilde{\sigma}_s^N := \sigma(s, X_{s-}^{t,x}) - \sigma(\kappa_N(s-), Y_{\kappa_N(s-)}^{t,x,N}), \quad \tilde{\sigma}_s^{N,ij} := \sigma^{ij}(s, X_{s-}^{t,x}) - \sigma^{ij}(\kappa_N(s-), Y_{\kappa_N(s-)}^{t,x,N}).$$

For $X^{t,x} - Y^{t,x,N} = (X^{t,x,1} - Y^{t,x,N,1}, \dots, X^{t,x,d} - Y^{t,x,N,d})$, the application of Itô's formula shows almost surely for all $s \in [t, T]$ that

$$\begin{aligned} \|X_s^{t,x} - Y_s^{t,x,N}\|^2 &= 2 \int_t^s \left(\langle (X_{r-}^{t,x} - Y_{r-}^{t,x,N}), \tilde{\mu}_r^N \rangle + \frac{1}{2} \|\tilde{\sigma}_r^N\|_F^2 \right) dr \\ &\quad + 2 \sum_{i=1}^d \sum_{j=1}^d \int_t^s (X_{r-}^{t,x,i} - Y_{r-}^{t,x,N,i}) \tilde{\sigma}_r^{N,ij} dW_r^j \\ &\quad + 2 \int_t^s \int_{\mathbb{R}^d} \langle (X_{r-}^{t,x} - Y_{r-}^{t,x,N}), \tilde{\eta}_r^N(z) \rangle \tilde{\pi}(dz, dr) \\ &\quad + \int_t^s \int_{\mathbb{R}^d} \|\tilde{\eta}_r^N(z)\|^2 \pi(dz, dr). \end{aligned}$$

Then by Davis' inequality we obtain for all $s \in [t, T]$ that

$$e_N(s) := \mathbb{E} \left[\sup_{r \in [t, s]} \|X_r^{t,x} - Y_r^{t,x,N}\|^2 \right] \leq \sum_{i=1}^4 E_i(s), \quad (3.23)$$

where

$$\begin{aligned} E_1(s) &:= 2\mathbb{E} \left[\int_t^s \left(\left| \langle (X_{r-}^{t,x} - Y_{r-}^{t,x,N}), \tilde{\mu}_r^N \rangle \right| + \frac{1}{2} \|\tilde{\sigma}_r^N\|_F^2 \right) dr \right], \\ E_2(s) &:= 8\mathbb{E} \left[\left(\int_t^s \left\| [\tilde{\sigma}_r^N]^T (X_{r-}^{t,x} - Y_{r-}^{t,x,N}) \right\|^2 dr \right)^{1/2} \right], \\ E_3(s) &:= 8\mathbb{E} \left[\left(\int_t^s \int_{\mathbb{R}^d} \left| \langle (X_{r-}^{t,x} - Y_{r-}^{t,x,N}), \tilde{\eta}_r^N(z) \rangle \right|^2 \pi(dz, dr) \right)^{1/2} \right], \\ E_4(s) &:= \mathbb{E} \left[\int_t^s \int_{\mathbb{R}^d} \|\tilde{\eta}_r^N(z)\|^2 \nu(dz) dr \right]. \end{aligned}$$

Moreover, by Lemma 3.3 we notice for all $s \in [t, T]$ that

$$\mathbb{E} \left[\left\| X_s^{t,x} - X_{\kappa_N(s)}^{t,x} \right\|^2 \right] \leq 2c(d^p + \|x\|^2)N^{-1}T(T+1), \quad (3.24)$$

where $c = c(T, L)$ is the positive constant introduced in (3.5) in Lemma 3.3. Then, by (2.3), (2.4), (2.10), and (2.11) we obtain for all $s \in [t, T]$ that

$$\begin{aligned} \|\tilde{\mu}_s^N\| &\leq \left\| \mu(s, X_{s-}^{t,x}) - \mu(s, X_{\kappa_N(s-)}^{t,x}) \right\| + \left\| \mu(s, X_{\kappa_N(s-)}^{t,x}) - \mu(\kappa_N(s-), X_{\kappa_N(s-)}^{t,x}) \right\| \\ &\quad + \left\| \mu(\kappa_N(s-), X_{\kappa_N(s-)}^{t,x}) - \mu(\kappa_N(s-), Y_{\kappa_N(s-)}^{t,x,N}) \right\| \\ &\leq L^{1/2} \left\| X_{s-}^{t,x} - X_{\kappa_N(s-)}^{t,x} \right\| + (L_1 N^{-1} T)^{1/2} + L^{1/2} \left\| X_{\kappa_N(s-)}^{t,x} - Y_{\kappa_N(s-)}^{t,x,N} \right\|, \end{aligned} \quad (3.25)$$

$$\|\tilde{\sigma}_s^N\|_F \leq L^{1/2} \left\| X_{s-}^{t,x} - X_{\kappa_N(s-)}^{t,x} \right\| + (L_1 N^{-1} T)^{1/2} + L^{1/2} \left\| X_{\kappa_N(s-)}^{t,x} - Y_{\kappa_N(s-)}^{t,x,N} \right\|, \quad (3.26)$$

and

$$\int_{\mathbb{R}^d} \|\tilde{\eta}_s^N(z)\|^2 \nu(dz) \leq 3L \left\| X_{s-}^{t,x} - X_{\kappa_N(s-)}^{t,x} \right\|^2 + 3L_2 N^{-1} T + 3L \left\| X_{\kappa_N(s-)}^{t,x} - Y_{\kappa_N(s-)}^{t,x,N} \right\|^2. \quad (3.27)$$

Thus, by (3.24), (3.25), (3.26), Cauchy-Schwarz inequality, and Young's inequality we obtain for every $s \in [t, T]$ that

$$\begin{aligned} E_1(s) &\leq 2\mathbb{E} \left[\int_t^s \left(\left\| X_r^{t,x} - Y_r^{t,x,N} \right\| \left((L_1 N^{-1} T)^{1/2} + L^{1/2} \left\| X_r^{t,x} - X_{\kappa_N(r)}^{t,x} \right\| \right) + L^{1/2} \left\| X_{\kappa_N(r)}^{t,x} - Y_{\kappa_N(r)}^{t,x,N} \right\| \right) \right. \\ &\quad \left. + \frac{1}{2} \left((L_1 N^{-1} T)^{1/2} + L^{1/2} \left\| X_r^{t,x} - X_{\kappa_N(r)}^{t,x} \right\| + L^{1/2} \left\| X_{\kappa_N(r)}^{t,x} - Y_{\kappa_N(r)}^{t,x,N} \right\| \right)^2 dr \right] \\ &\leq \mathbb{E} \left[\int_t^s \left(\left\| X_r^{t,x} - Y_r^{t,x,N} \right\|^2 + 6L_1 N^{-1} T + 6L \left\| X_r^{t,x} - X_{\kappa_N(r)}^{t,x} \right\|^2 + 6L \left\| X_{\kappa_N(r)}^{t,x} - Y_{\kappa_N(r)}^{t,x,N} \right\|^2 \right) dr \right] \end{aligned}$$

$$\leq 6T^2(L_1 + 2c(T+1)L)(d^p + \|x\|^2)N^{-1} + (1+6L) \int_t^s e_N(r) dr, \quad (3.28)$$

where we recall that $e_N(s) := \mathbb{E} \left[\sup_{r \in [t, s]} \|X_r^{t,x} - Y_r^{t,x,N}\|^2 \right]$ for each $s \in [t, T]$. Analogously, by (3.24), (3.26), Cauchy-Schwarz inequality, and Young's inequality we get for all $s \in [t, T]$ and $\varepsilon > 0$ that

$$\begin{aligned} E_2(s) &\leq 8\mathbb{E} \left[\left(\int_t^s \|X_r^{t,x} - Y_r^{t,x,N}\|^2 \|\tilde{\sigma}_r^N\|_F^2 dr \right)^{1/2} \right] \\ &\leq 8\mathbb{E} \left[\left(\sup_{r \in [t, s]} \|X_r^{t,x} - Y_r^{t,x,N}\| \right) \left(\int_t^s \|\tilde{\sigma}_r^N\|_F^2 dr \right)^{1/2} \right] \\ &\leq \frac{12}{\varepsilon} \mathbb{E} \left[\int_t^s \left(L_1 N^{-1} T + L \|X_r^{t,x} - X_{\kappa_N(r)}^{t,x}\|^2 + L \|X_{\kappa_N(r)}^{t,x} - Y_{\kappa_N(r)}^{t,x,N}\|^2 \right) dr \right] \\ &\quad + 4\varepsilon \mathbb{E} \left[\sup_{r \in [t, s]} \|X_r^{t,x} - Y_r^{t,x,N}\|^2 \right] \\ &\leq 4\varepsilon e_N(s) + \frac{12T^2(L_1 + 2c(T+1)L)(d^p + \|x\|^2)}{\varepsilon N} + \frac{12L}{\varepsilon} \int_t^s e_N(r) dr, \end{aligned} \quad (3.29)$$

and

$$\begin{aligned} E_3(s) &\leq 8\mathbb{E} \left[\left(\sup_{r \in [t, s]} \|X_r^{t,x} - Y_r^{t,x,N}\| \right) \left(\int_t^s \int_{\mathbb{R}^d} \|\tilde{\eta}_r^N(z)\|^2 \pi(dz, dr) \right)^{1/2} \right] \\ &\leq 4\varepsilon e_N(s) + \frac{4E_4(s)}{\varepsilon}. \end{aligned} \quad (3.30)$$

Moreover, by (3.24) and (3.27) we see for all $s \in [t, T]$ that

$$\begin{aligned} E_4(s) &\leq \mathbb{E} \left[\int_t^s \left(3L_2 N^{-1} T + 3L \|X_r^{t,x} - X_{\kappa_N(r)}^{t,x}\|^2 + 3L \|X_{\kappa_N(r)}^{t,x} - Y_{\kappa_N(r)}^{t,x,N}\|^2 \right) dr \right] \\ &\leq \frac{3T^2(L_2 + 2c(T+1)L)(d^p + \|x\|^2)}{N} + 3L \int_t^s e_N(r) dr. \end{aligned} \quad (3.31)$$

Then combining (3.23) and (3.28)-(3.31) yields for each $s \in [t, T]$ that

$$\begin{aligned} e_N(s) &\leq 8\varepsilon e_N(s) + \left(1 + 9L + \frac{24L}{\varepsilon} \right) \int_t^s e_N(r) dr \\ &\quad + \left[6T^2(L_1 + 2c(T+1)L) \left(1 + \frac{2}{\varepsilon} \right) + 3T^2(L_2 + 2c(T+1)L) \left(1 + \frac{4}{\varepsilon} \right) \right] \frac{(d^p + \|x\|^2)}{N}. \end{aligned}$$

Hence, by (2.9) and (3.15), taking $\varepsilon = 1/9$ and applying Grönwall's lemma gives

$$e_N(T) = \mathbb{E} \left[\sup_{s \in [t, T]} \|X_s^{t,x} - Y_s^{t,x,N}\|^2 \right] \leq C_2(d^p + \|x\|^2)N^{-1}e^{\rho_2(T-t)}, \quad (3.32)$$

where $C_2 := 27T^2(38L_1 + 37L_2 + 150c(T+1)L)$ and $\rho_2 := 9(1 + 225L)$. Therefore, (3.16) is proved, and we complete the proof of this lemma. \square

3.3. Bounds for Euler approximations with truncated jumps.

Lemma 3.7. *There exist positive constants $C_1 = C_1(T, L)$, $C_2 = C_2(T, L, K)$, $\rho_1 = \rho_1(T, L)$, and $\rho_2 = \rho_2(T, L)$ satisfying for each $(t, x) \in [0, T] \times \mathbb{R}^d$, $N \in \mathbb{N}$, and $\delta \in (0, 1)$ that*

$$\mathbb{E} \left[\sup_{s \in [t, T]} \|Y_s^{t,x,N,\delta}\|^2 \right] \leq C_1(d^p + \|x\|^2)e^{\rho_1(T-t)}, \quad (3.33)$$

and

$$\mathbb{E} \left[\sup_{s \in [t, T]} \|Y_s^{t,x,N,\delta} - Y_s^{t,x,N}\|^2 \right] \leq C_2\delta^q d^p (d^p + \|x\|^2)e^{\rho_2(T-t)}. \quad (3.34)$$

Proof. As before, we first fix $(t, x) \in [0, T] \times \mathbb{R}^d$, $N \in \mathbb{N}$, and $\delta \in (0, 1)$. One can follow the proof of Lemma 3.6 for (3.15) to obtain that

$$\mathbb{E} \left[\sup_{s \in [t, T]} \left\| Y_s^{t, x, N, \delta} \right\|^2 \right] \leq C_1 (d^p + \|x\|^2) e^{\rho_1(T-t)}$$

with positive constants $C_1 = C_1(T, L)$ and $\rho_1 = \rho_1(T, L)$ defined in Lemma 3.6, which proves (3.33). Next, by Davis' inequality, Hölder's inequality, and (2.3) we observe that

$$\begin{aligned} & \mathbb{E} \left[\sup_{u \in [t, s]} \left\| Y_u^{t, x, N, \delta} - Y_u^{t, x, N} \right\|^2 \right] \leq 3(s-t) \int_t^s \mathbb{E} \left[\left\| \mu(\kappa_N(r), Y_{\kappa_N(r)}^{t, x, N, \delta}) - \mu(\kappa_N(r), Y_{\kappa_N(r)}^{t, x, N}) \right\|^2 \right] dr \\ & \quad + 12 \int_t^s \mathbb{E} \left[\left\| \sigma(\kappa_N(r), Y_{\kappa_N(r)}^{t, x, N, \delta}) - \sigma(\kappa_N(r), Y_{\kappa_N(r)}^{t, x, N}) \right\|_F^2 \right] dr \\ & \quad + 12 \int_t^s \mathbb{E} \left[\int_{\mathbb{R}^d} \left\| \eta_{\kappa_N(r)}(Y_{\kappa_N(r)}^{t, x, N, \delta}, z) \mathbf{1}_{A_\delta}(z) - \eta_{\kappa_N(r)}(Y_{\kappa_N(r)}^{t, x, N}, z) \right\|^2 \nu(dz) \right] dr \\ & \leq 3(s-t) \int_t^s \mathbb{E} \left[\left\| \mu(\kappa_N(r), Y_{\kappa_N(r)}^{t, x, N, \delta}) - \mu(\kappa_N(r), Y_{\kappa_N(r)}^{t, x, N}) \right\|^2 \right] dr \\ & \quad + 12 \int_t^s \mathbb{E} \left[\left\| \sigma(\kappa_N(r), Y_{\kappa_N(r)}^{t, x, N, \delta}) - \sigma(\kappa_N(r), Y_{\kappa_N(r)}^{t, x, N}) \right\|_F^2 \right] dr \\ & \quad + 24 \int_t^s \mathbb{E} \left[\int_{\mathbb{R}^d} \left\| \eta_{\kappa_N(r)}(Y_{\kappa_N(r)}^{t, x, N, \delta}, z) \mathbf{1}_{A_\delta}(z) - \eta_{\kappa_N(r)}(Y_{\kappa_N(r)}^{t, x, N}, z) \mathbf{1}_{A_\delta}(z) \right\|^2 \nu(dz) \right] dr \\ & \quad + 24 \int_t^s \mathbb{E} \left[\int_{\mathbb{R}^d} \left\| \eta_{\kappa_N(r)}(Y_{\kappa_N(r)}^{t, x, N}, z) \mathbf{1}_{\mathbb{R}^d/A_\delta}(z) \right\|^2 \nu(dz) \right] dr \\ & \leq 3(T+12)L \int_t^s \mathbb{E} \left[\sup_{u \in [t, r]} \left\| Y_u^{t, x, N, \delta} - Y_u^{t, x, N} \right\|^2 \right] dr \\ & \quad + 24 \int_t^T \mathbb{E} \left[\int_{\mathbb{R}^d/A_\delta} \left\| \eta_{\kappa_N(r)}(Y_{\kappa_N(r)}^{t, x, N}, z) \right\|^2 \nu(dz) \right] dr. \end{aligned} \quad (3.35)$$

Furthermore, by (2.8) and (3.15) it holds that

$$\begin{aligned} \int_t^T \mathbb{E} \left[\int_{\mathbb{R}^d/A_\delta} \left\| \eta_{\kappa_N(r)}(Y_{\kappa_N(r)}^{t, x, N}, z) \right\|^2 \nu(dz) \right] dr & \leq K \delta^q d^p \int_t^T \mathbb{E} \left[1 + \left\| Y_{\kappa_N(r)}^{t, x, N} \right\|^2 \right] dr \\ & \leq (C_1 + 1) K T \delta^q d^p (d^p + \|x\|^2) e^{\rho_1 T}, \end{aligned} \quad (3.36)$$

where $C_1 = C_1(T, L)$ and $\rho_1 = \rho_1(L)$ are positive constants defined in (3.15) in Lemma 3.6. Combining (3.35) and (3.36), as well as taking into account (3.15) and (3.33), by Grönwall's lemma we obtain

$$\mathbb{E} \left[\sup_{s \in [t, T]} \left\| Y_s^{t, x, N, \delta} - Y_s^{t, x, N} \right\|^2 \right] \leq C_2 \delta^q d^p (d^p + \|x\|^2) e^{\rho_2(T-t)},$$

where $C_2 := 24(C_1 + 1) K T e^{\rho_1 T}$ and $\rho_2 := 3(T+12)L$. Thus, the proof of this lemma is completed. \square

3.4. Bounds for Euler approximations with truncated jumps and Monte Carlo approximations of compensator integrals.

Lemma 3.8. *There exist positive constants C_1, C_2, ρ_1 , and ρ_2 only depending on T and L satisfying for all $(t, x) \in [0, T] \times \mathbb{R}^d$, $\delta \in (0, 1)$, and $N, \mathcal{M} \in \mathbb{N}$ with $\mathcal{M} \geq \delta^{-2} K d^p$ that*

$$\mathbb{E} \left[\sup_{s \in [t, T]} \left\| Y_s^{t, x, N, \delta, \mathcal{M}} \right\|^2 \right] \leq C_1 (d^p + \|x\|^2) e^{\rho_1(T-t)} \quad (3.37)$$

and

$$\mathbb{E} \left[\sup_{s \in [t, T]} \left\| Y_s^{t, x, N, \delta, \mathcal{M}} - Y_s^{t, x, N, \delta} \right\|^2 \right] \leq C_2 \delta^{-2} K d^p \mathcal{M}^{-1} (d^p + \|x\|^2) e^{\rho_2(T-t)}. \quad (3.38)$$

Proof. We fix $(t, x) \in [0, T]$, $\delta \in (0, 1)$, and $N, \mathcal{M} \in \mathbb{N}$ with $\mathcal{M} \geq \delta^{-2} K d^p$ throughout this proof. To prove (3.37), we first notice that by (2.8) it holds that

$$\nu(A_\delta) = \int_{A_\delta} \frac{1 \wedge \|z\|^2}{1 \wedge \|z\|^2} \nu(dz) \leq \delta^{-2} \int_{\mathbb{R}^d} (1 \wedge \|z\|^2) \nu(dz) \leq \delta^{-2} K d^p, \quad (3.39)$$

which shows that ν is a finite measure on $(A_\delta, \mathcal{B}(A_\delta))$. Thus, by (2.15) we have almost surely for all $s \in [t, T]$ that

$$\begin{aligned} Y_s^{t,x,N,\delta,\mathcal{M}} &= x + \int_t^s \mu(\kappa_N(r-), Y_{\kappa_N(r-)}^{t,x,N,\delta,\mathcal{M}}) dr + \int_t^s \sigma(\kappa_N(r-), Y_{\kappa_N(r-)}^{t,x,N,\delta,\mathcal{M}}) dW_r \\ &+ \int_t^s \int_{A_\delta} \eta_{\kappa_N(r-)}(Y_{\kappa_N(r-)}^{t,x,N,\delta,\mathcal{M}}, z) \tilde{\pi}(dz, dr) + \int_t^s \int_{A_\delta} \eta_{\kappa_N(r-)}(Y_{\kappa_N(r-)}^{t,x,N,\delta,\mathcal{M}}, z) \nu(dz) dr \\ &- \int_t^s \left(\frac{\nu(A_\delta)}{\mathcal{M}} \sum_{j=1}^{\mathcal{M}} \eta_{\kappa_N(r-)}(Y_{\kappa_N(r-)}^{t,x,N,\delta,\mathcal{M}}, V_{(\kappa_N(r-)-t)N,j}^{N,\mathcal{M}}) \right) dr. \end{aligned} \quad (3.40)$$

By Hölder's inequality, Burkholder-Davis-Gundy inequality (see, e.g., Theorem VII.92 in [22]), Minkowski integral inequality, and the fact that $(\sum_{i=1}^5 a_i)^2 \leq 5 \sum_{i=1}^5 a_i^2$ for $a_i \in \mathbb{R}$, $i = 1, 2, \dots, 5$, we obtain for each $s \in [t, T]$ that

$$a_{N,\mathcal{M}}^\delta(s) := \mathbb{E} \left[\sup_{r \in [t,s]} \|Y_r^{t,x,N,\delta,\mathcal{M}}\|^2 \right] \leq 5 \left(\|x\|^2 + \sum_{i=1}^4 A_i(s) \right), \quad (3.41)$$

where

$$A_1(s) := (s-t) \mathbb{E} \left[\int_t^s \|\mu(\kappa_N(r), Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}})\|^2 dr \right], \quad (3.42)$$

$$A_2(s) := 8 \mathbb{E} \left[\int_t^s \|\sigma(\kappa_N(r), Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}})\|_F^2 dr \right], \quad (3.43)$$

$$A_3(s) := 8 \mathbb{E} \left[\int_t^s \int_{A_\delta} \|\eta_{\kappa_N(r)}(Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}}, z)\|^2 \nu(dz) dr \right], \quad (3.44)$$

$$\begin{aligned} A_4(s) &:= (s-t) \mathbb{E} \left[\int_t^s \left\| \int_{A_\delta} \eta_{\kappa_N(r)}(Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}}, z) \nu(dz) \right. \right. \\ &\quad \left. \left. - \left(\frac{\nu(A_\delta)}{\mathcal{M}} \sum_{j=1}^{\mathcal{M}} \eta_{\kappa_N(r)}(Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}}, V_{(\kappa_N(r)-t)N,j}^{N,\mathcal{M}}) \right) \right\|^2 dr \right]. \end{aligned} \quad (3.45)$$

Next, by the definition of $V_{i,j}^{N,\mathcal{M}}$, we observe for all $y \in \mathbb{R}^d$, $s \in [0, T]$, $i = 0, 1, \dots, N$, and $j = 1, 2, \dots, \mathcal{M}$ that

$$\nu(A_\delta) \mathbb{E} \left[\eta_s(y, V_{i,j}^{N,\mathcal{M}}) \right] = \int_{A_\delta} \eta_s(y, z) \nu(dz). \quad (3.46)$$

Then by the independence properties of $V_{i,j}^{N,\mathcal{M}}$, (2.3), (2.4), and (3.46) it holds for all $r \in [t, T]$ that

$$\begin{aligned} &\mathbb{E} \left[\left\| \int_{A_\delta} \eta_{\kappa_N(r)}(Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}}, z) \nu(dz) - \left(\frac{\nu(A_\delta)}{\mathcal{M}} \sum_{j=1}^{\mathcal{M}} \eta_{\kappa_N(r)}(Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}}, V_{(\kappa_N(r)-t)N,j}^{N,\mathcal{M}}) \right) \right\|^2 \right] \\ &= \mathbb{E} \left[\mathbb{E} \left[\left\| \int_{A_\delta} \eta_{\kappa_N(r)}(y, z) \nu(dz) - \left(\frac{\nu(A_\delta)}{\mathcal{M}} \sum_{j=1}^{\mathcal{M}} \eta_{\kappa_N(r)}(y, V_{(\kappa_N(r)-t)N,j}^{N,\mathcal{M}}) \right) \right\|^2 \right] \Big|_{y=Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}}} \right] \\ &= [\nu(A_\delta)]^2 \mathcal{M}^{-1} \mathbb{E} \left[\sum_{k=1}^d \mathbb{E} \left[\left| \mathbb{E} [\eta_{\kappa_N(r)}^k(y, V_{1,1}^{N,\mathcal{M}})] - \eta_{\kappa_N(r)}^k(y, V_{1,1}^{N,\mathcal{M}}) \right|^2 \right] \Big|_{y=Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}}} \right] \end{aligned}$$

$$\begin{aligned}
&\leq [\nu(A_\delta)]^2 \mathcal{M}^{-1} \mathbb{E} \left[\sum_{k=1}^d \mathbb{E} \left[\left| \eta_{\kappa_N(r)}^k(y, V_{1,1}^{N,\mathcal{M}}) \right|^2 \right] \Big|_{y=Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}}} \right] \\
&= \nu(A_\delta) \mathcal{M}^{-1} \mathbb{E} \left[\int_{A_\delta} \left\| \eta_{\kappa_N(r)}(Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}}, z) \right\|^2 \nu(dz) \right] \tag{3.47}
\end{aligned}$$

$$\begin{aligned}
&\leq 2\nu(A_\delta) \mathcal{M}^{-1} \mathbb{E} \left[\int_{A_\delta} \left\| \eta_{\kappa_N(r)}(Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}}, z) - \eta_{\kappa_N(r)}(0, z) \right\|^2 \nu(dz) \right] \\
&\quad + 2\nu(A_\delta) \mathcal{M}^{-1} \mathbb{E} \left[\int_{A_\delta} \left\| \eta_{\kappa_N(r)}(0, z) \right\|^2 \nu(dz) \right] \\
&\leq 2\nu(A_\delta) \mathcal{M}^{-1} L \left(\mathbb{E} \left[\left\| Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}} \right\|^2 \right] + d^p \right). \tag{3.48}
\end{aligned}$$

Thus, by (3.39) and the assumption that $\mathcal{M} \geq \delta^{-2} K d^p$, we have for all $s \in [t, T]$ that

$$\begin{aligned}
A_4(s) &\leq 2\nu(A_\delta) \mathcal{M}^{-1} L(s-t) \left(\int_t^s \mathbb{E} \left[\left\| Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}} \right\|^2 \right] dr + (s-t)d^p \right) \\
&\leq 2\delta^{-2} K d^p \mathcal{M}^{-1} L(s-t) \left(\int_t^s \mathbb{E} \left[\left\| Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}} \right\|^2 \right] dr + (s-t)d^p \right) \tag{3.49}
\end{aligned}$$

$$\leq 2L(s-t) \left(\int_t^s \mathbb{E} \left[\left\| Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}} \right\|^2 \right] dr + (s-t)d^p \right). \tag{3.50}$$

Furthermore, using (2.3) and (2.4) we obtain for all $s \in [t, T]$ that

$$\begin{aligned}
A_1(s) &\leq 2(s-t) \mathbb{E} \left[\int_t^s \left\| \mu(\kappa_N(r), Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}}) - \mu(\kappa_N(r), 0) \right\|^2 dr \right] \\
&\quad + 2(s-t) \mathbb{E} \left[\int_t^s \left\| \mu(\kappa_N(r), 0) \right\|^2 dr \right] \\
&\leq 2(s-t)L \int_t^s \mathbb{E} \left[\left\| Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}} \right\|^2 \right] dr + 2(s-t)^2 L d^p, \tag{3.51}
\end{aligned}$$

$$A_2(s) \leq 16L \int_t^s \mathbb{E} \left[\left\| Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}} \right\|^2 \right] dr + 2(s-t)8Ld^p, \tag{3.52}$$

$$A_3(s) \leq 16L \int_t^s \mathbb{E} \left[\left\| Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}} \right\|^2 \right] dr + 2(s-t)8Ld^p. \tag{3.53}$$

Then combining (3.41) and (3.50)-(3.53) yields for all $s \in [t, T]$ that

$$a_{N,\mathcal{M}}^\delta(s) \leq 5\|x\|^2 + 20LT(T+8)d^p + 20L(T+8) \int_t^s a_{N,\mathcal{M}}^\delta(r) dr. \tag{3.54}$$

Therefore, by the same stopping time argument used to prove (3.15) in Lemma 3.6 and Grönwall's lemma we get that

$$a_{N,\mathcal{M}}^\delta(T) \leq C_1(d^p + \|x\|^2)e^{\rho_1(T-t)}$$

with $C_1 := 5 \cdot \max\{1, 4LT(T+8)\}$ and $\rho_1 := 20L(T+8)$, which proves (3.37).

Next, we start to prove (3.38). By (2.15) and (3.40) we notice for all $s \in [t, T]$ that

$$\begin{aligned}
Y_s^{t,x,N,\delta,\mathcal{M}} - Y_s^{t,x,N,\delta} &= \int_t^s \left(\mu(\kappa_N(r-), Y_{\kappa_N(r-)}^{t,x,N,\delta,\mathcal{M}}) - \mu(\kappa_N(r-), Y_{\kappa_N(r-)}^{t,x,N,\delta}) \right) dr \\
&\quad + \int_t^s \left(\sigma(\kappa_N(r-), Y_{\kappa_N(r-)}^{t,x,N,\delta,\mathcal{M}}) - \sigma(\kappa_N(r-), Y_{\kappa_N(r-)}^{t,x,N,\delta}) \right) dW_r \\
&\quad + \int_t^s \int_{A_\delta} \left(\eta_{\kappa_N(r-)}(Y_{\kappa_N(r-)}^{t,x,N,\delta,\mathcal{M}}, z) - \eta_{\kappa_N(r-)}(Y_{\kappa_N(r-)}^{t,x,N,\delta}, z) \right) \tilde{\pi}(dz, dr) \\
&\quad + \int_t^s \int_{A_\delta} \eta_{\kappa_N(r-)}(Y_{\kappa_N(r-)}^{t,x,N,\delta,\mathcal{M}}, z) \nu(dz) dr \\
&\quad - \int_t^s \left(\frac{\nu(A_\delta)}{\mathcal{M}} \sum_{j=1}^{\mathcal{M}} \eta_{\kappa_N(r-)}(Y_{\kappa_N(r-)}^{t,x,N,\delta,\mathcal{M}}, V_{(\kappa_N(r-)-t)N,j}^{N,\mathcal{M}}) \right) dr.
\end{aligned}$$

Analogous to the argument to obtain (3.41), by Burkholder-Davis-Gundy inequality and Hölder's inequality, for each $s \in [t, T]$ it holds that

$$e_{N, \mathcal{M}}^\delta(s) := \mathbb{E} \left[\sup_{r \in [t, s]} \|Y_r^{t, x, N, \delta, \mathcal{M}} - Y_r^{t, x, N, \delta}\|^2 \right] \leq 4 \sum_{i=1}^3 E_i(s) + 4A_4(s), \quad (3.55)$$

where $A_4(s)$ is defined by (3.45), and

$$\begin{aligned} E_1(s) &:= (s-t) \mathbb{E} \left[\int_t^s \|\mu(\kappa_N(r), Y_{\kappa_N(r)}^{t, x, N, \delta, \mathcal{M}}) - \mu(\kappa_N(r), Y_{\kappa_N(r)}^{t, x, N, \delta})\|^2 dr \right], \\ E_2(s) &:= 8 \mathbb{E} \left[\int_t^s \|\sigma(\kappa_N(r), Y_{\kappa_N(r)}^{t, x, N, \delta, \mathcal{M}}) - \sigma(\kappa_N(r), Y_{\kappa_N(r)}^{t, x, N, \delta})\|_F^2 dr \right], \\ E_3(s) &:= 8 \mathbb{E} \left[\int_t^s \int_{A_\delta} \|\eta_{\kappa_N(r)}(Y_{\kappa_N(r)}^{t, x, N, \delta, \mathcal{M}}, z) - \eta_{\kappa_N(r)}(Y_{\kappa_N(r)}^{t, x, N, \delta}, z)\|^2 \nu(dz) dr \right] \end{aligned}$$

Moreover, by (2.3) we have for all $s \in [t, T]$ that

$$E_1(s) \leq 2(s-t)L \int_t^s \mathbb{E} \left[\|Y_{\kappa_N(r)}^{t, x, N, \delta, \mathcal{M}} - Y_{\kappa_N(r)}^{t, x, N, \delta}\|^2 \right] dr, \quad (3.56)$$

$$E_2(s) \leq 16L \int_t^s \mathbb{E} \left[\|Y_{\kappa_N(r)}^{t, x, N, \delta, \mathcal{M}} - Y_{\kappa_N(r)}^{t, x, N, \delta}\|^2 \right] dr, \quad (3.57)$$

$$E_3(s) \leq 16L \int_t^s \mathbb{E} \left[\|Y_{\kappa_N(r)}^{t, x, N, \delta, \mathcal{M}} - Y_{\kappa_N(r)}^{t, x, N, \delta}\|^2 \right] dr. \quad (3.58)$$

Hence, combining (3.55), (3.56)-(3.58), and (3.49) we obtain for all $s \in [t, T]$ that

$$e_{N, \mathcal{M}}^\delta(s) \leq 8L(16+T) \int_t^s e_{N, \mathcal{M}}^\delta(r) dr + 8L\delta^{-2}Kd^p\mathcal{M}^{-1}T \left(\int_t^s \mathbb{E} \left[\|Y_{\kappa_N(r)}^{t, x, N, \delta, \mathcal{M}}\|^2 \right] dr + (s-t)d^p \right).$$

Thus, by (3.37) we have for all $s \in [t, T]$ that

$$e_{N, \mathcal{M}}^\delta(s) \leq 8L(16+T) \int_t^s e_{N, \mathcal{M}}^\delta(r) dr + 16\delta^{-2}Kd^p\mathcal{M}^{-1}LT^2C_1(d^p + \|x\|^2)e^{\rho_1(T-t)}. \quad (3.59)$$

Furthermore, considering (3.33) and (3.37), we apply Grönwall's lemma to (3.59) to get

$$e_{N, \mathcal{M}}^\delta(T) \leq C_2\delta^{-2}d^pK\mathcal{M}^{-1}(d^p + \|x\|^2)e^{\rho_2(T-t)},$$

where $C_2 := 16LT^2C_1$ and $\rho_2 := 8L(16+T)$ are positive constants. Then, the proof of Lemma (3.8) is completed. \square

Lemma 3.9. *There exist positive constants C_1 and ρ_1 only depending on T and L satisfying for all $(t, x) \in [0, T] \times \mathbb{R}^d$, $s \in [t, T]$, $s' \in [s, T]$, $\delta \in (0, 1)$, and $N, \mathcal{M} \in \mathbb{N}$ with $\mathcal{M} \geq \delta^{-2}Kd^p$ that*

$$\mathbb{E} \left[\|Y_{s'}^{t, x, N, \delta, \mathcal{M}} - Y_s^{t, x, N, \delta, \mathcal{M}}\|^2 \right] \leq 8L(s'-s)(T+2+\nu(A_\delta)\mathcal{M}^{-1}) [d^p + C_1(d^p + \|x\|^2)e^{\rho_1 T}] \quad (3.60)$$

Proof. We fix $(t, x) \in [0, T] \times \mathbb{R}^d$, $s \in [t, T]$, $s' \in [s, T]$, $\delta \in (0, 1)$, and $N, \mathcal{M} \in \mathbb{N}$ with $\mathcal{M} \geq \delta^{-2}Kd^p$ throughout this proof. By (3.40), Itô's isometry, Hölder's inequality, and the fact that $(\sum_{i=1}^4 a_i)^2 \leq 4 \sum_{i=1}^4 a_i^2$ for $a_i \in \mathbb{R}$, $i = 1, \dots, 4$, we have

$$\begin{aligned} \mathbb{E} \left[\|Y_{s'}^{t, x, N, \delta, \mathcal{M}} - Y_s^{t, x, N, \delta, \mathcal{M}}\|^2 \right] &\leq 4(s'-s) \int_s^{s'} \mathbb{E} \left[\|\mu(\kappa_N(r), Y_{\kappa_N(r)}^{t, x, N, \delta, \mathcal{M}})\|^2 \right] dr \\ &\quad + 4 \int_s^{s'} \mathbb{E} \left[\|\sigma(\kappa_N(r), Y_{\kappa_N(r)}^{t, x, N, \delta, \mathcal{M}})\|_F^2 \right] dr \\ &\quad + 4 \int_s^{s'} \mathbb{E} \left[\int_{\mathbb{R}^d} \|\eta_{\kappa_N(r)}(Y_{\kappa_N(r)}^{t, x, N, \delta, \mathcal{M}}, z)\|^2 \nu(dz) \right] dr \\ &\quad + 4 \mathbb{E} \left[\left\| \int_s^{s'} \left(\int_{A_\delta} \eta_{\kappa_N(r)}(Y_{\kappa_N(r)}^{t, x, N, \delta, \mathcal{M}}, z) \nu(dz) \right) \right\|^2 \right] \end{aligned}$$

$$- \frac{\nu(A_\delta)}{\mathcal{M}} \sum_{j=1}^{\mathcal{M}} \eta_{\kappa_N(r)}(Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}}, V_{(\kappa_N(r)-t)N,j}^{N,\mathcal{M}}) dr \Big\|^2 \Big]. \quad (3.61)$$

Moreover, by Hölder's inequality, (3.48), and (3.37) it holds that

$$\begin{aligned} & \mathbb{E} \left[\left\| \int_s^{s'} \left(\int_{A_\delta} \eta_{\kappa_N(r)}(Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}}, z) \nu(dz) - \frac{\nu(A_\delta)}{\mathcal{M}} \sum_{j=1}^{\mathcal{M}} \eta_{\kappa_N(r)}(Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}}, V_{(\kappa_N(r)-t)N,j}^{N,\mathcal{M}}) \right) dr \right\|^2 \right] \\ & \leq 2(s' - s)\nu(A_\delta)\mathcal{M}^{-1}L \int_s^{s'} \left(d^p + \mathbb{E} \left[\|Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}}\|^2 \right] \right) dr \\ & \leq 2(s' - s)^2\nu(A_\delta)\mathcal{M}^{-1}L \left[d^p + C_1(d^p + \|x\|^2)e^{\rho_1 T} \right], \end{aligned} \quad (3.62)$$

where $C_1 = C_1(T, L)$ and $\rho_1 = \rho_1(T, L)$ are positive constants introduced in (3.37) in Lemma 3.8. Thus, by (3.61), (3.62), (2.3), and (2.4) we obtain

$$\begin{aligned} & \mathbb{E} \left[\left\| Y_{s'}^{t,x,N,\delta,\mathcal{M}} - Y_s^{t,x,N,\delta,\mathcal{M}} \right\|^2 \right] \\ & \leq 4(s' - s) \int_s^{s'} \mathbb{E} \left[2 \left\| \mu(\kappa_N(r), Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}}) - \mu(\kappa_N(r), 0) \right\|^2 + 2 \left\| \mu(\kappa_N(r), 0) \right\|^2 \right] dr \\ & \quad + 4 \int_s^{s'} \mathbb{E} \left[2 \left\| \sigma(\kappa_N(r), Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}}) - \sigma(\kappa_N(r), 0) \right\|_F^2 + 2 \left\| \sigma(\kappa_N(r), 0) \right\|^2 \right] dr \\ & \quad + 4 \int_s^{s'} \mathbb{E} \left[\int_{\mathbb{R}^d} \left(2 \left\| \eta_{\kappa_N(r)}(Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}}, z) - \eta_{\kappa_N(r)}(0, z) \right\|^2 + 2 \left\| \eta_{\kappa_N(r)}(0, z) \right\|^2 \right) \nu(dz) \right] dr \\ & \quad + 8(s' - s)^2\nu(A_\delta)\mathcal{M}^{-1}L \left[d^p + C_1(d^p + \|x\|^2)e^{\rho_1 T} \right] \\ & \leq 8L(s' - s) \int_s^{s'} \left(\mathbb{E} \left[\|Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}}\|^2 + d^p \right] \right) dr + 16L \int_s^{s'} \left(\mathbb{E} \left[\|Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}}\|^2 + d^p \right] \right) dr \\ & \quad + 8(s' - s)^2\nu(A_\delta)\mathcal{M}^{-1}L \left[d^p + C_1(d^p + \|x\|^2)e^{\rho_1 T} \right]. \end{aligned}$$

Hence, by (3.37) we obtain for all $s \in [t, T]$ and $s' \in [s, T]$ that

$$\begin{aligned} \mathbb{E} \left[\left\| Y_{s'}^{t,x,N,\delta,\mathcal{M}} - Y_s^{t,x,N,\delta,\mathcal{M}} \right\|^2 \right] & \leq 8L(s' - s)^2 \left[d^p + C_1(d^p + \|x\|^2)e^{\rho_1 T} \right] \\ & \quad + 16L(s' - s) \left[d^p + C_1(d^p + \|x\|^2)e^{\rho_1 T} \right] \\ & \quad + 8(s' - s)\nu(A_\delta)\mathcal{M}^{-1}L \left[d^p + C_1(d^p + \|x\|^2)e^{\rho_1 T} \right] \\ & \leq 8L(s' - s)(T + 2 + \nu(A_\delta)\mathcal{M}^{-1}) \left[d^p + C_1(d^p + \|x\|^2)e^{\rho_1 T} \right], \end{aligned}$$

which completes the proof of this lemma. \square

Lemma 3.10. *There exist constants C_1 and ρ_1 only depending on T and L satisfying for all $t \in [0, T]$, $t' \in [t, T]$, $s \in [t', T]$, $x, x' \in \mathbb{R}^d$, $\delta \in (0, 1)$, and $N, \mathcal{M} \in \mathbb{N}$ with $\mathcal{M} \geq \delta^{-2}Kd^p$ that*

$$\begin{aligned} e_s & := \sup_{u \in [t', s]} \mathbb{E} \left[\left\| Y_u^{t,x,N,\delta,\mathcal{M}} - Y_u^{t',x',N,\delta,\mathcal{M}} \right\|^2 \right] \\ & \leq 16 \exp \left\{ 4LT (2T + 8 + \nu(A_\delta)\mathcal{M}^{-1}T) \right\} \\ & \quad \cdot \left\{ (\|x - x'\| + |t' - t| + |t' - t|^{1/2}) [1 + 3L^{1/2}d^{p/2}(1 + \|x\|^2)^{1/2}]^2 \right. \\ & \quad \left. + 2L(t' - t)\nu(A_\delta)\mathcal{M}^{-1} \left[d^p + C_1(d^p + \|x\|^2)e^{\rho_1 T} \right] \right\}. \end{aligned} \quad (3.63)$$

Proof. We fix $x, x' \in \mathbb{R}^d$, $t \in [0, T]$, $t' \in [t, T]$, $\delta \in (0, 1)$, and $N, \mathcal{M} \in \mathbb{N}$ with $\mathcal{M} \geq \delta^{-2}Kd^p$ throughout this proof. One notices that almost surely for all $s \in [t', T]$

$$\begin{aligned} & Y_s^{t,x,N,\delta,\mathcal{M}} - Y_s^{t',x',N,\delta,\mathcal{M}} \\ & = Y_{t'}^{t,x,N,\delta,\mathcal{M}} - x' + \int_{t'}^s \left(\mu(\kappa_N(r-), Y_{\kappa_N(r-)}^{t,x,N,\delta,\mathcal{M}}) - \mu(\kappa_N(r-), Y_{\kappa_N(r-)}^{t',x',N,\delta,\mathcal{M}}) \right) dr \end{aligned}$$

$$\begin{aligned}
& + \int_{t'}^s \left(\sigma(\kappa_N(r-), Y_{\kappa_N(r-)}^{t,x,N,\delta,\mathcal{M}}) - \sigma(\kappa_N(r-), Y_{\kappa_N(r-)}^{t',x',N,\delta,\mathcal{M}}) \right) dW_r \\
& + \int_{t'}^s \int_{A_\delta} \left(\eta_{\kappa_N(r-)}(Y_{\kappa_N(r-)}^{t,x,N,\delta,\mathcal{M}}, z) - \eta_{\kappa_N(r-)}(Y_{\kappa_N(r-)}^{t',x',N,\delta,\mathcal{M}}, z) \right) \tilde{\pi}(dz, dr) \\
& + \int_{t'}^s \int_{A_\delta} \left(\eta_{\kappa_N(r-)}(Y_{\kappa_N(r-)}^{t,x,N,\delta,\mathcal{M}}, z) - \eta_{\kappa_N(r-)}(Y_{\kappa_N(r-)}^{t',x',N,\delta,\mathcal{M}}, z) \right) \nu(dz) dr \\
& - \int_{t'}^s \left[\frac{\nu(A_\delta)}{\mathcal{M}} \sum_{j=1}^{\mathcal{M}} \left(\eta_{\kappa_N(r-)}(Y_{\kappa_N(r-)}^{t,x,N,\delta,\mathcal{M}}, V_{(\kappa_N(r-)-t)N,j}^{N,\mathcal{M}} \right) \right. \\
& \quad \left. - \eta_{\kappa_N(r-)}(Y_{\kappa_N(r-)}^{t',x',N,\delta,\mathcal{M}}, V_{(\kappa_N(r-)-t)N,j}^{N,\mathcal{M}}) \right] dr.
\end{aligned}$$

Therefore, by Minkowski's inequality, Hölder's inequality, Itô's isometry, Hölder's inequality, and the analogous argument to get (3.47), we have for all $s \in [t', T]$ that

$$\begin{aligned}
e_s^{1/2} & \leq \left(\mathbb{E} \left[\left\| Y_{t'}^{t,x,N,\delta,\mathcal{M}} - x' \right\|^2 \right] \right)^{1/2} + L^{1/2} \left((s-t) \int_{t'}^s \mathbb{E} \left[\left\| Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}} - Y_{\kappa_N(r)}^{t',x',N,\delta,\mathcal{M}} \right\|^2 \right] dr \right)^{1/2} \\
& + 2L^{1/2} \left(\int_{t'}^s \mathbb{E} \left[\left\| Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}} - Y_{\kappa_N(r)}^{t',x',N,\delta,\mathcal{M}} \right\|^2 \right] dr \right)^{1/2} \\
& + \left(\nu(A_\delta) \mathcal{M}^{-1} T \int_{t'}^s \mathbb{E} \left[\int_{A_\delta} \left\| \eta_{\kappa_N(r)}(Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}}, z) - \eta_{\kappa_N(r)}(Y_{\kappa_N(r)}^{t',x',N,\delta,\mathcal{M}}, z) \right\|^2 \nu(dz) \right] dr \right)^{1/2}.
\end{aligned}$$

Thus, by (2.3) and the fact that $(\sum_{i=1}^4 a_i)^2 \leq 4 \sum_{i=1}^4 a_i^2$ for $a_i \in \mathbb{R}$, $i = 1, \dots, 4$, we have for all $s \in [t', T]$ that

$$e_s \leq 4\mathbb{E} \left[\left\| Y_{t'}^{t,x,N,\delta,\mathcal{M}} - x' \right\|^2 \right] + 4L(T+4) \int_{t'}^s e_r dr + 4L\nu(A_\delta) \mathcal{M}^{-1} T \int_{t'}^s e_r dr. \quad (3.64)$$

Hence, by (3.37) and Grönwall's lemma, we get for all $s \in [t', T]$ that

$$e_s \leq 4\mathbb{E} \left[\left\| Y_{t'}^{t,x,N,\delta,\mathcal{M}} - x' \right\|^2 \right] \cdot \exp \{ 4LT(T+4 + \nu(A_\delta) \mathcal{M}^{-1} T) \}. \quad (3.65)$$

Furthermore, we notice for all $s \in [t, t']$ almost surely that

$$\begin{aligned}
Y_s^{t,x,N,\delta,\mathcal{M}} - x' & = x - x' + \int_t^s \mu(\kappa_N(r-), Y_{\kappa_N(r-)}^{t,x,N,\delta,\mathcal{M}}) dr + \int_t^s \sigma(\kappa_N(r-), Y_{\kappa_N(r-)}^{t,x,N,\delta,\mathcal{M}}) dW_r \\
& + \int_t^s \int_{\mathbb{R}^d} \eta_{\kappa_N(r-)}(Y_{\kappa_N(r-)}^{t,x,N,\delta,\mathcal{M}}, z) \tilde{\pi}(dz, dr) \\
& + \int_t^s \int_{A_\delta} \eta_{\kappa_N(r-)}(Y_{\kappa_N(r-)}^{t,x,N,\delta,\mathcal{M}}, z) \nu(dz) dr \\
& - \int_t^s \left(\frac{\nu(A_\delta)}{\mathcal{M}} \sum_{j=1}^{\mathcal{M}} \eta_{\kappa_N(r-)}(Y_{\kappa_N(r-)}^{t,x,N,\delta,\mathcal{M}}, V_{(\kappa_N(r-)-t)N,j}^{N,\mathcal{M}}) \right) dr.
\end{aligned}$$

Then, by Minkowski's inequality, Hölder's inequality, Itô's isometry, (3.62), (2.3), and (2.4), we obtain for all $s \in [t, t']$ that

$$\begin{aligned}
& \left(\sup_{u \in [t, s]} \mathbb{E} \left[\left\| Y_u^{t,x,N,\delta,\mathcal{M}} - x' \right\|^2 \right] \right)^{1/2} \\
& \leq \|x - x'\| + \left((s-t) \int_t^s \|\mu(\kappa_N(r), x')\|^2 dr \right)^{1/2} \\
& + \left((s-t) \int_t^s \mathbb{E} \left[\left\| \mu(\kappa_N(r), Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}}) - \mu(\kappa_N(r), x') \right\|^2 \right] dr \right)^{1/2} \\
& + \left(\int_t^s \|\sigma(\kappa_N(r), x')\|_F^2 dr \right)^{1/2} + \left(\int_t^s \mathbb{E} \left[\left\| \sigma(\kappa_N(r), Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}}) - \sigma(\kappa_N(r), x') \right\|_F^2 \right] dr \right)^{1/2}
\end{aligned}$$

$$\begin{aligned}
& + \left(\int_t^s \mathbb{E} \left[\int_{\mathbb{R}^d} \left\| \eta_{\kappa_N(r)}(Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}}, z) - \eta_{\kappa_N(r)}(x', z) \right\|^2 \nu(dz) \right] dr \right)^{1/2} \\
& + \left(\int_t^s \int_{\mathbb{R}^d} \left\| \eta_{\kappa_N(r)}(x', z) \right\|^2 \nu(dz) dr \right)^{1/2} \\
& + [2(s-t)^2 \nu(A_\delta) \mathcal{M}^{-1} L(d^p + C_1(d^p + \|x\|^2) e^{\rho_1 T})]^{1/2} \\
\leq & \|x - x'\| + \left((s-t) \int_t^s \left\| \mu(\kappa_N(r), x') \right\|^2 dr \right)^{1/2} + L^{1/2} \left((s-t) \int_t^s \mathbb{E} \left[\left\| Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}} - x' \right\|^2 \right] dr \right)^{1/2} \\
& + 2L^{1/2} \left(\int_t^s \mathbb{E} \left[\left\| Y_{\kappa_N(r)}^{t,x,N,\delta,\mathcal{M}} - x' \right\|^2 \right] dr \right)^{1/2} + \left(\int_t^s \left\| \sigma(\kappa_N(r), x') \right\|_F^2 dr \right)^{1/2} \\
& + \left(\int_t^s \int_{\mathbb{R}^d} \left\| \eta_{\kappa_N(r)}(x', z) \right\|^2 \nu(dz) dr \right)^{1/2} + [2(s-t)^2 \nu(A_\delta) \mathcal{M}^{-1} L(d^p + C_1(d^p + \|x\|^2) e^{\rho_1 T})]^{1/2}.
\end{aligned}$$

Hence, by (2.4) and the fact that $(\sum_{i=1}^4 a_i)^2 \leq 4 \sum_{i=1}^4 a_i^2$ for $a_i \in \mathbb{R}$, $i = 1, \dots, 4$, it holds for all $s \in [t, t']$ that

$$\begin{aligned}
\sup_{u \in [t, s]} \mathbb{E} \left[\left\| Y_u^{t,x,N,\delta,\mathcal{M}} - x' \right\|^2 \right] & \leq 4L(T+4) \int_t^s \sup_{u \in [t, r]} \mathbb{E} \left[\left\| Y_u^{t,x,N,\delta,\mathcal{M}} - x' \right\|^2 \right] dr \\
& + 4 \left[\|x - x'\| + \left(|t' - t| \cdot \int_t^{t'} \left\| \mu(\kappa_N(r), x') \right\|^2 dr \right)^{1/2} \right. \\
& + \left(\int_t^{t'} \left\| \sigma(\kappa_N(r), x') \right\|_F^2 dr \right)^{1/2} \\
& + \left. \left(\int_t^{t'} \int_{\mathbb{R}^d} \left\| \eta_{\kappa_N(r)}(x', z) \right\|^2 \nu(dz) dr \right)^{1/2} \right]^2 \\
& + 8L(t' - t) \nu(A_\delta) \mathcal{M}^{-1} [d^p + C_1(d^p + \|x\|^2) e^{\rho_1 T}] \\
\leq & 4L(T+4) \int_t^s \sup_{u \in [t, r]} \mathbb{E} \left[\left\| Y_u^{t,x,N,\delta,\mathcal{M}} - x' \right\|^2 \right] dr \\
& + 4(\|x - x'\| + |t' - t| + |t' - t|^{1/2}) [1 + 3L^{1/2} d^{p/2} (1 + \|x\|^2)^{1/2}]^2 \\
& + 8L(t' - t) \nu(A_\delta) \mathcal{M}^{-1} [d^p + C_1(d^p + \|x\|^2) e^{\rho_1 T}].
\end{aligned}$$

Thus, by (3.37) and Grönwall's lemma, we obtain that

$$\begin{aligned}
& \sup_{u \in [t, t']} \mathbb{E} \left[\left\| Y_u^{t,x,N,\delta,\mathcal{M}} - x' \right\|^2 \right] \\
& \leq 4e^{4LT(T+4)} \left\{ (\|x - x'\| + |t' - t| + |t' - t|^{1/2}) [1 + 3L^{1/2} d^{p/2} (1 + \|x\|^2)^{1/2}]^2 \right. \\
& \quad \left. + 2L(t' - t) \nu(A_\delta) \mathcal{M}^{-1} [d^p + C_1(d^p + \|x\|^2) e^{\rho_1 T}] \right\}. \tag{3.66}
\end{aligned}$$

Finally, combining (3.65) and (3.66) yields (3.63). Thus, the proof of this lemma is completed. \square

Corollary 3.11. *Let $(t, x) \in [0, T] \times \mathbb{R}^d$, $s \in [t, T]$, and let $(t_k, s_k, x_k)_{k=1}^\infty$ be a sequence such that $(t_k, s_k, x_k) \in \Lambda \times \mathbb{R}^d$ for all $k \in \mathbb{N}$, and $\lim_{k \rightarrow \infty} (t_k, s_k, x_k) = (t, s, x)$. Then it holds for for all $\varepsilon > 0$, $\delta \in (0, 1)$, and $N, \mathcal{M} \in \mathbb{N}$ with $\mathcal{M} \geq \delta^{-2} K d^p$, that*

$$\lim_{k \rightarrow \infty} \mathbb{P} \left(\left\| Y_{s_k}^{t_k, x_k, N, \delta, \mathcal{M}} - Y_s^{t, x, N, \delta, \mathcal{M}} \right\| \geq \varepsilon \right) = 0. \tag{3.67}$$

Proof. By Lemma 3.9 and Lemma 3.10, one can precisely follow the proof of Corollary 3.5 to obtain (3.67). \square

4. STOCHASTIC FIXED-POINT EQUATIONS

The following results from [39] provide existence and uniqueness of solutions of stochastic fixed point equations. These results will play an important role in the proof of Theorem 2.9 in Section 7.

Proposition 4.1. Let $d \in \mathbb{N}$, $a, b, c, p, T \in (0, \infty)$, and let $Z_s^{t,x} : \Omega \rightarrow \mathbb{R}^d$ be a random variable for each $t \in [0, T]$, $s \in [t, T]$, and $x \in \mathbb{R}^d$. For every nonnegative Borel function $\varphi : [0, T] \times \Omega \rightarrow [0, \infty)$, we assume that the mapping $\{(t, s) \in [0, T]^2 : t \leq s\} \times \mathbb{R}^d \ni (t, s, x) \mapsto \mathbb{E}[\varphi(s, Z_s^{t,x})] \in [0, \infty)$ is measurable. Let $f : [0, T] \times \mathbb{R}^d \times \mathbb{R} \rightarrow \mathbb{R}$ and $g : \mathbb{R}^d \rightarrow \mathbb{R}$ be Borel functions, and assume for all $t \in [0, T]$, $s \in [t, T]$, $x \in \mathbb{R}^d$, and $y, y' \in \mathbb{R}$ that $|f(t, x, 0)| \leq a(d^p + \|x\|^2)^{1/2}$, $|g(x)| \leq a(d^p + \|x\|^2)^{1/2}$, $\mathbb{E}[(d^p + \|Z_s^{t,x}\|^2)^{1/2}] \leq b(d^p + \|x\|^2)^{1/2}$ and $|f(t, x, y) - f(t, x, y')| \leq c|y - y'|$.

Then there exists a unique Borel function $u : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$ satisfying for all $(t, x) \in [0, T] \times \mathbb{R}^d$ that

$$\mathbb{E} \left[|g(Z_T^{t,x})| \right] + \int_t^T \mathbb{E} \left[|f(s, Z_s^{t,x}, u(s, Z_s^{t,x}))| \right] ds + \sup_{y \in \mathbb{R}^d} \sup_{s \in [0, T]} \left(\frac{|u(s, y)|}{(d^p + \|y\|^2)^{1/2}} \right) < \infty$$

and

$$u(t, x) = \mathbb{E} \left[g(Z_T^{t,x}) \right] + \int_t^T \mathbb{E} \left[f(s, Z_s^{t,x}, u(s, Z_s^{t,x})) \right] ds.$$

Moreover, it holds for all $t \in [0, T]$ that

$$\sup_{x \in \mathbb{R}^d} \left(\frac{|u(t, x)|}{(d^p + \|x\|^2)^{1/2}} \right) \leq \left[\sup_{x \in \mathbb{R}^d} \left(\frac{|g(x)|}{(d^p + \|x\|^2)^{1/2}} \right) + \sup_{x \in \mathbb{R}^d} \sup_{s \in [t, T]} \left(\frac{T|f(s, x, 0)|}{(d^p + \|x\|^2)^{1/2}} \right) \right] e^{c(T-t)} < \infty.$$

Proof. For $b = 1$, the above proposition is proved in [39] as Proposition 2.2 (with $X_{t,s}^x \curvearrowright Z_s^{t,x}$, $\psi \curvearrowright \varphi$, $\mathcal{O} \curvearrowright \mathbb{R}^d$, $T \curvearrowright T$, $c \curvearrowright a$, $1 \curvearrowright b$, $L \curvearrowright c$, $V \curvearrowright (\mathbb{R}^d \ni x \mapsto (d^p + \|x\|^2)^{1/2} \in \mathbb{R})$, $u \curvearrowright u$, $f \curvearrowright f$, $g \curvearrowright g$ in the notation of Proposition 2.2 in [39]). For $b \in (0, \infty)$, one can precisely follow the proof of Proposition 2.2 in [39] to obtain the desired results. \square

Lemma 4.2. Let $d \in \mathbb{N}$, $c, \rho, \eta, L \in [0, \infty)$, $T, \delta \in (0, \infty)$, let $(X_s^{t,x,k})_{s \in [t, T]} : [t, T] \times \Omega \rightarrow \mathbb{R}^d$, $t \in [0, T]$, $x \in \mathbb{R}^d$, $k \in \{1, 2\}$, be $\mathcal{B}([0, T]) \otimes \mathcal{F}/\mathcal{B}(\mathbb{R}^d)$ -measurable, let $f : [0, T] \times \mathbb{R}^d \times \mathbb{R} \rightarrow \mathbb{R}$, $g : \mathbb{R}^d \rightarrow \mathbb{R}$, and $u_k : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$, $k \in \{1, 2\}$, be Borel functions. For all $t \in [0, T]$, $s \in [t, T]$, and all Borel functions $h : \mathbb{R}^d \times \mathbb{R}^d \rightarrow [0, \infty)$ assume that $\mathbb{R}^d \times \mathbb{R}^d \ni (y_1, y_2) \mapsto \mathbb{E}[h(X_s^{t,y_1,1}, X_s^{t,y_2,1})] \in [0, \infty)$ is measurable. Moreover, for all $x, y \in \mathbb{R}^d$, $t \in [0, T]$ and $s \in [t, T]$, $r \in [s, T]$, $v, w \in \mathbb{R}$, $k \in \{1, 2\}$, and all Borel functions $h : \mathbb{R}^d \times \mathbb{R}^d \rightarrow [0, \infty)$ assume that

$$X_t^{t,x,k} = x, \quad \mathbb{E} \left[d^p + \|X_s^{t,x,k}\|^2 \right] \leq \eta(d^p + \|x\|^2),$$

$$\mathbb{E} \left[\mathbb{E} \left[h(X_r^{s,x',1}, X_r^{s,y',1}) \right] \Big|_{(x',y')=(X_s^{t,x,1}, X_s^{t,y,1})} \right] = \mathbb{E} \left[h(X_r^{t,x,1}, X_r^{t,y,1}) \right],$$

$$\max\{T|f(t, x, v) - f(t, y, w)|, |g(x) - g(y)|\} \leq L \left(T|v - w| + T^{-1/2}(2d^p + \|x\|^2 + \|y\|^2)^{1/2} \|x - y\| \right),$$

$$\mathbb{E} \left[\mathbb{E} \left[\|X_r^{s,x',1} - X_r^{s,y',1}\|^2 \right] \Big|_{(x',y')=(X_s^{t,x,1}, X_s^{t,y,2})} \right] \leq \delta^2(d^p + \|x\|^2),$$

$$\mathbb{E} \left[|g(X_T^{t,x,k})| \right] + \int_t^T \mathbb{E} \left[|f(s, X_s^{t,x,k}, u_k(s, X_s^{t,x,k}))| \right] ds < \infty,$$

$$u_k(t, x) = \mathbb{E} \left[g(X_T^{t,x,k}) + \int_t^T f(s, X_s^{t,x,k}, u_k(s, X_s^{t,x,k})) ds \right],$$

and

$$\max \left\{ |u_k(t, x)|^2, e^{\rho(t-s)} \mathbb{E} \left[d^p + \|X_s^{t,x,k}\|^2 \right] \right\} \leq c(d^p + \|x\|^2). \quad (4.1)$$

Then for all $(t, x) \in [0, T] \times \mathbb{R}^d$ it holds that

$$|u_1(t, x) - u_2(t, x)| \leq 4c\delta T^{1/2} L(1 + LT)(d^p + \|x\|^2) \exp \left\{ (L + \rho/2 + (\eta c)^{1/2} L)(T - t) \right\}. \quad (4.2)$$

Proof. The above lemma is basically a special version of Lemma 2.3 in [39] (with $(X_{t,s}^{x,k})_{s \in [t, T]} \curvearrowright (X_s^{t,x,k})_{s \in [t, T]}$, $\eta \curvearrowright \eta$, $L \curvearrowright L$, $T \curvearrowright T$, $\psi \curvearrowright (\mathbb{R}^d \ni x \mapsto (d^p + \|x\|^2) \in \mathbb{R})$, $T^{1/2}$ in (27) in [39] $\curvearrowright LT^{1/2}$, $h \curvearrowright h$, $V \curvearrowright (\mathbb{R}^d \ni x \mapsto (d^p + \|x\|^2) \in \mathbb{R})$, $p \curvearrowright 2$, $q \curvearrowright 2$, $\delta \curvearrowright \delta$, $g \curvearrowright g$, $f \curvearrowright f$, $u_k \curvearrowright u_k$, $ce^{\rho(t-s)}$ in the last condition of Lemma 2.3 in [39] $\curvearrowright e^{\rho(t-s)}$ in the notation of Lemma 2.3 in [39]). One can follow the proof of Lemma 2.3 in [39] step by step to get (4.2). \square

5. EXISTENCE AND UNIQUENESS OF VISCOSITY SOLUTIONS OF PIDE (2.18) AND ITS FEYNMAN-KAC REPRESENTATION

There are many existence and uniqueness results for viscosity solutions of nonlinear second-order PIDEs in the literature (c.f. e.g., [2, 5, 6, 20, 43, 44]). For the comparison principle, we verify that the conditions of Theorem 3.1 in [44] are satisfied; see Proposition 5.4. Moreover, we show that there exists a viscosity solution of PIDE (2.18) which satisfies a Feynman-Kac representation; see Proposition 5.16. The idea is to first show for a corresponding linear PIDE having smooth coefficients that it possesses a classical solution satisfying a Feynman-Kac representation. Then, combining a limit argument involving mollifiers as well as a stochastic fixed point argument based on [12, 39] we can conclude the desired result.

We assume the settings in Section 2.1, fix d and θ , and omit the superscript d and θ in the notations. Then we introduce some notions and notations which will be used throughout this section. To ease notation we introduce some functionals. Let $H : (0, T) \times \mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d \times \mathbb{S}^d \rightarrow \mathbb{R}$ be a functional defined for each $(t, x, r, y, A) \in (0, T) \times \mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d \times \mathbb{S}^d$ by

$$H(t, x, r, y, A) := -\langle y, \mu(t, x) \rangle - \frac{1}{2} \text{Trace}(\sigma(t, x)[\sigma(t, x)]^T A) - f(t, x, r).$$

Let $G_0 : (0, T) \times \mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d \times \mathbb{S}^d \times C^2(\mathbb{R}^d) \rightarrow \mathbb{R}$ be a functional defined for all $(t, x, r, y, A, \phi) \in (0, T) \times \mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d \times \mathbb{S}^d \times C^2(\mathbb{R}^d)$ by

$$G_0(t, x, r, y, A, \phi) := G(t, x, r, y, A, \phi) + f(t, x, r) - h(t, x),$$

where G is defined by (2.17), and $h \in C([0, T] \times \mathbb{R}^d)$ is a fixed function satisfying the follows.

Assumption 5.1. *There exist a positive constant C_0 satisfying for all $(t, x) \in [0, T] \times \mathbb{R}^d$ that*

$$|h(t, x)|^2 \leq C_0(1 + \|x\|^2). \quad (5.1)$$

For every $\kappa \in (0, 1)$, let $G_\kappa : (0, T) \times \mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d \times \mathbb{S}^d \times SC((0, T) \times \mathbb{R}^d) \times C^{1,2}((0, T) \times \mathbb{R}^d) \rightarrow \mathbb{R}$ be a functional defined for each $(t, x, r, y, A, \psi, \varphi) \in (0, T) \times \mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d \times \mathbb{S}^d \times SC((0, T) \times \mathbb{R}^d) \times C^{1,2}((0, T) \times \mathbb{R}^d)$ by

$$\begin{aligned} G_\kappa(t, x, r, y, A, v, \varphi) := & H(t, x, r, y, A) \\ & - \int_{\|z\| \geq \kappa} (\psi(t, x + \eta_t(x, z)) - \psi(t, x) - \langle y, \eta_t(x, z) \rangle) \nu(dz) \\ & - \int_{\|z\| < \kappa} (\varphi(t, x + \eta_t(x, z)) - \varphi(t, x) - \langle \nabla_x \varphi(t, x), \eta_t(x, z) \rangle) \nu(dz). \end{aligned}$$

Lemma 5.2. *Let $u \in USC_1((0, T) \times \mathbb{R}^d)$ be a viscosity subsolution of PIDE (2.18). Let $\varphi \in C^{1,2}((0, T) \times \mathbb{R}^d)$, $(t_0, x_0) \in (0, T) \times \mathbb{R}^d$, and assume that $u - \varphi$ has a local maximum at (t_0, x_0) . Then*

$$-\frac{\partial}{\partial t} \varphi(t_0, x_0) + G(t_0, x_0, u(t_0, x_0), \nabla_x \varphi(t_0, x_0), \text{Hess}_x \varphi(t_0, x_0), \varphi(t, \cdot)) \leq 0.$$

Proof. First, notice that the upper semi-continuity of u ensures that there exist a non-empty open set $\mathcal{O} \subseteq (0, T) \times \mathbb{R}^d$ and $\phi \in C^{1,2}((0, T) \times \mathbb{R}^d)$ such that $(t_0, x_0) \in \mathcal{O}$, $\varphi(t, x) = \phi(t, x)$ for all $(t, x) \in \mathcal{O}$, and $u - \phi$ has a global maximum at (t_0, x_0) . Then we define the function

$$\phi_0(t, x) := \phi(t, x) - \phi(t_0, x_0) + u(t_0, x_0), \quad (t, x) \in (0, T) \times \mathbb{R}^d.$$

Note that $\phi_0 \in C^{1,2}((0, T) \times \mathbb{R}^d)$, $\phi_0(t_0, x_0) = u(t_0, x_0)$, and $u \leq \phi_0$. Thus, the fact $u \in USC_1((0, T) \times \mathbb{R}^d)$ is a viscosity subsolution of PIDE (2.18) implies that

$$-\frac{\partial}{\partial t} \phi_0(t_0, x_0) + G(t_0, x_0, \phi_0(t_0, x_0), \nabla_x \phi_0(t_0, x_0), \text{Hess}_x \phi_0(t_0, x_0), \phi_0(t, \cdot)) \leq 0.$$

Hence, by the definition of ϕ_0 , and the fact that $\phi_0(t_0, x_0) = u(t_0, x_0)$ and $\phi_0(t, x) = \varphi(t, x) - \phi(t_0, x_0) + u(t_0, x_0)$ for all $(t, x) \in \mathcal{O}$ we have

$$-\frac{\partial}{\partial t} \varphi(t_0, x_0) + G(t_0, x_0, u(t_0, x_0), \nabla_x \varphi(t_0, x_0), \text{Hess}_x \varphi(t_0, x_0), \varphi(t, \cdot)) \leq 0.$$

This completes the proof of Lemma 5.2. □

Lemma 5.3. *Let Assumptions 2.1, 2.2, and 5.1 hold. Then for every $\varphi \in C^{1,2}((0, T) \times \mathbb{R}^d)$, the mapping*

$$(0, T) \times \mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d \times \mathbb{S}^d \ni (t, x, r, y, A) \mapsto G_0(t, x, r, y, A, \varphi(t, \cdot)) \in \mathbb{R}$$

is continuous.

Proof. By the definition of G_0 , to prove this lemma we only need to show that for every $\varphi \in C^{1,2}((0, T) \times \mathbb{R}^d)$ the mapping

$$(0, T) \times \mathbb{R}^d \ni (t, x) \mapsto \int_{\mathbb{R}^d} \left(\varphi(t, x + \eta_t(x, z)) - \varphi(t, x) - \langle \nabla_x \varphi(t, x), \eta_t(x, z) \rangle \right) \nu(dz) \in \mathbb{R} \quad (5.2)$$

is continuous. To see this, note that by Taylor's formula we obtain for every $\varphi \in C^{1,2}((0, T) \times \mathbb{R}^d)$ and $(t, x) \in (0, T) \times \mathbb{R}^d$ that

$$\begin{aligned} & \int_{\mathbb{R}^d} \left(\varphi(t, x + \eta_t(x, z)) - \varphi(t, x) - \langle \nabla_x \varphi(t, x), \eta_t(x, z) \rangle \right) \nu(dz) \\ &= \int_{\mathbb{R}^d} \int_0^1 (1 - \alpha) \sum_{i,j=1}^d \left[\frac{\partial^2}{\partial x_i \partial x_j} \varphi(t, x + \alpha \eta_t(x, z)) \eta_t^i(x, z) \eta_t^j(x, z) \right] d\alpha \nu(dz) \end{aligned} \quad (5.3)$$

Moreover, by (2.6) and Cauchy-Schwarz inequality we have for all $\alpha \in [0, 1]$, $z \in \mathbb{R}^d$, $(t, x) \in (0, T) \times \mathbb{R}^d$, and $(t', x') \in (0, T) \times \mathbb{R}^d$ with $|t' - t| \leq [(T - t) \wedge t]/2$ and $\|x' - x\| \leq 1$ that

$$\begin{aligned} & \left| (1 - \alpha) \sum_{i,j=1}^d \left[\frac{\partial^2}{\partial x_i \partial x_j} \varphi(t', x' + \alpha \eta_{t'}(x', z)) \eta_{t'}^i(x', z) \eta_{t'}^j(x', z) \right] \right| \\ & \leq \sup_{s: |s-t| \leq \frac{(T-t) \wedge t}{2}} \sup_{y: \|y-x\| \leq 1 + C_d^{1/2}} \left[\|\text{Hess}_y \varphi(s, y)\|_F \right] \cdot \sum_{i,j=1}^d \left| \eta_{t'}^i(x', z) \eta_{t'}^j(x', z) \right| \\ & \leq \sup_{s: |s-t| \leq \frac{(T-t) \wedge t}{2}} \sup_{y: \|y-x\| \leq 1 + C_d^{1/2}} \left[\|\text{Hess}_y \varphi(s, y)\|_F \right] \cdot d \|\eta_{t'}(x', z)\|^2 \\ & \leq \sup_{s: |s-t| \leq \frac{(T-t) \wedge t}{2}} \sup_{y: \|y-x\| \leq 1 + C_d^{1/2}} \left[\|\text{Hess}_y \varphi(s, y)\|_F \right] \cdot d C_d (1 \wedge \|z\|^2). \end{aligned} \quad (5.4)$$

Thus, by (5.3) and Lebesgue's dominated convergence theorem we obtain that (5.2) is continuous. The proof of this lemma is therefore completed. \square

Proposition 5.4. *Let Assumptions 2.1 and 2.2 hold, and let $u_1, u_2 \in C_1([0, T] \times \mathbb{R}^d)$ be two viscosity solutions of PIDE (2.18) such that $u_1(T, x) = u_2(T, x) = g(x)$ for all $x \in \mathbb{R}^d$. Then we have for all $(t, x) \in [0, T] \times \mathbb{R}^d$ that $u_1(t, x) = u_2(t, x)$.*

Proof. By (2.2) and Rademacher's theorem (see, e.g., Theorem 3.1.6 in [29]) we first notice for almost all $x \in \mathbb{R}^d$ that $\nabla_x g(x)$ exists, and

$$\|\nabla_x g(x)\| \leq L^{1/2} T^{-1/2}. \quad (5.5)$$

To apply Theorem 3.1 in [44], it remains to verify that Assumptions (C1)-(C4) and (F0)-(F6) in [44] are satisfied. Note that by Remark 2.2 in [44], Assumptions (F0)-(F4) in [44] implies Assumptions (C1)-(C4) in [44]. Thus, we only need to verify Assumptions (F0)-(F6). In the rest of this proof, we fix $u, -v \in USC_1((0, T) \times \mathbb{R}^d)$, $w \in SC_1((0, T) \times \mathbb{R}^d)$, $\varphi, \psi \in C_1^{1,2}((0, T) \times \mathbb{R}^d)$. To ease notation, for each $\phi \in C^{1,2}((0, T) \times \mathbb{R}^d)$ and $(t, x) \in (0, T) \times \mathbb{R}^d$ we define

$$J\phi(t, x) := \int_{\mathbb{R}^d} \left(\phi(t, x + \eta_t(x, z)) - \phi(t, x) - \langle \nabla_x \phi(t, x), \eta_t(x, z) \rangle \right) \nu(dz).$$

By the definition of G and G_κ , we first notice that (F0) obviously holds (with $F \curvearrowright G$ and $F_\kappa \curvearrowright G_\kappa$ in the notation of [44]). Next, by Lemma 5.3 and the continuity of h and f , we notice that the mapping

$$(0, T) \times \mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d \times \mathbb{S}^d \ni (t, x, r, y, A) \mapsto G(t, x, r, y, A, \varphi(t, \cdot)) \in \mathbb{R} \quad (5.6)$$

is continuous. Furthermore, for each $k \in \mathbb{N}$ let $\varphi_k \in C^{1,2}((0, T) \times \mathbb{R}^d)$, $(t_k, x_k) \in (0, T) \times \mathbb{R}^d$, and let $(t_0, x_0) \in (0, T) \times \mathbb{R}^d$. Assume that $(t_k, x_k) \rightarrow (t_0, x_0)$ as $k \rightarrow \infty$, and

$$\varphi_k \rightarrow \varphi, \quad \nabla_x \varphi_k \rightarrow \nabla_x \varphi, \quad \text{Hess}_x \varphi_k \rightarrow \text{Hess}_x \varphi \quad \text{locally uniformly on } (0, T) \times \mathbb{R}^d \text{ as } k \rightarrow \infty. \quad (5.7)$$

By (2.6), (2.8), and the analogous argument to obtain (5.4), for all $k \in \mathbb{N}$ with

$$|t_k - t_0| \leq [(T - t_0) \wedge t_0]/2 \quad \text{and} \quad \|x_k - x_0\| \leq 1$$

we have that

$$\begin{aligned} & |J\varphi_k(t_k, x_k) - J\varphi(t_k, x_k)| \\ &= |J(\varphi_k - \varphi)(t_k, x_k)| \\ &\leq \left(\sup_{|s-t_0| \leq \frac{(T-t_0) \wedge t_0}{2}} \sup_{\|y-x_0\| \leq 1+C_d^{1/2}} \left[\|\text{Hess}_x(\varphi_k - \varphi)(s, y)\|_F \right] \right) \cdot \int_{\mathbb{R}^d} dC_d(1 \wedge \|z\|^2) \nu(dz) \\ &\leq K d^{p+1} C_d \cdot \left(\sup_{|s-t_0| \leq \frac{(T-t_0) \wedge t_0}{2}} \sup_{\|y-x_0\| \leq 1+C_d^{1/2}} \left[\|\text{Hess}_x(\varphi_k - \varphi)(s, y)\|_F \right] \right). \end{aligned}$$

Hence, (5.7) implies that

$$\lim_{k \rightarrow \infty} |J\varphi_k(t_k, x_k) - J\varphi(t_k, x_k)| = 0. \quad (5.8)$$

Then, by (5.6), (5.8), and the triangle inequality we have

$$\begin{aligned} & \lim_{k \rightarrow \infty} |J\varphi(t_0, x_0) - J\varphi_k(t_k, x_k)| \\ & \leq \lim_{k \rightarrow \infty} |J\varphi(t_0, x_0) - J\varphi(t_k, x_k)| + \lim_{k \rightarrow \infty} |J\varphi_k(t_k, x_k) - J\varphi(t_k, x_k)| = 0. \end{aligned} \quad (5.9)$$

This together with (5.6) ensure that (F1) holds (with $F \curvearrowright G$ in the notation of [44]). To verify (F2), let $(t_0, x_0) \in (0, T) \times \mathbb{R}^d$, and assume that the functions $(u - v)(t_0, \cdot)$ and $(\varphi - \psi)(t_0, \cdot)$ have global maxima at x_0 . Then for all $z \in \mathbb{R}^d$ we have

$$(u - v)(t_0, x_0 + \eta_{t_0}(x_0, z)) \leq (u - v)(t_0, x_0), \quad (\varphi - \psi)(t_0, x_0 + \eta_{t_0}(x_0, z)) \leq (\varphi - \psi)(t_0, x_0),$$

and

$$\nabla_x(\varphi - \psi)(t_0, x_0) = 0.$$

Hence, by the definition of G_κ we have for all $r \in \mathbb{R}$, $y \in \mathbb{R}^d$, and $A, B \in \mathbb{S}^d$ with $A \leq B$ that

$$\begin{aligned} & G_\kappa(t_0, x_0, r, y, A, u(t_0, \cdot), \varphi(t_0, \cdot)) - G_\kappa(t_0, x_0, r, y, B, v(t_0, \cdot), \psi(t_0, \cdot)) \\ &= -\frac{1}{2} \text{Trace}(\sigma(t_0, x_0)[\sigma(t_0, x_0)]^T(A - B)) \\ &\quad - \int_{\|z\| \geq \kappa} [(u - v)(t_0, x_0 + \eta_{t_0}(x_0, z)) - (u - v)(t_0, x_0)] \nu(dz) \\ &\quad - \int_{\|z\| < \kappa} [(\varphi - \psi)(t_0, x_0 + \eta_{t_0}(x_0, z)) - (\varphi - \psi)(t_0, x_0) - \langle \nabla_x(\varphi - \psi)(t_0, x_0), \eta_{t_0}(x_0, z) \rangle] \nu(dz) \\ &\geq \frac{1}{2} \sum_{i,j=1}^d \sum_{k=1}^d \sigma^{ik}(t_0, x_0) \sigma^{jk}(t_0, x_0) (B - A)^{ij} \geq 0 \end{aligned}$$

Thus, (F2) is verified (with $F_\kappa \curvearrowright G_\kappa$ in the notation of [44]). Next, by (2.2) we obtain for all $(t, x) \in (0, T) \times \mathbb{R}^d$, and $r, s \in \mathbb{R}$ that

$$\|f(t, x, r) - f(t, x, s)\| \leq L^{1/2}|r - s|.$$

This implies that for all $(t, x) \in (0, T) \times \mathbb{R}^d$, $y \in \mathbb{R}^d$, $A \in \mathbb{S}^d$, and $r, s \in \mathbb{R}$ with $r \leq s$

$$G(t, x, r, y, A, \varphi(t, \cdot)) - G(t, x, s, y, A, \varphi(t, \cdot)) \geq L^{1/2}(r - s).$$

Thus, we verify (F3) (with $F \curvearrowright G$ and $\gamma \curvearrowright L^{1/2}$ in the notation of [44]). Next, by the definition of G_κ it is easy to see that (F4) holds (with $F_\kappa \curvearrowright G_\kappa$ in the notation of [44]). To see that (F5) is satisfied, fix

$t_0 \in (0, T)$, and let $\phi_k \in C_1^{1,2}((0, T) \times \mathbb{R}^d)$ for all $k \in \mathbb{N}$ such that $\lim_{k \rightarrow \infty} \phi_k(t_0, x) = w(t_0, x)$ for almost all $x \in \mathbb{R}^d$. Moreover, assume that there is a constant $\mathfrak{C} > 0$ such that

$$|\phi_k(t_0, x)| \leq \mathfrak{C}(1 + \|x\|) \quad \text{for all } x \in \mathbb{R}^d. \quad (5.10)$$

Note that Assumption 2.2 ensures $\|\eta_{t_0}(x, z)\|^2 \leq C_d$ for all $x, z \in \mathbb{R}^d$. This together with (5.10) and the assumption $w \in SC_1((0, T) \times \mathbb{R}^d)$ imply that for all $x, z \in \mathbb{R}^d$

$$|(w - \phi_k)(t_0, x + \eta_{t_0}(x, z)) - (w - \phi_k)(t_0, x)| \leq \sup_{y: \|y-x_0\| \leq C_d^{1/2}} \{2|w(t_0, y)|\} + 2\mathfrak{C}(1 + \|x\| + C_d^{1/2}). \quad (5.11)$$

Moreover, ν being a Lévy measure on \mathbb{R}^d ensures that $\nu(\{z \in \mathbb{R}^d : \|z\| \geq \kappa\}) < \infty$ for all $\kappa \in (0, 1)$. Hence, (5.11) allows us to apply Lebesgue's dominated convergence theorem to show that for all $(t, x) \in (0, T) \times \mathbb{R}^d$ and $\kappa \in (0, 1)$

$$\left| \lim_{k \rightarrow \infty} \int_{\|z\| \geq \kappa} [(w - \phi_k)(t_0, x + \eta_{t_0}(x, z)) - (w - \phi_k)(t_0, x)] \nu(dz) \right| = 0.$$

This implies that for all $x \in \mathbb{R}^d$, $r \in \mathbb{R}$, $y \in \mathbb{R}^d$, $A \in \mathbb{S}^d$, and $\kappa \in (0, 1)$

$$\lim_{k \rightarrow \infty} G_\kappa(t_0, x, r, y, A, \phi_k(t_0 \cdot), \varphi(t_0, \cdot)) = G_\kappa(t_0, x, r, y, A, w(t_0 \cdot), \varphi(t_0, \cdot)).$$

Therefore, we verify (F5) (with $F_\kappa \curvearrowright G_\kappa$ in the notation of [44]).

Next, to verify (F6) let $\alpha, \varepsilon, \beta \in (0, \infty)$, and define

$$\phi(t, x, y) := e^{\beta t} \frac{\alpha}{2} \|x - y\|^2 + e^{\beta t} \frac{\varepsilon}{2} (\|x\|^2 + \|y\|^2), \quad (t, x, y) \in (0, T) \times \mathbb{R}^d \times \mathbb{R}^d.$$

Let (x_0, y_0, t_0) be a global maximum of

$$(0, T) \times \mathbb{R}^d \times \mathbb{R}^d \ni (t, x, y) \mapsto u(t, x) - v(t, y) - \phi(t, x, y) \in \mathbb{R}. \quad (5.12)$$

Moreover, let $X, Y \in \mathbb{S}^d$ satisfy

$$\begin{pmatrix} X & 0 \\ 0 & -Y \end{pmatrix} \leq 2e^{\beta t_0} \alpha \begin{pmatrix} \mathbf{I}_d & -\mathbf{I}_d \\ -\mathbf{I}_d & \mathbf{I}_d \end{pmatrix} + e^{\beta t_0} \varepsilon \begin{pmatrix} \mathbf{I}_d & 0 \\ 0 & \mathbf{I}_d \end{pmatrix}, \quad (5.13)$$

where we recall that \mathbf{I}_d denotes the $d \times d$ identity matrix. By (2.6) we first notice for all $\kappa \in (0, 1)$ that

$$\begin{aligned} & \int_{\|z\| < \kappa} [\phi(t_0, x_0, y_0 + \eta_{t_0}(y_0, z)) - \phi(t_0, x_0, y_0) - \langle \nabla_y \phi(t_0, x_0, y_0), \eta_{t_0}(y_0, z) \rangle] \nu(dz) \\ & + \int_{\|z\| < \kappa} [\phi(t_0, x_0 + \eta_{t_0}(x_0, z), y_0) - \phi(t_0, x_0, y_0) - \langle \nabla_x \phi(t_0, x_0, y_0), \eta_{t_0}(x_0, z) \rangle] \nu(dz) \\ & = \int_{\|z\| < \kappa} \int_0^1 (1 - \vartheta) \sum_{i,j=1}^d \left(\frac{\partial^2}{\partial y_i \partial y_j} \phi \right) (t_0, x_0, y_0 + \vartheta \eta_{t_0}(y_0, z)) \eta_{t_0}^i(y_0, z) \eta_{t_0}^j(y_0, z) d\vartheta \nu(dz) \\ & + \int_{\|z\| < \kappa} \int_0^1 (1 - \vartheta) \sum_{i,j=1}^d \left(\frac{\partial^2}{\partial x_i \partial x_j} \phi \right) (t_0, x_0 + \vartheta \eta_{t_0}(x_0, z), y_0) \eta_{t_0}^i(x_0, z) \eta_{t_0}^j(x_0, z) d\vartheta \nu(dz) \\ & = \int_{\|z\| < \kappa} \int_0^1 (1 - \vartheta) \sum_{i=1}^d \left[e^{\beta t_0} (-\alpha + \varepsilon) |\eta_{t_0}^i(y_0, z)|^2 + e^{\beta t_0} (\alpha + \varepsilon) |\eta_{t_0}^i(x_0, z)|^2 \right] d\vartheta \nu(dz) \\ & \leq \int_{\|z\| < \kappa} e^{\beta T} (\alpha + 2\varepsilon) C_d (1 \wedge \|z\|^2) \nu(dz). \end{aligned} \quad (5.14)$$

By (2.8), we obtain

$$\int_{\mathbb{R}^d} e^{\beta T} (\alpha + 2\varepsilon) C_d (1 \wedge \|z\|^2) \nu(dz) < \infty.$$

This together with Lebesgue's dominated convergence theorem imply that

$$\lim_{\kappa \rightarrow 0} \int_{\|z\| < \kappa} e^{\beta T} (\alpha + 2\varepsilon) C_d (1 \wedge \|z\|^2) \nu(dz) = 0. \quad (5.15)$$

Furthermore, note that for all $t \in (0, T)$ and $x, y \in \mathbb{R}^d$ we have

$$\nabla_x \phi(t, x, y) = e^{\beta t} \alpha(x - y) + e^{\beta t} \varepsilon x \quad \text{and} \quad \nabla_y \phi(t, x, y) = -e^{\beta t} \alpha(x - y) + e^{\beta t} \varepsilon y.$$

Hence, it holds for all $\kappa \in (0, 1)$ that

$$\begin{aligned} E_\kappa &:= - \int_{\|z\| \geq \kappa} \left[v(t_0, y_0 + \eta_{t_0}(y_0, z)) - v(t_0, y_0) \right. \\ &\quad \left. - \langle e^{\beta t_0} \alpha(x_0 - y_0) - e^{\beta t_0} \varepsilon y_0, \eta_{t_0}(y_0, z) \rangle \right] \nu(dz) \\ &\quad + \int_{\|z\| \geq \kappa} \left[u(t_0, x_0 + \eta_{t_0}(x_0, z)) - u(t_0, x_0) \right. \\ &\quad \left. - \langle e^{\beta t_0} \alpha(x_0 - y_0) + e^{\beta t_0} \varepsilon x_0, \eta_{t_0}(x_0, z) \rangle \right] \nu(dz) \\ &= - \int_{\|z\| \geq \kappa} \left[v(t_0, y_0 + \eta_{t_0}(y_0, z)) - v(t_0, y_0) + \langle \nabla_y \phi(t_0, x_0, y_0), \eta_{t_0}(y_0, z) \rangle \right] \nu(dz) \\ &\quad + \int_{\|z\| \geq \kappa} \left[u(t_0, x_0 + \eta_{t_0}(x_0, z)) - u(t_0, x_0) - \langle \nabla_x \phi(t_0, x_0, y_0), \eta_{t_0}(x_0, z) \rangle \right] \nu(dz). \end{aligned} \quad (5.16)$$

Moreover, (t_0, x_0, y_0) being a global maximum of the function defined by (5.12) ensures that for all $z \in \mathbb{R}^d$

$$\begin{aligned} &u(t_0, x_0) - v(t_0, y_0) - \phi(t_0, x_0, y_0) \\ &\geq u(t_0, x_0 + \eta_{t_0}(x_0, z)) - v(t_0, y_0 + \eta_{t_0}(y_0, z)) - \phi(t_0, x_0 + \eta_{t_0}(x_0, z), y_0 + \eta_{t_0}(y_0, z)). \end{aligned} \quad (5.17)$$

Then by (2.3), (2.4), (5.16), (5.17), and Taylor's formula we have for every $\kappa \in (0, 1)$ that

$$\begin{aligned} E_\kappa &\leq \int_{\|z\| \geq \kappa} \left[\phi(t_0, x_0 + \eta_{t_0}(x_0, z), y_0 + \eta_{t_0}(y_0, z)) - \phi(t_0, x_0, y_0) \right. \\ &\quad \left. - \langle \nabla_y \phi(t_0, x_0, y_0), \eta_{t_0}(y_0, z) \rangle - \langle \nabla_x \phi(t_0, x_0, y_0), \eta_{t_0}(x_0, z) \rangle \right] \nu(dz) \\ &= \int_{\|z\| \geq \kappa} \int_0^1 (1 - \vartheta) \left[\sum_{i=1}^d (e^{\beta t_0} \alpha + e^{\beta t_0} \varepsilon) |\eta_{t_0}^i(x_0, z)|^2 + \sum_{i=1}^d (e^{\beta t_0} \alpha + e^{\beta t_0} \varepsilon) |\eta_{t_0}^i(y_0, z)|^2 \right. \\ &\quad \left. - 2 \sum_{i=1}^d e^{\beta t_0} \alpha \eta_{t_0}^i(x_0, z) \eta_{t_0}^i(y_0, z) \right] d\vartheta \nu(dz) \\ &\leq \int_{\|z\| \geq \kappa} \left[e^{\beta t_0} \varepsilon (\|\eta_{t_0}(x_0, z)\|^2 + \|\eta_{t_0}(y_0, z)\|^2) + \sum_{i=1}^d e^{\beta t_0} \alpha |\eta_{t_0}^i(x_0, z) - \eta_{t_0}^i(y_0, z)|^2 \right] \nu(dz) \\ &\leq 2e^{\beta t_0} \varepsilon L d^p (1 + \|x_0\|^2 + \|y_0\|^2) + e^{\beta t_0} \alpha L \|x_0 - y_0\|^2. \end{aligned} \quad (5.19)$$

Furthermore, by (2.3), (2.4), and the Cauchy-Schwarz inequality it holds that

$$\begin{aligned} &-\langle \mu(t_0, y_0), e^{\beta t_0} \alpha(x_0 - y_0) - e^{\beta t_0} \varepsilon y_0 \rangle + \langle \mu(t_0, x_0), e^{\beta t_0} \alpha(x_0 - y_0) - e^{\beta t_0} \varepsilon x_0 \rangle \\ &\leq e^{\beta t_0} \alpha \|x_0 - y_0\| \cdot \|\mu(t_0, x_0) - \mu(t_0, y_0)\| + e^{\beta t_0} \varepsilon \|y_0\| \cdot \|\mu(t_0, y_0)\| + e^{\beta t_0} \varepsilon \|x_0\| \cdot \|\mu(t_0, x_0)\| \\ &\leq L^{1/2} e^{\beta t_0} \alpha \|x_0 - y_0\|^2 + 4L^{1/2} d^{p/2} e^{\beta t_0} \varepsilon (1 + \|x_0\|^2 + \|y_0\|^2). \end{aligned} \quad (5.20)$$

Next, we use a trick by Ishii (see, e.g., Example 3.6 in [20]) to get an estimate for

$$-\frac{1}{2} \text{Trace} \left(\Sigma_{y_0} \Sigma_{y_0}^T Y \right) + \frac{1}{2} \text{Trace} \left(\Sigma_{x_0} \Sigma_{x_0}^T X \right),$$

where

$$\Sigma_{x_0} := \sigma(t_0, x_0) \quad \text{and} \quad \Sigma_{y_0} := \sigma(t_0, y_0).$$

More precisely, left multiplying both sides of (5.13) by the symmetric matrix

$$\begin{pmatrix} \Sigma_{x_0} \Sigma_{x_0}^T & \Sigma_{y_0} \Sigma_{x_0}^T \\ \Sigma_{x_0} \Sigma_{y_0}^T & \Sigma_{y_0} \Sigma_{y_0}^T \end{pmatrix}$$

yields

$$\begin{aligned} \begin{pmatrix} \Sigma_{x_0} \Sigma_{x_0}^T X & -\Sigma_{y_0} \Sigma_{x_0}^T Y \\ \Sigma_{x_0} \Sigma_{y_0}^T X & -\Sigma_{y_0} \Sigma_{y_0}^T Y \end{pmatrix} &\leq 2e^{\beta t_0} \alpha \begin{pmatrix} \Sigma_{x_0} \Sigma_{x_0}^T - \Sigma_{y_0} \Sigma_{x_0}^T & -\Sigma_{x_0} \Sigma_{x_0}^T + \Sigma_{y_0} \Sigma_{x_0}^T \\ \Sigma_{x_0} \Sigma_{y_0}^T - \Sigma_{y_0} \Sigma_{y_0}^T & -\Sigma_{x_0} \Sigma_{y_0}^T + \Sigma_{y_0} \Sigma_{y_0}^T \end{pmatrix} \\ &+ e^{\beta t_0} \varepsilon \begin{pmatrix} \Sigma_{x_0} \Sigma_{x_0}^T & \Sigma_{y_0} \Sigma_{x_0}^T \\ \Sigma_{x_0} \Sigma_{y_0}^T & \Sigma_{y_0} \Sigma_{y_0}^T \end{pmatrix}. \end{aligned} \quad (5.21)$$

By Assumption 2.1, taking traces to (5.21) yields

$$\begin{aligned} &-\frac{1}{2} \text{Trace} \left(\Sigma_{y_0} \Sigma_{y_0}^T Y \right) + \frac{1}{2} \text{Trace} \left(\Sigma_{x_0} \Sigma_{x_0}^T X \right) \\ &\leq e^{\beta t_0} \alpha \text{Trace} \left\{ \left(\Sigma_{x_0} - \Sigma_{y_0} \right) \left(\Sigma_{x_0} - \Sigma_{y_0} \right)^T \right\} + \frac{1}{2} e^{\beta t_0} \varepsilon \text{Trace} \left\{ \Sigma_{x_0} \Sigma_{x_0}^T + \Sigma_{y_0} \Sigma_{y_0}^T \right\} \\ &= e^{\beta t_0} \alpha \left\| \Sigma_{x_0} - \Sigma_{y_0} \right\|_F^2 + \frac{1}{2} e^{\beta t_0} \varepsilon \left(\left\| \Sigma_{x_0} \right\|_F^2 + \left\| \Sigma_{y_0} \right\|_F^2 \right) \\ &\leq L e^{\beta t_0} \alpha \|x_0 - y_0\|^2 + L d^p e^{\beta t_0} \varepsilon (1 + \|x_0\|^2 + \|y_0\|^2). \end{aligned} \quad (5.22)$$

Moreover, by (2.2) it holds for all $r \in \mathbb{R}$ that

$$|f(t_0, x_0, r) - f(t_0, y_0, r)| \leq L^{1/2} T^{-3/2} \|x_0 - y_0\|. \quad (5.23)$$

Then using (5.14), (5.19), (5.20), (5.22), and (5.23) we verify condition (F6) on page 287 of [44] (with $m \curvearrowright 1$, $\alpha \curvearrowright \alpha$, $\varepsilon \curvearrowright \varepsilon$, $\lambda \curvearrowright \beta$, $p \curvearrowright 2$, $\bar{F}_\kappa \curvearrowright G_\kappa$, $F_\kappa \curvearrowright G_\kappa$, $\eta_1 = \eta_2 = \eta_3 = \eta_4 = 0$, $p_1 = p_2 = p_3 = p_4 = p_s = 0$, $K_1 = L^{1/2} T^{-3/2}$, $K_2 = e^{\beta t_0} L^{1/2} (1 + 2L^{1/2})$, $K_3 = L^{1/2} (3L^{1/2} d^p + 4d^{p/2})$, and $m_{\alpha, \varepsilon}$ being the integral term in (5.15) in the notation of condition (F6) in [44], noting that $m_{\alpha, \varepsilon} \equiv m_{\alpha, \varepsilon, \lambda}$ is allowed to depend on λ in the notation of condition (F6) in [44]; see p. 291-292 in [44]). Hence, by Remark 2.8 and Theorem 3.1 in [44] we obtain $u_1(t, x) = u_2(t, x)$ for all $(t, x) \in [0, T] \times \mathbb{R}^d$. \square

To show the existence of viscosity solutions of (2.18), we first consider the linear PIDE

$$-\frac{\partial}{\partial t} u(t, x) + G_0(t, x, u(t, x), \nabla_x u(t, x), \text{Hess}_x u(t, x), u(t, \cdot)) = 0 \quad \text{on } (0, T) \times \mathbb{R}^d, \quad (5.24)$$

$$u(T, x) = g(x) \quad \text{on } \mathbb{R}^d. \quad (5.25)$$

Lemma 5.5. *Let Assumptions 2.1, 2.2, 2.3, and 5.1 hold. Moreover, for every $t \in [0, T]$ and $z \in \mathbb{R}^d$ we assume that $\mu(t, \cdot)$, $\sigma(t, \cdot)$, $\eta_t(\cdot, z)$, $g(\cdot)$, and $h(t, \cdot)$ are compactly supported and infinitely differentiable. Then PIDE (5.24) has a unique classical solution $u \in C^{1,2}([0, T] \times \mathbb{R}^d)$ with $u(T, x) = g(x)$ for all $x \in \mathbb{R}^d$. Moreover, we have for all $(t, x) \in [0, T] \times \mathbb{R}^d$ that*

$$u(t, x) = \mathbb{E} \left[g(X_T^{t,x}) + \int_t^T h(s, X_s^{t,x}) ds \right]. \quad (5.26)$$

To prove the above lemma, we recall some notions and introduce some notations. For each $n \in \mathbb{N}$, a finite list $\alpha = \alpha_1 \alpha_2, \dots, \alpha_n$ of numbers $\alpha_i \in \{1, 2, \dots, d\}$ is called a multi-number of length $|\alpha| := n$, and we use the notation

$$D_\alpha := D_{\alpha_1} D_{\alpha_2} \dots D_{\alpha_n},$$

where for $i \in \{1, \dots, d\}$ and locally integrable functions $v : \mathbb{R}^d \rightarrow \mathbb{R}$, $D_i v$ refers to the weak derivative (see, e.g., 1.62 in [1] for the definition) of v with respect x_i . We also use the multi-number ϵ of length 0, and agree that D_ϵ means the identity operator. For a separable Banach space V we use the notation $L_2([0, T], V)$ for the space of Borel functions $v : [0, T] \rightarrow V$ such that

$$\int_0^T \|v(t)\|_V^2 dt < \infty.$$

We also denote by $L_2(\mathbb{R}^d)$ the space Borel functions $\varphi : \mathbb{R}^d \rightarrow \mathbb{R}$ such that

$$\|\varphi\|_{L_2(\mathbb{R}^d)}^2 := \int_{\mathbb{R}^d} |\varphi(x)|^2 dx < \infty.$$

For integers $m \in \mathbb{N}_0$, the notation W_2^m refers to the Sobolev space defined as the completion of $C_c^\infty(\mathbb{R}^d)$ in the norm

$$\|\varphi\|_{W_2^m(\mathbb{R}^d)} := \sum_{0 \leq |\alpha| \leq m} \|D_\alpha \varphi\|_{L_2(\mathbb{R}^d)}.$$

Moreover, for $\delta \in (0, 1)$ we use the notation $C_\delta^2(\mathbb{R}^d)$ to denote the Hölder space defined as the set of all functions $v : \mathbb{R}^d \rightarrow \mathbb{R}$ with continuous derivatives up to order 2 such that

$$\|v\|_{C_\delta^2} := \max_{0 \leq |\alpha| \leq 2} \sup_{x \in \mathbb{R}^d} |D_\alpha v(x)| + \max_{0 \leq |\alpha| \leq 2} \sup_{\substack{x, y \in \mathbb{R}^d \\ x \neq y}} \frac{|D_\alpha v(x) - D_\alpha v(y)|}{|x - y|^\delta} < \infty.$$

Proof of Lemma 5.5. We first introduce the notion of generalised solutions (cf., Definition 2.1 in [21]) of the linear PIDE

$$\frac{\partial}{\partial t} v(t, x) + G_0(t, x, v(t, x), \nabla_x v(t, x), \text{Hess}_x v(t, x), v(t, \cdot)) = 0 \quad \text{on } (0, T) \times \mathbb{R}^d, \quad (5.27)$$

$$v(0, x) = g(x) \quad \text{on } \mathbb{R}^d. \quad (5.28)$$

Note that by virtue of Remark 2.1 in [21], integrating by parts ensures for all $\varphi \in C_c^\infty(\mathbb{R}^d)$, $v \in W_2^1(\mathbb{R}^d)$, and $t \in [0, T]$ that

$$\begin{aligned} & \int_{\mathbb{R}^d} \varphi(x) \int_{\mathbb{R}^d} [v(x + \eta_t(x, z)) - v(t, x) - \langle \nabla_x v(x), \eta_t(x, z) \rangle] \nu(dz) dx \\ &= - \sum_{k=1}^d \int_{\mathbb{R}^d} [\mathcal{J}^k(t)v(x)] \cdot D_k \varphi(x) dx + \int_{\mathbb{R}^d} [\mathcal{J}^0(t)v(x)] \cdot \varphi(x) dx, \end{aligned} \quad (5.29)$$

where, by denoting $\eta_t(x, z) = (\eta_t^1(x, z), \dots, \eta_t^d(x, z))$

$$\mathcal{J}^k(t)v(x) := \int_0^1 \int_{\mathbb{R}^d} \eta_t^k(x, z) (v(x + \vartheta \eta_t(x, z)) - v(x)) \nu(dz) d\vartheta, \quad (t, x) \in [0, T] \times \mathbb{R}^d, \quad k = 1, \dots, d,$$

and

$$\begin{aligned} \mathcal{J}^0(t)v(x) := & - \int_0^1 \int_{\mathbb{R}^d} \left[\sum_{k=1}^d D_k \eta_t^k(x, z) (v(x + \vartheta \eta_t(x, z)) - v(x)) \right. \\ & \left. + \sum_{k, l=1}^d \vartheta \eta_t^k(x, z) (D_k \eta_t^l(x, z)) (D_l v)(x + \vartheta \eta_t(x, z)) \right] \nu(dz) d\vartheta, \quad (t, x) \in [0, T] \times \mathbb{R}^d. \end{aligned}$$

Define $a = (a^{i,j})_{i,j \in \{1, \dots, d\}} \in C([0, T] \times \mathbb{R}^d, \mathbb{S}^d)$ by

$$a(t, x) := \sigma(t, x) [\sigma(t, x)]^T, \quad (t, x) \in [0, T] \times \mathbb{R}^d.$$

Then an $L_2(\mathbb{R}^d)$ -valued continuous function $v = (v(t))_{t \in [0, T]}$, is called a generalised solution (see Definition 2.1 in [21]) to equation (5.27) with initial condition $v(0) = g$, if $v(t) \in W_2^1(\mathbb{R}^d)$ for dt -almost every $t \in [0, T]$, $v \in L_2([0, T], W_2^1(\mathbb{R}^d))$, and for every $\varphi \in C_c^\infty(\mathbb{R}^d)$ and $t \in [0, T]$

$$\begin{aligned} \int_{\mathbb{R}^d} v(t, x) \varphi(x) dx = & \int_{\mathbb{R}^d} g(x) \varphi(x) dx + \int_0^t \int_{\mathbb{R}^d} (h(s, x) \varphi(x) + [\mathcal{J}^0(s)v(s, x)] \cdot \varphi(x)) dx ds \\ & + \int_0^t \int_{\mathbb{R}^d} \left(\sum_{i,j=1}^d \left[- (a^{ij}(s, x) D_j v(s, x) + \mathcal{J}^i(s)v(s, x)) D_i \varphi(x) \right. \right. \\ & \left. \left. + [\mu^i(s, x) - D_j a^{ij}(s, x)] [D_i v(s, x)] \cdot \varphi(x) \right] \right) dx ds. \end{aligned} \quad (5.30)$$

By the assumptions of Lemma 5.5, for each $m \in \mathbb{N}$ Assumptions 2.1, 2.2, and 2.5 in [21] are satisfied (with $p = 2$, $m \curvearrowright m$, $a \curvearrowright a$, $b \curvearrowright \mu$, $\eta \curvearrowright \eta$, $\psi \curvearrowright g$, and $f \curvearrowright h$ in the notation of Assumptions 2.1, 2.2, and 2.5 in [21]). Thus, Theorem 2.1 in [21] ensures that PIDE (5.27) has a unique generalized solution $v = (v(t))_{t \in [0, T]}$ with $v(0) = g$, and for every $m \in \mathbb{N}_0$ the generalised solution v is strongly continuous as a $W_2^m(\mathbb{R}^d)$ -valued function of $t \in [0, T]$. Thus, by Sobolev embedding theorem (see, e.g., 4.12 Part II

in [1]) we have for each $\delta \in (0, 1)$ the generalised solution v is strongly continuous as a $C_\delta^2(\mathbb{R}^d)$ -valued function of $t \in [0, T]$. Hence, for every $(t_0, x_0) \in [0, T] \times \mathbb{R}^d$ and multi-number α with $|\alpha| \leq 2$ we have

$$\lim_{t \rightarrow t_0} \sup_{x \in \mathbb{R}^d} |D_\alpha v(t_0, x) - D_\alpha v(t, x)| = 0. \quad (5.31)$$

Moreover, the fact that $v(t, \cdot) \in C_\delta^2(\mathbb{R}^d)$ for all $t \in [0, T]$ and $\delta \in (0, 1)$ ensures that for every $(t_0, x_0) \in [0, T] \times \mathbb{R}^d$ and multi-number α with $|\alpha| \leq 2$

$$\lim_{x \rightarrow x_0} |D_\alpha v(t_0, x) - D_\alpha v(t_0, x_0)| = 0.$$

This together with (5.31) imply that for every $(t_0, x_0) \in [0, T] \times \mathbb{R}^d$ and multi-number α with $|\alpha| \leq 2$

$$\begin{aligned} \lim_{(t,x) \rightarrow (t_0,x_0)} |D_\alpha v(t, x) - D_\alpha v(t_0, x_0)| &\leq \lim_{(t,x) \rightarrow (t_0,x_0)} |D_\alpha v(t, x) - D_\alpha v(t_0, x)| \\ &\quad + \lim_{x \rightarrow x_0} |D_\alpha v(t_0, x) - D_\alpha v(t_0, x_0)| \\ &= 0. \end{aligned}$$

Thus, as a real-valued function on $[0, T] \times \mathbb{R}^d$, $v \in C^{1,2}([0, T] \times \mathbb{R}^d)$. Furthermore, (5.30), integrating by parts, and the fact that $v \in C^{1,2}([0, T] \times \mathbb{R}^d)$ ensures that the unique generalised solution v is the unique classical solution of (5.27) with $v(0, x) = g(x)$ for $x \in \mathbb{R}^d$. Then by time reversing, the function $u(t, x) := v(T - t, x)$, $(t, x) \in [0, T] \times \mathbb{R}^d$, is the unique classical solution of (5.24) with $u(T, x) = g(x)$ for $x \in \mathbb{R}^d$. Moreover, by Feynman-Kac formula (see, e.g., Theorem 17.4.10 in [16], or Theorem 285 in [55]) we verify (5.26). The proof of this lemma is therefore completed. \square

In the rest of this section, for every $n \in \mathbb{N}$ let $g^{(n)} \in C(\mathbb{R}^d, \mathbb{R})$, $h^{(n)} \in C([0, T] \times \mathbb{R}^d, \mathbb{R})$, $\mu^{(n)} \in C([0, T] \times \mathbb{R}^d, \mathbb{R}^d)$, $\sigma^{(n)} \in C([0, T] \times \mathbb{R}^d, \mathbb{R}^{d \times d})$, and $\eta^{(n)} = (\eta^{(n),1}, \dots, \eta^{(n),d}) \in C([0, T] \times \mathbb{R}^d \times \mathbb{R}^d, \mathbb{R}^d)$. Then we make the following assumptions.

Assumption 5.6. *It holds for all non-empty compact sets $\mathcal{K} \subseteq [0, T] \times \mathbb{R}^d$ that*

$$\begin{aligned} \lim_{n \rightarrow \infty} \left[\sup_{(t,x) \in \mathcal{K}} \left(|g^{(n)}(x) - g(x)| + |h^{(n)}(t, x) - h(t, x)| + \|\mu^{(n)}(t, x) - \mu(t, x)\| \right. \right. \\ \left. \left. + \|\sigma^{(n)}(t, x) - \sigma(t, x)\|_F \right) \right] = 0, \end{aligned} \quad (5.32)$$

and

$$\lim_{n \rightarrow \infty} \left[\sup_{(t,x) \in \mathcal{K}} \|\eta_t^{(n)}(x, z) - \eta_t(x, z)\| \right] = 0 \quad \nu(dz)\text{-almost everywhere.} \quad (5.33)$$

Assumption 5.7. *There exists a constant $C_1 > 0$ satisfying for all $n \in \mathbb{N}$, $x, y \in \mathbb{R}^d$, and $t \in [0, T]$ that*

$$|g^{(n)}(x) - g^{(n)}(y)|^2 + \|\mu^{(n)}(t, x) - \mu^{(n)}(t, y)\|^2 + \|\sigma^{(n)}(t, x) - \sigma^{(n)}(t, y)\|_F^2 \leq C_1 \|x - y\|^2, \quad (5.34)$$

and

$$|h^{(n)}(t, x)|^2 + |g^{(n)}(x)|^2 + \|\mu^{(n)}(t, x)\|^2 + \|\sigma^{(n)}(t, x)\|_F^2 \leq C_1 (1 + \|x\|^2). \quad (5.35)$$

Assumption 5.8. *There exists a constant $C_2 > 0$ such that for all $n \in \mathbb{N}$, $x, y \in \mathbb{R}^d$, $t \in [0, T]$, and $z \in \mathbb{R}^d$*

$$\|\eta_t^{(n)}(x, z)\|^2 \leq C_2 (1 \wedge \|z\|^2), \quad \|\eta_t^{(n)}(x, z) - \eta_t^{(n)}(y, z)\|^2 \leq C_2 \|x - y\|^2 (1 \wedge \|z\|^2). \quad (5.36)$$

Assumption 5.9. *There exists a constant $\delta > 0$ satisfying for all $n \in \mathbb{N}$, $(t, x) \in [0, T] \times \mathbb{R}^d$, $z \in \mathbb{R}^d$, and $\gamma \in [0, 1]$ that $D_x \eta_t^{(n)}(x, z)$ exists, and*

$$\delta \leq |\det(\mathbf{I}_d + \gamma D_x \eta_t^{(n)}(x, z))|. \quad (5.37)$$

Then for each $n \in \mathbb{N}$, let $G_0^{(n)} : (0, T) \times \mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d \times \mathbb{S}^d \times C^2(\mathbb{R}^d) \rightarrow \mathbb{R}$ be a functional defined as G_0 with $\mu^{(n)}$, $\sigma^{(n)}$, $\eta^{(n)}$, and $h^{(n)}$ in place of μ , σ , η , and h , respectively.

Lemma 5.10. *Let Assumptions 2.1, 2.2, and 5.1-5.8 hold. Then for every $\varphi \in C^{1,2}((0, T) \times \mathbb{R}^d)$ and every compact set $\mathcal{K} \subseteq (0, T) \times \mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d \times \mathbb{S}^d$ it holds that*

$$\lim_{n \rightarrow \infty} \left(\sup_{(t,x,r,y,A) \in \mathcal{K}} \left| G_0(t, x, r, y, A, \varphi(t, \cdot)) - G_0^{(n)}(t, x, r, y, A, \varphi(t, \cdot)) \right| \right) = 0. \quad (5.38)$$

Proof. By Assumption 5.6 and the definitions of G_0 and $G_0^{(n)}$, it is sufficient to show for every $\varphi \in C^{1,2}((0, T) \times \mathbb{R}^d)$ and compact set $\mathcal{K}_0 \subseteq (0, T) \times \mathbb{R}^d$ that

$$\lim_{n \rightarrow \infty} \left(\sup_{(t,x) \in \mathcal{K}_0} |\mathbb{J}^n \varphi(t, x)| \right) = 0, \quad (5.39)$$

where for all $(t, x) \in (0, T) \times \mathbb{R}^d$, $n \in \mathbb{N}$, and $\varphi \in C^{1,2}((0, T) \times \mathbb{R}^d)$,

$$\begin{aligned} \mathbb{J}^n \varphi(t, x) &:= \int_{\mathbb{R}^d} \left(\varphi(t, x + \eta_t^{(n)}(x, z)) - \varphi(t, x) - \langle \nabla_x \varphi(t, x), \eta_t^{(n)}(x, z) \rangle \right) \nu(dz) \\ &\quad - \int_{\mathbb{R}^d} \left(\varphi(t, x + \eta_t(x, z)) - \varphi(t, x) - \langle \nabla_x \varphi(t, x), \eta_t(x, z) \rangle \right) \nu(dz). \end{aligned}$$

To see this, note that by Taylor's formula and the triangle inequality we have for every $\varphi(t, x) \in C^{1,2}((0, T) \times \mathbb{R}^d)$, $n \in \mathbb{N}$, and $(t, x) \in (0, T) \times \mathbb{R}^d$ that

$$\begin{aligned} |\mathbb{J}^n \varphi(t, x)| &= \left| \int_{\mathbb{R}^d} \int_0^1 (1 - \alpha) \sum_{i,j=1}^d \left(\frac{\partial^2}{\partial x_i \partial x_j} \varphi(t, x + \alpha \eta_t^{(n)}(x, z)) \cdot \eta_t^{(n),i}(x, z) \eta_t^{(n),j}(x, z) \right. \right. \\ &\quad \left. \left. - \frac{\partial^2}{\partial x_i \partial x_j} \varphi(t, x + \alpha \eta_t(x, z)) \cdot \eta_t^i(x, z) \eta_t^j(x, z) \right) d\alpha \nu(dz) \right| \\ &\leq \int_0^1 \int_{\mathbb{R}^d} (1 - \alpha) \sum_{i,j=1}^d \left(\left| \frac{\partial^2}{\partial x_i \partial x_j} \varphi(t, x + \alpha \eta_t^{(n)}(x, z)) - \frac{\partial^2}{\partial x_i \partial x_j} \varphi(t, x + \alpha \eta_t(x, z)) \right| \right. \\ &\quad \left. \cdot |\eta_t^{(n),i}(x, z) \eta_t^{(n),j}(x, z)| \right) \nu(dz) d\alpha \\ &\quad + \int_0^1 \int_{\mathbb{R}^d} (1 - \alpha) \sum_{i,j=1}^d \left(\left| \eta_t^i(x, z) \eta_t^j(x, z) - \eta_t^{(n),i}(x, z) \eta_t^{(n),j}(x, z) \right| \right. \\ &\quad \left. \cdot \left| \frac{\partial^2}{\partial x_i \partial x_j} \varphi(t, x + \alpha \eta_t(x, z)) \right| \right) \nu(dz) d\alpha. \end{aligned} \quad (5.40)$$

Moreover, by Assumptions 2.2 and 5.8, the triangle inequality, and Cauchy Schwarz inequality we obtain for every $\varphi \in C^{1,2}((0, T) \times \mathbb{R}^d)$, $n \in \mathbb{N}$, $\alpha \in [0, 1]$, $z \in \mathbb{R}^d$, and $(t, x) \in (0, T) \times \mathbb{R}^d$ that

$$\begin{aligned} &\sum_{i,j=1}^d \left(\left| \frac{\partial^2}{\partial x_i \partial x_j} \varphi(t, x + \alpha \eta_t^{(n)}(x, z)) - \frac{\partial^2}{\partial x_i \partial x_j} \varphi(t, x + \alpha \eta_t(x, z)) \right| \cdot |\eta_t^{(n),i}(x, z) \eta_t^{(n),j}(x, z)| \right) \\ &\leq 2 \cdot \sup_{y: \|y-x\| \leq (C_d \vee C_2)^{1/2}} \left[\|\text{Hess}_x \varphi(t, y)\|_F \right] \cdot \sum_{i,j=1}^d |\eta_t^{(n),i}(x, z) \eta_t^{(n),j}(x, z)| \\ &\leq 2 \cdot \sup_{y: \|y-x\| \leq (C_d \vee C_2)^{1/2}} \left[\|\text{Hess}_x \varphi(t, y)\|_F \right] \cdot d \|\eta_t^{(n)}(x, z)\|^2 \\ &\leq 2 \cdot \sup_{y: \|y-x\| \leq (C_d \vee C_2)^{1/2}} \left[\|\text{Hess}_x \varphi(t, y)\|_F \right] \cdot d C_2 (1 \wedge \|z\|^2), \end{aligned}$$

and

$$\sum_{i,j=1}^d \left(\left| \eta_t^i(x, z) \eta_t^j(x, z) - \eta_t^{(n),i}(x, z) \eta_t^{(n),j}(x, z) \right| \cdot \left| \frac{\partial^2}{\partial x_i \partial x_j} \varphi(t, x + \alpha \eta_t(x, z)) \right| \right)$$

$$\begin{aligned}
&\leq \sup_{y: \|y-x\| \leq C_d^{1/2}} \left[\|\text{Hess}_x \varphi(t, y)\|_F \right] \cdot \sum_{i,j=1}^d \left[\left| \eta_t^i(x, z) \eta_t^j(x, z) \right| + \left| \eta_t^{(n),i}(x, z) \eta_t^{(n),j}(x, z) \right| \right] \\
&\leq \sup_{y: \|y-x\| \leq C_d^{1/2}} \left[\|\text{Hess}_x \varphi(t, y)\|_F \right] \cdot d(C_2 + C_d)(1 \wedge \|z\|^2),
\end{aligned}$$

which is integrable on \mathbb{R}^d with respect to $\nu(dz)$. Therefore, by (5.33), (5.40), and Lebesgue's dominated convergence theorem we get (5.39), which completes the proof of this lemma. \square

Lemma 5.11. *Let Assumptions 2.1, 2.2, and 5.1-5.8 hold. For every $n \in \mathbb{N}$, let $u \in C_1([0, T] \times \mathbb{R}^d)$, $u^{(n)} \in C_1([0, T] \times \mathbb{R}^d)$. Assume for all non-empty compact sets $\mathcal{K} \subseteq [0, T] \times \mathbb{R}^d$ that*

$$\lim_{n \rightarrow \infty} \left[\sup_{(t,x) \in \mathcal{K}} |u^{(n)}(t, x) - u(t, x)| \right] = 0. \quad (5.41)$$

Moreover, assume for all $n \in \mathbb{N}$ that $u^{(n)}$ is a viscosity solution of

$$-\frac{\partial}{\partial t} u^{(n)}(t, x) + G_0^{(n)}(t, x, u^{(n)}(t, x), \nabla_x u^{(n)}(t, x), \text{Hess}_x u^{(n)}(t, x), u^{(n)}(t, \cdot)) = 0 \text{ on } (0, T) \times \mathbb{R}^d \quad (5.42)$$

with $u^{(n)}(T, x) = g^{(n)}(x)$ for $x \in \mathbb{R}^d$. Then u is a viscosity solution of the linear PIDE (5.24) with $u(T, x) = g(x)$ for $x \in \mathbb{R}^d$.

Proof. This proof is analogous to the proof of Lemma 2.18 in [12]. We only prove the viscosity subsolution case, and the viscosity supersolution case can be proved analogously. First note that (5.32), (5.41), and the condition that $u^{(n)}(T, x) = g^{(n)}(x)$ for all $x \in \mathbb{R}^d$ and $n \in \mathbb{N}$ guarantees that $u(T, x) = g(x)$ for $x \in \mathbb{R}^d$. Throughout this proof, let $(t_0, x_0) \in (0, T) \times \mathbb{R}^d$, and let $\varphi \in C^{1,2}((0, T) \times \mathbb{R}^d)$ such that $\varphi(t_0, x_0) = u(t_0, x_0)$ and $\varphi(t, x) \geq u(t, x)$ for all $(t, x) \in [0, T] \times \mathbb{R}^d$. For every $\varepsilon \in (0, \infty)$ define $\varphi_\varepsilon \in C^{1,2}((0, T) \times \mathbb{R}^d)$ by

$$\varphi_\varepsilon(t, x) := \varphi(t, x) + \frac{\varepsilon}{2}(|t - t_0|^2 + \|x - x_0\|^2), \quad (t, x) \in (0, T) \times \mathbb{R}^d. \quad (5.43)$$

Moreover, let $\xi := \frac{(T-t_0) \wedge t_0}{2}$, and define the set

$$\mathcal{K}_0 := \{(t, x) \in [0, T] \times \mathbb{R}^d : \max\{|t - t_0|, \|x - x_0\|\} \leq \xi\}.$$

According to (5.41), we notice for each $\varepsilon \in (0, \infty)$ that there exists an integer $N_\varepsilon > 0$ such that for all integers $n \geq N_\varepsilon$

$$\sup_{(t,x) \in \mathcal{K}_0} \left(|u^{(n)}(t, x) - u(t, x)| \right) < \frac{\varepsilon \xi^2}{4}.$$

Thus, by (5.43) we obtain for all $\varepsilon \in (0, \infty)$, $n \in \mathbb{N} \cap [N_\varepsilon, \infty)$, and $(t, x) \in \mathcal{K}_0$ with $|t - t_0|^2 + \|x - x_0\|^2 \geq \xi^2$ that

$$\begin{aligned}
u^{(n)}(t_0, x_0) - \varphi_\varepsilon(t_0, x_0) &= u^{(n)}(t_0, x_0) - \varphi(t_0, x_0) = u^{(n)}(t_0, x_0) - u(t_0, x_0) > -\frac{\varepsilon \xi^2}{4} \\
&> u^{(n)}(t, x) - u(t, x) - \frac{\varepsilon}{2}(|t - t_0|^2 + \|x - x_0\|^2) \\
&\geq u^{(n)}(t, x) - \varphi(t, x) - \frac{\varepsilon}{2}(|t - t_0|^2 + \|x - x_0\|^2) \\
&= u^{(n)}(t, x) - \varphi_\varepsilon(t, x).
\end{aligned}$$

Since for every $\varepsilon \in (0, \infty)$ and $n \in \mathbb{N}$ the function $u^{(n)} - \varphi_\varepsilon$ is continuous, we obtain that for each $\varepsilon \in (0, \infty)$ there exist sequences $\{t_n^{(\varepsilon)}\}_{n=1}^\infty \subseteq \{t \in [0, T] : |t - t_0| \leq \xi\}$ and $\{x_n^{(\varepsilon)}\}_{n=1}^\infty \subseteq \{x \in \mathbb{R}^d : \|x - x_0\| \leq \xi\}$ satisfying for all $n \in \mathbb{N} \cap [N_\varepsilon, \infty)$ and $(t, x) \in \mathcal{K}_0$ that

$$u^{(n)}(t_n^{(\varepsilon)}, x_n^{(\varepsilon)}) - \varphi_\varepsilon(t_n^{(\varepsilon)}, x_n^{(\varepsilon)}) \geq u^{(n)}(t, x) - \varphi_\varepsilon(t, x). \quad (5.44)$$

Thus, taking into account that for each $n \in \mathbb{N}$ and $\varepsilon \in (0, \infty)$, $\varphi_\varepsilon \in C^{1,2}((0, T) \times \mathbb{R}^d)$, and that $u^{(n)}$ is a viscosity subsolution of (5.42), by Lemma 5.2 we have for all $\varepsilon \in (0, \infty)$ and $n \in \mathbb{N} \cap [N_\varepsilon, \infty)$ that

$$-\frac{\partial}{\partial t} \varphi_\varepsilon(t_n^{(\varepsilon)}, x_n^{(\varepsilon)})$$

$$+ G_0^{(n)}(t_n^{(\varepsilon)}, x_n^{(\varepsilon)}, u^{(n)}(t_n^{(\varepsilon)}, x_n^{(\varepsilon)}), \nabla_x \varphi_\varepsilon(t_n^{(\varepsilon)}, x_n^{(\varepsilon)}), \text{Hess}_x \varphi_\varepsilon(t_n^{(\varepsilon)}, x_n^{(\varepsilon)}), \varphi_\varepsilon(t_n^{(\varepsilon)}, \cdot)) \leq 0. \quad (5.45)$$

Furthermore, by (5.41), (5.43), and (5.44) it holds for all $\varepsilon \in (0, \infty)$ that

$$\begin{aligned} 0 &= \limsup_{n \rightarrow \infty} [u(t_0, x_0) - u^{(n)}(t_0, x_0)] \\ &= \limsup_{n \rightarrow \infty} [\varphi_\varepsilon(t_0, x_0) - u^{(n)}(t_0, x_0)] \\ &\geq \limsup_{n \rightarrow \infty} [\varphi_\varepsilon(t_n^{(\varepsilon)}, x_n^{(\varepsilon)}) - u^{(n)}(t_n^{(\varepsilon)}, x_n^{(\varepsilon)})] \\ &= \limsup_{n \rightarrow \infty} \left[\varphi(t_n^{(\varepsilon)}, x_n^{(\varepsilon)}) + \frac{\varepsilon}{2} \left(|t_n^{(\varepsilon)} - t_0|^2 + \|x_n^{(\varepsilon)} - x_0\|^2 \right) - u^{(n)}(t_n^{(\varepsilon)}, x_n^{(\varepsilon)}) \right] \\ &\geq \limsup_{n \rightarrow \infty} \left[u(t_n^{(\varepsilon)}, x_n^{(\varepsilon)}) - u^{(n)}(t_n^{(\varepsilon)}, x_n^{(\varepsilon)}) + \frac{\varepsilon}{2} \left(|t_n^{(\varepsilon)} - t_0|^2 + \|x_n^{(\varepsilon)} - x_0\|^2 \right) \right] \\ &= \limsup_{n \rightarrow \infty} \left[\frac{\varepsilon}{2} \left(|t_n^{(\varepsilon)} - t_0|^2 + \|x_n^{(\varepsilon)} - x_0\|^2 \right) \right]. \end{aligned}$$

Therefore, we see that for all $\varepsilon \in (0, \infty)$

$$\limsup_{n \rightarrow \infty} \left[\frac{\varepsilon}{2} \left(|t_n^{(\varepsilon)} - t_0|^2 + \|x_n^{(\varepsilon)} - x_0\|^2 \right) \right] = 0. \quad (5.46)$$

Moreover, the continuity of u , (5.41), and (5.46) ensure for all $\varepsilon \in (0, \infty)$ that

$$\begin{aligned} \lim_{n \rightarrow \infty} |u^{(n)}(t_n^{(\varepsilon)}, x_n^{(\varepsilon)}) - u(t_0, x_0)| &\leq \lim_{n \rightarrow \infty} |u^{(n)}(t_n^{(\varepsilon)}, x_n^{(\varepsilon)}) - u(t_n^{(\varepsilon)}, x_n^{(\varepsilon)})| \\ &\quad + \lim_{n \rightarrow \infty} |u(t_n^{(\varepsilon)}, x_n^{(\varepsilon)}) - u(t_0, x_0)| = 0. \end{aligned} \quad (5.47)$$

Then by (5.43), (5.46), (5.47), Lemma 5.3, and Lemma 5.10, it holds for all $\varepsilon \in (0, \infty)$ that

$$\lim_{n \rightarrow \infty} \left| \frac{\partial}{\partial t} \varphi_\varepsilon(t_n^{(\varepsilon)}, x_n^{(\varepsilon)}) - \frac{\partial}{\partial t} \varphi(t_0, x_0) \right| = 0,$$

and

$$\begin{aligned} &G_0 \left(t_0, x_0, \varphi(t_0, x_0), \nabla_x \varphi(t_0, x_0), \text{Hess}_x \varphi(t_0, x_0) + \varepsilon \mathbf{I}_d, \varphi(t_0, \cdot) \right) \\ &= G_0 \left(t_0, x_0, u(t_0, x_0), \nabla_x \varphi_\varepsilon(t_0, x_0), \text{Hess}_x \varphi_\varepsilon(t_0, x_0), \varphi_\varepsilon(t_0, \cdot) \right) \\ &= \lim_{n \rightarrow \infty} G_0 \left(t_n^{(\varepsilon)}, x_n^{(\varepsilon)}, u^{(n)}(t_n^{(\varepsilon)}, x_n^{(\varepsilon)}), \nabla_x \varphi_\varepsilon(t_n^{(\varepsilon)}, x_n^{(\varepsilon)}), \text{Hess}_x \varphi_\varepsilon(t_n^{(\varepsilon)}, x_n^{(\varepsilon)}), \varphi_\varepsilon(t_n^{(\varepsilon)}, \cdot) \right) \\ &= \lim_{n \rightarrow \infty} G_0^{(n)} \left(t_n^{(\varepsilon)}, x_n^{(\varepsilon)}, u^{(n)}(t_n^{(\varepsilon)}, x_n^{(\varepsilon)}), \nabla_x \varphi_\varepsilon(t_n^{(\varepsilon)}, x_n^{(\varepsilon)}), \text{Hess}_x \varphi_\varepsilon(t_n^{(\varepsilon)}, x_n^{(\varepsilon)}), \varphi_\varepsilon(t_n^{(\varepsilon)}, \cdot) \right). \end{aligned}$$

Hence, taking the limit as $n \rightarrow \infty$ in (5.45), and using the assumption $u(t_0, x_0) = \varphi(t_0, x_0)$ yields that for all $\varepsilon \in (0, \infty)$

$$-\frac{\partial}{\partial t} \varphi(t_0, x_0) + G_0 \left(t_0, x_0, \varphi(t_0, x_0), \nabla_x \varphi(t_0, x_0), \text{Hess}_x \varphi(t_0, x_0) + \varepsilon \mathbf{I}_d, \varphi(t_0, \cdot) \right) \leq 0.$$

Lemma 5.3 therefore ensures that

$$-\frac{\partial}{\partial t} \varphi(t_0, x_0) + G_0 \left(t_0, x_0, \varphi(t_0, x_0), \nabla_x \varphi(t_0, x_0), \text{Hess}_x \varphi(t_0, x_0), \varphi(t_0, \cdot) \right) \leq 0,$$

which shows that u is a viscosity subsolution of (5.24). The proof of this lemma is thus completed. \square

Lemma 5.12. *Let Assumptions 2.1, 2.2, and 5.1-5.9 hold. Assume that g , μ , σ , and h are compactly supported, and that $\eta(\cdot, z)$ is compactly supported for all $z \in \mathbb{R}^d$. Also assume for every $n \in \mathbb{N}$, $t \in [0, T]$, and $z \in \mathbb{R}^d$ that $\mu^{(n)}(t, \cdot)$, $\sigma^{(n)}(t, \cdot)$, $\eta_t^{(n)}(\cdot, z)$, $g^{(n)}(\cdot)$, and $h^{(n)}(t, \cdot)$ are compactly supported and infinitely differentiable. Moreover, we assume that there exists a non-empty compact set $\mathcal{K}_0 \in \mathbb{R}^d$ such that for all $n \in \mathbb{N}$, $t \in [0, T]$, and $z \in \mathbb{R}^d$*

$$\text{supp}\{\mu^{(n)}(t, \cdot)\} \cup \text{supp}\{\sigma^{(n)}(t, \cdot)\} \cup \text{supp}\{\eta_t^{(n)}(\cdot, z)\} \cup \text{supp}\{g^{(n)}(\cdot)\} \subseteq \mathcal{K}_0, \quad (5.48)$$

where $\text{supp}\{\mathfrak{f}\}$ denotes the support of the real-valued function \mathfrak{f} on \mathbb{R}^d . For every $n \in \mathbb{N}$, let $(X_s^{t,x,(n)})_{s \in [t, T]}$: $[t, T] \times \Omega \rightarrow \mathbb{R}^d$ be an \mathbb{F} -adapted càdlàg process satisfying that $X_t^{t,x,(n)} = x$ and almost surely for all

$s \in [0, T]$

$$dX_s^{t,x,(n)} = \mu^{(n)}(s, X_{s-}^{t,x,(n)}) ds + \sigma^{(n)}(s, X_{s-}^{t,x,(n)}) dW_s + \int_{\mathbb{R}^d} \eta_s^{(n)}(X_{s-}^{t,x,(n)}, z) \tilde{\pi}(dz, ds). \quad (5.49)$$

Then there is a constants $K_0 > 0$ only depending on T and L such that for all $n \in \mathbb{N}$ and $(t, x) \in [0, T] \times \mathbb{R}^d$

$$\mathbb{E} \left[\sup_{s \in [t, T]} \|X_s^{t,x,(n)} - X_s^{t,x}\|^2 \right] \leq K_0 \cdot \left[\sup_{(r,y) \in [0, T] \times \mathbb{R}^d} \left\{ \|\mu^{(n)}(r, y) - \mu(r, y)\|^2 + \|\sigma^{(n)}(r, y) - \sigma(r, y)\|_F^2 + \int_{\mathbb{R}^d} \|\eta_r^{(n)}(y, z) - \eta_r(y, z)\|^2 \nu(dz) \right\} \right] \quad (5.50)$$

Proof. We first notice for each $(t, x) \in [0, T] \times \mathbb{R}^d$, and $n \in \mathbb{N}$ that it holds almost surely for all $s \in [t, T]$ that

$$\begin{aligned} X_s^{t,x,(n)} - X_s^{t,x} &= \int_t^s \left(\mu^{(n)}(r, X_{r-}^{t,x,(n)}) - \mu(r, X_{r-}^{t,x}) \right) dr \\ &\quad + \int_t^s \left(\sigma^{(n)}(r, X_{r-}^{t,x,(n)}) - \sigma(r, X_{r-}^{t,x}) \right) dW_r \\ &\quad + \int_t^s \int_{\mathbb{R}^d} \left(\eta_r^{(n)}(X_{r-}^{t,x,(n)}, z) - \eta_r(X_{r-}^{t,x}, z) \right) \tilde{\pi}(dz, dr). \end{aligned}$$

Thus, by Jensen's inequality, Burkholder-Davis-Gundy inequality, and the fact that $(\sum_{i=1}^3 a_i)^2 \leq 3 \sum_{i=1}^3 a_i^2$ for $a_i \in \mathbb{R}$, $i = 1, 2, 3$, it holds for all $(t, x) \in [0, T] \times \mathbb{R}^d$, $s \in [t, T]$, and $n \in \mathbb{N}$ that

$$e_s^{(n)}(t, x) := \mathbb{E} \left[\sup_{r \in [t, s]} \|X_r^{t,x,(n)} - X_r^{t,x}\|^2 \right] \leq \sum_{i=1}^3 A_s^{(n),i}(t, x), \quad (5.51)$$

where

$$\begin{aligned} A_s^{(n),1}(t, x) &:= 3T \int_t^s \mathbb{E} \left[\|\mu^{(n)}(r, X_r^{t,x,(n)}) - \mu(r, X_r^{t,x})\|^2 \right] dr, \\ A_s^{(n),2}(t, x) &:= 24 \int_t^s \mathbb{E} \left[\|\sigma^{(n)}(r, X_r^{t,x,(n)}) - \sigma(r, X_r^{t,x})\|_F^2 \right] dr, \\ A_s^{(n),3}(t, x) &:= 24 \int_t^s \mathbb{E} \left[\int_{\mathbb{R}^d} \|\eta_r^{(n)}(X_r^{t,x,(n)}, z) - \eta_r(X_r^{t,x}, z)\|^2 \nu(dz) \right] dr. \end{aligned}$$

We notice that Assumptions 2.1, 5.7, and 5.8 implies for all $(t, x) \in [0, T] \times \mathbb{R}^d$, $s \in [0, T]$, and $n \in \mathbb{N}$ that

$$\begin{aligned} A_s^{(n),1}(t, x) &\leq 6T \int_t^s \mathbb{E} \left[\|\mu^{(n)}(r, X_r^{t,x,(n)}) - \mu(r, X_r^{t,x,(n)})\|^2 \right] dr \\ &\quad + 6T \int_t^s \mathbb{E} \left[\|\mu(r, X_r^{t,x,(n)}) - \mu(r, X_r^{t,x})\|^2 \right] dr \\ &\leq 6T^2 \cdot \left[\sup_{(r,y) \in [0, T] \times \mathbb{R}^d} \|\mu^{(n)}(r, y) - \mu(r, y)\|^2 \right] + 6TL \int_t^s e_r^{(n)}(t, x) dr, \end{aligned} \quad (5.52)$$

$$A_s^{(n),2}(t, x) \leq 48T \cdot \left[\sup_{(r,y) \in [0, T] \times \mathbb{R}^d} \|\sigma^{(n)}(r, y) - \sigma(r, y)\|_F^2 \right] + 48L \int_t^s e_r^{(n)}(t, x) dr, \quad (5.53)$$

and

$$A_s^{(n),3}(t, x) \leq 48T \cdot \left[\sup_{(r,y) \in [0, T] \times \mathbb{R}^d} \int_{\mathbb{R}^d} \|\eta_r^{(n)}(y, z) - \eta_r(y, z)\|^2 \nu(dz) \right] + 48L \int_t^s e_r^{(n)}(t, x) dr. \quad (5.54)$$

Moreover, by (2.9), and Assumptions 5.7 and 5.8 it holds for all $(t, x) \in [0, T] \times \mathbb{R}^d$ and $n \in \mathbb{N}$ that

$$\mathbb{E} \left[\sup_{s \in [t, T]} \|X_s^{t,x,(n)}\|^2 \right] < \infty. \quad (5.55)$$

Then by (2.9), (5.51)-(5.55), and Grönwall's lemma we obtain for all $(t, x) \in [0, T] \times \mathbb{R}^d$ and $n \in \mathbb{N}$ that

$$e_T^{(n)}(t, x) \leq 6T(T+16) \exp\{6L(T+16)(T-t)\} \cdot \left[\sup_{(r,y) \in [0,T] \times \mathbb{R}^d} \left\{ \left\| \mu^{(n)}(r, y) - \mu(r, y) \right\|^2 + \left\| \sigma^{(n)}(r, y) - \sigma(r, y) \right\|_F^2 + \int_{\mathbb{R}^d} \left\| \eta_r^{(n)}(y, z) - \eta_r(y, z) \right\|^2 \nu(dz) \right\} \right].$$

The proof of this lemma is thus completed. \square

Proposition 5.13. *Let Assumptions 2.1, 2.2, 2.3, and 5.1 hold. Moreover, we assume that g, μ, σ , and h are compactly supported, and that $\eta(\cdot, z)$ is compactly supported for all $z \in \mathbb{R}^d$. Let $u : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ satisfy for all $(t, x) \in [0, T] \times \mathbb{R}^d$ that*

$$u(t, x) = \mathbb{E} \left[g(X_T^{t,x}) + \int_t^T h(s, X_s^{t,x}) ds \right]. \quad (5.56)$$

Then u is a viscosity solution of the linear PIDE (5.24) with $u(T, x) = g(x)$ for $x \in \mathbb{R}^d$.

To prove the above proposition, we recall the notion of mollifications of functions. For $\varepsilon > 0$ and locally integrable functions $\phi : \mathbb{R}^d \rightarrow \mathbb{R}$ we use the notation $\phi^{(\varepsilon)}$ for the mollification of ϕ , defined by

$$\phi^{(\varepsilon)}(x) := \varepsilon^{-d} \int_{\mathbb{R}^d} \phi(y) k((x-y)/\varepsilon) dy = \int_{\mathbb{R}^d} \phi(x-\varepsilon z) k(z) dz, \quad x \in \mathbb{R}^d, \quad (5.57)$$

where $k : \mathbb{R}^d \rightarrow \mathbb{R}$ is a fixed nonnegative smooth function on \mathbb{R}^d such that $k(x) = 0$ for $|x| \geq 1$, $k(-x) = k(x)$ for $x \in \mathbb{R}^d$, and $\int_{\mathbb{R}^d} k(x) dx = 1$.

Proof of Proposition 5.13. Let $\{\varepsilon_n\}_{n=1}^\infty \subseteq (0, 1]$ such that $\lim_{n \rightarrow \infty} \varepsilon_n = 0$. Then for each $n \in \mathbb{N}$ we denote by $g^{(\varepsilon_n)}, h^{(\varepsilon_n)}, \mu^{(\varepsilon_n)}, \sigma^{(\varepsilon_n)}$, and $\eta^{(\varepsilon_n)}$ the mollifications (in $x \in \mathbb{R}^d$) of g, h, μ, σ , and η , respectively. For every $n \in \mathbb{N}$, let $(X_s^{t,x,(n)})_{s \in [t, T]} : [t, T] \times \Omega \rightarrow \mathbb{R}^d$ be an \mathbb{F} -adapted càdlàg process satisfying that $X_t^{t,x,(n)} = x$ and almost surely for all $s \in [0, T]$

$$dX_s^{t,x,(n)} = \mu^{(\varepsilon_n)}(s, X_{s-}^{t,x,(n)}) ds + \sigma^{(\varepsilon_n)}(s, X_{s-}^{t,x,(n)}) dW_s + \int_{\mathbb{R}^d} \eta_s^{(\varepsilon_n)}(X_{s-}^{t,x,(n)}, z) \tilde{\pi}(dz, ds). \quad (5.58)$$

For each $n \in \mathbb{N}, k \in \mathbb{N}$, and $(t, x) \in [0, T] \times \mathbb{R}^d$, we define

$$u^{n,k}(t, x) := \mathbb{E} \left[g^{(\varepsilon_k)}(X_T^{t,x,(n)}) + \int_t^T h^{(\varepsilon_k)}(s, X_s^{t,x,(n)}) ds \right], \quad (5.59)$$

and

$$u^{0,k}(t, x) := \mathbb{E} \left[g^{(\varepsilon_k)}(X_T^{t,x}) + \int_t^T h^{(\varepsilon_k)}(s, X_s^{t,x}) ds \right]. \quad (5.60)$$

We first notice that by (5.57) it holds for all $k \in \mathbb{N}$ and $(t, x) \in [0, T] \times \mathbb{R}^d$ that

$$\begin{aligned} |h^{(\varepsilon_k)}(t, x) - h(t, x)| &= \varepsilon_k^{-d} \left| \int_{\mathbb{R}^d} [h(t, y) - h(t, x)] k((x-y)/\varepsilon_k) dy \right| \\ &\leq \left[\sup_{s \in [0, T]} \sup_{\|y-y'\| \leq \varepsilon_k} (|h(t, y) - h(t, y')|) \right] \cdot \int_{\mathbb{R}^d} \varepsilon_k^{-d} k((x-y)/\varepsilon_k) dy \\ &= \sup_{s \in [0, T]} \sup_{\|y-y'\| \leq \varepsilon_k} (|h(s, y) - h(s, y')|). \end{aligned}$$

This together with the assumption that h is compactly supported imply

$$\lim_{k \rightarrow \infty} \left[\sup_{(t,x) \in [0, T] \times \mathbb{R}^d} |h^{(\varepsilon_k)}(t, x) - h(t, x)| \right] = 0. \quad (5.61)$$

Moreover, by (2.3), (5.57), and Jensen's inequality we first observe for all $(t, x) \in [0, T] \times \mathbb{R}^d, n \in \mathbb{N}$, and $z \in \mathbb{R}^d$ that

$$\left\| \mu^{(\varepsilon_n)}(t, x) - \mu(t, x) \right\|^2 = \left\| \int_{\mathbb{R}^d} \mu(t, x - \varepsilon_n y) k(y) dy - \int_{\mathbb{R}^d} \mu(t, x) k(y) dy \right\|^2$$

$$\leq \int_{\mathbb{R}^d} \|\mu(t, x - \varepsilon_n y) - \mu(t, x)\|^2 k(y) dy \leq \varepsilon_n^2 L, \quad (5.62)$$

$$\|\sigma^{(\varepsilon_n)}(t, x) - \sigma(t, x)\|_F^2 \leq \varepsilon_n^2 L, \quad \|\eta_t^{(\varepsilon_n)}(x, z) - \eta_t(x, z)\|^2 \leq \varepsilon_n^2 C_d(1 \wedge \|z\|^2), \quad (5.63)$$

and

$$|g^{(\varepsilon_n)}(x) - g(x)|^2 \leq \varepsilon_n^2 L T^{-1}.$$

Hence we have

$$\lim_{n \rightarrow \infty} \left[\sup_{(t,x) \in [0,T] \times \mathbb{R}^d} \left(\|\mu^{(\varepsilon_n)}(t, x) - \mu(t, x)\|^2 + \|\sigma^{(\varepsilon_n)}(t, x) - \sigma(t, x)\|^2 + |g^{(\varepsilon_n)}(x) - g(x)|^2 \right) \right] = 0, \quad (5.64)$$

and

$$\lim_{n \rightarrow \infty} \left[\sup_{(t,x) \in [0,T] \times \mathbb{R}^d} \|\eta_t^{(\varepsilon_n)}(x, z) - \eta_t(x, z)\|^2 \right] = 0 \quad \text{for all } z \in \mathbb{R}^d. \quad (5.65)$$

Hence by (5.61) and analogous argument to obtain (5.64) and (5.65), we have that Assumption 5.6 holds with $g^{(\varepsilon_n)}$, $h^{(\varepsilon_n)}$, $\mu^{(\varepsilon_n)}$, $\sigma^{(\varepsilon_n)}$, and $\eta^{(\varepsilon_n)}$ in place of $g^{(n)}$, $h^{(n)}$, $\mu^{(n)}$, $\sigma^{(n)}$, and $\eta^{(n)}$, respectively. Moreover, by (2.2), (2.4), (2.6), and Jensen's inequality we have for all $t \in [0, T]$, $x, x' \in \mathbb{R}^d$, $z \in \mathbb{R}^d$, and $n \in \mathbb{N}$ that

$$\begin{aligned} \|\eta_t^{(\varepsilon_n)}(x, z) - \eta_t^{(\varepsilon_n)}(x', z)\|^2 &= \left\| \int_{\mathbb{R}^d} \eta_t(x - \varepsilon_n y, z) k(y) dy - \int_{\mathbb{R}^d} \eta_t(x' - \varepsilon_n y, z) k(y) dy \right\|^2 \\ &\leq \int_{\mathbb{R}^d} \|\eta_t(x - \varepsilon_n y, z) - \eta_t(x' - \varepsilon_n y, z)\|^2 k(y) dy \\ &\leq C_d(1 \wedge \|z\|^2) \|x - x'\|^2 \int_{\mathbb{R}^d} k(y) dy \\ &= C_d(1 \wedge \|z\|^2) \|x - x'\|^2, \\ \|g^{(\varepsilon_n)}(x) - g^{(\varepsilon_n)}(x')\|^2 &\leq L T^{-1} \|x - x'\|^2, \end{aligned} \quad (5.66)$$

$$\begin{aligned} \|\eta_t^{(\varepsilon_n)}(x, z)\|^2 &\leq \int_{\mathbb{R}^d} \|\eta_t(x - \varepsilon_n y, z)\|^2 k(y) dy \leq C_d(1 \wedge \|z\|^2) \int_{\mathbb{R}^d} k(y) dy \\ &= C_d(1 \wedge \|z\|^2), \end{aligned} \quad (5.67)$$

$$\begin{aligned} \|g^{(\varepsilon_n)}(t, x)\|^2 &\leq \int_{\mathbb{R}^d} \|g(t, x - \varepsilon_n y)\|^2 k(y) dy \leq \int_{\mathbb{R}^d} L(d^p + \|x - \varepsilon_n y\|^2) k(y) dy \\ &\leq L(d^p + 2\|x\|^2) + \int_{\mathbb{R}^d} 2L\|\varepsilon_n y\|^2 k(y) dy \\ &\leq 2L(d^p + 1)(1 + \|x\|^2), \end{aligned} \quad (5.68)$$

and

$$\|h^{(\varepsilon_n)}(t, x)\|^2 \leq 2(C_0 + 1)(1 + \|x\|^2). \quad (5.69)$$

Thus, by analogous argument also for $\mu^{(\varepsilon_n)}$ and $\sigma^{(\varepsilon_n)}$ we get that Assumptions 5.7 and 5.8 hold with $h^{(\varepsilon_n)}$, $g^{(\varepsilon_n)}$, $\mu^{(\varepsilon_n)}$, $\sigma^{(\varepsilon_n)}$, and $\eta^{(\varepsilon_n)}$ in place of $h^{(n)}$, $g^{(n)}$, $\mu^{(n)}$, $\sigma^{(n)}$, and $\eta^{(n)}$, respectively. Furthermore, by virtue of Corollary 3.6 in [21] there exist constants $\delta' > 0$ and $\varepsilon_0 > 0$ such that for all $\varepsilon \in (0, \varepsilon_0)$, $(t, x) \in [0, T] \times \mathbb{R}^d$, $z \in \mathbb{R}^d$, and $\gamma \in [0, 1]$

$$\delta' \leq |\det(\mathbf{I}_d + \gamma D_x \eta_t^{(\varepsilon)}(x, z))|.$$

This verifies that Assumption 5.9 holds for sufficiently large n , with $\eta^{(\varepsilon_n)}$ in place of $\eta^{(n)}$. Moreover, the application of Lemma 5.12 yields for all $n \in \mathbb{N}$ and $(t, x) \in [0, T] \times \mathbb{R}^d$ that

$$\begin{aligned} \mathbb{E} \left[\sup_{s \in [t, T]} \|X_s^{t,x,(n)} - X_s^{t,x}\|^2 \right] &\leq c_0 \cdot \left[\sup_{(r,y) \in [0,T] \times \mathbb{R}^d} \left\{ \|\mu^{(\varepsilon_n)}(r, y) - \mu(r, y)\|^2 \right. \right. \\ &\quad \left. \left. + \|\sigma^{(\varepsilon_n)}(r, y) - \sigma(r, y)\|_F^2 + \int_{\mathbb{R}^d} \|\eta_r^{(\varepsilon_n)}(y, z) - \eta_r(y, z)\|^2 \nu(dz) \right\} \right], \end{aligned} \quad (5.70)$$

where c_0 a positive constant only depending on T and L . Hence, by (2.8) (5.62) and (5.63) we obtain for all $n \in \mathbb{N}$ that

$$\begin{aligned} \sup_{(t,x) \in [0,T] \times \mathbb{R}^d} \mathbb{E} \left[\sup_{s \in [t,T]} \|X_s^{t,x,(n)} - X_s^{t,x}\|^2 \right] &\leq c_0 \cdot \left[2\varepsilon_n^2 L + \varepsilon_n^2 C_d \int_{\mathbb{R}^d} (1 \wedge \|z\|^2) \nu(dz) \right] \\ &\leq c_0 \varepsilon_n^2 (2L + C_d K d^p), \end{aligned} \quad (5.71)$$

which implies

$$\lim_{n \rightarrow \infty} \left(\sup_{(t,x) \in [0,T] \times \mathbb{R}^d} \mathbb{E} \left[\sup_{s \in [t,T]} \|X_s^{t,x,(n)} - X_s^{t,x}\|^2 \right] \right) = 0. \quad (5.72)$$

Furthermore, by Assumptions 5.7-5.9, applying Lemma 5.5 (with $\mu \curvearrowright \mu^{(\varepsilon_n)}$, $\sigma \curvearrowright \sigma^{(\varepsilon_n)}$, $\eta \curvearrowright \eta^{(\varepsilon_n)}$, $g \curvearrowright g^{(\varepsilon_k)}$, $h \curvearrowright h^{(\varepsilon_k)}$ for $n, k \in \mathbb{N}$ in the notation of Proposition 5.5) yields that for all $k \in \mathbb{N}$ and sufficiently large $n \in \mathbb{N}$ the function $u^{n,k} \in C^{1,2}([0, T] \times \mathbb{R}^d)$ is a classical solution of

$$\begin{aligned} \frac{\partial}{\partial t} u^{n,k}(t, x) + \langle \nabla_x u^{n,k}(t, x), \mu^{(\varepsilon_n)}(t, x) \rangle + \frac{1}{2} \text{Trace} \left(\sigma^{(\varepsilon_n)}(t, x) [\sigma^{(\varepsilon_n)}(t, x)]^T \text{Hess}_x u^{n,k}(t, x) \right) \\ + h^{(\varepsilon_k)}(t, x) + \int_{\mathbb{R}^d} \left(u^{n,k}(t, x + \eta_t^{(\varepsilon_n)}(x, z)) - u^{n,k}(t, x) - \langle \nabla_x u^{n,k}(t, x), \eta_t^{(\varepsilon_n)}(x, z) \rangle \right) \nu(dz) = 0 \end{aligned} \quad (5.73)$$

on $(0, T) \times \mathbb{R}^d$ with $u^{n,k}(T, x) = g^{(\varepsilon_k)}(x)$ on \mathbb{R}^d .

Next, by well-known properties of derivatives of mollifications (see, e.g., Theorem 2.29 in [1]) and change of variables, we have for every $k \in \mathbb{N}$, $i \in \{1, \dots, d\}$, and $(t, x) \in [0, T] \times \mathbb{R}^d$ that

$$\begin{aligned} \frac{\partial}{\partial x_i} h^{(\varepsilon_k)}(t, x) &= \varepsilon_k^{-(d+1)} \int_{\mathbb{R}^d} h(t, y) \left(\frac{\partial}{\partial x_i} k \right) ((x - y)/\varepsilon_k) dy \\ &= \varepsilon_k^{-1} \int_{\mathbb{R}^d} h(t, x - \varepsilon_k z) \frac{\partial}{\partial x_i} k(z) dz. \end{aligned} \quad (5.74)$$

This together with the fact that the functions h and k are compactly supported and continuous imply for all $k \in \mathbb{N}$ and $i \in \{1, \dots, d\}$ that $\frac{\partial}{\partial x_i} h^{(\varepsilon_k)} \in C([0, T] \times \mathbb{R}^d)$ and $\frac{\partial}{\partial x_i} h^{(\varepsilon_k)}$ is compactly supported. Therefore, we obtain that for all $k \in \mathbb{N}$

$$\sup_{(t,x) \in [0,T] \times \mathbb{R}^d} \|\nabla_x h^{(\varepsilon_k)}(t, x)\| < \infty. \quad (5.75)$$

Furthermore, by (5.59), (5.60), (5.66), and the mean value theorem it holds for all $n, k \in \mathbb{N}$ and $x \in \mathbb{R}^d$ that

$$\begin{aligned} &|u^{n,k}(t, x) - u^{0,k}(t, x)| \\ &\leq \mathbb{E} \left[|g^{(\varepsilon_k)}(X_T^{t,x,(n)}) - g^{(\varepsilon_k)}(X_T^{t,x})| \right] + \int_t^T \mathbb{E} \left[|h^{(\varepsilon_k)}(s, X_s^{t,x,(n)}) - h^{(\varepsilon_k)}(s, X_s^{t,x})| \right] ds \\ &\leq \left(\mathbb{E} \left[LT^{-1} \|X_T^{t,x,(n)} - X_T^{t,x}\|^2 \right] \right)^{1/2} \\ &\quad + \int_t^T \mathbb{E} \left[\left(\sup_{(r,y) \in [0,T] \times \mathbb{R}^d} \|\nabla_y h^{(\varepsilon_k)}(r, y)\| \right) \cdot \|X_s^{t,x,(n)} - X_s^{t,x}\| \right] ds \\ &\leq \left(\mathbb{E} \left[LT^{-1} \|X_T^{t,x,(n)} - X_T^{t,x}\|^2 \right] \right)^{1/2} \\ &\quad + T \left(\sup_{(r,y) \in [0,T] \times \mathbb{R}^d} \|\nabla_y h^{(\varepsilon_k)}(r, y)\| \right) \cdot \left(\mathbb{E} \left[\sup_{s \in [t,T]} \|X_s^{t,x,(n)} - X_s^{t,x}\|^2 \right] \right)^{1/2} \end{aligned} \quad (5.76)$$

This together with (5.72) and (5.75) show for every $k \in \mathbb{N}$ that

$$\lim_{n \rightarrow \infty} \left[\sup_{(t,x) \in [0,T] \times \mathbb{R}^d} |u^{n,k}(t, x) - u^{0,k}(t, x)| \right] = 0. \quad (5.77)$$

Furthermore, by (2.9), (5.56), Assumptions 2.1 and 5.1, and Hölder's inequality we have for all $(t, x) \in [0, T]$ that

$$\begin{aligned} |u(t, x)| &\leq \left(\mathbb{E} \left[|g(X_T^{t,x})|^2 \right] \right)^{1/2} + \int_t^T \left(\mathbb{E} \left[|h(s, X_s^{t,x})|^2 \right] \right)^{1/2} ds \\ &\leq \left(\mathbb{E} \left[L(d^p + \|X_T^{t,x}\|^2) \right] \right)^{1/2} + \int_t^T \left(\mathbb{E} \left[C_0(1 + \|X_s^{t,x}\|^2) \right] \right)^{1/2} ds \\ &\leq [Ld^p + LC(1 + \|x\|^2)]^{1/2} + T [C_0 + C_0C(1 + \|x\|^2)]^{1/2}. \end{aligned}$$

This implies that

$$\sup_{(t,x) \in [0,T] \times \mathbb{R}^d} \frac{|u(t, x)|}{1 + \|x\|} < \infty. \quad (5.78)$$

Similarly, using Assumption 5.7 we also obtain for all $k, n \in \mathbb{N}$ that

$$\sup_{(t,x) \in [0,T] \times \mathbb{R}^d} \frac{|u^{n,k}(t, x)| + |u^{0,k}(t, x)|}{1 + \|x\|} < \infty. \quad (5.79)$$

Hence, by (5.77) and the fact that $u^{n,k} \in C_1([0, T] \times \mathbb{R}^d)$ for all $k \in \mathbb{N}$ and sufficiently large $n \in \mathbb{N}$ shows that $u^{0,k} \in C_1([0, T] \times \mathbb{R}^d, \mathbb{R})$ for all $k \in \mathbb{N}$. Therefore, by (5.73), (5.77), and Assumptions 5.6-5.8, applying Lemma 5.11 yields for all $k \in \mathbb{N}$ that the function $u^{0,k}$ is a viscosity solution of

$$\begin{aligned} &\frac{\partial}{\partial t} u^{0,k}(t, x) + \langle \nabla_x u^{0,k}(t, x), \mu(t, x) \rangle + \frac{1}{2} \text{Trace} \left(\sigma(t, x) [\sigma(t, x)]^T \text{Hess}_x u^{0,k}(t, x) \right) \\ &+ h^{(\varepsilon_k)}(t, x) + \int_{\mathbb{R}^d} \left(u^{0,k}(t, x + \eta_t(x, z)) - u^{0,k}(t, x) - \langle \nabla_x u^{0,k}(t, x), \eta_t(x, z) \rangle \right) \nu(dz) = 0 \end{aligned} \quad (5.80)$$

on $(0, T) \times \mathbb{R}^d$ with $u^{0,k}(T, x) = g^{(\varepsilon_k)}(x)$ on \mathbb{R}^d . Furthermore, by (5.56) and (5.60) we obtain for all $k \in \mathbb{N}$ and $(t, x) \in [0, T] \times \mathbb{R}^d$ that

$$\begin{aligned} |u(t, x) - u^{0,k}(t, x)| &\leq \mathbb{E} \left[|g^{(\varepsilon_k)}(X_T^{t,x}) - g(X_T^{t,x})| \right] + \int_t^T \mathbb{E} \left[|h^{(\varepsilon_k)}(s, X_s^{t,x}) - h(s, X_s^{t,x})| \right] ds \\ &\leq \sup_{y \in \mathbb{R}^d} \left\{ |g^{(\varepsilon_k)}(y) - g(y)| \right\} + T \cdot \sup_{(r,y) \in [0,T] \times \mathbb{R}^d} \left\{ |h^{(\varepsilon_k)}(s, y) - h(s, y)| \right\}. \end{aligned}$$

This together with (5.61) and (5.64) imply that

$$\lim_{n \rightarrow \infty} \sup_{(t,x) \in [0,T] \times \mathbb{R}^d} \left\{ |u(t, x) - u^{0,k}(t, x)| \right\} = 0. \quad (5.81)$$

Hence, (5.79) and the fact that $u^{0,k} \in C_1([0, T] \times \mathbb{R}^d)$ for all $k \in \mathbb{N}$ ensures $u \in C_1([0, T] \times \mathbb{R}^d)$. Therefore, by (5.61), (5.64), (5.80), Assumptions 2.1-2.3, and Assumption 5.1, the application of Lemma 5.11 yields that u is a viscosity solution of (5.24) with $u(T, x) = g(x)$ for all $x \in \mathbb{R}^d$. \square

Lemma 5.14. *Let $B \subseteq \mathbb{R}^d$ be a closed set. Let $\bar{\mu} \in C([0, T] \times \mathbb{R}^d, \mathbb{R}^d)$, $\bar{\sigma} \in C([0, T] \times \mathbb{R}^d, \mathbb{R}^{d \times d})$, and $\bar{\eta} \in C([0, T] \times \mathbb{R}^d \times \mathbb{R}^d, \mathbb{R}^d)$ such that for all $(t, x) \in [0, T] \times B$ and $z \in \mathbb{R}^d$*

$$\bar{\mu}(t, x) = \mu(t, x), \quad \bar{\sigma}(t, x) = \sigma(t, x), \quad \bar{\eta}_t(x, z) = \eta_t(x, z). \quad (5.82)$$

Assume that $\bar{\mu}$, $\bar{\sigma}$, and $\bar{\eta}$ satisfy Assumptions 2.1 and 2.2. Moreover, for each $(t, x) \in [0, T] \times \mathbb{R}^d$ let $(\bar{X}_s^{t,x})_{s \in [t, T]} : [t, T] \times \Omega \rightarrow \mathbb{R}^d$ be an \mathbb{F} -adapted càdlàg process satisfying that $\bar{X}_t^{t,x} = x$, and almost surely for all $s \in [t, T]$

$$d\bar{X}_s^{t,x} = \bar{\mu}(s, \bar{X}_{s-}^{t,x}) ds + \bar{\sigma}(s, \bar{X}_{s-}^{t,x}) dW_s + \int_{\mathbb{R}^d} \bar{\eta}_s(\bar{X}_{s-}^{t,x}, z) \tilde{\pi}(dz, ds).$$

For each $(t, x) \in [0, T] \times \mathbb{R}^d$, let $\tau^{t,x} : \Omega \rightarrow [0, T]$ be a stopping time defined by

$$\tau^{t,x} := \inf \{ s \geq t : X_s^{t,x} \notin B \text{ or } \bar{X}_s^{t,x} \notin B \} \wedge T. \quad (5.83)$$

Then it holds for all $(t, x) \in [0, T] \times \mathbb{R}^d$ that

$$\mathbb{P} \left[\mathbf{1}_{\{s \leq \tau\}} \|X_s^{t,x} - \bar{X}_s^{t,x}\| = 0 \text{ for all } s \in [t, T] \right] = 1. \quad (5.84)$$

Proof. Throughout this proof, we fix $(t, x) \in [0, T] \times \mathbb{R}^d$, and use the notation $\tau = \tau^{t,x}$. We first notice that almost surely for all $s \in [t, T]$

$$\begin{aligned} \bar{X}_{s \wedge \tau}^{t,x} - X_{s \wedge \tau}^{t,x} &= \int_t^{s \wedge \tau} \left(\bar{\mu}(r, \bar{X}_{r-}^{t,x}) - \mu(r, X_{r-}^{t,x}) \right) dr + \int_t^{s \wedge \tau} \left(\bar{\sigma}(r, \bar{X}_{r-}^{t,x}) - \sigma(r, X_{r-}^{t,x}) \right) dW_r \\ &\quad + \int_t^{s \wedge \tau} \int_{\mathbb{R}^d} \left(\bar{\eta}_r(\bar{X}_{r-}^{t,x}, z) - \eta_r(X_{r-}^{t,x}, z) \right) \tilde{\pi}(dz, dr). \end{aligned}$$

Hence, by Minkowski's inequality, Jensen's inequality, Itô's isometry, and (5.82) we have for all $s \in [t, T]$ that

$$\begin{aligned} \left(\mathbb{E} \left[\left\| \bar{X}_{s \wedge \tau}^{t,x} - X_{s \wedge \tau}^{t,x} \right\|^2 \right] \right)^{1/2} &\leq \left(\int_t^s (s-t) \mathbb{E} \left[\mathbf{1}_{\{r \leq \tau\}} \left\| \mu(r, \bar{X}_r^{t,x}) - \mu(r, X_r^{t,x}) \right\|^2 \right] dr \right)^{1/2} \\ &\quad + \left(\int_t^s \mathbb{E} \left[\mathbf{1}_{\{r \leq \tau\}} \left\| \sigma(r, \bar{X}_r^{t,x}) - \sigma(r, X_r^{t,x}) \right\|_F^2 \right] dr \right)^{1/2} \\ &\quad + \left(\int_t^s \mathbb{E} \left[\int_{\mathbb{R}^d} \mathbf{1}_{\{r \leq \tau\}} \left\| \eta_r(\bar{X}_r^{t,x}, z) - \eta_r(X_r^{t,x}, z) \right\|_F^2 \nu(dz) \right] dr \right)^{1/2} \\ &\leq \left(\int_t^s (s-t) \mathbb{E} \left[\left\| \mu(r, \bar{X}_{r \wedge \tau}^{t,x}) - \mu(r, X_{r \wedge \tau}^{t,x}) \right\|^2 \right] dr \right)^{1/2} \\ &\quad + \left(\int_t^s \mathbb{E} \left[\left\| \sigma(r, \bar{X}_{r \wedge \tau}^{t,x}) - \sigma(r, X_{r \wedge \tau}^{t,x}) \right\|_F^2 \right] dr \right)^{1/2} \\ &\quad + \left(\int_t^s \mathbb{E} \left[\int_{\mathbb{R}^d} \left\| \eta_r(\bar{X}_{r \wedge \tau}^{t,x}, z) - \eta_r(X_{r \wedge \tau}^{t,x}, z) \right\|_F^2 \nu(dz) \right] dr \right)^{1/2}. \end{aligned}$$

Thus, by (2.3) and the fact that $(a+b+c)^2 \leq 3(a^2+b^2+c^2)$ for $a, b, c \in \mathbb{R}$, we obtain for all $s \in [t, T]$ that

$$\mathbb{E} \left[\left\| \bar{X}_{s \wedge \tau}^{t,x} - X_{s \wedge \tau}^{t,x} \right\|^2 \right] \leq 3(T+2)L \int_t^s \mathbb{E} \left[\left\| \bar{X}_{r \wedge \tau}^{t,x} - X_{r \wedge \tau}^{t,x} \right\|^2 \right] dr. \quad (5.85)$$

Moreover, by (2.9) it holds that

$$\mathbb{E} \left[\sup_{s \in [t, T]} \left\| X_s^{t,x} \right\|^2 \right] + \mathbb{E} \left[\sup_{s \in [t, T]} \left\| \bar{X}_s^{t,x} \right\|^2 \right] < \infty. \quad (5.86)$$

Then by (5.85), (5.86), and Grönwall's lemma we have for every $s \in [t, T]$ that

$$\mathbb{E} \left[\left\| \bar{X}_{s \wedge \tau}^{t,x} - X_{s \wedge \tau}^{t,x} \right\|^2 \right] = 0.$$

Hence, considering that $(\bar{X}_s^{t,x})_{s \in [t, T]}$ and $(X_s^{t,x})_{s \in [t, T]}$ have càdlàg sample paths, we obtain (5.84). Thus, the proof of this lemma is completed. \square

Proposition 5.15. *Let Assumptions 2.1, 2.2, 2.3, and 5.1 hold, and let $u : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ satisfy for all $(t, x) \in [0, T] \times \mathbb{R}^d$ that*

$$u(t, x) = \mathbb{E} \left[g(X_T^{t,x}) + \int_t^T h(s, X_s^{t,x}) ds \right]. \quad (5.87)$$

Then u is a viscosity solution of the linear PIDE (5.24) with $u(T, x) = g(x)$ for all $x \in \mathbb{R}^d$.

Proof. Throughout this proof, let $\chi \in C_c^\infty(\mathbb{R}^d)$ such that $\chi(x) = 1$ for $\|x\| \leq 1$, $0 \leq \chi(x) \leq 1$ for $1 < \|x\| < 2$, and $\chi(x) = 0$ for $\|x\| \geq 2$. Then for each $n \in \mathbb{N}$, $(t, x) \in [0, T] \times \mathbb{R}^d$, and $z \in \mathbb{R}^d$ define $\chi_n(x) := \chi(x/n)$ and

$$\begin{aligned} \mu^{(n)}(t, x) &:= \mu(t, x) \chi_n(x), & \sigma^{(n)}(t, x) &:= \sigma(t, x) \chi_n(x), & \eta_t^{(n)}(x, z) &:= \eta_t(x, z) \chi_n(x), \\ g^{(n)}(x) &:= g(x) \chi_n(x), & h^{(n)}(t, x) &:= h(t, x) \chi_n(x). \end{aligned}$$

For every $n \in \mathbb{N}$, $(t, x) \in [0, T] \times \mathbb{R}^d$ let $(X_s^{t,x,(n)})_{s \in [t, T]} : [t, T] \times \Omega \rightarrow \mathbb{R}^d$ be an \mathbb{F} -adapted càdlàg process satisfying that $X_t^{t,x} = x$, and almost surely for all $s \in [t, T]$

$$dX_s^{t,x,(n)} = \mu^{(n)}(s, X_{s-}^{t,x,(n)}) ds + \sigma^d(s, X_{s-}^{t,x,(n)}) dW_s + \int_{\mathbb{R}^d} \eta_s^{(n)}(X_{s-}^{t,x,(n)}, z) \tilde{\pi}(dz, ds).$$

For each $n \in \mathbb{N}$, $k \in \mathbb{N}$, and $(t, x) \in [0, T] \times \mathbb{R}^d$, we define

$$u^{n,k}(t, x) := \mathbb{E} \left[g^{(k)}(X_T^{t,x,(n)}) + \int_t^T h^{(k)}(s, X_s^{t,x,(n)}) ds \right], \quad (5.88)$$

and

$$u^{0,k}(t, x) := \mathbb{E} \left[g^{(k)}(X_T^{t,x}) + \int_t^T h^{(k)}(s, X_s^{t,x}) ds \right]. \quad (5.89)$$

By construction, for every $n \in \mathbb{N}$ we have that $\mu^{(n)}$, $\sigma^{(n)}$, $\eta^{(n)}$, $g^{(n)}$, and $h^{(n)}$ satisfy Assumption 5.6. In addition, for all $n \in \mathbb{N}$, $t \in [0, T]$, and $x, x' \in \mathbb{R}^d$ with $\|x\| \wedge \|x'\| \geq 2n$ we have

$$\|\mu^{(n)}(t, x) - \mu^{(n)}(t, x')\| = 0.$$

Moreover, by (2.3), (2.4), and the mean value theorem we have for all $n \in \mathbb{N}$, $t \in [0, T]$, and $x, x' \in \mathbb{R}^d$ with $\|x'\| \leq 2n$ that

$$\begin{aligned} \|\mu^{(n)}(t, x) - \mu^{(n)}(t, x')\|^2 &= \|\mu(t, x)\chi_n(x) - \mu(t, x')\chi_n(x')\|^2 \\ &\leq 2\|\mu(t, x)\chi_n(x) - \mu(t, x')\chi_n(x)\|^2 + 2\|\mu(t, x')(\chi_n(x) - \chi_n(x'))\|^2 \\ &\leq 2L\|x - x'\|^2 \cdot \sup_{y \in \mathbb{R}^d} (|\chi(y)|^2) + 2 \cdot \sup_{\|y\| \leq 2n} (\|\mu(t, y)\|^2) \cdot \sup_{y \in \mathbb{R}^d} (\|\nabla \chi_n(y)\|^2) \cdot \|x - x'\|^2 \\ &\leq 2L\|x - x'\|^2 + \frac{2Ld^p(1 + 4n^2)}{n^2} \cdot \sup_{y \in \mathbb{R}^d} (\|\nabla \chi(y)\|^2) \cdot \|x - x'\|^2 \\ &\leq 2L\|x - x'\|^2 \cdot \left[1 + 5d^p \cdot \sup_{y \in \mathbb{R}^d} (\|\nabla \chi(y)\|^2) \right]. \end{aligned}$$

Furthermore, by (2.4) we have for all $n \in \mathbb{N}$ and $(t, x) \in [0, T] \times \mathbb{R}^d$ that

$$\|\mu^{(n)}(t, x)\|^2 \leq \|\mu(t, x)\|^2 \leq Ld^p(1 + \|x\|^2).$$

Hence, by analogous arguments we obtain that $\mu^{(n)}$, $\sigma^{(n)}$, $\eta^{(n)}$, $g^{(n)}$, and $h^{(n)}$ satisfy Assumptions 5.7 and 5.8. Furthermore, by the product rule it holds for all $n \in \mathbb{N}$, $\gamma \in [0, 1]$, $(t, x) \in [0, T] \times \mathbb{R}^d$, and $z \in \mathbb{R}^d$ that

$$\gamma D_x \eta_t^{(n)}(x, z) = \gamma D_x (\eta_t(x, z) \chi_n(x)) = \gamma \chi_n(x) D_x \eta_t(x, z) + \gamma \eta_t(x, z) [\nabla \chi_n(x)]^T, \quad (5.90)$$

where $\eta_t(x, z) [\nabla \chi_n(x)]^T$ is understood as the matrix product of the column vector $\eta_t(x, z)$ and the row vector $[\nabla \chi_n(x)]^T$. By the definition of $\chi_n(\cdot)$, $n \in \mathbb{N}$, and Assumption 2.2 we also notice that it holds for all $n \in \mathbb{N}$, $(t, x) \in [0, T] \times \mathbb{R}^d$, and $z \in \mathbb{R}^d$ that $0 \leq \chi_n(x) \leq 1$ and

$$\|\eta_t(x, z) [\nabla \chi_n(x)]^T\|_F \leq \|\eta_t(x, z)\| \cdot \|\nabla \chi_n(x)\| \leq \frac{C_d^{1/2}}{n} \cdot \sup_{y \in \mathbb{R}^d} (\|\nabla \chi(y)\|), \quad (5.91)$$

where C_d is the positive constant introduced in (2.6). Moreover, by Assumption 2.2 we have for all $(t, x) \in [0, T] \times \mathbb{R}^d$, $z \in \mathbb{R}^d$, and $\gamma \in [0, 1]$ that

$$\|\gamma D_x \eta_t(x, z)\|_F \leq d^{1/2} C_d^{1/2}. \quad (5.92)$$

We also notice that the mapping

$$\left\{ B \in \mathbb{R}^{d \times d} : \|B\|_F \leq C_d^{1/2} (d^{1/2} + \sup_{y \in \mathbb{R}^d} (\|\nabla \chi(y)\|)) \right\} \ni A \mapsto |\det(\mathbf{I}_d + A)| \in \mathbb{R}$$

is uniformly continuous. Hence, (2.7) and (5.90)-(5.92) ensure that there exists an integer $N > 0$ such that for all $n \geq N$, $(t, x) \in [0, T] \times \mathbb{R}^d$, $\gamma \in [0, 1]$, and $z \in \mathbb{R}^d$

$$\frac{\lambda}{2} \leq |\det(\mathbf{I}_d + \gamma D_x \eta_t^{(n)}(x, z))|.$$

This implies that $\eta^{(n)}$ satisfies Assumption 5.9 with $\delta = \lambda/2$ for sufficiently large $n \in \mathbb{N}$. By the above discussion, applying Proposition 5.13 (with $\mu \curvearrowright \mu^{(n)}$, $\sigma \curvearrowright \sigma^{(n)}$, $\eta \curvearrowright \eta^{(n)}$, $g \curvearrowright g^{(k)}$, $h \curvearrowright h^{(k)}$ for $n, k \in \mathbb{N}$ in the notation of Proposition 5.13) yields that for all $k \in \mathbb{N}$ and sufficiently large $n \in \mathbb{N}$ the function $u^{n,k} \in C_1([0, T] \times \mathbb{R}^d)$ is a viscosity solution of

$$\begin{aligned} & \frac{\partial}{\partial t} u^{n,k}(t, x) + \langle \nabla_x u^{n,k}(t, x), \mu^{(n)}(t, x) \rangle + \frac{1}{2} \text{Trace} \left(\sigma^{(n)}(t, x) [\sigma^{(n)}(t, x)]^T \text{Hess}_x u^{n,k}(t, x) \right) \\ & + h^{(k)}(t, x) + \int_{\mathbb{R}^d} \left(u^{n,k}(t, x + \eta_t^{(n)}(x, z)) - u^{n,k}(t, x) - \langle \nabla_x u^{n,k}(t, x), \eta_t^{(n)}(x, z) \rangle \right) \nu(dz) = 0 \end{aligned} \quad (5.93)$$

on $(0, T) \times \mathbb{R}^d$ with $u^{n,k}(T, x) = g^{(k)}(x)$ on \mathbb{R}^d .

Next, for each $n \in \mathbb{N}$ and $(t, x) \in [0, T] \times \mathbb{R}^d$, let $\tau_n^{t,x} : \Omega \rightarrow [0, T]$ be the stopping time defined by

$$\tau_n^{t,x} := \inf \{s \geq t : \max(\|X_s^{t,x,(n)}\|, \|X_s^{t,x}\|) > n\} \wedge T. \quad (5.94)$$

Then (5.94) and Lemma 5.14 ensures that for all $(t, x) \in [0, T] \times \mathbb{R}^d$ and $n \in \mathbb{N}$

$$\mathbb{P} \left(\mathbf{1}_{\{s \leq \tau_n^{t,x}\}} \|X_s^{t,x,(n)} - X_s^{t,x}\| = 0 \text{ for all } s \in [t, T] \right) = 1. \quad (5.95)$$

Hence, considering that for every $n \in \mathbb{N}$ and $(t, x) \in [0, T] \times \mathbb{R}^d$ the processes $(X_s^{t,x,(n)})_{s \in [t, T]}$ and $(X_s^{t,x})_{s \in [t, T]}$ have no fixed time of discontinuity, we have for all $n, k \in \mathbb{N}$, and $(t, x) \in [0, T] \times \mathbb{R}^d$ that

$$\begin{aligned} \mathbb{E} \left[|g^{(k)}(X_T^{t,x,(n)}) - g^{(k)}(X_T^{t,x})| \right] &= \mathbb{E} \left[\mathbf{1}_{\{\tau_n^{t,x} < T\}} |g^{(k)}(X_T^{t,x,(n)}) - g^{(k)}(X_T^{t,x})| \right] \\ &\leq 2 \left[\sup_{y \in \mathbb{R}^d} |g^{(k)}(y)| \right] \mathbb{P}(\tau_n^{t,x} < T), \end{aligned} \quad (5.96)$$

and

$$\begin{aligned} & \int_t^T \mathbb{E} \left[|h^{(k)}(s, X_s^{t,x,(n)}) - h^{(k)}(s, X_s^{t,x})| \right] ds \\ &= \int_t^T \mathbb{E} \left[\mathbf{1}_{\{\tau_n^{t,x} < T\}} |h^{(k)}(s, X_s^{t,x,(n)}) - h^{(k)}(s, X_s^{t,x})| \right] ds \\ &\leq 2T \left[\sup_{(s,y) \in [0, T] \times \mathbb{R}^d} |h^{(k)}(s, y)| \right] \mathbb{P}(\tau_n^{t,x} < T). \end{aligned} \quad (5.97)$$

Furthermore, by (2.9), the construction of $\mu^{(n)}$, $\sigma^{(n)}$, and $\eta^{(n)}$, and Assumptions 5.7 and 5.8, it holds for all $n \in \mathbb{N}$ that

$$\mathbb{E} \left[\sup_{s \in [t, T]} \|X_s^{t,x,(n)}\|^2 \right] \leq C_3(1 + \|x\|^2), \quad (5.98)$$

where C_3 is a positive constant only depending on L, d, p and T . Thus, by (5.94), Chebyshev's inequality, and (2.9) it holds for all $(t, x) \in [0, T] \times \mathbb{R}^d$ and $n \in \mathbb{N}$ that

$$\begin{aligned} \mathbb{P}(\tau_n^{t,x} < T) &\leq \mathbb{P} \left(\|X_{\tau_n^{t,x}}^{t,x} \| \geq n \right) + \mathbb{P} \left(\|X_{\tau_n^{t,x}}^{t,x,(n)} \| \geq n \right) \\ &\leq \frac{\mathbb{E} \left[\|X_{\tau_n^{t,x}}^{t,x} \|^2 \right] + \mathbb{E} \left[\|X_{\tau_n^{t,x}}^{t,x,(n)} \|^2 \right]}{n^2} \\ &\leq \frac{(C + C_3)(1 + \|x\|^2)}{n^2}. \end{aligned} \quad (5.99)$$

Combining (5.96), (5.97), and (5.99), we have for all $(t, x) \in [0, T] \times \mathbb{R}^d$ and $n, k \in \mathbb{N}$ that

$$|u^{n,k}(t, x) - u^{0,k}(t, x)| \leq \frac{2(C + C_3)(1 + \|x\|^2)}{n^2} \left[\sup_{y \in \mathbb{R}^d} (|g^{(k)}(y)|) + T \cdot \sup_{(s,y) \in [0, T] \times \mathbb{R}^d} (|h^{(k)}(s, y)|) \right].$$

This implies that for all $k \in \mathbb{N}$ and all compact sets $\mathcal{K} \subseteq [0, T] \times \mathbb{R}^d$ it holds that

$$\lim_{n \rightarrow \infty} \left[\sup_{(t,x) \in \mathcal{K}} |u^{n,k}(t, x) - u^{0,k}(t, x)| \right] = 0. \quad (5.100)$$

Moreover, by (2.5), (2.9), (5.1), (5.35), (5.88), (5.89), (5.98), and the fact that $(a + b)^{1/2} \leq a^{1/2} + b^{1/2}$ for $a, b \geq 0$, we notice for all $n, k \in \mathbb{N}$ and $(t, x) \in [0, T] \times \mathbb{R}^d$ that

$$\begin{aligned} |u^{n,k}(t, x)| &\leq \mathbb{E} \left[|g^{(k)}(X_T^{t,x,(n)})| \right] + \int_t^T \mathbb{E} \left[|h^{(k)}(s, X_s^{t,x,(n)})| \right] ds \\ &\leq \left(\mathbb{E} \left[L(d^p + \|X_T^{t,x,(n)}\|^2) \right] \right)^{1/2} + \int_t^T \left(\mathbb{E} \left[C_0(1 + \|X_s^{t,x,(n)}\|^2) \right] \right)^{1/2} ds \\ &\leq L^{1/2}d^{p/2} + L^{1/2}C_3^{1/2}(1 + \|x\|^2)^{1/2} + C_0^{1/2}T[1 + C_3^{1/2}(1 + \|x\|^2)^{1/2}] \\ &\leq L^{1/2}d^{p/2} + C_0^{1/2}T + C_3^{1/2}(L^{1/2} + C_0^{1/2}T)(1 + \|x\|^2)^{1/2}, \end{aligned} \quad (5.101)$$

and

$$|u^{0,k}(t, x)| \leq L^{1/2}d^{p/2} + C_0^{1/2}T + C^{1/2}(L^{1/2} + C_0^{1/2}T)(1 + \|x\|^2)^{1/2}. \quad (5.102)$$

This implies that for all $k, n \in \mathbb{N}$

$$\sup_{(t,x) \in [0,T] \times \mathbb{R}^d} \frac{|u^{n,k}(t, x)| + |u^{0,k}(t, x)|}{1 + \|x\|} < \infty.$$

Hence, (5.100) and the fact that $u^{n,k} \in C_1([0, T] \times \mathbb{R}^d)$ for all $k \in \mathbb{N}$ and sufficiently large $n \in \mathbb{N}$ ensure that $u^{0,k} \in C_1([0, T] \times \mathbb{R}^d, \mathbb{R})$ for all $k \in \mathbb{N}$. Therefore, by (5.93), (5.100), Assumptions 2.1, 2.2, and Assumptions 5.1-5.8, applying Lemma 5.11 yields that for all $k \in \mathbb{N}$ the function $u^{0,k}$ is a viscosity solution of

$$\begin{aligned} &\frac{\partial}{\partial t} u^{0,k}(t, x) + \langle \nabla_x u^{0,k}(t, x), \mu(t, x) \rangle + \frac{1}{2} \text{Trace} \left(\sigma(t, x) [\sigma(t, x)]^T \text{Hess}_x u^{0,k}(t, x) \right) \\ &+ h^{(k)}(t, x) + \int_{\mathbb{R}^d} \left(u^{0,k}(t, x + \eta_t(x, z)) - u^{0,k}(t, x) - \langle \nabla_x u^{0,k}(t, x), \eta_t(x, z) \rangle \right) \nu(dz) = 0 \end{aligned} \quad (5.103)$$

on $(0, T) \times \mathbb{R}^d$ with $u^{0,k}(T, x) = g^{(k)}(x)$ on \mathbb{R}^d . Furthermore, Lemma 3.1 implies for all $k \in \mathbb{N}$ and $(t, x) \in [0, T] \times \mathbb{R}^d$ that

$$\begin{aligned} &\frac{\mathbb{E} \left[\int_t^T |h^{(k)}(s, X_s^{t,x}) - h(s, X_s^{t,x})| ds \right]}{d^p + \|x\|^2} \\ &= \int_t^T \mathbb{E} \left[\frac{|h^{(k)}(s, X_s^{t,x}) - h(s, X_s^{t,x})|}{d^p + \|X_s^{t,x}\|^2} \cdot \frac{d^p + \|X_s^{t,x}\|^2}{d^p + \|x\|^2} \right] ds \\ &\leq \int_t^T \left[\sup_{(r,y) \in [0,T] \times \mathbb{R}^d} \left(\frac{|h^{(k)}(r, y) - h(r, y)|}{d^p + \|y\|^2} \right) \right] \cdot \frac{\mathbb{E} \left[d^p + \|X_s^{t,x}\|^2 \right]}{d^p + \|x\|^2} ds \\ &\leq c_1 T \left[\sup_{(s,y) \in [0,T] \times \mathbb{R}^d} \left(\frac{|h^{(k)}(s, y) - h(s, y)|}{d^p + \|y\|^2} \right) \right], \end{aligned} \quad (5.104)$$

where $c_1 := (6LT + 1)e^{(1+6L)T}$. Moreover, by Assumption 5.1 it holds for all $k \in \mathbb{N}$ and $(s, y) \in [0, T] \times \mathbb{R}^d$ that

$$\begin{aligned} \frac{|h^{(k)}(s, y) - h(s, y)|}{d^p + \|y\|^2} &= \frac{\mathbf{1}_{\{\|y\| \geq k\}} |h(s, y)(\chi_k(y) - 1)|}{d^p + \|y\|^2} \\ &\leq \frac{2C_0^{1/2}(d^p + \|y\|^2)^{1/2}}{(d^p + k^2)^{1/2}(d^p + \|y\|^2)^{1/2}} = 2C_0^{1/2}(d^p + k^2)^{-1/2}. \end{aligned}$$

Combining this with (5.104) implies that

$$\lim_{k \rightarrow \infty} \left[\sup_{(t,x) \in [0,T] \times \mathbb{R}^d} \left(\frac{\mathbb{E} \left[\int_t^T |h^{(k)}(s, X_s^{t,x}) - h(s, X_s^{t,x})| ds \right]}{d^p + \|x\|^2} \right) \right] = 0. \quad (5.105)$$

Analogously, by Lemma 3.1 and (2.5) we have

$$\lim_{k \rightarrow \infty} \left[\sup_{(t,x) \in [0,T] \times \mathbb{R}^d} \left(\frac{\mathbb{E} \left[|g^{(k)}(X_s^{t,x}) - g(X_s^{t,x})| \right]}{d^p + \|x\|^2} \right) \right] = 0. \quad (5.106)$$

Then (5.88), (5.89) together with (5.105) and (5.106) imply that

$$\lim_{k \rightarrow \infty} \left[\sup_{(t,x) \in [0,T] \times \mathbb{R}^d} \left(\frac{|u^{0,k}(t,x) - u(t,x)|}{d^p + \|x\|^2} \right) \right] = 0.$$

Hence, for all non-empty compact set $\mathcal{K} \subseteq [0, T] \times \mathbb{R}^d$ it holds that

$$\begin{aligned} & \lim_{k \rightarrow \infty} \left[\sup_{(t,x) \in \mathcal{K}} |u^{0,k}(t,x) - u(t,x)| \right] \\ & \leq \lim_{k \rightarrow \infty} \left[\sup_{(t,x) \in \mathcal{K}} \left(\frac{|u^{0,k}(t,x) - u(t,x)|}{d^p + \|x\|^2} \right) \right] \cdot \left[\sup_{(t,x) \in \mathcal{K}} (d^p + \|x\|^2) \right] = 0. \end{aligned} \quad (5.107)$$

By (2.5), (2.9), (5.87), and the analogous argument to obtain (5.101) and (5.102) we have that u has linear growth. Then by (5.107) and the fact that $u^{0,k} \in C_1([0, T] \times \mathbb{R}^d)$ for every $k \in \mathbb{N}$, we have that $u \in C_1([0, T] \times \mathbb{R}^d)$. Therefore, by (5.103), (5.107), and Assumptions 2.1, 2.2, and 5.1-5.9, we apply Lemma 5.11 to get that u is a viscosity solution of the linear PIDE (5.24) with $u(T, x) = g(x)$ for all $x \in \mathbb{R}^d$. The proof of this proposition is thus completed. \square

Proposition 5.16. *Let Assumptions 2.1, 2.2, and 2.3 hold. Then the following holds:*

(i) *There exists a unique Borel function $u : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$ satisfying*

$$u(t, x) := \mathbb{E} \left[g(X_T^{t,x}) + \int_t^T f(s, X_s^{t,x}, u(s, X_s^{t,x})) ds \right], \quad (t, x) \in [0, T] \times \mathbb{R}^d. \quad (5.108)$$

(ii) *Let $u : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$ satisfy (5.108). Then $u \in C_1([0, T] \times \mathbb{R}^d)$.*

(iii) *Let $u : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$ satisfy (5.108). Then u is a viscosity solution of PIDE (2.18) with $u(T, x) = g(x)$ for all $x \in \mathbb{R}^d$.*

Proof. First recall $\Lambda := \{(t, s) \in [0, T]^2 : t \leq s\}$. Then we notice that by Corollary 3.5, the mapping $\Lambda \times \mathbb{R}^d \ni (t, s, x) \mapsto (s, X_s^{t,x}) \in \mathcal{L}_0(\Omega, [0, T] \times \mathbb{R}^d)$ is continuous, where $\mathcal{L}_0(\Omega, [0, T] \times \mathbb{R}^d)$ denotes the space of all measurable functions from Ω to $[0, T] \times \mathbb{R}^d$ equipped with the metric deduced by convergence in probability. Moreover, notice that for every nonnegative Borel function $\varphi : [0, T] \times \mathbb{R}^d \rightarrow [0, \infty)$, the mapping $\mathcal{L}_0(\Omega, [0, T] \times \mathbb{R}^d) \ni Z \mapsto \mathbb{E}[\varphi(Z)] \in [0, \infty]$ is measurable. Hence for all $\theta \in \Theta$, $d \in \mathbb{N}$, and all nonnegative Borel functions $\varphi : [0, T] \times \mathbb{R}^d \rightarrow [0, \infty)$ it holds that the mapping

$$\Lambda \times \mathbb{R}^d \ni (t, s, x) \mapsto \mathbb{E}[\varphi(s, X_s^{t,x})] \in [0, \infty) \quad (5.109)$$

is measurable. Furthermore, by Lemma 3.1, we observe that for all $(t, x) \in [0, T] \times \mathbb{R}^d$ and $s \in [t, T]$

$$\mathbb{E} \left[d^p + \|X_s^{t,x}\|^2 \right] \leq c_1 (d^p + \|x\|^2) e^{\rho_1(s-t)}, \quad (5.110)$$

where $c_1 := 1 + 6LT$ and $\rho_1 := 1 + 6L$. Then, combining (2.2), (2.5), (5.109), (5.110), and applying Proposition 4.1 (with $Z_s^{t,x} \curvearrowright X_s^{t,x}$, $a \curvearrowright \max\{L^{1/2}, L^{1/2}T^{-1/2}\}$, $b \curvearrowright c_1^{1/2} e^{\rho_1 T/2}$, $c \curvearrowright L^{1/2}$, $f \curvearrowright f$, $g \curvearrowright g$, $u \curvearrowright u$ in the notation of Proposition 4.1), we obtain that there is a unique Borel function $u : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$ satisfying (5.108) as well as for all $t \in [0, T]$ that

$$\sup_{x \in \mathbb{R}^d} \frac{|u(t, x)|}{\sqrt{d^p + \|x\|^2}} \leq 2L^{1/2} \exp \{L^{1/2}(T-t)\}. \quad (5.111)$$

Hence, we finish the proof of (i). Furthermore, for every $r \in (0, \infty)$ let $B_r \subseteq \mathbb{R}^d$ be the set defined by $B_r := \{x \in \mathbb{R}^d : \|x\| \leq r\}$. Then by (2.5) and (5.111) we have for all $(s, y) \in [0, T] \times \mathbb{R}^d$ that

$$\frac{|f(s, y, 0)| + |u(s, y)| + g(y)}{d^p + \|y\|^2} \leq \frac{T^{-1}L^{1/2}(d^p + \|y\|^2)^{1/2} + 2L^{1/2} \exp\{L^{1/2}T\}(d^p + \|y\|^2)^{1/2}}{d^p + \|y\|^2}$$

$$\begin{aligned}
& + \frac{L^{1/2}(d^p + \|y\|^2)^{1/2}}{d^p + \|y\|^2} \\
& = \frac{L^{1/2}(T^{-1} + 2 \exp\{L^{1/2}T\} + 1)}{(d^p + \|y\|^2)^{1/2}}.
\end{aligned}$$

This implies that

$$\begin{aligned}
& \inf_{r \in (0, \infty)} \left[\sup_{s \in [0, T]} \sup_{y \in \mathbb{R}^d / B_r} \frac{|f(s, y, 0)| + |u(s, y)| + |g(y)|}{d^p + \|y\|^2} \right] \\
& \leq \inf_{r \in (0, \infty)} \left[\sup_{s \in [0, T]} \sup_{y \in \mathbb{R}^d / B_r} \frac{L^{1/2}(T^{-1} + 2 \exp\{L^{1/2}T\} + 1)}{(d^p + \|y\|^2)^{1/2}} \right] \\
& = \inf_{r \in (0, \infty)} \left[\frac{L^{1/2}(T^{-1} + 2 \exp\{L^{1/2}T\} + 1)}{(d^p + \|r\|)^{1/2}} \right] = 0. \tag{5.112}
\end{aligned}$$

Then, by (2.2), (2.5), (5.110), (5.112), and Corollary 3.5, we can apply Corollary 2.7 in [9] (with $X^{t,x} \curvearrowright X^{t,x}$, $f \curvearrowright f$, $g \curvearrowright g$, and $V(t, x) \curvearrowright (d^p + \|x\|^2)$ in the notation of Corollary 2.7 in [9]) to get that $u : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$ is continuous (Noting that the condition $\mathbb{E}[V(s, X_s^{t,x})] \leq V(t, x)$ in Corollary 2.7 in [9] can be relaxed to $\mathbb{E}[V(s, X_s^{t,x})] \leq cV(t, x)$ for any $c \geq 1$). Combining this with (5.111), we obtain that $u \in C_1([0, T] \times \mathbb{R}^d)$. Thus, we obtain (ii). Next, we define $\mathbf{h} : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$ by

$$\mathbf{h}(t, x) := f(t, x, u(t, x)), \quad (t, x) \in [0, T] \times \mathbb{R}^d.$$

By (2.2), (2.5), and (5.111) it holds for all $(t, x) \in [0, T] \times \mathbb{R}^d$ that

$$\begin{aligned}
|\mathbf{h}(t, x)| & = |f(t, x, u(t, x))| \leq |f(t, x, u(t, x)) - f(t, x, 0)| + |f(t, x, 0)| \\
& \leq L^{1/2}|u(t, x)| + L^{1/2}T^{-1/2}(d^p + \|x\|^2)^{1/2} \\
& \leq L^{1/2}d^{p/2}(2L^{1/2} \exp\{L^{1/2}T\} + T^{-1/2})(1 + \|x\|^2)^{1/2}. \tag{5.113}
\end{aligned}$$

This implies that \mathbf{h} satisfies Assumption 5.1. Thus, Assumptions 2.1-2.3 and the fact that $u \in C_1([0, T] \times \mathbb{R}^d)$ satisfies (5.108) allow us to apply Proposition 5.15 to obtain that u is a viscosity solution of PIDE (2.18) with $u(T, x) = g(x)$ for all $x \in \mathbb{R}^d$, which proves (iii). The proof of this proposition is therefore completed. \square

6. MULTILEVEL PICARD (MLP) APPROXIMATIONS NOT INVOLVING EULER SCHEMES

6.1. Setting. We assume the settings in Section 2.1. Moreover, for each $d \in \mathbb{N}$ and $(t, x) \in [0, T] \times \mathbb{R}^d$ let $(Z_s^{d, \theta, t, x})_{s \in [t, T]} : [t, T] \times \Omega \rightarrow \mathbb{R}^d$, $\theta \in \Theta$, be $\mathcal{B}([0, T]) \otimes \mathcal{F}/\mathcal{B}(\mathbb{R}^d)$ -measurable functions. Assume for all $(t, x) \in [0, T] \times \mathbb{R}^d$ and $s \in [t, T]$ that $Z_s^{d, \theta, t, x}$, $\theta \in \Theta$, are independent and identically distributed, and assume that $(Z_s^{d, \theta, t, x})_{(\theta, s) \in \Theta \times [t, T]}$ and $(\xi^\theta)_{\theta \in \Theta}$ are independent. For each $d \in \mathbb{N}$, $(t, x) \in [0, T] \times \mathbb{R}^d$, $s \in [t, T]$, and $\theta \in \Theta$, let $(Z_s^{(d, \theta, t, x, l, i)})_{(l, i) \in \mathbb{N} \times \mathbb{Z}}$ be independent copies of $Z_s^{d, \theta, t, x}$. For each $d \in \mathbb{N}$, $n \in \mathbb{N}_0$, $M \in \mathbb{N}$, and $\theta \in \Theta$, let $\mathcal{U}_{n, M}^{d, \theta} : [0, T] \times \mathbb{R}^d \times \Omega \rightarrow \mathbb{R}$ be a measurable function satisfying for all $(t, x) \in [0, T] \times \mathbb{R}^d$ that

$$\begin{aligned}
\mathcal{U}_{n, M}^{d, \theta}(t, x) & = \frac{\mathbf{1}_{\mathbb{N}}(n)}{M^n} \sum_{i=1}^{M^n} g^d \left(Z_T^{(d, \theta, t, x, 0, -i)} \right) \\
& + \sum_{l=0}^{n-1} \frac{(T-t)}{M^{n-l}} \left[\sum_{i=1}^{M^{n-l}} \left(F \left(U_{l, M}^{(d, \theta, l, i)} \right) - \mathbf{1}_{\mathbb{N}}(l) F \left(U_{l-1, M}^{(d, \theta, -l, i)} \right) \right) \left(\mathcal{R}_t^{(\theta, l, i)}, Z_{\mathcal{R}_t^{(\theta, l, i)}}^{(d, \theta, t, x, l, i)} \right) \right],
\end{aligned}$$

where $(\mathcal{U}_{n, M}^{(d, \theta, l, i)})_{(l, i) \in (\mathbb{Z} \setminus \{0\}) \times \mathbb{N}_0}$ are independent copies of $\mathcal{U}_{n, M}^{d, \theta}$. Furthermore, let $b, c, \rho \in (0, \infty)$, and let $u^d : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$, $d \in \mathbb{N}$, be measurable functions such that for all $d \in \mathbb{N}$, $(t, x) \in [0, T] \times \mathbb{R}^d$, and $s \in [t, T]$

$$\begin{aligned}
\mathbb{E} \left[|g^d(Z_T^{d, 0, t, x})| \right] + \int_t^T \mathbb{E} \left[|(F^d(u^d))(r, Z_r^{d, 0, t, x})| \right] dr & < \infty, \\
\mathbb{E} \left[d^p + \|Z_s^{d, 0, t, x}\|^2 \right] & \leq b e^{\rho(s-t)} (d^p + \|x\|^2),
\end{aligned}$$

$$u^d(t, x) = \mathbb{E} \left[g(Z_T^{d,0,t,x}) + \int_t^T (F^d(u^d))(r, Z_r^{d,0,t,x}) dr \right],$$

and

$$|(F^d(0))(t, x)|^2 + |g^d(x)|^2 + |u^d(t, x)|^2 \leq c(d^p + \|x\|^2). \quad (6.1)$$

6.2. Error bounds for MLP approximations. The following lemma is taken from [39].

Lemma 6.1. *Assume Setting 6.1. Then for all $d, n, M \in \mathbb{N}$ and $t \in [0, T]$ it holds that*

$$\begin{aligned} \sup_{x \in \mathbb{R}^d} \left(\frac{\mathbb{E} \left[|\mathcal{U}_{n,M}^{d,0}(t, x) - u^d(t, x)|^2 \right]}{d^p + \|x\|^2} \right)^{1/2} &\leq 2e^{M/2} M^{-n/2} (1 + 2TL^{1/2})^{n-1} b^{1/2} e^{\rho(T-t)/2} \\ &\cdot \left(\sup_{t \in [0, T]} \sup_{x \in \mathbb{R}^d} \left(\frac{\max \{ |T \cdot (F^d(0))(t, x)|, |g^d(x)| \}}{\sqrt{d^p + \|x\|^2}} \right) + TL^{1/2} \sup_{t \in [0, T]} \sup_{x \in \mathbb{R}^d} \left(\frac{|u^d(t, x)|}{\sqrt{d^p + \|x\|^2}} \right) \right) \end{aligned} \quad (6.2)$$

Proof. For $b = 1$, the above statement is proved in Corollary 3.12 in [39] (with $c \curvearrowright c, \rho \curvearrowright \rho, L \curvearrowright L^{1/2}, \tau \curvearrowright t, U_{n,M}^\theta \curvearrowright \mathcal{U}_{n,M}^{d,\theta}, Y_{t,s}^\theta(x) \curvearrowright Z_s^{d,\theta,t,x}, \varphi \curvearrowright (\mathbb{R}^d \ni x \mapsto (d^p + \|x\|^2) \in \mathbb{R}), F \curvearrowright F^d, g \curvearrowright g^d, u \curvearrowright u^d$ in the notation of Corollary 3.12 in [39]). For the general case $b \in (0, \infty)$, one can precisely follow the proof of Corollary 3.12 in [39] to obtain (6.2). \square

7. PROOF OF THE MAIN RESULTS

Proof of Theorem 2.9. (i) is well-known (see, e.g., Theorem 3.1 in [47] and Theorem 117 in [55]). To prove (ii) we notice that by Corollary 3.5, for all $\theta \in \Theta$ and $d \in \mathbb{N}$ the mapping $\Lambda \times \mathbb{R}^d \ni (t, s, x) \mapsto (s, X_s^{d,\theta,t,x}) \in \mathcal{L}_0(\Omega, [0, T] \times \mathbb{R}^d)$ is continuous, where $\mathcal{L}_0(\Omega, [0, T] \times \mathbb{R}^d)$ denotes the metric space of all measurable functions from Ω to $[0, T] \times \mathbb{R}^d$ equipped with the metric deduced by convergence in probability. Moreover, notice that for all $d \in \mathbb{N}$ and nonnegative Borel functions $\varphi : [0, T] \times \mathbb{R}^d \rightarrow [0, \infty)$, the mapping $\mathcal{L}_0(\Omega, [0, T] \times \mathbb{R}^d) \ni Z \mapsto \mathbb{E}[\varphi(Z)] \in [0, \infty)$ is measurable. Hence for all $\theta \in \Theta, d \in \mathbb{N}$, and all nonnegative Borel functions $\varphi : [0, T] \times \mathbb{R}^d \rightarrow [0, \infty)$ it holds that the mapping

$$\Lambda \times \mathbb{R}^d \ni (t, s, x) \mapsto \mathbb{E} \left[\varphi(s, X_s^{d,\theta,t,x}) \right] \in [0, \infty) \quad (7.1)$$

is measurable. Analogously, considering Corollary 3.11, we obtain for all $\theta \in \Theta, d \in \mathbb{N}, N \in \mathbb{N}$ and all nonnegative Borel functions $\varphi : [0, T] \times \mathbb{R}^d \rightarrow [0, \infty)$ that the mapping

$$\Lambda \times \mathbb{R}^d \ni (t, s, x) \mapsto \mathbb{E} \left[\varphi(s, Y_s^{d,\theta,t,x,N,\delta,\mathcal{M}}) \right] \in [0, \infty) \quad (7.2)$$

is measurable. Furthermore, by Lemma 3.1 and Lemma 3.2, we observe that for all $d \in \mathbb{N}, \theta \in \Theta, t \in [0, T], s \in [t, T]$, and $x, y \in \mathbb{R}^d$

$$\mathbb{E} \left[d^p + \|X_s^{d,\theta,t,x}\|^2 \right] \leq c_1(d^p + \|x\|^2) e^{\rho_1(s-t)}, \quad (7.3)$$

and

$$\mathbb{E} \left[\|X_s^{d,\theta,t,x} - X_s^{d,\theta,t,y}\|^2 \right] \leq 4\|x - y\|^2 e^{\rho_2(s-t)}, \quad (7.4)$$

where $c_1 := 1 + 6LT, \rho_1 := 1 + 6L$, and $\rho_2 := 4L(T + 2)$. Moreover, (3.37) in Lemma 3.8 ensures for all $\theta \in \Theta, d \in \mathbb{N}, \delta \in (0, 1), (t, x) \in [0, T] \times \mathbb{R}^d, s \in [t, T]$, and $N, \mathcal{M} \in \mathbb{N}$ with $\mathcal{M} \geq \delta^{-2} K d^p$ that

$$\mathbb{E} \left[d^p + \|Y_s^{d,\theta,t,x,N,\delta,\mathcal{M}}\|^2 \right] \leq c_3(d^p + \|x\|^2) e^{\rho_3(s-t)}, \quad (7.5)$$

where $c_3 := 5 \cdot \max\{1, 4LT(T + 8)\}$, and $\rho_3 := 20L(T + 8)$. Then, combining (2.2), (2.5), (7.1), (7.3), and applying Proposition 4.1 (with $Z_s^{t,x} \curvearrowright X_s^{d,0,t,x}, a \curvearrowright \max\{L^{1/2}, L^{1/2}T^{-1/2}\}, b \curvearrowright c_1^{1/2} e^{\rho_1 T/2}, c \curvearrowright L^{1/2}, f \curvearrowright f^d, g \curvearrowright g^d, u \curvearrowright u^d$ in the notation of Proposition 4.1), we obtain (ii) and

$$\frac{|u^d(t, x)|}{\sqrt{d^p + \|x\|^2}} \leq 2L^{1/2} \exp \{L^{1/2}(T - t)\}. \quad (7.6)$$

Analogously, combining (2.2), (2.5), (7.2), and (7.5), and applying Proposition 4.1 (with $Z_s^{t,x} \curvearrowright Y_s^{d,0,t,x,N,\delta,\mathcal{M}}, b \curvearrowright c_3^{1/2} e^{\rho_3 T/2}, c \curvearrowright L^{1/2}, f \curvearrowright f^d, g \curvearrowright g^d, u \curvearrowright u_{N,\delta,\mathcal{M}}^d$ in the notation of Proposition

4.1), we obtain for each $d \in \mathbb{N}$, $\delta \in (0, 1)$, and $N, \mathcal{M} \in \mathbb{N}$ with $\mathcal{M} \geq \delta^{-2} K d^p$ that there exists a unique Borel function $u_{N,\delta,\mathcal{M}}^d : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$ satisfying for all $(t, x) \in [0, T] \times \mathbb{R}^d$ that

$$\mathbb{E} \left[|g^d(Y_T^{d,0,t,x,N,\delta,\mathcal{M}})| \right] + \int_t^T \mathbb{E} \left[|f^d(s, Y_s^{d,0,t,x,N,\delta,\mathcal{M}}, u_{N,\delta,\mathcal{M}}^d(s, Y_s^{d,0,t,x,N,\delta,\mathcal{M}}))| \right] ds < \infty, \quad (7.7)$$

$$\sup_{y \in \mathbb{R}^d} \sup_{s \in [0, T]} \left(\frac{|u_{N,\delta,\mathcal{M}}^d(s, y)|}{\sqrt{d^p + \|y\|^2}} \right) < \infty, \quad (7.8)$$

$$u_{N,\delta,\mathcal{M}}^d(t, x) = \mathbb{E} \left[g^d(Y_T^{d,0,t,x,N,\delta,\mathcal{M}}) \right] + \int_t^T \mathbb{E} \left[f^d(s, Y_s^{d,0,t,x,N,\delta,\mathcal{M}}, u_{N,\delta,\mathcal{M}}^d(s, Y_s^{d,0,t,x,N,\delta,\mathcal{M}})) \right] ds, \quad (7.9)$$

$$\frac{|u_{N,\delta,\mathcal{M}}^d(t, x)|}{\sqrt{d^p + \|x\|^2}} \leq 2L^{1/2} \exp \{L^{1/2}(T - t)\}. \quad (7.10)$$

Furthermore, we notice that Proposition 5.4 ensures that for each $d \in \mathbb{N}$ there exists at most one viscosity solution $v^d \in C_1([0, T] \times \mathbb{R}^d)$. Hence, combining this with Proposition 5.16 we obtain (iii) and (iv).

Next, we start to prove (v). We first observe that by (3.16), (3.34) and (3.38), there exist a positive constant $c_4 = c_4(T, L, L_1, L_2, K)$ satisfying for all $\theta \in \Theta$, $d \in \mathbb{N}$, $\delta \in (0, 1)$, $(t, x) \in [0, T] \times \mathbb{R}^d$, $s \in [t, T]$, and $N, \mathcal{M} \in \mathbb{N}$ with $\mathcal{M} \geq \delta^{-2} K d^p$ that

$$\mathbb{E} \left[\sup_{s \in [t, T]} \left\| X_s^{d,\theta,t,x} - Y_s^{d,\theta,t,x,N,\delta,\mathcal{M}} \right\|^2 \right] \leq c_4(d^p + \|x\|^2) (N^{-1}T + \delta^q d^p + \delta^{-2} K d^p \mathcal{M}^{-1}). \quad (7.11)$$

Furthermore, we notice that it holds for all $d \in \mathbb{N}$, $\delta \in (0, 1)$, $n, M, N, \mathcal{M} \in \mathbb{N}$, and $(t, x) \in [0, T] \times \mathbb{R}^d$ that

$$\begin{aligned} \left(\mathbb{E} \left[|U_{n,M,N}^{d,0,\delta,\mathcal{M}}(t, x) - u^d(t, x)|^2 \right] \right)^{1/2} &\leq \left(\mathbb{E} \left[|U_{n,M,N}^{d,0,\delta,\mathcal{M}}(t, x) - u_{N,\delta,\mathcal{M}}^d(t, x)|^2 \right] \right)^{1/2} \\ &\quad + |u_{N,\delta,\mathcal{M}}^d(t, x) - u^d(t, x)|. \end{aligned} \quad (7.12)$$

Considering (2.5), (7.5), (7.7)-(7.10), and the fact that $1 + a \leq e^a$ for all $a \in \mathbb{R}$, and applying Lemma 6.1 (with $\mathcal{U}_{n,M}^{d,0} \curvearrowright U_{n,M,N}^{d,0,\delta,\mathcal{M}}$, $Z_s^{d,0,t,x} \curvearrowright Y_s^{d,0,t,x,N,\delta,\mathcal{M}}$, $b \curvearrowright c_3$, $\rho \curvearrowright \rho_3$, $u^d \curvearrowright u_{N,\delta,\mathcal{M}}^d$, $c \curvearrowright 3 \cdot (4L \exp\{2L^{1/2}T\} \vee T^{-2}L \vee L)$ in the notation of Lemma 6.1), we have for all $d \in \mathbb{N}$, $\delta \in (0, 1)$, $n \in \mathbb{N}_0$, $t \in [0, T]$, and $M, N, \mathcal{M} \in \mathbb{N}$ with $\mathcal{M} \geq \delta^{-2} K d^p$ that

$$\begin{aligned} \sup_{x \in \mathbb{R}^d} \left(\frac{\mathbb{E} \left[|U_{n,M,N}^{d,0,\delta,\mathcal{M}}(t, x) - u_{N,\delta,\mathcal{M}}^d(t, x)|^2 \right]}{d^p + \|x\|^2} \right)^{1/2} &\leq 2e^{M/2} M^{-n/2} (1 + 2TL^{1/2})^{n-1} c_3^{1/2} e^{\rho_3(T-t)/2} \\ &\cdot \left[\sup_{t \in [0, T]} \sup_{x \in \mathbb{R}^d} \left(\frac{\max\{T(F^d(0)(t, x)), |g^d(x)|\}}{\sqrt{d^p + \|x\|^2}} \right) + TL^{1/2} \sup_{t \in [0, T]} \sup_{x \in \mathbb{R}^d} \left(\frac{u_{N,\delta,\mathcal{M}}^d(t, x)}{\sqrt{d^p + \|x\|^2}} \right) \right] \\ &\leq 2e^{M/2} M^{-n/2} (1 + 2TL^{1/2})^{n-1} c_3^{1/2} e^{\rho_3(T-t)/2} \left(L^{1/2} + 2TL \exp\{L^{1/2}(T - t)\} \right) \\ &\leq 2e^{M/2} M^{-n/2} (1 + 2TL^{1/2})^{n-1} e^{\rho_3(T-t)/2} (c_3 L)^{1/2} \left(1 + 2TL^{1/2} \exp\{L^{1/2}(T - t)\} \right) \\ &\leq C_1 e^{M/2} M^{-n/2} e^{2nTL^{1/2}}, \end{aligned} \quad (7.13)$$

where $C_1 := 2e^{\rho_3(T-t)/2} (c_3 L)^{1/2} (1 + 2TL^{1/2} \exp\{L^{1/2}T\})$. To deal with the second term in (7.12), we notice that by (7.4) and (7.11), for all $d \in \mathbb{N}$, $\delta \in (0, 1)$, $(t, x) \in [0, T] \times \mathbb{R}^d$, $s \in [t, T]$, $r \in [s, T]$ and $N, \mathcal{M} \in \mathbb{N}$ with $\mathcal{M} \geq \delta^{-2} K d^p$ it holds that

$$\begin{aligned} &\mathbb{E} \left[\mathbb{E} \left[\left\| X_r^{d,0,s,x'} - X_r^{d,0,s,y'} \right\|^2 \middle| (x', y') = (X_s^{d,0,t,x}, Y_s^{d,0,t,x,N,\delta,\mathcal{M}}) \right] \right] \\ &\leq 4e^{\rho_2(r-s)} \mathbb{E} \left[\left\| X_s^{d,0,t,x} - Y_s^{d,0,t,x,N,\delta,\mathcal{M}} \right\|^2 \right] \\ &\leq 4c_4(d^p + \|x\|^2) e^{\rho_2(r-s)} (N^{-1}T + \delta^q d^p + \delta^{-2} K d^p \mathcal{M}^{-1}). \end{aligned}$$

Moreover, by (7.4) we notice that for each $d \in \mathbb{N}$, $s \in [0, T]$, and $r \in [s, T]$ the mapping $\mathbb{R}^d \times \mathbb{R}^d \ni (x, y) \mapsto (X_r^{d,0,s,x}, X_r^{d,0,s,y}) \in \mathcal{L}_0(\Omega, \mathbb{R}^d \times \mathbb{R}^d)$ is continuous and hence measurable, and we have for all nonnegative Borel functions $h : \mathbb{R}^d \times \mathbb{R}^d \rightarrow [0, \infty)$ that the mapping $\mathcal{L}_0(\Omega, \mathbb{R}^d \times \mathbb{R}^d) \ni Z \mapsto \mathbb{E}[\varphi(Z)] \in [0, \infty]$ is measurable. Hence, it holds for all $d \in \mathbb{N}$, $s \in [0, T]$, $r \in [s, T]$ and all nonnegative Borel functions $h : \mathbb{R}^d \times \mathbb{R}^d \rightarrow [0, \infty)$ that the mapping

$$\mathbb{R}^d \times \mathbb{R}^d \ni (x, y) \mapsto \mathbb{E}\left[h\left(X_r^{d,0,s,x}, X_r^{d,0,s,y}\right)\right] \in [0, \infty] \quad (7.14)$$

is measurable. Furthermore, Lemma 2.2 in [41] ensures that for all $d \in \mathbb{N}$, $t \in [0, T]$, $s \in [t, T]$, $r \in [s, T]$, $x, y \in \mathbb{R}^d$ and all nono-negative Borel functions $h : \mathbb{R}^d \times \mathbb{R}^d \rightarrow [0, \infty)$ it holds that

$$\mathbb{E}\left[\mathbb{E}\left[h\left(X_r^{d,0,s,x'}, X_r^{d,0,s,y'}\right)\right] \Big|_{(x',y')=(X_s^{d,0,t,x}, X_s^{d,0,t,y})}\right] = \mathbb{E}\left[h\left(X_r^{d,0,t,x}, X_r^{d,0,t,y}\right)\right]. \quad (7.15)$$

Then using (2.2), (7.3), (7.5), (7.11), (7.6), (7.7)-(7.10), (7.14), (7.15), and (iv), and applying Lemma 4.2 with $T \curvearrowright T$, $L \curvearrowright L^{1/2}$, $X^{t,x,1} \curvearrowright X^{d,0,t,x}$, $X^{t,x,2} \curvearrowright Y^{d,0,t,x,N,\delta,\mathcal{M}}$, $\eta \curvearrowright (c_1 \vee c_3)e^{(\rho_1 \vee \rho_3)T}$, $\delta \curvearrowright [4c_4e^{\rho_2 T}(N^{-1}T + \delta^q d^p + \delta^{-2}Kd^p\mathcal{M}^{-1})]^{1/2}$, $c \curvearrowright c_0 := c_1 \vee c_3 \vee (2L^{1/2} \exp\{L^{1/2}T\})$, $\rho \curvearrowright \rho_1 \vee \rho_3$ in the notation of Lemma 4.2, we obtain for all $d, N \in \mathbb{N}$, $\delta \in (0, 1)$, $(t, x) \in [0, T] \times \mathbb{R}^d$, and $\mathcal{M} \in \mathbb{N}$ with $\mathcal{M} \geq \delta^{-2}Kd^p$ that

$$\begin{aligned} & |u_{N,\delta,\mathcal{M}}^d(t, x) - u^d(t, x)| \\ & \leq 4c_0[4c_4e^{\rho_2 T}(N^{-1}T + \delta^q d^p + \delta^{-2}Kd^p\mathcal{M}^{-1})]^{1/2}(TL)^{1/2}(1 + L^{1/2}T)(d^p + \|x\|^2) \\ & \quad \cdot \exp\left\{\left(L^{1/2} + (\rho_1 \vee \rho_3)/2 + (Lc_0)^{1/2}(c_1 \vee c_3)^{1/2}e^{(\rho_1 \vee \rho_3)T/2}\right)(T - t)\right\} \\ & \leq C_2(d^p + \|x\|^2)(N^{-1}T + \delta^q d^p + \delta^{-2}Kd^p\mathcal{M}^{-1})^{-1/2}, \end{aligned} \quad (7.16)$$

where

$$\begin{aligned} C_2 := & \exp\left\{\left(L^{1/2} + (\rho_1 \vee \rho_3)/2 + (Lc_0)^{1/2}(c_1 \vee c_3)^{1/2}e^{(\rho_1 \vee \rho_3)T/2}\right)T\right\} \\ & \cdot 8c_0(c_4TLe^{\rho_2 T})^{1/2}(1 + L^{1/2}T). \end{aligned}$$

Hence, combining (7.12), (7.13), and (7.16) yields for for all $d \in \mathbb{N}$, $\delta \in (0, 1)$, $n \in \mathbb{N}_0$, $(t, x) \in [0, T] \times \mathbb{R}^d$, and $M, N, \mathcal{M} \in \mathbb{N}$ with $\mathcal{M} \geq \delta^{-2}Kd^p$ that

$$\begin{aligned} & \left(\mathbb{E}\left[|U_{n,M,N}^{d,0,\delta,\mathcal{M}}(t, x) - u^d(t, x)|^2\right]\right)^{1/2} \\ & \leq (C_1 \vee C_2)(d^p + \|x\|^2) \left[e^{M/2}M^{-n/2}e^{2nTL^{1/2}} + (N^{-1}T + \delta^q d^p + \delta^{-2}Kd^p\mathcal{M}^{-1})^{1/2}\right], \end{aligned}$$

which proves (v) with $c = C_1 \vee C_2$. Thus, the proof of this theorem is complete. \square

Proof of Theorem 2.10. Applying (v) in Theorem 4.2 in [39] with $n \curvearrowright n$, $M \curvearrowright M$, $\mathfrak{C}_{n,M} \curvearrowright \mathfrak{C}_{n,M}^{(d)}$, $\mathfrak{m} \curvearrowright \mathfrak{e}^{(d)}$, $\mathfrak{g} \curvearrowright \mathfrak{g}^{(d)}$, $\mathfrak{f} \curvearrowright \mathfrak{f}^{(d)}$ in the notation of Theorem 4.2 in [39], we obtain (2.22). Next, for each $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$, and $x \in \mathbb{R}^d$ define $\mathbf{n}^d(x, \varepsilon)$ by

$$\mathbf{n}^d(x, \varepsilon) := \inf\{n \in \mathbb{N} \cap [2, \infty) : \sup_{k \in [n, \infty) \cap \mathbb{N}} \sup_{t \in [0, T]} \mathbb{E}\left[|U_{(k)}^{d,0}(t, x) - u^d(t, x)|^2\right] < \varepsilon^2\}, \quad (7.17)$$

where we use the convention $\inf(\emptyset) = \infty$ (\emptyset denotes the empty set), and use the shorter notation

$$U_{(k)}^{d,0}(t, x) := U_{k,k,k^k}^{d,0,k^{-k/q},k^{k+2k/q}Kd^p}(t, x), \quad \forall d \in \mathbb{N}, k \in \mathbb{N}, (t, x) \in [0, T] \times \mathbb{R}^d.$$

Applying (v) in Theorem 2.9 (with $n \curvearrowright n$, $M \curvearrowright n$, $N \curvearrowright n^n$, $\delta \curvearrowright n^{-n/q}$, $\mathcal{M} \curvearrowright n^{n+2n/q}Kd^p$ in the notation of Theorem 2.9), we have for all $d \in \mathbb{N}$, $n \in \mathbb{N}$, and $(t, x) \in [0, T] \times \mathbb{R}^d$ that

$$\begin{aligned} & \left(\mathbb{E}\left[|U_{(n)}^{d,0}(t, x) - u^d(t, x)|^2\right]\right)^{1/2} \\ & \leq c(d^p + \|x\|^2) \left[e^{n/2}n^{-n/2}e^{2nTL^{1/2}} + (n^{-n} \cdot T + d^p n^{-n} + n^{-n})^{1/2}\right], \end{aligned} \quad (7.18)$$

where $c = c(T, L, L_1, L_2, K)$ is the constant introduced in (v) in Theorem 2.9. Moreover, we observe for all integers $n \geq 2 \exp\{4TL^{1/2} + 1\}$ that

$$e^{n/2} e^{2nTL^{1/2}} n^{-n/2} = \exp\left\{(4TL^{1/2} + 1) \cdot n/2\right\} \cdot n^{-n/2} \leq 2^{-n/2},$$

which implies that

$$\lim_{n \rightarrow \infty} e^{n/2} e^{2nTL^{1/2}} n^{-n/2} = 0.$$

Therefore, by (7.18) we have for all $d \in \mathbb{N}$, $n \in \mathbb{N}$, $\varepsilon \in (0, 1]$, and $(t, x) \in [0, T] \times \mathbb{R}^d$ that

$$\mathbf{n}^d(x, \varepsilon) < \infty \quad \text{and} \quad \sup_{n \in [\mathbf{n}^d(x, \varepsilon), \infty) \cap \mathbb{N}} \left(\mathbb{E} \left[|U_{(n)}^{d,0}(t, x) - u^d(t, x)|^2 \right] \right)^{1/2} < \varepsilon,$$

which establishes (2.23). Furthermore, by (2.22) and (7.18) we obtain for all $d, n \in \mathbb{N}$, $\gamma \in (0, 1]$, and $(t, x) \in [0, T] \times \mathbb{R}^d$ that

$$\begin{aligned} & \left(\sum_{k=1}^{n+1} \mathfrak{C}_{k,k}^{(d)} \right) \left(\mathbb{E} \left[|U_{(n)}^{d,0}(t, x) - u^d(t, x)|^2 \right] \right)^{\frac{\gamma+4}{2}} \\ & \leq 12(3\mathfrak{e}^{(d)} + \mathfrak{g}^{(d)} + 2\mathfrak{f}^{(d)}) 36^n n^{2n} [c(d^p + \|x\|^2)]^{\gamma+4} (e^{n/2} n^{-n/2} e^{2nTL^{1/2}} + (1+T+d^p)^{1/2} n^{-n/2})^{\gamma+4} \\ & = 12(3\mathfrak{e}^{(d)} + \mathfrak{g}^{(d)} + 2\mathfrak{f}^{(d)}) 36^n n^{-\gamma n/2} [c(d^p + \|x\|^2)]^{\gamma+4} (e^{n/2} e^{2nTL^{1/2}} + (1+T+d^p)^{1/2})^{\gamma+4}. \end{aligned} \quad (7.19)$$

Then (7.17) and (7.19) show for all $d \in \mathbb{N}$, $\varepsilon, \gamma \in (0, 1]$ and $(t, x) \in [0, T] \times \mathbb{R}^d$ that

$$\begin{aligned} \left(\sum_{k=1}^{\mathbf{n}^d(x, \varepsilon)} \mathfrak{C}_{k,k}^{(d)} \right) \varepsilon^{\gamma+4} & \leq \left(\sum_{k=1}^{\mathbf{n}^d(x, \varepsilon)} \mathfrak{C}_{k,k}^{(d)} \right) \left(\mathbb{E} \left[|U_{(\mathbf{n}^d(x, \varepsilon)-1)}^{d,0}(t, x) - u^d(t, x)|^2 \right] \right)^{\frac{\gamma+4}{2}} \\ & \leq 12(3\mathfrak{e}^{(d)} + \mathfrak{g}^{(d)} + 2\mathfrak{f}^{(d)}) [c(d^p + \|x\|^2)]^{\gamma+4} \\ & \quad \cdot \sup_{n \in \mathbb{N}} \left(36^n n^{-\gamma n/2} (e^{n/2} e^{2nTL^{1/2}} + (1+T+d^p)^{1/2})^{\gamma+4} \right). \end{aligned}$$

Moreover, note that we have for all $\gamma \in (0, 1]$ and $n \in \mathbb{N}$ satisfying

$$n \geq \left(2 \cdot 6^4 \cdot \exp\{(1 + 4TL^{1/2})(\gamma + 4)\} \right)^{1/\gamma}$$

that

$$\begin{aligned} 36^n n^{-\gamma n/2} (e^{n/2} e^{2nTL^{1/2}})^{\gamma+4} & = 6^{4n/2} \cdot \exp\{(1 + 4TL^{1/2})(\gamma + 4)n/2\} \cdot n^{-\gamma n/2} \\ & = \left[6^4 \cdot \exp\{(1 + 4TL^{1/2})(\gamma + 4)\} \right]^{n/2} \cdot n^{-\gamma n/2} \\ & \leq 2^{-n/2}, \end{aligned}$$

which implies for all $\gamma \in (0, 1]$ that

$$\sup_{n \in \mathbb{N}} \left(36^n n^{-\gamma n/2} (e^{n/2} e^{2nTL^{1/2}} + (1+T+d^p)^{1/2})^{\gamma+4} \right) < \infty.$$

Hence, the proof of this theorem is complete. \square

REFERENCES

- [1] R.A. Adams and J.J.F. Fourier. *Sobolev Spaces Second Edition*. Elsevier, 2003.
- [2] O. Alvarez and A. Tourin. Viscosity solutions of nonlinear integro-differential equations. *Annales de l'Institut Henri Poincaré C, Analyse non linéaire*, 13(3):293–317, 1996.
- [3] D. Applebaum. *Lévy Processes and Stochastic Calculus*. Cambridge University Press, 2009.
- [4] A. Al-Arabi, A. Correia, D.D.F. Naiff, G. Jardim, and Y. Saporito. Extensions of the deep galerkin method. *arXiv preprint arXiv:1912.01455*, 2019.
- [5] G. Barles, R. Buckdahn, and E. Pardoux. Backward stochastic differential equations and integral-partial differential equations. *Stochastics: An International Journal of Probability and Stochastic Processes*, 60(1-2):57–83, 1997.

- [6] G. Barles, and C. Imbert. Second-order elliptic integro-differential equations: viscosity solutions' theory revisited. *In Annales de l'IHP Analyse non linéaire*, 25(3):567–585, 2008.
- [7] C. Beck, S. Becker, P. Cheridito, A. Jentzen, and A. Neufeld. Deep splitting method for parabolic pdes. *SIAM Journal on Scientific Computing*, 43(5):A3135–A3154, 2021.
- [8] S. Becker, R. Braunwarth, M. Hutzenthaler, A. Jentzen, and P. von Wurstemberger. Numerical simulations for full history recursive multilevel picard approximations for systems of high-dimensional partial differential equations. *arXiv preprint arXiv:2005.10206*, 2020.
- [9] C. Beck, L. Gonon, M. Hutzenthaler, and A. Jentzen. On existence and uniqueness properties for solutions of stochastic fixed point equations. *arXiv preprint arXiv:1908.03382*, 2019.
- [10] C. Beck, L. Gonon, and A. Jentzen. Overcoming the curse of dimensionality in the numerical approximation of high-dimensional semilinear elliptic partial differential equations. *arXiv preprint arXiv:2003.00596*, 2020.
- [11] C. Beck, F. Hornung, M. Hutzenthaler, A. Jentzen, and T. Kruse. Overcoming the curse of dimensionality in the numerical approximation of allen–cahn partial differential equations via truncated full-history recursive multilevel picard approximations. *Journal of Numerical Mathematics*, 28(4):197–222, 2020.
- [12] C. Beck, M. Hutzenthaler, and A. Jentzen. On nonlinear Feynman-Kac formulas for viscosity solutions of semilinear parabolic partial differential equations. *arXiv preprint arXiv:2004.03389v2*, 2020.
- [13] V. Boussange, S. Becker, A. Jentzen, B. Kuckuck, and L. Pellissier. Deep learning approximations for non-local nonlinear pdes with neumann boundary conditions. *arXiv preprint arXiv:2205.03672*, 2022.
- [14] T.R. Cass and R.K. Friz. The Bismut-Elworthy-Li formula for jump-diffusions and applications to Monte Carlo methods in finance. *arXiv preprint arXiv:math/0604311*, 2007.
- [15] J. Castro. Deep learning schemes for parabolic nonlocal integro-differential equations. *arXiv preprint arXiv:2103.15008*, 2021.
- [16] S.N. Cohen and R.J. Elliott. *Stochastic calculus and applications* (Vol. 2). New York: Birkhäuser, 2015.
- [17] R. Cont and P. Tankov. *Financial modelling with jump processes*. Chapman & Hall/CRC Financial Mathematics Series. Chapman & Hall/CRC, Boca Raton, FL, 2004.
- [18] R. Cont and E. Voltchkova. Integro-differential equations for option prices in exponential Lévy models. *Finance and Stochastics*, 9(3):299–325, 2005.
- [19] P. Cont and E. Voltchkova. A finite difference scheme for option pricing in jump diffusion and exponential Lévy models. *SIAM Journal on Numerical Analysis*, 43(4):1596–1626, 2005.
- [20] M.G. Crandall., H. Ishii, and P.L. Lions. User's guide to viscosity solutions of second order partial differential equations. *Bulletin of the American mathematical society*, 27(1):1–67, 1992.
- [21] M. De León-Contreras, I. Gyöngy, and S. Wu. On Solvability of Integro-differential Equations. *Potential Analysis*, 55(3):443–475, 2021.
- [22] C. Dellacherie and P.A. Meyer, *Probabilities and Potential B: Theory of Martingales, Translation of Probabilités et Potentiel B*. North-Holland, 1982.
- [23] Ł. Delong. *Backward stochastic differential equations with jumps and their actuarial and financial applications*. Springer, 2013.
- [24] C. De Franco, P. Tankov, and X. Warin. Numerical methods for the quadratic hedging problem in Markov models with jumps. *arXiv preprint arXiv:1206.5393v3*, 2013.
- [25] K. Dareiotis. On finite difference schemes for partial integro-differential equations of Lévy type. *Journal of Computational and Applied Mathematics*, 368:112587, 2020.
- [26] W. E, M. Hutzenthaler, A. Jentzen, and T. Kruse. Multilevel picard iterations for solving smooth semilinear parabolic heat equations. *Partial Differential Equations and Applications*, 2(6):1–31, 2021.
- [27] W. E, J. Han, and A. Jentzen. Deep learning-based numerical methods for high-dimensional parabolic partial differential equations and backward stochastic differential equations. *Communications in Mathematics and Statistics*, 5(4):349–380, 2017.
- [28] W. E and B. Yu. The deep ritz method: a deep learning-based numerical algorithm for solving variational problems. *Communications in Mathematics and Statistics*, 6(1):1–12, 2018.

- [29] H. Federer. *Geometric Measure Theory, Reprint of the 1969 Edition*. Springer-Verlag Berlin Heidelberg, 1996.
- [30] R. Frey and V. Köck. Deep neural network algorithms for parabolic pides and applications in insurance mathematics. In *Methods and Applications in Fluorescence*, pages 272–277. Springer, 2022.
- [31] M. Germain, H. Pham, and X. Warin. Approximation error analysis of some deep backward schemes for nonlinear pdes. *SIAM Journal on Scientific Computing*, 44(1):A28–A56, 2022.
- [32] M.B. Giles, A. Jentzen, and T. Welti. Generalised multilevel picard approximations. *arXiv preprint arXiv:1911.03188*, 2019.
- [33] L. Gonon and C. Schwab. Deep ReLU network expression rates for option prices in high-dimensional, exponential Lévy models. *Finance and Stochastics*, 25(4):615–657, 2021.
- [34] L. Gonon and C. Schwab. Deep ReLU neural network approximation for stochastic differential equations with jumps. *arXiv Preprint, arXiv:2102.11707v1*, 2021.
- [35] I. Gyöngy and S. Wu. Itô’s formula for jump processes in L_p -spaces. *Stochastic Processes and their Application*, 131: 523-552, 2021
- [36] J. Han, A. Jentzen, and W. E. Solving high-dimensional partial differential equations using deep learning. *Proceedings of the National Academy of Sciences*, 115(34):8505–8510, 2018.
- [37] C. Huré, H. Pham, and X. Warin. Deep backward schemes for high-dimensional nonlinear pdes. *Mathematics of Computation*, 89(324):1547–1579, 2020.
- [38] M. Hutzenthaler, A. Jentzen, and T. Kruse. On multilevel picard numerical approximations for high-dimensional nonlinear parabolic partial differential equations and high-dimensional nonlinear backward stochastic differential equations. *Journal of Scientific Computing*, 79(3):1534–1571, 2019.
- [39] M. Hutzenthaler, A. Jentzen, T. Kruse, and T.A. Nguyen. Multilevel Picard approximations for high-dimensional semilinear second-order PDEs with Lipschitz nonlinearities. *arXiv preprint arXiv:2009.02484v4*, 2020.
- [40] M. Hutzenthaler and T. Kruse. Multilevel picard approximations of high-dimensional semilinear parabolic differential equations with gradient-dependent nonlinearities. *SIAM Journal on Numerical Analysis*, 58(2):929–961, 2020.
- [41] M. Hutzenthaler, A. Jentzen, T. Kruse, T.A. Nguyen, and P. Von Wurstemberger. Overcoming the curse of dimensionality in the numerical approximation of semilinear parabolic partial differential equations. *arXiv preprint arXiv:1807.01212*, 2018.
- [42] N. Ikeada and S. Watanabe. *Stochastic Differential Equations and Diffusion Processes Second Edition*. North-Holland/Kodansha, 1989.
- [43] K. Ishii and N. Yamada. Viscosity solutions of nonlinear second order elliptic PDEs involving nonlocal operators. *Osaka Journal of Mathematics*, 30(3):439-455, 1993.
- [44] E.R. Jakobsen and K.H. Karlsen. Continuous dependence estimates for viscosity solutions of integro-PDEs. *J. Differential Equations*, 212(2):278–318, 2005.
- [45] Y. Kwon and Y. Lee. A second-order finite difference method for option pricing under jump-diffusion models. *SIAM journal on numerical analysis*, 49(6):2598-2617, 2011
- [46] N.V. Krylov. *Introduction to the Theory of Random Processes*. American Mathematical Society, 2002.
- [47] H. Kunita. Stochastic Differential Equations Based on Lévy Processes and Stochastic Flows and Diffeomorphisms. In *Real and stochastic analysis*, pages 305-373, Birkhäuser Boston, 2004.
- [48] S. Larsson and V. Thomée. *Partial differential equations with numerical methods* (Vol. 45). Berlin: Springer, 2003.
- [49] L. Lu, X. Meng, Z. Mao, and G.E. Karniadakis. DeepXDE: a deep learning library for solving differential equations. *SIAM Rev.*, 63(1):208–228, 2021.
- [50] A.M. Matache, T. Von Petersdorff, and C. Schwab. Fast deterministic pricing of options on Lévy driven assets. *ESAIM: Mathematical Modelling and Numerical Analysis*, 38(1):37–71, 2004.
- [51] B. Øksendal and A. Sulem. *Stochastic Control of jump diffusions*. Springer, 2005.
- [52] E. Platen and N. Bruti-Liberati. *Numerical Solution of Stochastic Differential Equations with Jumps in Finance* (Vol. 64). Springer Science & Business Media, 2010.
- [53] M. Raissi, P. Perdikaris, and G. E. Karniadakis. Physics-informed neural networks: a deep learning framework for solving forward and inverse problems involving nonlinear partial differential

- equations. *J. Comput. Phys.*, 378:686–707, 2019.
- [54] Justin Sirignano and Konstantinos Spiliopoulos. DGM: a deep learning algorithm for solving partial differential equations. *J. Comput. Phys.*, 375:1339–1364, 2018.
- [55] R. Situ. *Theory of Stochastic Differential Equations with Jumps and Applications: Mathematical and Analytical Techniques with Applications to Engineering*. Springer Science & Business Media, 2006.
- [56] V. Thomée. *Galerkin finite element methods for parabolic problems*(Vol. 25). Springer Science & Business Media, 2007
- [57] J. Yong and X.Y. Zhou *Stochastic Controls Hamiltonian Systems and HJB Equations, Applications of Mathematics: Stochastic Modelling and Applied Probability* (Vol. 43). Springer Science & Business Media, 1999.
- [58] L. Yuan, Y.Q. Ni, X.Y. Deng, and S. Hao. A-pinn: Auxiliary physics informed neural networks for forward and inverse problems of nonlinear integro-differential equations. *Journal of Computational Physics*, page 111260, 2022.

Email address: ariel.neufeld@ntu.edu.sg

Email address: sizhou.wu@ntu.edu.sg