## Ground-state energy distribution of disordered many-body quantum systems

Wouter Buijsman,<sup>1</sup> Talía L. M. Lezama,<sup>2</sup> Tamar Leiser,<sup>2</sup> and Lea F. Santos<sup>2</sup>

<sup>1</sup>Department of Physics, Ben-Gurion University of the Negev, Beer-Sheva 84105, Israel

<sup>2</sup>Department of Physics, Yeshiva University, New York, New York 10016, USA

Extreme-value distributions are studied in the context of a broad range of problems, from the equilibrium properties of low-temperature disordered systems to the occurrence of natural disasters. Our focus here is on the ground-state energy distribution of disordered many-body quantum systems. We derive an analytical expression that, upon tuning a parameter, reproduces with high accuracy the ground-state energy distribution of all the systems that we consider. For several models with short-and long-range random couplings, the ground-state distribution agrees remarkably well with the Tracy-Widom distribution obtained from random matrices. They include transverse Ising models, the Sachdev-Ye model and variants, and a randomized version of the PXP model. In contrast, shapes at odds with the Tracy-Widom distribution are observed for disordered Bose-Hubbard models and the disordered spin-1/2 Heisenberg chain used to investigate many-body localization.

The level spacing distribution in the bulk of the spectrum of disordered many-body quantum systems and its comparison with full random matrices [1-4] has been intensively investigated, due to its relevance for the study of thermalization in isolated quantum systems [5, 6], many-body localization [7–10], and nonequilibrium quantum dynamics [11–15]. The present work focuses instead on the distribution of the lowest energy level of such systems and its relationship with random matrices, a subject that has received significantly less attention [16, 17].

Extreme-value statistics concerns the study of rare events, such as tsunamis, floods, earthquakes, and large variations in the stock market. It also involves the study of the fluctuations of the smallest (largest) eigenvalue of random matrices [18], which has applications in analyses of the stability of dynamical systems with interactions [19, 20] and of the equilibrium properties of disordered systems at low temperatures [20-22]. In the case of independent and identically distributed random variables, there are three universality classes for the distribution of the sample minimum (maximum). They are given by the Gumbel, Fréchet, or Weibull distribution, depending on the tail of the parent probability density of the variables [22]. However, when the random variables are correlated, there are very few cases for which the extreme-value distribution has been obtained [22], one being the distribution of the lowest (highest) eigenvalue of ensembles of large Gaussian random matrices derived by Tracy and Widom [23, 24].

The Tracy-Widom distribution arises in different theoretical and experimental contexts, such as in studies of mesoscopic fluctuations in quantum dots and of spatial correlations of non-interacting fermions at the edges of a trap (see [22] and references therein). Of particular interest to us is the verification that the ground-state energy distribution of even-even nuclei [25–27] agrees with the Tracy-Widom distribution obtained with Gaussian orthogonal ensembles (GOE) [28].

Nuclei are typical examples of interacting many-body quantum systems, so features found there may extend also to other many-body quantum systems. This prompts us to investigate in this work the distribution of the lowest level of different disordered many-body quantum systems.

We find excellent agreement with the GOE Tracy-Widom distribution for various spin models with shortand long-range random couplings, such as transverse Ising models [29], the Sachdev-Ye model [30], and a randomized version of the PXP model [31]. Many of these systems can be realized in experiments with cold atoms [32–34], ion traps [35, 36], and nuclear magnetic resonance [37, 38]. However, we also identify examples of disordered many-body quantum systems, where the ground-state energy distribution is other than the Tracy-Widom. These include spin-1/2 Heisenberg chains with onsite disorder and Bose-Hubbard models with random couplings. Using real-valued Wishart ensembles, we derive a general one-parameter analytical expression that reproduces up to high accuracy the ground-state energy distribution of all the models that we study, whether it is a Tracy-Widom distribution or not.

Agreement with random matrices. – The GOE consists of real-valued symmetric matrices with entries sampled independently from the Gaussian distribution with zero mean  $\mu = 0$  and off-diagonal (diagonal) components with variance  $\sigma^2 = 1/2$  ( $\sigma^2 = 1$ ) [39, 40]. The eigenvalues  $E_0, E_1, \ldots, E_{N-1}$  are distributed according to

$$P(E_0, E_1, \dots, E_{N-1}) = \frac{1}{\mathcal{Z}_N} \prod_{i=0}^{N-1} e^{-E_i^2/2} \prod_{j < k} |E_j - E_k|,$$
(1)

where  $Z_N$  is a normalization constant that fixes the probability integral to unity. This expression does not factorize in terms that uniquely depend on a single eigenvalue, indicating that the eigenvalues are (strongly) correlated. Sorting the eigenvalues in increasing order, the distribution of the smallest eigenvalue  $E_0$ , is obtained by integrating out all other eigenvalues,

$$P(E_0) = \int P(E_0, E_1, \dots, E_{N-1}) dE_1 dE_2 \dots dE_{N-1}.$$
(2)

For  $N \to \infty$ , the distribution of the smallest eigenvalue converges towards the Tracy-Widom distribution [23, 24]. This distribution does not have a closed-form expression, but it can be written in terms of the solution of the Painlevé II differential equation and evaluated numerically. After some manipulation [41], one finds that the distribution of the ground-state energy has the shape

$$P(E_0) = \sqrt{F_2(-E_0)} \exp\left[\frac{1}{2} \int_{-E_0}^{\infty} q(x) dx\right]$$
(3)

with

$$F_2(x) = \exp\left[-\int_x^\infty (z-x)q^2(x)dz\right],\qquad(4)$$

where q(x) is the solution of the Painlevé II differential equation  $q'' = xq + 2q^3$  subjected to the boundary condition  $q(x) \sim \operatorname{Ai}(x)$  for  $x \to \infty$ , with  $\operatorname{Ai}(x)$  denoting the Airy function.

We compare the Tracy-Widom distribution in Eq. (3) with the ground-state energy distribution of different disordered many-body quantum systems. For all the models considered in this work, the random parameters are independent real numbers sampled from the Gaussian distribution with mean  $\mu = 0$  and variance  $\sigma^2 = 1$ . The ground-state energy distributions are shifted and scaled such that their mean and variance are zero and unity, respectively.

In Fig. 1 (a), we plot  $P(E_0)$  for the Hamiltonian

$$H = \sum_{i=1}^{L} \frac{\epsilon_i}{2} \sigma_i^z + \sum_{i,j}^{L-1} \frac{J_{ij}}{4|j-i|^{\alpha}} \vec{\sigma}_i \cdot \vec{\sigma}_j, \tag{5}$$

where L is the system size,  $\epsilon_i$  and  $J_{ij} = J_{ji}$  are random numbers, and  $\sigma^{x,y,z}$  are the Pauli matrices. For all-to-all couplings ( $\alpha = 0$ ), the Hamiltonian in Eq. (5) is analogous to the two-body random ensembles studied in quantum chaos and nuclear physics [1, 4, 42–44] and coincides with the Sachdev-Ye model when  $\epsilon_i = 0$  [30]. We also study  $\alpha > 0$ , in which case we consider one-dimensional (1D) systems. Hamiltonian (5) conserves the total zmagnetization  $S_z = \sum_{i=1}^L S_i^z$ .

The distribution of the ground-state energy of H (5) is remarkably robust. As seen in Fig. 1 (a), the agreement with the Tracy-Widom distribution is excellent for  $\alpha = 0$ , and as  $\alpha$  increases from all-to-all to short-range couplings, the deviations remain minor. For the  $S_z = 0$ sector, the agreement holds for system sizes as small as L = 8; and for large system sizes, there is very good agreement for as few as 4 excitations.  $P(E_0)$  also matches the Tracy-Widom distribution for the Sachdev-Ye model, that is, for H (5) with  $\alpha = 0$  and  $\epsilon_i = 0$ .



FIG. 1. Ground-state energy distribution (a)-(d) of the systems defined in Eqs. (5)-(8), respectively. The data (symbols) consist of  $10^6$  realizations for (a) H in Eq. (5) for the zeromagnetization sector ( $S_z = 0$ ) and system size L = 16, (b) Hin Eq. (6) for L = 14, (c) H in Eq. (7) for L = 16, and (d) Hin Eq. (8) for  $S_z = 0$  and L = 16. The solid line represents the Tracy-Widom distribution.

The agreement with the Tracy-Widom distribution is also excellent in Fig. 1 (b) and Fig. 1 (c). In Fig. 1 (b), the comparison is made with the Ising model with longitudinal and transverse fields,

$$H = \sum_{i=1}^{L} \left( \frac{h_i^z}{2} \sigma_i^z + \frac{h_i^x}{2} \sigma_i^x \right) + \sum_{i,j}^{L-1} \frac{J_{ij}}{4|j-i|^{\alpha}} \sigma_i^z \sigma_j^z, \quad (6)$$

where  $h_i^z$ ,  $h_i^x$ , and  $J_{ij} = J_{ji}$  are random numbers. In the case of nearest-neighbor couplings, Eq. (6) agrees with the Hamiltonian used to study quantum phase transitions in Ref. [29]. We examine both short- and long-range interactions. In Fig. 1 (c), we consider the Hamiltonian

$$H = \sum_{i=1}^{L-2} J_{i+1} P_i \sigma_{i+1}^x P_{i+2} + J_1 \sigma_1^x P_2 + J_L P_{L-1} \sigma_L^x, \quad (7)$$

where  $P_i = (1 - \sigma_i^z)/2$  denotes the projection operator and  $J_i$  are random numbers. The main mechanism of this model is based on local constraints that forbid two adjacent spins to be simultaneously in the up-state. For  $J_i = 1$ , the Hamiltonian reduces to the PXP model [31, 32], which is paradigmatic in studies on quantum many-body scars and weak ergodicity breaking [45, 46]. For random parameters, model (7) is similar to the one introduced in [47].

Contrary to the previous models, the ground-state energy distribution of the disordered spin-1/2 Heisenberg Hamiltonian employed in studies of many-body localization [7–10],

$$H = W \sum_{i=1}^{L} \frac{h_i}{2} \sigma_i^z + \frac{J}{4} \sum_{i=1}^{L-1} \vec{\sigma}_i \cdot \vec{\sigma}_{i+1}, \qquad (8)$$

does not always agree with the Tracy-Widom distribution, as seen in Fig. 1 (d). In the equation above,  $h_i$ 



FIG. 2. Ground-state energy distribution of the Hamiltonian in Eq. (8), which is prototypical in studies of many-body localization. The distribution deviates from (approaches) the Tracy-Widom distribution for disorder strength  $0 < W \leq J$  (W > J). The data (shades) are obtained using 10<sup>6</sup> realizations for the zero-magnetization sector ( $S_z = 0$ ) and system size L = 16. Solid, dashed, and dot-dashed lines represent the Tracy-Widom,  $\chi^2$ -, and Weibull distribution, respectively. The values of M are rounded to their closest integers.

are random numbers, W is the disorder strength, and the coupling parameter J = 1. The properties of the bulk of the spectrum of this model are similar to those of GOE random matrices when  $0 < W \leq J$ , but differ when W > J [48, 49]. In stark contrast with the bulk of the spectrum, the ground-state energy distribution of H (8) differs from the GOE Tracy-Widom when  $0 < W \leq J$ and approaches it for W > J. To identify the distributions that best match those shown in Fig. 1 (d), we resort to real-valued Wishart ensembles, as described below.

General analytical expression.– The Wishart ensemble is one of the classical ensembles in random matrix theory and precedes the Gaussian ensembles [40]. It consists of matrices  $A = XX^T$ , where X is an  $N \times M$  matrix with (real-valued) entries sampled independently from the Gaussian distribution with mean  $\mu = 0$  and variance  $\sigma^2 = 1$ . The eigenvalues  $E_0, E_1, \ldots, E_{N-1}$  are distributed according to

$$P(E_0, E_1, \dots, E_{N-1}) = \frac{1}{Z_N} \prod_{i=0}^{N-1} e^{-E_i/2} \times \prod_i E_i^{(M-N-1)/2} \prod_{j < k} |E_j - E_k|, \quad (E_i \ge 0),$$
(9)

where, as in Eq. (1), the prefactor  $1/\mathcal{Z}_N$  is a normalization constant fixing the integral probability to unity.

For N = 1, we find that by varying M, Eq. (9) can capture the various levels of asymmetry exhibited by the ground-state energy distributions seen in Fig. 1 (d). In this case, the Wishart ensemble consists of samples given by the sum of M squared independent random numbers, so Eq. (9) reduces to the chi-squared  $(\chi^2-)$  distribution,

$$P(E_0) = \frac{1}{2^{M/2} \Gamma(M/2)} e^{-E_0/2} E_0^{M/2-1}, \quad (E_0 \ge 0),$$
(10)

with M degrees of freedom, mean  $\mu = M$ , and variance  $\sigma^2 = 2M$ . Since our focus is on minima, instead of maxima, we consider the distribution of  $-E_0$ .

To find the best-fitting value of M for a given groundstate energy distribution, we use the skewness  $\gamma = -\mu_3/\mu_2^{3/2}$  as a matching parameter, taking into account that the *n*-th moment is  $\mu_n = \int (x-\mu)^n P(x) dx$ . For the N = 1 Wishart ensemble,  $\gamma = -2\sqrt{2/M}$  and thus

$$M = \frac{8}{\gamma^2}.$$
 (11)

Agreement with the  $\chi^2$ -distribution. – In Fig. 2, we select some of the curves exhibited in Fig. 1 (d) to show that the analytical expression in Eq. (10) matches the numerical results for  $P(E_0)$  obtained with H (8) for any disorder strength W.

The best-fitting  $\chi^2$ -distributions for the cases with small disorder strength (W < 1) resemble the Weibull distribution,

$$P(E_0) = \frac{k}{\lambda} \left(\frac{E_0 - \theta}{\lambda}\right)^{k-1} e^{-\left(\frac{E_0 - \theta}{\lambda}\right)^k}, \qquad (E_0 > \theta).$$
(12)

The parameters  $\theta$ ,  $\lambda$ , and k are computed to ensure that the distribution has zero mean, unit variance, and the same skewness as that of the numerical data in



FIG. 3. In (a): The ground-state energy distribution for H (13) with  $N_b = L = 7$ . Symbols are numerical data, dashed lines represent the  $\chi^2$ -distribution, and the solid line is the Tracy-Widom distribution. Comparison between the Tracy-Widom [2 × 2 GOE ground-state energy] distribution and the  $\chi^2$ -distribution is shown in (b) [(c)].

Figs. 2 (a)-(c). As W increases above 1, the bestfitting  $\chi^2$ -distributions approach the Tracy-Widom distribution, the agreement being very good when W = 8in Fig. 2 (f).

For models that, similarly to H (8), have a fixed constant parameter, such as the Hamiltonians in [47, 50], we verify that the ground-state energy distribution also deviates from the Tracy-Widom distribution. For all of these cases,  $P(E_0)$  can be reproduced with an appropriate choice of M in Eq. (10). This equation also holds for the disordered 1D Bose-Hubbard model,

$$H = \sum_{i=1}^{L} \frac{U_i}{2} n_i \left( n_i - 1 \right) - \sum_{i,j}^{L-1} \frac{J_{i,j}}{|j-i|^{\alpha}} \left( a_i^{\dagger} a_j + a_j^{\dagger} a_i \right),$$
(13)

where  $a_i$   $(a_i^{\dagger})$  is the annihilation (creation) operator,  $n_i = a_i^{\dagger} a_i$ , and the parameters  $U_i$  and  $J_{i,j}$  are random numbers. When the total number of bosons,  $N_b$ , is small,  $P(E_0)$  shows good agreement with the Tracy-Widom distribution. However, for  $N_b = L$ , deviations occur for both short- and long-range couplings, as seen in Fig. 3 (a), and for any value of  $\alpha$ , the shapes are well captured by the  $\chi^2$ -distribution.

A natural question that emerges from our comparisons between Eq. (10) and the ground-state energy distribution of various disordered models is the value of M that leads to the agreement between the Tracy-Widom distribution and the  $\chi^2$ -distribution. The skewness of the GOE Tracy-Widom distribution has been obtained numerically,  $\gamma = -0.2934645240...$  [41]. Equating this value to the skewness of the  $\chi^2$ -distribution gives M = 93after rounding to the nearest integer. As shown in Fig. 3 (b), the two curves agree down to very small values. The parallel between these two distributions is useful, because a closed form for the Tracy-Widom distribution does not yet exist. One needs to evaluate it numerically, which requires solving the Painlevé II differential equation. Using instead the  $\chi^2$ -distribution is a simple and accurate alternative (see also Ref. [51]).

By further increasing M beyond the Tracy-Widom shape, the  $\chi^2$ -distribution eventually matches the analytical expression of  $P(E_0)$  obtained from  $2 \times 2$  GOE random matrices,

$$P(E_0) = \frac{1}{2\sqrt{\pi}} e^{-E_0^2} - \frac{1}{2\sqrt{2}} e^{-\frac{1}{2}E_0^2} E_0 \operatorname{erfc}\left(\frac{E_0}{\sqrt{2}}\right), \quad (14)$$

which, in turn, is comparable to the ground-state energy distribution of some nuclear and molecular models [28]. The skewness of the distribution in Eq. (14),  $\gamma = -[2(\pi - 3)\sqrt{\pi}/(6 - \pi)^{3/2}]$ , matches the skewness of the  $\chi^2$ -distribution for M = 742 after rounding to the nearest integer. The excellent agreement between the two curves is illustrated in Fig. 3 (c). Finally, for  $M \to \infty$ , the  $\chi^2$ -distribution converges towards a Gaussian.

Conclusions. – The use of the  $\chi^2$ -distribution to study the lowest energy level distribution of quantum systems is analogous to the use of the Brody distribution [1] (or the Izrailev distribution [52]) to study the level spacing distribution of the bulk of their spectra. By varying a fitting parameter, the Brody distribution changes from Poisson to Wigner-Dyson, covering all shapes in between. Here, we show that by varying the number of degrees of freedom of the  $\chi^2$ -distribution, we can reproduce with great accuracy the shape of the ground-state energy distribution for any of the disordered many-body quantum systems that we consider. In the case of the Heisenberg spin-1/2 chain with onsite disorder, in particular, the  $\chi^2$ -distribution captures the crossover from the Weibull distribution to the Tracy-Widom distribution as the disordered strength increases.

This research was supported by the United States National Science Foundation (NSF) Grant No. DMR-1936006. WB acknowledges support from the Kreitman School of Advanced Graduate Studies at Ben-Gurion University.

WB and TLML contributed equally to this work.

- T. A. Brody, J. Flores, J. B. French, P. A. Mello, A. Pandey, and S. S. M. Wong, Random-matrix physics – spectrum and strength fluctuations, Rev. Mod. Phys 53, 385 (1981).
- [2] V. Zelevinsky, B. A. Brown, N. Frazier, and M. Horoi, The nuclear shell model as a testing ground for manybody quantum chaos, Phys. Rep. 276, 85 (1996).
- [3] T. Guhr, A. Müller-Groeling, and H. A. Weidenmüller, Random matrix theories in quantum physics: Common concepts, Phys. Rep. 299, 189 (1998).
- [4] V. K. B. Kota, Embedded random matrix ensembles for

- [5] F. Borgonovi, F. M. Izrailev, L. F. Santos, and V. G. Zelevinsky, Quantum chaos and thermalization in isolated systems of interacting particles, Phys. Rep. 626, 1 (2016).
- [6] L. D'Alessio, Y. Kafri, A. Polkovnikov, and M. Rigol, From quantum chaos and eigenstate thermalization to statistical mechanics and thermodynamics, Adv. Phys. 65, 239 (2016).
- [7] L. F. Santos, G. Rigolin, and C. O. Escobar, Entanglement versus chaos in disordered spin systems, Phys. Rev. A 69, 042304 (2004).
- [8] R. Nandkishore and D. Huse, Many-body localization and thermalization in quantum statistical mechanics, Annu. Rev. Condens. Matter Phys. 6, 15 (2015).
- [9] F. Alet and N. Laflorencie, Many-body localization: An introduction and selected topics, C. R. Phys. 19, 498 (2018).
- [10] D. A. Abanin, E. Altman, I. Bloch, and M. Serbyn, Colloquium: Many-body localization, thermalization, and entanglement, Rev. Mod. Phys. 91, 021001 (2019).
- [11] G. P. Brandino, A. De Luca, R. M. Konik, and G. Mussardo, Quench dynamics in randomly generated extended quantum models, Phys. Rev. B 85, 214435 (2012).
- [12] E. J. Torres-Herrera, J. Karp, M. Távora, and L. F. Santos, Realistic many-body quantum systems vs. full random matrices: Static and dynamical properties, Entropy 18, 359 (2016).
- [13] J. Cotler, N. Hunter-Jones, J. Liu, and B. Yoshida, Chaos, complexity, and random matrices, J. High Energy Phys. 2017 (11), 48.
- [14] E. J. Torres-Herrera, A. M. García-García, and L. F. Santos, Generic dynamical features of quenched interacting quantum systems: Survival probability, density imbalance, and out-of-time-ordered correlator, Phys. Rev. B 97, 060303 (2018).
- [15] M. Schiulaz, E. J. Torres-Herrera, and L. F. Santos, Thouless and relaxation time scales in many-body quantum systems, Phys. Rev. B 99, 174313 (2019).
- [16] G. Gur-Ari, R. Mahajan, and A. Vaezi, Does the SYK model have a spin glass phase?, J. High Energy Phys. 2018 (11), 70.
- [17] J.-M. Stéphan, Free fermions at the edge of interacting systems, SciPost Phys. 6, 57 (2019).
- [18] P. Embrechts, C. Klüppelberg, and T. Mikosch, Modelling Extremal Events: for Insurance and Finance (Columbia University Press, New York, 2003).
- [19] R. M. May, Will a large complex system be stable?, Nature 238, 413 (1972).
- [20] S. N. Majumdar and G. Schehr, Top eigenvalue of a random matrix: large deviations and third order phase transition, J. Stat. Mech. 2014, P01012 (2014).
- [21] J.-P. Bouchaud and M. Mézard, Universality classes for extreme-value statistics, J. Phys. A 30, 7997 (1997).
- [22] S. N. Majumdar, A. Pal, and G. Schehr, Extreme value statistics of correlated random variables: A pedagogical review, Phys. Rep. 840, 1 (2020).
- [23] C. A. Tracy and H. Widom, Level-spacing distributions and the Airy kernel, Comm. Math. Phys. 119, 151 (1994).
- [24] C. A. Tracy and H. Widom, On orthogonal and symplectic matrix ensembles, Comm. Math. Phys. 177, 727 (1996).

- [25] C. W. Johnson, G. F. Bertsch, and D. J. Dean, Orderly spectra from random interactions, Phys. Rev. Lett. 80, 2749 (1998).
- [26] D. Kusnezov, Two-body random ensembles: From nuclear spectra to random polynomials, Phys. Rev. Lett. 85, 3773 (2000).
- [27] Y. Zhao, A. Arima, and N. Yoshinaga, Regularities of many-body systems interacting by a two-body random ensemble, Phys. Rep. 400, 1 (2004).
- [28] L. F. Santos, D. Kusnezov, and P. Jacquod, Ground state properties of many-body systems in the two-body random ensemble and random matrix theory, Phys. Lett. B. 537, 62 (2002).
- [29] Y.-P. Lin, Y.-J. Kao, P. Chen, and Y.-C. Lin, Griffiths singularities in the random quantum ising antiferromagnet: A tree tensor network renormalization group study, Phys. Rev. B 96, 064427 (2017).
- [30] S. Sachdev and J. Ye, Gapless spin-fluid ground state in a random quantum Heisenberg magnet, Phys. Rev. Lett. 70, 3339 (1993).
- [31] I. Lesanovsky and H. Katsura, Interacting Fibonacci anyons in a Rydberg gas, Phys. Rev. A 86, 041601 (2012).
- [32] H. Bernien, S. Schwartz, A. Keesling, H. Levine, A. Omran, H. Pichler, S. Choi, A. S. Zibrov, M. Endres, M. Greiner, V. Vuletić, and M. D. Lukin, Probing manybody dynamics on a 51-atom quantum simulator, Nature 551, 579 (2017).
- [33] I. Bloch, J. Dalibard, and W. Zwerger, Many-body physics with ultracold gases, Rev. Mod. Phys. 80, 885 (2008).
- [34] A. M. Kaufman, A. L. M. Eric Tai, M. Rispoli, R. Schittko, P. M. Preiss, and M. Greiner, Quantum thermalization through entanglement in an isolated many-body system, Science **353**, 794 (2016).
- [35] R. Blatt and C. F. Roos, Quantum simulations with trapped ions, Nat. Phys. 8, 277 (2012).
- [36] P. Richerme, Z.-X. Gong, A. Lee, C. Senko, J. Smith, M. Foss-Feig, S. Michalakis, A. V. Gorshkov, and C. Monroe, Non-local propagation of correlations in quantum systems with long-range interactions, Nature 511, 198 (2014).
- [37] K. X. Wei, C. Ramanathan, and P. Cappellaro, Exploring localization in nuclear spin chains, Phys. Rev. Lett. 120, 070501 (2018).
- [38] M. Niknam, L. F. Santos, and D. G. Cory, Sensitivity of quantum information to environment perturbations measured with a nonlocal out-of-time-order correlation function, Phys. Rev. Research 2, 013200 (2020).
- [39] M. L. Mehta, *Random Matrices* (Academic Press, Boston, 1991).
- [40] G. Livan, M. Novaes, and P. Vivo, *Introduction to Random Matrices: Theory and Practice* (Springer, New York, 2018).
- [41] S. R. Finch, *Mathematical Constants II* (Cambridge University Press, Cambridge, 2018).
- [42] J. B. French and S. S. M. Wong, Validity of random matrix theories for many-particle systems, Phys. Lett. B 33, 449 (1970).
- [43] O. Bohigas and J. Flores, Two-body random Hamiltonian and level density, Phys. Lett. B 34, 261 (1971).
- [44] T. Papenbrock and H. A. Weidenmüller, Two-body randem ensemble for nuclei, Int. J. Mod. Phys. E 15, 1885 (2006).
- [45] C. J. Turner, A. A. Michailidis, D. A. Abanin, M. Serbyn,

and Z. Papić, Weak ergodicity breaking from quantum many-body scars, Nat. Phys. 14, 745 (2018).

- [46] C. J. Turner, A. A. Michailidis, D. A. Abanin, M. Serbyn, and Z. Papić, Quantum scarred eigenstates in a Rydberg atom chain: Entanglement, breakdown of thermalization, and stability to perturbations, Phys. Rev. B 98, 155134 (2018).
- [47] C. Chen, F. Burnell, and A. Chandran, How does a locally constrained quantum system localize?, Phys. Rev. Lett. 121, 085701 (2018).
- [48] Y. Avishai, J. Richert, and R. Berkovitz, Level statistics in a Heisenberg chain with random magnetic field, Phys.

Rev. B 66, 052416 (2002).

- [49] L. F. Santos, Integrability of a disordered Heisenberg spin-1/2 chain, J. Phys. A 37, 4723 (2004).
- [50] L. Rademaker and D. A. Abanin, Slow nonthermalizing dynamics in a quantum spin glass, Phys. Rev. Lett. 125, 260405 (2020).
- [51] M. Chiani, Distribution of the largest eigenvalue for real Wishart and Gaussian random matrices and a simple approximation for the Tracy–Widom distribution, J. Multivariate Anal. **129**, 69 (2014).
- [52] F. M. Izrailev, Simple models of quantum chaos: Spectrum and eigenfunctions, Phys. Rep. 196, 299 (1990).