

Oscillations of High-Energy Cosmic Neutrinos in the Copious MeV Neutrino Background

Sajad Abbar,¹ Jose Alonso Carpio,² and Kohta Murase^{2,3}

¹*Max-Planck-Institut für Physik (Werner-Heisenberg-Institut), Föhringer Ring 6, 80805 München, Germany*

²*Department of Physics; Department of Astronomy and Astrophysics; Center for Multimessenger Astrophysics, Institute for Gravitation and the Cosmos, The Pennsylvania State University, University Park, Pennsylvania 16802, USA*

³*Center for Gravitational Physics, Yukawa Institute for Theoretical Physics, Kyoto, Kyoto 606-8502, Japan*

The core-collapse of massive stars and merger of neutron star binaries are among the most promising candidate sites for the production of high-energy cosmic neutrinos. We demonstrate that the high-energy neutrinos produced in such extreme environments can experience efficient flavor conversions on scales much shorter than those expected in vacuum, due to their coherent forward scatterings with the bath of decohered low-energy neutrinos emitted from the central engine. These low-energy neutrinos, which exist as mass eigenstates, provide a very special and peculiar dominant background for the propagation of the high-energy ones. We point out that the high-energy neutrino flavor ratio is modified to a value independent of neutrinos energies, which is distinct from the conventional prediction with the matter effect. We also suggest that the signals can be used as a novel probe of new neutrino interactions beyond the Standard Model. This is yet another context where neutrino-neutrino interactions can play a crucial role in their flavor evolution.

I. INTRODUCTION

Core-collapse supernovae (CCSNe) and neutron star mergers (NSMs) commonly lead to a burst of thermal neutrinos in the MeV range, with a very active literature on the physics of their oscillations. These environments are considered as the central engine of not only supernovae but also gamma-ray bursts (GRBs), and other energetic or transrelativistic supernovae driven by outflows such as jets and winds (i.e., engine-driven supernovae). It has been suggested that GeV-TeV neutrinos can be produced in such environments if neutron-loaded outflows are launched from a black hole with an accretion disk and/or a newborn magnetar [1–5]. Even TeV-PeV neutrinos can be generated inside the outflows through shock acceleration or magnetic reconnections [6–10].

The IceCube discovery of high-energy neutrinos (HE ν 's) has opened a new avenue to investigate the physics of neutrino oscillations and related neutrino physics (see Refs. [11–13] and references therein). In this article, we investigate a novel effect caused by the interplay between the HE ν 's produced in outflows and low-energy neutrinos (LE ν 's) directly from the central engine (see Fig. 1 for the schematic picture). Indeed, the decohered LE ν 's, which are in mass eigenstates, can provide a *dominant* unusual background for the propagation of the HE ones. In particular, we show that the resulting neutrino self-interactions (ν SI) leads to a very intriguing phenomenon, in which the HE ν 's experience short-scale flavor oscillations in such a way that on average, they end up in the mass eigenstates. This phenomenon is noncollective in spirit and differs remarkably from the well-known phenomenon of *collective oscillations* of MeV neutrinos occurring in dense neutrino environments such as CCSNe and NSM remnants [14–16].




FIG. 1. Schematic picture of HE ν (GeV-PeV) production and their interactions with LE ν 's (MeV-GeV) from the central engine such as a black hole with an accretion disk or a newborn magnetar. HE ν production occurs at $R_{\text{diss}} \gg R_{\text{eng}}$, which may be beamed with the opening angle $\sim 1/\Gamma$, with R_{diss} , R_{eng} , and Γ being the dissipation radius, the engine radius, and the outflow Lorentz factor. Note that the opening angle of the LE ν beams is exaggerated for illustration purposes.

II. HE NEUTRINO INTERACTIONS IN JETS OR WINDS

Various scenarios for HE ν production in GRBs, CCSNe and NSMs have been suggested. In this work, we are interested in the fate of HE ν 's so we assume that they are produced at the dissipation radius $R_{\text{diss}} \sim 10^8 - 10^{10}$ cm, which is much larger than the engine radius $R_{\text{eng}} \sim 10^6$ cm. Then an interesting interplay is particularly plausible when the HE ν production occurs within the duration of LE ν 's and outflow breakout time (that are longer than the light crossing time). Note that as seen below the effect on neutrino oscillation is largely model independent as long as R_{diss} is so small that LE ν 's govern the neutrino potential.

Although the proposed mechanism works in pretty general setups, for illustrative purposes, we consider models of GeV-TeV neutrinos. Quasithermal neutrinos can naturally be produced in the GeV-TeV range through inelastic neutron-proton collisions when neutrons decouple from protons or neutron-loaded outflows make collisions with the surrounding environment [1–3, 5], and higher-energy nonthermal neutrinos may also be produced through neutron-proton-converter acceleration [3, 4]. For these neutrinos, the dissipation may occur at $R_{\text{diss}} \sim 10^8 - 10^{10}$ cm [1, 5]. Protons could further be accelerated to higher energies via shock acceleration or magnetic reconnections, and nonthermal TeV neutrinos can be efficiently produced via inelastic pp and/or $p\gamma$ interactions [6–10]. These neutrinos are associated with the dissipation at internal, collimation, and termination shocks [5, 6, 9, 17, 18]. For example, the internal dissipation radius is estimated to be $R_{\text{diss}} \approx 2\Gamma^2 c\delta t \sim 6 \times 10^8$ cm $(\Gamma/3)^2 (\delta t/1 \text{ ms})$, where δt is the variability time.

The number density of $\text{LE}\nu$'s at R_{diss} (in the engine frame) is

$$n_{\text{LE}\nu} = \frac{L_{\nu_e}}{4\pi R_{\text{diss}}^2 c \langle E_\nu \rangle} \simeq 1.7 \times 10^{27} \text{ cm}^{-3} \left(\frac{L_{\nu_e}}{10^{52} \text{ erg s}^{-1}} \right) \times \left(\frac{R_{\text{diss}}}{10^9 \text{ cm}} \right)^{-2} \left(\frac{\langle E_\nu \rangle}{10 \text{ MeV}} \right)^{-1}, \quad (1)$$

which can be much larger than the expected number density of $\text{HE}\nu$'s therein, $n_{\text{HE}\nu} \lesssim 10^{24} \text{ cm}^{-3}$ at 10^9 cm. Here L_{ν_e} and $\langle E_\nu \rangle$ are the electron neutrino luminosity and average energy, respectively. In addition, the electron number density in the outflow is

$$n_e \approx \frac{\Gamma L}{4\pi R_{\text{diss}}^2 \Gamma^2 m_p c^3} \simeq 5.9 \times 10^{23} \text{ cm}^{-3} \left(\frac{L}{10^{52} \text{ erg s}^{-1}} \right) \times \left(\frac{R_{\text{diss}}}{10^9 \text{ cm}} \right)^{-2} \left(\frac{\Gamma}{30} \right)^{-1} \ll n_{\text{LE}\nu} \quad (2)$$

Unlike the flavor evolution of the $\text{LE}\nu$'s which is dominated by the mass Hamiltonian at such neutrino number densities, the evolution of $\text{HE}\nu$'s can be dominated by their coherent scattering with the bath of the $\text{LE}\nu$'s. This simply comes from the fact that for the $\text{HE}\nu$'s, the strength of νSI (see Eq. (5)),

$$\mu \approx \sqrt{2} G_F n_\nu \hbar^2 c^2 \xi \simeq 6.4 \times 10^{-6} \text{ cm}^{-1} \left(\frac{n_\nu}{10^{27} \text{ cm}^{-3}} \right) \xi, \quad (3)$$

can be much larger than their vacuum wavelength, $\omega \approx \Delta m_{\text{atm}}^2 c^3 / (2\hbar E_\nu) \simeq 6 \times 10^{-10} \text{ cm}^{-1}$ ($100 \text{ GeV}/E_\nu$), with G_F being the Fermi constant. In the above equation $\xi = 1 - \cos \Theta$, where Θ is the opening angle of the neutrino beams, which is determined here mainly by the opening angle of $\text{HE}\nu$'s. Note that as soon as the parameter μ is known, ξ and ν do not provide any more relevant information. For relativistic flows with $\Gamma \sim 2 - 100$, one has $\xi \approx \Theta^2/2 \sim 1/(2\Gamma^2)$. Note that the optical depth

to incoherent neutrino scatterings is so small that the electron-positron pair production is negligible. Moreover, given the fact that the number density of $\text{LE}\nu$'s is much larger than that of $\text{HE}\nu$'s, one can assume that n_ν is here exclusively determined by the $\text{LE}\nu$'s.

Although the number density of the $\text{LE}\nu$'s within the zones of interest is expected to be too small to allow for the νSI Hamiltonian to compete with or dominate their vacuum Hamiltonian, the evolution of $\text{HE}\nu$'s is almost completely governed by the interaction term for appropriate $\text{LE}\nu$ number densities ($\omega_{\text{HE}\nu} \ll \mu \lesssim \omega_{\text{LE}\nu}$).

III. TWO-BEAM MODEL

In order to demonstrate how the flavor content of $\text{HE}\nu$'s is impacted by their propagation in the bath of the $\text{LE}\nu$'s, we study neutrino flavor conversions in a one-dimensional two-beam model, which consists of *two* energy bins, and a *three-flavor* neutrino gas with two angular beams. The neutrino energies are taken to be $E_\nu = 10$ MeV and 100 GeV for the bins representing the $\text{LE}\nu$'s and the $\text{HE}\nu$'s, respectively, unless otherwise stated. Thus in brief, our model consists of two angle beams each including neutrinos and antineutrinos with two energies representing high- and low-energy neutrinos. We also assume that the neutrino density is constant within the bath of $\text{LE}\nu$'s.

In order to study the flavor evolution of neutrinos in our model, we solve the Liouville-von Neumann equation for the neutrino density matrix, ϱ ($c = \hbar = 1$) [19]

$$id_t \varrho_{\mathbf{p}} = \left[\frac{\mathbf{U} \mathbf{M}^2 \mathbf{U}^\dagger}{2E_\nu} + \mathbf{H}_m + \mathbf{H}_{\nu\nu, \mathbf{p}}, \varrho_{\mathbf{p}} \right], \quad (4)$$

with

$$\mathbf{H}_{\nu\nu, \mathbf{p}} = \sqrt{2} G_F \int \frac{d^3 p'}{(2\pi)^3} (1 - \mathbf{v} \cdot \mathbf{v}') (\varrho_{\mathbf{p}'} - \bar{\varrho}_{\mathbf{p}'}), \quad (5)$$

being the neutrino potential stemming from the neutrino-neutrino forward scattering [20–22]. Here \mathbf{p} is the neutrino momentum, $E_\nu = |\mathbf{p}|$, $\mathbf{v} = \mathbf{p}/E_\nu$, and \mathbf{M}^2 are the energy, velocity, and mass-square matrix of the neutrino, respectively, and \mathbf{U} is the Pontecorvo–Maki–Nakagawa–Sakata matrix. Moreover, \mathbf{H}_m is the contribution from the matter term which is proportional to matter (electron) density [23, 24], which is ignored in our calculations due to the relatively small matter density inside the outflow. Hence, there are only two nonzero terms in \mathbf{H} (vacuum and νSI) which are both diagonal in the mass basis and constant (see below), but with different eigenvalues.

As mentioned above, $\mathbf{H}_{\nu\nu}$ is almost exclusively determined by $\text{LE}\nu$'s here due to their much larger number densities. In this study, we assume that $\text{LE}\nu$'s are in mass eigenstates because they are expected to be already decohered within the zones of interests, which are very far from their emission region (with a typical coherence

length of a few 10^6 cm for the atmospheric mass difference) [25–27]. In addition, the $\text{HE}\nu$'s do not significantly disturb the flavor state of the $\text{LE}\nu$ bath due to their much smaller number densities. This implies that the $\text{LE}\nu$'s do not evolve since their evolution is dominated by the mass term and they are already in the mass eigenstates. Consequently, $H_{\nu\nu}$ remains approximately constant. (We will later discuss how the neutrino flavor evolution changes once the oscillations of $\text{LE}\nu$'s are taken into account.) On the other hand, the $\text{HE}\nu$'s find themselves in the bath of the $\text{LE}\nu$'s as soon as they are produced, and start flavor conversions on relatively short scales due to their interactions with the $\text{LE}\nu$'s. It should be kept in mind that the coherence length of $\text{HE}\nu$'s are expected to be much longer than those of the LE ones [28].

IV. RESULTS

In the upper panel of Fig. 2, we show the survival probabilities of $\text{HE}\nu_e$'s propagating in vacuum (dashed curve), and in the bath of the $\text{LE}\nu$'s (solid curve). As can be clearly seen, the oscillation scales of $\text{HE}\nu$'s can change by orders of magnitude when coherent scatterings with $\text{LE}\nu$'s are taken into account. As a matter of fact, the oscillation scale of $\text{HE}\nu$'s in a bath of the LE ones is determined by the number density of the $\text{LE}\nu$'s, namely $l_{\text{osc}} \sim |H_{\nu\nu}|^{-1} \sim \mu^{-1}$ ($\sim 10^5$ cm for this simulation). This scaling behavior can be immediately deduced from Eq. (4) given the fact that for the $\text{HE}\nu$'s, the dominant contribution to the Hamiltonian comes from coherent scatterings with $\text{LE}\nu$'s as long as $\omega_{\text{HE}\nu} \ll \mu$ (we indeed observe this behaviour for $10 \omega_{\text{HE}\nu} \lesssim \mu$). Note that here the only relevant physical parameter is $\mu/\omega_{\text{HE}\nu}$, therefore one can play with μ in Fig. 2 as long as this ratio is constant (provided that r is also appropriately rescaled for the upper panel).

The fact that $\text{HE}\nu$'s oscillate on scales $\sim \mu^{-1}$ might remind an astute reader of the phenomenon of *fast* flavor conversions occurring for MeV neutrinos in dense neutrino media [29, 30]. This similarity becomes more obvious once one notes that the oscillations of $\text{HE}\nu$'s can even occur when $\omega_{\text{HE}\nu} = 0$. However and in spite of this resemblance, it should be kept in mind that these two phenomena completely differ in spirit and *have nothing to do with each other*. Although for the occurrence of fast conversions certain criteria need to be fulfilled [31], the short-scale conversions of $\text{HE}\nu$'s in a bath of the LE ones is a generic phenomenon provided that there are two populations of neutrinos of which one is dominant.

Flavor conversions of $\text{HE}\nu$'s induced by the static bath of the $\text{LE}\nu$'s is also distinct from the phenomenon of ordinary collective oscillations in dense neutrino media. While the latter is a nonlinear phenomenon with a high level of coupling, the former is a linear phenomenon where $H_{\nu\nu}$ solely provides a constant background for the flavor evolution of $\text{HE}\nu$'s. This implies that such flavor conversions of $\text{HE}\nu$'s is a noncollective phenomenon.




FIG. 2. Upper panel: Survival probability of $\text{HE}\nu_e$'s propagating in vacuum (dashed red curve), and in the bath of the $\text{LE}\nu$'s (solid blue curve). $\text{HE}\nu$'s can experience flavor conversions on scales much shorter than those expected in vacuum. Here, for illustrative purposes we have assumed $\mu = 10^{-7} \text{ cm}^{-1}$, $E_{\text{LE}\nu} = 10 \text{ MeV}$ and $E_{\text{HE}\nu} = 100 \text{ GeV}$. For the $\text{LE}\nu$ bath, $n_{\bar{\nu}_e}/n_{\nu_e} = 1.3$ is also fixed, although the survival probabilities are independent of $n_{\bar{\nu}_e}/n_{\nu_e}$ as long as it is not too close to 1. Lower panel: The survival probabilities as a function of the $\text{HE}\nu$ energy, where the diamonds, points, and squares are the survival probabilities of ν_e , ν_μ , and ν_τ obtained from the simulations, respectively, and the black lines are the corresponding analytical solutions. We assume an initial flavor ratio of $\nu_e : \nu_\mu : \nu_\tau = 1 : 0 : 0$. We here set $\theta_{12} = 33.6^\circ$, $\theta_{23} = 47.2^\circ$, $\theta_{13} = 8.5^\circ$ and $\delta_{\text{CP}} = 0$. Antineutrinos behave exactly in the same manner.

Also note that the relevant n_ν 's can be many orders of magnitude smaller than the values for which collective oscillations of MeV neutrinos are expected, due to the much smaller $\omega_{\text{HE}\nu}$.

In order to see how the short-scale flavor conversions of $\text{HE}\nu$'s changes the expected $\nu_e : \nu_\mu : \nu_\tau$ ratio on earth, one can average the survival probabilities over a few oscillations. As indicated in our upcoming work [32], such

an averaging process in our two-beam model corresponds to averaging over the neutrino angular distribution in a more-realistic, multiangle neutrino gas. The average survival probabilities then reach a steady state which does not depend on the details of the simulation (apart from the neutrino mixing parameters as discussed in the following), shown in the lower panel of Fig. 2. This behaviour can be understood analytically as follows. In the mass basis, the ν SI Hamiltonian is diagonal with its kk -th component being $h_k \propto \sum_{\alpha} |U_{\alpha k}|^2 (\rho_{\alpha\alpha} - \bar{\rho}_{\alpha\alpha})$, where $\bar{\rho}_{\alpha\alpha}^{(-)}$ are the initial (anti)neutrino occupation numbers in flavor α . This comes from the fact that $H_{\nu\nu}$ is nearly determined only by the $\text{LE}\nu$'s which are in the mass eigenstates. Then the $\text{HE}\nu$ density matrix in the mass basis, $\tilde{\rho}$, evolves as,

$$\tilde{\rho}_{ij}(t) = \tilde{\rho}_{ij}(0) e^{-i(h_i - h_j)t}, \quad (6)$$

implying that the averaged flavor ratio, $\nu_e : \nu_{\mu} : \nu_{\tau}$, can be written as,

$$|U_{ek}|^2 |U_{\alpha k}|^2 f_{\alpha} : |U_{\mu k}|^2 |U_{\alpha k}|^2 f_{\alpha} : |U_{\tau k}|^2 |U_{\alpha k}|^2 f_{\alpha}, \quad (7)$$

where there is a summation over α and k , and $f_e : f_{\mu} : f_{\tau}$ is the initial flavor ratio at the production region. The black lines in the lower panel of Fig. 2 indicate the analytical flavor ratios in Eq. (7), which show a perfect agreement with the numerical results.

Note that the average density matrix in the flavor basis is equal to the one expected after the neutrino decoherence, and is also independent of the neutrino energy. This behaviour indeed results from the fact that the $\text{HE}\nu$'s oscillate very quickly about $H_{\nu\nu}$ and consequently, they end up in the mass eigenstates (on average). Hence, in summary, short-scale conversions of $\text{HE}\nu$'s induced by the ambient gas of the $\text{LE}\nu$'s lead to their decoherence on scales which can be shorter than their natural decoherence length [28] by *many orders of magnitude*.

Although such short-scale oscillations and the resulting decoherence of $\text{HE}\nu$'s is an interesting phenomenon by itself and could in principle impact the physics of their propagation by modifying their flavor ratio at the source, we here discuss a few important cases in which the induced conversions of $\text{HE}\nu$'s can be observable on Earth.

Once $\text{HE}\nu$'s leave the $\text{LE}\nu$ bath, they should propagate in the dense ejecta, where $n_e \gg n_{\nu}$ [33–37]. In particular, for engine-driven transients, the $\text{HE}\nu$ production region is surrounded with the stellar or merger ejecta, whose density is much larger than that in the production region. This means that in solving Eq. (4) for the neutrino propagation in this region, we ignore the $H_{\nu\nu}$ term. In order to account for the decoherence experienced by neutrinos in the $\text{LE}\nu$ bath, we start with an initial density matrix which is a time average of the one in Eq. (6). In addition, for the H_m term we consider a blue supergiant matter profile from Ref. [38], as an example (30 M_{\odot} BSG in Ref. [37]). Needless to say, the short-scale conversions of $\text{HE}\nu$'s will impact the outcome of the matter effect and correspondingly, their flavor ratio on Earth. This is illustrated clearly in Fig. 3 for a




FIG. 3. The expected $\nu_e : \nu_{\mu} : \nu_{\tau}$ ratio on Earth in the absence and presence of ν SI for different neutrino energies. Note that the matter effect is included in both cases, assuming the density profile of a blue supergiant, and the ratio with ν SI is very close to the total flavor equipartition. In addition, the green star indicates the total flavor equipartition expected from the propagation of $\text{HE}\nu$'s in a bath of *oscillating* $\text{LE}\nu$'s, as discussed in the text. Note that apart from the matter only case for the 1 TeV neutrinos, the other ones are almost on top of each other. It is also illuminating to keep in mind that the final flavor states are specific to the initial flavor composition of 1:2:0 and can vary under different circumstances.

case with the initial flavor ratio $\nu_e : \nu_{\mu} : \nu_{\tau} = 1 : 2 : 0$ in the normal mass ordering. This is particularly interesting considering the energy-independent nature of the $\text{LE}\nu$ -induced short-scale oscillations of $\text{HE}\nu$'s. Although the pure matter effect shows a clear sign of energy dependence here [37], it is almost independent of the neutrino energy in the presence of ν SI. This is even the case at high energies where the muon damping is expected to occur in such dense environments [39, 40]. This can provide one with a new observable indication of neutrino flavor mixing caused by ν SI.

So far we have assumed that the bath of the $\text{LE}\nu$'s is completely decohered. However, the situation could be different. On the one hand, since R_{diss} can be as low as 10^8 cm, the solar-mass channel $\text{LE}\nu$'s could be still in phase since $l_{\text{coh},\odot} \sim \sigma_x E_{\nu}^2 / \Delta m_{\text{sol}}^2 \gtrsim 10^8$ cm. Moreover, due to the possibility for the existence of $\mu \gtrsim 10^{-4} \text{ cm}^{-1}$, the decoherence of $\text{LE}\nu$'s *might* be suppressed [41] in the atmospheric channel as well and $\text{LE}\nu$'s can experience a sort of (partial) collective oscillations in the production region of HE ones. Such erratic conversions of $\text{LE}\nu$'s will lead to the total flavor equipartition of $\text{HE}\nu$'s regardless of their initial flavor content, as indicated by the green star in Fig. 3. This phenomenon will be discussed in more details in our upcoming work [32]¹.

In some of the beyond Standard Model (SM) theories of

¹ The only difference of this case with the results shown in Fig. 2

particle physics, neutrinos can experience neutrino non-standard self-interactions (ν NSSI) [42, 43]. Such ν NSSI modify Eq. (5) to [44–46]

$$H_{\nu\nu,\mathbf{p}} = \sqrt{2}G_F \int \frac{d^3p'}{(2\pi)^3} (1 - \mathbf{v} \cdot \mathbf{v}') \{ \hat{G}(\ell_{\mathbf{p}'} - \bar{\ell}_{\mathbf{p}'}) \hat{G} + \hat{G} \text{Tr}[(\ell_{\mathbf{p}'} - \bar{\ell}_{\mathbf{p}'}) \hat{G}] \}, \quad (8)$$

where \hat{G} contains information about ν NSSI ($\hat{G} = \mathbb{1}$ in SM). For example, in the vector mediator scenario, we may have $\mathcal{L}_{\text{eff}} \supset G_F [\hat{G}^{\alpha\beta} \bar{\nu}_\alpha \gamma^\mu \nu_\beta] [\hat{G}^{\xi\eta} \bar{\nu}_\xi \gamma_\mu \nu_\eta]$, and its ν NSSI components are related to the vector mediator mass m_V and the coupling strength g by $|\hat{G}^{\alpha\beta}| \propto g^2/m_V^2$. The current constraints on ν NSSI are model dependent and strong for the mediator mass below MeV energies. For heavier mediators, the constraints from the early universe are rather weak, e.g., $|\hat{G}^{\alpha\beta}| \lesssim 10^7$ [47], although laboratory constraints can be stronger in the limited parameter space [48]. It has been suggested that spectral modulations and time delays of $\text{HE}\nu$'s enable us to study the unexplored parameter space of ν NSSI [49–54]. We point out that coherent ν SI-induced oscillations of $\text{HE}\nu$'s can be used as a novel probe of ν NSSI. This is illustrated in Fig. 4 where the red region shows the impact of ν NSSI. The flavor content is expected to have observable sensitivity to ν NSSI, i.e., a $\sim 10\%$ change of flavor ratio is caused by $|\hat{G}^{\alpha\beta}| \sim 0.1$. This means that one could probe such weak couplings with this effect.




FIG. 4. The expected $\nu_e : \nu_\mu : \nu_\tau$ ratio after $\text{HE}\nu$'s escape their production region, in the presence of ν NSSI. The triangle, circle, and square indicate the ratio in SM while the red region shows how the ratio changes in the presence of ν NSSI for the 1 : 2 : 0 case. Here the red region is created by choosing a large set of randomly populated $\hat{G}^{\alpha\beta}$ assuming that $|\hat{G}^{\alpha\beta}| < 1$ (for $\alpha \neq \beta$). Via coherent ν SI, the final $\text{HE}\nu$ flavor ratio is very sensitive to the ν NSSI.

V. CONCLUSION

We have brought to light a novel phenomenon, in which a class of high-energy cosmic neutrino emission can ex-

perience flavor conversions induced by the copious $\text{LE}\nu$ background, on scales much shorter than their intrinsic vacuum oscillation wavelengths. Unlike the celebrated phenomenon of collective oscillations of MeV neutrinos in a dense neutrino medium, the unearthed flavor conversions of high-energy cosmic neutrinos is a noncollective phenomenon in spirit.

This intriguing phenomenon can occur when $\text{HE}\nu$'s from relativistic outflows launched at the core-collapse of massive stars or at the mergers propagate in the bath of the already-decohered lower energy neutrinos from the central engine. Despite the small number density of $\text{HE}\nu$'s which can be insufficient to result in their own collective oscillations, their presence can lead to short-scale conversions of $\text{HE}\nu$'s on scales determined by the density of $\text{LE}\nu$'s. The background-induced conversions of $\text{HE}\nu$'s change their flavor content in an energy-independent manner and takes the $\text{HE}\nu$ gas to a state which is diagonal in the mass basis. This way they cause an induced decoherence of $\text{HE}\nu$'s on scales which are many orders of magnitude shorter than their natural decoherence lengths. Such a modification of the $\text{HE}\nu$'s at source can impact the physics of the phenomena occurring during their propagation, such as neutrino decay, scattering, etc. We also point out a few possibilities where such short-scale induced decoherence can directly impact the flavor ratio of $\text{HE}\nu$ on Earth, including the matter effect of $\text{HE}\nu$'s, the ν NSSI, and the total flavor equipartition due to an oscillating ambient $\text{LE}\nu$ gas.

Our study provides the first step toward understanding this intriguing phenomenon and further exploration is needed to better understand its implications. This is yet another context where neutrino-neutrino interactions can play a crucial role in their flavor evolution, and also motivates further investigations into *multimessenger* high-energy emission from GRBs, CCSNe and NSMs.

ACKNOWLEDGMENTS

We are grateful to Georg Raffelt and Huaiyu Duan for very insightful discussions and their comments on the manuscript. S.A. acknowledges support by the German Research Foundation (DFG) through the Collaborative Research Centre “Neutrinos and Dark Matter in Astro- and Particle Physics (NDM),” Grant SFB-1258, and under Germany's Excellence Strategy through the Cluster of Excellence ORIGINS EXC-2094-390783311. This work of K.M. is supported by the NSF Grant No. AST-1908689, No. AST-2108466 and No. AST-2108467, and KAKENHI No. 20H01901 and No. 20H05852.

-
- [1] J. N. Bahcall, P. Meszaros, 5-GeV to 10-GeV neutrinos from gamma-ray burst fireballs, *Phys. Rev. Lett.* 85 (2000) 1362–1365. [arXiv:hep-ph/0004019](#), [doi:10.1103/PhysRevLett.85.1362](#).
- [2] P. Meszaros, M. J. Rees, Multi GeV neutrinos from internal dissipation in GRB fireballs, *Astrophys. J. Lett.* 541 (2000) L5–L8. [arXiv:astro-ph/0007102](#), [doi:10.1086/312894](#).
- [3] K. Murase, K. Kashiyama, P. Mészáros, Subphotospheric Neutrinos from Gamma-Ray Bursts: The Role of Neutrons, *Phys. Rev. Lett.* 111 (2013) 131102. [arXiv:1301.4236](#), [doi:10.1103/PhysRevLett.111.131102](#).
- [4] K. Kashiyama, K. Murase, P. Mészáros, Neutron-Proton-Converter Acceleration Mechanism at Subphotospheres of Relativistic Outflows, *Phys. Rev. Lett.* 111 (2013) 131103. [arXiv:1304.1945](#), [doi:10.1103/PhysRevLett.111.131103](#).
- [5] K. Murase, B. Dasgupta, T. A. Thompson, Quasithermal Neutrinos from Rotating Protoneutron Stars Born during Core Collapse of Massive Stars, *Phys. Rev. D* 89 (4) (2014) 043012. [arXiv:1303.2612](#), [doi:10.1103/PhysRevD.89.043012](#).
- [6] P. Mészáros, E. Waxman, TeV neutrinos from successful and choked gamma-ray bursts, *Phys. Rev. Lett.* 87 (2001) 171102. [arXiv:astro-ph/0103275](#), [doi:10.1103/PhysRevLett.87.171102](#).
- [7] S. Razzaque, P. Meszaros, E. Waxman, TeV neutrinos from core collapse supernovae and hypernovae, *Phys. Rev. Lett.* 93 (2004) 181101, [Erratum: *Phys. Rev. Lett.* 94, 109903 (2005)]. [arXiv:astro-ph/0407064](#), [doi:10.1103/PhysRevLett.94.109903](#).
- [8] S. Ando, J. F. Beacom, Revealing the supernova-gamma-ray burst connection with TeV neutrinos, *Phys. Rev. Lett.* 95 (2005) 061103. [arXiv:astro-ph/0502521](#), [doi:10.1103/PhysRevLett.95.061103](#).
- [9] K. Murase, K. Ioka, TeV-PeV Neutrinos from Low-Power Gamma-Ray Burst Jets inside Stars, *Phys. Rev. Lett.* 111 (12) (2013) 121102. [arXiv:1306.2274](#), [doi:10.1103/PhysRevLett.111.121102](#).
- [10] S. S. Kimura, K. Murase, I. Bartos, K. Ioka, I. S. Heng, P. Mészáros, Transejecta high-energy neutrino emission from binary neutron star mergers, *Phys. Rev. D* 98 (4) (2018) 043020. [arXiv:1805.11613](#), [doi:10.1103/PhysRevD.98.043020](#).
- [11] M. Ackermann, et al., Fundamental Physics with High-Energy Cosmic Neutrinos, *Bull. Am. Astron. Soc.* 51 (2019) 215. [arXiv:1903.04333](#).
- [12] M. Ackermann, et al., High-Energy and Ultra-High-Energy Neutrinos (3 2022). [arXiv:2203.08096](#).
- [13] C. A. Argüelles, et al., Snowmass White Paper: Beyond the Standard Model effects on Neutrino Flavor, in: 2022 Snowmass Summer Study, 2022. [arXiv:2203.10811](#).
- [14] S. Pastor, G. Raffelt, Flavor oscillations in the supernova hot bubble region: Nonlinear effects of neutrino background, *Phys. Rev. Lett.* 89 (2002) 191101. [arXiv:astro-ph/0207281](#).
- [15] H. Duan, G. M. Fuller, J. Carlson, Y.-Z. Qian, Simulation of Coherent Non-Linear Neutrino Flavor Transformation in the Supernova Environment. 1. Correlated Neutrino Trajectories, *Phys. Rev. D* 74 (2006) 105014. [arXiv:astro-ph/0606616](#), [doi:10.1103/PhysRevD.74.105014](#).
- [16] H. Duan, G. M. Fuller, J. Carlson, Y.-Z. Qian, Coherent Development of Neutrino Flavor in the Supernova Environment, *Phys. Rev. Lett.* 97 (2006) 241101. [arXiv:astro-ph/0608050](#), [doi:10.1103/PhysRevLett.97.241101](#).
- [17] K. Kashiyama, K. Murase, S. Horiuchi, S. Gao, P. Meszaros, High energy neutrino and gamma ray transients from relativistic supernova shock breakouts, *Astrophys. J. Lett.* 769 (2013) L6. [arXiv:1210.8147](#), [doi:10.1088/2041-8205/769/1/L6](#).
- [18] K. Murase, New Prospects for Detecting High-Energy Neutrinos from Nearby Supernovae, *Phys. Rev. D* 97 (8) (2018) 081301. [arXiv:1705.04750](#), [doi:10.1103/PhysRevD.97.081301](#).
- [19] G. Sigl, G. Raffelt, General kinetic description of relativistic mixed neutrinos, *Nucl. Phys. B* 406 (1993) 423–451. [doi:10.1016/0550-3213\(93\)90175-0](#).
- [20] G. M. Fuller, R. W. Mayle, J. R. Wilson, D. N. Schramm, Resonant neutrino oscillations and stellar collapse, *Astrophys. J.* 322 (1987) 795.
- [21] J. T. Pantaleone, Dirac neutrinos in dense matter, *Phys. Rev. D* 46 (1992) 510–523. [doi:10.1103/PhysRevD.46.510](#).
- [22] D. Nötzold, G. Raffelt, Neutrino dispersion at finite temperature and density, *Nucl. Phys. B* 307 (1988) 924.
- [23] L. Wolfenstein, Neutrino Oscillations in Matter, *Phys. Rev. D* 17 (1978) 2369–2374, [294(1977)]. [doi:10.1103/PhysRevD.17.2369](#).
- [24] S. P. Mikheyev, A. Yu. Smirnov, Resonance Amplification of Oscillations in Matter and Spectroscopy of Solar Neutrinos, *Sov. J. Nucl. Phys.* 42 (1985) 913–917, [305(1986)].
- [25] M. Beuthe, Oscillations of neutrinos and mesons in quantum field theory, *Phys. Rept.* 375 (2003) 105–218. [arXiv:hep-ph/0109119](#), [doi:10.1016/S0370-1573\(02\)00538-0](#).
- [26] E. K. Akhmedov, A. Y. Smirnov, Paradoxes of neutrino oscillations, *Phys. Atom. Nucl.* 72 (2009) 1363–1381. [arXiv:0905.1903](#), [doi:10.1134/S1063778809080122](#).
- [27] J. Kersten, A. Y. Smirnov, Decoherence and oscillations of supernova neutrinos, *Eur. Phys. J. C* 76 (6) (2016) 339. [arXiv:1512.09068](#), [doi:10.1140/epjc/s10052-016-4187-5](#).
- [28] Y. Farzan, A. Y. Smirnov, Coherence and oscillations of cosmic neutrinos, *Nucl. Phys. B* 805 (2008) 356–376. [arXiv:0803.0495](#), [doi:10.1016/j.nuclphysb.2008.07.028](#).
- [29] R. F. Sawyer, Speed-up of neutrino transformations in a supernova environment, *Phys. Rev. D* 72 (2005) 045003. [arXiv:hep-ph/0503013](#), [doi:10.1103/PhysRevD.72.045003](#).
- [30] R. F. Sawyer, Neutrino cloud instabilities just above the neutrino sphere of a supernova, *Phys. Rev. Lett.* 116 (8) (2016) 081101. [arXiv:1509.03323](#), [doi:10.1103/PhysRevLett.116.081101](#).

is that we here allow for flavor oscillations of the $LE\nu$ gas rather than fix it to be in the mass state.

- [31] T. Morinaga, Fast neutrino flavor instability and neutrino flavor lepton number crossings (3 2021). [arXiv:2103.15267](#).
- [32] S. Abbar, J. A. Carpio, K. Murase, In preparation.
- [33] O. Mena, I. Mocioiu, S. Razzaque, Oscillation effects on high-energy neutrino fluxes from astrophysical hidden sources, *Phys. Rev. D* 75 (2007) 063003. [arXiv:astro-ph/0612325](#), [doi:10.1103/PhysRevD.75.063003](#).
- [34] S. Sahu, B. Zhang, Effect of Resonant Neutrino Oscillation on TeV Neutrino Flavor Ratio from Choked GRBs, *Res. Astron. Astrophys.* 10 (2010) 943–949. [arXiv:1007.4582](#), [doi:10.1088/1674-4527/10/10/001](#).
- [35] S. Razzaque, A. Y. Smirnov, Flavor conversion of cosmic neutrinos from hidden jets, *JHEP* 03 (2010) 031. [arXiv:0912.4028](#), [doi:10.1007/JHEP03\(2010\)031](#).
- [36] D. Xiao, Z. G. Dai, TeV–PeV Neutrino Oscillation of Low-luminosity Gamma-ray Bursts, *Astrophys. J.* 805 (2) (2015) 137. [arXiv:1504.01603](#), [doi:10.1088/0004-637X/805/2/137](#).
- [37] J. Carpio, K. Murase, Oscillation of high-energy neutrinos from choked jets in stellar and merger ejecta, *Phys. Rev. D* 101 (12) (2020) 123002. [arXiv:2002.10575](#), [doi:10.1103/PhysRevD.101.123002](#).
- [38] S. E. Woosley, T. A. Weaver, The Evolution and explosion of massive stars. 2. Explosive hydrodynamics and nucleosynthesis, *Astrophys. J. Suppl.* 101 (1995) 181–235. [doi:10.1086/192237](#).
- [39] T. Kashti, E. Waxman, Flavoring astrophysical neutrinos: Flavor ratios depend on energy, *Phys. Rev. Lett.* 95 (2005) 181101. [arXiv:astro-ph/0507599](#), [doi:10.1103/PhysRevLett.95.181101](#).
- [40] S. Karmakar, S. Pandey, S. Rakshit, Astronomy with energy dependent flavour ratios of extragalactic neutrinos, *JHEP* 10 (2021) 004. [arXiv:2010.07336](#), [doi:10.1007/JHEP10\(2021\)004](#).
- [41] E. Akhmedov, J. Kopp, M. Lindner, Collective neutrino oscillations and neutrino wave packets, *JCAP* 09 (2017) 017. [arXiv:1702.08338](#), [doi:10.1088/1475-7516/2017/09/017](#).
- [42] Z. Bialynicka-Birula, Do Neutrinos Interact between Themselves?, *Nuovo Cim.* 33 (1964) 1484–1487. [doi:10.1007/BF02749481](#).
- [43] D. Y. Bardin, S. M. Bilenky, B. Pontecorvo, On the ν – ν interaction, *Phys. Lett. B* 32 (1970) 121–124. [doi:10.1016/0370-2693\(70\)90602-7](#).
- [44] G. Sigl, G. Raffelt, General kinetic description of relativistic mixed neutrinos, *Nuclear Physics B* 406 (1-2) (1993) 423–451.
- [45] M. Blennow, A. Mirizzi, P. D. Serpico, Nonstandard neutrino-neutrino refractive effects in dense neutrino gases, *Phys. Rev. D* 78 (2008) 113004. [arXiv:0810.2297](#), [doi:10.1103/PhysRevD.78.113004](#).
- [46] A. Das, A. Dighe, M. Sen, New effects of non-standard self-interactions of neutrinos in a supernova, *JCAP* 05 (2017) 051. [arXiv:1705.00468](#), [doi:10.1088/1475-7516/2017/05/051](#).
- [47] M. Archidiacono, S. Hannestad, Updated constraints on non-standard neutrino interactions from Planck, *JCAP* 07 (2014) 046. [arXiv:1311.3873](#), [doi:10.1088/1475-7516/2014/07/046](#).
- [48] J. M. Berryman, et al., Neutrino Self-Interactions: A White Paper, in: 2022 Snowmass Summer Study, 2022. [arXiv:2203.01955](#).
- [49] K. Ioka, K. Murase, IceCube PeV–EeV neutrinos and secret interactions of neutrinos, *PTEP* 2014 (6) (2014) 061E01. [arXiv:1404.2279](#), [doi:10.1093/ptep/ptu090](#).
- [50] K. C. Y. Ng, J. F. Beacom, Cosmic neutrino cascades from secret neutrino interactions, *Phys. Rev. D* 90 (6) (2014) 065035, [Erratum: *Phys. Rev. D* 90, 089904 (2014)]. [arXiv:1404.2288](#), [doi:10.1103/PhysRevD.90.065035](#).
- [51] M. Ibe, K. Kaneta, Cosmic neutrino background absorption line in the neutrino spectrum at IceCube, *Phys. Rev. D* 90 (5) (2014) 053011. [arXiv:1407.2848](#), [doi:10.1103/PhysRevD.90.053011](#).
- [52] K. Blum, A. Hook, K. Murase, High energy neutrino telescopes as a probe of the neutrino mass mechanism, *arXiv e-prints* (2014) [arXiv:1408.3799](#) [arXiv:1408.3799](#).
- [53] A. DiFranzo, D. Hooper, Searching for MeV-Scale Gauge Bosons with IceCube, *Phys. Rev. D* 92 (9) (2015) 095007. [arXiv:1507.03015](#), [doi:10.1103/PhysRevD.92.095007](#).
- [54] I. M. Shoemaker, K. Murase, Probing BSM Neutrino Physics with Flavor and Spectral Distortions: Prospects for Future High-Energy Neutrino Telescopes, *Phys. Rev. D* 93 (8) (2016) 085004. [arXiv:1512.07228](#), [doi:10.1103/PhysRevD.93.085004](#).