

Galaxy morpho-Z with neural Networks (GaZNets). I. Optimized accuracy and outlier fraction from Imaging and Photometry

Rui Li^{1,2}, Nicola R. Napolitano^{3,4}, Haicheng Feng^{1,5}, Ran Li^{1,2}, Valeria Amaro³, Linghua Xie³, Crescenzo Tortora⁶, Maciej Bilicki⁷, Massimo Brescia⁶, Stefano Caviuoti^{6,8}, and Mario Radovich⁹

¹ School of Astronomy and Space Science, University of Chinese Academy of Sciences, Beijing 100049, China
e-mail: liruiww@gmail.com

² National Astronomical Observatories, Chinese Academy of Sciences, 20A Datun Road, Chaoyang District, Beijing 100012, China

³ School of Physics and Astronomy, Sun Yat-sen University, Zhuhai Campus, 2 Daxue Road, Xiangzhou District, Zhuhai, P. R. China
e-mail: napolitano@mail.sysu.edu.cn

⁴ CSST Science Center for Guangdong-Hong Kong-Macau Great Bay Area, Zhuhai, China, 519082

⁵ Yunnan Observatories, Chinese Academy of Sciences, Kunming, 650011, Yunnan, People's Republic of China

⁶ INAF – Osservatorio Astronomico di Capodimonte, Salita Moiariello 16, 80131 - Napoli, Italy

⁷ Center for Theoretical Physics, Polish Academy of Sciences, Al. Lotników 32/46, 02-668 Warsaw, Poland

⁸ INFN - Sezione di Napoli, via Cinthia 9, 80126 Napoli, Italy

⁹ INAF - Osservatorio Astronomico di Padova, via dell'Osservatorio 5, 35122 Padova, Italy

Received September 15, 1996; accepted March 16, 1997

ABSTRACT

Aims. In the era of large sky surveys, photometric redshifts (photo- z) represent crucial information for galaxy evolution and cosmology studies. In this work, we propose a new Machine Learning (ML) tool called Galaxy morpho-Z with neural Networks (GaZNet-1), which uses both images and multi-band photometry measurements to predict galaxy redshifts, with accuracy, precision and outlier fraction superior to standard methods based on photometry only.

Methods. As a first application of this tool, we estimate photo- z of a sample of galaxies in the Kilo-Degree Survey (KiDS). GaZNet-1 is trained and tested on $\sim 140\,000$ galaxies collected from KiDS Data Release 4 (DR4), for which spectroscopic redshifts are available from different surveys. This sample is dominated by bright ($MAG_AUTO < 21$) and low redshift ($z < 0.8$) systems, however, we could use ~ 6500 galaxies in the range $0.8 < z < 3$ to effectively extend the training to higher redshift. The inputs are the r -band galaxy images plus the 9-band magnitudes and colors, from the combined catalogs of optical photometry from KiDS and near-infrared photometry from the VISTA Kilo-degree Infrared survey.

Results. By combining the images and catalogs, GaZNet-1 can achieve extremely high precision in normalized median absolute deviation (NMAD=0.014 for lower redshift and NMAD=0.041 for higher redshift galaxies) and low fraction of outliers (0.4% for lower and 1.27% for higher redshift galaxies). Compared to ML codes using only photometry as input, GaZNet-1 also shows a $\sim 10 - 35\%$ improvement in precision at different redshifts and a $\sim 45\%$ reduction in the fraction of outliers. We finally discuss that, by correctly separating galaxies from stars and active galactic nuclei, the overall photo- z outlier fraction of galaxies can be cut down to 0.3%.

Key words. Galaxies: distances and redshifts – Machine learning – Methods: data analysis

1. Introduction

In the last decade, the Stage III sky surveys, e.g., Kilo-Degree Survey (KiDS, de Jong et al. 2013), Hyper Suprime-Cam (HSC; Aihara et al. 2018), Dark Energy Survey (DES; The Dark Energy Survey Collaboration 2005), have provided images of hundreds of millions of galaxies at optical or near-infrared (NIR) wavelengths. These surveys have achieved significant advances in cosmology (e.g., Hildebrandt et al. 2017; Hikage et al. 2019; Abbott et al. 2022) and galaxy formation and evolution (e.g., Roy et al. 2018; Greco et al. 2018; Goulding et al. 2018; Adhikari et al. 2021), but, at the same time, have left many open questions about the overall cosmological model (Di Valentino et al. 2021).

In the next decade, the Stage IV surveys (Weinberg et al. 2013), e.g., Euclid (Laureijs et al. 2011a), Vera Rubin Legacy Survey in Space and Time (VR/LSST; Ivezić et al. 2019), China Space Station Telescope (CSST; Zhan 2018), will observe bil-

lions of galaxies with photometric bands ranging from the ultraviolet to the NIR. This unprecedented amount of data will help us to get a deeper insight into cosmology and galaxy evolution. For instance, we will be able to gain a more detailed understanding of the dark matter distribution in the universe, constrain the equation of state of the dark energy with weak lensing (e.g., Laureijs et al. 2011b; Hildebrandt et al. 2017; Abbott et al. 2018; Gong et al. 2019; Joachimi et al. 2021; Heymans et al. 2021), study the mass-size relation of galaxies at higher redshift ($z > 1.0$), and explore the stellar and dark matter assembly in galaxies and clusters (e.g., Yang et al. 2012; Moster et al. 2013; Behroozi et al. 2019, Tortora & Napolitano 2022) over enormous statistical samples.

To achieve real breakthroughs in these areas, accurate galaxy redshifts are essential, as, by providing object distances and lookback time, they permit to trace those objects back in time. Precise redshifts can only be estimated from galaxy spectra: current spectroscopic surveys, such as the Sloan Digital Sky Sur-

vey (SDSS, Ahumada et al. 2020), the Galaxy and Mass Assembly (GAMA Baldry et al. 2018a) and the Dark Energy Spectroscopic Instrument (DESI, DESI Collaboration et al. 2016) have collected data for millions of galaxies, while future surveys, e.g., the 4-meter Multi-Object Spectroscopic Telescope (4MOST, de Jong et al. 2019), plan to expand spectroscopic measurements to samples of hundreds of millions of galaxies. However, due to the limited observation depth and prohibitive exposure times, it is impossible to spectroscopically follow up the even larger and fainter samples of billions of galaxies expected in future imaging surveys.

A fast, low-cost alternative is offered by photometric redshifts (photo- z) estimated from deep, multi-band photometry. The idea of photo- z was initially proposed by Baum (1962), where they used a redshift-magnitude relation to predict the redshifts from the galaxy luminosities. Without spectroscopic observations and the knowledge of galaxy evolution, the relation could still provide acceptable redshifts, even using only a limited number of filters. Later, this method was adopted to extensively estimate galaxy redshifts (e.g., Couch et al. 1983; Koo 1985; Connolly et al. 1995; Connolly 1997; Wang et al. 1998). However, albeit straightforward, this method has some limitations: 1) the redshift-magnitude relation is inferred in advance from bright galaxies via spectroscopy, and 2) the relation is hard to extend to fainter galaxies. Another method used to determine photo- z is spectral-energy-distribution (SED, hereafter) fitting. This method is based on galaxy templates, both theoretical and empirical.

By fitting the observed multi-band photometry to the SED from galaxy templates, one can infer individual galaxy photo- z . With knowledge of galaxy types and their evolution with redshift, this method can be expanded to faint galaxies, and even extrapolated to redshifts higher than the spectroscopic limit. There is a variety of photo- z codes based on SED fitting. Among the most popular ones, there is HyperZ (Bolzonella et al. 2000), which makes use of multi-band magnitudes of galaxies and the corresponding errors to best fit the SED templates by minimizing a given χ^2 function. An extension of HyperZ, known as Bayesian photometric redshifts (BPZ, Benítez 2000), is another popular photo- z tool. Instead of simple χ^2 minimization, BPZ introduces the prior knowledge of the redshift distribution of magnitude-limited samples under a Bayesian framework, which effectively reduces the number of catastrophic outliers in the predictions.

Besides these fitting tools, Machine Learning (ML) algorithms, especially Artificial Neural Networks (ANNs), have started to be extensively used to determine galaxy photo- z (e.g., Collister et al. 2007; Abdalla et al. 2008; Banerji et al. 2008). Given a training sample of galaxies with spectroscopic redshifts, ML algorithms can learn the relationship between redshift and multi-band photometry. If the training sample covers a representative redshift range and the ML model is well trained, photo- z can be obtained with extremely high precision. Different tools for photo- z based on ML have been successfully tested on multi-band photometry data, for example estimating photo- z with ANNs (ANNz, Collister & Lahav 2004; ANNz2 Sadeh et al. 2016) or the Multi-Layer Perceptron trained with Quasi-Newton Algorithm (MLPQNA, Cavuoti et al. 2012, Amaro et al. 2021).

Accurate photometry measurements are extremely important for ML and SED fitting methods, as the presence of noisy or biased photometry would end up in large scatter and a high outlier fraction in the predicted values. For instance, biased photometry is typically produced in the case of close galaxy pairs or in the presence of bright neighbors. On top of that, there are well-

known degeneracies between colors and redshift plaguing late-type systems, in particular, as high- z star-forming galaxies can be confused with lower redshift ellipticals.

These examples suggest that there might be some crucial information encoded in images that can help solving typical systematics, affecting the methods based on photometry only.

ML has been shown to be able to learn galaxy properties like their size, morphology, and their environment from images. This information can help suppress catastrophic errors and improve the accuracy of the photo- z predictions. In recent years, many studies have been trying to estimate photo- z directly from multi-band images using deep learning. A first attempt was presented by Hoyle (2016), where they estimated photo- z with a Deep Neural Network (DNN) applied to full galaxy imaging data. Lately, a similar approach has been applied to data from the Sloan Digital Sky Survey and Hyper Suprime-Cam Subaru Strategic Program (e.g., D’Isanto & Polsterer 2018; Pasquet et al. 2019; Schuld et al. 2021; Dey et al. 2021). These analyses showed that unbiased photo- z can be estimated directly from multi-band images. A more simplistic method for taking morphology features into account - e.g. size, ellipticity and Sérsic index - has been proposed in Soo et al. (2018), where they added structural parameters to the photometric catalogs used in standard ANNs.

In this paper, we develop a new ML method to estimate the *morpho- z* , i.e. redshifts estimated from the combination of images and catalogs of photometry and color measurements. In the following, we distinguish these *morpho- z* from the redshift predicted from photometry only, the classical *photo- z* , and from the ones obtained from images only, which - for convenience - we call *morpho- z* . This is the first time such a technique has been developed and applied to real data: specifically, we will use optical images and optical+NIR multi-band photometry from the KiDS survey. Just before the submission of this paper, a similar approach was proposed by Zhou et al. (2021), but this latter work is based on (CSST) simulated data only.

This work is organized as follows. In §2, we describe how to build the ML models and to collect the training and testing samples. In §3, we train the networks and show the performance of the tools. In §4 and §5, we discuss the results and draw some conclusions.

2. The ML method

In this work, we intend to couple standard ML regression tools, usually applied to galaxy multi-band photometry, with Deep Learning techniques, to improve the estimates of galaxy redshifts using the information from features distilled from galaxy images.

In particular, we want to address the following questions: 1) can redshifts be estimated directly from multi-band images of KiDS galaxies, and how does typical accuracy compare to ML tools based on integrated photometry and color measurements only? 2) If any, how much improvement in precision and scatter can images and catalogs add to tools combining all together?

To answer these questions, we have developed and compared four ML tools to estimate the galaxy redshifts. These differ - among each other - for the type of input data they can work with. In this section, we start by describing the structures and the training of these four tools.

2.1. Network architectures

The ML parts of our networks are constituted by ANNs. These have been proved to work well on catalogs made of magnitudes




Fig. 1. The Machine Learning models are used in this work. Up: CNN structure of GaZNet-I4 with only galaxy images as input. Middle: ANN structure of GaZNet-C4 and GaZNet-C9, with only catalog as input; bottom: structure of GaZNet-1, fed by both galaxies images and the corresponding catalogs.

and color measurements (e.g., Collister et al. 2007; Abdalla et al. 2008; Banerji et al. 2008; Cavioti et al. 2015; Brescia et al. 2014; de Jong et al. 2017; Bilicki et al. 2018, 2021). A typical ANN structure consists of three main parts: input, hidden, and output layers. The input and the output layers are used to load the data in the network and to issue the predictions. The hidden layers, composed of fully connected artificial neurons in a sequence of multiple layers, are used to extract features. These features are subsequently abstracted to allow the networks to determine the final outputs. In redshift estimates, the inputs of the networks are catalogs of some form of multi-band aperture photometry of galaxies, i.e. a measurements of the total flux in different filters, usually from optical to the NIR wavelengths.

The Deep Learning components of the four tools are constituted by Convolutional Neural Networks (CNN, Cun et al. 1990), which are an effective family of algorithms for feature extraction from images. CNNs mimic the biological perception mechanisms with convolution operations. This makes them specially suitable for image processing, pattern recognition and other tasks relative to images (e.g., Domínguez Sánchez et al. 2018; Ackermann et al. 2018; Walmsley et al. 2020; Čiprijanović et al. 2020; Li et al. 2020, 2021a,b; Tohill et al. 2021). The CNNs have become popular years after its introduction, because of the significant progress in the graphics processing unit (GPU) technology.

Here below, we introduce the structures of the first series of algorithms for “Galaxy morpho-Z with neural Networks” (GaZNets, hereafter). These are introduced to perform galaxy

morpho/photo-zs using different combinations of inputs, including multi-band photometry and imaging (see Fig. 1). In details:

- **GaZNet-I4.** This is a CNN model, which makes use of 4 optical bands (u, g, r, i) galaxy images, with a cutout size of $8 \times 8''$ (corresponding to 40×40 pixels, see §2.2), as input. The model is a slightly modified architecture from VGGNet (Simonyan & Zisserman 2014). It is constituted of four blocks made of different numbers of convolutional layers. Each of the first two blocks contains two layers, and of the other two blocks contain three layers. After the four blocks, a flatten layer is used to transform the high dimensional features into one-dimensional features. Finally, we adopt 3 fully connected layers to combine the low-level features into higher-level ones and output the predicted redshift.
- **GaZNet-C4.** This is a simple ANN structure with 2 blocks made of 4 fully connected layers, separated by a flatten layer. The input is an optical 4-band (u, g, r, i) catalog of magnitude and color measurements. Since we use the information from the same bands, comparing GaZNet-I4 and GaZNet-C4 allows us to quantify the impact of the imaging and photometry on the redshift estimates.
- **GaZNet-C9.** It has the same structure as GaZNet-C4, but it is input with the 4-band optical catalogs from KiDS plus the 5-band catalogs from the VISTA Kilo-degree Infrared Galaxy survey (VIKING, Edge et al. 2014, see §2.2 for details). Using a broader wavelength baseline, GaZNet-C9 will allow us to estimate the impact of the multi-band coverage on the ANN redshift predictions.




Fig. 2. The distribution of some relevant parameters of the training and testing data. In the top row, number counts are in linear scale, while in the bottom row number counts are in logarithmic scale. The first panel on the left shows the original spectroscopic sample of the 148 521 galaxies collected in KiDS+VIKING. However, only the 134 148 galaxies located between the two vertical dashed lines ($\text{spec-}z = 0.04$ and $\text{spec-}z = 3$) are used in this work for training and testing the GaZNets. For these galaxies, we show the MAG_AUTO and SNRs in the second and third panels.

– **GaZNet-1.** This is the reference network we have developed: the input is the combination of the r -band images and the multi-band photometry catalogs. The GaZNet-1 has been designed to have a two-path structure. The first path comprises 4 blocks as GaZNet-I4, while the second is made of 8 fully connected layers as GaZNet-C4 and C9. After a flatten layer, the features from each path are concentrated together. Finally, 5 fully connected layers are added to combine the features from images and catalogs to generate the final redshift predictions.

Of the four tools illustrated above, the first three are mainly designed to test the impact of the different inputs on the final redshift estimates. Being constructed with the same structure assembled in the final GaZNet-1, i.e. the one to be used for science, they guarantee the homogeneity of the treatment of the input data (see Fig. 1).

In this first series of GaZNets, we do not consider the magnitude ratios between different bands as inputs, although there are experiments suggesting that they can improve the precision (see e.g., D’Isanto et al. 2018, Nakoneczny et al. 2019). We plan to implement this in future analyses, because here we are interested in checking the advantages of the combination of images and photometry compared to previous analyses made on the same data (see §3.2.3). Also, in this analysis, we focus on redshift point estimates. In the future, we plan to expand the capabilities of the GaZNets to estimate the probability density function $p(z)$ for each galaxy. This can be achieved by Mixture Density Networks (e.g., Rhea et al. 2021, Wang et al. 2022) or Bayesian Networks (Gal & Ghahramani 2015; Kendall & Gal 2017), and some works with these two networks have been done (e.g., D’Isanto & Polsterer 2018; Ramachandra et al. 2021; Podsztaev et al. 2022). We will also evaluate the performance of the $p(z)$ using cumulative distribution function (CDF) and CDF-based metrics, such as the Kolmogorov–Smirnov (KS) statistic,

the Cramer-von Mises statistic and the Anderson–Darling (AD) statistic (see details in Schmidt et al. 2020).

2.2. Training and testing data

The dataset used in this work is collected from KiDS and VIKING, two twin surveys covering the same 1350 deg^2 sky area, in optical and NIR bands, respectively. KiDS observations are carried out with the VST/Omegacam telescope (Capaccioli & Schipani 2011; Kuijken 2011) in 4 optical filters (u, g, r, i), with a spatial resolution of $0.2''/\text{pixel}$. The r -band images are observed with the best seeing (average FWHM $\sim 0.7''$), and its mean limiting AB magnitude (5σ in a $2''$ aperture) is 25.02 ± 0.13 . The seeing of the other 3 bands (u, g and i) is slightly worse than that of the r -band i.e. FWHMs $< 1.1''$, and the mean limiting AB magnitudes are also fainter, i.e. 24.23 ± 0.12 , 25.12 ± 0.14 , 23.68 ± 0.27 for u, g and i , respectively (Kuijken et al. 2019).

VIKING is carried out with the VISTA/VIRCAM (Sutherland et al. 2015) and aims at complementing KiDS observations with five NIR bands (Z, Y, J, H and K_s). The median value of the seeing in the images is $\sim 0.9''$ (Sutherland et al. 2015), and the AB magnitude depths are 23.1, 22.3, 22.1, 21.5 and 21.2 in the five bands (Edge et al. 2013), respectively.

In particular, the galaxy sample used in this work is made of 148 521 objects for which spectroscopic redshifts ($\text{spec-}z$ s, hereafter) are available from different surveys, such as the Galaxy And Mass Assembly survey (GAMA, Driver et al. 2011) data release 2 and 3, the zCOSMOS (Lilly et al. 2007), the Chandra Deep Field South (CDFS, Szokoly et al. 2004), and the DEEP2 Galaxy Redshift Survey (Newman et al. 2013). The $\text{spec-}z$ range of the galaxies covers quite a large baseline, between $\sim 0 - 7$, although the distribution is far to be uniform. Indeed, as shown in Fig. 2, the number of galaxies at higher redshift ($z \gtrsim 0.8$) is much smaller than the one at lower redshift. In the same figure, we can see a peak of distribution at $\text{spec-}z < 0.6$. It comes from the GAMA survey, which is the most complete spectro-

scopic surveys adopted, with $\sim 95.5\%$ completeness for r -band magnitude $\text{MAG_AUTO} < 19.8$ (Baldry et al. 2018b). Similarly, we see a second peak at $\text{spec-}z \sim 2.5$, due to the quite deep observations from zCOSMOS. Overall, this sample is dominated by bright and low redshift galaxies ($0.04 < z < 0.8$), however, between $0.8 < z < 3$, it still contains ~ 6500 galaxies with a quite uniform redshift distribution that can be used as training sample to extend the predictions to higher redshift. Due to the unbalanced redshift coverage, we expect the accuracy of the predictions to have a strong variation with redshift. However, we will check if the final estimates meet the accuracy and precision requirements for cosmological and galaxy formation studies (see e.g., LSST Science Collaboration et al. 2009). After this redshift cut, the final sample is made of 134 148 galaxies. The distributions of the r -band Kron-like magnitude, MAG_AUTO , obtained by SExtractor (Bertin & Arnouts 1996) for these galaxies, and their signal-to-noise ratio (SNR, defined as the inverse value of the error of MAG_AUTO , are also reported in Fig. 2. Finally, the 134 148 galaxies are divided into three datasets, 100 000 for training, 14 148 for validation, and 20,000 for testing and error statistical analysis.

The u, g, r, i band images, with size of $8 \times 8''$, are cutout from KiDS DR4 (Kuijken et al. 2019). The corresponding catalogs, made up of 9 Gaussian Aperture and point spread function (GAAP) magnitudes ($u, g, r, i, Z, Y, J, H, K_s$) and 8 derived colors (e.g., $u - g, g - r, r - i$ etc.), are directly selected from the KiDS public catalog¹. The GAAP magnitudes have been measured on Gaussian-weighted apertures, modified per-source and per-image, therefore providing seeing-independent flux estimates across different observations/bands, reducing the bias of colors (see detail in Kuijken et al. 2015, 2019). The extinction was also considered in the measurement of the GAAP magnitudes.

2.3. External photo- z catalog by MLPQNA

To test the performances of GaZNet-1 against other ML based photo- z methods, we have collected an external photo- z catalog obtained from MLPQNA for the same KiDS galaxies we have used as testing sample. This allows us to perform a quantitative comparison of diagnostics like accuracy, scatter, and fractions of outliers.

MLPQNA is an effective computing implementation of neural networks adopted for the first time to solve regression problems in the astrophysical context. A test on PHAT1 dataset (Hildebrandt et al. 2010) indicated that MLPQNA, with smaller bias and fewer outliers, performs better than most of the traditional standard SED fitting methods. This code has been used in some current sky surveys, e.g., KiDS (Cavuoti et al. 2015) and Sloan Digital Sky Survey (SDSS; Brescia et al. 2014). For our comparison, we adopt the MLPQNA photo- z catalog from Amaro et al. (2021), where they have used the same data presented in §2.2 to train and test their networks.

3. GaZNet training and testing

In §2.1 we have described the different GaZNets and anticipated that they accept either images or catalogs of galaxies as inputs, except the GaZNet-1, fed with both images and catalogs. In particular, for the first test of morpho- z predictions made with this latter, we choose only the r -band images, i.e. the ones with best

quality from KiDS, to combine with the 9-band photometry catalog. As we will demonstrate in §4, the multi-band imaging does not add sensitive improvements in the results for the higher computation time prize one has to pay.

In this section, we illustrate the procedures to train the networks and test their predicted photo- z s against the ground truth provided by the spec- z s of the test sample introduced in §2.2.

3.1. Training the networks

We train the networks by minimizing the “Huber” loss (see, Huber 1964; Friedman 1999) function with an “Adam” optimizer (Kingma & Ba 2014). The “Huber” loss is defined as

$$L_\delta(a) = \begin{cases} \frac{1}{2}(a)^2, & |a| \leq \delta \\ \delta \cdot (|a| - \frac{1}{2}\delta), & \text{otherwise.} \end{cases} \quad (1)$$

In which $a = y_{\text{true}} - y_{\text{pred}}$. y_{true} is the spec- z and y_{pred} is the predicted photo- z . δ is a parameter that can be pre-set. Given a δ (fixed to be 0.001 in this work), the loss will be a square error when the deviation of the prediction, $|a|$, is smaller than δ ; otherwise, the loss is reduced to a linear function. Compared to the commonly used Mean Square Error (MSE) or Mean Absolute Error (MAE) loss function defined as

$$\begin{aligned} \text{MAE} &= \frac{1}{n} \sum |z_{\text{pred}} - z_{\text{spec}}| \\ \text{MSE} &= \frac{1}{n} \sum (z_{\text{pred}} - z_{\text{spec}})^2. \end{aligned} \quad (2)$$

“Huber” loss is proved to be more accurate in such regression tasks (see detail discussion in Li et al. 2021a).

To guarantee a faster-reducing speed of the loss function, for each ML model, we set a larger learning rate of 0.001 at the beginning and train the networks for 30 epochs. In each epoch, the networks are trained on the training data and validated on the validation data to verify if further adjustments are needed to improve the overall accuracy. After the first training round, we reduce the learning rate to 0.0001, and load the pre-trained model with a “callback” operation. Then we train the networks for further 30 epochs. Changing the learning rate to a smaller value of 0.0001 can help the network converge to the global minimum, hence finding the best-trained model.

For the networks input with images, we also apply some data augmentations, including random shift, flip, and rotation (only 90° , 180° and 270°). We do not adopt any augmentation that needs interpolation algorithms², like crop, zoom, color-changing, and adding noise, since these operations would change the flux in the image pixels, affecting the magnitudes and colors of the galaxies.

Computing-time wise, with the NVIDIA RTX 2070 graphics processing unit (GPU), GaZNet-C4 and GaZNet-C9 need 28 minutes to complete the training and validation process, while GaZNet-1 takes about 134 minutes (including ~ 2.5 minutes for data reading) because of the time needed to process the r -band image data along with the magnitudes and colors. Compared with GaZNet-1, GaZNet-I4 does not significantly increase the time during the training and validation process. However, to deal with the 4 channel images, GaZNet need much larger memory of

¹ <https://kids.strw.leidenuniv.nl/DR4/access.php>

² Note that, even if a generic rotation does imply some interpolation due to pixel re-sampling, the adoption of $\pi/2$ multiples does not, because it preserves the overall geometry of the cutout, except the orientation.




Fig. 3. Comparison between the spectroscopic redshifts and the predicted photometric redshifts for different models. From top left to bottom right are the results from GaZNet-I, GaZNet-C, GaZNet and MLPQNA, respectively. Error bars represent the mean absolute errors (MAE), while the quoted numbers are the mean $|\delta z|$, in each bin.

GPU, and time of data reading increases significantly. Finally it needs ~ 4 times longer on data reading. We also estimated that to make predictions on 20 000 galaxies (see §3.2), in terms of the computing time for predictions, this is much shorter. GaZNet-C9 needs only ~ 6 seconds for the whole galaxies, while GaZNet-1 needs ~ 55 seconds, including ~ 30 seconds for data reading and ~ 25 seconds for prediction.

3.2. Testing the performance

After the training phase, we use the 20 000 testing galaxies to estimate the precision and the statistical errors of the redshift predictions from different GaZNets.

3.2.1. Statistical parameters

We define a series of statistical parameters to describe the overall performances: 1) the fraction of catastrophic outliers, 2) the mean bias, and 3) the normalized median absolute deviation (NMAD).

The fraction of the catastrophic outliers is defined as the fraction of galaxies with bias larger than 15% according to the following formula:

$$|\delta z| = \frac{|z_{\text{pred}} - z_{\text{spec}}|}{1 + z_{\text{spec}}} > 15\%. \quad (3)$$

where z_{spec} are the spec- z s of the test galaxies and z_{pred} are the predicted redshifts by the ML tools. This definition is usually

adopted for outliers in photo- z estimates (see details in e.g., Cavuoti et al. 2012, Amaro et al. 2021) and gives a measure of the fallibility of the method. In addition, the mean bias in this work is labeled as $\mu_{\delta z}$.

The normalized median absolute deviation (NMAD), between the predicted photo- z s and the true spec- z s, are defined as

$$\text{NMAD} = 1.4826 \times \text{median}(|\delta z - \text{median}(\delta z)|), \quad (4)$$

where δz comes from Eq. 3. NMAD allows us to quantify the scatter of the overall predictions in comparison to the ground truth, hence it is a measurement of the precision of the redshift estimates from the ML tools.

3.2.2. Predictions vs. ground truth

The testing results on 20 000 galaxies for the four GaZNets are shown in Fig. 3, where on the x-axis we plot the spec- z s as ground truth, and on the y-axis we plot the predicted redshifts. As a comparison, in the same figure, we also show the photo- z s estimated by the MLPQNA. We divided the galaxies into 6 redshift bins, and computed the mean absolute errors, shown as error bars, and the mean $|\delta z|$ defined in Eq. 3, reported as text. We use equally spaced bins to check the effect of the sampling as a function of the redshift.

From Fig. 3, a major feature one can see, at the first glance, is the odd coverage of the spec- z at high redshift ($z \gtrsim 0.8$), which we have also discussed in §2.2. This is a potential issue for all

methods, as a poor training set can introduce a large scatter in the predictions. Indeed, in Fig. 3 the δ_z tends to have an increasingly larger scatter at larger redshifts. This means that at $z \gtrsim 0.8$ the absolute scatters are dominated by the size of the training sample rather than the true intrinsic uncertainties of the methods. Unfortunately, this is a problem we cannot overcome with the current data and we need to wait for larger spec- z data samples to improve the precision at higher redshifts. However, given the current training set, we can still evaluate the relative performances of different methods and their ability to make accurate predictions even in the small training set regimes.

Given this necessary preamble, from Fig. 3 we see that unbiased photo- z can be obtained by GaZNet-I4, with only 4-band images as input, although it seems that this starts to deviate from the one-to-one relation at $z \gtrsim 1.5$. However, at these redshifts, a general trend of underestimating the ground truth is also shown by GaZNet-C4 and GaZNet-C9, although the latter uses the full photometry from the KiDS+VIKING dataset. Interestingly, looking at the scatter, GaZNet-I4 seems to perform better than GaZNet-C4 at all redshift bins and almost comparably to C9 in most cases.

The results from GaZNet-I4 demonstrate that morpho- z s from multi-band images are similar, if not potentially superior, to photo- z s from photometry in the same bands. This high-performance of morpho- z s is also confirmed by the noticeably smaller outlier fraction (1.5% for GaZNet-I4 and 2.2% for GaZNet-C4 in general). Even more interestingly, looking from the perspective of future surveys relying on a narrower wavelength baseline, such as the space missions Euclid and CSST, our results show that morpho- z s are not far from optical+NIR large photometric baselines in terms of accuracy, scatter, and the fraction of outliers. This is particularly true for $z < 1$, where, as seen in Fig. 3, the GaZNet-I4 shows less outliers than GaZNet-C9, while this latter shows a rather lower fraction of outliers at higher redshifts.

Moving to GaZNet-C9, the results show the impact of the broader wavelength baseline including the five NIR bands. Generally, photo- z determined by GaZNet-C9 show improved indicators in comparison to GaZNet-I4 and GaZNet-C4. From Fig. 3 we can see these coming from a better linear correlation, especially at $z > 1.5$, and smaller absolute errors. However, looking at the results in more detail, at $z > 1.5$ the presence of a rather large fraction of outliers causes the median values to diverge from the one-to-one relationship in a way similar to GaZNet-I4 and GaZNet-C4. It is hard to assess whether this is caused by the poor training sample, or it is an intrinsic shortcoming of the ML tool. In either cases, it is important to check whether using the information from images can improve this result.

Compared to GaZNet-C9, GaZNet-1 has overall better performances, with a tighter one-to-one relation, and smaller errors (by $\sim 10-35\%$) in all redshift bins. This is shown in the bottom-left panel of Fig 3. This result leads us to two main conclusions: 1) images (even a single high-quality band, see §4 for the test on multi-band imaging) provide crucial information to solve intrinsic issues related to the photometry only and improve all performances of the redshift predictions, in terms of accuracy, scatter, and outlier fraction. We will discuss the reason for this in §4; 2) due to the poor redshift coverage of the training sample at $z > 1$, the results we have obtained possibly represent a lower limit on the potential performances of the tool. No matter what, the GaZNet-1 reaches an excellent overall precision of $\delta_z = 0.038(1+z)$ up to $z = 3$, with an overall outlier fraction of 0.74%.

Table 1. Statistical properties of the predictions.

CNN model	Out. fr.	μ_{δ_z}	NMAD
Low redshift galaxies ($z \leq 0.8$)			
GaZNet-C9	0.007	0.0	0.016
GaZNet-1	0.004	0.0	0.014
MLPQNA	0.01	0.006	0.022
high redshift galaxies ($0.8 < z < 3$)			
GaZNet-C9	0.234	-0.067	0.073
GaZNet-1	0.127	-0.028	0.041
MLPQNA	0.216	-0.022	0.087

NOTE. —Outlier fraction, mean bias and NMAD (see §3.2.1) from different tools on lower ($z < 0.8$) and higher ($z > 0.8$) redshift galaxies.

3.2.3. Test vs. external catalogs

We can finally compare the performance of the four GaZNets versus the external catalogs. The MLPQNA is rather similar to the ANN method used for the GaZNets-C9, as it makes use of a similar algorithm and the same catalogs from KiDS DR4. From Fig. 3 we see that MLPQNA performs similar to GaZNet-I4 and better than GaZNet-C4. This is not surprising as the MLPQNA uses a larger wavelength baseline. This is particularly visible at higher redshift, where the predictions from MLPQNA are tighter distributed around the one-to-one relationship with the ground truth than GaZNet-I4 and GaZNet-C4.

3.2.4. Performance in space of redshift and magnitude

In the last subsections, we have shown that GaZNet-1 shows better performances than others tools, in terms of accuracy and precision. However, in Fig. 3, we have also seen a variation of these performances as a function of the redshift. Here, we want to more detailedly quantify this effect as well as the dependence on the magnitudes of the same performances. The reason for this diagnostics is to assess the impact of selection effects on the overall performances (see e.g. van den Busch et al. 2020). E.g., in §3.2.2, we have anticipated that the redshift sampling by the training sample can be one source of degradation of the performances at $z > 1$.

In Fig. 4 we plot the outlier fraction (out. fr.), mean bias (μ_{δ_z}) and scatter (NMAD) as functions of spec- z , photo- z , and r -band magnitudes. As comparison we plot the same relations for GaZNet-C9 and MLPQNA, the other two tools showing comparable performances to the GaZNet-1. The bottom row of the same Fig. 4, finally shows the distribution of the training sample in the same parameter space.

From this figure, the overall impression is that GaZNet-1 performs generally better than the other two tools in most, if not all, redshift and magnitude bins, with lower outlier fraction, smaller mean bias and scatter. We also see a clear correlation of the performances of all tools, included GaZNet-1, with the size and magnitude of the training sample in different redshift bins and the magnitudes of the training galaxies. All of the 3 tools performs quite well in the range of $z \lesssim 0.8$, where the training sample is about one order of magnitude larger, resulting on a more accurate training. To quantify the overall performance in this redshift range, in Tab 1 we report the global statistical parameters for these galaxies. The three tools all can achieve quite small outlier fractions ($\lesssim 0.01$), mean bias (close to 0) and scatters ($\lesssim 0.022$). Compared to the other two tools, GaZNet-1




Fig. 4. Outlier fraction (out. fr.), mean bias ($\mu_{\delta z}$), and Scatter (NMAD) as functions of spec- z , photo- z , and magnitudes in 20 bins. In each panel, blue line is for GaZNet-1, orange is for GaZNet-C9 and green is for MLPQNA. In the last row we also present the number distribution in the corresponding parameter space.

shows the best performance. In particular, its outlier fraction is 43% smaller than GaZNet-C9 and 60% smaller than MLPQNA.

At $z \gtrsim 0.8$, in Fig. 4 we see that the number of galaxies decreases rapidly, which produces a degradation of the performances of all tools. Interestingly, looking at the central panels, after $z \sim 1.5$, where the COSMOS spec- z sample is concentrated, the performances, especially in terms of scatter (NMAD), show a significant improvement up to $z \gtrsim 2.6$, where the spec- z of the training sample quickly drop in number again. This is also quantified in Tab 1 by the global statistical parameters for these higher redshift galaxies. Compared to GaZNet-C9, all the indicators from GaZNet-1 are significantly improved. The fraction of outliers, the mean bias and the scatter are decreased by 46%, 58%, and 44%, respectively. On the other hand, MLPQNA remains the tool performing worse.

A similar behavior of the performances is seen also as a function of the photo- z , as these latter closely follow the spec- z (see Fig. 3).

Going to the magnitude space, we find that all indicators shows very small values at $\text{MAG_AUTO} \lesssim 21$, which means that the redshift estimates for brighter galaxies are highly reliable. After r -band $\text{MAG_AUTO} \sim 21$, all indicators degrade, showing a worsening of the accuracy, precision and outlier fraction. Among the three tools, the GaZNet-1 is the one with better per-

formances, though. In particular, after $\text{MAG_AUTO} \sim 22$ there is a peak at 30%, which can be either driven by the poorer SNR of the systems, but more likely by the smaller statistics. In general, though, the GaZNet-1 has still a quite small outlier fraction ($\sim 1.0\%$).

Overall, it is clear that collecting more galaxies covering higher redshift ($z \gtrsim 0.8$) and fainter magnitudes ($\text{MAG_AUTO} \gtrsim 22$) will be essential for improving the performance of these ML tools, and the results collected here represents just a lower limit of the real performances that these tools, especially GaZNet-1, can achieve. However, even with the current training set, GaZNet-1 can provide even for high redshift galaxies results that satisfy the requirements for weak gravitational lensing studies in next generation ground-based surveys (e.g., $\text{NMAD}=0.05$ in VR/LSST, LSST Science Collaboration et al. 2009), although this has been currently tested only on a relatively bright sample with AB magnitude $\text{MAG_AUTO} \leq 22$ (see 4). For the low-redshift samples the GaZNet-1 is already well within the requirements for the same surveys and it is virtually science ready. In the future, we will look for more higher redshift galaxies from different spectroscopic surveys to build a less biased training sample and improve the performances at $z > 0.8$.

4. Discussion

In the previous section, we have compared the performances of the different GaZNets based on different architectures, including or not deep learning. We have also compared the GaZNets against external catalogs of photo- z based on traditional machine learning algorithms. The reason for us to develop different tools with an increasing degree of complexity, is to understand the impact of the different features in the final predictions. The main conclusion of this comparison is that GaZNet-1, using the combination of 9-band photometry and r -band imaging, clearly over-perform all the other tools, either developed by us or taken from literature, based on photometry only and no use of deep learning. We have also seen how deep learning only, applied to only 4-band optical images, can produce morpho- z that are more accurate than the photo- z from the corresponding photometry and equalize the performance of the 9-band photometry, except for redshift larger than $z > 0.8$. Overall, we have discussed that part of the over-performance of deep learning is concentrated on the outlier fraction. In this section, we investigate the reasons for these results and discuss the impact of some choices we have made in the set-up of the GaZNets presented in this first paper.

4.1. The outliers

From Table 1, it is evident that the major advantage from deep learning, applied to high-quality imaging, resides in the low outlier fraction. For GaZNet-1, this is smaller than the one of GaZNet-C9 by $\sim 43\%$ for low redshift galaxies ($z < 0.8$) and 46% for higher redshift galaxies ($z > 0.8$). Understanding the reasons for these results is important to figure out the source of systematics and plan next developments for more accurate morpho- z estimates.

To investigate the genesis of these outliers, we start by checking the galaxies for which the ML tools fail to obtain accurate photo- z s. In Fig. 5, we show the optical *gri* color-composed and the r -band images of representative outliers from GaZNet-C9, which are not outliers anymore for GaZNet-1. In each r -band image, we report the spec- z on the top, the GaZNet-C9 photo- z on the bottom-left and the GaZNet-1 morpho- z on the bottom-right. As a comparison, in Fig. 5, we also show outliers from GaZNet-C9 that are still outliers for GaZNet-1. The color images in this figure give a fair idea of the galaxy SEDs, while the r -band images illustrate the corresponding “morphological” features that the GaZNet-1 uses to improve the overall predictions.

From Fig. 5 we can distinguish four kinds of outliers for GaZNet-C9. The first one (A-row) is made of galaxies that are close to bright, often saturated, stars or bright large galaxies. In some cases, GaZNet-1 can improve the predictions and solve the discrepancy with the ground truth values (A-row). However, in some other cases, the environment is too confused to allow the CNN to guess correctly, despite the CNN can deblend the embedded source (E-row, see discussion below). The second type of outliers (B-row) are irregular galaxies or, generally, diffuse nearby systems. These systems are generally starforming and blue, similarly to the majority of high redshift galaxies. Thus, they typically have a higher chance to be confused with higher- z systems. In this case, the GaZNet-1 can recognize the complex morphology (knots, substructures, pseudo-arms etc), or a noisier surface brightness distribution, which are typical features of closer galaxies³. The third type of outliers (C-row) is made of

merging/interacting systems. For these systems, GaZNet-1 can solve the discrepancy using the information of the size of the two systems and the degree of details of the substructures, making these systems rather accurate to predict. The fourth type of outliers (D-row) is made of blue objects, generally high- z compact systems, sometimes also at low- z . In this case, again, the GaZNet-1 can make more accurate predictions, from the size and the round morphology.

With this insight on the way Deep Learning help improve the predictions of photo- z , we can now check where it still fails. This might give us valuable indications on how we can improve the GaZNet-1 performances in future analyses. In Fig. 5 we see three types of outliers also for the GaZNet-1.

The first one (E-row), similarly to the ones of GaZNet-C9 in the A-row above, is caused by the presence of large, bright systems. In these cases, GaZNet-1 has difficulties in either correctly deblending the source or correctly evaluating the size, especially if very compact. We stress here, though, that these outliers are generally fewer than all of the other kinds ($\sim 5\%$ of the total outliers for both GaZNet-1 and C9), and in the case of bright stars, can often be automatically masked out from catalogs. The second type (F-row) is made of galaxies that have odd sizes for their redshifts, e.g., small-sized low- z objects (ultra-compact galaxies? misclassified stars? etc.) or even large-sized high- z systems (very massive/luminosity systems? galaxies with large diffuse haloes? etc.). As for the previous type, these systems are also quite limited in number ($\sim 4\%$), and their failure also depends on the poor training sample. In general, these outliers do not represent a significant issue. The third type (G-row) is made of extremely compact, almost point-like and generally blue sources. These are the most abundant sample of outliers ($\sim 58\%$). Although their redshift distribution is relatively sparse, they have a very similar appearance, being mainly concentrated at $z > 1$ but with cases even at $z < 0.5$. There is a little chance that all these systems are misclassified as stars or very compact blue galaxies (e.g., blue nuggets), although we cannot exclude some of them to be either case. The only possible option is that these are galaxies hosting active galactic nuclei (AGNs) or quasars. If so, these represent a marginal fraction of the training sample, and for this reason, they are not accurately predicted.

The SED of a quasar is different from that of a typical galaxy (e.g., Feng et al. 2021), and a small fraction of quasars may not provide enough training samples. Besides, most quasars can present strong variability, since they are observed in different bands at different times. This introduces fictitious color terms that increase their uncertainties in photo- z measurements. To verify this assumption, we check the star/galaxy/quasar separation in Khramtsov et al. (2019), based on an ML classifier. We find that only $\sim 35\%$ of the 147 outliers are classified as galaxies. For the remaining $\sim 65\%$, about half are classified as quasars and half as stars. Regardless of the accuracy of the ML method to classify stars and quasars, this analysis confirms that only a minority of the catastrophic events are made of galaxies, consistently with our guess based on the visual inspection above. In particular, the three objects in Fig. 5-G, are all quasars in the ML classification.

If indeed, the outliers are dominated by misclassified stars and AGNs/quasars, we can easily figure that optimizing the classification of these groups of contaminants would reduce the overall outlier fraction down to a tiny value, $\sim 0.3\%$.

³ We can guess, here, that the CNN can learn the surface brightness fluctuation (SBF) of galaxies, which is a notorious distance indicator (see e.g., Cantiello et al. 2005).




Fig. 5. The g, r, i color-composited color images ($20'' \times 20''$) and the corresponding r -band images for some representative outliers. Rows A, B, C and D (blue framed) show the outliers in GaZNet-C9, which are no longer outliers in GaZNet-1 predictions. Rows E, F and G (red framed) show the objects that remain outliers both for GaZNet-1 and GaZNet-C9. In the r -band images we report the spec- z on the top, the GaZNet-C9 photo- z on the bottom left and the GaZNet-1 morpho- z on the bottom right.

4.2. Other tests

The four GaZNets illustrated in §2.1 and discussed in §3.2.2 and §4 have been distilled by a number of other models we have tested, with different kinds of inputs and different ML structures. Among these, we focus on two other experiments where we have tested two set-ups that, in principle, can impact the final results. The parameters describing the performances of these two further

configurations are shown in Table 2. Here below, we summarize their properties and the major results:

1. **GaZNet-81pix to test the cutout size.** The GaZNets work on images with a size of $8'' \times 8''$, which might be too small to collect the light of the whole galaxies and their environments, thus leaving some important features that the CNNs can not see. In order to check this, we have tested the GaZNet-1 on images with twice the size of each side

Table 2. Statistical properties of the predictions.

CNN model	Out. fr.	μ_{δ_z}	NMAD
Lower redshift galaxies			
GaZNet-81pix	0.004	0.001	0.015
GaZNet-C9I4	0.004	-0.002	0.015
higher redshift galaxies			
GaZNet-81pix	0.15	-0.039	0.045
GaZNet-C9I4	0.124	-0.033	0.033

NOTE. — Outlier fraction, mean bias and NMAD (see §3.2.1) from the further tools tested in §4.2.

($16'' \times 16''$). Compared with the previous result in Table 2, the parameters remain almost unchanged. A possible reason is that the features that the CNNs extract from images are concentrated in the high SNR regions of the galaxies, while the outer regions bring little information, about both the galaxy properties and the environment, to improve the redshift estimates. Using these arguments, one can ask whether the standard $8'' \times 8''$ cutouts are too large and one can use a smaller cutout. To check this, we also tested $4'' \times 4''$ and found slightly worse results, so we kept the $8'' \times 8''$ as the best choice.

- GaZNet-C9I4 to test the 9-band catalogs plus 4-band images:** GaZNet-1 makes use of 9-band catalogs and only r -band images. To check if the addition of images on other bands can produce a sensitive improvement, we train a GaZNet with the 4 optical band images plus the 9-band catalogs. We report the statistical parameters obtained with this new GaZNet in Table 2: generally speaking, there is no obvious improvement. Some tiny differences are compatible with the random statistical effects in the training process. Even taking these results at face values, compared to the GaZNet-1, the computing time registered by the GaZNet-C9I4 is almost 3 times longer. For these reasons, we can discard this solution for the poor cost-benefit ratio.

5. Conclusions

Several millions of galaxies have been observed in the third generation wide-field sky surveys, and tens of billions of galaxies will be observed in the next ten years by the fourth generation surveys from ground and space. This enormous amount of data provides an unprecedented opportunity to study in detail the evolution of galaxies, and constrain the cosmological parameters with unprecedented accuracy. To fully conduct these studies over the expected gigantic datasets, fast and accurate photo- z are indispensable. In this work, we have explored the feasibility of determining the redshift with ML by combining the images and photometry catalogs. We designed 4 ML tools, named GaZNet-I4, GaZNet-C4, GaZNet-C9 and GaZNet-1. The inputs for these tools are 4-band images, 4-band catalogs, 9-band catalogs, and r -band images plus 9-band catalogs, respectively. We have trained using a sample of $\sim 140,000$ spectra from different spectroscopic surveys. The training sample is dominated by bright ($MAG_AUTO < 21$) and low redshift ($z < 0.8$) galaxies, which provides a quite accurate knowledge base in this parameter space. On the other hand, the higher- z and fainter magnitudes are poorly covered by the training set. Despite that, we have shown that the four tools, especially GaZNet-1, still return accurate predictions also at $z > 0.8$.

In more details, our tests show that accurate morpho- z can be directly obtained from the multi-band images (u, g, r, i) by GaZNet-I4, with less outliers and smaller scatters than GaZNet-C4 using only four-band optical aperture photometry. We have also seen that the combination of optical and NIR photometry in 9-band catalogs, by GaZNet-C9, can provide a much better determination of photo- z . However, the information added by even one single-band high-quality images, as tested with our GaZNet-1, can achieve noticeable performances that highly overtake the ones registered for GaZNet-C9. The statistical errors are $\sim 10 - 35\%$ smaller at different redshift bins, while the outlier fraction reduces by 43% for lower redshift galaxies and 46% for higher redshift galaxies. We have estimated the variation of the scatter as a function of the redshift, over the range of $z = 0 - 3$, of the order of $\delta_z = 0.038(1 + z)$. This is heavily affected by the poor coverage of the training base at large redshifts and we expect to improve significantly this prediction by adding a few thousands more galaxies in this redshift range.

By visually inspecting the images of all outliers produced by the GaZNet-C9 and GaZNet-1, we have confidently demonstrated that the largest portion of the catastrophic estimates correspond to systems that are AGNs/quasars. This is corroborated by an independent ML classification from Khramtsov et al. (2019). If these contaminants are correctly separated from galaxies, the overall outlier fraction of GaZNet-1 can reduce to 0.3%. This is potentially an impressive result, that, combined with the rather high precision and small δ_z , will make the GaZNets performance close to the requirements for galaxy evolution and cosmology studies from the 4th generation surveys (e.g., Ivezić et al. 2019; Laureijs et al. 2011a, Zhan 2018).

Acknowledgements. Rui Li and Ran Li acknowledges the support of National Nature Science Foundation of China (Nos 11988101, 11773032, 12022306), the science research grants from the China Manned Space Project (No CMS-CSST-2021-B01, CMS-CSST-2021-A01) and the support from K.C.Wong Education Foundation. NRN acknowledge financial support from the “One hundred top talent program of Sun Yat-sen University” grant N. 71000-18841229. MB is supported by the Polish National Science Center through grants no. 2020/38/E/ST9/00395, 2018/30/E/ST9/00698, 2018/31/G/ST9/03388 and 2020/39/B/ST9/03494, and by the Polish Ministry of Science and Higher Education through grant DIR/WK/2018/12. This work is based on observations made with ESO Telescopes at the La Silla Paranal Observatory under programme IDs 177.A-3016, 177.A-3017, 177.A-3018 and 179.A-2004, and on data products produced by the KiDS consortium.

References

- Abbott, T. M. C., Abdalla, F. B., Alarcon, A., et al. 2018, Phys. Rev. D, 98, 043526
- Abbott, T. M. C., Aguena, M., Alarcon, A., et al. 2022, Phys. Rev. D, 105, 023520
- Abdalla, F. B., Amara, A., Capak, P., et al. 2008, MNRAS, 387, 969
- Ackermann, S., Schawinski, K., Zhang, C., Weigel, A. K., & Turp, M. D. 2018, MNRAS, 479, 415
- Adhikari, S., Shin, T.-h., Jain, B., et al. 2021, ApJ, 923, 37
- Ahumada, R., Prieto, C. A., Almeida, A., et al. 2020, ApJS, 249, 3
- Aihara, H., Arimoto, N., Armstrong, R., et al. 2018, PASJ, 70, S4
- Amaro, V., Cavauoti, S., Brescia, M., et al. 2021, Rejection Criteria Based on Outliers in the KiDS Photometric Redshifts and PDF Distributions Derived by Machine Learning, ed. I. Zelinka, M. Brescia, & D. Baron, Vol. 39, 245–264
- Baldry, I. K., Liske, J., Brown, M. J. I., et al. 2018a, MNRAS, 474, 3875
- Baldry, I. K., Liske, J., Brown, M. J. I., et al. 2018b, MNRAS, 474, 3875
- Banerji, M., Abdalla, F. B., Lahav, O., & Lin, H. 2008, MNRAS, 386, 1219
- Baum, W. A. 1962, in Problems of Extra-Galactic Research, ed. G. C. McVittie, Vol. 15, 390
- Behroozi, P., Wechsler, R. H., Hearin, A. P., & Conroy, C. 2019, MNRAS, 488, 3143
- Benítez, N. 2000, ApJ, 536, 571
- Bertin, E. & Arnouts, S. 1996, A&AS, 117, 393

- Bilicki, M., Dvornik, A., Hoekstra, H., et al. 2021, A&A, 653, A82
- Bilicki, M., Hoekstra, H., Brown, M. J. I., et al. 2018, A&A, 616, A69
- Bolzonella, M., Miralles, J. M., & Pelló, R. 2000, A&A, 363, 476
- Brescia, M., Cavuoti, S., Longo, G., & De Stefano, V. 2014, A&A, 568, A126
- Cantiello, M., Blakeslee, J. P., Raimondo, G., et al. 2005, ApJ, 634, 239
- Capaccioli, M. & Schipani, P. 2011, The Messenger, 146, 2
- Cavuoti, S., Brescia, M., Longo, G., & Mercurio, A. 2012, A&A, 546, A13
- Cavuoti, S., Brescia, M., Tortora, C., et al. 2015, MNRAS, 452, 3100
- Čiprijanović, A., Snyder, G. F., Nord, B., & Peek, J. E. G. 2020, Astronomy and Computing, 32, 100390
- Collister, A., Lahav, O., Blake, C., et al. 2007, MNRAS, 375, 68
- Collister, A. A. & Lahav, O. 2004, PASP, 116, 345
- Connolly, A. 1997, The Properties of High Redshift Galaxies : A Near-Infrared Redshift Survey at $1 < z < 2$, HST Proposal
- Connolly, A. J., Csabai, I., Szalay, A. S., et al. 1995, AJ, 110, 2655
- Couch, W. J., Ellis, R. S., Godwin, J., & Carter, D. 1983, MNRAS, 205, 1287
- Cun, Y. L., Boser, B., Denker, J. S., Henderson, D., & Jackel, L. D. 1990, Advances in neural information processing systems, 2, 396
- de Jong, J. T. A., Verdoes Kleijn, G. A., Erben, T., et al. 2017, A&A, 604, A134
- de Jong, J. T. A., Verdoes Kleijn, G. A., Kuijken, K. H., & Valentijn, E. A. 2013, Experimental Astronomy, 35, 25
- de Jong, R. S., Agertz, O., Berbel, A. A., et al. 2019, The Messenger, 175, 3
- DESI Collaboration, Aghamousa, A., Aguilar, J., et al. 2016, arXiv e-prints, arXiv:1611.00036
- Dey, B., Andrews, B. H., Newman, J. A., et al. 2021, arXiv e-prints, arXiv:2112.03939
- Di Valentino, E., Anchordoqui, L. A., Akarsu, Ö., et al. 2021, Astroparticle Physics, 131, 102606
- D’Isanto, A., Cavuoti, S., Gieseke, F., & Polsterer, K. L. 2018, A&A, 616, A97
- D’Isanto, A. & Polsterer, K. L. 2018, A&A, 609, A111
- Domínguez Sánchez, H., Huertas-Company, M., Bernardi, M., Tuccillo, D., & Fischer, J. L. 2018, MNRAS, 476, 3661
- Driver, S. P., Hill, D. T., Kelvin, L. S., et al. 2011, MNRAS, 413, 971
- Edge, A., Sutherland, W., Kuijken, K., et al. 2013, The Messenger, 154, 32
- Edge, A., Sutherland, W., & The Viking Team. 2014, VizieR Online Data Catalog, 2329, 0
- Feng, H.-C., Liu, H. T., Bai, J. M., et al. 2021, ApJ, 912, 92
- Friedman, J. H. 1999, Annals of Statistics, 29, 1189
- Gal, Y. & Ghahramani, Z. 2015, arXiv e-prints, arXiv:1506.02158
- Gong, Y., Liu, X., Cao, Y., et al. 2019, ApJ, 883, 203
- Goulding, A. D., Greene, J. E., Bezanson, R., et al. 2018, PASJ, 70, S37
- Greco, J. P., Greene, J. E., Strauss, M. A., et al. 2018, ApJ, 857, 104
- Heymans, C., Tröster, T., Asgari, M., et al. 2021, A&A, 646, A140
- Hikage, C., Oguri, M., Hamana, T., et al. 2019, PASJ, 71, 43
- Hildebrandt, H., Arnouts, S., Capak, P., et al. 2010, A&A, 523, A31
- Hildebrandt, H., Viola, M., Heymans, C., et al. 2017, MNRAS, 465, 1454
- Hoyle, B. 2016, Astronomy and Computing, 16, 34
- Huber, P. J. 1964, The Annals of Mathematical Statistics, 35, 73
- Ivezić, Ž., Kahn, S. M., Tyson, J. A., et al. 2019, ApJ, 873, 111
- Joachimi, B., Lin, C. A., Asgari, M., et al. 2021, A&A, 646, A129
- Kendall, A. & Gal, Y. 2017, arXiv e-prints, arXiv:1703.04977
- Khramtsov, V., Sergeyev, A., Spiniello, C., et al. 2019, A&A, 632, A56
- Kingma, D. P. & Ba, J. 2014, arXiv e-prints, arXiv:1412.6980
- Koo, D. C. 1985, AJ, 90, 418
- Kuijken, K. 2011, The Messenger, 146, 8
- Kuijken, K., Heymans, C., Dvornik, A., et al. 2019, A&A, 625, A2
- Kuijken, K., Heymans, C., Hildebrandt, H., et al. 2015, MNRAS, 454, 3500
- Laureijs, R., Amiaux, J., Arduini, S., et al. 2011a, arXiv e-prints, arXiv:1110.3193
- Laureijs, R., Amiaux, J., Arduini, S., et al. 2011b, arXiv e-prints, arXiv:1110.3193
- Li, R., Napolitano, N. R., Roy, N., et al. 2021a, arXiv e-prints, arXiv:2111.05434
- Li, R., Napolitano, N. R., Spiniello, C., et al. 2021b, ApJ, 923, 16
- Li, R., Napolitano, N. R., Tortora, C., et al. 2020, ApJ, 899, 30
- Lilly, S. J., Le Fèvre, O., Renzini, A., et al. 2007, ApJS, 172, 70
- LSST Science Collaboration, Abell, P. A., Allison, J., et al. 2009, arXiv e-prints, arXiv:0912.0201
- Moster, B. P., Naab, T., & White, S. D. M. 2013, MNRAS, 428, 3121
- Nakoneczny, S., Bilicki, M., Solarz, A., et al. 2019, A&A, 624, A13
- Newman, J. A., Cooper, M. C., Davis, M., et al. 2013, ApJS, 208, 5
- Pasquet, J., Bertin, E., Treyer, M., Arnouts, S., & Fouchez, D. 2019, A&A, 621, A26
- Podsztaevak, O., Škoda, P., & Tvrdík, P. 2022, Bayesian SZNet: Bayesian deep learning to predict redshift with uncertainty, Astrophysics Source Code Library, record ascl:2204.004
- Ramachandra, N., Chaves-Montero, J., Alarcon, A., et al. 2021, arXiv e-prints, arXiv:2111.12118
- Rhea, C., Hlavacek-Larrondo, J., Rousseau-Nepton, L., & Prunet, S. 2021, Research Notes of the American Astronomical Society, 5, 276
- Roy, N., Napolitano, N. R., La Barbera, F., et al. 2018, MNRAS, 480, 1057
- Sadeh, I., Abdalla, F. B., & Lahav, O. 2016, PASP, 128, 104502
- Schmidt, S. J., Malz, A. I., Soo, J. Y. H., et al. 2020, MNRAS, 499, 1587
- Schuldt, S., Suyu, S. H., Cañameras, R., et al. 2021, A&A, 651, A55
- Simonyan, K. & Zisserman, A. 2014, arXiv e-prints, arXiv:1409.1556
- Soo, J. Y. H., Moraes, B., Joachimi, B., et al. 2018, MNRAS, 475, 3613
- Sutherland, W., Emerson, J., Dalton, G., et al. 2015, A&A, 575, A25
- Szokoly, G. P., Bergeron, J., Hasinger, G., et al. 2004, ApJS, 155, 271
- The Dark Energy Survey Collaboration. 2005, arXiv e-prints, astro
- Tohill, C., Ferreira, L., Conselice, C. J., Bamford, S. P., & Ferrari, F. 2021, ApJ, 916, 4
- Tortora, C. & Napolitano, N. R. 2022, Frontiers in Astronomy and Space Sciences, 8, 197
- van den Busch, J. L., Hildebrandt, H., Wright, A. H., et al. 2020, A&A, 642, A200
- Walmsley, M., Smith, L., Lintott, C., et al. 2020, MNRAS, 491, 1554
- Wang, G.-J., Cheng, C., Ma, Y.-Z., & Xia, J.-Q. 2022, arXiv e-prints, arXiv:2207.00185
- Wang, Y., Bahcall, N., & Turner, E. L. 1998, AJ, 116, 2081
- Weinberg, D. H., Mortonson, M. J., Eisenstein, D. J., et al. 2013, Phys. Rep., 530, 87
- Yang, X., Mo, H. J., van den Bosch, F. C., Zhang, Y., & Han, J. 2012, ApJ, 752, 41
- Zhan, H. 2018, in 42nd COSPAR Scientific Assembly, Vol. 42, E1.16–4–18
- Zhou, X., Gong, Y., Meng, X.-M., et al. 2021, arXiv e-prints, arXiv:2112.08690