

Fully symmetric nonequilibrium response of stochastic systems

David Andrieux

We show that the nonequilibrium response of Markovian dynamics can be made fully symmetric, both near and far from equilibrium. This is achieved by varying the affinities along equivalence classes in the space of stochastic dynamics.

CONTEXT AND OBJECTIVES

Nonequilibrium response theory has long been studied near equilibrium, where general results such as the fluctuation-dissipation theorem or the Onsager reciprocity relations hold. Far from equilibrium, relations for the response coefficients have been derived as consequences of the fluctuation theorem for currents [1].

However, these nonequilibrium relations are much more intricate than their near equilibrium counterparts. In particular, they are not fully symmetric and involve combinations of multiple response coefficients.

Recently, the author derived a mapping between nonequilibrium fluctuations and an associated equilibrium dynamics [2, 3]. Building on this result, we demonstrate that the nonequilibrium response of currents is fully symmetric within an equivalence class of stochastic dynamics. To this end, we show that:

1. The space of stochastic dynamics can be structured into equivalence classes. Each equivalence class is determined by the composition of an equilibrium dynamics and all possible affinities
2. The equilibrium dynamics of an equivalence class determines the fluctuations of all (nonequilibrium) dynamics in that class
3. Within an equivalence class, the nonequilibrium response of currents is symmetric at all orders and for any values of the affinities (i.e., both near and far from equilibrium)

I. THE SPACE OF STOCHASTIC DYNAMICS IS STRUCTURED INTO EQUIVALENCE CLASSES DETERMINED BY EQUILIBRIUM DYNAMICS

We consider a Markov chain characterized by a transition matrix $P = (P_{ij}) \in \mathbb{R}^{N \times N}$ on a finite state space [4]. We assume that the Markov chain is primitive, i.e., there exists an n_0 such that P^{n_0} has all positive entries.

The space of stochastic dynamics can be partitioned into equivalence classes. We define the operator

$$\bar{P}_{ij} = \sqrt{P_{ij}P_{ji}} \quad (1)$$

and the corresponding *equivalence relation*


$$P \sim H \quad \text{if there exists a factor } \gamma \text{ such that } \bar{P} = \gamma \bar{H}. \quad (2)$$

This relation is reflexive, symmetric, and transitive. It thus forms a partition of the space of stochastic dynamics: every dynamics belongs to one and only one equivalence class.

An equivalence class $[P] = \{P' : P' \sim P\}$ is determined by the composition of an equilibrium dynamics E and the set of all possible affinities \mathbf{A} (Fig. 1, see Ref. [2] for a detailed construction). In particular, to any P we can associate an equilibrium dynamics $E[P]$ as follows. Let χ denote the Perron root of \bar{P} and its right Perron vector by x . Then if $D = \text{diag}(x_1, \dots, x_N)$ we have that

$$E[P] = \frac{1}{\chi} D^{-1} \bar{P} D \quad (3)$$

defines an equilibrium stochastic dynamics.

(a) Equivalence classes are parametrized by an equilibrium dynamics E and the set of affinities \mathbf{A} 

(b) Dynamics composing a given equivalence class




FIG. 1. **Equivalence classes and their elements.** (a) The equivalence class $[P]$ is parametrized by the equilibrium dynamics $E[P]$ and the set of all affinities \mathbf{A} [5]. (b) Dynamics composing an equivalence class typically depend non-linearly on the affinity. As an illustration, we consider a random walk on a ring composed of 3 sites. The equivalence class is defined by $\bar{E}_{1,2} = \bar{E}_{2,1} = 0.2$, $\bar{E}_{2,3} = \bar{E}_{3,2} = 0.4$, $\bar{E}_{3,1} = \bar{E}_{1,3} = 0.3$. The affinity A then fully defines each dynamics within $[E]$ and the corresponding transition probabilities $P_{1,2} = 1 - P_{1,3}$, $P_{2,3} = 1 - P_{2,1}$, $P_{3,1} = 1 - P_{3,2}$.

II. THE EQUILIBRIUM DYNAMICS DETERMINE THE FLUCTUATIONS OF ALL THE NONEQUILIBRIUM DYNAMICS WITHIN ITS EQUIVALENCE CLASS

We consider the thermodynamic currents (see Ref. [6] for a definition) and their fluctuations. The current fluctuations for a dynamics P are characterized by the generating function

$$Q_P(\boldsymbol{\lambda}) = \lim_{n \rightarrow \infty} -\frac{1}{n} \ln \left\langle \exp \left(-\boldsymbol{\lambda} \cdot \sum_{k=1}^n \mathbf{J}(k) \right) \right\rangle_P. \quad (4)$$

All cumulants can then be obtained by successive derivations of Q with respect to the counting parameters $\boldsymbol{\lambda}$:

$$Q_P(\boldsymbol{\lambda}) = \sum_{m=1}^{\infty} \frac{1}{m!} K_{\alpha_1 \dots \alpha_m}[P] \lambda_{\alpha_1} \cdots \lambda_{\alpha_m}. \quad (5)$$

In particular, the first-order cumulant equals the mean current, $K_{\alpha} = \partial Q / \partial \lambda_{\alpha}(\mathbf{0}) = \langle J_{\alpha} \rangle$.

We now consider the fluctuations of dynamics within the equivalence class $[P]$. We denote by $Q_{[P]}(\boldsymbol{\lambda}, \mathbf{A})$ the generating functions of the elements $(E[P], \mathbf{A})$ within $[P]$. We then have the

Theorem. *Let $[P]$ be an equivalence class and $E[P]$ its associated equilibrium dynamics. Then their current generating functions are related as*


$$Q_{[P]}(\boldsymbol{\lambda}, \mathbf{A}) = Q_E(\boldsymbol{\lambda} - \mathbf{A}/2) - Q_E(-\mathbf{A}/2). \quad (6)$$

DEMONSTRATION: In Ref. [3], we showed that the nonequilibrium current fluctuations of an arbitrary nonequilibrium dynamics P are determined by its corresponding equilibrium dynamics $E[P]$ according to

$$Q_P(\boldsymbol{\lambda}) = Q_E(\boldsymbol{\lambda} - \mathbf{A}/2) - Q_E(-\mathbf{A}/2). \quad (7)$$

By construction, all dynamics in the same equivalence class $P' \sim E$ will satisfy the relation (6) with the same function Q_E . As a result, relation (6) holds for all dynamics $P' \sim P$ with affinities \mathbf{A} . Conversely, for any affinity \mathbf{A} there exists a corresponding dynamics P' in the equivalence class. The relation (6) therefore holds for all \mathbf{A} within the same equivalence class. \square

(a) Generating functions within an equivalence class



(b) Generating functions are identical after translation




FIG. 2. **Generating functions for different dynamics within an equivalence class.** (a) The generating functions are numerically calculated for different values of the affinity A within the equivalence class. (b) The generating functions are identical images of each other and related according to (6). All parameters are identical to Fig. 1.

In other words, the generating function of the current Q_P is entirely determined by the equilibrium fluctuations Q_E (Fig 2) [7]. Therefore, $E[P]$ acts as a reference equilibrium dynamics that is intrinsic to P . We also note that the generating functions Q_P is directly related to the factor γ defining the equivalence relation (2). If $\bar{P} = \gamma \bar{E}$ then $Q_P = -2 \log \gamma$ [2].

III. THE NONEQUILIBRIUM RESPONSE IS SYMMETRIC WITHIN AN EQUIVALENCE CLASS

We now explore the consequences of Eq. (6) on the response theory. As the right-hand side of Eq. (6) only depends on the affinities through the translation $\lambda - \mathbf{A}/2$, the first order response of the mean current $\langle J_\alpha \rangle = K_\alpha(\mathbf{A})$ reads

$$\frac{\partial \langle J_\alpha \rangle}{\partial A_\beta}(\mathbf{A}) = -\frac{1}{2} \frac{\partial^2 Q_E}{\partial \lambda_\alpha \partial \lambda_\beta}(-\mathbf{A}/2) \quad (8)$$

which is fully symmetric in (α, β) for any \mathbf{A} (Fig. 3). When $\mathbf{A} = 0$ we recover the Onsager reciprocity relations and the Green-Kubo formula. Also, since as $Q_E(\lambda) = Q_E(-\lambda)$, the current is antisymmetric with respect to \mathbf{A} , $\langle J_\alpha \rangle(-\mathbf{A}) = -\langle J_\alpha \rangle(\mathbf{A})$, while its response $\partial \langle J_\alpha \rangle / \partial A_\beta$ is symmetric in \mathbf{A} .

More generally, the response of any cumulant is expressed as

$$\frac{\partial^l K_{\alpha_1 \dots \alpha_m}(\mathbf{A})}{\partial A_{\beta_1} \dots \partial A_{\beta_l}} = \left(-\frac{1}{2}\right)^l \frac{\partial^{m+l} Q_E}{\partial \lambda_{\beta_1} \dots \partial \lambda_{\beta_l} \partial \lambda_{\alpha_1} \dots \partial \lambda_{\alpha_m}}(-\mathbf{A}/2), \quad (9)$$

which is fully symmetric in $(\alpha_1, \dots, \alpha_m, \beta_1, \dots, \beta_l)$ for all \mathbf{A} . The cumulant response is symmetric in $\pm \mathbf{A}$ if $m + l$ is even, and antisymmetric otherwise.

This result is complementary to the traditional response theory. In the traditional response theory, the resulting dynamics shifts across both the affinity space and the equivalence classes as the affinities are varied [8]. In contrast, here the dynamics remains in its original equivalence class when varying the affinities (Fig. 4).

Note, however, that varying affinities while staying in a given equivalence class typically requires changing multiple transition probabilities at the same time (Fig 1b), which might not be feasible in all experimental systems. In any case, the present results offer a new way to explore nonequilibrium systems by leveraging their intrinsic dynamical characteristics.





FIG. 3. **Symmetry of the nonequilibrium current response.** We consider a 4-state system defined by the equivalence class $\bar{E}_{1,2} = \bar{E}_{2,1} = 0.2$, $\bar{E}_{2,3} = \bar{E}_{3,2} = 0.4$, $\bar{E}_{3,1} = \bar{E}_{1,3} = 0.5$, $\bar{E}_{3,4} = \bar{E}_{4,3} = 0.1$, $\bar{E}_{4,1} = \bar{E}_{1,4} = 0.1$. This system has 2 independent currents and affinities, here measured along the cycles (1, 2, 3) and (1, 3, 4). The symmetry $\partial J_1/\partial A_2 = \partial J_2/\partial A_1$ is verified for multiple values of the affinities (for visual clarity, we depict $-\partial J_2/\partial A_1$ and verify that it's the mirror image of $\partial J_1/\partial A_2$).

(a) Traditional response theory



(b) Present approach




FIG. 4. **Schematic comparison of the traditional and the present approach to the response theory.** In both cases, the system traces the same path in the space of affinities. (a) In the traditional response theory, both the affinities and equivalence classes change. (b) In the present approach, the system remains in its original equivalence class when varying the affinities.

Disclaimer. This paper is not intended for journal publication.

-
- [1] D. Andrieux and P. Gaspard, J. Chem. Phys. **48**, 571 (2004).
 [2] D. Andrieux, arXiv 1208.5699 (2012).
 [3] D. Andrieux, arXiv 1212.1807 (2012).
 [4] The same results can be readily obtained for continuous time Markov processes
 [5] For simplicity we depict the space of equilibrium dynamics as 1-dimensional.
 [6] J. Schnakenberg, Rev. Mod. Phys. **48**, 571 (1976).
 [7] The fluctuation theorem $Q(\boldsymbol{\lambda}) = Q(\mathbf{A} - \boldsymbol{\lambda})$ is satisfied as $Q_E(\boldsymbol{\lambda}) = Q_E(-\boldsymbol{\lambda})$ for an equilibrium dynamics.
 [8] As a result, the shape of the generating function changes in addition to being translated as the affinities are varied. This leads to additional contributions of the form $\partial Q/\partial A_\alpha \partial \lambda_\beta$ in the response theory, breaking the symmetry (9).