arXiv:2205.10994v1 [hep-ph] 23 May 2022

Prediction of a narrow exotic hadronic state with quantum numbers $J^{PC} = 0^{--}$

Teng Ji,^{1,2,*} Xiang-Kun Dong,^{1,2,†} Feng-Kun Guo,^{1,2,‡} and Bing-Song Zou^{1,2,3,§}

¹CAS Key Laboratory of Theoretical Physics, Institute of Theoretical Physics,

Chinese Academy of Sciences, Beijing 100190, China

²School of Physical Sciences, University of Chinese Academy of Sciences, Beijing 100049, China

³School of Physics, Central South University, Changsha 410083, China

Lots of charmonium-like structures have been observed. Most of them share the same quantum numbers with conventional charmonium states, with exceptions of those with an electric charge and/or strangeness. We show that a neutral and zero-strangeness charmonium-like exotic state with quantum numbers $J^{PC} = 0^{--}$, denoted as $\psi_0(4360)$, is a robust prediction in the hadronic molecular scenario, where the $\psi(4230), \psi(4360)$ and $\psi(4415)$ are identified as $D\bar{D}_1, D^*\bar{D}_1$ and $D^*\bar{D}_2^*$ bound states, respectively; the mass and width are predicted to be (4366 ± 18) MeV and less than 10 MeV, respectively. The interactions are calculated by the *t*-channel vector and pseudoscalar meson exchanges assisted by heavy quark spin symmetry. The coupled-channel effects and the *u*-channel pion exchange including full 3-body effects of the $D^*\bar{D}^*\pi$ intermediate states are carefully examined. The $\psi_0(4360)$ is significant in two folds: no 0^{--} hadron has been observed so far, and a study of this state will enlighten the understanding of the mysterious vector mesons between 4.2 and 4.5 GeV. We propose that such an exotic state can be searched for and uniquely identified in $e^+e^- \to \eta\psi_0(4360)$ by measuring the angular distribution of the outgoing η meson.

Introduction.—The study on exotic states beyond the conventional quark model [1, 2], where mesons and baryons are composed of a pair of quark-antiquark $(q\bar{q})$ and three quarks (qqq), has been a focus of hadron physics in the last two decades. Quantum chromodynamics (QCD), the underlying theory that guides the formation of hadrons from quarks and gluons, does not forbid the existence of exotic configurations, such as multiquark states (with more than 3 valence (anti)quarks), hybrid states (with gluonic excitations in addition to the valence quarks), glueballs (only gluons) and so on. Many candidates of such exotic states have been observed in experiments, see Refs. [3–16] for recent reviews on the experimental and theoretical status of exotic states. However, fundamental questions still remain unanswered, such as whether there is a dominant configuration for the excited hadrons and what that configuration (if any) should be.

Among the exotic states, those with exotic J^{PC} are of special interests since they cannot be ordinary $q\bar{q}$ mesons, such as $0^{--}, 0^{+-}, 1^{-+}$ and so on. Although dozens of exotic candidates have been observed in experiments, only a few of them have exotic J^{PC} , including $\pi_1(1400), \pi_1(1600)$ [17] and the most recently observed $\eta_1(1855)$ [18, 19], all of which lie in the light quark sector and have $J^{PC} = 1^{-+}$. Up to now, no signal of 0^{-} states occurs although many theoretical investigations predict the existence of such states as compact tetraquark states, hybrid states, glueballs or hadronic molecules. In Refs. [20, 21] the 0^{--} light flavor tetraquark states are predicted to be located at 1.35 GeV for a molecular configuration and 2.15 GeV for a compact state in the QCD Coulomb gauge approach. However, the QCD sum rules concluded that no such states exist below 2 GeV [22, 23]. For the 0^{--} hybrid states, the constituent gluon model [24], QCD Coulomb gauge approach [25] and QCD sum rules [26–28] all predict the existence but the resulting masses scatter between 1.6 and 3.3 GeV. The quenched lattice QCD predicted the ground state $0^{--} c\bar{c}g$ hybrid to be located at around 5.9 GeV [29]. Glueballs consisting of 3 gluons can have $J^{PC} = 0^{--}$ and the masses are predicted to be around 4 GeV [30], while the results in Refs. [31–33] are around 6 or 7 GeV. A hadronic molecule composed of $D\bar{D}_0^*$ with $J^{PC} = 0^{--}$ was predicted in Ref. [34] with a mass of around 4.25 GeV but note that the D_0^* is too wide to form a bound state [35, 36] and its mass listed in the Review of Particle Physics (RPP) [37] is too high (see Ref. [38] and references therein). In the tetraquark model of Ref. [39], a 0^{--} hidden-charm state was predicted below 4.4 GeV, which is accompanied by 2^{--} and 3^{--} states with masses below 4.1 GeV [40].

Hadronic molecules have a unique feature that they are close to the thresholds of their components, which results in the much less uncertainties of their masses and distinguishes them from other exotic configurations. Most of the experimental candidates of exotic states with small widths are found to be possible hadronic molecules, such as the $\chi_{c1}(3872)$ [41], the $Z_c(3900)^{\pm}$ [42–44], the P_c states [45] and the T_{cc}^+ [46, 47], see Refs. [7, 16, 48, 49 for reviews and general discussions. Besides, the $\psi(4230), \psi(4360)$ and $\psi(4415)$ are good candidates of hadronic molecules of 1^{--} $D\bar{D}_1$, $D^*\bar{D}_1$ and $D^*\bar{D}_2^*$, respectively [40, 50–53] (throughout this Letter, the D_1 refers to the narrow $D_1(2420)$ listed in the RPP [37], and $D\bar{D}_1$ means a linear combination of $D\bar{D}_1$ and their antiparticles with certain C-parity and similar for others), especially after noticing the remarkable feature that

$$m_{\psi(4360)} - m_{\psi(4230)} \approx m_{D^*} - m_D,$$
 (1)

$$m_{\psi(4415)} - m_{\psi(4360)} \approx m_{D_2^*} - m_{D_1},$$
 (2)

TABLE I. Hadronic molecules considered in this work and their possible experimental candidates. The binding energies of the 1⁻⁻ states are obtained by the experimental masses of their candidates and that of the ψ_0 is the prediction in this Letter. The values of the thresholds and E_B are in units of MeV.

Molecule	Components	J^{PC}	Threshold	E_B
$\psi(4230)$	$\frac{1}{\sqrt{2}}(D\bar{D}_1 - \bar{D}D_1)$	1	4287	-67 ± 15
$\psi(4360)$	$\frac{1}{\sqrt{2}}(D^*\bar{D}_1-\bar{D}^*D_1)$	1	4429	-62 ± 14
$\psi(4415)$	$\frac{1}{\sqrt{2}}(D^*\bar{D}_2-\bar{D}^*D_2)$	1	4472	-49 ± 4
ψ_0	$\frac{1}{\sqrt{2}}(D^*\bar{D}_1+\bar{D}^*D_1)$	0	4429	-63 ± 18

which is a natural consequence of heavy quark spin symmetry (HQSS) where the low energy interaction between hadrons is independent of the spins of heavy quarks. In fact, the interactions in these channels are the most attractive ones among all the narrow charm-(anti)charm meson pairs from exchanging the light vector mesons [48, 54], and thus these states could be the deepest bound hadronic molecules in the hidden-charm and double-charm meson-meson sectors.

In the heavy quark limit of $m_c \to \infty$, D and D^{*} belong to the same spin multiplet H with the angular momentum of the light degrees of freedom $s_{\ell} = 1/2$, and D_1, D_2^* belong to the multiplet T with $s_\ell = 3/2$. Heavy quark spin partners of the $\psi(4230)$ have been estimated using a constant interaction from the vector-meson dominance model in the exploratory study of Ref. [48], among which there are four isoscalar states with exotic quantum numbers: a $0^{--} D^* \overline{D}_1$ molecule (denoted as ψ_0) around 4.4 GeV, and three $1^{-+} D\bar{D}_1$, $D^*\bar{D}_1$ and $D^*\bar{D}_2^*$ molecules (denoted as η_{c1}) from about 4.3 to 4.5 GeV. They can be searched for in hadron and e^+e^- collisions. In $e^+e^$ collisions below 5 GeV, within the current reach of the BESIII experiment, the 1^{-+} can be produced through $e^+e^- \rightarrow \gamma \eta_{c1}$, while the ψ_0 can be produced in reactions with hadronic final states $e^+e^- \rightarrow \eta \psi_0$. Therefore, it is timely to carefully investigate the ψ_0 , which does not mix with ordinary charmonia and provides a unique portal to understand the vector states in the energy region between 4.2 and 4.5 GeV. In this Letter, we show that the existence of the explicitly exotic ψ_0 is robust in the molecular picture of the vector states $\psi(4230)$, $\psi(4360)$ and $\psi(4415)$, and it can be searched for in electron-positron collisions with an unambiguous signature.

Framework.—The flavor wave functions of the $\psi(4230)$, $\psi(4360)$ and $\psi(4415)$ as 1⁻⁻ molecules, and ψ_0 as a 0⁻⁻ molecule are listed in Table I, where we have adopted the following charge conjugation conventions,

$$C|D\rangle = |\bar{D}\rangle, \quad C|D^*\rangle = -|\bar{D}^*\rangle, C|D_1\rangle = |\bar{D}_1\rangle, \quad C|D_2^*\rangle = -|\bar{D}_2^*\rangle.$$
(3)

In the near-threshold energy region, the interactions between charmed mesons can be described with a nonrelativistic effective field theory, and at leading order there are four independent constant contact terms for the Swave interactions between the H and T multiplets (for each possible isospin) [7]. In the lack of data to fix these contact terms, their values may be estimated with the resonance saturation model by considering the exchange of light mesons [55, 56].

In the following we first focus on the t-channel exchanges and then discuss the u-channel pion-exchange corrections (the contribution of the u-channel exchanges of other mesons are much weaker than the t-channel ones [48, 57]). We consider the exchange of light vector (V) and pseudoscalar (P) mesons by keeping the momentum dependence in the Yukawa potentials, which can be regarded as a way of resumming part of higher order contributions in the momentum expansion. The three meson-meson 1^{--} channels listed in the second column of Table I can couple with one another and the scattering amplitudes by the t-channel V and P exchanges can be expressed as

$$\mathcal{M}_{ij}^{V} = A_{ij}^{V} \frac{1}{q^2 + m_V^2} + B_{ij}^{V}, \tag{4}$$

$$\mathcal{M}_{ij}^{P} = A_{ij}^{P} \frac{1}{q^{2} + m_{P}^{2}} + B_{ij}^{P}, \qquad (5)$$

where q is the transferred 3-momentum and i, j = 1, 2, 3denote channels. They are derived using Lagrangians satisfying HQSS, SU(3) flavor symmetry and chiral symmetry (and hidden local symmetry for light vectors), constructed in Refs. [58–63] and collected in Ref. [48]. The coefficients $A_{ij}^{V,P}, B_{ij}^{V,P}$ can be expressed in terms of coupling constants with phenomenologically known values [64–67], see the Supplemental Material. The amplitude of the 0⁻⁻ $D^* \bar{D}_1$ scattering by the *t*-channel V and P exchanges is similar to the 1⁻⁻ ones but with different coefficients $A_0^{V,P}, B_0^{V,P}$.

The first term of these amplitudes corresponds to Yukawa potentials contributing to the long and midrange interaction, while the second term is a short-range constant contact term. It turns out that the t- and uchannel exchanges of V and P mesons produce four different contact terms; the number matches that of the leading order terms from HQSS analysis [7] mentioned above. Since the contact terms produce ultraviolet (UV) divergence in the Lippmann-Schwinger equation (LSE), they receive scale dependence from renormalization. Therefore, we introduce two scale-dependent factors $c_V(\Lambda)$ and $c_P(\Lambda)$ to the constant terms in Eqs. (4) and (5) serving as counterterms (the constant terms from the u-channel exchanges produce another two). The nonrelativistic potential in momentum space reads

$$V_{ij} = -\frac{1}{\prod_{\alpha=1}^{4} \sqrt{2m_{\alpha}}} \mathcal{M}_{ij},\tag{6}$$

with m_{α} the mass of the initial or final particle and

$$\mathcal{M}_{ij} = \frac{A_{ij}^V}{q^2 + m_V^2} + \frac{A_{ij}^P}{q^2 + m_P^2} + c_V B_{ij}^V + c_P B_{ij}^P.$$
(7)

The potential for the 0^{--} system is similar with the same parameters c_V and c_P due to HQSS.

Bound states are obtained by solving the LSE,

$$T_{ij}(E; \mathbf{k}', \mathbf{k}) = V_{ij}(\mathbf{k}', \mathbf{k}) + \sum_{n} \int \frac{\mathrm{d}^{3} \mathbf{l}}{(2\pi)^{3}} \frac{V_{in}(\mathbf{k}', \mathbf{l}) T_{nj}(E; \mathbf{l}, \mathbf{k})}{E - \mathbf{l}^{2} / (2\mu_{n}) - \Delta_{n1} + i\epsilon}, \quad (8)$$

where \mathbf{k} and $\mathbf{k'}$ are the 3-momenta of the initial and final states in the center-of-mass (c.m.) frame, μ_n is the reduced mass of the *n*-th channel, E is the energy relative to threshold of the first channel and Δ_{n1} is the difference between the *n*-th threshold and the first one. A Gaussian form factor is introduced to regularize the UV divergence,

$$V_{ij}\left(\boldsymbol{k}',\boldsymbol{k}\right) \to V_{ij}\left(\boldsymbol{k}',\boldsymbol{k}\right)e^{-\boldsymbol{q}^{2}/\Lambda^{2}}$$
(9)

with $\boldsymbol{q} = \boldsymbol{k} - \boldsymbol{k}'$ and Λ the cutoff.

The parameters $c_V(\Lambda)$ and $c_P(\Lambda)$ at a given Λ are determined by reproducing the binding energies of the three 1^{--} molecular candidates, as shown in the last column in Table I. We define the χ^2 function,

$$\chi^2 = \sum_{i} \left(\frac{E_{B,ii} - E_{\exp,ii}^{\text{cen}}}{E_{\exp,ii}^{\text{err}}} \right)^2, \qquad (10)$$

where $E_{B,ii}$ is the binding energy of *i*-th channel bound state depending on $c_V(\Lambda)$ and $c_P(\Lambda)$, and $E_{\exp,ii}^{\text{cen, err}}$ is the corresponding experimental center value and error.

t-channel results.—Let us first focus on the singlechannel case by turning off the off-diagonal elements of the potential matrix V_{ij} . By minimizing the χ^2 in Eq. (10) for a given Λ , which is chosen in the phenomenologically reasonable range of $0.8 \sim 1.5$ GeV, we obtain the results shown in the top plot of Fig. 1. It is clear that when $\Lambda \approx 1.2$ GeV, we can find suitable $c_V = 0.50, c_P = 0.18$ reproducing the experimental central values and the corresponding binding energy of ψ_0 is (-72.4 ± 17.4) MeV, where the error is estimated by setting $\chi^2 = 1$ for $\Lambda = 1.2$ GeV.

Coupled-channel effects.—After turning on the offdiagonal V_{ii} , the poles corresponding to the $\psi(4360)$ and $\psi(4415)$ will move to the complex plane on the unphysical Riemann sheets (RSs) due to the opening of the lower $D_1 \overline{D}$ channel. However, it turns out that the coupled-channel effects are negligible and the pole locations are very close to the real axis (the imaginary part at $\Lambda = 1.2$ GeV is less than 1 MeV). We take the real parts of the complex pole locations as the corresponding binding energies and the results are shown in the bottom plot of Fig. 1. The best solution is still located at $\Lambda \approx 1.2$ GeV and the predicted binding energy of $\psi_0(4360)$ is -72.4 MeV. The difference from the singlechannel result, ~ 0.1 MeV, is much less than the estimated uncertainty from the experimental errors, see the Supplemental Material for more comparisons. Therefore,



3

FIG. 1. The minimized χ^2 as a function of Λ and the corresponding $c_{V,P}(\Lambda)$ for the single-channel (top) and coupledchannel (bottom) cases.

we conclude that the coupled-channel effects are negligible.

u-channel pion exchange and 3-body effects.— Although the contribution from the *u*-channel exchange is usually small, the *u*-channel exchanged pion can go on-shell, which means that it contributes to the longestrange interaction and the intermediate 3-body channel will introduce additional cuts to the scattering amplitude and result in nonzero decay widths of the predicted molecules. Thus, we take the $D^*\bar{D}_1$ single channel as an example to carefully investigate such 3-body effects to the pole positions.

It is known that for the $D_1(2420)$, although dominated by $s_{\ell} = 3/2$ state, the S-wave contribution to the decay width of $D_1 \rightarrow D^*\pi$ is sizeable [59, 68]. The $D_1 D^*\pi$ coupling reads

$$\mathcal{L}_{D_1 D^* P} = g_S \sqrt{m_{D_1} m_{D^*}} D_1^{\mu} \partial_{\nu} P D_{\mu}^{*\dagger} v^{\nu} + g_D \sqrt{m_{D_1} m_{D^*}} \left[3 D_1^{\mu} \partial_{\mu} \partial_{\nu} P D^{*\dagger \nu} - D_1^{\alpha} \left(\partial^2 P - \partial_{\mu} \partial_{\nu} P v^{\mu} v^{\nu} \right) D_{\alpha}^{*\dagger} \right] + \text{h.c.}, \quad (11)$$

with $g_S = 2.0 \text{ GeV}^{-1}$ and $g_D = 4.9 \text{ GeV}^{-2}$, where the latter is fixed from D_2^* width and the former from reproducing the D_1 width together with the g_D term. Correspondingly, the S- and D-wave decay widths of $D_1(2420)$ are around 12 and 19 MeV, respectively.

For the S-wave coupling in Eq. (11), the transition

amplitude of $D^*\bar{D}_1 \to D_1\bar{D}^*$ reads

$$\mathcal{M}_u = \frac{g_S^2}{4} (m_{D_1}^2 - m_{D^*}^2)^2 \frac{1}{q^2 - m_{\pi}^2 + i\epsilon}.$$
 (12)

As for the *D*-wave $D_1 D^* \pi$ coupling, its inclusion would lead to new UV divergence and calls for more counterterms. To avoid this issue, we consider only the *S*wave coupling with two different values of $g_{S:} g_{S0} =$ 2.0 GeV^{-1} as given above and $g_{S1} = \sqrt{31/12} g_{S0}$ to mimic the total width of D_1 . As the *D*-wave vertex is of higher order in the momentum expansion than the *S*wave one, the real *u*-channel pion-exchange contribution should live between these two extreme cases.

Note that the exchanged pion can be on-shell, i.e., Eq. (12) can diverge in the physical energy region. Therefore, the $D^*\bar{D}^*\pi$ 3-body channel enters the problem in two aspects [69, 70]. First, the propagator of the exchanged pion in LSE reads

$$\frac{1}{q^2 - m_\pi^2 + i\epsilon} \to \frac{1}{2E(m_\pi, \boldsymbol{q})} \left(\frac{1}{d_1} + \frac{1}{d_2}\right) \qquad (13)$$

where

$$d_i = \sqrt{s} - E(m_\pi, \boldsymbol{q}) - E(m_i, \boldsymbol{k}) - E(m_i, \boldsymbol{k'}) \qquad (14)$$

with $m_1 = m_{D^*}, m_2 = m_{D_1}, \sqrt{s} = E + m_{D^*} + m_{D_1}$ and $E(m, \mathbf{p}) = \sqrt{m^2 + \mathbf{p}^2}$. Second, the $D^*\pi$ loop contributes to the D_1 self-energy, leading to an energy-dependent width of D_1 ,

$$\Gamma_{D_1}(E, \boldsymbol{l}) = \frac{g_S^2}{4} (m_{D_1}^2 - m_{D^*}^2)^2 \frac{p_{\rm cm}}{8\pi m_{D^*\pi}^2}, \qquad (15)$$

where $m_{D^*\pi}$ is the invariant mass of $D^*\pi$ and $p_{\rm cm}$ is the magnitude of the 3-momentum of D^*/π in their c.m. frame, determined by

$$\sqrt{m_{D^*\pi}^2 + l^2} + \sqrt{m_{D^*}^2 + l^2} = E + m_{D^*} + m_{D_1}, \quad (16)$$

$$\sqrt{m_{D^*}^2 + p_{\rm cm}^2} + \sqrt{m_{\pi}^2 + p_{\rm cm}^2} = m_{D^*\pi}.$$
(17)

The $i\epsilon$ term in Eq. (8) should be replaced with $\frac{i}{2}\Gamma_{D_1}(E, \boldsymbol{l})$.

Equations (13) and (15) introduce additional cuts to the amplitude [71], which should be properly treated when searching for poles on the unphysical RS. The details can be found in the Supplemental Material and here we only show the final results, as listed in Table II. We find that the D_1 self-energy gives the molecule a width smaller than that of D_1 and has little influence on the binding energy, and the *u*-channel pion exchange has influence on both the real and imaginary parts of the pole position: the imaginary parts from the above two contributions interfere constructively for the $\psi(4360)$ and destructively for the ψ_0 ; the binding energies change by $\lesssim 10$ MeV, within the errors of the *t*-channel results. As discussed above, the real 3-body effects should live between those of $g_S = g_{S0}$ and g_{S1} since the *D*-wave coupling in Eq. (11) is of higher order in the derivative expansion than the *S*-wave one. Thus, we conclude that the ψ_0 has a mass of (4366 ± 18) MeV, where the central value is obtained by averaging the results of ψ_0 with $g_S = g_{S0}$ and g_{S1} and the uncertainty sums in quadrature half their difference and the one from the *t*-channel fitting (that in the second row of Table II).

The existence of lower channels which are not considered here can increase the widths, which are twice the absolute values of the imaginary parts of the poles listed in Table II. In particular, the S-wave $J/\psi(\psi')\pi\pi$ and P-wave $D^{(*)}\bar{D}^{(*)}$ should be crucial to bring the width to (96 ± 7) MeV [37] measured by experiments [72–75]. On the contrary, the ψ_0 cannot decay into $J/\psi(\psi')\pi^+\pi^-$ or $D\bar{D}$, and its width should be significantly smaller than that of the $\psi(4360)$. An estimate of the decay width by considering $\psi_0 \to D^*\bar{D}_1 \to D^{(*)}\bar{D}^*$ through the V and P exchanges leads to about $1 \sim 5$ MeV for $\bar{D}^*\bar{D}^*$ and $\lesssim 1$ MeV for $D\bar{D}^*$. Consequently, we expect the total width of the ψ_0 to be $\lesssim 10$ MeV.

Experimental search.—The exotic $0^{--} \psi_0$ can be searched for in electron-positron collisions in final states such as $D^{(*)}\bar{D}^*$ and $J/\psi(\psi')\eta$. At $\sqrt{s} \approx 5$ GeV, reachable by the BESIII experiment, the only production channel is $e^+e^- \rightarrow \eta\psi_0(4360)$ in *P*-wave. Given that the $e^+e^- \rightarrow \pi^+ D^0 D^{*-}$ cross section is as large as about 0.4 nb at 4.6 GeV, and the integrated luminosity of BE-SIII at 4.95 GeV is 0.16 fb⁻¹ [76], there is a high chance for the $\psi_0(4360)$ to be found in the $D^{(*)}\bar{D}^*$ final state of $e^+e^- \rightarrow \eta D^{(*)}\bar{D}^*$.

However, this process may always be accompanied by $e^+e^- \rightarrow \eta \psi(4360)$. The decay channels of ψ_0 , such as $D^*\bar{D}^*$ and $J/\psi\eta$, are also shared by the $\psi(4360)$. Furthermore, the masses of these two resonances are similar. Therefore, we need to identify an observable that is unique in distinguishing the ψ_0 from the $\psi(4360)$, and the distribution of the angle between the outgoing η and the e^+e^- beam in the laboratory frame, denoted as θ , fulfills the requirement.

For $e^+e^- \to \gamma^* \to \eta(p_1)\psi_0(p_2)$ and $\eta(p_1)\psi(p_2)$, the amplitudes \mathcal{M}_0 and \mathcal{M}_1 have the following forms,

$$\mathcal{M}_0 \propto \epsilon(\gamma^*) \cdot q,\tag{18}$$

$$\mathcal{M}_1 \propto \epsilon_{\alpha\beta\gamma\delta} \epsilon^{\alpha}(\gamma^*) \epsilon^{*\beta}(\psi) P^{\gamma} q^{\delta}, \qquad (19)$$

where $P = p_1 + p_2$ and $q = p_1 - p_2$. Because the intermediate virtual photon γ^* from e^+e^- annihilations at high energy is transversely polarized, summing over the initial and final polarizations leads to totally different angular distributions for $e^+e^- \rightarrow \eta\psi$ and $e^+e^- \rightarrow \eta\psi_0$, as shown in Fig. 2, and the ψ_0 signal can be clearly distinguished from that of the $\psi(4360)$.

Conclusion and outlook.—The existence of a 0^{--} $D^*\overline{D}_1$ bound state ψ_0 is a natural consequence in the

TABLE II. Pole positions relative to the $D^* \bar{D}_1$ threshold in units of MeV with $c_V = 0.50, c_P = 0.18$ from the single *t*-channel fitting. The uncertainties of *t*-channel E_B inherit from the experimental errors of E_B . "Only Γ_{D_1} " means the D_1 self-energy considered while the *u*-channel pion exchange not and "Full 3-body" means both contributions included.

System	1		0		
<i>t</i> -channel	-63.5 ± 13.8		-72.4 ± 17.4		
g_S	g_{S0}	g_{S1}	g_{S0}	g_{S1}	
Only Γ_{D_1}	-61.5 - 3.5i	-61.5 - 9.2i	-70.0 - 3.5i	-70.0 - 8.9i	
Full 3-body	-65.8 - 6.6i	-73.1 - 14.2i	-65.8 - 0.30i	-59.4 - 1.1i	



FIG. 2. Angular distributions of $e^+e^- \rightarrow \eta \psi_0$ and $\eta \psi$. θ is the angle between the outgoing η and initial e^+e^- beam. The distributions are in arbitrary units and the maxima are normalized to 1.

molecular scenario of the $\psi(4230), \psi(4360)$ and $\psi(4415)$. Being explicitly exotic, it does not mix with charmonium states. We have shown that the existence of the ψ_0 is robust against coupled-channel and 3-body pion-exchange effects. The mass and width of the ψ_0 are predicted to be (4366 ± 18) MeV and $\lesssim 10$ MeV, respectively. We may denote such a state as $\psi_0(4360)$.

It is promising to search for the $\psi_0(4360)$ in e^+e^- collisions through the process $e^+e^- \rightarrow \eta\psi_0(4360)$. The angular distribution provides an unambiguous signature to distinguish the explicitly exotic $\psi_0(4360)$ from states of other possible quantum numbers, such as a vector state in the same mass range. Moreover, the width of the $\psi_0(4360)$ is expected to be much smaller than that of the $\psi(4360)$.

So far no 0^{--} meson has been observed. Being a robust prediction of the hadronic molecular model, the $\psi_0(4360)$ will provide a unique opportunity to infer the internal structure of the vector mesons in the mass range between 4.2 and 4.5 GeV, which has been a riddle since the discovery of the $\psi(4260)$ [77].

We are grateful to the fruitful discussions with Vadim Baru, Meng-Lin Du, Jia-Jun Wu and Mao-Jun Yan. This project is supported in part by the Chinese Academy of Sciences under Grants No. XDB34030000 and No. XDPB15; by the National Natural Science Foundation of China (NSFC) under Grants No. 12125507, No. 11835015, No. 12047503, and No. 11961141012; and by the NSFC and the Deutsche Forschungsgemeinschaft (DFG) through the funds provided to the Sino-German Collaborative Research Center TRR110 "Symmetries and the Emergence of Structure in QCD" (NSFC Grant No. 12070131001, DFG Project-ID 196253076).

- * jiteng@itp.ac.cn
- [†] dongxiangkun@itp.ac.cn
- [‡] fkguo@itp.ac.cn
- § zoubs@itp.ac.cn
- M. Gell-Mann, A Schematic Model of Baryons and Mesons, Phys. Lett. 8, 214 (1964).
- [2] G. Zweig, An SU(3) model for strong interaction symmetry and its breaking. Version 2, in *Developments in the Quark Theory of Hadrons. VOL. 1. 1964 - 1978*, edited by D. B. Lichtenberg and S. P. Rosen (1964) pp. 22–101.
- [3] H.-X. Chen, W. Chen, X. Liu, and S.-L. Zhu, The hiddencharm pentaquark and tetraquark states, Phys. Rept. 639, 1 (2016), arXiv:1601.02092 [hep-ph].
- [4] A. Hosaka, T. Iijima, K. Miyabayashi, Y. Sakai, and S. Yasui, Exotic hadrons with heavy flavors: X, Y, Z, and related states, PTEP **2016**, 062C01 (2016), arXiv:1603.09229 [hep-ph].
- [5] R. F. Lebed, R. E. Mitchell, and E. S. Swanson, Heavy-Quark QCD Exotica, Prog. Part. Nucl. Phys. 93, 143 (2017), arXiv:1610.04528 [hep-ph].
- [6] A. Esposito, A. Pilloni, and A. D. Polosa, Multiquark Resonances, Phys. Rept. 668, 1 (2017), arXiv:1611.07920 [hep-ph].
- [7] F.-K. Guo, C. Hanhart, U.-G. Meißner, Q. Wang, Q. Zhao, and B.-S. Zou, Hadronic molecules, Rev. Mod. Phys. **90**, 015004 (2018), arXiv:1705.00141 [hep-ph].
- [8] A. Ali, J. S. Lange, and S. Stone, Exotics: Heavy Pentaquarks and Tetraquarks, Prog. Part. Nucl. Phys. 97, 123 (2017), arXiv:1706.00610 [hep-ph].
- [9] S. L. Olsen, T. Skwarnicki, and D. Zieminska, Nonstandard heavy mesons and baryons: Experimental evidence, Rev. Mod. Phys. **90**, 015003 (2018), arXiv:1708.04012 [hep-ph].
- [10] W. Altmannshofer *et al.* (Belle-II), The Belle II Physics Book, PTEP **2019**, 123C01 (2019), [Erratum: PTEP 2020, 029201 (2020)], arXiv:1808.10567 [hep-ex].
- [11] Y.-R. Liu, H.-X. Chen, W. Chen, X. Liu, and S.-L. Zhu, Pentaquark and Tetraquark states, Prog. Part. Nucl. Phys. 107, 237 (2019), arXiv:1903.11976 [hep-ph].
- [12] N. Brambilla, S. Eidelman, C. Hanhart, A. Nefediev, C.-P. Shen, C. E. Thomas, A. Vairo, and C.-Z. Yuan,

The XYZ states: experimental and theoretical status and perspectives, Phys. Rept. 873, 1 (2020), arXiv:1907.07583 [hep-ex].

- [13] F.-K. Guo, X.-H. Liu, and S. Sakai, Threshold cusps and triangle singularities in hadronic reactions, Prog. Part. Nucl. Phys. **112**, 103757 (2020), arXiv:1912.07030 [hepph].
- [14] G. Yang, J. Ping, and J. Segovia, Tetra- and penta-quark structures in the constituent quark model, Symmetry 12, 1869 (2020), arXiv:2009.00238 [hep-ph].
- [15] H.-X. Chen, W. Chen, X. Liu, Y.-R. Liu, and S.-L. Zhu, An updated review of the new hadron states, (2022), arXiv:2204.02649 [hep-ph].
- [16] L. Meng, B. Wang, G.-J. Wang, and S.-L. Zhu, Chiral perturbation theory for heavy hadrons and chiral effective field theory for heavy hadronic molecules, (2022), arXiv:2204.08716 [hep-ph].
- [17] C. A. Meyer and E. S. Swanson, Hybrid Mesons, Prog. Part. Nucl. Phys. 82, 21 (2015), arXiv:1502.07276 [hepph].
- [18] M. Ablikim *et al.* (BESIII), Observation of an isoscalar resonance with exotic $J^{PC} = 1^{-+}$ quantum numbers in $J/\psi \rightarrow \gamma \eta \eta'$, (2022), arXiv:2202.00621 [hep-ex].
- [19] M. Ablikim *et al.*, Partial wave analysis of $J/\psi \to \gamma \eta' \eta'$, Phys. Rev. D **105**, 072002 (2022), arXiv:2201.09710 [hepex].
- [20] S. R. Cotanch, I. J. General, and P. Wang, QCD Coulomb Gauge Approach to Exotic Hadrons, Eur. Phys. J. A 31, 656 (2007), arXiv:hep-ph/0610071.
- [21] I. J. General, P. Wang, S. R. Cotanch, and F. J. Llanes-Estrada, Light 1⁻⁺ exotics: Molecular resonances, Phys. Lett. B 653, 216 (2007), arXiv:0707.1286 [hep-ph].
- [22] H.-J. Lee, Discussion on the $J^{PC} = 0^{--}$ Tetraquark of u and d Quarks within the QCD sum rule, New Phys. Sae Mulli **70**, 836 (2020).
- [23] C.-K. Jiao, W. Chen, H.-X. Chen, and S.-L. Zhu, The Possible $J^{PC} = 0^{--}$ Exotic State, Phys. Rev. D **79**, 114034 (2009), arXiv:0905.0774 [hep-ph].
- [24] S. Ishida, H. Sawazaki, M. Oda, and K. Yamada, Decay properties of hybrid mesons with a massive constituent gluon and search for their candidates, Phys. Rev. D 47, 179 (1993).
- [25] I. J. General, S. R. Cotanch, and F. J. Llanes-Estrada, QCD Coulomb gauge approach to hybrid mesons, Eur. Phys. J. C 51, 347 (2007), arXiv:hep-ph/0609115.
- [26] J. Govaerts, F. de Viron, D. Gusbin, and J. Weyers, QCD Sum Rules and Hybrid Mesons, Nucl. Phys. B 248, 1 (1984).
- [27] K. G. Chetyrkin and S. Narison, Light hybrid mesons in QCD, Phys. Lett. B 485, 145 (2000), arXiv:hepph/0003151.
- [28] Z.-R. Huang, W. Chen, T. G. Steele, Z.-F. Zhang, and H.-Y. Jin, Investigation of the light four-quark states with exotic $J^{PC} = 0^{--}$, Phys. Rev. D **95**, 076017 (2017), arXiv:1610.02081 [hep-ph].
- [29] Y. Liu and X.-Q. Luo, Estimate of the charmed 0⁻⁻ hybrid meson spectrum from quenched lattice QCD, Phys. Rev. D 73, 054510 (2006), arXiv:hep-lat/0511015.
- [30] C.-F. Qiao and L. Tang, Finding the 0⁻⁻ Glueball, Phys. Rev. Lett. **113**, 221601 (2014), arXiv:1408.3995 [hep-ph].
- [31] A. Pimikov, H.-J. Lee, N. Kochelev, and P. Zhang, Is the exotic 0⁻⁻ glueball a pure gluon state ?, Phys. Rev. D 95, 071501 (2017), arXiv:1611.08698 [hep-ph].
- [32] A. Pimikov, H.-J. Lee, N. Kochelev, P. Zhang, and

V. Khandramai, Exotic glueball $0^{\pm -}$ states in QCD sum rules, Phys. Rev. D **96**, 114024 (2017), arXiv:1708.07675 [hep-ph].

- [33] A. Pimikov, H.-J. Lee, and N. Kochelev, Comment on "Finding the 0⁻⁻ Glueball", Phys. Rev. Lett. **119**, 079101 (2017), arXiv:1702.06634 [hep-ph].
- [34] L.-L. Shen, X.-L. Chen, Z.-G. Luo, P.-Z. Huang, S.-L. Zhu, P.-F. Yu, and X. Liu, The Molecular systems composed of the charmed mesons in the H\$\overline{S}\$ + h.c. doublet, Eur. Phys. J. C 70, 183 (2010), arXiv:1005.0994 [hep-ph].
- [35] A. A. Filin, A. Romanov, V. Baru, C. Hanhart, Y. S. Kalashnikova, A. E. Kudryavtsev, U.-G. Meißner, and A. V. Nefediev, Comment on 'Possibility of Deeply Bound Hadronic Molecules from Single Pion Exchange', Phys. Rev. Lett. **105**, 019101 (2010), arXiv:1004.4789 [hep-ph].
- [36] F.-K. Guo and U.-G. Meißner, More kaonic bound states and a comprehensive interpretation of the D_{sJ} states, Phys. Rev. D 84, 014013 (2011), arXiv:1102.3536 [hepph].
- [37] P. A. Zyla *et al.* (Particle Data Group), Review of Particle Physics, PTEP **2020**, 083C01 (2020), and 2021 update.
- [38] M.-L. Du, F.-K. Guo, C. Hanhart, B. Kubis, and U.-G. Meißner, Where is the lightest charmed scalar meson?, Phys. Rev. Lett. **126**, 192001 (2021), arXiv:2012.04599 [hep-ph].
- [39] L. Maiani, F. Piccinini, A. D. Polosa, and V. Riquer, The Z(4430) and a new paradigm for spin interactions in tetraquarks, Phys. Rev. D 89, 114010 (2014), arXiv:1405.1551 [hep-ph].
- [40] M. Cleven, F.-K. Guo, C. Hanhart, Q. Wang, and Q. Zhao, Employing spin symmetry to disentangle different models for the XYZ states, Phys. Rev. D 92, 014005 (2015), arXiv:1505.01771 [hep-ph].
- [41] S. K. Choi *et al.* (Belle), Observation of a narrow charmonium-like state in exclusive $B^{\pm} \rightarrow K^{\pm}\pi^{+}\pi^{-}J/\psi$ decays, Phys. Rev. Lett. **91**, 262001 (2003), arXiv:hep-ex/0309032.
- [42] M. Ablikim *et al.* (BESIII), Observation of a Charged Charmoniumlike Structure in $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ at \sqrt{s} =4.26 GeV, Phys. Rev. Lett. **110**, 252001 (2013), arXiv:1303.5949 [hep-ex].
- [43] Z. Q. Liu *et al.* (Belle), Study of $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ and Observation of a Charged Charmoniumlike State at Belle, Phys. Rev. Lett. **110**, 252002 (2013), [Erratum: Phys.Rev.Lett. 111, 019901 (2013)], arXiv:1304.0121 [hep-ex].
- [44] M. Ablikim *et al.* (BESIII), Observation of a charged $(D\bar{D}^*)^{\pm}$ mass peak in $e^+e^- \rightarrow \pi D\bar{D}^*$ at $\sqrt{s} = 4.26$ GeV, Phys. Rev. Lett. **112**, 022001 (2014), arXiv:1310.1163 [hep-ex].
- [45] R. Aaij *et al.* (LHCb), Observation of a narrow pentaquark state, $P_c(4312)^+$, and of two-peak structure of the $P_c(4450)^+$, Phys. Rev. Lett. **122**, 222001 (2019), arXiv:1904.03947 [hep-ex].
- [46] R. Aaij *et al.* (LHCb), Study of the doubly charmed tetraquark T_{cc}^+ , (2021), arXiv:2109.01056 [hep-ex].
- [47] R. Aaij *et al.* (LHCb), Observation of an exotic narrow doubly charmed tetraquark, (2021), arXiv:2109.01038 [hep-ex].
- [48] X.-K. Dong, F.-K. Guo, and B.-S. Zou, A survey of heavy-antiheavy hadronic molecules, Progr. Phys. 41, 65 (2021), arXiv:2101.01021 [hep-ph].

- [49] X.-K. Dong, F.-K. Guo, and B.-S. Zou, Explaining the Many Threshold Structures in the Heavy-Quark Hadron Spectrum, Phys. Rev. Lett. **126**, 152001 (2021), arXiv:2011.14517 [hep-ph].
- [50] Q. Wang, C. Hanhart, and Q. Zhao, Decoding the riddle of Y(4260) and Z_c(3900), Phys. Rev. Lett. **111**, 132003 (2013), arXiv:1303.6355 [hep-ph].
- [51] Q. Wang, M. Cleven, F.-K. Guo, C. Hanhart, U.-G. Meißner, X.-G. Wu, and Q. Zhao, Y(4260): hadronic molecule versus hadro-charmonium interpretation, Phys. Rev. D 89, 034001 (2014), arXiv:1309.4303 [hep-ph].
- [52] L. Ma, X.-H. Liu, X. Liu, and S.-L. Zhu, Strong decays of the XYZ states, Phys. Rev. D 91, 034032 (2015), arXiv:1406.6879 [hep-ph].
- [53] C. Hanhart and E. Klempt, Are the XYZ states unconventional states or conventional states with unconventional properties?, Int. J. Mod. Phys. A 35, 2050019 (2020), arXiv:1906.11971 [hep-ph].
- [54] X.-K. Dong, F.-K. Guo, and B.-S. Zou, A survey of heavy-heavy hadronic molecules, Commun. Theor. Phys. 73, 125201 (2021), arXiv:2108.02673 [hep-ph].
- [55] G. Ecker, J. Gasser, A. Pich, and E. de Rafael, The Role of Resonances in Chiral Perturbation Theory, Nucl. Phys. B 321, 311 (1989).
- [56] E. Epelbaum, U.-G. Meißner, W. Glöckle, and C. Elster, Resonance saturation for four nucleon operators, Phys. Rev. C 65, 044001 (2002), arXiv:nucl-th/0106007.
- [57] X.-K. Dong, Y.-H. Lin, and B.-S. Zou, Prediction of an exotic state around 4240 MeV with $J^{PC} = 1^{-+}$ as C-parity partner of Y(4260) in molecular picture, Phys. Rev. D **101**, 076003 (2020), arXiv:1910.14455 [hep-ph].
- [58] M. B. Wise, Chiral perturbation theory for hadrons containing a heavy quark, Phys. Rev. D 45, R2188 (1992).
- [59] A. F. Falk and M. E. Luke, Strong decays of excited heavy mesons in chiral perturbation theory, Phys. Lett. B 292, 119 (1992), arXiv:hep-ph/9206241.
- [60] B. Grinstein, E. E. Jenkins, A. V. Manohar, M. J. Savage, and M. B. Wise, Chiral perturbation theory for f_{D_S}/f_D and B_{B_S}/B_B , Nucl. Phys. B **380**, 369 (1992), arXiv:hepph/9204207.
- [61] R. Casalbuoni, A. Deandrea, N. Di Bartolomeo, R. Gatto, F. Feruglio, and G. Nardulli, Light vector resonances in the effective chiral Lagrangian for heavy mesons, Phys. Lett. B 292, 371 (1992), arXiv:hepph/9209248.
- [62] R. Casalbuoni, A. Deandrea, N. Di Bartolomeo, R. Gatto, F. Feruglio, and G. Nardulli, Effective Lagrangian for heavy and light mesons: Semileptonic decays, Phys. Lett. B 299, 139 (1993), arXiv:hepph/9211248.
- [63] R. Casalbuoni, A. Deandrea, N. Di Bartolomeo, R. Gatto, F. Feruglio, and G. Nardulli, Phenomenology of heavy meson chiral Lagrangians, Phys. Rept. 281, 145 (1997), arXiv:hep-ph/9605342.
- [64] M. Bando, T. Kugo, and K. Yamawaki, Nonlinear Realization and Hidden Local Symmetries, Phys. Rept. 164, 217 (1988).
- [65] C. Isola, M. Ladisa, G. Nardulli, and P. Santorelli, Charming penguins in $B \to K^*\pi, K(\rho, \omega, \phi)$ decays, Phys. Rev. D 68, 114001 (2003), arXiv:hep-ph/0307367.
- [66] R. Chen, Z.-F. Sun, X. Liu, and S.-L. Zhu, Strong LHCb evidence supporting the existence of the hiddencharm molecular pentaquarks, Phys. Rev. D 100, 011502 (2019), arXiv:1903.11013 [hep-ph].

- [67] F.-L. Wang and X. Liu, Investigating new type of doubly charmed molecular tetraquarks composed of charmed mesons in the H and T doublets, Phys. Rev. D 104, 094030 (2021), arXiv:2108.09925 [hep-ph].
- [68] F.-K. Guo, Triangle Singularities and Charmoniumlike XYZ States, Nucl. Phys. Rev. 37, 406 (2020), arXiv:2001.05884 [hep-ph].
- [69] V. Baru, A. A. Filin, C. Hanhart, Y. S. Kalashnikova, A. E. Kudryavtsev, and A. V. Nefediev, Three-body $D\bar{D}\pi$ dynamics for the X(3872), Phys. Rev. D 84, 074029 (2011), arXiv:1108.5644 [hep-ph].
- [70] M.-L. Du, V. Baru, X.-K. Dong, A. Filin, F.-K. Guo, C. Hanhart, A. Nefediev, J. Nieves, and Q. Wang, Coupled-channel approach to T⁺_{cc} including three-body effects, Phys. Rev. D 105, 014024 (2022), arXiv:2110.13765 [hep-ph].
- [71] M. Döring, C. Hanhart, F. Huang, S. Krewald, and U.-G. Meißner, Analytic properties of the scattering amplitude and resonances parameters in a meson exchange model, Nucl. Phys. A 829, 170 (2009), arXiv:0903.4337 [nucl-th].
- [72] J. P. Lees *et al.* (BaBar), Study of the reaction $e^+e^- \rightarrow \psi(2S)\pi^-\pi^-$ via initial-state radiation at BaBar, Phys. Rev. D **89**, 111103 (2014), arXiv:1211.6271 [hep-ex].
- [73] X. L. Wang *et al.* (Belle), Measurement of $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$ via Initial State Radiation at Belle, Phys. Rev. D **91**, 112007 (2015), arXiv:1410.7641 [hep-ex].
- [74] M. Ablikim *et al.* (BESIII), Precise measurement of the $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ cross section at center-of-mass energies from 3.77 to 4.60 GeV, Phys. Rev. Lett. **118**, 092001 (2017), arXiv:1611.01317 [hep-ex].
- [75] M. Ablikim *et al.* (BESIII), Measurement of $e^+e^- \rightarrow \pi^+\pi^-\psi(3686)$ from 4.008 to 4.600[°]GeV and observation of a charged structure in the $\pi^\pm\psi(3686)$ mass spectrum, Phys. Rev. D **96**, 032004 (2017), [Erratum: Phys.Rev.D 99, 019903 (2019)], arXiv:1703.08787 [hep-ex].
- [76] M. Ablikim *et al.* (BESIII), Luminosities and energies of e^+e^- collision data taken between \sqrt{s} =4.612 GeV and 4.946 GeV at BESIII, (2022), arXiv:2205.04809 [hep-ex].
- [77] B. Aubert *et al.* (BaBar), Observation of a broad structure in the $\pi^+\pi^- J/\psi$ mass spectrum around 4.26-GeV/c², Phys. Rev. Lett. **95**, 142001 (2005), arXiv:hep-ex/0506081.

Supplemental Material

t-channel potential

The amplitudes for $D^{(*)}\overline{D}_{1,2}^{(*)}$ scattering via the V and P exchanges can be expressed in Eqs. (4) and (5) and the coefficients take the following expressions,

$$A_{11}^V = -4\beta\beta_2 g_V^2 m_D m_{D_1} - \frac{20}{9}\beta\lambda_2 g_V^2 m_D m_\rho^2, \qquad (20)$$

$$B_{11}^V = \frac{20}{9} \beta \lambda_2 g_V^2 m_D, \tag{21}$$

$$A_{11}^P = B_{11}^P = 0, (22)$$

Æ

$$B_{22}^{V} = \frac{4}{9}g_{V}^{2}(6\beta_{2}\lambda m_{D_{1}} + 5\beta\lambda_{2}m_{D^{*}} + 20\lambda\lambda_{2}m_{D_{1}}m_{D^{*}}),$$
(24)

$$A_{22}^{P} = -\frac{3}{2} \frac{10gk}{9f_{\pi}^{2}} m_{D_{1}} m_{D^{*}} m_{\pi}^{2} = -\frac{3}{4} m_{\pi}^{2} B_{22}^{P}, \qquad (25)$$

$$A_{33}^{V} = -4\beta\beta_2 g_V^2 m_{D^*} m_{D_2} - \frac{8}{3} g_V^2 m_\rho^2 (\beta \lambda_2 m_{D^*} + \beta_2 \lambda m_{D_2} - 6\lambda \lambda_2 m_{D^*} m_{D_2}), \qquad (26)$$

$$B_{33}^{V} = \frac{8}{3}g_{V}^{2}(\beta\lambda_{2}m_{D^{*}} + \beta_{2}\lambda m_{D_{2}} - 6\lambda\lambda_{2}m_{D^{*}}m_{D_{2}}), \quad (27)$$

$$A_{33}^{P} = -\frac{3}{2} \frac{2gk}{f_{\pi}^{2}} m_{D_{2}} m_{D^{*}} m_{\pi}^{2} = -\frac{3}{4} m_{\pi}^{2} B_{33}^{P}, \qquad (28)$$

$$A_{12}^{V} = \frac{20\sqrt{2}}{9}g_{V}^{2}\lambda\lambda_{2}m_{D_{1}}(m_{D^{*}} + m_{D})m_{\rho}^{2}$$

= $-m_{\rho}^{2}B_{12}^{V},$ (29)

$$A_{12}^{P} = -\frac{3}{2} \frac{10\sqrt{2}gk}{9f_{\pi}^{2}} m_{D_{1}}\sqrt{m_{D^{*}}m_{D}}m_{\pi}^{2}$$
$$= -\frac{3}{4}m_{\pi}^{2}B_{12}^{P}, \qquad (30)$$

$$A_{13}^{V} = \frac{8\sqrt{10}}{3} g_{V}^{2} \lambda \lambda_{2} m_{D_{1}} m_{D} m_{\rho}^{2} = -m_{\rho}^{2} B_{13}^{V}, \qquad (31)$$

$$A_{13}^{P} = -\frac{3}{2} \frac{2\sqrt{10}gk}{9f_{\pi}^{2}} \sqrt{m_{D}m_{D^{*}}m_{D_{1}}m_{D_{2}}} m_{\pi}^{2}$$
$$= -\frac{3}{4}m_{\pi}^{2}B_{32}^{P}, \qquad (32)$$

$$A_{23}^{V} = -\frac{16\sqrt{5}}{9}g_{V}^{2}\lambda\lambda_{2}\sqrt{m_{D_{1}}m_{D_{2}}}m_{D^{*}}m_{\rho}^{2} = -m_{\rho}^{2}B_{23}^{V},$$
(33)

$$A_{23}^{P} = \frac{3}{2} \frac{2\sqrt{5}gk}{9f_{\pi}^{2}} \sqrt{m_{D_{1}}m_{D_{2}}} m_{D^{*}}m_{\pi}^{2} = -\frac{3}{4}m_{\pi}^{2}B_{32}^{P}, \quad (34)$$

$$A_{0}^{V} = -4\beta\beta_{2}g_{V}^{2}m_{D}m_{D^{*}} - \frac{4}{9}g_{V}^{2}m_{\rho}^{2}(6\beta_{2}\lambda m_{D_{1}} + 5\beta\lambda_{2}m_{D^{*}} + 40\lambda\lambda_{2}m_{D_{1}}m_{D^{*}}), \qquad (35)$$

$$B_0^V = \frac{4}{9}g_V^2(6\beta_2\lambda m_{D_1} + 5\beta\lambda_2 m_{D^*} + 40\lambda\lambda_2 m_{D_1} m_{D^*}),$$
(36)

$$A_0^P = A_{22}^P, \quad B_0^P = B_{22}^P. \tag{37}$$

Note that the isospin factors for isospin-0 have been taken into account. For the P exchange, the Yukawa term contains only the pion-exchange contributions while the corresponding counterterm contains contributions from both the pion and the $\eta^{(\prime)}$ mesons. The numerical values of coupling constants are g = -k = 0.59 [59, 65], $\beta = -\beta_2 = 0.9$ [65], $g_V = 5.8$ [64], $\lambda = -\lambda_2 =$ 0.56 [66, 67] and $f_{\pi} = 132$ MeV is the pion decay constant. The masses of the involved particles are taken from the RPP [37].

Analytical properties of amplitudes

The on-shellness of the propagating intermediate particles will introduce cuts to the scattering amplitudes, which are analytical functions of the complex energy except for these cuts and possible poles. Usually, the cut introduced by the intermediate component particles, say $D^*\bar{D}_1$ in the $\psi(4360)$ case, is chosen from the threshold to positive infinity, called the right-hand cut (RHC) while those from the exchanged particles are called the left-hand cuts (LHCs). Here we give detailed discussions on the cuts in the $D^*\bar{D}_1$ scattering amplitude including the $D^*\bar{D}^*\pi$ 3-body channel.

The RHC in the $D^*D^*\pi$ (with $\bar{D}^*\pi$ from \bar{D}_1) channel originates from the $p_{\rm cm}$ in $\Gamma_{D_1}(E, \mathbf{l})$ (see Eq.(15)) and starts from $s = s_0(|\mathbf{l}|)$ to positive infinity by choosing a cut of the square root function in $p_{\rm cm}$ so that

$$\operatorname{Im}[p_{\mathrm{cm}}] \ge 0 \quad \text{on 1st RS,} \\
\operatorname{Im}[p_{\mathrm{cm}}] < 0 \quad \text{on 2nd RS.}$$
(38)

Note that the branch point $s_0(|\boldsymbol{l}|)$ is a moving point from $s_0(0) = (2m_{D^*} + m_{\pi})^2$ to the right as $|\boldsymbol{l}|$ increases. See the red-dashed line in Fig. 3 for the cut with $|\boldsymbol{l}| = 0$. Due to the finite decay width of D_1 , one of the branch points of the $D^*\bar{D}_1$ RHC is located at $\sqrt{s} = m_{D^*} + m_{D_1} - \frac{i}{2}\Gamma_{D_1}$ on the 2nd RS, instead of at the nominal threshold on the real axis. See the green line in Fig. 3.

To see the LHCs, we need to analyze the poles of the meson-exchange potentials (both on-shell and off-shell for the initial and final scattered particles) before the partial wave projection. For the on-shell *t*-channel one, the cut lies at $s \in (-\infty, s_1]$ with $s_1 < (m_{D^*} + m_{D_1})^2$. While for the off-shell one with $|\mathbf{k}|, |\mathbf{k'}| \ge 0$, there is no cut. The branch points for the *u*-channel pion exchange are determined by $d_1 = 0$ at $z \equiv \cos \mathbf{\hat{k} \cdot k'} = \pm 1$ with d_1 given in Eq. (14), which translates to

$$\sqrt{m_{D_1}^2 + \mathbf{k}^2} = \sqrt{m_{D^*}^2 + \mathbf{k'}^2} + \sqrt{m_{\pi}^2 + \mathbf{k'}^2 \pm 2|\mathbf{k}||\mathbf{k'}|}.$$
 (39)

For the on-shell case where $|\mathbf{k}| = |\mathbf{k}'|$, it is actually a RHC (still called "L"HC in the following) from a point s_2 above threshold to positive infinity. For the off-shell case with $|\mathbf{k}|, |\mathbf{k}'| \ge 0$, one of the branch points, $s_3(|\mathbf{k}|, |\mathbf{k}'|)$, is moving from $s_3(0,0) = (2m_{D^*} + m_{\pi})^2$ to the right as $|\mathbf{k}|$ and $|\mathbf{k}'|$ increase and the other is positive infinity, again a RHC ("L"HC). See the blue line in Fig. 3 for the cut with $|\mathbf{k}|, |\mathbf{k}'| = 0$.

To search for poles corresponding to the $\psi_0(4360)$ and $\psi(4360)$, which are now located on the unphysical RS marked by the red \otimes in Fig. 3 and close to the physical axis on the first RS, one has to perform the analytical continuation properly.

• *t*- and *u*-channel cuts when the scattered particles are on-shell: They will not contribute by construc-



FIG. 3. The cuts encountered in the $D^*\bar{D}_1$ system. The off-shell *u*-channel "L"HC is located on the real axis and moved away slightly for better illustration. For $D^*\bar{D}^*\pi$ RHCs and off-shell *u*-channel "L"HCs, only that with $|\boldsymbol{l}| = 0$ or $|\boldsymbol{k}|, |\boldsymbol{k'}| \geq 0$ is shown.

tion when solving LSE and therefore, they are not shown in Fig. 3.

- The off-shell *u*-channel "L"HCs with $|\mathbf{k}|, |\mathbf{k}'| \ge 0$: When performing the *S*-wave projection, i.e. the integral over *z* from -1 to 1 along the real axis, the pole of the integrand, $z_0(E, |\mathbf{k}|, |\mathbf{k}'|)$, will move from the lower half *z*-plane to the upper one when *E* moves from physical axis to the unphysical region where the pole of *T* matrix is located. If $z_0(E, |\mathbf{k}|, |\mathbf{k}'|)$ cross over the integral path, the result will have discontinuity. To avoid crossing over these cuts, we deform the integral path of *z* from a straight line $-1 \rightarrow 1$ to a polyline $-1 \rightarrow -1 + ia \rightarrow$ $1 + ia \rightarrow 1$ with *a* a sufficiently large positive number.
- The RHC of $D^*\bar{D}_1$: This one is on the unphysical sheet, beyond the region of the possible pole and hence of no interest here.
- The $D^* \bar{D}^* \pi$ RHCs: If we use the cuts defined in Eq. (38), it is found that the values of |1 - VG| (an intermediate function in solving LSE, whole roots are the pole positions of the T matrix) in the 1stupper and 2nd-lower RSs do not match. The reason is that when integrating |l| from 0 to $+\infty$ along the real axis, the argument of the square root function in $p_{\rm cm}$ would go across its branch point from right to left, see the Fig. 7 in Ref [71]. Therefore we have to change the integral path to avoid this discontinuation. In our calculation, we have chosen a proper cut of the square root which lies parallel to the negative imaginary axis to realize a smooth continuation [70, 71]. See the red-solid line in Fig. 3 for the modified cut with |l| = 0.

After the above treatments, the values of |1 - VG| are continued smoothly from the physical axis to the 2nd RS, see Fig. 4 for illustration.



FIG. 4. An illustration of the connection between the 1stupper $(E + i10^{-7} \text{ GeV})$ and 2nd-lower $(E - i10^{-7} \text{ GeV})$ RSs where the solid and dashed (dotted and dotdashed) lines are the real (imaginary) parts. "No *u*" means only *t*-channel interaction considered and in turn no cuts while "Full *u*" means that the *u*-channel interaction and D_1 self-energy are both included. The blue curves are covered by the red ones and hence they are connected correctly.



FIG. 5. Comparison between $c_{P,V}$ of single-channel (solid) and coupled-channel (dashed) results with only *t*-channel meson-exchange interaction.

Comparisons between the single- and coupled-channel results

Here we compare the values of $c_{V,P}$ explicitly in Fig. 5. We can see that for $\Lambda = 1.0 \sim 1.3$ GeV, there are almost no differences between single-channel and coupledchannel results. The binding energies corresponding to these parameters are shown in Fig. 6 and the predicted binding energy of $\psi_0(4360)$ are robust against Λ .



FIG. 6. Comparison of the single-channel (s.c.) and coupled-channel (c.c.) results for binding energies of the $\psi(4230)$, $\psi(4360)$, $\psi(4415)$ and $\psi_0(4360)$ with only the *t*-channel meson-exchange interaction. The dashed lines and shadows represent the experimentally measured central values and errors.