arXiv:2205.11190v1 [hep-th] 23 May 2022

Boundary conditions and anomalies of conformal field theories in 1+1 dimensions

Linhao Li,¹ Chang-Tse Hsieh,^{2,3,*} Yuan Yao,^{4,†} and Masaki Oshikawa^{1,5,6}

¹Institute for Solid State Physics, The University of Tokyo. Kashiwa, Chiba 277-8581, Japan

²Department of Physics and Center for Theoretical Physics,

National Taiwan University, Taipei 10607, Taiwan

³Physics Division, National Center for Theoretical Science,

⁴Condensed Matter Theory Laboratory, RIKEN CPR, Wako, Saitama 351-0198, Japan

⁵Kavli Institute for the Physics and Mathematics of the Universe (WPI),

The University of Tokyo, Kashiwa, Chiba 277-8583, Japan

⁶ Trans-scale Quantum Science Institute, University of Tokyo, Bunkyo-ku, Tokyo 113-0033, Japan

(Dated: May 24, 2022)

We study a relationship between conformally invariant boundary conditions and anomalies of conformal field theories (CFTs) in 1+1 dimensions. For a given CFT with a global symmetry, we consider symmetric "gapping potentials" which are relevant perturbations to the CFT. If a gapping potential is introduced only in a subregion of the system, it provides a certain boundary condition to the CFT. From this equivalence, if there exists a Cardy boundary state which is invariant under a symmetry, then the CFT can be gapped with a unique ground state by adding the corresponding gapping potential. This means that the symmetry of the CFT is anomaly free. Using this approach, we systematically deduce the anomaly-free conditions for various types of CFTs with several different symmetries. They include the free compact boson theory, Wess-Zumino-Witten models, and unitary minimal models. When the symmetry of the CFT is anomalous, it implies a Lieb-Schultz-Mattis type ingappability of the system. Our results are consistent with, where available, known results in the literature. Moreover, we extend the discussion to other symmetries including spin groups and generalized time-reversal symmetries.

CONTENTS

I.	Introduction	2
II.	A review on Boundary Conformal Field Theory (BCFT)A. BCFT on the Upper Half PlaneB. BCFT on the annulus	333
III.	Boundary Conformal Field Theories (BCFTs) and quantum anomalies	5
IV.	 Multicomponent U(1) compactified boson theory A. Canonical quantization B. Anomaly free condition and Haldane gapping potential C. Anomaly of U(1) × Z₂ symmetry D. Anomaly of SU(2) × Z₂ symmetry E. Anomaly of SU(N) × Z_N symmetry 	6 6 7 8 9
V.	 Anomaly of WZW model A. Introduction to WZW model and global symmetry B. Action of global symmetry on Cardy state C. Mixed anomaly of WZW model 1. Invarinat boundary states of center symmetry and vector rotation symmetry 2. Invariant boundary states of time 	10 10 10 10
	reversal and vector rotation symmetry	11

	D. Application to a spin chain with the triple product interactions	12
VI.	Discrete symmetries of minimal models A. Invariant boundary states of \mathbb{Z}_2 symmetry B. Invariant boundary states of S_3 symmetry	$13 \\ 13 \\ 13 \\ 13$
VII.	T-duality symmetry of $SU(2)_1$ WZW model A. T-duality symmetry of $SU(2)_1$ WZW	14
	model	14
	B. Anomaly of T-duality symmetry and center	14
	C Pure anomaly of T duality symmetry	14
	C. I use anomary of 1 duanty symmetry	10
VIII.	Conclusion and discussion	16
	Acknowledgements	16
	References	17
А.	$\mathrm{SU}(N)$ symmetry invariant boundary states of free boson model	18
В.	Invariant boundary state of vector rotation	
	symmetry and center symmetry	19
	1. B_N affine Lie algebra	19
	2. D_{2l} affine Lie algebra	20
	3. C_N affine Lie algebra	20
	4. D_{2l+1} affine Lie algebra	20
	5. E_6 affine Lie algebra	21
	6. E_7 affine Lie algebra	21

National Taiwan University, Taipei 10607, Taiwan

C. Invariant boundary state of vector rotation symmetry and CP_2 symmetry

21

- D. Symmetric interaction of $SU(2)_{2k}$ WZW model 22
- E. The LSM ingappability of the spin chain with triple-product interactions 23
- F. Vector rotation on the *y*-direction 24

I. INTRODUCTION

Symmetry often plays an essential role in study of quantum many-body physics. In classical physics, it implies various conservation laws and corresponding conserved current. However, on the quantum level, there is a potential obstruction in promoting a global symmetry to a gauge symmetry which is called 't Hooft anomaly¹. Recent works on topological phases of matter have revealed that a *d*-dimensional quantum field theory with a global symmetry having an 't Hooft anomaly can be regarded as the boundary theory of a nontrivial symmetry protected topological (SPT) phases^{2–7} in one higher dimension and the anomaly of the boundary theory corresponds to certain topological property of the bulk theory^{8–12}.

For a (1+1)d conformal field theory (CFT) with an anomalous global symmetry G, the anomaly can be detected by examining modular invariance of the partition function(s) of the CFT orbifolded by $G^{13,14}$. (In a Gorbifold CFT, states except for G singlets are projected out and G-twisted sectors are included when evaluating the partition function.) If G has an 't Hooft anomaly, it is impossible to construct a modular invariant G-orbifold partition function from the the original CFT.

Another probe of anomaly is edgeability^{15,16}. A CFT with a global symmetry G is edgeable means it can be cut open into a boundary conformal field theory (BCFT) where G is preserved both in the bulk and on the boundary. The connection between edgeability and anomaly relies on the observation that a (1+1)d CFT with an anomalous symmetry lives on the boundary of a (2+1)d nontrivial SPT phase and thus can not be formulated consistently on a one-dimensional space with a boundary (as the boundary of a boundary vanishes)¹⁷. That is, a (1+1)d CFT with an anomalous symmetry-preserving manner to a BCFT.

In this work, we further investigate the correspondence between boundary conditions and anomalies of CFTs from a different aspect. We consider adding spatially dependent relevant perturbations to a CFT such that the perturbations are present only outside a finite length segment of the one-dimensional space \mathbb{R}^1 . The boundary conditions at the interfaces between the regions with and without perturbations are determined by the forms of the added relevant operators and undergo renormalization group (RG) flow with fixed points corresponding $\mathbf{2}$

to conformal boundary conditions. It turns out that the theory on the finite segment will become a BCFT with these conformal boundary conditions¹⁸.

We will focus on the uniqueness of ground state after adding particular space-dependent perturbation. More precisely, in the limit length $L \rightarrow 0$, the multiplicity of the identity operator contained in partition function equals ground-state degeneracy. In order to have a unique ground state, the identity operator should appear in the partition function just once which means the partition function goes to 1 in this limit. We will show this requirement is equivalent to finding a symmetric boundary state or boundary condition. On the contrary, if all the possible boundary amplitudes (partition function) with an imposed symmetry exhibit a degeneracy, it suggests that the CFT cannot be gapped with a unique ground state, i.e., ingappability. The latter property implies an anomaly of the CFT.

If we use the gapping-potential argument directly to deduce whether a symmetry is anomaly-free, we need to first construct a symmetric gapping potential and then check whether the gapped ground state(s) does not break the symmetry spontaneously. This procedure can be systematically implemented on free compact boson theories but is usually not easy for most other CFTs, such as WZW models and minimal models. On the other hand, the BCFT formulation and the construction of certain boundary states of several CFTs, including the two mentioned above, have been known to us, so we can check the existence of symmetric boundary states systematically and quickly using the BCFT appraoch. For example, for the $SU(2)_1$ WZW model with the T-duality symmetry, we will show the boundary state approach can give the symmetric gapping potentials which are beyond the usual Haldane null gapping potentials¹⁹. Moreover, the boundary states approach can also imply the anomaly with respect to time reversal symmetry of WZW models, which are usually more difficult to be proved by other approaches.

An analogous application of BCFT was discussed in Ref. 18. There, the SPT phases in 1+1 dimensions were related to the spectrum of the "mother CFT" with distinct boundary conditions corresponding to the trivial and SPT phases imposed at the two ends. In the present paper, we rather discuss the anomaly of the CFT, which is related to the Lieb-Schultz-Mattis (LSM) type "ingappability"^{20–23}, by imposing the "same" boundary conditions at the two ends. Thus our discussion is complementary to Ref. 18. (For previous works on anomaly-based arguments for the LSM type ingappability, see Ref. 24–29)

Furthermore, a similar idea was used to discuss the anomaly of WZW models between the center symmetry and the spacetime (large) diffeomorphism³⁰. While this was useful in revealing the boundary ingappability of SPT phases, its application on LSM-type ingappability — ingappability induced from some internal symmetry and lattice symmetry — was still unclear. In contrast, in our work, we provide a classification of mixed anomalies of e.g., a large class of Lie groups and their centers, where the Lie groups can be the internal symmetry and the center symmetry is expected to realize the lattice symmetry. Thus, our results are more directly relevant to LSM-type ingappability of lattice models. Indeed, our results are consistent with known ingappabilities found on lattices. Moreover, our argument covers a broad class of 1d spin systems. One class of such systems that has been rarely studied is the SO(n) spin chains, whose associated LSM theorem was discussed in Ref. 31 and 32 and is consistent with the results presented here. For more general cases that have not been explored, our work could lead to novel predictions.

The organization of the rest of the paper is as follows. In section II, a brief introduction to BCFT is provided. In section III, we will discuss the correspondence between boundary conditions and quantum anomaly of CFTs in 1+1 dimensions. As an application and illustration of our framework, we study some concrete examples. In section IV, the free boson theory in 1+1 dimensions with PSU(N) and \mathbb{Z}_N symmetry is considered. In section V, we will apply our framework to anomalies of WZW models with center symmetry, vector rotation symmetry and time reversal symmetries of minimal models are all anomaly-free. In section VII, we will discuss the anomaly of free boson theory with respect to the T-duality symmetry. Our main results are summarized in TABLE I.

II. A REVIEW ON BOUNDARY CONFOMRAL FIELD THEORY (BCFT)

In this section, we will give a basic review of the Boundary Conformal Field Theory (BCFT) on the upper half plane and annulus, which is an important basis for our formulation.

A. BCFT on the Upper Half Plane

On the complex plane, the analytic coordinate transformations form the conformal group. If we put the CFT on the Upper Half Plane (UHP), the conformal transformation should also map UHP to itself^{33–35}:

$$(x,y) \to (x,y) + (\epsilon_x, \epsilon_y)$$
 and $\epsilon_y(x,0) = 0.$ (1)

To see the effect of boundary condition, we can calculate the expectation value of changes of primary fields X transformed by the above conformal transformation:

$$\int d\Phi \delta X e^{-S[\Phi]} = \int d\Phi X \delta S e^{-S[\Phi]}$$
$$= -\int dx \epsilon_x(x,0) \langle T^{xy}(x,0)X \rangle$$
$$+ \int dx dy \epsilon_\mu \partial_\nu \langle T^{\mu\nu}(x,y)X \rangle.$$
(2)

The second term generates the conformal transformation on X producing $\langle \delta X \rangle$, which demands the first term to vanish, i.e., $T^{xy}(x,0) = 0$. This condition can be interpreted as the absence of energy flow across the boundary.

In the complex coordinate, this boundary condition can be written as:

$$T(z) = \bar{T}(\bar{z}), \quad z \in \mathbb{R}.$$
(3)

Thus from the definition of the generators of the conformal group expanding the Laurent series of energy momentum tensors, we obtain

$$T(x) = L_n x^{-n-2} = \bar{T}(x) = \bar{L}_n x^{-n-2} \Rightarrow L_n = \bar{L}_n.$$
 (4)

Therefore, in contrast to the case of CFTs on full complex plane, only one copy of Virasoro algebra acts on the Hilbert space $\oplus n_h V_h$. If we see the *y*-axis as the time direction, then the Hilbert space is defined on the *x*-axis.

B. BCFT on the annulus

We can also define CFT on an annulus. The boundary can be put perpendicular to the spatial direction (open string) or to the time direction (closed string but with time open).

For the closed string, such a boundary state is represented by a state in the Hilbert space of a CFT defined on a circle and boundary A is on the time t = 0 and boundary B is on t = L. For spatial dimension, we identify $x + it \sim x + \beta + it$. Then we let $\omega = x + it$ and map the cylinder into the plane: $z = \exp(-i2\pi\omega/\beta)$. The boundary is mapped to the circle |z| = 1 and $|z| = \exp(2\pi L/\beta)$. From the calculation in the above section, we can see the conformal boundary conditions should be: $(c = \bar{c})$

$$z^{2}T(z) - c/24 = \bar{z}^{2}\bar{T}(\bar{z}) - \bar{c}/24 , \quad |z| = 1, \exp(2\pi L/\beta).$$
(5)

More precisely, after quantization, the above equation satisfies when it acts on the boundary state in the Heisenberg picture. Using the mode expansion of the stressenergy tensor, we can obtain

$$(L_n - \bar{L}_{-n})|a\rangle\rangle = 0.$$
(6)

Thus the physical boundary condition that we put on the boundary should satisfy the conformal boundary conditions. The physical boundary condition describes an automorphism from the holomorphic sector to antiholomorphic sector

$$[S(z) - \rho_{\rm A/B}(\bar{S}(\bar{z}))]|a\rangle\rangle = 0.$$
(7)

where S belongs to some symmetry algebra and $\rho_{A/B}$ denotes an automorphism of the algebra of fields. For example, the U(1) symmetry in free boson theory has the current $i\partial\varphi$ and the boundary condition can be: $\partial_z\varphi_z =$ $\pm \partial_{\bar{z}}\varphi_{\bar{z}}|_{|z|=1,\exp(2\pi L/\beta)}$.

Model	Symmetry	Invariant boundary states
$\mathrm{SU}(N)_k$ WZW model		$k \in \! N\mathbb{N}$
$\operatorname{Spin}(2N+1)_k$ WZW model		$k\in\mathbb{N}$
$\mathrm{Usp}(N)_k$ WZW model	diagonal rotation symmetry	$k\in\!\!2\mathbb{N}$ or $N\in\!\!2\mathbb{N}$
$\operatorname{Spin}(4N+2)_k$ WZW model	&	$k\in\!\!4\mathbb{N}$
$\operatorname{Spin}(4N)_k$ WZW model	center symmetry	$k\in\!\!2\mathbb{N}$
E_6 WZW model		$k \in 3\mathbb{N}$
E_7 WZW model		$k\in\!\!2\mathbb{N}$
$\mathrm{SU}(N)_k$ WZW model		$k\in\!\!2\mathbb{N}$ or $N\in\!\!4\mathbb{N}$
$\operatorname{Spin}(2N+1)_k$ WZW model		$k\in\mathbb{N}$
$\mathrm{Usp}(N)_k$ WZW model	diagonal rotation symmetry	$k\in\!\!2\mathbb{N}$ or $N\in\!\!2\mathbb{N}$
$\operatorname{Spin}(4N+2)_k$ WZW model	&	$k\in\mathbb{N}$
$\operatorname{Spin}(4N)_k$ WZW model	time-reversal symmetry \mathcal{T}_2	$k\in\!\!2\mathbb{N}$
E_7 WZW model		$k\in\!\!2\mathbb{N}$
$SU(2)_1$ WZW model	T-duality	yes
(self-dual compact boson)	T-duality extended by center symmetry	no
unitary (Virasoro) minimal models	any finite symmetry i.e. \mathbb{Z}_2 or S_3	yes

TABLE I. Summary of the conditions for the existence of symmetry invariant boundary states in various CFTs. Here we denote the G_k -WZW models at the level k by the associated Lie groups G.

A kind of quantum states $|a\rangle\rangle$ satisfying the equation (7) are called Ishibashi states³⁶. They have the following properties:

$$\langle \langle b|e^{-2\pi\frac{L}{\beta}(\hat{H}_L+\hat{H}_R)}|a\rangle \rangle = \delta_{ba}\chi_a(\tilde{q} = e^{-4\pi\frac{L}{\beta}}).$$
(8)

Here χ_a is the character of an irreducible representation of the CFT.

The boundary states can be expanded in terms of Ishibashi states:

$$|A\rangle = A_a |a\rangle\rangle. \tag{9}$$

The partition function can be written as an amplitude of boundary states:

$$Z_{\rm AB} = \langle \Theta A | e^{-2\pi \frac{L}{\beta} (\hat{H}_L + \hat{H}_R)} | B \rangle.$$
 (10)

Now β is the circumference of space direction with periodic boundary conditions (a circle) and L is propagating time. Here Θ is a CPT operator since these two boundaries have opposite orientation:

$$\Theta c \Theta^{-1} = c^*, \quad \Theta |B\rangle = (B_a)^* |a^+\rangle\rangle, \tag{11}$$

where $|a^+\rangle\rangle = C|a\rangle\rangle$ and C is the charge conjugation operator. In the most CFTs, the CPT operator Θ is just the identity operator.

Then, the partition function can be written as

$$Z_{\rm AB} = \sum_{a} A_a B_a \chi_a (\tilde{q} = e^{-4\pi \frac{L}{\beta}}).$$
(12)

In the open string picture, BCFT is a CFT with boundary conditions specified by A(x = 0) and B(x = L) at these two conformal boundaries in spatial direction and with $t+ix \sim t+\beta+ix$ identified. Since these two pictures can be related by the S-modular transformation — a 90degree rotation of the space-time manifold, the symmetry algebra will transform in the following way:

$$\ln z = i \ln z' , S(z) = i^h S(z') \frac{z'}{z}, \bar{S}(\bar{z}) = i^{-h} S(\bar{z}') \frac{\bar{z}'}{\bar{z}}.$$
(13)

Then the boundary condition in the open chain is obtained from that of closed string picture under the Smodular transformation:

$$S(z) = \rho_{AB}(\bar{S}(\bar{z})) \longrightarrow S(z') = (-1)^h \rho_{AB}(\bar{S}(\bar{z}')),$$
(14)

where $|z'| = 1, \exp(2\pi L/\beta)$. In this picture, the boundary condition constrains the Hibert space of quantum states on the interval which is denoted by \mathcal{H}_{AB} . The partition function (at inverse temperature β) is written as a trace of the Hamiltonian \hat{H}_{AB}^{open} of the finite interval of length with boundary conditions A and B at the two boundaries:

$$Z_{\rm AB} = \operatorname{tr}_{\mathcal{H}_{\rm AB}} e^{-\beta H_{\rm AB}^{\rm open}}.$$
 (15)

The partition function can be rewritten using the Hamiltonian defined purely in the holomorphic sector

since only holomorphic (left-moving) degrees of freedom contributes:

$$Z_{\rm AB} = \operatorname{tr}_{\mathcal{H}_{\rm AB}} q^{\hat{H}^L}.$$
 (16)

All terms in the partition function are powers of $q = e^{-\pi\beta/L}$ related to the length L and inverse temperature β .

Since only holomorphic (left-moving) degrees of freedom appear in the BCFT, the partition function can be decomposed into characters of different irreducible representations ϕ_a of the holomorphic Virasoro algebra³⁷:

$$Z_{\rm AB} = \sum_{a} n^a_{\rm AB} \chi_a(q). \tag{17}$$

The non-negative integers n_{AB}^a represent the multiplicity with which the irreducible representations appear in the Hibert space \mathcal{H}_{AB} .

These characters in the open string and closed string are related by a modular transformation of the spacetime torus:

$$\chi_a(q) = \sum_b S_{ab} \chi_b(\tilde{q}). \tag{18}$$

Now the Cardy condition requires this expression in the closed picture can be interpreted as that in the open picture. That is, the non-negative integers and the expansion coefficients can be related via³⁷:

$$n_{\rm AB}^a = \sum_b A_b B_b S_{ab}.$$
 (19)

III. BOUNDARY CONFORMAL FIELD THEORIES (BCFTS) AND QUANTUM ANOMALIES

In this section, we give a set of arguments which support the advocated relation between BCFTs and gappability of CFTs.

We start by giving a brief overview of gappability of CFTs. A given CFT in (1+1) dimensions can be gapped by a "massive deformation" of a CFT,

$$S \to S - \lambda \int dt dx V(\phi(x)),$$
 (20)

where $V(\phi(x))$ is a relevant operator, and $\lambda \in \mathbb{R}$ is the coupling constant.

To construct boundary conditions, one can also consider a relevant operator whose form depends on the space coordinate. More precisely, one can consider such relevant operator:

$$V(x) = \begin{cases} V_{\rm A}(\phi(x)), & x < 0; \\ 0, & 0 \le x < L; \\ V_{\rm B}(\phi(x)), & L \le x. \end{cases}$$
(21)

If both $V_{\rm A}(\phi)$ and $V_{\rm B}(\phi)$ can gap the system with a unique ground state, only the middle region is still gapless in the sense that the bulk gapless modes have nonzero amplitude only there. At the low energy, the boundaries between the CFT and any of the neighborhood gapped phases are expected to be renormalized into conformal invariant boundary conditions¹⁸. In other words, this implies that there is a relationship between relevant operators and conformal invariant boundary conditions.

For our interest, the interactions are assumed to be related by a symmetry G. That is, in the setup (21), we choose

$$V_{\rm B} = g V_{\rm A}(\phi(x)) g^{-1},$$
 (22)

where g is an element of the symmetry group G, as shown in Fig. 1.

$$V = V_{\mathcal{A}}(\phi) \qquad \qquad V = 0 \qquad \qquad V = gV_{\mathcal{A}}(\phi)g^{-1}$$
$$x = 0 \qquad \qquad x = L$$

FIG. 1. space-dependent interaction with respect to transformation g.

The corresponding boundary conditions should also be related via the same transformation. Now if $V_{\rm A}(\phi)$ is symmetric, the interactions in the region x < 0 and x > Lare same. In the limit $L \to 0$ $(q \to 0)$, the theory is fully gapped and the partition function of BCFT is just that of this gapped phase which is 1:

$$Z_{\rm AB}(q) \to 1, \quad L \to 0.$$
 (23)

If we can find a boundary condition A satisfying Eq. (23) for every $g \in G$, we can argue that the symmetry G is anomaly-free according to the gapping potential argument.

On the other hand, in the closed string picture, the length L plays a role of time. Thus the equation (23) implies the boundary state $|A\rangle$ is invariant under each symmetry transformation g:

$$Z_{AB}(L \to 0) = \langle \Theta A | g | A \rangle = 1, g \in G.$$
(24)

Here we use the fact that the CPT operator is an identity operator for most CFTs. Actually, the equation (24) means the boundary $|A\rangle$ is symmetric under G. Therefore the boundary condition A and interaction $V_{\rm A}(\phi)$ should be also symmetric.

In fact, from an argument about the close relationship between boundary state and ground state after adding gapping potential, we can think the smeared boundary state $e^{-\tau H_{\rm CFT}}|A\rangle$ as the possible ground state after adding the relevant operator $V_{\rm A}(\phi)^{38}$. Since Hamiltonian of the CFT commutes with symmetry G, this possible ground state is symmetric if and only if the boundary state is symmetric.

For example, the Ising CFT has a \mathbb{Z}_2 symmetry which flips every spin and three boundary states $|\uparrow\rangle$, $|\downarrow\rangle$, $|\text{free}\rangle$. Since the boundary state $|\text{free}\rangle$ is symmetric, the corresponding smeared boundary state is a symmetric ground state and the corresponding perturbation is $V = t \int dx \ \hat{\epsilon}(x), \ t > 0$ where ϵ is energy density³⁸.

IV. MULTICOMPONENT U(1) COMPACTIFIED BOSON THEORY

In this section, we will use the boundary state approach to calculate the anomaly of the U(1) compactified boson theory with respect to some global symmetries.

A. Canonical quantization

Multicomponent compactified boson theories are a general class of CFTs which can be realized at the edge of (2+1) dimensional gapped many-body systems with SPT phase¹²:

$$S = \frac{1}{8\pi} \int dx dt [K_{IJ}\partial_t \phi^I \partial_x \phi^J - V_{IJ}\partial_x \phi^I \partial_x \phi^J],$$
(25)

where K matrix is $\bigoplus_{i=1}^{N} \sigma^{x}$ and $I, J = 1, \cdots, 2N$. We can see that this theory has N copies of non-chiral bosons, V is a $2N \times 2N$ symmetric, positive definite real matrix and contains the information like the velocities of the edge modes. Here we choose the V matrix to be an identity matrix in this paper since it does not influence the topological properties of this theory. If this CFT is realized on the cylinder of circumference L, the compact U(1) boson fields ϕ^{I} should satisfy the compact condition with radius $R = \sqrt{2}$:

$$\phi^{I}(x+L,t) = \phi^{I}(x,t) + 2\pi n^{I}\sqrt{2}, \quad n^{I} \in \mathbb{Z}.$$
 (26)

We can also write the action above in the usual way:

$$S = \frac{1}{8\pi} \int dx dt \sum_{I=1}^{I=N} \partial_{\mu} \phi^{2I-1} \partial^{\mu} \phi^{2I-1}.$$
 (27)

We define variable ϕ^{2I} as the dual variable θ^{I} .

It is more convenient to carry out the quantization in the chiral basis φ_I which we define by diagonalizing the K matrix as:

$$P\phi = \varphi, \quad PKP^T = 2\eta.$$
 (28)

where P is $2N \times 2N$ orthogonal matrix and η is a diagonal matrix $(1,-1,1,-1,\cdots)$.

The action can be rewritten in the chiral basis as follows:

$$S = \frac{1}{4\pi} \int dx^2 [\partial_t \varphi_L^I x \varphi_L^I - \partial_t \varphi_R^I x \varphi_R^I - \partial_x \varphi_L^I \partial_x \varphi_L^I - \partial_x \varphi_R^I \partial_x \varphi_R^I].$$
(29)

In this picture, chiral bosons satisfy the compactified condition:

$$\varphi_L^I(x+L,t) = \varphi_L^I(x,t) + 2\pi n^I R, \quad n^I \in \mathbb{Z},
\varphi_R^I(x+L,t) = \varphi_R^I(x,t) + 2\pi m^I R, \quad m^I \in \mathbb{Z}.$$
(30)

After canonical quantization, the boson fields satisfy the commutation relations:

$$\begin{aligned} [\varphi_L^I(x), \partial_x \varphi_L^J(x')] &= 2\pi i \delta^{IJ} \delta(x - x'), \\ [\varphi_R^I(x), \partial_x \varphi_R^J(x')] &= -2\pi i \delta^{IJ} \delta(x - x'). \end{aligned}$$
(31)

The equation of motion is

$$\partial_t \varphi_L^I - \partial_x \varphi_L^I = 0, \partial_t \varphi_R^I + \partial_x \varphi_R^I = 0.$$
 (32)

As a result, the mode expansion is

$$\varphi^{I}(x,t)_{L/R} = \varphi_{0} + \frac{2\pi}{L}(t\pm x)a_{0}^{I} + \sum_{r\neq 0}a_{r}^{I}e^{-\frac{2\pi ri}{L}(t\pm x)}.$$
(33)

The canonical commutation relations and the compactified conditions for the operators are

$$[a_{r,L/R}^{I}, a_{s,L/R}^{J}] = r\delta^{IJ}\delta_{r+s,0},$$

$$\varphi_{0,L/R}^{I} \sim \varphi_{0,L/R}^{I} + 2\pi n^{I}R,$$

$$[\varphi_{0,L/R}^{I}, a_{0,L/R}^{J}] = \pm i\delta^{IJ}.$$
(34)

And the eigenvalue a_0^I is quantized to be m^I , where $\sqrt{2}m^I \in \mathbb{Z}$.

B. Anomaly free condition and Haldane gapping potential

In the boson CFTs, the conformal boundary conditions for Ishibashi states is:

$$(L_r - \bar{L}_{-r})|K\rangle\rangle = 0.$$
(35)

Here L and \overline{L} are Virasoro generators in each chiral sector. A special solution for this equation is given by³⁹:

$$\left(a_{r,L}^{I} - D_{IJ}a_{-r,R}^{J}\right)\left|v\right\rangle = 0.$$
(36)

The Ishibashi states for this condition are:

$$|v\rangle\rangle = \exp\left(\sum_{r=1}^{\infty} \frac{1}{r} a^{I}_{-r,,L} D_{IJ} a^{J}_{-r,,R}\right) |v\rangle.$$
(37)

where $|v\rangle$ are eigenstates of $a_{0,L}^{I}$. And its eigenvalues v^{I} satisfy $\sqrt{2}v^{I} \in \mathbb{Z}$. This eigenstate can written as a coherent state, namely,

$$|v\rangle = e^{i\sum_{I}v^{I}[\varphi_{0,L}^{I} + (D\varphi_{0,R})^{I})]}|0\rangle.$$
(38)

The matrix D can be any $N \times N$ orthogonal matrix with eigenvalue $d^{I} = 1$ or -1. The eigenvector is denoted as e_{I} .

The Cardy states can be constructed as the superposition of Ishibashi states:

$$|B, \{\alpha^I\}\rangle = \otimes_I |B, \alpha^I\rangle, \ |B, \alpha^I\rangle = \sum_{v^I} e^{iv^I \alpha^I} |v^I\rangle\rangle.(39)$$

Such Cardy states can be rewritten as follows:

$$|B, \alpha_{j}\rangle = e^{\sum_{I,J=1}^{N} \sum_{r=1}^{\infty} \frac{1}{r} a_{-r,,L}^{I} D_{IJ} a_{-r,,R}^{J}}} \sum_{v^{I}} \cos[v^{I} (\varphi_{0,L}^{I} + (D\varphi_{0,R})^{I} + \alpha^{I})]|0\rangle.$$
(40)

Here the most relevant operators $\sum_{j=1}^{N} \cos[(\varphi_{0,L}^{I} + (D\varphi_{0,R})^{I} + \alpha^{I})/\sqrt{2}]$ in the cosine term are gapping potential which can condense the value of boson fields. If this interaction satisfies the Haldane condition¹⁹ for chiral Tomonaga-Luttinger liquids

$$e_I^T K e_J = 0, (41)$$

It can gap the boson the theory when the coupling constant is large.

In the open string picture, the Cardy state correspond to the following boundary condition:

$$\varphi_L^I = (D\varphi_R)^I + \alpha^I. \tag{42}$$

So the computation above just implies the information on gapping potential from Cardy states or boundary conditions.

Now for our interest, the boundary states on t = 0 and t = L are assumed to be $|\{\alpha_1^I\}, D_1\rangle$, $|\{\alpha_2^I\}, D_2\rangle$. These two boundary states are related by a symmetry transformation $g \in G$:

$$|\{\alpha_2^I\}, D_2\rangle = g|\{\alpha_1^I\}, D_1\rangle.$$

$$(43)$$

Firstly we assume these two boundary stats have the same D matrix. Then the amplitude is given by:

$$Z(q) = (\eta(q))^{-N} \sum q^{u^2} \exp(-\frac{\beta}{8\pi L}(\alpha_2 - \alpha_1)^2), (44)$$

where u is in the Bravais lattice and $\eta(q)$ is the Dedekind function. Only the $\alpha_2 = \alpha_1$, the partition survives in the limit $L \to 0$.

For the boundary states with the different D matrices, we can assume the number of eigenvalues -1 of D_1D_2 is k (k > 0) in the Ishibashi condition. The partition function is

$$Z(q) \propto (\eta(q))^{-N} (\theta_2(q))^k \sum q^{u^2}.$$
 (45)

It is easy to see that the partition function goes to zero in the limit $q \rightarrow 0$ if the two boundary states have different D matrix.

Therefore we conclude only $\alpha_2 = \alpha_1$ and $D_2 = D_1$, which is equivalent to the boundary state $|A\rangle$ is symmetric, the partition function goes to 1 in the limit $L \to 0$. This property of the partition function is just the special case of the anomaly free condition of our argument in the previous section.

C. Anomaly of $U(1) \times \mathbb{Z}_2$ symmetry

A simple example is the free boson theory with U(1) and \mathbb{Z}_2 symmetry. The element in U(1) and \mathbb{Z}_2 symmetry acts on the ϕ with radius $R = \sqrt{2}$:

$$h^{-1}\phi h = \phi + \pi R,$$

$$h^{-1}\theta h = \theta + \pi R,$$

$$U^{-1}\theta U = \theta + \delta\theta,$$

$$U^{-1}\phi U = \phi.$$
(46)

On the lattice, this boson theory can be realized as the antiferromagnetic Heisenberg spin-1/2 chain. The U(1) symmetry is spin rotation on the z-axis and \mathbb{Z}_2 is the IR emergent \mathbb{Z}_2 partner of translation symmetry. The LSM theorem claims that there is no gapped ground state keeping spin rotation on z-axis and translation symmetry if the number of spin in the unit cell is odd. As claimed above, if one can find a symmetric gapping potential with a unique ground state, there will exist symmetry invariant boundary states. Therefore we expect there will be symmetric boundary states only we consider even copies of boson theory.

For one copy of boson theory, the boundary state is just the Dirichlet boundary state or Neumann boundary state:

$$|B,\phi_{0}\rangle_{D} = \sum_{v \in \frac{\sqrt{2}}{2}\mathbb{Z}} e^{iv\phi_{0}}|v\rangle\rangle_{D},$$
$$|B,\theta_{0}\rangle_{N} = \sum_{v \in \frac{\sqrt{2}}{2}\mathbb{Z}} e^{iv\theta_{0}}|v\rangle\rangle_{N}.$$
(47)

Here

$$|v\rangle\rangle_{D} = e^{\sum_{r>0} \frac{1}{r}a_{-r,L}a_{-r,R}} |a_{0,L} = a_{0,R} = v\rangle,$$

$$|v\rangle\rangle_{N} = e^{\sum_{r>0} -\frac{1}{r}a_{-r,L}a_{-r,R}} |a_{0,L} = -a_{0,R} = v\rangle.$$

(48)

are Ishibashi states satisfying the conformal boundary condition.

Each boundary state can not be symmetric since the U(1) $\times \mathbb{Z}_2$ symmetry changes both ϕ_0 and θ_0 .

However, for two copies of the above theory, $U(1) \times \mathbb{Z}_2$ symmetry is expected to be anomaly-free. The variables of two copies theory are noted by (ϕ_1, θ_1) , (ϕ_1, θ_2) .

We can construct a symmetric boundary state satisfying the constraint:

$$(\theta_1 - \theta_2)|B\rangle = \alpha_1|B\rangle, \quad (\phi_1 + \phi_2)|B\rangle = \alpha_2|B\rangle.$$
(49)

For simplify, we redefine the boson fields¹⁷:

$$\Phi_1 = \frac{1}{\sqrt{2}}(\theta_1 - \theta_2), \quad \Phi_2 = \frac{1}{\sqrt{2}}(\phi_1 + \phi_2).$$
 (50)

The mode expansion is given by:

$$\Phi_{i,L} = \Phi_{i,0,L} + \frac{2\pi}{L}(t+x)c_{i,0} + \sum_{r\neq 0} c_{i,r}e^{-\frac{2\pi i}{L}(t+x)},$$

$$\Phi_{i,R} = \Phi_{i,0,R} + \frac{2\pi}{L}(t-x)\bar{c}_{i,0} + \sum_{r\neq 0} \bar{c}_{i,r}e^{-\frac{2\pi i}{L}(t-x)}(51)$$

The symmetric boundary state for the redefined boson variables is:

$$|B\rangle = \left(\frac{1}{\mathcal{N}_N} \sum_{n_1 \in \mathbb{Z}} e^{iv_1 \alpha_1} |v_1\rangle\rangle_N\right) \otimes \left(\frac{1}{\mathcal{N}_D} \sum_{v_2 \in \mathbb{Z}} e^{in_2 \alpha_2} |v_2\rangle\rangle_D\right).$$
(52)

And the symmetric gapping potential is

$$H' = U[\cos(\frac{1}{R}(\theta_1 - \theta_2 + \alpha_1)) + \cos(\frac{1}{R}(\phi_1 + \phi_2 + \alpha_2))].$$
(53)

For such gapping potential, it satisfies the Haldane condition (41) under which H' is guaranteed to gap the bosons if U is large.

D. Anomaly of $SU(2) \times \mathbb{Z}_2$ symmetry

Actually, when the boson system is on the self-dual point, i.e., the radius is $\sqrt{2}$, the U(1) × \mathbb{Z}_2 symmetry can be extended into a the SU(2)× \mathbb{Z}_2 symmetry and it is equivalent to SU(2)₁ WZW model. Here the \mathbb{Z}_2 is still the symmetry mentioned before and SU(2) symmetry is vector rotation subgroup of SU(2)_L× SU(2)_R with generators

$$S^{z} = \partial_{x}\phi,$$

$$\hat{S}^{+} = \exp(i\frac{1}{\sqrt{2}}(\phi+\theta)) + \exp(i\frac{1}{\sqrt{2}}(-\phi+\theta)),$$

$$\hat{S}^{-} = \exp(-i\frac{1}{\sqrt{2}}(\phi+\theta)) + \exp(-i\frac{1}{\sqrt{2}}(-\phi+\theta)).(54)$$

On the lattice, SU(2) symmetry is the projective representation of SO(3) rotation and \mathbb{Z}_2 symmetry is still translation symmetry.

From the previous analysis, there does not exist boundary states invariant under $SU(2) \times \mathbb{Z}_2$ symmetry in the free boson theory.

Similarly, for two copies of the boson theory, there is no anomaly for $SU(2) \times \mathbb{Z}_2$ symmetry. Thus we expect there will be a symmetric boundary state. Now the generators of SU(2) vector rotation are

$$\hat{S}^{z} = \partial_{x}(\phi_{1} + \phi_{2}),$$

$$\hat{S}^{+} = \sum_{i=1}^{2} \exp(i\frac{1}{\sqrt{2}}(\phi_{i} + \theta_{i})) + \sum_{i} \exp(i\frac{1}{\sqrt{2}}(-\phi_{i} + \theta_{i})),$$

$$\hat{S}^{-} = \sum_{i=1}^{2} \exp(-i\frac{1}{\sqrt{2}}(\phi_{i} + \theta_{i})) + \sum_{i} \exp(i\frac{1}{\sqrt{2}}(\phi_{i} - \theta_{i})).$$
(55)

The symmetric Cardy states satisfy the constraint:

$$\hat{S}^{z}|B\rangle = 0, \quad \hat{S}^{+}|B\rangle = 0,$$
$$\hat{S}^{-}|B\rangle = 0, \quad h|B\rangle = |B\rangle.$$
(56)

A special solution of these equations is

$$|B\rangle = \left(\frac{1}{\mathcal{N}_N} \sum_{v_1 \in \frac{\sqrt{2}}{2} \mathbb{Z}} |v_1\rangle\rangle_N\right) \otimes \left(\frac{1}{\mathcal{N}_D} \sum_{v_2 \in \frac{\sqrt{2}}{2} \mathbb{Z}} e^{iv_2\sqrt{2}\pi} |v_2\rangle\rangle_D\right).$$
(57)

Here $|v_1\rangle\rangle_N$ and $|v_2\rangle\rangle_D$ are the same Ishibashi state mentioned before. The corresponding gapping potential is

$$H' = U(\cos(\frac{\sqrt{2}}{2}(\theta_1 - \theta_2)) + \cos(\frac{\sqrt{2}}{2}(\phi_1 + \phi_2 + \sqrt{2}\pi)).$$
(58)

Moreover, the ground state is a trivial state when U is positive whereas it is the Haldane state when U is negative⁴⁰⁻⁴².

For two copies of the boson system on self-dual point, it can be realized as antiferromagnetic SU(2) spin ladder in 1+1 dimension:

$$H = J \sum_{i} \vec{S}_{i}^{1} \cdot \vec{S}_{i+1}^{1} + \vec{S}_{i}^{2} \cdot \vec{S}_{i+1}^{2}, \quad J > 0.$$
(59)

Here \vec{S} is spin 1/2 operator. In the continuum limit, the spin operators can be expressed by bosonization:

$$S^{x} = \hat{S}^{x} + (-1)^{x/a} \frac{\lambda}{\pi a} \cos(\frac{\sqrt{2}}{2}\theta),$$

$$S^{y} = \hat{S}^{y} + (-1)^{x/a} \frac{\lambda}{\pi a} \sin(\frac{\sqrt{2}}{2}\theta),$$

$$S^{z} = \hat{S}^{z} + (-1)^{x/a} \frac{\lambda}{\pi a} \sin(\frac{\sqrt{2}}{2}\phi).$$
 (60)

The gapping potential (58) can be realized as the following interaction on the lattice:



FIG. 2. SU(2) and translation invariant gapping potential

$$H' = U\lambda_1^2 \sum_i \vec{S}_i^1 \cdot \vec{S}_i^2 - U\lambda_2^2 \sum_i (\vec{S}_i^1 \cdot \vec{S}_{i+1}^1) (\vec{S}_i^2 \cdot \vec{S}_{i+1}^2).$$
(61)

Here λ_1 and λ_2 are nonuniversal constants⁴³. On the lattice, the SU(2) × Z₂ symmetry is spin rotation and IR emergent Z₂ partner of translation symmetry. This interaction is invariant under the spin rotation and translation transformation. To see this lattice model having a unique gapped ground state, we can do the Jordan Wigner transformation and this potential will give a unique ground state and fermionic mass $m \propto 3U^{44}$. This result agrees with the LSM theorem which claims there will be a gapped ground state keeping SO(3) and translation symmetry if the number of spins in each unit cell is even.

Indeed, the standard Abelian bosonization approach⁴⁵ to Haldane gap problem is based on the description of the spin-S antiferromagnetic chain as 2S coupled spin-1/2 chains, which is mapped to 2S-component boson field theory (Tomonaga-Luttinger Liquid) with possible interactions. Often we can focus on one particular linear combination corresponding to the "center-of-mass" field, in the low energy limit. However, when 2S is even, the translation symmetry acts trivially on this center-ofmass field and thus we do not expect anomaly. Thus we generically expect the "Haldane gap". In Ref. 18, the degeneracy in the BCFT spectrum similar to what we discuss in this paper was used as a probe of the Haldane SPT phase. In their construction, the boundary conditions imposed on the two sides correspond to different phases (trivial and SPT), and not related by a symmetry transformation of the system. Despite these differences, it would be interesting to find more deeper connections between their analysis and ours.

E. Anomaly of $SU(N) \times \mathbb{Z}_N$ symmetry

The discussion on the SU(2)₁ WZW model can be generalized to SU(N)₁ WZW model with SU(N) × \mathbb{Z}_N symmetry. For SU(N)₁WZW model, the generators of vector SU(N) transformation is:

$$H^{i} = i\partial_{x}\phi^{i},$$

$$E^{\alpha} = e^{i\alpha\cdot\varphi_{L}} + e^{-i\alpha\cdot\varphi_{R}}.$$
(62)

where α are root vectors with N-1 components, H are generators of the Cartan subgroups and E are other vertex generators. Since the SU(N) symmetry has N-1 H generators, the SU(N)₁ WZW model can be described by N-1 free massless bosons. We should note here the radius of bosons is not $\sqrt{2}$ but depends on the root of SU(N) Lie algebra.

The \mathbb{Z}_N symmetry only acts on the last massless boson:

$$g\varphi_L^{N-1}g^{-1} = \varphi_L^{N-1} + 4\pi \sqrt{\frac{N-1}{N}},$$

$$g\varphi_R^{N-1}g^{-1} = \varphi_R^{N-1}.$$
 (63)

The condition for symmetric boundary states is:

$$H^{i}|B\rangle = 0, \quad E^{\alpha}|B\rangle = 0,$$

$$g|B\rangle = |B\rangle.$$
(64)

For one copy of the $SU(N)_1$ WZW model, in analogy to the previous analysis for the SU(2) case, there does not exist a symmetric boundary state.

However, as the index of LSM anomaly shows, one can find a trivial gapped ground state for N copies of $SU(N)_1$ WZW model²⁹. More precisely, there exists SU(N) and translational invariant gapped interaction with the unique ground state. Therefore we expect to construct a symmetric boundary state for N copies of $SU(N)_1$ WZW model.

Firstly, it is natural to ask how many there are independent constraints for boundary state. To solve this question, we should notice when α is a root, then $-\alpha$ is also a root. Therefore, we only consider positive roots where its first non-zero component is positive. Another helpful information is the communication relations between vertex generators with positive roots:

$$[E^{\alpha}, E^{\beta}] \propto \frac{E^{\alpha+\beta}}{x_1 - x_2}, \quad \alpha + \beta \in \Delta_+,$$
$$[E^{\alpha}, E^{\beta}] = 0, \quad \alpha + \beta \notin \Delta_+, \tag{65}$$

where Δ_+ is the set of positive roots.

As a result, independent positive roots can not be written as the sum or minus of two different positive roots. Actually, there are N-1 independent positive roots for SU(N) group:

$$m_{1} = \frac{\sqrt{2}}{2}(1, 0, 0, \dots, 0) ,$$

$$m_{2} = \frac{\sqrt{2}}{2}(\frac{1}{2}, \frac{\sqrt{3}}{2}, 0, \dots, 0),$$

$$\vdots$$

$$m_{k} = \frac{\sqrt{2}}{2}(\frac{1}{2}, \frac{1}{2\sqrt{3}}, \dots, \sqrt{\frac{1}{2k(k-1)}}, \sqrt{\frac{k+1}{2k}}, 0, \dots, 0),$$

$$\vdots$$

$$m_{N-1} = \frac{\sqrt{2}}{2}\left(\frac{1}{2}, \frac{1}{2\sqrt{3}}, \dots, \sqrt{\frac{N}{2(N-1)}}\right).$$
(66)

So for N copies of $SU(N)_1$ WZW model, the independent constraints for symmetric boundary states can be written in the form of vertex operators with m vector:

$$\sum_{j=1}^{N} E_{j}^{m} |B\rangle = 0, \quad \sum_{j=1}^{N} H_{j}^{i} |B\rangle = 0,$$
$$h|B\rangle = |B\rangle. \tag{67}$$

Here j is the index for the jth copy.

We find a special solution for these equations:

$$\varphi_{j,L}^{1} + \varphi_{j+1,R}^{1} = \sqrt{2}\pi,
\varphi_{j,L}^{2} + \varphi_{j+1,R}^{2} = \frac{2\pi}{\sqrt{6}},
\vdots
\varphi_{j,L}^{k} + \varphi_{j+1,R}^{k} = 2\pi\sqrt{\frac{1}{(k+1)k}},
\vdots
\varphi_{j,L}^{N-1} + \varphi_{j+1,R}^{N-1} = 2\pi\sqrt{\frac{1}{(N-1)N}}.$$
(68)

V. ANOMALY OF WZW MODEL

In this section, we will discuss the WZW models with center symmetry, vector rotation symmetry and time reversal symmetry. We will show our argument is still valid for these models.

A. Introduction to WZW model and global symmetry

The WZW model for the level k is defined with the action:

$$S = \frac{k}{16\pi} \int_{M_2} dx dt \ Tr(\partial_{\mu}g^{-1}\partial^{\mu}g) + \frac{k}{24\pi} \int_{B_3} d^3x \ Tr[(\tilde{g}^{-1}d\tilde{g})^3],$$
(69)

where g is a smooth map from the spacetime to the group manifold G and the second term is defined on a 3-dimensional manifold B_3 whose boundary is the original spacetime. \tilde{g} is denoted to be an extension of g field to B_3 and k is a non-negative integer.

There are three kinds of global symmetry:

1. Center symmetry Γ is the center of the group G which acts as multiplying a U(1) phase:

$$g \to hg, \quad h \in \Gamma.$$
 (70)

2. Vector rotation symmetry acts as the adjoint representation of G:

$$g \to V g V^{-1}$$
, $V \in G$. (71)

3. Charge conjugation symmetry maps the group element to its complex conjugation (without transpose):

$$g \to g^*$$
. (72)

B. Action of global symmetry on Cardy state

For the WZW model, there is a set of special solutions for the Cardy condition³³:

$$n^{a}_{\alpha\beta} = \sum_{i} \frac{S_{\alpha i} S_{\beta i} S_{ia}}{S_{0i}} \in \mathbb{Z}^{+}, \quad |B_{\alpha}\rangle = \sum_{i} \frac{S_{\alpha i}}{\sqrt{S_{0i}}} |i\rangle\rangle,$$
(73)

where $S_{\alpha i}$ is elements of S matrix of the WZW model and $|i\rangle$ is Ishibashi state.

The action of center symmetry on the Cardy states is same as the action of outer automorphism A on the Dynkin label. More precisely, the action of center symmetry on the Ishibashi state is given by

$$h|i\rangle\rangle = e^{-2\pi(A\omega_0,i)}|i\rangle\rangle. \tag{74}$$

Then we can obtain the action on Cardy states³⁰:

$$h|B_{\alpha}\rangle = \sum_{i} \frac{S_{\alpha i}}{\sqrt{S_{0i}}} e^{-2\pi (A\omega_{0},i)} |i\rangle\rangle$$
$$= \sum_{i} \frac{(AS)_{\alpha i}}{\sqrt{S_{0i}}} |i\rangle\rangle = |B_{A\alpha}\rangle.$$
(75)

Here we use the relationship:

$$AS)_{\alpha i} = S_{\alpha i} e^{-2\pi (A\omega_0, i)}.$$
(76)

The charge conjugation will act on the Cardy states as follows:

$$C|B_{\alpha}\rangle = \sum_{i} \frac{S_{\alpha i}}{\sqrt{S_{0i}}} |i^{+}\rangle\rangle,$$

= $C_{\alpha\beta}|B_{\beta}\rangle.$ (77)

where C martix is the charge conjugation operator on the Dynkin label.

C. Mixed anomaly of WZW model

1. Invarinat boundary states of center symmetry and vector rotation symmetry

Let's firstly consider $\mathrm{SU}(N)_k$ WZW model with the center symmetry \mathbb{Z}_N and vector $\mathrm{PSU}(N)$ symmetry.

The generators of PSU(N) symmetry can be written in the chiral form:

$$S^a = J_0^a + \bar{J}_0^a. (78)$$

Here J_0^a and \bar{J}_0^a are zero modes of holomorphic and antiholomorphic affine Lie group.

As we discussed in the previous section, if the WZW model is anomaly free with respect to $PSU(N) \times \mathbb{Z}_N$ symmetry, we expect to construct the symmetric boundary state:

$$V|B\rangle = |B\rangle, \quad h|B\rangle = |B\rangle.$$
 (79)

To construct such a boundary state, we should know how the PSU(N) symmetry acts on the Ishibashi state. In analogy to the boundary state of free boson, we denote an orthogonal basis in the holomorphic and antiholomorphic sector:

$$\begin{aligned} |\phi_i^{\lambda}, m\rangle &= \prod_{k=1,a} \frac{1}{\sqrt{m_k!}} (\frac{j_{-k}^a}{\sqrt{k}})^{m_k} |\phi_i^{\lambda}, 0\rangle, \\ |\bar{\phi}_i^{\lambda}, m\rangle &= \prod_{k=1,a} \frac{1}{\sqrt{m_k!}} (\frac{\bar{j}_{-k}^a}{\sqrt{k}})^{m_k} |\bar{\phi}_i^{\lambda}, 0\rangle. \end{aligned}$$
(80)

Here $|\phi_i^{\lambda}, 0\rangle$ is vacua in different positive representations λ of the Virasoro algebra. Now we can construct the Ishibashi state:

$$|\lambda\rangle\rangle = \sum_{m} |\phi_{i}^{\lambda}, m\rangle \otimes U |\bar{\phi}_{i}^{\lambda}, \bar{m}\rangle, \qquad (81)$$

where U operator is an anti-unitary operator acting on the chiral generators as follows:

$$U^{-1}\bar{J}_n^a U = -(\bar{J}_{-n}^a)^+ = -(\bar{J}_n^a).$$
(82)

The vacuum of $SU(N)_k$ WZW model will form an irreducible representation of vector PSU(N) symmetry:

$$J^{a}|\phi_{i}^{\lambda},m\rangle = \sum_{j} T_{ij}^{\lambda,a}|\phi_{j}^{\lambda},m\rangle,$$

$$\bar{J}^{a}|\bar{\phi}_{i}^{\lambda},m\rangle = \sum_{j} T_{ij}^{\lambda,a}|\bar{\phi}_{j}^{\lambda},m\rangle.$$
 (83)

Here $T_{ij}^{\lambda,a}$ are generators of $\mathrm{SU}(N)$ in the λ representation.

From the above equation, we can show the Ishibashi state is invariant under vector PSU(N) transformation. The detail is shown in Appendix B. As a result, the Cardy state is also invariant under vector PSU(N) transformation which means we only need to find Cardy state satisfying the following conditions:

$$h|B\rangle = |B\rangle. \tag{84}$$

If we denote the Cardy state using the affine Dynkin labels $[\lambda_0; \lambda_1, \cdots, \lambda_{N-1}]$, the action of the center symmetry is given by

$$A: [\lambda_0; \lambda_1, \cdots, \lambda_{N-1}] \to [\lambda_{N-1}; \lambda_0, \cdots, \lambda_{N-2}], (85)$$

The solution for invariant condition is:

$$\lambda_0 = \lambda_1 = \dots = \lambda_{N-1},\tag{86}$$

$$k = \lambda_0 + \lambda_1 + \dots + \lambda_{N-1} = N\lambda_0.$$
(87)

Therefore only when k is a multiple of N, there exists a symmetric Cardy state. Since the index of mixed anomaly with respect to $PSU(N) \times \mathbb{Z}_N$ symmetry is k mod N, there exists a translation and PSU(N) symmetric gapped ground state only when the level k is a multiple of N^{29} . So the result of BCFT approach agrees with this classification of mixed anomaly and the symmetric Cardy state corresponds to the translation and SU(N) invariant ground state.

Moreover, for the SU(2)_{2k} WZW model, we can show the symmetric boundary state corresponds to the massive phase after adding symmetric gapping potential $\lambda \operatorname{tr}(g^2)$ $(\lambda > 0)$. When k is even, the ground state is trivial state and when k odd, the ground state is Haldane state⁴⁶. This can also be seen from the partition function of an open string. The detail of calculation is shown in the Appendix D.

We can also apply this argument to the WZW model with other simple Lie algebras. The results are listed in TABLE II. The details of calculation are shown in the Appendix B. The anomaly-free conditions on the level k for most Lie algebras there implies the classification of LSM-type anomaly, e.g., $k \in N\mathbb{N}$ for SU(N) consistent with the corresponding \mathbb{Z}_N classification of LSM anomaly index²⁹ and the projective representation argument⁴⁷ on lattices.

Moreover, our results for the Spin(n) WZW models including B and D series are also consistent with a generalized LSM theorem for SO(n) spin chains proposed in Ref. 31 and 32, which claims that an SO(n) spin chain in the *n*-dimensional vector representation with even n has either gapless excitations or degenerate gapped ground states with broken translational symmetry. In particular, the model studied in Ref. 31 is the SO(n) bilinearbiquadratic spin chain whose low-energy effective field theory is described a $\text{Spin}(n)_1$ WZW model. According to our results in TABLE II (or TABLE I), such a lowenergy theory indeed has an LSM-type anomaly if n is even.

2. Invariant boundary states of time reversal and vector rotation symmetry

We can also consider WZW models with vector rotation and time-reversal symmetries. There are two types of time-reversal symmetry: one acts as $\mathcal{T}_1: g(x,t) \rightarrow g^{-1}(x,-t)$ and the other is given by the combination of \mathcal{T}_1 and any order 2 element h_2 (if exists) of the center symmetry group, $\mathcal{T}_2 = \mathcal{T}_1 h_2: g(x,t) \rightarrow -g^{-1}(x,-t)$; either one squares to the identity, i.e. $\mathcal{T}_1^2 = \mathcal{T}_2^2 = 1$. As we will seen in the following discussion, however, it is only \mathcal{T}_2 which has a mixed anomaly with the vector rotation symmetry.

To construct the Cardy states, we work in Euclidean signature by performing the Wick rotation $t = -i\tau$. And we should consider the boundary condition in the closed channel where the space-time cylinder has been rotated by $\pi/2$, namely, $(x', \tau') = (\tau, -x)$. Due to the Lorentz symmetry of CFTs, the original time-reversal symmetry \mathcal{T}_{η} , which is an anti-unitary operator in the Lorentz signature, becomes the unitary \mathcal{CP}_{η} symmetry in the Euclidean signature:

$$C\mathcal{P}_1: g(x', \tau') \to (g^*)^{-1}(-x', \tau'),$$
 (88)

$$C\mathcal{P}_2: g(x', \tau') \to -(g^*)^{-1}(-x', \tau').$$
 (89)

Here we can see \mathcal{CP}_1 is the combination of charge conjugation C and spatial reflection P.

In the first case, the Cardy state should satisfy the constraint as follows:

$$C\mathcal{P}_1|B\rangle = |B\rangle,$$

$$V|B\rangle = |B\rangle.$$
(90)

Since the spatial reflection maps x' to -x', it exchanges the state of left and right handed section. Thus the Ishibashi states (81) and Cardy states are invariant under the spatial reflection symmetry.

Thus we only need to find the invariant Cardy state under the charge conjugation which acts on the Cardy states as follows:

$$C: [\lambda_0; \lambda_1, \cdots, \lambda_{N-1}] \to [\lambda_0; \lambda_{N-1}, \cdots, \lambda_2, \lambda_1].$$
(91)

Cartan matrix	Group	Action of h	Invariant Boundary states
A_{N-1}	$\mathrm{SU}(N)$	$[\lambda_{N-1};\lambda_0,,\lambda_{N-2}]$	$k\in N\mathbb{N}$
B_N	$\operatorname{Spin}(2N+1)$	$[\lambda_1;\lambda_0,,\lambda_{N-1},\lambda_N]$	$k\in\mathbb{N}$
C_N	$\mathrm{USp}(N)$	$[\lambda_N;\lambda_{N-1},,\lambda_1,\lambda_0]$	$k\in\!\!2\mathbb{N}$ or $N\in\!\!2\mathbb{N}$
D_{2N+1}	$\operatorname{Spin}(4N+2)$	$[\lambda_{2N};\lambda_{2N+1},\cdots,\lambda_1,\lambda_0]$	$k\in\!\!4\mathbb{N}$
D_{2N}	$\operatorname{Spin}(4N)$	$[\lambda_1;\lambda_0,\lambda_2,\cdots,\lambda_{2N},\lambda_{2N-1}]$	$k\in\!\!2\mathbb{N}$
D_{2N}	$\operatorname{Spin}(4N)$	$[\lambda_{2N};\lambda_{2N-1},\lambda_{2N-2},\cdots,\lambda_0]$	_
E_6	E_6	$[\lambda_1;\lambda_5,\lambda_4,\lambda_3,\lambda_6,\lambda_0,\lambda_2]$	$k \in 3\mathbb{N}$
E_7	E_7	$[\lambda_6;\lambda_5,\lambda_4,\lambda_3,\lambda_2,\lambda_1,\lambda_0,\lambda_7]$	$k\in\!\!2\mathbb{N}$

12

TABLE II. The action of the center symmetry on Cardy state in the WZW model and condition for the existing of symmetric Cardy state

The roots of boundary state should satisfy :

$$\lambda_i = \lambda_{N-i}, \quad i > 0. \tag{92}$$

When N is even, k is given by

$$k = \lambda_0 + 2(\lambda_1 + \dots + \lambda_{\frac{N}{2}-1}) + \lambda_{\frac{N}{2}}.$$
(93)

On the other hand, when N is odd, k is given by

$$k = \lambda_0 + 2(\lambda_1 + \dots + \lambda_{\frac{N-1}{2}}). \tag{94}$$

In each condition, k can take an arbitrary natural number. Therefore, there is no mixed anomaly between the \mathcal{T}_1 and $\mathrm{PSU}(N)$ symmetry.

In the second case, since the \mathcal{T}_2 symmetry needs a order 2 element of center symmetry, it only exists when N is even. The condition for an invariant boundary state is:

$$C\mathcal{P}_2|B\rangle = C\mathcal{P}_1 h_2 |B\rangle = |B\rangle,$$

$$V|B\rangle = |B\rangle.$$
(95)

Since Cardy states are invariant under the spatial reflection symmetry, we only need to find the invariant Cardy state under the combination of charge conjugation and the order 2 element of the center symmetry group.

The Dynkin roots of such boundary state should satisfy

$$[\lambda_0; \lambda_1, ..., \lambda_{N-1}] = [\lambda_{N/2}; \lambda_{N/2-1}, \cdots, \lambda_{N/2+2}, \lambda_{N/2+1}].$$
(96)

Thus the relations $\lambda_i = \lambda_{N/2-i}$ if $0 \leq i \leq N/2$ and $\lambda_i = \lambda_{3N/2-i}$ if $N/2 < i \leq N-1$) hold for symmetric boundary states. When $N \in 4\mathbb{Z} + 2$, the level k is given by

$$k = 2(\lambda_0 + \dots + \lambda_{\frac{N-2}{4}} + \lambda_{\frac{N+2}{2}} + \dots + \lambda_{\frac{3N-2}{4}}).$$
(97)

And when $N \in 4\mathbb{N}_+$, k is given by

$$k = 2(\lambda_0 + \dots + \lambda_{\frac{N-4}{4}} + \lambda_{\frac{N+2}{2}} + \dots + \lambda_{\frac{3N-4}{4}}) + \lambda_{\frac{N}{4}} + \lambda_{\frac{3N}{4}}.$$
(98)

So a symmetric boundary state exists only when k is even or when N is a multiple of 4. As a result, there should be no mixed anomaly between PSU(N) and \mathcal{T}_2 symmetry under the same condition.

D. Application to a spin chain with the triple product interactions

On the other hand, when $N \in 4\mathbb{N} + 2$, $\mathrm{PSU}(N)$ and \mathcal{T}_2 have a mixed anomaly. The anomaly implies a LSM-type ingappability under the $\mathrm{PSU}(N) \times \mathcal{T}_2$ symmetry. Let us illustrate the point with the simplest case with N = 2. A lattice model which reflect the ingappability of the $\mathrm{SU}(2)_1$ WZW theory is given by the spin-1/2 chain with $\mathrm{SO}(3)$ -invariant three-spin (triple-product) interactions⁴⁸

$$H = \sum_{j} (-1)^{j} \mathbf{S}_{j} \cdot (\mathbf{S}_{j+1} \times \mathbf{S}_{j+2}).$$
(99)

Although this model is invariant under the PSU(2) symmetry, it is invariant only under two-site translation and not one-site translation. Thus, the standard LSM theorem does not give any constraint on this model. Nevertheless, in Ref. 48 the system was found to be gapless and described by the SU(2)₁ WZW theory. To our knowledge, the mechanism behind this phenomenon has not been clarified. Here we point out that the gapless nature of the model is not accidental but rather reflects the ingappability of the SU(2)₁ WZW theory due to the mixed anomaly of PSU(2) and \mathcal{T}_2 symmetries.

On the lattice, \mathcal{T}_2 corresponds to the combination of the time reversal $\mathcal{T}_1 : \mathbf{S}_j \to -\mathbf{S}_j$ and the one-site translation. Thus, the mixed anomaly implies an LSM-type ingappability of one-dimensional PSU(2)-symmetric spin systems which are invariant under the combined operation. This is a field-theory derivation of a simple onedimensional case of the LSM-type ingappability due to magnetic space group symmetries⁴⁹.

Indeed, we can observe that the model (99) has this symmetry, since the Hamiltonian is odd under both the time reversal \mathcal{T}_1 and the one-site translation. Therefore, the model (99) can be gapped only if the ground states are at least doubly degenerate. This was a background why it was gapless without any fine-tuning of parameters⁴⁸.

In fact, this ingappability does not require the full PSU(2) symmetry and can be protected by the smaller symmetry

$$\mathcal{T}_2: T_2 = TKU_T = TK \prod_{j=1}^{j=L} i\sigma_j^y,$$
 (100)

$$U(1)_z: U_\theta = \exp(i\theta \sum_{j=1}^L \mathbf{S}_j^z), \qquad (101)$$

$$\mathbb{Z}_{2}^{y}: R_{y}^{\pi} = \prod_{j=1}^{L} i\sigma_{j}^{y}.$$
 (102)

We present a simple proof in Appendix E, which is analogous to the original proof of the LSM theorem.

Moreover, we can obtain the similar argument of the WZW models with other simple Lie groups. On the one hand, the result for \mathcal{T}_2 and vector rotation symmetry is listed in the Table III. The detail of calculation is shown in the Appendix C. On the other hand, there is always no mixed anomaly between \mathcal{T}_1 and vector rotation symmetry.

As mentioned in the Introduction, anomaly-free conditions of the WZW model was discussed earlier in Ref. 30 using a similar formulation. However, they discuss the anomaly-free condition for the center symmetry with the charge conjugation. In contrast, we have studied the mixed anomaly of the Lie Group and center symmetries, which is more relevant to the LSM-type ingappability of lattice models.

VI. DISCRETE SYMMETRIES OF MINIMAL MODELS

In this section, we will discuss the global discrete symmetries of minimal models.

The classification of c < 1 minimal models is given in terms of a pair of Dynkin diagrams (A_h,G) where G is A - D - E type. The boundary states of (A_h,G) models are labeled by pair (r,a) of nodes of (A_h,G) graph with the identification⁵⁰:

$$(r, a) = (h + 1 - r, \gamma(a)).$$
(103)

Here γ is an automorphism of the *G* graph. Thus independent boundary states are half of nodes of the (A_h, G) graph.

These minimal models have a unique and maximal \mathbb{Z}_2 symmetry except six cases⁵¹. The (A_4, D_4) (the critical 3-state Potts model) and (A_6, D_4) (the tricritical 3-state Potts model) have a \mathbb{Z}_2 and \mathbb{Z}_3 symmetry which combine to an S_3 symmetry. The four models related to the E_7 and E_8 have no symmetry.

A. Invariant boundary states of \mathbb{Z}_2 symmetry

Let's firstly consider the invariant boundary state of minimal models with unique \mathbb{Z}_2 symmetry. Since this \mathbb{Z}_2 symmetry is anomaly free⁵², we expect there are always invariant boundary states.

For (A_{m-1}, A_m) models, the Cardy state (a,b) is labeled by points of its Dynkin graph. When m is odd, the \mathbb{Z}_2 symmetry maps the (a,b) to (a,m+1-b). Thus the invariant boundary state is $(a, \frac{m+1}{2})$. On the other hand, when m is even, the \mathbb{Z}_2 symmetry maps the (a,b) to (m-a,b). And the invariant boundary state is $(\frac{m}{2},b)$.

For (A_{4l}, D_{2l+2}) models, the \mathbb{Z}_2 symmetry exchanges two Cardy states if they correspond the same point of A graph and two endpoints of D graph. Thus the other Cardy state is invariant under \mathbb{Z}_2 symmetry. Since the action of \mathbb{Z}_2 symmetry of the (A_{4l+2}, D_{2l+2}) , (A_{4l+2}, D_{2l+3}) and (A_{4l+4}, D_{2l+3}) models, is same as above, the calculation for the invariant boundary state is also similar.

For (A_{10}, E_6) and (A_{11}, E_6) models, the \mathbb{Z}_2 symmetry is the reflection symmetry of E_6 graph. Thus the invariant boundary states correspond to nodes of reflection axis in the E_6 graph.

B. Invariant boundary states of S_3 symmetry

Now we consider the S_3 symmetry of (A_4, D_4) (the critical 3-state Potts model) and (A_6, D_4) (the tricritical 3-state Potts model).

In these two models, the S_3 symmetry is permutation of three outside nodes of the D_4 graph. Thus the invariant boundary states correspond to the only one inside node of the D_4 graph which implies the S_3 symmetry should be anomaly free.

For example, there are eight boundary states for (A_4,D_4) model. The first three states $|A\rangle$, $|B\rangle$, $|C\rangle$ describe fixed boundary conditions where the spin on the boundary takes one of three possible values. The mixed boundary states $|AB\rangle$, $|BC\rangle$, $|AC\rangle$ describe boundary conditions where the spin on the boundary can take on two values independently. These six boundary states correspond to six outside nodes of (A_4,D_4) graph. The remain two boundary states $|ABC\rangle$, $|N\rangle$ correspond to free boundary conditions and are invariant under the S_3 symmetry.

Actually, the \mathbb{Z}_3 subgroup is \mathbb{Z}_3 rotation of 3-state Potts model on the lattice, thus it is anomaly-free as gauging it results in the same theory of the original one⁵³. This is nothing but the Kramers-Wannier duality of the 3-state Potts model.

Cartan matrix	Group	action of CP_2	Invariant Boundary states
A_{N-1}	$\mathrm{SU}(N)$	$[\lambda_{N/2};\lambda_{N/2-1},,\lambda_{N/2+1}]$	$k\in\!\!2\mathbb{N}$ or $N\in\!\!4\mathbb{N}$
B_N	$\operatorname{Spin}(2N+1)$	$[\lambda_1;\lambda_0,,\lambda_{N-1},\lambda_N]$	$k\in\mathbb{N}$
C_N	$\mathrm{USp}(N)$	$[\lambda_N;\lambda_{N-1},,\lambda_1,\lambda_0]$	$k\in\!\!2\mathbb{N}$ or $N\in\!\!2\mathbb{N}$
D_{2N+1}	$\operatorname{Spin}(4N+2)$	$[\lambda_1;\lambda_0,\lambda_2,,\lambda_{2N},\lambda_{2N+1}]$	$k\in\mathbb{N}$
D_{2N}	$\operatorname{Spin}(4N)$	$[\lambda_1;\lambda_0,\lambda_2,,\lambda_{2N},\lambda_{2N-1}]$	$k\in\!\!2\mathbb{N}$
D_{2N}	$\operatorname{Spin}(4N)$	$[\lambda_{2N};\lambda_{2N-1},\lambda_{2N-2},,\lambda_1,\lambda_0]$	_
E_7	E_7	$[\lambda_6;\lambda_5,\lambda_4,\lambda_3,\lambda_2,\lambda_1,\lambda_0,\lambda_7]$	$k\in\!\!2\mathbb{N}$

TABLE III. The action of \mathcal{CP}_2 symmetry on Cardy state in the WZW model and condition for the existing of symmetric Cardy state

VII. T-DUALITY SYMMETRY OF $SU(2)_1$ WZW MODEL

In the previous sections, we consider two kinds of symmetries that act on the zero modes of the primary fields of CFTs. The constructing of symmetric Cardy states implies that the potential can gap a CFT without spontaneous symmetry breaking. The gapping potentials appearing in the free boson theory are all Haldane null vectors. A natural question to be asked is whether symmetric Cardy states can imply a gapping potential beyond the usual Haldane null-vectors.

In this section, we will discuss a related example: the $SU(2)_1$ WZW model with T-duality symmetry.

A. T-duality symmetry of $SU(2)_1$ WZW model

For a U(1) boson theory with a compacting radius, there is a duality relating theories of radius 2/R and R. Only the theory with the self-dual radius which is exactly $SU(2)_1$ WZW model, this duality becomes a symmetry which exchanges of two fields

$$T\phi T^{-1} = \theta, \quad T\theta T^{-1} = \phi. \tag{104}$$

This T-duality symmetry not only acts on the zero modes of fields, but also changes every mode in the expansion. In the chiral representation, it acts as follows:

$$T\varphi_{0,R}T^{-1} = -\varphi_{0,R}, \quad Ta_{r,R}T^{-1} = -a_{r,R}.$$
 (105)

B. Anomaly of T-duality symmetry and center symmetry

For our interest, we firstly consider Cardy states invariant under both T-duality symmetry and center symmetry h. The SU(2)₁ WZW model has SU(2)_L × SU(2)_R transformation and the generators can be written in terms of boson field:

$$J^{z}(z) = \partial_{z}(\phi + \theta) ,$$

$$J^{+}(z) = \exp\left[i\frac{\sqrt{2}}{2}(\phi + \theta)\right] ,$$

$$J^{-}(z) = \exp\left[-i\frac{\sqrt{2}}{2}(\phi + \theta)\right] ,$$

$$\bar{J}^{z}(\bar{z}) = \partial_{\bar{z}}(-\phi + \theta) ,$$

$$\bar{J}^{+}(\bar{z}) = \exp\left[i\frac{\sqrt{2}}{2}(-\phi + \theta)\right] ,$$

$$\bar{J}^{-}(z) = \exp\left[-i\frac{\sqrt{2}}{2}(-\phi + \theta)\right] .$$
 (106)

In this representation, the center symmetry is the minus identity operator $h = -\mathbb{I}$ and the T-duality symmetry acts as π rotation along the *x*-axis on the antiholomorphic boson:

$$T J^{z} T^{-1} = J^{z},$$

$$T J^{\pm} T^{-1} = J^{\pm},$$

$$T \bar{J}^{z} T^{-1} = -\bar{J}^{z},$$

$$T \bar{J}^{\pm} T^{-1} = \bar{J}^{\mp}.$$
(107)

Since the 4π rotation is identity in the SU(2) rotation, the T-duality symmetry is a \mathbb{Z}_4 symmetry in the spin representation. This extension from \mathbb{Z}_2 to \mathbb{Z}_4 is because of the center symmetry acting like minus identity⁵⁴. In the next section, we will move to the case with only Tduality symmetry where it is a \mathbb{Z}_2 symmetry.

Because $T^2 = h$ after extension, we only need Cardy states invariant under the extended T-duality symmetry:

$$T|B\rangle = |B\rangle. \tag{108}$$

To construct such Cardy states, we need to do vector $\pi/2$ rotation along the *y*-axis:

$$J^{z} \to J^{\prime x}, \quad J^{\prime x} \to -J^{\prime z}, \quad J^{\prime y} \to J^{\prime y}, \bar{J}^{z} \to \bar{J}^{\prime x}, \quad \bar{J}^{x} \to -\bar{J}^{\prime z}, \quad \bar{J}^{t} \to \bar{J}^{\prime y}.$$
(109)

In the new coordinate, the T-duality symmetry acts as π rotation along the z-axis on the right-handed sector:

$$T J'^{z} T^{-1} = J^{z},$$

$$T J'^{\pm} T^{-1} = J^{\pm},$$

$$T \bar{J}'^{z} T^{-1} = \bar{J}'^{z},$$

$$T \bar{J}'^{\pm} T^{-1} = -\bar{J}'^{\pm}.$$
(110)

We can also represent these generators in terms of new boson fields ϕ' and θ' . Therefore the new T-duality symmetry acts on the new boson fields as follows:

$$T\phi'T^{-1} = \phi' + \frac{3\pi\sqrt{2}}{2},$$

$$T\theta'T^{-1} = \theta' + \frac{\pi\sqrt{2}}{2}.$$
 (111)

For one boson theory, there is no symmetric boundary state. However, for 4 copies of $SU(2)_1$ WZW model, we can redefine the boson fields:

$$\phi_1'' = \frac{1}{\sqrt{4}} (\phi_1' + \phi_2' + \phi_3' + \phi_4'),
\theta_2'' = \frac{1}{\sqrt{2}} (\theta_1' - \theta_2'),
\theta_3'' = \frac{1}{\sqrt{2}} (\theta_2' - \theta_3'),
\theta_4'' = \frac{1}{\sqrt{2}} (\theta_3' - \theta_4').$$
(112)

The symmetric Cardy state takes the form

$$|B\rangle = \otimes_{i=1}^{4} \left(\frac{1}{\mathscr{N}} \sum_{v_i} |v_i\rangle \right), \qquad (113)$$

where the v_i is Ishibashi state for the boson ϕ_i . The first boson takes the Dirichlet boundary state and the others take Neumann boundary states.

The corresponding gapping potential can be given from the Cardy state

$$H' = U\left[\cos\left(\frac{\sqrt{2}}{2}(\phi_1' + \phi_2' + \phi_3' + \phi_4')\right) + \cos\left(\frac{\sqrt{2}}{2}(\theta_1' - \theta_2')\right) + \cos\left(\frac{\sqrt{2}}{2}(\theta_2' - \theta_3')\right) + \cos\left(\frac{\sqrt{2}}{2}(\theta_3' - \theta_4')\right)\right].$$
 (114)

After vector $-\pi/2$ rotation along the *y*-axis, we get the invariant gapping potential under original symmetry:

$$H' = U \sum_{j=1}^{4} \cos(\frac{\sqrt{2}}{2}\phi_j) \cos(\frac{\sqrt{2}}{2}\phi_{j+1}) + U \sum_{j=1}^{4} \cos(\frac{\sqrt{2}}{2}\theta_j) \cos(\frac{\sqrt{2}}{2}\theta_{j+1}) + U \operatorname{Re} \prod_{j=1}^{4} \left[\cos(\frac{\sqrt{2}}{2}\phi_j) + i\cos(\frac{\sqrt{2}}{2}\theta_j) \right].$$
(115)

Where Re means taking the real part of the third term.

This potential can also be realized as a SU(2) spin ladder model:

$$H' = U(\lambda_2)^2 \sum_{j=1}^{4} \sum_k S_{j,k}^z S_{j,k+1}^z + S_{j,k}^y S_{j,k+1}^y + U(\lambda_1)^4 \sum_k \operatorname{Re} \prod_{j=1}^{4} (S_{j,k} \cdot S_{j+1,k} + iS_{j,k}^x), \quad (116)$$

where λ_1 and λ_2 are nonuniversal constants.

This potential satisfies the translation symmetry which is reduced to the center symmetry in the low energy. However, it remains unclear how to realize the T-duality symmetry in the lattice model, so we conclude that the mixed anomaly between the T-duality and center symmetry is at least stable in a perturbative manner around this critical point in the field theory. For the perturbation on the lattice, it is left for the future research.

C. Pure anomaly of T duality symmetry

Now we can consider the case of $SU(2)_1$ WZW model with only T-duality symmetry. Since the corresponding SPT phase is trivial⁵⁵, we expect the boson CFT can be trivially gapped with persevering T-duality symmetry and has invariant Cardy states.

When there is no \mathbb{Z}_2 center symmetry, we do not need to see it as a subgroup of $\mathrm{SU}(2)_R$. To construct the symmetric boundary state, we also apply the vector $\mathrm{SU}(2)$ transformation. The fundamental operator is the vertex operator $\exp(\pm i\theta/\sqrt{2})$ and $\exp(\pm i\phi/\sqrt{2})$. This is equivalent to cosine terms $\cos(\theta/\sqrt{2})$ and $\cos(\phi/\sqrt{2})$ and similar sine terms. Under vector $\pi/2$ rotation along y-axis, these terms transform as follows:

$$\cos(\frac{\sqrt{2}}{2}\theta') = -\sin(\frac{\sqrt{2}}{2}\phi),$$

$$\sin(\frac{\sqrt{2}}{2}\theta') = \sin(\frac{\sqrt{2}}{2}\theta),$$

$$\cos(\frac{\sqrt{2}}{2}\phi') = \cos(\frac{\sqrt{2}}{2}\phi),$$

$$\sin(\frac{\sqrt{2}}{2}\phi') = \cos(\frac{\sqrt{2}}{2}\theta).$$
(117)

In this new coordinate, the T-duality symmetry acts like:

$$T \cos(\frac{\sqrt{2}}{2}\theta')T^{-1} = -\sin(\frac{\sqrt{2}}{2}\theta'),$$

$$T \sin(\frac{\sqrt{2}}{2}\theta')T^{-1} = -\cos(\frac{\sqrt{2}}{2}\theta'),$$

$$T \cos(\frac{\sqrt{2}}{2}\phi')T^{-1} = \sin(\frac{\sqrt{2}}{2}\phi'),$$

$$T \sin(\frac{\sqrt{2}}{2}\phi')T^{-1} = \cos(\frac{\sqrt{2}}{2}\phi').$$
 (118)

This transformation is equivalent to a new \mathbb{Z}_2 symmetry:

$$\phi' \to \frac{\pi\sqrt{2}}{2} - \phi', \quad \theta' \to \frac{3\sqrt{2}\pi}{2} - \theta'.$$
 (119)

We can find a symmetric Cardy states:

$$|B\rangle = \sum_{v \in \frac{\sqrt{2}}{2}\mathbb{Z}} e^{-i\frac{\sqrt{2}\pi}{4}v} |v\rangle\rangle_D.$$
(120)

And the gapping potential is given by:

$$H' = \sqrt{2}U\cos(\frac{\sqrt{2}}{2}\phi' - \frac{\pi}{4}).$$
 (121)

After rotating back, the corresponding gapping potential is given by:

$$H' = U\left[\cos(\frac{\sqrt{2}}{2}\theta) + \cos(\frac{\sqrt{2}}{2}\phi)\right].$$
 (122)

On the lattice, it can be realized as a spin 1/2 antiferromagnetic Heisenberg chain in a staggered magnetic field along the *x*-direction⁵⁶:

$$H = H_0 + H' = J \sum_{i} S_i \cdot S_{i+1} + U\lambda_1 \sum_{i} (-1)^i S_i \cdot S_{i+1} + U\lambda_2 \sum_{i} (-1)^i S_i^x,$$
(123)

where λ_1 and λ_2 are nonuniversal constants.

Since the gapping potential (121) can condense the boson field and have a unique ground state, the gapping potential (122) will also gap the boson theory with a unique ground state after rotating back. As a result, the ground state of corresponding Hamiltonian on the lattice should be unique.

Now if we also consider the center symmetry:

$$\phi' \to \phi' + \sqrt{2}\pi, \quad \theta' \to \theta' + \sqrt{2}\pi, \quad (124)$$

there will be an invariant boundary state under these symmetry only for 4k copies of $SU(2)_1$ WZW model. And the boundary state for 4 copies theory is same as the equation (113). Therefore we obtain the same result whether we do extension of T-duality symmetry or not.

On the lattice, the T-duality symmetry is not an exact symmetry, but can be an emergent symmetry in the lowenergy limit. It would be interesting to study the consequence of the emergent T-duality on the lattice model.

VIII. CONCLUSION AND DISCUSSION

In this work, we discuss the relationship between global anomaly and boundary states of CFTs from the perspective of massive deformation. That is if a 1+1d CFT with symmetry G is anomaly-free, there will be symmetric boundary conditions where the partition function of BCFT converges to one when the length goes to zero. For the closed string, symmetric boundary conditions imply there will be symmetric boundary states which can describe the symmetric gapped ground state.

Then we apply our approach on several examples and show the anomaly-free condition derived from the existence of symmetric boundary conditions is consistent with the result from the 't Hooft anomaly-free condition obtained directly from the bulk CFT. In the multicomponent U(1) boson theory, the existence of symmetric boundary conditions can imply the symmetric gapping potentials which belong to Haldane null vectors. We also show this relationship between global anomaly and boundary states can be generalized to WZW models. As the last example, for SU(2)₁ WZW model with the Tduality symmetry, we use our approach to find the symmetric gapping potentials which are beyond the Haldane null vectors. This result coincides with the fact that there is no anomaly for the T-duality symmetry.

We should note there is slightly different context discussing the relation between BCFT and 1+1d SPT phases or anomalous 0+1d critical theory¹⁸. In this reference, it is found that boundary state $|B\rangle_h$ in the sector twisted by $h \in G$ can defect the anomalous phase $\epsilon(g|h)$ which is related to the cocycles in $H^2(G, U(1))$:

$$g|B\rangle_h = \epsilon_B(g|h)|B\rangle_h.$$
 (125)

More precisely, this correspondence detects the action of symmetry operation on the twisted boundary condition or twisted boundary state. However in our approach, we are looking for boundary states in the untwisted sector which remain invariant under the global symmetry G.

As an outlook, it is interesting to consider the relationship between boundary condition and quantum anomaly in the fermionic CFTs. Here symmetry should include global symmetry G and the fermionic parity symmetry $(-1)^{F}$. The key to solving this problem is searching for the boundary states of a fermionic minimal model. For the bulk fermionic CFTs, they can be constructed from A-type bosonic $CFTs^{52}$ and two exceptional $CFTs^{57}$ attached with a Kitaev chain after \mathbb{Z}_2 orbifolding^{58,59} with a parafermionic generalization⁶⁰. On the lattice model, this transformation is called the Jordan-Wigner transformation and, more generally, the Fradkin-Kadanoff transformation⁶¹. Naturally, there is a correspondence between boundary states of bosonic CFTs and fermionic CFTs^{62,63}. This correspondence can help us find symmetric boundary states of fermionic CFTs from that of related bosonic CFTs.

ACKNOWLEDGEMENTS

L. L. is supported by Global Science Graduate Course (GSGC) program at the University of Tokyo. This work was supported in part by MEXT/JSPS KAKENHI Grants No. JP17H06462 and No. JP19H01808, and by JST CREST Grant No. JPMJCR19T2.

- * cthsieh@phys.ntu.edu.tw
- [†] yuan.yao@riken.jp
- ¹ G. Hooft, Recent developments in gauge theories , 135 (1980).
- ² Z.-C. Gu and X.-G. Wen, Physical Review B 80, 155131 (2009).
- ³ X. Chen, Z.-C. Gu, and X.-G. Wen, Physical review b 83, 035107 (2011).
- ⁴ X. Chen, Z.-C. Gu, and X.-G. Wen, Physical review b 84, 235128 (2011).
- ⁵ M. Z. Hasan and C. L. Kane, Reviews of modern physics 82, 3045 (2010).
- ⁶ X.-L. Qi and S.-C. Zhang, Reviews of Modern Physics 83, 1057 (2011).
- ⁷ L. Fu, Physical Review Letters **106**, 106802 (2011).
- ⁸ X. Chen, Z.-C. Gu, Z.-X. Liu, and X.-G. Wen, Physical Review B 87, 155114 (2013).
- ⁹ X. Chen, Z.-C. Gu, Z.-X. Liu, and X.-G. Wen, Science 338, 1604 (2012).
- ¹⁰ F. Pollmann, E. Berg, A. M. Turner, and M. Oshikawa, Phys. Rev. B 85, 075125 (2012).
- ¹¹ L. Fidkowski and A. Kitaev, Physical review b 83, 075103 (2011).
- ¹² Y.-M. Lu and A. Vishwanath, Phys. Rev. B 86, 125119 (2012).
- ¹³ S. Ryu and S.-C. Zhang, Phys. Rev. B **85**, 245132 (2012).
- ¹⁴ O. M. Sule, X. Chen, and S. Ryu, Phys. Rev. B **88**, 075125 (2013).
- ¹⁵ A. Vishwanath and T. Senthil, Physical Review X 3, 011016 (2013).
- ¹⁶ X. Chen, A. Tiwari, and S. Ryu, Physical Review B 94, 045113 (2016).
- ¹⁷ B. Han, A. Tiwari, C.-T. Hsieh, and S. Ryu, Physical Review B **96**, 125105 (2017).
- ¹⁸ G. Y. Cho, K. Shiozaki, S. Ryu, and A. W. Ludwig, Journal of Physics A: Mathematical and Theoretical **50**, 304002 (2017).
- ¹⁹ F. D. M. Haldane, Phys. Rev. Lett. **74**, 2090 (1995).
- ²⁰ E. Lieb, T. Schultz, and D. Mattis, Annals of Physics **16**, 407 (1961).
- ²¹ I. Affleck and E. H. Lieb, Lett. Math. Phys. **12**, 57 (1986).
- ²² Oshikawa and Masaki, Physical Review Letters 84, 1535 (2000).
- ²³ M. B. Hastings, Phys. Rev. B **69**, 104431 (2004).
- ²⁴ M. Cheng, M. Zaletel, M. Barkeshli, A. Vishwanath, and P. Bonderson, Phys. Rev. X 6, 041068 (2016).
- ²⁵ S. C. Furuya and M. Oshikawa, Phys. Rev. Lett. **118**, 021601 (2017).
- ²⁶ G. Y. Cho, C.-T. Hsieh, and S. Ryu, Phys. Rev. B **96**, 195105 (2017).
- ²⁸ Y. Tanizaki and T. Sulejmanpasic, Phys. Rev. B 98, 115126 (2018).
 ²⁹ V. Verse, G. T. Heich, and M. Ochikawa,
- ²⁹ Y. Yao, C.-T. Hsieh, and M. Oshikawa, Phys. Rev. Lett. **123**, 180201 (2019).
- ³⁰ T. Numasawa and S. Yamaguchi, Journal of High Energy Physics **2018**, 202 (2018).
- ³¹ H.-H. Tu and R. Orus, Physical review letters **107** 7, 077204 (2011).

- ³² C.-M. Jian, Z. Bi, and C. Xu, Phys. Rev. B **97**, 054412 (2018).
- ³³ J. Cardy, arXiv preprint hep-th/0411189 (2004).
- ³⁴ S. Kawai, *Boundary conformal field theory in free-field representation*, Ph.D. thesis, University of Oxford (2002).
- ³⁵ A. Recknagel and V. Schomerus, Boundary conformal field theory and the worldsheet approach to D-branes (Cambridge University Press, 2013).
- ³⁶ N. Ishibashi, Modern Physics Letters A **04**, 251 (1989).
- ³⁷ J. L. Cardy, Nuclear Physics B **324**, 581 (1989).
- ³⁸ J. Cardy, SciPost Phys. **3**, 011 (2017).
- ³⁹ M. Oshikawa, arXiv preprint arXiv:1007.3739 (2010).
- ⁴⁰ Y. Fuji, Phys. Rev. B **93**, 104425 (2016).
- ⁴¹ Y. Fuji, F. Pollmann, and M. Oshikawa, Phys. Rev. Lett. **114**, 177204 (2015).
- ⁴² L. H. Santos, Phys. Rev. B **91**, 155150 (2015).
- ⁴³ I. Affleck, arXiv preprint cond-mat/9802045 (1998).
- ⁴⁴ J.-H. Chen, C. Mudry, C. Chamon, and A. Tsvelik, Physical Review B **96**, 224420 (2017).
- ⁴⁵ H. J. Schulz, Phys. Rev. B **34**, 6372 (1986).
- ⁴⁶ P. Lecheminant, Nucl. Phys. B **901**, 510 (2015).
- ⁴⁷ Y. Yao and M. Oshikawa, arXiv preprint arXiv:2010.09244 (2020).
- ⁴⁸ P. Schmoll, A. Haller, M. Rizzi, and R. Orús, Physical Review B **99**, 205121 (2019).
- ⁴⁹ H. Watanabe, Phys. Rev. B **97**, 165117 (2018).
- ⁵⁰ R. Behrend, P. Pearce, V. Petkova, and J. Zuber, arXiv: High Energy Physics - Theory (1999).
- ⁵¹ P. Ruelle and O. Verhoeven, Nuclear Physics B **535**, 650 (1998),
- arXiv:hep-th/9803129 [hep-th].
- ⁵² C.-T. Hsieh, Y. Nakayama, and Y. Tachikawa, Phys. Rev. Lett. **126**, 195701 (2021).
- ⁵³ K. Kikuchi and Y. Zhou, "Two-dimensional anomaly, orbifolding, and boundary states," (2019), arXiv:1908.02918 [hep-th].
- ⁵⁴ J. A. Harvey and G. W. Moore, Journal of High Energy Physics **2018**, 145 (2018).
- ⁵⁵ C. Heinrich and M. Levin, Physical Review B 98, 035101 (2018).
- ⁵⁶ P. Lecheminant, A. O. Gogolin, and A. A. Nersesyan, Nuclear Physics B **639**, 502 (2002).
- ⁵⁷ J. Kulp, JHEP **03**, 124 (2021), arXiv:2003.04278 [hep-th].
- ⁵⁸ A. Karch, D. Tong, and C. Turner, SciPost Phys 7 (2019).
- ⁵⁹ Y. Yao and Y. Fukusumi, Phys. Rev. B **100** (2019).
- ⁶⁰ Y. Yao and A. Furusaki, JHEP **2021**, 285 (2021).
- ⁶¹ E. Fradkin and L. P. Kadanoff, Nucl. Phys. B **170**, 1 (1980).
- ⁶² Y. Fukusumi, Y. Tachikawa, and Y. Zheng, arXiv preprint arXiv:2103.00746 (2021).
- ⁶³ H. Ebisu and M. Watanabe, arXiv preprint arXiv:2103.01101 (2021).
- ⁶⁴ J. Cardy, SciPost Phys **3**, 1706 (2017).

Appendix A: SU(N) symmetry invariant boundary states of free boson model

In this appendix, we will use the affine Lie algebra to show the boundary states (57)(66) are invariant under vector SU(N) spin rotation.

Firstly, we prove the result for the case of SU(2). For simplify, the boundary state is written in the language of chiral operators:

$$\begin{split} |B\rangle &= \frac{e^{\sum_{r\in\mathbb{N}_{+}}\frac{-1}{r}\left(a_{-r,L}^{1}a_{-r,R}^{1}+a_{-r,L}^{2}a_{-r,R}^{2}\right)}}{\mathcal{N}_{N}\mathcal{N}_{D}}\sum_{m,n}\left(e^{in\frac{\sqrt{2}}{2}\left(\theta_{1}^{0}-\theta_{2}^{0}\right)}-e^{im\frac{\sqrt{2}}{2}\left(\phi_{1}^{0}+\phi_{2}^{0}\right)}\right)e^{-\frac{1}{r}a_{-r,L}^{1}a_{-r,R}^{1}a_{-r,R}^{1}a_{-r,L}^{2}a_{-r,R}^{2}}|0\rangle\\ &= \frac{e^{\sum_{r\in\mathbb{N}_{+}}\frac{-1}{r}\left(a_{-r,L}^{1}a_{-r,R}^{1}+a_{-r,L}^{2}a_{-r,R}^{2}\right)}}{\mathcal{N}_{N}\mathcal{N}_{D}}\sum_{m,n}\left(e^{in\frac{\sqrt{2}}{2}\left(\varphi_{1,L}^{0}-\varphi_{1,R}^{0}-\varphi_{2,L}^{0}+\varphi_{2,R}^{0}\right)}-e^{im\frac{\sqrt{2}}{2}\left(\varphi_{1,L}^{0}+\varphi_{1,R}^{0}+\varphi_{2,L}^{0}+\varphi_{2,R}^{0}\right)}\right)|0\rangle. \end{split}$$
(A1)

where $\varphi_L^0 / \varphi_R^0$ is zero mode of primary fields $\varphi_L(\omega = 0) / \varphi_R(\bar{\omega} = 0)$. The affine SU(2) generators are given by:

$$H = i\partial\varphi_L, \quad E^{\pm} = e^{\pm i\sqrt{2}\varphi_L}, \\ \bar{H} = -i\bar{\partial}\varphi_R, \quad \bar{E}^{\pm} = e^{\pm i\sqrt{2}\varphi_R}.$$
(A2)

The OPE between affine SU(2) generators and primary fields are:

$$H(z)e^{\pm i\frac{1}{\sqrt{2}}\varphi_{L}^{0}} = \frac{\pm e^{\pm\frac{1}{\sqrt{2}}\varphi_{L}^{0}}}{\sqrt{2}z}, \quad E^{\pm}(z)e^{\mp i\frac{1}{\sqrt{2}}\varphi_{L}^{0}} = \frac{e^{\pm i\frac{1}{\sqrt{2}}\varphi_{L}^{0}}}{z},$$
$$\bar{H}(\bar{z})e^{\pm i\frac{1}{\sqrt{2}}\varphi_{R}^{0}} = \frac{\mp e^{\pm\frac{1}{\sqrt{2}}\varphi_{R}^{0}}}{\sqrt{2}\bar{z}}, \quad \bar{E}^{\pm}(\bar{z})e^{\pm i\frac{1}{\sqrt{2}}\varphi_{R}^{0}} = \frac{e^{\mp i\frac{1}{\sqrt{2}}\varphi_{R}^{0}}}{\bar{z}}.$$
(A3)

Since the generators of SU(2) rotation are zero modes of affine SU(2) generators, they satisfy the following OPE with primary fields:

$$H^{0}e^{\pm i\frac{1}{\sqrt{2}}\varphi_{L}^{0}} = \pm e^{\pm\frac{1}{\sqrt{2}}\varphi_{L}^{0}}, \quad E^{0,\pm}e^{\mp i\frac{1}{\sqrt{2}}\varphi_{L}^{0}} = e^{\pm i\frac{1}{\sqrt{2}}\varphi_{L}^{0}},$$

$$\bar{H}^{0}e^{\pm i\frac{1}{\sqrt{2}}\varphi_{R}^{0}} = \mp e^{\pm\frac{1}{\sqrt{2}}\varphi_{R}^{0}}, \quad \bar{E}^{0,\pm}e^{\pm i\frac{1}{\sqrt{2}}\varphi_{R}^{0}} = e^{\mp i\frac{1}{\sqrt{2}}\varphi_{R}^{0}}.$$
 (A4)

Since the other OPEs are zero, the SU(2) rotation only acts on the following parts of the boundary state:

$$\begin{split} & \left[e^{i\frac{\sqrt{2}}{2}(\varphi_{1,L}^{0}-\varphi_{1,R}^{0}-\varphi_{2,L}^{0}+\varphi_{2,R}^{0})}-e^{i\frac{\sqrt{2}}{2}(\varphi_{1,L}^{0}+\varphi_{1,R}^{0}+\varphi_{2,L}^{0}+\varphi_{2,R}^{0})}\right.\\ & \left.+e^{-i\frac{\sqrt{2}}{2}(\varphi_{1,L}^{0}-\varphi_{1,R}^{0}-\varphi_{2,L}^{0}+\varphi_{2,R}^{0})}-e^{-i\frac{\sqrt{2}}{2}(\varphi_{1,L}^{0}+\varphi_{1,R}^{0}+\varphi_{2,L}^{0}+\varphi_{2,R}^{0})}\right]\left|0\right\rangle. \end{split}$$
(A5)

For the vector SU(2) rotation, the generators is $S^z = H^0 + \overline{H}^0$, $S^{\pm} = E^{0,\pm} + \overline{E}^{0,\pm}$. We can see the right-hand boson is the conjugated representation of affine SU(2) Lie algebra. For simplify, we can do the charge conjugation on the right hand bosons since there is an extra phase -1 for antiholomorphic sector in Eq. (A3):

$$\begin{aligned} \varphi_{i,R}' &= -\varphi_{i,R} ,\\ \left[e^{i\frac{\sqrt{2}}{2}} (\varphi_{1,L}^{0} + \varphi_{1,R}^{\prime 0} - \varphi_{2,L}^{0} - \varphi_{2,R}^{\prime 0}) - e^{i\frac{\sqrt{2}}{2}} (\varphi_{1,L}^{0} - \varphi_{1,R}^{\prime 0} + \varphi_{2,L}^{0} - \varphi_{2,R}^{\prime 0}) \right. \\ \left. + e^{i\frac{\sqrt{2}}{2}} (-\varphi_{1,L}^{0} - \varphi_{1,R}^{\prime 0} + \varphi_{2,L}^{\prime 0} + \varphi_{2,R}^{\prime 0}) - e^{i\frac{\sqrt{2}}{2}} (-\varphi_{1,L}^{0} + \varphi_{1,R}^{\prime 0} - \varphi_{2,L}^{\prime 0} + \varphi_{2,R}^{\prime 0})} \right] |0\rangle. \end{aligned}$$
(A6)

In this representation, the SU(2) generators are the sum of four spin-1/2 generators. The boundary state can be written in the spin language since each primary fields transform like a spinor:

$$|\uparrow\uparrow\downarrow\downarrow\rangle\rangle - |\uparrow\downarrow\uparrow\downarrow\rangle\rangle + |\downarrow\downarrow\uparrow\uparrow\rangle\rangle - |\downarrow\uparrow\downarrow\uparrow\rangle\rangle = (|\uparrow\downarrow\rangle\rangle - |\downarrow\uparrow\rangle\rangle) \otimes (|\uparrow\downarrow\rangle\rangle - |\downarrow\uparrow\rangle\rangle).$$
 (A7)

Here in the second arrow, the first two spins are $\exp(\pm i\varphi_{1,L}^0/\sqrt{2})|0\rangle$ and $\exp(\mp i\varphi_{2,R}^{\prime 0}/\sqrt{2})|0\rangle$ which are first and fourth spins in the first arrow. The other two spins are $\exp(\pm i\varphi_{1,R}^{\prime 0}/\sqrt{2})|0\rangle$ and $\exp(\mp i\varphi_{2,L}^0/\sqrt{2})|0\rangle$. So it is easy to see the boundary state is invariant under vector SU(2) transformation since it is direct product of two spin singlet states.

For general vector SU(N) transformation, the Lie algebra can be composed into N(N-1)/2 SU(2) Lie subalgebras. For each SU(2) Lie subalgebras, we can do similar calculations to show the eigenvalue of SU(2) generators is zero. So the boundary state is invariant under vector SU(N) spin rotation.

Appendix B: Invariant boundary state of vector rotation symmetry and center symmetry

In this appendix, we will give a detailed calculation on invariant boundary states of vector rotation symmetry and center symmetry in the WZW model.

The Ishibashi state is defined as follows:

$$|\lambda\rangle\rangle = \sum_{m} |\phi_i^{\lambda}, m\rangle \otimes U |\bar{\phi}_i^{\lambda}, \bar{m}\rangle.$$
(B1)

To show the Ishibashi state is invariant under the vector rotation symmetry, we only need to consider the action of generators on Cardy state is zero. Since the holomorphic and antiholomorphic sector is tensor product, the action of a generator is a summation of action on each part : $S^a = J_0^a + \bar{J}_0^a$. The vacuum states of each sector form different irreducible representations of Lie algebra labeling by λ :

$$J_0^a |\phi_i^{\lambda}, m\rangle = \sum_j T_{ij}^{\lambda, a} |\phi_j^{\lambda}, m\rangle,$$

$$\bar{J}_0^a |\bar{\phi}_i^{\lambda}, m\rangle = \sum_j T_{ij}^{\lambda, a} |\bar{\phi}_j^{\lambda}, m\rangle.$$
 (B2)

Since U operator is an anti-unitary operator, it will anti-commute with Lie algebra:

$$U^{-1}\bar{J}_n^a U = -(\bar{J}_{-n}^a)^+ = -(\bar{J}_n^a).$$
(B3)

Now we can show the action of generators on each vacuum in the summation is zero:

$$\sum_{m} \langle \phi_{j}^{\lambda}, n | \otimes \langle U \bar{\phi_{k}}^{\lambda}, n' | S^{a} | \phi_{i}^{\lambda}, m \rangle \otimes U | \bar{\phi_{i}}^{\lambda}, m \rangle$$

$$= \sum_{m} \delta_{ik} \delta_{n'm} \langle \phi_{j}^{\lambda}, n | J_{0}^{a} | \phi_{i}^{\lambda}, m \rangle + \delta_{ij} \delta_{nm} \langle U \bar{\phi}_{k}^{\lambda}, n' | \bar{J}_{0}^{a} U | \bar{\phi}_{i}^{\lambda}, m \rangle$$

$$= \delta_{ik} \delta_{ij} (\langle \phi_{i}^{\lambda}, n | J_{0}^{a} | \phi_{i}^{\lambda}, n' \rangle + \langle U \bar{\phi}_{i}^{\lambda}, n' | U^{+} \bar{J}_{0}^{a} U | \bar{\phi}_{i}^{\lambda}, n \rangle)$$

$$= \delta_{ik} \delta_{jk} (\langle \phi_{i}^{\lambda}, n | J_{0}^{a} | \phi_{i}^{\lambda}, n' \rangle - \langle \bar{\phi}_{i}^{\lambda}, n' | \bar{J}_{0}^{a} | \bar{\phi}_{i}^{\lambda}, n \rangle^{*})$$

$$= 0. \qquad (B4)$$

Thus the Cardy state is also invariant under vector rotation symmetry since it is linear summation of Ishibashi states. The condition for invariant Cardy states is now:

$$h|B\rangle = |B\rangle. \tag{B5}$$

1. B_N affine Lie algebra

For B_N type Lie algebra, the action of the outer automorphism is

$$A: [\lambda_0; \lambda_1, \cdots, \lambda_{N-1}, \lambda_N] \to [\lambda_1; \lambda_0, \cdots, \lambda_{N-1}, \lambda_N] .$$
(B6)

Thus the affine Dynkin labels of an invariant boundary state satisfy

$$\lambda_0 = \lambda_1. \tag{B7}$$

Since the comarks of B_N are:

$$(a_0^{\vee}; a_1^{\vee}, \cdots, a_N^{\vee}) = (1; 1, 2, \cdots, 2, 2, 1).$$
 (B8)

We can obtain the level of \boldsymbol{k}

$$k = \sum_{i=0}^{N} a_i^{\vee} \lambda_i = 2(\lambda_1 + \dots + \lambda_{N-1}) + \lambda_N.$$
(B9)

Thus we can always find a Cardy state invariant under the vector SO(2l+1) symmetry and center symmetry.

2. D_{2l} affine Lie algebra

For D_{2l} type Lie algebra, there are two outer automorphisms:

$$A_{1}: [\lambda_{0}; \lambda_{1}, \lambda_{2}, \cdots, \lambda_{2l-1}, \lambda_{2l}] \rightarrow [\lambda_{1}; \lambda_{0}, \lambda_{2}, \cdots, \lambda_{2l}, \lambda_{2l-1}], A_{2}: [\lambda_{0}; \lambda_{1}, \lambda_{2}, \cdots, \lambda_{2l-2}, \lambda_{2l-1}, \lambda_{2l}] \rightarrow [\lambda_{2l}; \lambda_{2l-1}, \lambda_{2l-2}, \cdots, \lambda_{2}, \lambda_{1}, \lambda_{0}].$$
(B10)

So the affine Dynkin labels of an invariant boundary state satisfy

$$\lambda_1 = \lambda_0 = \lambda_{2l} = \lambda_{2l-1},$$

$$\lambda_j = \lambda_{2l-j} \ (j \ge 2).$$
(B11)

The comarks of D_N are:

$$(a_0^{\vee}; a_1^{\vee}, \cdots, a_N^{\vee}) = (1; 1, 2, \cdots, 2, 1, 1).$$
 (B12)

As a result, for the existing of an invariant boundary state, k satisfies

$$k = 4\lambda_0 + 4(\lambda_2 \dots + \lambda_{l-1}) + 2\lambda_l. \tag{B13}$$

Therefore, there exists an invariant Cardy state under vector SO(4l) symmetry and center symmetry only for even k.

3. C_N affine Lie algebra

For C_N type Lie algebra, the action of the outer automorphism is

$$4: [\lambda_0; \lambda_1, \cdots, \lambda_{N-1}, \lambda_N] \to [\lambda_N; \lambda_{N-1}, \cdots, \lambda_1, \lambda_0].$$
(B14)

Thus the affine Dynkin labels of an invariant boundary state satisfy

$$\lambda_j = \lambda_{N-j}.\tag{B15}$$

Since the comarks of C_N are:

$$(a_0^{\vee}; a_1^{\vee}, \cdots, a_N^{\vee}) = (1; 1, \cdots, 1).$$
 (B16)

We can obtain the level of \boldsymbol{k}

$$k = \begin{cases} 2(\lambda_0 + \dots + \lambda_{\frac{N}{2}-1}) + \lambda_{\frac{N}{2}}, & \text{if } N \text{ is even;} \\ 2(\lambda_0 + \dots + \lambda_{\frac{N-1}{2}}), & \text{if } N \text{ is odd.} \end{cases}$$
(B17)

Therefore only k is even or N is even, there will be a Cardy state invariant under the vector Usp(N) symmetry and center symmetry.

4. D_{2l+1} affine Lie algebra

For D_{2l+1} type Lie algebra, the action of the outer automorphism is

$$A: [\lambda_0; \lambda_1, \cdots, \lambda_{2l}, \lambda_{2l+1}] \to [\lambda_{2l}; \lambda_{2l+1}, \cdots, \lambda_1, \lambda_0].$$
(B18)

The affine Dynkin labels of an invariant boundary state satisfy

$$\lambda_0 = \lambda_{2l} = \lambda_1 = \lambda_{2l+1}, \lambda_j = \lambda_{2l+1-j} \quad (2 \le j \le 2l-1).$$
(B19)

Since the comarks of D_N are:

$$(a_0^{\vee}; a_1^{\vee}, \cdots, a_N^{\vee}) = (1; 1, 2, \cdots, 2, 1, 1).$$
(B20)

We can obtain the level of \boldsymbol{k}

$$k = 4(\lambda_0 + \lambda_2 + \dots + \lambda_l). \tag{B21}$$

Therefore only k is a multiple of 4, there will be a Cardy state invariant under the vector SO(4l+2) symmetry and center symmetry.

5. E_6 affine Lie algebra

For E_6 type Lie algebra, the action of the outer automorphism is

$$A: [\lambda_0; \lambda_1, \cdots, \lambda_6] \to [\lambda_1; \lambda_5, \lambda_4, \lambda_3, \lambda_6, \lambda_0, \lambda_2].$$
(B22)

The affine Dynkin labels of an invariant boundary state satisfy

$$\lambda_0 = \lambda_5 = \lambda_1, \quad \lambda_2 = \lambda_4 = \lambda_6. \tag{B23}$$

Since the comarks of E_6 are:

$$(a_0^{\vee}; a_1^{\vee}, \cdots, a_6^{\vee}) = (1; 1, 2, 3, 2, 1, 2).$$
 (B24)

Thus the level k is given by

$$k = \lambda_0 + \lambda_1 + 2\lambda_2 + 3\lambda_3 + 2\lambda_4 + \lambda_5 + 2\lambda_6 = 3(\lambda_0 + \lambda_3 + 2\lambda_2).$$
(B25)

This implies only if k is multiple of 3, there will be a Cardy state invariant under the vector E_6 symmetry and center symmetry.

6. E_7 affine Lie algebra

For
$$E_7$$
 type Lie algebra, the action of the outer automorphism is

$$A: [\lambda_0; \lambda_1, \cdots, \lambda_7] \to [\lambda_6; \lambda_5, \lambda_4, \lambda_3, \lambda_2, \lambda_1, \lambda_0, \lambda_7].$$
(B26)

The affine Dynkin labels of an invariant boundary state satisfy

$$\lambda_0 = \lambda_6, \quad \lambda_1 = \lambda_5, \quad \lambda_2 = \lambda_4 = \lambda_4. \tag{B27}$$

Since the comarks of E_7 are:

$$(a_0^{\vee}; a_1^{\vee}, \cdots, a_7^{\vee}) = (1; 2, 3, 4, 3, 2, 1, 2),$$
(B28)

the level k is given by

$$k = 2\lambda_0 + 4\lambda_1 + 6\lambda_2 + 4\lambda_3 + 2\lambda_7. \tag{B29}$$

This implies only if k is even, there will be a Cardy state invariant under the vector E_7 symmetry and center symmetry.

Appendix C: Invariant boundary state of vector rotation symmetry and CP_2 symmetry

In this appendix, we will give a detailed derivation on the invariant boundary state of vector rotation symmetry and $C\mathcal{P}_2$ symmetry in the WZW model. As shown in the appendix B, the Cardy states have been invariant under vector rotation and spatial reflection symmetry, we only need to consider $\mathbb{Z}_2^{Ch_2}$ symmetry where h_2 is a order 2 element of center symmetry. Since the charge conjugation acts as an identity on Cardy states for B_N , C_N , D_{2l} , E_7 type Lie group and center symmetry is a \mathbb{Z}_2 group, the result is the same as that of appendix B. Moreover, the center symmetry of E_6 WZW model is \mathbb{Z}_3 group which implies we can't construct \mathcal{T}_2 symmetry. Thus we only need to calculate the D_{2l+1} WZW model where the order 2 element is A^2 .

For D_{2l+1} type Lie algebra, the action of the $\mathbb{Z}_2^{Ch_2}$ symmetry is

$$CA^{2}: [\lambda_{0}; \lambda_{1}, \cdots, \lambda_{2l}, \lambda_{2l+1}] \to [\lambda_{1}; \lambda_{0}, \cdots, \lambda_{2l}, \lambda_{2l+1}].$$
(C1)

The affine Dynkin labels of an invariant boundary state satisfy

$$\lambda_0 = \lambda_1. \tag{C2}$$

Then when there is an invariant boundary state, k satisfies:

$$k = 2\lambda_0 + 2(\lambda_2 + \dots + \lambda_{2l-1}) + \lambda_{2l} + \lambda_{2l+1}.$$
(C3)

Therefore, there always exists an invariant boundary state for arbitrary k.

Appendix D: Symmetric interaction of $SU(2)_{2k}$ WZW model

The calculation in this appendix is based on an $\operatorname{argument}^{64}$ for a perturbed CFT with the hamiltonian:

$$H = H_{CFT} + \sum_{j} \lambda_j \int \Phi^j(x) dx.$$
 (D1)

Here Φ^j are relevant operators (primiary fields). It claims that the smeared boundary state $e^{-\tau_a H}|B_a\rangle$ with the lowest variational energy can be very close to the ground state. This variational energy is given by:

$$E_a = \frac{\pi c}{96\tau_a^2} + \sum_{j\neq 0} \lambda_j \frac{S_a^j}{S_a^0} \left(\frac{S_0^0}{S_0^j}\right)^{\frac{1}{2}} \frac{\pi^{\Delta_j}}{(2\tau_a)^{\Delta_j}}.$$
 (D2)

For the SU(2)_{2k} WZW model, the relevant operator $\lambda \operatorname{tr}(g^2)$ is the primary field $\lambda \Phi^1$. We have 2k+1 boundary states $|B_a\rangle$ for $2a \in \mathbb{Z}$ and $0 \le a \le k$. Thus the variational energy of the 'smeared' boundary state $e^{-\tau_a H}|B_a\rangle$ is:

$$E_a = \frac{\pi c}{96\tau_a^2} + \lambda \frac{S_a^1}{S_a^0} \left(\frac{S_0^0}{S_0^1}\right)^{\frac{1}{2}} \left(\frac{\pi}{2\tau_a}\right)^{\frac{2}{k+1}}.$$
 (D3)

When $\lambda > 0$, we can vary τ_a and obtain the minimal of E_a :

$$min(E_a) = \begin{cases} 0, & \text{if } \frac{S_a^1}{S_a^0} \ge 0; \\ -\lambda b \left| \frac{S_a^1}{S_a^0} \right|^{\frac{k+1}{k}}, & \text{if } \frac{S_a^1}{S_a^0} < 0. \end{cases}$$
(D4)

where b is a positive number only depending on k.

The modular S matrix of $SU(2)_{2k}$ WZW model are given by:

$$S_j^{j'} = \sqrt{\frac{1}{k+1}} \sin \frac{\pi (2j+1)(2j'+1)}{2k+2}.$$
 (D5)

Thus the minimal of E_a can be rewritten as:

$$min(E_a) = \begin{cases} 0, & \text{if } j \le \frac{2k-1}{6} \text{ or } j \ge \frac{4k+1}{6}; \\ -\lambda b \left| 4\cos^2(\frac{2j+1}{2k+2}\pi) - 1 \right|^{\frac{k+1}{k}}, & \text{if } \frac{2k-1}{6} \le j \le \frac{4k+1}{6}. \end{cases}$$
(D6)

It is easy to see when a = k/2, the minimal of variational energy is lowest. In the language of Dykin label, this boundary state is [k; k] which is the only invariant state under the center symmetry.

When $\lambda < 0$, the similar calculation can show when a = 0 or a = k, the variational energy is lowest. These two states correspond to the Valence Bond Solid (VBS) phase which breaks translation symmetry.

We can also show the massive phase with $\lambda > 0$ is Haldane phase if k is odd and is trivial phase if k is even. To see this, we consider such massive deformation:

$$V(x) = \begin{cases} \lambda \operatorname{tr}(g^2), & x < 0; \\ 0, & 0 \le x < L; \\ -\lambda \operatorname{tr}(g^2), & L \le x. \end{cases}$$

The partition function of the middle gapless region is given by

$$Z_{\frac{k}{2}0} = Z_{\frac{k}{2}k} = \chi_{\frac{k}{2}}(q) = (k+1)q^{\frac{k}{2}} + \cdots$$
(D7)

Here \cdots is polynomials of q whose degree is large than $\triangle_{k/2}$.

Thus when L goes to zero, there will be k + 1 states on the interface which are equivalent to states of spin k/2. When k is even, the spin on the interface is an integer which can be gapped without degeneracy. When k is odd, the spin is a half integer which is always degenerate. We show the lattice version of the interface for $SU(2)_2$ WZW in the figure 3. Here dots are the effective spin 1/2 decoupled from the spin 1 on each site. Thus the unpaired spin 1/2 on the interface implies the symmetry protected degeneracy.



FIG. 3. The lattice version of interface between Haldane phase and VBS phase of $SU(2)_2$ WZW model.

Appendix E: The LSM ingappability of the spin chain with triple-product interactions

In this appendix, we will prove the LSM ingappability of the Hamiltonian (99) with the following symmetry:

$$\mathcal{T}_2: T_2 = TKU_T = TK \prod_{j=1}^{j=L} i\sigma_j^y, \tag{E1}$$

$$U(1)_z : U_\theta = \exp(i\theta \sum_{j=1}^{L} \mathbf{S}_j^z), \tag{E2}$$

$$\mathbb{Z}_2^y : R_y^\pi = \prod_{j=1}^L i\sigma_j^y.$$
(E3)

Similar to the original LSM theorem, the twisting operator U_{twist} of $U(1)_z$ is given by:

$$U_{\text{twist}} = \exp(\frac{2\pi i}{L} \sum_{j=1}^{L} j \mathbf{S}_j^z) = \exp(\frac{\pi i}{L} \sum_{j=1}^{L} j \sigma_j^z).$$
(E4)

Let $|G.S.\rangle$ is a ground state of the Hamiltonian (99) with length L (periodic boundary condition). To show the $U_{twist}|G.S.\rangle$ has low energy, we consider the term $H_{j,j+1,j+2}$ in Hamiltonian (99) involving the three neighbouring sites, j, j + 1 and j + 2. Since $[\sigma_j^z + \sigma_{j+1}^z + \sigma_{j+2}^z, H_{j,j+1,j+2}] = 0$, and others commute with $H_{j,j+1,j+2}$,

$$U_{\text{twist}}^{+}(H_{j,j+1,j+2})U_{\text{twist}}$$

$$= e^{(-\pi i/L)(\sigma_{j}^{z}+2\sigma_{j+1}^{z}+3\sigma_{j+2}^{z})}H_{j,j+1,j+2}e^{(\pi i/L)(\sigma_{j}^{z}+2\sigma_{j+1}^{z}+3\sigma_{j+2}^{z})}$$

$$= H_{j,j+1,j+2} - \frac{\pi i}{L}[\sigma_{j}^{z}+2\sigma_{j+1}^{z}+3\sigma_{j+2}^{z},H_{j,j+1,j+2}] + O(\frac{1}{L^{2}})$$

$$= H_{j,j+1,j+2} - [\frac{\pi i}{L}\sum_{j=1}^{L}j\sigma_{j}^{z},H_{j,j+1,j+2}] + O(\frac{1}{L^{2}}).$$
(E5)

We can sum over all terms in (E5):

$$\langle \mathbf{G.S.} | U_{\mathbf{twist}}^{+} H U_{\mathbf{twist}} | \mathbf{G.S.} \rangle = E_{\mathrm{gs}} + O(\frac{1}{L}).$$
 (E6)

The reminder $O(\frac{1}{L})$ comes from third term in (E5) and \mathcal{T}_2 symmetry ensures contributions from $H_{j,j+1,j+2}$ doesn't depend on site. Therefore we have $U_{\text{twist}}|\text{G.S.}\rangle = e^{i\alpha}|\text{G.S.}\rangle + O(\frac{1}{L})|\psi\rangle$.

Now, let's assume there is an unique ground state which possess all symmetries. Therefore it is invariant under the combination of \mathcal{T}_2 and \mathbb{Z}_2^y :

$$T_2 R_q^{\pi} | \text{G.S.} \rangle = T K | \text{G.S.} \rangle = | \text{G.S.} \rangle$$
 (E7)

Under periodic boundary condition, we can obtain

$$(TK)^{-1}U_{\text{twist}}TK = K^{-1}U_{\text{twist}}\exp\left(\frac{\pi i}{L}\sum_{j=1}^{L}\sigma_{j}^{z}\right)\exp(i\pi\sigma_{1}^{z})K$$
$$= -U_{\text{twist}}^{+}\exp\left(-\frac{\pi i}{L}\sum_{j=1}^{L}\sigma_{j}^{z}\right).$$
(E8)

then

$$(TK)^{-1}U_{\mathbf{twist}}TK|\mathbf{G.S.}\rangle = -U_{\mathbf{twist}}^{+}|\mathbf{G.S.}\rangle = -e^{-i\alpha}|\mathbf{G.S.}\rangle - O(\frac{1}{L})|\psi'\rangle.$$
(E9)

The first equation comes from the fact $R_y^{\pi}U_{\theta} = U_{\theta}^+ R_y^{\pi}$ and $U_0 = \mathbb{I}$.

On the other hand, we can calculate this equation directly:

$$(TK)^{-1}U_{\mathbf{twist}}TK|\mathbf{G.S.}\rangle = (TK)^{-1}(e^{i\alpha}|\mathbf{G.S.}\rangle + O(\frac{1}{L})|\psi\rangle)$$
$$= e^{-i\alpha}|\mathbf{G.S.}\rangle + O(\frac{1}{L})TK|\psi\rangle.$$
(E10)

When L goes to infinity, there will be a conflict between (E9) and (E10). Therefore, the ground state can not be unique.

Appendix F: Vector rotation on the y-direction

In this appendix, we will use the affine SU(2) Lie algebra to show how the primary terms transform after vector SU(2) rotation. For U(1) boson theory with the self-dual radius, the primary terms are just $\exp(\pm i\theta/\sqrt{2})$ and $\exp(\pm i\phi/\sqrt{2})$. For simplify, we also written them in the chiral language: $\exp(\pm i(\varphi_L + \varphi_R)/\sqrt{2})$ and $\exp(\pm i(\varphi_L - \varphi_R)/\sqrt{2})$.

The zero modes of affine SU(2) algebra can be rewritten as $J_a^0 = \oint dz J_a(z)$. The commutation relationship between zero modes and other fields are related to the OPEs as follows:

$$[J_a^0, b(w)] = \oint_w dz J_a(z) b(w).$$
(F1)

Therefore the commutation relationship between zero modes of affine SU(2) algebra and chiral primary field can be calculated using OPEs and the result is given by:

$$[H^{0}, e^{\pm i\frac{\sqrt{2}}{2}\varphi_{L}}] = \pm e^{\pm i\frac{\sqrt{2}}{2}\varphi_{L}}, \quad [E^{0,\pm}, e^{\mp i\frac{\sqrt{2}}{2}\varphi_{L}}] = 2ie^{\pm i\frac{\sqrt{2}}{2}\varphi_{L}}, \quad [E^{0,\pm}, e^{\pm i\frac{\sqrt{2}}{2}\varphi_{L}}] = 0,$$

$$[\bar{H}^{0}, e^{\pm i\frac{\sqrt{2}}{2}\varphi_{R}}] = \mp e^{\pm i\frac{\sqrt{2}}{2}\varphi_{R}}, \quad [\bar{E}^{0,\pm}, e^{\pm i\frac{\sqrt{2}}{2}\varphi_{R}}] = 2ie^{\mp i\frac{\sqrt{2}}{2}\varphi_{R}}, \quad [\bar{E}^{0,\pm}, e^{\mp i\frac{\sqrt{2}}{2}\varphi_{R}}] = 0.$$

$$(F2)$$

Now we can perform the vector SU(2) rotation along y-axis whose generator is given by:

$$S^{y} = \frac{i}{2} (E^{0,+} - E^{0,-} + \bar{E}^{0,+} - \bar{E}^{0,-}).$$
(F3)

For $\pi/2$ rotation, the transformation can be written in the language of the generator:

$$U(\frac{\pi}{2}) = \frac{\sqrt{2}}{2}(1+iS^y).$$
 (F4)

For simplify, we consider the real part of primary fields: $\cos(\phi/\sqrt{2}), \sin(\phi/\sqrt{2}), \cos(\theta/\sqrt{2}), \sin(\theta/\sqrt{2})$. There are two primary fields invariant under rotation since they commute with S^y :

$$U^{-1}\cos(\frac{\sqrt{2}}{2}\phi)U = \cos(\frac{\sqrt{2}}{2}\phi), U^{-1}\sin(\frac{\sqrt{2}}{2}\theta)U = \sin(\frac{\sqrt{2}}{2}\theta).$$
(F5)

The other two primary fields satisfy the following commutation relationship:

$$[S^{y}, \cos(\frac{\sqrt{2}}{2}\theta)] = 2i\sin(\frac{\sqrt{2}}{2}\phi) , [S^{y}, \sin(\frac{\sqrt{2}}{2}\phi)] = -2i\cos(\frac{\sqrt{2}}{2}\theta),$$
(F6)

thus they transform under the $\pi/2$ rotation as follows:

$$U^{-1}\sin(\frac{\sqrt{2}}{2}\phi)U = \cos(\frac{\sqrt{2}}{2}\theta), U^{-1}\cos(\frac{\sqrt{2}}{2}\theta)U = -\sin(\frac{\sqrt{2}}{2}\phi).$$
 (F7)