Capacity Bounds for the DeepONet Method of Solving Differential Equations

Pulkit Gopalani¹, Sayar Karmakar², and Anirbit Mukherjee³

¹Department of Electrical Engineering, IIT Kanpur, India ²Department of Statistics, University of Florida, US ³Department of Computer Science, The University of Manchester, UK

May 24, 2022

Abstract

In recent times machine learning methods have made significant advances in becoming a useful tool for analyzing physical systems. A particularly active area in this theme has been "physics informed machine learning" [15] which focuses on using neural nets for numerically solving differential equations. Among all the proposals for solving differential equations using deep-learning, in this paper we aim to advance the theory of generalization error for DeepONets - which is unique among all the available ideas because of its particularly intriguing structure of having an inner-product of two neural nets.

Our key contribution is to give a bound on the Rademacher complexity for a large class of DeepONets. Our bound does not explicitly scale with the number of parameters of the nets involved and is thus a step towards explaining the efficacy of overparameterized DeepONets. Additionally, a capacity bound such as ours suggests a novel regularizer on the neural net weights that can help in training DeepONets – irrespective of the differential equation being solved.

Keywords: Rademacher complexity, DeepONets, Physics-Inspired ML, Operator Learning

1 Introduction

In recent times there have been multiple methodological advances about using neural nets for solving differential equations, the deep-BSDE method [9] [13], Deep Ritz Method [35], Deep Galerkin Method [31], PINNs (Physics-Informed Neural Nets) [30], Neural Operators [16], DeepONets [20] and many more. In this paper we focus on DeepONets whose foundations were laid in [5], where the authors had pointed out how outputs of pairs of neural nets can be multiplied to approximate 'nice' operators between function spaces. But it was not until recently when "Deep Operator Nets" (DeepONet) [20] were introduced, that the full potential of this idea was realized.

It is increasingly becoming evident that DeepONets constitute a particularly powerful technique for solving differential equations – more so when one needs a "one shot" solution for a family of differential equations, with a fixed differential operator and different "source / forcing" functions. For example, consider solving a pendulum O.D.E., $\mathbb{R}^2 \ni \frac{\mathrm{d}(y,v)}{\mathrm{d}t} = (v,-k\cdot\sin(y)+f(t)) \in \mathbb{R}^2$ for different forcing functions f(t). Then the DeepONet setup can be trained only once, and then repeatedly be used for inference on new forcing functions to estimate their corresponding solutions. This approach is fundamentally unlike traditional numerical methods where one needs to rerun the optimization algorithm afresh for every new source function. In a recent study [21], the authors showed how the DeepONet setup has significant advantages over Fourier Neural Operators (FNO, [19]) in solving various differential equations of popular industrial use.

Over the past few years, many novel generalization bounds for neural nets have been established. Many of these works have computed bounds on Rademacher complexity for various classes of nets, to show how

different norm combinations of the involved weight matrices affect generalization performance, [10], [4], [27]. For shallow neural nets such methods have also been useful in explaining the benefits of dropout [1] and overparameterization [29]. However, for state-of-the-art neural nets, other approaches [8], [2], [28], [25] have given tighter generalization bounds. A thorough discussion of the reach and limitations of uniform convergence based bounds can be found in [26].

In this work we take some of the first steps towards putting the training procedure of DeepONets into the framework of statistical learning theory. Most importantly we compute their Rademacher complexity and give the first bounds on their generalization error which does not explicitly scale with the number of parameters – and this in turn identifies novel complexity measures for this architecture.

1.1 Overview of Training DeepONets & Our Main Result (Theorem 15)

We refer to the schematic in Fig.1 below for the DeepONet architecture and its constituent matrices,




Figure 1: A Sketch of the DeepONet Architecture

In the above diagram the Branch Network and the Trunk Network are neural nets with a common output dimension. $\mathbf{x}_B(f) \in \mathbb{R}^{d_1}$, the input to the branch net is an "encoded" version of a function f i.e a discretization of f onto a d_1 -sized grid of "sensor points" in its input domain. $\mathbf{x}_T \in \mathbb{R}^{d_2}$ is the trunk input. If the activation is σ at all layers and the branch net and the trunk net's layers are named $\mathbf{B}_k, k = 1, 2, \ldots, q_B$ and $\mathbf{T}_k, k = 1, 2, \ldots, q_T$ respectively, then the above architecture implements the map,

$$\mathbb{R}^{d_1} \times \mathbb{R}^{d_2} \ni (\mathbf{x}_B(f), \mathbf{x}_T) \mapsto \text{DeepONet}(\mathbf{x}_B(f), \mathbf{x}_T) := \left(\mathbf{B}_{q_B}(\sigma(\mathbf{B}_{q_{B}-1}(\dots \sigma(\mathbf{B}_1(\mathbf{x}_B(f))) \dots))) \right)^{\mathsf{T}} \left(\mathbf{T}_{q_T}(\sigma(\mathbf{T}_{q_{T}-1}(\dots \sigma(\mathbf{T}_1(\mathbf{x}_T)) \dots))) \right)$$
(1)

For a concrete example of using the above architecture, consider the task of solving the pendulum O.D.E from the previous section, $\mathbb{R}^2 \ni \frac{\mathrm{d}(y,v)}{\mathrm{d}t} = (v,-k\cdot\sin(y)+f(t))\in\mathbb{R}^2$. For a fixed initial condition, here the training/test data sets would be 3-tuples of the form, $(\mathbf{x}_B(f),x_T,y)$ where $y\in\mathbb{R}$ is the angular position of the pendulum at time $t=x_T$ for the forcing function f. Typically y is a standard O.D.E. solver's approximate solution. Given m such training data samples, the ℓ_2 empirical loss would be,

$$\hat{\mathcal{L}}_{DON} := \frac{1}{2m} \sum_{i=1}^{m} (y_i - \text{DeepONet}(\mathbf{x}_B(f_i), x_{T,i}))^2.$$
 (2)

The rationale for this loss function above comes from the universal approximation property of DeepONets which we shall quickly review in Section 1.2. Our key Theorem 15, applied to a DeepONet class with absolute value activations, would give that its average Rademacher complexity, for a training sample of size m, is bounded by,

$$\mathcal{O}\left(\frac{2^{n-1}\mathcal{C}_{n,n-1}}{\sqrt{m}} \left(\prod_{j=2}^{n-1} \mathcal{C}_{-j,-j}\right) M_{x,B} M_{x,T}\right)$$
(3)

with branch and trunk network both being of depth n (i.e $q_B = q_T = n$ in equation 1) and the inputs to them being bounded in expectation by $M_{x,B}$ and $M_{x,T}$ respectively. (For any weight matrix say A_i we shall let $A_{i,j}$ denote its j-th row.) Then the constants $C_{n,n-1}$ and $C_{-k,-k}$, k = 2, 3, ..., n-1 are defined in

terms of weight matrices of the branch and the trunk as follows,

$$\sum_{k_{1}=1}^{b_{-1}} \sum_{k_{2}=1}^{t_{-1}} \left[\sum_{j=1}^{p} (\mathbf{B}_{n,j} \mathbf{T}_{n,j}^{\mathsf{T}}) \right]_{k_{1},k_{2}} \cdot \|\mathbf{B}_{n-1,k_{1}}\| \cdot \|\mathbf{T}_{n-1,k_{2}}\| \leq C_{n,n-1},$$

$$\sup_{(\mathbf{v},\mathbf{w})} \sum_{j_{1}=1}^{b_{-k}} \sum_{j_{2}=1}^{t_{-k}} \left| (\mathbf{v}\mathbf{w}^{\mathsf{T}})_{j_{1},j_{2}} \right| \cdot \|\mathbf{B}_{n-k,j_{1}}\| \|\mathbf{T}_{n-k,j_{2}}\| \leq C_{-k,-k} \qquad \forall k = 2, \dots, n-1, \tag{4}$$

where in each sum, \mathbf{v}, \mathbf{w} are on the unit spheres of the same dimensionality as the number of rows in \mathbf{B}_{n-k} and \mathbf{T}_{n-k} respectively.

Towards making the above measures computationally more tractable, in Appendix \mathbf{H} we have shown that one can choose, $\forall k = 2, ..., n-1$, $C_{-k,-k} = \|\mathbf{X}\|$ where $\mathbf{X} \in \mathbb{R}^{b_{-k} \times t_{-k}}$ with $\mathbf{X}_{j_1,j_2} \coloneqq \|\mathbf{B}_{n-k,j_1}\| \cdot \|\mathbf{T}_{n-k,j_2}\|$

At the very outset, we discuss the *practical applicability* and *qualitative significance* of such a bound on the Rademacher complexity.

Choosing the regularizer: Since low Rademacher complexity implies better generalization (Theorem 16), our bound on the Rademacher complexity can be seen as suggesting the right regularization to be used in training DeepONets. Our bound in Theorem 15, suggests this novel combination of the layer norms of a DeepONet $(C_{n,n-1} \prod_{j=2}^{n-1} C_{-j,-j})$ – to be a good regularizer irrespective of the differential equation being solved.

Additionally, note that, DeepONets without biases and with positively homogeneous activation, are invariant to scaling the branch net's layers \mathbf{B}_k and trunk net's layers \mathbf{T}_k by any μ_k & $\lambda_k > 0$ respectively $\forall k \text{ s.t } \prod_k (\mu_k \cdot \lambda_k) = 1$ This is a larger symmetry than for usual nets but our complexity measure mentioned above is invariant under this combined scaling. This can be seen as being strongly suggestive of our result being a step in the right direction.

Explaining overparameterization: Since our generalization error bound has no explicit dependence on the width (or the number of parameters), it constitutes a step towards explaining the benefits of overparameterization in this new setup.

Outline of the Proof Technique Derivation of the main Theorem 15 involves 3 key steps: (a) formulating a variation of the standard Talagrand contraction (Lemma 14) (b) using this to bound the Rademacher complexity of a class of DeepONets with certain activations (e.g. absolute value) by the Rademacher complexity for a class of DeepONets having 1-lower depth and 1-dimensional outputs, for both the branch and the trunk (Lemma 17) and lastly (c) uncovering a recursive structure for the Rademacher complexity across depths between special DeepONet classes having 1-dimensional outputs for both the branch and trunk. (Lemma 18).

Lemma 17 removes the last 4 matrices from the DeepONet (2 each from the branch and the trunk) leading to one-dimensional output branch and trunk nets. Lemma 18 removes 2 matrices (1 each from branch and trunk) – by an entirely different argument than needed in the former. Lemma 17 is invoked only once at the beginning, while Lemma 18 is repeatedly used for each remaining layer of the DeepONet.

We would also like to emphasize that both our "peeling" lemmas above are structurally very different from the one in [10] - where the last layer of a standard net gets peeled in every step.

1.2 Review of the Universal Approximation Property of DeepONets

We follow the setup in [18] for a brief review of its key theorem about DeepONets.

Definition 1 (Solution Operator) Given $U \subset \mathbb{R}^n$ and $D \subset \mathbb{R}^d$, suppose we have a differential operator $\mathcal{L}: L^2(U) \to C(D)$. Denote the functions in the range of \mathcal{L} to be "forcing" / "input" functions and those in the domain of \mathcal{L} to be the "output" functions. Let ∂U denote the boundary of U. Then given $g \in L^2(\partial U)$ and $f \in C(D)$ a solution to the differential system (g, f, \mathcal{L}) is a function $u^* \in L^2(U)$ s.t.

$$\mathcal{L}u^* = f \ s.t \ u^* = g \ on \ \partial U.$$

At a fixed "boundary condition" g for such a differential system, we assume that the solutions for the above system for different forcing functions f are given via an operator/map, \mathcal{G} s.t.

$$\mathcal{G}: C(D) \to L^2(U)$$
 s.t $\mathcal{L} \circ \mathcal{G}(f) = f$ and $\mathcal{G}(f) = g$ on ∂U .

Further, assume that μ is a probability measure on C(D) s.t $\mathcal{G} \in L^2(\mu)$ i.e $\int_{C(D)} \|\mathcal{G}(f)\|_{L^2(U)}^2 d\mu(f) < \infty$.

Multiple examples of \mathcal{G} have been discussed in [18], and bounds on these operators evaluated – for instance, in Lemma 4.1 therein one can see the bounds on the \mathcal{G} that corresponds to the the case of a forced pendulum that we used as a demonstrative example in Section 1.1

Definition 2 (DeepONet (Version 1)) Suppose $\forall f \in C(D), \mathbf{x}_B(f)$ is a discretization of f. Further suppose \mathcal{A} is a branch net mapping to \mathbb{R}^p and $\tau : \mathbb{R}^n \mapsto \mathbb{R}^{p+1}$ is a trunk net. Then a "DeepONet" $(\mathcal{N}: L^2(D) \mapsto C(U))$ is defined as,

$$\forall \mathbf{x}_T \in U \subset \mathbb{R}^n, \quad \mathcal{N}(f)(\mathbf{x}_T) \coloneqq \tau_0(\mathbf{x}_T) + \sum_{k=1}^p \mathcal{A}_k(\mathbf{x}_B(f))\tau_k(\mathbf{x}_T).$$

Theorem 3 (Restatement of a key result from [18]) Let μ be as in Definition 1. Let $\mathcal{G}: C(D) \to L^2(U)$ be a Borel measurable mapping, with $\mathcal{G} \in L^2(\mu)$, then for every $\epsilon > 0$, there exists an operator network $\mathcal{N}: C(D) \to L^2(U)$, such that

$$\|\mathcal{G} - \mathcal{N}\|_{L^{2}(\mu)} = \left(\int_{C(D)} \|\mathcal{G}(f) - \mathcal{N}(f)\|_{L^{2}(U)}^{2} d\mu(f)\right)^{1/2} < \epsilon.$$

Remark 4 Henceforth $\mathbf{x}_B = \mathbf{x}_B(f)$ for any function f, and similarly $\mathbf{x}_{B,i}$ for a function f_i .

The above approximation guarantee between DeepONets (\mathcal{N}) and solution operators of differential equations (\mathcal{G}) clearly motivates the use of DeepONets for solving differential equations.

1.3 Related Work

To the best of our knowledge, Theorem 5.3 in [18] is the only existing result on generalization error bounds for DeepONets. However, this bound has an explicit dependence on the total number of parameters (the parameter d_{θ} there) in the DeepONet. Such a bound is not expected to explain the benefits of overparameterization well, which is one of the key features of modern deep learning. [7], [36]. We note that for usual implementations for DeepONets, where depths are typically small and the layers are wide, for a class at any fixed value of our complexity measure i.e $C_{n,n-1}\left(\prod_{j=2}^{n-1} C_{-j,-j}\right)$, a Rademacher complexity based bound that scales exponentially in depth (as ours) will be better than one that scales with the total number of parameters. We recall that PINNs are an alternative method to solve differential equations using a single neural net, for which generalization properties have been analysed in [14].

Organization In Section 2 we describe the mathematical setup needed for our main results. In Section 3 we give the formal statements of the central results in this paper: Lemma 14 (Proof in Appendix F.1) being the new variation of the Talagrand contraction lemma that we need, Theorem 15 (Proof in Section 5) being the main theorem giving a bound on the Rademacher complexity of a class of DeepONets and in Theorem 16 (Proof in Appendix E.1) we use the former to give a generalization error bound on the same.

In Section 4 we state the "peeling" lemmas required for proving Theorem 15: Lemma 17 (Proof in Appendix C) and Lemma 18 (Proof in Appendix D). The argument in Section 5 shows how these two lemmas combine to give the proof of Theorem 15. We conclude in Section 6 by discussing some open questions.

Notation Given any $U \subset \mathbb{R}^n$, denote as $L^2(U)$ the set of all functions $f: U \to \mathbb{R}$ s.t $\int_U f^2(\mathbf{x}) \, \mathrm{d}\mu(\mathbf{x}) < \infty$ where μ is the standard Lebesgue measure on \mathbb{R}^n . And we denote as C(U), the set of all real valued continuous functions on U. The unit sphere in \mathbb{R}^k is denoted by $S^{k-1} \coloneqq \left\{ \mathbf{x} \in \mathbb{R}^k \mid \|x\|_2 = 1 \right\}$. For any matrix \mathbf{A} , $\|\mathbf{A}\| = \sup_{\mathbf{v} \neq 0} \frac{\|\mathbf{A}\mathbf{v}\|_2}{\|\mathbf{v}\|_2}$ denotes its spectral norm.

2 Mathematical Setup

We define a few specific DeepONet architectures, which will be our focus in the subsequent work.

Definition 5 (A DeepONet (Version 2)) Let p be the common output dimension of the "branch net" and the "trunk net". Let q_B be the depth of the branch network and q_T be the depth of trunk net. Let $\mathbf{B}_1, \mathbf{B}_2, \ldots, \mathbf{B}_{q_B}$ be a valid set of the branch net's weight matrices (i.e the product $\mathbf{B}_{q_B} \cdot \ldots \cdot \mathbf{B}_1$ is well-defined and \mathbf{B}_{q_B} has p rows). Let $\mathbf{T}_1, \mathbf{T}_2, \ldots, \mathbf{T}_{q_T}$ be a valid set of trunk net's weight matrices (i.e the product $\mathbf{T}_{q_T} \cdot \ldots \cdot \mathbf{T}_1$ is well-defined and \mathbf{T}_{q_T} has p rows).

Let σ_1 and σ_2 be two $\mathbb{R} \to \mathbb{R}$ functions (applied pointwise on vector inputs), d_1 be the domain dimension of \mathbf{B}_1 and d_2 be the domain dimension of \mathbf{T}_1 . Then we define the corresponding DeepONet as the map,

$$\mathbb{R}^{d_1} \times \mathbb{R}^{d_2} \ni (\mathbf{x}_B, \mathbf{x}_T) \mapsto DeepONet(\mathbf{x}_B, \mathbf{x}_T) = \langle f_B(\mathbf{x}_B), f_T(\mathbf{x}_T) \rangle \in \mathbb{R},$$

where,

$$f_B(\mathbf{x}_B) \coloneqq \mathbf{B}_{q_B}(\sigma_1(\mathbf{B}_{q_B-1}(\dots \sigma_1(\mathbf{B}_1(\mathbf{x}_B))\dots))),$$

$$f_T(\mathbf{x}_T) \coloneqq \mathbf{T}_{q_T}(\sigma_2(\mathbf{T}_{q_T-1}(\dots \sigma_2(\mathbf{T}_1(\mathbf{x}_T))\dots))).$$

Remark 6 To emphasize some constraint on the weights of a DeepONet, we will denote DeepONet($\mathbf{x}_B, \mathbf{x}_T$) by DeepONet_{\mathbf{w}}($\mathbf{x}_B, \mathbf{x}_T$), where \mathbf{w} collectively stands for all the weight matrices in branch and trunk networks

Remark 7 The Rademacher complexity bound in Theorem 13 will be specifically for a class of Deep ONets with $q_B = q_T = 2$ and $\sigma_1 = \sigma_2 = \text{ReLU}$. The stronger bound in the main Theorem 15 and the supporting lemmas 14, 17 and 18 will apply for a class of activation functions that include the absolute value function non-linearity i.e $\mathbb{R} \ni x \mapsto |x| \in \mathbb{R}$. In Appendix G we will indicate why this is sufficient to capture the core mathematical challenges of the usual ReLU based implementations. Now we formalize the setup of the training data and the loss function for training Deep ONets.

Definition 8 (A Loss function for DeepONets) We continue in the same setup as Definition 1 and define a function class of allowed forcing functions $\mathcal{F} \subset L^2(D)$. Further, we consider DeepONet maps as given in Definition 5, mapping as DeepONet: $\mathbb{R}^{d_1} \times \mathbb{R}^{d_2} \to \mathbb{R}$ and consider an instance of the training data given as,

$$\{(f_i, \mathbf{x}_{T,i}) \in \mathcal{F} \times \mathbb{R}^{d_2} \mid i = 1, \dots, \mathcal{S}\}.$$

Then the corresponding DeepONet loss function is given by,

$$\hat{\mathcal{L}}_{\text{DON}} = \frac{1}{2m} \sum_{i=1}^{m} (\mathcal{G}(f_i)(\mathbf{x}_{T,i}) - \text{DeepONet}(\mathbf{x}_{B,i}, \mathbf{x}_{T,i}))^2.$$

Here we assume a fixed grid of size d_1 on which the function f_i gets discretized, to get $\mathbf{x}_{B,i} \in \mathbb{R}^{d_1}$.

It can be seen that the loss function in the experiment described in Section 1.1 was a special case of the above loss function – if we assume that the numerical solver exactly solved the forced pendulum O.D.E.

Definition 9 (\mathcal{H} function class) Given an operator $\mathcal{G}: C(D) \to L^2(U)$ as in Definition 1 (with say $U \subset \mathbb{R}^{d_2}$), a class $\mathcal{F} \subset C(D)$ of "forcing functions", and maps DeepONet_w: $\mathbb{R}^{d_1} \times \mathbb{R}^{d_2} \to \mathbb{R}$ as in Definition 5 and a set \mathcal{W} of allowed \mathbf{w} , we define the class $\mathcal{H}(\mathcal{F}, \mathcal{G}, \mathcal{W})$ as,

$$\mathcal{H}(\mathcal{F}, \mathcal{G}, \mathcal{W}) \coloneqq \{ \mathcal{F} \times \mathbb{R}^{d_2} \ni (f, \mathbf{x}_T) \mapsto \mathcal{G}(f)(\mathbf{x}_T) - \text{DeepONet}_{\mathbf{w}}(\mathbf{x}_B(f), \mathbf{x}_T) \in \mathbb{R} \mid \mathbf{w} \in \mathcal{W} \}.$$

Here we assume that there is a fixed grid of size d_1 on which the function f gets discretized to create the vector $\mathbf{x}_B(f) \in \mathbb{R}^{d_1}$.

Now, for some R > 0, we consider the map $\ell_2 := [-R, R] \ni x \mapsto \frac{1}{2} \cdot x^2$. Referring to Definition 8, for any $h \in \mathcal{H}(\mathcal{C})$, the DeepONet mean-squared loss can be rewritten as

$$\hat{\mathcal{L}}_{\text{DON}} = \frac{1}{m} \sum_{i=1}^{m} \ell_2 \circ h(f_i, \mathbf{x}_{T,i}).$$
 (5)

Definition 10 (Empirical and Average Rademacher complexity) For a function class \mathcal{F} , suppose being given a m-sized data-set \mathcal{D} of points $\{\mathbf{x}_i \mid i=1,\ldots,m\}$ in the domain of the elements in \mathcal{F} . For $\epsilon_i \sim \pm 1$ with equal probability, the corresponding empirical Rademacher complexity is given by,

$$\hat{\mathcal{R}}_m(\mathcal{F}) = \mathbb{E}\left[\sup_{\epsilon \in \mathcal{F}} \frac{1}{m} \sum_{i=1}^m \epsilon_i f(\mathbf{x}_i)\right].$$

If the elements of \mathcal{D} above are sampled from a distribution P, then the average Rademacher complexity is given by,

$$\mathcal{R}(\mathcal{F}) = \underset{\mathbf{x}_1, \dots, \mathbf{x}_m \sim P}{\mathbb{E}} \left(\mathbb{E} \left[\sup_{f \in \mathcal{F}} \frac{1}{m} \sum_{i=1}^m \epsilon_i f(\mathbf{x}_i) \right] \right).$$

We will uncover a recursive structure for the Rademacher complexity of DeepONets. Towards stating these results, we define a class of sub-DeepONets (Equation 9). Corresponding to any given DeepONet (Definition 5), these sub-DeepONets would themselves be one depth lower in the branch and the trunk network, and would always have one dimensional outputs for the branch and the trunk.

Definition 11 (Certain Classes of sub-DeepONets & Width Parameters for DeepONets)

- For $k = 1, ..., q_B 1$ and $\ell = 1, ..., q_T 1$, define b_{-k} , &t_{-\ell} to be the number of rows in the weight matrices \mathbf{B}_{q_B-k} and $\mathbf{T}_{q_T-\ell}$ respectively.
- In the setup of Definition 5, we define the functions \mathbf{f}_B' and \mathbf{f}_T' s.t given any m inputs to a DeepONet as, $\{(\mathbf{x}_{B,i},\mathbf{x}_{T,i}) \mid i=1,\ldots,m\}$ we have the following equalities,

$$f_B(\mathbf{x}_{B,i}) = \mathbf{B}_{q_B} \sigma_1 \left(\mathbf{B}_{q_B-1} \mathbf{f}_B'(\mathbf{x}_{B,i}) \right), \ f_T(\mathbf{x}_{T,i}) = \mathbf{T}_{q_T} \sigma_2 \left(\mathbf{T}_{q_T-1} \mathbf{f}_T'(\mathbf{x}_{T,i}) \right). \tag{6}$$

- Let W_{rest} be a set of allowed matrices for nets f'_B and f'_T .
- Further, given a constant $C_{q_B,q_T,q_{B}-1,q_{T}-1} > 0$ we define the following set of 4-tuples of outermost layer of matrices in the DeepONet as,

$$\mathcal{W}(\mathcal{C}_{q_{B},q_{T},q_{B}-1,q_{T}-1}) \coloneqq \left\{ (\mathbf{B}_{q_{B}}, \mathbf{B}_{q_{B}-1}, \mathbf{T}_{q_{T}}, \mathbf{T}_{q_{T}-1}) \right| \\
\sum_{k_{1}=1}^{b_{-1}} \sum_{k_{2}=1}^{t_{-1}} \|\mathbf{B}_{q_{B}-1,k_{1}}\| \cdot \|\mathbf{T}_{q_{T}-1,k_{2}}\| \cdot \left[\sum_{j=1}^{p} (\mathbf{B}_{q_{B},j} \mathbf{T}_{q_{T},j}^{\mathsf{T}}) \right]_{k_{1},k_{2}} \le \mathcal{C}_{q_{B},q_{T},q_{B}-1,q_{T}-1} \right\}.$$
(7)

• Corresponding to the above we define the following class of DeepONets,

DeepONet(
$$\mathcal{W}(\mathcal{C}_{q_B,q_T,q_{B}-1,q_T-1}), \mathcal{W}_{rest}$$
) :=
$$\left\{ \text{DeepONet}_{\mathbf{w}} \ (as \ in \ Definition \ 5) \ \middle| \ \mathbf{w} \in (\mathcal{W}(\mathcal{C}_{q_B,q_T,q_{B}-1,q_T-1}), \mathcal{W}_{rest}) \right\}. \tag{8}$$

• Further, given f'_B and f'_T as defined in Equation 6 and $(\mathbf{x}_B, \mathbf{x}_T) \in \mathbb{R}^{d_1} \times \mathbb{R}^{d_2}$, we define a class of DeepONets which is of one lower depth on its branch as well as the trunk than in Equation 8.

DeepONet(
$$\mathcal{W}_{rest}$$
) := $\left\{ (\mathbf{x}_B, \mathbf{x}_T) \mapsto \mathbf{v}^{\mathsf{T}} \mathbf{f}_B'(\mathbf{x}_B) \cdot \mathbf{w}^{\mathsf{T}} \mathbf{f}_T'(\mathbf{x}_T) \in \mathbb{R} \right\}$ (9)
$$\left\{ \text{the set of allowed matrices for nets } f_B' \& f_T' \text{ are in } \mathcal{W}_{rest} ; (\mathbf{v}, \mathbf{w}) \in S^{b_{-2}-1} \times S^{t_{-2}-1} \right\}.$$

3 Results

Our central result about Rademacher complexity of general DeepONets will be stated in the two theorems in Section 3.2. Towards that, we start building intuition with two relatively simple classes of DeepONets. A trivial DeepONet is where both the nets are linear transformations and its Rademacher complexity bound has been given in Appendix A. In the following we give the results for when both the branch and the trunk network have a single layer of ReLU activations.

3.1 Capacity Bounds for (2,2) – DeepONets

Definition 12 (A (2,2) – (ReLU) **DeepONet Class)** Given a constant C > 0, we define a (2,2) – (ReLU) DeepONet class where the branch (of width b_1) and the trunk (of width t_1) are mapping $\mathbb{R}^{d_1} \to \mathbb{R}^p$ and $\mathbb{R}^{d_2} \to \mathbb{R}^p$ respectively,

$$\mathcal{F}_{(2,2)-(\text{ReLU})}(\mathcal{C}) \coloneqq \left\{ \mathbb{R}^{d_1} \times \mathbb{R}^{d_2} \ni (\mathbf{x}_1, \mathbf{x}_2) \to \left\langle \mathbf{B}_2 \operatorname{ReLU}(\mathbf{B}_1 \mathbf{x}_1), \mathbf{T}_2 \operatorname{ReLU}(\mathbf{T}_1 \mathbf{x}_2) \right\rangle \in \mathbb{R} \right.$$

$$\left. \left| \max_{\substack{k_1 = 1, \dots, b_1 \\ k_2 = 1, \dots, t_1}} \| \mathbf{B}_{1,k_1} \| \| \mathbf{T}_{1,k_2} \| \left(\sum_{j=1}^p (\mathbf{B}_{2,j} \mathbf{T}_{2,j}^\top)_{k_1,k_2} \right) \le \mathcal{C} \right.$$

$$\left. \left| \mathbf{B}_2 \in \mathbb{R}^{p \times b_1}, \mathbf{B}_1 \in \mathbb{R}^{b_1 \times d_1}, \mathbf{T}_2 \in \mathbb{R}^{p \times t_1}, \mathbf{T}_1 \in \mathbb{R}^{t_1 \times d_2} \right\}.$$

Here $\mathbf{B}_{i,k}$ is the k^{th} – row of \mathbf{B}_i and similarly for the \mathbf{T}_i s.

Theorem 13 (Rademacher complexity of (2,2) – (ReLU)DeepONets) In the setup of Definition 10, suppose the given data-set \mathcal{D} of size m is $\{(\mathbf{x}_B^i, \mathbf{x}_I^i) \in \mathbb{R}^{d_1} \times \mathbb{R}^{d_2} \mid i = 1, ..., m\}$.

Then we have for the function class given in Definition 12,

$$\hat{\mathcal{R}}_m\Big(\mathcal{F}_{(2,2)-(\text{ReLU})}(\mathcal{C})\Big) \leq \frac{\mathcal{C}}{m} \, b_1 t_1 \left(\sqrt{\sum_{i=1}^m \left\|\mathbf{x}_B^i\right\|^2} \cdot \sqrt{\sum_{i=1}^m \left\|\mathbf{x}_T^i\right\|^2} \right).$$

The proof of the above has been given in Appendix B.3 The key upshot from the above theorem is to identify a capacity measure for the (2,2) – (ReLU) DeepONet as,

$$\max_{\substack{k_1 = 1, \dots, b_1 \\ k_2 = 1, \dots, t_1}} \|\mathbf{B}_{1, k_1}\| \|\mathbf{T}_{1, k_2}\| \left(\sum_{j=1}^p (\mathbf{B}_{2, j} \mathbf{T}_{2, j}^{\mathsf{T}})_{k_1, k_2}\right).$$

3.2 Main Results About General DeepONets

We note that the bound on the Rademacher complexity of (2,2)–(ReLU) DeepONet obtained in Theorem 13 has the lacuna of having explicit dependence on width. In the next (main) result we see how this can be improved upon for a certain class of activations, including the absolute value function. This main result will also hold for DeepONets of arbitrary depth – which leads us to defining a complexity measure for each layer. Towards stating Theorem 13, we need the following lemma which can be seen as a variation of the standard Talagrand contraction lemma,

Lemma 14 Let \mathcal{F} and \mathcal{G} be two sets of functions valued in \mathbb{R} . Let $\phi_F, \phi_G : \mathbb{R} \to \mathbb{R}$ be s.t $\exists L > 0$ s.t $\forall (f,g), (f',g') \in \mathcal{F} \times \mathcal{G}$ and $\forall (\mathbf{x},\mathbf{y}) \in \text{Domain}(\mathcal{F}) \times \text{Domain}(\mathcal{G})$,

$$|\phi_F(f(\mathbf{x}))\phi_G(g(\mathbf{y})) - \phi_F(f'(\mathbf{x}))\phi_G(g'(\mathbf{y}))| \le L|f(\mathbf{x})g(\mathbf{y}) - f'(\mathbf{x})g'(\mathbf{y})|. \tag{10}$$

Given points $\{\mathbf{x}_i \mid i=1,\ldots,m\}$, $\{\mathbf{y}_i \mid i=1,\ldots,m\}$ in the domain of the functions in $\mathcal{F} \times \mathcal{G}$ we have the following inequality of Rademacher complexities - where both the sides are being evaluated on this same set of points,

$$\mathcal{R}_{\mathrm{m}}(\phi_F \circ \mathcal{F} \cdot \phi_G \circ \mathcal{G}) \leq L\mathcal{R}_{\mathrm{m}}(\mathcal{F} \cdot \mathcal{G}).$$

The above lemma has been proven in Appendix F.1.

Theorem 15 (Rademacher Complexity of Special Symmetric DeepONets) We consider a special case of Definition 5, with (a) $q_B = q_T = n$, and (b) σ_1, σ_2 satisfying Lemma 14 and being positively homogeneous. Further, let b_{-i} and t_{-i} be as in Definition 11 for $i \ge 1$ and recall from Definition 5 that p is the number of rows of \mathbf{B}_n and \mathbf{T}_n . Then given a class of DeepONet maps as above, we define n-1 constants: $C_{n,n-1} > 0$ and $C_{-k,-k} > 0$, $k = 2, \ldots, n-1$, such that the following n-1 inequalities hold,

$$(\mathcal{S}_k \coloneqq S^{b_{-k}-1} \times S^{t_{-k}-1})$$

$$\sum_{k_{1}=1}^{b_{-1}} \sum_{k_{2}=1}^{t_{-1}} \|\mathbf{B}_{n-1,k_{1}}\| \cdot \|\mathbf{T}_{n-1,k_{2}}\| \cdot \left[\sum_{j=1}^{p} (\mathbf{B}_{n,j} \mathbf{T}_{n,j}^{\mathsf{T}})\right]_{k_{1},k_{2}} \leq C_{n,n-1},$$

$$\sup_{(\mathbf{v},\mathbf{w}) \in \mathcal{S}_{k}} \sum_{j_{1}=1}^{b_{-k}} \sum_{j_{2}=1}^{t_{-k}} \left| (\mathbf{v} \mathbf{w}^{\mathsf{T}})_{j_{1},j_{2}} \right| \|\mathbf{B}_{n-k,j_{1}}\| \|\mathbf{T}_{n-k,j_{2}}\| \leq C_{-k,-k}, \ \forall k = 2, \ldots, n-1.$$

Then given training data as in Definition 11, the empirical Rademacher complexity is bounded as,

$$\hat{\mathcal{R}}_{m} \leq \frac{(2L)^{n-1} \mathcal{C}_{n,n-1}}{m} \left(\prod_{j=2}^{n-1} \mathcal{C}_{-j,-j} \right) \sqrt{\sum_{i=1}^{m} \|\mathbf{x}_{B,i}\|_{2}^{2} \|\mathbf{x}_{T,i}\|_{2}^{2}}.$$

Further assuming that the input distribution is bounded as $\mathbb{E}\left[\|\mathbf{x}_B\|_2^2\right] \leq M_{x,B}^2$, $\mathbb{E}\left[\|\mathbf{x}_T\|_2^2\right] \leq M_{x,T}^2$, we have the average Rademacher complexity bounded as:

$$\mathcal{R} \leq \frac{(2L)^{n-1} \mathcal{C}_{n,n-1}}{\sqrt{m}} \left(\prod_{j=2}^{n-1} \mathcal{C}_{-j,-j} \right) M_{x,B} M_{x,T}.$$

The proof for above can be found in Appendix 5. Further note that, (a) It follows from previous definitions that the rows of $\mathbf{B}_{n-(i-1)}$ and $\mathbf{T}_{n-(i-1)}$ are vectors in \mathbb{R}^{b-i} and \mathbb{R}^{t-i} respectively. (b) For $\sigma_1(x) = \sigma_2(x) = |x|$ i.e for DeepONets with absolute value activations, Lemma 14 holds with L = 1 and hence the subsequent simplification also happens in the result in Theorem 15.

Towards deriving a generalization error bound we need the following assumption that is motivated from [18].

Assumption 1 We recall assumption 5.1 from [18] that for all $f \in C(D)$, $\mathbf{x}_T \in U$, there exist constants $C_f, \kappa > 0$ such that,

$$|\mathcal{G}(f)(\mathbf{x}_T)| \le C_f (1 + ||f||_{L^2})^{\kappa}.$$
 (11)

Further, given \mathcal{F} as in Definition 8, we assume that the following is finite, $\sup_{f \in \mathcal{F}} \left(C_f \left(1 + \|f\|_{L^2} \right)^{\kappa} \right)$

We note that often in DeepONet experimental setups the function space \mathcal{F} is chosen as the set of all linear combinations of a finite set of functions with the coefficients coming from specified compact intervals of \mathbb{R} . Given the continuity of the arguments of the sup above, from the perspective of many usual experimental setups, it follows that it is natural to assume the finiteness of the sup required above.

Theorem 16 (Generalization Error Bound for DeepONet) Given \mathcal{H} as in Definition 9, define $h^* \in \mathcal{H}$ as the minimizer for the empirical loss $\hat{\mathcal{L}}_{DON}$ in Equation 5. We assume that \mathcal{G} satisfies Assumption 1. Then, $\forall \delta > 0$, w.p at least $1 - \delta$ over sampling the training data, the difference between true risk of this minimizer and the minimum possible true risk can be bounded as,

$$\mathbb{E}_{(f,\mathbf{x}_T)\sim\mathcal{D}}\left[\ell_2(h^*(f,\mathbf{x}_T)) - \min_{h\in\mathcal{H}}\left[\mathbb{E}_{(f,\mathbf{x}_T)\sim\mathcal{D}}\left[\ell_2(h(f,\mathbf{x}_T))\right]\right] \leq 2B\cdot\mathcal{R} + B^2\sqrt{\frac{\ln(1/\delta)}{2m}},$$

where \mathcal{R} is as in Theorem 15. \mathcal{F} is as in Definition 8, L_1, L_2 are the Lipschitz constants of activation functions $\sigma_1(\cdot), \sigma_2(\cdot)$ respectively and we have defined B as,

$$B = \sup_{f \in \mathcal{F}} \left(C_f \left(1 + \| f \|_{L^2} \right)^{\kappa} \right) + L_1^n L_2^n m_{x,B} m_{x,T} P,$$

where we assume $\|\mathbf{x}_B\|_2 \le m_{x,B}$, $\|\mathbf{x}_T\|_2 \le m_{x,T}$ ($\mathbf{x}_B, \mathbf{x}_T$ are inputs to branch and trunk respectively), and $P = \sup \left(\prod_{i=1}^n \|\mathbf{B}_i\| \|\mathbf{T}_i\|\right)$, where the sup is over the same set of DeepONets as considered in Theorem 15.

Proof of the above can be found in Appendix E.1. Note that for absolute value activations - which would satisfy the condition for Theorem 15., the quantity B above would simplify to,

$$B = \sup_{f \in \mathcal{F}} \left(C_f \left(1 + ||f||_{L^2} \right)^{\kappa} \right) + m_{x,B} \, m_{x,T} \, P,$$

4 Key Lemmas Needed for Theorem 15

We would like to emphasize that the subsequent lemmas 17 and 18 hold in more generality than Theorem 15, because they do not need the branch and the trunk nets to be of equal depth.

Lemma 17 (Removal of the last 4 matrices of a DeepONet) We continue to be in the setup of Definition 11 and assume that σ_1, σ_2 satisfy Lemma 14 and are positively homogeneous. Then given the definitions of DeepONet($W(C_{q_B,q_T,q_{B-1},q_{T-1}}), W_{\text{rest}}$) (Eq. 8) and DeepONet(W_{rest}) (Eq. 9) in Definition 11, we have the following upperbound on emoirical Rademacher complexity of a DeepONet, (here $S_k := S^{b_{-k}-1} \times S^{t_{-k}-1}$)

$$\hat{\mathcal{R}}_{m}\left(\text{DeepONet}(\mathcal{W}(\mathcal{C}_{q_{B},q_{T},q_{B}-1,q_{T}-1}),\mathcal{W}_{\text{rest}})\right) \\
\leq 2L \cdot \mathcal{C}_{q_{B},q_{T},q_{B}-1,q_{T}-1} \cdot \mathbb{E}_{\epsilon}\left[\sup_{\substack{(\mathbf{v},\mathbf{w}) \in \mathcal{S}_{2} \\ \mathcal{W}_{\text{rest}}}} \left(\sum_{i=1}^{m} \epsilon_{i} \mathbf{v}^{\mathsf{T}} \mathbf{f}_{B}'(\mathbf{x}_{B,i}) \cdot \mathbf{w}^{\mathsf{T}} \mathbf{f}_{T}'(\mathbf{x}_{T,i})\right)\right] \\
\leq 2L \cdot \mathcal{C}_{q_{B},q_{T},q_{B}-1,q_{T}-1} \cdot \hat{\mathcal{R}}_{m}\left(\text{DeepONet}(\mathcal{W}_{\text{rest}})\right).$$

Note that both sides of the above are computed for the same data $\{(\mathbf{x}_{B,i},\mathbf{x}_{T,i}) \mid i=1,\ldots,m\}$ as given in Definition 11. The proof of the above lemma is given in Appendix C.

By referring to the definitions of the DeepONet classes on the L.H.S. and the R.H.S. of the above, as given in equations 8 and 9 respectively, we see that the above lemma upperbounds the Rademacher complexity of a DeepONet class (whose individual nets can have multi-dimensional outputs) by the Rademacher complexity of a simpler DeepONet class. The DeepONet class in the R.H.S. is simpler because here the last layer of each of the individual nets is constrained to be an unit vector of appropriate dimensions (and thus the individual nets here are always of 1 dimensional output) – and whose both branch and the trunk are shorter in depth by 1 activation and 1 linear transform than the L.H.S.

Lemma 18 (Peeling for DeepONets) We continue in the setup of Lemma 17 and define the functions \mathbf{f}_B'' and \mathbf{f}_T'' s.t we have the following equalities, $f_B' = \sigma_1 \left(\mathbf{B}_{q_B-2} \mathbf{f}_B'' \right) \& f_T' = \sigma_2 \left(\mathbf{T}_{q_T-2} \mathbf{f}_T'' \right)$ Further, given a constant $\mathcal{C}_{-2,-2} > 0$, we define \mathcal{W}'_{rest} as the union of (a) the set of weights that are allowed in the \mathcal{W}_{rest} set for the matrices $\mathbf{B}_{q_B-3}, \mathbf{B}_{q_B-4}, \ldots, \mathbf{B}_1$ and $\mathbf{T}_{q_T-3}, \mathbf{T}_{q_T-4}, \ldots, \mathbf{T}_1$ and (b) the subset of the weights for $\mathbf{B}_{q_B-2} \& \mathbf{T}_{q_T-2}$ that are allowed by \mathcal{W}_{rest} which also additionally satisfy the constraint, (here $\mathcal{S}_k \coloneqq S^{b_{-k}-1} \times S^{t_{-k}-1}$)

$$\sup_{(\mathbf{v}, \mathbf{w}) \in \mathcal{S}_2} \sum_{j_1=1}^{b-2} \sum_{j_2=1}^{t-2} \left| (\mathbf{v} \mathbf{w}^{\mathsf{T}})_{j_1, j_2} \right| \| \mathbf{B}_{q_B - 2, j_1} \| \| \mathbf{T}_{q_T - 2, j_2} \| \le \mathcal{C}_{-2, -2}. \tag{12}$$

Then we get the following inequality between Rademacher complexities,

$$\mathbb{E}_{\epsilon} \left[\sup_{(\mathbf{v}, \mathbf{w}) \in \mathcal{S}_{2}} \left(\sum_{i=1}^{m} \epsilon_{i} \mathbf{v}^{\mathsf{T}} \mathbf{f}_{B}'(\mathbf{x}_{B,i}) \cdot \mathbf{w}^{\mathsf{T}} \mathbf{f}_{T}'(\mathbf{x}_{T,i}) \right) \right]$$

$$\leq 2LC_{-2,-2} \mathbb{E}_{\epsilon} \left(\sup_{\substack{(\mathbf{v}, \mathbf{w}) \in \mathcal{S}_{3} \\ \mathcal{W}_{rest}'}} \sum_{i=1}^{m} \epsilon_{i} (\mathbf{v}^{\mathsf{T}} \mathbf{f}_{B}''(\mathbf{x}_{B,i})) (\mathbf{w}^{\mathsf{T}} \mathbf{f}_{T}''(\mathbf{x}_{T,i})) \right),$$

where b_{-k}, t_{-k} are as in Definition 11.

Proof of the above lemma is in Appendix D.

5 Proof of Theorem 15

Proof For each k = 2, 3, ..., n-1, we define a product of unit-spheres, $S_k := S^{b_{-k}-1} \times S^{t_{-k}-1}$ and let $S_n := S^{d_1-1} \times S^{d_2-1}$. Now we define,

$$\mathcal{W}_{\text{rest}}^{1} \coloneqq \bigg\{ \big(\mathbf{B}_{n}, \mathbf{B}_{n-1}, \mathbf{T}_{n}, \mathbf{T}_{n-1} \big) \; \bigg| \; \sum_{k_{1}=1}^{b-1} \sum_{k_{2}=1}^{t-1} \big\| \mathbf{B}_{n-1, k_{1}} \big\| \cdot \big\| \mathbf{T}_{n-1, k_{2}} \big\| \cdot \bigg[\sum_{j=1}^{p} \big(\mathbf{B}_{n, j} \mathbf{T}_{n, j}^{\mathsf{T}} \big) \bigg]_{k_{1}, k_{2}} \le \mathcal{C}_{n, n-1} \bigg\}.$$

Next, for each i = 2, 3, ..., n-1, we define

$$\mathcal{W}_{\mathrm{rest}}^{i} \coloneqq \bigcup_{k=1}^{n-i} \left\{ (\mathbf{B}_{k}, \mathbf{T}_{k}) \mid \sup_{(\mathbf{v}, \mathbf{w}) \in \mathcal{S}_{k}} \sum_{j_{1}=1}^{b_{-k}} \sum_{j_{2}=1}^{t_{-k}} \left| (\mathbf{v} \mathbf{w}^{\mathsf{T}})_{j_{1}, j_{2}} \right| \cdot \| \mathbf{B}_{n-k, j_{1}} \| \cdot \| \mathbf{T}_{n-k, j_{2}} \| \le \mathcal{C}_{-k, -k} \right\}.$$

Thus we have,

$$m\hat{\mathcal{R}}_{m} = \mathbb{E}_{\epsilon} \left[\sup_{(\mathcal{W}_{\text{rest}}^{1}, \mathcal{W}_{\text{rest}}^{2})} \left(\sum_{i=1}^{m} \epsilon_{i} \left\langle \mathbf{B}_{n} \sigma_{1} \left(\mathbf{B}_{n-1} \mathbf{f}_{B}'(\mathbf{x}_{B,i}) \right), \mathbf{T}_{n} \sigma_{2} \left(\mathbf{T}_{n-1} \mathbf{f}_{T}'(\mathbf{x}_{T,i}) \right) \right) \right) \right].$$

Then we can invoke Lemma 17 on the above to get,

$$m\hat{\mathcal{R}}_{m} = \mathbb{E}_{\epsilon} \left[\sup_{(\mathcal{W}_{\text{rest}}^{1}, \mathcal{W}_{\text{rest}}^{2})} \left(\sum_{i=1}^{m} \epsilon_{i} \left\langle \mathbf{B}_{n} \sigma_{1} \left(\mathbf{B}_{n-1} \mathbf{f}_{B}'(\mathbf{x}_{B,i}) \right), \mathbf{T}_{n} \sigma_{2} \left(\mathbf{T}_{n-1} \mathbf{f}_{T}'(\mathbf{x}_{T,i}) \right) \right\rangle \right) \right]$$

$$\leq 2LC_{n,n-1} \mathbb{E}_{\epsilon} \left[\sup_{\substack{\mathcal{W}_{\text{rest}}^{2} \\ \mathbf{v}_{1}, \mathbf{v}_{2} \in \mathcal{S}_{2}}} \left(\sum_{i=1}^{m} \epsilon_{i} \mathbf{v}_{1}^{\mathsf{T}} \mathbf{f}_{B}'(\mathbf{x}_{B,i}) \mathbf{v}_{2}^{\mathsf{T}} \mathbf{f}_{T}'(\mathbf{x}_{T,i}) \right) \right].$$

Now, using Lemma 18 repeatedly on the R.H.S above, and defining in a natural fashion the subsequent branch and trunk sub-networks as $\mathbf{f}^{(i)}(\cdot)$ we have,

$$m\hat{\mathcal{R}}_{m} \leq (2L)^{2}\mathcal{C}_{n,n-1}\mathcal{C}_{-2,-2}\mathbb{E}_{\epsilon} \left(\sup_{\substack{\mathcal{W}_{rest}^{3} \\ (\mathbf{v}_{1},\mathbf{v}_{2}) \in \mathcal{S}_{3}}} \sum_{i=1}^{m} \epsilon_{i}(\mathbf{v}_{1}^{\mathsf{T}}\mathbf{f}_{B}^{"}(\mathbf{x}_{B,i}))(\mathbf{v}_{2}^{\mathsf{T}}\mathbf{f}_{T}^{"}(\mathbf{x}_{T,i})) \right)$$

$$\cdots$$

$$\leq (2L)^{i}\mathcal{C}_{n,n-1} \left(\prod_{j=2}^{i} \mathcal{C}_{-j,-j} \right) \mathbb{E}_{\epsilon} \left(\sup_{\substack{\mathcal{W}_{rest}^{i+1} \\ (\mathbf{v}_{1},\mathbf{v}_{2}) \in \mathcal{S}_{i+1}}} \sum_{i=1}^{m} \epsilon_{i}(\mathbf{v}_{1}^{\mathsf{T}}\mathbf{f}_{B}^{(i)}(\mathbf{x}_{B,i}))(\mathbf{v}_{2}^{\mathsf{T}}\mathbf{f}_{T}^{(i)}(\mathbf{x}_{T,i})) \right)$$

$$\cdots$$

$$\leq (2L)^{n-1}\mathcal{C}_{n,n-1} \left(\prod_{j=2}^{n-1} \mathcal{C}_{-j,-j} \right) \mathbb{E}_{\epsilon} \left(\sup_{(\mathbf{v}_{1},\mathbf{v}_{2}) \in \mathcal{S}_{n}} \sum_{i=1}^{m} \epsilon_{i}(\mathbf{v}_{1}^{\mathsf{T}}\mathbf{x}_{B,i})(\mathbf{v}_{2}^{\mathsf{T}}\mathbf{x}_{T,i}) \right).$$

Using Lemma 19, (and its defintion of $\tilde{\mathbf{x}}^i = \mathbf{x}_{B,i} \mathbf{x}_{T,i}^{\mathsf{T}}$), the final bound on the empirical Rademacher complexity becomes,

$$\hat{\mathcal{R}}_m \leq \frac{(2L)^{n-1} \mathcal{C}_{n,n-1}}{m} \left(\prod_{j=2}^{n-1} \mathcal{C}_{-j,-j} \right) \sqrt{\sum_{i=1}^{m} \left\| \tilde{\mathbf{x}}^i \right\|_2^2}.$$

Invoking the assumption that the input is bounded s.t $\mathbb{E}\left[\|\tilde{\mathbf{x}}_B\|_2^2\right] \leq M_{x,B}^2$ and $\mathbb{E}\left[\|\tilde{\mathbf{x}}_T\|_2^2\right] \leq M_{x,T}^2$, the average Rademacher complexity can be bounded as

$$\mathcal{R} \leq \frac{(2L)^{n-1}\mathcal{C}_{n,n-1}}{\sqrt{m}} \left(\prod_{j=2}^{n-1} \mathcal{C}_{-j,-j} \right) M_{x,B} M_{x,T}.$$

Lemma 19 Using $S_n = S^{d_1-1} \times S^{d_2-1}$, and $\tilde{\mathbf{x}}^i = \mathbf{x}_{B,i} \mathbf{x}_{T,i}^{\mathsf{T}}$

$$\mathbb{E}_{\epsilon} \left(\sup_{(\mathbf{v}_1, \mathbf{v}_2) \in \mathcal{S}_n} \sum_{i=1}^m \epsilon_i(\mathbf{v}_1^{\mathsf{T}} \mathbf{x}_{B,i}) (\mathbf{v}_2^{\mathsf{T}} \mathbf{x}_{T,i}) \right) \leq \sqrt{\sum_{i=1}^m \|\mathbf{x}_{B,i}\|_2^2 \|\mathbf{x}_{T,i}\|_2^2}.$$

Proof Denote by $S_n = S^{d_1-1} \times S^{d_2-1}$. Define $\tilde{\mathbf{W}} = (\mathbf{v}_1 \mathbf{v}_2^{\mathsf{T}})$ and $\tilde{\mathbf{x}}_i = (\mathbf{x}_{B,i} (\mathbf{x}_{T,i})^{\mathsf{T}})$. Then

$$\begin{split} &= \mathbb{E}_{\epsilon} \left(\sup_{(\mathbf{v}_{1}, \mathbf{v}_{2}) \in \mathcal{S}_{n}} \sum_{i=1}^{m} \epsilon_{i} \left(\sum_{k_{1}=1}^{d_{1}} v_{1, k_{1}} x_{B, i, k_{1}} \right) \left(\sum_{k_{2}=1}^{d_{2}} v_{2, k_{2}} x_{T, i, k_{2}} \right) \right) \\ &= \mathbb{E}_{\epsilon} \left(\sup_{(\mathbf{v}_{1}, \mathbf{v}_{2}) \in \mathcal{S}_{n}} \sum_{i=1}^{m} \sum_{k_{1}=1}^{d_{1}} \sum_{k_{2}=1}^{d_{2}} \epsilon_{i} \left(v_{1, k_{1}} x_{B, i, k_{1}} \right) \left(v_{2, k_{2}} x_{T, i, k_{2}} \right) \right) \\ &= \mathbb{E}_{\epsilon} \left(\sup_{(\mathbf{v}_{1}, \mathbf{v}_{2}) \in \mathcal{S}_{n}} \sum_{k_{1}=1}^{d_{1}} \sum_{k_{2}=1}^{d_{2}} v_{1, k_{1}} v_{2, k_{2}} \left(\sum_{i=1}^{m} \epsilon_{i} \left(x_{B, i, k_{1}} \right) \left(x_{T, i, k_{2}} \right) \right) \right) \\ &\text{Note that,} (\mathbf{v}_{1}, \mathbf{v}_{2}) \in \mathcal{S}_{n} \implies \left\| \tilde{\mathbf{W}} \right\|_{2} = \sqrt{\sum_{k_{1}=1}^{d_{1}} \sum_{k_{2}=1}^{d_{2}} v_{1, k_{1}}^{2} v_{2, k_{2}}^{2}} = \left\| \mathbf{v}_{1} \right\|_{2} \left\| \mathbf{v}_{2} \right\|_{2} = 1. \end{split}$$

Hence, we can enlarge the domain to get,

$$\leq \mathbb{E}_{\epsilon} \left(\sup_{\|\tilde{\mathbf{W}}\|_{2} \leq 1} \sum_{k_{1}=1}^{d_{1}} \sum_{k_{2}=1}^{d_{2}} \tilde{W}_{(k_{1},k_{2})} \left(\sum_{i=1}^{m} \epsilon_{i} \, \tilde{x}_{i,k_{1},k_{2}} \right) \right).$$

The above sum can be viewed as an inner product of 2 vectors in $\mathbb{R}^{d_1 \times d_2}$. We have,

$$= \mathbb{E}_{\epsilon} \left(\sup_{\left\|\tilde{\mathbf{W}}\right\|_{2} \le 1} \left\langle \tilde{\mathbf{W}}, \sum_{i=1}^{m} \epsilon_{i} \tilde{\mathbf{x}}_{i} \right\rangle \right) \le \sqrt{\sum_{i=1}^{m} \left\|\tilde{\mathbf{x}}_{i}\right\|_{2}^{2}} = \sqrt{\sum_{i=1}^{m} \left\|\mathbf{x}_{B,i}\right\|_{2}^{2} \left\|\mathbf{x}_{T,i}\right\|_{2}^{2}},$$

where the inequality follows from Theorem 5.5 in [22].

6 Conclusion & Future Work

We would like to point out that our work immediately suggests an exciting line of future experimental studies investigating the optimal weighting for the regularizer $(C_{n,n-1}\prod_{j=2}^{n-1}C_{-j,-j})$ (equation 4) implied by our result, to gain the most from using it while training DeepONets. Different variations of the DeepONet architecture have also been proposed – (Fig 5, [33]), [21]. We anticipate that our mathematical analysis can be built upon to similarly derive regularizers for these architectures too.

In the context of understanding the generalization error of Siamese networks, the authors in [6] dealt with a certain product of neural outputs structure (where the nets share weights). In their analysis, the authors bound the Rademacher complexity via covering numbers. Since we try to directly bound the Rademacher complexity for DeepONets, it would be interesting to investigate if our methods can be adapted to improve such results about Siamese nets.

We recall that in works like [23], [17], [12] the authors were able to show that single neural nets can be constructed for efficient representation of solutions of certain classes of P.D.Es. Such P.D.E. class specific generalization bounds in the setup of Physics Inspired Neural Nets were given in [24]. We anticipate that similar analysis might be possible for DeepONets too giving tighter generalization bounds tuned to specific types of P.D.Es.

Further, in [34], an unsupervised variation of the loss function of a DeepONet was shown to give better performance. In [11], authors employ a variational framework for solving differential equations using DeepONets, through a novel loss function. Understanding precisely as to when these variations give advantages over the basic DeepONet setup is another interesting direction for future research.

Lastly, we note that the existing bounds on the Rademacher complexity of nets have typically been proven by making ingenious use of the algebraic identities satisfied by Rademacher complexity (Theorem 12 in [3]). But to the best of our knowledge, we are not aware of any general results on how Rademacher complexity of a product of function spaces can be written in terms of individual Rademacher complexities. Consequently, we are not aware of any method which can directly use the existing state-of-the-art bounds on Rademacher complexities for standard nets (in particular with a ReLU activation) to obtain the same for DeepONets. We posit that such a mathematical development if achieved can be a significant advance in the theory of DeepONets.

References

- [1] R. Arora, P. Bartlett, P. Mianjy, and N. Srebro. Dropout: Explicit forms and capacity control. In M. Meila and T. Zhang, editors, *Proceedings of the 38th International Conference on Machine Learning, ICML 2021, 18-24 July 2021, Virtual Event*, volume 139 of *Proceedings of Machine Learning Research*, pages 351–361. PMLR, 2021. URL http://proceedings.mlr.press/v139/arora21a.html.
- [2] S. Arora, R. Ge, B. Neyshabur, and Y. Zhang. Stronger generalization bounds for deep nets via a compression approach. In J. G. Dy and A. Krause, editors, *Proceedings of the 35th International Conference on Machine Learning, ICML 2018, Stockholmsmässan, Stockholm, Sweden, July 10-15, 2018*, volume 80 of *Proceedings of Machine Learning Research*, pages 254–263. PMLR, 2018. URL http://proceedings.mlr.press/v80/arora18b.html.
- [3] P. L. Bartlett and S. Mendelson. Rademacher and gaussian complexities: Risk bounds and structural results. *J. Mach. Learn. Res.*, 3(null):463–482, mar 2003. ISSN 1532-4435.
- [4] P. L. Bartlett, D. J. Foster, and M. Telgarsky. Spectrally-normalized margin bounds for neural networks. In I. Guyon, U. von Luxburg, S. Bengio, H. M. Wallach, R. Fergus, S. V. N. Vishwanathan, and R. Garnett, editors, Advances in Neural Information Processing Systems 30: Annual Conference on Neural Information Processing Systems 2017, December 4-9, 2017, Long Beach, CA, USA, pages 6240–6249, 2017. URL https://proceedings.neurips.cc/paper/2017/hash/b22b257ad0519d4 500539da3c8bcf4dd-Abstract.html.
- [5] T. Chen and H. Chen. Universal approximation to nonlinear operators by neural networks with arbitrary activation functions and its application to dynamical systems. *IEEE Transactions on Neural Networks*, 6(4):911–917, 1995. DOI: 10.1109/72.392253.
- [6] K. Dahiya, A. Agarwal, D. Saini, G. K, J. Jiao, A. Singh, S. Agarwal, P. Kar, and M. Varma. Siamesexml: Siamese networks meet extreme classifiers with 100m labels. In M. Meila and T. Zhang, editors, Proceedings of the 38th International Conference on Machine Learning, volume 139 of Proceedings of Machine Learning Research, pages 2330-2340. PMLR, 18-24 Jul 2021. URL https://proceedings.mlr.press/v139/dahiya21a.html.
- [7] Y. Dar, V. Muthukumar, and R. G. Baraniuk. A farewell to the bias-variance tradeoff? an overview of the theory of overparameterized machine learning. *CoRR*, abs/2109.02355, 2021. URL https://arxiv.org/abs/2109.02355.
- [8] G. K. Dziugaite and D. M. Roy. Computing nonvacuous generalization bounds for deep (stochastic) neural networks with many more parameters than training data. In G. Elidan, K. Kersting, and A. T. Ihler, editors, *Proceedings of the Thirty-Third Conference on Uncertainty in Artificial Intelligence, UAI 2017, Sydney, Australia, August 11-15, 2017.* AUAI Press, 2017. URL http://auai.org/uai 2017/proceedings/papers/173.pdf.
- [9] W. E, J. Han, and A. Jentzen. Algorithms for solving high dimensional PDEs: from nonlinear monte carlo to machine learning. *Nonlinearity*, 35(1):278–310, dec 2021. DOI: 10.1088/1361-6544/ac337f. URL https://doi.org/10.1088/1361-6544/ac337f.
- [10] N. Golowich, A. Rakhlin, and O. Shamir. Size-independent sample complexity of neural networks. In *Conference On Learning Theory*, pages 297–299. PMLR, 2018.
- [11] S. Goswami, M. Yin, Y. Yu, and G. E. Karniadakis. A physics-informed variational DeepONet for predicting crack path in quasi-brittle materials. *Computer Methods in Applied Mechanics and Engineering*, 391:114587, mar 2022. DOI: 10.1016/j.cma.2022.114587. URL https://doi.org/10.1016%2Fj.cma.2022.114587.
- [12] P. Grohs, F. Hornung, A. Jentzen, and P. von Wurstemberger. A proof that artificial neural networks overcome the curse of dimensionality in the numerical approximation of black-scholes partial differential equations, 2018. URL https://arxiv.org/abs/1809.02362.
- [13] J. Han, A. Jentzen, and W. E. Solving high-dimensional partial differential equations using deep learning. *Proceedings of the National Academy of Sciences*, 115(34):8505-8510, 2018. DOI: 10.1073/pnas.1718942115. URL https://www.pnas.org/doi/abs/10.1073/pnas.1718942115.

- [14] Z. Hu, A. D. Jagtap, G. E. Karniadakis, and K. Kawaguchi. When do extended physics-informed neural networks (xpinns) improve generalization? arXiv preprint arXiv:2109.09444, 2021.
- [15] G. E. Karniadakis, I. G. Kevrekidis, L. Lu, P. Perdikaris, S. Wang, and L. Yang. Physics-informed machine learning. *Nature Reviews Physics*, 3(6):422–440, 2021.
- [16] N. B. Kovachki, Z. Li, B. Liu, K. Azizzadenesheli, K. Bhattacharya, A. M. Stuart, and A. Anandkumar. Neural operator: Learning maps between function spaces. CoRR, abs/2108.08481, 2021. URL https://arxiv.org/abs/2108.08481.
- [17] G. Kutyniok, P. Petersen, M. Raslan, and R. Schneider. A theoretical analysis of deep neural networks and parametric pdes. CoRR, abs/1904.00377, 2019. URL http://arxiv.org/abs/1904.00377.
- [18] S. Lanthaler, S. Mishra, and G. E. Karniadakis. Error estimates for DeepONets: a deep learning framework in infinite dimensions. *Transactions of Mathematics and Its Applications*, 6(1), 03 2022. ISSN 2398-4945. DOI: 10.1093/imatrm/tnac001. URL https://doi.org/10.1093/imatrm/tnac001. tnac001.
- [19] Z. Li, N. B. Kovachki, K. Azizzadenesheli, B. Liu, K. Bhattacharya, A. M. Stuart, and A. Anand-kumar. Fourier neural operator for parametric partial differential equations. In 9th International Conference on Learning Representations, ICLR 2021, Virtual Event, Austria, May 3-7, 2021. Open-Review.net, 2021. URL https://openreview.net/forum?id=c8P9NQVtmn0.
- [20] L. Lu, P. Jin, G. Pang, Z. Zhang, and G. E. Karniadakis. Learning nonlinear operators via DeepONet based on the universal approximation theorem of operators. *Nature Machine Intelligence*, 3(3):218–229, 2021.
- [21] L. Lu, X. Meng, S. Cai, Z. Mao, S. Goswami, Z. Zhang, and G. E. Karniadakis. A comprehensive and fair comparison of two neural operators (with practical extensions) based on fair data. Computer Methods in Applied Mechanics and Engineering, 393:114778, 2022. ISSN 0045-7825. DOI: https://doi.org/10.1016/j.cma.2022.114778. URL https://www.sciencedirect.com/science/article/pii/S0045782522001207.
- [22] T. Ma. STATS214 / CS229M: Machine Learning Theory, 2021. URL http://web.stanford.edu/class/stats214/.
- [23] T. Marwah, Z. C. Lipton, and A. Risteski. Parametric complexity bounds for approximating PDEs with neural networks. In A. Beygelzimer, Y. Dauphin, P. Liang, and J. W. Vaughan, editors, Advances in Neural Information Processing Systems, 2021. URL https://openreview.net/forum?id=HbViCqfbd7.
- [24] S. Mishra and R. Molinaro. Estimates on the generalization error of physics-informed neural networks for approximating a class of inverse problems for PDEs. *IMA Journal of Numerical Analysis*, 06 2021. ISSN 0272-4979. DOI: 10.1093/imanum/drab032. URL https://doi.org/10.1093/imanum/drab032. drab032.
- [25] A. Mukherjee. A study of the mathematics of deep learning. *CoRR*, abs/2104.14033, 2021. URL https://arxiv.org/abs/2104.14033.
- [26] V. Nagarajan. Explaining generalization in deep learning: progress and fundamental limits. *CoRR*, abs/2110.08922, 2021. URL https://arxiv.org/abs/2110.08922.
- [27] B. Neyshabur, R. Tomioka, and N. Srebro. Norm-based capacity control in neural networks. In P. Grünwald, E. Hazan, and S. Kale, editors, Proceedings of The 28th Conference on Learning Theory, COLT 2015, Paris, France, July 3-6, 2015, volume 40 of JMLR Workshop and Conference Proceedings, pages 1376-1401. JMLR.org, 2015. URL http://proceedings.mlr.press/v40/Neys habur15.html.
- [28] B. Neyshabur, S. Bhojanapalli, and N. Srebro. A pac-bayesian approach to spectrally-normalized margin bounds for neural networks. In 6th International Conference on Learning Representations, ICLR 2018, Vancouver, BC, Canada, April 30 May 3, 2018, Conference Track Proceedings. Open-Review.net, 2018. URL https://openreview.net/forum?id=Skz_WfbCZ.

- [29] B. Neyshabur, Z. Li, S. Bhojanapalli, Y. LeCun, and N. Srebro. The role of over-parametrization in generalization of neural networks. In *International Conference on Learning Representations*, 2019. URL https://openreview.net/forum?id=BygfghAcYX.
- [30] M. Raissi, P. Perdikaris, and G. E. Karniadakis. Physics-informed neural networks: A deep learning framework for solving forward and inverse problems involving nonlinear partial differential equations. *Journal of Computational Physics*, 378:686–707, 2019.
- [31] J. Sirignano and K. Spiliopoulos. Dgm: A deep learning algorithm for solving partial differential equations. *Journal of Computational Physics*, 375:1339-1364, 2018. ISSN 0021-9991. DOI: https://doi.org/10.1016/j.jcp.2018.08.029. URL https://www.sciencedirect.com/science/article/pii/S0021999118305527.
- [32] A. Tewari and S. Kakade. Lecture 9, Topics in Artificial Intelligence (Learning Theory) Spring 2008, 2008. URL https://home.ttic.edu/~tewari/lectures/lecture9.pdf.
- [33] S. Wang, H. Wang, and P. Perdikaris. Improved architectures and training algorithms for deep operator networks, 2021. URL https://arxiv.org/abs/2110.01654.
- [34] S. Wang, H. Wang, and P. Perdikaris. Learning the solution operator of parametric partial differential equations with physics-informed DeepONets. *Science Advances*, 7(40):eabi8605, 2021. DOI: 10.1126/sciadv.abi8605.
- [35] E. Weinan and B. Yu. The deep ritz method: A deep learning-based numerical algorithm for solving variational problems. *Communications in Mathematics and Statistics*, 6(1), 2018.
- [36] Z. Yang, Y. Yu, C. You, J. Steinhardt, and Y. Ma. Rethinking bias-variance trade-off for generalization of neural networks. In Proceedings of the 37th International Conference on Machine Learning, ICML 2020, 13-18 July 2020, Virtual Event, volume 119 of Proceedings of Machine Learning Research, pages 10767-10777. PMLR, 2020. URL http://proceedings.mlr.press/v119/yang20j.html.

Appendix

A (1,1) – DeepONets

Definition 20 (A (1,1)-(Linear) **DeepONet Class)** Given a constant C > 0, we define a (1,1) linear DeepONet class as follows,

$$\mathcal{F}_{(1,1)}(\mathcal{C}) \coloneqq \left\{ \mathbb{R}^{d_1} \times \mathbb{R}^{d_2} \ni (\mathbf{x}_1, \mathbf{x}_2) \to \langle \mathbf{W}_B(\mathbf{x}_1), \mathbf{W}_T(\mathbf{x}_2) \rangle \in \mathbb{R} \right.$$

$$\left. \left\| \mathbf{W}_B \in \mathbb{R}^{p \times d_1}, \mathbf{W}_T \in \mathbb{R}^{p \times d_2}, \left\| \sum_{j=1}^p \tilde{\mathbf{W}}_j \right\|_2 \le \mathcal{C}, \text{where, } (\tilde{W}_j)_{(k_1, k_2)} = W_{B, j, k_1} W_{T, j, k_2} \right\}.$$

Here, $\tilde{\mathbf{W}}_j$ are $d_1 \cdot d_2$ dimensional vectors indexed by 2-tuples as, $(\tilde{W}_j)_{(k_1,k_2)} = W_{B,j,k_1}W_{T,j,k_2}$ for $k_1 = 1, \ldots, d_1$ and $k_2 = 1, \ldots, d_2$.

Theorem 21 (Rademacher complexity for (1,1)-(Linear) **DeepONets)** In the setup of Definition 10, suppose the given data-set \mathcal{D} of size m is $\{(\mathbf{x}_B^i, \mathbf{x}_T^i) \in \mathbb{R}^{d_1} \times \mathbb{R}^{d_2} \mid i = 1, \dots, m\}$. Then corresponding to the i^{th} sample in \mathcal{D} , we define $\tilde{\mathbf{x}}^i \in \mathbb{R}^{d_1 \times d_2}$ s.t $(\tilde{\mathbf{x}}^i)_{k_1,k_2} = (\mathbf{x}_B^i)_{k_1}(\mathbf{x}_T^i)_{k_2}$.

Then we have for the function class given in Definition 20,

$$\hat{\mathcal{R}}_m\Big(\mathcal{F}_{(1,1)}(\mathcal{C})\Big) \leq \frac{\mathcal{C}}{m} \sqrt{\sum_{i=1}^m \left\|\tilde{\mathbf{x}}^i\right\|_F^2}.$$

Further suppose that the $d_1 \times d_2$ matrices $\tilde{\mathbf{x}}^i$ are samples from a matrix distribution P, such that $\mathbb{E}_P\left[\|\tilde{\mathbf{x}}\|_F^2\right] \leq M^2$. Then, the average Rademacher complexity is bounded as

$$\mathcal{R}\left(\mathcal{F}_{(1,1)}(\mathcal{C})\right) \leq \frac{\mathcal{C}M}{\sqrt{m}}.$$

The above has been proven in Section B.1. The essential gist of the above theorem is to identify a capacity measure for the (1,1)-(Linear) DeepONet as,

$$\sqrt{\sum_{k_1=1}^{d_1}\sum_{k_2=1}^{d_2}\left(\sum_{j=1}^p \mathbf{W}_{B,j,k_1}\mathbf{W}_{T,j,k_2}\right)^2} = \left\|\mathbf{W}_B^{\intercal}\mathbf{W}_T\right\|_F$$

For better intuition, we state here a consequence of Theorem 21 when the function class is defined in terms of bounds given on the branch and the trunk network matrices individually.

Lemma 22 (2-norm bounded weight matrices) Given M as in Theorem 21, suppose that spectral norms of weights \mathbf{W}_B , \mathbf{W}_T are bounded as $\|\mathbf{W}_B\| \leq M_B$, $\|\mathbf{W}_T\| \leq M_T$. In this case, the average Rademacher complexity is bounded as follows

$$\hat{\mathcal{R}} \leq \frac{pM_B M_T M}{\sqrt{m}}.$$

The above has been proven in Section B.2.

B Proofs of Rademacher Complexity for Shallow Operator Nets

B.1 Proof of Theorem 21

Proof In the context of the class as given in Definition 20 we define a vector $\mathbf{w} := \sum_{j=1}^{p} \tilde{\mathbf{W}}_{j}$. Then invoking the definition of empirical Rademacher complexity for this class $\mathcal{F}_{(1,1)}(\mathcal{C})$ for the given data-set we get,

$$\hat{\mathcal{R}}_{m}\left(\mathcal{F}_{(1,1)}(\mathcal{C})\right) = \mathbb{E}_{\epsilon} \left(\sup_{\|\mathbf{w}\|_{2} \leq \mathcal{C}} \frac{1}{m} \sum_{i=1}^{m} \epsilon_{i} \left\langle \mathbf{W}_{B} \mathbf{x}_{B}^{i}, \mathbf{W}_{T} \mathbf{x}_{T}^{i} \right\rangle \right)$$

$$= \mathbb{E}_{\epsilon} \left(\sup_{\|\mathbf{w}\|_{2} \leq \mathcal{C}} \frac{1}{m} \sum_{i=1}^{m} \sum_{j=1}^{p} \epsilon_{i} \left(\sum_{k_{1}=1}^{d_{1}} \mathbf{W}_{B,j,k_{1}} x_{B,k_{1}}^{i}\right) \left(\sum_{k_{2}=1}^{d_{2}} \mathbf{W}_{T,j,k_{2}} x_{T,k_{2}}^{i}\right) \right)$$

$$= \mathbb{E}_{\epsilon} \left(\sup_{\|\mathbf{w}\|_{2} \leq \mathcal{C}} \frac{1}{m} \sum_{i=1}^{m} \sum_{j=1}^{p} \sum_{k_{1}=1}^{d_{1}} \sum_{k_{2}=1}^{d_{2}} \epsilon_{i} \left(\mathbf{W}_{B,j,k_{1}} x_{B,k_{1}}^{i}\right) \left(\mathbf{W}_{T,j,k_{2}} x_{T,k_{2}}^{i}\right) \right)$$

$$= \mathbb{E}_{\epsilon} \left(\sup_{\|\mathbf{w}\|_{2} \leq \mathcal{C}} \sum_{j=1}^{p} \sum_{k_{1}=1}^{d_{1}} \sum_{k_{2}=1}^{d_{2}} \mathbf{W}_{B,j,k_{1}} \mathbf{W}_{T,j,k_{2}} \left(\frac{1}{m} \sum_{i=1}^{m} \epsilon_{i} \left(x_{B,k_{1}}^{i}\right) \left(x_{T,k_{2}}^{i}\right) \right) \right)$$

$$= \mathbb{E}_{\epsilon} \left(\sup_{\|\mathbf{w}\|_{2} \leq \mathcal{C}} \sum_{j=1}^{p} \sum_{k_{1}=1}^{d_{1}} \sum_{k_{2}=1}^{d_{2}} \tilde{\mathbf{W}}_{j,(k_{1},k_{2})} \left(\frac{1}{m} \sum_{i=1}^{m} \epsilon_{i} \tilde{x}_{k_{1},k_{2}}^{i}\right) \right).$$

The above triple-sum is an inner-product of two vectors in $\mathbb{R}^{d_1 \times d_2}$. Therefore,

$$= \mathbb{E}_{\epsilon} \left(\sup_{\|\mathbf{w}\|_{2} \leq \mathcal{C}} \left\langle \sum_{j=1}^{p} \tilde{\mathbf{W}}_{j}, \frac{1}{m} \sum_{i=1}^{m} \epsilon_{i} \tilde{\mathbf{x}}^{i} \right\rangle \right)$$

$$= \frac{1}{m} \mathbb{E}_{\epsilon} \left(\sup_{\|\mathbf{w}\|_{2} \leq \mathcal{C}} \left\langle \mathbf{w}, \sum_{i=1}^{m} \epsilon_{i} \tilde{\mathbf{x}}^{i} \right\rangle \right)$$

$$\leq \frac{\mathcal{C}}{m} \sqrt{\sum_{i=1}^{m} \left\| \tilde{\mathbf{x}}^{i} \right\|_{2}^{2}} \qquad (Eq. (5.23) in [22]).$$

$$\implies \mathcal{R}\Big(\mathcal{F}_{(1,1)}(\mathcal{C})\Big) = \mathbb{E}[\hat{\mathcal{R}}_m\Big(\mathcal{F}_{(1,1)}(\mathcal{C})\Big)]$$

$$= \frac{\mathcal{C}}{m}\mathbb{E}\left[\sqrt{\sum_{i=1}^{m} \left\|\tilde{\mathbf{x}}^i\right\|_F^2}\right] \leq \frac{\mathcal{C}}{m}\sqrt{\sum_{i=1}^{m} \mathbb{E}\left[\left\|\tilde{\mathbf{x}}^i\right\|_F^2\right]} \qquad \text{(Jensen's Inequality)}$$

$$\leq \frac{\mathcal{C}M}{\sqrt{m}}.$$

B.2 Proof of Lemma 22

Proof Due to constraints on spectral norms of $\mathbf{W}_B, \mathbf{W}_T$ as defined earlier, we have

$$\begin{split} \left\| \sum_{j=1}^{p} \tilde{\mathbf{W}}_{j} \right\|_{2} &\leq \sum_{j=1}^{p} \left\| \tilde{\mathbf{W}}_{j} \right\|_{2} & \text{(Triangle Inequality)} \\ &\leq \sum_{j=1}^{p} M_{B} M_{T} \leq p M_{B} M_{T}, \end{split}$$

where we have used,

$$\|\tilde{\mathbf{W}}_{j}\|_{2} = \sqrt{\sum_{k_{1}=1}^{d_{1}} \sum_{k_{2}=1}^{d_{2}} W_{B,j,k_{1}}^{2} W_{T,j,k_{2}}^{2}} = \|\mathbf{W}_{B,j}\|_{2} \|\mathbf{W}_{T,j}\|_{2} \le \|\mathbf{W}_{B}\| \|\mathbf{W}_{T}\| \le M_{B}M_{T},$$

where $\|\mathbf{A}\|$ denotes spectral norm of \mathbf{A} . Hence, invoking the definition of \mathcal{C} the upper bound on average Rademacher complexity becomes,

 $\mathcal{R} \leq \frac{pM_B M_T M}{\sqrt{m}}.$

B.3 Proof of Theorem 13

Proof For ease of notation we define a capacity variable as,

$$w_{2,1} = w(\mathbf{B}_2, \mathbf{B}_1, \mathbf{T}_2, \mathbf{T}_1) \coloneqq \max_{\substack{k_1 = 1, \dots, b_1 \\ k_2 = 1, \dots, t_1}} \|\mathbf{B}_{1,k_1}\| \|\mathbf{T}_{1,k_2}\| \left(\sum_{j=1}^p (\mathbf{B}_{2,j} \mathbf{T}_{2,j}^{\mathsf{T}})_{k_1,k_2}\right).$$

Hence we have,

$$\begin{split} &\hat{\mathcal{R}}_{m}\Big(\mathcal{F}_{(2,2)-(\text{ReLU})}(\mathcal{C})\Big) \\ &= \mathbb{E}_{\epsilon} \left[\sup_{w_{2,1} \leq c} \frac{1}{m} \sum_{i=1}^{m} \epsilon_{i} \left\langle \mathbf{B}_{2} \operatorname{ReLU}\left(\mathbf{B}_{1} \mathbf{x}_{B}^{i}\right), \mathbf{T}_{2} \operatorname{ReLU}\left(\mathbf{T}_{1} \mathbf{x}_{T}^{i}\right) \right\rangle \right] \\ &= \mathbb{E}_{\epsilon} \left[\sup_{w_{2,1} \leq c} \frac{1}{m} \sum_{i=1}^{m} \epsilon_{i} \sum_{j=1}^{p} \left\langle \mathbf{B}_{2,j}, \operatorname{ReLU}\left(\mathbf{B}_{1} \mathbf{x}_{B}^{i}\right) \right\rangle \cdot \left\langle \mathbf{T}_{2,j}, \operatorname{ReLU}\left(\mathbf{T}_{1} \mathbf{x}_{T}^{i}\right) \right\rangle \right] \\ &= \mathbb{E}_{\epsilon} \left[\sup_{w_{2,1} \leq c} \frac{1}{m} \sum_{i=1}^{m} \sum_{j=1}^{p} \sum_{k_{1}=1}^{b_{1}} \sum_{k_{2}=1}^{t_{1}} \epsilon_{i} B_{2,j,k_{1}} \operatorname{ReLU}\left(\mathbf{B}_{1,k_{1}}^{\mathsf{T}} \mathbf{x}_{B}^{i}\right) T_{2,j,k_{2}} \operatorname{ReLU}\left(\mathbf{T}_{1,k_{2}}^{\mathsf{T}} \mathbf{x}_{T}^{i}\right) \right] \\ &= \mathbb{E}_{\epsilon} \left[\sup_{w_{2,1} \leq c} \frac{1}{m} \sum_{j=1}^{p} \sum_{k_{1}=1}^{b_{1}} \sum_{k_{2}=1}^{t_{1}} B_{2,j,k_{1}} T_{2,j,k_{2}} \sum_{i=1}^{m} \epsilon_{i} \operatorname{ReLU}\left(\mathbf{B}_{1,k_{1}}^{\mathsf{T}} \mathbf{x}_{B}^{i}\right) \operatorname{ReLU}\left(\mathbf{T}_{1,k_{2}}^{\mathsf{T}} \mathbf{x}_{T}^{i}\right) \right] \\ &= \mathbb{E}_{\epsilon} \left[\sup_{w_{2,1} \leq c} \frac{1}{m} \sum_{j=1}^{p} \sum_{k_{1}=1}^{b_{1}} \sum_{k_{2}=1}^{t_{1}} \left(\mathbf{B}_{2,j} \mathbf{T}_{2,j}^{\mathsf{T}}\right)_{k_{1},k_{2}} \sum_{i=1}^{m} \epsilon_{i} \operatorname{ReLU}\left(\mathbf{B}_{1,k_{1}}^{\mathsf{T}} \mathbf{x}_{B}^{i}\right) \operatorname{ReLU}\left(\mathbf{T}_{1,k_{2}}^{\mathsf{T}} \mathbf{x}_{T}^{i}\right) \right]. \end{split}$$

We introduce two vectors in $\mathbb{R}^{b_1 \times t_1}$ **v**, **w** which are indexed by a tuple (k_1, k_2) such that

$$v_{k_{1},k_{2}} = \|\mathbf{B}_{1,k_{1}}\| \|\mathbf{T}_{1,k_{2}}\| \left(\sum_{j=1}^{p} (\mathbf{B}_{2,j} \mathbf{T}_{2,j}^{\mathsf{T}})_{k_{1},k_{2}} \right).$$

$$w_{k_{1},k_{2}} = \frac{1}{m} \cdot \sum_{i=1}^{m} \epsilon_{i} \operatorname{ReLU} \left(\frac{\mathbf{B}_{1,k_{1}}^{\mathsf{T}}}{\|\mathbf{B}_{1,k_{1}}\|} \mathbf{x}_{B}^{i} \right) \operatorname{ReLU} \left(\frac{\mathbf{T}_{1,k_{2}}^{\mathsf{T}}}{\|\mathbf{T}_{1,k_{2}}\|} \mathbf{x}_{T}^{i} \right).$$

Then we can rewrite the above expression as,

$$\hat{\mathcal{R}}_{m}\Big(\mathcal{F}_{(2,2)-(\text{ReLU})}(\mathcal{C})\Big) = \mathbb{E}_{\epsilon} \left[\sup_{w_{2,1} \leq \mathcal{C}} \frac{1}{m} \sum_{k_{1}=1}^{d_{1}} \sum_{k_{2}=1}^{d_{2}} v_{k_{1},k_{2}} w_{k_{1},k_{2}} \right].$$

Note that for any $\alpha, \beta \in \mathbb{R}^k$ we have, $(\alpha, \beta) \leq (\max_{p \in 1, ..., k} |\beta_p|) \cdot \sum_{j=1}^k |\alpha_j|$. Hence,

$$\hat{\mathcal{R}}_{m}\Big(\mathcal{F}_{(2,2)-(\text{ReLU})}(\mathcal{C})\Big) \leq \mathbb{E}_{\epsilon} \left[\sup_{w_{2,1} \leq \mathcal{C}} \left(\max_{k_{1},k_{2}} |v_{k_{1},k_{2}}| \right) \sum_{k_{1}=1}^{b_{1}} \sum_{k_{2}=1}^{t_{1}} |w_{k_{1},k_{2}}| \right].$$

We recall the definition of \mathcal{C} to upperbound the above as,

$$\begin{split} \hat{\mathcal{R}}_{m}\Big(\mathcal{F}_{(2,2)-(\text{ReLU})}(\mathcal{C})\Big) &\leq \mathcal{C}\mathbb{E}_{\epsilon}\left[\sup_{w_{2,1} \leq \mathcal{C}} \sum_{k_{1}=1}^{d_{1}} \sum_{k_{2}=1}^{d_{2}} |w_{k_{1},k_{2}}|\right] \\ &\leq \frac{\mathcal{C}}{m} \,\mathbb{E}_{\epsilon}\left[\sum_{k_{1}=1}^{b_{1}} \sum_{k_{2}=1}^{t_{1}} \sup_{w_{2,1} \leq \mathcal{C}} \left|\sum_{i=1}^{m} \epsilon_{i} \operatorname{ReLU}\left(\frac{\mathbf{B}_{1,k_{1}}^{\mathsf{T}}}{\|\mathbf{B}_{1,k_{1}}\|} \mathbf{x}_{B}^{i}\right) \operatorname{ReLU}\left(\frac{\mathbf{T}_{1,k_{2}}^{\mathsf{T}}}{\|\mathbf{T}_{1,k_{2}}\|} \mathbf{x}_{T}^{i}\right)\right]\right]. \end{split}$$

Now, we can define $(\mathbf{v}_1, \mathbf{v}_2) \in S^{b_1-1} \times S^{t_1-1}$ s.t

$$\mathbf{v}_1 \coloneqq \frac{\mathbf{B}_{1,k_1}}{\|\mathbf{B}_{1,k_1}\|}, \quad \mathbf{v}_2 \coloneqq \frac{\mathbf{T}_{1,k_2}}{\|\mathbf{T}_{1,k_2}\|}.$$

And hence we have,

$$\begin{split} &\left| \sum_{i=1}^{m} \epsilon_{i} \operatorname{ReLU}\left(\mathbf{v}_{1}^{\intercal} \mathbf{x}_{B}^{i}\right) \operatorname{ReLU}\left(\mathbf{v}_{2}^{\intercal} \mathbf{x}_{T}^{i}\right) \right| \\ &\leq \sqrt{\sum_{i=1}^{m} \left(\epsilon_{i} \operatorname{ReLU}\left(\mathbf{v}_{1}^{\intercal} \mathbf{x}_{B}^{i}\right)\right)^{2}} \cdot \sqrt{\sum_{i=1}^{m} \left(\operatorname{ReLU}\left(\mathbf{v}_{2}^{\intercal} \mathbf{x}_{T}^{i}\right)\right)^{2}} \leq \sqrt{\sum_{i=1}^{m} \left\|\mathbf{x}_{B}^{i}\right\|^{2}} \cdot \sqrt{\sum_{i=1}^{m} \left\|\mathbf{x}_{T}^{i}\right\|^{2}}. \end{split}$$

Putting the above back into the original expression we have.

$$\hat{\mathcal{R}}_{m}\left(\mathcal{F}_{(2,2)-(\text{ReLU})}(\mathcal{C})\right) \leq \frac{\mathcal{C}}{m} b_{1} t_{1}\left(\sqrt{\sum_{i=1}^{m} \left\|\mathbf{x}_{B}^{i}\right\|^{2}} \cdot \sqrt{\sum_{i=1}^{m} \left\|\mathbf{x}_{T}^{i}\right\|^{2}}\right).$$

C Proof of Lemma 17

Proof We recall from the setup of Definition 11 that b_{-1} and t_{-1} are the number of rows of the matrices \mathbf{B}_{q_B-1} and \mathbf{T}_{q_T-1} - and b_{-2} and t_{-2} are the output dimensions of the nets \mathbf{f}_B' and \mathbf{f}_T' . From Definition 11 we further recall the definitions of the 2 sets of matrices $\mathcal{W}(\mathcal{C}_{q_B,q_T,q_B-1,q_T-1})$ and $\mathcal{W}_{\text{rest}}$ and we simplify the required empirical Rademacher complexity as,

$$m\hat{\mathcal{R}}_{m} = \mathbb{E}\left[\sup_{\left(\mathcal{W}(\mathcal{C}_{q_{B},q_{T},q_{B}-1,q_{T}-1}),\mathcal{W}_{\text{rest}}\right)} \sum_{i=1}^{m} \epsilon_{i} \left\langle \mathbf{f}_{B}(\mathbf{x}_{B,i}), \mathbf{f}_{T}(\mathbf{x}_{T,i}) \right\rangle \right]$$

$$= \mathbb{E}_{\epsilon}\left[\sup_{\left(\mathcal{W}(\mathcal{C}_{q_{B},q_{T},q_{B}-1,q_{T}-1}),\mathcal{W}_{\text{rest}}\right)} \left(\sum_{i=1}^{m} \epsilon_{i} \left\langle \mathbf{B}_{q_{B}} \sigma_{1} \left(\mathbf{B}_{q_{B}-1} \mathbf{f}_{B}'(\mathbf{x}_{B,i})\right), \mathbf{T}_{q_{T}} \sigma_{2} \left(\mathbf{T}_{q_{T}-1} \mathbf{f}_{T}'(\mathbf{x}_{T,i})\right) \right\rangle \right)\right]. \quad (13)$$

Here, we have substituted the definitions of the functions \mathbf{f}'_B and the \mathbf{f}'_T as given in the statement of Lemma 17. To ease notation now we define the following vectors,

$$\mathbf{b}_{i} \coloneqq \sigma_{1}\left(\mathbf{B}_{q_{B}-1}\mathbf{f}_{B}'(\mathbf{x}_{B,i})\right), \ \mathbf{t}_{i} \coloneqq \sigma_{2}\left(\mathbf{T}_{q_{T}-1}\mathbf{f}_{T}'(\mathbf{x}_{T,i})\right).$$

Then we can rewrite the above as,

$$\begin{split} m\hat{\mathcal{R}}_{m} &= \mathbb{E}_{\epsilon} \left[\sup_{\left(\mathcal{W}\left(\mathcal{C}_{q_{B},q_{T},q_{B}-1,q_{T}-1}\right),\mathcal{W}_{\mathrm{rest}}\right)} \left(\sum_{i=1}^{m} \epsilon_{i} \left\langle \mathbf{B}_{q_{B}} \mathbf{b}_{i}, \mathbf{T}_{q_{T}} \mathbf{t}_{i} \right\rangle \right) \right] \\ &= \mathbb{E}_{\epsilon} \left[\sup_{\left(\mathcal{W}\left(\mathcal{C}_{q_{B},q_{T},q_{B}-1,q_{T}-1}\right),\mathcal{W}_{\mathrm{rest}}\right)} \left(\sum_{i=1}^{m} \epsilon_{i} \sum_{j=1}^{p} \mathbf{B}_{q_{B},j}^{\mathsf{T}} \mathbf{b}_{i} \cdot \mathbf{T}_{q_{T},j}^{\mathsf{T}} \mathbf{t}_{i} \right) \right] \\ &= \mathbb{E}_{\epsilon} \left[\sup_{\left(\mathcal{W}\left(\mathcal{C}_{q_{B},q_{T},q_{B}-1,q_{T}-1}\right),\mathcal{W}_{\mathrm{rest}}\right)} \left(\sum_{i=1}^{m} \epsilon_{i} \sum_{j=1}^{p} \left(\sum_{k_{1}=1}^{b-1} B_{q_{B},j,k_{1}} b_{i,k_{1}} \right) \cdot \left(\sum_{k_{2}=1}^{t-1} T_{q_{T},j,k_{2}} t_{i,k_{2}} \right) \right) \right] \\ &= \mathbb{E}_{\epsilon} \left[\sup_{\left(\mathcal{W}\left(\mathcal{C}_{q_{B},q_{T},q_{B}-1,q_{T}-1}\right),\mathcal{W}_{\mathrm{rest}}\right)} \left(\sum_{j=1}^{p} \sum_{k_{1}=1}^{b-1} \sum_{k_{2}=1}^{t-1} B_{q_{B},j,k_{1}} T_{q_{T},j,k_{2}} \sum_{i=1}^{m} \epsilon_{i} b_{i,k_{1}} \cdot t_{i,k_{2}} \right) \right] \\ &= \mathbb{E}_{\epsilon} \left[\sup_{\left(\mathcal{W}\left(\mathcal{C}_{q_{B},q_{T},q_{B}-1,q_{T}-1}\right),\mathcal{W}_{\mathrm{rest}}\right)} \left(\sum_{j=1}^{p} \sum_{k_{1}=1}^{b-1} \sum_{k_{2}=1}^{t-1} \left(\mathbf{B}_{q_{B},j} \mathbf{T}_{q_{T},j}^{\mathsf{T}}\right)_{k_{1},k_{2}} \right. \\ &\left. \cdot \sum_{i=1}^{m} \epsilon_{i} \sigma_{1} \left(\mathbf{B}_{q_{B}-1} \mathbf{f}_{B}^{\prime}(\mathbf{x}_{B,i})\right)_{k_{1}} \cdot \sigma_{2} \left(\mathbf{T}_{q_{T}-1} \mathbf{f}_{T}^{\prime}(\mathbf{x}_{T,i})\right)_{k_{2}} \right) \right] \\ &= \mathbb{E}_{\epsilon} \left[\sup_{\mathcal{W}\left(\mathcal{C}_{q_{B},q_{T},q_{B}-1,q_{T}-1}\right)} \left(\sum_{j=1}^{p} \sum_{k_{1}=1}^{b-1} \sum_{k_{2}=1}^{t-1} \left(\mathbf{B}_{q_{B},j} \mathbf{T}_{q_{T},j}^{\mathsf{T}}\right)_{k_{1},k_{2}} \cdot \left(\mathbf{T}_{q_{T}-1} \mathbf{f}_{T}^{\prime}(\mathbf{x}_{T,i})\right)_{k_{2}} \right) \right] \\ &= \mathbb{E}_{\epsilon} \left[\sup_{\mathcal{W}\left(\mathcal{C}_{q_{B},q_{T},q_{B}-1,q_{T}-1}\right)} \left(\sum_{j=1}^{p} \sum_{k_{1}=1}^{b-1} \sum_{k_{2}=1}^{t-1} \left(\mathbf{B}_{q_{B},j} \mathbf{T}_{q_{T},j}^{\mathsf{T}}\right)_{k_{1},k_{2}} \cdot \left(\mathbf{B}_{q_{B}-1,k_{1}} \right) \cdot \left(\mathbf{T}_{q_{T}-1,k_{2}}^{\mathsf{T}} \mathbf{f}_{T}^{\prime}(\mathbf{x}_{T,i})\right)_{k_{2}} \right) \right] \\ &= \mathbb{E}_{\epsilon} \left[\sup_{\mathcal{W}\left(\mathcal{C}_{q_{B},q_{T},q_{B}-1,q_{T}-1}\right)} \left(\sum_{j=1}^{p} \sum_{k_{1}=1}^{b-1} \sum_{k_{2}=1}^{b-1} \left(\mathbf{B}_{q_{B},j} \mathbf{T}_{q_{T},j}^{\mathsf{T}}\right)_{k_{1},k_{2}} \cdot \left(\mathbf{B}_{q_{B}-1,k_{1}} \right) \cdot \left(\mathbf{T}_{q_{T}-1,k_{2}}^{\mathsf{T}} \mathbf{f}_{T}^{\prime}(\mathbf{x}_{T,i})\right) \right) \right] \\ &= \mathbb{E}_{\epsilon} \left[\sup_{\mathcal{W}\left(\mathcal{C}_{q_{B},q_{T},q_{B}-1,q_{T}-1}\right)} \left(\sum_{j=1}^{p} \sum_{k_{1}=1}^{b-1} \sum_{k_{2}=1}^{b-1} \left(\mathbf{B}_{q_{B},j} \mathbf{T}_$$

In the last line above we have invoked the positive homogeneity of σ_1 and σ_2 .

Now assume 2 vectors \mathbf{v} and \mathbf{w} of dimensions $b_{-1} - 1 \times t_{-1} - 1$ s.t they are indexed by the tuple (k_1, k_2) for $k_1 \in \{1, \ldots, -1 + b_{-1}\}$ and $k_2 \in \{1, \ldots, -1 + t_{-1}\}$ as follows,

$$v_{(k_1,k_2)} \coloneqq \|\mathbf{B}_{q_B-1,k_1}\| \cdot \|\mathbf{T}_{q_T-1,k_2}\| \cdot \sum_{j=1}^{p} (\mathbf{B}_{q_B,j} \mathbf{T}_{q_T,j}^{\mathsf{T}})_{k_1,k_2},$$

$$w_{(k_1,k_2)} \coloneqq \sum_{i=1}^{m} \epsilon_i \sigma_1 \left(\hat{\mathbf{B}}_{q_B-1,k_1}^{\mathsf{T}} \mathbf{f}_B'(\mathbf{x}_{B,i}) \right) \cdot \sigma_2 \left(\hat{\mathbf{T}}_{q_T-1,k_2}^{\mathsf{T}} \mathbf{f}_T'(\mathbf{x}_{T,i}) \right).$$

Then we have,

$$m\hat{\mathcal{R}}_{m} = \mathbb{E}_{\epsilon} \left[\sup_{\left(\mathcal{W}(\mathcal{C}_{q_{B}, q_{T}, q_{B}-1, q_{T}-1}), \mathcal{W}_{\text{rest}} \right)} \left(\sum_{k_{1}=1}^{b_{-1}} \sum_{k_{2}=1}^{t_{-1}} v_{(k_{1}, k_{2})} \cdot w_{(k_{1}, k_{2})} \right) \right].$$

Note that for any $\alpha, \beta \in \mathbb{R}^k$ we have, $\langle \alpha, \beta \rangle \leq (\max_{p \in 1, \dots, k} |\beta_p|) \cdot \sum_{j=1}^k |\alpha_j|$

$$\leq \mathbb{E}_{\epsilon} \left[\sup_{\left(\mathcal{W}(C_{q_B, q_T, q_{B}-1, q_{T}-1}), \mathcal{W}_{\text{rest}} \right)} \left(\sum_{k_1=1}^{b_{-1}} \sum_{k_2=1}^{t_{-1}} \left| v_{(k_1, k_2)} \right| \cdot \max_{\substack{k_1=1, \dots, b_{-1}-1 \\ k_2=1, \dots, t_{-1}-1}} \left| w_{(k_1, k_2)} \right| \right) \right].$$

Now we define a set of weights $W_{\text{except-outer}}$ which is an union of all possible weight matrices that are allowed in W_{rest} and all possible choices of $(\mathbf{B}_{q_B-1}, \mathbf{T}_{q_T-1})$ that are allowed in the set $W(\mathcal{C}_{q_B,q_T,q_B-1,q_T-1})$.

Then we recall the definition of $W(C_{q_B,q_T,q_{B}-1,q_{T}-1})$, to get,

$$m\hat{\mathcal{R}}_{m} \leq \mathbb{E}_{\epsilon} \left[\sup_{\mathcal{W}_{\text{except-outer}}} \left(\mathcal{C}_{q_{B},q_{T},q_{B}-1,q_{T}-1} \cdot \max_{\substack{k_{1}=1,\dots,b_{-1}-1\\k_{2}=1,\dots,t_{-1}-1}} \left| \sum_{i=1}^{m} \epsilon_{i} \sigma_{1} \left(\hat{\mathbf{B}}_{q_{B}-1,k_{1}}^{\intercal} \mathbf{f}_{B}'(\mathbf{x}_{B,i}) \right) \cdot \sigma_{2} \left(\hat{\mathbf{T}}_{q_{T}-1,k_{2}}^{\intercal} \mathbf{f}_{T}'(\mathbf{x}_{T,i}) \right) \right| \right) \right].$$

Note that any pair of row directions in the pair of matrices $(\mathbf{B}_{q_B-1}, \mathbf{T}_{q_T-1})$ is in $\in S^{b-2-1} \times S^{t-2-1}$. So a sup over the set of \mathbf{B}_{q_B-1} & \mathbf{T}_{q_T-1} that is allowed by the constraint of C_{q_B,q_T,q_B-1,q_T-1} and a subsequent max over the pairs of row directions can be upperbounded by a single sup over $S^{b-2-1} \times S^{t-2-1}$. Thus we have,

$$m\hat{\mathcal{R}}_{m} \leq \mathbb{E}_{\epsilon} \left[\sup_{\substack{(\mathbf{v}, \mathbf{w}) \in S^{b-2-1} \times S^{t-2-1} \\ \mathcal{W}_{cont}}} \left(\mathcal{C}_{q_{B}, q_{T}, q_{B}-1, q_{T}-1} \cdot \left| \sum_{i=1}^{m} \epsilon_{i} \sigma_{1} \left(\mathbf{v}^{\mathsf{T}} \mathbf{f}_{B}'(\mathbf{x}_{B, i}) \right) \cdot \sigma_{2} \left(\mathbf{w}^{\mathsf{T}} \mathbf{f}_{T}'(\mathbf{x}_{T, i}) \right) \right| \right) \right].$$

Invoking Lemma 28, we have

$$m\hat{\mathcal{R}}_{m} \leq \mathbb{E}_{\epsilon} \left[2 \times \sup_{\left(\mathbf{v},\mathbf{w}\right) \in S^{b-2^{-1}} \times S^{t-2^{-1}}} \left(\mathcal{C}_{q_{B},q_{T},q_{B}-1,q_{T}-1} \cdot \sum_{i=1}^{m} \epsilon_{i} \sigma_{1} \left(\mathbf{v}^{\mathsf{T}} \mathbf{f}_{B}'(\mathbf{x}_{B,i}) \right) \cdot \sigma_{2} \left(\mathbf{w}^{\mathsf{T}} \mathbf{f}_{T}'(\mathbf{x}_{T,i}) \right) \right) \right].$$

Now we invoke Lemma 14 using the assumption about σ_1 and σ_2 to get,

$$m\hat{\mathcal{R}}_{m} \leq 2L \cdot \mathcal{C}_{q_{B},q_{T},q_{B}-1,q_{T}-1} \cdot \mathbb{E}_{\epsilon} \left[\sup_{\substack{(\mathbf{v},\mathbf{w}) \in S^{b-2-1} \times S^{t-2-1} \\ \mathcal{W}_{\text{nest}}}} \left(\sum_{i=1}^{m} \epsilon_{i} \mathbf{v}^{\mathsf{T}} \mathbf{f}_{B}'(\mathbf{x}_{B,i}) \cdot \mathbf{w}^{\mathsf{T}} \mathbf{f}_{T}'(\mathbf{x}_{T,i}) \right) \right].$$

The above inequality is exactly what we set out to prove.

D Proof of Lemma 18

Proof We start with the expression in the R.H.S. of the Lemma 17 and simplify it similarly as was done in its proof in the previous appendix. Denote $S_2 = S^{b-2-1} \times S^{t-2-1}$.

$$\mathbb{E}_{\epsilon} \left[\sup_{\substack{(\mathbf{v}, \mathbf{w}) \in \mathcal{S}_{2} \\ \mathcal{W}_{rest}}} \left(\sum_{i=1}^{m} \epsilon_{i} \mathbf{v}^{\mathsf{T}} \mathbf{f}_{B}'(\mathbf{x}_{B,i}) \cdot \mathbf{w}^{\mathsf{T}} \mathbf{f}_{T}'(\mathbf{x}_{T,i}) \right) \right]$$

$$= \mathbb{E}_{\epsilon} \left[\sup_{\substack{(\mathbf{v}, \mathbf{w}) \in \mathcal{S}_{2} \\ \mathcal{W}_{rest}}} \left(\sum_{i=1}^{m} \epsilon_{i} \mathbf{v}^{\mathsf{T}} \sigma_{1} (\mathbf{B}_{q_{B}-2} \mathbf{f}_{B}''(\mathbf{x}_{B,i})) \cdot \mathbf{w}^{\mathsf{T}} \sigma_{2} (\mathbf{T}_{q_{T}-2} \mathbf{f}_{T}''(\mathbf{x}_{T,i})) \right) \right]$$

$$= \mathbb{E}_{\epsilon} \left[\sup_{\substack{(\mathbf{v}, \mathbf{w}) \in \mathcal{S}_{2} \\ \mathcal{W}}} \sum_{j_{1}=1}^{b-2} \sum_{j_{2}=1}^{t-2} (\mathbf{v} \mathbf{w}^{\mathsf{T}})_{j_{1}, j_{2}} \|\mathbf{B}_{q_{B}-2, j_{1}}\| \|\mathbf{T}_{q_{T}-2, j_{2}}\| \cdot \sum_{i=1}^{m} \epsilon_{i} \sigma_{1} (\hat{\mathbf{B}}_{q_{B}-2, j_{1}}^{\mathsf{T}} \mathbf{f}_{B}''(\mathbf{x}_{B,i})) \sigma_{2} (\hat{\mathbf{T}}_{q_{T}-2, j_{2}}^{\mathsf{T}} \mathbf{f}_{T}''(\mathbf{x}_{T,i})) \right].$$

Define

$$\begin{split} \tilde{v}_{j_1,j_2} &\coloneqq (\mathbf{v}\mathbf{w}^{\mathsf{T}})_{j_1,j_2} \|\mathbf{B}_{q_B-2,j_1}\| \|\mathbf{T}_{q_T-2,j_2}\| \\ \tilde{w}_{j_1,j_2} &\coloneqq \sum_{i=1}^m \epsilon_i \sigma_1(\hat{\mathbf{B}}_{q_B-2,j_1}^{\mathsf{T}} \mathbf{f}_B''(\mathbf{x}_{B,i})) \sigma_2(\hat{\mathbf{T}}_{q_T-2,j_2}^{\mathsf{T}} \mathbf{f}_T''(\mathbf{x}_{T,i})). \end{split}$$

Thus we get,

$$\mathbb{E}_{\epsilon} \left[\sup_{\substack{(\mathbf{v}, \mathbf{w}) \in \mathcal{S}_{2} \\ \mathcal{W}_{rest}}} \left(\sum_{i=1}^{m} \epsilon_{i} \mathbf{v}^{\mathsf{T}} \mathbf{f}_{B}'(\mathbf{x}_{B,i}) \cdot \mathbf{w}^{\mathsf{T}} \mathbf{f}_{T}'(\mathbf{x}_{T,i}) \right) \right] = \mathbb{E}_{\epsilon} \left[\sup_{\substack{(\mathbf{v}, \mathbf{w}) \in \mathcal{S}_{2} \\ \mathcal{W}_{rest}}} \sum_{j_{1}=1}^{b_{-2}} \sum_{j_{2}=1}^{t_{-2}} \tilde{v}_{j_{1}, j_{2}} \tilde{w}_{j_{1}, j_{2}} \right]$$

$$\leq \mathbb{E}_{\epsilon} \left[\sup_{\substack{(\mathbf{v}, \mathbf{w}) \in \mathcal{S}_{2} \\ \mathcal{W}_{rest}}} \left(\sum_{j_{1}=1}^{b_{-2}} \sum_{j_{2}=1}^{t_{-2}} |\tilde{v}_{j_{1}, j_{2}}| \cdot \max_{j_{1}, j_{2}} |\tilde{w}_{j_{1}, j_{2}}| \right) \right].$$

For ease of notation we define the set $W_{-2,-2}$ as the set of matrices allowed in W_{rest} but with those with indices $q_B - 2$ and $q_T - 2$ additionally satisfying the constraint in Equation 12 in the statement of the

lemma.

$$\begin{split} & \mathbb{E}_{\epsilon} \left[\sup_{(\mathbf{v}, \mathbf{w}) \in \mathcal{S}_{2}} \left(\sum_{i=1}^{m} \epsilon_{i} \mathbf{v}^{\mathsf{T}} \mathbf{f}_{B}'(\mathbf{x}_{B,i}) \cdot \mathbf{w}^{\mathsf{T}} \mathbf{f}_{T}'(\mathbf{x}_{T,i}) \right) \right] \\ & \leq \mathcal{C}_{-2,-2} \mathbb{E}_{\epsilon} \left[\sup_{\mathcal{W}_{-2,-2}} \left(\max_{\substack{j_{1}=1, \dots, q_{B}-2 \\ j_{2}=1, \dots, q_{T}-2}} |\tilde{w}_{j_{1},j_{2}}| \right) \right] \\ & \leq \mathcal{C}_{-2,-2} \mathbb{E}_{\epsilon} \left(\sup_{\mathcal{W}_{-2,-2}} \left(\max_{\substack{j_{1}=1, \dots, q_{B}-2 \\ j_{2}=1, \dots, q_{T}-2}} \left| \sum_{i=1}^{m} \epsilon_{i} \sigma_{1} (\hat{\mathbf{B}}_{q_{B}-2,j_{1}}^{\mathsf{T}} \mathbf{f}_{B}''(\mathbf{x}_{B,i})) \sigma_{2} (\hat{\mathbf{T}}_{q_{T}-2,j_{2}}^{\mathsf{T}} \mathbf{f}_{T}''(\mathbf{x}_{T,i})) \right| \right) \right). \end{split}$$

We note that any pair of row directions in the pair of matrices $(\mathbf{B}_{q_B-2}, \mathbf{T}_{q_T-2})$ is $\in S^{b_{-3}-1} \times S^{t_{-3}-1}$. So a sup over the set of $\mathbf{B}_{q_B-2} \& \mathbf{T}_{q_T-2}$ that is allowed by the constraint of $\mathcal{C}_{-2,-2}$ and a subsequent max over the pairs of row directions can be upperbounded by a single sup over $S^{b_{-3}-1} \times S^{t_{-3}-1}$.

We recall the definition of \mathcal{W}'_{rest} given in the lemma to conclude, $(S_3 = S^{b_{-3}-1} \times S^{t_{-3}-1})$

$$\mathbb{E}_{\epsilon} \left[\sup_{\substack{(\mathbf{v}, \mathbf{w}) \in \mathcal{S}_{2} \\ \mathcal{W}_{\text{rest}}}} \left(\sum_{i=1}^{m} \epsilon_{i} \mathbf{v}^{\mathsf{T}} \mathbf{f}_{B}'(\mathbf{x}_{B,i}) \cdot \mathbf{w}^{\mathsf{T}} \mathbf{f}_{T}'(\mathbf{x}_{T,i}) \right) \right]$$

$$\leq C_{-2,-2} \mathbb{E}_{\epsilon} \left(\sup_{\substack{(\mathbf{v}, \mathbf{w}) \in \mathcal{S}_{3} \\ \mathcal{W}_{rest}'}} \left| \sum_{i=1}^{m} \epsilon_{i} \sigma_{1}(\mathbf{v}^{\mathsf{T}} \mathbf{f}_{B}''(\mathbf{x}_{B,i})) \sigma_{2}(\mathbf{w}^{\mathsf{T}} \mathbf{f}_{T}''(\mathbf{x}_{T,i})) \right| \right).$$

Invoking Lemma 28,

$$\leq 2C_{-2,-2}\mathbb{E}_{\epsilon} \left(\sup_{\substack{(\mathbf{v},\mathbf{w}) \in \mathcal{S}_{3} \\ \mathcal{W}'_{rest}}} \sum_{i=1}^{m} \epsilon_{i} \sigma_{1}(\mathbf{v}^{\mathsf{T}} \mathbf{f}''_{B}(\mathbf{x}_{B,i})) \sigma_{2}(\mathbf{w}^{\mathsf{T}} \mathbf{f}''_{T}(\mathbf{x}_{T,i})) \right)$$

$$\leq 2LC_{-2,-2}\mathbb{E}_{\epsilon} \left(\sup_{\substack{(\mathbf{v},\mathbf{w}) \in \mathcal{S}_{3} \\ \mathcal{W}'_{rest}}} \sum_{i=1}^{m} \epsilon_{i}(\mathbf{v}^{\mathsf{T}} \mathbf{f}''_{B}(\mathbf{x}_{B,i}))(\mathbf{w}^{\mathsf{T}} \mathbf{f}''_{T}(\mathbf{x}_{T,i})) \right).$$

In the last step above we have invoked Lemma 14 using the assumption about σ_1 and σ_2 as in Lemma 17.

E Generalization Bound

We begin by recalling the following two standard results,

Theorem 23 (Theorem 2.4, [32]) Let $\phi_{\mathcal{F}}$ denote the loss class associated with \mathcal{F} . Then, we have, with probability at least $1-2\delta$,

$$L_{\phi}\left(\hat{f}_{\phi}^{*}\right) - \min_{f \in \mathcal{F}} L_{\phi}(f) \leq 2\mathcal{R}\left(\phi_{\mathcal{F}}\right) + 2\sqrt{\frac{\ln(1/\delta)}{2m}},$$

where $L_{\phi}(f)$ is the expected risk for a function $f \in \mathcal{F}$, m is the number of samples used in the empirical risk, the minimizer for empirical risk is given by $f_{\phi}^* \in \mathcal{F}$, and \mathcal{R} denotes the average Rademacher complexity of $\phi_{\mathcal{F}}$.

Lemma 24 (Lipschitz composition in Rademacher complexity) Assume that $\phi : \mathbb{R} \to \mathbb{R}$ is a L_{ϕ} Lipschitz continuous function, such that $|\phi(t) - \phi(s)| \leq L_{\phi}|t - s|$. Suppose $\mathcal{F} \subset \mathbb{R}^m$. Then,

$$\mathbb{E}_{\epsilon} \left[\sup_{f \in \mathcal{F}} \frac{1}{m} \sum_{i=1}^{m} \epsilon_{i} \phi(f_{i}) \right] \leq L_{\phi} \, \mathbb{E}_{\epsilon} \left[\sup_{f \in \mathcal{F}} \frac{1}{m} \sum_{i=1}^{m} \epsilon_{i} f_{i} \right].$$

Lemma 25 $(\mathcal{R}(\ell(\mathcal{H})))$ in terms of $\mathcal{R}(\mathcal{H})$ Given $\mathcal{Q} \subset \mathcal{H}$ as in Definition 9, if $\ell_2(\cdot)$ is R-Lipschitz,

$$\mathbb{E}_{\epsilon} \left[\sup_{h \in \mathcal{Q}} \frac{1}{m} \sum_{i=1}^{m} \epsilon_{i} \ell(h(f_{i}, \mathbf{x}_{T, i})) \right] = \mathcal{R}(\ell(\mathcal{H})) \leq R \cdot \mathcal{R}(\mathcal{H}) = R \cdot \mathbb{E}_{\epsilon} \left[\sup_{h \in \mathcal{Q}} \frac{1}{m} \sum_{i=1}^{m} \epsilon_{i} h(f_{i}, \mathbf{x}_{T, i}) \right].$$

Proof Any function $h \in \mathcal{H}$ maps from $\mathcal{F} \times \mathbb{R}^{d_2} \mapsto \mathbb{R}$. Hence, training data of size m for the DeepONet is of the form $\{(f_i, \mathbf{x}_{T,i}) \in \mathcal{F} \times \mathbb{R}^{d_2}, i = 1, 2, ..., m\}$.

Consider the set $\mathcal{K} = \{(h(f_1, \mathbf{x}_1), \dots, h(f_m, \mathbf{x}_m)) \mid h \in \mathcal{Q}\} \subset \mathbb{R}^m$. Then we can re-write,

$$\mathcal{R}(\ell(\mathcal{H})) = \mathbb{E}_{\epsilon} \left[\sup_{k \in \mathcal{K}} \frac{1}{m} \sum_{i=1}^{m} \epsilon_i \, \ell(k_i) \right].$$

Invoking that the Lipschitz constant for $\ell(\cdot)$ is R, and Lemma 24 we have,

$$\mathcal{R}(\ell(\mathcal{H})) \leq R \, \mathbb{E}_{\epsilon} \left[\sup_{k \in \mathcal{K}} \frac{1}{m} \sum_{i=1}^{m} \epsilon_i \, k_i \right],$$

which we recognize as the inequality we set out to prove

$$\mathcal{R}(\ell(\mathcal{H})) \leq R \cdot \mathcal{R}(\mathcal{H}).$$

Lemma 26 ($\mathcal{R}(\mathcal{H})$ in terms of $\mathcal{R}(\text{DeepONet})$) If \mathcal{W} is the set of allowed weights, and $\text{DeepONet}_{\mathbf{w}}$ be the DeepONet with weight \mathbf{w} then the following holds,

$$\mathcal{R}(\text{DeepONet}) \coloneqq \mathbb{E}_{\epsilon} \left[\sup_{\mathbf{w} \in \mathcal{W}} \frac{1}{m} \sum_{i=1}^{m} \epsilon_{i} \text{DeepONet}_{\mathbf{w}}(f_{i}, \mathbf{x}_{T, i}) \right] = \mathcal{R}(\mathcal{H}).$$

Proof If \mathcal{W} is the set of possible weights, then a corresponding \mathcal{Q} (and consequently \mathcal{K}) from Lemma 25 can be defined to be constrained only by the weights $\mathbf{w} \in \mathcal{W}$ of the DeepONet DeepONet, taking a sup over $h \in \mathcal{Q}$ and $\mathbf{w} \in \mathcal{W}$ are equivalent, as below:

$$\mathbb{E}_{\epsilon} \left[\sup_{h \in \mathcal{Q}} \frac{1}{m} \sum_{i=1}^{m} \epsilon_{i} h(f_{i}, \mathbf{x}_{T, i}) \right] = \mathbb{E}_{\epsilon} \left[\sup_{\mathbf{w} \in \mathcal{W}} \frac{1}{m} \sum_{i=1}^{m} \epsilon_{i} \left(\mathcal{G}(f_{i}, \mathbf{x}_{T, i}) - \text{DeepONet}_{\mathbf{w}}(f_{i}, \mathbf{x}_{T, i}) \right) \right],$$

where we have used $h(f_i, \mathbf{x}_{T,i}) = \mathcal{G}(f_i, \mathbf{x}_{T,i})$ – DeepONet_w $(f_i, \mathbf{x}_{T,i})$. We can write $\mathcal{R}(\mathcal{H})$ as

$$\mathcal{R}(\mathcal{H}) = \mathbb{E}_{\epsilon} \left[\sup_{\mathbf{w} \in \mathcal{W}} \frac{1}{m} \sum_{i=1}^{m} \epsilon_{i} \left(\mathcal{G}(f_{i}, \mathbf{x}_{T, i}) - \text{DeepONet}_{\mathbf{w}}(f_{i}, \mathbf{x}_{T, i}) \right) \right]$$

$$= \mathbb{E}_{\epsilon} \left[\sup_{\mathbf{w} \in \mathcal{W}} \left\{ \frac{1}{m} \sum_{i=1}^{m} \epsilon_{i} \mathcal{G}(f_{i}, \mathbf{x}_{T, i}) - \frac{1}{m} \sum_{i=1}^{m} \epsilon_{i} \text{DeepONet}_{\mathbf{w}}(f_{i}, \mathbf{x}_{T, i}) \right\} \right]$$

$$= \mathbb{E}_{\epsilon} \left[\frac{1}{m} \sum_{i=1}^{m} \epsilon_{i} \mathcal{G}(f_{i}, \mathbf{x}_{T, i}) + \sup_{\mathbf{w} \in \mathcal{W}} \frac{1}{m} \sum_{i=1}^{m} (-\epsilon_{i}) \text{DeepONet}_{\mathbf{w}}(f_{i}, \mathbf{x}_{T, i}) \right]$$

$$= 0 + \mathbb{E}_{\epsilon} \left[\sup_{\mathbf{w} \in \mathcal{W}} \frac{1}{m} \sum_{i=1}^{m} \epsilon_{i} \text{DeepONet}_{\mathbf{w}}(f_{i}, \mathbf{x}_{T, i}) \right]$$

$$= \mathcal{R}(\text{DeepONet}).$$

since, $\epsilon = -\epsilon$ in distribution.

Theorem 27 $(\mathcal{R}(\ell(\mathcal{H})))$ in terms of $\mathcal{R}(\text{DeepONet})$) Let $\ell(\cdot)$ be R-Lipschitz. Then,

$$\mathcal{R}(\ell(\mathcal{H})) \leq R \cdot \mathcal{R}(\text{DeepONet}).$$

E.1 Proof of Theorem 16

Proof With \mathbf{x}_B as in Remark 4.

$$\begin{aligned} |\text{DeepONet}(f, \mathbf{x}_T)| &= |\langle \mathbf{B}(\mathbf{x}_B), \mathbf{T}(\mathbf{x}_T) \rangle| \\ &\leq \|\mathbf{B}(\mathbf{x}_B)\|_2 \|\mathbf{T}(\mathbf{x}_T)\|_2 \\ &= \|\mathbf{B}_{q_B}(\sigma_1(\mathbf{B}_{q_{B}-1}(\dots \sigma_1(\mathbf{B}_1(\mathbf{x}_B))\dots)))\|_2 \|\mathbf{T}_{q_T}(\sigma_2(\mathbf{T}_{q_{T}-1}(\dots \sigma_2(\mathbf{T}_1(\mathbf{x}_T))\dots)))\|_2 \\ &\leq L_1^{q_B} L_2^{q_T} \|\mathbf{x}_B\|_2 \|\mathbf{x}_T\|_2 \left(\prod_{i=1}^{q_B} \|\mathbf{B}_i\|\right) \left(\prod_{i=1}^{q_T} \|\mathbf{T}_i\|\right). \end{aligned}$$

Invoking Assumption 1 and $h \in \mathcal{H}$ from Definition 9, in addition to the above bound, we get,

$$h(f, \mathbf{x}_T) \leq \sup_{f \in \mathcal{F}} \left(C_f (1 + \|f\|_{L^2})^{\kappa} \right) + L_1^{q_B} L_2^{q_T} \|\mathbf{x}_B\|_2 \|\mathbf{x}_T\|_2 \left(\prod_{i=1}^{q_B} \|\mathbf{B}_i\| \right) \left(\prod_{i=1}^{q_T} \|\mathbf{T}_i\| \right) = B.$$

Using Theorem 23 with the fact that $\ell_2(h(f, \mathbf{x}_T)) \in [0, \frac{1}{2}B^2]$, we have,

$$\mathbb{E}_{(\mathbf{f},\mathbf{x}_T)\sim\mathcal{D}}\left[\ell_2(h^*(f,\mathbf{x}_T)) - \min_{h\in\mathcal{H}}\left[\mathbb{E}_{(f,\mathbf{x}_T)\sim\mathcal{D}}\left[\ell_2(h(f,\mathbf{x}_T))\right]\right] \leq 2B\cdot\mathcal{R} + B^2\sqrt{\frac{\ln(1/\delta)}{2m}}.$$

Note that \mathcal{R} has an additional factor of B, due to Lipschitz composition of $\ell_2(\cdot)$ with \mathcal{H} .

F Contraction Lemmas

Lemma 28 (From [22]) Let $\epsilon \sim \text{Uniform}(\{1,-1\}^m)$ and suppose we have functions \mathbf{f}_{θ} and f'_{θ} parameterized by θ s.t, $(\mathbb{R}^k)^m \ni \mathbf{x} \mapsto \mathbf{f}_{\theta}(\mathbf{x}) = (f'_{\theta}(x_1), \dots, f'_{\theta}(x_m)) \in \mathbb{R}^m$. Suppose that for any $\epsilon \in \{\pm 1\}^m$, $\sup_{\theta} \langle \epsilon, \mathbf{f}_{\theta}(x) \rangle \geq 0$. Then,

$$\mathbb{E}_{\epsilon} \left[\sup_{\theta} |\langle \epsilon, \mathbf{f}_{\theta}(\mathbf{x}) \rangle| \right] \leq 2 \mathbb{E}_{\epsilon} \left[\sup_{\theta} \langle \epsilon, \mathbf{f}_{\theta}(\mathbf{x}) \rangle \right].$$

Proof Letting ϕ be the ReLU function, the lemma's assumption implies that $\sup_{\theta} \phi(\langle \epsilon, f_{\theta}(x) \rangle) = \sup_{\theta} \langle \epsilon, f_{\theta}(x) \rangle$ for any $\epsilon \in \{\pm 1\}^n$. Observing that $|z| = \phi(z) + \phi(-z)$,

$$\sup_{\theta} |\langle \epsilon, \mathbf{f}_{\theta}(x) \rangle| = \sup_{\theta} \left[\phi \left(\langle \epsilon, \mathbf{f}_{\theta}(x) \rangle \right) + \phi \left(\langle -\epsilon, \mathbf{f}_{\theta}(x) \rangle \right) \right]$$

$$\leq \sup_{\theta} \phi \left(\langle \epsilon, \mathbf{f}_{\theta}(x) \rangle \right) + \sup_{\theta} \phi \left(\langle -\epsilon, \mathbf{f}_{\theta}(x) \rangle \right)$$

$$= \sup_{\theta} \langle \epsilon, \mathbf{f}_{\theta}(x) \rangle + \sup_{\theta} \langle -\epsilon, \mathbf{f}_{\theta}(x) \rangle.$$

Taking the expectation over ϵ (and noting that ϵ and $-\epsilon$ have the same distribution) we get the desired conclusion.

F.1 Proof of Lemma 14

Proof

$$\mathcal{R}_{\mathrm{m}}(\phi_F \circ \mathcal{F} \cdot \phi_G \circ \mathcal{G}) = \frac{1}{m} \mathbb{E}_{\epsilon \sim U(\{-1,1\}^m)} \left[\sup_{(f,g) \in \mathcal{F} \times \mathcal{G}} \sum_{i=1}^m \epsilon_i \cdot \phi_F(f(\mathbf{x}_i)) \phi_G(g(\mathbf{y}_i)) \right].$$

We explicitly open up the expectation on ϵ_1 to get,

$$= \frac{1}{2m} \cdot \mathbb{E}_{(\epsilon_{2},...,\epsilon_{m}) \sim U(\{-1,1\}^{m-1})} \left[\sup_{f,g \in \mathcal{F} \times \mathcal{G}} \left(\phi_{F}(f(\mathbf{x}_{1})) \phi_{G}(g(\mathbf{y}_{1})) + \sum_{i=2}^{m} \epsilon_{i} \cdot \phi_{F}(f(\mathbf{x}_{i})) \phi_{G}(g(\mathbf{y}_{i})) \right) + \sum_{i=2}^{m} \epsilon_{i} \cdot \phi_{F}(f'(\mathbf{x}_{i})) \phi_{G}(g'(\mathbf{y}_{1})) + \sum_{i=2}^{m} \epsilon_{i} \cdot \phi_{F}(f'(\mathbf{x}_{i})) \phi_{G}(g'(\mathbf{y}_{i})) \right) \right]$$

$$\leq \frac{1}{2m} \cdot \mathbb{E}_{(\epsilon_{2},...,\epsilon_{m})} \left[\sup_{((f,g),(f',g')) \in (\mathcal{F} \times \mathcal{G})^{2}} \left(L \cdot |\phi_{F}(f(\mathbf{x}_{1})) \phi_{G}(g(\mathbf{y}_{1})) - \phi_{F}(f'(\mathbf{x}_{1})) \phi_{G}(g'(\mathbf{y}_{1})) | + \sum_{i=2}^{m} \epsilon_{i} \left(\phi_{F}(f(\mathbf{x}_{i})) \phi_{G}(g(\mathbf{y}_{i})) + \phi_{F}(f'(\mathbf{x}_{i})) \phi_{G}(g'(\mathbf{y}_{i})) \right) \right]$$

By invoking the the assumption in Equation 10 we get,

$$\leq \frac{1}{2m} \cdot \mathbb{E}_{(\epsilon_2, \dots, \epsilon_m)} \left[\sup_{((f,g), (f',g')) \in (\mathcal{F} \times \mathcal{G})^2} \left(L \cdot |f(\mathbf{x}_1)g(\mathbf{y}_1) - f'(\mathbf{x}_1)g'(\mathbf{y}_1)| + \sum_{i=2}^m \epsilon_i \left(\phi_F(f(\mathbf{x}_i)) \phi_G(g(\mathbf{y}_i)) + \phi_F(f'(\mathbf{x}_i)) \phi_G(g'(\mathbf{y}_i)) \right) \right]$$

Now we invoke the fact that inside a supremum, an absolute value function is redundant, since anyway the higher combination will get picked. Thus we can rearrange the above to get,

$$\leq \frac{1}{2m} \cdot \mathbb{E}_{(\epsilon_{2},...,\epsilon_{m})} \left[\sup_{(f,g)\in\mathcal{F}\times\mathcal{G}} \left(L \cdot f(\mathbf{x}_{1})g(\mathbf{y}_{1}) \right) + \sum_{i=2}^{m} \epsilon_{i} \cdot \phi_{F}(f(\mathbf{x}_{i}))\phi_{G}(g(\mathbf{y}_{i})) \right) \\
+ \sup_{(f',g')\in\mathcal{F}\times\mathcal{G}} \left(-L \cdot f'(\mathbf{x}_{1})g'(\mathbf{y}_{1}) + \sum_{i=2}^{m} \epsilon_{i} \cdot \phi_{F}(f'(\mathbf{x}_{i}))\phi_{G}(g'(\mathbf{y}_{i})) \right) \right] \\
\leq \frac{1}{m} \cdot \mathbb{E}_{\epsilon} \left[\sup_{(f,g)\in\mathcal{F}\times\mathcal{G}} \left(\epsilon_{1} \cdot L \cdot f(\mathbf{x}_{1})g(\mathbf{y}_{1}) + \sum_{i=2}^{m} \epsilon_{i} \cdot \phi(f(\mathbf{x}_{i}))\phi_{G}(g(\mathbf{y}_{i})) \right) \right]$$

Re-iterating the above argument for $i = 2, 3, \ldots, m$ we get,

$$= \frac{1}{m} \mathbb{E}_{\epsilon \sim U(\{-1,1\}^m)} \sup_{(f,g) \in \mathcal{F} \times \mathcal{G}} \left[\sum_{i=1}^m \epsilon_i L \cdot f(\mathbf{x}_i) g(\mathbf{y}_i) \right]$$
$$= L \mathcal{R}_{\mathbf{m}}(\mathcal{F} \cdot \mathcal{G}).$$

F.2 A Contraction Lemma with Biased Absolute Functions

Lemma 29 Let \mathcal{F} and \mathcal{G} be set of functions valued in \mathbb{R} closed under negation. Given points $\{\mathbf{x}_i \mid i=1,\ldots,m\}$, $\{\mathbf{y}_i \mid i=1,\ldots,m\}$ in the domain of the functions in we have the following inequality of Rademacher complexities - where both the sides are being evaluated on this same set of points,

$$\mathcal{R}_{\mathrm{m}}(|\mathcal{F} + B_1| \cdot |\mathcal{G} + B_2|) \leq \mathcal{R}_{\mathrm{m}}(\mathcal{F} \cdot \mathcal{G}) + |B_2|\mathcal{R}_{\mathrm{m}}(\mathcal{F}) + |B_1|\mathcal{R}_{\mathrm{m}}(\mathcal{G})$$

Proof Denote the uniform distribution over $\{-1,1\}^m$ by $U(\{-1,1\}^m)$. Then,

$$\mathcal{R}_{\mathrm{m}}(|\mathcal{F} + B_1| \cdot |\mathcal{G} + B_2|) \leq \frac{1}{m} \mathbb{E}_{\epsilon \sim U(\{-1,1\}^m)} \sup_{f,g \in \mathcal{F} \times \mathcal{G}} \left[\sum_{i=1}^m \epsilon_i \cdot |f(\mathbf{x}_i) + B_1| |g(\mathbf{y}_i) + B_2| \right].$$

We explicitly open up the expectation on ϵ_1 to get,

$$= \frac{1}{2m} \cdot \mathbb{E}_{(\epsilon_{2},...,\epsilon_{m}) \sim U(\{-1,1\}^{m-1})} \left[\sup_{f,g \in \mathcal{F} \times \mathcal{G}} \left(|f(\mathbf{x}_{1}) + B_{1}||g(\mathbf{y}_{1}) + B_{2}| + \sum_{i=2}^{m} \epsilon_{i} \cdot \phi_{F}(f(\mathbf{x}_{i}))\phi_{G}(g(\mathbf{y}_{i})) \right) + \sup_{f',g' \in \mathcal{F} \times \mathcal{G}} \left(-|f'(\mathbf{x}_{1}) + B_{1}||g'(\mathbf{y}_{1}) + B_{2}| + \sum_{i=2}^{m} \epsilon_{i} \cdot \phi_{F}(f'(\mathbf{x}_{i}))\phi_{G}(g'(\mathbf{y}_{i})) \right) \right]$$

$$\leq \frac{1}{2m} \cdot \mathbb{E}_{(\epsilon_{2},...,\epsilon_{m})} \left[\sup_{f,f',g,g' \in \mathcal{F} \times \mathcal{F} \times \mathcal{G} \times \mathcal{G}} \left(|f(\mathbf{x}_{1})g(\mathbf{y}_{1}) - f'(\mathbf{x}_{1})g'(\mathbf{y}_{1})| + |B_{2}||f(\mathbf{x}_{1}) - f'(\mathbf{x}_{1})| + |B_{1}||g(\mathbf{y}_{1}) - g'(\mathbf{y}_{1})| + \sum_{i=2}^{m} \epsilon_{i} \left(\phi_{F}(f(\mathbf{x}_{i}))\phi_{G}(g(\mathbf{y}_{i})) + \phi_{F}(f'(\mathbf{x}_{i}))\phi_{G}(g'(\mathbf{y}_{i})) \right) \right] \right]$$

$$= \frac{1}{2m} \cdot \mathbb{E}_{(\epsilon_{2},...,\epsilon_{m})} \left[\sup_{f,g \in \mathcal{F} \times \mathcal{G}} \left(f(\mathbf{x}_{1})g(\mathbf{y}_{1}) + |B_{2}|f(\mathbf{x}_{1}) + |B_{1}|g(\mathbf{y}_{1}) + \sum_{i=2}^{m} \epsilon_{i} \cdot \phi_{F}(f(\mathbf{x}_{i}))\phi_{G}(g'(\mathbf{y}_{i})) \right) \right] + \sup_{f',g' \in \mathcal{F} \times \mathcal{G}} \left(-f'(\mathbf{x}_{1})g'(\mathbf{y}_{1}) - |B_{2}|f'(\mathbf{x}_{1}) - |B_{1}|g'(\mathbf{y}_{1}) + \sum_{i=2}^{m} \epsilon_{i} \cdot \phi_{F}(f'(\mathbf{x}_{i}))\phi_{G}(g'(\mathbf{y}_{i})) \right) \right]$$

$$\leq \frac{1}{m} \cdot \mathbb{E}_{\epsilon} \left[\sup_{f,g \in \mathcal{F} \times \mathcal{G}} \left(\epsilon_{1} \cdot \left(f(\mathbf{x}_{1})g(\mathbf{y}_{1}) + |B_{2}|f(\mathbf{x}_{1}) + |B_{1}|g(\mathbf{y}_{1}) \right) + \sum_{i=2}^{m} \epsilon_{i} \cdot \phi_{F}(f(\mathbf{x}_{i}))\phi_{G}(g'(\mathbf{y}_{i})) \right) \right]$$

Iterating this, we get

$$\leq \frac{1}{m} \mathbb{E}_{\epsilon \sim U(\{-1,1\}^m)} \sup_{f,g \in \mathcal{F} \times \mathcal{G}} \left[\sum_{i=1}^m \epsilon_i \cdot (f(\mathbf{x}_i)g(\mathbf{y}_i) + |B_2|f(\mathbf{x}_i) + |B_1|g(\mathbf{y}_i) \right]$$

$$\leq \mathcal{R}_{\mathbf{m}}(\mathcal{F} \cdot \mathcal{G}) + |B_2|\mathcal{R}_{\mathbf{m}}(\mathcal{F}) + |B_1|\mathcal{R}_{\mathbf{m}}(\mathcal{G}).$$

In the second inequality we have used

$$\left| |f(\mathbf{x}_1) + B_1| \cdot |g(\mathbf{y}_1) + B_2| - |f'(\mathbf{x}_1) + B_1| \cdot |g'(\mathbf{y}_1) + B_2| \right| \le \left| (f(\mathbf{x}_1) + B_1)(g(\mathbf{y}_1) + B_2) - (f'(\mathbf{x}_1) + B_1)(g'(\mathbf{y}_1) + B_2) \right|.$$

In the next line we could open up the $|\cdot|$ since the supremum would be positive and \mathcal{F} and \mathcal{G} are closed under multiplication by -1.

G Converting ReLU-DeepONets to abs-DeepONets

Firstly, we recall that the map $\mathbb{R}^q \ni \mathbf{x} \mapsto \mathbf{x} \in \mathbb{R}^q$ is an exact depth 2 ReLU net – one which passes every coordinate of the input through the ReLU net $\mathbb{R} \ni z \mapsto \max\{0, z\} - \max\{0, -z\} \in \mathbb{R}$. Using this, given any ReLU DeepONet, we can symmetrize the depths between the branch and the trunk by attaching the required number of identity computing layers at the input to the shorter side.

Secondly, it is natural to assume for typical DeepONet experiments that the set of all possible input data is bounded. Combining this with the assumption of boundedness of the allowed matrices, we conclude that $\exists \mathcal{B} > 0$ s.t the input to any ReLU gate in any DeepONet in the given class is bounded by \mathcal{B} . Now we observe that $\forall |z| \leq \mathcal{B}$, we can rewrite the map $z \mapsto \max\{0,z\}$ as, $z \mapsto \frac{1}{2}|z| + \frac{1}{4}|z + \mathcal{B}| - \frac{1}{4}|z - \mathcal{B}|$.

Hence, doing the above replacement at every gate we can rewrite any ReLU DeepONet (without biases) as a DeepONet using only absolute value activations but with biases – and computing the same function on the same bounded domain, at the cost of increasing the size of the branch and the trunk net by a factor of 3. Thirdly, a similar result as in Lemma 14 continues to hold for this setup as given in Lemma 29. Note that, Lemma 29 bounds the Rademacher complexity of a DeepONet class with the branch and trunk depths (k, k) by a linear sum of that of (k-1, k-1) and sum of a (k, 0) and a (0, k) DeepONet. While the first term can be recursed on again (using identical techniques as used to prove Theorem 15), for the last two one can invoke Theorem 1 of [10].

Thus, we observe that following the same arguments as in the proof for Theorem 15 we can derive an analogous bound for arbitrary ReLU DeepONet classes - but with twice the depth of the DeepONet

number of extra terms in the R.H.S. These extra terms would come in pairs – each pair consisting of a Rademacher complexity bound on a standard net, one from the branch side and one from the trunk side and of decreasing depths.

H Choosing $C_{-k,-k}$

Defining $\begin{bmatrix} X_{j_1,j_2} \end{bmatrix}_{\substack{j_1=1,\ldots,b_{-k}\\k_1=1,\ldots,t_{-k}}} \coloneqq \mathbf{X} \in \mathbb{R}^{b_{-k} \times t_{-k}} \text{ s.t } X_{j_1,j_2} \coloneqq \|\mathbf{B}_{n-k,j_1}\| \cdot \|\mathbf{T}_{n-k,j_2}\|, \text{ we can simplify as follows the}$

LHS of the inequality defining $C_{-k,-k}$ in equation 4,

$$\sup_{(\mathbf{v}, \mathbf{w}) \in S^{-1+b_{-k}} \times S^{-1+t_{-k}}} \sum_{j_1=1}^{b_{-k}} \sum_{j_2=1}^{t_{-k}} |(\mathbf{v} \mathbf{w}^{\mathsf{T}})_{j_1, j_2}| \|\mathbf{B}_{n-k, j_1}\| \|\mathbf{T}_{n-k, j_2}\|$$

$$= \sup_{(\mathbf{v}, \mathbf{w}) \in S^{-1+b_{-k}} \times S^{-1+t_{-k}}} \sum_{j_1=1}^{b_{-k}} \sum_{j_2=1}^{t_{-k}} |v_{j_1}| \|\mathbf{B}_{n-k, j_1}\| \|\mathbf{T}_{n-k, j_2}\| |w_{j_2}|$$

$$= \sup_{(\mathbf{v}, \mathbf{w}) \in S^{-1+b_{-k}} \times S^{-1+t_{-k}}} \sum_{j_1=1}^{b_{-k}} \sum_{j_2=1}^{t_{-k}} |v_{j_1}| \cdot \mathbf{X}_{j_1, j_2} \cdot |w_{j_2}|$$

$$= \sup_{(\mathbf{v}, \mathbf{w}) \in S^{-1+b_{-k}} \times S^{-1+t_{-k}}} \tilde{\mathbf{v}}^{\mathsf{T}} \mathbf{X} \tilde{\mathbf{w}}$$

In the last line above we defined vectors $\tilde{\mathbf{v}}$ and $\tilde{\mathbf{w}}$ of appropriate dimensions s.t $\tilde{v}_i = |v_i|$ and $\tilde{w}_i = |w_i|$. Then note that, $\tilde{\mathbf{v}}^{\top} \mathbf{X} \tilde{\mathbf{w}} \leq ||\tilde{\mathbf{v}}||_2 ||\mathbf{X} \tilde{\mathbf{w}}||_2$ (Cauchy Schwarz) $\leq ||\tilde{\mathbf{v}}||_2 ||\mathbf{X}|| ||\tilde{\mathbf{w}}||_2$

In above $\|\mathbf{X}\|$ is the spectral norm of \mathbf{X} . Therefore we have the following upperbound,

$$\sup_{\mathbf{v},\mathbf{w}} \|\tilde{\mathbf{v}}\|_2 \|\mathbf{X}\| \|\tilde{\mathbf{w}}\|_2 \leq \sup_{\mathbf{v}} \|\tilde{\mathbf{v}}\|_2 \cdot \sup_{\mathbf{w}} \|\tilde{\mathbf{w}}\|_2 \cdot \|\mathbf{X}\|$$

Since $\|\tilde{\mathbf{v}}\| = \|\mathbf{v}\| \le 1$ and $\|\tilde{\mathbf{w}}\| = \|\mathbf{w}\| \le 1$ we have,

$$\sup_{(\mathbf{v}, \mathbf{w}) \in S^{-1+b_{-k}} \times S^{-1+t_{-k}}} \tilde{\mathbf{v}}^{\top} \mathbf{X} \tilde{\mathbf{w}} \leq \left\| \left[\| \mathbf{B}_{n-k, j_1} \| \cdot \| \mathbf{T}_{n-k, j_2} \| \right]_{\substack{j_1 = 1, \dots, b_{-k} \\ k_1 = 1, \dots, t_{-k}}} \right\|$$

Thus it follows that the RHS of the above inequality gives an intuitive candidate for the quantity $C_{-k,-k}$.